

Supplement to,

Trends in crop phenology and terrestrial fluxes in major Indian agroecosystems using Community Land Model (CLM5)

K. N. Reddy and S. Baidya Roy

Text S1: Carbon fluxes in CLM5 for C3 crops

Photosynthesis in C₃ plants is based on the model of Farquhar et al. (1980). In its simplest form, leaf net photosynthesis after accounting for dark respiration (R_d) is:

$$A_n = \min(A_c, A_j, A_p) - R_d$$

$$A_c = \frac{V_{c\max}(c_i - \Gamma)}{c_i + K_c(1 + o_i/K_o)}$$

$$A_j = \frac{J_x(c_i - \Gamma)}{4c_i + 8\Gamma}$$

$$A_p = 3T_p$$

where A_c ($\mu\text{ mol CO}_2\text{ m}^{-2}\text{ s}^{-1}$) is the RuBP carboxylase (Rubisco) limited rate of carboxylation, A_j ($\mu\text{ mol CO}_2\text{ m}^{-2}\text{ s}^{-1}$) is the maximum rate of carboxylation allowed by the capacity to regenerate RuBP (i.e., the light-limited rate), and A_p ($\mu\text{ mol CO}_2\text{ m}^{-2}\text{ s}^{-1}$) is the product-limited rate of carboxylation for C₃ plants. In these equations, c_i is the internal leaf CO₂ partial pressure (Pa), and $o_i = 0.20P_{\text{atm}}$ is the O₂ partial pressure (Pa). K_c and K_o are the Michaelis-Menten constants (Pa) for CO₂ and O₂. Γ (Pa) is the CO₂ compensation point. $V_{c\max}$ is the maximum rate of carboxylation ($\mu\text{mol m}^{-2}\text{ s}^{-1}$), and J_x is the electron transport rate ($\mu\text{mol m}^{-2}\text{ s}^{-1}$). T_p is the triose phosphate utilization rate ($\mu\text{mol m}^{-2}\text{ s}^{-1}$), taken as $T_p = 0.167V_{c\max}$ so that $A_p = 0.5V_{c\max}$ for C₃ plants (as in Collatz et al. 1992). For more details on the calculation of $V_{c\max}$, J_x , T_p , K_c , K_o , and Γ at various temperatures, please refer to CLM5 Tech Note: Chapter 9, and for information on respiration, refer to CLM5 Tech Note: Chapter 17 (Lawrence et al., 2018).




Figure S1: Comparison of monthly water and energy fluxes simulated by CLM5 CTRL run against the FLUXCOM data during the rice growing season, July to October. The area shown here is a region with more than 10% of the land area covered by rice. The second and fourth columns show the difference between CLM5 and FLUXCOM latent and sensible heat, respectively. The monthly mean fluxes over the period 1970-2014 are shown here. Pearson r in the plots is tested for significance, and values that are significant have an asterisk ‘*’.




Figure S2: Comparison of monthly water and energy fluxes simulated by CLM5 CTRL run against the FLUXCOM data during the wheat growing season, December to March. The area shown here is a region with more than 10% of the land area covered by wheat. The second and fourth columns show the difference between CLM5 and FLUXCOM latent and sensible heat, respectively. The monthly mean fluxes over the period 1970-2014 are shown here. Pearson r in the plots is tested for significance, and values that are significant have an asterisk “*”.




Figure S3: Spatial trend observed in (a) max. LAI, (b) Yield, and (c) Dry Matter of wheat-growing regions in the (i) CTRL simulation. (ii) to (v) shows the impact of each driver. Regions stippled implies the trend is statistically significant at $p < 0.05$.

Table S1: Comparing GPP simulated by CLM5 for rice and wheat against other studies

	Location	GPP (gC/m²)	AR/GPP	Reference
Rice	International Rice Research Institute, Philippines	1192 1464	0.67	Alberto et al., 2009
	Mase paddy site, Japan	1140	0.65	Saito et al., 2005
	Central Rice Research Institute, Cuttak, India	811	0.44	Bhattacharya et al., 2013
	Bangladesh Agricultural University Farmland, Dhaka, Bangladesh	1312 1574	0.58 0.52	Hossen et al., 2011
	North China Plain, China	987 966	0.56 0.56	Chen et al., 2015
	Indian Agricultural Research Institute, New Delhi, India	753	0.51	Kumar et al., 2021
	Rice growing region of India	1528.3 ± 561	0.53 ± 0.05	This study
	Selhausen test site, Germany (First Year)	1304	0.51	Schmidt et al., 2012
Wheat	Selhausen test site, Germany (Second Year)	1067	0.49	Schmidt et al., 2012
	North China Plain, China	1220 1135 859	0.52 0.54 0.53	Chen et al., 2015
	Saharanpur, India	1024	-	Patel et al., 2021
	Indian Agricultural Research Institute, New Delhi, India	888	0.34	Kumar et al., 2021
	Wheat growing regions of India	862.6 ± 438.6	0.42 ± 0.04	This study

References:

- Alberto, M. C. R., Wassmann, R., Hirano, T., Miyata, A., Kumar, A., Padre, A., and Amante, M.: CO₂/heat fluxes in rice fields: Comparative assessment of flooded and non-flooded fields in the Philippines, *Agr. Forest Meteorol.*, 149, 1737–1750, <https://doi.org/10.1016/j.agrformet.2009.06.003>, 2009.
- Bhattacharyya P., Neogi S., Roy K. S., Dash P. K., Tripathi R., and Rao K. S.: Net ecosystem CO₂ exchange and carbon cycling in tropical lowland flooded rice ecosystem. *Nutr. Cycl. Agroecosyst.*, 95:133–144, <https://doi.org/10.1007/s10705-013-9553-1>, 2013.
- Chen, C., Dan, L.I., Zhiqiu, G., Tang, J., Xiaofeng, G., Linlin, W., and Bingcheng, W.: Seasonal and interannual variations of carbon exchange over a rice-wheat rotation system on the north china plain, *Advan. Atmos. Sci.*, 32, 1365–1380, <https://doi.org/10.1007/s00376-015-4253-1>, 2015.
- Farquhar, G.D., von Caemmerer, S.V. and Berry, J.A.: A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *planta*, 149, pp.78-90, <https://doi.org/10.1007/BF00386231>, 1980.
- Hossen, M. S., Mano, M., Miyata, A., Baten, M. A., and Hiyama, T.: Seasonality of ecosystem respiration in a double-cropping paddy field in Bangladesh, *Biogeosciences Discuss.*, 8, 8693–8721, <https://doi.org/10.5194/bgd-8-8693-2011>, 2011.
- Kumar, A., Bhatia, A., Sehgal, V. K., Tomer, R., Jain, N., and Pathak, H.: Net Ecosystem Exchange of Carbon Dioxide in Rice-Spring Wheat System of Northwestern Indo-Gangetic Plains, *Land*, 10, 701, <https://doi.org/10.3390/land10070701>, 2021.
- Lawrence, D. M., Fisher, R., Koven, C., Oleson, K., Svenson, S., Vertenstein, M. (coordinating lead authors), Andre, B., Bonan, G., Ghimire, B., van Kampenhout, L., Kennedy, D., Kluzek, E., Knox, R., Lawrence, P., Li, F., Li, H., Lombardozzi, D., Lu, Y., Perket, J., Riley, W., Sacks, W., Shi, M., Wieder, W., Xu, C. (lead authors), Ali, A., Badger, A., Bisht, G., Broxton, P., Brunke, M., Buzan, J., Clark, M., Craig, T., Dahlin, K., Drewniak, B., Emmons, L., Fisher, J., Flanner, M., Gentine, P., Lenaerts, J., Levis, S., Leung, L. R., Lipscomb, W., Pelletier, J., Ricciuto, D., M., Sanderson, B., Shuman, J., Slater, A., Subin, Z., Tang, J., Tawfik, A., Thomas, Q., Tilmes, S., Vitt, F., and Zeng, X.: Technical Description of version 5.0 of the Community Land Model (CLM), Natl. Cent. Atmospheric Res. (NCAR), available at http://www.cesm.ucar.edu/models/cesm2/land/CLM50_Tech_Note.pdf (last access: 08 July 2024), 2018.
- Patel, N. R., Pokhariyal, S., Chauhan, P., and Dadhwal, V. K.: Dynamics of CO₂ fluxes and controlling environmental factors in sugarcane (C₄)–wheat (C₃) ecosystem of dry sub-humid region in India, *Int. J. Biometeorol.*, 65, 1069–1084, <https://doi.org/10.1007/s00484-021-02088-y>, 2021.
- Saito, M., Miyata, A., Nagai, H. and Yamada, T.: Seasonal variation of carbon dioxide exchange in rice paddy field in Japan. *Agric. For. Meteorol.*, 135(1-4), pp.93-109, <https://doi.org/10.1016/j.agrformet.2005.10.007>, 2005.
- Schmidt, M., Reichenau, T.G., Fiener, P. and Schneider, K.: The carbon budget of a winter wheat field: An eddy covariance analysis of seasonal and inter-annual variability, *Agric. For. Meteorol.*, 165, pp.114-126, <https://doi.org/10.1016/j.agrformet.2012.05.012>, 2012.