

The Influence of Glacial Northern Hemisphere Ice Sheets on Atmospheric Circulation

Himadri Saini^{1,2,3,4}, David K. Hutchinson^{1,2,5}, Josephine R. Brown^{3,4}, Russell N. Drysdale³, Yanxuan Du^{3,4}, and Laurie Menviel^{1,2}

¹Climate Change Research Centre, University of New South Wales, Sydney, New South Wales, Australia

²The Australian Centre for Excellence in Antarctic Science, University of New South Wales, Sydney, New South Wales, Australia

³School of Geography, Earth and Atmospheric Sciences, University of Melbourne, Melbourne, Victoria, Australia

⁴Australian Research Council Centre of Excellence for Weather of the 21st Century, University of Melbourne, Melbourne, Victoria, Australia

⁵Australian Research Council Centre of Excellence for Weather of the 21st Century, University of New South Wales, Sydney, New South Wales, Australia

Correspondence: Himadri Saini (himadri.saini@unimelb.edu.au)

Abstract. ~~During the The Laurentide ice-sheet affected the North Atlantic Ocean during the last glacial period, Northern Hemisphere (NH) ice sheets significantly influenced atmospheric circulation, however most studies primarily focused on the North Atlantic but its impact on the global atmospheric circulation remains unclear. Here, we investigate the influence of ice sheets on global climate during use the Australian Earth System Model (ACCESS-ESM1.5) to investigate the relative~~
5 ~~roles of~~ Marine Isotope Stage 3 (MIS 3, 65-25 ka) ~~, a period marked by prominent millennial-scale variability. Using the Australian Earth System Model (ACCESS-ESM1.5), we simulate the glacial climate around boundary conditions in shaping global climate ~49,000 years ago (49 ka). Our findings demonstrate that the NH ice sheets induced a~~, a period marked by prominent millennial-scale variability. Our simulations show that Northern Hemisphere (NH) ice sheets were the primary driver of large-scale circulation changes. In particular, NH ice-sheet topography induced a 6° and 4° southward shift of the
10 NH westerlies during boreal ~~summer and winter~~, ~~winter and summer, respectively~~, increasing rainfall over Eurasia during summer ~~by 31%~~ but reducing it in winter. ~~The influence of orbital parameters and greenhouse gases alone produces only minor circulation changes. Adding ice sheet albedo intensifies the NH cooling and strengthens the westerlies but does not significantly alter their latitudinal position. The ice sheets' influence also extends across the tropics and the Southern Hemisphere (SH). The most pronounced southward displacements~~ In contrast, orbital forcing and greenhouse gas (GHG)
15 ~~changes did not lead to a significant NH westerly shift, while ice-sheet albedo strengthened NH westerlies (10–14%) through enhanced cooling (by 3–4°C) without altering their position in both seasons. NH ice sheets topography also affected the global atmospheric circulation, leading to an additional 0.9° southward shift of the Intertropical Convergence Zone (ITCZ) and 1.5° southward displacement of the NH Hadley cell during austral summer—associated with enhanced Australian rainfall—occur when all ice sheet-related boundary conditions are included. In the SH, the presence of ice sheets leads to an~~ in austral summer,
20 ~~relative to changes simulated due to orbital forcing plus GHG (0.6° and 2° southward shifts for ITCZ and NH Hadley cell, respectively), and albedo (0.4° and 0.1°, respectively). The full glacial boundary conditions, including changes in Antarctic~~

ice-sheet topography, also lead to a 2° equatorward shift of the SH Hadley cell during JJA, with the largest displacement occurring when all boundary conditions are combined. Additionally, SH westerlies show no latitudinal shift in response to orbital parameters, greenhouse gases, or albedo alone, but exhibit a pronounced equatorward shift when ice-sheet topography is included. These findings highlight the non-linear interactions between ice sheets, large-scale atmospheric circulation, and precipitation patterns. Southern Hemisphere (SH) Hadley cell, and a 2.5° equatorward shift of SH westerlies during austral winter. Orbital forcing plus GHG and albedo primarily modulating the strength of the SH westerlies and tropical atmospheric circulation. These results highlight the roles of ice-sheet topography in controlling shifts in the atmospheric circulation and surface albedo in modulating atmospheric circulation intensity through radiative cooling.

30 1 Introduction

Earth's climate history has been shaped by the expansion and retreat of Northern Hemisphere (NH) ice sheets, particularly during the last glacial cycle. The expansion of large NH ice sheets has long been associated with global and regional cooling, particularly over the North Atlantic, which is directly influenced by the ice sheet extent and height (Kageyama et al., 2020). Ice sheet growth affects land-ocean geography by lowering sea levels (Lambeck et al., 2014), exposing land bridges, and altering local heating patterns due to the differing heat capacities of land and ocean (Byrne and O'Gorman, 2013). Additionally, changes in land bridges and oceanic gateways, such as the Bering Strait, influence large-scale ocean circulation, including the Atlantic Meridional Overturning Circulation (AMOC) (Hu et al., 2015). These factors have significant implications for the climate system, particularly through their impact on the atmospheric circulation and ocean currents.

In today's climate, warm water in the North Atlantic is transported northward via the Gulf Stream, and atmospheric Rossby wave propagation results in a northeastward-tilted Atlantic jet stream. This configuration contributes to the relatively mild winters in Europe compared to regions at similar latitudes in North America (Seager et al., 2002; Brayshaw et al., 2009). However, during past glacial periods, and particularly the Last Glacial Maximum (LGM; 19-21 ka), thick ice sheets covered northern North America and may have reached up to 5 km in height (Peltier, 2004). This so-called Laurentide Ice Sheet (LIS) caused a pronounced southward shift of the jet stream, and in some cases, a more zonal orientation compared to today's tilted pattern (Manabe and Broccoli, 1985; Pausata et al., 2011; Löfverström et al., 2014; Ullman et al., 2014). This southward shift of the jet stream, which was primarily driven by ice sheet topography, rather than changes in albedo or greenhouse gas (GHG) forcing (Pausata et al., 2011), had important consequences for European climate, impacting winter storm tracks, precipitation, and seasonal temperatures (Löfverström et al., 2014).

Previous research on understanding the impact of NH ice sheets on climate (Broccoli and Manabe, 1987; Pausata et al., 2011; Löfverström et al., 2014; Ullman et al., 2014; Wang et al., 2018; Izumi et al., 2023) primarily focused on the LGM, an important period due to its extreme climate conditions, including the coldest global temperatures (Kageyama et al., 2020), lowest atmospheric CO₂ levels (Köhler et al., 2017), and most extensive ice sheets (Peltier, 2004; Clark et al., 2009; Kageyama et al., 2017; Gowan et al., 2021). Most of these studies examined the North Atlantic region, assessing the impact of ice sheets on NH westerlies and the Atlantic jet, with limited investigation into regions beyond this sector. An exception from this list is

55 ~~the work of Löffverström et al. (2014), who investigated~~ Löffverström et al. (2014) investigated both the LGM as well as pre-
LGM ice sheets of Marine Isotope Stage (MIS) 5b (~88 ka) and MIS 4 (~66 ka). ~~Their study and~~ found that the westerlies
shifted southward in all ice sheet configurations, but a fully zonal jet orientation emerged only during the LGM. They also
suggested that the ice sheet topography dynamically induced warming in Alaska and central Asia, potentially contributing to
the westward expansion of the Eurasian ice sheets from MIS 4 to the LGM. Other studies, such as Ullman et al. (2014) and
60 Zhang et al. (2014a), further explored the impact of ~~varying LIS height on the atmospheric-oceanic system~~ different LIS heights
on both the atmosphere and the ocean.

In contrast, the climate of MIS 3 (65–25 ka) has received comparatively less attention in the context of ice sheet–atmosphere
interactions. Existing studies of MIS 3 have largely concentrated on large-scale oceanic responses, such as AMOC variability,
surface temperature patterns, and sea ice changes (~~Malmierca-Vallet and Sime, 2023; Brandefelt et al., 2011; Merkel et al., 2010; Guo et al.~~
65 ~~(Merkel et al., 2010; Brandefelt et al., 2011; Guo et al., 2019; Malmierca-Vallet and Sime, 2023)~~). Some have investigated trop-
ical climate variability, including changes in ENSO El Niño–Southern Oscillation (ENSO) behaviour during this period (Br-
andefelt et al., 2011; Merkel et al., 2010), while others ~~briefly mention reduced global precipitation, reported~~ a southward-shifted
ITCZ ~~; intensified and associated changes in tropical precipitation and~~ Southern Hemisphere (SH) ~~westerly wind stress, and~~
~~strengthened NH trade winds alongside weakened SH trade winds circulation~~ (Guo et al., 2019). However, the detailed, mech-
70 anistic impacts of NH ice sheets on atmospheric circulation during MIS 3—particularly beyond the North Atlantic—remain
underexplored. This gap may be partly due to the significant millennial-scale climate variability that characterizes MIS 3—in-
cluding abrupt temperature fluctuations in Greenland of 8–10°C (Huber et al., 2006) and multiple episodes of AMOC weaken-
ing (Meniel et al. (2020) and references therein)—which ~~make makes~~ it challenging to isolate the equilibrium climate response
to boundary conditions. Here, we employ quasi-equilibrium climate simulations with prescribed
49 ka boundary conditions and
75 systematically isolate the effects of orbital parameters and GHGs, ice-sheet albedo, and ice-sheet topography. This framework
enables us to quantify the mechanistic contribution of each forcing to atmospheric circulation and hydroclimate changes during
MIS 3.

~~To address this, we focus on the~~ The period around 49 ka ~~within MIS 3, which~~ stands out due to one of the highest obliquity
values (24.43°) of the last glacial cycle (Berger, 1978), resulting in greater summer insolation in both hemispheres compared
80 to the preindustrial (PI). Variations in obliquity alter the distribution of incoming solar radiation across latitudes at the top
of the atmosphere. High obliquity increases incoming solar insolation at high latitudes in summer, while decreasing it in
winter, leading to enhanced seasonal contrast with warmer summers and colder winters in both hemispheres. These changes
in radiative forcing influence the large-scale atmospheric circulation and can affect the position and strength of the Hadley
circulation and the ITCZ (Mantsis et al., 2014; Liu et al., 2015; Zhang et al., 2022). In particular, 49 ka exhibits an obliquity
85 value higher than some interglacial periods, such as the Last Interglacial (LIG, 127 ka), yet features a climate state marked by
low CO₂ (~199 ppm, similar to LGM (Köhler et al., 2017)) and extensive Laurentide and Scandinavian ice sheets (Gowan
et al., 2021), leading to a colder-than-PI climate. Furthermore, global sea levels were approximately 60–65 m lower than PI
(Shakun et al., 2015). This ~~combination of factors makes 49 ka an intriguing period for assessing how ice sheets influenced~~
~~atmospheric circulation and hydroclimate~~ unusual combination of strong seasonal insolation forcing, associated with high

90 obliquity and low atmospheric CO₂ and extensive ice sheets, provides an opportunity to investigate how competing climate forcings interact to shape atmospheric teleconnections and hydroclimate responses on a planetary scale.

Here, we use the Australian Earth System Model (ACCESS-ESM1.5) to examine the influence of glacial ice sheets on global atmospheric circulation, focusing on their cascading effects across Eurasia, the tropics, and the SH. Our analysis emphasizes the role of the ice sheets in modulating Rossby waves, Hadley circulation, and the Intertropical Convergence Zone (ITCZ),
95 shaping rainfall patterns in both hemispheres, and influencing regional climates, such as over Eurasia and northern Australia. In addition, we examine how the ~~inclusion of the~~ larger MIS 3 ice-sheets ~~compared to PI influence~~, relative to PI, influences SH westerlies.

2 Methods

2.1 Model description

100 We employ the Australian Community Climate and Earth System Simulator-Earth System Model version 1.5 (ACCESS-ESM1.5), a comprehensive Earth system model developed for the Coupled Model Intercomparison Project Phase 6 (CMIP6) (Ziehn et al., 2020). The atmospheric model is the UK Met Office Unified Model version 7.3 (UM7.3) (Martin et al., 2010; The HadGEM2 Development Team et al., 2011) with a horizontal resolution of $1.875^\circ \times 1.25^\circ$ and 38 vertical levels. The land surface is represented by the Community Atmosphere Biosphere Land Exchange (CABLE) model version 2.4 (Kowalczyk et al., 2013), which includes biogeochemical processes through the CASA-CNP module (Wang et al., 2010). The ocean
105 component consists of the NOAA/GFDL Modular Ocean Model version 5 (MOM5) (Griffies, 2012), featuring a horizontal resolution of $1^\circ \times 1^\circ$, with the meridional spacing refined to 0.4° in the Southern Ocean and 0.33° near the equator, and with 50 vertical levels. Furthermore, sea ice processes are simulated using the Los Alamos National Laboratory Consortium Model for Sea Ice Development (LANL CICE) version 4.1 (Hunke et al., 2010). The coupling between the atmosphere, ocean and
110 sea ice components is achieved through the Ocean Atmosphere Sea Ice Soil-Model Coupling Toolkit (OASIS-MCT) (Craig et al., 2017). The ocean carbon cycle ~~dynamics are is~~ modeled using the ~~nutrient-phytoplankton-zooplankton-detritus (NPZD) scheme in the~~ World Ocean Model of Biogeochemistry and Trophic dynamics (WOMBAT) model for ocean biogeochemistry which includes a nutrient-phytoplankton-zooplankton-detritus (NPZD) scheme (Oke et al., 2013).

This model has previously been used to simulate the ~~last interglacial~~ Last Interglacial (lig127k) time slice (127 ka) as part of
115 the Tier 1 Paleoclimate Model Intercomparison Project 4 (PMIP4)-CMIP6 experiments (Yeung et al. (2021), MIS9e (~~~~~336-321 ka; Duboc et al. (2025)), and mid-Holocene (6 ka; Mackallah et al. (2022)).

2.1.1 Experimental setup

~~The~~

2.2 Experimental setup and diagnostics

	PI	49 ka
Orbital parameters		
Eccentricity	0.01674	0.01292
Obliquity (°)	23.459	24.435
Perihelion-180 (°)	100.33	62.451
Greenhouse Gases		
CO ₂	284.3 (ppm)	199 (ppm)
N ₂ O	273 (ppb)	237 (ppb)
CH ₄	808.2 (ppb)	432 (ppb)

Table 1. Orbital parameters and Greenhouse gas concentrations for PI and 49 ka climate.

Experiment	Model years	Years analysed	Orbital parameters and GHGs	Albedo and vegetation	Salinity	Topography and runoff
49ka-co	0 to 618	569 to 618	49ka <u>49 ka</u>	PI	PI	PI
49ka-alb	297 to 569	520 to 569	49ka <u>49 ka</u>	49ka <u>52.5 ka</u>	PI	PI
49ka-full	824 to 1584	1485 to 1584	49ka <u>49 ka</u>	49ka <u>52.5 ka</u>	49ka <u>52.5 ka</u>	49ka <u>52.5 ka</u>

Table 2. Boundary conditions used in the different 49 ka experiments. Model years indicate the total number of years that the corresponding experiment has run for as shown in the time series of Figure A1.

120 As a reference for our paleoclimate simulations, we use the 1000-year pre-industrial (PI) ~~simulation is the 1000-year piControl~~run
control simulation (piControl) based on the year 1850 CE ~~as described~~ by Ziehn et al. (2020). We analyse the average of the
last 100 years of this simulation (Figure A1, black line). Orbital parameters and GHG concentrations for the PI configuration
are provided in Table 1.

To investigate the climate around 49 ka, the boundary conditions are changed in a step-wise manner using the ACCESS-
125 ESM1.5 model. The first experiment, 49ka-co, simulates a baseline climate with orbital parameters (Berger, 1978) and GHG
concentrations (Köhler et al., 2017) being set to 49 ka values (Table 1). 49ka-co is run for 618 years (Figure A1, cyan line).

The model uses prescribed Plant Functional Types (PFTs) that remain fixed during each individual simulation. We modified
the PFT distributions in selected simulations to reflect the altered boundary conditions. ~~For instance~~Therefore, in a second
step, 49ka-alb, we ~~incorporate the ice sheet extent~~modify the land surface properties by incorporating only the ice-sheet extent

130 ~~(no topography yet)~~ from Gowan et al. (2021) corresponding to 52.5 ka, ~~adopting~~. We adopted the maximal scenario with an ice-capped Hudson Bay (Figure 1d). This time period was chosen because the simulated ice sheet at 52.5 ka most closely matches the sea-level estimates for 48–52 ka, based on global seawater $\delta^{18}\text{O}$ stack records (Shakun et al., 2015). The NH ice sheets at ~~49 ka~~ this time period include two separate ice sheets over the North American continent—Laurentide in the east and Cordilleran in the west—as well as the Greenland Ice Sheet and Fennoscandian and Eurasian ice sheets (Figure 1d). Their extent and height are smaller than during the LGM (not shown), which featured a single, extensive LIS over North America and continuous ice sheet coverage from Eurasia to Scandinavia, fully covering the Barents Sea. The Antarctic Ice Sheet (AIS) at ~~49–52.5~~ ka also has a smaller extent compared to that of the LGM, especially over the Ross and Weddell Seas. At ~~49–52.5~~ ka, the LIS was up to 3 km thick, while the AIS was up to 3.6 km thick. In the present day, permanent ice cover in the NH is restricted to Greenland (Figure 1c).

140 In the 49ka-alb experiment, additional modifications are introduced at different stages (Figure A1, orange line). The vegetation cover is modified to reflect ice coverage by removing vegetation in ice-covered regions and converting needle leaf evergreen forests in areas between the Cordilleran and LIS to bare soil (Figure 1a,b). These modifications align with the vegetation reconstruction from Allen et al. (2020) for 52 ka. Additionally, C3 crops are removed, and replaced by the next two most dominant vegetation types in the corresponding grid cell. For example, in India, C3 crop values are replaced by broadleaf deciduous type and C4 grass. In Europe and eastern North America, they are replaced by broadleaf deciduous type and C3 grass, while they are replaced by broadleaf deciduous over China. 49ka-alb (Figure A1, orange line) starts from year 297 of 49ka-co, and was run for a total of 272 years with the above mentioned albedo and vegetation changes (Table 2). As can be seen in Figure A1, 49ka-alb is not in complete equilibrium. To this 49ka-alb experiment, we add a gradual increase in surface salinity from year 547 (Figure A1, magenta line) to account for the 44 m sea level drop relative to PI, based on the prescribed ice sheet, achieving a global average increase of +0.33 psu from PI levels. At year 729, the Bering Strait is closed.

Our final experiment, referred to as 49ka-full (Figure A1, blue line, Table 2), starts at year 824, when the topography is adjusted to reflect conditions at ~~49–52.5~~ ka, based on Gowan et al. (2021), including an increase in ice sheet height and changes to the land-sea mask, such as the closure of the Bering Strait (done in the previous step), Hudson Bay, the Sahul and Sunda shelves, and the Tasman Gateway (Figure 1). Nearest-neighbour interpolation is applied to determine vegetation types on newly exposed land. The river runoff is also adjusted at year 1258, to reflect the ~~49–52.5~~ ka topography. In total, 49ka-full is run for 760 years, with 325 years incorporating all the boundary conditions. In the Results section, we compare the ~~average-averages~~ of the last 100 years of 49ka-co, last 50 years of 49ka-alb, and the ~~average-of-the~~ last 100 years of the 49ka-full experiment (Table 2, Figure A1, darker colors). Although the deep ocean has not fully equilibrated in the 49ka-co and 49ka-alb simulations, our comparison with the 49ka-full simulation focuses primarily on atmospheric circulation responses, which adjust to changes in boundary conditions much more rapidly than the ocean. Surface and atmospheric variables exhibit no systematic drift over the analysis period, supporting the robustness of the diagnostics presented here.

160 The ITCZ position is determined using the precipitation centroid index (Braconnot et al., 2007), which identifies the latitude that evenly divides the tropical precipitation amount between 20°S and 20°N. This approach effectively represents the center of mass of tropical rainfall distribution. The strength of the northern Hadley cell ~~is~~ is determined by the maximum value of




Figure 1. (a,b) Vegetation and ice mask changes at 49 ka compared to PI. The color indicates the dominant Plant Functional Type (PFT) of each grid cell. (c,d) Ice sheet thickness (m) and extent from (Gowan et al., 2021) for PI and 49 ka (represented by 52.5 ka as described (refer to the description in the text for the choice of 52.5 ka ice sheets))

165 the atmospheric mass streamfunction, ψ , within the 400–600 hPa pressure range, while the strength of the southern Hadley cell is determined by its minimum value over the same vertical pressure range (Equation 1).

$$\psi = \frac{2\pi R \cos \phi}{g} \int_0^p v dp \quad (1)$$

where, R is the radius of the earth, ϕ is latitude, g is gravitational acceleration, v is the meridional wind and p is pressure. ψ is expressed in Sverdrups (Sv), where $1 \text{ Sv} = 10^9 \text{ kg s}^{-1}$

170 3 Results

3.1 Simulated glacial climate at 49 ka compared to PI

We first describe the 49ka-full experiment in comparison to PI. At 49 ka, the higher obliquity (24.43°) compared to PI (23.45°) results in slightly increased summer insolation in both hemispheres (Figure A2). Despite the higher obliquity, the glacial simulation (49ka-full) exhibits substantial oceanic and atmospheric cooling due to lower GHG concentrations and extensive ice sheet expansion at this time (Figure 2). Global annual mean surface air temperatures (SATs) at 49 ka are 2.7°C lower than

175

PI (Table A1 [and Figure 2a](#)), with the NH cooling by 4.9°C and the SH by 2°C ([Figure 2a](#) ~~not listed~~). The most pronounced cooling is simulated in ice sheet regions and northern high latitudes. Seasonal variations are evident in the NH (Table A1), where temperatures drop by 6.4°C in winter (DJF) and 2.6°C in summer (JJA). Over SH, DJF (austral summer) SATs are 1.7°C lower, while JJA (austral winter) SATs are 2.4°C lower (Table A1).

180 Global mean ocean temperature is 1°C lower, and global annual mean sea surface temperature (SST) (Figure 2b) is reduced by 1.2°C (Table A1). The simulated SATs and SSTs anomalies show a good agreement with proxy records indicating a qualitative estimate of the changes between Greenland Interstadial (GI 13; 49ka-50ka) and PI climate (Figure 2a,b). Higher SSTs in the Labrador region are ~~linked to~~ [associated with](#) a stronger AMOC in 49ka-full, with a transport rate of 31 Sv (about 10 Sv greater than the PI value). This strengthening is attributed to topographical changes and the presence of ice sheets, which enhance westerly winds over Baffin Bay and weaken them over the Labrador Sea (Figure A3, [middle panels](#)). These wind changes modify the North Atlantic gyres, leading to an overall strengthening while shifting the separation between the subpolar and polar gyres southward and orienting the subpolar gyre in a more zonal direction (Figure A3, [bottom panels](#)). A strengthening of the cyclonic circulation in the Labrador Sea is also simulated, which increases the westward transport of warm water from south of Iceland to the Labrador Sea and further strengthens the AMOC ([reflected by the increase in mixed layer depth in this](#)

185 [region in 49ka-full compared to PI, Figure A3, top panels](#)). The stronger AMOC also influences sea ice extent. During boreal winter, the sea-ice edge at 49 ka expands in the North Pacific and Nordic Sea sectors but retreats in the Labrador Sea due to the influx of warm water (Figure 2b, orange dashed line compared to black dashed line). In boreal summer, the 49 ka sea-ice edge is similar to PI across most regions, except in the Nordic seas, where the sea-ice edge extends further south than in PI by 5 to 6° latitude (Figure 2b, orange solid line compared to black solid line). In the SH, austral summer sea ice retreats in the

190 South Atlantic as stronger North Atlantic Deep Water (NADW) formation leads to a deep-ocean warming of about 0.3°C (not shown), and this warm water gets upwelled in the South Atlantic (Figure 2b, orange dashed line compared to black dashed line). However, the austral winter sea ice edge remains similar to PI in the South Atlantic while expanding in the Indian and Pacific sectors of the Southern Ocean (Figure 2b, orange solid line compared to black solid line).

Drier conditions are simulated at 49 ka compared to PI over NH high-latitude land regions, particularly over the LIS, due to colder conditions and reduced atmospheric moisture availability (Figure 2c). However, precipitation increases in the North Atlantic and Eurasia, between 30°N and 60°N, as well as over India and East Asia. In contrast, most land areas in the SH display decreased precipitation, except over northern Australia, parts of Southeast Africa, and over eastern Brazil.

3.2 Impact of 49 ka boundary conditions on the circulation ~~changes~~ over North Atlantic and Eurasia

At 49 ka (experiment 49ka-full), the presence of large ice sheets over North America significantly alters the atmospheric circulation, particularly in the North Atlantic region. In DJF, the westerlies over the North Atlantic strengthen due to the strong meridional temperature gradient and shift by ~6° southward compared to PI (Figures 3a,c and 4a), with a more zonal flow replacing the northeast-southwest tilt of the ~~jet stream~~ [wind field](#) (Figure 5). This pattern is evident both at the surface and [at the jet stream level](#) in the upper troposphere (Figure A4). Geopotential height (GH) differences (Figure 3e) indicate that the LIS induces a high-pressure system (not shown) over North America, initiating a Rossby wave response. This Rossby wave causes a




Figure 2. Annual mean anomalies of surface air temperature (SAT), sea surface temperature (SST), and precipitation for 49 ka climate (exp 49ka-full) compared to PI. Contours show seasonal mean 15% sea ice concentration for (orange) 49 ka and (black) PI during DJF (dashed) and JJA (solid). Stippling indicates non-significant changes based on a Student's t-test at [the](#) 95% confidence level. Stars in subpanels (a) and (b) indicate qualitative estimates of the changes between Greenland Interstadial (GI 13) and PI climate, as inferred from the proxy records (MDO1-2444,ODP-977A,MD95-2043,MD01-2443,MD95-2042 (Martrat et al., 2007); ODP-1233 (Kaiser et al., 2005); TR163-19 (Lea et al., 2000); RC13-110 (Feldberg and Mix, 2003); ODP846B (Martínez et al., 2003); TG-7 (Calvo et al., 2001); MD97-2120 (Pahnke et al., 2003); MD07-3128 (Caniupán et al., 2011); EPICA Dronning Maud Land (EPICA community members, 2006); WAIS (WAIS Divide Project Members, 2015); NGRIP (Huber et al., 2006; Kindler et al., 2014); SO188-17286-1 (Lauterbach et al., 2020). Dark (light) blue [stars](#) represent much colder (slightly colder) climate [in proxy records](#).

210 southward shift of the Icelandic low and a weakening of the Azores High, contributing to the zonal orientation of the westerlies (Figures 3c, 5, and A5). The southward shift and ~~zonal~~more zonal orientation of the jet alter moisture-bearing storm tracks, which in PI were tilted southwest-northeast, to a more ~~zonal configuration~~southerly pathway at 49 ka. This reorganization of pressure systems (not shown), with a negative GH anomaly extending from the North Atlantic to eastern Eurasia (Figure 3e), alters precipitation patterns, enhancing precipitation over the North Atlantic and western Europe (Figure 3g). However, colder
215 conditions over Eurasian landmasses lead to drier conditions inland in DJF, despite ~~the persistence of~~ negative GH anomalies (Figure 3a,e,g).

The westerlies are also shifted southward in JJA ~~;~~(Figures 3d, and 4b), with the flow remaining more zonal in 49 ka than during PI (~~Figures 3d, 5, and 4b~~Figure 5, right panels), coinciding with a positive GH anomaly over the Labrador Sea region while a negative GH anomaly forms over Europe (Figure 3f). These GH anomalies enhance the westerly flow and contribute
220 to moisture transport convergence (not shown) toward the Eurasian landmass, increasing precipitation along the ~~westerly band~~westerlies in the North Atlantic Ocean and over parts of Eurasia (Figure 3d,f,h). In addition, ~~mild temperature conditions~~minor cooling over Eurasia (Figure 3b) further ~~contribute~~contributes to the increased precipitation. Precipitation also increases north of 60°N in Asia (Figure 3h) due to higher specific humidity (not shown) resulting from warmer conditions and the development of a negative GH anomaly (Figure 3b,f). The warming in this region arises from the topographical changes corresponding to 49
225 ka, where ocean points have been converted to land. Since land has a lower heat capacity than the ocean, it warms more rapidly in summer, amplifying the temperature-driven moisture increase. Conversely, north of 60°N in Europe precipitation decreases. This is due to the ~~influx of~~ colder north-easterly winds and the presence of a blocking high over the Labrador-Greenland sector, which suppresses rainfall in these areas.

The role of individual boundary conditions becomes evident in the 49ka-co experiment, which includes only orbital parameters and GHGs. During DJF, strong cooling in the 49ka-co experiment reduces moisture availability, leading to widespread drying over NH landmasses (Figure 6a,c). The westerlies are ~~slightly enhanced and shift southward by only~~enhanced and slightly shift southward (~1°) (Figure 4a, 9a). This shift is accompanied by a low-pressure anomaly over the North Atlantic and a high-pressure system over Europe, which enhances subsidence and suppresses precipitation over Europe (Figure 6b,c). This pattern is consistent with the overall drying trend in the NH, except for a small increase in precipitation over the North Atlantic
235 Ocean (Figure 6c), likely due to storm activity in this region. In 49ka-alb, changing the vegetation and albedo has little impact compared to 49ka-co, except over northern North America (Figure 6d,f), where cooling is enhanced by 8°C and drying increases by 25% due to the albedo change over the LIS. The westerlies intensify further compared to 49ka-co, but their position shifts back to the PI configuration (Figure 4a). The GH response features a low-pressure anomaly over the Hudson Bay and Labrador Sea, which is consistent with some increased precipitation in this region (Figure 6e,f).

240 In JJA, the 49ka-co experiment shows no change in westerly strength and position over the North Atlantic (Figure 7b, 4b 9b) and modest temperature and little significant precipitation changes over the NH landmass (Figure 7a,c). In contrast, the 49ka-alb experiment exhibits pronounced drying over the Laurentide region and northern Europe, driven by intense cooling from albedo changes and persistent high pressure over these regions (Figure 7d-f). While the westerlies strengthen in 49ka-alb,




Figure 3. Anomalies between 49ka-full and PI for (a,b) SAT (°C), (c,d) zonal winds at 850 hPa (m/s) overlaid with 850 hPa wind vectors (m/s), (e,f) geopotential height at 850 hPa (m) with zonal mean removed and overlaid with 850 hPa wind vectors (m/s), and (g,h) precipitation (mm/yr) during (left) DJF and (right) JJA. Stippling in subpanels (g,a,b) and (g,h) indicates non-significant changes based on a Student's t-test at the 95% confidence level. The white space in subpanels c-f indicates the ice mask at the 850 hPa level.




Figure 4. Zonal winds at the 850 hPa level (m/s) averaged (top) over the North Atlantic (80W:0E) and (bottom) averaged over all longitudes representing westerlies over the SH during (a,c) DJF and (b,d) JJA seasons.

their position remains unchanged (Figure 4b, 9b). This intensification leads to slightly wetter conditions inland over Eurasia
245 in the 49ka-alb scenario.

~~Figures 6, 7, 4a, and 9 show that~~ In summary, only the full implementation of the ice sheet, including its topography, can induce a significant planetary wave response in the atmosphere, with southward-shifted and intensified North Atlantic westerlies ~~-(Figures 4a, 6, 7, and 9)~~. This response results in substantial drying over the northern high-latitude landmasses, including the Laurentide region during DJF, along with increased milder and wetter conditions over Eurasia during JJA.




Figure 5. Zonal winds at 850 hPa (m/s) overlaid with [850 hPa](#) wind vectors (m/s) in (left) DJF and (right) JJA for (top) PI and (bottom) 49ka-full. [The white space the ice mask at the 850 hPa level.](#)

250 3.3 Impact of 49 ka boundary conditions on tropical and SH atmospheric circulation.

In DJF, the presence of the LIS forces a southward shift of the Ferrell cell. This displacement further leads to a southward migration of the NH Hadley cell's southern edge, shifting from 7.5°S in PI to 11°S in 49ka-full (Figure 8a, see zero contour), and results in a globally averaged ITCZ shift of approximately 1.5° southward in DJF (Figure 3g, 9a). Additionally, the NH Hadley cell strengthens by 20 Sv relative to PI ([refer to section 2.2 for diagnostics](#)). The southward shift of the NH Hadley cell and ITCZ during DJF is also captured in 49ka-co (Figure 8c), where the NH Hadley cell strengthens by +11 Sv relative to PI. The shift in the ITCZ is driven by peak summer insolation in the SH due to higher obliquity at 49 ka. However, the magnitude of change in 49ka-co is smaller than in 49ka-full (Figure 8a, 9a), with an ITCZ shift of 0.6°. Implementing albedo changes in 49ka-alb enhances the Laurentide cooling, further strengthening the NH Hadley cell (by +8 Sv), albeit without additional displacement (9.5°S in 49ka-co vs. 9.6°S in 49ka-alb) (Figure 8c,e). The minor shift in the NH Hadley cell and ITCZ position explains why lower latitude precipitation changes remain insignificant between 49ka-alb and 49ka-co (Figure 6c,f). However, the inclusion of topography in 49ka-full results in a slight additional strengthening of the NH Hadley cell and a more substantial southward ITCZ shift (by 1.5° relative to 49ka-alb), due to the stronger temperature gradient between the two hemispheres (Figure 9a, Table A1).




Figure 6. DJF (a,d,g) SAT (°C), (b,e,h) geopotential height (m) overlaid with winds-wind vector anomalies at 850 hPa and precipitation (mm/yr) anomalies for (a-c) 49ka-co minus PI, (d-f) 49ka-alb minus 49ka-co, (g-i) and 49ka-full minus 49ka-alb. Stippling indicates non-significant changes based on a Student's t-test at the 95% confidence level. The white space in subpanels (b,e,h) indicates the ice mask at the 850 hPa level.

49ka-co - PI




Figure 7. Same as Figure 6 but for JJA.

During JJA, the southern edge of the SH Hadley cell shifts northward by 2° in 49ka-full, moving from 32°S in PI to 30°S , along with a northward shift of the Southern Ferrel cell (Figure 8b, see zero contour). This shift causes a slight northward migration of the ~~globally-averaged-global~~ ITCZ (Figure 3h), while the SH Hadley cell strengthens by 35 Sv. In 49ka-co, a slight warming over the North Atlantic and stronger cooling in the SH shifts the SH Hadley cell's southern edge to 31°S , accompanied by a strengthening of +22 Sv (Figure 8d, 9b). The changes in 49ka-alb do not alter the SH Hadley cell's latitudinal position significantly but weaken its strength to PI levels (Figure 8f, 9b), due to intensified Laurentide cooling. As a result, precipitation changes in the lower latitudes remain small (Figure 7c,f). Temperature contrasts between hemispheres are reduced by ice sheet topography (~~Tabel A1~~[Table A1, comparison of JJA NH minus SH SAT differences between 49ka-full, 49ka-alb and 49ka-co](#)), which induces localized warming in Siberia and strong Antarctic cooling, and by changes in the land-sea mask with the replacement of ocean grid cells by land.




Figure 8. Atmospheric mass streamfunction (Sv) anomalies (shading) between (a,b) 49ka-full and PI, (c,d) ~~49ka-ic~~49ka-co and PI, and (e,f) ~~49ka-ice~~49ka-alb and PI, for (left) DJF and (right) JJA. Contours (solid=positive; dashed=negative) are PI (black) absolute streamfunction values. The thick black and blue lines represent the zero contour for PI and 49ka experiments, respectively.

This reduced thermal gradient with warming in the NH leads to a slight northward ITCZ shift, further strengthening the SH
 275 Hadley cell and shifting it an additional 1° northward, toward the warmer hemisphere, relative to 49ka-alb.

The SH westerlies also exhibit distinct responses across the simulations. In DJF, they weaken slightly in 49ka-co ~~but weaken further in and~~ 49ka-alb, accompanied by a small northward shift (Figure 4c). In contrast, in 49ka-full, they remain in the same position as in PI but strengthen due to the increased height and extent of the AIS. During JJA, the SH westerlies weaken across all 49 ka simulations compared to PI, with the ~~strongest-most pronounced~~ weakening occurring in the experiment where albedo is changed (Figure 4d). However, in 49ka-full, they shift northward by 2.5° , a displacement linked to the northward shift of the SH Hadley cell in 49ka-full (Figure 4d, 9b).

3.4 Impact of 49 ka boundary conditions on Australian rainfall

~~As shown in Figures 3, 9a, the~~ The implementation of 49 ka boundary conditions leads to a notable increase in Australian rainfall during the monsoon season (Figures 3g and 9a). This section focuses on the large-scale processes driving this regional rainfall response. While a detailed analysis of monsoon changes in other regions lies beyond the scope of this study, our results underscore the ~~broader~~ influence of NH ice sheets on hemispheric-scale atmospheric dynamics.

The presence of NH and larger AISs induces substantial atmospheric circulation changes by modifying the Hadley and Ferrel cells and influencing the westerlies in both hemispheres. These shifts propagate through the atmosphere, altering surface pressure systems and triggering a hemispheric-scale response.

During DJF, the ~~NH ice sheet~~ full boundary conditions in 49ka-full forces a southward shift of the Hadley and Ferrel cells (Figure 8a), enhancing subsidence over East Asia around 40°N and leading to drier conditions (Figure 3e,g). This southward displacement of the NH Hadley cell extends across hemispheres, impacting Australian rainfall by shifting convergence southwards along with the ITCZ (Figure 8a). These changes explain the wetter conditions in northern Australia and drier conditions over the Maritime Continent (Figure 3g). A comparable but weaker response is simulated in the 49ka-co experiment (Figure 6c). In contrast, changes in albedo do not impact Australian precipitation significantly (Figure 6f). However, modifying the topography induces wetter conditions in northern Australia driven by shifts in the Hadley cell and ITCZ, while southern Australia experiences drier conditions (Figure 6h,i).

During JJA, in response to the 49 ka boundary conditions, most of Australia experiences drying (Figure 3h), primarily due to the establishment of blocking high-pressure systems south of the continent (Figure 3f). These high-pressure systems reinforce subsidence and restrict moisture transport ~~into the~~ from the Southern Ocean into the Australian region (not shown). However, the drying response is not significant over central Australia and remains less pronounced along the eastern and western edges of the continent. The southwestern part of Australia becomes wetter, while the southeastern region experiences drying, due to the northward displacement of the westerlies, resembling a negative winter Southern Annular Mode (SAM). Significant drying also occurs across northern Australia and much of the Maritime Continent south of the equator.

When only orbital parameters and GHGs are modified, the SH Hadley cell shifts northward and strengthens. However, in this scenario, the weakening of the SH westerlies and the southward displacement of subtropical high-pressure systems result in increased rainfall over southern Australia (Figure 7b,c). Introducing albedo changes alone has little impact on the Australian climate (Figure 7d-f). However, when ice sheet topography is incorporated, the northward shift in SH westerlies linked with the strengthening and equatorward displacement of the SH Hadley cell causes enhanced subsidence and thus drier conditions

310 over Australia, coinciding with the northward ITCZ migration (Figure 7i, 4d). Compared to the PI climate, this drying effect is somewhat mitigated, as the westerlies — despite their northward shift— remain weaker overall.

4 Discussion

Our simulations of the 49 ka glacial climate reveal a complex interplay of ice sheet topography, GHG forcing, and orbital configurations. The results underscore how large-scale climatic features, such as atmospheric circulation patterns, precipitation
315 distribution, and ocean circulation, respond to glacial boundary conditions. Additionally, much of the existing research has focused primarily on the North Atlantic region, driven by the greater availability of paleoclimate data from this area and the presence of extensive North American ice sheets that strongly influenced both regional and global climate. Our results add to this literature by demonstrating that the influence of glacial ice sheets extends well beyond the North Atlantic, with significant climatic impacts also occurring in other regions, including the SH. Although the present experimental design does
320 not fully isolate the individual contributions of the Laurentide and AISs, sensitivity experiments and previous studies indicate that the NH ice sheet topography, dominated by the LIS, provides the primary control on midlatitude circulation changes. SH circulation changes arise from the combined influence of both ice sheets, with Antarctic topography acting primarily through modulation of inter-hemispheric temperature gradients.

Here, the simulated state in the 49ka-full experiment represents an interstadial climate. The global annual mean cooling in
325 49ka-full relative to PI (a 2.7°C reduction in global SATs) aligns with the expected glacial forcing associated with lower GHG concentrations and extensive ice sheet expansion. The pronounced winter cooling in the NH, exceeding 6°C, is ~~consistent with similar to~~ PMIP4 simulations of LGM climate, with ice sheet-driven albedo changes amplifying high-latitude cooling during glacial periods (Kageyama et al., 2020). This magnitude of cooling from PI is also broadly consistent with other MIS 3 modelling studies, though direct comparisons are not possible as applied boundary conditions between other MIS 3 studies
330 and ours were different. ~~For example, Guo et al. (2019) found a 2.9°C~~ cooling-for-globally averaged annual mean cooling over a 38 ka time slice using the Norwegian Earth System Model (NorESM)~~model~~. Simulations using the Community Climate System Model version 3 (CCSM3) model suggest stronger cooling, with Merkel et al. (2010) reporting a 3.4°C drop at 35 ka, and Zhang et al. (2014b) estimating 3.5°C at 38 ka.

Changes in AMOC

335 Proxy records suggest significant AMOC variability during the MIS 3 period, with stronger AMOC during the onset of interstadials (Böhm et al., 2015; Henry et al., 2016; Menviel et al., 2020), and weak AMOC during stadials. The processes driving this AMOC variability ~~is-are~~ still debated (Menviel et al., 2020). Model simulations ~~also show considerable spread in AMOC strength across different of~~ MIS 3 ~~timeslices and model setups. For instance, the~~ climates show a range of AMOC responses depending on the prescribed boundary conditions, particularly the configuration of Northern Hemisphere ice sheets.
340 For example, CCSM3 simulation simulations at 35 ka, initialized from the LGM, showed a and 44 ka show AMOC reductions of approximately 40% (Merkel et al., 2010) and 50% (Brandefelt et al., 2011), respectively, while a University of Victoria (UVic)-ESM simulation at ~40 % reduction in AMOC strength (Merkel et al., 2010), while the same model produced ka

produces a ~33% reduction relative to PI (Saini et al., 2024). In contrast, other simulations show little change or a modest strengthening of the AMOC. For example, Zhang et al. (2014b) simulated a ~50% reduction at 44 ka, also initialised from the LGM state (Brandefelt et al., 2011). In contrast, NorESM simulations indicate 7% strengthening at 38 ka using CCSM3, while Guo et al. (2019) reported a ~13% strengthening of the AMOC at 38 ka (Guo et al., 2019), using NorESM. Similarly, Van Meerbeeck et al. (2009) simulated a 22% increase in AMOC strength during MIS 3 compared to PI (estimated from Menviel et al., 2008) using the LOVECLIM model.

The simulated AMOC in our 49ka-49ka-full simulation is strong (31 Sv) with a ~47% increase relative to PI. This strengthening is driven by changes in the LIS, which enhances westerly winds over Baffin Bay while weakening them over the Labrador Sea, intensifying the cyclonic circulation in the Labrador Sea and facilitating the transport of warm, saline waters to the region. This process is also associated with sea ice retreat in polar waters, further reinforcing the AMOC. These findings align with earlier modelling studies emphasizing the role of topography and ocean-atmosphere interactions in AMOC enhancement (Pausata et al., 2011; Zhang et al., 2014a; Hu et al., 2015; Brown and Galbraith, 2016; Klockmann et al., 2018; Sherriff-Tadano et al., 2018; Guo et al., 2019). For example, Pausata et al. (2011) demonstrated that the presence of an extensive ice sheet during the LGM enhanced AMOC by altering atmospheric circulation patterns. However, their simulations also showed that the cooling effects from lower GHG concentrations and increased albedo suppressed AMOC strength. In our 49ka-full experiment, these cooling effects appear to be offset by high insolation state at 49 ka, allowing topographical impacts to dominate.

The simulated AMOC in 49ka-full is stronger than that reported in most previous MIS 3 simulations. However, the spread among MIS 3 simulations highlights the sensitivity of AMOC strength to the prescribed LIS configuration. Previous studies focusing on LGM climate have shown that changes in North Atlantic westerlies induced by the presence of the LIS can strengthen the AMOC through enhanced wind-driven circulation and deep-water formation (Muglia and Schmittner, 2015). Consistent with this mechanism, the stronger AMOC in our 49ka-full simulation is associated with LIS-driven changes in atmospheric circulation and enhanced westerlies over the North Atlantic. Conversely, changes in LIS height have also been shown to strongly affect AMOC stability and can trigger transitions between different overturning states (Zhang et al., 2014a), indicating that different ice-sheet reconstructions can produce markedly different AMOC responses. Since the LIS evolved substantially throughout MIS 3 and remains poorly constrained (Malmierca-Vallet and Sime, 2023), alternative but equally plausible ice-sheet configurations may yield weaker AMOC states than simulated here. This sensitivity is consistent with evidence for large AMOC variability during MIS 3, including rapid transitions and possible overshoots following Heinrich stadials. We therefore interpret the strong AMOC in 49ka-full as a response to the specific ice-sheet configuration prescribed in our experiments rather than as a robust feature of all MIS 3 climates.

Evidence based on foraminifer assemblages and mineral grains in ice-rafted debris also suggests a northward retreat of sea ice and higher temperatures in the Labrador Sea between ~56 and 49 ka, potentially driven by the insolation maxima (Griem et al., 2019). After 49 ka, sea ice variability increased, with near-perennial sea ice cover during most stadials and seasonal ice-free conditions during interstadials. However, it remains uncertain whether interstadials reflect a mean state or are primarily linked to ~~DO~~ Dansgaard-Oeschger (D-O) variability.

Changes in NH westerlies

The simulated changes in the North Atlantic westerlies and associated precipitation patterns during DJF highlight the critical role of ice sheet topography in driving atmospheric circulation (Figure 9a). At 49ka, the surface westerlies shift southward and adopt a more zonal flow (Figure 3), aligning with previous studies (Löfverström et al., 2014; Ullman et al., 2014; Pausata et al., 2011; Kageyama et al., 2020; Stadelmaier et al., 2024), which emphasize the influence of glacial ice sheet topography on jet stream reorganization. Our results further expand on this, demonstrating that the high-pressure system induced by the LIS initiates a Rossby wave response, shifting the Icelandic low southward and weakening the Azores High. These circulation changes lead to reduced precipitation over NH landmasses during DJF.

During JJA, the changes in westerlies and pressure result in increased rainfall over Eurasia (Figure 9b), with the jet stream extending further inland. Additionally, this rainfall increase is supported by milder land temperatures and enhanced moisture availability (not shown). Our findings on seasonal precipitation changes are consistent with those of Löfverström et al. (2014), as the ice sheet configuration used in our study closely resembles their MIS 4 ice sheet setup. However, while their study reports a ~~full~~ fully zonal orientation of the jet stream only during the LGM, our results demonstrate that this change can also occur with a MIS 3 ice sheet configuration, suggesting a dynamic response to glacial ice sheets earlier in the glacial period.

Changes in tropical atmospheric circulation and SH westerlies

The presence of ice sheet topography leads to a more pronounced southward shift of the NH Hadley cell during DJF and a stronger northward shift of the SH Hadley cell during JJA, compared to our simulations that include only GHG concentrations, orbital parameters, and an ice mask at 49 ka (Figure 9a,b). This enhanced Hadley cell shift in 49ka-full drives a more substantial southward displacement of the ITCZ during DJF (Figure 9a). Consequently, the northern tropics experience drier conditions, while precipitation increases in the southern tropics, consistent with the mechanisms proposed by Chiang and Bitz (2005).

During JJA, LIS topography induces warming over Siberia, in agreement with previous studies (Liakka and Lofverstrom, 2018; Bakker et al., 2020), which attribute this warming to large-scale atmospheric circulation changes driven by pressure differences linked to LIS height. In contrast, AIS topography enhances cooling over Antarctica during austral winter. This interhemispheric contrast, i.e. relatively warmer conditions in the NH and colder conditions in the SH, pushes the ITCZ northward and strengthens the SH Hadley cell. Additionally, the SH Hadley cell northward shift is associated with the northward displacement of the SH westerlies (Figure 9b), consistent with Ceppi et al. (2013), who demonstrated that interhemispheric teleconnections link ITCZ migration with changes in Hadley cell strength and SH jet positioning. Likewise, NH cooling during DJF drives a southward shift of the North Atlantic westerlies, influencing the Hadley cell strength and position, and pushing the ITCZ closer to the equator.

These circulation shifts generate regional impacts. Increased subsidence over East Asia creates a high-pressure anomaly, leading to drier conditions, while the southward-shifted ITCZ enhances convergence over Australia, increasing rainfall. The larger shift in the ITCZ and the NH Hadley cell coincides with the maximum increase in Australian rainfall (Figure 9a).

The weakening of the SH westerlies during both DJF and JJA under a high-obliquity climate (49ka-co and 49ka-alb) compared to a low-obliquity climate (PI) aligns with findings from Timmermann et al. (2014). However, the increased height and extent of the AIS counteract this weakening during DJF, leading to SH westerly intensification in 49ka-full. Additionally, our




Figure 9. Summary of key atmospheric responses across experiments. (a) DJF anomalies: (left) NH SAT (°C); (middle) latitudinal shifts (°) in westerlies (calculated as the latitude of the maximum wind speed averaged over the North Atlantic (80°W:0°E, 30°N:90°N)), NH Hadley cell's southern edge, and globally averaged ITCZ position; and (right) northern Australian precipitation (116°E:152°E, 25°S:12°S, mm/day). (b) JJA anomalies: (left) NH SAT (°C); (middle) latitudinal shifts (°) in westerlies averaged over the North Atlantic, SH Hadley cell's southern edge (°), westerlies averaged over the SH; and (right) precipitation changes over Eurasia (60°W:80°E, 48°N:60°N). Results are shown for the 49ka-co (cyan), 49ka-alb (orange), and 49ka-full (blue) experiments, with all values presented as anomalies relative to PI. Negative values in latitudinal shifts indicate an equatorward shift.

results indicate that NH westerlies strengthen with albedo changes relative to orbital parameter and GHG adjustments, whereas SH westerlies weaken under the same conditions (Figure 4). Introducing topographical changes induces an equatorward shift in both NH and SH westerlies during their respective winter seasons (Figure 4a,d).

Overall, our study reveals nonlinear interactions between large-scale atmospheric circulation, surface westerlies, and precipitation patterns under different ~~climate combinations of~~ boundary conditions (Figure 9). Orbital forcing, GHGs, albedo, and ice sheet topography collectively shape these interactions in a nonlinear manner.

MIS 3 hydroclimate in Australia

Proxy records from marine sediments, lake sediments, and river systems, show that Australia experienced spatially variable climates during MIS 3, with greater-than-modern mean water availability peaking between ~49 and ~40 ka (Kemp et al., 2019). Wet conditions, especially in central and northern Australia between ~50 and ~45 ka, are linked to an intensified Australian monsoon driven by the summer insolation peak. Our model results align with these observations, showing increased annual precipitation over much of Australia at 49 ka, with particularly intensified monsoonal rainfall in the northern regions (Figure 2c, 3g). ~~This increase in rainfall~~ The increase in monsoonal rainfall during austral summer is consistent with a southward displacement of the ITCZ due to the intensified NH Hadley cell, driven by increased SH insolation ~~during in response to~~ high obliquity. However, our results also reveal that the southward shift of the ITCZ and Hadley cell ~~occur~~ occurs in the 49ka-full simulation (which includes ice sheet topography) as well as in the 49ka-co and 49ka-alb simulations (which lack ice sheet topography). Despite this, the intensified Australian monsoon rainfall is more pronounced in the 49ka-full experiment, as the presence of ice sheets induces larger shifts of the Hadley cell and ITCZ.

Our simulations generally show an insignificant response in southern ~~Australia~~ Australian annual rainfall at 49 ka, ~~with some drying, which contrasts with the increased rainfall seen in southern Australia indicated by paleo-records from a variety of archives, including speleothems, lake sediments, river systems, and offshore sediments (Kemp et al., 2019). In their study, the strengthening or northward movement of the southwesterly winds is linked to increased rainfall in southern Australia, a response that does not align with our simulation. However, our results are somewhat consistent with Weij et al. (2024), although austral winter exhibits some drier conditions. These results are consistent with those of Weij et al. (2024), in which speleothem growth rates from southwest and southeast Australia indicate drier-than-modern conditions around 49 ka-ka. In contrast, paleo records from Kemp et al. (2019) suggest increased rainfall in southern Australia. The wetter signal in that study is interpreted as a response to changes in the SH westerlies, involving either strengthening or an equatorward shift. While the wind response is consistent with the equatorward shift simulated in the 49ka-full experiment, the rainfall response is not reproduced, indicating that southern Australian precipitation is not solely controlled by latitudinal shifts of the westerlies in ACCESS-ESM1.5. Instead, rainfall anomalies appear to be more strongly influenced by persistent subtropical high-pressure anomalies and associated subsidence over the Australian continent, which suppress moisture transport convergence despite the shift in westerlies. This may indicate limitations in the simulated coupling between storm-track variability and regional precipitation under glacial boundary conditions. However, the contrasting interpretations from available proxy records also suggest that hydroclimate conditions in southern Australia during MIS 3 remain uncertain~~

5 Conclusions

Our results ~~underscore the far-reaching effects~~, based on a stepwise set of MIS 3 ~~ice sheets~~ boundary condition experiments, ~~underscore the impacts of orbital forcing, greenhouse gases, surface albedo, and ice-sheet topography~~ on global atmospheric circulation and regional hydroclimate. ~~We demonstrate that the~~ Orbital parameters and greenhouse gases alone produce limited dynamical shifts, while the addition of ice-sheet albedo primarily enhances high-latitude cooling without substantially altering circulation patterns. In contrast, the presence of NH ice sheets during MIS 3 ~~strengthened and shifted~~ strengthens and shifts the North Atlantic westerlies equatorward by $\sim 6^\circ$ in DJF ~~and 4° in JJA~~, leading to increased precipitation over the North Atlantic ~~and Eurasian landmass in both seasons, while over the Eurasian landmass precipitation increases~~ in boreal summer ~~but decreases in winter due to colder conditions~~. These ice sheet-induced changes ~~extend meridionally~~ affect the global atmospheric circulation, with a southward shift of the NH Hadley cell and ITCZ during DJF, resulting in ~~increased Australian rainfall~~, enhanced monsoonal rainfall over northern Australia, while producing a comparatively weak hydroclimatic response in southern Australia. Regional differences, particularly between northern and southern Australia, reflect contrasting dynamical controls, with northern Australia dominated by monsoon and ITCZ-related moisture convergence, and southern Australia more strongly influenced by subtropical high-pressure anomalies and associated subsidence. These patterns are broadly consistent with proxy evidence indicating spatially heterogeneous hydroclimate conditions during MIS 3, with records suggesting both wetter and drier conditions across different regions and archives.

Although our experiments do not isolate the individual roles of the Laurentide and AISs, ~~simulations with MIS 3 topography suggest they show~~ that the combined ice sheet configuration ~~also~~ contributes to an equatorward shift of the SH westerlies and SH Hadley cell. Overall, our findings ~~underscore~~ highlight the importance of ~~including realistic MIS the LIS in~~ modulating atmospheric circulation and climate dynamics, noting that its geometry and extent varied substantially during MIS 3 ~~boundary conditions when assessing ice sheet-atmosphere interactions on atmospheric circulation and the deglaciation, likely contributing to diverse AMOC responses and associated climate changes~~.

Data availability. All final data from the modelling simulations will be published on the DRYAD repository under the corresponding author's account (<https://datadryad.org/search?utf8=%E2%9C%93&q=himadri+saini>) and made publicly available upon acceptance of the manuscript.

Appendix A




Figure A1. Time series of anomalies of annual mean (left) globally, (middle) southern hemisphere (SH), and (right) northern hemisphere (NH) averaged surface air temperature (SAT) and sea surface temperature (SST) for (black) PI, (cyan) 49ka-co, (orange) 49ka-alb, and (blue) 49ka-full compared to PI. 49ka-co is orbital parameters+GHGs, 49ka-alb is 49ka-co with albedo and vegetation changes. Additional changes to 49ka-alb are shown in magenta colors (49ka-alb+). At year 547, surface salinity is gradually increased to account for the 44 m sea level drop relative to PI, based-on-to account for the prescribed-increase of continental ice sheet mass, achieving a global average increase of +0.33 psu from PI levels. At year 729, the Bering Strait is closed. ~~These experiments are~~ The 49ka-alb+ experiment is not analysed in the manuscript. At year 824, the 49 ka topography is implemented. River runoff is adjusted to reflect the 49 ka topography at year 1258. Darker colors indicate the time-averaged-period-for-each-experiment-periods used in-for the analysis of time averages from each experiment. The horizontal red line is the zero line for each diagnostic shown.




Figure A2. Insolation (Shortwave downward radiation at the top of the atmosphere) anomalies (W/m^2) between 49 ka and PI as a function of latitude and month.




Figure A3. (top) Winter mixed layer depth (m), (middle) Zonal winds (m/s) at 850 hPa, and (bottom) North Atlantic gyre strength ([calculated from the barotropic streamfunction](#), N/m²) for (left) PI and (right) 49ka-full compared to PI. The white areas in the middle panel mark grid cells with continental ice, and for the 49ka-full simulation, continental outlines are shown based on the adjusted land-sea mask.




Figure A4. Zonal winds at 200 hPa (m/s) overlaid with 200 hPa wind vectors (m/s) in (left) DJF and (right) JJA for (top) PI and (middle) 49ka-full, and (bottom) 49ka-full minus PI (overlaid with 200 hPa wind vectors anomalies (m/s)).




Figure A5. Geopotential height (m) at 850 hPa with zonal mean removed and overlaid with 850 hPa wind vectors in (left) DJF and (right) JJA for (top) PI and (bottom) 49ka-full.

Experiments	PI	49ka-full	49ka-alb	49ka-ie 49ka-co
Annual mean SAT (°C)	15.41	12.71 (2.7 -2.7)	13.11 (-2.3)	13.05 (-2.36)
DJF NH SAT (°C)	-4.75	-11.15 (6.4 -6.4)	-9.25 (-4.5)	-7.75 (-3)
JJA NH SAT (°C)	14.95	12.35 (2.6 -2.6)	12.15 (-2.8)	14.25 (-0.7)
DJF SH SAT (°C)	2.85	1.15 (1.7 -1.7)	2.15 (-0.7)	1.95 (-0.9)
JJA SH SAT (°C)	-4.85	-7.25 (2.4 -2.4)	-6.45 (-1.6)	-6.45 (-1.6)
Annual mean Ocean Temp (°C)	4.32	3.3 (1.02 -1.02)	3.49 (-0.83)	3.62 (-0.7)
Annual mean SST (°C)	18.95	17.75 (1.2 -1.2)	17.65 (-1.3)	17.75 (-1.2)

Table A1. Global mean values for PI, 49ka-full, 49ka-alb, and ~~49ka-ie~~49ka-co experiments. The values in the brackets show differences from PI.

Author contributions. HS, LM, JRB, and RND designed the study. HS performed the modelling simulations with assistance from DKH. HS conducted the analysis, interpreted the results and wrote the manuscript with input from LM. JRB, RND, DKH and YD provided valuable
475 comments on the manuscript.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Climate of the Past

Acknowledgements. Himadri Saini, Josephine R. Brown, Russell N. Drysdale, Yanxuan Du, and Laurie Menviel would like to acknowledge funding from the Australian Research Council (ARC) grant DP220102134. Laurie Menviel acknowledges support from ARC grant SR200100008. David K. Hutchinson acknowledges support from ARC grant DE220100279. Josephine R. Brown received support from
480 ARC Centre of Excellence for Weather of the 21st Century (CE230100012).

This research was supported by the Australian Government's National Collaborative Research Infrastructure Strategy (NCRIS), with access to computational resources provided by the National Computational Infrastructure (NCI) through the National Computational Merit Allocation Scheme. The authors thank Commonwealth Scientific and Industrial Research Organisation (CSIRO) for developing the ACCESS-ESM1.5 model configuration and making it freely available to researchers. This research used the ACCESS-ESM1.5 model infrastructure provided by ACCESS-NRI, which is enabled by the Australian Government's National Collaborative Research Infrastructure Strategy (NCRIS).
485 The high performance computing (HPC) resources required to run the model are available through UNSW Resource Allocation Scheme doi:10.26190/PMN5-7J50.

References

- Allen, J. R., Forrest, M., Hickler, T., Singarayer, J. S., Valdes, P. J., and Huntley, B.: Global vegetation patterns of the past 140,000 years, *Journal of Biogeography*, 47, 2073–2090, 2020.
- Bakker, P., Rogozhina, I., Merkel, U., and Prange, M.: Hypersensitivity of glacial summer temperatures in Siberia, *Climate of the Past*, 16, 371–386, 2020.
- Berger, A.: Long-term variations of caloric insolation resulting from the Earth’s orbital elements, *Quaternary research*, 9, 139–167, 1978.
- Böhm, E., Lippold, J., Gutjahr, M., Frank, M., Blaser, P., Antz, B., Fohlmeister, J., Frank, N., Andersen, M., and Deininger, M.: Strong and deep Atlantic meridional overturning circulation during the last glacial cycle, *Nature*, 517, 73–76, 2015.
- Braconnot, P., Otto-Bliesner, B., Harrison, S., Joussaume, S., Peterchmitt, J.-Y., Abe-Ouchi, A., Crucifix, M., Driesschaert, E., Fichefet, T., Hewitt, C., et al.: Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum–Part 2: feedbacks with emphasis on the location of the ITCZ and mid-and high latitudes heat budget, *Climate of the Past*, 3, 279–296, 2007.
- Brandefelt, J., Kjellström, E., Näslund, J.-O., Strandberg, G., Voelker, A., and Wohlfarth, B.: A coupled climate model simulation of Marine Isotope Stage 3 stadial climate, *Climate of the Past*, 7, 649–670, 2011.
- Brayshaw, D. J., Hoskins, B., and Blackburn, M.: The basic ingredients of the North Atlantic storm track. Part I: Land–sea contrast and orography, *Journal of the Atmospheric Sciences*, 66, 2539–2558, 2009.
- Broccoli, A. and Manabe, S.: The influence of continental ice, atmospheric CO₂, and land albedo on the climate of the last glacial maximum, *Climate dynamics*, 1, 87–99, 1987.
- Brown, N. and Galbraith, E. D.: Hosed vs. unhosed: interruptions of the Atlantic Meridional Overturning Circulation in a global coupled model, with and without freshwater forcing, *Climate of the Past*, 12, 1663–1679, 2016.
- Byrne, M. P. and O’Gorman, P. A.: Land–ocean warming contrast over a wide range of climates: Convective quasi-equilibrium theory and idealized simulations, *Journal of Climate*, 26, 4000–4016, 2013.
- Calvo, E., Pelejero, C., Herguera, J. C., Palanques, A., and Grimalt, J. O.: Insolation dependence of the southeastern Subtropical Pacific sea surface temperature over the last 400 kyrs, *Geophysical Research Letters*, 28, 2481–2484, 2001.
- Caniupán, M., Lamy, F., Lange, C. B., Kaiser, J., Arz, H., Kilian, R., Baeza Urrea, O., Aracena, C., Hebbeln, D., Kissel, C., et al.: Millennial-scale sea surface temperature and Patagonian Ice Sheet changes off southernmost Chile (53 S) over the past 60 kyr, *Paleoceanography*, 26, 2011.
- Ceppi, P., Hwang, Y.-T., Liu, X., Frierson, D. M., and Hartmann, D. L.: The relationship between the ITCZ and the Southern Hemispheric eddy-driven jet, *Journal of Geophysical Research: Atmospheres*, 118, 5136–5146, 2013.
- Chiang, J. C. and Bitz, C. M.: Influence of high latitude ice cover on the marine Intertropical Convergence Zone, *Climate Dynamics*, 25, 477–496, 2005.
- Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., Mitrovica, J. X., Hostetler, S. W., and McCabe, A. M.: The Last Glacial Maximum, *Science*, 325, 710–714, 2009.
- Craig, A., Valcke, S., and Coquart, L.: Development and performance of a new version of the OASIS coupler, OASIS3-MCT_3. 0, *Geoscientific Model Development*, 10, 3297–3308, 2017.
- Duboc, B., Meissner, K. J., Menviel, L., Yeung, N. K., Hoogakker, B., Ziehn, T., and Chamberlain, M.: Simulated ocean oxygenation during the interglacials MIS 5e and MIS 9e, *Climate of the Past*, 21, 1093–1122, 2025.
- EPICA community members: One-to-one coupling of glacial climate variability in Greenland and Antarctica, *Nature*, 444, 195–198, 2006.

- 525 Feldberg, M. J. and Mix, A. C.: Planktonic foraminifera, sea surface temperatures, and mechanisms of oceanic change in the Peru and south equatorial currents, 0–150 ka BP, *Paleoceanography*, 18, 2003.
- Gowan, E. J., Zhang, X., Khosravi, S., Rovere, A., Stocchi, P., Hughes, A. L., Gyllencreutz, R., Mangerud, J., Svendsen, J.-I., and Lohmann, G.: A new global ice sheet reconstruction for the past 80 000 years, *Nature communications*, 12, 1199, 2021.
- Griem, L., Voelker, A. H., Berben, S. M., Dokken, T. M., and Jansen, E.: Insolation and glacial meltwater influence on sea-ice and circulation
530 variability in the northeastern Labrador Sea during the Last Glacial period, *Paleoceanography and Paleoclimatology*, 34, 1689–1709, 2019.
- Griffies, S. M.: Elements of the modular ocean model (MOM), GFDL Ocean Group Tech. Rep, 7, 47, 2012.
- Guo, C., Nisancioglu, K. H., Bentsen, M., Bethke, I., and Zhang, Z.: Equilibrium simulations of marine isotope stage 3 climate, *Climate of the Past*, 15, 1133–1151, 2019.
- Henry, L., McManus, J., Curry, W., Roberts, N., Piotrowski, A., and Keigwin, L.: North Atlantic ocean circulation and abrupt climate change
535 during the last glaciation, *Science*, 353, 470–474, 2016.
- Hu, A., Meehl, G. A., Han, W., Otto-Blietner, B., Abe-Ouchi, A., and Rosenbloom, N.: Effects of the Bering Strait closure on AMOC and global climate under different background climates, *Progress in Oceanography*, 132, 174–196, 2015.
- Huber, C., Leuenberger, M., Spahni, R., Flückiger, J., Schwander, J., Stocker, T. F., Johnsen, S., Landais, A., and Jouzel, J.: Isotope calibrated Greenland temperature record over Marine Isotope Stage 3 and its relation to CH₄, *Earth and Planetary Science Letters*, 243, 504–519,
540 2006.
- Hunke, E. C., Lipscomb, W. H., Turner, A. K., Jeffery, N., and Elliott, S.: Cice: the los alamos sea ice model documentation and software user’s manual version 4.1 la-cc-06-012, T-3 Fluid Dynamics Group, Los Alamos National Laboratory, 675, 500, 2010.
- Izumi, K., Valdes, P., Ivanovic, R., and Gregoire, L.: Impacts of the PMIP4 ice sheets on Northern Hemisphere climate during the last glacial period, *Climate Dynamics*, 60, 2481–2499, 2023.
- 545 Kageyama, M., Albani, S., Braconnot, P., Harrison, S. P., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Marti, O., Peltier, W. R., Peterschmitt, J.-Y., et al.: The PMIP4 contribution to CMIP6–Part 4: Scientific objectives and experimental design of the PMIP4-CMIP6 Last Glacial Maximum experiments and PMIP4 sensitivity experiments, *Geoscientific Model Development*, 10, 4035–4055, 2017.
- Kageyama, M., Harrison, S. P., Kapsch, M.-L., Löfverström, M., Lora, J. M., Mikolajewicz, U., Sherriff-Tadano, S., Vadsaria, T., Abe-Ouchi, A., Bouttes, N., et al.: The PMIP4-CMIP6 Last Glacial Maximum experiments: preliminary results and comparison with the
550 PMIP3-CMIP5 simulations, *Climate of the Past Discussions*, 2020, 1–37, 2020.
- Kaiser, J., Lamy, F., and Hebbeln, D.: A 70-kyr sea surface temperature record off southern Chile (Ocean Drilling Program Site 1233), *Paleoceanography*, 20, 2005.
- Kemp, C., Tibby, J., Arnold, L., and Barr, C.: Australian hydroclimate during Marine Isotope Stage 3: a synthesis and review, *Quaternary Science Reviews*, 204, 94–104, 2019.
- 555 Kindler, P., Guillevic, M., Baumgartner, M., Schwander, J., Landais, A., and Leuenberger, M.: Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core, *Climate of the Past*, 10, 887–902, 2014.
- Klockmann, M., Mikolajewicz, U., and Marotzke, J.: Two AMOC states in response to decreasing greenhouse gas concentrations in the coupled climate model MPI-ESM, *Journal of Climate*, 31, 7969–7984, 2018.
- Köhler, P., Nehrbass-Ahles, C., Schmitt, J., Stocker, T. F., and Fischer, H.: A 156 kyr smoothed history of the atmospheric greenhouse gases
560 CO₂, CH₄, and N₂O and their radiative forcing, *Earth System Science Data*, 9, 363–387, 2017.

- Kowalczyk, E., Stevens, L., Law, R., Dix, M., Wang, Y., Harman, I., Haynes, K., Srbinovsky, J., Pak, B., and Ziehn, T.: The land surface model component of ACCESS: description and impact on the simulated surface climatology, *Australian Meteorological and Oceanographic Journal*, 63, 65–82, 2013.
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambridge, M.: Sea level and global ice volumes from the Last Glacial Maximum to the Holocene, *Proceedings of the National Academy of Sciences*, 111, 15 296–15 303, 2014.
- Lauterbach, S., Andersen, N., Wang, Y. V., Blanz, T., Larsen, T., and Schneider, R. R.: An ~ 130 kyr record of surface water temperature and $\delta^{18}\text{O}$ from the northern Bay of Bengal: Investigating the linkage between Heinrich events and Weak Monsoon Intervals in Asia, *Paleoceanography and Paleoclimatology*, 35, e2019PA003 646, 2020.
- Lea, D. W., Pak, D. K., and Spero, H. J.: Climate impact of late Quaternary equatorial Pacific sea surface temperature variations, *science*, 289, 1719–1724, 2000.
- Liakka, J. and Lofverstrom, M.: Arctic warming induced by the Laurentide Ice Sheet topography, *Climate of the Past*, 14, 887–900, 2018.
- Liu, Y., Lo, L., Shi, Z., Wei, K.-Y., Chou, C.-J., Chen, Y.-C., Chuang, C.-K., Wu, C.-C., Mii, H.-S., Peng, Z., et al.: Obliquity pacing of the western Pacific Intertropical Convergence Zone over the past 282,000 years, *Nature communications*, 6, 10 018, 2015.
- Löfverström, M., Caballero, R., Nilsson, J., and Kleman, J.: Evolution of the large-scale atmospheric circulation in response to changing ice sheets over the last glacial cycle, *Climate of the Past*, 10, 1453–1471, 2014.
- Mackallah, C., Chamberlain, M., Law, R., Dix, M., Ziehn, T., Bi, D., Bodman, R., Brown, J., Dobrohotoff, P., Druken, K., et al.: ACCESS datasets for CMIP6: methodology and idealised experiments, *Journal of Southern Hemisphere Earth Systems Science*, 72, 93–116, 2022.
- Malmierca-Vallet, I. and Sime, L. C.: Dansgaard–Oeschger events in climate models: review and baseline Marine Isotope Stage 3 (MIS3) protocol, *Climate of the Past*, 19, 915–942, 2023.
- Manabe, S. and Broccoli, A.: The influence of continental ice sheets on the climate of an ice age, *Journal of Geophysical Research: Atmospheres*, 90, 2167–2190, 1985.
- Mantsis, D. F., Lintner, B. R., Broccoli, A. J., Erb, M. P., Clement, A. C., and Park, H.-S.: The response of large-scale circulation to obliquity-induced changes in meridional heating gradients, *Journal of Climate*, 27, 5504–5516, 2014.
- Martin, G., Milton, S., Senior, C., Brooks, M., Ineson, S., Reichler, T., and Kim, J.: Analysis and reduction of systematic errors through a seamless approach to modeling weather and climate, *Journal of Climate*, 23, 5933–5957, 2010.
- Martínez, I., Keigwin, L., Barrows, T. T., Yokoyama, Y., and Southon, J.: La Niña-like conditions in the eastern equatorial Pacific and a stronger Choco jet in the northern Andes during the last glaciation, *Paleoceanography*, 18, 2003.
- Martrat, B., Grimalt, J. O., Shackleton, N. J., de Abreu, L., Hutterli, M. A., and Stocker, T. F.: Four climate cycles of recurring deep and surface water destabilizations on the Iberian margin, *science*, 317, 502–507, 2007.
- Menviel, L. C., Skinner, L. C., Tarasov, L., and Tzedakis, P. C.: An ice–climate oscillatory framework for Dansgaard–Oeschger cycles, *Nature Reviews Earth & Environment*, 1, 677–693, 2020.
- Merkel, U., Prange, M., and Schulz, M.: ENSO variability and teleconnections during glacial climates, *Quaternary Science Reviews*, 29, 86–100, 2010.
- Muglia, J. and Schmittner, A.: Glacial Atlantic overturning increased by wind stress in climate models, *Geophysical Research Letters*, 42, 9862–9868, 2015.
- Oke, P. R., Griffin, D. A., Schiller, A., Matear, R., Fiedler, R., Mansbridge, J., Lenton, A., Cahill, M., Chamberlain, M., and Ridgway, K.: Evaluation of a near-global eddy-resolving ocean model, *Geoscientific model development*, 6, 591–615, 2013.

- Pahnke, K., Zahn, R., Elderfield, H., and Schulz, M.: 340,000-year centennial-scale marine record of Southern Hemisphere climatic oscillation, *Science*, 301, 948–952, 2003.
- 600 Pausata, F. S., Li, C., Wettstein, J., Kageyama, M., and Nisancioglu, K. H.: The key role of topography in altering North Atlantic atmospheric circulation during the last glacial period, *Climate of the Past*, 7, 1089–1101, 2011.
- Peltier, W. R.: Global glacial isostasy and the surface of the ice-age Earth: the ICE-5G (VM2) model and GRACE, *Annu. Rev. Earth Planet. Sci.*, 32, 111–149, 2004.
- Saini, H., Meissner, K. J., Menviel, L., and Kvale, K.: Transient response of Southern Ocean ecosystems during Heinrich stadials, *Paleo-*
605 *ceanography and Paleoclimatology*, 39, e2023PA004 754, 2024.
- Seager, R., Battisti, D. S., Yin, J., Gordon, N., Naik, N., Clement, A. C., and Cane, M. A.: Is the Gulf Stream responsible for Europe’s mild winters?, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 128, 2563–2586, 2002.
- Shakun, J. D., Lea, D. W., Lisiecki, L. E., and Raymo, M. E.: An 800-kyr record of global surface ocean $\delta^{18}\text{O}$ and implications for ice
610 volume-temperature coupling, *Earth and Planetary Science Letters*, 426, 58–68, 2015.
- Sherriff-Tadano, S., Abe-Ouchi, A., Yoshimori, M., Oka, A., and Chan, W.-L.: Influence of glacial ice sheets on the Atlantic meridional overturning circulation through surface wind change, *Climate dynamics*, 50, 2881–2903, 2018.
- Stadella, K. H., Ludwig, P., Pinto, J. G., and Újvári, G.: Changes in Atmospheric Dynamics Over Dansgaard-Oeschger Climate Oscillations Around 40 ka and Their Impact on Europe, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD040 247, 2024.
- 615 The HadGEM2 Development Team, Bellouin, N., Collins, W., Culverwell, I., Halloran, P., Hardiman, S., Hinton, T., Jones, C., McDonald, R., McLaren, A., O’Connor, F., et al.: The HadGEM2 family of met office unified model climate configurations, *Geoscientific Model Development*, 4, 723–757, 2011.
- Timmermann, A., Friedrich, T., Timm, O. E., Chikamoto, M. O., Abe-Ouchi, A., and Ganopolski, A.: Modeling obliquity and CO₂ effects on Southern Hemisphere climate during the past 408 ka, *Journal of Climate*, 27, 1863–1875, 2014.
- 620 Ullman, D., LeGrande, A., Carlson, A. E., Anslow, F., and Licciardi, J.: Assessing the impact of Laurentide Ice Sheet topography on glacial climate, *Climate of the Past*, 10, 487–507, 2014.
- Van Meerbeeck, C. J., Renssen, H., and Roche, D. M.: How did Marine Isotope Stage 3 and Last Glacial Maximum climates differ?—perspectives from equilibrium simulations, *Climate of the Past*, 5, 33–51, 2009.
- WAIS Divide Project Members: Precise interglacial phasing of abrupt climate change during the last ice age, *Nature*, 520, 661–665, 2015.
- 625 Wang, N., Jiang, D., and Lang, X.: Northern westerlies during the Last Glacial Maximum: Results from CMIP5 simulations, *Journal of Climate*, 31, 1135–1153, 2018.
- Wang, Y., Law, R., and Pak, B.: A global model of carbon, nitrogen and phosphorus cycles for the terrestrial biosphere, *Biogeosciences*, 7, 2261–2282, 2010.
- Weij, R., Sniderman, J. K., Woodhead, J. D., Hellstrom, J. C., Brown, J. R., Drysdale, R. N., Reed, E., Bourne, S., and Gordon, J.: Elevated
630 Southern Hemisphere moisture availability during glacial periods, *Nature*, 626, 319–326, 2024.
- Yeung, N. K.-H., Menviel, L., Meissner, K. J., Taschetto, A. S., Ziehn, T., and Chamberlain, M.: Land–sea temperature contrasts at the Last Interglacial and their impact on the hydrological cycle, *Climate of the Past*, 17, 869–885, 2021.
- Zhang, P., Xu, J., Holbourn, A., Kuhnt, W., Xiong, Z., and Li, T.: Obliquity induced latitudinal migration of the intertropical convergence zone during the past 410 kyr, *Geophysical Research Letters*, 49, e2022GL100 039, 2022.

- 635 Zhang, X., Lohmann, G., Knorr, G., and Purcell, C.: Abrupt glacial climate shifts controlled by ice sheet changes, *Nature*, 512, 290–294, 2014a.
- Zhang, X., Prange, M., Merkel, U., and Schulz, M.: Instability of the Atlantic overturning circulation during Marine Isotope Stage 3, *Geophysical Research Letters*, 41, 4285–4293, 2014b.
- Ziehn, T., Chamberlain, M. A., Law, R. M., Lenton, A., Bodman, R. W., Dix, M., Stevens, L., Wang, Y.-P., and Srbinovsky, J.: The Australian
640 earth system model: ACCESS-ESM1. 5 doi:10.22033/ESGF/CMIP6.4248, *Journal of Southern Hemisphere Earth Systems Science*, 70, 193–214, 2020.