

Projected climate change in Fennoscandia — and its relation to ensemble spread and global trends

Gustav Strandberg^{1,2}, August Thomasson^{1,3}, Lars Bärring¹, Erik Kjellström¹, Michael Sahlin¹, Renate Anna Irma Wilcke¹, Grigory Nikulin¹

5 ¹Rossby Centre, Swedish Meteorological and Hydrological Institute, Norrköping, SE-601 76, Sweden

²Centre for Climate Science and Policy Research, Linköping University, Linköping SE-581 83, Sweden

³Department of Physics, Lund University, Lund, SE-223 64, Sweden

Correspondence to: Gustav Strandberg (gustav.strandberg@smhi.se)

Abstract. The need for information about climate change is great. This information is usually based on climate model data, which often have systematic biases. Furthermore, climate information is based on ensembles of climate models, which raises the question about how such ensembles are affected by the choice of models and emission scenarios. Here, we aim to describe give a description of climate change in Sweden and neighbouring countries and discuss how local changes relate to global warming, as well as a discussion on how local climate change relates to global warming. We present climate change projections based on bias adjusted Euro-CORDEX (Coordinated Regional Downscaling Experiment) regional climate model data centred over Sweden. Global warming results in higher temperature, more warm days, and fewer cold days in Sweden. The regional climate models replicate the signal of the driving global models. Yet, the model spread is smaller than in the full CMIP5 ensemble, which means that the RCMs do not fully represent the potential model spread. The choice of emission scenario has minimal effect on the calculation of mean climate change at a global warming level of 2 degrees. This implies that it would be safe to mix emission scenarios in calculations of global warming levels, at least up to +2°C, and as long as mean values are concerned. Moreover, the differences in local and global warming rates seem to decrease with time, suggesting that climate change in Sweden may currently be at its fastest.

1 Introduction

Unless strong reductions in greenhouse gas emissions of greenhouse gases are implemented, global warming is likely to reach +2 °C above compared to pre-industrial levels within the 21st century (Forster et al.,

25 2024). The temperature response in Europe correlates is correlated with global temperature change but increases at a faster rate to, but warms at stronger rates, than the global temperature change (IPCC, 2021). Since 1850-1900, a the global temperature-increase of 1.3°C has translated into a warming of 2.3°C in Europe and 3.3°C in the Arctic (C3S, 2024).

The current rapid strong global warming calls for climate adaptation in all parts of society. Adaptation measures must be based on informed decisions to be cost efficient and to avoid maladaptation (IPCC, 2022). Thus, there is a great need for 30 climate data to support decision making and adaptation.

A way to avoid the discussion on which emission scenario to use and which scenario is the most likely — a discussion that is sometimes heated (Hausfather & Peters, 2020; Schwalm et al., 2020) — is ~~to apply to use~~ global warming levels (GWL). Instead of a fixed period of time in a certain scenario GWLs focus on the period when a particular level of global warming is reached. For example, GWL2 is the period when +2°C global warming is reached compared to pre-industrial times. This 35 period may occur at different times in different models — instead of consistency in time between the members of the ensemble there is thus a consistency in the magnitude of temperature increase. In that way, using GWLs is a powerful method since it is possible to mix simulations using different scenarios to create larger ensembles; and since it reduces the uncertainty around the choice of emission scenarios (Maule et al., 2017). One example of how to use GWLs for regional data is found in Strandberg et al. (2024b). The mixing of emission scenarios in GWLs can nevertheless be criticised because the 40 trends are different between scenarios (Bärring & Strandberg, 2018); a GWL based on RCP2.6 does not have the same characteristics as a GWL based on RCP8.5. This means that also a GWL ensemble is sensitive to how it is constructed with regards to which models and scenarios that are used as input. We want to investigate the robustness of the ensembles and how the simulated climate at a specific GWL is affected by the choice of emissions scenario, and global and regional models.

45 Climate models are our main tool ~~for to make~~ projecting ~~ings~~ ~~of~~ future climate change. Climate modelling is computationally expensive, which means that global climate models (GCMs) ~~are~~ usually run on relatively coarse horizontal resolutions (typically 100 - 300 km). ~~In contrast~~ ~~On the other hand~~, regional climate models (RCMs) can ~~be~~ run at higher resolutions (typically 5-20 km) ~~because~~ ~~since~~ they cover ~~smaller domains~~ ~~a smaller part of the globe~~. ~~As a result~~, ~~Therefore~~, RCMs can provide ~~additional~~ ~~new~~ information despite being governed by the driving GCM (e.g. Vautard et al., 2020; Strandberg & 50 Lind, 2020). Topographical features, such as coastlines or mountains, are better described with higher model resolution. Furthermore, RCM simulations ~~offer~~ ~~give~~ more details and a better representation of physical processes, especially local events like convective rain and short-duration extreme events (e.g. Olsson et al., 2015; Prein et al., 2015; Rummukainen, 2016; Lind et al., 2020).

CORDEX (Coordinated Regional climate Downscaling Experiment, Jacob et al., 2024) provides the most comprehensive 55 ~~high-resolution~~ RCM ensemble for Europe ~~on a high resolution~~. A key advantage of using climate model ensembles, like the CORDEX ensemble, is that they allow for a probabilistic assessment of potential changes, uncertainty estimations and a wider set of statistical tests (Déqué et al., 2012; Coppola et al., 2021). ~~By~~ ~~relying~~ on only one or very few model simulations ~~there is a risks~~ ~~sampling that you~~ only ~~sample~~ a small part of the possible outcomes. ~~Moreover, a~~ ~~single~~ ~~Furthermore, one~~ simulation is not enough to estimate model sensitivity to emissions of greenhouse gases, model 60 uncertainty, or natural variability (e.g. von Trentini et al., 2019; Christensen and Kjellström, 2020; 2021).

Since all parts of society are affected by climate change, it is crucial to have a well-founded description of it—
~~particularly especially~~ given the significant economic investments that will ~~rely be based~~ on climate projections. By “~~a good~~ description”,¹ we mean an ensemble that is both accurate and representative, and, not least, large enough to enable the

65 assessment of the significance and robustness of simulated climate change. In a~~Additionally~~, a general understanding of ensembles is necessary:~~i.~~ It is important to know how an ensemble's characteristics is shaped by the models and scenarios that compose it.

Here we present a new dynamically downscaled and bias~~–~~adjusted ensemble of climate projections for Sweden. Compared to the previous ensemble (Kjellström et al., 2016), improvements include higher horizontal resolution in the RCMs, bias 70 adjustment of ~~the~~ results, more ensemble members, and more indicators developed in dialogue with users to meet their needs. Climate model projections are~~is~~ an important tool for illustrating various aspects of climate change and its potential impacts on how it could impact society. These~~is~~ data is used to support decision~~–~~makers² in their work on climate ~~change~~ adaptation in Sweden. Rather than relying solely on standard climatological variables, inclusion of climate indicators in the assessment enables insights into~~regarding~~ impacts that are more directly relevant to society. These indicators aim to~~should~~ support climate adaptation~~–~~ by serving as decision support and informing the general public.

75 Since these data cover Fennoscandia and the Baltic States, they may also be applicable to surrounding countries. They are based on RCP (Representative Concentration Pathways) scenarios and CMIP5 (Coupled Model Intercomparison Project Phase 5; Taylor et al., 2022) global models. The Swedish climate service (SMHI, 2025) relies on these data, and at least~~at least~~ until a CMIP6-based downscaled ensemble becomes available, they will continue to be used. Since this RCM ensemble is already existing and used it is important to also discuss how the ensemble is constructed and how that influence the characteristics of the ensemble. ¶

80 The RCM ensemble presented here is already existing and used, making it. Therefore, it is important to also discuss how the ensemble is constructed and how that influence its~~the~~ characteristics~~—~~ of the ensemble; ~~as a~~ serving~~ee~~ to all users. This study addresses~~ims at~~ four main~~general~~ topics:

85 i) Projected climate change in Fennoscandia. This paper provides~~serves as~~ a general overview of projected climate change in Sweden based on the best available material, making it~~this~~ the currently most comprehensive projection for the region to date~~of climate change in the region~~ and a foundation~~basis~~ for further research and decision-making.

90 ii) How local trends in climate relate to global warming. Fennoscandia is known to have a warming trend that greatly exceeds the global trend, but still with a relatively linear relationship (C3S, 2024). It is, however, unknown whether~~if~~ this relationship will persist in the future.

95 iii) Model spread in the RCM ensemble compared to the spread of the~~the~~ larger CMIP5 ensemble. Since the RCM ensemble is forced by a sub-set of available GCMs, the model spread may be reduced, potentially resulting in a loss of information~~is potentially reduced. This would mean that information is lost in the RCM ensemble.~~

iv) The role of climate model and emission scenario selection in projected changes in temperature and precipitation at +2°C global warming. This is particularly important because the Paris Agreement (UNFCCC, 2015) aims to keep temperature rise well below 2 °C. Consequently, descriptions of projected climate change naturally focus on a two-degree warmer world.~~Since it is likely that global warming will reach +2 °C within~~

100 this century, and since the Paris Agreement (UNFCCC, 2015) speaks of keeping the temperature rise to well below 2 °C, it is natural that descriptions of projected climate change are formed around a two-degree warmer world. The question is how such ‘global warming levels’ are influenced by the climate models and emission scenarios used to calculate them.¶

iv)

2 Methods

105 2.1 The Euro-CORDEX ensemble

The presented data describing simulated present and future climates are based on the Euro-CORDEX ensemble covering Europe with a grid spacing of 0.11°, which approximately equals 12.5×12.5 km (Jacob et al., 2014). Within CORDEX several global climate models (GCMs) are used to force a number of regional climate models (RCMs). Every six hours the RCMs read data from the GCMs on the boundary of their model domains. These boundary conditions include temperature, 110 pressure, humidity and wind at ~~multiple~~several vertical levels, as well as sea surface temperature and sea ice conditions.

The Euro-CORDEX RCMs used ~~in this study~~here are forced by a subset of GCMs from ~~the~~CMIP5. The RCMs ~~have~~were ~~been~~ evaluated ~~against~~using observations and were judged to generally perform well in the historical climate of the late 20th century (Vautard et al., 2021). ~~However, t~~This does not mean that the CORDEX simulations are ~~free from~~without systematic errors. Vautard et al (2021) conclude that the simulations are generally too wet, too cold and too windy compared to 115 observations. Some of the discrepancies between GCMs and RCMs, as well as the weak warming trend, ~~may~~could be explained by ~~an overly-tee~~ simplified description of aerosol forcing (Boé et al., 2021; Katragkou et al., 2024). Projections for the 21st century from the RCMs have previously been assessed for Europe by Coppola et al. (2021). ¶

The simulations and their combinations of GCMs, RCMs and RCPs are listed in Table 1. As this study is based on an ~~already~~ existing ensemble ~~that is~~ already ~~in~~being used (SMHI, 2025), we have not ~~made any choices of~~ excluded ~~any~~ simulations.

120 ~~Adding or removing~~To add or exclude members would mean that we investigate another ensemble than the one used ~~in-for~~ example the SMHI climate service. ~~T~~When the ensemble was created ~~following~~it was created after a “the more the better”- approach, ~~which~~ meanings that as many simulations as possible were included~~are used~~.

Table 1 The simulations used in this study and the GCMs, RCMs and RCPs that they consist of. Members that are part of an ensemble consistent across all RCPs (RCM17) are marked with an ‘*’.

Driving GCM	GCM	No.	RCM	Scenario		
				RCP2.6	RCP4.5	RCP8.5
CNRM-CERFACS-	r1i1p1		CLMcom-ETH-COSMO-		X	
CNRM-CM5			crCLIM-v1-1	X	X	X *
			CNRM-ALADIN63			

		DMI-HIRHAM5		X	
		GERICS-REMO2015	X		X
		IPSL-WRF381P		X	
		KNMI-RACMO22E	X	X	X *
ICHEC-EC-EARTH	r1i1p1	CLMcom-ETH-COSMO-crCLIM-v1-1		X	
		DMI-HIRHAM5		X	
		KNMI-RACMO22E		X	X
		SMHI-RCA4		X	
	r3i1p1	CLMcom-ETH-COSMO-crCLIM-v1-1		X	
		DMI-HIRHAM5	X	X	X *
		KNMI-RACMO22E		X	
		SMHI-RCA4		X	
	r12i1p1	CLMcom-CCLM4-8-17	X	X	X *
		CLMcom-ETH-COSMO-crCLIM-v1-1		X	
		DMI-HIRHAM5		X	
		ICTP-RegCM4-6		X	
		GERICS-REMO2015	X	X	X *
		KNMI-RACMO22E	X	X	X *
		MOHC-HadREM3-GA7-05	X		X
		SMHI-RCA	X	X	X *
		IPSL-WRF381P		X	
IPSL-IPSL-CM5A-MR	r1i1p1	DMI-HIRHAM5		X	
		GERICS-REMO2015		X	
		KNMI-RACMO22E		X	
		SMHI-RCA4		X	X
		IPSL-INNERIS-WRF331P	X	X	
MIROC-MIROC5	r1i1p1	CLMcom-CCLM4-8-17	X		X

		GERICS-REMO2015	X	X	
MOHC-HadGEM2-ES	r1i1p1	CLMcom-CCLM4-8-17		X	X
		CLMcom-ETH-COSMO-crCLIM			X
		CNRM-ALADIN63			X
		DMI-HIRHAM5	X	X	X *
		GERICS-REMO2015	X	X	X *
		ICTP-RegCM4-6	X		X
		KNMI-RACMO22E	X	X	X *
		MOHC-HadREM3-GA7-05	X		X
		SMHI-RCA4	X	X	X *
		IPSL-WRF381P			X
MPI-M-MPI-ESM-LR	r1i1p1	CLMcom-CCLM4-8-17	X	X	X *
		CLMcom-ETH-COSMO-crCLIM-v1-1			X
		CNRM-ALADIN63			X
		DMI-HIRHAM5			X
		MPI-CSC-REMO2009	X	X	X *
		ICTP-RegCM4-6	X		X
		KNMI-RACMO22E	X		X
		MOHC-HadREM3-GA7-05			X
		SMHI-RCA4	X	X	X *
		IPSL-WRF381P			X
r2i1p1	CLMcom-ETH-COSMO-crCLIM-v1-1				X
		MPI-CSC-REMO2009	X	X	X *
		SMHI-RCA4			X
NCC-NorESM1-M	r1i1p1	CLMcom-ETH-COSMO-crCLIM-v1-1			X
		CNRM-ALADIN63			X
		DMI-HIRHAM5		X	X
		GERICS-REMO2015	X	X	X *
		ICTP-RegCM4-6	X		X
		KNMI-RACMO22E	X		X

MOHC-HadREM3-GA7-05		X		X
SMHI-RCA4	X	X	X	*
IPSL-WRF381P			X	

125

2.2 Bias adjustment

To minimise systematic errors, the Euro-CORDEX ensemble was bias-adjusted using the method “Multi-scale Bias Adjustment” method available in MIdAS (Berg et al., 2022). MIdAS is based on quantile mapping ‘day-of-year’ adjustments (Themeßl et al 2011; Wilcke et al., 2013), meaning. This means that the distribution used for to adjust the data is different for each day of the year. MIdAS aims is aiming to at preserv ing the trends in future projections and does performs similarly to methods that explicitly preserve trends (Berg et al., 2022). As reference data, the SMHI gridded climatology (SMHIGridClim) data set (SMHIGridClim; Andersson et al., 2021) was used. SMHIGridClim covers Fennoscandia and the Baltic states (region A in Fig 1), which means that the bias-adjusted ensemble covers a smaller domain centred over Sweden, instead of the entire European domain. The bias adjustment was applied made using the period 1980-2000 as a reference. The variables tas, tasmin, tasmax and pr (see Table 2 for explanations) were adjusted in all grid points within the domain. Hereafter, Below, any further mentions of the CORDEX RCMs refers to this bias-adjusted ensemble covering Fennoscandia and the Baltic states (region A in Fig 1).

2.3 Calculation of indicators

To assess climate change, a set of climate indicators are calculated using the software package Climix (Bärring et al., 2024). 140 A number of indicators were identified, building on the work of Kjellström et al (2016), and in collaboration together with the Swedish County Administrative Boards and other governmental agencies, to that can describe relevant changes in climate. The indicators are meant to be relevant for large parts of society, but agriculture (Strandberg et al., 2024a) and the energy sector (Strandberg et al., 2024b) have also been specifically targeted. The indicators used in this study are listed in Table 2. 145 The indicators are presented as averages for the 30-year periods used in the SMHI web service (SMHI, 2025): the reference period 1971-2000 and the future periods 2011-2040, 2041-2070, and 2071-2100. While WMO recommends 1961-1990 as the reference period for describing options of climate change (WMO, 2017), but since several RCM simulations begin in start 1971, making 1971-2000 a practical a compromise, is to use 1971-2000.

2.4 Definition of global warming levels

The GWLs are calculated for each driving GCM and based on the global mean surface temperature (GMST) using the period 1850-1900 as the a reference period, following the protocol in the IPCC-WG1 Atlas protocol (Iturbide et al., 2022). A GWL is reached when the GMST for a moving 20-year time window first exceeds for the first time passes that level. For

example: GWL2 occurs when the GMST for the first time is 2°C highermore than duringin the reference period. The timing of a GWL is represented by a central year. In this study we use 30-year periods for each GWL stretching from 15 years before the central year to 14 years after.¶

155 We analysed GWL1.5 and GWL2. GWL1.5 is reached in all scenarios, while GWL2 is reached in RCP4.5 and RCP8.5, but not in RCP2.6. Already at GWL3 most of the RCP4.5 simulations are excluded because they do not reach that level of warming. The limited number of scenarios and the smaller ensemble size makes GWL3 less interesting and less useful for this analysis. ¶

160

Table 2 Definitions and short names of indicators.

Indicator	Name	Definition	Unit
Average temperature	tas	The daily average temperatures	°C
Minimum temperature	tasmin	The daily minimum temperatures averaged over a selected period	°C
Maximum temperature	tasmax	The daily maximum temperatures averaged over a selected period	°C
Frost days	fd	Number of days with daily minimum temperature < 0°C	days
Summer days	su	Number of days with daily maximum temperature above 20°C	days
Consecutive summer days	csu	Longest period with consecutive days with daily maximum temperature above 20°C	days
Days with zero crossings	nzero	Number of days over with daily maximum temperature above 0°C and daily minimum temperature below 0°C	days
Precipitation	pr	Average precipitation amount	mm/ mon
Days with heavy precipitation	r10mm	Number of days with precipitation amount of more than 10 mm	days
Dry days	dd	Number of days with precipitation less than 1 mm	days

2.54 GCM ensembles

The bias adjusted CORDEX RCMs are compared to two GCM ensembles.

165 CORDEX GCMs: This ensemble consisting of the GCMs actually used to drive the RCMs (leftmost column in Table 1).
(Ensemble sizes are 5, 9 and 9 for scenarios RCP2.6, RCP4.5, and RCP8.5, respectively.). This ensemble includes several realisations for some GCMs since they are used to force RCMs.

CMIP5 GCMs: This ensemble includes consisting of all CMIP5 models available on the Earth System Grid Federation, but restricted to one realisation per GCM to avoid overweight on certain GCMs. Ensemble sizes are 24, 28 and 34 for scenarios RCP2.6, RCP4.5, and RCP8.5, respectively.)

170 The GCMs are not bias-adjusted. For all GCMs, the grid points falling within the Fennoscandian region (A in Figure 1) are used to calculate ensemble mean and spread for the region. For both GCM ensembles, the GMST values are calculated as 30-year averages for the reference period 1971-2000 and the future periods 2011-2040, 2041-2070, and 2071-2100, are
175 calculated¶

2.65 Selection and analyses of sub-ensembles

180 We aim to study the relative importance of the choice of RCP, GCM, and RCM at a specific GWL. To create a consistent ensemble across RCPs, we select only the combinations of GCMs and RCMs that simulated all three of the scenarios—RCP2.6, RCP4.5, and RCP8.5 (indicated with '*' in Table 1). From these 17 combinations of GCMs, RCMs, and RCPs (i.e., 51 RCM simulations), we construct sub-ensembles where all 17 members share the same RCP, GCM, or RCM. We refer to call this ensemble as RCM17. Using If we would use the full CORDEX RCM ensemble it would make it be difficult to separate the effects of different ensemble sizes and the effects of models or scenarios. The RCM17 ensemble
185 is used in section 3.5.

190 To illustrate the procedure, consider we make the hypothetical case with of three GCMs (GCM1-3) and three RCMs (RCM1-3) combined in different ways (Table 3). A sub-ensemble using only using GCM1 would include all RCMs forced by GCM1, i.e., the simulations in row R1 in Table 3 (i.e. three simulations). Similarly, In the same way the sub-ensemble based on GCM2 consists of two simulations. Sub-ensembles using only one RCM include use all simulations with that one RCM forced by different GCMs, i.e., one of the columns C1-3. For example, the sub-ensemble based on RCM1 has three simulations. Sub-ensembles based on a single emission one scenario include use all simulations run with that scenario.

Table 3 Hypothetical sketch of how three GCMs (GCM1-3) could be downscaled by three RCMs (RCM1-3) and how the sub-ensemble strategy works

	C1	C2	C3
	RCM1	RCM2	RCM3
R1	GCM1	X	X

R2	GCM2	X	X
R3	GCM3	X	

195

The GWLs are calculated based on the GMST using the period 1850–1900 as a reference, following the protocol in the IPCC-WG1 Atlas (Iturbide et al., 2022). A GWL is reached when the GMST for a moving 20-year time window for the first time passes that level. For example: GWL2 occurs when GMST for the first time is 2°C more than in the reference period.

200 The timing of a GWL is represented by a central year. In this study we use 30-year periods for each GWL stretching from 15 years before the central year to 14 years after. We analysed GWL1.5 and GWL2. GWL1.5 is reached in all scenarios, while GWL2 is reached in RCP4.5 and RCP8.5, but not in RCP2.6. Already at GWL3 most of the RCP4.5 are discarded since they do not reach that level of warming. The lack of different scenarios and the smaller ensemble size makes GWL3 a less interesting, and less useful, case for this analysis. ¶

205 Seven GCMs are combined in various ways in different ways combined with seven different RCMs, resulting in. This means that we have 2 RCP-based, 7 GCM-based, and 7 RCM-based sub-ensembles. In order to determine whether if any of the model combinations differs significantly different from the others, we perform two statistical tests under, with the null hypothesis that any given two ensembles have the same average. We use a one-way ANOVA (Analysis of variance) (Press, 1972) test, which tests whether two or more groups share have the same average or not. If the number of groups is equal to 2, as in which is the case of when we compare RCP-based ensembles, a one-way ANOVA is the same as a student's t-test. ¶

The ANOVA test does not indicate specify which sub-ensemble is different from the others, if any. Therefore, we apply a To find the different sub-ensemble(s) a post-hoc test: is used, the so-called Tukey's honestly significant difference (Tukey, 1949). This test is performed after a successful ANOVA test and compares all sub-ensembles pair-wise using with a 215 studentised q distribution. The tests are conducted done for two regions representing different climatic conditions in Sweden: one a region in the northern and one in the south Sweden and a region in southern Sweden representing different climatic conditions in Sweden (regions C and D in figure 1). A significance level with a family-wise error of 95 % is used, meaning. This means that the probability of one or more false positives among all points grid points cells is 5 % instead of a 5 % false positive rate in each individual grid point, if no correction is applied. The analyses were made for tas, csu, tasmin, 220 tasmax, pr, dd, cdd, su, r20mm and nzero.




Fig 1: Maps of regions used in analyses in this paper. A) Fennoscandian region (black, full line) is the domain on which bias adjustment is applied, B) Scandinavia (dash dotted orange line), C) northern Sweden (dotted blue line), D) southern Sweden (dashed red line).

225 3 Results and discussion

We begin here, we start by describing average climate changes according to the CORDEX RCM ensemble. To better understand these trends, we relate them they are then put in relation to the trend in GMST in the driving GCMs (CORDEX GCMs). This is followed by a comparison of the ensemble spread between CORDEX RCMs, CORDEX GCMs and a larger ensemble of CMIP5 GCMs to assess see how much of the potential spread that is lost by not using all available GCMs. 230 Section 3 is concluded by an investigation of how the description of a GWL based on the RCM17 ensemble is influenced by the GCMs, RCMs and RCPs of which it is constructed.

3.1 Projected change in temperature and temperature-based indicators




Figure 2: Ensemble mean temperatures of the CORDEX RCMs (°C) in winter (DJF, first column), spring (MAM, second column), summer (JJA, third column) and autumn (SON, fourth column). First row shows absolute values for 1971-2000. Rows 2 to 4 show anomalies from 1971-2000 to 2071-2100 according to scenarios RCP2.6, RCP4.5 and RCP8.5, respectively.

240 The mean temperatures (tas) are projected to increase in all seasons and underin all emission scenarios across the domain (Fig. 2). By the end of the century, t~~The increase in~~ annual mean temperature in Sweden is expected to rise by~~to the end of the century is~~ 1-2°C underin RCP2.6, 2-4°C in RCP4.5, and 4-6°C underin RCP8.5, with larger increasesdifferences in the north than in the south. The changes scale consistently in such a way that RCP8.5 in the nearelose future shows similar warming to RCP2.6 in the mid-dle of the century, and RCP8.5 in the mid-dle of the century is similar to RCP4.5 at the 245 end of the century (Strandberg et al., 2024a).

250 In Fennoscandia, we highlight two climate change patterns for temperature: winter is the season with the fastest warming rate, and the northern parts of the region areis warming faster than the southern parts. UnderIn RCP2.6, the warming in winter is 1.5-3.5°C from south to north (Fig. 2e) and 1.5-2°C in summer (Fig. 2g). UnderIn RCP8.5, the corresponding numbers are 4.5-8°C in winter (Fig. 2m), and 4-5°C in summer (Fig. 2o). This means that the warming is larger in winter, but also the difference between north and south.

255 The temperature change is especially large for the daily minimum temperature (tasmin) (Fig. 3a). For example, underin RCP4.5, the increase in tasmin is 3-6.5°C, to be compared to an increase in annual tas of 2-4°C. The increase in daily maximum temperature (tasmax) is comparable to tas, 2-3.5°C (Fig. 3b). A warmer climate means fewer cold days and more warm days. Accordingly, the number of frost days (fd) is projected to decrease, though relatively uniformly across the domain (Fig. 3c). RCP4.5 gives a reduction of 40-50 days in most of Fennoscandia and the Baltic countries. The change is somewhat smaller in parts of the Scandinavian mountain chain (a decrease in fd with 30-40 days), and larger over the Bothnian Sea and Bothnian Bay (a reduction of 65 days or more). See figure S1 for absolute values of the indicators in 1971-2000 and figures S2-4 for climate anomalies in all scenarios RCP2.6, RCP4.5 and RCP8.5.

260 Under RCP4.5, the ~~The~~ increase in the number of summer days (su) ranges from zero—according to RCP4.5 stretches from ~~zero, or just a few days, in large parts of the mountain chain and most sea areas and over most of the sea in the domain~~ to 20-24 days in southern Sweden and Denmark (Fig. 3d). The number of days with zero crossings (nzero) shows a general decrease on the annual scale (Fig. 3e). In winter, however, nzero increases in most of the domain, except for Denmark, southern Sweden and the Baltic countries (Fig. S5). In these areas, the temperatures will not drop below zero degrees as often, whereas in parts of northern Sweden the increase is as much as around 10 days (roughly corresponding to an increase 265 of 50 %).




Figure 3: Annual climate change anomalies in the CORDEX RCMs between 1971-2000 and 2071-2100 according to scenario RCP4.5. The maps show ensemble means of a) daily minimum temperature (tasmin, $^{\circ}\text{C}$), b) daily maximum temperature (tasmax, $^{\circ}\text{C}$), c) number of frost days (fd, days), d) number of summer days (su, days), e) number of days with zero crossings (nzero, days), f) mean precipitation (pr, mm mon^{-1}), g) number of days with heavy precipitation (r10mm, days) and h) dry days (dd, days). See table 1 for definitions of the indicators.

270

3.2 Projected change in precipitation and precipitation-based indicators

The annual average precipitation shows a general increase in the future (Fig. 3f). According to RCP4.5 the increase in
 275 annual average daily precipitation is $5\text{-}10 \text{ mm mon}^{-1}$ in large parts of the domain, with the increases along the Norwegian

west coast ~~of is~~ up to 15 mm mon⁻¹. Under ~~In~~ RCP2.6, the increase is smaller, 2-6 mm mon⁻¹ (fig. S2), and in RCP8.5 larger, 8-15 mm mon⁻¹ (fig. S4). For most of the domain, the increase is larger in winter and smaller in summer compared to the annual change (Figs S5-S8). Denmark and southern Sweden show ~~ehanges in~~ summer precipitation ~~changes~~ close to zero. On the annual scale, all models agree on the sign of change in most of the domain and all RCPs (Fig 3., Figs S2-S4). The 280 signal is least robust in RCP2.6 ~~because since~~ the change is smaller ~~there~~, and ~~since~~ precipitation ~~has~~ generally ~~has~~ large variability. The number of days with heavy precipitation (r10mm) is projected to increase with 3-5 in most of the domain (~~to be~~ compared ~~to with~~ 10-12 days in the reference period) (Fig. 3g). The change is smaller in RCP2.6 (up to +2 days ~~increase~~) and larger in RCP8.5 (+4-8 days) (Figs S1g & S2g). The number of dry days is projected to decrease ~~by with~~ 1-8 days (Fig 3h). ~~However, t~~he signal is not robust; half of the ensemble members ~~project an increase in dry days, while the other half~~ 285 ~~project a decrease give increasing number of dry days, and half of the members deereasing.~~


3.3 Local trends in climate indicators related to global warming

Climate change is unevenly distributed across the globe. In Scandinavia, like most of Europe, the overall warming since pre-industrial times ~~has been was~~ about twice the global mean at the end of the 20th century (Schimanke et al., 2022; WMO, 2023). In this section, we take a look at how specific features of local climate change in the CORDEX RCMs relates to the 290 change in global mean surface temperature (GMST) in the CMIP5 GCMs (Fig. 4).

The almost two-to-one relationship between global and local temperature is seen for mean, minimum and maximum temperatures ~~in the early parts of the 21st century~~ until the period 2011-2040 (Fig. 4 a-c). Within this period, the ratio between regional and global warming is 1.6-1.8. With increasing global warming, this relationship weakens and approaches a one-to-one relationship between change in global and local temperatures (i.e. parallel to the dotted lines in Figs 4a-c). In 295 RCP4.5 and RCP8.5 the trend from 2041-2070 to 2071-2100 is roughly one to one (1.1-1.2), suggesting that the faster warming in Scandinavia will slow down as GMST increases. A conclusion of this could be that the ratio between warming in Scandinavia and global warming is at its ~~maximum largest~~ in the beginning of the 21st century.

For indicators representing cold conditions, the trend gets flatter in RCP8.5, reflecting that the potential for change decreases. For example: the number of frost days cannot be less than zero. For warm indicators, the trend instead 300 ~~steepens gets steeper~~. The number of summer days is based on a temperature threshold, which means that there is a sudden effect when temperatures exceed the threshold. Consequently, the increase may be limited if the number of days above the threshold is already large.

Indicators for precipitation shows continued increase under global warming. Here, results both for pr and r10mm show a slightly weaker trend in RCP8.5 than in the other two scenarios.



305


Fig 4: Changes relative to 1971-2000 for the Fennoscandian domain (region A in Fig 1) in the CORDEX RCMs (y-axes) against that in global annual temperature in the driving CORDEX GCMs (x-axes), relative to the period 1971-2000. Different indicators are calculated based on RCM data: a) mean temperature (tas, $^{\circ}\text{C}$), b) minimum temperature (tasmin, $^{\circ}\text{C}$), c) maximum temperature (tasmax, $^{\circ}\text{C}$), d) no. of frost days (fd, days), e) no. of summer days (su, days), f) no. of days with zero crossings (nzero, days) g) precipitation (pr, mm mon⁻¹) h) no. of days with heavy precipitation (r10mm, days). Markers represent the periods 1971-2000 (cross), 2011-2040 (triangle), 2041-2070 (square), 2071-2100 (circle) for emissions scenarios RCP2.6 (green), RCP4.5 (orange) and RCP8.5 (light blue). In panels a-c the one-to-one relationship is shown with a dashed line, and the two-to-one with a dotted line.

310

315 **3.4 Model spread in the CORDEX RCM and CORDEX GCM ensembles compared to the spread in the CMIP5 GCM ensemble**

Even though the CORDEX RCM ensemble consists of several simulations using different GCM-RCM combinations, it may not represent the full potential spread of the climate change signal. To investigate how well the CORDEX RCMs capture the variability within the greater CMIP5 GCM ensemble, the average changes in temperature and precipitation over the
320 Fennoscandian domain (region A in Fig. 1) ~~were~~^{are} calculated. Figure 5 shows that the ensemble spread in the CMIP5 ensembles is larger than in the CORDEX RCM ensemble. ~~In particular~~^{Especially}, the difference between the minimum and maximum is larger in the CMIP5 GCMs than in the CORDEX RCMs. This could not entirely be explained by differences in ensemble sizes. ~~F~~See for example, ~~see~~ the numbers for RCP8.5 in Fig. 5c, where the CMIP5 GCMs and the CORDEX RCMs show large differences in spread although the ensembles are of comparable sizes. In the case of RCP8.5, the 6²⁷
325 members in the CORDEX RCM ensemble ~~use~~^{are} only ~~using~~ 7 unique GCMs and 11 RCMs, which is much less than the 34²⁵ unique GCMs in the full CMIP5 ensemble. When ~~considering~~^{looking} at an ensemble ~~just~~^{consisting} ~~only~~ of the 9 GCMs (including different realisations) used to force the RCMs, the spread is much smaller. ¶
The CORDEX RCM ensemble is compared to its raw equivalent, where no bias adjustment has been performed, to assess the impact of bias adjustment on the climate change signal. The means and spreads are similar in both RCM ensembles, but
330 the raw ensemble systematically shows smaller changes. Although small, these differences are significant in DJF, and in JJA under RCP8.5.

The ensemble means, however, are quite similar. In general, ~~all~~^{the} ~~two~~ ensembles agree on the large-scale differences, and the choice of emission scenario is of greater importance than the construction of the ensemble (Fig 5). The result is the same even when ~~examining~~^{looking} at smaller regions within the domain (e.g. regions B, C and D in Fig 1). ~~In~~^A conclusion, ~~is that~~ the Euro-CORDEX ensemble well captures the mean climate change signal, but that the spread is limited compared to the CMIP5 ensemble.



340 **Figure 5:** Temperature (tas, $^{\circ}\text{C}$) (a, b) and precipitation (pr, %) (c, d) anomalies in Fennoscandia 1971-2000 to 2071-2100 for
 340 winter (a, c) and summer (b, d) according to the scenarios RCP2.6 (yellowgreen), RCP4.5 (blueyellow) and RCP8.5 (redblue). The
 340 CMIP5 GCMs are represented by circles, the CORDEX GCMs by triangles, the unadjusted raw CORDEX RCMs by diamonds
 340 and the CORDEX RCMs by squares. The central marker represents the ensemble mean, the line spans between the 10th and 90th
 340 percentiles, open markers show ensemble minima and maxima. Panel b) also shows the number of members in the respective
 340 ensembles.

345

3.5 How the simulated GWL climate is influenced by the choice of GCMs, RCMs and RCPs

Here, we investigate how the characteristics of a certain GWL are influenced by the models and scenarios it is made of. Are all GWL2 the same, even if different models and scenarios are used to calculate them? First, we look at sub-ensembles based on GCMs (all members in a sub-ensemble are forced with the same GCM). Then we examine look at sub-ensembles

350 based on RCM and RCP (all members of a sub-ensemble used the same RCM and RCP, respectively). Statistically significant differences are assessed using an ANOVA analysis (see section 2.5).

3.5.1 Sub-ensembles based on driving GCMs

The results for sub-ensembles forced by the same GCM (all members of a sub-ensemble are forced with the same GCM, see Methods) are exemplified by temperature (tas) and annual number of summer days (su, see table 2 for definitions). Figure 6 355 shows which sub-ensembles that are significantly different from each other in the case of tas. All sub-ensembles from 1 to 7 are compared pairwise to see if they are significantly different or not. As an example, a green box at row 5 and column 1 means that sub-ensembles 5 and 1 are significantly different. In winter, the average temperature change at GWL2 is +1.5- 2.8 °C in the south and +1.7-4.2 °C in the north, depending on the chosen sub-ensemble (Fig. S9). Despite the rather large spread in warming the significant differences between sub-ensembles are not systematic in winter. However, in summer, 360 where the temperature change is +1.0-2.5 °C in the south and 1.3-2.9 °C in the north (Fig. S9), there are systematic significant differences between sub-ensembles. The two sub-ensembles with the largest warming, labelled 4 & 7, are significantly different from the other sub-ensembles (green boxes at lines 4 and 7, and columns 4 and 7 in Fig. 5). This pattern is also, to some extent, seen for su (Fig. 7). In the south, sub-ensemble 7 is significantly different from 5 of the other sub-ensembles; in the north sub-ensemble 4 is significantly different from 5 other. For precipitation, the difference at GWL2 365 is small compared to the variability within each sub-ensemble. Only a few pairs of sub-ensembles are significantly different (none in summer in the north), but not in a systematic way (Fig. S10).

The choice of GCM can have a large impact on the ensemble. The difference in simulated change in tas can be up to 2 °C depending on the driving GCM; this does however, transfer into consistent significant differences for only two sub-ensembles.

370




Fig 6: Matrix of significant differences in temperature (tas) between GCM-based sub-ensembles within RCM17, for southern Sweden (South, region C in Fig. 1) and northern Sweden (North, region D in Fig. 1). Green colours indicate significant differences between two sub-ensembles and pink non-significant differences. White colours indicate that an ensemble is compared with itself. Numbers indicate sub-ensemble numbers, with the number of members in parenthesis.

375





Figure 7: Same as Figure 6 but for annual number of summer days (su, see table 2 for definitions)

3.5.2 Sub-ensembles based on RCMs


380 Next, we examine Then, we proceed looking at sub-ensembles where the same RCM is used (all members of a sub-ensemble use the same RCM). Figure 8 shows which sub-ensembles that are significantly different from each other with regards to tas. The difference in projected change is about 1 °C between the sub-ensemble with the smallest and the largest change. Still, sub-ensemble no. 7 is the only sub-ensemble with systematically significant differences; in winter in the northern region and in summer it's different to all, or all but one, of the other ensembles. Sub-ensemble no. 7 is the sub-ensemble with the

385 smallest temperature increase. For su, there are more significant differences in the southern region than in the northern, reflecting the larger variability in su in the south (Fig 9). There are however, only two sub-ensembles that are significantly different from ~~to~~ three other sub-ensembles. Again, sub-ensembles 4 and 7, with a low number of su. For precipitation, the difference at GWL2 is small compared to the variability within each sub-ensemble. Only a few pairs of sub-ensembles are significantly different (in winter in the north one), but not in a systematic way (Fig. S11).



390

Figure 8 Matrix of significant differences in temperature (tas) between RCM-based sub-ensembles within RCM17, for southern Sweden (South, region C in Fig. 1) and northern Sweden (North, region D in Fig. 1.). Green colours indicate significant differences. Numbers indicate sub-ensemble numbers, with the number of members in parenthesis.



395 **Fig 9 Same as Fig 8, but for annual su.**

3.5.3 Sub-ensembles based on RCPs

As a last step, we ~~examine~~~~look at~~ sub-ensembles using the same RCPs. This analysis ~~addresses~~~~answers the question~~ whether ~~the choice of it~~~~matters which RCP~~ ~~affects the description of you use to describe~~ a GWL climate. ~~Here, In this case there are~~ 400 only two sub-ensembles ~~are to be~~ compared. ~~D~~~~The~~ differences between the ensembles based on RCP4.5 and RCP8.5 are generally small and not ~~statistically~~ significant (see ~~F~~fig. S12 for tas). RCP8.5 gives larger anomalies in tas, tasmin and tasmax in summer in all regions. The difference compared to RCP4.5 is around 0.15°C and just below the 95 % confidence ~~threshold~~~~interval~~. The difference in all other indicators are insignificant on the 99 % level.

405 Inevitably, the characteristics of a climate model ensemble ~~are~~~~is~~ determined by the simulations ~~that~~~~it~~ ~~comprises~~~~consists of~~. Using other models will not ~~yield identical~~~~give the same~~ results. These differences are however not systematic in any way, and mostly not significant. Even though an ensemble should be constructed with care, the role of the composition should not be exaggerated.

4 Discussion

4.1 The role of the models used on projected climate change

The projections of future climate presented here are consistent with other studies of the European climate (e.g. Coppola et al., 2021; Ranasinghe et al., 2021) and the climate in the Nordic region (e.g. Christensen et al., 2022). ~~The ensemble used here is an unbalanced ‘ensemble of opportunity’, as no pre-selection of models was applied. No pre-selection of models was made, which makes the ensemble used here an unbalanced ‘ensemble of opportunity’.~~ In such cases there is a risk that some 415 models are under- or over-represented, which influences the ensemble mean (Evin et al., 2021; Sobolowski et al., 2025). On the other hand, information is lost when simulations are discarded, and natural variability is best sampled by single-model large ensembles (e.g. von Trentini et al., 2019; Maher et al., 2020). Furthermore, we note that different selections of individual GCM-RCM-RCP-combinations can have significant impact ~~on~~~~f~~ the resulting ensemble as illustrated above. In the end, it is difficult to say that there is one approach that is always the most suitable. Different choices in the construction of an 420 ensemble can be made and ~~justified~~~~motivated~~ depending on the aim.

Insufficient aerosol forcing is proposed as a reason for the observed underestimation of the trend in summer temperature in RCMs over central Europe compared to observations (e.g. Boé et al., 2020; Schumacher et al., 2024). However, ~~the~~ the difference in summer warming between CORDEX and ERA5 is small in southern Sweden and Finland, and actually positive in Norway and northern Sweden (Schumacher et al., 2024). Bias adjustment may ~~alter~~~~change~~ the climate change signal, ~~but~~ 425 ~~this.~~ ~~This is,~~ ~~however,~~ generally seen as an improvement of the signal (Gobiet et al., 2015). MidAS, the bias adjustment method used here, is shown to add a small increase in the climate change signal for both temperature and precipitation in

Europe (Berg et al., 2022). The effect of bias adjustment on indicators ~~remains uncertain is unknown~~ and should be studied in the future.

A notable feature ~~of in~~ the scaling between local and global climate change is seen for the precipitation indicators (Figs 4g & 430 h). Here, there are clear differences between RCP4.5 and RCP8.5 even at the same level of global warming. It ~~has previously been is previously~~ shown on the global scale that the response in precipitation depends on both surface warming and radiative effect of increased amounts of greenhouse gases (Pendergrass et al., 2015). The net effect of these depends on the RCP scenario. Furthermore, the aerosol forcing is different in the different scenarios. This would make GWLs less suitable for precipitation. On the European scale this is further complicated by local features. The weaker response in precipitation could 435 be a consequence of drier conditions over the European continent leading to excessive evaporation and soil drying (e.g. Tuel and Eltahir, 2021).

4.2 Difference in model spread between GCM and RCM ensembles

In this study we show that the spread between the driving GCMs ~~wasere~~ larger than the spread between RCMs, even ~~in the eases~~ when the RCM ensemble ~~containedhad~~ more members. This is supported by Kjellström et al. (2018). A potential 440 explanation is that number of members is not the same as number of models. Previous studies show that multi-model ensembles have larger spread than single-model ensembles of similar, or even larger, sizes (von Trentini et al., 2019; Maher et al., 2021), ~~which~~. ~~This~~ is perhaps not surprising ~~given that~~. ~~As~~ different models have different physics. ~~Consequently~~, a multi-model ensemble can ~~provideoffer~~ a wider response to forcing and natural variability than ~~what~~ a single-model ensemble ~~can~~. ~~This isA~~ supported by the observation to this is that the ensemble means in the CORDEX GCM ensemble is 445 not affected in any major way when ~~additional realisations from the same GCM are included we include more realisations with some GCMs. ALikely, adding more realisations likely improves gives a better estimates of natural variability and extremes, but does not influence the mean values as much, assince all realisations simulate the same climate (as opposed to simulations with different physics or forcing).~~

In this study, bias-adjusted RCMs are compared to non-adjusted GCMs. Bias adjustment may reduce ~~the~~ model spread in 450 absolute values since systematic biases are minimised and all models are forced towards the reference data. ~~Here, it systematically increases the climate change signal in the RCM ensemble. Although this increase is in many cases significant, it is relatively small, and the raw RCM ensemble is more similar to the bias-adjusted RCM ensemble than to any of the GCM ensembles. The model spread in the climate change signal would, however, not be affected, assuming that bias adjustment with MIdAS preserves the climate change signal (Berg et al., 2022). The analysis of model spread in Section 3.5 and Figure 455 5 builds on the spread in climate change signal.~~ Consequently, the differences between GCMs and RCMs are likely not explained by the application of bias adjustment.

Another explanation for differences in model spread ~~areis~~ inconsistencies in forcing between the RCMs and the driving GCMs, where aerosol forcing probably is the most prominent factor in the context of this study (Taranu et al., 2023). ~~This problem is indeed seen in both GCMs and RCMs, but only for summer in central Europe (Schumacher et al., 2024).~~

460 4.3 On the characteristics of GWL ensembles

SinceAs GWLs ~~are~~ in fact ~~are~~ used for many different purposes it is necessary to investigate the characteristics of GWL ensembles~~s~~.~~e~~Especially how RCPs influence the GWL climate. Our study shows, for a broad range of indicators, that the choice of RCPs ~~used~~has minimal effect on the GWL climate~~created~~. Furthermore, it is difficult to ~~demonstrate that including show that the inclusion of~~ specific GCMs ~~or and~~ RCMs influence the GWL climate in a significant way. This is 465 perhaps expected considering that GCMs and RCMs are not independent (Sørland et al., 2018) and that the uncertainty in climate change due to GCMs can be as large as the uncertainty due to RCMs (Evin et al., 2021).

A caveat to our findings relates to the small number of members in the sub-ensembles. ~~Sub-ensemble~~ sizes of 2-8 make it difficult to draw robust conclusions. Small samples reduce the power of the ANOVA test to detect differences between sub-ensembles and ~~are~~ more likely to fail to reject a false null hypothesis. In any case, this—and similar—ensemble is what is 470 used to create GWL ensembles, and they must therefore be evaluated as much as possible. Adding more members would increase the statistical power, but would also ~~alter change~~ the ensemble's composition~~to something else~~. We just have to do what we can with the ensemble at hand. A more solid evaluation could perhaps be achieved if AI or emulators were first used to fill all gaps in the matrix. That would enable a balanced comparison across GCMs and RCMs.

We performed our analysis on GWL1.5 and GWL2₁ and our conclusions only apply to these specific GWLs. It would be 475 interesting to expand the analysis to more GWLs, but there are practical limitations to this. Smaller GWL increments would mean larger overlap between GWLs₁ making it difficult to draw robust conclusions about the differences between GWLs. Furthermore, most RCP4.5 simulations do not reach GWL3 which means that the ensemble size would be heavily reduced, making the statistical analysis less solid. Also, if only one RCP reaches GWL3₁ it is not possible to investigate the role of RCPs in the construction of a GWL~~—; arguably the most relevant aspect to understand perhaps the most relevant thing to know~~. ~~Studying a broader~~~~To study~~ a range of GWLs in an RCM ensemble ~~would require a separate~~~~is another~~ study, a study that would require other simulations, and maybe simulations that do not exist (for example more scenarios that reach 480 GWL3).

5 Summary and conclusions

485 Global warming ~~in means for~~ Fennoscandia ~~means~~ higher temperatures₁, more warm days₁ and fewer cold days. In southern Sweden the number of summer days is doubled until the end of the century according to RCP4.5. At the same time₁, the number of frost days decreases ~~by with~~ 20-50 %. Precipitation increases generally₁, this shows in increasing mean precipitation, increasing number of days with heavy precipitation and decreasing number of dry days.

The RCM ensemble used here captures₁, on average, the change pattern from the CMIP5 GCM ensemble. ~~However, t~~The 490 ensemble spread,~~however~~, is larger in the CMIP5 ensemble.

The choice of RCP has minimal influencesignificance on the GWL2 ensembles. This implies that it would be safe to mix RCPs in the construction of GWL ensembles in order to increase ensemble size, and that a GWL could be based on only one RCP. It should be noted, however, that we only look at mean changes. Trends within a GWL period do indeed depend on the RCP, and this could influence extremes. For example: the last years within a GWL period based on RCP8.5 may be warmer than the last years within a GWL period based on RCP2.6. The largest difference between GWL2 sub-ensembles, regardless of how they are constructed in terms of combining GCMs and RCMs, is seen for temperature-based indices.H; however, it remainsis difficult to say whether the choice of GCM or RCM contributes most to these variations.

All studied climate indicators scale approximatelysomewhat linearly to the change in GMST. For indicators based on temperature thresholds, trend slopes may shift when there may be a shift in trend slope when the temperatures exceed certain levelsrise above a certain level. Currently the regional temperature change in Sweden is almost twice as large as the global trend. This ratio will decrease as GMST increases, to more and more approach a one-to-one relationship. This suggests that there is a limit to the feedback mechanisms that now accelerates the warming in Sweden. aAnd indicates that the ratio between local and global warming currently may be at its maximumlargest. Furthermore, this means that the steady relationship between global and regional warming that is sometimes assumed in weather attribution and regional warming levels may not remain validhold in the future.

Acknowledgements

We acknowledge Euro-CORDEX and all groups that contribute with simulations. Analyses were performed on the Swedish climate computing resources Bi and Freja at the Swedish National Supercomputing Centre (NSC), Linköping University. This is a contribution to the strategic research areas MERGE (Modelling the Regional and Global Earth system) and the 510 Bolin Centre for Climate Research.

Funding statement

This work is partially funded by Sweden's innovation agency, Vinnova, to the project titled "Adapting Urban Rail Infrastructure to Climate Change (AdaptUrbanRail)" (grant no. 2021-02456). This work was also supported via Horizon Europe IA project Precilience (Grant Agreement 101157094).

515 Author contributions

GS – Conceptualization, Methodology, Formal analysis, Visualization, Project administration, Writing – Original Draft; **AT** – Methodology, Formal analysis, Visualization, Writing – Review & Editing; **LB** – Software, Data Curation, Writing –

520 **Data availability**

Ensemble means of 30 year periods (absolute values and anomalies) can be viewed and downloaded at:

<https://www.smhi.se/en/climate/tools-and-inspiration/climate-change-scenario/climate-change-scenario-tool>

Competing interests

The authors declare that they have no conflict of interest.

525 **References**

Andersson, S., Bärring, L., Landelius, T., Samuelsson, P. and Schimanke, S.: SMHI Gridded Climatology, Report Meteorology and Climatology No. 118, 2021, SMHI, Norrköping, Sweden, 2021.

Bärring, L. and Strandberg, G.: Does the projected pathway to global warming targets matter? Environ. Res. Lett. 13, <https://doi.org/10.1088/1748-9326/aa9f7>, 2018.

530 **Software:** Bärring, L. Zimmermann, K., Löw, J. and Nilsson, C: Climix, <https://climix.readthedocs.io/>, 2024

Berg, P., Bosshard, T., Yang, W., and Zimmermann, K.: MIdASv0.2.1 – Multi-scale bias Adjustment, Geosci. Model Dev., 15, 6165–6180, <https://doi.org/10.5194/gmd-15-6165-2022>, 2022.

Boé, J., Somot, S., Corre, L. And Nabat, P.: Large discrepancies in summer climate change over Europe as projected by global and regional climate models: causes and consequences Clim. Dyn. 54 2981–3002, <https://doi.org/10.1007/s00382-020-05153-1>, 2020

535 Christensen, O. B. and Kjellström, E.: Partitioning uncertainty components of mean climate and climate change in a large ensemble of European regional climate model projections Clim. Dyn. 54 4293–308, 2020.

Christensen, O. B. and Kjellström, E.: Filling the matrix: an ANOVA-based method to emulate regional climate model simulations for equally-weighted properties of ensembles of opportunity Clim. Dyn. 58 2371–85, 2021.

540 Copernicus Climate Change Service (C3S): European State of the Climate 2023, Summary: <https://doi.org/10.24381/bs9v-8c66>, 2024

Christensen, O. B., Kjellström, E., Dieterich, C., Gröger, M., and Meier, H. E. M.: Atmospheric regional climate projections for the Baltic Sea region until 2100, Earth Syst. Dynam., 13, 133–157, <https://doi.org/10.5194/esd-13-133-2022>, 2022.

545 Coppola, E., Nogherotto, R., Ciarlo', J. M., Giorgi, F., van Meijgaard, E., Kadygov, N., Iles, C., Corre, L., Sandstad, M., Somot, S., Nabat, P., Vautard, R., Levavasseur, G., Schwingshakl, C., Jana Sillmann, J., Kjellström, E., Nikulin, G.,

Aalbers, E., Lenderink, G., Christensen, O. B., Boberg, F., Lund Sørland, S., Demory, M.-E., Bülow, K., Teichmann, C., Warrach-Sagi, K. and Wulfmeyer, V.: Assessment of the European climate projections as simulated by the large EURO-CORDEX regional climate model ensemble *J. Geophys. Res.: Atmospheres* 126 e2019JD032356, 2021.

Déqué, M., Somot, S., Sanchez-Gomez, E., Goodess, C. M., Jacob, D., Lenderink, G. and Christensen, O. B.: The spread amongst ENSEMBLES regional scenarios: regional climate models, driving general circulation models and interannual variability *Climate Dynamics*, 951, 964, 38, 5, 1432-0894, 10.1007/s00382-011-1053-x, 2012

Forster, P. M., Smith, C., Walsh, T., Lamb, W. F., Lamboll, R., Hall, B., Hauser, M., Ribes, A., Rosen, D., Gillett, N. P., Palmer, M. D., Rogelj, J., von Schuckmann, K., Trewin, B., Allen, M., Andrew, R., Betts, R. A., Borger, A., Boyer, T., Broersma, J. A., Buontempo, C., Burgess, S., Cagnazzo, C., Cheng, L., Friedlingstein, P., Gettelman, A., Gütschow, J., Ishii, M., Jenkins, S., Lan, X., Morice, C., Mühle, J., Kadow, C., Kennedy, J., Killick, R. E., Krummel, P. B., Minx, J. C., Myhre, G., Naik, V., Peters, G. P., Pirani, A., Pongratz, J., Schleussner, C.-F., Seneviratne, S. I., Szopa, S., Thorne, P., Kovilakam, M. V. M., Majamäki, E., Jalkanen, J.-P., van Marle, M., Hoesly, R. M., Rohde, R., Schumacher, D., van der Werf, G., Vose, R., Zickfeld, K., Zhang, X., Masson-Delmotte, V., and Zhai, P.: Indicators of Global Climate Change 2023: annual update of key indicators of the state of the climate system and human influence, *Earth Syst. Sci. Data*, 16, 2625–2658, <https://doi.org/10.5194/essd-16-2625-2024>, 2024.

Gobiet, A., Suklitsch, M., and Heinrich, G.: The effect of empirical-statistical correction of intensity-dependent model errors on the temperature climate change signal, *Hydrol. Earth Syst. Sci.*, 19, 4055–4066, <https://doi.org/10.5194/hess-19-4055-2015>, 2015.

Hausfather, Z. and Peters, G.P.: RCP8.5 is a problematic scenario for near-term emissions, *Proc. Natl. Acad. Sci. U.S.A.* 117 (45) 27791-27792, <https://doi.org/10.1073/pnas.2017124117>, 2020.

IPCC: Summary for Policymakers. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 3–32, doi:10.1017/9781009157896.001, 2021.

IPCC: Summary for Policymakers [H.-O. Pörtner, D.C. Roberts, E.S. Poloczanska, K. Mintenbeck, M. Tignor, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem (eds.)]. In: *Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 3-33, doi:10.1017/9781009325844.001, 2022.

Iturbide, M., Fernández, J., Gutiérrez, J. M., Pirani, A., Huard, D., Al Khourdajie, A., Baño-Medina, J., Bedia, J., Casanueva, A., Cimadevilla, E., Cofiño, A. S., De Felice, M., Diez-Sierra, J. García-Díez, M., Goldie, J., Herrera, D. A., Herrera, S., Manzanas, R., Milovac, J., Radhakrishnan, A., San-Martín, D., Spinuso, A., Thyng, K. M., Trenham, C. and Yelekçi, Ö.:

580 Implementation of FAIR principles in the IPCC: the WGI AR6 Atlas repository, *Scientific Data*, 629, 9, 1, 2052-4463, <https://doi.org/10.1038/s41597-022-01739-y>, 2022.

Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O. B., Bouwer, L. M., Braun, A., Colette, A., Déqué, M., Georgievski, G., Georgopoulou, E., Gobiet, A., Menut, L., Nikulin, G., Haensler, A., Hempelmann, N., Jones, C., Keuler, K., Kovats, S., Kröner, N., Kotlarski, S., Kriegsmann, A., Martin, E., van Meijgaard, E., Moseley, C., Pfeifer, S., Preuschmann, 585 S., Radermacher, C., Radtke, K., Rechid, D., Rounsevell, M., Samuelsson, P., Somot, S., Soussana, J.-F., Teichmann, C., Valentini, R., Vautard, R., Weber, B. and Pascal Yiou, P.: EURO-CORDEX: new high-resolution climate change projections for European impact research *Reg. Environ. Change* 14 563–78, 2014.

Katragkou, E., Sobolowski, S. P., Teichmann, C., Solmon, F., Pavlidis, V., Rechid, D., Hoffmann, P., Fernandez, J., Nikulin, G. and D. Jacob, D.: Delivering an Improved Framework for the New Generation of CMIP6-Driven EURO-CORDEX 590 Regional Climate Simulations. *Bull. Amer. Meteor. Soc.*, 105, E962–E974, <https://doi.org/10.1175/BAMS-D-23-0131.1>, 2024.

Kjellström, E., Bärring, L., Nikulin, G., Nilsson, C., Persson, N. and Strandberg, G: Production and use of regional climate model projections – A Swedish perspective on building climate services, *Clim. Serv.* 2-3, 15-29, <http://dx.doi.org/10.1016/j.ciser.2016.06.004>, 2016.

595 Kjellström, E., Nikulin, G., Strandberg, G., Bøssing Christensen, O., Jacob, D., Keuler, K., Lenderink, G., van Meijgaard, E., Schär, C., Somot, S., Lund Sørland, S., Teichmann, C. and Vautard, R.: European climate change at global mean temperature increases of 1.5 and 2 °C above pre-industrial conditions as simulated by the EURO-CORDEX regional climate models. *Earth Syst. Dynam.*, 9, 459-478, 2018.

Lind, P., Belušić, D., Christensen, O B., Dobler, A., Kjellström, E., Landgren, O., Lindstedt, D., Matte, D., Pedersen, R. A., 600 Toivonen, E. and Wang, F.: Benefits and added value of convection-permitting climate modeling over Fennoscandia, *Climate Dynamics*, 1893-1912, 55, 7, <https://doi.org/10.1007/s00382-020-05359-3>, 2020.

Maher, N., Lehner, F. and Marotzke, J.: Quantifying the role of internal variability in the temperature we expect to observe in the coming decades *Environmental Research Letters*, Volume 15, Number 5, DOI 10.1088/1748-9326/ab7d02, 2020

605 Maher, N., Power, S.B. and Marotzke, J.: More accurate quantification of model-to-model agreement in externally forced climatic responses over the coming century *Nat Commun* 12, 788, <https://doi.org/10.1038/s41467-020-20635-w>, 2021.

Maule, C. F., Mendlík, T. and Christensen, O. B.: IMPACT2C - Quantifying projected impacts under 2°C warming *Climate Services*, 7, 3, 11, 2405-8807, <https://doi.org/10.1016/j.ciser.2016.07.002>, 2017

Olsson J., Berg., P. and Kawamura, A.: Impact of RCM spatial resolution on the reproduction of local, subdaily precipitation 610 *J Hydrometeorol* 16 534–47, 2015.

Pendergrass, A. G., F. Lehner, B. M. Sanderson, and Y. Xu: Does extreme precipitation intensity depend on the emissions scenario?, *Geophys. Res. Lett.*, 42 8767–8774, doi:10.1002/2015GL065854, 2025

Prein, A. F., Langhans, W., Fosser, G., Ferrone, A., Ban, N., Goergen, K., Keller, M., Tölle, M., Gutjahr, O., Feser, F.,
 615 Brisson, E., Kollet, S., Schmidli, J., van Lipzig, N. P. M. and Leung, R.: A review on regional convection-permitting climate modeling: Demonstrations, prospects, and challenges *Rev. Geophys.* 53 323–61, 2015.

Press, S. J.: *Applied Multivariate Analysis*. Holt, Rinehart and Winston, 521 pp., 1972.

Ranasinghe, R., A.C. Ruane, R. Vautard, N. Arnell, E. Coppola, F.A. Cruz, S. Dessai, A.S. Islam, M. Rahimi, D. Ruiz Carrascal, J. Sillmann, M.B. Sylla, C. Tebaldi, W. Wang, and R. Zaaboul, 2021: Climate Change Information for Regional
 620 Impact and for Risk Assessment. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1767–1926, doi:10.1017/9781009157896.014.

625 Rummukainen, M.: Added value in regional climate modeling *Wire Clim. Change* 7 145–59, 2016.

Schimanke, S., Joelsson, M., Andersson, S., Carlund, T., Wern, L., Hellström, S. and Kjellström, E.: *Observerad klimatförändring i Sverige, 1860–2021* SMHI Climatol. Rep. 69 89, 2022.

Schumacher, D. L., Singh, J., Hauser, J., Fischer, E. M., Wild, M., and Seneviratne, S. I.: Exacerbated summer European warming not captured by climate models neglecting long-term aerosol changes, *Communications Earth & Environment*, 182,
 630 5 1 2662-4435, <https://doi.org/10.1038/s43247-024-01332-8>, 2024

SMHI: <https://www.smhi.se/en/climate/tools-and-inspiration/climate-change-scenario/climate-change-scenario-tool/>, last access: 10 July 2025

Strandberg, G. and Lind, P.: The importance of horizontal model resolution on simulated precipitation *Weather Clim. Dynam.* 2 181–204, 2021.

635 Strandberg, G., Andersson, B. and Berlin A.: Plant pathogen infection risk and climate change in the Nordic and Baltic countries, *Environ. Res. Commun.* 6 031008, DOI 10.1088/2515-7620/ad352a, 2024a.

Strandberg, G, Blomqvist, P., Fransson, N., Göransson, L., Hansson, J., Hellsten, S., Kjellström, E., Lin, C., Löfblad, E., Montin, S., Nyholm, E., Sandgren, A., Unger, T., Walter, V. and Westerberg, J.: Bespoke climate indicators for the Swedish energy sector – a stakeholder focused approach, *Climate Services*, Volume 34, 100486, ISSN 2405-8807,
 640 <https://doi.org/10.1016/j.cliser.2024.100486>, 2024b.

Schwalm, C.R., Glendon, S., Duffy, P. B.: Reply to Hausfather and Peters: RCP8.5 is neither problematic nor misleading, *Proc. Natl. Acad. Sci. U.S.A.* 117 (45) 27793–27794, <https://doi.org/10.1073/pnas.2018008117>, 2020.

Taranu, I. S., Somot, S., Alias, A., Boé, J. and Delire, C.: Mechanisms behind large-scale inconsistencies between regional and global climate model-based projections over Europe *Climate Dynamics*, 3813–3838, 60, 11, 10.1007/s00382-022-06540-
 645 6, 2023

Taylor, K. E., Stouffer, R. J. and Meehl, G. A.: An overview of CMIP5 and the experiment design *Bull. Am. Meteorol. Soc.* 93 485–98, 2012.

650 Themeßl, M. J., Gobiet, A. and Heinrich, G.: Empirical-statistical downscaling and error correction of regional climate models and its impact on the climate change signal, *Climatic Change*, 112, 449–468, <https://doi.org/10.1007/s10584-011-0224-4>, 2011.

655 von Trentini, F., Leduc, M. and Ludwig, R.: Assessing natural variability in RCM signals: comparison of a multi model EURO-CORDEX ensemble with a 50-member single model large ensemble *Clim Dyn* 53, 1963–1979, <https://doi.org/10.1007/s00382-019-04755-8>, 2019

Tuel, A., and Eltahir, E. A. B., 2021: Mechanisms of European Summer Drying under Climate Change. *J. Climate*, 34, 8913–8931, <https://doi.org/10.1175/JCLI-D-20-0968.1>, 2021

Tukey, J. W.: Comparing Individual Means in the Analysis of Variance. *Biometrics*, vol. 5, no. 2, pp. 99-114. JSTOR, www.jstor.org/stable/3001913, 1949.

660 UNFCCC: The Paris Agreement. United Nations Framework Convention on Climate Change, http://unfccc.int/paris_agreement/items/9485.php, 2015

Vautard, R., Kadygrov, N., Iles, C., Boberg, F., Buonomo, E., Bülow, K., Coppola, E., Lola Corre, L., van Meijgaard, E., Nogherotto, R., Sandstad, M., Schwingshakl, C., Somot, S., Aalbers, E., Christensen, O. B., Ciarlo, J. M., Demory, M.-E., Giorgi, F., Jacob, D., Jones, R. G., Keuler, K., Kjellström, E., Lenderink, G., Levavasseur, G., Nikulin, G., Sillmann, J., Solidoro, C., Lund Sørland, S., Steger, C., Teichmann, C., Warrach-Sagi, K. and Wulfmeyer, V.: Evaluation of the large EURO-CORDEX regional climate model ensemble *J. Geophys. Res.* 126 e2019JD032344, 2020.

665 Wilcke, R.A.I., Mendlik, T. and Gobiet, A.: Multi-variable error correction of regional climate models. *Climatic Change* 120, 871–887. <https://doi.org/10.1007/s10584-013-0845-x>, 2013.

670 WMO: WMO Guidelines on the Calculation of Climate Normals, WMO-No 1203, ISBN 978-92-63-11203-3, https://library.wmo.int/viewer/55797?medianame=1203_en, 2017. Last access 2 Jul, 2025.

WMO: State of the global climate 2022, WMO-No 1316 55 ISBN 978-92-63-11316-0 <https://library.wmo.int/records/item/66214-state-of-the-global-climate-2022>, 2023. last access 18 Feb, 2025