

Role of aerosol–cloud–radiation interactions in modulating summertime quasi-biweekly rainfall intensity over South China

Hongli Chen¹, Pang-Chi Hsu¹, Anbao Zhu^{1,2}, and Xiaoyan Ma³

¹State Key Laboratory of Climate System Prediction and Risk Management/Key Laboratory of Meteorological Disaster, Ministry of Education/Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing University of Information Science and Technology, Nanjing, China

²Joint International Research Laboratory of Atmospheric and Earth System Sciences, School of Atmospheric Sciences, Nanjing University, Nanjing, China

³Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, School of Atmospheric Physics, Nanjing University of Information Science and Technology, Nanjing, China

Correspondence to: Pang-Chi Hsu (pangchi@nuist.edu.cn)

Abstract. Persistent heavy rainfall events over densely populated South China are closely linked to the intensification of quasi-biweekly (8–30-day) oscillations. This study examines whether and how aerosols influence quasi-biweekly oscillations using observational analyses and model experiments. Statistical analysis reveals a significant phase-leading relationship between increased aerosol loadings, quantified by aerosol optical depth, and subsequent enhancement of 8–30-day rainfall anomalies. At the 8–30-day timescale, aerosols primarily influence rainfall intensity through cloud microphysical processes, with radiative effects playing a secondary role. Approximately four days before enhanced rainfall events, positive aerosol anomalies contribute to increased low-level cloud water content, leading to condensational latent heat release. This low-level latent heating strengthens low-level moisture convergence and ascending motion, which uplifts cloud droplets above the freezing level. Subsequently, additional latent heat release from mixed-phase processes (freezing/deposition) further intensifies vertical motion, amplifying precipitation anomalies. Once deep convection develops, clouds absorb longwave radiation, sustaining precipitation intensification. Sensitivity experiments using the Weather Research and Forecasting model coupled with Chemistry (WRF-Chem) confirm these mechanisms, demonstrating that anthropogenic aerosol enhancement intensifies precipitation anomalies through both aerosol-cloud microphysical interactions and longwave cloud-radiative effects, with the former being more dominant. Quantitatively, aerosol-induced latent heating exceeds aerosol-induced longwave radiative heating by a factor of ~4–7 in both observations and model simulations. These findings highlight the need to improve aerosol-cloud microphysical parameterizations in operational models to enhance the accuracy of extended-range heavy rainfall predictions in South China.

1 Introduction

Aerosols influence clouds and precipitation through two primary mechanisms: one involves directly modifying radiation, while the other acts through their role as cloud condensation nuclei (CCN) or ice nuclei (IN) (e.g., Koren et al., 2004; Tao et al., 2012; Li et al., 2016, 2019; Zhu et al., 2022; Zhao et al., 2023; Stier et al., 2024). These are referred to as the radiative effect and the microphysical effect, respectively. The radiative effect involves the scattering and absorption of solar radiation by aerosols (i.e., the so-called “direct effect”), which commonly leads to atmospheric heating, surface cooling, stabilization of atmospheric stratification, and suppression of precipitation (Bollasina et al., 2011), but can also enhance local or remote precipitation under favorable conditions (e.g., Fan et al., 2015; Zhu et al., 2022; Wei et al., 2023). In particular, absorbing aerosols within clouds enhance cloud evaporation, thereby inhibiting cloud and precipitation formation, a phenomenon known as the semi-direct effect (Ackerman et al., 2000). The microphysical effect occurs when aerosols serve as CCN or IN (referred to as the “indirect effect”), increasing cloud droplet number concentration, enhancing cloud albedo (Twomey, 1977), reducing precipitation efficiency, and prolonging cloud lifetime (Albrecht, 1989). Additionally, the aerosol indirect effect can invigorate deep convective cloud through freezing-induced intensification (Rosenfeld et al., 2008) and enhanced condensational heating (Fan et al., 2018) precipitation, the so-called invigoration effect (Rosenfeld et al., 2008; Li et al., 2011; Fan et al., 2025), though its significance mechanisms remains debated qualitatively and quantitatively poorly understood.

A recent review by Zhao et al. (2024) summarized that aerosol effects on precipitation are complex and highly variable, depending on factors such as aerosol type and concentration, precipitation characteristics and meteorological conditions, leading to substantial regional disparities. Additional uncertainties in the aerosol–precipitation relationship arise across when examining different regions and timescales due to variations in dynamic and thermodynamic conditions, aerosol properties, and multi-scale interactions. While the impacts of aerosols on short-term weather phenomena (i.e., diurnal cycles and synoptic variability of rainfall) and long-term climate variations (i.e., seasonal and interannual-decadal changes in precipitation) have been well documented (e.g., Li et al., 2016; Chen et al., 2018; Sun and Zhao, 2021; Zhao et al., 2024; Chen et al., 2025), their influences on intraseasonal rainfall variability—closely linked to persistent extreme precipitation events in the Asian monsoon regions (Hsu et al., 2016; Chen et al., 2024; Xie et al., 2024)—remain poorly understood.

Prior research has primarily focused on the effects of aerosols on the intraseasonal evolution of the Indian summer monsoon; however, no consensus has yet been reached. Some studies suggest that aerosols suppress Indian monsoon rainfall (Bhattacharya et al., 2017; Dave et al., 2017; Arya et al., 2021; Surendran et al., 2022; Debnath et al., 2023), whereas others highlight their positive contribution to

enhanced monsoon precipitation variations (Manoj et al., 2011; Hazra et al., 2013; Vinoj et al., 2014; Singh et al., 2019). These contrasting findings may stem from differences in datasets and models or, more fundamentally, from uncertainties in subseasonal aerosol–cloud–precipitation interactions (Li et al., 2016; Wang et al., 2022). In particular, the competing effects of radiative and microphysical processes may vary spatially and temporally, depending on aerosol type and emission sources (local vs. remote). Among these effects, the positive contribution of aerosol indirect effects on intraseasonal oscillations of the Indian monsoon has been emphasized (Hazra et al., 2013), mainly through the enhancement of cold-rain processes involving increased ice hydrometeors and strengthened high-level latent heating.

Influenced by active intraseasonal oscillations, persistent heavy precipitation frequently strikes densely populated southeastern China (Hsu et al., 2016), posing increasingly severe threats to socioeconomic development and the livelihoods of billions. Research on aerosol–precipitation interactions over South China in summer has predominantly examined diurnal precipitation shifts (Guo et al., 2016; Lee et al., 2016; Sun and Zhao, 2021), mesoscale rainfall intensity (Zhang et al., 2020; Xiao et al., 2023a), synoptic-scale rainfall variability (Liu et al., 2020; Guo et al., 2022), and seasonal-to-climatological rainfall changes (Wang et al., 2011; Yang and Li, 2014; Zhu et al., 2023). However, despite the importance of intraseasonal oscillations in regulating regional rainfall, few studies have examined aerosol impacts on intraseasonal variability of rainfall intensity. ~~Focusing on densely populated South China, where nonabsorbing aerosols and intraseasonal oscillations are active during the summer monsoon season (Lee et al., 2007; Huang et al., 2014; Yang et al., 2016),~~ Thus, this study explores the role of aerosols in modulating rainfall anomalies at the quasi-biweekly (8–30-day) timescale, which is a critical factor in persistent heavy rainfall in this region (Chen et al., 2024). The relative contributions of cloud microphysical and radiative processes to rainfall intensification are examined through observational analysis and model experiments. The remainder of the paper is structured as follows. Section 2 describes the datasets and methodologies used in this study. Section 3 analyzes the observed aerosol impacts on quasi-biweekly precipitation anomalies. Section 4 presents the experimental results from the Weather Research and Forecasting model coupled with Chemistry (WRF-Chem). Concluding remarks and discussion are provided in Section 5.

2 Data and methods

2.1 Datasets

Daily precipitation data were obtained from the National Oceanic and Atmospheric Administration (NOAA) Climate Prediction Center (CPC) dataset (Chen et al., 2008) with a horizontal resolution of 0.5°. Deep convection activity was analyzed using daily interpolated outgoing longwave radiation (OLR) data from NOAA at a 2.5° grid resolution (Liebmann and Smith, 1996).

To investigate the effects of aerosols on radiative and moisture processes, ~~daily meteorological and aerosol data~~ mean aerosol optical depth (AOD), radiative fluxes, specific humidity and circulation variables from the Modern-Era Retrospective Analysis for Research and Applications version 2 (MERRA-2), ~~produced provided~~ by the National Aeronautics and Space Administration (NASA) Global Modeling and Assimilation Office (Gelaro et al., 2017), were collected at a ~~spatial~~horizontal resolution of $0.5^\circ \times 0.625^\circ$ (~~longitude $\times$ latitude $\times$ longitude~~) on 42 vertical levels. ~~Daily mean aerosol optical depth (AOD), which shows good agreement with observations (Buchard et al., 2017), was used to represent aerosol pollution conditions. Radiation variables were employed to assess the atmospheric radiative budget. MERRA-2 provides the complete set of variables required for atmospheric radiation and moisture budget quantifications, whereas other reanalyses and observations lack some of these key variables. Even so, to ensure that our conclusions are not dataset-dependent, we compared budget analysis results with those derived from the fifth-generation European Center for Medium-Range Weather Forecasts (ECMWF) atmospheric reanalysis (ERA5; Hersbach et al., 2020), which has a spatial resolution of $0.25^\circ \times 0.25^\circ$ and 37 vertical levels. To further reduce uncertainties inherent in reanalyses, we also employed radiative fluxes from Clouds and the Earth's Radiant Energy System Synoptic products (CERES-SYN; Rutan et al., 2015) at 1° resolution, and AOD from the Moderate Resolution Imaging Spectroradiometer (MODIS) Collection 6 Level-3 aerosol product onboard the Terra satellite (Levy et al., 2013) at 1° resolution.~~

~~To reduce uncertainties inherent in reanalysis data, we also used the Moderate Resolution Imaging Spectroradiometer (MODIS) Collection 6 Level-3 aerosol product onboard the Terra satellite, with a horizontal resolution of $1^\circ \times 1^\circ$ (Levy et al., 2013). To examine aerosol influences on cloud microphysical processes, daily two-dimensional~~ cloud parameters, including liquid cloud fraction, ice cloud fraction, cloud top pressure, cloud droplet effective radius and ice water path, were obtained from the MODIS Collection 6 Level-3 cloud product (Platnick et al., 2017). These cloud properties have been widely used to analyze aerosol indirect effects (e.g., Zhou et al., 2020; Jia et al., 2024). ~~Although CloudSat provides three-dimensional cloud products (Austin et al., 2009), substantial temporal gaps prevent its use for analyzing continuous sequences of aerosol-cloud-precipitation interactions at intraseasonal timescales. Additionally, Thus, three-dimensional liquid and ice cloud water contents from the fifth-generation European Centre for Medium-Range Weather Forecasts (ECMWF) atmospheric reanalysis (ERA5; Hersbach et al., 2020), with a spatial resolution of $1^\circ \times 1^\circ$ on 37 vertical levels, was collected were instead taken from MERRA-2 and ERA5 to evaluate vertical cloud structures. Specific humidity and circulation fields from ERA5 were used to conduct a moisture budget diagnostic analysis.~~

All datasets cover the period from 2000 to 2021, with ~~the analysis~~is focuseding on the boreal summer season (May–September, MJJAS).

2.2 Diagnostic method

To investigate the processes regulating precipitation anomalies, we diagnosed the budget equation of column-integrated moisture perturbation:

$$\left\langle \frac{\partial q}{\partial t} \right\rangle' = -\langle \mathbf{V} \cdot \nabla q \rangle' - \langle q \nabla \cdot \mathbf{V} \rangle' - \left\langle \frac{\partial}{\partial p} (\omega q) \right\rangle' - \left\langle \frac{Q_2}{L} \right\rangle', \quad (1)$$

where the primes denote the 8–30-day component derived using the Lanczos filtering method (Duchon, 1979), angle brackets indicate a mass-weighted vertical integration from 1000 to 100 hPa level, $\mathbf{V}$ is the horizontal wind field, q is the specific humidity, ω is vertical pressure velocity, ∇ is the horizontal gradient operator, p is pressure, L is the latent heat of condensation, and Q_2 represents the latent heating due to condensation/evaporation processes and subgrid-scale moisture flux convergences ~~is the apparent moisture sink as the residual of the moisture budget~~ (Yanai et al., 1973). The terms on the right-hand side of Eq. (1) correspond to the horizontal moisture advection, moisture convergence, vertical moisture flux, and the moisture source or sink associated with phase transitions (i.e., precipitation and evaporation) at the 8–30-day timescale.

The radiative effect is quantified using MERRA-2 data. The difference between net surface radiation and net top-of-atmosphere (TOA) radiation is defined as the atmospheric net radiative change, which includes both longwave and shortwave components. Longwave and shortwave radiation can be further categorized into clear-sky radiation and cloud radiative effects. Following Lin and Chen (2023), the radiative effects driven by aerosols and greenhouse gases are isolated linearly under assumed clear-sky and/or no-aerosol conditions:

$$\begin{cases} Cld_RE = Rad - Rad_{clr} \\ GHG_RE = Rad_{clr+no_aer} \\ Aer_RE = Rad_{clr} - Rad_{clr+no_aer} \end{cases}, \quad (2)$$

where Rad represents the atmospheric net radiation flux, Cld_RE is the cloud radiative effect, Aer_RE is the aerosol radiative effect, and GHG_RE is the greenhouse gas radiative effect. The subscripts “ clr ” and “ $clr+no_aer$ ” denote radiation fluxes under clear sky conditions and clear-sky/aerosol-free conditions, respectively. A positive value indicates a downward flux, contributing to atmospheric warming. Note that the Aer_RE in Eq. (2) captures only the direct radiative effect, as this method does not account for indirect aerosol effects on cloud properties and droplets formation (Lin and Chen, 2023).

The bootstrap method (Mudelsee et al., 2014) is applied to assess the statistical significance of results from observations and model experiments. The procedure consists of the following steps: first, paired resampled datasets are generated through random sampling with replacement from both experimental groups. Next, target statistical metrics are computed for each resampled pair. Finally, this process is

repeated 1,000 times, and the 90% confidence interval is determined by selecting the 5th and 95th percentile values.

2.3 WRF-Chem experiments

The covariation of aerosols and meteorological conditions complicates the attribution of causality between aerosol–cloud–radiation interactions and precipitation in southern China through observational analyses. To address this, we conducted a series of experiments using the WRF-Chem version 4.2.2 (Grell et al., 2005; Fast et al., 2006) to [verify-support](#) the observed mechanisms responsible for aerosol impacts on clouds and precipitation, [although uncertainties remain due to the dependence on emission inventories, physical parameterizations, and initial and boundary conditions](#). To increase computational efficiency, we adopted a single-domain configuration, consistent with previous studies focusing on aerosol–cloud–radiation feedbacks beyond weather timescales (e.g., Zhang et al., 2010; Chen et al., 2018). The model domain covers most of China (approximately 8°–43°N, 89°–140°E), with a horizontal resolution of 20 km and 40 vertical levels extending from the surface to 50 hPa.

The meteorological initial and boundary conditions are obtained from the National Centers for Environmental Prediction Final Analysis (NCEP-FNL) dataset, which is provided on a 1° × 1° grid and updated every six hours. Sea surface temperature (SST) data are sourced from the NCEP real-time global SST analysis. [Simulations using NCEP-FNL data as WRF-Chem input better capture the AOD evolution compared to those driven by higher-resolution ERA5 data \(figure not shown\)](#). To better reproduce the observed circulation [and aerosol](#) pattern, grid analysis nudging is applied ~~to constrain meteorological variables only~~ during the spin-up period [\(Abida et al., 2022\), following Yun et al. \(2024\), allowing meteorological fields to freely interact with aerosols during the analysis period. While nudging could potentially constrain aerosol feedbacks \(He et al., 2017\), our sensitivity tests confirm that it does not affect the main conclusions \(figure not shown\)](#).

The primary parameterization schemes used in the simulations are listed in Table 1, following the model configurations of Zhang et al. (2021). To comprehensively investigate aerosol indirect effects, the Morrison double-moment microphysics scheme (Morrison et al., 2009) and the Grell–Freitas cumulus scheme (Grell and Freitas, 2014) are employed. Shortwave and longwave radiation are parameterized using the Rapid Radiative Transfer Model for General Circulation Models (RRTMG) (Iacono et al., 2008) with aerosol radiative effects included, as the Morrison scheme enables direct communication of cloud droplet, ice, and snow effective radii to the radiation module.

Key chemistry options include the Carbon-Bond Mechanism version Z (CBMZ) scheme (Zaveri and Peters, 1999) for gas-phase chemistry, the Model for Simulating Aerosol Interactions and Chemistry (MOSAIC) 4-bin scheme (Zaveri et al., 2008) for aerosols, and the Fast-J scheme (Fast et al., 2006) for

196 photolysis. Anthropogenic emissions are derived from the Multi-resolution Emission Inventory for China
 197 (Li et al., 2017a) and the MIX inventory (Li et al., 2017b) for regions outside China. Biogenic emissions
 198 are calculated online using the Model of Emissions of Gases and Aerosols from Nature (MEGAN)
 199 (Guenther et al., 2012). Dust emissions are simulated using the original Goddard Chemistry Aerosol
 200 Radiation and Transport (GOCART) dust emission scheme (Ginoux et al., 2001). [Although the GOCART
 201 scheme may underestimate dust aerosol concentrations in Asia \(Zhao et al., 2020\), no emission scaling
 202 or tuning was applied in this study because our simulations focus on anthropogenic aerosols and summer
 203 rainfall in South China, where dust emissions contribute minimally.](#) Biomass burning emissions are
 204 obtained from the high-resolution fire emissions dataset based on the Fire Inventory from National Center
 205 for Atmospheric Research (NCAR) version 1.5 (Wiedinmyer et al., 2011).

206 Table 1. Main parameterization schemes used in the WRF-Chem simulations


Option name	Schemes	References
Microphysics	Morrison double-moment	Morrison et al. (2009)
Cumulus	Grell-Freitas scheme	Grell and Freitas (2014)
Longwave radiation	RRTMG	Iacono et al. (2008)
Shortwave radiation	RRTMG	Iacono et al. (2008)
Surface layer	Monin-Obukhov	Pahlow et al. (2001)
Land surface	Unified Noah	Chen et al. (2010)
Boundary layer	Yonsei University	Hong et al. (2006)
Aerosol chemistry	MOSAIC	Zaveri et al. (2008)
Gas chemistry	CBMZ	Zaveri and Peters (1999)
Photolysis	Fast-J	Fast et al. (2006)

207 The case selected for simulation is a persistent heavy rainfall event over South China in July 2015,
 208 lasting over five days and characterized by high aerosol loading before the onset of heavy precipitation.
 209 To minimize uncertainties in the modeling outcomes, we perform six ensemble simulations for each
 210 experiment by perturbing the initial and boundary conditions. Specifically, the ensemble members start
 211 at one-day intervals from 4 to 9 July 2015, with all simulations ending on 6 August 2015. [To allow locally
 212 emitted aerosols to become sufficiently mixed and reach a quasi-equilibrated distribution, we adopted
 213 spin-up times of 1–6 days, consistent with previous studies \(e.g., Zhu et al., 2022\).](#) The first few days of
 214 each run (4–9 July) are discarded ~~as spin-up time~~, and the analysis focuses on [10 July–6 August 2015](#).
 215 ~~the period from 10 July to 6 August 2015.~~

216 3 Observed aerosol effects on enhanced 8–30-day rainfall anomalies

217 In the East Asian monsoon region, the quasi-biweekly (8–30-day) oscillation is particularly strong
 218 in southern China (18°–32°N, 108°–123°E; box in Fig. 1a) and has been identified as a key trigger of
 219 persistent heavy rainfall events (Chen et al., 2024). Meanwhile, aerosol emissions, measured by AOD,
 220 also exhibit significant quasi-biweekly variability (Fig. 1b), despite their maximum concentrations
 221 occurring in North China.

222 To examine whether aerosols in southern China are correlated with local extreme precipitation, we
 223 analyze the evolution of AOD (blue curve in Fig. 1c) in relation to persistent heavy rainfall events (blue
 224 bars in Fig. 1c), which are defined as daily precipitation exceeding the 90th percentile for at least three
 225 consecutive days. An increase in AOD is observed before the onset of persistent heavy rainfall, followed
 226 by a decline during the heavy rainfall period due to wet deposition. The evolution of both the heavy
 227 rainfall and AOD closely follow their respective 8–30-day components (red and purple curves in Fig. 1c),
 228 indicating that both fields are strongly modulated by the quasi-biweekly oscillation. In contrast, neither
 229 AOD nor rainfall displays quasi-biweekly variability in composites of normal rainfall cases, where
 230 amplitudes remain around the climatological mean (45th–55th percentiles). No phase-leading relationship
 231 between AOD and rainfall is observed in these cases (Fig. 1d). The distinct evolution of AOD associated
 232 with different rainfall intensities suggests that the phase-leading increase in AOD could play a role in
 233 triggering persistent heavy rainfall events, with their connections primarily manifesting at the quasi-
 234 biweekly timescale.



235
 236 Figure 1. Spatial distributions of the standard deviation of (a) 8–30-day-filtered precipitation anomalies (mm d⁻¹) and (b)
 237 8–30-day-filtered aerosol optical depth anomalies (AOD; unitless) during MJJAS from 2000 to 2021, based on CPC
 238 rainfall data and MERRA-2 AOD data, respectively. The black boxes indicate the study domain of southern China (18°–
 239 32°N, 108°–123°E). (c) Composites of daily precipitation (light blue bars, left y-axis in light blue, mm d⁻¹), 8–30-day-

240 filtered precipitation (red curve, left y-axis in red, mm d^{-1}), AOD (blue curve, right y-axis in blue, unitless), and 8–30-
 241 day-filtered AOD (purple curve, right y-axis in purple, unitless) associated with persistent heavy rainfall events over
 242 South China. The x-axis is centered at “0”, representing the median timing of individual rainfall events, with negative
 243 and positive values indicating periods before and after the event. (d) As in (c) but for the composites of normal-intensity
 244 rainfall events.

245 Focusing on the quasi-biweekly timescale, we further examine whether the leading phase of AOD is
 246 consistently present and essential for enhanced 8–30-day rainfall anomalies, based on statistical analyses
 247 (Fig. 2). If not, we explore under what conditions AOD contributes positively to rainfall intensification.
 248 ~~Based on e~~Composite analysis; ~~shows that AOD exhibits positive anomalies beginning about~~starting six
 249 days before the ~~occurrence peak~~ of 8–30-day rainfall events ~~(red curve in Fig. 2a); defined as periods with~~
 250 ~~positive rainfall anomalies (Day 0; red curve in Fig. 2a; blue curve for AOD); AOD shows positive~~
 251 ~~anomalies (blue curve in Fig. 2a).~~ Among all 8–30-day rainfall events (249 cases), the majority (about
 252 87%) are preceded by a leading phase of AOD anomalies, supporting a potential causal role of antecedent
 253 AOD anomalies in subsequent precipitation enhancement. The consistent evolution of AOD from MODIS
 254 and ERA5 datasets (Figs. S1a–b) demonstrates the robustness of intraseasonal AOD variations associated
 255 with heavy rainfall events. Validation ~~using multiple~~with long-term reconstructed gridded datasets of
 256 particulate matter ~~with aerodynamic diameter less~~smaller than 2.5 ~~micrometer~~ (PM_{2.5}) confirms these
 257 findings ~~(Fig. S1c)(figure not shown)~~, indicating that the observed leading phase of aerosols is robust,
 258 regardless of whether AOD or PM_{2.5} is usedusing AOD as ~~the~~a proxy for aerosol concentrations ~~variations~~.
 259 ~~While the two AOD datasets show high consistency (blue shading in Fig. 2a), MERRA-2 AOD is used~~
 260 ~~in the following observational analysis, which can capture the dynamic variations in AOD (Shi et al.,~~
 261 ~~2019; Choudhury et al., 2020), since MODIS AOD is only available under cloud-free conditions.~~

262 The composite analysis in Fig. 2a illustrates the overall lead-lag phase relationship between AOD
 263 and rainfall anomalies but does not directly quantify their correlation. To further assess this correlation,
 264 Fig. 2b examines how subsequent rainfall anomalies vary with the amplitude of preceding AOD
 265 anomalies during Days –6 to –1, considering all 8–30-day rainfall events. Although the correlation
 266 coefficient between the two parameters is relatively low (0.12; red line in Fig. 2b), it is statistically
 267 significant at the 90% confidence level due to the large sample size. A visual inspection suggests that this
 268 positive correlation is not well established when rainfall anomalies are weak (symbols to the left of the
 269 dashed line in Fig. 2b). However, when focusing on rainfall events with greater amplitude, the correlation
 270 becomes stronger. Specifically, for events with rainfall amplitude exceeding 0.25 standard deviation (σ)
 271 (blue line, symbols to the right of the dashed line in Fig. 2b), the correlation coefficient increases to 0.25,
 272 which is significant at the 95% confidence level. These results indicate that the effect of preceding AOD
 273 increases on subsequent rainfall enhancement is not linear.

To determine the threshold of rainfall amplitude significantly modulated by AOD magnitudes, we further compute the correlation coefficients between AOD anomalies preceding rainfall events and the amplitude of subsequent rainfall anomalies by categorizing events into different amplitude bins (x -axis in Fig. 2c). The results show that the correlation coefficient reaches 0.2, becoming significant at the 90% confidence level, when rainfall anomalies exceed the mean of all events (i.e., pink bar on the x -axis is at 0). The coefficient further increases to 0.25, surpassing the 95% confidence level, for events with rainfall amplitudes above 0.25σ . The correlation is even stronger, reaching 0.39, for heavy rainfall events with amplitudes exceeding 1σ (rightmost pink bar in Fig. 2c). These findings suggest that higher aerosol concentrations before rainfall occurrence are conducive to rainfall amplification, particularly for events with above-average amplitude. This behavior aligns with previous studies at the synoptic and decadal timescales (Wang et al., 2011; Yang et al., 2018; [Su et al., 2020](#); Shao et al., 2022; Xiao et al., 2023b), which emphasize that the aerosols tend to suppress light rainfall while enhancing heavy convective precipitation.

The statistical results in Fig. 2 indicate that AOD anomalies have a more pronounced impact on 8–30-day precipitation events exceeding a certain intensity threshold (e.g., 0.25σ), with stronger antecedent AOD anomalies leading to amplified subsequent precipitation anomalies, [although the correlation coefficients are modest \(\$r=0.25\text{--}0.39\$ \). These modest values highlight the complexity of rainfall intensification mechanisms, which involve circulation anomalies that may be induced by, or independent of, AOD variations](#). To investigate the physical processes through which AOD contributes to rainfall intensification, we compared enhanced 8–30-day rainfall cases (amplitude greater than 0.25σ) under varying AOD intensities prior to the event, categorized as High AOD–Strong Precipitation (HA–SP) and Low AOD–Strong Precipitation (LA–SP) cases. Thus, preceding AOD cases are classified as high or low based on whether the standardized AOD anomaly is positive or negative. [To distinguish clean and polluted conditions and ensure balanced sample sizes](#)~~To ensure balanced sample sizes~~, LA–SP cases with AOD anomalies between ~~0 and~~ -0.4σ [and 0 \(approximately 40th–60th percentiles\)](#) were excluded (Fig. 2b).




Figure 2. (a) Evolution of 8–30-day precipitation (red curve, mm d^{-1}) and 8–30-day AOD (blue curve, unitless) associated with quasi-biweekly precipitation events, defined as periods with positive 8–30-day rainfall anomalies over South China. [These results are based on CPC rainfall data and MERRA-2 AOD data](#). Day 0 ~~represents~~[denotes](#) the

occurrence peak of rainfall events, while negative and positive values on the x-axis indicate periods days before and after the event peak, respectively. Light blue shading marks the range of one standard deviation (σ) across the two AOD datasets from MERRA-2 and MODIS. (b) Scatterplot of rainfall intensity (i.e., 8–30-day-filtered rainfall anomalies at Day 0; x-axis, mm d⁻¹) versus preceding AOD intensity (represented by the peak value of 8–30-day-filtered AOD anomalies during Days –6 to –1; y-axis, unitless), with both variables normalized by their climatology. The red line represents the linear regression fitted to all cases, while the blue line corresponds to events where rainfall intensity exceeds 0.25σ . The correlation coefficients between the two variables are shown in the bottom right corner, with single and double asterisks indicating significance at the 90% and 95% confidence levels, respectively. Enhanced 8–30-day rainfall events ($>0.25\sigma$) are classified into High AOD–Strong Precipitation (HA–SP; cross sign) and Low AOD–Strong Precipitation (LA–SP; plus sign) categories, with case counts displayed in each quadrant. To maintain nearly equal sample sizes for these two groups, the LA–SP classification is determined using an AOD anomaly threshold of -0.4σ (blue dashed line and blue text). (c) Correlation coefficients between preceding AOD intensity and rainfall intensity at Day 0 for different rainfall amplitude thresholds (bins with 0.25σ intervals). Pink bars represent correlations when 8–30-day rainfall exceeds a given threshold, while blue bars show correlation coefficients for the remaining events. Hatching indicates correlations significant at the 90% confidence level.

Composite patterns of convection and circulation during the early and peak phases of HA–SP and LA–SP events, along with their differences, are shown in Fig. 3 based on MERRA-2 data, and are consistent with those derived from ERA5 (Fig. S2). At four days before the occurrence of heavy rainfall events (Day –4), both types of events exhibit suppressed convection of similar magnitude but with distinct wind anomalies (contours and vectors in Figs. 3a, d, g), which likely explain their differing AOD behaviors. HA–SP events are associated with anomalous northerlies that transport aerosols from North China to South China, increasing regional AOD, whereas LA–SP events exhibit an opposite transport pattern. Moreover, the anomalous moisture convergence (vectors in Fig. 3g) coincides with significant AOD enhancement (shading in Fig. 3g) and intensified convection over coastal Southern China (contours in Fig. 3g), implying the aerosol effects on moistening process. Approximately two days before the heavy rainfall events, convective anomalies (indicated by negative OLR anomalies) were observed over South China in both cases (contours in Figs. 3b, e). However, in high AOD cases (HA–SP), convection and moisture convergence signals are noticeably stronger than in low AOD cases (LA–SP) (Fig. 3h). At the peak of the heavy rainfall event (Day 0), strong convection, cyclonic circulation anomalies, and significant moisture convergence dominate South China (Figs. 3c, f), coinciding with AOD reductions due to wet scavenging. Notably, convective anomalies at Day 0 are more intense in HA–SP events than in LA–SP events (Fig. 3i), suggesting that preceding positive AOD anomalies may amplify subsequent convective activity through enhanced moisture convergence.

Based on the moisture budget diagnosis, we further examine the processes responsible for increasing moisture and enhancing rainfall amplitude in HA–SP cases, comparing the budget terms with those in LA–SP events (Figs. 4a–c). In both cases, moisture convergence ($-\langle q\nabla \cdot \mathbf{V} \rangle'$, green curves in Figs. 4a–b)

demonstrated a growth of $2\text{--}3\text{ mm d}^{-1}$ at Day 0, accounting for $\sim 50\%$ of the positive rainfall anomalies
 (red curves in Figs. 4a–b) and ~~serv~~inges as the primary moisture source. Moreover, moisture convergence
 also explains~~also accounting for~~ the differences in rainfall amplitude between HA–SP and LA–SP events
 (green curve in Fig. 4c), which exhibited a significant increase of $1\text{--}1.8\text{ mm d}^{-1}$ during Days -4 to 0 . The
 moisture sink associated with latent heating is in phase with rainfall and offsets the moisture source from
 convergence with a reduction of $0.9\text{--}1.6\text{ mm d}^{-1}$ during Days -3 to 1 (cyan curve in Fig. 4c). Nevertheless,
 their combined effect still contributes positively to heavy rainfall occurrence (red curve in Fig. 4c). The
 temporal evolution of these key terms reveals their sequential influence. ~~The increase in AOD~~ An
increment of ~ 0.1 in AOD at Day -4 slightly precedes the enhancement of moisture convergence (blue
 and green curves in Fig. 4c), indicating that the aerosols could play roles in moistening process, which
 may subsequently lead to intensification of $0.7\text{--}1\text{ mm d}^{-1}$ in ~~intensified~~ rainfall during Days -2 to 4 (red
 curve in Fig. 4c). At Day 0, the intensity of quasi-biweekly precipitation is increased by $\sim 20\%$. The
horizontal moisture advection and vertical moisture flux (magenta curves and pink curves in Figs. 4a–b)
~~Other moisture processes have~~make relatively minor contributions, and their differences between HA–
 SP and LA–SP events are not statistically significant. Overall, the key moisture processes associated with
quasi-biweekly precipitation events and their modulation by aerosols are consistently shown in the ERA5
reanalysis (Figs. S3a–c).





Figure 3. (a)–(c) Composite evolution of 8–30-day outgoing longwave radiation (OLR; purple contours, W m^{-2}), AOD (shading, unitless), and column-integrated moisture flux (vectors, $\text{kg m}^{-1} \text{s}^{-1}$) at 4, 2, and 0 days before the occurrence of HA–SP events, respectively. [The results are based on NOAA OLR data, MERRA-2 AOD and meteorological data.](#) (d)–(f) and (g)–(i) Similar to (a)–(c), but for the composite results of LA–SP events and the differences between HA–SP and LA–SP events, respectively. Only AOD and moisture flux anomalies with statistically significant changes at the 90% confidence level are shown. The red box outlines the study domain of South China.

Building on the observed lead-lag relationship between aerosols and moistening process (Figs. 4a–c), the next question is: through what physical processes do aerosols enhance moisture convergence and rainfall? Previous studies have shown that aerosols primarily influence summer precipitation in South China through cloud microphysical effects, as clouds in this region are readily invigorated by the aerosols which are less absorbing and highly hygroscopic (Fan et al., 2008; Yang et al., 2016). To investigate this mechanism, we analyzed several key cloud-related parameters for HA–SP and LA–SP events (Figs. 4d–f). Although the temporal evolution of cloud properties is similar in both cases (Figs. 4d–e), their magnitudes exhibit notable differences (Fig. 4f). [At Day –4, the anomalous liquid cloud fraction increased significantly \(purple curve in Fig. 4f\).](#) The most ~~intensesignificant~~ increase prior to heavy rainfall in HA–SP events is observed in the ice water path (orange curve in Fig. 4f), [which rises by 22.2–26.8 \$\text{g m}^{-2}\$](#)

during Days -3 to -1. This magnitude of increase is comparable to the cloud water path enhancements under pollution reported in Zhou et al. (2020). following an increase in liquid cloud fraction (purple curve in Fig. 4f). These changes are likely linked to precursor moisture convergence anomalies (green curve in Fig. 4c), which enhance latent heating in the lower troposphere. The resulting low-level warming destabilizes atmospheric stratification, facilitating the uplift of abundant cloud droplets above the freezing level. Consequently, the ice water path increases markedly (orange curve and gray asterisk in Fig. 4f), whereas the ice cloud fraction shows a modest enhancement of ~ 0.01 at Day -3. The statistically while the modest and insignificant increase in ice cloud fraction may reflect a limited reduction in the liquid cloud fraction and persistent supercooled droplets. These conditions favor interactions between supercooled droplets and cloud ice particles.

Subsequently, deep convective clouds develop, characterized by a reduction in cloud top pressure (magenta curve in Fig. 4f) and an increase in ice cloud fraction (light blue curve in Fig. 4f), alongside a gradual weakening of the liquid cloud fraction (purple curve in Fig. 4f). Additional latent heat released from freezing and deposition could further enhance upward motion, collectively amplifying the intensity of 8–30-day heavy precipitation events. Note that both cases exhibit positive anomalies in the effective radius of cloud droplets before the onset of heavy precipitation (cyan curves in Figs. 4d–e), although the difference between them is not statistically significant. This suggests that aerosol impacts on the warm-rain processes are not the primary driver of the 8–30-day rainfall enhancement, with aerosol indirect effects likely exerted through ice-phase microphysical processes.

In addition to aerosol–cloud interactions, changes in liquid and ice cloud fractions also influence atmospheric radiation variations. The right panels of Fig. 4 illustrate the longwave and shortwave radiative effects induced by clouds, aerosols, and greenhouse gases. During HA–SP events, the dominant longwave cloud-radiative effect closely follows the temporal evolution of 8–30-day precipitation anomalies (magenta curve in Fig. 4g), supporting the maintenance of heavy precipitation intensity (Chen et al., 2024). This behavior is consistent with estimates from CERES-SYN (Fig. S4). Although the amplitude is relatively small, the significant increase in GHG-induced longwave heating prior to heavy rainfall (orange curve in Fig. 4i) may be associated with the preceding phase of moistening driven by moisture convergence anomalies (green curve in Fig. 4c). Focusing on the aerosol effect, the aerosol-induced direct shortwave atmospheric heating (light blue curve in Fig. 4i) is small but in phase with AOD anomalies (blue curve in Fig. 4c). This suggests that some absorbing aerosols, such as anthropogenic black carbon, may contribute to shortwave radiation absorption over southern China. However, due to the predominance of nonabsorbing aerosols in this region (Lee et al., 2007; Huang et al., 2014; Yang et al., 2016), the resultant aerosol shortwave radiative forcing (Aer_SW) remains relatively weak (light blue curve in Fig. 4i).



408

409 Figure 4. As in Fig. 2a, but for the composite evolution of 8–30-day (a) precipitation (red curve; left y-axis, mm d⁻¹),
 410 AOD (blue curve; right y-axis in blue, unitless), and individual moisture budget terms (various colored curves
 411 representing each budget term in Eq. (1); left y-axis, mm d⁻¹) associated with HA–SP events. (d, g) As in (a), except that
 412 (d) shows the 8–30-day evolution of cloud top pressure (CTP; magenta curve; left y-axis, hPa), ice water path (IWP;
 413 orange curve; left y-axis, g m⁻²), cloud droplet effective radius (CER; orange curve; right y-axis, μm), liquid cloud
 414 fraction (CF_Liquid; purple curve; right y-axis, unitless), and ice cloud fraction (CF_Ice; light blue curve; right y-axis).
 415 Panel (g) presents 8–30-day column-integrated latent heat heating (Q_2 ; green curve; left y-axis in green, W m⁻²) and 8–
 416 30-day radiative budget terms calculated from Eq. (2) (right y-axis, W m⁻²), including longwave/shortwave cloud
 417 radiative effects (Cld_LW/Cld_SW; magenta curve/pink curve), longwave/shortwave aerosol direct radiative effects
 418 (Aer_LW/Aer_SW; cyan curve/light blue curve), and longwave/shortwave greenhouse gas radiative effects
 419 (GHG_LW/GHG_SW; orange curve/gray curve). (b, e, h) and (c, f, i) are similar to (a, d, g), but represent the composite
 420 results for LA–SP events and the differences between HA–SP and LA–SP events, respectively. In panels (c) and (i), only
 421 terms with statistically significant differences at the 90% confidence level are shown, with significant periods marked
 422 by gray asterisks. [These results are based on MERRA-2 budget variables and MODIS cloud parameters.](#)

423 [Latent heating, a direct product of phase transition processes, serves as an indicator of aerosol effects](#)
 424 [on cloud microphysical properties \(Zhu et al., 2024; Fan et al., 2025\).](#) Both the aerosol-induced direct
 425 radiative effects (light blue curve in Fig. 4i) and longwave cloud-radiative effects (magenta curve in Fig.
 426 4i), with magnitudes of approximately 2–5.5 W m⁻², are significantly smaller than atmospheric latent
 427 heating associated with moisture processes, which exceeds 430 W m⁻² (green curve in Fig. 4i).
 428 [Quantitatively, aerosol-induced latent heating is approximately seven times greater than aerosol-induced](#)

longwave radiative heating. The relative contributions and quantitative ratios between the two variables are similarly shown in the ERA5 data (Figs. S3d–f). This indicates that aerosol–cloud microphysical effect plays a dominant role in enhancing heavy precipitation at the 8–30-day timescale, as demonstrated by the diagnostic frameworks of moisture and radiative budget analyses. The latent heating magnitude ($\sim 40 \text{ W m}^{-2}$) is also comparable to values reported in previous studies of tropical intraseasonal convection ($\sim 60 \text{ W m}^{-2}$; Fu et al., 2020) and monsoonal extreme precipitation ($\sim 50 \text{ W m}^{-2}$; Chyi et al., 2023).

To confirm the stepwise contribution of aerosol-induced low-level moistening to high-level freezing and deposition, ultimately leading to deep convection and rainfall intensification, as discussed in Fig. 4, we further analyze the time-height evolution of latent heating, vertical velocity, and liquid/ice cloud water content differences between HA–SP and LA–SP events (Fig. 5). As shown in Fig. 5a, low-level liquid cloud water content increases below 500 hPa, coinciding with peak AOD at four days (Day –4) before the occurrence of heavy rainfall events (Fig. 4). The condensation process in shallow convection releases latent heat, forming a positive latent heating anomaly centered at 850–700 hPa (Fig. 5c), which enhances low-level moisture convergence and upward motion (Fig. 5d). This process provides favorable conditions for the uplift of cloud droplets above the freezing level. From Days –3 to –1, strong latent heating (Fig. 5c) and ascending motion anomalies (Fig. 5d) develop above 500 hPa, corresponding to the maxima of increased ice cloud water content in the middle and upper troposphere (Fig. 5b). The enhanced intensified conversion of supercooled cloud water to ice hydrometeors through freezing and deposition further releases latent heat, reinforcing vertical motion and convection. Ultimately, these processes ultimately lead to the intensification of intensify rainfall anomalies, a finding that is also confirmed by ERA5 (Fig. S5).




Figure 5. Height–time evolution of 8–30-day anomalies in (a) liquid cloud water content (CLWC; 10^{-6} kg kg $^{-1}$), (b) ice cloud water content (CIWC; 10^{-6} kg kg $^{-1}$), (c) latent heat heating (K d^{-1})($\text{J kg}^{-1} \text{h}^{-1}$), and (d) vertical velocity anomalies (-10^{-2} Pa s $^{-1}$) between the HA–SP and LA–SP events, based on the MERRA-2 data. Stippling denotes differences statistically significant at the 90% confidence level. Vertical blue lines denote the peak timing of heavy rainfall event occurrence (Day 0).

4 Support for Verification of mechanisms using WRF-Chem experiments

Due to the complex interactions between aerosols and rainfall, as well as the limited availability of observational data on cloud microphysical properties, we employ the WRF-Chem model to verify the mechanisms identified in Sect. 3. Following the methodology of Guo et al. (2022) and Yun et al. (2024), we conduct two sets of simulations to assess the effects of anthropogenic aerosols. In the control experiment (CTRL), anthropogenic aerosol emissions are kept unchanged, while in the sensitivity experiment (CLEAN), emissions are scaled to 10% of those in the CTRL run. Hence, we examine differences in cloud microphysics, radiation, and surface rain rates between the CTRL and CLEAN simulations.




Figure 6. Normalized occurrence frequency Probability density functions (PDFs) of (a) daily precipitation (mm d^{-1}) and (b) AOD (unitless) from observations (pink curves, CPC rainfall and MODIS AOD data) and the CTRL experiment (black curves) during the model integration period over the key region of interest (21° – 24°N , 111° – 116°E) in South China.

Before conducting sensitivity experiments, we evaluate the reliability of the model simulations by comparing the occurrence frequencyprobability density functions of precipitation and AOD from the CTRL experiment with observations over the key region of South China (21° – 24°N , 111° – 116°E), where the primary 8–30-day heavy precipitation event occurred in July 2015 (Fig. 6). The precipitation distribution from the CTRL simulation (black bars in Fig. 6a) generally resembles the observed distribution (pink bars in Fig. 6a).However, CTRL yields a mean precipitation of $\sim 10.2 \text{ mm d}^{-1}$ compared with $\sim 7.7 \text{ mm d}^{-1}$ in CPC over the key region, corresponding to an overestimation of about

32%.although it tends to slightly overestimate precipitation intensity. This bias is common in regional climate models, including WRF, and is attributable to limitations in convective cloud and microphysical parameterizations (Caldwell et al., 2009; Argüeso et al., 2012). Additionally, the CTRL simulation captures the overall pattern of the observed AOD distribution (Fig. 6b), though it underestimates the mean AOD by ~35%the simulated peaks are underestimated (Fig. 6b). This discrepancy is likely due to uncertainties in emission inventories and biases in meteorological fields (Huang and Ding, 2021). Despite these biases, the CTRL experiment demonstrates reasonable skill in reproducing the observed precipitation and AOD distributions over the study area.




Figure 7. Evolution of (a) precipitation (mm d^{-1}) and (b) AOD (unitless) anomalies averaged over the Pearl River Delta (21° – 24°N , 111° – 116°E) in the observations (black curves, CPC rainfall and MODIS AOD data), CTRL (blue curves) and CLEAN (red curves) experiments. Colored shading represents the range of 0.5σ across the six ensemble members.

By comparing rainfall and AOD anomalies—both subjected to a 5-day running mean to remove high-frequency synoptic signals and focus on the subseasonal timescale—between the CTRL and CLEAN experiments, the observed relationship between preceding AOD and heavy rainfall is confirmed (Fig. 7). Notably, the CTRL experiment captures the temporal evolution of precipitation and AOD anomalies reasonably well, despite slight biases in the timing of their peaks. Quantitatively, the simulated precipitation anomalies are overestimated by about 6.5 mm d^{-1} relative to CPC observations during the heavy rainfall case on 21–23 July 2015 (blue curve and shading in Fig. 7a), and the model fails to reproduce the observed precipitation peak on 19 July. Since the analyses are based on composites aligned to the precipitation peak (Day 0), the results are not sensitive to small timing offsets of 1–2 days between model and observations. The robust feature is that positive aerosol anomalies consistently precede

enhanced quasi-biweekly precipitation (black and blue curves in Figs. 7a–b), supporting our main conclusions. An increase in AOD precedes the intensification of rainfall (blue curves in Figs. 7a–b), consistent with the observations (black curves in Figs. 7a–b). This further supports and the reliability of the model simulations employed in this study. When AOD is suppressed in the CLEAN experiment, rainfall amplitude decreases accordingly (red curves in Figs. 7a–b). The most significant reduction in rainfall amplitude occurs when AOD declines notably between 21–23 July, which will be the focus of the following analyses.

To verify whether aerosols drive rainfall intensification through the aerosol–cloud interactions identified in Sect. 3, we compare these cloud-related variables between the CTRL and CLEAN simulations (Fig. 8). When aerosols are enhanced (transition from CLEAN to CTRL), the key region exhibits an increase in column cloud fraction (Fig. 8a), with distinct positive anomaly centers in the lower troposphere (925 hPa) and upper troposphere (200 hPa). The increase in low-level cloud fraction is attributed to aerosol-induced enhancement in cloud droplet number concentration (figure not shown) and liquid cloud water content (Fig. 8b). Between 700–400 hPa, higher supersaturation promotes increased cloud water formation (Fig. 8b), consistent with observational findings (Fig. 5b). The enhanced upper-level cloud fraction is associated with an increase in ice-phase hydrometeors (Figs. 8d–f), where cloud ice likely forms through cloud water freezing and vapor deposition. Meanwhile, snow growth appears to be driven by depositional processes and the riming of abundant cloud droplets. Although graupel mixing ratios also increase, their magnitude is much smaller compared to the dominant snow mixing ratios. Overall, cold-phase and mixed-phase processes amplify atmospheric rain mixing ratios (Fig. 8c), confirming the observed aerosol-mediated precipitation intensification through enhanced cloud water and cloud ice pathways (Figs. 4–5).




Figure 8. Vertical profiles of (a) cloud fraction anomalies (unitless) and (b)–(f) hydrometeor mixing ratio anomalies ($10^{-6} \text{ kg kg}^{-1}$) averaged over the key region for the CTRL and CLEAN experiments during 21–23 July 2015: (b) cloud

524 water anomalies, (c) raindrop anomalies, (d) cloud ice anomalies, (e) snow anomalies, and (f) graupel anomalies. Colored
525 shading represents $\pm 0.5\sigma$ across the six ensemble members.

526 To examine the stepwise modulation of heavy rainfall occurrence by aerosol-induced low-level
527 moistening and deep convection development, we compare the vertical profiles of latent heating
528 anomalies, vertical velocity anomalies, and moisture convergence anomalies between the CTRL and
529 CLEAN simulations (Figs. 9a–c). When aerosols are enhanced, the increased latent heating rate exhibits
530 two distinct centers at 850 hPa and 400 hPa (Fig. 9a), consistent with observed profiles (Fig. 5a). These
531 heating centers correspond to enhanced low-level cloud water and increased high-level cloud ice and
532 snow concentrations (Fig. 8), where latent heat is released through condensation at lower levels and
533 freezing and riming processes at upper levels. Thus, aerosols intensify latent heating through cold-phase
534 and mixed-phase processes, strengthening upward motion throughout the atmosphere (Fig. 9b), similar
535 to the observed patterns (Fig. 5c). Simultaneously, aerosol-induced enhanced low-level moisture
536 convergence anomalies (Fig. 9c) further increase supersaturation and latent heating, amplifying the
537 intensity of rainfall anomalies.

538 Based on observational diagnostics, we suspect that aerosol–cloud microphysical processes
539 contribute more significantly to rainfall intensification than aerosol–cloud radiative effects (Fig. 5i). To
540 verify this, we analyze the vertical profiles of longwave radiative heating anomalies (Fig. 9d) in the CTRL
541 and CLEAN experiments, along with changes in longwave and shortwave radiation fluxes (Figs. 9e–f).
542 ~~Compared with CLEAN experiment~~~~In the CTRL experiment~~, longwave radiative heating exhibits
543 ~~negligible~~~~minimal~~ changes below 925 hPa, but ~~generally~~ increases by $0.1\text{--}0.3\text{ K d}^{-1}$ between 850–500
544 hPa and above 400 hPa in the CTRL experiment (Fig. 9d), consistent with ~~the vertical profile of~~ enhanced
545 cloud fractions (Fig. 8a). To quantify cloud contributions to atmospheric longwave heating, we calculate
546 the longwave cloud-radiative effect using all-sky and clear-sky radiative fluxes based on Eq. (2) (gray
547 bars in Fig. 9e). The development of deep convection triggered by enhanced aerosol emissions reduces
548 OLR by $\sim 10\text{ W m}^{-2}$ at the TOA, ~~while~~ increases atmospheric longwave radiation by $\sim 9\text{ W m}^{-2}$, and
549 raises downward longwave radiation by $\sim 1\text{ W m}^{-2}$ at the surface. For shortwave radiation, the increased
550 cloud cover reflects more shortwave radiation back to the TOA (gray bars in Fig. 9f). Additionally, the
551 dominance of scattering aerosols ~~(e.g., sulfate aerosols)~~ during this rainfall event, as indicated by the high
552 single-scatter albedo, contributes to a small shortwave radiation flux of $\sim 2\text{ W m}^{-2}$ through atmosphere
553 (blue bars in Fig. 9f).

554 The reduction in latent heating is much larger than that in radiative heating, with values of $\sim 0.8\text{ K}$
555 d^{-1} for vertical-mean (1000–100 hPa) latent heating compared to $\sim 0.2\text{ K d}^{-1}$ for column longwave heating
556 in the CLEAN experiment relative to CTRL. These modeling results align with observations, showing
557 that aerosol-induced latent heating rates are stronger than aerosol-induced radiative heating rates (Figs.

9a, d). This suggests that aerosol-driven cloud microphysical processes dominate the total heating contribution, while radiative processes play a secondary role (Fig. 4). [To test the sensitivity of our results to nesting and spatial resolution, we conducted additional high-resolution experiments with nested domains of 20 km and 4 km \(Fig. S6a\). These nested simulations consistently demonstrate that aerosols enhance rainfall intensity \(Figs. S6b–c\) and that aerosol–cloud microphysical effects remain dominant \(Fig. S7\). Importantly, the ratio of aerosol-induced latent heating to longwave radiative heating remains close to 4:1 \(Figs. S7e–f\). However, larger biases in AOD and precipitation were evident in the nested simulation \(Figs. S6b–c\), likely due to error transmission from the parent grid and accumulated uncertainties introduced at finer resolution during long-term integrations \(Baklanov et al., 2014; Wang et al., 2016\). The main conclusions remained robust in high-resolution simulations with two nested domains \(figure not shown\).](#)




Figure 9. As in Fig. 8, but for the vertical profiles of (a) latent heat flux anomalies (K d^{-1}), (b) vertical velocity anomalies (m s^{-1}), (c) moisture convergence anomalies ($10^{-2} \text{ kg kg}^{-1} \text{ s}^{-1}$), and (d) longwave radiative heating rate anomalies (K d^{-1}) in the CTRL and CLEAN experiments. (e) Anomalous longwave radiation fluxes (W m^{-2}) averaged over the key region at the top of the atmosphere (TOA), within the whole atmosphere (ATM), and at the surface (SFC) under all-sky (pink bars), clear-sky (blue bars), and cloudy-sky (gray bars) conditions. (f) Similar to (e), but for anomalous shortwave radiation fluxes. Colored bars represent the ensemble mean from six simulations, while whiskers indicate the range of $\pm 0.5\sigma$ across the six ensemble members.


5 Summary and discussion

South China, a densely populated region, frequently experiences persistent heavy rainfall events that are closely linked to 8–30-day rainfall anomalies. This study investigates how anthropogenic aerosols intensify these quasi-biweekly rainfall anomalies using observational diagnostics and model simulations. Statistical analysis reveals that aerosol concentrations tend to increase approximately four days before the occurrence of rainfall anomalies. The phase-leading increase in AOD is positively correlated with subsequent rainfall intensity, primarily by enhancing moisture convergence, which strengthens deep

584 convection and precipitation. The evolution of moisture sources, cloud properties, and radiative processes
585 suggests that aerosol–cloud microphysical effects play a dominant role in rainfall enhancement, while
586 radiative feedbacks have a secondary influence. These processes are summarized in Fig. 10.

587 For aerosol–cloud microphysical effects (red and pink sectors in Fig. 10), increased aerosol
588 concentrations enhance CCN activation, promoting the accumulation of low-level cloud water. Latent
589 heat release from the condensation of low-level cloud water induces localized warming, which reduces
590 surface pressure anomalies, strengthens upward motion, and enhances moisture convergence. This
591 facilitates the uplift of cloud droplets above the freezing level, where additional latent heat release from
592 cold-phase and mixed-phase processes (freezing and deposition) further amplifies ascending motion,
593 leading to stronger deep convection and intensified rainfall anomalies. In addition to microphysical effects,
594 aerosols in southern China can absorb shortwave radiation, while aerosol-induced changes in water vapor
595 and cloud cover also contribute to increased longwave radiation. These radiative processes are illustrated
596 on the right side of Fig. 10 (blue and light blue sectors). The resulting radiative heating provides additional
597 energy for deep convection and enhances moisture flux, further supporting rainfall development.

598 WRF-Chem experiments for a heavy rainfall event in July 2015, conducted under different aerosol
599 emission conditions, confirm the observational findings. Under CLEAN conditions, where anthropogenic
600 aerosol concentrations are ~~greatly~~ reduced by 90%, precipitation anomalies ~~decrease by ~7 mm d⁻¹~~ weaken
601 due to suppressed cold-phase and mixed-phase processes during the heavy rainfall case on 21–23 July
602 2015 over the key region. In contrast, in the CTRL experiment, increased aerosols appear before enhanced
603 rainfall occurrence, consistent with observations. Enhanced low-level cloud water and mid-to-upper-level
604 ice hydrometeors lead to the formation of dual latent heating centers at 850 hPa and 400 hPa, which drive
605 stronger low-level moisture convergence and column-wide upward motion, ultimately intensifying
606 precipitation anomalies. While longwave cloud-radiative effects marginally increase atmospheric moist
607 static energy, their contribution to precipitation anomalies is relatively weaker. Quantitatively, the column
608 latent heating rate is reduced by ~0.8 K d⁻¹ when anthropogenic aerosols are removed, approximately four
609 times greater than the reduction in column longwave radiative heating (~0.2 K d⁻¹). These modeling
610 results further corroborate the aerosol-driven intensification of quasi-biweekly precipitation observed in
611 South China.



612

613 Figure 10. Schematic diagram illustrating the mechanisms of aerosol effects on heavy rainfall events at the 8–30-day
614 timescale. The symbol “↑” denotes the enhancement of the corresponding processes, “ATM” refers to the whole
615 atmosphere, and “>” indicates stronger contributions of the process to heavy precipitation events.

616 Numerous studies on synoptic and long-term aerosol effects have reported that cloud liquid
617 accumulated by aerosols is converted into ice hydrometeors above the freezing level, thereby invigorating
618 deep convection and intensifying heavy precipitation, the so-called invigoration effect (e.g., Rosenfeld et
619 al., 2008; Li et al., 2011; Hazra et al., 2013; Guo et al., 2014). Aligning with this framework, our
620 observational analysis shows that aerosols similarly enhance cold-phase and mixed-phase cloud
621 development at the 8–30-day timescale, leading to further intensification of heavy precipitation. Our
622 findings contribute to a deeper mechanistic understanding of aerosol–cloud–radiation interactions in
623 modulating precipitation anomalies at the intraseasonal timescale over southern China. Note that the
624 positive contribution of increased AOD to rainfall intensification contrasts with previous studies on the
625 Indian summer monsoon, which reported a link between increased 20–100-day-filtered AOD and the
626 intensification of monsoon breaks (Arya et al., 2021; Surendran et al., 2022). This discrepancy may be
627 attributed to differences in aerosol types and concentrations between the two monsoon regions. In the
628 Indian monsoon region, rainfall suppression has been associated with the indirect effect of dust aerosols
629 through reduced cloud effective radius and decreased precipitation efficiency, whereas nonabsorbing
630 aerosols are more prevalent in southern China (Lee et al., 2007; Huang et al., 2014). Furthermore, Singh
631 et al. (2019) showed that a simultaneous enhancement of dust emissions from West Asia and the Tibetan
632 Plateau intensify the Indian monsoon rainfall with 10–20-day periodicity, but remarkably decrease the
633 spatial scale of the 30–60-day rainfall. Hence, it would be valuable to investigate whether aerosol impacts

634 vary across intraseasonal timescales in southern China, such as the 8–30-day and 20–100-day bands,
635 which we leave for future study.

636 Given that intraseasonal variability serves as a key source of subseasonal predictability for extreme
637 events (e.g., Wei et al., 2024; Xie et al., 2024), which remains a challenge for both the scientific and
638 operational communities, these results provide valuable insights for improving subseasonal prediction
639 skill by refining aerosol–cloud microphysical processes in prediction models. However, the extent to
640 which aerosols influence subseasonal prediction skill for extreme events requires further investigation.
641 Modulated by intraseasonal convective and circulation anomalies, aerosols exhibit significant variability
642 at the subseasonal timescale (e.g., Tian et al., 2011; Reid et al., 2015; Yu and Ginoux, 2021). However,
643 the factors controlling aerosol loading at the quasi-biweekly timescale in South China remain an open
644 question. ~~At this timescale, the quantitative contributions of aerosol radiation and aerosol–cloud~~
645 ~~interactions to heavy rainfall events also warrant further in-depth exploration (e.g., Liu et al., 2020; Zhang~~
646 ~~et al., 2021).~~ Another unresolved issue concerns quantitative attribution. This study provides an initial
647 estimate, but further refined experiments with different aerosol emission scenarios, parameterization
648 schemes, and background conditions are needed for more precise results. In particular, the relative
649 contributions of distinct aerosol pathways to intraseasonal precipitation should be determined by
650 selectively deactivating aerosol–radiation and aerosol–cloud interactions in WRF-Chem simulations (e.g.,
651 Liu et al., 2020; Zhang et al., 2021). These aspects are the focus of our ongoing research.

652

653 **Data Availability.** The source codes of the WRF-Chem model are available on the University
654 Corporation for Atmospheric Research (UCAR) website at
655 https://www2.mmm.ucar.edu/wrf/users/download/get_source.html (UCAR, 2025a). The FNL data are
656 available at <https://rda.ucar.edu/datasets/ds083.2/> (NCEP, NWS, NOAA, U.S. DOC, 2000). The biomass
657 burning emission data of FINN version 1.5 can be obtained at <https://www.aom.ucar.edu/Data/fire/>
658 (UCAR, 2025b). The MEIC and MIX anthropogenic emissions are available at <http://meicmodel.org.cn/>
659 (Tsinghua University, CEADs, CAEP, 2025). The ERA5 data are available at
660 <https://cds.climate.copernicus.eu/datasets> (last access: 2027 SeptemberApril 2025),
661 <https://doi.org/10.24381/cds.bd0915c6> (Hersbach et al., 2023a), and
662 <https://doi.org/10.24381/cds.adbb2d47> (Hersbach et al., 2023b). MERRA-2 ~~radiation reanalysis~~ data
663 (<https://doi.org/10.5067/Q9QMY5PBNV1T>, GMAO, 2015a), and aerosol data
664 (<https://doi.org/10.5067/KLICLTZ8EM9D>, GMAO, 2015b), and meteorological data
665 (<https://doi.org/10.5067/QBZ6MG944HW0>, GMAO, 2015c) are available at
666 <https://disc.gsfc.nasa.gov/datasets> (last access: 2027 SeptemberApril 2025). The MODIS data is available
667 at https://doi.org/10.5067/MODIS/MOD08_D3.061 (Platnick et al., 2015). The CERES data are from

668 https://doi.org/10.5067/Terra-Aqua-NOAA20/CERES/SYN1degDay_L3.004B (NASA, 2017). The CPC
669 precipitation and OLR data are openly available from NOAA at
670 <https://psl.noaa.gov/data/gridded/data.cpc.globalprecip.html> (NOAA, 2025a) and
671 <https://psl.noaa.gov/data/gridded/data.olrcdr.interp.html> (NOAA, 2025b), respectively.

672

673 **Author contributions.** PCH and HC conceptualized the research goals and aims. HC performed the
674 analysis and wrote the manuscript draft. AZ and HC ran the simulations. PCH, AZ, and XM reviewed
675 and edited the manuscript.

676

677 **Competing interests.** The contact author has declared that none of the authors has any competing
678 interests.

679

680 **Acknowledgments.** We appreciate the anonymous reviewers for their constructive comments, which
681 greatly improved the manuscript. This work was supported by the National Natural Science Foundation
682 of China (42225502). We acknowledge the High Performance Computing Center of Nanjing University
683 of Information Science and Technology for their support of this study. We also thank all the
684 corresponding institutions for providing their data for this study.

685

686 **Financial support.** This research has been supported by the National Natural Science Foundation of
687 China (42225502).

688 References

689 [Abida, R., Addad, Y., Francis, D., Temimi, M., Nelli, N., Fonseca, R., Nesterov, O., and Bosc, E.: Evaluation of the](#)
690 [Performance of the WRF Model in a Hyper-Arid Environment: A Sensitivity Study, Atmosphere, 13, 985,](#)
691 <https://doi.org/10.3390/atmos13060985>, 2022.

692 Ackerman, A. S., Toon, O. B., Stevens, D. E., Heymsfield, A. J., Ramanathan, V., and Welton, E. J.: Reduction of
693 Tropical Cloudiness by Soot, Science, 288, 1042-1047, <https://doi.org/10.1126/science.288.5468.1042>, 2000.

694 Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, Science, 245, 1227-1230,
695 <https://doi.org/10.1126/science.245.4923.1227>, 1989.

696 Argüeso, D., Hidalgo-Muñoz, J. M., Gámiz-Fortis, S. R., Esteban-Parra, M. J., and Castro-Díez, Y.: High-resolution
697 projections of mean and extreme precipitation over Spain using the WRF model (2070–2099 versus 1970–1999), J.
698 Geophys. Res.-Atmos., 117, <https://doi.org/10.1029/2011JD017399>, 2012.

- Arya, V. B., Surendran, S., and Rajendran, K.: On the build-up of dust aerosols and possible indirect effect during Indian summer monsoon break spells using recent satellite observations of aerosols and cloud properties, *J. Earth Syst. Sci.*, 130, 42, <https://doi.org/10.1007/s12040-020-01526-6>, 2021.
- [Austin, R. T., Heymsfield, A. J., and Stephens, G. L.: Retrieval of ice cloud microphysical parameters using the CloudSat millimeter-wave radar and temperature, *J. Geophys. Res.-Atmos.*, 114, <https://doi.org/10.1029/2008JD010049>, 2009.](#)
- [Baklanov, A., Schlünzen, K., Suppan, P., Baldasano, J., Brunner, D., Aksoyoglu, S., Carmichael, G., Douros, J., Flemming, J., Forkel, R., Galmarini, S., Gauss, M., Grell, G., Hirtl, M., Joffre, S., Jorba, O., Kaas, E., Kaasik, M., Kallos, G., Kong, X., Korsholm, U., Kurganskiy, A., Kushta, J., Lohmann, U., Mahura, A., Manders-Groot, A., Maurizi, A., Moussiopoulos, N., Rao, S. T., Savage, N., Seigneur, C., Sokhi, R. S., Solazzo, E., Solomos, S., Sørensen, B., Tsegas, G., Vignati, E., Vogel, B., and Zhang, Y.: Online coupled regional meteorology chemistry models in Europe: current status and prospects, *Atmos. Chem. Phys.*, 14, 317-398, <https://doi.org/10.5194/acp-14-317-2014>, 2014.](#)
- Bhattacharya, A., Chakraborty, A., and Venugopal, V.: Role of aerosols in modulating cloud properties during active–break cycle of Indian summer monsoon, *Clim. Dynam.*, 49, 2131-2145, <https://doi.org/10.1007/s00382-016-3437-4>, 2017.
- Bollasina, M. A., Ming, Y., and Ramaswamy, V.: Anthropogenic Aerosols and the Weakening of the South Asian Summer Monsoon, *Science*, 334, 502-505, <https://doi.org/10.1126/science.1204994>, 2011.
- ~~[Buchard, V., Randles, C., Da Silva, A., Darmenov, A., Colarco, P., Govindaraju, R., Ferrare, R., Hair, J., Beyersdorf, A., and Ziemba, L.: The MERRA-2 aerosol reanalysis, 1980 onward. Part II: Evaluation and case studies, *J. Climate*, 30, 6851–6872, <https://doi.org/10.1175/jcli-d-16-0613.1>, 2017.](#)~~
- Caldwell, P., Chin, H.-N. S., Bader, D. C., and Bala, G.: Evaluation of a WRF dynamical downscaling simulation over California, *Climatic Change*, 95, 499-521, <https://doi.org/10.1007/s10584-009-9583-5>, 2009.
- Chen, F., Yang, Y., Yu, L., Li, Y., Liu, W., Liu, Y., and Lolli, S.: Distinct effects of fine and coarse aerosols on microphysical processes of shallow-precipitation systems in summer over southern China, *Atmos. Chem. Phys.*, 25, 1587-1601, <https://doi.org/10.5194/acp-25-1587-2025>, 2025.
- [Chen, G., Yang, J., Bao, Q., and Wang, W.-C.: Intraseasonal responses of the East Asia summer rainfall to anthropogenic aerosol climate forcing, *Clim. Dynam.*, 51, 3985-3998, <https://doi.org/10.1007/s00382-017-3691-0>, 2018.](#)
- Chen, H., Hsu, P.-C., and Hu, S.: Role of Quasi-Biweekly Cloud-Radiative Feedback in Modulating and Simulating Extreme Rainfall Intensity Over Asian Monsoon Regions, *Geophys. Res. Lett.*, 51, e2024GL111671, <https://doi.org/10.1029/2024GL111671>, 2024.
- [Chen, M., Shi, W., Xie, P., Silva, V. B. S., Kousky, V. E., Wayne Higgins, R., and Janowiak, J. E.: Assessing objective techniques for gauge-based analyses of global daily precipitation, *J. Geophys. Res.-Atmos.*, 113, <https://doi.org/10.1029/2007JD009132>, 2008.](#)
- ~~[Chen, G., Yang, J., Bao, Q., and Wang, W.-C.: Intraseasonal responses of the East Asia summer rainfall to anthropogenic aerosol climate forcing, *Clim. Dynam.*, 51, 3985–3998, <https://doi.org/10.1007/s00382-017-3691-0>, 2018.](#)~~

- Chen, Y., Yang, K., Zhou, D., Qin, J., and Guo, X.: Improving the Noah Land Surface Model in Arid Regions with an Appropriate Parameterization of the Thermal Roughness Length, *J. Hydrol.*, 11, 995-1006, <https://doi.org/10.1175/2010JHM1185.1>, 2010.
- ~~Chen, M., Shi, W., Xie, P., Silva, V. B. S., Kousky, V. E., Wayne Higgins, R., and Janowiak, J. E.: Assessing objective techniques for gauge-based analyses of global daily precipitation, *J. Geophys. Res. Atmos.*, 113, <https://doi.org/10.1029/2007JD009132>, 2008.~~
- ~~Choudhury, G., Tyagi, B., Vissa, N. K., Singh, J., Sarangi, C., Tripathi, S. N., and Tesche, M.: Aerosol-enhanced high precipitation events near the Himalayan foothills, *Atmos. Chem. Phys.*, 20, 15389-15399, <https://doi.org/10.5194/acp-20-15389-2020>, 2020.~~
- ~~Chyi, D., Wang, X., Yu, X., and Zhang, J.: Synoptic-Scale Analysis on Development and Maintenance of the 19–21 July 2021 Extreme Heavy Rainfall in Henan, Central China, *J. Meteorol. Res.*, 37, 174-191, <https://doi.org/10.1007/s13351-023-2914-z>, 2023.~~
- Dave, P., Bhushan, M., and Venkataraman, C.: Aerosols cause intraseasonal short-term suppression of Indian monsoon rainfall, *Sci. Rep.*, 7, 17347, <https://doi.org/10.1038/s41598-017-17599-1>, 2017.
- Debnath, S., Govardhan, G., Saha, S. K., Hazra, A., Pohkrel, S., Jena, C., Kumar, R., and Ghude, S. D.: Impact of dust aerosols on the Indian Summer Monsoon Rainfall on intra-seasonal time-scale, *Atmos. Environ.*, 305, 119802, <https://doi.org/10.1016/j.atmosenv.2023.119802>, 2023.
- Duchon, C. E.: Lanczos Filtering in One and Two Dimensions, *J. Appl. Meteorol. Clim.*, 18, 1016-1022, [https://doi.org/10.1175/1520-0450\(1979\)018<1016:LFIOAT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1979)018<1016:LFIOAT>2.0.CO;2), 1979.
- Fan, J., Zhang, R., Tao, W.-K., and Mohr, K. I.: Effects of aerosol optical properties on deep convective clouds and radiative forcing, *J. Geophys. Res.-Atmos.*, 113, <https://doi.org/10.1029/2007JD009257>, 2008.
- ~~Fan, J., Rosenfeld, D., Yang, Y., Zhao, C., Leung, L. R., and Li, Z.: Substantial contribution of anthropogenic air pollution to catastrophic floods in Southwest China, *Geophys. Res. Lett.*, 42, 6066-6075, <https://doi.org/10.1002/2015GL064479>, 2015.~~
- Fan, J., Rosenfeld, D., Zhang, Y., Giangrande, S. E., Li, Z., Machado, L. A. T., Martin, S. T., Yang, Y., Wang, J., Artaxo, P., Barbosa, H. M. J., Braga, R. C., Comstock, J. M., Feng, Z., Gao, W., Gomes, H. B., Mei, F., Pöhlker, C., Pöhlker, M. L., Pöschl, U., and de Souza, R. A. F.: Substantial convection and precipitation enhancements by ultrafine aerosol particles, *Science*, 359, 411-418, <https://doi.org/10.1126/science.aan8461>, 2018.
- ~~Fan, J., Zhang, Y., Li, Z., Yan, H., Prabhakaran, T., Rosenfeld, D., and Khain, A.: Unveiling Aerosol Impacts on Deep Convective Clouds: Scientific Concept, Modeling, Observational Analysis, and Future Direction, *J. Geophys. Res.-Atmos.*, 130, e2024JD041931, <https://doi.org/10.1029/2024JD041931>, 2025.~~
- Fast, J. D., Gustafson Jr., W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model, *J. Geophys. Res.-Atmos.*, 111, <https://doi.org/10.1029/2005JD006721>, 2006.

- Fu, Z., Hsu, P.-C., and Liu, F.: Factors Regulating the Multidecadal Changes in MJO Amplitude over the Twentieth Century, *J. Climate*, 33, 9513-9529, <https://doi.org/10.1175/JCLI-D-20-0111.1>, 2020.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419-5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Global Modeling and Assimilation Office (GMAO): MERRA-2 tavg1_2d_aer_Nx: 2d,1-Hourly, Time-averaged, Single-Level, Assimilation, Aerosol Diagnostics V5.12.4, Goddard Earth Sciences Data and Information Services Center (GES DISC) [data set], <https://doi.org/10.5067/KLICLTZ8EM9D>, 2015a.
- Global Modeling and Assimilation Office (GMAO): MERRA-2 tavg1_2d_rad_Nx: 2d,1-Hourly, Time-Averaged, Single-Level, Assimilation, Radiation Diagnostics V5.12.4, Goddard Earth Sciences Data and Information Services Center (GES DISC) [data set]. <https://doi.org/10.5067/Q9QMY5PBNV1T>, 2015b.
- Global Modeling and Assimilation Office (GMAO): MERRA-2 inst3_3d_asm_Np: 3d,3-Hourly, Instantaneous, Pressure-Level, Assimilation, Assimilated Meteorological Fields V5.12.4, Goddard Earth Sciences Data and Information Services Center (GES DISC) [data set]. <https://doi.org/10.5067/QBZ6MG944HW0>, 2015c.
- Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S.-J.: Sources and distributions of dust aerosols simulated with the GOCART model, *J. Geophys. Res.-Atmos.*, 106, 20255-20273, <https://doi.org/10.1029/2000JD000053>, 2001.
- Grell, G. A. and Freitas, S. R.: A scale and aerosol aware stochastic convective parameterization for weather and air quality modeling, *Atmos. Chem. Phys.*, 14, 5233-5250, <https://doi.org/10.5194/acp-14-5233-2014>, 2014.
- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmos. Environ.*, 39, 6957-6975, <https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471-1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- Guo, J., Deng, M., Lee, S. S., Wang, F., Li, Z., Zhai, P., Liu, H., Lv, W., Yao, W., and Li, X.: Delaying precipitation and lightning by air pollution over the Pearl River Delta. Part I: Observational analyses, *J. Geophys. Res.-Atmos.*, 121, 6472-6488, <https://doi.org/10.1002/2015JD023257>, 2016.
- Guo, J., Luo, Y., Yang, J., Furtado, K., and Lei, H.: Effects of anthropogenic and sea salt aerosols on a heavy rainfall event during the early-summer rainy season over coastal Southern China, *Atmos. Res.*, 265, 105923, <https://doi.org/10.1016/j.atmosres.2021.105923>, 2022.
- Guo, X., Fu, D., Guo, X., and Zhang, C.: A case study of aerosol impacts on summer convective clouds and precipitation over northern China, *Atmos. Res.*, 142, 142-157, <https://doi.org/10.1016/j.atmosres.2013.10.006>, 2014.

807 Hazra, A., Goswami, B. N., and Chen, J.-P.: Role of Interactions between Aerosol Radiative Effect, Dynamics, and
808 Cloud Microphysics on Transitions of Monsoon Intraseasonal Oscillations, *J. Atmos. Sci.*, 70, 2073-2087,
809 <https://doi.org/10.1175/JAS-D-12-0179.1>, 2013.

810 [He, J., Glotfelty, T., Yahya, K., Alapaty, K., and Yu, S.: Does temperature nudging overwhelm aerosol radiative effects](#)
811 [in regional integrated climate models?, *Atmos. Environ.*, 154, 42-52,](#)
812 <https://doi.org/10.1016/j.atmosenv.2017.01.040>, 2017.

813 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R.,
814 Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J.,
815 Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes,
816 M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P.,
817 Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global
818 reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999-2049, <https://doi.org/10.1002/qj.3803>, 2020.

819 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R.,
820 Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J. N.: ERA5 hourly data on pressure levels
821 from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set],
822 <https://doi.org/10.24381/cds.bd0915c6>, 2023a.

823 [Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R.,](#)
824 [Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J.-N.: ERA5 hourly data on single levels from](#)
825 [1940 to present. Copernicus Climate Change Service \(C3S\) Climate Data Store \(CDS\) \[data set\],](#)
826 <https://doi.org/10.24381/cds.adbb2d47>, 2023b.

827 Hong, S.-Y., Noh, Y., and Dudhia, J.: A New Vertical Diffusion Package with an Explicit Treatment of Entrainment
828 Processes, *Mon. Weather Rev.*, 134, 2318-2341, <https://doi.org/10.1175/MWR3199.1>, 2006.

829 Hsu, P.-C., Lee, J.-Y., and Ha, K.-J.: Influence of boreal summer intraseasonal oscillation on rainfall extremes in
830 southern China, *Int. J. Climatol*, 36, 1403-1412, <https://doi.org/10.1002/joc.4433>, 2016.

831 [Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M.,](#)
832 [Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A.,](#)
833 [Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad,](#)
834 [I. E., and Prévôt, A. S. H.: High secondary aerosol contribution to particulate pollution during haze events in China,](#)
835 [Nature, 514, 218-222, https://doi.org/10.1038/nature13774](#), 2014.

836 Huang, X. and Ding, A.: Aerosol as a critical factor causing forecast biases of air temperature in global numerical weather
837 prediction models, *Sci. Bull.*, 66, 1917-1924, <https://doi.org/10.1016/j.scib.2021.05.009>, 2021.

838 ~~[Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M.,](#)~~
839 ~~[Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A.,](#)~~
840 ~~[Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad,](#)~~
841 ~~[I. E., and Prévôt, A. S. H.: High secondary aerosol contribution to particulate pollution during haze events in China,](#)~~
842 ~~[Nature, 514, 218-222, https://doi.org/10.1038/nature13774](#), 2014.~~

Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *J. Geophys. Res.-Atmos.*, 113, <https://doi.org/10.1029/2008JD009944>, 2008.

Jia, H., Hasekamp, O., and Quaas, J.: Revisiting Aerosol–Cloud Interactions From Weekly Cycles, *Geophys. Res. Lett.*, 51, e2024GL108266, <https://doi.org/10.1029/2024GL108266>, 2024.

Koren, I., Kaufman, Y. J., Remer, L. A., and Martins, J. V.: Measurement of the Effect of Amazon Smoke on Inhibition of Cloud Formation, *Science*, 303, 1342-1345, <https://doi.org/10.1126/science.1089424>, 2004.

Lee, K. H., Li, Z., Wong, M. S., Xin, J., Wang, Y., Hao, W.-M., and Zhao, F.: Aerosol single scattering albedo estimated across China from a combination of ground and satellite measurements, *J. Geophys. Res.-Atmos.*, 112, <https://doi.org/10.1029/2007JD009077>, 2007.

[Lee, S. S., Guo, J., and Li, Z.: Delaying precipitation by air pollution over the Pearl River Delta: 2. Model simulations, J. Geophys. Res.-Atmos., 121, 11,739-711,760, https://doi.org/10.1002/2015JD024362, 2016.](#)

Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, *Atmos. Meas. Tech.*, 6, 2989-3034, <https://doi.org/10.5194/amt-6-2989-2013>, 2013.~~[Li, Z., Niu, F., Fan, J., Liu, Y., Rosenfeld, D., and Ding, Y.: Long-term impacts of aerosols on the vertical development of clouds and precipitation, Nat. Geosci., 4, 888-894, https://doi.org/10.1038/ngeo1313, 2011.](#)~~

~~[Li, Z., Lau, W. K. M., Ramanathan, V., Wu, G., Ding, Y., Manoj, M. G., Liu, J., Qian, Y., Li, J., Zhou, T., Fan, J., Rosenfeld, D., Ming, Y., Wang, Y., Huang, J., Wang, B., Xu, X., Lee, S. S., Cribb, M., Zhang, F., Yang, X., Zhao, C., Takemura, T., Wang, K., Xia, X., Yin, Y., Zhang, H., Guo, J., Zhai, P. M., Sugimoto, N., Babu, S. S., and Brasseur, G. P.: Aerosol and monsoon climate interactions over Asia, Rev. Geophys., 54, 866-929, https://doi.org/10.1002/2015RG000500, 2016.](#)~~

Li, M., Liu, H., Geng, G., Hong, C., Liu, F., Song, Y., Tong, D., Zheng, B., Cui, H., Man, H., Zhang, Q., and He, K.: Anthropogenic emission inventories in China: a review, *Natl. Sci. Rev.*, 4, 834-866, <https://doi.org/10.1093/nsr/nwx150>, 2017a.

Li, M., Zhang, Q., Kurokawa, J. I., Woo, J. H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935-963, <https://doi.org/10.5194/acp-17-935-2017>, 2017b.

[Li, Z., Niu, F., Fan, J., Liu, Y., Rosenfeld, D., and Ding, Y.: Long-term impacts of aerosols on the vertical development of clouds and precipitation, Nat. Geosci., 4, 888-894, https://doi.org/10.1038/ngeo1313, 2011.](#)

[Li, Z., Lau, W. K.-M., Ramanathan, V., Wu, G., Ding, Y., Manoj, M. G., Liu, J., Qian, Y., Li, J., Zhou, T., Fan, J., Rosenfeld, D., Ming, Y., Wang, Y., Huang, J., Wang, B., Xu, X., Lee, S.-S., Cribb, M., Zhang, F., Yang, X., Zhao, C., Takemura, T., Wang, K., Xia, X., Yin, Y., Zhang, H., Guo, J., Zhai, P. M., Sugimoto, N., Babu, S. S., and Brasseur, G. P.: Aerosol and monsoon climate interactions over Asia, Rev. Geophys., 54, 866-929, https://doi.org/10.1002/2015RG000500, 2016.](#)

878 [Li, Z., Wang, Y., Guo, J., Zhao, C., Cribb, M. C., Dong, X., Fan, J., Gong, D., Huang, J., Jiang, M., Jiang, Y., Lee, S.-](#)
879 [S., Li, H., Li, J., Liu, J., Qian, Y., Rosenfeld, D., Shan, S., Sun, Y., Wang, H., Xin, J., Yan, X., Yang, X., Yang, X.-](#)
880 [q., Zhang, F., and Zheng, Y.: East Asian Study of Tropospheric Aerosols and their Impact on Regional Clouds,](#)
881 [Precipitation, and Climate \(EAST-AIRCPC\), J. Geophys. Res.-Atmos., 124, 13026-13054,](#)
882 <https://doi.org/10.1029/2019JD030758>, 2019.

883 Liebmann, B. and Smith, C. A.: Description of a complete (interpolated) outgoing longwave radiation dataset, Bull. Am.
884 Meteorol. Soc., 77, 1275-1277, 1996.

885 Lin, Z. and Chen, G.: The Role of Anthropogenic Forcings on the Regional Climate of Summertime Diurnal Variations
886 over North China, J. Climate, 36, 4491-4509, <https://doi.org/10.1175/JCLI-D-22-0498.1>, 2023.

887 Liu, Z., Ming, Y., Zhao, C., Lau, N. C., Guo, J., Bollasina, M., and Yim, S. H. L.: Contribution of local and remote
888 anthropogenic aerosols to a record-breaking torrential rainfall event in Guangdong Province, China, Atmos. Chem.
889 Phys., 20, 223-241, <https://doi.org/10.5194/acp-20-223-2020>, 2020.

890 Manoj, M. G., Devara, P. C. S., Safai, P. D., and Goswami, B. N.: Absorbing aerosols facilitate transition of Indian
891 monsoon breaks to active spells, Clim. Dynam., 37, 2181-2198, <https://doi.org/10.1007/s00382-010-0971-3>, 2011.

892 Morrison, H., Thompson, G., and Tatarskii, V.: Impact of Cloud Microphysics on the Development of Trailing Stratiform
893 Precipitation in a Simulated Squall Line: Comparison of One- and Two-Moment Schemes, Mon. Weather Rev., 137,
894 991-1007, <https://doi.org/10.1175/2008MWR2556.1>, 2009.

895 Mudelsee, M.: Bootstrap confidence intervals, in: Climate Time Series Analysis, Atmospheric and Oceanographic
896 Sciences Library (Vol. 51), Springer, Cham, Germany, 61-104, <https://doi.org/10.1007/978-3-319-04450-7>, 2014.

897 [National Aeronautics and Space Administration \(NASA\). CERES and GEO-Enhanced TOA, Within-Atmosphere and](#)
898 [Surface Fluxes, Clouds and Aerosols Daily Terra-Aqua-NOAA20 Edition4B. NASA Langley Atmospheric Science](#)
899 [Data Center DAAC \[data set\]. \[https://doi.org/10.5067/Terra-Aqua-NOAA20/CERES/SYN1degDay_L3.004B\]\(https://doi.org/10.5067/Terra-Aqua-NOAA20/CERES/SYN1degDay_L3.004B\),](#)
900 [2017.](#)

901 National Centers for Environmental Prediction (NCEP), National Weather Service (NWS), NOAA, U.S. Department of
902 Commerce (DOC): NCEP FNL Operational Model Global Tropospheric Analyses, continuing from July 1999,
903 Research Data Archive at the NCEP, Computational and Information Systems Laboratory [data set],
904 <https://doi.org/10.5065/D6M043C6>, 2000.

905 National Oceanic and Atmospheric Administration (NOAA): CPC Global Unified Gauge-Based Analysis of Daily
906 Precipitation data [data set], <https://psl.noaa.gov/data/gridded/data.cpc.globalprecip.html>, last access: 27 April
907 2025a.

908 National Oceanic and Atmospheric Administration (NOAA): NOAA Interpolated Outgoing Longwave Radiation (OLR)
909 data [data set], <https://psl.noaa.gov/data/gridded/data.olrcdr.interp.html>, last access: 27 April 2025b.

910 Pahlow, M., Parlange, M. B., and Porté-Agel, F.: On Monin–Obukhov Similarity In The Stable Atmospheric Boundary
911 Layer, Bound.-Lay. Meteorol., 99, 225-248, <https://doi.org/10.1023/A:1018909000098>, 2001.

912 Platnick, S., Hubanks, P., Meyer, K., and King, M. D.: MODIS Atmosphere L3 Daily Product, NASA MODIS Adaptive
 913 Processing System, Goddard Space Flight Center, ~~U.S.A.~~ [data set],
 914 https://doi.org/10.5067/MODIS/MOD08_MD3.061, 2015.

915 Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks,
 916 P. A., Holz, R. E., Yang, P., Ridgway, W. L., and Riedi, J.: The MODIS Cloud Optical and Microphysical Products:
 917 Collection 6 Updates and Examples from Terra and Aqua, IEEE T. Geosci. Remote, 55, 502-525,
 918 <https://doi.org/10.1109/TGRS.2016.2610522>, 2017.

919 Reid, J. S., Lagrosas, N. D., Jonsson, H. H., Reid, E. A., Sessions, W. R., Simpas, J. B., Uy, S. N., Boyd, T. J., Atwood,
 920 S. A., Blake, D. R., Campbell, J. R., Cliff, S. S., Holben, B. N., Holz, R. E., Hyer, E. J., Lynch, P., Meinardi, S.,
 921 Posselt, D. J., Richardson, K. A., Salinas, S. V., Smirnov, A., Wang, Q., Yu, L., and Zhang, J.: Observations of the
 922 temporal variability in aerosol properties and their relationships to meteorology in the summer monsoonal South
 923 China Sea/East Sea: the scale-dependent role of monsoonal flows, the Madden–Julian Oscillation, tropical cyclones,
 924 squall lines and cold pools, Atmos. Chem. Phys., 15, 1745-1768, <https://doi.org/10.5194/acp-15-1745-2015>, 2015.

925 Rosenfeld, D., Lohmann, U., Raga, G. B., O'Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A., and Andreae, M. O.:
 926 Flood or Drought: How Do Aerosols Affect Precipitation?, Science, 321, 1309-1313,
 927 <https://doi.org/10.1126/science.1160606>, 2008.

928 [Rutan, D. A., Kato, S., Doelling, D. R., Rose, F. G., Nguyen, L. T., Caldwell, T. E., and Loeb, N. G.: CERES Synoptic](https://doi.org/10.1175/JTECH-D-14-00165.1)
 929 [Product: Methodology and Validation of Surface Radiant Flux, J. Atmos. Oceanic Technol., 32, 1121-1143,](https://doi.org/10.1175/JTECH-D-14-00165.1)
 930 <https://doi.org/10.1175/JTECH-D-14-00165.1>, 2015.

931 Shao, T., Liu, Y., Wang, R., Zhu, Q., Tan, Z., and Luo, R.: Role of anthropogenic aerosols in affecting different-grade
 932 precipitation over eastern China: A case study, Sci. Total Environ., 807, 150886,
 933 <https://doi.org/10.1016/j.scitotenv.2021.150886>, 2022.

934 ~~Shi, H., Xiao, Z., Zhan, X., Ma, H., and Tian, X.: Evaluation of MODIS and two reanalysis aerosol optical depth products~~
 935 ~~over AERONET sites, Atmos. Res., 220, 75–80, <https://doi.org/10.1016/j.atmosres.2019.01.009>, 2019.~~

936 Singh, C., Ganguly, D., and Sharma, P.: Impact of West Asia, Tibetan Plateau and local dust emissions on intra-seasonal
 937 oscillations of the South Asian monsoon rainfall, Clim. Dynam., 53, 6569-6593, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-019-04944-5)
 938 [019-04944-5](https://doi.org/10.1007/s00382-019-04944-5), 2019.

939 Stier, P., van den Heever, S. C., Christensen, M. W., Gryspeerdt, E., Dagan, G., Saleeby, S. M., Bollasina, M., Donner,
 940 L., Emanuel, K., Ekman, A. M. L., Feingold, G., Field, P., Forster, P., Haywood, J., Kahn, R., Koren, I., Kummerow,
 941 C., L'Ecuyer, T., Lohmann, U., Ming, Y., Myhre, G., Quaas, J., Rosenfeld, D., Samset, B., Seifert, A., Stephens, G.,
 942 and Tao, W.-K.: Multifaceted aerosol effects on precipitation, Nat. Geosci., 17, 719-732,
 943 <https://doi.org/10.1038/s41561-024-01482-6>, 2024.

944 [Su, Y., Zhao, C., Wang, Y., and Ma, Z.: Spatiotemporal Variations of Precipitation in China Using Surface Gauge](https://doi.org/10.3390/atmos11030303)
 945 [Observations from 1961 to 2016, Atmosphere, 11, 303, <https://doi.org/10.3390/atmos11030303>, 2020.](https://doi.org/10.3390/atmos11030303)

946 Sun, Y. and Zhao, C.: Distinct impacts on precipitation by aerosol radiative effect over three different megacity regions
 947 of eastern China, Atmos. Chem. Phys., 21, 16555-16574, <https://doi.org/10.5194/acp-21-16555-2021>, 2021.

948 Surendran, S., Ajay Anand, K. V., Ravindran, S., and Rajendran, K.: Exacerbation of Indian Summer Monsoon Breaks
 949 by the Indirect Effect of Regional Dust Aerosols, *Geophys. Res. Lett.*, 49, e2022GL101106,
 950 <https://doi.org/10.1029/2022GL101106>, 2022.

951 Tao, W.-K., Chen, J.-P., Li, Z., Wang, C., and Zhang, C.: Impact of aerosols on convective clouds and precipitation,
 952 *Rev. Geophys.*, 50, <https://doi.org/10.1029/2011RG000369>, 2012.

953 Tian, B., Waliser, D. E., Kahn, R. A., and Wong, S.: Modulation of Atlantic aerosols by the Madden-Julian Oscillation,
 954 *J. Geophys. Res.-Atmos.*, 116, <https://doi.org/10.1029/2010JD015201>, 2011.

955 Tsinghua University, China Carbon Emission Accounts and Datasets (CEADs) team, Chinese Academy of
 956 Environmental Planning of the Ministry of Ecology and Environment (CAEP): Multi-resolution Emission Inventory
 957 for China and MIX-Asia dataset, MEIC Data Platform [data set], <http://meicmodel.org.cn/>, last access: 27 April
 958 2025.

959 Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *J. Atmos. Sci.*, 34, 1149-1152,
 960 [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.

961 University Corporation for Atmospheric Research (UCAR): WRF Source Codes and Graphics Software Downloads,
 962 UCAR [code], https://www2.mmm.ucar.edu/wrf/users/download/get_source.html, last access: 27 April 2025a.

963 University Corporation for Atmospheric Research (UCAR): Fire Emission Factors and Emission Inventories, UCAR
 964 [data set], <https://www.aom.ucar.edu/Data/fire/>, last access: 27 April 2025b.

965 Vinoj, V., Rasch, P. J., Wang, H., Yoon, J.-H., Ma, P.-L., Landu, K., and Singh, B.: Short-term modulation of Indian
 966 summer monsoon rainfall by West Asian dust, *Nat. Geosci.*, 7, 308-313, <https://doi.org/10.1038/ngeo2107>, 2014.

967 [Wang, L., Zhang, Y., Wang, K., Zheng, B., Zhang, Q., and Wei, W.: Application of Weather Research and Forecasting](#)
 968 [Model with Chemistry \(WRF/Chem\) over northern China: Sensitivity study, comparative evaluation, and policy](#)
 969 [implications, *Atmos. Environ.*, 124, 337-350, <https://doi.org/10.1016/j.atmosenv.2014.12.052>, 2016.](#)

970 Wang, Y., Wan, Q., Meng, W., Liao, F., Tan, H., and Zhang, R.: Long-term impacts of aerosols on precipitation and
 971 lightning over the Pearl River Delta megacity area in China, *Atmos. Chem. Phys.*, 11, 12421-12436,
 972 <https://doi.org/10.5194/acp-11-12421-2011>, 2011.

973 Wang, Z., Xue, L., Liu, J., Ding, K., Lou, S., Ding, A., Wang, J., and Huang, X.: Roles of Atmospheric Aerosols in
 974 Extreme Meteorological Events: a Systematic Review, *Curr. Pollut. Rep.*, 8, 177-188,
 975 <https://doi.org/10.1007/s40726-022-00216-9>, 2022.

976 [Wei, J., Mao, Q., Shan, Y., Jin, Q., Yang, Y., and Chen, H.: Spring biomass burning in Indochina enhances summer](#)
 977 [Yangtze River Valley rainfall through land-atmosphere interactions, *npj Clim. Atmos. Sci.*, 6, 183,](#)
 978 [https://doi.org/10.1038/s41612-023-00514-z, 2023.](#)

979 Wei, S., Hsu, P.-C., and Xie, J.: Effects of Intraseasonal Oscillation on Timing and Subseasonal Predictability of Mei-
 980 yu Onset over the Yangtze River Basin, *J. Climate*, 37, 2277-2295, <https://doi.org/10.1175/JCLI-D-23-0504.1>, 2024.

981 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire
 982 INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci.*
 983 *Model Dev.*, 4, 625-641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.

984 [Xiao, H., Liu, X., Li, H., Yue, Q., Feng, L., and Qu, J.: Extent of aerosol effect on the precipitation of squall lines: A](#)
985 [case study in South China, Atmos. Res., 292, 106886, <https://doi.org/10.1016/j.atmosres.2023.106886>, 2023a.](#)

986 Xiao, Z., Yu, Y., Miao, Y., Zhu, S., He, H., Wang, Y., and Che, H.: Impact of Aerosols on Convective System Over the
987 North China Plain: A Numerical Case Study in Autumn, J. Geophys. Res.-Atmos., 128, e2022JD037465,
988 <https://doi.org/10.1029/2022JD037465>, 2023b.

989 Xie, J., Hsu, P.-C., Lee, J.-Y., Wang, L., and Turner, A. G.: Tropical intraseasonal oscillations as key driver and source
990 of predictability for the 2022 Pakistan record-breaking rainfall event, npj Clim. Atmos. Sci., 7, 256,
991 <https://doi.org/10.1038/s41612-024-00809-9>, 2024.

992 Yanai, M., Esbensen, S., and Chu, J.-H.: Determination of Bulk Properties of Tropical Cloud Clusters from Large-Scale
993 Heat and Moisture Budgets, J. Atmos. Sci., 30, 611-627, <https://doi.org/10.1175/1520->
994 0469(1973)030<0611:DOBPOT>2.0.CO;2, 1973.

995 [Yang, X. and Li, Z.: Increases in thunderstorm activity and relationships with air pollution in southeast China, J. Geophys.](#)
996 [Res.-Atmos., 119, 1835-1844, <https://doi.org/10.1002/2013JD021224>, 2014.](#)

997 Yang, X., Zhou, L., Zhao, C., and Yang, J.: Impact of aerosols on tropical cyclone-induced precipitation over the
998 mainland of China, Climatic Change, 148, 173-185, <https://doi.org/10.1007/s10584-018-2175-5>, 2018.

999 Yang, X., Li, Z., Liu, L., Zhou, L., Cribb, M., and Zhang, F.: Distinct weekly cycles of thunderstorms and a potential
1000 connection with aerosol type in China, Geophys. Res. Lett., 43, 8760-8768, <https://doi.org/10.1002/2016GL070375>,
1001 2016.

1002 Yu, Y. and Ginoux, P.: Assessing the contribution of the ENSO and MJO to Australian dust activity based on satellite-
1003 and ground-based observations, Atmos. Chem. Phys., 21, 8511-8530, <https://doi.org/10.5194/acp-21-8511-2021>,
1004 2021.

1005 Yun, Y., Zhang, D.-L., Gao, W., Yin, J., Zhao, C., Li, J., Guo, J., and Liu, H.: Spatiotemporal Variations of the Effects
1006 of Aerosols on Clouds and Precipitation in an Extreme-Rain-Producing MCS in South China, J. Geophys. Res.-
1007 Atmos., 129, e2023JD040014, <https://doi.org/10.1029/2023JD040014>, 2024.

1008 Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, J.
1009 Geophys. Res.-Atmos., 104, 30387-30415, <https://doi.org/10.1029/1999JD900876>, 1999.

1010 Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry
1011 (MOSAIC), J. Geophys. Res.-Atmos., 113, <https://doi.org/10.1029/2007JD008782>, 2008.

1012 [Zhang, L., Fu, T.-M., Tian, H., Ma, Y., Chen, J.-P., Tsai, T.-C., Tsai, I.-C., Meng, Z., and Yang, X.: Anthropogenic](#)
1013 [Aerosols Significantly Reduce Mesoscale Convective System Occurrences and Precipitation Over Southern China](#)
1014 [in April, Geophys. Res. Lett., 47, e2019GL086204, <https://doi.org/10.1029/2019GL086204>, 2020.](#)

1015 Zhang, Y., Gao, Y., Guo, L., and Zhang, M.: Numerical analysis of aerosol direct and indirect effects on an extreme
1016 rainfall event over Beijing in July 2016, Atmos. Res., 264, 105871, <https://doi.org/10.1016/j.atmosres.2021.105871>,
1017 2021.

1018 Zhang, Y., Wen, X. Y., and Jang, C. J.: Simulating chemistry–aerosol–cloud–radiation–climate feedbacks over the
 1019 continental U.S. using the online-coupled Weather Research Forecasting Model with chemistry (WRF/Chem),
 1020 Atmos. Environ., 44, 3568–3582, <https://doi.org/10.1016/j.atmosenv.2010.05.056>, 2010.

1021 [Zhao, C., Yang, Y., Chi, Y., Sun, Y., Zhao, X., Letu, H., and Xia, Y.: Recent progress in cloud physics and associated
 1022 radiative effects in China from 2016 to 2022, Atmos. Res., 293, 106899,
 1023 <https://doi.org/10.1016/j.atmosres.2023.106899>, 2023.](#)

1024 Zhao, C., Sun, Y., Yang, J., Li, J., Zhou, Y., Yang, Y., Fan, H., and Zhao, X.: Observational evidence and mechanisms
 1025 of aerosol effects on precipitation, Sci. Bull., 69, 1569–1580, <https://doi.org/10.1016/j.scib.2024.03.014>, 2024.

1026 [Zhao, J., Ma, X., Wu, S., and Sha, T.: Dust emission and transport in Northwest China: WRF-Chem simulation and
 1027 comparisons with multi-sensor observations, Atmos. Res., 241, 104978,
 1028 <https://doi.org/10.1016/j.atmosres.2020.104978>, 2020.](#)

1029 Zhou, S., Yang, J., Wang, W. C., Zhao, C., Gong, D., and Shi, P.: An observational study of the effects of aerosols on
 1030 diurnal variation of heavy rainfall and associated clouds over Beijing–Tianjin–Hebei, Atmos. Chem. Phys., 20,
 1031 5211–5229, <https://doi.org/10.5194/acp-20-5211-2020>, 2020.

1032 Zhu, A., Xu, H., Deng, J., Ma, J., and Hua, S.: Instant and delayed effects of March biomass burning aerosols over the
 1033 Indochina Peninsula, Atmos. Chem. Phys., 22, 15425–15447, <https://doi.org/10.5194/acp-22-15425-2022>, 2022.

1034 [Zhu, H., Li, R., Yang, S., Zhao, C., Jiang, Z., and Huang, C.: The impacts of dust aerosol and convective available
 1035 potential energy on precipitation vertical structure in southeastern China as seen from multisource observations,
 1036 Atmos. Chem. Phys., 23, 2421–2437, <https://doi.org/10.5194/acp-23-2421-2023>, 2023.](#)

1037 [Zhu, H., Yang, S., Zhao, H., Wang, Y., and Li, R.: Complex interplay of sulfate aerosols and meteorology conditions on
 1038 precipitation and latent heat vertical structure, npj Clim. Atmos. Sci., 7, 191, \[https://doi.org/10.1038/s41612-024-
 00743-w\]\(https://doi.org/10.1038/s41612-024-

 1039 00743-w\), 2024.](#)