

Impacts of shipping emissions on ozone pollution in China

Zhenyu Luo^{#,1}, Li Peng^{#,1}, Zhaofeng Lv¹, Junchao Zhao², Tingkun He¹, Wen Yi¹, Yongyue Wang¹, Kebin He¹, Huan Liu^{*,1}

¹State Key Laboratory of Regional Environment and Sustainability, School of Environment, Tsinghua University, Beijing 100084, China

²Key Laboratory of Vehicle Emission Control and Simulation of Ministry of Ecology and Environment, Vehicle Emission Control Center, Chinese Research Academy of environmental Sciences, Beijing, China

[#] These authors contributed equally to this work.

* Correspondence: Huan Liu (liu_env@tsinghua.edu.cn)

Abstract

With the Two Phases of Clean Air Actions in China, the shipping sector has emerged as a significant source with substantial emission reduction potential compared to land-based anthropogenic sectors. Therefore, understanding the contribution of shipping emissions to ozone (O₃) pollution is therefore essential for advancing China's air pollution control efforts. In this study, a coupled framework including a chemical transport model with machine learning techniques was developed to systematically investigate the interannual and seasonal impacts of shipping emissions on O₃ concentrations across China during the period from 2016 to 2020, and explore mechanisms ~~by which~~ shipping emissions influence O₃ formation. Results indicate that shipping emissions increases O₃ concentrations by a five-year average of 3.5 ppb nationwide, exhibiting significant spatial and temporal heterogeneity across different regions and seasons. Although significant differences exist between the emissions of ocean vessels and inland vessels, their contributions to O₃ formation are becoming increasingly comparable. Solely controlling shipping emissions may ~~has limited impact on O₃ mitigation~~~~not necessarily result in effective O₃ mitigation~~. Instead, coordinated reductions targeting both shipping and land-based anthropogenic sources, along with region-specific and targeted emission control strategies, are critical for achieving substantial improvements in O₃ pollution mitigation.

1 Introduction

Over the past decades, China's rapid industrialization and urbanization have boost the economy but also exacerbated air pollution (Zhang et al., 2019). To address air pollution and the health burden from anthropogenic emissions, China introduced the Air Pollution Prevention and Control Action Plan in 2013 (Zheng et al., 2018). Although China has implemented synergistic control of VOC and NO_x emissions, the warm-season mean maximum daily 8 h average ozone (MDA8 O₃) increased by 2.6 µg m⁻³ yr⁻¹ in China between 2013 and 2020. Although a series of stringent measures have greatly improved PM_{2.5} pollution in China, ozone (O₃) pollution has become increasingly prominent—especially in urban areas where declining PM_{2.5} levels offset gains in O₃ mitigation (Liu et al., 2023). Numerous epidemiological studies have shown that ground-level O₃ pollution leads to a range of adverse health effects, including increased incidence and mortality of respiratory diseases (Ito et al., 2005; Jerrett et al., 2009; Tao et al., 2012). Therefore, the health benefits achieved by reducing PM_{2.5} pollution are partially offset by the increase in O₃ pollution (Wang et al., 2021a; Xie et al., 2019), indicating a need to explore strategies to mitigate O₃ pollution in China.

~~With~~ During the promotion of China's emission control actions, emissions from the industry and power sectors declined substantially, with NO_x reductions exceeding 50%, while the transportation sector still retains significant potential for further cuts~~the potential for further reductions from land anthropogenic emissions has diminished, while the shipping sector has shown considerable potential~~ (Liu et al., 2023). Ships emit both gaseous and particulate pollutants, including sulfur dioxide (SO₂), nitrogen oxides (NO_x), particulate matter, and volatile organic compounds (VOC). As the largest maritime trading nation, China has a higher share of shipping emissions among anthropogenic sources compared to international levels (Fu et al., 2017; Yi et al., 2025). ~~Currently~~ From 2016 to 2019, shipping emission controls in China focused on reducing SO₂ and PM emissions through the adoption of low-sulfur fuels, while NO_x and VOC emissions from shipping continued to rise by approximately 13% due to increasing trade volumes, shipping emission controls in China focus on reducing SO₂ and PM emissions through the adoption of low-sulfur fuels (Wang et al., 2021). With increasing trade volumes, however, shipping NO_x and VOC emissions continue to rise— (Wang et al., 2021b). Additionally, ~~and~~ the use of low-sulfur fuel may further increase VOC emissions (Wu et al., 2020). Therefore, clarifying the historical and current contribution of shipping emissions to the formation of O₃ is critically important for further pollution control in China.

Previous studies have quantified the impacts of shipping emissions on O₃ pollution in China. In the southern coastal region, shipping emissions contributed approximately 0.9 µg/m³ to annual O₃ pollution (Cheng et al., 2023), with a peak winter contribution of up to 10% (Feng et al., 2023). In the eastern coastal region during summer, the shipping-related O₃ concentration ranged from -15 to 15 ppb (Fu et al., 2023; Wang et al., 2019a). In the Bohai Rim Area (BRA), shipping emissions showed a maximum annual negative contribution of 0.5 µg/m³ (Wan et al., 2023), while summer O₃ concentration in Shandong Province increased by up to 10 ppb due to shipping (Wang et al., 2022), particularly in the Yangtze River Delta (YRD), Pearl River Delta (PRD), and Bohai Rim Area (BRA) (Wang et al., 2019; Fu et al., 2023; Zheng et al., 2024). However, these studies predominantly focused on coastal areas, such as Yangtze River Delta

(YRD), Pearl River Delta (PRD), and BRA, while the potential inland impacts are not well studied. However, recent Research has studies have indicated that the air quality effects of inland shipping should not be neglected (Huang et al., 2022; Luo et al., 2024). The formation of O₃ exhibits strong spatial heterogeneity and is influenced by multiple factors, such as meteorological conditions and the emission intensities from both shipping and land-based sources. Currently, there is a lack of comprehensive studies employing a unified methodology to assess the nationwide impact of shipping emissions on ~~ozone~~O₃. Furthermore, previous studies were limited to restricted ~~timeframes~~periods, resulting in a deficiency in comprehensive assessments of multi-year scenarios that encompass coordinated variations in both land and shipping emissions.

O₃ is generated by photochemical reactions between NO_x and VOC under solar radiation, thus, the impacts of shipping emissions on O₃ concentrations are attributed by the nonlinear response of O₃ to changes in NO_x and VOC emissions (Wang et al., 2019a, 2017). Additionally, the titration of O₃ by NO from shipping emissions, particularly within a few kilometers of ship tracks, can further complicate the simulation and interpretation of O₃ concentrations at the local scale (Merico et al., 2016). However, previous studies commonly used the zero-out method to assess ship's impacts by comparing scenario differences simulated by chemical transport models, which does not fully involve the nonlinear response of O₃ to its precursors and would result in considerable uncertainty in the evaluations (Cheng et al., 2023; Feng et al., 2023; Fu et al., 2023; Wan et al., 2023; Wang et al., 2022, 2019a). Furthermore, although model-based assessments can generate large amounts of simulation data to investigate the impacts of shipping emissions, the number of scenarios that can be simulated by chemical transport models remains limited due to computational constraints. As a result, current analyses struggle to struggles the mechanism of how shipping emissions contribute to ~~O₃ozone~~formation from these discrete scenarios. Recently, the advancement of machine learning techniques, with strong capabilities in capturing nonlinear relationships, provides a valuable approach for uncovering underlying patterns in such datasets from these simulation big data (Luo et al., 2025).

In this study, we conducted source-oriented chemical transport model with a spatial resolution of 36 km×36 km, to investigate the annual and seasonal impacts of shipping emissions on O₃ concentrations in China, especially for key coastal and inland regions from 2016 to 2020. We also ~~allocate apportion culpabilities of the contribution of~~ shipping emissions from ocean-going vessels (OGVs), coastal vessels (CVs), and river vessels (RVs) ~~on to~~ O₃ pollution to identify the influences of regionally differentiated shipping emission control policies. Furthermore, an explainable machine learning model was applied to explore investigate the potential source-receptor relationships between how shipping emissions affect and the ~~formation of~~O₃ formation based on five-years simulated data. Our study provides a nationwide and long-term analysis of the impacts shipping emissions on China's O₃ pollution, and provide new insights for shipping control measures in the future.

2 Methods

2.1 Shipping emissions

The Shipping Emission Inventory Model (SEIM v2.0) is a disaggregate dynamic method

(Wang et al., 2021b)(Luo et al., 2025)—driven by ~~driven~~ (a) the high-frequency ship Automatic Identification System (AIS) data, including signal time, coordinate location, navigational speed, and operating status, and (b) the integrated Ship Technical Specifications Database (STSD) (updated to 2020), which describes ship static properties, including vessel type, maximum designed speed, DWT and engine power. First, the originally collected raw AIS data and ship profile data from multiple sources are combined to form a ship activity database and STSD. Second, a route restoration module is applied for cross-land trajectory with a long distance in the AIS data, in which the 10 min linear interpolation will be applied on the shorted paths instead. Third, the instantaneous emission along with the movement of the ship's trajectory will be calculated based on the ship's static technical parameters, dynamic load changes, and extra parameters and factors. ~~Finally, shipping emission inventory datasets will be established and used for visualization and analyses from multiple perspectives.~~ In the SEIM, shipping emissions for both air pollutants (e.g., SO₂, PM, NO_x, CO and VOCHC) and greenhouse gases (e.g., CO₂, CH₄ and N₂O) from the main engines, auxiliary engines and boilers were calculated, detailed information of SEIM is described in our previous study (Wang et al., 2021b). Here, emissions beyond 200 nautical miles from the Chinese mainland's territorial sea baseline were excluded from the domain by applying GIS-based spatial processing to the global shipping emission inventory, and only the annual shipping emissions from 2016 to 2020 within 200 nautical miles were used in the simulation.~~Here, we calculated annual shipping emissions from 2016 to 2020 within 200 Nm from the Chinese mainland's territorial sea baseline.~~

In this study, vessels were classified as ocean-going vessels (OGVs), coastal vessels (CVs) and river vessels (RVs) for emission estimation according the following rules: (a) OGVs were identified by both valid International Maritime Organization (IMO) numbers and the Maritime Mobile Service Identity (MMSI) numbers, since they are mostly engaged in international trade following the management of the IMO; (b) RVs were identified by frequency distribution method based on the navigation trajectories for each vessel. Vessels with more than 50 % of the AIS signals throughout the entire year occurring on inland rivers (14–43° N, 104–130° E) were considered as RVs; ~~(c) Finally and;~~ (c) vessels that are not identified as OGVs or RVs are regarded as CVs. **Figure 1** shows the interannual variation of shipping NO_x and VOC emissions from 2016 to 2020. Overall, the increasing-growing trade demands and total cargo throughput of Chinese ports ~~led to contributed to increased elevated levels of shipping activities, which in turn resulted in a steady rise and consequently a sustained rise~~ in shipping NO_x and VOC emissions, especially for OGVs. Additionally, following the implementation of the global sulfur cap (IMO, 2018), the shift to low-sulfur fuels, which are typically richer in short-chain hydrocarbons (Wu et al., 2020), has contributed to a rise in shipping VOC emissions. It is worth noting that changes in vessel operating conditions, such as idling time and engine load, also influenced emissions. Although the COVID-19 pandemic had a temporary effect on maritime activity, its impact on annual shipping emissions was relatively minor due to the rapid rebound in trade during the second half of the year (Yi et al., 2024).




Figure 1. The interannual variation of shipping (a) NO_x and (b) VOC emissions from 2016 to 2020.

2.2 Air quality model

The Weather Research and Forecasting (WRF, version 3.8.1, using meteorological fields from 2018, as detailed later)–Community Multiscale Air Quality (CMAQ, version 5.4) model was applied to simulate the air quality in China during January, April, July and November from 2016 to 2020. Considering the relatively stable monthly anthropogenic emissions, this study simulated the O₃ concentrations during January, April, July, and November to represent winter, spring, summer, and fall, respectively, for the calculation of annual and seasonal mean values, which represented winter, spring, summer and fall, respectively, with 3 days of spin-up time for each run. As shown in **Figure S1S2**, the modeling domain covered all of China and some parts of East Asia with a horizontal resolution of 36 km × 36 km. Here, we have defined four key regions: BRA, YRD, PRD and inland river areas (Moirangthem, 2016), in which we focus on shipping-related O₃ pollution.

Here, we primarily focused on examining the impact of anthropogenic emission changes on shipping-related O₃ from a historical perspective. To eliminate the impact of interannual meteorological variability, we used meteorological field of 2018 (Zhao et al., 2022), which simulated by WRF and identified as a typical meteorological year due to its relatively stable climate conditions, to drive the CMAQ simulations for the period 2016-2020. Here the year of 2018 was regarded as a typical meteorological year since the climate for this year was relatively stable, and we used the meteorological filed of 2018 simulated by WRF in our previous study (Zhao et al., 2022) to drive the CMAQ simulation from 2016 to 2020. The first guess field and boundary conditions for WRF were generated from the 6 h NCEP FNL Operational Model Global Tropospheric Analyses dataset. The four-dimensional data assimilation (FDDA) was enabled using the NCEP ADP global surface and upper air observational weather data (<http://rda.ucar.edu>, last access: 25 March 2023). WRF and CMAQ used 32 vertical layers up to 100 hPa, and the lowest layer had a thickness of approximately 37 m. The major physical options in WRF included a Morrison two-moment microphysics scheme (Morrison et al., 2009), a Kain–Fritsch cumulus cloud parameterization (Kain, 2004), the Rapid Radiative Transfer Model (RRTM) longwave and shortwave radiation scheme (Iacono et al., 2008), the Pleim–Xiu

Land Surface Model (Xiu and Pleim, 2001), and the Asymmetric Convective Model version 3.0 for the PBL parameterization (Pleim, 2007). The distribution of meteorological stations for validation of and WRF performance is shown in Figure S3 and Table S1, respectively.

Atmospheric gas-phase chemistry in the CMAQ was simulated with the SAPRC07tic chemical mechanism, and aerosols were predicted using the AERO7. The chemical boundary conditions of CMAQ inputs, corresponding to each simulation period, were collected from the Community Atmosphere Model with Chemistry (CAM-chem) simulation output of global tropospheric and stratospheric compositions (Buchholz et al., 2019). Each run included a 3-day spin-up period. In this study, the Integrated Source Apportionment Method (ISAM) was applied to determine the source contribution to the ambient O₃ concentrations. Here, ISAM-OP3 was applied to attribute all secondary products to sources emitting NO_x or reactive VOC species and radicals when present in the parent reactants, and otherwise assign them based on stoichiometric reaction rates (Shu et al., 2023). We divided the emissions into five groups to trace them separately in the ISAM, including the land-based anthropogenic emission (the mobiles, industry, power, domestic, and agriculture) from the MEIC and the open burning emissions from Cai's study et al (Cai et al., 2017), the RVs' emission, the CVs' emission, the OGVs' emission, and the other emission (the nature sources emission from the MEGANv3 and the anthropogenic emission from other countries within the modeling domain from the MIX (Li et al., 2017), details of emissions are shown in **Table S2**.

We evaluated the simulated MDA8 O₃ concentrations for the of 2018 against 1455 available ground-based observations– (Figure S3) for model validation. ~~Until the simulation time, a total of 1455 available observation sites in China were used for O₃ validation.~~ As shown in **Table S3**, the simulated MDA8 O₃ agreed well with observations, with the overall model performance within the performance criteria suggested by Boylan and Russell (Boylan and Russell, 2006) (mean fractional bias (MFB) $\leq \pm 60\%$ and mean fractional error (MFE) $\leq \pm 75\%$), while the model overestimated O₃ a little, mainly due to uncertainties in emission inventory and unavoidable deficiencies during meteorological and air quality simulation. Meteorological performance for simulated periods was described in our previous study (Zhao et al., 2022).

2.3 Explainable machine learning model

Although CMAQ-ISAM can generate large amounts of simulation data to investigate the impacts of shipping emissions, the number of scenarios remains limited due to computational constraints. As a result, current analyses struggle to elucidate the mechanisms by which shipping emissions contribute to O₃ formation from these discrete scenarios. In particular, capturing nonlinear interactions between emission sources, meteorological conditions, and chemical processes is challenging when only a limited number of emission perturbations are available. Recently, the advancement of machine learning techniques, especially explainable models, has provided a promising complementary approach (Liu et al., 2025; Yao et al., 2024a, b). These models can learn from existing models to approximate the source-receptor relationships embedded in the simulation results. By identifying key emission drivers, quantifying their nonlinear contributions to O₃, and revealing latent patterns across spatiotemporal scales.

Here, based on the simulated data from 2016 to 2020 using WRF-CMAQ-ISAM, the RF model was used to simulated the monthly average O₃ concentration. In the RF model, the input predictor variables included relative humidity, temperature, wind speed, wind direction, solar radiation, land anthropogenic NO_x emissions, land anthropogenic VOC emissions, shipping NO_x emission and shipping VOC emission. Notably, the emissions selected were the sum of emissions from each grid and its eight neighboring grids. ~~To investigate the impact of shipping emissions on O₃ formation in different regions, w~~We trained four RF models specifically for the BRA, YRD, PRD, and IRA regions, respectively. The simulation data from 2016 to 2019 were used as the training samples, while the simulation data from 2020, under the scenario of the most significant changes in shipping emissions, used as the test samples to validate the generalization capability of the RF models. By comparing the root mean squared error (RMSE) for testing datasets across models with candidate parameter combinations, we set mtry and NumTrees as 6 and 200 in RF, respectively. Additionally, the 10-fold cross-validation repeated 10 times was considered to evaluate the prediction performance of our models. The total dataset was randomly divided into 10 subsets, where 9 subsets was used to train the model and another was applied for validation. As shown in **Figure 2**, averages of RMSE and correlation coefficient (R²) in the CV of the RF models were 2.12~2.47 ppb, and 0.90~0.98, respectively, indicating an acceptable performance.




Figure 2. Performances of RF models for (a) BRA, (b) YRD, (c) PRD and (d) IRA. Each point represents the monthly average O₃ concentration at each CMAQ grid cell.

In order to identify the sensitivity and response relationship between prediction variables and results in the RF models, the SHapley Additive exPlanations (SHAP) technique, a game-theoretic framework introduced by Lundberg et al. (Lundberg et al., 2020; Lundberg and Lee, 2017), was employed to interpret the pattern learned from the 2016-2019 simulation data by the RF models using the Python scikit-learn library. The SHAP approach enables the quantification of both global and local influences of input variables on the model's predictions, thereby improving the interpretability of factors contributing to air pollution. Additionally, the study examined feature interactions, which can affect the model's predictive accuracy, to gain a more comprehensive understanding of the intricate relationships among the variables.

2.4 Limitations

In this study, the spatial resolution of 36 km × 36 km may not fully capture the fine-scale spatial heterogeneity of O₃ concentrations, particularly in coastal urban areas where emissions and photochemical reactions exhibit strong spatial variability. This resolution is relatively coarse for accurately representing O₃ exceedances and local photochemical processes, which often occur at much finer spatial scales. Consequently, localized O₃ peaks and gradients may be underestimated or smoothed in the model outputs. Despite this limitation, the selected resolution represents a practical compromise that enables multi-year simulations across the national domain.

Only four representative months were simulated each year to reflect annual and seasonal patterns. While this captures broad seasonal variability, it may overlook intra-seasonal fluctuations and short-term anomalies. Using these months to estimate annual and seasonal means introduces uncertainty, especially for sources with stronger monthly variation. Although monthly changes in anthropogenic and shipping emissions are generally modest (except in 2020), future work could benefit from higher temporal resolution to improve accuracy.

Anthropogenic emissions from other countries within the modeling domain were held fixed at 2010 levels, and open burning emissions were fixed at 2015 levels throughout the simulation period (2016–2020). Although this assumption simplifies the modeling framework and is unlikely to significantly alter the relative changes in shipping-related O₃ assessed in this work, it may still introduce some degree of uncertainty, particularly in regions where long-range transport or fire-related emissions could have contributed more dynamically during specific years. Future studies could benefit from incorporating temporally varying background emissions to further reduce potential uncertainties and improve the representation of external influences.

Explainable machine learning model relies on the structure and quality of the input dataset and cannot account for unmeasured or omitted variables, such as hemispheric background O₃ concentrations. As a result, the derived feature importance reflects statistical associations rather than causal relationships. It should be noted that if one seeks to determine whether a given variable promotes or suppresses O₃ pollution using machine learning methods, additional field observations, experimental data, and corresponding simulation results may be required as supporting evidence. Considering the interactions among variables, even if individual contributions are small, the SHAP estimates for each explanatory variable are unlikely to perfectly reflect their actual contributions in

the underlying physical processes. Furthermore, in the presence of strong collinearity or complex nonlinear interactions, SHAP values may not fully disentangle overlapping influences among features.

In this study, monthly and annual mean O₃ concentrations were derived from hourly model outputs, rather than the widely used MDA8 O₃. While this approach is consistent with the study's focus on long-term trends and average responses, it may introduce bias due to the well-known overestimation of nighttime ozone in chemical transport models. A sensitivity test comparing shipping-related O₃ contributions based on hourly averages and MDA8 revealed that over oceanic areas, the difference may reach 2-5 ppb, while over land, it remains within 2 ppb. Given that the relative contribution of shipping emissions to total O₃ is generally low, the impact of this bias is expected to be limited.

3 Results and discussions

3.1 Annual O₃ impact from shipping emissions

Figure 3a shows the five-year average of shipping-related O₃ calculated based on hourly values, which is defined as the sum of O₃ concentration caused by emissions of OGVs, CVs, and RVs traced by CMAQ-ISAM. Overall, the shipping emissions increases O₃ concentrations by 3.5 ppb nationwide (Table S4), showing a decrease trend from the southeast coast toward inland areas. This result is greater than the findings in other countries such as 1.97 ppb in Europe, and 2.08 ppb in the United States under ambient temperature and pressure (Sun et al., 2024). Due to the high coastal shipping emissions, the shipping-related O₃ could exceed 15.0 ppb in southeast coastal regions, especially in YRD and PRD, where maximum values reach 25.4 ppb and 26.3 ppb, respectively. For the regions with low shipping emissions, the shipping-related O₃ is relatively low, not exceeding 5 ppb. For example, in IRA and BRA, the shipping-related O₃ is 3.9 ppb and 4.5 ppb respectively, but is slightly higher than the national average. Meteorological factors are just as important as anthropogenic emission influences in O₃ production (Liu et al., 2023; Zhang et al., 2024). For example, the PRD region is characterized by a persistently warm and humid climate, which provides favorable conditions for ozone formation. In contrast, although the BRA is also a coastal region, The BRA and IRA have it experiences lower temperatures and weaker solar radiation-compared to the YRD and PRD (Figure SXS4), which reduces the formation of hydroxyl radicals, weakening the atmospheric photochemical oxidizing capacity and ultimately limiting the O₃ production.

Figure 3b shows the five-year average of the relative contribution of shipping emissions to O₃. Nationwide, the shipping emissions accounts for a 9.98.6% increase in O₃, showing a similar decreasing trend of from the southeast coast to inland regions. This result is higher than 3.7% reported for the Mediterranean region in 2015 (Fink et al., 2023), but lower than the 12-21% reported in another European study for 2010 (Lupascu and Butler, 2019). Notably, some coastal cities exhibit particularly high values. For example, in PRD region, such as in Shenzhen, Guangdong Province, the relative shipping-related O₃ exceeds 30.4%-in Shenzhen, Guangdong Province, due to -the high intensity of shipping emissions combined with relatively low emissions from land-based anthropogenic sources such as industry and power generation. Similarly, tThe IRA is relatively low at 10.4%, mainly because of the higher background O₃ concentrations from land-based anthropogenic sources-or O₃ already formed by chemical reactions at sea and subsequently

transported inland by onshore winds (Cheng et al., 2023).




Figure 3. Contributions of shipping emissions to the five-year average O₃ pollution, including (a) absolute contributions and (b) relative contributions. Maps created with MeteoInfoMap (<http://www.meteothink.org>).

Figure 4 illustrates the interannual trend in shipping-related O₃ in key regions from 2016 to 2020. Overall Nationwide, the shipping-related O₃ shows a slight upward trend, with an average annual growth rate of 0.541.7%, — primarily observed in coastal regions. This trend aligns with the changes in shipping NO_x and VOC emissions, especially in 2020 when a 0.2-0.3 ppb rise in shipping-related O₃ was observed, partly attributable to the notable increase in VOC emissions following the implementation of the global sulfur cap. However, this increase in O₃ concentrations is substantially lower than the growth in precursor emissions, highlighting the complex and nonlinear response of O₃ formation to changes in shipping emissions which is lower than the increases in shipping-related NO_x and VOC (Figure 1). This finding suggests that O₃ formation is not highly sensitive to ship emissions. — This is because the formation of O₃ depends on photochemical reactions involving NO_x and VOC under solar radiation, and is influenced not only by the level of shipping emissions but also by land-based anthropogenic emissions, meteorological conditions, and long-range transport. Because O₃ is produced by photochemical reactions of NO_x and VOC in the presence of solar radiation, influenced by factors such as meteorological conditions, anthropogenic emissions, and long-range transport (Ye et al., 2023). Its response to shipping emissions is complex. Therefore, changes in shipping-related O₃ do not scale linearly with the changes in shipping NO_x and VOC emissions. — The Chinese government's two phases of clean air actions (Phase I, 2013–2017; Phase II, 2018–2020) resulted in increasing trend of O₃ nationwide (Liu et al., 2023), and the relative contribution of shipping emissions to O₃ also rose slightly during the same ~~period.~~ period. It is worth noting that, despite continuous increases in shipping NO_x and VOC emissions, their relative contributions to O₃ decreased in 2018 and 2020. This pattern may result from simultaneous land-based emission reductions, which can affect atmospheric oxidizing capacity (Lv et al., 2020).




Figure 4. The interannual trend in shipping-related O₃, including (a) absolute contributions and (b) relative contributions, in key regions from 2016 to 2020.

3.2 Contribution of different types of vessels

We further investigated the relationship between O₃ pollution and shipping emissions from sub-ship sectors. **Figure 5 and Table S4** shows ~~the spatial distribution of~~ the five-year average contribution of emissions from different ship types to the shipping-related O₃ and the total O₃, respectively. Nationwide, OGVs, CVs, and RVs contributed 2.6%, 2.6%, and 3.3% to the total O₃, respectively. In coastal regions, OGVs contribute more than 50% to most to the shipping-related O₃, accounting for more than 50% of such O₃ pollution and 9.7% to the total O₃. CVs are the second-largest contributor to O₃ pollution with an average contribution of 20-30%, and contribute up to 40% in the southern coastal regions near Zhejiang and Fujian Provinces. Notably, RVs can still contribute over 30% to shipping-related O₃ pollution in some coastal regions, particularly in the YRD and PRD. This is primarily due to the presence of major inland waterways such as the lower Yangtze River and the Pearl River, respectively, as well as the influence of regional pollutant transport that enhances the impact of RVs' emissions in these areas. ~~Due to the regional transport and some RVs sail on coastal regions, contributions from RVs can also reach 20-30%.~~ In inland regions, RVs remain the main source of shipping-related O₃, with contributions in the middle reaches of the Yangtze River reaching 50%.




Figure 5. Contributions of emissions from (a) OGVs, (b) CVs, and (c) RVs to the five-year average shipping-related O₃ pollution. Maps created with MeteoInfoMap (<http://www.meteothink.org>).

The interannual variation in the contribution of the different types of ship to shipping-related O₃ between 2016 and 2020 for different regions is illustrated in **Figure 6**. Nationwide, the contribution of OGVs and RVs increased by 2.7% and 0.6%, respectively, while the contribution of

CVs decreased by 3.3%. This pattern was observed in all coastal regions and IRA. The interannual variation in the contribution of the different types of ship to shipping-related O₃ follows a similar pattern to that of shipping-related NO_x and VOC (**Figure 1**), particularly NO_x emissions, which shows an upward trend for OGVs and RVs but a downward trend for CVs. As a result, the difference in the contribution of different types of ships to air quality is gradually narrowing. Notably, although RVs emissions significantly less than those of OGVs and CVs, its contribution to O₃ is comparable to that of other ship types, even exceeds that of CVs in some coastal regions. In addition, although China has required certain categories of ships to install AIS equipment since 2010, a large part of small RVs in China have not been equipped with AIS (Zhang et al., 2017), which is not considered in this study. This result suggests the importance of paying greater attention to RVs in future emission control strategies.




Figure 6. The interannual trend in Contributions of emissions from OGVs, CVs, and RVs to shipping-related O₃ in key regions from 2016 to 2020.

3.3 Seasonal O₃ impact from shipping emissions

The five-years-average seasonal variations in the contribution of shipping emissions to O₃ concentrations across different regions ~~are the continent is~~ shown in **Figure 7** and **Tabel 1**, with January, April, July, and November representing winter, spring, summer, and fall, respectively. ~~The relative contributions of ship emissions to O₃ in different seasons are presented in Figure S3. Overall, the contributions of ship emissions to O₃ concentrations in China were 1.53 ppb (5.6%), 4.24 ppb (9.8%), 4.77 ppb (13.7%), and 2.41 ppb (7.9%) for January, April, July, and November, respectively, which affected by the prevailing wind direction (Figure S4) and atmospheric oxidizing properties.~~

For cold seasons, including winter and fall, ~~the shipping-related O₃ remains relatively low, primarily~~ due to ~~less weaker solar irradiation~~ and lower temperatures ~~that limit O₃ formation~~ (**Figure S5S4**), ~~the shipping-related O₃ remains relatively lower than warm seasons (spring and summer), with national average and relative contribution of 1.53 ppb (5.6%) and 2.41 ppb (7.9%), respectively (Figure S5) Consequently, most regions remain under 5 ppb. However, except in the south of~~

~~YRAPRD~~, especially Guangdong and Hainan Provinces (**Figure 7a, 7d, and Table 1**), ~~where the average and maximum of seasonal shipping-related O₃ exceeds 5 ppb and 21 ppb, respectively. Notably, fall pollution even severer than that in summer. This is mainly because the PRD remains warm and humid in fall, and prevailing monsoon winds are more likely to transport ship-borne pollutants from the sea to inland areas (Figure S4, S6, and S7). Another distinct pattern is observed in BRA, where shipping-related O₃ formation tends to be more localized during the cold seasons, as indicated by a larger difference between the median and average values (Table 1). During this period, mainland China is under the influence of the Mongolian High Pressure System, and continental winds generally suppress the inland transport of ship-related O₃ (Cheng et al., 2023; Zhao et al., 2023). Therefore, significant shipping-related O₃ pollution only appears in major port cities with intensive maritime activity. (Figure 7a and 7d).~~

~~During winter and fall, when the wind direction is mainly north and northeast, shipping-related pollutants are transported to the sea, limiting the accumulation of O₃ on land (Cheng et al., 2023). However, shipping-related O₃ in fall is slightly higher than in winter, mainly because the persistent and intense solar radiation after the rainy summer season enhances O₃ formation. Another reason for this is that during the winter months, mainland China is under the influence of the Mongolian High Pressure System, where continental winds reduce the shipping-related O₃ (Cheng et al., 2023; Zhao et al., 2023).~~

~~Warm seasons generally include spring and summer. In spring, shipping-related O₃ reached its peak is highest in YRD and PRD in coastal areas, particularly across Guangdong, Fujian, and Zhejiang Provinces, where it with the maximum value exceedings 15-30 ppb (Figure 7b and Table 1). During this season, increasing solar radiation and moderate temperatures can enhance photochemical reactions, thereby promoting O₃ formation in these coastal areas, consistent with the results of previous studies (Cheng et al., 2023; Schwarzkopf et al., 2022). Although spring is generally less favorable for O₃ formation compared to summer in terms of temperature and humidity, strong onshore winds may play an important role in reduce the influence of shipping emissions (Cheng et al., 2023; Ma et al., 2022) (Figure S4, S6, and S7). In addition, more complex physicochemical interactions may drive springtime O₃ (Cao et al., 2024; Zhang et al., 2024), which needs further investigation. In summer, shipping emissions significantly increased O₃ concentrations nationwide by 4.77 ppb and responsible for 13.7% of national O₃ pollution (Figure 7c and Figure S5). Notably, even in IRA, where shipping emissions are much lower than in coastal regions, shipping-related O₃ were comparable to those along the coast. Most coastal regions near the BRA and the PRD recorded levels above 10 ppb, while central China, located hundreds of kilometers from the coast, was still significantly impacted with average level of around 10 ppb. This is primarily because central China lies in a perennial monsoon region, where summer monsoons can carry shipping-related air pollutants inland from coastal cities (Zheng et al., 2024). Notably, the impact of ship emissions on O₃ in PRD reaches its highest contribution in spring rather than summer, consistent with the results of previous studies (Cheng et al., 2023; Schwarzkopf et al., 2022). The possible reason for this is that in PRD, particularly Fujian and Guangdong Provinces, experiences high humidity and frequent precipitation in summer (Figure S6), which decreases both O₃ concentrations and the impact of shipping emissions, even though high temperatures can promote~~

photochemical activity. Strong onshore winds also play a role in reducing the influence of shipping emissions (Cheng et al., 2023; Ma et al., 2022).

These findings indicate that seasonal variations for shipping-related O₃ are driven by meteorological factors, particularly changes in prevailing wind direction, which are crucial for the diffusion and long-range transport of shipping emissions. Researchers suggest that the mixing emissions between shipping and local anthropogenic emissions can amplify complicated O₃ chemical formation in coastal cities (Wang et al., 2019b). Therefore, seasonal mitigation strategies and a better understanding of regional monsoon dynamics and their interaction with local anthropogenic emissions are crucial for effectively reducing shipping-related O₃ pollution.




Figure 7. Contributions of shipping emissions to the seasonal mean O₃ concentrations for (a) winter (JAN), (b) spring (APR), (c) summer (JUL), and (d) fall (NOV). Maps created with MeteoInfoMap (<http://www.meteothink.org>).

Table 1 Seasonal ranges of shipping-related O₃ across BRA, YRD, PRD, and IRA. (unit: ppb)

Region	Metric	Winter	Spring	Summer	Fall
BRA	Minimum	0.01	0.83	1.25	0.06
	25% Quartile	0.05	3.09	7.13	0.39
	Median	0.11	4.02	9.32	0.78
	75% Quartile	0.91	6.55	11.90	1.63

	<u>Maximum</u>	<u>6.59</u>	<u>23.22</u>	<u>32.40</u>	<u>14.39</u>
	<u>Mean</u>	<u>0.71</u>	<u>5.36</u>	<u>10.13</u>	<u>1.74</u>
<u>YRD</u>	<u>Minimum</u>	<u>0.20</u>	<u>1.92</u>	<u>1.45</u>	<u>0.55</u>
	<u>25% Quartile</u>	<u>1.01</u>	<u>5.02</u>	<u>6.11</u>	<u>1.91</u>
	<u>Median</u>	<u>1.79</u>	<u>6.64</u>	<u>7.29</u>	<u>3.23</u>
	<u>75% Quartile</u>	<u>2.93</u>	<u>8.35</u>	<u>9.03</u>	<u>5.14</u>
	<u>Maximum</u>	<u>16.32</u>	<u>31.47</u>	<u>25.86</u>	<u>24.59</u>
	<u>Mean</u>	<u>2.47</u>	<u>7.39</u>	<u>7.92</u>	<u>4.41</u>
<u>PRD</u>	<u>Minimum</u>	<u>1.11</u>	<u>3.91</u>	<u>0.11</u>	<u>1.91</u>
	<u>25% Quartile</u>	<u>2.79</u>	<u>7.94</u>	<u>3.19</u>	<u>4.85</u>
	<u>Median</u>	<u>5.19</u>	<u>10.07</u>	<u>5.65</u>	<u>7.97</u>
	<u>75% Quartile</u>	<u>7.68</u>	<u>15.17</u>	<u>7.77</u>	<u>11.25</u>
	<u>Maximum</u>	<u>21.98</u>	<u>33.46</u>	<u>26.19</u>	<u>28.78</u>
	<u>Mean</u>	<u>5.96</u>	<u>11.91</u>	<u>5.77</u>	<u>8.79</u>
<u>IRA</u>	<u>Minimum</u>	<u>0.17</u>	<u>0.91</u>	<u>2.32</u>	<u>0.10</u>
	<u>25% Quartile</u>	<u>1.19</u>	<u>2.88</u>	<u>5.65</u>	<u>1.44</u>
	<u>Median</u>	<u>1.61</u>	<u>4.85</u>	<u>9.07</u>	<u>2.85</u>
	<u>75% Quartile</u>	<u>2.29</u>	<u>5.73</u>	<u>11.31</u>	<u>4.16</u>
	<u>Maximum</u>	<u>5.53</u>	<u>11.91</u>	<u>26.19</u>	<u>7.52</u>
	<u>Mean</u>	<u>1.76</u>	<u>4.58</u>	<u>9.03</u>	<u>2.87</u>

1

2 3.4 Effects of shipping emission on O₃ formation

3 Although the features in the RF model include both meteorological factors and emissions,
4 this section focuses on impacts of anthropogenic emissions, especially shipping emissions, on
5 O₃ pollution. **Figure 8** presents the SHAP summary plots for selected features in the RF model
6 for BRA, YRD, PRD, and IRA, showing the magnitude, prevalence, and direction of each
7 feature's impact on the model output (O₃ concentration). In the summary plot, the further a
8 feature's SHAP value is from zero, the greater its influence; positive SHAP values indicate a
9 positive contribution, while negative values indicate a negative effect. For example, in **Figure**
10 **8a**, land-based NO_x and VOC emissions both have a significant impact on O₃ formation in the
11 BRA, with contributions of 16.8% and 11.0%, respectively, suggesting that the atmospheric
12 chemistry in this area is significantly affected by land-based anthropogenic emissions.

Moreover, lower NO_x and VOC emissions leads to higher SHAP values, indicating a negative correlation between land-based anthropogenic emissions and O₃ pollution. In the coastal areas of BRA, shipping emissions are much smaller than land-based emissions, therefore, contribute only approximately 2.9% to O₃ formation, and exhibit a similar negative correlation as land-based sources.

As the share of shipping emissions increases within total anthropogenic emissions in the coastal areas of YRD and PRD, the difference between the contributions of shipping and land-based emissions to O₃ pollution regions decreases. Especially in the PRD, shipping NO_x contributes up to 9.7%, exceeding the land-based VOC contribution of 5.3%. In the IRA, emissions from inland vessels are much lower than those from ocean-going and coastal ships, thus, shipping NO and VOC contributions only 1.5% and 0.8% to O₃ pollution, respectively.

For meteorological factors, the contributions of solar radiation (25.2%), temperature (20.0%), wind speed (11.5%), and wind direction (4.9%), relative humidity (4.8%) to O₃ formation exhibit clear regional heterogeneity. Overall, meteorological influences are greater than those of shipping emissions. This may be attributed to the highly complex physical and chemical processes involved, including cloud–radiation interactions, air mass transport, and water-related reactions.




Figure 8. Feature importance results of the random forest regression model for (a) BRA, (b) YRD, (c) PRD, and (d) IRA. The x-axis shows SHAP values representing the impact of each feature on O₃ predictions (positive: increasing O₃; negative: decreasing O₃). Each dot is a grid-month sample, with color indicating the feature value. Instances with identical x-values are stacked, and the stack height signifies the density.

The dependence plot (**Figure. 9**) further quantifies how changes in shipping emissions affect O₃ concentrations. Notably, due to the uneven distribution of shipping emissions, the x-axis in **Figure. 9** is set to non-uniform intervals to better illustrate their impact. Overall, as previously mentioned, changes in shipping NO_x and VOC emissions have a relatively minor effect on regional O₃ levels—approximately between –2.5 ppb to 8 ppb—although there is clear regional heterogeneity. In the BRA region, shipping emissions are negatively correlated with O₃, with increases in shipping NO_x and VOC emissions leading to a reduction of about 1–2 ppb in O₃. Differently, in the YRD and PRD, increases in shipping NO_x emissions slightly promote O₃ formation—especially in the PRD region, where O₃ may increase by up to 3 ppb, which is opposite to the impact of land-based NO_x emissions (**Figure. S7**). This difference may be because, in the YRD and PRD, land-based NO_x emissions do not dominate the overall anthropogenic emissions as they do in the BRA, allowing shipping NO_x to also influence atmospheric chemistry. Furthermore, changes in shipping VOC emissions have almost no impact, consistent with the effect of changes in land-based VOC emissions (**Figure. S7**). Only when shipping VOC emissions increase dramatically in the YRD region is O₃ formation suppressed—though such cases are rare, as reflected by the partially negative SHAP values for shipping VOC emissions in **Figure. 8b**. In the IRA, similar to the BRA region, shipping emissions are very small compared to land-based sources, so that changes in shipping NO_x and VOC emissions have a similar negative effect on O₃ as those from land-based sources.

~~It should be noted that if one seeks to determine whether a given variable promotes or suppresses O₃ pollution using machine learning methods, additional field observations, experimental data, and corresponding simulation results may be required as supporting evidence. Considering the interactions among variables, even if individual contributions are small, the SHAP estimates for each explanatory variable are unlikely to perfectly reflect their actual contributions in the underlying physical processes.~~




Figure 9. SHAP dependence plot for (a) BRA, (b) YRD, (c) PRD, and (d) IRA.

4 Conclusion

In this study, we conducted multi-year CMAQ-ISAM simulations to investigate the how shipping emissions impacted O_3 across China, with a focus on three coastal regions and a inland region. From 2016 to 2020, shipping emissions increased national average O_3 concentrations by 3.5 ppb, accounting for 8.6% of total O_3 , with a spatial gradient decreasing from coastal to inland regions. Despite the increasing intensity of shipping activity and the implementation of the global sulfur cap, shipping NO_x and VOCs emissions rose significantly during this period. However, the national average shipping-related O_3 increased by only 0.23 ppb, while the relative contribution of shipping emissions to O_3 pollution rose by approximately 0.5%. Notably, this relative contribution did not increase continuously; instead, a decline was observed in 2018 and 2020. This non-linear response, under conditions of simultaneous changes in multiple pollutants from different sectors, highlights the complexity and need for further investigation of attribution of O_3 pollution. For the four focus regions, the contribution of shipping to O_3 levels exceeded the national average, with more pronounced interannual increases.

We further disaggregate ship types to OGVs, CVs, and RVs. The result revealed that OGVs were the dominant contributors to shipping-related O_3 in coastal areas, followed by CVs, whereas RVs were the main source in inland river areas. Although OGVs, CVs, and RVs differ significantly in their emission magnitudes, the difference in their contributions to O_3 pollution is gradually narrowing. This trend suggests that the influence of RVs on regional O_3 levels should no longer be overlooked and that emission control efforts for RVs deserve renewed attention. However, from the perspective of sulfur emission control, RVs in China had already reached the final stage of sulfur regulation by 2018 under the implementation of domestic emission control policies. In contrast, NO_x control for inland vessels remains largely unaddressed. Globally, there is limited precedent or experience in regulating NO_x emissions from inland waterways, leaving China without a clear reference framework for RVs NO_x mitigation. Future control of shipping NO_x emissions needs to take into account both inland

1 waterways and coastal areas.

2 Overall, the shipping emissions increases O_3 concentrations by 3.5 ppb nationwide, with
3 a significant spatial and temporal heterogeneity. The air quality in Chinese coastal cities is
4 highly impacted by shipping emissions. For example, the relative shipping-related O_3 exceeds
5 26.3 ppb (30.4%) in PRD, while is relatively low at 3.9 ppb (10.4%) in IRA. Such pollution is
6 more serious in warm seasons with an average of 4.77 ppb (13.7%) nationwide in summer, due
7 to the increased solar radiation and moderate temperatures. although OGVs, CVs, and RVs
8 exhibit significant differences in their emissions, the difference in their contributions to O_3
9 pollution are diminishing and becoming relatively comparable on a national scale. The impacts
10 of shipping emissions on O_3 also exhibited significant seasonal and regional characteristics.
11 While shipping-related O_3 levels were generally lower in colder seasons, fall pollution in
12 southern coastal regions exceeded that of summer due to favorable land-sea monsoon transport.
13 Peak shipping-related O_3 levels occurred in spring over YRD and PRD, and in summer over
14 inland areas. These patterns highlight the importance of implementing seasonal and region-
15 specific control strategies to mitigate shipping-related O_3 pollution effectively. In particular,
16 quality-oriented management policies such as seasonal routing adjustments, port operation
17 scheduling, or dynamic emission monitoring, may play a more immediate role than emission
18 control policies, which are typically less adaptable to seasonal variability and require long-term
19 infrastructure or regulatory changes. Therefore, combining flexible operational measures with
20 long-term emission reduction plans could enhance the overall effectiveness of O_3 mitigation.

21 Interpretable machine learning analysis further revealed significant spatial differences in
22 the contribution of shipping emissions to O_3 . In BRA and IRA, O_3 formation was primarily
23 driven by land-based NO_x and VOC emissions, with shipping emissions playing a minor role
24 and even showing a suppressive effect on O_3 formation. In contrast, in coastal regions such as
25 YRD and PRD, the increasing share of shipping emissions in the total anthropogenic emissions
26 enhanced their contribution to O_3 , with shipping NO_x emissions showing a slight promoting
27 effect on O_3 formation. To mitigate O_3 pollution, it is important to coordinate the control of
28 shipping emissions with land-based emission sources. The results show that the impact of
29 shipping NO_x and VOC emissions on O_3 formation exhibits significant regional differences. In
30 areas where land-based emissions dominate, such as the BRA region, the impact of shipping
31 emissions on O_3 pollution is relatively small and shows a negative correlation. In contrast, in
32 coastal regions such as the YRD and PRD, where the share of land-based emissions has
33 decreased, the contribution of shipping emissions to O_3 formation becomes more significant.
34 In particular, s shipping NO_x emissions have become an important source of O_3 precursors in
35 the PRD region. This regional difference suggests that solely controlling shipping emissions
36 may lead to unexpected atmospheric chemical responses and, under certain conditions, could
37 even cause an increase in O_3 concentrations. Therefore, effective O_3 pollution control requires
38 a coordinated reduction of both land-based and shipping emissions, based on regional emission
39 structures and atmospheric oxidation characteristics. Further analysis reveals that
40 meteorological conditions play an even more critical role in regulating O_3 formation. Variables
41 such as solar radiation, temperature, wind speed, and relative humidity show strong sensitivity
42 to changes in O_3 concentration and also display clear regional differences. Therefore, it is

~~recommended that differentiated and refined joint control strategies be developed for different regions based on their specific emission characteristics and meteorological conditions.~~

Data availability

Data used during the current study are available from the corresponding author upon reasonable request.

Code availability

Codes used during the current study are available from the corresponding author upon reasonable request.

Acknowledgments

This research has been supported by: National Natural Science Foundation of China (grant nos. 42325505 to H.L.), National Key Research and Development Program of China (grant no. 2022YFC3704200 to H.L.), the China National Postdoctoral Program for Innovative Talents (No. BX20250327 to Z.Luo), the Shuimu Tsinghua Scholar Program (no. 2024SM028 to Z.Luo), National Natural Science Foundation of China (grant no. 42405097 to Z.Lv).

Author Contributions

Conceptualization: Z.Luo, L.P., H.L.; Methodology: Z.Luo, Z.Lv, J.Zhao; Investigation: W.Y., T.H., Y.W; Visualization: Z.Luo L.P.; Supervision: H.L., K.H.; Writing—original draft: Z.Luo, L.P.; Writing—review & editing: Z.Luo, H.L.

Declaration of interests

The authors declare no competing interests.

Reference

- Boylan, J. W. and Russell, A. G.: PM and light extinction model performance metrics, goals, and criteria for three-dimensional air quality models, *Atmos. Environ.*, 40, 4946–4959, <https://doi.org/10.1016/j.atmosenv.2005.09.087>, 2006.
- Buchholz, R. R., Emmons, L. K., Tilmes, S., and The CESM2 Development Team: CESM2.1/CAM-chem instantaneous output for boundary conditions, <https://doi.org/10.5065/NMP7-EP60>, 2019.
- Cai, S., Wang, Y., Zhao, B., Wang, S., Chang, X., and Hao, J.: The impact of the “air pollution prevention and control action plan” on PM_{2.5} concentrations in jing-jin-ji region during 2012–2020, *Sci. Total Environ.*, 580, 197–209,

- 1 <https://doi.org/10.1016/j.scitotenv.2016.11.188>, 2017.
- 2 Cao, T., Wang, H., Chen, X., Li, L., Lu, X., Lu, K., and Fan, S.: Rapid increase in spring ozone
- 3 in the pearl river delta, china during 2013-2022, *npj Clim. Atmos. Sci.*, 7, 309,
- 4 <https://doi.org/10.1038/s41612-024-00847-3>, 2024.
- 5 Cheng, Q., Wang, X., Chen, D., Ma, Y., Zhao, Y., Hao, J., Guo, X., Lang, J., and Zhou, Y.:
- 6 Impact of ship emissions on air quality in the guangdong-hong kong-macao greater bay
- 7 area (GBA): With a particular focus on the role of onshore wind, *Sustainability*, 15, 8820,
- 8 <https://doi.org/10.3390/su15118820>, 2023.
- 9 Feng, X., Ma, Y., Lin, H., Fu, T.-M., Zhang, Y., Wang, X., Zhang, A., Yuan, Y., Han, Z., Mao,
- 10 J., Wang, D., Zhu, L., Wu, Y., Li, Y., and Yang, X.: Impacts of ship emissions on air quality
- 11 in southern China: Opportunistic insights from the abrupt emission changes in early 2020,
- 12 *Environ. Sci. Technol.*, 57, 16999–17010, <https://doi.org/10.1021/acs.est.3c04155>, 2023.
- 13 Fink, L., Karl, M., Matthias, V., Oppo, S., Kranenburg, R., Kuenen, J., Moldanova, J.,
- 14 Jutterstrom, S., Jalkanen, J.-P., and Majamaki, E.: Potential impact of shipping on air
- 15 pollution in the Mediterranean region - a multimodel evaluation: comparison of
- 16 photooxidants NO₂ and O₃, *Atmos. Chem. Phys.*, 23, 1825–1862,
- 17 <https://doi.org/10.5194/acp-23-1825-2023>, 2023.
- 18 Fu, M., Liu, H., Jin, X., and He, K.: National- to port-level inventories of shipping emissions
- 19 in China, *Environ. Res. Lett.*, 12, 114024, <https://doi.org/10.1088/1748-9326/aa897a>,
- 20 2017.
- 21 Fu, X., Chen, D., Wang, X., Li, Y., Lang, J., Zhou, Y., and Guo, X.: The impacts of ship
- 22 emissions on ozone in eastern China, *Sci. Total Environ.*, 903, 166252,
- 23 <https://doi.org/10.1016/j.scitotenv.2023.166252>, 2023.
- 24 Huang, H., Zhou, C., Huang, L., Xiao, C., Wen, Y., Li, J., and Lu, Z.: Inland ship emission
- 25 inventory and its impact on air quality over the middle yangtze river, china, *Sci. Total*
- 26 *Environ.*, 843, 156770, <https://doi.org/10.1016/j.scitotenv.2022.156770>, 2022.
- 27 Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W.
- 28 D.: Radiative forcing by long-lived greenhouse gases: calculations with the AER radiative
- 29 transfer models, *J. Geophys. Res.: Atmos.*, 113, <https://doi.org/10.1029/2008JD009944>,
- 30 2008.
- 31 IMO, 2018. Resolution MEPC.305(73), Amendments to the Annex of the Protocol of 1997 to
- 32 Amend the International Convention for the Prevention of Pollution from Ships.
- 33 Ito, K., De Leon, S. F., and Lippmann, M.: Associations between ozone and daily mortality:
- 34 analysis and meta-analysis, *Epidemiol. (Camb. Mass.)*, 16, 446–457,
- 35 <https://doi.org/10.1097/01.ede.0000165821.90114.7f>, 2005.
- 36 Jerrett, M., Burnett, R. T., Pope, C. A., Ito, K., Thurston, G., Krewski, D., Shi, Y., Calle, E., and
- 37 Thun, M.: Long-term ozone exposure and mortality, *N. Engl. J. Med.*, 360, 1085–1095,
- 38 <https://doi.org/10.1056/NEJMoa0803894>, 2009.
- 39 Kain, J. S.: The kain–fritsch convective parameterization: an update, 2004.
- 40 Li, M., Zhang, Q., Kurokawa, J., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G.,
- 41 Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and
- 42 Zheng, B.: MIX: a mosaic asian anthropogenic emission inventory under the international
- 43 collaboration framework of the MICS-asia and HTAP, *Atmos. Chem. Phys.*, 17, 935–963,

- 1 <https://doi.org/10.5194/acp-17-935-2017>, 2017.
- 2 Liu, J., Chen, M., Chu, B., Chen, T., Ma, Q., Wang, Y., Zhang, P., Li, H., Zhao, B., Xie, R.,
3 Huang, Q., Wang, S., and He, H.: Assessing the significance of regional transport in ozone
4 pollution through machine learning: a case study of hainan island, *ACS ES&T Air*, 2, 416–
5 425, <https://doi.org/10.1021/acsestair.4c00297>, 2025.
- 6 Liu, Y., Geng, G., Cheng, J., Liu, Y., Xiao, Q., Liu, L., Shi, Q., Tong, D., He, K., and Zhang,
7 Q.: Drivers of increasing ozone during the two phases of clean air actions in China 2013–
8 2020, *Environ. Sci. Technol.*, 57, 8954–8964, <https://doi.org/10.1021/acs.est.3c00054>,
9 2023.
- 10 Lundberg, S. M. and Lee, S.-I.: A unified approach to interpreting model predictions, in:
11 *Proceedings of the 31st International Conference on Neural Information Processing*
12 *Systems*, Red Hook, NY, USA, 4768–4777, 2017.
- 13 Lundberg, S. M., Erion, G., Chen, H., DeGrave, A., Prutkin, J. M., Nair, B., Katz, R.,
14 Himmelfarb, J., Bansal, N., and Lee, S.-I.: From local explanations to global
15 understanding with explainable AI for trees, *Nat. Mach. Intell.*, 2, 56–67,
16 <https://doi.org/10.1038/s42256-019-0138-9>, 2020.
- 17 Luo, Z., Lv, Z., Zhao, J., Sun, H., He, T., Yi, W., Zhang, Z., He, K., and Liu, H.: Shipping-
18 related pollution decreased but mortality increased in Chinese port cities, *Nat Cities*, 1,
19 295–304, <https://doi.org/10.1038/s44284-024-00050-8>, 2024.
- 20 Luo, Z., He, T., Lv, Z., Zhao, J., Zhang, Z., Wang, Y., Yi, W., Lu, S., He, K., and Liu, H.: Insights
21 into transportation CO₂ emissions with big data and artificial intelligence, *Patterns*,
22 101186, <https://doi.org/10.1016/j.patter.2025.101186>, 2025.
- 23 Lupascu, A. and Butler, T.: Source attribution of european surface O₃ using a tagged O₃
24 mechanism, *Atmos. Chem. Phys.*, 19, 14535–14558, [https://doi.org/10.5194/acp-19-](https://doi.org/10.5194/acp-19-14535-2019)
25 14535-2019, 2019.
- 26 Lv, Z., Wang, X., Deng, F., Ying, Q., Archibald, A. T., Jones, R. L., Ding, Y., Cheng, Y., Fu, M.,
27 Liu, Y., Man, H., Xue, Z., He, K., Hao, J., and Liu, H.: Source–receptor relationship
28 revealed by the halted traffic and aggravated haze in beijing during the COVID-19
29 lockdown, *Environ. Sci. Technol.*, 54, 15660–15670,
30 <https://doi.org/10.1021/acs.est.0c04941>, 2020.
- 31 Ma, Y., Chen, D., Fu, X., Shang, F., Guo, X., Lang, J., and Zhou, Y.: Impact of sea breeze on
32 the transport of ship emissions: A comprehensive study in the bohai rim region, China,
33 *Atmosphere*, 13, 1094, <https://doi.org/10.3390/atmos13071094>, 2022.
- 34 Merico, E., Donato, A., Gambaro, A., Cesari, D., Gregoris, E., Barbaro, E., Dinoi, A.,
35 Giovanelli, G., Masieri, S., and Contini, D.: Influence of in-port ships emissions to gaseous
36 atmospheric pollutants and to particulate matter of different sizes in a Mediterranean
37 harbour in Italy, *Atmos. Environ.*, 139, 1–10,
38 <https://doi.org/10.1016/j.atmosenv.2016.05.024>, 2016.
- 39 Moirangthem, K., 2016. Alternative fuels for marine and inland waterways: An exploratory
40 study.
- 41 Morrison, H., Thompson, G., and Tatarskii, V.: Impact of cloud microphysics on the
42 development of trailing stratiform precipitation in a simulated squall line: comparison of
43 one- and two-moment schemes, <https://doi.org/10.1175/2008MWR2556.1>, 2009.

- 1 Pleim, J. E.: A combined local and nonlocal closure model for the atmospheric boundary layer.
2 Part I: model description and testing, <https://doi.org/10.1175/JAM2539.1>, 2007.
- 3 Schwarzkopf, D. A., Petrik, R., Matthias, V., Quante, M., Yu, G., and Zhang, Y.: Comparison
4 of the impact of ship emissions in northern europe and eastern China, *Atmosphere*, 13,
5 894, <https://doi.org/10.3390/atmos13060894>, 2022.
- 6 Shu, Q., Napelenok, S. L., Hutzell, W. T., Baker, K. R., Henderson, B. H., Murphy, B. N., and
7 Hogrefe, C.: Comparison of ozone formation attribution techniques in the northeastern
8 united states, *Geosci. Model Dev.*, 16, 2303–2322, [https://doi.org/10.5194/gmd-16-2303-](https://doi.org/10.5194/gmd-16-2303-2023)
9 2023, 2023.
- 10 Sun, W., Jiang, W., and Li, R.: Global health benefits of shipping emission reduction in early
11 2020, *Atmos. Environ.*, 333, 120648, <https://doi.org/10.1016/j.atmosenv.2024.120648>,
12 2024.
- 13 Tao, Y., Huang, W., Huang, X., Zhong, L., Lu, S.-E., Li, Y., Dai, L., Zhang, Y., and Zhu, T.:
14 Estimated Acute Effects of Ambient Ozone and Nitrogen Dioxide on Mortality in the Pearl
15 River Delta of Southern China, *Environ Health Perspect*, 120, 393–398,
16 <https://doi.org/10.1289/ehp.1103715>, 2012.
- 17 Wan, Z., Cai, Z., Zhao, R., Zhang, Q., Chen, J., and Wang, Z.: Quantifying the air quality impact
18 of ship emissions in China's bohai bay, *Mar. Pollut. Bull.*, 193, 115169,
19 <https://doi.org/10.1016/j.marpolbul.2023.115169>, 2023.
- 20 Wang, F., Qiu, X., Cao, J., Peng, L., Zhang, N., Yan, Y., and Li, R.: Policy-driven changes in
21 the health risk of PM_{2.5} and O₃ exposure in China during 2013–2018, *Sci. Total Environ.*,
22 757, 143775, <https://doi.org/10.1016/j.scitotenv.2020.143775>, 2021a.
- 23 Wang, H., Gao, Y., Sheng, L., Wang, Y., Zeng, X., Kou, W., Ma, M., and Cheng, W.: The impact
24 of meteorology and emissions on surface ozone in shandong province, china, during
25 summer 2014–2019, *International Journal of Environmental Research and Public Health*,
26 19, 6758, <https://doi.org/10.3390/ijerph19116758>, 2022.
- 27 Wang, P., Chen, Y., Hu, J., Zhang, H., and Ying, Q.: Attribution of tropospheric ozone to NO_x
28 and VOC emissions: considering ozone formation in the transition regime, *Environ. Sci.*
29 *Technol.*, 53, 1404–1412, <https://doi.org/10.1021/acs.est.8b05981>, 2019a.
- 30 Wang, R., Tie, X., Li, G., Zhao, S., Long, X., Johansson, L., and An, Z.: Effect of ship emissions
31 on O₃ in the Yangtze River delta region of China: analysis of WRF-chem modeling, *Sci.*
32 *Total Environ.*, 683, 360–370, <https://doi.org/10.1016/j.scitotenv.2019.04.240>, 2019b.
- 33 Wang, T., Xue, L., Brimblecombe, P., Lam, Y. F., Li, L., and Zhang, L.: Ozone pollution in
34 China: a review of concentrations, meteorological influences, chemical precursors, and
35 effects, *Sci. Total Environ.*, 575, 1582–1596,
36 <https://doi.org/10.1016/j.scitotenv.2016.10.081>, 2017.
- 37 Wang, X., Yi, W., Lv, Z., Deng, F., Zheng, S., Xu, H., Zhao, J., Liu, H., and He, K.: Ship
38 emissions around china under gradually promoted control policies from 2016 to 2019,
39 *Atmos. Chem. Phys.*, 21, 13835–13853, <https://doi.org/10.5194/acp-21-13835-2021>,
40 2021b.
- 41 Wu, Z., Zhang, Y., He, J., Chen, H., Huang, X., Wang, Y., Yu, X., Yang, W., Zhang, R., Zhu, M.,
42 Li, S., Fang, H., Zhang, Z., and Wang, X.: Dramatic increase in reactive volatile organic
43 compound (VOC) emissions from ships at berth after implementing the fuel switch policy

- in the pearl river delta emission control area, *Atmos. Chem. Phys.*, 20, 1887–1900, <https://doi.org/10.5194/acp-20-1887-2020>, 2020.
- Xie, Y., Dai, H., Zhang, Y., Wu, Y., Hanaoka, T., and Masui, T.: Comparison of health and economic impacts of PM_{2.5} and ozone pollution in China, *Environ. Int.*, 130, 104881, <https://doi.org/10.1016/j.envint.2019.05.075>, 2019.
- Xiu, A. and Pleim, J. E.: Development of a land surface model. Part I: application in a mesoscale meteorological model, 2001.
- Yao, L., Han, Y., Qi, X., Huang, D., Che, H., Long, X., Du, Y., Meng, L., Yao, X., Zhang, L., and Chen, Y.: Determination of major drive of ozone formation and improvement of O₃ prediction in typical north China plain based on interpretable random forest model, *Sci. Total Environ.*, 934, 173193, <https://doi.org/10.1016/j.scitotenv.2024.173193>, 2024a.
- Yao, T., Lu, S., Wang, Y., Li, X., Ye, H., Duan, Y., Fu, Q., and Li, J.: Revealing the drivers of surface ozone pollution by explainable machine learning and satellite observations in hangzhou bay, china, *J. Cleaner Prod.*, 440, 140938, <https://doi.org/10.1016/j.jclepro.2024.140938>, 2024b.
- Ye, C., Lu, K., Ma, X., Qiu, W., Li, S., Yang, X., Xue, C., Zhai, T., Liu, Y., Li, X., Li, Y., Wang, H., Tan, Z., Chen, X., Dong, H., Zeng, L., Hu, M., and Zhang, Y.: HONO chemistry at a suburban site during the EXPLORE-YRD campaign in 2018: formation mechanisms and impacts on O₃ production, *Atmos. Chem. Phys.*, 23, 15455–15472, <https://doi.org/10.5194/acp-23-15455-2023>, 2023.
- Yi, W., He, T., Wang, X., Soo, Y. H., Luo, Z., Xie, Y., Peng, X., Zhang, W., Wang, Y., Lv, Z., He, K., and Liu, H.: Ship emission variations during the COVID-19 from global and continental perspectives, *Sci. Total Environ.*, 954, 176633, <https://doi.org/10.1016/j.scitotenv.2024.176633>, 2024.
- Yi, W., Wang, X., He, T., Liu, H., Luo, Z., Lv, Z., and He, K.: The high-resolution global shipping emission inventory by the shipping emission inventory model (SEIM), *Earth Syst. Sci. Data*, 17, 277–292, <https://doi.org/10.5194/essd-17-277-2025>, 2025.
- Zhang, L., Wang, L., Ji, D., Xia, Z., Nan, P., Zhang, J., Li, K., Qi, B., Du, R., Sun, Y., Wang, Y., and Hu, B.: Explainable ensemble machine learning revealing the effect of meteorology and sources on ozone formation in megacity hangzhou, china, *Sci. Total Environ.*, 922, 171295, <https://doi.org/10.1016/j.scitotenv.2024.171295>, 2024.
- Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu, W., Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L., Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu, B., Liu, Z., Yang, F., He, K., and Hao, J.: Drivers of improved PM_{2.5} air quality in China from 2013 to 2017, *Proc. Natl. Acad. Sci. U. S. A.*, 116, 24463–24469, <https://doi.org/10.1073/pnas.1907956116>, 2019.
- Zhang, Y., Yang, X., Brown, R., Yang, L., Morawska, L., Ristovski, Z., Fu, Q., and Huang, C.: Shipping emissions and their impacts on air quality in China, *Sci. Total Environ.*, 581–582, 186–198, <https://doi.org/10.1016/j.scitotenv.2016.12.098>, 2017.
- Zhao, C.-B., Li, Q.-Q., Nie, Y., Wang, F., Xie, B., Dong, L.-L., and Wu, J.: The reversal of surface air temperature anomalies in China between early and late winter 2021/2022: Observations and predictions, *Advances in Climate Change Research*, 14, 660–670,

1 <https://doi.org/10.1016/j.accre.2023.09.004>, 2023.

2 Zhao, J., Lv, Z., Qi, L., Zhao, B., Deng, F., Chang, X., Wang, X., Luo, Z., Zhang, Z., Xu, H.,

3 Ying, Q., Wang, S., He, K., and Liu, H.: Comprehensive assessment for the impacts of

4 S/IVOC emissions from mobile sources on SOA formation in China, *Environ. Sci.*

5 *Technol.*, 56, 16695–16706, <https://doi.org/10.1021/acs.est.2c07265>, 2022.

6 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L.,

7 Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic

8 emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18,

9 14095–14111, <https://doi.org/10.5194/acp-18-14095-2018>, 2018.

10 Zheng, S., Jiang, F., Feng, S., Liu, H., Wang, X., Tian, X., Ying, C., Jia, M., Shen, Y., Lyu, X.,

11 Guo, H., and Cai, Z.: Impact of marine shipping emissions on ozone pollution during the

12 warm seasons in China, *J. Geophys. Res.: Atmos.*, 129, e2024JD040864,

13 <https://doi.org/10.1029/2024JD040864>, 2024.