

Response to reviewers

Global Inductive Magnetosphere-Ionosphere-Thermosphere Coupling

We sincerely thank the reviewers for very constructive and important comments. We have prepared a revised manuscript that includes a new appendix to describe how the 2D equations can be derived by integrating over the ionospheric Ohm's law. We have also made a number of smaller additions and changes. The reviewers' text is quoted below in black, with our response to each point in blue. New text that is added to the manuscript is shown in dark red. A version of the manuscript where all the proposed changes are highlighted is attached.

Arthur D. Richmond

This is an excellent description and analysis of magnetic induction effects on global ionospheric electrodynamics. It describes how the electroquasistatic assumption commonly used to model global electrodynamics can be avoided in order to examine the physical processes by which ionospheric electric fields and currents are established. A numerical model is developed to demonstrate and analyze the effects of induction. A surprising prediction of the modeling is that fully steady-state electrodynamic conditions within the ionosphere may require several minutes to establish, even without considering magnetospheric feedback, instead of tens of seconds that simple order-of-magnitude estimates typically predict. The article is properly placed in the context of prior work. Most of the limitations and potential future developments are well described.

A point requiring further clarification is the significance of height variations of winds, conductivities, and current densities within the ionosphere for the modeling. The mathematical development essentially ignores variations of these quantities with height, treating the quantities as existing only at a single height in an infinitesimally thin layer. The numerical modeling uses winds at 110 km from the Horizontal Wind Model and height-integrated conductivities. In reality winds and conductivities vary strongly with altitude in the lower ionosphere. Height integration of Equation (6) does not yield Equation (7) as the result. Instead, Equation (3) needs to be integrated in height and then solved for E in order to get something similar to Equation (7), except that the $u \times B$ term becomes terms involving height-averaged winds weighted by the Pedersen and Hall conductivities.

We thank the reviewer for this important comment. We agree that our description of the relationship between the 3D and 2D quantities was lacking. We have kept the model as is, but done two things: 1) We have modified the statements in section 3 to avoid the indication that we integrate the Generalized Ohm's law, but instead project the 3D equation on a spherical shell. 2) We have added a new appendix to justify our use of height-integrated conductivities and current density, and to outline an approach for a more generally applicable model with better treatment of neutral winds, and possible inclusion of field-aligned conductivity. The appendix gives a more complete description of the relationship between the 3D equations and the conductance/resistance tensors in the Ohm's laws.

We include a screenshot of the new appendix below:

Appendix A: Derivation of the Thin-Sheet Conductivity and Resistivity Tensors

In Section 3 we present a two-dimensional version of the Generalized Ohm's law for a collisional ionosphere (7). The equation is a projection of the corresponding 3D equation (3) onto a spherical shell and so it is valid for an ionosphere that is truly two-dimensional. Nevertheless, as indicated by the choice of notation, we interpret $\mathbf{J}_S$ and $\Sigma_{P,H}$ as the height-integrated versions of the corresponding 3D quantities $\mathbf{j}$ and $\sigma_{P,H}$. If $\mathbf{j}$ or $\sigma_{P,H}$ vary with height, which they do in the realistic ionosphere, Equation (7) is not simply a height integral of Equation (2), so this interpretation requires more justification.

The justification is provided by using the ionospheric Ohm's law, and then solve this equation for $\mathbf{E}_S$. In the dynamo region of the ionosphere (primarily the E-region), $\mathbf{E}$ and $\mathbf{B}$ vary only slowly with altitude, whereas the conductivities vary sharply. Because of this scale separation, the altitude dependence is carried almost entirely by the conductivities, so integrating the ionospheric Ohm's law yields well-defined conductances. For generality we retain a finite field-aligned conductivity $\sigma_{\parallel}$ (though elsewhere in the paper we assume it to be infinite), and write the ionospheric Ohm's law as

$$\mathbf{j} = \sigma_P \mathbf{E}'_{\perp} + \sigma_H \hat{\mathbf{b}} \times \mathbf{E}' + \sigma_{\parallel} E_{\parallel} \hat{\mathbf{b}}. \quad (\text{A1})$$

Here, $\mathbf{E}' = \mathbf{E} + \mathbf{u} \times \mathbf{B}$ is the electric field in the neutral frame of reference. If we assume that both $\mathbf{B}$ (which is dominated by the main field, $\mathbf{B}_0$) and $\mathbf{E}$ vary little over the height of the ionosphere, we can integrate this equation to get a 3D height-integrated current:

$$\mathbf{J} = \Sigma_P \mathbf{E}_{\perp} + \Sigma_H \hat{\mathbf{b}} \times \mathbf{E} + \Sigma_{\parallel} E_{\parallel} \hat{\mathbf{b}} + \mathbf{Q}. \quad (\text{A2})$$

Here, $\Sigma_{P,H,\parallel}$ are the height-integrated $\sigma_{P,H,\parallel}$, and $\mathbf{Q} = \mathbf{Q}_P + \mathbf{Q}_H$, where

$$\mathbf{Q}_P = \int \sigma_P(r) (\mathbf{u}(r) \times \mathbf{B}) dr \quad (\text{A3})$$

$$\mathbf{Q}_H = \int \sigma_H(r) (\mathbf{u}(r) \times \mathbf{B}) \times \hat{\mathbf{b}} dr \quad (\text{A4})$$

are integrated wind terms weighted by the Pedersen and Hall conductivities (Richmond, 1995; Hatch et al., 2024).

We use that $\mathbf{E}_{\perp} = \mathbf{E} - E_{\parallel} \hat{\mathbf{b}}$ and decompose $\mathbf{J}$ in radial (subscript r) and tangential/surface components (subscript S):

$$\begin{pmatrix} J_r \\ \mathbf{J}_S \end{pmatrix} = \begin{pmatrix} \Sigma_P E_r + \Sigma_H (\hat{\mathbf{b}} \times \mathbf{E})_r + (\Sigma_{\parallel} - \Sigma_P) E_{\parallel} b_r + Q_r \\ \Sigma_P \mathbf{E}_S + \Sigma_H (\hat{\mathbf{b}} \times \mathbf{E})_S + (\Sigma_{\parallel} - \Sigma_P) E_{\parallel} \hat{\mathbf{b}}_S + \mathbf{Q}_S \end{pmatrix}. \quad (\text{A5})$$

In our 2D/thin-sheet formulation the discontinuity of the horizontal magnetic field across the ionosphere is described by the Ampère jump condition (10). This relation requires a surface current confined to the sheet: $\mathbf{J}_S$ in the equation above. Because the discontinuity only involves the horizontal magnetic field above and below, there can be no independent horizontal perturbation field inside the sheet itself. Consequently the sheet current is purely tangential: $J_r = 0$. The horizontal divergence of $\mathbf{J}_S$ is in general nonzero and is balanced by the radial current carried in the field-aligned system above. We therefore impose $J_r = 0$ in the radial component of (A5) and solve for E_r :

$$E_r = - \frac{\Sigma_H (\hat{\mathbf{b}} \times \mathbf{E})_r + (\Sigma_{\parallel} - \Sigma_P) b_r (\hat{\mathbf{b}}_S \cdot \mathbf{E}_S) + Q_r}{\Sigma_P (b_{\theta}^2 + b_{\phi}^2) + \Sigma_{\parallel} b_r^2} \quad (\text{A6})$$

720 This can be inserted in the expression for $\mathbf{J}_S$ to get an equation for the surface current that only depends on the horizontal components of the electric field. This is a version of the height-integrated ionospheric Ohm's law that explicitly states the dependence on the conductance and main magnetic field geometry through the tensor $\mathbb{A}_{\text{cond}}(\Sigma, \hat{\mathbf{b}})$:

$$\mathbf{J}_S = \mathbb{A}_{\text{cond}}(\Sigma, \hat{\mathbf{b}})\mathbf{E}_S + \mathbf{Q}_{\text{eff}}, \quad (\text{A7})$$

where

$$725 \quad \mathbf{Q}_{\text{eff}} = \mathbf{Q}_S - \frac{(\Sigma_{\parallel} - \Sigma_P)b_r\mathbf{b}_S Q_r - \Sigma_H(\hat{\mathbf{r}} \times \hat{\mathbf{b}})_S Q_r}{\Sigma_P(b_{\theta}^2 + b_{\phi}^2) + \Sigma_{\parallel}b_r^2}. \quad (\text{A8})$$

is the effective wind, a combination of the tangential/surface component of the integrated wind terms and the contribution that comes from substituting E_r in the surface component. The conductance tensor can be expressed as

$$\mathbb{A}_{\text{cond}}(\Sigma, \hat{\mathbf{b}}) = \frac{1}{\Delta} \begin{pmatrix} Kb_{\phi}^2 + M(1 - b_{\phi}^2) & (M - K)b_{\theta}b_{\phi} - \Sigma_H\Sigma_{\parallel}b_r \\ (M - K)b_{\theta}b_{\phi} + \Sigma_H\Sigma_{\parallel}b_r & Kb_{\theta}^2 + M(1 - b_{\theta}^2) \end{pmatrix}. \quad (\text{A9})$$

where

$$730 \quad \Delta \equiv \Sigma_P(b_{\theta}^2 + b_{\phi}^2) + \Sigma_{\parallel}b_r^2, \quad K \equiv \Sigma_P^2 + \Sigma_H^2, \quad M \equiv \Sigma_P\Sigma_{\parallel}. \quad (\text{A10})$$

This conductance tensor reduces to the expression given by Amm (1996); Rishbeth and Garriott (1969) for $b_{\phi} = 0$.

We can solve Equation (A7) for $\mathbf{E}_S$ to get the corresponding 2D *Generalized* Ohm's law:

$$\mathbf{E}_S = \mathbb{A}_{\text{cond}}^{-1}(\Sigma, \hat{\mathbf{b}})(\mathbf{J}_S - \mathbf{Q}_{\text{eff}}) = \mathbb{A}_{\text{res}}(\eta, \hat{\mathbf{b}})(\mathbf{J}_S - \mathbf{Q}_{\text{eff}}) \quad (\text{A11})$$

where the inverse of the conductance tensor is the resistance tensor, $\mathbb{A}_{\text{res}}(\eta, \hat{\mathbf{b}})$:

$$735 \quad \mathbb{A}_{\text{res}}(\eta, \hat{\mathbf{b}}) = \begin{pmatrix} \eta_P + (\eta_{\parallel} - \eta_P)b_{\theta}^2 & (\eta_{\parallel} - \eta_P)b_{\theta}b_{\phi} + \eta_Hb_r \\ (\eta_{\parallel} - \eta_P)b_{\theta}b_{\phi} - \eta_Hb_r & \eta_P + (\eta_{\parallel} - \eta_P)b_{\phi}^2 \end{pmatrix}, \quad (\text{A12})$$

where η_P and η_H are defined in terms of the conductances as in (8) and (9), and $\eta_{\parallel} = 1/\Sigma_{\parallel}$. We emphasize that $\eta_{P,H}$ are *not* height-integrated resistances, but quantities derived from the height-integrated conductivities.

The resistance tensor (A12) is equivalent to the matrix defined in equation (11) of O'Sullivan and Downes (2006), which is written in Cartesian coordinates, and it reduces to the matrix in equation (11) of this paper in the limit $\eta_{\parallel} = 0$. We have
740 thus shown that the expression that we get by simply projecting the 3D Generalized Ohm's law to 2D can also be derived by height-integrating the Ionospheric Ohm's law and solving for $\mathbf{E}$. The advantage of this derivation is that it quantifies the relationship between the 2D and 3D quantities. For example, it provides a consistent way to include altitude-dependent winds in the thin-sheet formulation, by calculating the height integrals in (A3) and (A4) to give $\mathbf{Q}_H$ and $\mathbf{Q}_P$.

Another point requiring further clarification is the nature of winds responsible for Sq-like geomagnetic variations. The statement "Sq currents will only form if the winds imply an interhemispheric imbalance [of winds] at conjugate points" (Line 575) is incorrect. Winds at conjugate points can be balanced and still create a curl in $\mathbf{u} \times \mathbf{B}_0$ that is balanced at conjugate

points such that no interhemispheric magnetic tension is created, only a balanced change in magnetic pressure.

Our original wording was imprecise and we did not mean to suggest that neutral winds must be imbalanced at conjugate points in order for Sq currents to arise. As the reviewer correctly notes, winds that are balanced at conjugate points can still produce a curl in $\mathbf{u} \times \mathbf{B}_0$, which drives currents without creating interhemispheric tension. What matters is not the wind itself but the resulting Lorentz force. We have therefore revised the sentence to read: "Sq currents will only form if there is an interhemispheric imbalance in Lorentz force at conjugate points, mainly arising from neutral winds."

Minor comments

1. Add j_{parallel} to the right-hand side of (3).

Done

2. Line 104: F can also be important at low altitudes, where E_{parallel} may be non-negligible.

We changed the sentence to

Since $\mathbf{F}$ in this equation only becomes important outside the E-region (Brekke, 2013), we will neglect it here.

3. In the caption for Figure 1 all the mathematical quantities should be defined.

We have simplified the figure and defined all the mathematical quantities. Below is a screenshot of the figure with the new caption




Figure 1. Cross section of the Earth, ionosphere, and inner magnetosphere (gap region) shown to scale. The ionosphere appears as a blue line but is in fact a linear color scale contour plot of a typical height profile of the Pedersen conductivity, highlighting the appropriateness of a 2D treatment. We simulate the radial component of the magnetic field $\mathbf{B}$ on the spherical surface at radius $r = R$, but the magnetic field is defined everywhere using the assumptions indicated in the figure. Above the ionosphere, we assume that the magnetic field is force-free ($\mathbf{B} \times \nabla \times \mathbf{B} = 0$) and that it is frozen-in with the plasma, which moves at velocity $\mathbf{v}$. Below the ionosphere, the magnetic field is assumed to be curl-free.

4. Lines 136-140: The reason why the radial component of the current does not appear in (7) has nothing to do with high field-aligned electron mobility, but rather to the fact that current density appears in (6) and (7) only in the form $\mathbf{j} \times \mathbf{b}$, and therefore excludes $\mathbf{j}_{\parallel}$.

Thank you for catching this. We have changed the last sentence of the paragraph to:

In this expression, $\mathbf{j}_{\parallel}$ appears only in the last term, which represents the magnetic field-aligned component of $\mathbf{j}$, which is not included in (7) because of the cross product with $\mathbf{b}$.

5. Concerning notation, $\mathbf{B}$ is the total magnetic field in (2),(3),(6),(7), but is only the perturbation field starting at Line 151. Perhaps different symbols can be used.

We agree that this is unfortunate, but it is difficult to adopt an alternative symbol for the perturbation field, since most common choices (e.g., ΔB) are already in use. To address this, we

have revised the text to make the change in notation explicit. We now write
From this point onward, we refer to the main field as $\mathbf{B}_0$, [...]

6. Line 162: Is Equation (11) meant instead of Equation (10)?

To avoid confusion we changed the sentence to

It remains to specify the components of $\Delta\mathbf{B}$, which consists of the magnetic field immediately above ($\mathbf{B}^+$) and below ($\mathbf{B}^-$) the ionosphere, according to Equation (10).

7. Lines 294-295: It is not obvious that the deformation can be neglected if $d\mathbf{B}/dt$ is non-zero.

We changed the sentence to say

We make the physical assumption that the induction electric field is small enough that this deformation is negligible

8. Lines 313-314 and Line 561: The force-free assumption applies only above the current layer, not within it. Currents can cross magnetic field lines everywhere in the current layer.

Our point was not precisely stated. It is now changed to

We note that our 2D ionosphere is likely incompatible with the physics of the equatorial electrojet: If the 2D sheet current has a non-zero divergence at the dip equator, it implies a radial current above the sheet which crosses magnetic field lines, in violation of the force-free assumption.

Stephan C. Buchert

The authors skillfully condense into the paper a relatively large amount of work and impressive progress on modeling the thermosphere-ionosphere-magnetosphere. I have only a few minor comments in parts of section 5 "Discussion"

"Sq currents will only form if the winds imply an interhemispheric imbalance [of winds] at conjugate points" (Line 575) is a bit vague about what interhemispheric (im)balance means, but essentially correct, in my opinion. It is the $\mathbf{u} \times \mathbf{B}$ term (integrated over the dynamo region in the 2D model) which' mismatch at conjugate points forms Sq. In order to balance, i.e. suppress Sq completely, the neutral wind vectors would need to compensate also for any conjugate asymmetries of the magnetic field, including asymmetric magnitudes $|\mathbf{B}_N|$ and $|\mathbf{B}_S|$. This is implied by the statement "2) if $\mathbf{u} \times \mathbf{B}_0$ at conjugate points imply different electric fields, ..." in lines 570-572. I'm not sure whether this is relevant for Art Richmond's comment on the issue in his review.

Thank you for this comment. See response to Art Richmond's comment, above. We changed our phrasing to: "Sq currents will only form if there is an interhemispheric imbalance in Lorentz force at conjugate points, mainly arising from neutral winds."

Regarding altitude dependence of the neutral wind, rather of the $\mathbf{u} \times \mathbf{B}$ term within the ionosphere (in each hemisphere), an imbalance would form intra-ionosphere current systems, i.e current loops closing within different layers. I'm not aware of any experimental evidence of such currents which is probably owing to the absence of satellite data from the lower-middle

thermosphere-ionosphere. According to Fukushima's theorem the effects would be almost undetectable both below/on the ground as well as above in low Earth orbits. A future 3D version of the B,v model presented here could demonstrate the existence of such currents/magnetic tensions between thermospheric layers in the presence of horizontal neutral wind with vertical gradients. This is meant as a contribution to the discussion, not necessarily a suggestion for additions to the paper, which might rather focus on describing what has been achieved so far.

We thank the reviewer for this interesting and insightful comment. The relationship between 3D and 2D formulations is now discussed in more detail in the new appendix. We agree that a full treatment of intra-ionospheric current loops driven by vertical wind gradients would require a dedicated 3D framework, which is beyond the scope of the present paper.

Lines 579-580, "In steady state the electric field is such that no further deformation happens, as the $\mathbf{j} \times \mathbf{B}$ force and momentum transfer due to collisions with neutrals counterbalance each other.": The $\mathbf{j} \times \mathbf{B}$ force and momentum transfer by collisions are different views of the same forcing, the former in the B,v paradigm, the latter in E,j. They do not counterbalance. $\mathbf{j} \times \mathbf{B}$ (or collisional transfer in E,j) rather is counterbalanced by the inertia of the atmosphere. Thus strictly only a quasi-steady state is reached, with generally long atmospheric time scales. Rapid changes as mentioned in lines 494-500 would so be able to excite atmospheric acoustic/gravity waves, probably of relatively low amplitudes, Parker's (1996) long paper includes an example calculation of the neutral atmosphere forcing by magnetic stress (i.e. closure of FACs). This has been studied/simulated within the E,j paradigm modeling the frictional forcing of magnetosphere ion convection, for example, one simulated effect has been described as "flywheel" (Deng et al., JGR< 1991). But an inductive treatment has advantages. I suggest to briefly mention the potential to couple dynamic models of the atmosphere with an inductive Magnetosphere-Ionosphere-Thermosphere (B,v) description in a conceptually relatively straight forward manner (i.e. propagate forward in time $du/dt = \dots + \mathbf{j} \times \mathbf{B}$ and $d\mathbf{B}/dt = -(\nabla \times \mathbf{E}(u, \mathbf{B}))$, Faraday with GOL, equation (2)). The treatment of the inductive Magnetosphere-Ionosphere-Thermosphere would so become more analogue to that of the Earth core and relatively recently also Ocean dynamos.

We agree with this point, but we have not added anything in the paper since the possibility of including fluid dynamics is discussed in Section 5.4.3, and also in the first paragraph of Section 3.

Section "Conclusions", especially paragraph lines "677-683": The high-latitude field-aligned currents are described as being prescribed and an input into the coupled Magnetosphere-Ionosphere-Thermosphere model of this work. I would suggest to mention (at this paragraph and/or elsewhere in the paper) that the neutral wind in the dynamo region is the other 2nd input to the 2D model. Here it is prescribed using the HWM by Drob et al. (2017). In reality the neutral wind varies with time. Precise and spatially sufficiently dense magnetic measurements on the ground and in space at mid-latitudes would so carry information on neutral wind differences at magnetically conjugate points.

Thank you for this comment. We agree that this point could be emphasized more. In the paragraph that is referred to, we have added a mention of given neutral wind pattern to one of the sentences:

At high latitudes, the imposed magnetic field is defined to be consistent with a prescribed pattern of field-aligned currents; at low latitudes it is calculated by assuming that the magnetic field adapts to preserve current continuity between hemispheres and matching electric potentials along magnetic field lines, given a neutral wind pattern.

We have also included mention of global circulation models in the first sentence in the next paragraph:

The model presented here can in principle be incorporated into global magnetosphere simulations and upper-atmospheric circulation models to provide a more complete description of magnetosphere–ionosphere interactions.

Global Inductive Magnetosphere-Ionosphere-Thermosphere Coupling

Karl M. Laundal^{1,2}, Andreas S. Skeidsvoll², Beatrice Popescu Braileanu², Spencer M. Hatch², Nils Olsen¹, and Heikki Vanhamäki³

¹Division of Geomagnetism, DTU Space, Technical University of Denmark

²Department of Physics and Technology, University of Bergen, Norway

³Space Physics and Astronomy Research Unit, University of Oulu, Finland

Correspondence: Karl M. Laundal (kalau@space.dtu.dk)

Abstract. The ionosphere forms the inner boundary of near-Earth space, where collisionless space plasma transitions into a partially ionized gas that interacts with the neutral atmosphere through collisions. Conventional models for magnetosphere-ionosphere (MI) coupling use an electric circuit framework, where an electric potential is calculated from the current continuity equation on a thin spherical shell that represents the ionosphere. This approach, founded in the $\mathbf{E}, \mathbf{j}$ (electric field and current density) paradigm, contrasts with the approach used to study plasmas in other regions of cosmos, where the magnetic field $\mathbf{B}$ and plasma velocity $\mathbf{v}$ are treated as fundamental variables (the $\mathbf{B}, \mathbf{v}$ paradigm). Since traditional MI coupling models also neglect induction by setting $\partial \mathbf{B} / \partial t = 0$, they omit the dynamic processes by which $\mathbf{B}$ evolves, leaving the global MI coupling process arguably poorly understood. To advance our understanding of MI coupling, we present a new global model of the 2D ionosphere that incorporates induction, with $\mathbf{B}$ as the primary variable. This model accommodates arbitrary ionospheric conductance, neutral wind patterns, and realistic main magnetic field geometries. Simulations reveal the complex nature of the induction process over a few seconds to several minutes. The induction timescales depend on the magnitudes and spatial scales of conductance, neutral wind, imposed magnetic field perturbations, and main magnetic field geometry. We simulate for the first time how low-latitude Sq currents and electric fields emerge through induction. Our model has the potential to replace existing MI coupling modules in magnetospheric simulation codes, offering both a truly global solution, and the inclusion of induction in the coupled system dynamics.

1 Introduction

The magnetosphere constitutes the vast region of space where Earth's magnetic field has a dominating influence, a comet-shaped region that extends to about $10R_E$ in the sunward direction, with a much longer tail on the nightside. Beyond the magnetosphere's outer boundary is the solar wind, which shapes it and powers most of its dynamics, including phenomena such as geomagnetic storms and auroral displays. The inner boundary of the magnetosphere is the ionosphere, where space plasma overlaps and interacts with the neutral atmosphere. Collisions between the charged particles of the plasma and neutrals in the ionosphere lead to exchanges of momentum and energy when the two fluids move relative to each other.

To our knowledge, all global simulations of the magnetosphere (Tóth et al., 2012; Zhang et al., 2019; Sorathia et al., 2020; Von Althaus et al., 2014) use the same principle for coupling to the ionosphere: Field-aligned currents (FACs) are mapped
 25 along the main magnetic field of the Earth from the inner boundary of the magnetospheric simulation domain, typically a few Earth radii (R_E , set to 6371.2 km in this work), to a 2D spherical shell that represents the ionosphere. The incident FAC is used in the current continuity equation, along with prescribed electric conductivities and neutral winds, to solve for an electric potential (Merkin and Lyon, 2010; Ridley et al., 2004; Ganse et al., 2025) which serves as the inner boundary condition for the magnetosphere simulations. Earthward of the inner boundary, the potential depends on the neutral winds and assumptions
 30 about interhemispheric coupling (Maute et al., 2021; Richmond and Maute, 2014). In existing implementations, the potential is treated independently in the two polar hemispheres, and separated from the low latitudes by a boundary condition at some fixed latitude. This circuit-based approach, relating electric fields, conductivities, and currents via the so-called Ionospheric Ohm’s law, has a long tradition in ionospheric physics (see, e.g., reviews by Richmond, 2016; Leake et al., 2014), and is derived from the momentum equations of electrons, ions, and neutrals, together with important assumptions for each of them.

In the conventional magnetosphere-ionosphere coupling schemes described above, the electric field $\mathbf{E}$ is represented as the gradient of a scalar potential, which implies that it is curl-free, and that $\partial \mathbf{B} / \partial t = 0$ according to Faraday’s law. In this work, we do not make this assumption, but we still neglect displacement currents, $\partial \mathbf{E} / \partial t = 0$. While it is true that the rotational part of $\mathbf{E}$ is often small compared to the potential part of the field, it has been observed to be significant in some cases (Madelaine et al., 2024). Importantly, the lack of induction conceals the physical processes that drive ionospheric electrodynamics. The
 40 widespread use of this approximation has led to the idea that field-aligned currents set up an electric field, or that electric fields *penetrate* from higher altitudes, and *drive* convection and currents across magnetic field lines. This concept has been challenged in a series of papers arguing that instead of treating $\mathbf{E}$ and $\mathbf{j}$ as primary (the $\mathbf{E}, \mathbf{j}$ paradigm), $\mathbf{B}$ and $\mathbf{v}$ should be seen as the primary variables, while $\mathbf{E}$ and $\mathbf{j}$ are derived (Vasyliūnas, 2001, 2012; Vasyliūnas and Song, 2005; Vasyliūnas, 2005; Parker, 1996). In the $\mathbf{B}, \mathbf{v}$ paradigm the electric field and currents are understood as natural consequences of plasma
 45 convection and magnetic field deformations propagating from the magnetosphere, emphasizing the dynamic coupling between the magnetosphere and the ionosphere rather than viewing electric fields as externally imposed drivers.

Laundal et al. (2024) recently presented a simple conceptual model, a thought experiment in which the ionosphere is flat, the main magnetic field is vertical, the neutral winds are zero, and the dynamics is resolved in a single horizontal dimension. Their purpose was to elucidate the physical process of how the magnetosphere drives ionospheric dynamics, but the restrictive
 50 simplifications make it difficult to directly apply their model in a realistic scenario. In this paper, we extend the model by treating the ionosphere as a 2D spherical shell, and allowing arbitrary main magnetic field geometry, neutral wind, and horizontal gradients in all quantities. The resulting model can, in principle, replace the magnetostatic magnetosphere-ionosphere coupling scheme in conventional global magnetospheric models. Incidentally, our $\mathbf{B}, \mathbf{v}$ -based approach can also be solved in steady state, in which case it reduces to the conventional magnetostatic approach. However, unlike conventional magnetostatic
 55 solvers, our model is truly global, with no boundaries that disconnect high- and low-latitude regions.

In Section 2 we explain the basic concept of our approach, starting from Maxwell’s equations and the Generalized Ohm’s law in 3D. In Section 3 we discuss simplifying assumptions, including the projection onto a 2D spherical shell, that reduces the

computational complexity. Special attention is given to the boundary conditions, which are defined by the magnetic field above and below the ionosphere, represented with spherical harmonics. Section 4 presents example simulation results. We discuss the results, implications, and limitations in Section 5. Section 6 concludes the paper.

2 Basic Theory of Inductive Ionospheric Dynamics

In this section we present the basic equations of the inductive magnetosphere-ionosphere coupling scheme used in this paper. Our approach is inspired by Vasyliūnas (2012) and Chapter 9.5 in Parker (2007). It is related to a number of previous simulation studies in which the equations given below are solved in various 1D and 2D geometries (Tu and Song, 2016, 2019; Dreher, 1997; Otto and Zhu, 2003; Laundal et al., 2024). Here we present the equations for a 3D ionosphere. In the next section, we present the simplified equations that model the ionosphere as a 2D spherical shell, along with the approach used to define the boundary conditions.

The induction equation, or Faraday's law, is

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E}. \quad (1)$$

To integrate this equation in time, the electric field $\mathbf{E}$ must be defined. We consider it to be given by the Generalized Ohm's law (Leake et al., 2014),

$$\mathbf{E} = \frac{\sigma_H \mathbf{j} \times \hat{\mathbf{b}} + \sigma_P \hat{\mathbf{b}} \times (\mathbf{j} \times \hat{\mathbf{b}})}{\sigma_H^2 + \sigma_P^2} - \mathbf{u} \times \mathbf{B} + \mathbf{F}, \quad (2)$$

where $\mathbf{u}$ is the neutral wind, $\hat{\mathbf{b}}$ is the unit vector in the direction of the magnetic field, $\mathbf{j}$ is the electric current density, and σ_P and σ_H are Pedersen and Hall conductivities, respectively. This equation is derived from the momentum equations for ions and electrons. The vector $\mathbf{F}$ includes contributions from inertia, pressure gradients, gravity, and the magnetic field-aligned component of $\mathbf{E}$. For a detailed derivation of this equation, see Leake et al. (2014). When $\mathbf{F} = \mathbf{0}$, the remaining terms in the momentum equations represent the Lorentz force and the momentum exchange with neutrals via collisions; and Equation (2) represents the relationship between $\mathbf{E}$ and $\mathbf{j}$ when these two forces balance. In the following, we will assume that $\mathbf{F} = \mathbf{0}$. Pressure gradients and inertia are important to consider at high altitudes but become relatively less important in the E-region, where the collisional coupling between plasma and neutrals is strongest (Brekke, 2013). Furthermore, the conductivity along the magnetic field lines can be assumed to be large, implying that any electric field in the direction of the field lines is quickly equilibrated, which implies that $\mathbf{B} \cdot \mathbf{E} = 0$. The effects of pressure gradients and gravity are generally considered to be small, although their impacts are sometimes detectable (Alken, 2016; Laundal et al., 2019).

The Generalized Ohm's law and the Ionospheric Ohm's law are fundamentally the same equation, differing only in the variable being solved for: $\mathbf{E}$ for the Generalized Ohm's law or $\mathbf{j}$ for the Ionospheric Ohm's law. For $\mathbf{F} = \mathbf{0}$ the Ionospheric Ohm's law is (e.g., Brekke, 2013)

$$\mathbf{j} = \sigma_P (\mathbf{E} + \mathbf{u} \times \mathbf{B}) + \sigma_H \hat{\mathbf{b}} \times (\mathbf{E} + \mathbf{u} \times \mathbf{B}) + \underbrace{j_{\parallel}}_{\text{parallel current}} \hat{\mathbf{b}}. \quad (3)$$

The conductivities primarily depend on plasma density and on the collision frequencies between the plasma species and neutrals (e.g., Strangeway (2012)). The collision frequencies, in turn, depend on temperature (Schunk and Nagy, 2009). The time evolution of density, momentum, and temperature are described by the continuity, momentum, and energy equations, respectively (Dreher, 1997; Otto and Zhu, 2003; Tu and Song, 2016; Laundal et al., 2024). In this paper, we assume that the density and temperature, and hence conductivity, are given, and that both ion and electron inertia can be neglected. Thus, the only equation we need to integrate in time is Faraday’s law.

We obtain a closed set of equations by replacing the current density using Ampère’s law in the quasi-static approximation,

$$\mathbf{j} = \frac{1}{\mu_0} \nabla \times \mathbf{B}, \quad (4)$$

and choosing appropriate boundary conditions for $\mathbf{B}$. Equations (1), (2), and (4) can then be combined to describe the evolution of the magnetic field as a function of the magnetic field itself. The electric field can be retrieved at any time from Equation (2) and the currents from Equation (4), but, as pointed out by Parker (1996), they do not dictate the dynamics. It is possible but computationally expensive to evolve Faraday’s law in 3D. In the next section, we derive the ionospheric induction equation for a 2D spherical shell, along with other simplifying assumptions that we use in our numerical simulation.

3 Dynamic MIT Coupling With 2D Ionosphere

In this section, we describe how we use Equations (1), (2), and (4) for a 2D spherical shell at radius $r = R$ that represents the ionosphere. Since the collisional interaction between ions and neutrals ~~peak-peaks~~ in the E-region, this region ~~will dominate in a height-integral of Equation (2) over the ionosphere~~ dominates the ionospheric response in any height-integrated description.

Since $\mathbf{F}$ in this equation only becomes important ~~at higher altitudes outside the E-region~~ (Brekke, 2013), we will neglect it here. In addition we will neglect the continuity and energy equations, which means that, as in conventional magnetosphere-ionosphere coupling schemes, the conductivities, which are functions of density and temperature (Schunk and Nagy, 2009), must be considered as given. Future iterations of the model may include all these effects as it is conceptually straightforward but computationally more expensive to evolve the fluid moments in time together with Faraday’s law (Tu and Song, 2016).

Above the 2D ionosphere, we assume ideal MHD and that the magnetic field is force-free, which means that $\mathbf{B} \times (\nabla \times \mathbf{B}) = 0$. That is, all currents are field-aligned in this region. Below the ionosphere, we assume that there are no currents, so that the magnetic field is curl-free and can be expressed as a gradient of a scalar potential. These assumptions lead to a 3D description of the magnetic field above and below the ionosphere, described in more detail later in this section. The geometry is illustrated in Figure 1. In this figure, which is to scale, the ionosphere looks like a thin blue line at about 110 km altitude, but is actually a contour plot of a typical Pedersen conductivity. Here, the Pedersen conductivity is based on measurements taken at a single geographic location using the EISCAT incoherent scatter radar on Svalbard, but the values are plotted across all latitudes to illustrate the radial dependence relative to a 2D cross-section of the Earth. This shows that the 2D assumption is reasonable on large scales.

From magnetic flux conservation, the radial component of the magnetic field must be continuous across the ionosphere, which means that the radial component B_r must be the same as the ionosphere (located at $r = R$) is approached from above or




Figure 1. Cross section of the Earth, ionosphere, and inner magnetosphere (gap region) shown to scale. The ionosphere appears as a blue line but is in fact a linear color scale contour plot of a typical height profile of the Pedersen conductivity, highlighting the appropriateness of a 2D treatment. We simulate B_r —the radial component of the magnetic field $\mathbf{B}$ on the spherical surface at radius $r = R$, but the magnetic field is defined everywhere using the assumptions indicated in the figure. Above the ionosphere, we assume that the magnetic field is force-free ($\mathbf{B} \times \nabla \times \mathbf{B} = 0$) and that it is frozen-in with the plasma, which moves at velocity $\mathbf{v}$. Below the ionosphere, the magnetic field is assumed to be curl-free.

from below. In our 2D model it will only be necessary to consider the radial component of Faraday's law,

$$\frac{\partial B_r}{\partial t} = -(\nabla \times \mathbf{E}) \cdot \hat{\mathbf{r}}. \quad (5)$$

The horizontal part of the electric field $\mathbf{E}_S$ can be expressed by projecting Equation (2) on the horizontal plane by making the cross product with $\hat{\mathbf{r}}$ from the left and right: $\mathbf{E}_S = \hat{\mathbf{r}} \times \mathbf{E} \times \hat{\mathbf{r}}$. The symmetry of the operation allows us to omit brackets, as
 125 the result is independent of the order of the cross products. Neglecting $\mathbf{F}$ we get

$$\mathbf{E}_S = \hat{\mathbf{r}} \times \left[\frac{\sigma_H \mathbf{j} \times \hat{\mathbf{b}} + \sigma_P \hat{\mathbf{b}} \times (\mathbf{j} \times \hat{\mathbf{b}})}{\sigma_P^2 + \sigma_H^2} - \mathbf{u} \times \mathbf{B} \right] \times \hat{\mathbf{r}}. \quad (6)$$

~~If we consider the radial dependence of the ionospheric conductivities to be given by~~ To represent the ionosphere as a thin sheet at $r = R$, we treat its radial dependence as a delta function centered at $r = R$, it is trivial to integrate this equation over the height of the ionosphere. To signify that we are working with height-integrated quantities, we simply replace the conductivities $\sigma_{H,P}$ with the conductances $\Sigma_{H,P}$, and introduce the height-integrated horizontal we introduce the corresponding sheet current density $\mathbf{J}_S$. A detailed derivation of this thin-sheet formulation from the ionospheric Ohm's law is given in Appendix A.
 130 by height-integrated quantities, we simply replace the conductivities $\sigma_{H,P}$ with the conductances $\Sigma_{H,P}$, and introduce the height-integrated horizontal we introduce the corresponding sheet current density $\mathbf{J}_S$. A detailed derivation of this thin-sheet formulation from the ionospheric Ohm's law is given in Appendix A.

$$\begin{aligned} \mathbf{E}_S &= \hat{\mathbf{r}} \times \left[\frac{\Sigma_H \mathbf{J}_S \times \hat{\mathbf{b}} + \Sigma_P \hat{\mathbf{b}} \times (\mathbf{J}_S \times \hat{\mathbf{b}})}{\Sigma_P^2 + \Sigma_H^2} - \mathbf{u} \times \mathbf{B} \right] \times \hat{\mathbf{r}} \\ &= \hat{\mathbf{r}} \times [\eta_H \mathbf{J}_S \times \hat{\mathbf{b}} + \eta_P \hat{\mathbf{b}} \times (\mathbf{J}_S \times \hat{\mathbf{b}}) - \mathbf{u} \times B_r \hat{\mathbf{r}}] \times \hat{\mathbf{r}}. \end{aligned} \quad (7)$$

135 In the last line, we introduced the height-integrated *resistances*

$$\eta_P = \Sigma_P / (\Sigma_H^2 + \Sigma_P^2) \quad (8)$$

$$\eta_H = \Sigma_H / (\Sigma_H^2 + \Sigma_P^2). \quad (9)$$

The radial component of the current does not appear in our 2D projection of $\mathbf{E}$. This can be understood by expressing the 3D current system in our model as $\mathbf{j} = \delta(r - R) \mathbf{J}_S + H(r - R) j_{\parallel} \hat{\mathbf{b}}$, where δ is the Dirac delta function and H is the Heaviside step function (1 for $r \geq R$ and 0 for $r < R$). In this expression, j_r appears only in the last term, which represents the magnetic field-aligned component of $\mathbf{j}$, which we neglect due to the high field-aligned electron mobility throughout the ionosphere (Brekke, 2013) is not included in (7) because of the cross product with $\hat{\mathbf{b}}$.
 140 step function (1 for $r \geq R$ and 0 for $r < R$). In this expression, j_r appears only in the last term, which represents the magnetic field-aligned component of $\mathbf{j}$, which we neglect due to the high field-aligned electron mobility throughout the ionosphere (Brekke, 2013) is not included in (7) because of the cross product with $\hat{\mathbf{b}}$.

To replace the sheet current density, we use the jump condition for the magnetic field across the spherical shell sheet current (or 2D Ampère's law),

$$145 \quad \mathbf{J}_S = \frac{1}{\mu_0} \hat{\mathbf{r}} \times \Delta \mathbf{B}, \quad \Delta \mathbf{B} = \mathbf{B}^+ - \mathbf{B}^-, \quad (10)$$

where the superscript "+" signifies the limit of the magnetic field as r approaches R from above, and superscript "-" signifies the limit from below. The cross product with $\hat{\mathbf{r}}$ implies that only the horizontal components of the magnetic field are involved in this equation.

Equations (5), (7), and (10) form, with the boundary conditions and input discussed below, a closed set of equations, in which B_r evolves as a function of the magnetic field itself, similar to the 3D equations presented in Section 2. In combining these equations we introduce another simplifying and very useful assumption: That the perturbation magnetic field is so small that it does not significantly change the geometry of the magnetic field. This corresponds to a linearization of the equations. ~~We~~ From this point onward, we refer to the main field as $\mathbf{B}_0$, with unit vector components b_r, b_θ , and b_ϕ , and perturbations as $\mathbf{B}$ with components B_r, B_θ , and B_ϕ . Equation (5) can then be written as

$$\frac{\partial B_r}{\partial t} = \frac{1}{r \sin \theta} \left(\frac{\partial}{\partial \phi} - \frac{\partial}{\partial \theta} (\sin \theta \cdot) \right) \left[\frac{1}{\mu_0} \begin{pmatrix} \eta_P(b_\phi^2 + b_r^2) & -\eta_P b_\theta b_\phi + \eta_H b_r \\ -\eta_P b_\theta b_\phi - \eta_H b_r & \eta_P(b_\theta^2 + b_r^2) \end{pmatrix} \begin{pmatrix} -\Delta B_\phi \\ \Delta B_\theta \end{pmatrix} - B_0 \begin{pmatrix} -b_\phi u_r + b_r u_\phi \\ -b_r u_\theta + b_\theta u_r \end{pmatrix} \right], \quad (11)$$

where we have used Equations (7) – (10) to express the evolution of B_r in terms of the magnetic field, resistances, magnetic field geometry, and neutral wind. In the neutral wind term, we neglect the magnetic field perturbation since its contribution is very small compared to the main field contribution. This is the equation that we integrate in time to model the ionospheric dynamics. If needed, the horizontal electric field $\mathbf{E}_S$ can be retrieved by evaluating the expression in square brackets (equivalent to Equation (7)). The top row of the expression in square brackets refers to the θ component and the bottom row refers to the ϕ component.

It remains to specify the components of $\Delta \mathbf{B}$, which consists of the magnetic field immediately above ($\mathbf{B}^+$) and below ($\mathbf{B}^-$) the ionosphere ~~to satisfy the boundary condition in,~~ according to Equation (10). Below the ionosphere, this is straightforward: Since there are no currents there, the magnetic field is a potential field, and its potential satisfies Laplace's equation. To determine the solution, we integrate Equation (5) in time to obtain $B_r(R, \theta, \phi)$, which then serves as a boundary condition for the Laplace equation. We return to this in more detail in Section 3.1. Above the ionosphere, the situation is more complex. In addition to an *induced* part similar to the one below, there is also an *imposed* magnetic field. The imposed field is determined by solving the steady-state ideal MHD equations under the assumption of a force-free magnetic field, with boundary conditions set by specified field-aligned currents at high latitudes and constraints from interhemispheric coupling along magnetic field lines at low latitudes. This is discussed in detail in Section 3.2. We thus decompose the magnetic field into *induced* and *imposed* parts. The imposed field is zero below the ionosphere, while both parts contribute above it. Each part can be described in terms of different current systems: The induced part corresponds to a horizontal, divergence-free sheet current at $r = R$, referred to as the equivalent current. The imposed part, on the other hand, is associated with field-aligned currents. These relationships are discussed in more detail below.

In the above equations, we need only the horizontal component of the electric field $\mathbf{E}_S$. At $r \geq R$, the vertical component of the electric field can be retrieved from the assumption that the parallel component of $\mathbf{E}$ is zero:

$$\mathbf{E} \cdot \mathbf{B}_0 = 0 \Rightarrow E_r = -(E_\theta b_\theta + E_\phi b_\phi)/b_r. \quad (12)$$

Likewise, the sheet current at $r = R$ (Equation (10)) can be connected to the full 3D current density at $r > R$ by first using the
180 current continuity equation to determine $j_r = -\nabla_S \cdot \mathbf{J}_S$, with ∇_S the horizontal (surface) gradient defined as

$$\nabla_S = \nabla - \hat{\mathbf{r}} \frac{\partial}{\partial r}, \quad (13)$$

and then the force-free assumption to reconstruct the full current vector at $r > R$:

$$\mathbf{j} \times \mathbf{B}_0 = \mathbf{0} \Rightarrow \mathbf{j} = j_r / b_r \hat{\mathbf{b}}. \quad (14)$$

In the next subsections we define the induced and imposed parts of the magnetic field, together with their corresponding cur-
185 rents, using spherical harmonics. We also describe how this decomposition is related to the more conventional decomposition into poloidal and toroidal parts.

3.1 The Induced Magnetic Field

Below the ionosphere $\mathbf{j} = 0$, which means that the magnetic field is curl-free and can be expressed as $\mathbf{B}(r < R) = -\nabla V$,
where the magnetic potential V obeys the Laplace equation $\nabla^2 V = 0$. The solution can be expanded in spherical harmonics
190 (see e.g., Sabaka et al. (2013); Olsen (1997))

$$\mathbf{B}_{\text{ind}}^e = -\nabla V^e = -\nabla \left(R \sum_{n,m} \left(\frac{r}{R} \right)^n P_n^m(\theta) [q_n^m \cos(m\phi) + s_n^m \sin(m\phi)] \right), \quad (15)$$

where the superscript e signifies that the magnetic field corresponds to a current that is *external* with respect to the observation
radius r . Above the ionosphere, the situation is more complicated, because there the magnetic field is a superposition of
induced and imposed parts. The imposed part is discussed in detail below. The induced part of the magnetic field at $r > R$ is,
195 by definition, the part that satisfies the Laplace equation. The solution is

$$\mathbf{B}_{\text{ind}}^i = -\nabla V^i = -\nabla \left(R \sum_{n,m} \left(\frac{R}{r} \right)^{n+1} P_n^m(\theta) [g_n^m \cos(m\phi) + h_n^m \sin(m\phi)] \right), \quad (16)$$

where the superscript i signifies that the field corresponds to an *internal* current. For B_r , which defines the induced part of the
magnetic field, to be continuous across $r = R$ we must have that

$$\lim_{r \rightarrow R^-} \frac{\partial}{\partial r} V^e = \lim_{r \rightarrow R^+} \frac{\partial}{\partial r} V^i \quad (17)$$

200 which leads to the following relationships between the coefficients q_n^m, s_n^m and g_n^m, h_n^m (Sabaka et al., 2013)

$$nq_n^m = -(n+1)g_n^m, \quad ns_n^m = -(n+1)h_n^m. \quad (18)$$

These relationships are enforced by defining the coefficients k_n^m and l_n^m ,

$$\{k_n^m, l_n^m\} \equiv -\frac{1}{n+1} \{q_n^m, s_n^m\} = \frac{1}{n} \{g_n^m, h_n^m\}, \quad (19)$$

with these forms chosen to simplify subsequent mathematical relationships. Inserting Equation (19) into Equations (15) and
 205 (16) we obtain that the total induced magnetic field $\mathbf{B}_{\text{ind}}$ can be written as $\mathbf{B}_{\text{ind}} = \nabla V$, where

$$V = \begin{cases} -R \sum_{n,m} (n+1) \left(\frac{r}{R}\right)^n P_n^m(\theta) [k_n^m \cos(m\phi) + l_n^m \sin(m\phi)], & r < R, \\ R \sum_{n,m} n \left(\frac{R}{r}\right)^{n+1} P_n^m(\theta) [k_n^m \cos(m\phi) + l_n^m \sin(m\phi)], & r > R. \end{cases} \quad (20)$$

Note that V is undefined at $r = R$, but Equation (17) implies that its radial derivative is defined also at this point. In other words, the radial component B_r of the magnetic field at $r = R$ can be obtained through

$$B_r(R, \theta, \phi) = \sum_{n,m} n(n+1) P_n^m(\theta) [k_n^m \cos(m\phi) + l_n^m \sin(m\phi)]. \quad (21)$$

210 Equations (15) – (18) describe a one-to-one relationship between B_r and the induced part of the magnetic field on both sides of the ionosphere. Given an initial condition for B_r , the coefficients k_n^m and l_n^m can be determined by integrating Equation (11) and inverting the spherical harmonic representation in Equation (21). Once these coefficients are known, the magnetic field immediately below the ionosphere ($\mathbf{B}^-$) and the induced part of the magnetic field immediately above the ionosphere ($\mathbf{B}_{\text{ind}}^+$) can be reconstructed using Equation (20). The imposed part of the magnetic field immediately above the ionosphere is yet to
 215 be defined.

The horizontal sheet current at $r = R$ that is equivalent with $\mathbf{B}_{\text{ind}}$ can be found with Ampère's law, Equation (10). Inserting the magnetic field we get the equivalent current, which can be expressed as

$$\mathbf{J}_{\text{eq}} = \hat{\mathbf{r}} \times \nabla_S \Psi, \quad (22)$$

where the equivalent current function Ψ can be identified from $\mathbf{J}_{\text{eq}} = \frac{1}{\mu_0} \hat{\mathbf{r}} \times \Delta \mathbf{B}_{\text{ind}}$, giving (see, e.g., Laundal et al. (2018))

$$220 \quad \Psi = -\frac{R}{\mu_0} \sum_{n,m} (2n+1) P_n^m(\theta) [k_n^m \cos(m\phi) + l_n^m \sin(m\phi)]. \quad (23)$$

The equivalent current is important because it can be derived from ground magnetometer measurements of magnetic field perturbations (Kamide et al., 1981; Untiedt and Baumjohann, 1993; Laundal et al., 2015). If the main magnetic field is assumed to be radial ($b_\theta = b_\phi = 0$), the equivalent current equals the divergence-free part of $\mathbf{J}_S$ (Vasyliūnas, 2007).

3.2 The Imposed Magnetic Field

225 While the induced magnetic field has a one-to-one relationship with the equivalent current, the imposed magnetic field is a force-free magnetic field confined to $r > R$ and has a one-to-one relationship to the field-aligned currents. As in the conventional MI coupling approach, we make the simplifying assumption that field-aligned currents map instantaneously between the magnetosphere and the upper boundary of the ionosphere, through what Merkin and Lyon (2010) term the *gap region*. At low latitudes, we assume that field-aligned currents instantaneously connect the hemispheres along magnetic field lines such that
 230 the electric field matches at conjugate points, in keeping with the assumption of a force-free magnetic field and ideal MHD, $\mathbf{E} + \mathbf{v} \times \mathbf{B} = 0$, above the ionosphere (Hesse et al., 1997). An equivalent statement is that the Alfvén speed is so high that any

$\mathbf{j} \times \mathbf{B}$ force is immediately removed, or shifted to the magnetospheric and ionospheric boundaries, so that the magnetic field in the gap region at all times is relaxed in a force-free configuration. This is of course a simplification, and a physical description of the Alfvén wave propagation through the gap region has been studied by e.g. Lotko (2004) and Wright (1996).

235 As for any divergence-free vector field we can write the imposed magnetic field in terms of poloidal and toroidal parts (Sabaka et al., 2013; Olsen, 1997; Backus, 1986),

$$\mathbf{B}_{\text{imp}} = \mathbf{P} + \mathbf{T}, \quad (24)$$

where the toroidal part $\mathbf{T} = \mathbf{r} \times \nabla T$ is related to the radial current density and the poloidal part $\mathbf{P}$ is related to the horizontal part of the field-aligned current density. The toroidal scalar T can be expanded in terms of spherical harmonics (e.g., Olsen
240 (1997); Laundal et al. (2016); Fillion et al. (2023)):

$$T = \begin{cases} \sum_{n,m} P_n^m(\theta) \left(\frac{R}{r}\right) [\psi_n^m \cos(m\phi) + \xi_n^m \sin(m\phi)], & r > R, \\ 0, & r < R \end{cases} \quad (25)$$

and the associated radial current density immediately above the ionosphere is:

$$j_r = -\frac{1}{R\mu_0} \sum_{n,m} n(n+1) P_n^m(\theta) [\psi_n^m \cos(m\phi) + \xi_n^m \sin(m\phi)]. \quad (26)$$

Thus, if we know $j_{\parallel}$ globally, we can find $j_r = b_r j_{\parallel}$ and ψ_n^m and ξ_n^m , and hence T .

245 There are two aspects that make it somewhat complicated to calculate $\mathbf{B}_{\text{imp}}$: 1) In MI coupling applications $j_{\parallel}$ is given at high latitudes, but not at low latitudes. The global pattern of $j_{\parallel}$, and thus the coefficients ψ_n^m and ξ_n^m , must instead be specified through a combination of prescribed current patterns at high latitudes and the set of interhemispheric coupling constraints at low latitudes mentioned previously. We return to this in detail in Section 3.2.1. 2) The poloidal magnetic field $\mathbf{P}$ is a function of the horizontal part of the field-aligned current density in the entire volume at $r > R$, even if we only evaluate it immediately
250 above R . In our model, this field encompasses the contribution from the horizontal part of the field-aligned currents at $r > R$ and a divergence-free horizontal shielding current in the ionosphere that ensures that $\mathbf{P}$ is zero below the ionosphere. ~~The shielding current ensures that the magnetic field of FACs is initially contained~~ This shielding current is constructed as part of the imposed system in our decomposition and modifies the poloidal component of the imposed magnetic field above the ionosphere, and that it only penetrates the ionosphere via induction so that its radial component is zero at the ionosphere boundary.
255 boundary. In Section 3.2.2 we describe how the shielding current can be calculated from the coefficients ψ_n^m and ξ_n^m .

Before proceeding, we clarify a potential source of confusion in our terminology. We assume that the imposed magnetic field is contained above the ionosphere and describe induction as the process that *allows* magnetic field disturbances to penetrate through the ionosphere. An alternative and perhaps more intuitive interpretation is to view induction as the mechanism that *prevents* the external magnetic field from directly penetrating the ionosphere. In conventional MI coupling approaches, where
260 induction is ignored, the magnetic field generated by currents in and above the ionosphere would be immediately observable at ground level.

It is also important to clarify our usage of the term *induced* field. While the term “inductive fields” sometimes refers to the transient quantities $\nabla \times \mathbf{E}$ or $\partial \mathbf{B} / \partial t$ associated with the induction process, we use “induced” to denote the resultant magnetic field produced by these transient processes, a field that can persist in a steady state, effectively representing the integral of inductive fields.

3.2.1 Specifying the Imposed Magnetic Field

As discussed above, the imposed magnetic field has a one-to-one relationship with the field-aligned currents. At high latitudes, the field-aligned current density is assumed to be given. At low latitudes, we make two assumptions about the field-aligned current density:

1. Since the magnetic field is force-free for $r > R$, there can be no currents perpendicular to the magnetic field in this region. Current continuity therefore implies that any current that leaves one hemisphere must enter the other at the conjugate footpoint. Because the main magnetic field strength is not necessarily equal at the two footpoints, the current density can differ between them. The preserved quantity, as defined by Richmond (1995), is J_{e3} (his notation, Equation 4.14), which represents $j_{\parallel}$ divided by a geometric factor that varies along the main magnetic field in proportion to its field strength. Thus, we require that $J_{e3}(\lambda_m, \phi_m) = J_{e3}(-\lambda_m, \phi_m)$, where (λ_m, ϕ_m) are the Modified Apex magnetic latitude and longitude, respectively, which are constant along main magnetic field lines (Richmond, 1995). This constraint can be expressed in terms of ψ_n^m and ξ_n^m using Equation (26).
2. Ideal MHD applies at $r > R$ so that $\mathbf{E}$ maps along field lines between hemispheres. We seek an imposed magnetic field that – given the neutral wind, ionospheric conductivity, main magnetic field geometry, and the induced magnetic field – produces an electric field that maps along the main magnetic field lines to the conjugate points. In practice, we transform Equation (7) into a form describing the Modified Apex electric field components. These components, which Richmond (1995) denotes by E_{d1} and E_{d2} , are assumed to be equal along magnetic field lines and thus also at conjugate points (Richmond, 1995; Laundal and Richmond, 2017). For a potential field, this assumption is obviously true according to Equations (4.8) and (4.9) in Richmond (1995). However, we use it for an electric field that includes a rotational part, which requires further justification. Consider a Faraday loop, a closed line integral of $\mathbf{E}$ that includes a line segment $\Delta \mathbf{l}_N$ connecting two points $(\lambda_m^{1,2}, \phi_m^{1,2})$ horizontally in the ionosphere, which is further connected along paths that follow $\mathbf{B}_0$ to another horizontal segment in the opposite hemisphere, $\Delta \mathbf{l}_S$, connected by the points $(-\lambda_m^{1,2}, \phi_m^{1,2})$. Faraday’s law gives (with $\mathbf{E} \cdot \mathbf{B}_0 = 0$),

$$\mathbf{E}_N \cdot \Delta \mathbf{l}_N + \mathbf{E}_S \cdot \Delta \mathbf{l}_S = - \int_A \frac{\partial \mathbf{B}}{\partial t} \cdot d\mathbf{n} \quad (27)$$

where $d\mathbf{n}$ is a vector that points perpendicular to an integration surface A enclosed by the described path. Now we make a small adjustment in these segments so that they, and the surface, move with the rate at which the surface deforms because of induction, to make the right-hand side zero. However this requires an adjustment to the left hand side to

transform into the new reference frame.

$$\begin{aligned}
 & (\mathbf{E}_N + \delta \mathbf{v}_n \times \mathbf{B}_0) \cdot \Delta \mathbf{l}_N + (\mathbf{E}_S + \delta \mathbf{v}_S \times \mathbf{B}_0) \cdot \Delta \mathbf{l}_S = 0 \\
 295 \quad & \Rightarrow \mathbf{E}_N \cdot \Delta \mathbf{l}_N + \mathbf{E}_S \cdot \Delta \mathbf{l}_S = -\delta \mathbf{E}_N \cdot \Delta \mathbf{l}_N - \delta \mathbf{E}_S \cdot \Delta \mathbf{l}_S,
 \end{aligned} \tag{28}$$

where $\delta \mathbf{v}$ is the velocity that describes the deformation of the surface A in the ionosphere and $\delta \mathbf{E}$ is the corresponding electric field. The terms on the right-hand side were introduced to make the surface integral zero, and therefore correspond to the deformation of the total magnetic field by the induction. We make the physical assumption that the induction electric field is small enough that this deformation is negligible, reducing the equation to

$$300 \quad \mathbf{E}_N \cdot \Delta \mathbf{l}_N + \mathbf{E}_S \cdot \Delta \mathbf{l}_S = 0, \tag{29}$$

which is the same relation that is satisfied by potential electric fields. Under this assumption, the total electric field $\mathbf{E}$ thus maps between hemispheres as if it was a potential, allowing us to use the equations by Richmond (1995) to perform the mapping.

These constraints, involving the parallel current density and two components of the electric field, may seem to overdetermine the system, as we have only two unknowns: the value of the toroidal scalar T , which describes the imposed magnetic field, in each of the two hemispheres. However, Hesse et al. (1997) show that mapping the vector electric field is equivalent to mapping a single scalar, even when induction is included, though this scalar is not necessarily equal to the electric potential.

In summary, we seek coefficients (ψ_n^m, ξ_n^m) of a magnetic field that is simultaneously representative of 1) the prescribed $j_{\parallel}$ at high latitudes and 2) low-latitude interhemispheric currents and an electric field that maps between hemispheres. We describe the details of the numerical implementation of this in C.

The interhemispheric constraints are consistent with the idea by Buchert (2020) about the Sq currents being driven by differences in the neutral winds at conjugate points. We show below that the interhemispheric constraints, together with a neutral wind pattern from the empirical model by Drob et al. (2015) are sufficient to develop Sq currents. To our knowledge, this is the first inductive explanation of Sq currents. A detailed discussion of this alternative view of the mechanism for Sq current production is given in Section 5.2.

We note that our 2D ionosphere is likely incompatible with the physics of the equatorial electrojet: ~~At the dip equator, the 2D sheet current has a non-zero divergence at the dip equator, it implies a radial current above the sheet which crosses magnetic field lines are horizontal at altitudes where plasma-neutral collisions are important. Therefore, currents can cross magnetic field lines, which is,~~ in violation of ~~our~~ the force-free assumption. We discuss the limitations of our 2D approach in more detail in Section 5.

3.2.2 The Poloidal Part of the Imposed Magnetic Field

When the field-aligned currents have a horizontal component, there will be an associated poloidal magnetic field. Unlike the simple relationship between the toroidal magnetic field and the radial current density, the poloidal magnetic field is determined by a volume integral of the horizontal field-aligned current density (Backus, 1986; Engels and Olsen, 1998; Olsen, 1997).

Engels and Olsen (1998) presented a method for calculating the poloidal magnetic field of field-aligned currents at some given radius. Their method yields a representation in terms of magnetic potentials for internal and external sources, similar to Equations (16) and (15). Here we use their technique for $r \leq R$, where all field-aligned currents are external. We get a poloidal magnetic field of the field-aligned currents that can be written in terms of a potential with external origin, $\mathbf{P}_0 = -\nabla P_0$, where

$$P_0 = R \sum_{n,m} \left(\frac{r}{R}\right)^n P_n^m(\theta) [q_n^m \cos(m\phi) + s_n^m \sin(m\phi)], \quad (30)$$

similar to Equation (15). Since the field-aligned currents associated with P_0 are volumetric, the poloidal field is continuous and can therefore be evaluated immediately above the ionosphere, where it equals its value at $r = R$. We will return to how the coefficients q_n^m and s_n^m are related to ψ_n^m and ξ_n^m shortly, but first we introduce the shielding effect.

Since the imposed magnetic field does not penetrate the ionosphere immediately, there must be an associated divergence-free sheet current on $r = R$ that exactly negates the radial magnetic field implied by P_0 below the ionosphere. The magnetic field of such a divergence-free current can be written as $\mathbf{P}_{\text{shield}} = -\nabla P_{\text{shield}}$, where P_{shield} is a magnetic potential for a source at $r = R$. This magnetic field can be represented with an expansion as in Equation (20),

$$P_{\text{shield}} = \begin{cases} -R \sum_{n,m} (n+1) \left(\frac{r}{R}\right)^n P_n^m(\theta) [\kappa_n^m \cos(m\phi) + \iota_n^m \sin(m\phi)], & r < R, \\ R \sum_{n,m} n \left(\frac{R}{r}\right)^{n+1} P_n^m(\theta) [\kappa_n^m \cos(m\phi) + \iota_n^m \sin(m\phi)], & r > R. \end{cases} \quad (31)$$

This magnetic potential has a continuous radial derivative, but its horizontal gradient is in general discontinuous across $r = R$. The shielding current ensures that $B_r^{\text{imp}} = -\partial P_0 / \partial r - \partial P_{\text{shield}} / \partial r = 0$ for all $r < R$. Equations (30) and (31) then imply that

$$\{q_n^m, s_n^m\} = (n+1) \{\kappa_n^m, \iota_n^m\}. \quad (32)$$

Inserting this equation for the coefficients into Equation (30) and combining the result with Equation (31) gives a total magnetic potential $P = P_0 + P_{\text{shield}}$ for the horizontal magnetic field immediately above the ionosphere:

$$P(r = R^+, \theta, \phi) = R \sum_{n,m} (2n+1) P_n^m(\theta) [\kappa_n^m \cos(m\phi) + \iota_n^m \sin(m\phi)]. \quad (33)$$

The horizontal components of $-\nabla P$ define the poloidal part of the imposed magnetic field $\mathbf{P}$ in Equation (24).

The question of how to relate the coefficients κ_n^m and ι_n^m to ψ_n^m and ξ_n^m remains. Ultimately, each κ_n^m and ι_n^m is a linear combination of the coefficients ψ_n^m and ξ_n^m . We use the approach by Engels and Olsen (1998) to find this linear combination. Their approach is essentially to sum the magnetic field of the divergence-free part of the horizontal current density at each r . This gives the following equation for the coefficients κ_n^m, ι_n^m , which corresponds to a Biot-Savart integral:

$$\{\kappa_n^m, \iota_n^m\} = -\frac{1}{2n+1} \frac{\mu_0}{R} \int_R^\infty \{q_n^m, s_n^m\} \left(\frac{R}{r'}\right)^{n-1} dr', \quad (34)$$

where $q_n^m(r)$ and $s_n^m(r)$ are spherical harmonic coefficients that describe the divergence-free part of the horizontal components of the current density at radius $r > R$. Note that we need to scale this expression with $-(n+1)$ to obtain the expressions

in Engels and Olsen (1998), due to our scaling of the external magnetic potentials. The coefficients $q_n^m(r)$ and $s_n^m(r)$ are the coefficients of a surface spherical harmonic expansion of an equivalent current function $\Psi'(r)$:

$$355 \quad \mathbf{j}_h(r) = \hat{\mathbf{r}} \times \nabla_S \Psi'(r) \quad (35)$$

To find the coefficients $q_n^m(r)$ and $s_n^m(r)$ we in principle need to do a spherical harmonic analysis at all $r > R$. The horizontal current density at r can be found by mapping $j_{\parallel}$ from R to r , again assuming that the currents are weak, so that $\mathbf{B}_0$ can be used to specify the field geometry. We get

$$360 \quad \begin{aligned} \mathbf{j}(r, \theta_m, \phi_m) &= j_{\parallel}(R, \theta_m, \phi_m) \frac{\mathbf{B}_0(r', \theta_m, \phi_m)}{B_0(r', \theta_m, \phi_m)} \frac{B_0(r', \theta_m, \phi_m)}{B_0(R, \theta_m, \phi_m)} \\ &= j_{\parallel}(R, \theta_m, \phi_m) \frac{\mathbf{B}_0(r', \theta_m, \phi_m)}{B_0(R, \theta_m, \phi_m)}, \end{aligned} \quad (36)$$

where θ_m and ϕ_m refer to coordinates that map along magnetic field lines such as Modified Apex coordinates (Richmond, 1995).

The horizontal components of Equation (36) relates $\mathbf{j}_h(r')$ and $j_{\parallel}(r = R)$, which in turn is described by the coefficients ψ_n^m and ξ_n^m . This can be used to relate q_n^m and s_n^m to ψ_n^m and ξ_n^m , and Equation (34) provides the connection to κ_n^m and l_n^m . The numerical implementation of this is explained in detail in C.

The end result of this is that all the terms in Equation (24), that describes $\mathbf{B}_{\text{imp}}$, are given by ψ_n^m and ξ_n^m .

3.3 Summary of the Magnetic Field Description

In summary, we have a decomposition of the full magnetic field that is

$$\mathbf{B} = -\nabla V^e(k_n^m, l_n^m) - \nabla V^i(k_n^m, l_n^m) - \nabla P(\psi_n^m, \xi_n^m) + \mathbf{r} \times \nabla T(\psi_n^m, \xi_n^m), \quad (37)$$

370 where the two sets of coefficients refer to imposed and induced parts, rather than the more conventional decomposition in terms of poloidal and toroidal parts (e.g., Tamao, 1986). Here, the toroidal magnetic field is part of the imposed magnetic field, while the poloidal magnetic field (the potential fields) is partly in the imposed and partly in the induced parts of the magnetic field. In principle, this equation is valid everywhere, but our description of P in the previous section is only valid at $r < R$, where all FACs and the shielding current are external, and immediately above the ionosphere, where the shielding current is internal and the FACs are external. It could easily be generalized by including internal sources, using Equations (14)–(16) of Engels and Olsen (1998).

Table 1 summarizes the decomposition of the magnetic field into induced and imposed parts, their relationships to currents, and to the poloidal-toroidal decomposition.

380 The time evolution of Faraday's law, Equation (11), can be represented by the temporal integration of the induced magnetic field coefficients k_n^m, l_n^m , along with the application of boundary conditions involving the imposed magnetic field coefficients ψ_n^m and ξ_n^m . This evolution is governed by the resistances η_P and η_H , the magnetic field geometry $\mathbf{B}_0$, the neutral winds $\mathbf{u}$, and the high-latitude FAC density $j_{\parallel}$. The electric field can be calculated at any time by evaluating the expression in square

Table 1. A summary of the magnetic field decomposition used in this work.

$\mathbf{B} =$	$-\nabla V^e$	$-\nabla V^i$	$-\nabla P$	$+\mathbf{r} \times \nabla T$
	Potential magnetic field for external source	Potential magnetic field for internal source	Potential magnetic field of horizontal component of FACs and its divergence-free horizontal shielding current at $r = R$	Magnetic field of radial component of FACs and its curl-free horizontal closure current at $r = R$
	Induced magnetic field, spherical harmonic coefficients k_n^m, l_n^m Related to the equivalent current Ψ		Imposed magnetic field, spherical harmonic coefficients ψ_n^m, ξ_n^m Related to the field-aligned current density $j_{\parallel}$	
	Poloidal magnetic field			Toroidal magnetic field
	Zero at $r > R$		Zero at $r < R$	

brackets. A steady-state solution can be obtained by setting $\partial B_r / \partial t = 0$ and solving for k_n^m and l_n^m . We present a numerical scheme in C, and example results in the following section.

385 4 Example Simulations

In this section, we present example simulations to illustrate the impact of incorporating induction in magnetosphere-ionosphere coupling. For simplicity, we hold the spatial patterns of conductance, wind, and high-latitude field-aligned currents constant. We explore two cases:

- 390 1. Simulations where the high-latitude field-aligned currents are set to zero, while the conductances or wind patterns (either would produce the same result) increase from zero as a step function. This demonstrates how Sq currents emerge via induction.
2. From the steady state established in case 1, we introduce a step function increase in the high-latitude field-aligned currents. The resulting evolution of the polar magnetic and electric fields demonstrates how the introduction of polar FACs changes the low-latitude currents and produces what is commonly referred to as penetration electric field.

395 Figures 2 and 3 show the input parameters and corresponding steady-state solutions for the two cases. The two top rows show the input patterns of FACs (a and b), winds (c), and conductance (d and e). The two bottom rows show the steady-state electric potential for this input. We will show below the inductive part of the electric field that is present before steady state is reached. The global maps show the Earth on 1 June 2001, 00:00 UT with local noon at the center. The main magnetic field is given by the International Geomagnetic Reference Field (IGRF) (Alken et al., 2021) for the time of our simulation. The dashed
400 blue lines in the global maps show the dip equator, where $b_r = 0$, and the boundaries between high and low latitudes, set here to 45° Modified Apex latitude (Richmond, 1995). The coordinates in these maps refer to geographic coordinates, while the




Figure 2. Simulation inputs and steady-state solutions for simulation case 1, where input high-latitude FACs are zero. Polar maps show parameters in Modified Apex latitude and magnetic local time, while global maps are in geographic coordinates with local noon at the center. The input parameters are the field-aligned currents in the Northern (a) and Southern Hemispheres (b); the horizontal winds (c); and the Hall (d) and Pedersen (e) conductance. All input parameters are represented using spherical harmonics, which may differ slightly from their empirical model origins (Drob et al., 2015; Hardy et al., 1987); the spherical harmonic representations shown in the figure are used in our simulations. Based on these inputs, the steady-state solution for the radial magnetic field at $r = R$ is shown in panels f (Northern hemisphere), g (Southern hemisphere), and h, with colors representing the field magnitude. Black contours in these panels indicate the equivalent current, spaced at intervals of 20 kA. Panels i (Northern hemisphere), j (Southern hemisphere), and k show the steady-state electric potential, with contours spaced at 3 kV. The induction electric field is zero in the steady state and therefore does not appear in the figure. The blue lines in the global plots represent the magnetic dip equator (dashed) and $\pm 45^\circ$ Modified Apex latitude, which is the boundary between high and low latitudes used in our simulations.




Figure 3. Simulation inputs and steady-state solutions for simulation case 2. The format of the figure is the same as in Figure 2. The input high-latitude FACs are from the AMPS model (Laundal et al., 2018).

coordinates in the polar maps refer to Modified Apex latitude and magnetic local time (Laundal and Richmond, 2017). Our simulations are carried out in geocentric spherical coordinates. We truncate the spherical harmonic expansion at $n, m = 90$, corresponding to a spatial resolution of about 400 – 500 km. The color scales and contour spacing are the same across both
 405 Figures 2 and 3 and throughout the subsequent simulation result figures.

The Hall and Pedersen conductances are given as $\sqrt{\Sigma_{\text{aurora}}^2 + \Sigma_{\text{EUV}}^2 + \Sigma_{\text{background}}^2}$ on a grid, and then represented with surface spherical harmonics. Σ_{aurora} is the conductance produced by auroral precipitation, here defined by the precipitation model by Hardy et al. (1987) for $K_p = 4$ converted to conductance using the equations by Robinson et al. (1987). Σ_{EUV} is the solar EUV-produced conductance, which is based on a simple empirical model by Moen and Brekke (1993) but rigorously accounts
 410 for the circular shape of Earth (Laundal et al., 2022b). This model approximately captures the conductance’s dependence on plasma density resulting from ionization by solar EUV radiation, but it does not account for the conductance’s dependence on magnetic field strength (Richmond, 2016). The main magnetic field varies significantly across the Earth, and the resulting conductance variations may influence the induction process. This effect is not addressed here and may be explored in future work. $\Sigma_{\text{background}}$ is a constant set to 2 mho in our simulations, that represents a background conductance, sometimes referred
 415 to as starlight conductance. The three contributions are combined using quadratic addition due to the mainly quadratic density dependence of the plasma decay in the E-region (Robinson et al., 2021).

The horizontal wind vectors are from the Horizontal Wind Model (HWM) by Drob et al. (2015), evaluated with code by Ilma (2017) at 110 km altitude and with an Ap index of 35. Although vertical winds should ideally be included in our simulation (u_r in Equation (11)), their contribution is minimal outside the equatorial region because of the main field geometry, and since
 420 the HWM only provides horizontal winds, we have opted to neglect them in this analysis.

The field-aligned current patterns are given by the Average Magnetic Field and Polar Current System (AMPS) model (Laundal et al., 2018) for a solar wind speed of 400 m/s, IMF $B_y = 4$ nT, $B_z = -5$ nT, $F_{10.7} = 100$ sfu, and a dipole tilt angle calculated to be 18.3° for the time chosen for our simulation.

4.1 Inductive Formation of Sq Currents

425 Figure 4 shows global maps of the radial magnetic field B_r at $r = R$ and the equivalent current function Ψ (Eq 23) at different simulation times following a step-function increase at $t = 0$ in winds or conductance (both yielding the same result) to the patterns in Figure 2c–e. We see that during the seven minutes included here, a mid/low-latitude current pattern develops on the dayside (center of the maps) with a clockwise cell in the Northern hemisphere and an anti-clockwise cell in the Southern hemisphere. This is the Sq current system, which is driven by neutral winds (Yamazaki and Maute, 2017). In this model, the Sq
 430 current system results from the magnetic field deformation implied by the induction equation, driven by a static wind pattern and the hemispheric linkage described in Section 3.2.1, rather than being derived from electrostatic potentials obtained from solving the current continuity equation as in previous models (Richmond, 2016). Current continuity is always maintained in our simulations since we use Ampère’s law without displacement currents (Equation (10)).




Figure 4. The radial magnetic field (shown in colors) and equivalent current (black contours, spaced at 20 kA intervals) are plotted at 16 different simulation times, following a step increase in winds and/or conductance from zero to the values shown in Figure 2. The simulation time in seconds is indicated above each panel. The plots are presented in geographic coordinates with noon at the center (consistent with Figure 2). The blue dashed lines are the magnetic dip equator and contours of $\pm 45^\circ$ Modified Apex latitude. The color scale for $B_r(r = R)$ is identical to that in Figure 2.

Figure 5 shows maps for the same simulation times as in Figure 4, but this time of the surface electric field (Equation (7)).
 435 The surface electric field can be written as

$$\mathbf{E}_S = -\nabla_S \Phi + \hat{\mathbf{r}} \times \nabla_S W, \quad (38)$$

where Φ and W are scalar fields. Φ is the electric potential, and its isocontours, spaced by 3 kV, are shown in black in Figure 5. W is the electric field stream function, and its isocontours, also spaced by 3 kV, are shown in orange. While the contour lines of Φ are perpendicular to the potential electric field, the contours of W are parallel to the rotational part of the electric field.
 440 The corresponding plasma flow above the ionosphere, where ideal MHD holds, will be perpendicular to the induction contours, which means that it describes a compression or expansion of the plasma. Only W contributes to $(\nabla \times \mathbf{E}) \cdot \hat{\mathbf{r}}$ and therefore only W is directly related to temporal variations in the magnetic field. Even though there are no contour lines of W visible after 50 s, its impact remains visible in the changing Φ and in the changing magnetic field in Figure 4. The magnitude and scale sizes of the Sq currents in Figure 4 and low-latitude electric field in Figure 5 are consistent with typical patterns reported in the
 445 literature (e.g., Yamazaki and Maute (2017); Richmond et al. (1980)).

4.2 Dynamic Response to FAC Increase

Figures 6 – 9 show the response in the magnetic and electric fields, respectively, during the seven minutes following a step increase in high-latitude field-aligned currents, from zero to the patterns shown in Figure 3a and b. The wind and conductance is the same as in the no-FAC case, and the initial condition for this simulation is the steady state solution for the no-FAC case
 450 shown in Figure 2f–k.

Figures 6 and 7 focus on the polar regions and display the response in the magnetic and electric fields, respectively, as functions of magnetic latitude and local time. These figures show that the smaller scale sizes emerge quickly, within few seconds, while the large-scale field continues to change slowly for several tens of seconds. The induction electric field is strong in the beginning, but decreases to a fraction of the potential electric field within about 10 s. It reduces faster in the Southern
 455 hemisphere than in the Northern hemisphere because η_P and η_H are higher there due to the lower conductance. We stress that even though the induction electric field quickly becomes small, careful inspection of the last snapshots in Figures 6 and 7 reveal subtle differences which imply that induction continues to be active for several minutes.

Figures 8 and 9 show the magnetic and electric fields, respectively, on a global scale. In Figure 8 the first snapshot is identical to Figure 2h and represents the steady-state solution in the absence of FACs, but with the winds and conductance of Figures 3c–
 460 e. The subsequent snapshots show how B_r and the equivalent current Ψ develop as a result of the step increase in high-latitude FACs. The clearest change is the appearance of the polar currents, the details of which were discussed above. In Figure 8 we see their equatorward extension, emerging on time scales of several tens of seconds. Another clear change is the emergence of a region with negative B_r (blue) east of the main Sq current cell in the Northern hemisphere, which takes several minutes to develop. This difference in response time at high and low latitudes to a rapid increase in polar FACs is not predicted by
 465 conventional models, and might be measurable with for example high time-resolution ground magnetometer networks.




Figure 5. The electric potential Φ (black contours) and electric stream function W (orange contours) at various simulation times, following a step increase in the winds and/or conductance from zero to the values shown in Figure 2. Φ and W are defined in Equation (38). The contour spacing is 3 kV for both scalar fields. The simulation times and map projection are the same as in Figure 4.




Figure 6. The radial magnetic field (colors) and equivalent current (black contours, spaced at 20 kA intervals) at 16 simulation times, following a step increase in polar field-aligned currents from zero to the values shown in Figure 3. At each simulation time, a pair of polar plots is displayed, with the northern hemisphere on the left and the southern hemisphere on the right. The initial condition for this simulation is the steady-state solution from Figure 2 with no FAC but with the same winds and conductance. The simulation time in seconds is indicated above each panel. The plots are presented on Modified Apex latitude and local time grids. The color scale for $r = R$ is identical to that in Figures 2 and 3.




Figure 7. The electric potential (black contours) and electric stream function (orange contours) at various simulation times following a step increase of the high-latitude field-aligned currents from zero to the values shown in Figure 3. The contour spacing is 3 kV for both scalar fields. The simulation times and map projection are the same as in Figure 6.




Figure 8. The radial magnetic field and equivalent current from Figure 6 shown in geographic coordinates. The simulation times, scales, and format are the same as in previous figures.

Figure 9 shows the global evolution of the electric field following the step increase in polar FACs. Ten seconds after the increase, we see that a local maximum emerges near midnight (left and right edges of the plots), and continues to increase for 20–30 seconds, before it decreases and disappears completely after three minutes. This can be interpreted as a transient penetration electric field that corresponds to upward plasma flow in the post-midnight region and downward plasma flow in the pre-midnight region. After seven minutes (the last snapshot) a low-latitude electric field is established that is much stronger and significantly different from the pattern seen in Figure 5, prior to the increase in high-latitude FACs. The difference can be interpreted as a slowly varying penetration electric field that eventually reaches the steady state shown in Figure 3k. It is interesting to note that even after seven minutes, we see subtle differences from Figure 3k, which means that it takes even longer to reach a steady state.




Figure 9. The electric potential and stream function from Figure 7 shown in geographic coordinates. The simulation times and format are the same as in previous figures.

475 In the simulations presented here, the FAC, wind, and conductance patterns are held fixed in a geographic coordinate system, instead of slowly rotating them to be fixed with respect to the Sun. The rotation would be slow compared to the time scales of most processes seen in Figures 4–9, but given that some changes take place over several minutes, induction could nevertheless play a non-negligible role even in the dynamics that results from the rotations of static patterns.

5 Discussion

480 The model and simulation results presented in this study explain and demonstrate the dynamic interplay between the magnetosphere, ionosphere, and neutral atmosphere. By implementing a global 2D ionospheric model based on the theoretical framework outlined in Sections 2 and 3, we have demonstrated the effect of induction, as captured by Faraday’s law and the

Generalized Ohm’s law, which combine to Equation (11). In our model, the electromagnetic fields respond dynamically to changes in the driving parameters – the magnetic field of high-latitude field-aligned currents, neutral winds, or ionospheric conductivity – in contrast to the instant magnetostatic response implied by conventional models for large-scale magnetosphere-ionosphere coupling. While capturing this dynamic response for the first time on a global scale, our model also successfully reproduces key steady-state features of the ionospheric current systems and magnetic field perturbations observed in previous studies, but using a fundamentally different approach. Instead of treating the electric field and current density as the primary variables, as is common in ionospheric physics, the primary parameter of our model is the magnetic field. The velocity could be included as a primary variable by evolving the ion momentum equation in time (Laundal et al., 2024), but for simplicity, we use a steady-state solution here. If needed, the steady-state ion and electron velocities can be calculated using equations from (Brekke, 2013, Chapter 5.1), and the height-integration approach of Laundal et al. (2024). In this way, we offer a framework within the $\mathbf{B}, \mathbf{v}$ paradigm (Parker, 1996; Vasyliūnas, 2012) to explain the magnetosphere-ionosphere coupling process, capable of accounting for its dynamics.

The simulation results shown in Sections 4.1 and 4.2 are intentionally idealized, with step changes in winds/conductance and high-latitude field-aligned currents, respectively, to demonstrate the dynamic transition of the ionosphere between states. While these abrupt transitions are artificial, large-scale sudden changes do occur in nature, and our simulations highlight the necessity of accounting for inductive effects when analyzing variations on time scales of tens of seconds. Example of such large-scale rapid changes are solar flares or solar eclipses, which can cause sudden changes in ionospheric conductivity. The inductive effects of these changes have been previously considered (Nagata, 1966), but not with the level of realism achieved by our model. Future work could apply our model with more realistic time profiles of conductance changes associated with solar flares to explore how geomagnetic crochets (e.g., Yamauchi et al. (2020)) arise through inductive processes. Another large-scale disturbance where inductive effects should be considered is the sudden increase in solar wind pressure (Madelaire et al., 2022), which propagate through the magnetosphere in seconds and cause magnetic field variations on the ground that imply the presence of significant induction electric fields (Madelaire et al., 2024).

In the rest of this section, we discuss the novel aspects of our model in more detail. Section 5.1 compares our work with previous studies on inductive magnetosphere-ionosphere coupling. Section 5.2 discusses how the perspective on low-latitude wind-driven magnetic field disturbances differs from conventional perspectives. Section 5.3 investigates induction time scales based on an analytical solution to a highly simplified version of the induction equation. Section 5.4 discusses the main limitations of our model and avenues for improvements in future iterations.

5.1 Comparison With Previous Work

Many previous studies of magnetosphere-ionosphere coupling have concentrated on the “gap region” (Merkin and Lyon, 2010), often neglecting the effects of induction at the lower boundary, which is the focus of our investigation. Previous models of magnetosphere-ionosphere coupling typically employ 1D frameworks that consider only the vertical direction (Tu et al., 2011, 2014; Wright, 1996) or 2D models that account for both the vertical dimension and one horizontal dimension (Tu and Song, 2016, 2019; Lotko, 2004; Dreher, 1997; Otto and Zhu, 2003). These studies, many of which also include more

detailed treatments of fluid dynamics, have demonstrated that our assumption about the gap region – namely, that the magnetic field rapidly reaches a force-free state – is overly simplistic, particularly on the short time scales we consider. However, all these studies overlook the role of induction at the lower boundary, which we account for in this work. Our results shows the importance of including this lower-boundary induction.

Over the past decades, starting with Yoshikawa and Itonaga (1996), several papers have investigated the inductive response of the ionosphere in terms of Alfvén wave reflection (Buchert and Budnik, 1997), and how it affects ultra-low frequency (ULF) wave propagation (Sciffer et al., 2004; Waters and Sciffer, 2008; Lysak, 2004; Lysak and Song, 2006; Lysak et al., 2020). These studies provide valuable insights into the ionospheric response over different time scales, and clearly demonstrate the need to account for induction in the ionospheric boundary. However, by focusing on wave solutions, they are complicated to use for capturing rapid changes as in the simulations described in Sections 4.1 and 4.2, or to the magnetosphere-ionosphere coupling schemes required in global magnetospheric simulations. In addition, our approach is arguably more straightforward to expand to capture non-linear behavior.

Vanhamäki (2011) presented an approach for inductive magnetosphere-ionosphere coupling that could overcome these issues. His method shares similarities with ours, using a set of basis functions for the magnetic field to solve for the electric field given the FAC. Instead of spherical harmonics, he used Spherical Elementary Current Systems (SECS) (Amm, 1997; Vanhamäki and Juusola, 2020). The SECS functions are global, but have short reach, which makes them ideal for regional analyses, but less suitable for large-scale domains. Vanhamäki (2011) therefore only demonstrated the technique in a limited region, studying inductive effects in the vicinity of an auroral omega band. In Section 5.4.4 we discuss how the approach by Vanhamäki (2011) could be integrated with ours in the future to achieve higher spatial resolution in certain regions.

5.2 Low-latitude Ionospheric Electrodynamics

The global maps in Section 4 show that induction processes can play a role for a long time after a step change in winds or FACs, even beyond the seven minutes displayed in the figures.

The slow changes seen at low latitudes can be understood by inspecting the ionospheric induction equation, Equation (11). The time scale of magnetic field variations (left-hand side) depends on the spatial derivatives and, consequently, the scale sizes of specific combinations of conductance, neutral wind, magnetic field disturbance, and main magnetic field (right-hand side). At low latitudes, the conductance is mainly determined by solar EUV radiation, and gradients are expected to be small. The drivers of the induction at low latitudes are the wind, the poloidal field associated with field-aligned currents, and the corresponding shielding current. Since the field-aligned currents are remote, the structure of this poloidal magnetic field is expected to be dominated by large scales. It may therefore take a long time (tens of seconds to minutes) before the magnetic field of field-aligned currents become visible in low-latitude ground magnetometers.

Our model offers an alternative perspective on low-latitude ionospheric dynamics relative to conventional textbook descriptions. For example, (Kelley, 2009, Chapter 2.6) argues, in response to the $\mathbf{B}, \mathbf{v}$ -centered view advanced by Vasyliūnas and Song (2005), that “[...]decades of successful application of electroquasistatics in the ionosphere should not be replaced when it is applicable”. While the historical success of electroquasistatics is undisputed in ionospheric physics, our simulations show

that important dynamics will be missing without considering how the magnetic field changes through induction. Our results show that even as the system converges toward an electroquasistatic equilibrium, it can take a long time, and involve large-scale intermittent structures in the electric field.

The electroquasistatic approach is computationally efficient, and often yields results that are nearly identical with results from our inductive model that is by comparison more computationally expensive. We nevertheless argue that induction is needed to answer the question of *how* electric fields and currents change. For example, in our model we see a strong dependence in the low-latitude electric field on the pattern of field-aligned currents at high latitudes. This is the so-called penetration electric field. The term penetration electric field stems from an idea that charges in the inner magnetosphere produces a large-scale electric field (e.g. Kelley, 2009). This contradicts basic facts about space plasmas, since any such charge distribution would quickly (on plasma frequency time scales) be canceled in a frame of reference moving with the plasma (Parker, 2007; Vasyliūnas, 2005, 2012; Parker, 1996). In our model, on the other hand, the electric field is produced in response to a deformation of the magnetic field. The magnetic field deformation changes the Lorentz force, and hence the force balance between the Lorentz force and collisional friction. The ion and electron motion in the ionosphere changes accordingly, and their motion gives the electric field.

In our approach, we assume that the magnetic field deformation related to field-aligned currents occurs instantaneously – an unrealistic simplification – by assuming the magnetic field is perfectly force-free. Simulations by Tu and Song (2019) offer a complementing perspective, resolving plasma dynamics in the vertical dimension and in only a single dimension on the sphere. The propagation of magnetic disturbances from the magnetosphere to the ionosphere as explained by Tu and Song (2019), and the dynamic response of the ionosphere as presented in this paper explains the apparent penetration electric field in terms of $\mathbf{B}$ and $\mathbf{v}$. Our simulations also show transient features of the penetration electric field that are impossible to predict with conventional theory, but might be possible to observe in radar measurements.

The traditional explanation of the Sq current system is that neutral winds set up a polarization electric field to ensure charge neutrality, and that this electric field drives currents, and the magnetic field is perturbed accordingly. The inductive formation shown in Section 4.1 also starts with the neutral winds, but yields an entirely different description of the chain of causality. The winds do two things: 1) If $\mathbf{u} \times \mathbf{B}_0$ has a curl, it directly induces a magnetic field perturbation; 2) if $\mathbf{u} \times \mathbf{B}_0$ at conjugate points imply different electric fields, the magnetic field lines at conjugate points move in different directions and cause magnetic tension to build up. Alfvén waves between hemispheres communicate the imbalance, relaxing the magnetic tension. In reality, this would take some time, but in our study we assume that the balance is reached immediately, and the (imposed) magnetic field adjusts such that the electric potential maps between hemispheres, without violating the assumption of a force-free magnetic field, and current continuity. Sq currents will only form if ~~the winds imply there is~~ an interhemispheric imbalance in Lorentz force at conjugate points, mainly arising from neutral winds. The formation of the Sq currents is a manifestation of the magnetic field deforming, building up the Lorentz force so that (mainly) ions are pushed through the moving neutral atmosphere while (mainly) electrons act as a neutralizing fluid, moving to preserve charge neutrality. The electric field is determined by this motion. In steady state the electric field is such that no further deformation happens, as the $\mathbf{j} \times \mathbf{B}$ force and momentum transfer

585 due to collisions with neutrals counterbalance each other. At this point, the electric field is purely a potential field, and there is no induction as per Faraday’s law.

The critical importance of imbalanced winds at conjugate points agrees with the conclusion by Buchert (2020), who investigated this from an $\mathbf{E}, \mathbf{j}$ perspective. He describes the low-latitude ionosphere as an *entangled dynamo* since the winds at conjugate points are coupled.

590 The equatorial electrojet (EEJ), a narrow channel of intense eastward currents along the dayside dip equator, is missing from the patterns in Section 4. This is presumably because the formation of the EEJ involves 3-dimensional structures, as discussed by Yamazaki and Maute (2017), neglected in our 2D approach. In addition, the horizontal resolution of our simulations (≈ 450 km) may anyway be too crude to resolve the EEJ, which extends $\approx 3^\circ$ to either side of the dip equator (Yamazaki and Maute, 2017).

595 5.3 Approximating Induction Time Scales With Analytical Solution

It is instructive to consider the highly simplified case of a constant and radial main magnetic field, zero wind, and conductances that are uniform, conditions under which Fukushima’s theorem holds (Fukushima, 1976). In that case, we show in B that Equation (11) reduces to a differential equation for the coefficients of the induced field (q_n^m, s_n^m) , as defined in Equation (15), as functions of the coefficients of the imposed field (ψ_n^m, ξ_n^m) . For constant imposed field coefficients, the solution to this
600 equation is

$$\begin{pmatrix} q_n^m(t) \\ s_n^m(t) \end{pmatrix} = \frac{\Sigma_H}{\Sigma_P} \frac{n+1}{2n+1} \begin{pmatrix} \psi_n^m \\ \xi_n^m \end{pmatrix} \left[1 - e^{-\frac{\eta_P(2n+1)}{R\mu_0}t} \right] + \begin{pmatrix} q_n^m(t=0) \\ s_n^m(t=0) \end{pmatrix} e^{-\frac{\eta_P(2n+1)}{R\mu_0}t}. \quad (39)$$

In this simplified case, there is no coupling between scale sizes. The exponents show that the characteristic time scales of magnetic field variations depend on scale size (spherical harmonic degree n), the Pedersen resistance η_P , and the radius of the ionosphere R . Figure 10 shows contours of constant e-folding time for different n and Σ_P , with Σ_H held fixed at 5 mho. We see
605 that variations are quickest when $\Sigma_P = \Sigma_H$, ranging from about one min for global scale sizes, to 1 s for $n \sim 80 - 90$, which corresponds to scale sizes of about 500 km. We stress that realistic conditions can yield very different results, as illustrated in the previous section.

5.4 Limitations and Potential Enhancements

While the model presented here clearly has limitations, the approach opens several opportunities to study aspects of magnetosphere-
610 ionosphere-thermosphere coupling that have so far not been possible. Here we discuss some of the limitations and how they could be eliminated in future iterations.

5.4.1 From 2D to 3D

Our model is two-dimensional and therefore by definition does not capture dynamics in the vertical direction, including inherently 3D phenomena such as the EEJ. The spherical harmonic representations used here may be difficult to generalize to a 3D




Figure 10. Characteristic time of magnetic field variations at different scale sizes for different Pedersen resistances. In this plot, the Pedersen resistance is varied by setting the Hall conductance to 5 mho and varying the Pedersen conductance.

615 ionosphere. An alternative approach could be to use a finite difference or finite volume approach inside a 3D ionosphere, and use our spherical harmonic approach to specify upper and lower boundary conditions. We note that excellent 3D models of the ionosphere already exist (Qian et al., 2014; Huba et al., 2008; Zettergren and Snively, 2015; Ridley et al., 2006), but none of them include induction.

5.4.2 Coupling With Global Magnetosphere Simulations

620 Our model uses as input the patterns of high-latitude FACs, 2D horizontal wind, and ionospheric conductance, and can output the electric field. These input and output types are also used in magnetosphere-ionosphere coupling modules of global magnetospheric simulations (Merkin and Lyon, 2010; Ridley et al., 2004; Ganse et al., 2025), suggesting that these modules could be replaced with ones based on the approach presented here. This would allow us to study the impact of the magnetosphere on ionospheric induction, and the impact of ionospheric induction on the global magnetosphere.

625 Our global solver also addresses an inconsistency that can arise in existing models: In conventional approaches, the current continuity equation is solved independently for the electric potential in each hemisphere. This approach does not guarantee consistent mapping of the potential between conjugate points, violating the assumptions of ideal MHD in the magnetosphere and no induction in the ionosphere (Hesse et al., 1997). In contrast, our approach includes induction in the ionosphere, which means that an electric potential mismatch between hemispheres is not in violation of the assumptions of the model. In addition, 630 in our approach, the ionosphere behaves more like a dynamic fluid cell at the inner edge of the magnetosphere rather than a rigid boundary. As a result, abrupt changes in ionospheric conductance, neutral winds, or high-latitude magnetic field perturbations (or field-aligned currents) do not produce the same sudden, potentially unphysical changes in the ionospheric electric field seen in conventional magnetostatic MI coupling schemes. Instead, the electric field evolves smoothly through induction, reflecting the motion of magnetic field lines. This motion may be asymmetric between hemispheres, influencing the displacement field 635 as described by Laundal et al. (2022a). Although including induction does not ensure that the potential maps along magnetic field lines, as noted by Hesse et al. (1997), the total electric field will map in regions where ideal MHD holds.

5.4.3 Inclusion of Fluid Equations

In this paper we only evolve B_r in time, while the wind, conductance, and high-latitude FACs are given. Evolving the FACs would amount to running a global magnetosphere simulation in parallel, as discussed above. The winds and conductance, 640 however, have their own dynamics in the ionosphere, which we neglect here for simplicity. The conductance depends mainly on the plasma density, which is described by the continuity equation; but it also depends on the collision frequencies which depend on temperature (Schunk and Nagy, 2009), which is described by the energy equation. Including plasma fluid moments in the simulation, as in the 1D simulation presented by Laundal et al. (2024), is likely to reveal more complex dynamics. For example, frictional heating, caused by differential motion between ions and neutrals, raises the temperature. This increase alters 645 collision frequencies and conductances, which, in turn, affect the induction process. The modified induction process further influences frictional heating, creating a nonlinear feedback loop. These complex interactions will be explored in future studies.

5.4.4 Refinement

Our use of spherical harmonics is well suited for global analyses. However, this method becomes increasingly ineffective with increasing spatial resolution due to the rapidly growing number of terms required in the spherical harmonic expansion. To address this, our model could be improved by combining spherical harmonics with Spherical Elementary Current Systems (SECS). In this hybrid approach, spherical harmonics would provide the background global representation, while SECS could be used to superimpose a high-resolution magnetic field on a fine grid in specific regions of interest. To implement this consistently and account for the poloidal magnetic field of field-aligned currents, the standard SECS functions would need to be modified. Specifically, the poloidal field of the field-aligned currents must be incorporated along realistic magnetic field geometries, following an approach similar to Vanhamäki et al. (2020) using dipole field lines.

5.4.5 Inclusion of Telluric Currents

In this paper we have neglected the conductivity of the Earth. In reality, the magnetic field induced in the ionosphere leads to currents flowing in the interior of the Earth. Accounting for these induced currents is important for the interpretation of ground magnetometer data, since ground magnetometers observe a combination of ionospheric and telluric magnetic fields. While this issue has been discussed extensively in the literature (e.g., Juusola et al., 2016, 2020; Olsen et al., 2010), the influence of ground induction on ionospheric dynamics is much less studied. Vanhamäki et al. (2005) made simple order-of-magnitude estimates of this effect by using 1D ground conductivity and complex image method with realistic ionospheric sources, and concluded that the effect of ground induction on ionospheric dynamics is significantly smaller than the effect of ionospheric self-induction. However, recently Juusola et al. (2024) made more detailed calculations with 3D ground conductivity and showed that telluric induction can significantly modify the inductive electric field in the ionosphere and should not be ignored.

Including ground induction in our model would change the lower boundary condition for the ionosphere, $\mathbf{B}^-$. To account for ground induction, the Laplace equation must be solved for the potential magnetic field below the ionosphere V^e with more accurate lower boundary conditions. With spherical harmonics, this can be done with transfer functions that depend on ground conductivity, and map between the ionospheric magnetic field observable from the ground (described here by coefficients k_n^m and l_n^m) and spherical harmonic coefficients that describe the induced magnetic field in the ground (e.g., Grayver et al., 2021). While this is feasible, it is beyond the scope of this paper.

6 Conclusions

We have presented a new model to describe magnetosphere-ionosphere-thermosphere coupling on a global scale, that takes into account the induction equation, Faraday's law. Our approach treats the magnetic field as the primary variable, while currents and electric fields are evaluated from Ampère's law and the Generalized Ohm's law, respectively. In this way, our work deviates from the $\mathbf{E}, \mathbf{j}$ -centered view that is conventionally taken in ionospheric physics, but aligns with how space plasmas are handled elsewhere (Leake et al., 2014; Parker, 2007).

Results from numerical simulations, using spherical harmonic representations of the various quantities (see C), were presented in Section 4. The results show that induction takes place on a broad range of time scales, from fractions of seconds to minutes. The time scale depends on the magnitudes and spatial scales of conductances, imposed magnetic field, neutral wind, and on the main magnetic field geometry (Equation (11)). Large-scale structures at low latitudes take particularly long to adapt to sudden changes in polar FACs, winds, or conductance.

Our model is two-dimensional, and uses as boundary condition an imposed magnetic field that is in reality a result of an inductive process above the ionosphere that we neglect here for simplicity. At high latitudes, the imposed magnetic field is defined to be consistent with a prescribed pattern of field-aligned currents; at low latitudes it is calculated by assuming that the magnetic field adapts to preserve current continuity between hemispheres and matching electric potentials along magnetic field lines, given a neutral wind pattern. These assumptions lead to Sq currents and low-latitude electric fields appearing through induction in our simulations, representing, to our knowledge, the first time that this has been demonstrated.

The model presented here can in principle be ~~directly applied in~~ incorporated into global magnetosphere simulations ~~to obtain~~ and upper-atmospheric circulation models to provide a more complete description of ~~how the magnetosphere and ionosphere interact~~ magnetosphere-ionosphere interactions. Steady-state solutions for the electric field can also be obtained, see C for implementation details. This gives a truly global alternative to prevailing magnetostatic models, which invoke boundary conditions that decouple high and low latitudes. The numerical implementation of the model, used to produce the simulation results presented here, is discussed in C.

Code availability. The model is implemented in Python, in a package called PynaMIT, which is available at <https://github.com/DynaMIT-uib/PynaMIT>.

PynaMIT is highly flexible, and can be run with any resolution (limited by computing power). It can be run with different main field models: Radial field, dipole field, or IGRF (as demonstrated in Section 4).

Appendix A: Derivation of the Thin-Sheet Conductivity and Resistivity Tensors

700 In Section 3 we present a two-dimensional version of the Generalized Ohm's law for a collisional ionosphere (7). The equation is a projection of the corresponding 3D equation (3) onto a spherical shell and so it is valid for an ionosphere that is truly two-dimensional. Nevertheless, as indicated by the choice of notation, we interpret $\mathbf{J}_S$ and $\Sigma_{P,H}$ as the height-integrated versions of the corresponding 3D quantities $\mathbf{j}$ and $\sigma_{P,H}$. If $\mathbf{j}$ or $\sigma_{P,H}$ vary with height, which they do in the realistic ionosphere, Equation (7) is not simply a height integral of Equation (2), so this interpretation requires more justification.

705 The justification is provided by using the ionospheric Ohm's law, and then solve this equation for $\mathbf{E}_S$. In the dynamo region of the ionosphere (primarily the E-region), $\mathbf{E}$ and $\mathbf{B}$ vary only slowly with altitude, whereas the conductivities vary sharply. Because of this scale separation, the altitude dependence is carried almost entirely by the conductivities, so integrating the ionospheric Ohm's law yields well-defined conductances. For generality we retain a finite field-aligned conductivity $\sigma_{\parallel}$ (though elsewhere in the paper we assume it to be infinite), and write the ionospheric Ohm's law as

$$710 \quad \mathbf{j} = \sigma_P \mathbf{E}'_{\perp} + \sigma_H \hat{\mathbf{b}} \times \mathbf{E}' + \sigma_{\parallel} E_{\parallel} \hat{\mathbf{b}}. \quad (\text{A1})$$

Here, $\mathbf{E}' = \mathbf{E} + \mathbf{u} \times \mathbf{B}$ is the electric field in the neutral frame of reference. If we assume that both $\mathbf{B}$ (which is dominated by the main field, $\mathbf{B}_0$) and $\mathbf{E}$ vary little over the height of the ionosphere, we can integrate this equation to get a 3D height-integrated current:

$$\mathbf{J} = \Sigma_P \mathbf{E}_{\perp} + \Sigma_H \hat{\mathbf{b}} \times \mathbf{E} + \Sigma_{\parallel} E_{\parallel} \hat{\mathbf{b}} + \mathbf{Q}. \quad (\text{A2})$$

715 Here, $\Sigma_{P,H,\parallel}$ are the height-integrated $\sigma_{P,H,\parallel}$, and $\mathbf{Q} = \mathbf{Q}_P + \mathbf{Q}_H$, where

$$\mathbf{Q}_P = \int \sigma_P(r) (\mathbf{u}(r) \times \mathbf{B}) dr \quad (\text{A3})$$

$$\mathbf{Q}_H = \int \sigma_H(r) (\mathbf{u}(r) \times \mathbf{B}) \times \hat{\mathbf{b}} dr \quad (\text{A4})$$

are integrated wind terms weighted by the Pedersen and Hall conductivities (Richmond, 1995; Hatch et al., 2024).

We use that $\mathbf{E}_{\perp} = \mathbf{E} - E_{\parallel} \hat{\mathbf{b}}$ and decompose $\mathbf{J}$ in radial (subscript r) and tangential/surface components (subscript S):

$$720 \quad \begin{pmatrix} J_r \\ \mathbf{J}_S \end{pmatrix} = \begin{pmatrix} \Sigma_P E_r + \Sigma_H (\hat{\mathbf{b}} \times \mathbf{E})_r + (\Sigma_{\parallel} - \Sigma_P) E_{\parallel} b_r + Q_r \\ \Sigma_P \mathbf{E}_S + \Sigma_H (\hat{\mathbf{b}} \times \mathbf{E})_S + (\Sigma_{\parallel} - \Sigma_P) E_{\parallel} \hat{\mathbf{b}}_S + \mathbf{Q}_S \end{pmatrix}. \quad (\text{A5})$$

In our 2D/thin-sheet formulation the discontinuity of the horizontal magnetic field across the ionosphere is described by the Ampère jump condition (10). This relation requires a surface current confined to the sheet: $\mathbf{J}_S$ in the equation above. Because the discontinuity only involves the horizontal magnetic field above and below, there can be no independent horizontal perturbation field inside the sheet itself. Consequently the sheet current is purely tangential: $J_r = 0$. The horizontal divergence of $\mathbf{J}_S$ is in general nonzero and is balanced by the radial current carried in the field-aligned system above. We therefore impose

725

$J_r = 0$ in the radial component of (A5) and solve for E_r :

$$E_r = - \frac{\Sigma_H(\hat{\mathbf{b}} \times \mathbf{E})_r + (\Sigma_{\parallel} - \Sigma_P)b_r(\hat{\mathbf{b}}_S \cdot \mathbf{E}_S) + Q_r}{\Sigma_P(b_\theta^2 + b_\phi^2) + \Sigma_{\parallel}b_r^2} \quad (\text{A6})$$

This can be inserted in the expression for $\mathbf{J}_S$ to get an equation for the surface current that only depends on the horizontal components of the electric field. This is a version of the height-integrated ionospheric Ohm's law that explicitly states the dependence on the conductance and main magnetic field geometry through the tensor $\mathbb{A}_{\text{cond}}(\Sigma, \hat{\mathbf{b}})$:

$$\mathbf{J}_S = \mathbb{A}_{\text{cond}}(\Sigma, \hat{\mathbf{b}})\mathbf{E}_S + \mathbf{Q}_{\text{eff}}, \quad (\text{A7})$$

where

$$\mathbf{Q}_{\text{eff}} = \mathbf{Q}_S - \frac{(\Sigma_{\parallel} - \Sigma_P)b_r\mathbf{b}_S Q_r - \Sigma_H(\hat{\mathbf{r}} \times \hat{\mathbf{b}})_S Q_r}{\Sigma_P(b_\theta^2 + b_\phi^2) + \Sigma_{\parallel}b_r^2}. \quad (\text{A8})$$

is the effective wind, a combination of the tangential/surface component of the integrated wind terms and the contribution that comes from substituting E_r in the surface component. The conductance tensor can be expressed as

$$\mathbb{A}_{\text{cond}}(\Sigma, \hat{\mathbf{b}}) = \frac{1}{\Delta} \begin{pmatrix} Kb_\phi^2 + M(1 - b_\phi^2) & (M - K)b_\theta b_\phi - \Sigma_H \Sigma_{\parallel} b_r \\ (M - K)b_\theta b_\phi + \Sigma_H \Sigma_{\parallel} b_r & Kb_\theta^2 + M(1 - b_\theta^2) \end{pmatrix}. \quad (\text{A9})$$

where

$$\Delta \equiv \Sigma_P(b_\theta^2 + b_\phi^2) + \Sigma_{\parallel}b_r^2, \quad K \equiv \Sigma_P^2 + \Sigma_H^2, \quad M \equiv \Sigma_P \Sigma_{\parallel}. \quad (\text{A10})$$

This conductance tensor reduces to the expression given by Amm (1996); Rishbeth and Garriott (1969) for $b_\phi = 0$.

We can solve Equation (A7) for $\mathbf{E}_S$ to get the corresponding 2D Generalized Ohm's law:

$$\mathbf{E}_S = \mathbb{A}_{\text{cond}}^{-1}(\Sigma, \hat{\mathbf{b}})(\mathbf{J}_S - \mathbf{Q}_{\text{eff}}) = \mathbb{A}_{\text{res}}(\eta, \hat{\mathbf{b}})(\mathbf{J}_S - \mathbf{Q}_{\text{eff}}) \quad (\text{A11})$$

where the inverse of the conductance tensor is the resistance tensor, $\mathbb{A}_{\text{res}}(\eta, \hat{\mathbf{b}})$:

$$\mathbb{A}_{\text{res}}(\eta, \hat{\mathbf{b}}) = \begin{pmatrix} \eta_P + (\eta_{\parallel} - \eta_P)b_\theta^2 & (\eta_{\parallel} - \eta_P)b_\theta b_\phi + \eta_H b_r \\ (\eta_{\parallel} - \eta_P)b_\theta b_\phi - \eta_H b_r & \eta_P + (\eta_{\parallel} - \eta_P)b_\phi^2 \end{pmatrix}, \quad (\text{A12})$$

where η_P and η_H are defined in terms of the conductances as in (8) and (9), and $\eta_{\parallel} = 1/\Sigma_{\parallel}$. We emphasize that $\eta_{P,H}$ are *not* height-integrated resistances, but quantities derived from the height-integrated conductivities.

The resistance tensor (A12) is equivalent to the matrix defined in equation (11) of O'Sullivan and Downes (2006), which is written in Cartesian coordinates, and it reduces to the matrix in equation (11) of this paper in the limit $\eta_{\parallel} = 0$. We have thus shown that the expression that we get by simply projecting the 3D Generalized Ohm's law to 2D can also be derived by height-integrating the Ionospheric Ohm's law and solving for $\mathbf{E}$. The advantage of this derivation is that it quantifies the relationship between the 2D and 3D quantities. For example, it provides a consistent way to include altitude-dependent winds in the thin-sheet formulation, by calculating the height integrals in (A3) and (A4) to give $\mathbf{Q}_H$ and $\mathbf{Q}_P$.

Appendix B: Analytical Solution for Radial Magnetic Field, No wind, and Uniform conductance

In this appendix we derive Equation (39), the analytical solution discussed in Section 5.3. This solution applies under some very restrictive assumptions: The main magnetic field $\mathbf{B}_0$ is radial and constant everywhere, the resistances η_P and η_H are uniform, and there are no neutral winds. Then Faraday's law, Equation (5) reduces to

$$\frac{\partial B_r}{\partial t} = -\eta_P (\nabla_S \times \mathbf{J}_S) \cdot \hat{\mathbf{r}} \mp \eta_H \nabla_S \times (\mathbf{J}_S \times \hat{\mathbf{r}}) \quad (\text{B1})$$

where Equation (7) was used to replace $\mathbf{E}_S$. The second term is negative when $\mathbf{B}_0$ points downward (as in the Northern hemisphere), and positive when it points upward. In the following we proceed only with the negative sign.

The sheet current can always be written as a sum of divergence-free and curl-free parts. In this case, where the main field is radial, the divergence-free part of the current is equal to the equivalent current (Equation (22)). The curl-free part can be written in terms of a potential α , so that we get

$$\mathbf{J}_S = \hat{\mathbf{r}} \times \nabla_S \Psi - \nabla_S \alpha. \quad (\text{B2})$$

where α describes the divergent part of $\mathbf{J}_S$. With this representation of $\mathbf{J}_S$ we can write Faraday's law as (e.g., Backus, 1986)

$$\frac{\partial B_r}{\partial t} = -\eta_P \nabla_S^2 \Psi - \eta_H \nabla_S^2 \alpha, \quad (\text{B3})$$

We can represent α in terms of the spherical harmonic coefficients ψ_n^m and ξ_n^m . The expression for α is (e.g., Laundal et al., 2016)

$$\alpha = -\frac{R}{\mu_0} \sum_{n,m} P_n^m(\theta) [\psi_n^m \cos(m\phi) + \xi_n^m \sin(m\phi)]. \quad (\text{B4})$$

Both Ψ and B_r can be expanded in terms of spherical harmonics with expansion coefficients q_n^m and s_n^m , by replacing k_n^m and l_n^m in Equations (23) and (21), respectively, using Equation (19). With these spherical harmonic representations, and the property that $\nabla_S^2 = -n(n+1)/R^2$, Equation (B3) becomes an equation that relates three spherical harmonic series. These series depend on q_n^m and s_n^m (the Ψ term), their time derivatives (the $\partial B_r / \partial t$ term), and on ψ_n^m and ξ_n^m (the α term). If we match the different terms in these series, we get the following set of differential equations:

$$\frac{\partial}{\partial t} \begin{pmatrix} q_n^m \\ s_n^m \end{pmatrix} = -\frac{1}{R\mu_0} \begin{pmatrix} \eta_P(2n+1)q_n^m + \eta_H(n+1)\psi_n^m \\ \eta_P(2n+1)s_n^m + \eta_H(n+1)\xi_n^m \end{pmatrix}. \quad (\text{B5})$$

For constant ψ_n^m and ξ_n^m and specified initial conditions, $q_n^m(t)$ and $s_n^m(t)$ for time $t > 0$ are given by Equation (39).

Appendix C: Discretization

The numerical implementation of the model described in this work involves discretization in terms of a combination of spherical harmonics and grid points. Here, we describe how this is done.

C1 Spherical Harmonic Representation of Scalar and Horizontal vector fields

To perform the spherical harmonic decomposition of scalar fields, we use a spherical harmonic transform approach of solving
 780 the linear system

$$\mathbb{G}_s \tilde{\mathbf{s}} = \mathbf{s}, \quad (\text{C1})$$

where the forward operator $\mathbb{G}_s$ evaluates vectors of spherical harmonic coefficients $\tilde{\mathbf{s}}$ at a given set of grid points, giving the grid-resolved data vector $\mathbf{s}$ representing the scalar field s .

Similarly, the spherical harmonic decomposition of horizontal vector fields is done by using a spherical harmonic transform
 785 approach of solving the linear system

$$\mathbb{G}_t \tilde{\mathbf{t}} = \mathbf{t}. \quad (\text{C2})$$

In this case, the field is two-dimensional, and the model vector is resolved into curl-free and divergence-free parts,

$$\tilde{\mathbf{t}} = \begin{pmatrix} \tilde{\mathbf{t}}_{\text{cf}} \\ \tilde{\mathbf{t}}_{\text{df}} \end{pmatrix}, \quad (\text{C3})$$

and the grid-resolved data vector into θ and ϕ components

$$\mathbf{t} = \begin{pmatrix} \mathbf{t}_\theta \\ \mathbf{t}_\phi \end{pmatrix}. \quad (\text{C4})$$

The forward operator can be expressed as

$$\mathbb{G}_t = \begin{pmatrix} (\mathbb{G}_t)_{\theta,\text{cf}} & (\mathbb{G}_t)_{\theta,\text{df}} \\ (\mathbb{G}_t)_{\phi,\text{cf}} & (\mathbb{G}_t)_{\phi,\text{df}} \end{pmatrix}. \quad (\text{C5})$$

$\mathbb{G}_{\theta,\text{cf}}$ and $\mathbb{G}_{\phi,\text{cf}}$ give the grid-resolved θ - and ϕ -components, respectively, of $(-\nabla_S)$ applied to the spherical harmonics used to represent the scalar potential of the curl-free part of the horizontal vector field. Likewise, $\mathbb{G}_{\theta,\text{df}}$ and $\mathbb{G}_{\phi,\text{df}}$ give the grid-resolved
 795 θ - and ϕ -components, respectively, of $(\hat{\mathbf{r}} \times \nabla_S)$ applied to the spherical harmonics used to represent the scalar potential of the divergence-free part of the horizontal vector field.

Matrices are denoted by blackboard bold. We use $\mathbb{G}_X$ to denote matrices that transform from spherical harmonics to grid points for the quantity X . All other matrices either transform between different spherical harmonic coefficients or between different quantities on the grid. Vectors and matrices associated with spherical harmonics are marked with a tilde, while those
 800 associated with grid points are not.

C2 Matrix formulation of Faraday's Law and the Steady-State Condition

From the two-dimensional Faraday's law in Equation (5), the the time derivative of the spherical harmonic coefficients $\tilde{\mathbf{B}}_{\text{ind}}$ of the induced magnetic field, which corresponds to the radial component of $\mathbf{B}$, can be written as

$$\frac{\partial \tilde{\mathbf{B}}_{\text{ind}}}{\partial t} = \tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t} \tilde{\mathbf{E}}_S \quad (\text{C6})$$

805 where $\tilde{\mathbf{E}}_S$ is a spherical harmonic representation of the electric field tangential to the ionosphere, and $\tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t}$ is the spherical harmonic representation of $(-\hat{\mathbf{r}} \times \nabla_S)$. In Section C3, we show how $\tilde{\mathbf{E}}_S$ depends linearly on the spherical harmonic coefficients of the induced magnetic field, $\tilde{\mathbf{B}}_{\text{ind}}$, the parameter simulated by Equation (C6), as well as on the input parameters: the horizontal neutral wind, $\tilde{\mathbf{u}}$ and the radial component of the field-aligned current density, $\tilde{\mathbf{j}}_r$. We get

$$\tilde{\mathbf{E}}_S = \tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}} \tilde{\mathbf{B}}_{\text{ind}} + \tilde{\mathbb{E}}_{\mathbf{u}} \tilde{\mathbf{u}} + \tilde{\mathbb{E}}_{j_r} \tilde{\mathbf{j}}_r, \quad (\text{C7})$$

810 where the matrices $\tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}}$, $\tilde{\mathbb{E}}_{\mathbf{u}}$, and $\tilde{\mathbb{E}}_{j_r}$ are given in Equations (C34) – (C36). The coefficients for the radial component of the field-aligned current density is a solution of the linear system

$$\mathbb{G}_{j_r} \tilde{\mathbf{j}}_r = \begin{pmatrix} \mathbf{j}_r^{\text{hl}} \\ \mathbf{0} \end{pmatrix}, \quad (\text{C8})$$

defined in accordance with Equation (C1). In this equation, the upper block of the vector (superscript "hl") refers to grid points at high latitudes, while the lower block refers to grid points at low latitudes, where the hemispheres are connected as described
815 in Section 3.2.1. The grid values $\mathbf{j}_r^{\text{hl}}$ are the user-specified input to the simulation, taken from the AMPS model (Laundal et al., 2018) in the simulations presented in this work. The minimum-norm least-squares solution of the linear system is found by calculating

$$\tilde{\mathbf{j}}_r = \mathbb{G}_{j_r}^\dagger \begin{pmatrix} \mathbf{j}_r^{\text{hl}} \\ \mathbf{0} \end{pmatrix} \quad (\text{C9})$$

where $\mathbb{G}_{j_r}^\dagger$ denotes the Moore-Penrose inverse of $\mathbb{G}_{j_r}$. Similarly, the coefficients for the horizontal neutral winds is a solution
820 of the linear system

$$\mathbb{G}_{\mathbf{u}} \tilde{\mathbf{u}} = \mathbf{u} \quad (\text{C10})$$

defined in accordance with Equation (C2). The grid values $\mathbf{u}$ are the user-specified input, taken from the Horizontal Wind Model (Drob et al., 2015) in the simulations presented in this work. The minimum-norm least-squares solution of the linear system is found by calculating

$$825 \quad \tilde{\mathbf{u}} = \mathbb{G}_{\mathbf{u}}^\dagger \mathbf{u}, \quad (\text{C11})$$

where $\mathbb{G}_{\mathbf{u}}^\dagger$ denotes the Moore-Penrose inverse of $\mathbb{G}_{\mathbf{u}}$.

To obtain final expressions for Faraday's law and the steady-state condition, we note that the surface electric field coefficients can be separated as

$$\tilde{\mathbf{E}}_S = \begin{pmatrix} \tilde{\mathbf{E}}_{\text{cf}} \\ \tilde{\mathbf{E}}_{\text{df}} \end{pmatrix} = \begin{pmatrix} (\tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}} \tilde{\mathbf{B}}_{\text{ind}} + \tilde{\mathbb{E}}_{\mathbf{u}} \tilde{\mathbf{u}} + \tilde{\mathbb{E}}_{j_r} \tilde{\mathbf{j}}_r)_{\text{cf}} \\ (\tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}} \tilde{\mathbf{B}}_{\text{ind}} + \tilde{\mathbb{E}}_{\mathbf{u}} \tilde{\mathbf{u}} + \tilde{\mathbb{E}}_{j_r} \tilde{\mathbf{j}}_r)_{\text{df}} \end{pmatrix} \quad (\text{C12})$$

830 in accordance with Equation (C2). The curl of the curl-free part is by definition zero, implying that we can express $\tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t}$ as

$$\tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t} = \begin{pmatrix} \mathbf{0} & \tilde{\mathbb{T}}_{\mathbf{E}_{\text{df}} \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t} \end{pmatrix}, \quad (\text{C13})$$

where $\tilde{\mathbb{T}}_{\mathbf{E}_{\text{df}} \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t}$ is the diagonal matrix given in Table A1, representing the application of $(-\hat{\mathbf{r}} \times \nabla_S)$ on the divergence-free part of $\mathbf{E}_S$. Inserting Equations (C12) and (C13) into Equation (C6), we obtain the following matrix formulation of Faraday's law,

$$\frac{\partial \tilde{\mathbf{B}}_{\text{ind}}}{\partial t} = \tilde{\mathbb{T}}_{\mathbf{E}_{\text{df}} \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t} (\tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}} \tilde{\mathbf{B}}_{\text{ind}} + \tilde{\mathbb{E}}_{\mathbf{u}} \tilde{\mathbf{u}} + \tilde{\mathbb{E}}_{j_r} \tilde{\mathbf{j}}_r)_{\text{df}}. \quad (\text{C14})$$

When simulating the temporal evolution of the induced magnetic field, we perform a discretized integration of Equation (C14) using Euler's method,

$$\tilde{\mathbf{B}}_{\text{ind},n+1} \approx \tilde{\mathbf{B}}_{\text{ind},n} + \frac{\partial \tilde{\mathbf{B}}_{\text{ind},n}}{\partial t} \Delta t, \quad (\text{C15})$$

840 where we use the time step $\Delta t = 5 \times 10^{-4}$ s for all simulations in this work. The error of this method is of the order $O(\Delta t^2)$.

The steady state of the ionosphere is reached when $\frac{\partial \tilde{\mathbf{B}}_{\text{ind}}}{\partial t} = \mathbf{0}$. Inserting this constraint into Equation (C14), we obtain

$$\tilde{\mathbb{T}}_{\mathbf{E}_{\text{df}} \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t} (\tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}} \tilde{\mathbf{B}}_{\text{ind}} + \tilde{\mathbb{E}}_{\mathbf{u}} \tilde{\mathbf{u}} + \tilde{\mathbb{E}}_{j_r} \tilde{\mathbf{j}}_r)_{\text{df}} = \mathbf{0}. \quad (\text{C16})$$

Since $\tilde{\mathbb{T}}_{\mathbf{E}_{\text{df}} \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t}$ is a diagonal and invertible matrix, we can multiply Equation (C16) from the left by its inverse, giving

$$(\tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}} \tilde{\mathbf{B}}_{\text{ind}})_{\text{df}} = -(\tilde{\mathbb{E}}_{\mathbf{u}} \tilde{\mathbf{u}} + \tilde{\mathbb{E}}_{j_r} \tilde{\mathbf{j}}_r)_{\text{df}}. \quad (\text{C17})$$

845 This linear system can be solved for the steady state $\tilde{\mathbf{B}}_{\text{ind}}$. Note that it is also possible to solve the equation for $\tilde{\mathbf{u}}$ or $\tilde{\mathbf{j}}_r$ given the other vectors, which could give alternative approaches to determining horizontal neutral winds or field-aligned currents experimentally, assuming that the system is in steady state.

Thus, it is possible to find both time-dependent and steady-state solutions for the model parameters $\tilde{\mathbf{B}}_{\text{ind}}$ by solving equations involving conductance-dependent matrices $\tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}}$, $\tilde{\mathbb{E}}_{\mathbf{u}}$ and $\tilde{\mathbb{E}}_{j_r}$, obtained from Equations (C34) – (C36), respectively, and the spherical harmonic coefficient vectors $\tilde{\mathbf{j}}_r$ and $\tilde{\mathbf{u}}$, obtained from Equations (C8) and (C10) with grid-resolved input from external models, respectively.

855 As a side note, Equation (C14) represents a linear and time-invariant (LTI) system during periods when the conductance, field-aligned current, and neutral wind are constant. In such cases, solutions at any time within the period can be computed using matrix exponentials (Chen, 2014), bypassing numerical integration. This approach could improve simulation efficiency, particularly in scenarios where these quantities vary slowly compared to the timescales of induction, a common assumption in ionospheric modeling.

C3 Matrix Formulation of the Generalized Ohm's Law

Here, we derive an expression for linearly constructing the spherical harmonic coefficients of the horizontal electric field from those of the induced magnetic field, neutral winds, and field-aligned currents, while accounting for boundary conditions

860 through the imposed magnetic field. This expression, presented in Equation (C33), is derived from the 2D Generalized Ohm's law in Equation (7), which can be discretized in terms of grid points and spherical harmonics as

$$\mathbb{G}_{\mathbf{E}_S} \tilde{\mathbf{E}}_S = \mathbb{T}_{\mathbf{J}_S \mapsto \mathbf{E}_S} \mathbb{G}_{\mathbf{J}_S} \tilde{\mathbb{T}}_{\mathbf{B} \mapsto \mathbf{J}_S} \tilde{\mathbf{B}} + \mathbb{T}_{\mathbf{u} \mapsto \mathbf{E}_S} \mathbb{G}_{\mathbf{u}} \tilde{\mathbf{u}}, \quad (\text{C18})$$

where $\tilde{\mathbf{B}}$, $\tilde{\mathbf{E}}_S$ and $\tilde{\mathbf{u}}$ are vectors containing the spherical harmonic coefficients of the magnetic field, horizontal electric field and horizontal neutral wind, respectively. $\tilde{\mathbb{T}}_{\mathbf{B} \mapsto \mathbf{J}_S}$ is the representation of the jump condition $\mathbf{J}_S = \frac{1}{\mu_0} \hat{\mathbf{r}} \times \Delta \mathbf{B}$ in the basis of
865 the spherical harmonics used for representing $\mathbf{B}$. The evaluation matrices $\mathbb{G}_{\mathbf{E}_S}$, $\mathbb{G}_{\mathbf{J}_S}$ and $\mathbb{G}_{\mathbf{u}}$ evaluate the spherical harmonic expansions of the horizontal electric field, current and neutral wind, respectively, on a common set of grid points in accordance with Equation (C2). In the simulations presented in this work, we use a cubed-sphere grid with 100×100 grid points in each of the six cubed-sphere blocks. $\mathbb{T}_{\mathbf{J}_S \mapsto \mathbf{E}_S} = \mathbb{T}_{\mathbf{J}_S \mapsto \mathbf{E}_S}(\mathbf{B}_0, \eta_P, \eta_H)$ and $\mathbb{T}_{\mathbf{u} \mapsto \mathbf{E}_S} = \mathbb{T}_{\mathbf{u} \mapsto \mathbf{E}_S}(\mathbf{B}_0)$ represent the current-dependent and neutral wind-dependent terms in the 2D Generalized Ohm's law, respectively, expressed on the same grid points.

870 We can obtain the minimum-norm least-squares solution for the spherical harmonic coefficients $\tilde{\mathbf{E}}_S$ by multiplying Equation (C18) from the left by the Moore-Penrose inverse of $\mathbb{G}_{\mathbf{E}_S}$, denoted by $\mathbb{G}_{\mathbf{E}_S}^\dagger$, giving

$$\tilde{\mathbf{E}}_S = \mathbb{G}_{\mathbf{E}_S}^\dagger \mathbb{T}_{\mathbf{J}_S \mapsto \mathbf{E}_S} \mathbb{G}_{\mathbf{J}_S} \tilde{\mathbb{T}}_{\mathbf{B} \mapsto \mathbf{J}_S} \tilde{\mathbf{B}} + \mathbb{G}_{\mathbf{E}_S}^\dagger \mathbb{T}_{\mathbf{u} \mapsto \mathbf{E}_S} \mathbb{G}_{\mathbf{u}} \tilde{\mathbf{u}}. \quad (\text{C19})$$

Continuing, we decompose the magnetic field $\mathbf{B}$ into the sum of its induced part $\mathbf{B}_{\text{ind}}$ and its imposed part $\mathbf{B}_{\text{imp}}$, in accordance with Section 3, implying that $\tilde{\mathbf{B}}$ can be written as

$$875 \quad \tilde{\mathbf{B}} = \begin{pmatrix} \tilde{\mathbf{B}}_{\text{ind}} \\ \tilde{\mathbf{B}}_{\text{imp}} \end{pmatrix}. \quad (\text{C20})$$

Similarly, the matrix evaluating $\mathbf{J}_S$ from the model vector $\tilde{\mathbf{B}}$ can be separated into blocks referencing the induced and imposed parts of the magnetic field,

$$\tilde{\mathbb{T}}_{\mathbf{B} \mapsto \mathbf{J}_S} = \begin{pmatrix} \tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{J}_S} & \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{J}_S} \end{pmatrix}. \quad (\text{C21})$$

where the spherical harmonic transformation matrices $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{J}_S}$ and $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{J}_S}$ are given in Table A1. The matrix $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{J}_S}$
880 includes the contribution to the poloidal field coming from the toroidal $\mathbf{B}_{\text{imp}}$, representing the Biot-Savart integral of the horizontal part of the field-aligned currents above the ionosphere, as discussed in Section 3.2.2. The discretization procedure is discussed in Section C4. Inserting Equations (C20) and (C21) into Equation (C19), we obtain

$$\tilde{\mathbf{E}}_S = \tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{E}_S} \tilde{\mathbf{B}}_{\text{ind}} + \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{E}_S} \tilde{\mathbf{B}}_{\text{imp}} + \mathbb{T}_{\mathbf{u} \mapsto \mathbf{E}_S} \tilde{\mathbf{u}}, \quad (\text{C22})$$

where the spherical harmonic coefficient transformation matrices $\tilde{\mathbb{T}}_{X \mapsto \mathbf{E}_S}$ are given in Table A1

885 $\tilde{\mathbf{B}}_{\text{imp}}$ depends on the specified field-aligned currents at high latitudes and on the constraints that the current density and electric field map between hemispheres at low latitudes, as discussed in Section 3.2.1. These three constraints, which together specify $\tilde{\mathbf{B}}_{\text{imp}}$, can be expressed by a linear system of equations,

$$\mathbb{G}_{\mathbf{c}} \tilde{\mathbf{B}}_{\text{imp}} = \mathbf{c} \quad (\text{C23})$$

which can be solved for $\tilde{\mathbf{B}}_{\text{imp}}$ at each time step. In the following, we go through the details of how this is done, and how this
 890 expression can be used to eliminate $\tilde{\mathbf{B}}_{\text{imp}}$ from Equation (C22). The forward operator in Equation (C23) can be written as

$$\mathbb{G}_{\mathbf{c}} = \mathbb{A} \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto j_r, \mathbf{E}_S}, \quad (\text{C24})$$

where the matrix transforming from $\tilde{\mathbf{B}}_{\text{imp}}$ to corresponding spherical harmonic coefficients for the radial component of the field-aligned current density (j_r) and for the horizontal electric field ($\mathbf{E}_S$) is given by

$$\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto j_r, \mathbf{E}_S} = \begin{pmatrix} \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto j_r} \\ \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{E}_S} \end{pmatrix}, \quad (\text{C25})$$

895 and $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto j_r}$ and $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{E}_S}$ are given in Table A1. The matrix $\mathbb{A}$ can be written as

$$\mathbb{A} = \begin{pmatrix} \mathbb{A}_{j_r}^{\text{hl}} & \mathbf{0} \\ \mathbb{A}_{j_r}^{\text{ll}} - \mathbb{A}_{j_r}^{\text{ll,cp}} & \mathbf{0} \\ \mathbf{0} & \mathbb{A}_{\mathbf{E}_S}^{\text{ll}} - \mathbb{A}_{\mathbf{E}_S}^{\text{ll,cp}} \end{pmatrix}, \quad (\text{C26})$$

where the top part represents the constraint that $\tilde{\mathbf{B}}_{\text{imp}}$ should describe the given high-latitude (superscript "hl" refers to high-latitude grid points) FAC, the middle part represents current continuity between hemispheres at low latitudes (superscript "ll" refers to low-latitude grid points and superscript "ll,cp" refers to their conjugate points), and the bottom part represents the
 900 mapping of electric fields between hemispheres at low latitudes.

$\mathbb{A}_{j_r}^{\text{ll}}$ and $\mathbb{A}_{j_r}^{\text{hl}}$ evaluate the spherical harmonic expansion of j_r in accordance with Equation (C1) at the high- and low-latitude grid points immediately above the ionosphere, respectively, construct the corresponding $\mathbf{j}$ vectors from the force-free condition

$$\mathbf{j} \times \mathbf{B}_0 = 0 \quad \Rightarrow \quad \begin{aligned} j_\theta &= j_r b_\theta / b_r, \\ j_\phi &= j_r b_\phi / b_r, \end{aligned} \quad (\text{C27})$$

905 and project these vectors onto the field-aligned Modified Apex unit vectors $\hat{\mathbf{d}}_3$ (defined in Richmond (1995)) at the same grid points, giving Modified Apex components J_{e3} that are constant along the main field lines (Richmond, 1995). $\mathbb{A}_{j_r}^{\text{ll,cp}}$ performs the same operation, but for the field-conjugated low-latitude grid points, which are determined by tracing the main field lines that intersect the low-latitude grid points outward until they intersect the targeted points on the opposite hemisphere of the ionospheric shell. In practice, we find the conjugate points using Modified Apex coordinate conversions (van der Meeren et al.,
 910 2025; Emmert et al., 2010). The term $\mathbb{A}_{\mathbf{E}_S}^{\text{ll}}$ performs the evaluation of the spherical harmonic expansion of $\mathbf{E}_S = E_\theta \hat{\boldsymbol{\theta}} + E_\phi \hat{\boldsymbol{\phi}}$ (projection of $\mathbf{E}$ onto $\hat{\boldsymbol{\theta}}$ and $\hat{\boldsymbol{\phi}}$) at low-latitude grid points immediately above the ionosphere in accordance with Equation (C2), retrieves the third (radial) component of the corresponding $\mathbf{E}$ vectors from the ideal MHD condition

$$\mathbf{E} \cdot \mathbf{B}_0 = 0 \quad \Rightarrow \quad E_r = -(E_\theta b_\theta + E_\phi b_\phi) / b_r, \quad (\text{C28})$$

and projects these vectors onto the field-perpendicular Modified Apex unit vectors $\hat{\mathbf{e}}_1$ and $\hat{\mathbf{e}}_2$ (see Richmond (1995)) at the
 915 same grid points, giving Modified Apex components E_{d1} and E_{d2} that are constant along the main field lines. The matrix $\mathbb{A}_{\mathbf{E}_S}^{\text{ll,cp}}$ performs the same operation, but for the field-conjugated low-latitude grid points.

The constraint vector $\mathbf{c}$ can be written as

$$\mathbf{c} = \mathbb{A} \begin{pmatrix} \tilde{\mathbf{j}}_r \\ -\tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{E}_S} \tilde{\mathbf{B}}_{\text{ind}} - \tilde{\mathbb{T}}_{\mathbf{u} \mapsto \mathbf{E}_S} \tilde{\mathbf{u}} \end{pmatrix}. \quad (\text{C29})$$

The spherical harmonic coefficient vectors $\tilde{\mathbf{j}}_r$ and $\tilde{\mathbf{u}}$ are found from Equations (C8) and (C10), respectively. $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{E}_S}$ and $\tilde{\mathbb{T}}_{\mathbf{u} \mapsto \mathbf{E}_S}$ relate the spherical harmonic coefficients for the induced magnetic field and horizontal neutral wind, respectively, to the corresponding coefficients for the horizontal electric field, and are given in Table A1. $\mathbb{A}$ is given in Equation (C26). The negative sign ensures that the imposed field cancels any difference between the conjugate point E_{d1} and E_{d2} Modified Apex components of the electric field at low latitudes, ensuring that $(\mathbb{A}_{\mathbf{E}_S}^{\text{ll}} - \mathbb{A}_{\mathbf{E}_S}^{\text{ll, cp}}) \tilde{\mathbf{E}}_S = \mathbf{0}$ for the total electric field coefficients $\tilde{\mathbf{E}}_S$.

We can obtain the minimum-norm least-squares solution for the spherical harmonic coefficients $\tilde{\mathbf{B}}_{\text{imp}}$ by multiplying Equation (C23) from the left by the Moore-Penrose inverse of $\mathbb{G}_{\mathbf{c}}$, denoted by $\mathbb{G}_{\mathbf{c}}^\dagger$, giving

$$\begin{aligned} \tilde{\mathbf{B}}_{\text{imp}} &= \mathbb{G}_{\mathbf{c}}^\dagger \mathbf{c} \\ &= \mathbb{G}_{\mathbf{c}}^\dagger \mathbb{A} \begin{pmatrix} \tilde{\mathbf{j}}_r \\ -\tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{E}_S} \tilde{\mathbf{B}}_{\text{ind}} - \tilde{\mathbb{T}}_{\mathbf{u} \mapsto \mathbf{E}_S} \tilde{\mathbf{u}} \end{pmatrix}. \end{aligned} \quad (\text{C30})$$

Now, we can represent the matrix product $\mathbb{G}_{\mathbf{c}}^\dagger \mathbb{A}$ in terms of the block that references the $\tilde{\mathbf{j}}_r$ coefficients and the one that references the $\tilde{\mathbf{E}}_S$ coefficients, and name them $\tilde{\mathbb{T}}_{j_r \mapsto \mathbf{B}_{\text{imp}}}$ and $\tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \mathbf{B}_{\text{imp}}}$, respectively, giving

$$\mathbb{G}_{\mathbf{c}}^\dagger \mathbb{A} = \begin{pmatrix} \tilde{\mathbb{T}}_{j_r \mapsto \mathbf{B}_{\text{imp}}} & \tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \mathbf{B}_{\text{imp}}} \end{pmatrix}. \quad (\text{C31})$$

This implies that Equation (C30) can be written as

$$\tilde{\mathbf{B}}_{\text{imp}} = \tilde{\mathbb{T}}_{j_r \mapsto \mathbf{B}_{\text{imp}}} \tilde{\mathbf{j}}_r - \tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \mathbf{B}_{\text{imp}}} \tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{E}_S} \tilde{\mathbf{B}}_{\text{ind}} - \tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \mathbf{B}_{\text{imp}}} \tilde{\mathbb{T}}_{\mathbf{u} \mapsto \mathbf{E}_S} \tilde{\mathbf{u}}. \quad (\text{C32})$$

Inserting this expression into Equation (C22), we can write $\tilde{\mathbf{E}}_S$ as

$$\tilde{\mathbf{E}}_S = \tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}} \tilde{\mathbf{B}}_{\text{ind}} + \tilde{\mathbb{E}}_{\mathbf{u}} \tilde{\mathbf{u}} + \tilde{\mathbb{E}}_{j_r} \tilde{\mathbf{j}}_r, \quad (\text{C33})$$

where the matrices giving the different contributions to the horizontal electric field are given by

$$\tilde{\mathbb{E}}_{\mathbf{B}_{\text{ind}}} = (\mathbb{1} - \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{E}_S} \tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \mathbf{B}_{\text{imp}}}) \tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{E}_S}, \quad (\text{C34})$$

$$\tilde{\mathbb{E}}_{\mathbf{u}} = (\mathbb{1} - \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{E}_S} \tilde{\mathbb{T}}_{\mathbf{E}_S \mapsto \mathbf{B}_{\text{imp}}}) \tilde{\mathbb{T}}_{\mathbf{u} \mapsto \mathbf{E}_S}, \quad (\text{C35})$$

$$\tilde{\mathbb{E}}_{j_r} = \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{E}_S} \tilde{\mathbb{T}}_{j_r \mapsto \mathbf{B}_{\text{imp}}}. \quad (\text{C36})$$

The spherical harmonic coefficient transformation matrices $\tilde{\mathbb{T}}_{X \mapsto \mathbf{E}_S}$ are given in Table A1 and the matrices $\tilde{\mathbb{T}}_{X \mapsto \mathbf{B}_{\text{imp}}}$ are, as illustrated in Equation (C31), blocks of the product of the Moore-Penrose inverse of $\mathbb{G}_{\mathbf{c}}$ with $\mathbb{A}$, which are specified in Equations (C24) – (C26). Note that the coefficients of the neutral winds and of the induced magnetic field contribute to the total $\tilde{\mathbf{E}}_S$ both directly and through the imposed magnetic field, while the coefficients of the radial component of the field-aligned current density contribute only through the imposed magnetic field.

C4 Matrix Formulation of the Poloidal Magnetic Field of Field-Aligned Currents

In the numerical implementation, we represent Equation (34), based on Engels and Olsen (1998), as a matrix $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{P}^e}$ that
 945 transforms vectors of toroidal spherical harmonic coefficients to corresponding vectors of spherical harmonic coefficients of external poloidal fields. This is done by performing the discretized integration of poloidal magnetic field contributions from the horizontal components of the field-aligned currents at $r > R$ via the midpoint rule

$$\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{P}^e} = -\Delta r \sum_{i=1}^N \tilde{\mathbb{T}}_{\mathbf{P}^e(r_i) \mapsto \mathbf{P}^e} \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{P}^e(r_i)} \quad (\text{C37})$$

where the $r_i = R + \Delta r/2$ are the integration radii. The matrix

$$950 \quad \tilde{\mathbb{T}}_{\mathbf{P}^e(r_i) \mapsto \mathbf{P}^e} = \text{diag} \left[\left(\frac{R}{r_i} \right)^{n-1} \right] \quad (\text{C38})$$

represents the conversion of the spherical harmonic coefficients of an external poloidal field at radius r_i to radius R . The matrix $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{P}^e(r_i)}$ constructs the external poloidal field associated with the horizontal components of the field-aligned current density at r_i , and can be written as

$$\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{P}^e(r_i)} = \tilde{\mathbb{T}}_{\mathbf{j}_h(r_i) \mapsto \mathbf{P}^e(r_i)} \mathbb{G}_{\mathbf{j}_h(r_i)}^\dagger \mathbb{T}_{j_r \mapsto \mathbf{j}_h(r_i)} \mathbb{G}_{j_r} \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto j_r}. \quad (\text{C39})$$

955 $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto j_r}$ transforms spherical harmonic coefficients of the imposed magnetic field to coefficients for the scalar j_r immediately above the ionosphere, and is given in Table A1. $\mathbb{G}_{j_r}$ evaluates the spherical harmonic expansion of j_r at cubed-sphere grid points according to Equation (C1). In this work, we use the same $6 \times 100 \times 100$ cubed-sphere grid for this calculation as used when calculating the matrices in Section C3. $\mathbb{T}_{j_r \mapsto \mathbf{j}_h(r_i)}$ represents the mapping of the grid-evaluated j_r to the corresponding grid-evaluated $\mathbf{j}_h$ at $r = r_i$ via Equation (36). The Moore-Penrose inverse $\mathbb{G}_{\mathbf{j}_h(r_i)}^\dagger$ performs a minimum-norm least-squares
 960 spherical harmonic decomposition of the horizontal current density at the mapped grid points at $r = r_i$ according to Equation (C2), and $\tilde{\mathbb{T}}_{\mathbf{j}_h(r_i) \mapsto \mathbf{P}^e(r_i)} = \tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{J}_S}^{-1}$ converts the coefficients into corresponding coefficients of an external poloidal magnetic field with r_i as the reference radius, where $\tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{J}_S}$ is given in Table A1.

Table A1. A summary of the matrices transforming between different spherical harmonic vectors. The integer n is the spherical harmonic degree, μ_0 is the vacuum permeability, and R is the ionospheric radius. The matrix $\tilde{\mathbb{T}}_{\nabla_S^2}$ is a spherical harmonic representation of the surface Laplacian, and the matrix $\tilde{\mathbb{T}}_{\mathbf{P}^e \mapsto \Delta \mathbf{P}}$ represents the mapping of the coefficients of external poloidal fields to the corresponding potential difference across the 2D ionosphere. The matrices $\mathbb{G}_{\mathbf{J}_S}$, $\mathbb{G}_{\mathbf{E}_S}$ and $\mathbb{G}_{\mathbf{u}}$ evaluate the spherical harmonic representations of the horizontal current, electric field and wind, respectively, at a given set of grid points, in accordance with Equation (C2). $\mathbb{T}_{\mathbf{J}_S \mapsto \mathbf{E}_S}$ and $\mathbb{T}_{\mathbf{u} \mapsto \mathbf{E}_S}$ are the horizontal wind- and current-dependent terms in the 2D Generalized Ohm's law, respectively, discretized on the same grid points.

$\tilde{\mathbb{T}}_{\mathbf{E}_{\text{df}} \mapsto \partial \mathbf{B}_{\text{ind}} / \partial t}$	$\frac{1}{R} \mathbb{1}$
$\tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto B_r}$	$-R^2 \tilde{\mathbb{T}}_{\nabla_S^2}$
$\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto j_r}$	$\frac{R}{\mu_0} \tilde{\mathbb{T}}_{\nabla_S^2}$
$\tilde{\mathbb{T}}_{\mathbf{P}^e \mapsto \mathbf{J}_{\text{df}}}$	$-\frac{R}{\mu_0} \tilde{\mathbb{T}}_{\mathbf{P}^e \mapsto \Delta \mathbf{P}}$
$\tilde{\mathbb{T}}_{\mathbf{T} \mapsto \mathbf{J}_{\text{cf}}}$	$\frac{R}{\mu_0} \mathbb{1}$
$\tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{J}_S}$	$\begin{pmatrix} \mathbf{0} \\ \tilde{\mathbb{T}}_{\mathbf{P}^e \mapsto \mathbf{J}_{\text{df}}} \end{pmatrix}$
$\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{J}_S}$	$\begin{pmatrix} \tilde{\mathbb{T}}_{\mathbf{T} \mapsto \mathbf{J}_{\text{cf}}} \\ \tilde{\mathbb{T}}_{\mathbf{P}^e \mapsto \mathbf{J}_{\text{df}}} \tilde{\mathbb{T}}_{\mathbf{T} \mapsto \mathbf{P}^e} \end{pmatrix}$
$\tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{E}_S}$	$\mathbb{G}_{\mathbf{E}_S}^\dagger \mathbb{T}_{\mathbf{J}_S \mapsto \mathbf{E}_S} \mathbb{G}_{\mathbf{J}_S} \tilde{\mathbb{T}}_{\mathbf{B}_{\text{ind}} \mapsto \mathbf{J}_S}$
$\tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{E}_S}$	$\mathbb{G}_{\mathbf{E}_S}^\dagger \mathbb{T}_{\mathbf{J}_S \mapsto \mathbf{E}_S} \mathbb{G}_{\mathbf{J}_S} \tilde{\mathbb{T}}_{\mathbf{B}_{\text{imp}} \mapsto \mathbf{J}_S}$
$\tilde{\mathbb{T}}_{\mathbf{u} \mapsto \mathbf{E}_S}$	$\mathbb{G}_{\mathbf{E}_S}^\dagger \mathbb{T}_{\mathbf{u} \mapsto \mathbf{E}_S} \mathbb{G}_{\mathbf{u}}$
$\tilde{\mathbb{T}}_{\nabla_S^2}$	$\text{diag}\left(-\frac{n(n+1)}{R^2}\right)$
$\tilde{\mathbb{T}}_{\mathbf{P}^e \mapsto \Delta \mathbf{P}}$	$\text{diag}(2n+1)$

Author contributions. KML conceived and designed the concept, wrote simulation and visualization software, and drafted the manuscript except for Appendix B. ASS led the development of the simulation software and drafted Appendix B. ASS, BPB, and SMH contributed to the model concept through active discussions and significantly revised the manuscript. HV helped inspire the original idea and significantly revised the manuscript. NO conceived a critical component of the model and significantly revised the manuscript. All authors approved the submitted version of the manuscript.

Competing interests. The authors declare no conflicts of interest relevant to this study.

Acknowledgements. We thank Michael Madelaire and Fasil Tesema Kebede for their constructive input during the development of this article. We thank Lotte van Hazendonk at the University Centre in Svalbard for providing the EISCAT plasma density height profile used in Figure 1. We thank Stephan Buchert, Slava Merkin, and Kareem Sorathia for their inspiring and insightful discussions, which have contributed to the development of this work. [We thank the reviewers for their careful reading and valuable feedback, which helped strengthen the manuscript.](#)

This work was funded by the European Union (ERC, DynaMIT, 101086985). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. The work was also funded by the Research Council of Norway, contract and 300844/F50, and by the Trond Mohn Research Foundation. The work of HV was supported by the Research Council of Finland project 354521.

References

- Alken, P.: Observations and modeling of the ionospheric gravity and diamagnetic current systems from CHAMP and Swarm measurements,
980 Journal of Geophysical Research: Space Physics, 121, 589–601, <https://doi.org/10.1002/2015JA022163>, 2016.
- Alken, P., Thébault, E., Beggan, C. D., Amit, H., Aubert, J., Baerenzung, J., Bondar, T. N., Brown, W. J., Califf, S., Chambodut, A., Chulliat, A., Cox, G. A., Finlay, C. C., Fournier, A., Gillet, N., Grayver, A., Hammer, M. D., Holschneider, M., Huder, L., Hulot, G., Jager, T., Kloss, C., Korte, M., Kuang, W., Kuvshinov, A., Langlais, B., Léger, J.-M., Lesur, V., Livermore, P. W., Lowes, F. J., Macmillan, S., Magnes, W., Manda, M., Marsal, S., Matzka, J., Metman, M. C., Minami, T., Morschhauser, A., Mound, J. E., Nair, M., Nakano, S.,
985 Olsen, N., Pavón-Carrasco, F. J., Petrov, V. G., Ropp, G., Rother, M., Sabaka, T. J., Sanchez, S., Saturnino, D., Schnepf, N. R., Shen, X., Stolle, C., Tangborn, A., Tøffner-Clausen, L., Toh, H., Torta, J. M., Varner, J., Vervelidou, F., Vigneron, P., Wardinski, I., Wicht, J., Woods, A., Yang, Y., Zeren, Z., and Zhou, B.: International Geomagnetic Reference Field: the thirteenth generation, Earth, Planets and Space, 73, 49, <https://doi.org/10.1186/s40623-020-01288-x>, 2021.
- Amm, O.: Comment on “A three-dimensional, iterative mapping procedure for the implementation of an ionosphere–magnetosphere
990 anisotropic Ohm’s law boundary condition in global magnetohydrodynamic simulations” by Michael L. Goodman, Annales Geophysicae, 14, 773–774, letter to the Editor, 1996.
- Amm, O.: Ionospheric Elementary Current Systems in Spherical Coordinates and Their Application., Journal of geomagnetism and geoelectricity, 49, 947–955, <https://doi.org/10.5636/jgg.49.947>, 1997.
- Backus, G.: Poloidal and toroidal fields in geomagnetic field modeling, Reviews of Geophysics, 24, 75–109,
995 <https://doi.org/10.1029/RG024i001p00075>, 1986.
- Brekke, A.: Physics of the Upper Polar Atmosphere, Springer Atmospheric Sciences, Springer Berlin Heidelberg, Berlin, Heidelberg, ISBN 978-3-642-27400-8 978-3-642-27401-5, <https://doi.org/10.1007/978-3-642-27401-5>, 2013.
- Buchert, S. C.: Entangled dynamos and Joule heating in the Earth’s ionosphere, Annales Geophysicae, 38, 1019–1030, <https://doi.org/10.5194/angeo-38-1019-2020>, 2020.
- 1000 Buchert, S. C. and Budnik, F.: Field-aligned current distributions generated by a divergent Hall current, Geophysical Research Letters, 24, 297–300, <https://doi.org/10.1029/97GL00064>, 1997.
- Chen, C.-T.: Linear system theory and design, Oxford University Press, New York, international 4. ed edn., ISBN 978-0-19-996454-3, 2014.
- Dreher, J.: On the self-consistent description of dynamic magnetosphere-ionosphere coupling phenomena with resolved ionosphere, Journal of Geophysical Research: Space Physics, 102, 85–94, <https://doi.org/10.1029/96JA02800>, 1997.
- 1005 Drob, D. P., Emmert, J. T., Meriwether, J. W., Makela, J. J., Doornbos, E., Conde, M., Hernandez, G., Noto, J., Zawdie, K. A., McDonald, S. E., Huba, J. D., and Klenzing, J. H.: An update to the Horizontal Wind Model (HWM): The quiet time thermosphere, Earth and Space Science, 2, 301–319, <https://doi.org/10.1002/2014EA000089>, 2015.
- Emmert, J. T., Richmond, A. D., and Drob, D. P.: A computationally compact representation of Magnetic-Apex and Quasi-Dipole coordinates with smooth base vectors, Journal of Geophysical Research: Space Physics, 115, 2010JA015 326, <https://doi.org/10.1029/2010JA015326>,
1010 2010.
- Engels, U. and Olsen, N.: Computation of magnetic fields within source regions of ionospheric and magnetospheric currents, Journal of Atmospheric and Solar-Terrestrial Physics, 60, 1585–1592, [https://doi.org/10.1016/S1364-6826\(98\)00094-7](https://doi.org/10.1016/S1364-6826(98)00094-7), 1998.

- Fillion, M., Hulot, G., Alken, P., and Chulliat, A.: Modeling the Climatology of Low- and Mid-Latitude F-Region Ionospheric Currents Using the Swarm Constellation, *Journal of Geophysical Research: Space Physics*, 128, e2023JA031344, <https://doi.org/10.1029/2023JA031344>, 2023.
- Fukushima, N.: Generalized theorem of no ground magnetic effect of vertical current connected with Pedersen currents in the uniform conductivity ionosphere, *Rep. Ionos. Space Res. Jpn*, 30, 35–40, 1976.
- Ganse, U., Pfau-Kempf, Y., Zhou, H., Juusola, L., Workayehu, A., Kebede, F., Papadakis, K., Grandin, M., Alho, M., Battarbee, M., Dubart, M., Kotipalo, L., Lalagüe, A., Suni, J., Horaites, K., and Palmroth, M.: The Vlasiator 5.2 ionosphere – coupling a magnetospheric hybrid-Vlasov simulation with a height-integrated ionosphere model, *Geoscientific Model Development*, 18, 511–527, <https://doi.org/10.5194/gmd-18-511-2025>, 2025.
- Grayver, A. V., Kuvshinov, A., and Werthmüller, D.: Time-Domain Modeling of Three-Dimensional Earth’s and Planetary Electromagnetic Induction Effect in Ground and Satellite Observations, *Journal of Geophysical Research: Space Physics*, 126, e2020JA028672, <https://doi.org/10.1029/2020JA028672>, 2021.
- Hardy, D. A., Gussenhoven, M. S., Raistrick, R., and McNeil, W. J.: Statistical and functional representations of the pattern of auroral energy flux, number flux, and conductivity, *Journal of Geophysical Research: Space Physics*, 92, 12275–12294, <https://doi.org/10.1029/JA092iA11p12275>, 1987.
- Hatch, S. M., Burchill, J. K., Vanhamäki, H., de Mesquita, R. L. A., Laundal, K., and Knudsen, D. J.: What is the neutral wind in height-integrated ionospheric electrodynamics?, <https://doi.org/10.22541/essoar.171472681.15570950/v1>, 2024.
- Hesse, M., Birn, J., and Hoffman, R. A.: On the mapping of ionospheric convection into the magnetosphere, *Journal of Geophysical Research: Space Physics*, 102, 9543–9551, <https://doi.org/10.1029/96JA03999>, 1997.
- Huba, J. D., Joyce, G., and Krall, J.: Three-dimensional equatorial spread *F* modeling, *Geophysical Research Letters*, 35, 2008GL033509, <https://doi.org/10.1029/2008GL033509>, 2008.
- Ilma, R.: rilma/pyHWM14: Official release of the HWM14 wrapper in Python, <https://doi.org/10.5281/ZENODO.240890>, 2017.
- Juusola, L., Kauristie, K., Vanhamäki, H., Aikio, A., and Van De Kamp, M.: Comparison of auroral ionospheric and field-aligned currents derived from Swarm and ground magnetic field measurements, *Journal of Geophysical Research: Space Physics*, 121, 9256–9283, <https://doi.org/10.1002/2016JA022961>, 2016.
- Juusola, L., Vanhamäki, H., Viljanen, A., and Smirnov, M.: Induced currents due to 3D ground conductivity play a major role in the interpretation of geomagnetic variations, *Annales Geophysicae*, 38, 983–998, <https://doi.org/10.5194/angeo-38-983-2020>, 2020.
- Juusola, L., Vanhamäki, H., Marshalko, E., Kruglyakov, M., and Viljanen, A.: Estimation of the 3-D geoelectric field at the Earth’s surface using Spherical Elementary Current Systems, <https://doi.org/10.5194/egusphere-2024-2831>, 2024.
- Kamide, Y., Richmond, A. D., and Matsushita, S.: Estimation of ionospheric electric fields, ionospheric currents, and field-aligned currents from ground magnetic records, *Journal of Geophysical Research: Space Physics*, 86, 801–813, <https://doi.org/10.1029/JA086iA02p00801>, 1981.
- Kelley, M. C.: The earth’s ionosphere: plasma physics and electrodynamics, no. v. 96 in International geophysics series, Academic Press, Amsterdam Boston, 2nd ed edn., ISBN 978-0-12-088425-4 978-0-08-091657-6, 2009.
- Laundal, K. M. and Richmond, A. D.: Magnetic Coordinate Systems, *Space Science Reviews*, 206, 27–59, <https://doi.org/10.1007/s11214-016-0275-y>, 2017.

- Laundal, K. M., Haaland, S. E., Lehtinen, N., Gjerloev, J. W., Østgaard, N., Tenfjord, P., Reistad, J. P., Snekvik, K., Milan, S. E., Ohtani, S., and Anderson, B. J.: Birkeland current effects on high-latitude ground magnetic field perturbations, *Geophysical Research Letters*, 42, 7248–7254, <https://doi.org/10.1002/2015GL065776>, 2015.
- Laundal, K. M., Finlay, C. C., and Olsen, N.: Sunlight effects on the 3D polar current system determined from low Earth orbit measurements, *Earth, Planets and Space*, 68, 142, <https://doi.org/10.1186/s40623-016-0518-x>, 2016.
- Laundal, K. M., Finlay, C. C., Olsen, N., and Reistad, J. P.: Solar Wind and Seasonal Influence on Ionospheric Currents From Swarm and CHAMP Measurements, *Journal of Geophysical Research: Space Physics*, 123, 4402–4429, <https://doi.org/10.1029/2018JA025387>, 2018.
- Laundal, K. M., Hatch, S. M., and Moretto, T.: Magnetic Effects of Plasma Pressure Gradients in the Upper *F* Region, *Geophysical Research Letters*, 46, 2355–2363, <https://doi.org/10.1029/2019GL081980>, 2019.
- Laundal, K. M., Madelaire, M., Ohma, A., Reistad, J., and Hatch, S.: The relationship between interhemispheric asymmetries in polar ionospheric convection and the magnetic field line footpoint displacement field, *Frontiers in Astronomy and Space Sciences*, 9, 957 223, <https://doi.org/10.3389/fspas.2022.957223>, 2022a.
- Laundal, K. M., Reistad, J. P., Hatch, S. M., Madelaire, M., Walker, S., Hovland, A. O., Ohma, A., Merkin, V. G., and Sorathia, K. A.: Local Mapping of Polar Ionospheric Electrodynamics, *Journal of Geophysical Research: Space Physics*, 127, e2022JA030 356, <https://doi.org/10.1029/2022JA030356>, 2022b.
- Laundal, K. M., Hatch, S. M., Reistad, J. P., Ohma, A., Tenfjord, P., and Madelaire, M.: How the Ionosphere Responds Dynamically to Magnetospheric Forcing, *Geophysical Research Letters*, 51, e2024GL108 695, <https://doi.org/10.1029/2024GL108695>, 2024.
- Leake, J. E., DeVore, C. R., Thayer, J. P., Burns, A. G., Crowley, G., Gilbert, H. R., Huba, J. D., Krall, J., Linton, M. G., Lukin, V. S., and Wang, W.: Ionized Plasma and Neutral Gas Coupling in the Sun’s Chromosphere and Earth’s Ionosphere/Thermosphere, *Space Science Reviews*, 184, 107–172, <https://doi.org/10.1007/s11214-014-0103-1>, 2014.
- Lotko, W.: Inductive magnetosphere–ionosphere coupling, *Journal of Atmospheric and Solar-Terrestrial Physics*, 66, 1443–1456, <https://doi.org/10.1016/j.jastp.2004.03.027>, 2004.
- Lysak, R. and Song, Y.: Magnetosphere–ionosphere coupling by Alfvén waves: Beyond current continuity, *Advances in Space Research*, 38, 1713–1719, <https://doi.org/10.1016/j.asr.2005.08.038>, 2006.
- Lysak, R. L.: Magnetosphere-ionosphere coupling by Alfvén waves at midlatitudes, *Journal of Geophysical Research: Space Physics*, 109, 2004JA010 454, <https://doi.org/10.1029/2004JA010454>, 2004.
- Lysak, R. L., Song, Y., Waters, C. L., Sciffer, M. D., and Obana, Y.: Numerical Investigations of Interhemispheric Asymmetry due to Ionospheric Conductance, *Journal of Geophysical Research: Space Physics*, 125, e2020JA027 866, <https://doi.org/10.1029/2020JA027866>, 2020.
- Madelaire, M., Laundal, K. M., Reistad, J. P., Hatch, S. M., and Ohma, A.: Transient high latitude geomagnetic response to rapid increases in solar wind dynamic pressure, *Frontiers in Astronomy and Space Sciences*, 9, 953 954, <https://doi.org/10.3389/fspas.2022.953954>, 2022.
- Madelaire, M., Laundal, K., Hatch, S., Vanhamäki, H., Reistad, J., Ohma, A., Merkin, V., and Lin, D.: Estimating the Ionospheric Induction Electric Field Using Ground Magnetometers, *Geophysical Research Letters*, 51, e2023GL105 443, <https://doi.org/10.1029/2023GL105443>, 2024.
- Maute, A., Richmond, A. D., Lu, G., Knipp, D. J., Shi, Y., and Anderson, B.: Magnetosphere-Ionosphere Coupling via Prescribed Field-Aligned Current Simulated by the TIEGCM, *Journal of Geophysical Research: Space Physics*, 126, e2020JA028 665, <https://doi.org/10.1029/2020JA028665>, 2021.

- Merkin, V. G. and Lyon, J. G.: Effects of the low-latitude ionospheric boundary condition on the global magnetosphere, *Journal of Geophysical Research: Space Physics*, 115, 2010JA015461, <https://doi.org/10.1029/2010JA015461>, 2010.
- Moen, J. and Brekke, A.: The solar flux influence on quiet time conductances in the auroral ionosphere, *Geophysical Research Letters*, 20, 971–974, <https://doi.org/10.1029/92GL02109>, 1993.
- 1090 Nagata, T.: Solar Flare Effect on the Geomagnetic Field, *Journal of geomagnetism and geoelectricity*, 18, 197–219, <https://doi.org/10.5636/jgg.18.197>, 1966.
- Olsen, N.: Ionospheric *F* region currents at middle and low latitudes estimated from Magsat data, *Journal of Geophysical Research: Space Physics*, 102, 4563–4576, <https://doi.org/10.1029/96JA02949>, 1997.
- Olsen, N., Glassmeier, K.-H., and Jia, X.: Separation of the Magnetic Field into External and Internal Parts, *Space Science Reviews*, 152, 135–157, <https://doi.org/10.1007/s11214-009-9563-0>, 2010.
- 1095 O’Sullivan, S. and Downes, T. P.: An explicit scheme for multifluid magnetohydrodynamics, *Monthly Notices of the Royal Astronomical Society*, 366, 1329–1336, <https://doi.org/10.1111/j.1365-2966.2005.09898.x>, 2006.
- Otto, A. and Zhu, H.: Fluid Plasma Simulation of Coupled Systems: Ionosphere and Magnetosphere, in: *Space Plasma Simulation*, edited by Büchner, J., Scholer, M., and Dum, C. T., vol. 615, pp. 193–211, Springer Berlin Heidelberg, Berlin, Heidelberg, ISBN 978-3-540-00698-5, https://doi.org/10.1007/3-540-36530-3_10, series Title: Lecture Notes in Physics, 2003.
- 1100 Parker, E. N.: The alternative paradigm for magnetospheric physics, *Journal of Geophysical Research: Space Physics*, 101, 10 587–10 625, <https://doi.org/10.1029/95JA02866>, 1996.
- Parker, E. N.: Conversations on electric and magnetic fields in the cosmos, no. 11 in Princeton series in astrophysics, Princeton University Press, Princeton, ISBN 978-0-691-12840-5 978-0-691-12841-2, oCLC: ocm70707945, 2007.
- 1105 Qian, L., Burns, A. G., Emery, B. A., Foster, B., Lu, G., Maute, A., Richmond, A. D., Roble, R. G., Solomon, S. C., and Wang, W.: The NCAR TIE-GCM: A Community Model of the Coupled Thermosphere/Ionosphere System, in: *Geophysical Monograph Series*, edited by Huba, J., Schunk, R., and Khazanov, G., pp. 73–83, Wiley, 1 edn., ISBN 978-0-87590-491-7 978-1-118-70441-7, <https://doi.org/10.1002/9781118704417.ch7>, 2014.
- Richmond, A. D.: Ionospheric Electrodynamics Using Magnetic Apex Coordinates., *Journal of geomagnetism and geoelectricity*, 47, 191–212, <https://doi.org/10.5636/jgg.47.191>, 1995.
- 1110 Richmond, A. D.: Ionospheric Electrodynamics, in: *Space Weather Fundamentals*, edited by Khazanov, G. V., pp. 245–259, CRC Press, 1 edn., ISBN 978-1-315-36847-4, <https://doi.org/10.1201/9781315368474-15>, 2016.
- Richmond, A. D. and Maute, A.: Ionospheric Electrodynamics Modeling, in: *Geophysical Monograph Series*, edited by Huba, J., Schunk, R., and Khazanov, G., pp. 57–71, Wiley, 1 edn., ISBN 978-0-87590-491-7 978-1-118-70441-7, <https://doi.org/10.1002/9781118704417.ch6>, 2014.
- 1115 Richmond, A. D., Blanc, M., Emery, B. A., Wand, R. H., Fejer, B. G., Woodman, R. F., Ganguly, S., Amayenc, P., Behnke, R. A., Calderon, C., and Evans, J. V.: An empirical model of quiet-day ionospheric electric fields at middle and low latitudes, *Journal of Geophysical Research: Space Physics*, 85, 4658–4664, <https://doi.org/10.1029/JA085iA09p04658>, 1980.
- Ridley, A., Deng, Y., and Tóth, G.: The global ionosphere–thermosphere model, *Journal of Atmospheric and Solar-Terrestrial Physics*, 68, 839–864, <https://doi.org/10.1016/j.jastp.2006.01.008>, 2006.
- 1120 Ridley, A. J., Gombosi, T. I., and DeZeeuw, D. L.: Ionospheric control of the magnetosphere: conductance, *Annales Geophysicae*, 22, 567–584, <https://doi.org/10.5194/angeo-22-567-2004>, 2004.

- Rishbeth, H. and Garriott, O. K.: Introduction to Ionospheric Physics, vol. 14 of *International Geophysics Series*, Academic Press, New York, ISBN 0125889402, 1969.
- 1125 Robinson, R. M., Vondrak, R. R., Miller, K., Dabbs, T., and Hardy, D.: On calculating ionospheric conductances from the flux and energy of precipitating electrons, *Journal of Geophysical Research: Space Physics*, 92, 2565–2569, <https://doi.org/10.1029/JA092iA03p02565>, 1987.
- Robinson, R. M., Zanetti, L., Anderson, B., Vines, S., and Gjerloev, J.: Determination of Auroral Electrodynamic Parameters From AMPERE Field-Aligned Current Measurements, *Space Weather*, 19, e2020SW002 677, <https://doi.org/10.1029/2020SW002677>, 2021.
- 1130 Sabaka, T. J., Hulot, G., and Olsen, N.: Mathematical Properties Relevant to Geomagnetic Field Modeling, in: *Handbook of Geomathematics*, edited by Freeden, W., Nashed, M. Z., and Sonar, T., pp. 1–37, Springer Berlin Heidelberg, Berlin, Heidelberg, ISBN 978-3-642-27793-1, https://doi.org/10.1007/978-3-642-27793-1_17-2, 2013.
- Schunk, R. W. and Nagy, A.: *Ionospheres: physics, plasma physics, and chemistry*, Cambridge University Press, Cambridge, UK, 2nd edn., ISBN 978-0-511-63489-5, oCLC: 503049988, 2009.
- 1135 Sciffer, M. D., Waters, C. L., and Menk, F. W.: Propagation of ULF waves through the ionosphere: Inductive effect for oblique magnetic fields, *Annales Geophysicae*, 22, 1155–1169, <https://doi.org/10.5194/angeo-22-1155-2004>, 2004.
- Sorathia, K. A., Merkin, V. G., Panov, E. V., Zhang, B., Lyon, J. G., Garretson, J., Ukhorskiy, A. Y., Ohtani, S., Sitnov, M., and Wiltberger, M.: Ballooning-Interchange Instability in the Near-Earth Plasma Sheet and Auroral Beads: Global Magnetospheric Modeling at the Limit of the MHD Approximation, *Geophysical Research Letters*, 47, e2020GL088 227, <https://doi.org/10.1029/2020GL088227>, 2020.
- 1140 Strangeway, R. J.: The equivalence of Joule dissipation and frictional heating in the collisional ionosphere, *Journal of Geophysical Research: Space Physics*, 117, 2011JA017 302, <https://doi.org/10.1029/2011JA017302>, 2012.
- Tamao, T.: Direct contribution of oblique field-aligned currents to ground magnetic fields, *Journal of Geophysical Research: Space Physics*, 91, 183–189, <https://doi.org/10.1029/JA091iA01p00183>, 1986.
- Tu, J. and Song, P.: A two-dimensional global simulation study of inductive-dynamic magnetosphere-ionosphere coupling, *Journal of Geophysical Research: Space Physics*, 121, <https://doi.org/10.1002/2016JA023393>, 2016.
- 1145 Tu, J. and Song, P.: On the Momentum Transfer From Polar to Equatorial Ionosphere, *Journal of Geophysical Research: Space Physics*, 124, 6064–6073, <https://doi.org/10.1029/2019JA026760>, 2019.
- Tu, J., Song, P., and Vasyliūnas, V. M.: Ionosphere/thermosphere heating determined from dynamic magnetosphere-ionosphere/thermosphere coupling: BRIEF REPORT, *Journal of Geophysical Research: Space Physics*, 116, n/a–n/a, <https://doi.org/10.1029/2011JA016620>, 2011.
- 1150 Tu, J., Song, P., and Vasyliūnas, V. M.: Inductive-dynamic magnetosphere-ionosphere coupling via MHD waves, *Journal of Geophysical Research: Space Physics*, 119, 530–547, <https://doi.org/10.1002/2013JA018982>, 2014.
- Tóth, G., Van Der Holst, B., Sokolov, I. V., De Zeeuw, D. L., Gombosi, T. I., Fang, F., Manchester, W. B., Meng, X., Najib, D., Powell, K. G., Stout, Q. F., Gloer, A., Ma, Y.-J., and Opher, M.: Adaptive numerical algorithms in space weather modeling, *Journal of Computational Physics*, 231, 870–903, <https://doi.org/10.1016/j.jcp.2011.02.006>, 2012.
- 1155 Untiedt, J. and Baumjohann, W.: Studies of polar current systems using the IMS Scandinavian magnetometer array, *Space Science Reviews*, 63, 245–390, <https://doi.org/10.1007/BF00750770>, 1993.
- van der Meeren, C., Laundal, K. M., Burrell, A. G., Lamarche, L. L., Starr, G., Reimer, A. S., Morschhauser, A., Michaelis, I., and Klenzing, J.: aburrell/apexpy: ApexPy Version 2.1.0, <https://doi.org/10.5281/ZENODO.1214206>, 2025.
- Vanhamäki, H.: Inductive ionospheric solver for magnetospheric MHD simulations, *Annales Geophysicae*, 29, 97–108, <https://doi.org/10.5194/angeo-29-97-2011>, 2011.
- 1160

- Vanhamäki, H. and Juusola, L.: Introduction to Spherical Elementary Current Systems, in: *Ionospheric Multi-Spacecraft Analysis Tools*, edited by Dunlop, M. W. and Lühr, H., pp. 5–33, Springer International Publishing, Cham, ISBN 978-3-030-26731-5 978-3-030-26732-2, https://doi.org/10.1007/978-3-030-26732-2_2, 2020.
- 1165 Vanhamäki, H., Viljanen, A., and Amm, O.: Induction effects on ionospheric electric and magnetic fields, *Annales Geophysicae*, 23, 1735–1746, <https://doi.org/10.5194/angeo-23-1735-2005>, 2005.
- Vanhamäki, H., Maute, A., Alken, P., and Liu, H.: Dipolar elementary current systems for ionospheric current reconstruction at low and middle latitudes, *Earth, Planets and Space*, 72, 146, <https://doi.org/10.1186/s40623-020-01284-1>, 2020.
- Vasyliūnas, V. M.: Electric field and plasma flow: What drives what?, *Geophysical Research Letters*, 28, 2177–2180, <https://doi.org/10.1029/2001GL013014>, 2001.
- 1170 Vasyliūnas, V. M.: Relation between magnetic fields and electric currents in plasmas, *Annales Geophysicae*, 23, 2589–2597, <https://doi.org/10.5194/angeo-23-2589-2005>, 2005.
- Vasyliūnas, V. M.: The mechanical advantage of the magnetosphere: solar-wind-related forces in the magnetosphere-ionosphere-Earth system, *Annales Geophysicae*, 25, 255–269, <https://doi.org/10.5194/angeo-25-255-2007>, 2007.
- Vasyliūnas, V. M.: The physical basis of ionospheric electrodynamics, *Annales Geophysicae*, 30, 357–369, [https://doi.org/10.5194/angeo-](https://doi.org/10.5194/angeo-30-357-2012)
- 1175 30-357-2012, 2012.
- Vasyliūnas, V. M. and Song, P.: Meaning of ionospheric Joule heating, *Journal of Geophysical Research: Space Physics*, 110, 2004JA010615, <https://doi.org/10.1029/2004JA010615>, 2005.
- Von Alfthan, S., Pokhotelov, D., Kempf, Y., Hoilijoki, S., Honkonen, I., Sandroos, A., and Palmroth, M.: Vlasiator: First global hybrid-Vlasov simulations of Earth’s foreshock and magnetosheath, *Journal of Atmospheric and Solar-Terrestrial Physics*, 120, 24–35, <https://doi.org/10.1016/j.jastp.2014.08.012>, 2014.
- 1180 Waters, C. L. and Sciffer, M. D.: Field line resonant frequencies and ionospheric conductance: Results from a 2-D MHD model, *Journal of Geophysical Research: Space Physics*, 113, 2007JA012822, <https://doi.org/10.1029/2007JA012822>, 2008.
- Wright, A. N.: Transfer of magnetosheath momentum and energy to the ionosphere along open field lines, *Journal of Geophysical Research: Space Physics*, 101, 13 169–13 178, <https://doi.org/10.1029/96JA00541>, 1996.
- 1185 Yamauchi, M., Johnsen, M. G., Enell, C.-F., Tjulin, A., Willer, A., and Sormakov, D. A.: High-latitude crochet: solar-flare-induced magnetic disturbance independent from low-latitude crochet, *Annales Geophysicae*, 38, 1159–1170, <https://doi.org/10.5194/angeo-38-1159-2020>, 2020.
- Yamazaki, Y. and Maute, A.: Sq and EEJ—A Review on the Daily Variation of the Geomagnetic Field Caused by Ionospheric Dynamo Currents, *Space Science Reviews*, 206, 299–405, <https://doi.org/10.1007/s11214-016-0282-z>, 2017.
- 1190 Yoshikawa, A. and Itonaga, M.: Reflection of shear Alfvén waves at the ionosphere and the divergent Hall current, *Geophysical Research Letters*, 23, 101–104, <https://doi.org/10.1029/95GL03580>, 1996.
- Zettergren, M. D. and Snively, J. B.: Ionospheric response to infrasonic-acoustic waves generated by natural hazard events, *Journal of Geophysical Research: Space Physics*, 120, 8002–8024, <https://doi.org/10.1002/2015JA021116>, 2015.
- Zhang, B., Sorathia, K. A., Lyon, J. G., Merkin, V. G., Garretson, J. S., and Wiltberger, M.: GAMERA: A Three-dimensional Finite-volume MHD Solver for Non-orthogonal Curvilinear Geometries, *The Astrophysical Journal Supplement Series*, 244, 20, <https://doi.org/10.3847/1538-4365/ab3a4c>, 2019.