

Biases in estimated vegetation indices from observations under cloudy conditions

Kevin Wolf¹, Evelyn Jäkel¹, André Ehrlich¹, Michael Schäfer¹, Hannes Feilhauer^{2,3,4}, Andreas Huth^{4,5,6}, and Manfred Wendisch¹

¹Leipzig Institute for Meteorology (LIM), Leipzig University, Leipzig, Germany

²Institute for Earth System Science & Remote Sensing, Leipzig University, Leipzig, Germany.

³Remote Sensing Centre for Earth System Research, Leipzig University, Leipzig, Germany.

⁴iDiv German Centre for Integrative Biodiversity Research Halle-Jena-Leipzig, Leipzig, Germany.

⁵Department of Ecological Modelling, Helmholtz Centre for Environmental Research–UFZ Leipzig, Leipzig, Germany.

⁶Institute for Environmental Systems Research, University of Osnabrück, Osnabrück, Germany.

Correspondence: Kevin Wolf (kevin.wolf@uni-leipzig.de)

Abstract. Field observations of vegetation indices (VIs) ~~from~~ are derived from ratios of spectral reflectance data that are collected by drones and aircraft ~~provide~~ providing higher spatial resolution than satellites. ~~Vegetation indices are derived from ratios of spectral reflectivity measurements. The reflectivity is measured in a relative way by~~ These reflectance data require periodic reference measurements over ~~reflectance panels. This requires calibrated reflectance panels under~~ cloud-free or at
5 ~~least stable cloud conditions between reflectance panel measurements. This assumption is often violated,~~ conditions. However,
the reference measurements are partly performed in cloudy situations with the effect that wavelength-dependent scattering and absorption of solar radiation by clouds ~~lead to a distortion of the below-cloud spectral downward irradiance $F^{\downarrow}(\lambda)$ and thus affects estimates of VIs.~~

~~This paper presents~~ affects the subsequently derived VIs. This paper quantifies these effects using combined atmosphere-
10 ~~vegetation radiative transfer (RT) simulations to systematically investigate cloud-induced biases in remotely sensed VIs derived from below-cloud measurements. The biases in VIs have been investigated for~~ We study the general case ~~of two-band VIs when VIs are obtained from reflectance ratios of two wavelengths,~~ and for the special cases of the normalized difference vegetation index (NDVI), the normalized difference water index (NDWI), and the enhanced vegetation index (EVI). For the general case of two-band VIs the lowest sensitivity to cloud changes was found for wavelength combinations below 1400 nm and outside
15 ~~the water vapor absorption bands. The NDVI was found to be almost insensitive to changes in cloud conditions, while greater biases were identified for the NDWI. The EVI was also found to be most~~ susceptible to cloud changes, leading to biases of
0.36 with biases of 0.2 in the selected example ~~with~~ . This lead to biases in the estimated leaf area index of ~~1.30.9. Biophysical properties derived from EVI, such as gross primary product, are also affected with variations of up to $\pm 2 \text{ gC m}^{-2} \text{ d}^{-1}$ in the selected cases.~~

20 1 Introduction




Figure 1. (A) Schematic illustration of measured spectral upward radiance $I^\uparrow$ under cloud-free conditions. (B) and (C) Schematic illustration of below cloud measurements that can be subject to a change in cloud conditions from τ_{cal} (during reflectance panel (RP) calibration) to cloud conditions τ_{mea} (during actual measurements). The spectral total downward irradiance $F^\downarrow$ and spectral upward radiance $I^\uparrow$ are affected by the cloud conditions. The ~~reflectivity-reflectance~~ ρ_{RP} of the RP panel is known from its specifications and ρ_{mea} is the ~~inherent-spectral reflectivity-reflectance~~ of the vegetation. The transfer ~~factors-functions~~ κ_{cal} and κ_{mea} ~~relate-connect~~ measured counts and ρ .

~~Since the advent of satellite-based vegetation remote sensing, numerous~~ Numerous vegetation indices (VIs) have been developed (Xue and Su, 2017). These indices are designed to remotely characterize the ~~vegetation properties in terms of~~ vegetation. They are based on ratios of vegetation-reflected radiation in wavelength regions that are sensitive to chlorophyll and liquid water absorption, thus indirectly providing information on plant vitality, productivity, and photosynthetic activity, and to monitor anthropogenic degradation (Myneni et al., 1997; Saleska et al., 2007; Jiang et al., 2008; Mura et al., 2018) (Collins, 1978; Horler et al., 1983). Established VIs are, for example, the normalized difference vegetation index (NDVI), the normalized difference water index (NDWI), and the enhanced vegetation index (EVI) (Kriegler et al., 1969; Huete et al., 1994; Myneni et al., 1995; Gao, 1996; Chen et al., 2005; Jiang et al., 2008; Jones and Vaughan, 2010). ~~In general, VIs are based on the ratios of vegetation-reflected radiation in wavelength regions that are sensitive to chlorophyll and water absorption, thus indirectly providing information on plant productivity and vitality~~ (Bowker, 1985; Collins, 1978; Horler et al., 1983; Knyazikhin et al., 2013; Richardson et al., 2021). ~~Vegetation indices~~ They were originally developed for satellite observations and ~~were~~ customized to spectral bands of satellites such as Landsat (Wulder et al., 2019), the Moderate Resolution Imaging Spectroradiometer (MODIS; Salomonson et al., 1989), or Sentinel-2 (Spoto et al., 2012). ~~During the development of VIs~~ In the course of the time, the available measurement platforms diversified, ~~now including satellite, aircraft,~~ additionally including aircraft and unmanned aerial vehicles as ~~measurement platforms~~ (UAVs, Matese et al., 2015; Duan et al., 2017; Singh and Frazier, 2018; Jiang et al., 2020).

All remotely sensed VIs rely on the spectral ~~reflectivity-reflectance~~ $\rho(\lambda)$, ~~which is formalized by~~ $\rho(\lambda)$. In its most general form, $\rho(\lambda)$ is defined as the ratio of the surface-reflected radiance from within an infinite solid angle to the incoming radiance from within an infinite solid angle (Nicodemus, 1977; Martonchik et al., 2000; Schaepman-Strub et al., 2006). The amount of radiation reflected into a given solid angle is defined by the surface-specific bidirectional reflectance distribution function (BRDF, Schaepman-Strub et al., 2006). Furthermore, the surface reflected radiation depends on topography (Matsushita et al., 2007) and on changes in illumination conditions determined by solar zenith angle, aerosol particles, and clouds (Pinty et al., 2005; Singh and Frazier

Definitions of $\rho(\lambda)$ based on first principals are given by Nicodemus (1977) and Schaepman-Strub et al. (2006). For brevity and simplicity, we restrict the definition of $\rho(\lambda)$, assuming idealized conditions of pure diffuse illumination and Lambertian reflection at the surface, which results in:

$$\rho(\lambda) = \frac{I^\uparrow(\lambda) \cdot \pi \cdot \text{sr}}{F^\downarrow(\theta, \lambda)} \frac{I^\uparrow(\lambda) \cdot \pi \cdot \text{sr}}{F^\downarrow(\theta, \lambda)}, \quad (1)$$

where λ symbolizes the wavelength, sr the unit steradian, $I^\uparrow(\lambda)$ ~~represents~~ the spectral, upward (reflected) radiance in units of $\text{W m}^{-2} \text{nm}^{-1} \text{sr}^{-1}$, and $F^\downarrow(\theta, \lambda)$ ~~represents~~ the downward spectral irradiance in units of $\text{W m}^{-2} \text{nm}^{-1}$, which also depends on the solar zenith angle θ . ~~At the top of the atmosphere, $F^\downarrow(\theta, \tau=0, \lambda)$ is solely defined by direct solar irradiance $F_{\text{dir}}^\downarrow(\theta, \tau, \lambda)$, which can partially reach the surface without any interaction in the atmosphere.~~ The cloud optical thickness $\tau(\lambda)$ is a measure of the extinction of radiation for a vertical path through the cloud, serving as vertical coordinate. Radiation that has never been scattered represents the direct radiation component $F_{\text{dir}}^\downarrow(\theta, \lambda)$. The spectral reflectance ranges between 0 and 1, except under broken cloud conditions where radiation enhancement occurs, leading to reflectance values exceeding unity (Schaepman-Strub et al., 2006; Marshak et al., 2008). Only in heterogeneous cloud cases, where three-dimensional scattering effects occur, are reflectance values greater than one observable. In neighboring regions, reflectance is reduced. On average, this results in energy conservation. Radiation that has undergone at least one scattering event contributes to diffuse irradiance $F_{\text{dif}}^\downarrow(\theta, \tau, \lambda)$. Thus, $F^\downarrow(\theta, \tau, \lambda)$ is decomposed into:

$$F^\downarrow(\theta, \tau, \lambda) = F_{\text{dir}}^\downarrow(\theta, \tau, \lambda) + F_{\text{dif}}^\downarrow(\theta, \tau, \lambda). \quad (2)$$

The related direct fraction $f_{\text{dir}}(\theta, \tau, \lambda)$ is defined by:

$$f_{\text{dir}}(\theta, \tau, \lambda) = \frac{F_{\text{dir}}^\downarrow(\theta, \tau, \lambda)}{F^\downarrow(\theta, \tau, \lambda)}. \quad (3)$$

~~The spectral reflectivity ranges between 0 and 1, except under broken cloud conditions where radiation enhancement occurs, leading to values exceeding 1.~~

Remote sensing techniques to ~~retrieve estimate~~ VIs from satellite measurements require cloud-free conditions. Cloud masks are used to flag cloud-contaminated pixels, which are excluded from the VI calculation. The remaining pixels are assumed to be unbiased by clouds. Figure 1 (A) schematically illustrates the measurement principle. However, for all types of airborne measurements, ~~such as drone or aircraft measurements that are from drones or aircraft that are maybe~~ performed below clouds, see Fig. 1 (B and C), the downward spectral irradiance $F^\downarrow(\lambda)$ ~~depends on the cloud optical thickness τ and the cloud particle size (Twomey and Cocks, 1989). The cloud optical thickness $\tau(\lambda)$ is a measure of the extinction of radiation for a vertical path through the cloud that depends on the volumetric extinction coefficient.~~

Regardless of the platform, the derived VIs are affected by external parameters, such as topography, due to angle-dependent reflectivity properties of the terrain (Matsushita et al., 2007), and in particular by changes in illumination conditions determined by solar zenith angle, aerosol particles, and clouds (Singh and Frazier, 2018). Clouds alter $F^\downarrow(\theta, \tau, \lambda)$ spectrally as a result of spectral-dependent scattering of radiation ~~is altered by the presences of clouds. The impact of clouds depends on τ , droplet/ice~~

crystal size, and wavelength (Twomey and Cocks, 1989). The contribution of scattering and absorption is spectrally dependent, with scattering most pronounced in the visible near-infrared (VNIR, 0.3–1.0 μm) part of the solar spectrum (Mie, 1908; van de Hulst, 1981; Dubovik et al., 2002) and absorption in the shortwave infrared (SWIR, 1.0–2.5 μm). ~~Scattering and absorption lead wavelength range. The spectral dependency leads~~ to a spectral slope in $F^\downarrow(\lambda)$, $I^\uparrow(\lambda)$, and ~~finally in~~ $\rho(\lambda)$ (Wiscombe and Warren, 1980; Grenfell and Perovich, 2008). Consequently, changes in cloud properties and illumination ~~conditions~~ lead to biases in estimated VIs.

To obtain $\rho(\lambda)$ ~~and retrieve VIs for VI retrievals~~ during field studies, calibrated $I^\uparrow(\tau, \lambda)$ and $F^\downarrow(\theta, \tau, \lambda)$ are required, which ~~are~~ sometimes measured with dedicated, calibrated sensors (Hakala et al., 2013; Honkavaara et al., 2013; Miyoshi et al., 2018). Alternatively, $F^\downarrow(\theta, \tau, \lambda)$ is simulated using radiative transfer (RT) models. In either case, absolute measurements are complicated and simulations introduce uncertainties. For practical applications, absolute measurements are often avoided by performing relative measurements, using ~~measurements of~~ reflectance panels (RPs) characterized by a ~~well-known reflectivity~~ well-known reflectance $\rho_{\text{RP}}(\lambda)$ (Burkart et al. (2014); Aasen et al. (2015); Aasen and Bolten (2018); Hakala et al. (2018); Fawcett et al. (2020); Köppl et al., 2021)

Using a RP ~~attempts allows~~ a transfer calibration. Digital spectrometers register digital counts S , which are related to $I^\uparrow(\tau, \lambda)$ by a calibration factor $\mathcal{C}(\lambda)$. Including this calibration factor in Eq. 1 yields:

$$\rho(\lambda) = \frac{S^\uparrow(\tau, \lambda) \cdot \mathcal{C}^\uparrow(\lambda) \cdot \pi \cdot \text{sr}}{F^\downarrow(\theta, \tau, \lambda)} \frac{S^\uparrow(\tau, \lambda) \cdot \mathcal{C}^\uparrow(\lambda) \cdot \pi \cdot \text{sr}}{F^\downarrow(\theta, \tau, \lambda)}, \quad (4)$$

where $S^\uparrow(\tau, \lambda)$, ~~and therefore~~ $I^\uparrow(\tau, \lambda)$, and $F^\downarrow(\theta, \tau, \lambda)$ below clouds ~~are additionally~~ dependent on τ . Equation 4 can be simplified to:

$$\rho(\lambda) = S^\uparrow(\tau, \lambda) \cdot \kappa(\theta, \tau, \lambda), \quad (5)$$

with the transfer ~~factor function~~ $\kappa(\theta, \tau, \lambda)$ given by:

$$\kappa(\lambda, \tau, \theta) = \frac{\mathcal{C}^\uparrow(\lambda) \cdot \pi \cdot \text{sr}}{F^\downarrow(\tau, \lambda)} \frac{\mathcal{C}^\uparrow(\lambda) \cdot \pi \cdot \text{sr}}{F^\downarrow(\tau, \lambda)}. \quad (6)$$

In a first step during the relative measurements, the RP is overflowed and the signal is taken for calibration (Fig. 1, B), where $\rho(\lambda)$ in Eq. 5 is set to the well-defined $\rho_{\text{RP}}(\lambda)$ and $S^\uparrow(\lambda)$ is the number of counts registered by the radiance sensor above the RP. The transfer ~~factor function~~ $\kappa_{\text{cal}}(\theta, \tau_{\text{cal}}, \lambda)$ determined during the calibration procedure now represents the relationship between the recorded digital counts and the surface ~~reflectivity reflectance~~ under the illumination conditions at that time. During ~~the~~ actual measurements over vegetation (Fig. 1, C), the ~~reflectivity reflectance~~ $\rho_{\text{mea}}(\lambda)$ of the vegetated surface is determined ~~by applying~~ $\kappa(\theta, \tau_{\text{cal}}, \lambda)$:

$$\rho_{\text{mea}}(\lambda) = \kappa_{\text{cal}}(\theta, \tau_{\text{cal}}, \lambda) \cdot S^\uparrow(\lambda). \quad (7)$$

Although RP calibration avoids the determination of the individual factors included in $\kappa_{\text{cal}}(\theta, \tau_{\text{cal}}, \lambda)$, the dependence of $\kappa(\theta, \tau, \lambda)$ on θ and τ , requires frequent ~~transfer calibrations with the RP~~ RP overflights to account for illumination changes and to obtain updated transfer functions.

105 Lohmann (2018) presented example time series of $F^\downarrow$ from a field campaign (Madhavan et al., 2016; Macke et al., 2017) and compared them with cloud-free simulations. They determined single-point enhancements of $F^\downarrow$ up to 50 % due to cloud-side enhancements. Short-term fluctuations of $F^\downarrow$ have been found on time scales down to 300 s (Jurado et al., 1995), 60 s (van Haaren et al., 2014), 20 s (Perez et al., 2011), and also below 1 s (Calif et al., 2013). Consequently, $\kappa(\theta, \tau, \lambda)$ and thus $\rho_{\text{mea}}(\lambda)$ vary on the same time scales. Therefore, even regular sequences of RP measurements seem insufficient to capture
110 fluctuations caused by broken clouds, either by the clouds themselves or by their shadows, which may lead to potential errors in the retrieved VIs (Burkart et al., 2014; Behmann et al., 2015; Köppl et al., 2021).

Even when $F^\downarrow(\lambda)$ is measured by a dedicated sensor or frequent measurements of the RP are performed, the sole presence of clouds causes a change in $\rho_{\text{mea}}(\lambda)$, since clouds change the ratio of direct and diffuse radiation, which determines how radiation is reflected by a non-isotropic surface (Schaeppman-Strub et al., 2006).

115 Here we present coupled simulations using the atmospheric RT model "library for Radiative transfer" (libRadtran Emde et al., 2016) and the vegetation RT model "Soil Canopy Observation of Photosynthesis and Energy fluxes" (~~SCOPE2.0 Yang et al., 2021; ?~~) (SCOPE2.0 Yang et al., 2021; Wolf et al., 2025a). The iterative model coupling allows to account for mutual influences between cloud and canopy. The coupled simulations ~~were~~ are used to systematically investigate the impact of different cloud conditions on general two-band VIs, and in particular on NDVI, NDWI, and EVI. The aim is to quantify the impact of clouds on
120 below-cloud airborne observations that are based on relative measurements using RP calibration thus providing approximate associated errors in estimated VI. The VIs ~~were~~ are calculated using the wavelength bands of the multispectral instrument (MSI) onboard the Sentinel-2 satellite. This introduction is followed by a brief overview of the main terminology, the applied RT models, and a short review of the model coupling in Section 2. This is followed by the results in Section 3, where the effect of clouds and changing cloud conditions on estimated VIs are presented. Section 4 provides a summary of the results
125 and outlines the limitations of the simulations.

2 Vegetation indices and radiative transfer simulations

2.1 Definition of vegetation indices

~~Sentinel-2 wavelength bands from the multispectral instrument (MSI) following Drusch et al. (2012): Band number-Center wavelength-nmSpectral width-nmB2 490 ±33 B3 560 ±35 B4 665 ±15 B8 842 ±56 B8a 865 ±10 B9 945 ±10 B10 1375 ±15 B11 1610 ±45 B12 2190 ±180~~ Vegetation indices are based on ratios of $\rho(\lambda)$ at several wavelengths (mostly pairs). The exact center wavelength and width used in the calculation of VIs depend on the characteristics of the observing instrument. In the case of the Sentinel-2 satellites, several wavelength combinations are suitable for VI retrievals (Kriegler et al., 1969; Carlson and Ripley, 1997; Mura et al., 2018). A subset of the Sentinel-2 wavelength bands are listed in Table 1. Unless otherwise noted, the Sentinel-2 wavelength bands were used to calculate VIs throughout this paper by applying a boxcar-like spectral response

Table 1. Sentinel-2 wavelength bands from the multispectral instrument (MSI) following Drusch et al. (2012).

Band number	Center wavelength [nm]	Spectral width [nm]
B2	490	±33
B3	560	±35
B4	665	±15
B8	842	±56
B8a	865	±10
B9	945	±10
B10	1375	±15
B11	1610	±45
B12	2190	±180

function. The Sentinel-2 wavelength bands are considered representative of other measurement platforms and sensors. UAVs may be equipped with spectral sensors that cover similar wavelength bands, or in the case of multispectral sensors, spectral integration is performed over similar wavelength bands (Jones et al., 2012).

Multiple two-band VIs with the index value γ exist that follow the general form of a band transformation:

$$\gamma = \frac{\rho(\lambda_1) - \rho(\lambda_2)}{\rho(\lambda_1) + \rho(\lambda_2)}, \tag{8}$$

where λ_1 and λ_2 represent a pair of single-individual wavelengths or narrow bands with band centers between 400 and 2400 nm wavelengths.

An example of a two-band VI is the normalized difference water index (NDWI). Two versions of the NDWI exist. The NDWI proposed by Gao (1996) uses a wavelength combination of 980 and 1240 nm; it is subsequently labeled with NDWI₁₂₄₀. Chen et al. (2005) provided the alternative NDWI₁₆₄₀, which was designed to be less sensitive to saturation with respect to the water content in plant matter compared to NDWI₁₂₄₀.

Another commonly used VI that follows Eq. 8 is the normalized difference vegetation index (NDVI; Kriebler et al., 1969), with $\rho(\lambda_1)$ from B4 and $\rho(\lambda_2)$ from B8 from the MSI Sentinel-2 channels. By construction, the NDVI ranges between -1 and 1 . An NDVI close to $+1$ indicates vital and productive vegetation, while stressed vegetation results in NDVI values around 0.2 . Measurements above bare soil will give a NDVI around 0 , while measurements over water will yield negative NDVI. Even though the NDVI proved to be useful it also showed sensitivity to surface brightness (Huete et al., 1985; Bausch, 1993) and scattering and absorption by atmospheric constituents (Kaufman and Tanre, 1992; Miura et al., 1998) (Huete et al., 1985; Kaufman and Tanre, 1992). The NDVI also becomes saturated and insensitive to vegetation structure when the canopies become dense, i.e., high leaf density (Ünsalan and Boyer, 2011).

These issues led to the development of the enhanced vegetation index (EVI; Huete et al., 1994), which was derived to overcome the shortcomings of the NDVI (Boegh et al., 2002; Huete et al., 2006). The EVI attempts to account for aerosol loading, surface reflectivityreflectance, and other factors by including spectral information from the "blue" region (ρ_{B2}) of the

VNIR spectrum. The EVI is calculated for MSI Sentinel-2 bands by:

$$\text{EVI} = G \cdot \frac{\rho_{B8} - \rho_{B4}}{\rho_{B8} + C_1 \cdot \rho_{B4} - C_2 \cdot \rho_{B2} + L}, \quad (9)$$

with the scaling factors $G = 2.5$, $C_1 = 6$, $C_2 = 7.5$, and $L = 1$ (Liu and Huete, 1995).

160 2.2 Radiative transfer simulations

Equation 1 shows that the ~~reflectivity~~ reflectance of a surface is determined by several factors, including θ , the observation geometry, as well as the direct and diffuse components of $F^\downarrow(\lambda)$ that are determined by τ (see Section B in the appendix). Furthermore, radiation interactions may occur between the surface and the cloud, which can be accounted for by iterative coupling of the RT models of the atmosphere and vegetation ~~(?)~~ (Wolf et al., 2025a). In the present paper we use the same model coupling setup introduced ~~by ?~~ and described by Wolf et al. (2025a).

2.2.1 Atmospheric radiative transfer model libRadtran

The atmospheric RT above the canopy was simulated with the library for Radiative transfer (libRadtran, Emde et al., 2016). The RT equation was calculated with the one-dimensional solver "Discrete-Ordinate-Method Radiative Transfer" (DISORT, Stamnes et al., 1988; Buras et al., 2011) using 12 streams. Clouds were assumed to be homogeneous. A low-level and mid-level warm stratus or altostratus were included by liquid water clouds between 3 and 3.5 km altitude with a fixed droplet effective radius of 10 μm (Stephens, 1994; Frisch et al., 2002; Aebi et al., 2020). A high-level cirrostratus was included by ice water clouds between 11 and 11.5 km altitude with a fixed ice effective radius of 85 μm (Freudenthaler et al., 1995; Sassen and Campbell, 2001; Noël and Haeffelin, 2007; Iwabuchi et al., 2012; Luebke et al., 2016; Krämer et al., 2016). Aggregated ice particles with moderate surface roughness are assumed to represent mature ice crystals (Liu et al., 2014; Holz et al., 2016; Järvinen et al., 2018). The ice particle parametrization after Yang et al. (2013) was applied. The liquid water and ice water path were scaled so that a specific $\tau(\lambda = 550\text{nm})$ at 550 nm wavelength was reached, with all other wavelengths being scaled considering the wavelength dependence of τ . For simplicity, $\tau(\lambda = 550\text{nm})$ is referred to as τ in the following. The incoming spectral irradiance at the top of atmosphere was represented by the solar reference spectrum provided by Coddington et al. (2021). Molecular absorption was considered using the parameterization of the "medium" resolution of Gasteiger et al. (2014). A default aerosol distribution after Shettle (1989) was applied, representing rural type aerosol in the boundary layer, background aerosol above 2 km during spring-summer, and a visibility of 50 km. Atmospheric profiles of air temperature, humidity, and gas concentrations were taken from the mid-latitude summer profile 'afglms' (Anderson et al., 1986). Absorption by water vapor and other atmospheric trace gases was accounted for in the simulations (Anderson et al., 1986; Emde et al., 2016). ~~As~~ The iteration process was first started by running libRadtran with an initial guess ~~of for~~ for the surface albedo ~~in libRadtran, the~~ . The "mixed-forest" albedo was taken from the IGBP database (Loveland and Belward, 1997). After one iteration cycle, the surface albedo determined during the iterative model coupling process was used (Wolf et al., 2025a). The RT simulations of $F^\uparrow(\lambda)$, $F^\downarrow(\lambda)$, and $I^\uparrow(\lambda)$ spanned a wavelength range from 0.4 to 2.4 μm . The output of libRadtran was specified to be at an

altitude of 40 m above the canopy to be representative for UAV measurements. The radiances were simulated for a nadir sensor geometry.

190 2.2.2 Vegetation radiative transfer model SCOPE2.0

The solar RT within vegetation was simulated with the Soil Canopy Observation of Photosynthesis and Energy fluxes version 2 (SCOPE2.0, Yang et al., 2017, 2020, 2021). The surface albedo was determined using the Brightness-Shape-Moisture (BSM) model (Verhoef et al., 2018; Yang et al., 2020). ~~Simulations Irradiance and radiance simulations~~ in the solar part of the spectrum have been performed for a wavelength range from 0.4 to 2.4 μm . The angular dependence of $I^\uparrow(\lambda)$ is considered for by the
195 actual illumination and observation geometries, the direct and diffuse $F^\downarrow(\lambda)$, and the parameterized intrinsic reflectivity within SCOPE2.0. An important parameter to describe the radiative properties of a canopy is the leaf area index (LAI, Watson, 1947; Asner, 1998; Jones and Vaughan, 2010), ~~ranging typically~~ typically ranging between 0 and 12. It provides a measure of the accumulated, one-sided area of leaves per unit of ground area given in units of $\text{m}^2 \text{m}^{-2}$. A constant LAI of $3 \text{ m}^2 \text{m}^{-2}$ was used in the simulations, corresponding to the typical LAI of temperate vegetation types such as pine forests or various grasslands.
200 The leaf angle distribution (LAD) is another important parameter controlling the RT in a canopy (Baldocchi et al., 2002; Jones and Vaughan, 2010; Verrelst et al., 2015; Yang et al., 2023). It is a measure of the overall orientation of the leaf ensemble of a tree. Therefore, it influences the area of a leaf illuminated by the Sun with respect to the total one-sided leaf area (Asner, 1998; Stuckens et al., 2009; Vicari et al., 2019). The simulations were primarily considered for the general case of a spherical LAD, where all leaf angles have a similar probability (Goel, 1988). Further simulations have been performed to also include
205 the erectophile and the planophile LADs. The static LAD values obviously ignores that short term changes in weather and illumination conditions affect the ~~leaf-angle-distribution-LAD~~ (Kattenborn et al., 2022), which may likewise have significant effects on VIs (Kattenborn et al., 2024), but is nevertheless common practice in many RTM parameterizations. Table 2 provides an overview of the selected parameters for the vegetation RT simulations~~in the visible-near-infrared wavelength range that were kept constant in the simulations.~~ The parameters were selected on the basis of their relevance to surface reflectance
210 within the visible and near-infrared wavelength ranges. Their individual relevance was estimated in a sensitivity study by Wolf et al. (2025a).

3 Results and discussion

The one-dimensional simulations combining libRadtran and SCOPE2.0 are to be interpreted as synthetic measurements of $F^\downarrow(\lambda)$, $I^\uparrow(\lambda)$, and $\rho(\lambda)$ under varying atmospheric conditions, ranging from cloud-free to stratiform clouds, with values of
215 τ between 0 and 40, and θ between 25° and 70° . By assuming various combinations of τ_{cal} , which prevailed during calibration and τ_{mea} , present during actual measurements, the effects of changes in cloud conditions between RP measurements on $\rho_{\text{mea}}(\lambda)$ and estimated VIs were determined. This was ~~realized-performed~~ using simulated $I^\uparrow(\tau_{\text{mea}}, \lambda)$ and $F^\downarrow(\tau_{\text{cal}}, \lambda)$ in Eq. 1, which is synonymous with ~~assuming the assumption of~~ constant values of κ_{cal} . The values of simulated τ_{cal} and τ_{mea} range from 0 to 40 for the liquid water cloud and the ice water cloud. Two aspects influenced the chosen ~~rangeranges~~. First,

Table 2. Selected configuration of the SCOPE2.0 simulations.

Description	Symbol	Setting	Unit
Leaf chlorophyll concentration	C_{ab}	40	$\mu\text{g cm}^{-2}$
Leaf carotenoid concentration	C_{ca}	10	$\mu\text{g cm}^{-2}$
Leaf water equivalent layer	C_w	0.009	cm
Leaf structure parameter	N	2.1	Unitless
BSM model parameter for soil brightness	B	0.5	Unitless
Vegetation height	h_c	20	m
Output height	h_{out}	40	m

the simulated ice clouds with τ up to 6 can be considered as a high-level cirrus. The simulated ice clouds with $\tau > 6$ can be regarded as mid-level ice clouds. This flexible interpretation is feasible since cloud altitude is of minor importance for the RT simulations in the solar wavelength range. However, the shape information of the natural variation in non-spherical ice particles and the absorption coefficient of ice are important particle shapes was taken into account by selecting aggregated ice particles (see Sec.2.2.1). The second aspect is that the bias in $\rho(\lambda)$ and the estimated VIs depend on the difference between τ_{cal} and τ_{mea} but also on the absolute value. To cover even optically thick clouds, values of τ up to 40 were included. Second, liquid water clouds were considered by simulating mid-level, continental stratus clouds, which are known to often reach values of up to $\tau = 40$ (King et al., 1993; Lu et al., 2008). The subsequent analysis focuses primarily on the spherical LAD, which is considered the general case. The effect of the selected LAD is discussed in Appendix Sec. A.

3.1 Influence of clouds on the spectral reflectivityreflectance

Pinty et al. (2005) have shown the influence of clouds molecules and aerosols on direct and diffuse $F^\downarrow(\lambda)$, the associated effects on $F^\uparrow(\lambda)$, and the resulting albedo effects over vegetated areas using based on analytical equations. Wolf et al. (2025a) used coupled atmosphere–vegetation radiative transfer models. They to investigate these effects, explicitly considering clouds and spectral depending scattering and absorption in the atmosphere. An increase in τ leads to a decrease in $F^\downarrow_{dir}(\lambda)$, while the response of $F^\downarrow_{dif}(\lambda)$ depends on τ . For values of τ less than 4 to 6, $F^\downarrow_{dif}(\lambda)$ first increases and then decreases as τ increases further. The total $F^\downarrow(\lambda)$ and $f^\downarrow_{dir}(\lambda)$ both continuously decrease as τ increases. In addition, $F^\uparrow(\lambda)$ became less sensitive to changes in θ . Lastly, the presence of clouds modulates the incoming radiation spectrally by shifting the incoming radiation towards shorter wavelengths, as clouds primarily scatter radiation at shorter wavelengths and absorb radiation at longer wavelengths. Wolf et al. (2025a) also showed that radiative interactions between the canopy and the cloud base increase $F^\downarrow_{dif}(\lambda)$ and albedo compared to cloud-free conditions. The present paper focuses on the related effects on $I^\uparrow(\lambda)$ and $\rho(\lambda)$.




Figure 2. (a) Simulated spectral ~~reflectivity~~-reflectance $\rho(\lambda)$ caused by an ice cloud for a solar zenith angle θ of 25°, and constant cloud optical thickness during calibration and ~~the~~ actual ~~synthetic~~ measurements ($\tau_{cal} = \tau_{mea}$). ~~The simulations base on a spherical LAD.~~ (b) Ratios of spectral ~~reflectivity~~-reflectance with $\tau_{cal} = \tau_{cal} > 0$ with respect to the spectral ~~reflectivity~~-reflectance under cloud-free conditions. ~~Panels (a) and (b) share the same legend. The gray marked areas highlight the Sentinel-2 bands B2, B4, B8, B8a, and B11.~~

240 3.1.1 Effect of diffuse radiation on spectral ~~reflectivity~~reflectance above vegetation

First, we consider situations, where the cloud conditions are similar during the RP calibration and the measurements ($\tau_{\text{cal}} = \tau_{\text{mea}}$). Figure 2a shows $\rho(\lambda)$ for $\theta = 25^\circ$ with the lowest $\rho(\lambda)$ under cloud-free conditions ($\tau_{\text{cal}} = \tau_{\text{mea}} = 0$, black line). With increasing values of τ , $\rho(\lambda)$ also increases, especially between 750 and 1300 nm ~~wavelength~~wavelengths, where vegetation is generally characterized by high ~~reflectivity~~and reflectance. The sensitivity of $\rho(\lambda)$ to τ is highest for small values of τ and quickly approaches an asymptotic value when $F_{\text{dif}}^\downarrow(\lambda)$ dominates the radiation field. Figure 2b shows the ratio of $\rho(\lambda)$ between cloudy and cloud-free conditions. Independent of τ , the ratio has a distinct spectral dependency with pronounced water absorption bands. Under cloudy conditions ($\tau = 10$) and for $\theta = 25^\circ$, $\rho(\lambda)$ decreases by up to 30 % for wavelength below 750 nm, while for longer wavelengths $\rho(\lambda)$ increases by up to 8 % at about 780 nm. An exception is the dip in $\rho(\lambda)$ at about 1450 nm wavelength and wavelengths greater than 1800 nm. For $\theta > 60^\circ$, an increase in τ leads to an increase in $\rho(\lambda)$ throughout the simulated wavelength range.

The response of $\rho(\lambda)$ with increasing τ results from the transition from only direct to only diffuse radiation, which is controlled by the combination of θ and τ . Two theoretical extremes are distinguished: i) only direct radiation, also called "black-sky", and ii) only diffuse radiation, also called "white-sky". Natural conditions typically lie between these two extremes; they are referred to as "blue-sky" (Lucht et al., 2000). The amount of direct radiation ~~has two main controls~~two effects. The first effect acts on the canopy level, where diffuse radiation can penetrate deeper into the canopy and interacts with leaves that would be shaded under black-sky conditions. Consequently, for the same LAD and LAI, a greater total leaf area interacts with the incoming radiation (Jarvis et al., 1985; Freedman et al., 2001). Model simulations by ~~?~~Wolf et al. (2025a) showed an increase in broadband solar canopy albedo with increasing f_{dir} . Furthermore, the extinction of radiation is sensitive to the incident angle, which is equal to θ for direct radiation and approaches an effective value of about 60° under overcast conditions, due to the increasing contribution from diffuse radiation (~~Gardner and Sharp, 2010~~). ~~Therefore, $\rho(\lambda)$ is also sensitive to θ (see ?, Appendix D)~~(Pinty et al., 2005; Gardner and Sharp, 2010). The second factor acts on the individual leaf level, where the inherent directional reflection of radiation on surfaces is relevant. ~~While in the case of non-isotropic surfaces, direct radiation is reflected into a narrow solid angle, diffuse radiation is scattered over a wider solid angle (Schaepman-Strub et al., 2006). The angular-dependent scattering on surfaces under black-sky conditions is described by the bidirectional reflectance distribution function (BRDF, Schaepman-Strub et al., 2006). Under scattered in all directions with a significant portion of the radiation being scattered in a specific solid angle. Diffuse radiation is scattered almost equally over the entire hemisphere. (Asner, 1998; Schaepman-Strub et al., 2006). Both effects are directly linked to the directional reflectance under blue-sky conditions, the directional reflectivity conditions that~~ is described by the hemispherical-directional reflectance factor (HDRF, Schaepman-Strub et al., 2006), which includes the influence of the diffuse component in $F^\downarrow(\lambda)$. The reflection on the leaf-level combined with the LAD then determines the HDRF of the entire canopy. An ~~exemplary example of~~a HDRF for a spherical LAD is given in ~~the~~ Appendix B.




Figure 3. Ratio of downward irradiance $F_{\text{mea}}^{\downarrow}(\lambda)$ present during the synthetic measurements and downward irradiance $F_{\text{cal}}^{\downarrow}(\lambda)$ present during the calibration. The ratio is calculated for ice clouds, where $\tau_{\text{cal}} = 1$ and τ_{mea} is set to 0, 4, and 10. An exemplary As an example, a solar zenith angle θ of 25° is selected. The gray marked areas highlight the Sentinel-2 bands B2, B4, B8, B8a, and B11.

3.1.2 Effect of cloud changes on spectral reflectivity-reflectance above vegetation

The effects presented above, do not included-include changes in cloud conditions between RP calibration and actual measurement over vegetated surfaces, where $\tau_{\text{cal}} \neq \tau_{\text{mea}}$ and $F_{\text{cal}}^{\downarrow}(\lambda) \neq F_{\text{mea}}^{\downarrow}(\lambda)$. The effects of cloud changes between RP measurements to $F_{\text{mea}}^{\downarrow}(\lambda)$ are quantified and shown in Fig. 3as-, providing the illumination ratio $F_{\text{mea}}^{\downarrow}(\lambda)/F_{\text{cal}}^{\downarrow}(\lambda)$. The illumination ratio is calculated between the irradiance $F_{\text{cal}}^{\downarrow}(\lambda)$ that prevailed during the RP calibration and the irradiance $F_{\text{mea}}^{\downarrow}(\lambda)$ that is present during the above-vegetation measurements. The given illumination ratio represents the conditions below an ice cloud with $\tau_{\text{cal}} = 1$ and three value-values of τ_{mea} of 0, 4, and 10. In general, measurements where $\tau_{\text{cal}} < \tau_{\text{mea}}$ yield an illumination ratio less than one, because the reflectivity-reflectance at cloud top and the absorption inside the cloud become more intense, reducing the total $F^{\downarrow}(\lambda)$ below the cloud. For opposite conditions, where $\tau_{\text{cal}} > \tau_{\text{mea}}$, the illumination ratio is greater than one, since scattering and absorption are less during actual measurement compared to the RP calibration. Clearly visible are the ice absorption features, for example at 1500 nm wavelength, and a generally larger sensitivity of the illumination ratio towards longer wavelengths. Similar illumination ratios are calculated for liquid water clouds but with lower sensitivity for same values of τ . This is due to the smaller cloud droplet size of 10 μm compared to the ice particle size used in the simulations and the difference in the single-scattering phase function between liquid water droplets and ice crystals. In addition, differences in the imaginary part of the refractive indices lead to a slight spectral shift in the absorption features (Pilewskie and Twomey, 1987).




Figure 4. Panels (a) and (c): Synthetic spectral ~~reflectivity~~ reflectance $\rho(\lambda)$ measurements, when cloud conditions change from $\tau_{cal} = 0$ to $\tau_{mea} = 1$ and 4, respectively. Panels (b) and (d): Same as (a) and (b) but for $\tau_{cal} = 4$ and values of τ_{mea} of 1 and 4, respectively. In all panels: The black line represents $\rho(\lambda)$ under cloud-free conditions ($\tau_{cal} = \tau_{mea} = 0$) and the gray line represents $\rho(\lambda)$ under constant cloudy conditions ($\tau_{cal} = \tau_{mea} \neq 0$). Red lines represent ice water clouds and blue lines represent liquid water clouds. The gray marked areas highlight the Sentinel-2 bands B2, B4, B8, B8a, and B11.

The spectral distortion in $F^\downarrow(\lambda)$ due to cloud changes has an immediate effect on $\rho(\lambda)$ that is calculated with Eq. 1. For illustration, Fig. 4 shows $\rho(\lambda)$ of a vegetated surface for an intermediate value θ of 45° and for four different combinations of τ_{cal} and τ_{mea} . The ground truth $\rho(\lambda)$, as it would be obtained from satellites, with $\tau_{cal} = \tau_{mea} = 0$, is given by the black line. Only in the trivial case, when the airborne observations are performed under the same cloud-free conditions, the same value of $\rho(\lambda)$ would be measured. For reference, $\rho(\lambda)$ under constant cloud conditions during RP calibration and measurement, where $\tau_{cal} = \tau_{mea} \neq 0$, is indicated by the gray line.

The cases shown in Fig. 4a and c represent cloud-free conditions during the RP calibration ($\tau_{cal} = 0$), while clouds are present during the actual measurements with $\tau_{mea} = 1$ and $\tau_{mea} = 4$, respectively. Due to the cloud change with $\tau_{cal} > \tau_{mea}$, the amused $\rho(\lambda)$ is lower compared to the reference. The difference between measured $\rho(\lambda)$ and the reference increases with increasing difference $\Delta\tau = \tau_{mea} - \tau_{cal}$ and is more pronounced for ice clouds than for liquid water clouds of the same τ . For the ice cloud, $\rho(\lambda)$ at about 842 nm wavelength (Sentinel-2 B8) is reduced from about 0.38 ($\tau_{cal} = \tau_{mea} = 0$) to about

0.33 ($\tau_{\text{cal}} = 0, \tau_{\text{mea}} = 1$) and 0.26 ($\tau_{\text{cal}} = 0, \tau_{\text{mea}} = 4$). Similarly, the opposite situation is possible, where τ decreases after
 300 calibration ($\tau_{\text{cal}} > \tau_{\text{mea}}$), which is shown in the right column in Fig. 4. Since $\tau_{\text{cal}} > \tau_{\text{mea}}$, the measured $\rho(\lambda)$ overestimates the
 expected ground truth $\rho(\lambda)$, with the greatest bias in $\rho(\lambda)$ at about 1610 nm wavelength (Sentinel-2 B10) in the case of an ice
 cloud.

The examples given in Fig. 4b and d show that the estimated $\rho(\lambda)$ under constant cloud conditions (gray lines) are slightly
 enhanced compared to $\rho(\lambda)$ under cloud-free conditions but the change in $\rho(\lambda)$ due to differences between τ_{cal} and τ_{mea} are
 305 much greater and dominating. Therefore, the subsequent analysis primarily focuses on the contribution of changing cloud
 conditions.

3.2 Effect of clouds and cloud changes on two-band vegetation indices

Figure 4 showed that the spectral distortion due to differences in τ_{cal} and τ_{mea} ~~affect~~affects certain wavelengths stronger
 than others. Subsequently, all wavelength combinations of $\rho(\lambda_1)$ and $\rho(\lambda_2)$, with $\lambda_1, \lambda_2 \in [400, 2400]$, that might be used in
 310 two-band VIs following Eq. 8 are examined with respect to their sensitivity to cloud changes. A similar approach was used, for
 example, by Werner et al. (2013) to determine the effect of cirrus clouds on upward radiances used for cloud remote sensing.

Figure 5 shows the effect of cloud changes on γ expressed as $\Delta\gamma = \gamma_{\text{mea}} - \gamma$, with γ_{mea} obtained over vegetation using
 the latest RP calibration that was performed under τ_{cal} . The expected ground truth γ is the value that would be expected if
 an immediate estimate of κ would be available or cloud conditions would not have changed ($\tau_{\text{cal}} = \tau_{\text{mea}} \neq 0$). A negative
 315 $\Delta\gamma$ therefore indicates an underestimation of the true γ and vice versa. The combinations of τ_{cal} and τ_{mea} were chosen to
 represent optically thin cirrus with changes in τ that could occur between RP calibration measurements, e.g. 10 minutes apart.
 Each panel in Fig. 5 is divided into four quadrants, starting with the first quadrant (Q1) in the lower right part and turning
 counter-clockwise. It follows from Eq. 8 that $\Delta\gamma$ is point-symmetric with respect to the diagonal from the origin to the upper
 right corner. Thus, for any combination of λ_1 and λ_2 that causes an overestimation or underestimation, the inverse wavelength
 320 combination λ_2 and λ_1 will cause the same bias $\Delta\gamma$, but with opposite sign. In addition to NDVI and NDWI, **exemplary**
 selected two-band VIs that follow Eq. 8 and use SWIR wavelengths were added to the plot, namely: the modified normalized
 difference water index (MNDVI; Jurgens, 1997), the modified normalized difference water index (MNDWI; Xu, 2006), and
 the normalized difference infrared index (NDII; Hardisky et al., 1983). The list is not comprehensive and many more VIs exist,
 e.g., ~~listed~~tabulated by Jones et al. (2012), Zeng et al. (2022), and Montero et al. (2023). Wavelengths affected by strong water
 325 vapor absorption between 1350 and 1400 nm as well as 1830 and 1920 nm wavelengths have been masked because $F^\downarrow(\lambda)$ and
 $F^\uparrow(\lambda)$ are close to zero causing numerical unstable values of $\rho(\lambda)$.

The largest absolute values of $|\Delta\gamma|$, indicated by dark red or dark blue colors, occur for wavelength combinations with the
 greatest spectral distance $|\Delta\lambda| = |\lambda_1 - \lambda_2|$. Also wavelength combinations with one wavelength greater than 1400 nm, i.e., in
 Q1 and Q3, are affected by cloud transitions, since the spectral slope in $F^\downarrow(\lambda)$ is most pronounced towards longer wavelengths
 330 (see Fig. 3).

In contrast, independent of the selected combination of τ_{cal} , τ_{mea} , and θ generally low values of $|\Delta\gamma|$ are found for wave-
 length combinations with $\lambda < 1400$ nm (Q4). Additionally, small values of $\Delta\gamma$ also occur along the diagonal from the origin




Figure 5. Absolute difference $\Delta\gamma = \gamma_{\text{mea}} - \gamma$ for change in illumination conditions, when cloud optical thickness is different during the reflectance panel measurement (τ_{cal}) and actual condition during the measurement (τ_{mea}). Absolute differences in γ are given for combinations of λ_1 (x -axis) and λ_2 (y -axis) simulated for an ice cloud, and for solar zenith angles θ of 25° (top) and 50° (bottom). Left column: Absolute differences, when $\tau_{\text{cal}} = 2$ and $\tau_{\text{mea}} = 0$. Right column: Absolute differences, when $\tau_{\text{cal}} = 2$ and $\tau_{\text{mea}} = 4$. Red areas show an overestimation of γ , while blue areas indicate an underestimation of γ . Wavelength combinations of γ affected by strong water vapor absorption between 1350 and 1400 nm as well as 1830 and 1920 nm have been masked.

Table 3. Difference $\Delta\gamma = \gamma_{\text{mea}} - \gamma$ between estimated vegetation index γ_{mea} and ground truth vegetation index γ for four vegetation indices for the four example cases given in Fig. 5a–d. The calculations assume a spherical LAD. Sentinel-2 band ratios were taken from Montero et al. (2023).

Vegetation index	Sentinel-2 bands ratios	$\Delta\gamma$			
		Case (a)	Case (b)	Case (c)	Case (d)
NDVI	$(B8 - B4) / (B8 + B4)$	0.043	−0.011	0.009	−0.007
NDWI ₁₂₄₀	$(B3 - B8) / (B3 + B8)$	−0.048	0.018	−0.022	0.015
NDWI ₁₆₄₀	$(B8a - B11) / (B8a + B11)$	0.184	−0.212	0.216	−0.267
MNDVI	$(B8 - B12) / (B8 + B12)$	0.111	−0.127	0.112	−0.162
MNDWI	$(B3 - B11) / (B3 + B11)$	0.08	−0.113	0.156	−0.148
NDII	$(B8 - B11) / (B8 + B11)$	0.173	−0.205	0.208	−0.264

to the upper right corner, i.e., the smaller $|\Delta\lambda|$ becomes. While small $|\Delta\lambda|$ would minimize the cloud effect, the proximity of λ_1 and λ_2 limits the information content that can be extracted from spectral ratios. A trade-off between information content
335 and small cloud influence is required and Fig. 5 provides guidance to choose suitable wavelength combinations.

Figure 5a ~~exemplary shows~~ illustrates $\Delta\gamma$ for $\theta = 25^\circ$, where $\tau_{\text{cal}} = 2$ decreases to $\tau_{\text{mea}} = 0$. The decrease in τ after the RP calibration leads to predominantly negative $\Delta\gamma$ (blue colors) in Q3 and positive $\Delta\gamma$ (red colors) in Q2, indicating a respective underestimation and overestimation of the true γ . In the selected case, $\Delta\gamma$ of NDVI (black circle) is small compared to the full potential range of NDVI (see Table 3). Greater values of $\Delta\gamma$ are calculated for NDWI₁₆₄₀ with up to 0.184, resulting from the
340 second wavelength being located in the SWIR region that is subject to the spectral slope in $F^\downarrow(\lambda)$.

The second example in Fig. 5b shows the opposite transition from an optically thin cloud during the RP calibration with $\tau_{\text{cal}} = 2$ to an optically thicker cloud during measurement $\tau_{\text{mea}} = 4$. This leads to an inverted pattern of $\Delta\gamma$. Although the change in cloudiness $|\Delta\tau| = \tau_{\text{cal}} - \tau_{\text{mea}}$ is similar to the example in Fig. 5a, the magnitude of $|\Delta\gamma|$ for all wavelength combinations in Q2 and Q4 are slightly smaller, while the effect is greater for wavelength combinations in Q1 and Q3. This shows
345 that the bias $|\Delta\gamma|$ from changing cloud conditions is affected by $\Delta\tau$ but also depends on the absolute value of τ_{cal} during calibration. For this example, NDVI and NDWI₁₆₄₀ are subject to biases $\Delta\gamma$ of 0.011 and −0.212, respectively.

Figure 5c shows $\Delta\gamma$ for the same combination of τ_{cal} and τ_{mea} that is given in Figure 5a but for a greater value of θ of 50° . This leads to greater absolute values of $|\Delta\gamma|$ in Q1–Q3. An exception is Q4, which is characterized by $\Delta\gamma$ around 0 for all wavelengths with less pronounced water vapor absorption features compared to θ of 25° . This indicates that with increasing
350 θ , changes in τ lead to reduced biases in γ for wavelengths less than 1400 nm, while larger biases in γ are expected for VIs using wavelengths beyond 1400 nm. In general, the effects of changing cloud conditions on $\rho(\lambda)$ and $\Delta\gamma$ are less pronounced for the planophile LAD, while for the erectophile LAD greater effects were determined. This results primarily from the higher sensitivity of $\rho(\lambda)$ on θ in case of the erectophile LAD (Wolf et al., 2025a). An overview of $\Delta\gamma$ for all example two-band VIs derived for the four cases that are marked in Fig. 5 are given in Table 3.




Figure 6. Top row: Absolute values of NDVI as obtained for solar zenith angles θ of 25° , 50° , and 70° . Subsequent rows same as top row but for NDWI₁₆₄₀ and EVI. Color-coded is the cloud optical thickness τ_{cal} that was present during the reflectance panel measurement. The cloud optical thickness τ_{mea} during the measurement is given on the x -axis. Simulations for the liquid water cloud are given by solid lines and simulations for the ice cloud are given by the dotted lines. [All simulations base on the assumption of a spherical LAD.](#) The variability in VI due to the presence of clouds ($\tau_{cal} = \tau_{mea}$) and the associated change in the surface [reflectivity](#)[reflectance](#), is highlighted in gray.

355 Subsequently, the effects of changing τ between RP calibration and measurement on three selected VIs are investigated.

3.2.1 Effect of cloud changes on the normalized differential vegetation index (NDVI)

The estimated NDVI for three values of θ depending on the combination of τ_{cal} and τ_{mea} is shown in the top row of Fig. 6.

First, we consider the trivial case of cloud-free conditions with $\tau_{\text{cal}} = \tau_{\text{mea}} \approx 0$, where NDVI of about 0.87, 0.9, and 0.92 are calculated for θ of 25° , 50° , and 70° , respectively. These cases are marked by the dark-blue dots and represent the reference
360 NDVI. The increase of NDVI with increasing θ is related to scattering and absorption at gas molecules and aerosol particles. Next, cloudy conditions are considered, where $\tau_{\text{cal}} \approx \tau_{\text{mea}} \neq 0$ (colored dots), showing that the NDVI is enhanced by the presence of the cloud (gray highlighted area), which influences $f_{\text{dir}}(\lambda)$ and $\rho(\lambda)$. The ~~largest-greatest~~ variability is found for $\theta = 25^\circ$, where NDVI increases from 0.87 to 0.91 with increasing τ , ~~indicating a tendency to overestimate~~ which could be interpreted as an overestimation of vegetation health. Smaller effects are found for $\theta = 50^\circ$ and for $\theta = 70^\circ$ the NDVI even
365 decreases. It ~~is noteworthy-should be noted~~ that measuring NDVI under cloud-free conditions at different times of day, i.e., different θ , causes the same variability as measuring NDVI at a fixed time with $\theta = 25^\circ$ but under different cloud conditions, for example, on consecutive days.

For cases where ~~the~~ cloud conditions differ between τ_{cal} and τ_{mea} , the change of τ_{mea} for fixed τ_{cal} is given by the colored lines. Moving to the right along lines of same τ_{cal} can be understood as the advection of an optically thicker cloud during
370 the measurement, while moving left represents the advection of an optically thinner cloud. The largest differences between measured and reference NDVI generally occur for $\theta = 25^\circ$, and become successively smaller with increasing θ . Thus, the NDVI is subject to the largest biases from cloud transitions when θ is small. The advection of an optically thicker cloud after the RP measurement ($\tau_{\text{cal}} < \tau_{\text{mea}}$) results in a decrease in NDVI and an underestimation of the expected value. For $\tau_{\text{cal}} = 0$ an extreme increase of τ_{mea} from 0 to 40 results in a decrease in NDVI from 0.87 to 0.84, ~~indicating a tendency to underestimate~~
375 ~~the-vegetation-healthstatus~~ which could be interpreted as an underestimation of vegetation health. The advection of an optically thinner cloud after the RP measurement results in an overestimation of the actual NDVI value. For example, the combination of $\tau_{\text{cal}} = 40$ and $\tau_{\text{mea}} = 0$ increases the NDVI from 0.92 to 0.94. The aforementioned extreme changes in τ from 0 to 40, and vice versa, between RP overflights are unlikely in the case of stratiform clouds. These extremes of τ have been selected to estimate the maximum envelope for biases in NDVI. Biases in NDVI, retrieved under more homogeneous conditions, will be
380 smaller. However, even seemingly stratiform clouds vary in τ and therefore bias the retrieved values of NDVI. These variations have the greatest impact, when the RP overflight is performed under cloud-free conditions ($\tau_{\text{cal}} = 0$) or under conditions with small values of τ_{cal} , since the slope of the curves is greatest in these situations (see, for example, the blue lines in Fig. 6a–c).

Comparing the responses of NDVI obtained below liquid and ice water clouds shows that both cloud types cause similar biases. However, the magnitude is generally greater for the ice cloud with the largest difference for small values of θ and for
385 τ_{mea} between 10 and 30. However, it is acknowledged that the calculated deviations between the measured and expected NDVI are small, considering the selected differences between τ_{cal} and ~~τ_{cal}~~ τ_{mea} , and the typical range of NDVI between 0 and 1.

3.2.2 Effect of cloud changes on the normalized differential water index (NDWI)

Remote sensing of NDWI_{1640} is based on the Sentinel-2 bands B8a and B11, a combination that is more sensitive to changes in τ compared to the NDVI, because one wavelength is located in the SWIR (Q3 in Fig. 5). The middle row in Fig. 6 shows the change of NDWI induced by cloud changes.

Regardless of θ , the bias in NDWI_{1640} is under cloudy conditions with $\tau_{\text{cal}} = \tau_{\text{mea}}$ is of similar magnitude compared to the NDVI. A variation in NDWI_{1640} with maximal values of ± 0.02 (gray highlighted area) is calculated, which is small compared to the influence of changes in cloudiness between RP calibrations. Here, the change in cloud conditions is ~~exemplarily~~ discussed for $\theta = 25^\circ$. When the RP measurements was performed for $\tau_{\text{cal}} = 0$ the advection of an optically thicker cloud, represented by an increase in τ_{cal} from 0 to 40, leads to an increase in NDWI_{1640} from 0.28 to 0.89 ($\Delta\gamma = 0.61$), which is relevant considering the typical range between ~~-1~~ and 1. In such scenarios, the interpreted true health status would be well overestimated using NDWI_{1640} . For the same θ , a calibration under cloudy sky with $\tau_{\text{cal}} = 40$ and a subsequent decrease in τ_{mea} from 40 to 0 results in a decrease in NDWI_{1640} from 0.59 to -0.43 ($\Delta\gamma = -1.02$). Even though $\Delta\tau$ is equal in both cases, the resulting $\Delta\gamma$ differs, which emphasizes the dependence of the bias $\Delta\gamma$ on the absolute τ_{cal} . As for the NDVI, the transitions in τ from 0 to 40, or vice versa, are extreme cases. For practical applications, variations in τ_{mea} for a given τ_{cal} are more relevant. For the NDWI_{1640} , the slopes in Fig. 6d-f are shifted in absolute terms but are constant for the same values of θ . Therefore, variations in τ_{mea} between RP overflights will cause the same bias in NDWI_{1640} , independent of τ_{cal} . Similar to NDVI, the effect of ice clouds on NDWI_{1640} is qualitatively similar to the effect from liquid water clouds, however the magnitude is almost twice as large. ~~While the chosen values of τ_{cal} and τ_{mea} represent extreme cases, they can occur under heterogeneous cloud conditions. Considering homogeneous cloud transitions with $\tau_{\text{cal}} = 4$ and a subsequent decrease to $\tau_{\text{mea}} = 0$, NDWI_{1640} still decreases from 0.29 to 0.2. Even for the latter example, the resulting biases in NDWI_{1640} are considered as relevant, considering the typical range between -1 and 1 , and considering that these biases become even larger with increasing values of θ . Consequently, care must be taken when estimating NDWI_{1640} or other VIs with wavelength combinations in the SWIR when clouds are present. Unlike the NDVI, the bias in NDWI_{1640} under constant cloud conditions is negligible compared to the variation caused by cloud changes.~~

3.3 Effect of clouds and cloud changes on the three-band enhanced vegetation index (EVI)

As an example for a three-band VI, the effects of changing cloud conditions between RP calibration and measurement are shown for the EVI (see Fig. 6g-i). The effect of ~~a change~~ changes in $f_{\text{dir}}(\lambda)$ under constant conditions ($\tau_{\text{cal}} = \tau_{\text{mea}}$) is generally small (gray area) compared to the variation in EVI that is associated with the mismatch between τ_{cal} and τ_{mea} . Irrespective of θ , the advection of an optically thicker cloud ($\tau_{\text{cal}} < \tau_{\text{mea}}$) causes an increase in estimated EVI, which leads to an overestimation of the true EVI. Conversely, the advection of an optically thinner cloud ($\tau_{\text{cal}} > \tau_{\text{mea}}$) leads to and underestimation of the true EVI. The bias gets more pronounced with increasing difference between τ_{cal} and τ_{mea} , and with increasing absolute value of τ_{cal} . The magnitude of these biases are more pronounced for the ice cloud than for the liquid water cloud. Furthermore, the biases become larger with increasing θ , as well as with increasing difference between τ_{cal} and τ_{mea} , which is particularly

420 pronounced in combination with small τ_{cal} . Thus, estimated EVI are less susceptible to changes in cloud conditions at low values of θ , e.g., around noon or at low latitudes, compared to cloud changes during measurements with the Sun close to the horizon or generally at higher latitudes. As an example, a calibration performed under $\theta = 25^\circ$ and cloud-free conditions followed by the advection of an optically thicker ice cloud with $\tau_{\text{mea}} = 4$ results in an increase in EVI from the expected value of 0.67 to 0.75. For the same transition in τ but for $\theta = 70^\circ$ an increase from 0.67 to 1.04 is estimated. In both cases, the ~~actual~~
425 inferred ground-truth vegetation health would be overestimated.

The different response of EVI to cloud changes compared to the NDVI is related to two factors. First, the equation to calculate the EVI is fundamentally different from the two-band VIs. The use of an additional third spectral band at 490 nm wavelength makes the EVI generally more responsive to the spectral slope in $F^\downarrow(\lambda)$ caused by absorption. Second, the spectral slope effect is amplified by the pre-factors in Eq. 9, since re-writing Eq. 9 results in:

$$430 \quad \text{EVI} = G \cdot \frac{a - b}{a + C_1 \cdot b - C_2 + L/\rho_{\text{B2}}}, \quad (10)$$

with the ~~reflectivity~~-reflectance ratios $a = \rho_{\text{B8}}/\rho_{\text{B2}}$ and $b = \rho_{\text{B4}}/\rho_{\text{B2}}$ at Sentinel-2 bands B2, B4, and B8 (see Table 1). The ratios a and b are affected by the spectral slope in the illumination ratio (see Fig. 3). The spectral slope is greater for a than for b , since $\Delta\lambda$ is greater in a than in b . The differences between a and b do not cancel out and are amplified by the pre-factor C_1 . The term L/ρ_{B2} also contributes, since the constant L is inversely scaled with ρ_{B2} and band B2 is sensitive to atmospheric
435 scattering.

~~The bias in EVI has potential implications for other estimated canopy parameters. For example, the EVI is frequently used for the remote sensing of the LAI.~~

3.4 Implication of biases in vegetation indices on estimated biophysical properties

In vegetation remote sensing, VIs are used to estimate biophysical properties such as LAI, gross primary production (GPP),
440 fresh and dry biomass, and vegetation water content (Gitelson et al., 2006; Hong et al., 2007; Ahmadian et al., 2016). For example, Boegh et al. (2002) proposed an empirical linear regression given by:

$$\text{LAI} = 3.618 \cdot \text{EVI} - 0.118, \quad (11)$$

where the LAI scales linearly with the EVI. Due to the direct relationship between EVI and LAI, biases in EVI are linearly scaled by the pre-factor 3.618 and lead to biased estimates of LAI. For example, when a calibration for ~~$\theta = 50^\circ$~~ $\theta = 50^\circ$ was
445 performed under cloud conditions with $\tau_{\text{cal}} = 0$ but the measurement is influenced by an ice cloud with $\tau_{\text{mea}} = 10$, ~~is introduced~~ This results in a bias in EVI of 0.25 ~~is introduced~~, which corresponds to a ~~bis-bias~~ in LAI of about $\Delta\text{LAI} = 0.25 \cdot 3.618 = 0.9$.

Similarly, the gross primary product is an important measure in the context of the global carbon cycle, since it indicates how quickly an ecosystem accumulates biomass (Gitelson et al., 2006). Several attempts have been made to estimate GPP based on NDVI and EVI; for example Rahman et al. (2005), Wu et al. (2009), or Zhou et al. (2014). Correlations have been
450 estimated using in-situ carbon uptake flux measurements and remotely sensed NDVI or EVI. Although the correlations between GPP and EVI or NDVI vary by site and crop type, stable correlations could be identified. Using the correlations

455 from Rahman et al. (2005) or Zhou et al. (2014), a variation in EVI of $\Delta\text{EVI} = 0.2$ would yield variations in estimated GPP of about $2.6 \text{ gC m}^{-2} \text{ d}^{-1}$. Zhou et al. (2014) also provided relationships for NDVI, where a variation of $\Delta\text{NDVI} = 0.2$ would result in a variation in GPP of about $0.51 \text{ gC m}^{-2} \text{ d}^{-1}$. Considering that the total range of GPP spans from $0 \text{ gC m}^{-2} \text{ d}^{-1}$, when the NDVI or EVI is zero, to $8 \text{ gC m}^{-2} \text{ d}^{-1}$, when NDVI or EVI is close to one, the bias in estimated GPP from biases in EVI is of relevance.

460 Several studies have attempted to estimate vegetation water content (VegWC) based on NDVI (Hong et al., 2007). The relationships derived between NDVI and VegWC vary greatly depending on the year, location, and crop type. Therefore, the identified cloud-induced biases in NDVI are expected to play only a minor role in the total uncertainty of estimated VegWC. Ahmadian et al. (2016) aimed to estimate fresh and dry biomass, and LAI based on derived NDVI and EVI. These relationships are strongly dependent on vegetation type. Therefore, cloud-induced variations in NDVI are expected to contribute only a small amount to the total uncertainty in estimated fresh and dry biomass. However, since EVI is more sensitive to cloud-induced biases than NDVI, the uncertainty in EVI can be of a similar order of magnitude compared to the uncertainty in the relationships.

465 4 Summary and conclusions

This paper presented results of coupled atmosphere-vegetation radiative transfer (RT) simulations, using the coupled RT models libRadtran and SCOPE2.0, to systematically investigate biases in remotely sensed vegetation indices (VIs) due to changing cloud conditions. Simulations were performed for a stratiform liquid water cloud and an ice cloud representative of high-level cirrus. The ~~cloud optical thickness~~ optical thickness of the cloud τ ~~was~~ varied between 0 and 40, and ~~the~~ solar zenith angles θ ~~ranging~~ ranged from 25° to 70° ~~were covered~~. The optical properties were represented by a spherical leaf angle distribution (LAD) and a leaf area index (LAI) of 3. The simulations were designed to resemble below-cloud measurements of downward irradiance $F^\downarrow(\lambda)$ and spectral surface ~~reflectivity~~ reflectance $\rho(\lambda)$ used for remote sensing of vegetation from airborne observations. The synthetic measurements mimic the typical observation strategy during field measurements, where reflectance panels (RPs) are used to calibrate the surface ~~reflectivity~~ reflectance measurements.

475 Field measurements can be performed below clouds, which reduce the direct fraction $f_{\text{dir}}(\lambda)$ in $F^\downarrow(\lambda)$, which is a controlling factor of the surface ~~reflectivity~~ reflectance. Clouds also change the downward irradiance $F^\downarrow(\lambda)$ spectrally and in absolute terms through spectral dependent scattering and absorption. Changes in cloud conditions, expressed as cloud optical thickness τ , between periodic RP calibrations are expected to cause spectral distortions in the estimated $\rho(\lambda)$ and thus introduce biases in VIs estimated in the presence of clouds.

480 The synthetic observations allowed to separate the effect from changes in $f_{\text{dir}}(\lambda)$ from the changes in τ between RP calibrations. For a solar zenith angle $\theta = 25^\circ$, a reduction in $f_{\text{dir}}(\lambda)$ led to an increase in $\rho(\lambda)$ for wavelengths greater than 650 nm compared to cloud-free conditions, while for wavelengths below 650 nm $\rho(\lambda)$ was reduced. For $\theta > 55^\circ$, $\rho(\lambda)$ below 650 nm wavelengths was also increased with decreasing $f_{\text{dir}}(\lambda)$. The change in $\rho(\lambda)$ under cloudy conditions therefore depends on the

combination of θ and τ , and the resulting $f_{\text{dir}}(\lambda)$. The effect of changes in $f_{\text{dir}}(\lambda)$ on estimated VIs was found to be small
485 compared to the effect of changes in τ between RP calibrations.

The influence of cloud changes on two-band VIs was investigated for the normalized difference vegetation index (NDVI) and the normalized water index (NDWI). Other examples of two-band VIs were provided. The influence of cloud changes on the enhanced vegetation index (EVI), which is representative for three-band VIs, was also investigated. The effect of cloud changes on narrow-band VIs was found to be small for VIs using wavelength combinations below 1400 nm and for decreasing
490 spectral distance between wavelength combinations. An optimum must be found between the information content obtained from the wavelength ratio and the sensitivity of the respective wavelengths pair to cloud changes. Guidance was provided by presenting the cloud-induced bias for potential two-band wavelength combinations between 400 and 2400 nm. For wavelengths greater than 1400 nm, the sensitivity to cloud changes was found to increase and is particularly pronounced in the proximity and within the water absorption bands. Wavelengths smaller than 1400 nm, were found to be less impacted. For the NDVI,
495 a generally low sensitivity was found. For an intermediate value of $\theta = 50^\circ$, the transition in τ from 0 to 10 led to a bias of about -0.035 . With increasing θ and for same transition in τ , the bias in NDVI increased, while a decrease in θ resulted in lower biases. The NDWI ~~is~~ subject to greater biases of up to 0.26 for an intermediate $\theta = 50^\circ$ and a transition in τ from 0 to 10. ~~Biases in the~~ The biases in NDWI generally increase with increasing θ . The EVI was also found to be sensitive to changes in τ . A transition in τ from 0 to 10 in combination with $\theta = 50^\circ$ led to an overestimation of the EVI by 0.25. For the same
500 transitions in τ , the bias in EVI decreases with decreasing θ . The leaf area index estimated from EVI using empirical equations is directly affected by potential biases in EVI. For ~~the exemplary example, the~~ biases in EVI of 0.25 would ~~case aa cause a bias~~ in LAI of 0.9. Similarly, biases in EVI and NDVI of about ± 0.2 can lead to biases in estimated gross primary product of up to $\pm 2 \text{ gC m}^{-2} \text{ d}^{-1}$. The impact for other LADs, such as the erectophile and planophile, has been investigated. The erectophile LAD is more susceptible to biases in VIs under constant cloud conditions than the spherical LAD, while the bias in VIs with
505 underlying planophile LAD is almost unaffected. Although the absolute value of the VI biases are determined by the LAD, the relative change due to a transition in τ between RP measurements was found to be constant among the three LADs.

The presented analysis showed that the practice of using relative measurements with RPs is prone to uncertainties. With the improvement of drone technology and their ability to carry heavier payloads, and in combination with the advancements in sensor technology, it would be advantageous to measure $F^\downarrow(\lambda)$ directly instead of relying on relative measurements of $\rho(\lambda)$
510 using RPs. Uncertainties associated with changing cloud conditions could be minimized.

The simulations presented here are limited in their representation because the full natural variability in vegetation and canopy types could not be covered. Furthermore, the assumed default values used in the atmosphere and vegetation RT simulations can influence the results. In particular, the assumed LAI, LAD, plant dry matter, and soil properties in the vegetation RT simulations, and the assumed atmospheric profile in the atmosphere RT simulations can influence the results ~~(?)~~ (Wolf et al., 2025a). Since
515 the natural variability cannot be covered in a single study, the presented work is to be interpreted as a conceptual study and to highlight the potential impacts of clouds during field observations. It is also emphasized that the presented simulations are based on one-dimensional RT only and lack a more detailed representation of the three-dimensional nature of RT below heterogeneous cloud fields. This is particularly problematic in the vicinity of clouds, where a nearby cloud casts a shadow




Figure A1. Same as Fig. 6 but for the erectophyll leaf angle distribution.

while no cloud is in the zenith. Nevertheless, we argue that the presented study can be used as a first approximation for the transition between cloud-free and cloudy regions.

Data availability. The simulated spectra of radiance, irradiance, and vegetation albedo are made available via NetCDF files. The data are available on the Zenodo platform via <https://doi.org/10.5281/zenodo.15275610> (Wolf et al., 2025b).

Appendix A: Effect of cloud changes on the NDVI, $NDWI_{1640}$, and EVI for erectophyll and planophyll LADs




Figure A2. Same as Fig. 6 but for the planophile leaf angle distribution.

525 This section provides an overview of how an erectophile or planophile leaf angle distribution (LAD) affects biases in normalized
differential vegetation index (NDVI), normalized difference water index (NDWI₁₆₄₀), and the enhanced vegetation index
(EVI), compared to the spherical LAD presented in the main text. First, we consider constant cloud optical thickness during
the reflectance panel (RP) overflight with τ_{cal} and the actual measurement with τ_{mea} , such that $\tau_{cal} = \tau_{mea}$. Comparing Fig. 6
and Fig. A1 reveals a greater variation in the NDVI bias for the erectophile LAD than for the spherical LAD (gray highlighted
area). The biases are greatest for small solar zenith angle θ , when incoming radiation can penetrate deep into the canopy. In
530 the case of $\theta = 25^\circ$, the NDVI bias varies between 0.67 and 0.89 for the erectophile LAD, compared to a variation between
0.75 and 0.91 in case of the spherical LAD. In this particular case, the variation in NDVI exceeds the bias that results from a
transition in cloud conditions, when $\tau_{cal} \neq \tau_{mea}$. However, when θ is greater than 25° and the penetration depth of the radiation

into the canopy is reduced, the bias from changing cloud conditions between RP overflights again dominates the NDVI bias. Comparing Fig. 6 and Fig. A2 shows that for the planophile LAD and $\tau_{\text{cal}} = \tau_{\text{mea}}$, the variation in NDVI is negligible, with a nearly constant value of around 0.89. Similar effects are found for NDWI₁₆₄₀ and EVI. In general, the greatest biases occur for the erectophile LAD, independent of θ . Within on LAD the greatest biases occur for the lowest values of θ . Taking into account changes in cloud conditions, i.e., when $\tau_{\text{cal}} \neq \tau_{\text{mea}}$, the comparison of Fig. 6, Fig. A1, and Fig. A2 shows that for the same values of θ , the relative variations in VI due to cloud changes are the same for all three LADs and VIs, which is indicated by a constant sensitivity of the VIs on τ_{mea} independent of the value of τ_{cal} . However, the absolute values in the VIs are shifted along the y -axis depending on τ_{cal} and the selected LAD.

Appendix B: Simulated reflectance factors and reflectance functions

The intrinsic reflectivity properties of a surface are given by its bidirectional reflectance distribution function (BRDF). The BRDF quantifies the reflection and scattering of incident radiation on the surface from one direction of the hemisphere to another. The spectral BRDF f_{BRDF} , in units of sr^{-1} , gives the ratio between the reflected radiance $I_{\text{r}}^{\uparrow}(\theta_{\text{i}}, \varphi_{\text{i}}; \theta_{\text{r}}, \varphi_{\text{r}}; \lambda)$ with respect to the incident irradiance $F_{\text{i}}^{\downarrow}(\theta_{\text{i}}, \varphi_{\text{i}}; \lambda)$. The resulting f_{BRDF} depends on the zenith (θ_{i}) and the azimuth angle (φ_{i}) of the incoming radiation, and the zenith (θ_{r}) and azimuth angle (φ_{r}) of the reflected radiation. The spectral BRDF f_{BRDF} also depends on the wavelength λ and is given by:

$$f_{\text{BRDF}} = \frac{dI_{\text{r}}^{\uparrow}(\theta_{\text{i}}, \varphi_{\text{i}}; \theta_{\text{r}}, \varphi_{\text{r}}; \lambda)}{dF_{\text{i}}^{\downarrow}(\theta_{\text{i}}, \varphi_{\text{i}}; \lambda)}, \quad (\text{B1})$$

given in the unit steradian (sr^{-1}).

Under atmospheric conditions, where $F^{\downarrow}(\lambda)$ is composed of a direct and a diffuse fraction, the BRDF cannot be measured (Pinty et al., 2005; Schaepman-Strub et al., 2006). What can be observed is the spectral ~~reflectivity~~ reflectance $\rho(\lambda)$ and the hemispherical-directional reflectance factor (HDRF), both of which include radiation from the entire hemisphere and not just from a single direction. The hemispherical-directional reflectance factor is formalized by:

$$\begin{aligned} R_{\text{HDRF}} &= \frac{dI_{\text{r}}^{\uparrow}(\theta_{\text{i}}, \varphi_{\text{i}}, 2\pi; \theta_{\text{r}}, \varphi_{\text{r}})}{dF_{\text{o}}^{\downarrow}(\theta_{\text{i}}, \varphi_{\text{i}}, 2\pi)} \\ &= R_{\text{HDRF}}(\theta_{\text{i}}, \varphi_{\text{i}}; \theta_{\text{r}}, \varphi_{\text{r}}) \cdot f_{\text{dir}} \\ &\quad + R(2\pi; \theta_{\text{r}}, \varphi_{\text{r}}) \cdot (1 - f_{\text{dir}}). \end{aligned} \quad (\text{B2})$$

Thus, R_{HDRF} represents the ratio of the reflected incident radiation from a surface in relation to an ideal, lambertian surface. Furthermore, in Eq. B2 it is assumed that the incident, diffuse radiation is isotropic (Schaepman-Strub et al., 2006).

Figure B1a and e show R_{HDRF} at 550 nm wavelength for cloud-free conditions with $\tau = 0$ for θ of 25° and 70° , respectively. Scattering by gas molecules and aerosol particles lead to direct fraction $f_{\text{dir}}(\lambda) = 0.94$ just below 1. Under these conditions the first term in Eq. B2 dominates and the directional effect of the incoming radiation on the reflected radiation is most pronounced leading to the clear development of the hot spot that follows the solar zenith angle of the Sun (see white markers). With




Figure B1. All sub-panels show the normalized hemispherical-directional reflectance factor R_{HDRF} at 550 nm wavelength. Panels (a), (b), (e), and (f) present polar plots of R_{HDRF} for combinations of solar zenith angle θ and cloud optical thickness τ . Panels (c), (d), (g), and (h) present cross-sections of R_{HDRF} along lines of 0°–180° and 90°–270° azimuth.

increasing values of τ and decreasing $f_{\text{dir}}(\lambda)$, the second term in Eq. B2 dominates since the radiation reaching the surface is predominantly diffuse and almost isotropic. The directional component of the surface ~~reflectivity~~ reflectance vanishes, causing the hot spot to disappear and leading to a general smoothing of R_{HDRF} , which is also confirmed in the cross-sections in panels

565 c, d, g, and h.

Author contributions. **KW** designed and implemented the model coupling, performed the simulations, and drafted the manuscript. **EJ**, **AE**, and **MW**, contributed to the preparation and revisions of the manuscript. **MS**, **HF**, and **AHu** supported during the model set-up and provided suggestions for the manuscript.

Competing interests. The authors declare no competing interest.

570 *Acknowledgements.* We thank the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, which is a research center of the Deutsche Forschungsgemeinschaft (DFG). We also thank the Saxon State Ministry for Science, Culture and Tourism (SMWK) for funding through grant 3-7304/44/4-2023/8846. We would also like to acknowledge the three anonymous reviewers whose comments helped to improve the final manuscript.

References

- 575 Aasen, H. and Bolten, A.: Multi-temporal high-resolution imaging spectroscopy with hyperspectral 2D imagers – From theory to application, *Remote Sens. Environ.*, 205, 374–389, <https://doi.org/10.1016/j.rse.2017.10.043>, 2018.
- Aasen, H., Burkart, A., Bolten, A., and Bareth, G.: Generating 3D hyperspectral information with lightweight UAV snapshot cameras for vegetation monitoring: From camera calibration to quality assurance, *ISPRS*, 108, 245–259, <https://doi.org/10.1016/j.isprsjprs.2015.08.002>, 2015.
- 580 Aebi, C., Gröbner, J., Kazadzis, S., Vuilleumier, L., Gkikas, A., and Kämpfer, N.: Estimation of cloud optical thickness, single scattering albedo and effective droplet radius using a shortwave radiative closure study in Payerne, *Atmos. Meas. Tech.*, 13, 907–923, <https://doi.org/10.5194/amt-13-907-2020>, 2020.
- Ahmadian, N., Ghasemi, S., Wigneron, J.-P., and Zölitz, R.: Comprehensive study of the biophysical parameters of agricultural crops based on assessing Landsat 8 OLI and Landsat 7 ETM+ vegetation indices, *GISci. Remote Sens.*, 53, 337–359, <https://doi.org/10.1080/15481603.2016.1155789>, 2016.
- 585 Anderson, G. P., Clough, S. A., Kneizys, F. X., Chetwynd, J. H., and Shettle, E. P.: AFGL atmospheric constituent profiles, *Environ. Res. Pap.*, 954, 1–46, 1986.
- Asner, G. P.: Biophysical and Biochemical Sources of Variability in Canopy Reflectance, *Remote Sens. Environ.*, 64, 234–253, [https://doi.org/10.1016/S0034-4257\(98\)00014-5](https://doi.org/10.1016/S0034-4257(98)00014-5), 1998.
- 590 Baldocchi, D. D., Wilson, K. B., and Gu, L.: How the environment, canopy structure and canopy physiological functioning influence carbon, water and energy fluxes of a temperate broad-leaved deciduous forest—an assessment with the biophysical model CANOAK, *Tree Physiol.*, 22, 1065–1077, <https://doi.org/10.1093/treephys/22.15-16.1065>, 2002.
- Bausch, W. C.: Soil background effects on reflectance-based crop coefficients for corn, *Remote Sens. Environ.*, 46, 213–222, 1993.
- Behmann, J., Mahlein, A.-K., Paulus, S., Kuhlmann, H., Oerke, E.-C., and Plümer, L.: Calibration of hyperspectral close-range pushbroom cameras for plant phenotyping, *ISPRS J. Photogramm. Remote Sens.*, 106, 172–182, <https://doi.org/10.1016/j.isprsjprs.2015.05.010>, 2015.
- 595 Boegh, E., Soegaard, H., Broge, N., Hasager, C. B., Jensen, N. O., Schelde, K., and Thomsen, A.: Airborne multispectral data for quantifying leaf area index, nitrogen concentration, and photosynthetic efficiency in agriculture, *Remote Sens. Environ.*, 81, 179–193, [https://doi.org/10.1016/S0034-4257\(01\)00342-X](https://doi.org/10.1016/S0034-4257(01)00342-X), 2002.
- Bowker, D. E.: Spectral reflectances of natural targets for use in remote sensing studies, vol. 1139, NASA, 1985.
- 600 Buras, R., Dowling, T., and Emde, C.: New secondary-scattering correction in DISORT with increased efficiency for forward scattering, *J. Quant. Spectrosc. Radiat. Transfer*, 112, 2028–2034, <https://doi.org/10.1016/j.jqsrt.2011.03.019>, 2011.
- Burkart, A., Cogliati, S., Schickling, A., and Rascher, U.: A novel UAV-based ultra-light weight spectrometer for field spectroscopy, *IEEE Sens. J.*, 14, 62–67, <https://doi.org/10.1109/JSEN.2013.2279720>, 2014.
- Calif, R., Schmitt, F. G., Huang, Y., and Soubdhan, T.: Intermittency study of high frequency global solar radiation sequences under a tropical climate, *Sol. Energy*, 98, 349–365, <https://doi.org/10.1016/j.solener.2013.09.018>, 2013.
- 605 Carlson, T. N. and Ripley, D. A.: On the relation between NDVI, fractional vegetation cover, and leaf area index, *Remote Sens. Environ.*, 62, 241–252, [https://doi.org/10.1016/S0034-4257\(97\)00104-1](https://doi.org/10.1016/S0034-4257(97)00104-1), 1997.
- Chen, D., Huang, J., and Jackson, T. J.: Vegetation water content estimation for corn and soybeans using spectral indices derived from MODIS near- and short-wave infrared bands, *Remote Sens. Environ.*, 98, 225–236, <https://doi.org/10.1016/j.rse.2005.07.008>, 2005.

- 610 Coddington, O. M., Richard, E. C., Harber, D., Pilewskie, P., Woods, T. N., Chance, K., Liu, X., and Sun, K.: The TSIS-1 Hybrid Solar Reference Spectrum, *Geophys. Res. Lett.*, 48, e2020GL091709, <https://doi.org/10.1029/2020GL091709>, 2021.
- Collins, W.: Remote sensing of crop type and maturity, *Photogramm. Eng. Remote Sens.*, 44, 43–55, 1978.
- Drusch, M., Del Bello, U., Carlier, S., Colin, O., Fernandez, V., Gascon, F., Hoersch, B., Isola, C., Laberinti, P., Martimort, P., Meygret, A., Spoto, F., Sy, O., Marchese, F., and Bargellini, P.: Sentinel-2: ESA's optical high-resolution mission for GMES operational services, 615 *Remote Sens. Environ.*, 120, 25–36, <https://doi.org/10.1016/j.rse.2011.11.026>, the Sentinel Missions - New Opportunities for Science, 2012.
- Duan, T., Chapman, S. C., Guo, Y., and Zheng, B.: Dynamic monitoring of NDVI in wheat agronomy and breeding trials using an unmanned aerial vehicle, *Field Crops Res.*, 210, 71–80, <https://doi.org/10.1016/j.fcr.2017.05.025>, 2017.
- Dubovik, O., Holben, B. N., Lapyonok, T., Sinyuk, A., Mishchenko, M. I., Yang, P., and Slutsker, I.: Non-spherical aerosol retrieval method 620 employing light scattering by spheroids, *Geophys. Res. Lett.*, 29, 54–1–54–4, <https://doi.org/10.1029/2001GL014506>, 2002.
- Emde, C., Buras-Schnell, R., Kylling, A., Mayer, B., Gasteiger, J., Hamann, U., Kylling, J., Richter, B., Pause, C., Dowling, T., and Bugliaro, L.: The libRadtran software package for radiative transfer calculations (version 2.0.1), *Geosci. Model Dev.*, 9, 1647–1672, <https://doi.org/10.5194/gmd-9-1647-2016>, 2016.
- Fawcett, D., Panigada, C., Tagliabue, G., Boschetti, M., Celesti, M., Evdokimov, A., Biriukova, K., Colombo, R., Miglietta, F., Rascher, U., 625 and Anderson, K.: Multi-scale evaluation of drone-based multispectral surface reflectance and vegetation indices in operational conditions, *Remote Sens.*, 12, <https://doi.org/10.3390/rs12030514>, 2020.
- Freedman, J. M., Fitzjarrald, D. R., Moore, K. E., and Sakai, R. K.: Boundary layer clouds and vegetation–atmosphere feedbacks, *J. Climate*, 14, 180 – 197, [https://doi.org/10.1175/1520-0442\(2001\)013<0180:BLCAVA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)013<0180:BLCAVA>2.0.CO;2), 2001.
- Freudenthaler, V., Homburg, F., and Jäger, H.: Contrail observations by ground-based scanning lidar: Cross-sectional growth, *Geophys. Res. 630 Lett.*, 22, 3501–3504, <https://doi.org/10.1029/95GL03549>, 1995.
- Frisch, S., Shupe, M., Djalalova, I., Feingold, G., and Poellot, M.: The retrieval of stratus cloud droplet effective radius with cloud radars, *J. Atmos. Oceanic Technol.*, 19, 835 – 842, [https://doi.org/10.1175/1520-0426\(2002\)019<0835:TROSCD>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0835:TROSCD>2.0.CO;2), 2002.
- Gao, B.-C.: NDWI—A normalized difference water index for remote sensing of vegetation liquid water from space, *Remote Sens. Environ.*, 58, 257–266, [https://doi.org/10.1016/S0034-4257\(96\)00067-3](https://doi.org/10.1016/S0034-4257(96)00067-3), 1996.
- 635 Gardner, A. S. and Sharp, M. J.: A review of snow and ice albedo and the development of a new physically based broadband albedo parameterization, *J. Geophys. Res. Earth Surf.*, 115, F01009, <https://doi.org/10.1029/2009JF001444>, 2010.
- Gasteiger, J., Emde, C., Mayer, B., Buras, R., Buehler, S., and Lemke, O.: Representative wavelengths absorption parameterization applied to satellite channels and spectral bands, *J. Quant. Spectrosc. Radiat. Transfer*, 148, 99–115, <https://doi.org/10.1016/j.jqsrt.2014.06.024>, 2014.
- 640 Gitelson, A. A., Viña, A., Verma, S. B., Rundquist, D. C., Arkebauer, T. J., Keydan, G., Leavitt, B., Ciganda, V., Burba, G. G., and Suyker, A. E.: Relationship between gross primary production and chlorophyll content in crops: Implications for the synoptic monitoring of vegetation productivity, *J. Geophys. Res. Atmos.*, 111, <https://doi.org/10.1029/2005JD006017>, 2006.
- Goel, N. S.: Models of vegetation canopy reflectance and their use in estimation of biophysical parameters from reflectance data, *Remote Sens. Rev.*, 4, 1–212, <https://doi.org/10.1080/02757258809532105>, 1988.
- 645 Grenfell, T. C. and Perovich, D. K.: Incident spectral irradiance in the Arctic Basin during the summer and fall, *J. Geophys. Res. Atmos.*, 113, <https://doi.org/10.1029/2007JD009418>, 2008.

- Hakala, T., Honkavaara, E., Saari, H., Mäkynen, J., Kaivosoja, J., Pesonen, L., and Pölönen, I.: Spectral imaging From UAVs under varying illumination conditions, *ISPRS, XL-1/W2*, 189–194, <https://doi.org/10.5194/isprsarchives-XL-1-W2-189-2013>, 2013.
- Hakala, T., Markelin, L., Honkavaara, E., Scott, B., Theocharous, T., Nevalainen, O., Näsi, R., Suomalainen, J., Viljanen, N., Greenwell, C.,
650 and Fox, N.: Direct reflectance measurements from drones: Sensor absolute radiometric calibration and system tests for forest reflectance characterization, *Sensors*, 18, <https://doi.org/10.3390/s18051417>, 2018.
- Hardisky, M. A., Klemas, V., and Smart, R. M.: The influence of soil salinity, growth form, and leaf moisture on the spectral radiance of *Spartina alterniflora* canopies, *Photogramm. Eng. Remote Sens.*, 49, 77–83, 1983.
- Holz, R. E., Platnick, S., Meyer, K., Vaughan, M., Heidinger, A., Yang, P., Wind, G., Dutcher, S., Ackerman, S., Amarasinghe, N., Nagle, F.,
655 and Wang, C.: Resolving ice cloud optical thickness biases between CALIOP and MODIS using infrared retrievals, *Atmos. Chem. Phys.*, 16, 5075–5090, <https://doi.org/10.5194/acp-16-5075-2016>, 2016.
- Hong, S., Lakshmi, V., and Small, E. E.: Relationship between vegetation biophysical properties and surface temperature using multisensor satellite data, *J. Climate*, 20, 5593 – 5606, <https://doi.org/10.1175/2007JCLI1294.1>, 2007.
- Honkavaara, E., Saari, H., Kaivosoja, J., Pölönen, I., Hakala, T., Litkey, P., Mäkynen, J., and Pesonen, L.: Processing and assessment of
660 spectrometric, stereoscopic imagery collected using a lightweight UAV spectral camera for precision agriculture, *Remote Sens.*, 5, 5006–5039, <https://doi.org/10.3390/rs5105006>, 2013.
- Horler, D. N. H., Dockray, M., and Berber, J.: The red edge of plant leaf reflectance, *Int. J. Remote Sens.*, 4, 273–288, <https://doi.org/10.1080/01431168308948546>, 1983.
- Huete, A., Justice, C., and Liu, H.: Development of vegetation and soil indices for MODIS-EOS, *Remote Sens. Environ.*, 49, 224–234,
665 [https://doi.org/10.1016/0034-4257\(94\)90018-3](https://doi.org/10.1016/0034-4257(94)90018-3), 1994.
- Huete, A. R., Jackson, R. D., and Post, D. F.: Spectral response of a plant canopy with different soil backgrounds, *Remote Sens. Environ.*, 17, 37–53, 1985.
- Huete, A. R., Didan, K., Shimabukuro, Y. E., Ratana, P., Saleska, S. R., Hutyra, L. R., Yang, W., Nemani, R. R., and Myneni, R.: Amazon rainforests green-up with sunlight in dry season, *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2005GL025583>, 2006.
- Iwabuchi, H., Yang, P., Liou, K. N., and Minnis, P.: Physical and optical properties of persistent contrails: Climatology and interpretation, *J. Geophys. Res. Atmos.*, 117, D06 215, <https://doi.org/10.1029/2011JD017020>, 2012.
- Järvinen, E., Jourdan, O., Neubauer, D., Yao, B., Liu, C., Andreae, M. O., Lohmann, U., Wendisch, M., McFarquhar, G. M., Leisner, T., and Schnaiter, M.: Additional global climate cooling by clouds due to ice crystal complexity, *Atmos. Chem. Phys.*, 18, 15 767–15 781, <https://doi.org/10.5194/acp-18-15767-2018>, 2018.
- Jarvis, P. G., Miranda, H. S., and Muetzelfeldt, R. I.: Modelling canopy exchanges of water vapor and carbon dioxide in coniferous forest plantations, pp. 521–542, Springer Netherlands, Dordrecht, ISBN 978-94-009-5305-5, https://doi.org/10.1007/978-94-009-5305-5_31, 1985.
- Jiang, R., Wang, P., Xu, Y., Zhou, Z., Luo, X., Lan, Y., Zhao, G., Sanchez-Azofeifa, A., and Laakso, K.: Assessing the operation parameters of a low-altitude UAV for the collection of NDVI values over a paddy rice field, *Remote Sens.*, 12, <https://doi.org/10.3390/rs12111850>,
680 2020.
- Jiang, Z., Huete, A. R., Didan, K., and Miura, T.: Development of a two-band enhanced vegetation index without a blue band, *Remote Sens. Environ.*, 112, 3833–3845, <https://doi.org/10.1016/j.rse.2008.06.006>, 2008.
- Jones, H. G. and Vaughan, R. A.: Remote sensing of vegetation: principles, techniques, and applications, Oxford University Press, USA, ISBN 978-0199207794, 2010.

- 685 Jones, H. M., Haywood, J., Marengo, F., O'Sullivan, D., Meyer, J., Thorpe, R., Gallagher, M. W., Krämer, M., Bower, K. N., Rädcl, G., Rap, A., Woolley, A., Forster, P., and Coe, H.: A methodology for in-situ and remote sensing of microphysical and radiative properties of contrails as they evolve into cirrus, *Atmos. Chem. Phys.*, 12, 8157–8175, <https://doi.org/10.5194/acp-12-8157-2012>, 2012.
- Jurado, M., Caridad, J. M., and Ruiz, V.: Statistical distribution of the clearness index with radiation data integrated over five minute intervals, *Sol. Energy*, 55, 469–473, [https://doi.org/10.1016/0038-092X\(95\)00067-2](https://doi.org/10.1016/0038-092X(95)00067-2), 1995.
- 690 Jurgens, C.: The modified normalized difference vegetation index (mNDVI) a new index to determine frost damages in agriculture based on Landsat TM data, *Int. J. Remote Sens.*, 18, 3583–3594, <https://doi.org/10.1080/014311697216810>, 1997.
- Kattenborn, T., Richter, R., Guimarães Steinicke, C., Feilhauer, H., and Wirth, C.: AngleCam: Predicting the temporal variation of leaf angle distributions from image series with deep learning, *Methods Ecol. Evol.*, 13, 2531–2545, <https://doi.org/10.1111/2041-210X.13968>, 2022.
- Kattenborn, T., Wieneke, S., Montero, D., Mahecha, M. D., Richter, R., Guimarães-Steinicke, C., Wirth, C., Ferlian, O., Feilhauer, H.,
- 695 Sachsenmaier, L., et al.: Temporal dynamics in vertical leaf angles can confound vegetation indices widely used in Earth observations, *Commun. Earth Environ.*, 5, 550, 2024.
- Kaufman, Y. J. and Tanre, D.: Atmospherically resistant vegetation index (ARVI) for EOS-MODIS, *IEEE Trans. Geosci. Remote Sens.*, 30, 261–270, 1992.
- King, M. D., Radke, L. F., and Hobbs, P. V.: Optical properties of marine stratocumulus clouds modified by ships, *J. Geophys. Res. Atmos.*,
- 700 98, 2729–2739, <https://doi.org/10.1029/92JD02082>, 1993.
- Knyazikhin, Y., Schull, M. A., Stenberg, P., Möttus, M., Rautiainen, M., Yang, Y., Marshak, A., Carmona, P. L., Kaufmann, R. K., Lewis, P., Disney, M. I., Vanderbilt, V., Davis, A. B., Baret, F., Jacquemoud, S., Lyapustin, A., and Myneni, R. B.: Hyperspectral remote sensing of foliar nitrogen content, *PNAS*, 110, E185–E192, <https://doi.org/10.1073/pnas.1210196109>, 2013.
- Köppl, C. J., Malureanu, R., Dam-Hansen, C., Wang, S., Jin, H., Barchiesi, S., Serrano Sandí, J. M., Muñoz-Carpena, R., Johnson, M., Durán-Quesada, A. M., Bauer-Gottwein, P., McKnight, U. S., and Garcia, M.: Hyperspectral reflectance measurements from UAS under intermittent clouds: Correcting irradiance measurements for sensor tilt, *Remote Sens. Environ.*, 267, 112719, <https://doi.org/10.1016/j.rse.2021.112719>, 2021.
- 705 Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Meyer, J., Zöger, M., Smith, J., Herman, R. L., Buchholz, B., Ebert, V., Baumgardner, D., Borrmann, S., Klingebiel, M., and Avallone, L.: A microphysics guide to cirrus clouds – Part 1: Cirrus types, *Atmos. Chem. Phys.*, 16, 3463–3483, <https://doi.org/10.5194/acp-16-3463-2016>, 2016.
- 710 Kriegler, F. J., Malila, W. A., Nalepka, R. F., and Richardson, W.: Preprocessing transformations and their effects on multispectral recognition, in: *Remote Sens. Environ.*, VI, pp. 97–131, 1969.
- Liu, C., Yang, P., Minnis, P., Loeb, N., Kato, S., Heymsfield, A., and Schmitt, C.: A two-habit model for the microphysical and optical properties of ice clouds, *Atmos. Chem. Phys.*, 14, 13719–13737, <https://doi.org/10.5194/acp-14-13719-2014>, 2014.
- 715 Liu, H. Q. and Huete, A.: A feedback based modification of the NDVI to minimize canopy background and atmospheric noise, *IEEE Trans. Geosci. Remote Sens.*, 33, 457–465, 1995.
- Lohmann, G. M.: Irradiance variability quantification and small-scale averaging in space and time: A short review, *Atmosphere*, 9, <https://doi.org/10.3390/atmos9070264>, 2018.
- Loveland, T. R. and Belward, A. S.: The IGBP-DIS global 1km land cover data set, DISCover: First results, *Int. J. Remote Sens.*, 18,
- 720 3289–3295, <https://doi.org/10.1080/014311697217099>, 1997.
- Lu, M.-L., Feingold, G., Jonsson, H. H., Chuang, P. Y., Gates, H., Flagan, R. C., and Seinfeld, J. H.: Aerosol-cloud relationships in continental shallow cumulus, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009354>, 2008.

- Lucht, W., Schaaf, C., and Strahler, A.: An algorithm for the retrieval of albedo from space using semiempirical BRDF models, *IEEE Trans. Geosci. Remote Sens.*, 38, 977–998, <https://doi.org/10.1109/36.841980>, 2000.
- 725 Luecke, A. E., Afchine, A., Costa, A., Grob, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Krämer, M.: The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, *Atmos. Chem. Phys.*, 16, 5793–5809, <https://doi.org/10.5194/acp-16-5793-2016>, 2016.
- Macke, A., Seifert, P., Baars, H., Barthlott, C., Beekmans, C., Behrendt, A., Bohn, B., Brueck, M., Bühl, J., Crewell, S., Damian, T., Deneke, H., Düsing, S., Foth, A., Di Girolamo, P., Hammann, E., Heinze, R., Hirsikko, A., Kalisch, J., Kalthoff, N., Kinne, S., Kohler, M., Löhnert, U., Madhavan, B. L., Maurer, V., Muppa, S. K., Schween, J., Serikov, I., Siebert, H., Simmer, C., Späth, F., Steinke, S., Träumner, K., Trömel, S., Wehner, B., Wieser, A., Wulfmeyer, V., and Xie, X.: The HD(CP)² Observational Prototype Experiment (HOPE) – an overview, *Atmos. Chem. Phys.*, 17, 4887–4914, <https://doi.org/10.5194/acp-17-4887-2017>, 2017.
- 730 Madhavan, B. L., Kalisch, J., and Macke, A.: Shortwave surface radiation network for observing small-scale cloud inhomogeneity fields, *Atmos. Meas. Tech.*, 9, 1153–1166, <https://doi.org/10.5194/amt-9-1153-2016>, 2016.
- 735 Marshak, A., Wen, G., C. Jr., J. A., Remer, L. A., Loeb, N. G., and Cahalan, R. F.: A simple model for the cloud adjacency effect and the apparent bluing of aerosols near clouds, *J. Geophys. Res. Atmos.*, 113, <https://doi.org/10.1029/2007JD009196>, 2008.
- Martonchik, J. V., Bruegge, C. J., and Strahler, A. H.: A review of reflectance nomenclature used in remote sensing, *Remote Sensing Reviews*, 19, 9–20, <https://doi.org/10.1080/02757250009532407>, 2000.
- Matese, A., Toscano, P., Di Gennaro, S. F., Genesio, L., Vaccari, F. P., Primicerio, J., Belli, C., Zaldei, A., Bianconi, R., and Gioli, B.: Intercomparison of UAV, Aircraft and Satellite Remote Sensing Platforms for Precision Viticulture, *Remote Sens.*, 7, 2971–2990, <https://doi.org/10.3390/rs70302971>, 2015.
- 740 Matsushita, B., Yang, W., Chen, J., Onda, Y., and Qiu, G.: Sensitivity of the Enhanced Vegetation Index (EVI) and Normalized Difference Vegetation Index (NDVI) to topographic effects: A case study in high-density cypress forest, *Sensors*, 7, 2636–2651, <https://doi.org/10.3390/s7112636>, 2007.
- 745 Mie, G.: Beiträge zur Optik trüber Medien, speziell kolloidaler Metallösungen, *Ann. Phys.*, 330, 377–445, <https://doi.org/10.1002/andp.19083300302>, 1908.
- Miura, T., Huete, A. R., Van Leeuwen, W. J. D., and Didan, K.: Vegetation detection through smoke-filled AVIRIS images: An assessment using MODIS band passes, *J. Geophys. Res. Atmos.*, 103, 32 001–32 011, 1998.
- Miyoshi, G. T., Imai, N. N., Tommaselli, A. M. G., Honkavaara, E., Näsi, R., and Moriya, E. A. S.: Radiometric block adjustment of hyperspectral image blocks in the Brazilian environment, *Int. J. Remote Sens.*, 39, 4910–4930, <https://doi.org/10.1080/01431161.2018.1425570>, 2018.
- 750 Montero, D., Aybar, C., Mahecha, M. D., Martinuzzi, F., Söchting, M., and Wieneke, S.: A standardized catalogue of spectral indices to advance the use of remote sensing in Earth system research, *Sci. Data*, 10, 197, 2023.
- Mura, M., Bottalico, F., Giannetti, F., Bertani, R., Giannini, R., Mancini, M., Orlandini, S., Travaglini, D., and Chirici, G.: Exploiting the capabilities of the Sentinel-2 multi spectral instrument for predicting growing stock volume in forest ecosystems, *Int. J. Appl. Earth Obs. Geoinf.*, 66, 126–134, <https://doi.org/10.1016/j.jag.2017.11.013>, 2018.
- 755 Myneni, R. B., Maggion, S., Iaquinta, J., Privette, J. L., Gobron, N., Pinty, B., Kimes, D. S., Verstraete, M. M., and Williams, D. L.: Optical remote sensing of vegetation: Modeling, caveats, and algorithms, *Remote Sens. Environ.*, 51, 169–188, [https://doi.org/10.1016/0034-4257\(94\)00073-V](https://doi.org/10.1016/0034-4257(94)00073-V), remote Sensing of Land Surface for Studies of Global Change, 1995.

- 760 Myneni, R. B., Keeling, C. D., Tucker, C. J., Asrar, G., and Nemani, R. R.: Increased plant growth in the northern high latitudes from 1981 to 1991, *Nature*, 386, 698–702, 1997.
- Nicodemus, F. E.: Geometrical considerations and nomenclature for reflectance, vol. 160, US Department of Commerce, National Bureau of Standards Washington, DC, USA, <https://graphics.stanford.edu/courses/cs448-05-winter/papers/nicodemus-brdf-nist.pdf>, 1977.
- Noël, V. and Haeffelin, M.: Midlatitude cirrus clouds and multiple tropopauses from a 2002–2006 climatology over the SIRTa observatory, *J. Geophys. Res. Atmos.*, 112, <https://doi.org/10.1029/2006JD007753>, 2007.
- 765 Perez, P., Kivalov, S., Schlemmer, J., Hemker, K., and Hoff, T.: Parameterization of site-specific short-term irradiance variability, *Sol. Energy*, 85, 1343–1353, <https://doi.org/10.1016/j.solener.2011.03.016>, 2011.
- Pilewskie, P. and Twomey, S.: Discrimination of ice from water in clouds by optical remote sensing, *Atmos. Res.*, 21, 113–122, [https://doi.org/10.1016/0169-8095\(87\)90002-0](https://doi.org/10.1016/0169-8095(87)90002-0), 1987.
- 770 Pinty, B., Lattanzio, A., Martonchik, J. V., Verstraete, M. M., Gobron, N., Taberner, M., Widlowski, J.-L., Dickinson, R. E., and Govaerts, Y.: Coupling diffuse sky radiation and surface albedo, *J. Atmos. Sci.*, 62, 2580 – 2591, <https://doi.org/10.1175/JAS3479.1>, 2005.
- Rahman, A. F., Sims, D. A., Cordova, V. D., and El-Masri, B. Z.: Potential of MODIS EVI and surface temperature for directly estimating per-pixel ecosystem C fluxes, *Geophys. Res. Lett.*, 32, <https://doi.org/10.1029/2005GL024127>, 2005.
- Richardson, A. D., Aubrecht, D. M., Basler, D., Hufkens, K., Muir, C. D., and Hanssen, L.: Developmental changes in the reflectance spectra of temperate deciduous tree leaves and implications for thermal emissivity and leaf temperature, *New Phytol.*, 229, 791–804, <https://doi.org/10.1111/nph.16909>, 2021.
- 775 Saleska, S. R., Didan, K., Huete, A. R., and Da Rocha, H. R.: Amazon forests green-up during 2005 drought, *Science*, 318, 612–612, 2007.
- Salomonson, V., Barnes, W., Maymon, P., Montgomery, H., and Ostrow, H.: MODIS: advanced facility instrument for studies of the Earth as a system, *IEEE Trans. Geosci. Remote Sens.*, 27, 145–153, <https://doi.org/10.1109/36.20292>, 1989.
- 780 Sassen, K. and Campbell, J. R.: A midlatitude cirrus cloud climatology from the facility for atmospheric remote sensing. Part I: macrophysical and synoptic properties, *J. Atmos. Sci.*, 58, 481–496, [https://doi.org/10.1175/1520-0469\(2001\)058<0481:AMCCCF>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<0481:AMCCCF>2.0.CO;2), 2001.
- Schaepman-Strub, G., Schaepman, M. E., Painter, T. H., Dangel, S., and Martonchik, J. V.: Reflectance quantities in optical remote sensing—definitions and case studies, *Remote Sen. Environ.*, 103, 27–42, <https://doi.org/10.1016/j.rse.2006.03.002>, 2006.
- Shettle, E.: Models of aerosols, clouds and precipitation for atmospheric propagation studies, in: *Atmospheric propagation in the uv, visible, ir and mm-region and related system aspects*, no. 454 in AGARD Conference Proceedings, 1989.
- 785 Singh, K. K. and Frazier, A. E.: A meta-analysis and review of unmanned aircraft system (UAS) imagery for terrestrial applications, *Int. J. Remote Sens.*, 39, 5078–5098, <https://doi.org/10.1080/01431161.2017.1420941>, 2018.
- Spoto, F., Sy, O., Laberinti, P., Martimort, P., Fernandez, V., Colin, O., Hoersch, B., and Meygret, A.: Overview Of Sentinel-2, in: *2012 IEEE International Geoscience and Remote Sensing Symposium*, pp. 1707–1710, <https://doi.org/10.1109/IGARSS.2012.6351195>, 2012.
- 790 Stamnes, K., Tsay, S.-C., Wiscombe, W., and Jayaweera, K.: Numerically stable algorithm for discrete-ordinate-method radiative transfer in multiple scattering and emitting layered media, *Appl. Opt.*, 27, 2502–2509, <https://doi.org/10.1364/AO.27.002502>, 1988.
- Stephens, G. L.: *Remote sensing of the lower atmosphere, an introduction*, Oxford University Press, New York, ISBN 978-0195081889, 1994.
- Stuckens, J., Somers, B., Delalieux, S., Verstraeten, W. W., and Coppin, P.: The impact of common assumptions on canopy radiative transfer simulations: A case study in Citrus orchards, *J. Quant. Spectrosc. Radiat. Transfer*, 110, 1–21, <https://doi.org/10.1016/j.jqsrt.2008.09.001>, 2009.
- 795

- Twomey, S. and Cocks, T.: Remote sensing of cloud parameters from spectral reflectance in the near-infrared, *Beitr. Phys. Atmos.*, 62, 172–179, 1989.
- Ünsalan, C. and Boyer, K. L.: Linearized Vegetation Indices, *Multispectral satellite image understanding: From land classification to building and road detection*, pp. 19–39, 2011.
- van de Hulst, H. C.: *Light scattering by small particles*, Courier Corporation, first edn., 1981.
- van Haaren, R., Morjaria, M., and Fthenakis, V.: Empirical assessment of short-term variability from utility-scale solar PV plants, *Prog. Photovoltaics Res. Appl.*, 22, 548–559, <https://doi.org/10.1002/pip.2302>, 2014.
- Verhoef, W., van der Tol, C., and Middleton, E. M.: Hyperspectral radiative transfer modeling to explore the combined retrieval of biophysical parameters and canopy fluorescence from FLEX – Sentinel-3 tandem mission multi-sensor data, *Remote Sens. Environ.*, 204, 942–963, <https://doi.org/10.1016/j.rse.2017.08.006>, 2018.
- Verrelst, J., Rivera, J. P., van der Tol, C., Magnani, F., Mohammed, G., and Moreno, J.: Global sensitivity analysis of the SCOPE model: What drives simulated canopy-leaving sun-induced fluorescence?, *Remote Sens. Environ.*, 166, 8–21, <https://doi.org/10.1016/j.rse.2015.06.002>, 2015.
- Vicari, M. B., Pisek, J., and Disney, M.: New estimates of leaf angle distribution from terrestrial LiDAR: Comparison with measured and modelled estimates from nine broadleaf tree species, *Agric. For. Meteorol.*, 264, 322–333, <https://doi.org/10.1016/j.agrformet.2018.10.021>, 2019.
- Watson, D. J.: Comparative physiological studies on the growth of field crops: I. Variation in net assimilation rate and leaf area between species and varieties, and within and between years, *Ann. Bot.*, 11, 41–76, 1947.
- Werner, F., Siebert, H., Pilewskie, P., Schmeissner, T., Shaw, R. A., and Wendisch, M.: New airborne retrieval approach for trade wind cumulus properties under overlying cirrus, *J. Geophys. Res. Atmos.*, 118, 3634–3649, <https://doi.org/https://doi.org/10.1002/jgrd.50334>, 2013.
- Wiscombe, W. J. and Warren, S. G.: A model for the spectral albedo of snow. I: pure snow, *J. Atmos. Sci.*, 37, 2712 – 2733, [https://doi.org/10.1175/1520-0469\(1980\)037<2712:AMFTSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2712:AMFTSA>2.0.CO;2), 1980.
- Wolf, K., Jäkel, E., Ehrlich, A., Schäfer, M., Feilhauer, H., Huth, A., Weigelt, A., and Wendisch, M.: Impact of stratiform liquid water clouds on vegetation albedo quantified by coupling an atmosphere and a vegetation radiative transfer model, *Biogeosciences*, 22, 2909–2933, <https://doi.org/10.5194/bg-22-2909-2025>, 2025a.
- Wolf, K., Jäkel, E., Ehrlich, A., Schäfer, M., Feilhauer, H., Huth, A., and Wendisch, M.: Simulated spectral irradiances, radiances, and vegetation albedo obtained from coupling libRadtran and SCOPE2.0, <https://doi.org/10.5281/zenodo.15275610>, 2025b.
- Wu, C., Niu, Z., Tang, Q., Huang, W., Rivard, B., and Feng, J.: Remote estimation of gross primary production in wheat using chlorophyll-related vegetation indices, *Agric. For. Meteorol.*, 149, 1015–1021, <https://doi.org/10.1016/j.agrformet.2008.12.007>, 2009.
- Wulder, M. A., Loveland, T. R., Roy, D. P., Crawford, C. J., Masek, J. G., Woodcock, C. E., Allen, R. G., Anderson, M. C., Belward, A. S., Cohen, W. B., et al.: Current status of Landsat program, science, and applications, *Remote Sens Environ.*, 225, 127–147, 2019.
- Xu, H.: Modification of normalised difference water index (NDWI) to enhance open water features in remotely sensed imagery, *Int. J. Remote Sens.*, 27, 3025–3033, <https://doi.org/10.1080/01431160600589179>, 2006.
- Xue, J. and Su, B.: Significant remote sensing vegetation indices: A review of developments and applications, *J. Sens.*, 2017, 2017.
- Yang, P., Bi, L., Baum, B. A., Liou, K.-N., Kattawar, G. W., Mishchenko, M. I., and Cole, B.: Spectrally consistent scattering, absorption, and polarization properties of atmospheric ice crystals at wavelengths from 0.2 to 100 μm , *J. Atmos. Sci.*, 70, 330–347, <https://doi.org/10.1175/JAS-D-12-039.1>, 2013.

- 835 Yang, P., Verhoef, W., and van der Tol, C.: The mSCOPE model: A simple adaptation to the SCOPE model to describe reflectance, fluorescence and photosynthesis of vertically heterogeneous canopies, *Remote Sens. Environ.*, 201, 1–11, <https://doi.org/10.1016/j.rse.2017.08.029>, 2017.
- Yang, P., van der Tol, C., Yin, T., and Verhoef, W.: The SPART model: A soil-plant-atmosphere radiative transfer model for satellite measurements in the solar spectrum, *Remote Sens. Environ.*, 247, 111 870, <https://doi.org/10.1016/j.rse.2020.111870>, 2020.
- 840 Yang, P., Prikaziuk, E., Verhoef, W., and van der Tol, C.: SCOPE 2.0: a model to simulate vegetated land surface fluxes and satellite signals, *Geosci. Model Dev.*, 14, 4697–4712, <https://doi.org/10.5194/gmd-14-4697-2021>, 2021.
- Yang, X., Li, R., Jablonski, A., Stovall, A., Kim, J., Yi, K., Ma, Y., Beverly, D., Phillips, R., Novick, K., Xu, X., and Lerdau, M.: Leaf angle as a leaf and canopy trait: Rejuvenating its role in ecology with new technology, *Ecol. Lett.*, 26, 1005–1020, <https://doi.org/10.1111/ele.14215>, 2023.
- 845 Zeng, Y., Hao, D., Huete, A., Dechant, B., Berry, J., Chen, J. M., Joiner, J., Frankenberg, C., Bond-Lamberty, B., Ryu, Y., et al.: Optical vegetation indices for monitoring terrestrial ecosystems globally, *Nat. Rev. Earth Environ.*, 3, 477–493, 2022.
- Zhou, Y., Zhang, L., Xiao, J., Chen, S., Kato, T., and Zhou, G.: A comparison of satellite-derived vegetation indices for approximating gross primary productivity of grasslands, *Rangeland Ecol. Manage.*, 67, 9–18, <https://doi.org/10.2111/REM-D-13-00059.1>, 2014.