



Fresh tephra deposits from the Tajogaite Volcano boost thermophile proliferation and soil organic matter recovery

Sara Gutiérrez-Patricio¹, Alba Gómez-Arias¹, Pedro Nolasco-Jiménez¹, Jorge Mataix-Solera², Javier Martínez-Martínez³,
5 Bruno Martínez-Haya⁴, Juana Vegas³, Nicasio T. Jiménez-Morillo¹, Ana Z. Miller^{1,5}

¹ Instituto de Recursos Naturales y Agrobiología de Sevilla (IRNAS-CSIC), Avda. Reina Mercedes 10, 41012, Seville, Spain

² Grupo de Edafología y Tecnologías del Medio Ambiente GETECMA. Departamento de Agroquímica y Medio Ambiente, Universidad Miguel Hernández, 03202 Elche, Alicante, Spain.

10 ³ Instituto Geológico y Minero de España (IGME-CSIC), Ríos Rosas 23, 28003 Madrid, Spain.

⁴ BIO-MS group, Department of Physical, Chemical and Natural Systems, Universidad Pablo de Olavide, 41013, Seville, Spain.

⁵ Hercules Laboratory, University of Évora, Palacio do Vimioso, 7000-809, Évora, Portugal.

Correspondence to: Nicasio T. Jimenez-Morillo (ntjm@imas.csic.es) / Ana Z. Miller (anamiller@imas.csic.es)

15


Abstract. Tephra fallout deposition during volcanic eruptions overlays existing soils, profoundly altering their physical, chemical, and biological properties. This study investigates the impact of the newly deposited tephra blanket from the 2021 Tajogaite eruption (La Palma Island) on the molecular composition of soil organic matter and microbial diversity across different soil horizons. A combination of 16S and 18S rRNA gene sequencing, pyrolysis-gas chromatography/mass
20 spectrometry (Py-GC/MS), and elemental and isotope ratio mass spectrometry (EA/IRMS) was employed. Our results demonstrate that tephra deposits significantly modify the organic matter composition of the underlying soils, promoting microbial activity linked to the degradation and transformation of organic carbon and nitrogen compounds. The soil horizon directly beneath the tephra layer (horizon O) displayed a higher abundance of labile organic compounds and a reduced presence of recalcitrant compounds compared to the deeper horizons (A and Bw). This pattern is strongly associated with the



25 predominance of thermophilic bacteria, which contribute actively to the breakdown of complex organic materials such as lignin and hydrocarbons, and drive key biogeochemical processes including nitrogen and carbon cycling. The continuous geothermal influence of nearby fumaroles further supports the persistence and ecological success of thermophilic communities in these volcanic soils. These findings underscore the critical role of volcanic activity not only in reshaping soil structure but also in enhancing soil fertility and resilience through microbial-mediated processes. Understanding these dynamics is essential for

30 soil management and ecosystem recovery strategies in volcanic regions, providing new insights into the long-term effects of tephra deposition on soil health and the carbon cycle.

Graphical abstract





1. Introduction

The Tajogaite volcanic eruption, which began in September 2021 on La Palma Island (Canary Islands, Spain), was an extraordinary geological event that significantly transformed the local environment (Birnbbaum et al., 2023; Troll et al., 2024).

40 Over 85 days, the eruption released over 200 million cubic meters of lava and pyroclasts, burying over 1,200 hectares of land and displacing 7000 residents. The voluminous ashfall, with a complex tephra blanket reaching up to 2.5 m thick, profoundly impacted the ecosystem of La Palma and posed substantial challenges for local agriculture, infrastructure, and human health (Craig et al., 2016; Bonadonna et al., 2022, 2023; Birnbbaum et al., 2023; Sánchez-España et al., 2023; Biass et al., 2024). According to these authors, the grain size of the tephra blanket varied during the eruption from lapilli >6 cm to very fine ash 45 (<2 mm) that spread to La Gomera, El Hierro, Tenerife and Gran Canaria islands during the more explosive phases. Ashfall significantly impacted agricultural productivity (Sánchez-España et al., 2023; Taddeucci et al., 2023). The loss of fertile topsoil and the alteration of soil properties created long-term challenges for agricultural recovery and land use (Ustiatik et al., 2023). Volcanic tephra is a fine-grained unconsolidated deposit, highly reactive material that can considerably modify soil structure, chemistry, and biology (Nanzyo et al., 1993; Arnalds, 2013; Wygel et al., 2019). When deposited on existing soil layers, tephra 50 can create a new surface horizon that introduces novel mineral substrates and interacts with soil organic matter (SOM), triggering complex chemical and biological transformations (Hayakawa et al., 2020; Wijesinghe et al., 2020; Peng et al., 2021). These changes can influence soil fertility, stability, and overall health by modifying nutrient availability, pH, and soil aggregation (Peng et al., 2021; De la Rosa et al., 2023). Additionally, the ash can act as a catalyst for the formation of new organic-mineral complexes, altering SOM decomposition pathways and the long-term sequestration of carbon and other 55 essential nutrients (Nierop and Buurman, 2007; Hernández et al., 2012; Iwasaki et al., 2021). Understanding these processes is crucial, particularly in volcanic regions where soil resilience is key to ecosystem recovery.

Soil health is intrinsically linked to the composition and stability of SOM (Arias, 2005), which consists of a diverse array of plant-derived, microbial, and fungal compounds (González-Pérez et al., 2004). Fresh tephra deposition in regions affected by volcanic activity can significantly alter the soil capacity to retain nutrients and support plant and microbial communities, 60 ultimately influencing the trajectory of ecosystem recovery (Hernández et al., 2012; Yokobe et al., 2020; Muñoz et al., 2021).



Furthermore, the interactions between volcanic ashfall and SOM can lead to the formation of polycyclic aromatic hydrocarbons (PAHs) and other condensed compounds (Nanzyo et al., 1993; Tomašek et al., 2021).

The effects of volcanic ash extend beyond physical and chemical changes; they also shape the **microbial communities in soils** (Zeglin et al., 2016; Yokobe et al., 2020; Chen et al., 2021). Microorganisms are highly sensitive to changes in soil properties, and ash deposition can create new niches that favour specific microbial taxa adapted to the altered conditions (Chen et al., 2021). This alteration in microbial diversity can have profound implications for soil function, influencing processes such as SOM decomposition, biogeochemical cycle of carbon and nitrogen, nutrient cycling, and the production of secondary metabolites (Campos et al., 2020; Gutierrez-Patricio et al., 2024).

This work aimed to investigate the impact of the Tajogaite volcanic eruption on the molecular and microbial diversity of an existing soil profile that has been covered by tephra deposits. Using advanced molecular and microbiological techniques, including pyrolysis-gas chromatography/mass spectrometry (Py-GC/MS), stable isotope analysis, and 16S and 18S rRNA gene-sequencing, soil profiles under recent ash deposition were analysed to unravel how volcanic ash alters SOM quality and microbial community structure. The findings provide key insights into the mechanisms driving soil transformation following volcanic events and highlight the critical role of SOM and microbial dynamics in soil resilience and ecosystem recovery.

75

2 Material and Methods

2.1 Study site and sampling

The soil profile studied is located in the *Las Manchas* area of La Palma Island, within the region most affected by the 2021 Tajogaite volcanic eruption. La Palma Island, part of the Canary Islands archipelago and located approximately 400 km west of the African coast, has a total area of 708 km² and is one of the youngest in the archipelago, formed during the last 1.8 million years (Hoernle et al., 2009). **La Palma Island shows the most complete record of the evolution of an oceanic volcanic island, starting from the Pliocene submarine basal complex, the metamorphism originated by magmatic intrusion, the building of a Quaternary volcanic edifice, and giant landslides and intense erosion that defines the final morphology of Caldera de Taburiente.** Volcanism displaced to a N-S elongated polygenetic Cumbre Nueva volcanic ridge that has developed over the

80



85 last c. 125 ka (Carracedo et al., 2001; Sánchez-España et al., 2023) being the most active area of this island (Longpre & Felpeto, 2021).

The 2021 Tajogaite volcanic eruption, which began on September 19 and lasted for 85 days, emitted over 200 million cubic meters of lava and pyroclastic materials, creating extensive lava flows that covered approximately 1,200 hectares and extended into the Atlantic Ocean, forming two new lava deltas (Ferrer et al., 2023). The pyroclastic materials deposited during this
90 eruption ranged in size from fine ash to large lapilli and created a stark contrast with the pre-existing landscape (Bonadonna et al., 2022; Ferrer et al., 2023). The eruption also introduced a tephra blanket, up to 3 m thick in areas close to the volcanic cone, that overlies the native soils (Bonadonna et al., 2023).

The study site is located in the *Las Manchas* area (Los Llanos de Aridane municipality), situated on the western slopes of the Cumbre Nueva Ridge (Fig. 1A). Before the eruption, the soils in this area were classified as Typic Ustivitrands (Soil Survey
95 Staff, 2022). These soils typically exhibit a sandy loam texture, low bulk density, and a high degree of aggregation, particularly in the surface horizons, which are enriched with organic matter (see Table S1).

Following the eruption, the pre-existing soils in the study area were buried under a 20 cm thick layer of tephra and ash. Two years after the event, mosses colonized the tephra deposit surface, and biofilms developed at the soil-tephra interface (Fig. 1B). Sampling included the collection of moss growing on the fresh tephra, the tephra layer directly beneath the moss cover
100 (moss substrate, MS), a dark green biofilm on horizon O, and three soil horizons O (0–2 cm), A (2–10 cm) and Bw (10–20 cm) (Fig. 1B). This volcanic ash-derived soil profile was located near steam-emitting fumaroles, in a sector with a high temperature anomaly in the soil. These fumaroles vent water vapour as the main component and other chemical species, such as SO₂, H₂S, HCl, and CO gases up to 20.7, 17.7, >30.0 and 19.5 ppm, respectively, at temperatures higher than the ambient (Campeny et al., 2023; Martínez-Martínez et al., 2023).


105 This site provides a unique opportunity to study the interactions between fresh tephra deposits and the underlying soil, particularly their effects on soil structure, organic matter composition, and microbial communities.

All samples were collected using sterile tools, such as spatulas or individually wrapped disposable scoops, to avoid contamination, and all samples were immediately placed into sterile Whirl-Pak bags. The samples were kept on ice during



transportation to the laboratory for subsequent analysis. Subsamples of each material were used for DNA-based analysis,
110 carbon and nitrogen elemental and isotope analysis, and molecular composition.

The environmental conditions of the sampling area were monitored by installing a temperature and relative humidity sensor (HOBO MX2301A, ONSET®, Fig. S1A and B) near the fumarole vent to assess the temperature of the water vapour emissions from the fumarole. This sensor took measurements every hour for one year.



115

Figure 1: A) Location of the soil profile under tephra blanket in *Las Manchas* (La Palma, Spain), and B) sampling area of the volcanic ash-derived soil profile comprising moss, moss substrate (MS), biofilm located in the horizon O, and three horizons (O, A and Bw).

2.2 Carbon and nitrogen elemental and isotope analysis

120 The compositional analysis of organic carbon and nitrogen in both pyroclastic materials and soil samples was carried out using a Flash 2000 HT Elemental Micro-analyzer connected to a Thermal Conductivity Detector (EA/TCD; Thermo Scientific in Bremen, Germany). Samples, ranging in weight from 5 to 10 milligrams, were processed in duplicate ($n = 2$) following a **decarbonization** treatment as outlined by Jiménez-Morillo et al., (2016, 2020a). The calibration of carbon and nitrogen content employed certified standards such as acetanilide, nicotinamide, and aspartic acid, provided by Thermo Scientific, Bremen,



125 Germany. The carbon and nitrogen isotope ratios ($^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$) were determined using an Elemental Analyzer
connected to a Thermo Fisher Delta V Advantage Isotope Ratio Mass Spectrometer (EA/IRMS), following the method
described by (Jiménez-Morillo et al., 2020b). These procedures were conducted at the Stable Isotope Laboratory - MOSS at
IRNAS-CSIC (Spain). Isotope values are expressed in delta (δ) notation in parts per thousand (‰) relative to certified standards
set by the International Atomic Energy Agency (IAEA), with a standard deviation for bulk $\delta^{13}\text{C}$ of ± 0.05 ‰ and $\delta^{15}\text{N}$ of ± 0.1
130 ‰. Duplicate analyses were performed for samples and standards ($n = 2$).

2.3 Analytical pyrolysis (Py-GC/MS)

The analytical pyrolysis (Py-GC/MS) technique was composed of i) a double-shot pyrolyzer, model 3030D (Frontier
Laboratories, Japan), set at 500 °C; ii) an Agilent 6890N gas chromatograph, and iii) an Agilent 5973 MSD (mass selective
135 detector) mass spectrometer, operating at an electron impact energy of 70 eV. The chromatographic separation involved an
HP-5MS UI capillary column (30 m length, 0.25 mm internal diameter, 0.25 μm film thickness) with a helium flow rate of 1
mL min^{-1} . The sample injection was conducted in splitless mode at 250 °C, and the temperature program was adopted from
protocols described by Jiménez-Morillo et al., (2020a). Ion fragments ranging from 50 to 550 m/z were recorded, and
compound identification was conducted by comparison with virtual spectral libraries, such as the NIST20 and Wiley7. The
140 molecular compounds released by Py-GC/MS were categorized into the eight biogenic families proposed by Jiménez-Morillo
et al (2016a): unspecific aromatic compounds (ARO), polysaccharides (POL), peptides (PEP), lignin (LIG), lipids (LIP),
nitrogen-containing compounds (N-comp), polycyclic aromatic hydrocarbons (PAHs), and steranes (STR). Chromatographic
peak area $>0.2\%$ of the total chromatographic area was not taken into account. For a detailed study of homologous series of n -
alkanes and n -alken-2-ones, single ion monitoring of specific fragments - specifically ions with m/z 57 and m/z 59, respectively
145 - was adopted.

2.4 DNA extraction and sequencing

Genomic DNA was extracted from 250 mg of each of the five samples using the DNeasy PowerSoil Pro Kit (Qiagen, Germany)
following the manufacturer's instructions. The DNA was quantified using a Qubit 4.0 fluorometer (Invitrogen).



150 The extracted DNA was analyzed through next-generation sequencing (NGS), targeting the V3-V4 hypervariable region of the 16S rRNA gene for prokaryotes. Amplicons were sequenced using the Illumina MiSeq platform to generate 250bp paired-end raw reads, performed by Novogene Sequencing Services (China).

Raw amplicon sequence data were processed using QIIME 2 version 2024.5 (Bolyen et al., 2019). The table of Amplicon Sequence Variants (ASVs) was generated using DADA2, following its instructions (Callahan et al., 2016). Taxonomic classification was assigned using the SILVA reference database, version 138 (Quast et al., 2013). Alpha diversity analysis was calculated in QIIME2 version 2024.5 (Bolyen et al., 2019) to quantify the microbial diversity within each sample.

The raw reads were deposited in the NCBI Sequence Read Archive (SRA) database under accession number PRJNA1192049.



3 Results and discussion

160 3.1 Elemental and isotope composition of carbon and nitrogen

Table 1: Isotope and elemental composition of nitrogen (N) and carbon (C) of moss, moss substrate (MS), biofilm and three soil horizons (O, A and Bw). Significant ($p < 0.05$) differences between samples are indicated by different letters ("one-way ANOVA"; means compared to the Tukey test, $p = 0.05$).

Sample	$\delta^{15}\text{N}$ (‰, V-Air)	$\delta^{13}\text{C}$ (‰, VPDB)	N (%)	C (%)
Moss	-5.1±0.4d	-28.5±0.1b	0.62±0.12b	12.53±1.69c
MS*	5.0±2.9bc	-24.3±1.6ab	3.17±0.80ab	28.71±4.53bc
Biofilm	1.9±0.9c	-28.4±3.5ab	4.37±1.86a	38.59±16.65ab
Horizon O	7.9±0.5b	-29.2±1.6b	4.99±0.78a	55.27±8.09a
Horizon A	6.0±0.4bc	-24.8±1.3ab	1.60±0.51b	17.60±3.39bc
Horizon Bw	16.6±3.1a	-23.5±0.4a	0.95±0.87b	8.65±4.48c

165 *MS = moss substrate



Table 1 depicts the average values and standard deviations of the elemental and isotope composition of carbon (C) and nitrogen (N) in moss, moss substrate (MS), biofilm, and three soil horizons (Horizon O, A, and Bw). Horizon O samples exhibited the highest average of C and N content (55.3 ± 8.1 and 5.0 ± 0.8 , respectively), which may be associated with a noticeable root layer (Fig. 1). The biofilm also shows notable C and N content (38.59% and 4.37% , respectively) and is located in Horizon O. These values are typical of biofilms, and its ratio C/N close to 9 (8.83 on average) indicates that the biofilm was found in a healthy state. Previous studies have shown that biofilms with C/N ratio close to 9 have higher concentrations of carbohydrates and proteins in the biofilm and a significant upregulation of the Quorum Sensing genes (Ramos et al., 2023).

In contrast, the C and N values in the horizons A (17.6 ± 3.39 and 1.6 ± 0.51 , respectively) and Bw (8.65 ± 4.48 and 0.95 ± 0.87 , respectively) were consistent with those reported in different soil types, the concentration of both elements in horizon A is three-fold of those found in B (Girona-García et al., 2018; Jiménez-Morillo et al., 2020a). These findings align with previous studies indicating a decline in C and N content with increasing soil depth. For moss and moss substrate (MS), the higher C content in MS compared to moss could be attributed to the presence of numerous small moss roots, which form a three-dimensional network that aggregates and stabilizes lapilli grains. Additionally, the presence of root exudates, rich in C and N compounds, may contribute to this remarkable concentration of organic compounds. This is the first indication of soil regeneration after the volcanic eruption of Tajogaite, which reveals the importance of moss in this process. Previous studies about the colonization of soils affected by volcanic activity highlight the abundance of photosynthetic organisms, including moss, as pioneer colonizers (Smith, 1984; Ingimundardóttir et al., 2014; Vilmundardóttir et al., 2018; Dragone et al., 2023).

Regarding the C isotope composition, all samples exhibited $\delta^{13}\text{C}$ values within the typical range for C_3 plants (between -25‰ and -35‰ (Katsumi et al., 2015; Ghezzi and Jimenez-Morillo et al., 2024). However, significant differences were observed among the samples. Moss, Horizon O, and biofilm displayed the most negative $\delta^{13}\text{C}$ values ($-28.5\text{‰} \pm 0.1$, $-28.4\text{‰} \pm 3.5$, and $-29.2\text{‰} \pm 1.6$, respectively), which reflected the presence of fresh organic matter (Jiménez-Morillo et al., 2020b). In contrast, samples from the A and Bw soil horizons and the moss substrate (MS) were enriched in ^{13}C (Table 1). This trend has also been observed by Girona-García et al., (2018) in soil profiles under pine and beech forests, where deeper horizons exhibited higher $\delta^{13}\text{C}$ values. Such ^{13}C -enrichment in soil samples is often associated with more humified organic matter (Jiménez-Morillo et al., 2020b). For the MS ($\delta^{13}\text{C} = -24.3\text{‰} \pm 1.6$), the observed ^{13}C -enrichment may be attributed to i) the heavier isotope



signature of root exudates compared to other organic compounds commonly found in soil samples (Girona-García et al., 2018), and ii) microbial activity linked to diverse families of microorganisms (Jiménez-Morillo et al., 2020b; Miller et al., 2022; San-Emeterio et al., 2023). In addition, Horizons A and Bw have similar $\delta^{13}\text{C}$ values (-24.8 ± 1.3 and -23.5 ± 0.4 , respectively) to 195 the MS, which is mainly composed of tephra. This could be indicative of the volcanic origins of the paleo soil (Horizons A and B) (Carracedo et al., 2022) and the importance of lapilli, among other volcanic materials from previous eruptions, in the carbon cycle during the soil formation (Matus et al., 2014 and references therein).

A key finding is the strong negative correlation observed between $\delta^{13}\text{C}$ values and C content across the different soil horizons and the biofilm (Fig. 2). This trend is particularly pronounced in the soil horizon samples ($R^2 = 0.9991$; solid line). However, 200 when the biofilm sample is included, the correlation weakens slightly ($R^2 = 0.9572$; dashed line).




Figure 2: Plot of average ($n=3$) $\delta^{13}\text{C}$ vs %C of moss (red triangle), moss substrate (MS, red square), biofilms (black dashed circle) and soil horizons O, A and Bw (open black circles) with linear regression of horizons (solid line) and including biofilms (dashed line).

205 This negative correlation between $\delta^{13}\text{C}$ and C content has been previously described in soils worldwide. In all cases, this trend has been attributed to either the decomposition of organic matter, the fractionation by microbial activity or the shift of a



predominant contribution from plants to microorganisms in the residual organic matter as it progresses through the soil horizons (Ågren et al., 1996; Volk et al., 2018; Ehleringer et al., 2000; Krüger et al., 2024). The microbial activity may process fresh organic matter, leaving behind evolved (humified) material with a heavier isotope. This alteration may also reduce the C
210 content due to microbial respiration. A similar pattern has been observed in the isotope composition of highly humified (evolved) organic matter associated with very fine sand particles in shallow soil layer (Jiménez-Morillo et al., 2020a), which may confirm the influence of microorganisms in the humification process of fresh organic matter. However, the moss and the tephra collected under the moss (MS) did not correlate with the soil. This is probably because the samples were collected shortly after the volcanic eruption that accumulated the tephra over the previous soil (Horizon O). Therefore, the C cycle of
215 this interface was still in an early stage of the andisol formation process, preventing the observation of statistical correlations between the soil samples and the tephra.

Regarding the nitrogen isotope composition ($\delta^{15}\text{N}$), the moss sample exhibited the lowest $\delta^{15}\text{N}$ value (Table 1), a typical characteristic for mosses (Zechmeister et al., 2008). In contrast, the other samples displayed positive $\delta^{15}\text{N}$ values (Table 1). For the MS sample, the ^{15}N -enrichment and their statistical similarity with $\delta^{15}\text{N}$ values observed in the soil and biofilm samples
220 suggest that it could be derived from the activity of certain microorganisms, such as cyanobacteria, that utilize nitrogen-derived compounds excreted by the moss roots (Carrell et al., 2022). In addition, the deepest soil sample (Horizon Bw) showed the highest $\delta^{15}\text{N}$ enrichment, which could be associated with denitrification processes (Girona-García et al., 2018). While humification is well known to cause ^{15}N -enrichment, particularly in forest soils (Szpak, 2014), ^{15}N -depletion can occur during the uptake of inorganic N by vegetation (Högberg, 1997). Therefore, the observed trend can be best explained by the alteration
225 of biomass by heterotrophic organisms, which produce ^{15}N -depleted compounds while progressively enriching the remaining biomass with ^{15}N over time (Billings and Richter, 2006; Makarov, 2009; Szpak, 2014; Girona-García et al., 2018).

3.2 Molecular composition of organic fraction

The use of analytical pyrolysis (Py-GC/MS) has revealed remarkable molecular differences between soil, tephra, biofilm, and
230 moss samples (Fig. 3). The moss sample is rich in organic compounds, particularly lipids (31.56%) and polysaccharides (29.75%), aligning with findings from previous studies (Klavina et al., 2015; Petkova et al., 2023). Among the lipid compounds



identified were essential oils, such as α -Terpinene and Phytol, *n*-alkanes (e.g., heptadecane, heneicosane; Klegin et al., 2023) and *n*-alkanoic acids (e.g., 9,12,15-Octadecatrienoic acid, 13-Eicosenoic acid, 22-Tricosenoic acid; Yucel, 2021; Klegin et al., 2023). Polysaccharides in bryophytes play an important role in the chemical evolution of vegetation and assist as structural components, which are also assumed to contribute to bryophyte stress tolerance (Klavina et al., 2015). In this study, the identified polysaccharides were primarily furans (e.g., 2(5H)-Furanone, 5-methyl-, 2-Furancarboxylic acid, methyl ester, 3-Furaldehyde) and levoglucosan (β -D-Glucopyranose, 1,6-anpah). However, due to dehydration and molecular restructuring during pyrolysis, complete identification of polysaccharides is not feasible (González-Pérez et al., 2016). The molecular analysis also revealed a significant presence of non-specific aromatic compounds (13.60%) and peptide compounds (11.69%).

Among the aromatic compounds, phenol derivatives such as Phenol, 2-methyl-, Phenol, 2,3-dimethyl-, 1,2-Benzenediol, and Phenol, 2,3,5-trimethyl were prominent. These phenolic compounds are considered critical in early plant evolution, particularly in moss growth and cuticle permeability, which were key for adaptation to terrestrial environments (Renault et al., 2017). Such phenolic compounds are precursors to lignin, synthesized by the phenylpropanoid pathway, which strengthens cell walls, aids in water transport, and supports upright growth (Fraser and Chapple, 2011). Peptide-derived compounds were also found, with their primary function being the defence of mosses against environmental stressors such as drought and high temperatures (Ren et al., 2024). Additionally, these peptides provide resistance against microbial pathogens (Valeeva et al., 2022). Plants produce a range of bioactive metabolites, peptides, and small molecules that contribute to defence mechanisms and microbial resistance (Valeeva et al., 2022). In addition, peptide compounds are crucial regulators of ecosystem function because they may provide as direct supplies of nutrients for soil microorganisms and plants (Wang et al., 2023).

Only a few organic compounds were identified in the MS samples, mainly high molecular weight alkyl lipids (*n*-alkanes) and sterols, being the second the dominant group (65%). Key sterols detected include Stigmastan-3,5-dien, Campesterol, and γ -Sitosterol, also found in the moss samples (Table S2). The presence of Campesterol and γ -Sitosterol may indicate an input of organic matter from the moss via rhizoid exudates. Sterols are crucial for various plant physiological processes, not only supporting cell membrane structure and fluidity but also acting as hormone precursors and playing a role in responses to biotic and abiotic stress (Valitova et al., 2016; Cabianna et al., 2021). In addition, the presence of high molecular weight alkyl compounds, originating from epicuticular waxes, supports the hypothesis that there is a direct input of organic material from



the moss to the substrate (tephra). The $\delta^{13}\text{C}$ value observed in this sample does not align with the molecular composition identified by Py-GC/MS, as the detected compounds usually exhibit a markedly negative isotope signature (Park and Epstein, 1961; San-Emeterio et al., 2023). However, the observed ^{13}C -enrichment in this sample against the moss could be attributed
260 to low-molecular-weight polysaccharide-derived compounds undetected by this analytical technique.


The biofilm sample collected from Horizon O consists mainly of lipids (40.54%), non-specific aromatic compounds (38.36%), polysaccharides in the form of furans (13.08%), and peptides (8.02%). This molecular composition aligns perfectly with the main molecular composition found in biofilm samples analyzed in different volcanic regions (Miller et al., 2020; Ghezzi and Jiménez-Morillo et al., 2024; Palma et al., 2024). The lipid composition is mainly dominated by *n*-alkane/alkene pairs with
265 different proportions, which will be discussed in the section 3.3.1.

The Horizon O sample is dominated by lipid compounds (42.43%), mainly alkyl compounds (*n*-alkane/alkene pairs and *n*-alkanoic acids), followed by aromatic moieties (17.80%) and polysaccharides (12.57%). In addition, this is the only sample with a relatively high concentration of methoxyphenols, i.e., lignin-derived compounds (12.18%). The abundance of alkyl compounds suggests the existence of different organic fractions from distinct origins: i) organic matter from pre-eruption
270 vegetation, also indicated by the $\delta^{13}\text{C}$ of the Horizon O sample (-29.2 ± 1.6) typical of C_3 plants, ii) the degradation of lipid membranes from microorganisms induced by the deposition of tephra during the Tajogaite volcanic eruption, and iii) highly evolved organic matter from post-eruption microbial activity in the soil (Jiménez-Morillo et al., 2017; Bull et al., 2000; Jiménez-Morillo et al., 2016b; Leal and Jimenez-Morillo et al., 2023). In contrast, the presence of lignin- and polysaccharide-derived compounds probably indicates the recent input of herbaceous/shrubby plant material (Fig. 1). It could also stem from
275 root exudates of vegetation present before the ash deposition, supported by the high concentration of roots in the Horizon O (Fig. 1). Lignin compounds, particularly from the guaiacol and syringol groups, confirm a contribution of tree-derived organic matter (San-Emeterio et al., 2023). However, the greater abundance of guaiacol over syringol suggests ongoing microbial degradation activity, as syringol is more easily degraded by microorganisms (San-Emeterio et al., 2023). Additionally, condensed compounds (PAHs) were found in the shallow horizon. This could be generated by: i) ancient fires with particles
280 trapped in the soil layers (Jiménez-Morillo et al., 2016b), ii) high-temperature vapour emissions, from volcanic fracture, leading to defunctionalization and condensation of polar aromatic compounds (Capaccioni et al., 1995) or, iii) microbial



activity. Several pathways have been described for hydrocarbon synthesis by microorganisms (Ladygyna et al., 2006 and references therein). Some of these biogenic PAHs have been detected in forest soil (Kraus et al., 2005) with high degradation of lignin, in peatlands and wetlands (Wakeham et al., 2016), where organic matter accumulates and transforms slowly, and in
285 lake and marine sediments (Love et al., 2021), where microbial activity and anoxic conditions promote the polymerization of aromatic compounds.

The two deepest Horizons (A and Bw) of the soil profile share a similar molecular composition, dominated by non-specific aromatic compounds, lipid compounds (mainly *n*-alkane/alkene pairs), polysaccharides and peptides, with a complete absence of lignin molecules. This pattern has been observed in forest soils across the Iberian Peninsula (Aznar et al., 2016; Jiménez-Morillo et al., 2016a; De la Rosa et al., 2019), indicating the presence of microbial-derived compounds and linked to high
290 microbial activity (Jiménez-Morillo et al., 2018). The abundance of non-specific aromatic compounds may also result from fungal degradation of plant lignins (Kellner et al., 2014). However, important differences include a higher proportion of condensed compounds (PAH = 15.17%) in the upper Horizon, while the deeper horizon has more sterane compounds (10.76%). In the Horizon A, highly condensed PAHs, such as phenanthrene and pyrene suggest the presence of pyrogenic material,
295 usually associated to organic matter combustion (González-Pérez et al., 2014; Miller et al., 2020, 2022). In contrast, the deeper Horizon only contains branched naphthalene compounds, possibly due to lower-intensity fires or microbial degradation of condensed compounds by thermophiles (Musat et al., 2009; Mohapatra and Phale, 2021). In the sterane family, the deeper Horizon (Bw) contains Stigmastan-3,5,22-triene, Ergosta-4,6,22-trien-3-ol, and Friedelan-3-one. These plant- and fungi-synthesized compounds are well-preserved in deeper layers due to their stability (Kristan and Rižner, 2012), with clear
300 accumulation in the lower horizons, while they are almost absent in the Horizon O and the biofilm (Fig. 3).




305 **Figure 3: Relative abundance (%) of the main organic families identified by Py/GC/MS from moss, moss substrate (MS), biofilm, and soil horizons (O, A, Bw) samples. The main organic families are: unspecific aromatic compounds (ARO), polysaccharides (POL), peptides (PEP), lignin (LIG), lipids (LIP), nitrogen-containing compounds (N-comp), polycyclic aromatic hydrocarbons (PAHs), and steranes (STR).**



3.3. Lipid analysis by Py-GC/MS

3.3.1. *n*-Alkanes



310

Figure 4: *n*-alkane (m/z 57) distribution obtained by direct pyrolysis (Py-GC/MS) of moss, moss substrate (MS), biofilms and soil horizons (O, A and Bw) samples.



The *n*-alkane series were identified using m/z 57 in the selected-ion-monitoring (SIM) chromatogram (Fig. 4). Chain lengths
315 ranged from C_{10} to C_{35} , reflecting various organic matter sources (Jiménez-Morillo et al., 2017). In the moss sample, a bimodal
distribution was observed with peaks at *n*-pentadecane (C_{15}) and *n*-hentriacontane (C_{31}), with odd-chain *n*-alkanes
predominating, especially in longer chains ($>C_{25}$). This suggests *n*-alkanes from moss epicuticular waxes, with a peak at *n*-
hentriacontane (Schwark et al., 2002). The presence of low-molecular-weight (LMW) *n*-alkanes ($<C_{20}$) may point to associated
microorganisms, as LMW *n*-alkanes typically originate from microbial cell walls (Jiménez-Morillo et al., 2017, 2022).

320 In the MS sample, the dominance of odd-chain high-molecular-weight (HMW) *n*-alkanes supported direct organic matter input
from moss. However, the abundance of *n*-tritriacontane (C_{33}) and *n*-pentatriacontane (C_{35}) *n*-alkanes may also indicate C4
vegetation (Vogts et al., 2009; Feakins et al., 2016) or the adaptation of pre-eruption vegetation to water stress caused by
volcanic ash.

In Horizon O, a bimodal *n*-alkane distribution with peaks at C_{15} and *n*-docosane (C_{22}) suggested the presence of organic matter
325 from soil microorganisms or the degradation of plant biomass by microorganisms over time (Jiménez-Morillo et al., 2016b,
2017). The gradual decline in HMW *n*-alkanes and the absence of a clear odd-over-even pattern indicates microbial degradation
processing of organic matter (De Blas et al., 2013; Jiménez-Morillo et al., 2022; Leal and Jimenez-Morillo et al., 2023). The
higher root density in this layer could also contribute to the observed *n*-alkane pattern.

The biofilm sample was dominated by LMW *n*-alkanes, confirming its microbial origin (Jiménez-Morillo et al., 2017). The
330 presence of branched alkanes like neophytadiene, 2,6,10-trimethyltridecane, and 2,6,11-trimethyldodecane further supports a
microbial contribution (Palma et al., 2024).

In deeper horizons (A and Bw), there is a clearer differentiation between LMW and HMW *n*-alkanes, indicating two carbon
pools from microorganisms (LMW *n*-alkanes) and plant vegetation biomass (HMW *n*-alkanes). The microbial fraction may
stem from lipidic membranes (Miller et al., 2016) or microbial degradation of plant biomass, such as alkyl compounds from
335 lignocellulose breakdown (Rabbi et al., 2014; Jiménez-Morillo et al., 2022; Miller et al., 2022). The nonspecific aromatic
compounds previously discussed also suggest the lignin breakdown process. However, the most significant differences among
horizons are seen in HMW *n*-alkanes. The Horizon A peaks at *n*-nonacosane (C_{29}), indicating tree or shrub vegetation, while

<https://doi.org/10.5194/egusphere-2025-2086>

Preprint. Discussion started: 15 May 2025

© Author(s) 2025. CC BY 4.0 License.



the Horizon Bw peaks at *n*-hentriacontane (C_{31}), suggesting herbaceous vegetation (Bull et al., 2000). Therefore, results indicate a change in vegetation probably due to the alteration of the environmental conditions.

340



3.3.2. *n*-Alkane-2-ones




Figure 5: *n*-alkane-2-one (m/z 59) distribution was obtained by direct pyrolysis (Py-GC/MS) of moss, moss substrate (MS), biofilms and soil horizons (O, A and Bw) samples.



345 The pyrolyzed samples from La Palma, except for the biofilm sample, revealed a series of *n*-alkane-2-ones (m/z 59 ion chromatograms) ranging from C₁₇ to C₃₁, with a general odd-over-even chain-length predominance (Fig. 5). However, each sample showed distinct patterns. In the moss sample, the maximum was at *n*-alkane-2-one C₂₃, without a clear parity pattern, though the odd-over-even trend was evident in LMW *n*-alkane-2-ones. This suggests a mixture of lipids from both the moss and its associated microbiome, as also seen in the *n*-alkanes (Fig. 4).

350 The MS sample showed a *n*-alkane-2-ones distribution similar to that observed for *n*-alkanes (Fig. 4), being dominated by C₂₇ and C₃₁ molecules, which supports the idea of a direct contribution of organic matter from the moss.

The soil horizons showed differences in the *n*-alkane-2-one distribution (Fig. 5). In Horizon O, the maximum was at *n*-alkane-2-one C₂₁, while in horizons A and Bw, the maxima were at C₂₅ and C₂₇, respectively. The Horizon O had a range of *n*-alkane-2-ones from C₁₇ to C₂₉, suggesting more degraded organic matter, possibly due to fine ash deposition affecting organic matter or microorganisms. In Horizon A, the *n*-alkane-2-one distribution resembled typical soils, with two carbon pools: i) LMW *n*-alkane-2-ones from microbial activity, and ii) HMW ones from plant-derived organic matter (Jaffé et al., 1996; Kumar et al., 2020). This aligns with the *n*-alkane distribution (Fig. 4) and overall molecular analysis (Fig. 3). In contrast, the Horizon Bw contained only HMW *n*-alkane-2-ones (>C₂₅), considered biomarkers of vegetation (Kumar et al., 2020; Palma et al., 2024). This aligns with the isotope carbon signature observed in the soil (Table 1 and Fig. 2) that indicates the influence of microbial processes at the top layer (Horizon O) and a gradual shift to plant-influenced processes towards the bottom (Horizon Bw). The absence of LMW *n*-alkane-2-ones in Horizon Bw could be due to concentrations below detection limits, as this horizon had low TOC (Fig. 1). Environmental factors such as moisture and nutrient availability may affect *n*-alkane-2-one formation by altering the degradation of *n*-alkanes and *n*-alkanoic acids (Lehtonen and Ketola, 1990; Zhang et al., 2004). Zhang et al. (2004) identified potential sources of *n*-alkane-2-ones, including: (i) epicuticular waxes from plants; (ii) microbial oxidation of *n*-alkanes; (iii) β -oxidation and decarboxylation of *n*-alkanoic acids; and (iv) microbial cracking of HMW *n*-alkane-2-ones into LMW ones. The predominance of LMW *n*-alkane-2-ones in Horizon O further supports an active microbial contribution to organic matter transformation.

360

365



370 3.4 Bacterial community composition

A total of 198.129 ASVs (amplicon sequence variants) were analysed from 16S rRNA gene sequencing. The richness and evenness of microbial communities were estimated using Chao1, Shannon and Simpson indices (Table 2). The biofilm was the most biodiverse sample, with the highest species richness, and Shannon and Simpson indices of 7.860 and 0.989, respectively, followed by horizons Bw and A and moss. In contrast, Horizon O showed a lower diversity, with Shannon and Simpson indices of 4.166 and 0.689, respectively. According to the Chao1 richness estimator, Horizon O and the biofilm showed the highest predicted microbial richness, indicating that most of the species present in these samples were successfully detected, with minimal undetected diversity. It is noteworthy that the microbial community in the tephra collected beneath the moss (MS) reached a substantial diversity and richness within just two years following the deposition of the Tajogaite volcanic tephra.

380


Table 2: Alfa diversity 16S of moss, moss substrate (MS), biofilms and soil horizons samples.

Sample	Shannon	Simpson	chao1	Observed
Moss	6.359	0.973	426.143	426
MS	5.522	0.950	255.500	254
Biofilm	7.860	0.989	832.783	832
Horizon O	4.166	0.689	833.948	829
Horizon A	6.531	0.972	510.315	473
Horizon Bw	7.321	0.972	814.125	812

Most of the sequences identified belong to the Bacteria kingdom; however, Archaea, was also found, primarily in the moss, biofilm and MS samples, comprising 2.63%, 2.48% and 1.31%, respectively (Fig. 6A). They have been reported as early colonizers in samples from the volcanic island *Hunga Tonga Hunga Ha'apai*, in the Kingdom of Tonga (Fujimura et al., 2016; Dragone et al., 2023). Archaea are known for their ability to inhabit extreme environments and play critical roles in biogeochemical cycles (Offre et al., 2013; Martínez-Espinosa, 2020). The main archaeal class detected in these samples was *Nitrososphaeria*, belonging to the phylum *Nitrososphaerota*, which plays a key role in the nitrogen cycle by transforming ammonia into nitrite, thereby contributing to the nitrogen availability in the soil. This process indirectly supports plant nutrition



390 (Leininger et al., 2006; Lehtovirta-Morley et al., 2024) and influences the distribution of N concentration described in the biofilm and MS samples. The presence of N in these samples infers that the identified genus *Nitrososphaeria* is oxidizing the ammonium to nitrite, which requires the presence of oxygen. This might explain the presence of this archaea only in the moss, MS and biofilm samples, which are in contact with the atmosphere.



395 **Figure 6: Relative abundances of microbial phyla (A) and genera (B) across different samples collected from moss, moss substrate (MS), biofilm, and soil horizons (O, A and Bw) samples.**

The bacterial phylum *Pseudomonadota* was found in all samples, with relative abundances decreasing with depth, ranging from 58.1% (moss) to 5.6% (horizon Bw). Members of this phylum included *Acidibacter*, the *Craurococcus-Caldovatus* bacteria group, and the *Methyloligellaceae* family (Fig. 6B). The acidophilic, ferric iron-reducing, heterotrophic bacterium *Acidibacter*, previously isolated from the Guadiana pit lake, currently has only one described species (Falagán and Johnson, 2014). This bacterium was found in all three horizons, although its abundance decreased with depth, indicating its adaptation to metabolize fresh organic matter, which is more abundant in the Horizon O. The *Caldovatus* genus, a thermophilic bacterium, was found in the MS sample (6.29%). This genus includes only two described species, *C. aquaticus* and *C. sediminis* (Habib



405 et al., 2017; Hu et al., 2022). The *Methyloligellaceae* family is characterised by its ability to use methane as a carbon and energy source, playing a key role in the carbon cycle (Bryukhanov et al., 2024), dominating the Horizon A, MS and biofilm samples.

Other bacterial genera identified greater than 1% of relative abundance were *Aureimonas*, *Bosea*, *Hyphomonas*, *Legionella*, *Methylorubrum* and *Nordella* in the moss sample; *Candidatus Halysiosphaera*, *Limnobacter*, and *Chthonobacter* in the MS; 410 *Dongia* in the biofilm; and *Bauldia* in both the biofilm and Horizon A, and *Brevundimonas* in both the moss and MS samples. Notably, the genus *Bosea*, detected in the moss sample, is known for its ability to obtain energy through the oxidation of reduced sulfur compounds, a process highly relevant in volcanic environments where sulfur-rich fumaroles, such as those produced during the Tajogaite eruption, have been documented (Martínez-Martínez et al., 2023). Furthermore, this metabolic activity relies on the availability of organic carbon, which was confirmed in all samples through EA/IRMS and Py-GC/MS 415 analyses, highlighting the active coupling between volcanic sulfur emissions and early microbial colonization of the tephra substrate.

The phylum *Chloroflexota* was also present in all samples and was particularly abundant in the O, A and Bw horizons, with relative abundances of 27.23%, 26.77% and 38.92%, respectively. The most abundant members of this phylum were TK10, S085 and KD4-96, present in all samples along with the thermophilic bacteria *Thermogemmatispora* and *Litorilinea*, found 420 predominantly in horizon Bw and the biofilm, respectively. Species of *Thermogemmatispora* are involved in carbon monoxide oxidation in geothermal environments (King and King, 2014). The *Candidatus Chloroploca*, found in the moss, is capable of using sulfide oxidation as an electron donor (Gorlenko et al., 2014).

The *Actinomycetota* was the most abundant phylum in the MS sample, with the following genera being particularly prominent: *Nocardioides* (9.33%), IMCC26256 (2.02%), 67-14 (1.64%), *Conexibacter* (1.23%) and *Gaiella* (1.01%). In the biofilm, the 425 *Micronomospora*, *Actinophytocola*, and 67-14 genera were the most abundant, with a relative abundance of 4.16%, 1.81%, and 2.94%, respectively. In the soil horizons, thermophilic bacteria such as *Thermasporomyces* and *Thermopolyspora* (Horizon O) and *Thermoleophilum* (dominant in the Horizon Bw) were prevalent. Species of the *Thermopolyspora* genus are capable of degrading lignocellulosic compounds (Yang et al., 2022), which aligns with the detection of lignin exclusively in the Horizon O sample by Py-GC/MS. In contrast, members of the genus *Thermoleophilum* can grow exclusively on C₁₂–C₁₉ n-alkanes,



430 cyclohexane, cycloheptane, C₁₂–C₁₈ alcohols and C₁₃–C₁₉ ketones (Yakimov et al., 2003), all of which were identified in the soil samples through Py-GC/MS (Table S2).

The *Bacillota* was a dominant phylum in Horizons Bw (26.29%) and O (14.28%), as well as in the MS sample (6.90%). The most abundant genera within this phylum in the soil horizon samples were *Brockia*, *Thermaerobacter*, *Thermobacillus*, and *Lihuaxuella* (dominant in the Horizon Bw). The *Limnochorda* was found primarily in all three horizons, while *Planifilum*,
435 dominant in Horizons O and Bw, plays a key role in the degradation of organic matter and harbors potential antibiotic-resistance genes (Tseng et al., 2024). This genus was also found in the basaltic subsurface of young volcanic islands in Iceland (Bergsten et al., 2022). Non-thermophilic genera of the *Bacillota* phylum, such as *Exiguobacterium* and *Bacillus*, were also identified in all three sample types.

The *Thermomicrobiota* phylum was present in all samples, particularly in horizons O (13.63%) and A (12.53%), mainly
440 represented by the *Sphaerobacter* genus, which was found in the three soil horizons, with particularly high dominance in Horizon O (20.51%). This bacterium was originally isolated from thermophilically treated sludge, and proteogenomic analysis indicated that it contains numerous aromatic degradation pathways, such as those involved in lignin degradation (D'haeseleer et al., 2013), suggesting its role in the breakdown of complex organic matter. Notably, lignin was detected in the Horizon O sample (Fig. 3), while aromatic compounds have been detected in all samples (Fig. 3).

445 The hyperthermophilic family *Thermomicrobiaceae* was also found, predominantly in horizon A (15.11%), represented by the 67-14 bacterium.

The *Cyanobacteriota* phylum was mainly found in the biofilm (20.64%), MS (3.07%), and moss (3.04%) samples. Members of this phylum play a crucial role as nitrogen fixers and contribute significantly to the primary production of organic compounds (Rippin et al., 2018), as represented by genera such as *Arthronema* and *Nodosilinea*. This is consistent with the
450 detection of nitrogen in the biofilm, MS, and moss samples, and supported by isotope ratios that predicted the presence of photosynthetic microorganisms. *Arthronema* is a filamentous cyanobacterium highly adaptive, typical of extreme desert environments, where it contributes to the nutritional chain and soil formation. Additionally, it exhibits antitumor and antioxidant properties and produces phycobiliproteins (Iliev and Andreeva, 2006; Chaneva et al., 2007; Gardeva et al., 2014;



Petrova et al., 2020). Markers of proteins (peptides) were detected by Py-GC/MS in all samples, except MS. Although this
455 technique cannot identify specific proteins, the presence of phycobiliproteins cannot be ruled out.

Other phyla identified in these samples included *Acidobacteriota*, mainly found in the biofilm, MS and the A and Bw horizons;
Gemmatimonadota, predominantly present in the biofilm and soil horizon samples; and *Deinococcota* and *Bacteroidota*,
primarily in the MS sample, represented by *Meiothermus* and *Persicitalea*. The *Deinococcota* phylum is known as one of the
most extremophilic phyla of bacteria, due to its resistance to extreme stressors such as oxidation, radiation, desiccation, and
460 high temperature (Vilhelmsson et al., 2023).

 In general, the sampling area exhibited a significant presence of thermophilic bacteria, particularly in Horizons O and Bw, and
in the biofilm. This abundance is likely related to the proximity of steam-emitting fumaroles near the sampling site. These
thermophilic microorganisms, which thrive at elevated temperatures, are highly adapted to the geothermal conditions generated
by the fumaroles. Thermophilic genera such as *Thermasporomyces*, *Thermopolyspora*, *Thermobacillus*, *Planifium* and
465 *Pyrinomonas*, were observed in the soil horizons. Notably, the only described species of the genus *Pyrinomonas*, *Pyrinomonas*
methylaliphatogenes, was originally isolated from geothermally heated soils associated with fumaroles (Crowe et al., 2014).
Members of these genera are recognized for their ability to degrade complex organic materials, such as lignocellulose (Yang
et al., 2022), detected in these soil samples, and for their active role in the degradation of organic matter (Tseng et al., 2024).
This enzymatic capability is particularly beneficial in volcanic soils, where organic matter turnover is critical for early soil
470 development and nutrient cycling. The dominance of *Thermoleophilum* in the Horizon Bw highlights its ecological role in
metabolizing hydrocarbons, such as PAHs, present in all the soil samples analysed. The hyperthermophilic family
Thermomicrobiaceae, predominantly found in Horizon A, and the genus *Sphaerobacter*, highly abundant in the Horizon O, as
well as the thermophilic bacteria *Thermogemmatispora* and *Litorilinea*, predominantly in the deeper soil horizons and the
biofilm, further underline the influence of geothermal activity on microbial composition.

475 The presence of thermophiles across all collected samples, along with their co-occurrence with non-thermophilic genera,
suggests an ecological adaptation of the microbial community to thermal disturbances associated with volcanic activity and
fumarolic emissions. Furthermore, the relatively lower abundance of thermophiles in horizon A compared to horizons O and
Bw suggests that thermal events occurred more frequently or with great intensity during the formation of horizons O and Bw.





480 The adaptation of thermophiles to high temperatures enables them to dominate during periods of elevated temperatures (e.g.,
in the presence of fumaroles), while mesophilic counterparts persist and thrive in cooler periods. Temperature and humidity
were continuously monitored at the sampling site using a HOBO sensor over one year. During this period, temperatures
generally fluctuated between 20°C and 30°C, with the highest temperature (51.33°C) recorded on August 12 and the lowest
(13.76°C) recorded on February 24 (Fig. S1B). The highest temperatures always coincided with the periods of lowest recorded
humidity.

485



4 Conclusions

This study demonstrates that the moss colonizing the upper layer of the tephra deposits releases organic compounds that serve
as carbon sources for microorganisms, facilitating rapid microbial colonization of the inert volcanic substrate within two years
after the Tajogaite eruption. This eruption produced a complex tephra blanket exceeding 50 cm in thickness, composed of
490 lapilli and fine ash, which accumulated over the pre-existing soil (Horizon O).

Our analyses reveal that Horizon O exhibits a distinctive hydrocarbon signature, indicative of both high-temperature thermal
processes and active microbial transformation of organic matter. In contrast, the deeper soil layers (Horizons A and Bw) display
alkane patterns characteristic of plant-derived organic matter inputs. Furthermore, $\delta^{13}\text{C}$ isotopic data strongly suggest intense
microbial processing of organic carbon in Horizon O, reinforcing the hypothesis that microbial activity significantly
495 contributes to the formation of hydrocarbons in this horizon.

Importantly, microbial sequencing revealed the presence of taxa capable of hydrocarbon metabolism, including the family
Methylobacteriaceae, supporting the hypothesis that some of the detected polycyclic aromatic hydrocarbons (PAHs) could be
of biogenic origin.

In addition, our results indicate that the deposition of fresh tephra materials from the Tajogaite Volcano has the potential to
500 alter soil organic matter (SOM) dynamics, favouring the accumulation of labile organic compounds and reducing the
abundance of recalcitrant components such as lignin, which are typically more resistant to degradation.

The dominance of thermophilic bacteria, particularly in Horizons O and Bw, highlights their key ecological role in the
breakdown of recalcitrant organic matter, the facilitation of nitrogen and carbon cycling, and the stabilization of the evolving



soil structure. Their metabolic versatility, including the ability to degrade lignin-derived compounds and synthesize
505 hydrocarbons, positions thermophiles as critical drivers in the early formation of andisols and the recovery of soils disturbed
by volcanic activity.

Finally, the persistent presence of thermophilic microorganisms across the studied profile reflects the influence of volcanic
and fumarolic activity in shaping the microbial community structure in the post-eruptive landscape of the Tajogaite Volcano.
Given the recurrent volcanic activity on La Palma Island, particularly along the Cumbre Vieja Volcanic Ridge, thermophilic
510 communities are likely an enduring and essential component of soil ecosystem dynamics in this region.

Author contributions

The manuscript was written through the contributions of all authors. All authors have approved the final version of the
manuscript. S.G-P: data curation, formal analysis, methodology, visualization, writing—original draft. A.G.A.: data curation,
515 formal analysis, methodology, writing—original draft. P.N-J.: methodology, validation, writing—original draft. J.M.-S.:
methodology, validation, writing—review and editing. J.M-M.: methodology, validation, writing—review and editing. B.M-H.:
methodology, writing—review and editing. J.V.: methodology, validation, writing—review and editing. N.T.J.-M.: data curation,
formal analysis, methodology, visualization, validation, writing—original draft, writing—review and editing. A.Z.M.: data
curation, formal analysis, methodology, validation, visualization, supervision, funding acquisition, writing—original draft,
520 writing—review and editing.

Competing interests

The authors declare that they have no conflict of interest.

525 Acknowledgements

This work received support from the Spanish Ministry of Science, Innovation and Universities (MICIU) under the research
project HIRES-SOM (ref. TED2021-130683B-C21/C22) funded by MCIN/AEI/10.13039/501100011033 and the European
Union “NextGenerationEU”/PRTR. The research project MICROLAVA (ref. PROYEXCEL_00185) funded by Junta de

Andalucía is also acknowledged. N.T. Jiménez-Morillo acknowledges the “Ramón y Cajal” contract (ref. RYC2021-031253-
530 I) funded by MCIN/AEI/10.13039/501100011033 and the European Union “NextGenerationEU”/PRTR. A. Gómez-Arias
acknowledges the “Juan de la Cierva” contract (ref. JDC2022-049199-I). The authors are grateful to José Heriberto López,
forest officer at the Caldera de Taburiente National Park (La Palma, Spain) for providing logistical support during the sampling
campaign.

535 **References**

- Ågren, G. I., Bosatta, E., and Balesdent, J.: Isotope discrimination during decomposition of organic matter: A theoretical
analysis, *Soil Sci. Soc. Am. J.*, 60, 1121–1126, <https://doi.org/10.2136/sssaj1996.03615995006000040023x>, 1996.
- Allaguvatova, R. Z., Nikulin, A. Y., Nikulin, V. Y., Bagmet, V. B., and Gaysina, L. A.: Study of biodiversity of algae and
cyanobacteria of Mutnovsky and Gorely volcanoes soils (Kamchatka Peninsula) using a polyphasic approach, *Diversity*, 14,
540 375, <https://doi.org/10.3390/d14050375>, 2022.
- Arias, M. E., González-Pérez, J. A., González-Vila, F. J., and Ball, A. S.: Soil health—a new challenge for microbiologists
and chemists, *Int. Microbiol.*, 8, 13–21, 2005.
- Arnalds, O.: The influence of volcanic tephra (ash) on ecosystems, in: *Advances in Agronomy*, by Sparks, D. L., Academic
Press 121, 331–380, <https://doi.org/10.1016/B978-0-12-407685-3.00006-2>, 2013.
- 545 Aznar, J. M., González-Pérez, J. A., Badía, D., and Martí, C.: At what depth are the properties of a gypseous forest topsoil
affected by burning? *Land. Degrad. Dev.*, 27, 1344–1353, <https://doi.org/10.1002/ldr.2258>, 2016.
- Bergsten, P., Vannier, P., Frion, J., Mougeolle, A., and Marteinsson, V. Þ.: Culturable bacterial diversity from the basaltic
subsurface of the young volcanic island of Surtsey, Iceland, *Microorganisms*, 10, 1177,
<https://doi.org/10.3390/microorganisms10061177>, 2022.
- 550 Biass, S., Reyes-Hardy, M. P., Gregg, C., Di Maio, L.S., Dominguez, L., Frischknecht, C., Bonadonna, C., and Perez, N.: The
spatiotemporal evolution of compound impacts from lava flow and tephra fallout on buildings: lessons from the 2021 Tajogaite
eruption (La Palma, Spain). *B. Volcanol.*, 86, 10, <https://doi.org/10.1007/s00445-023-01700-w>, 2024.
- Billings, S. A. and Richter, D. D.: Changes in stable isotopic signatures of soil nitrogen and carbon during 40 years of forest
development, *Oecologia*, 148, 325–333, <https://doi.org/10.1007/s00442-006-0366-7>, 2006.
- 555 Birnbaum, J., Lev, E., Hernandez, P. A., Barrancos, J., Padilla, G. D., Asensio-Ramos, M., Calvo, D., Rodríguez, F., Pérez, N.
M., D’Auria, L., and Calvari, S.: Temporal variability of explosive activity at Tajogaite volcano, Cumbre Vieja (Canary



- Islands), 2021 eruption from ground-based infrared photography and videography, *Front. Earth Sci.*, 11, <https://doi.org/10.3389/feart.2023.1193436>, 2023.
- de Blas, E., Almendros, G., and Sanz, J.: Molecular characterization of lipid fractions from extremely water-repellent pine and eucalyptus forest soils, *Geoderma*, 206, 75–84, <https://doi.org/10.1016/j.geoderma.2013.04.027>, 2013.
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., Cope, E. K., Da Silva, R., Diener, C., Dorrestein, P. C., Douglas, G. M., Durall, D. M., Duvall, C., Edwardson, C. F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J. M., Gibbons, S. M., Gibson, D. L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H., Huttenhower, C., Huttley, G. A., Janssen, S., Jarmusch, A. K., Jiang, L., Kaehler, B. D., Kang, K. Bin, Keefe, C. R., Keim, P., Kelley, S. T., Knights, D., Koester, I., Kosciulek, T., Kreps, J., Langille, M. G. I., Lee, J., Ley, R., Liu, Y. X., Lofffield, E., Lozupone, C., Maher, M., Marotz, C., Martin, B. D., McDonald, D., McIver, L. J., Melnik, A. V., Metcalf, J. L., Morgan, S. C., Morton, J. T., Naimey, A. T., Navas-Molina, J. A., Nothias, L. F., Orchanian, S. B., Pearson, T., Peoples, S. L., Petras, D., Preuss, M. L., Pruesse, E., Rasmussen, L. B., Rivers, A., Robeson, M. S., Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A., Sinha, R., Song, S. J., Spear, J. R., Swafford, A. D., Thompson, L. R., Torres, P. J., Trinh, P., Tripathi, A., Turnbaugh, P. J., Ul-Hasan, S., van der Hooft, J. J. J., Vargas, F., Vázquez-Baeza, Y., Vogtmann, E., von Hippel, M., et al.: Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2, *Nat. Biotechnol.* 37, 852–857, <https://doi.org/10.1038/S41587-019-0209-9>, 2019.
- Bonadonna, C., Pistolesi, M., Biass, S., Voloschina, M., Romero, J., Coppola, D., Folch, A., D`Auria, L., Marrin-Lorenzo, A., Dominguez, L., Pastore, C., Reyes Hardy, M. P., and Rodriguez, F.: Physical characterization of long-lasting hybrid eruptions: The 2021 tajogaite eruption of cumbre vieja (La Palma, Canary Islands). *J. Geophys. Res-Solid Ea.* 127, 11, <https://doi.org/10.1029/2022JB025302>, 2022.
- Bonadonna, C., Pistolesi, M., Dominguez L., Freret-Lorgeril, V., Rossi, E., Fries, A., Biass, S., Voloschina, M., Lemus, J., Romero, J. E., Zanon V., Pastore C., Reyes Hardy M-P., Di Maio LS., Gabellini P., Martin-Lorenzo A., Rodriguez F., and Perez, N. M.: Tephra sedimentation and grainsize associated with pulsatory activity: the 2021 Tajogaite eruption of Cumbre Vieja (La Palma, Canary Islands, Spain). *Front. Earth Sci.* 11:1166073, <https://doi.org/10.3389/feart.2023.1166073>, 2023.
- Bryukhanov, A. L., Sevastyanov, V. S., Kravchishina, M. D., Voropaev, S. A., Dushenko, N. V., Kurakov, A. V., and Fedulova, V. Y.: Composition of methane cycle microbial communities in the upper layers of bottom sediments of the Kara sea, *Geochem. Int.*, 62, 609–617, <https://doi.org/10.1134/S0016702924700277>, 2024.
- Bull, I. D., Nott, C. J., Van Bergen, P. F., Poulton, P. R., and Evershed, R. P.: Organic geochemical studies of soils from the Rothamsted classical experiments VI. The occurrence and source of organic acids in an experimental grassland soil, *Soil Biol. Biochem.*, 32, 1367–1376, 2000.



- Cabianca, A., Müller, L., Pawlowski, K., and Dahlin, P.: Changes in the plant β -sitosterol/stigmasterol ratio caused by the plant parasitic nematode meloidogyne incognita, *Plants*, 10, 1–15, <https://doi.org/10.3390/plants10020292>, 2021.
- 590 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., and Holmes, S. P.: DADA2: High-resolution sample inference from Illumina amplicon data, *Nat. Methods*, 13, 581–583, <https://doi.org/10.1038/nmeth.3869>, 2016.
- Campeny, M., Menéndez, I., Ibáñez-Insa, J., Rivera-Martínez, J., Yepes, J., Álvarez-Pousa, S., Méndez-Ramos, J., and Mangas, J.: The ephemeral fumarolic mineralization of the 2021 Tajogaite volcanic eruption (La Palma, Canary Islands, Spain). *Sci. Rep.*, 13, 6336, <https://doi.org/10.1038/s41598-023-33387-6>, 2023.
- 595 Capaccioni, B., Martini, M., and Mangani, F.: Light hydrocarbons in hydrothermal and magmatic fumaroles: hints of catalytic and thermal reactions, *B. Volcanol.*, 56, 593-600, 1995.
- Carracedo, J.C., Badiola, E.R., Guillou, H., de la Nuez, J., and Pérez-Torrado, F.J.: Geology and volcanology of la Palma and El Hierro. *Estud. Geol.*, 57, 175-273, 2001.
- Carracedo, J. C., Troll, V. R., Day, J. M. D., Geiger, H., Aulinas, M., Soler, V., Deegan, F. M., Perez-Torrado, F. J., Gisbert, 600 G., Gazel, E., Rodriguez-Gonzalez, A., and Albert, H.: The 2021 eruption of the Cumbre Vieja volcanic ridge on La Palma, Canary Islands, *Geology Today*, 38, 94–107, <https://doi.org/10.1111/gto.12388>, 2022.
- Carrell, A. A., Veličković, D., Lawrence, T. J., Bowen, B. P., Louie, K. B., Carper, D. L., Chu, R. K., Mitchell, H. D., Orr, G., Markillie, L. M., Jawdy, S. S., Grimwood, J., Shaw, A. J., Schmutz, J., Northen, T. R., Anderton, C. R., Pelletier, D. A., and Weston, D. J.: Novel metabolic interactions and environmental conditions mediate the boreal peatmoss-cyanobacteria 605 mutualism, *ISME J.*, 16, 1074–1085, <https://doi.org/10.1038/s41396-021-01136-0>, 2022.
- Chaneva, G., Furnadzhieva, S., Minkova, K., and Lukavsky, J.: Effect of light and temperature on the cyanobacterium *Arthronema africanum* - A prospective phycobiliprotein-producing strain, *J. Appl. Phycol.*, 19, 537–544, <https://doi.org/10.1007/s10811-007-9167-6>, 2007.
- Chen, J., Zheng, Y., Guo, Y., Li, F., Xu, D., Chao, L., Qu, H., Wang, B., Ma, X., Wang, S., and Bao, Y.: Differences in 610 microbial communities from Quaternary volcanic soils at different stages of development: Evidence from Late Pleistocene and Holocene volcanoes, *Catena (Amst)*, 201, <https://doi.org/10.1016/j.catena.2021.105211>, 2021.
- Craig, H., Wilson, T., Stewart, C., Outes, V., Villarosa, G., and Baxter, P.: Impacts to agriculture and critical infrastructure in Argentina after ashfall from the 2011 eruption of the Cordon Caulle volcanic complex: An assessment of published damage and function thresholds, *J. Appl. Volcanol.*, 5, 7, <https://doi.org/10.1186/s13617-016-0046-1>, 2016.
- 615 Crowe, M. A., Power, J. F., Morgan, X. C., Dunfield, P. F., Lagutin, K., Rijpstra, W. I. C., Rijpstra, I. C., Sinnighe Damste, J. S., Houghton, K. M., Ryan, J. L. J., and Stott, M. B.: *Pyrinomonas methylaliphatogetes* gen. nov., sp. nov., a novel group 4



- thermophilic member of the phylum Acidobacteria from geothermal soils, *Int. J. Syst. Evol. Micr.*, 64, 220–227, <https://doi.org/10.1099/ijs.0.055079-0>, 2014.
- Das, S. K.: *Bosea*, in: *Bergey's Manual of Systematics of Archaea and Bacteria*, edited by Whitman, W. B., Wiley, J., and Sons, Ltd., Hoboken, N.J., USA, 1–4, <https://doi.org/10.1002/9781118960608.gbm00801>, 2015.
- De la Rosa, J. M., Jiménez-Morillo, N. T., González-Pérez, J. A., Almendros, G., Vieira, D., Knicker, H. E., and Keizer, J.: Mulching-induced preservation of soil organic matter quality in a burnt eucalypt plantation in central Portugal, *J. Environ. Manage.*, 231, 1135–1144, <https://doi.org/10.1016/j.jenvman.2018.10.114>, 2019.
- De la Rosa, J. M., Pérez-Dalí, S. M., Campos, P., Sánchez-Martín, Á., González-Pérez, J. A., and Miller, A. Z.: Suitability of volcanic ash, rice husk ash, green compost and biochar as amendments for a Mediterranean alkaline soil, *Agronomy*, 13, 4, <https://doi.org/10.3390/agronomy13041097>, 2023.
- Dhabalia Ashok, A., De Vries, S., Darienko, T., Irisarri, I., and De Vries, J.: Evolutionary assembly of the plant terrestrialization toolkit from protein domains, *P. Roy. Soc. B-Biol. Sci.*, 291, <https://doi.org/10.1098/rspb.2024.0985>, 2024.
- D'haeseleer, P., Gladden, J. M., Allgaier, M., Chain, P. S. G., Tringe, S. G., Malfatti, S. A., Aldrich, J. T., Nicora, C. D., Robinson, E. W., Paša-Tolić, L., Hugenholtz, P., Simmons, B. A., and Singer, S. W.: Proteogenomic analysis of a thermophilic bacterial consortium adapted to deconstruct switchgrass, *PLOS One*, 8, <https://doi.org/10.1371/journal.pone.0068465>, 2013.
- Dragone, N. B., Whittaker, K., Lord, O. M., Burke, E. A., Dufel, H., Hite, E., Miller, F., Page, G., and Slayback, D.: The early microbial colonizers of a short-lived volcanic island in the Kingdom of Tonga, *mBio*, 14, 1, <https://doi.org/10.1128/mbio.03313-22>, 2023.
- Ehleringer, J. R., Buchmann, N., and Flanagan, L. B.: Carbon isotope ratios in belowground carbon cycle processes, *Ecol. Appl.*, 10, 412–422, <https://doi.org/10.2307/2641103>, 2000.
- Falagán, C. and Johnson, D. B.: *Acidibacter ferrireducens* gen. nov., sp. nov.: an acidophilic ferric iron-reducing gammaproteobacterium. *Extremophiles*, 18, 1067–1073, <https://doi.org/10.1007/s00792-014-0684-3>, 2014.
- Feakins, S. J., Peters, T., Wu, M. S., Shenkin, A., Salinas, N., Girardin, C. A. J., Bentley, L. P., Blonder, B., Enquist, B. J., Martin, R.E., Asner, G. P., and Malhi, Y.: Production of leaf wax n-alkanes across a tropical forest elevation transect. *Org. Geochem.*, 100, 89–100, <https://doi.org/10.1016/j.orggeochem.2016.07.004>, 2016.
- Ferrer, N., Marrero-Rodríguez, N., Sanromualdo-Collado, A., Vegas, J., and García-Romero, L.: Early morphodynamics of the sudden formation of beaches during the 2021 volcanic eruption of La Palma, *Geomorphology*, 436, <https://doi.org/10.1016/j.geomorph.2023.108779>, 2023.
- Fraser, C. M. and Chapple, C.: The phenylpropanoid pathway in *Arabidopsis*, *Arabidopsis Book*, 9, e0152, <https://doi.org/10.1199/tab.0152>, 2011.



- Fujimura, R., Kim, S. W., Sato, Y., Oshima, K., Hattori, M., Kamijo, T., and Ohta, H.: Unique pioneer microbial communities exposed to volcanic sulfur dioxide, *Sci. Rep.*, 6, 19687, <https://doi.org/10.1038/srep19687>, 2016.
- García-Carmona, M., García-Orenes, F., Arcenegui, V., and Mataix-Solera, J.: The recovery of mediterranean soils after post-
650 fire management: The role of biocrusts and soil microbial communities, *Span. J. Soil Sci.*, 13, <https://doi.org/10.3389/sjss.2023.11388>, 2023.
- Gardeva, E. G., Toshkova, R. A., Yossifova, L. S., Minkova, K., Ivanova, N. Y., and Gigova, L. G.: Antitumor activity of C-phycocyanin from *Arthonema africanum* (Cyanophyceae), *Braz. Arch. Biol. Techn.*, 57, 675–684, <https://doi.org/10.1590/S1516-89132014005000018>, 2014.
- 655 Ghezzi, D., Jiménez-Morillo, N. T., Foschi, L., Donini, E., Chiarini, V., De Waele, J., Miller, A. Z., and Cappelletti, M.: The microbiota characterizing huge carbonatic moonmilk structures and its correlation with preserved organic matter, *Environ. Microbiome*, 19, 25, <https://doi.org/10.1186/s40793-024-00562-9>, 2024.
- Girona-García, A., Badía-Villas, D., Jiménez-Morillo, N. T., de la Rosa, J. M., and González-Pérez, J. A.: Soil C and N isotope composition after a centennial Scots pine afforestation in podzols of native European beech forests in NE-Spain, *Catena*
660 (Amst), 165, 434–441, <https://doi.org/10.1016/j.catena.2018.02.023>, 2018.
- González-Pérez, J. A., González-Vila, F. J., Almendros, G., and Knicker, H.: The effect of fire on soil organic matter - A review, *Environ. Int.*, 30, 855-870, <https://doi.org/10.1016/j.envint.2004.02.003>, 2004.
- González-Pérez, J. A., Almendros, G., De La Rosa, J. M., and González-Vila, F. J.: Appraisal of polycyclic aromatic hydrocarbons (PAHs) in environmental matrices by analytical pyrolysis (Py-GC/MS), *J. Anal. Appl. Pyrol.*, 109,1-8,
665 <https://doi.org/10.1016/j.jaap.2014.07.005>, 2014.
- González-Pérez, J. A., Jiménez-Morillo, N. T., de la Rosa, J. M., Almendros, G., and González-Vila, F. J.: Compound-specific stable carbon isotopic signature of carbohydrate pyrolysis products from C3 and C4 plants, *J. Sci. Food. Agric.*, 96, 948–953, <https://doi.org/10.1002/jsfa.7169>, 2016.
- Gorlenko, V. M., Bryantseva, I. A., Kalashnikov, A. M., Gaisin, V. A., Sukhacheva, M. V., Gruzdev, D. S., and Kuznetsov,
670 B. B.: *Candidatus 'Chloroploca asiatica'* gen. nov., sp. nov., a new mesophilic filamentous anoxygenic phototrophic bacterium, *Microbiology (Russian Federation)*, 83, 838–848, <https://doi.org/10.1134/S0026261714060083>, 2014.
- Guarro, J., Abdullah, S. K., Al-Bader, S. M., Figueras, M. J., and Gené, J.: The genus *Melanocarpus*, *Mycol. Res.*, 100, 75–78, [https://doi.org/10.1016/S0953-7562\(96\)80103-1](https://doi.org/10.1016/S0953-7562(96)80103-1), 1996.
- Gutierrez-Patricio, S., Osman, J. R., Luis Gonzalez-Pimentel, J., Jurado, V., Laiz, L., Laínez Concepción, A., Saiz-Jimenez, C., and Miller, A. Z.: Microbiological exploration of the Cueva del Viento lava tube system in Tenerife, Canary Islands, *Env. Microbiol. Rep.*, 16, e13245, <https://doi.org/10.1111/1758-2229.13245>, 2024.



- Habib, N., Khan, I. U., Hussain, F., Zhou, E. M., Xiao, M., Ahmed, I., Zhi, X. Y., and Li, W. J.: *Caldovatus sediminis* gen. nov., sp. nov., a moderately thermophilic bacterium isolated from a hot spring, *Int. J. Syst. Evol. Microbiol.*, 67, 4716–4721, <https://doi.org/10.1099/ijsem.0.002363>, 2017.
- 680 Hadland, N., Hamilton, C. W., and Duhamel, S.: Young volcanic terrains are windows into early microbial colonization, *Commun. Earth Environ.*, 5, 114, <https://doi.org/10.1038/s43247-024-01280-3>, 2024.
- Hernández, Z., Almendros, G., Carral, P., Álvarez, A., Knicker, H., and Pérez-Trujillo, J. P.: Influence of non-crystalline minerals in the total amount, resilience and molecular composition of the organic matter in volcanic ash soils (Tenerife Island, Spain), *Eur. J. Soil Sci.*, 63, 603–615, <https://doi.org/10.1111/j.1365-2389.2012.01497.x>, 2012.
- 685 Hoernle, K. A. J., Carracedo, J. C., Canary Islands, geology. University of California Press, 2009.
- Högberg, P.: Tansley review no. 95 natural abundance in soil-plant systems, *New Phytol.*, 137, 179–203, <https://doi.org/10.1046/j.1469-8137.1997.00808.x>, 1997.
- Hu, C. J., Xian, W. D., Lv, Y. Q., Peng, C. X., Shan, R. X., Cheng, Z. C., Lv, Q., Tian, Y., Jiao, J. Y., Tan, S., and Li, W. J.: *Caldovatus aquaticus* sp. nov., a moderately thermophilic bacterium isolated from hot spring microbial mat. *Int. J. Syst. Evol. Microbiol.*, 72, 5627, <https://doi.org/10.1099/ijsem.0.005627>, 2022.
- 690 Iliev, I. P. G. D. and Andreeva, R.: Membrane metabolites of *Arthronema africanum* strains from extreme habitats, *Gen. Appli. Plant. Physiology*, 2006, 117–123, 2006.
- Ingimundardóttir, G. V., Weibull, H., and Cronberg, N.: Bryophyte colonization history of the virgin volcanic Island Surtsey, Iceland, *Biogeosciences*, 11, 4415–4427, <https://doi.org/10.5194/bg-11-4415-2014>, 2014.
- 695 Iwasaki, S., Endo, Y., Hatano, R., and Toma, Y.: Factors impacting soil organic carbon pool in different types of Andosols in Toya, Hokkaido, Japan, *Soil Sci. Plant. Nutr.*, 67, 594–605, <https://doi.org/10.1080/00380768.2021.1959836>, 2021.
- Jaffé, R., Elismé, T., and Cabrera, A. C.: Organic geochemistry of seasonally flooded rain forest soils: molecular composition and early diagenesis of lipid components, *Org. Geochem.*, 25, 9–17, 1996.
- 700 Jiménez-Morillo, N. T., de la Rosa, J. M., Waggoner, D., Almendros, G., González-Vila, F. J., and González-Pérez, J. A.: Fire effects in the molecular structure of soil organic matter fractions under *Quercus suber* cover, *Catena*, 145, 266–273, <https://doi.org/10.1016/j.catena.2016.06.022>, 2016a.
- Jiménez-Morillo, N. T., González-Pérez, J. A., Jordán, A., Zavala, L. M., de la Rosa, J. M., Jiménez-González, M. A., and González-Vila, F. J.: Organic matter fractions controlling soil water repellency in sandy soils from the Doñana national park (Southwestern Spain), *Land. Degrad. Dev.*, 27, 1413–1423, <https://doi.org/10.1002/ldr.2314>, 2016b.



- 705 Jiménez-Morillo, N. T., Spangenberg, J. E., Miller, A. Z., Jordán, A., Zavala, L. M., González-Vila, F. J., and González-Pérez, J. A.: Wildfire effects on lipid composition and hydrophobicity of bulk soil and soil size fractions under *Quercus suber* cover (SW-Spain), *Environ. Res.*, 159, 394–405, <https://doi.org/10.1016/j.envres.2017.08.022>, 2017.
- Jiménez-Morillo, N. T., González-Pérez, J. A., Almendros, G., De la Rosa, J. M., Waggoner, D. C., Jordán, A., Zavala, L. M., González-Vila, F. J., and Hatcher, P. G.: Ultra-high resolution mass spectrometry of physical speciation patterns of organic matter in fire-affected soils, *J. Environ. Manage.*, 225, 139–147, <https://doi.org/10.1016/j.jenvman.2018.07.069>, 2018.
- 710 Jiménez-Morillo, N. T., Almendros, G., De la Rosa, J. M., Jordán, A., Zavala, L. M., Granged, A. J. P., and González-Pérez, J. A.: Effect of a wildfire and of post-fire restoration actions in the organic matter structure in soil fractions, *Sci. Total Environ.*, 728, 138715, <https://doi.org/10.1016/j.scitotenv.2020.138715>, 2020a.
- Jiménez-Morillo, N. T., Almendros, G., González-Vila, F. J., Jordán, A., Zavala, L. M., de la Rosa, J. M., and González-Pérez, J. A.: Fire effects on C and H isotopic composition in plant biomass and soil: Bulk and particle size fractions, *Sci. Total Environ.*, 749, 141417, <https://doi.org/10.1016/j.scitotenv.2020.141417>, 2020b.
- 715 Jiménez-Morillo, N. T., Almendros, G., Miller, A. Z., Hatcher, P. G., and González-Pérez, J. A.: Hydrophobicity of soils affected by fires: An assessment using molecular markers from ultra-high resolution mass spectrometry, *Sci. Total Environ.*, 817, 152957, <https://doi.org/10.1016/j.scitotenv.2022.152957>, 2022.
- 720 Katsumi, N., Yonebayashi, K., Fujitake, N., and Okazaki, M.: Relationship between stable carbon and nitrogen isotope ratios of humic acids extracted from Andisols and non-Andisols, *Catena (Amst)*, 127, 214–221, <https://doi.org/10.1016/j.catena.2015.01.005>, 2015.
- Kellner, H., Luis, P., Pecyna, M. J., Barbi, F., Kapturska, D., Kružer, D., Zak, D. R., Marmeisse, R., Vandenbol, M., and Hofrichter, M.: Widespread occurrence of expressed fungal secretory peroxidases in forest soils, *PLOS One*, 9, e95557, <https://doi.org/10.1371/journal.pone.0095557>, 2014.
- 725 King, C. E. and King, G. M.: Description of *Thermogemmatispora carboxidivorans* sp. nov., a carbon-monoxideoxidizing member of the class Ktedonobacteria isolated from a geothermally heated biofilm, and analysis of carbon monoxide oxidation by members of the class Ktedonobacteria, *Int. J. Syst. Evol. Microbiol.*, 64, 1244–1251, <https://doi.org/10.1099/ijs.0.059675-0>, 2014.
- 730 Klavina, L., Springe, G., Nikolajeva, V., Martsinkevich, I., Nakurte, I., Dzabijeva, D., and Steinberga, I.: Molecules chemical composition analysis, antimicrobial activity and cytotoxicity screening of moss extracts (Moss Phytochemistry), *Molecules*, 20, 17221–17243, <https://doi.org/10.3390/molecules200917221>, 2015.
- Klegin, C., Ethur, E. M., Bordin, J., and Baia Figueiredo, P. L.: Chemical composition of essential oil from mosses from the brazilian atlantic forest, *Chem. Biodivers.*, 20, e202300286, <https://doi.org/10.1002/cbdv.202300286>, 2023.



- 735 Krauss, M., Wilcke, W., Martius, C., Bandeira, A. G., Garcia, M. V. B., Amelung, W.: Atmospheric versus biological sources of polycyclic aromatic hydrocarbons (PAHs) in a tropical rain forest environment. *Environ. Pollut.*, 135, 143-154, <https://doi.org/10.1016/j.envpol.2004.09.012>, 2005.
- Kristan, K. and Rižner, T. L.: Steroid-transforming enzymes in fungi, *J. Steroid. Biochem. Mol. Biol.*, 129, 79-91, <https://doi.org/10.1016/j.jsbmb.2011.08.012>, 2012.
- 740 Krüger, N., Finn, D. R., and Don, A.: Soil depth gradients of organic carbon-13 – A review on drivers and processes, *Plant Soil*, 495, 113-136, <https://doi.org/10.1007/s11104-023-06328-5>, 2024.
- Kumar, M., Boski, T., González-Vila, F. J., Jiménez-Morillo, N. T., and González-Pérez, J. A.: Characteristics of organic matter sources from Guadiana Estuary salt marsh sediments (SW Iberian Peninsula), *Cont. Shelf Res.*, 197, 104076, <https://doi.org/10.1016/j.csr.2020.104076>, 2020.
- 745 Ladygina, N., Dedyukhina, E. G., and Vainshtein, M. B.: A review on microbial synthesis of hydrocarbons. *Process Biochem.*, 41, 1001–1014. <https://doi.org/10.1016/j.procbio.2005.12.007>, 2006.
- Leal, O. dos A., Jiménez-Morillo, N. T., González-Pérez, J. A., Knicker, H., de Souza Costa, F., Jiménez-Morillo, P. N., de Carvalho Júnior, J. A., dos Santos, J. C., and Pinheiro Dick, D.: Soil organic matter molecular composition shifts driven by forest regrowth or pasture after slash-and-burn of amazon forest, *Int. J. Environ. Res. Public Health*, 20, 3485, <https://doi.org/10.3390/ijerph20043485>, 2023.
- 750 Lehtonen, K. and Ketola, M.: Occurrence of long-chain acyclic methyl ketones in *Sphagnum* and *Carex* peats of various degrees of humification, *Org. Geochem.*, 15, 275–280, [https://doi.org/10.1016/0146-6380\(90\)90005-K](https://doi.org/10.1016/0146-6380(90)90005-K), 1990.
- Lehtovirta-Morley, L. E., Ge, C., Ross, J., Yao, H., Hazard, C., Gubry-Rangin, C., Prosser, J. I., and Nicol, G. W.: *Nitrosotalea devaniterrae* gen. nov., sp. nov. and *Nitrosotalea sinensis* sp. nov., two acidophilic ammonia oxidising archaea isolated from acidic soil, and proposal of the new order Nitrosotaleales ord. nov. within the class Nitrososphaeria of the phylum Nitrososphaerota, *Int. J. Syst. Evol. Microbiol.*, 74, <https://doi.org/10.1099/ijsem.0.006387>, 2024.
- 755 Leininger, S., Urich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G. W., Prosser, J. I., Schuster, S. C., and Schleper, and C.: Archaea predominate among ammonia-oxidizing prokaryotes in soils, *Nature*, 442, 806-809, <https://doi.org/10.1038/nature04983>, 2006.
- 760 Longpré, M.-A., Felpeto, A.: Historical volcanism in the Canary Islands; part 1: A review of precursory and eruptive activity, eruption parameter estimates, and implications for hazard assessment, *J. Volcanol. Geoth. Res.*, 419, 107363, <https://doi.org/10.1016/j.jvolgeores.2021.107363>, 2021.



- 765 Love, C. R., Arrington, E. C., Gosselin, K. M., Reddy, C. M., Van Mooy, B. A. S., Nelson, R. K., and Valentine, D. L.:
Microbial production and consumption of hydrocarbons in the global ocean. *Nat. Microbiol.* 6, 489–498,
<https://doi.org/10.1038/s41564-020-00859-8>, 2021.
- Nanzyo, M., Dahlgren, R. and Shoji, S.: Chemical characteristics of volcanic ash soils. In: Shoji, S., Nanzyo, M. and Dahlgren,
R., edited by *Volcanic Ash Soils—Genesis, Properties and Utilization, Developments in Soil Science*, Elsevier, Amsterdam,
145-188, [http://dx.doi.org/10.1016/S0166-2481\(08\)70267-8](http://dx.doi.org/10.1016/S0166-2481(08)70267-8), 1993.
- Makarov, M. I.: The nitrogen isotopic composition in soils and plants: Its use in environmental studies (A Review), *Soil*
770 *Chemistry*, 42, 1335-1347, <https://doi.org/10.1134/S1064229309120035>, 2009.
- Martínez-Espinosa, R. M.: Microorganisms and their metabolic capabilities in the context of the biogeochemical nitrogen cycle
at extreme environments, *Int. J. Mol. Sci.*, 21, 4228, <https://doi.org/10.3390/ijms21124228>, 2020.
- Martínez-Martínez, J., Mediato, J. F., Mata, M. P., Ordóñez, B., del Moral, B., Bellido, E., Pérez-López, R., Rodríguez-Pascua,
775 M. A., Vegas, J., Lozano Otero, G., Mateos, R. M., Sánchez, N., and Galindo, I.: Early fumarolic minerals from the Tajogaite
volcanic eruption (La Palma, 2021), *J. Volcanol. Geotherm. Res.*, 435, 107771,
<https://doi.org/10.1016/j.jvolgeores.2023.107771>, 2023.
- Matus, F., Rumpel, C., Neculman, R., Panichini, M., and Mora, M. L.: Soil carbon storage and stabilisation in andic soils: A
review, *Catena*, 120, 102-110, <https://doi.org/10.1016/j.catena.2014.04.008>, 2014.
- 780 Miller, A. Z., De la Rosa, J. M., Jiménez-Morillo, N. T., Pereira, M. F. C., González-Pérez, J. A., Calaforra, J. M., Saiz-
Jimenez, C.: (2016). Analytical pyrolysis and light stable isotope analyses reveal environmental changes in coralloid
speleothems from Easter Island (Chile). *J. Chromatogr. A.* 1461, 144–152, <https://doi.org/10.1016/j.chroma.2016.07.038>,
2016.
- 785 Miller, A. Z., De la Rosa, J. M., Jiménez-Morillo, N. T., Pereira, M. F. C., Gonzalez-Perez, J. A., Knicker, H., and Saiz-
Jimenez, C.: Impact of wildfires on subsurface volcanic environments: New insights into speleothem chemistry, *Science of*
the Total Environment, 698, <https://doi.org/10.1016/j.scitotenv.2019.134321>, 2020.
- Miller, A. Z., Jiménez-Morillo, N. T., Coutinho, M. L., Gazquez, F., Palma, V., Sauro, F., Pereira, M. F. C., Rull, F.,
Toulkeridis, T., Caldeira, A. T., Forti, P., and Calaforra, J. M.: Organic geochemistry and mineralogy suggest anthropogenic
impact in speleothem chemistry from volcanic show caves of the Galapagos, *iScience*, 25, 104556,
<https://doi.org/10.1016/j.isci.2022.104556>, 2022.
- 790 Mohapatra, B. and Phale, P. S.: Microbial degradation of naphthalene and substituted naphthalenes: Metabolic diversity and
genomic insight for bioremediation, *Front. Bioeng. Biotechnol.*, 9, <https://doi.org/10.3389/fbioe.2021.602445>, 2021.



- Muneer, M. A., Huang, X., Hou, W., Zhang, Y., Cai, Y., Munir, M. Z., Wu, L., and Zheng, C.: Response of fungal diversity, community composition, and functions to nutrients management in red soil, *J. Fungi*, 7, 554, <https://doi.org/10.3390/jof7070554>, 2021.
- 795 Muñoz, G., Orlando, J., and Zuñiga-Feest, A.: Plants colonizing volcanic deposits: root adaptations and effects on rhizosphere microorganisms, *Plant Soil*, 461, 265-279, <https://doi.org/10.1007/s11104-020-04783-y>, 2021.
- Musat, F., Galushko, A., Jacob, J., Widdel, F., Kube, M., Reinhardt, R., Wilkes, H., Schink, B., and Rabus, R.: Anaerobic degradation of naphthalene and 2-methylnaphthalene by strains of marine sulfate-reducing bacteria, *Environ. Microbiol.*, 11, 209–219, <https://doi.org/10.1111/j.1462-2920.2008.01756.x>, 2009.
- 800 Nierop, K. G. J. and Buurman, P.: Thermally assisted hydrolysis and methylation of organic matter in two allophanic volcanic ash soils from the Azores Islands, in: *Soils of Volcanic Regions in Europe*, edited by: Arnalds, O., óskarsson, H., Bartolini, F., Buurman, P., Stoops, G., García-Rodeja, E. Springer, Berlin, Heidelberg, https://doi.org/10.1007/978-3-540-48711-1_30, 2007.
- Offre, P., Spang, A., and Schleper, C.: Archaea in biogeochemical cycles, *Annu. Rev. Microbiol.*, 67, 437–457, 805 <https://doi.org/10.1146/annurev-micro-092412-155614>, 2013.
- Palma, V., De la Rosa, J. M., Onac, B. P., Sauro, F., Martínez-Frías, J., Caldeira, A. T., González-Pérez, J. A., Jiménez-Morillo, N. T., and Miller, A. Z.: Decoding organic compounds in lava tube sulfates to understand potential biomarkers in the Martian subsurface, *Commun. Earth Environ.*, 5, 530, <https://doi.org/10.1038/s43247-024-01673-4>, 2024.
- Peng, X., Tamura, K., Asano, M., Takano, A., Kawagoe, M., and Kamijo, T.: Changes in soil physical and chemical properties 810 during vegetation succession on miyake-jima island, *Forests*, 12, 1435, <https://doi.org/10.3390/f12111435>, 2021.
- Petkova, Z., Teneva, O., Antova, G., Angelova-Romova, M., Gecheva, G., and Dimitrova-Dyulgerova, I.: Chemical composition, lipid-soluble bioactive compounds and potential health benefits of the moss *Hypnum cupressiforme* Hedw, *Plants*, 12, 4190, <https://doi.org/10.3390/plants12244190>, 2023.
- Petrova, D. H., Yocheva, L., and Petrova, M.: Antimicrobial and antioxidant activities of microalgal extracts, *Oxid. Commun.*, 815 43, 103–116, <https://doi.org/10.13140/RG.2.2.20509.84963>, 2020.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools, *Nucleic Acids Res.*, 41, D590-D596, <https://doi.org/10.1093/nar/gks1219>, 2013.
- Rabbi, S. M. F., Wilson, B. R., Lockwood, P. V., Daniel, H., and Young, I. M.: Soil organic carbon mineralization rates in 820 aggregates under contrasting land uses, *Geoderma*, 216, 10–18, <https://doi.org/10.1016/j.geoderma.2013.10.023>, 2014.



- Ramos, P., Honda, R., Hoek, E. M. V., and Mahendra, S.: Carbon/nitrogen ratios determine biofilm formation and characteristics in model microbial cultures, *Chemosphere*, 313, <https://doi.org/10.1016/j.chemosphere.2022.137628>, 2023.
- Ren, G., Zhang, Y., Chen, Z., Xue, X., and Fan, H.: Research progress of small plant peptides on the regulation of plant growth, development, and abiotic stress, *Int. J. Mol. Sci.*, 25, 4114, <https://doi.org/10.3390/ijms25074114>, 2024.
- 825 Renault, H., Alber, A., Horst, N. A., Lopes, A. B., Fich, E. A., Kriegshauser, L., Wiedemann, G., Ullmann, P., Herrgott, L., Erhardt, M., Pineau, E., Ehlting, J., Schmitt, M., Rose, J. K. C., Reski, R., and Werck-Reichhart, D.: A phenol-enriched cuticle is ancestral to lignin evolution in land plants, *Nat. Commun.*, 8, 14713, <https://doi.org/10.1038/ncomms14713>, 2017.
- Rippin, M., Borchhardt, N., Williams, L., Colesie, C., Jung, P., Büdel, B., Karsten, U., and Becker, B.: Genus richness of microalgae and Cyanobacteria in biological soil crusts from Svalbard and Livingston Island: morphological versus molecular approaches, *Polar Biol.*, 41, 909–923, <https://doi.org/10.1007/s00300-018-2252-2>, 2018.
- 830 Roderic Park, S. E.: Metabolic fractionation of C13 and C12 in plants, *Plant Physiol.*, 36, 133, 1961.
- San-Emeterio, L. M., Zavala, L. M., Jiménez-Morillo, N. T., Pérez-Ramos, I. M., and González-Pérez, J. A.: Effects of climate change on soil organic matter C and H isotope composition in a Mediterranean Savannah (Dehesa): An assessment using Py-CSIA, *Environ. Sci. Technol.*, 57, 13851–13862, <https://doi.org/10.1021/acs.est.3c01816>, 2023.
- 835 Sánchez-España, J., Mata, M. P., Vegas, J., Lozano, G., Mediato, J., Martínez Martínez, J., Galindo, I., Sánchez, N., del Moral, B., Ordóñez, B., de Vergara, A., Nieto, A., Andrés, M., Vázquez, I., Bellido, E., and Castillo-Carrión, M.: Leaching tests reveal fast aluminum fluoride release from ashfall accumulated in La Palma (Canary Islands, Spain) after the 2021 Tajogaite eruption. *J. Volcanol. Geoth. Res.*, 444, 107959, <https://doi.org/10.1016/j.jvolgeores.2023.107959>, 2023.
- Schwark, L., Zink, K., and Lechterbeck, J.: Reconstruction of postglacial to early Holocene vegetation history in terrestrial Central Europe via cuticular lipid biomarkers and pollen records from lake sediments, *Geology*, 30, 463–466, [https://doi.org/10.1130/0091-7613\(2002\)030<0463:ROPTEH>2.0.CO;2](https://doi.org/10.1130/0091-7613(2002)030<0463:ROPTEH>2.0.CO;2), 2002.
- 840 La Scola, B., Barrassi, L., and Raoult, D.: A novel alpha-Proteobacterium, *Nordella oligomobilis* gen. nov., sp. nov., isolated by using amoebal co-cultures, *Res. Microbiol.*, 155, 47–51, <https://doi.org/10.1016/j.resmic.2003.09.012>, 2004.
- Smith, R. J. L.: Colonization by bryophytes following recent volcanic activity on an Antarctic island, *J. Hattori Bot. Lab.*, 56, 845 53–63, 1984.
- Survey Staff, S.: Keys to Soil Taxonomy, 13th ed., edited by: USDA Natural Resources Conservation Service, 2022.
- Szpak, P.: Complexities of nitrogen isotope biogeochemistry in plant-soil systems: Implications for the study of ancient agricultural and animal management practices, *Front. Plant Sci.*, 5, <https://doi.org/10.3389/fpls.2014.00288>, 2014.



- Taddeucci, J., Scarlato, P., Andronico, D., Ricci, T., Civico, R., Del Bello, E., Spina, L., D'Auria, L., Asensio-Ramos, M.,
850 Calvo, D., Padrón, E., Hernández, P. A., and Pérez, N. M.: The explosive activity of the 2021 Tajogaite Eruption (La Palma, Canary Islands, Spain), *Geochem., Geophys., Geosyst.*, 24, <https://doi.org/10.1029/2023GC010946>, 2023.
- Tomašek, I., Damby, D. E., Andronico, D., Baxter, P. J., Boonen, I., Claeys, P., Denison, M. S., Horwell, C. J., Kervyn, M., Kueppers, U., Romanias, M. N., and Elskens, M.: Assessing the biological reactivity of organic compounds on volcanic ash: implications for human health hazard, *Bull. Volcanol.*, 83, <https://doi.org/10.1007/s00445-021-01453-4>, 2021.
- 855 Troll, V. R., Aulinas, M., Carracedo, J. C., Geiger, H., Perez-Torrado, F. J., V. Soler, Deegan, F. M., Bloszies, C., Weis, F., Albert, H., Gisbert, G., Day, J. M. D., Rodríguez-Gonzalez, A., Gazel, E., and Dayton, K.: The 2021 La Palma eruption: social dilemmas resulting from life close to an active volcano. *Geol. Today*, 40, 96-111, <https://doi.org/10.1111/gto.12472>, 2024.
- Tseng, H. C., Matsutani, M., Fujimoto, N., and Ohnishi, A.: Draft genome sequence and morphological data of *Planifilum fimeticola* PLACP1, a thermophilic chloramphenicol-resistant bacterium isolated from thermophilic sludge, *Data Brief*, 54,
860 <https://doi.org/10.1016/j.dib.2024.110447>, 2024.
- Ustiatik, R., Ariska, A. P., Hakim, Q. L., Wicaksono, K. S., and Utami, S. R.: Volcanic deposits thickness and distance from mt Semeru crater strongly affected phosphate solubilizing bacteria population and soil organic carbon, *J. Ecol. Eng.*, 24, 360–368, <https://doi.org/10.12911/22998993/170860>, 2023.
- Valeeva, L. R., Dague, A. L., Hall, M. H., Tikhonova, A. E., Sharipova, M. R., Valentovic, M. A., Bogomolnaya, L. M., and
865 Shakirov, E. V.: Antimicrobial activities of secondary metabolites from model mosses, *Antibiotics*, 11, 1004, <https://doi.org/10.3390/antibiotics11081004>, 2022.
- Valitova, J. N., Sulkarnayeva, A. G., and Minibayeva, F. V.: Plant sterols: Diversity, biosynthesis, and physiological functions, *Biochemistry (Moscow)*, 81, 819-834, <https://doi.org/10.1134/S0006297916080046>, 2016.
- Vilhelmsson, O., Sigurbjörnsdóttir, M. A., Thorsteinsdóttir, G. V., Cascone, M., Corso, D., Toniatti, L., Migliaccio, F., Nappi,
870 N., Ricciardelli, A., Selci, M., Montemagno, F., Barosa, B., Bastoni, D., Bastianoni, A., Cordone, A., and Giovannelli, D.: Diversity of thermophilic prokaryotes, in: *Thermophilic anaerobes. Grand Challenges in Biology and Biotechnology*, edited by: Scully S.M., 21–90, https://doi.org/10.1007/978-3-031-41720-7_2, 2023.
- Vilmundardóttir, O. K., Sigurmundsson, F. S., Møller Pedersen, G. B., Belart, J. M. C., Kizel, F., Falco, N., Benediktsson, J. A., and Gísladóttir, G.: Of mosses and men: Plant succession, soil development and soil carbon accretion in the sub-Arctic
875 volcanic landscape of Hekla, Iceland, *Prog. Phys. Geogr.*, 42, 765–791, <https://doi.org/10.1177/0309133318798754>, 2018.
- Vogts, A., Moossen, H., Rommerskirchen, F., and Rullkötter, J.: Distribution patterns and stable carbon isotopic composition of alkanes and alkan-1-ols from plant waxes of African rain forest and savanna C3 species. *Org. Geochem.*, 40, 1037-1054, <https://doi.org/10.1016/j.orggeochem.2009.07.011>, 2009.



- Volk, M., Bassin, S., Lehmann, M. F., Johnson, M. G., and Andersen, C. P.: ^{13}C isotopic signature and C concentration of soil density fractions illustrate reduced C allocation to subalpine grassland soil under high atmospheric N deposition, *Soil Biol. Biochem.*, 125, 178–184, <https://doi.org/10.1016/j.soilbio.2018.07.014>, 2018.
- 880 Wang, H., Wang, J., Xiao, M., Ge, T., Gunina, A., and Jones, D. L.: The fate of amino acid and peptide as affected by soil depth and fertilization regime in subtropical paddies, *Sci. Total Environ.*, 889, <https://doi.org/10.1016/j.scitotenv.2023.164245>, 2023.
- 885 Wakeham, S. G., Canuel, E. A. Biogenic polycyclic aromatic hydrocarbons in sediments of the San Joaquin River in California (USA), and current paradigms on their formation. *Environ. Sci. Pollut. Res.* 23, 10426–10442, <https://doi.org/10.1007/s11356-015-5402-x>, 2016.
- Wijesinghe, J. N., Koarashi, J., Atarashi-Andoh, M., Saito-Kokubu, Y., Yamaguchi, N., Sase, T., Hosono, M., Inoue, Y., Mori, Y., and Hiradate, S.: Formation and mobility of soil organic carbon in a buried humic horizon of a volcanic ash soil, *Geoderma*, 890 374, <https://doi.org/10.1016/j.geoderma.2020.114417>, 2020.
- Wygel, C. M., Peters, S. C., McDermott, J. M., and Sahagian, D. L.: Bubbles and Dust: Experimental results of dissolution rates of metal salts and glasses from volcanic ash deposits in terms of surface area, chemistry, and human health impacts, *Geohealth*, 3, 338–355, <https://doi.org/10.1029/2018GH000181>, 2019.
- Yakimov, M. M., Lünsdorf, H., and Golyshin, P. N.: *Thermoleophilum album* and *Thermoleophilum minutum* are culturable representations of group 2 of the Rubrobacteridae (Actinobacteria), *Int. J. Syst. Evol. Microbiol.*, 53, 377–380, <https://doi.org/10.1099/ijs.0.02425-0>, 2003.
- 895 Yang, Y. R., Guo, Y. X., Wang, Q. Y., Hu, B. Y., Tian, S. Y., Yang, Q. Z., Cheng, Z. A., Chen, Q. J., and Zhang, G. Q.: Impacts of composting duration on physicochemical properties and microbial communities during short-term composting for the substrate for oyster mushrooms, *Sci. Total Environ.*, 847, <https://doi.org/10.1016/j.scitotenv.2022.157673>, 2022.
- 900 Yokobe, T., Hyodo, F., and Tokuchi, N.: Volcanic deposits affect soil nitrogen dynamics and fungal–bacterial dominance in temperate forests, *Soil Biol. Biochem.*, 150, <https://doi.org/10.1016/j.soilbio.2020.108011>, 2020.
- Yucel, T. B.: Chemical composition and antimicrobial and antioxidant activities of essential oils of *Polytrichum commune* (Hedw.) and *Antitrichia curtispindula* (Hedw.) Brid. grown in Turkey, *Int. J. Secondary Metabolite*, 8, 272, <https://doi.org/10.21448/ijsm.945405>, 2021.
- 905 Zechmeister, H. G., Richter, A., Smidt, S., Hohenwallner, D., Roder, I., Maringer, S., and Wanek, W.: Total nitrogen content and $\delta^{15}\text{N}$ signatures in moss tissue: indicative value for nitrogen deposition patterns and source allocation on a nationwide scale, *Environ. Sci. Technol.*, 42, 8661–8667, <https://doi.org/10.1021/es801865d>, 2008.

<https://doi.org/10.5194/egusphere-2025-2086>

Preprint. Discussion started: 15 May 2025

© Author(s) 2025. CC BY 4.0 License.



Zeglin, L. H., Wang, B., Waythomas, C., Rainey, F., and Talbot, S. L.: Organic matter quantity and source affects microbial community structure and function following volcanic eruption on Kasatochi Island, Alaska, *Environ. Microbiol.*, 18, 146–158, 910 <https://doi.org/10.1111/1462-2920.12924>, 2016.

Zhang, J., Cai, L., Yuan, D., and Chen, M.: Distribution and sources of polynuclear aromatic hydrocarbons in Mangrove surficial sediments of Deep Bay, China, *Mar Pollut. Bull.*, 49, 479–486, <https://doi.org/10.1016/j.marpolbul.2004.02.030>, 2004.