

Distinct effects of several ice production processes on thunderstorm electrification and lightning activity

Inès Vongpaseut¹ and Christelle Barthe¹

¹LAERO, Université de Toulouse, CNRS, UT3, IRD, Toulouse, France

Correspondence: Inès Vongpaseut (ines.vongpaseut@univ-tlse3.fr)

Abstract. Ice particles play a crucial role in shaping cloud electrification, affecting the intensity of lightning activity. Previous studies have found a change of electric activity with varying aerosols concentration or active secondary ice production processes (SIP). However, the electric response to those parameters can differ with different cloud conditions and interact between themselves. The Meso-NH model was used with the two-moment microphysics scheme LIMA coupled with an explicit electrical scheme. Three idealized storms with varying warm-phase thicknesses were simulated to examine their response to aerosol concentrations and SIP mechanisms.

Increasing the cloud condensation nuclei (CCN) or the ice nucleating particle (INP) concentration increases ice crystal concentration, non-inductive charging and lightning activity up to a threshold. The main ice production processes (heterogeneous, homogeneous nucleation or Hallett-Mossop mechanism) depend on the cloud base temperature, and the aerosol concentration. CCN concentration thresholds ($1000\text{--}8000\text{ cm}^{-3}$) differ across all storms due to cloud base temperature, while the threshold for INP concentration is generally $\sim 100\text{ L}^{-1}$. Higher CCN concentrations increase cloud water content, affecting charge polarity, but ~~graupel mass has a smaller impact on electrification~~ has a relatively limited impact on graupel mass.

SIP mechanisms significantly enhance ~~electrical~~ non-inductive charging and lightning activity by increasing ice crystal concentrations, particularly at low altitudes where primary ice production is inactive. This promotes ice-graupel collisions and amplifies charge exchange in each grid cell. The intensity of SIP processes varies with the thickness of the warm-phase region. Raindrop shattering freezing is the most sensitive and requires a deep warm-phase, while Hallett-Mossop and collisional ice break-up produce abundant ice crystals in all storms.

1 Introduction

Cloud electrification processes are a key element in understanding and anticipating the electrical structure of thunderstorms and their electrical activity. The non-inductive charging mechanism refers to charge separation during rebounding collisions between ice crystals and graupel in the presence of supercooled liquid water ([Reynolds et al., 1957](#); [Takahashi, 1978](#)). It is known to be the leading process of deep convective cloud electrification ([Norville et al., 1991](#); [Helsdon Jr. et al., 2001](#)). Although all laboratory studies agree on the dependence of the sign and amplitude of the separated charge on temperature and supercooled liquid water content (e.g., Takahashi, 1978; Jayaratne et al., 1983; Saunders and Brooks, 1992; Saunders and Peck, 1998), they can strongly differ in the position of the charge reversal line (see figure 1 in Takahashi et al. (2017) or figure 2 in

Emersic and Saunders (2020)). This charge reversal line represents the temperature and liquid water content conditions where the sign acquired by the hydrometeors reverses. These conflicting laboratory results may be the consequence of difficulties in reproducing natural conditions in a cloud chamber (Takahashi et al., 2017). According to the theory of relative diffusional growth rate (RDGR) (Baker et al., 1987), the temperature at which the exchanged electric charge reverses sign depends on the vapor diffusion growth rate of ice particles. The particle that grows faster by water vapor diffusion charges positively during collision. By examining the factors influencing the rate of vapor deposition growth of pristine ice and graupel, Emersic and Saunders (2020) concluded from their laboratory experiments that, among other factors, the size of ice crystals and the cloud supersaturation should be better characterized. Glassmeier et al. (2018) have performed calculations of the RDGR theory as post-processing of the COSMO model and explored its sensitivity to numerous parameters. They identified ice crystal size as the most important parameter of RDGR, followed by graupel size.

While ice crystals are a key element in cloud electrification, their formation is complex and can follow different pathways. Ice crystals can be formed via homogeneous freezing of supercooled droplets at temperatures below -35°C . At warmer temperatures, aerosols acting as ice nucleating particles (INP) are required to form ice crystals through heterogeneous nucleation. Consequently, aerosols are indirectly involved in cloud electrification via the formation of ice crystals and cloud droplets.

The influence of aerosols acting as cloud condensation nuclei (CCN) on thunderstorm electrification and subsequent lightning activity has been examined in various observation-based and numerical modeling studies. In general, observation-based studies indicate a correlation between increased total lightning or cloud-to-ground lightning activity and increased Aerosol Optical Depth (AOD) in different regions of the world (Shi et al., 2020; Proestakis et al., 2016; Dayeh et al., 2021; Wang et al., 2023; Altaratz et al., 2010). However, studies have observed a decline in lightning activity when the AOD exceeds a threshold value, which ranges from 0.25 to 1, depending on the study. The same behavior is obtained in modeling studies in which the total lightning activity increases with the CCN concentration (Mansell and Ziegler, 2013; Sun et al., 2021; Tan et al., 2017). All these studies argue that an increase in CCN concentration increases the concentration of cloud droplets while reducing their size. Collision-coalescence processes are thus diminished in favor of droplet transport in the mixed phase of the cloud. The release of additional latent heat leads to increased vertical velocities and ice crystal concentrations, which in turn promotes charge separation via the non-inductive process ([van den Heever et al., 2006](#); [Rosenfeld et al., 2008](#); [Sun et al., 2021](#)). Mansell and Ziegler (2013) also detected an optimal CCN concentration of approximately 2000 cm^{-3} , at which total lightning activity is maximized. They attributed the sharp decline in lightning activity at very high CCN concentration to the reduced efficiency of the Hallett-Mossop ice multiplication process, as the size of cloud droplets becomes too small for effective rime splintering. As for the effect of INP concentration on cloud electrification, it has received limited investigation. Using a 1.5D aerosol-cloud bin model, Yang et al. (2020) showed that increasing INP concentration from ~~0.3 to 1.3~~ [300 to 1300](#) L^{-1} results in larger ice particles and enhanced charging rate. However, as shown by Fuchs et al. (2015) and Phillips and Patade (2022), the aerosol effect on cloud electrification is modulated by the temperature at the cloud base or by the warm-phase thickness of the cloud.

Secondary ice production (SIP) processes are recognized as major contributors to ice particle concentrations (e.g., Field et al., 2016; Korolev and Leisner, 2020). Several SIP processes have been identified from laboratory experiments and in situ measurements, and some of them are now parameterized in microphysical schemes enabling the study of their im-

pact on the development and microphysical structure of deep convective clouds (e.g., Sullivan et al., 2018; Qu et al., 2022; Grzegorzczuk et al., 2025a). However, up to now, few studies have focused on their effect on cloud electrification. Yang et al. (2024) simulated a cold-season thunderstorm with four different SIP processes. They showed that the SIP processes are active at different times in the storm lifecycle and at different altitudes, and that the rime splintering mechanism produces the higher ice crystal concentration in this case study. Through an enhancement of graupel and ice crystal production, an increase in the low-level positive charging rate on graupel is simulated, and the modeled flash rate shows better agreement with observations. In a simulation of a cold-based continental thunderstorm, Phillips and Patade (2022) found that the most active SIP process was breakup during ice-ice collisions. This process, acting as a sink of liquid water content, has the ability to alter the polarity of the charge ~~Graupel~~graupel acquires and, consequently, the electric charge structure. They also stated that the cold cloud base temperature makes SIP processes less sensitive to CCN concentration. Several studies have shown that cloud electrification is sensitive to aerosol concentration and SIP processes, but the impact of CCN, INP, and SIP processes on cloud electrification has been studied separately. Moreover, their impact can be modulated by the cloud base temperature and the warm-phase thickness, while most studies have focused on a single case study ([Mansell and Ziegler, 2013](#); [Tan et al., 2017](#); [Yang et al., 2020](#); [Sun et al., 2021](#); [Yang et al., 2024](#); [Huang et al., 2025](#)). Therefore, to assess the impact of ice production on cloud electrification and lightning activity, three idealized thunderstorms with different cloud base temperatures are simulated using the 3D cloud-resolving model Meso-NH (Lac et al., 2018) with a quasi two-moment microphysical scheme (Vié et al., 2016) coupled to an explicit electrical scheme (Barthe et al., 2012). The simultaneous variation of CCN and INP concentrations enables the evaluation of their combined contribution to ice production and their subsequent impact on cloud electrification. The study also tests the sensitivity to three SIP processes. This paper is organized as follows. The model set-up and the methodology are presented in Section 2. Section 3 presents the results on aerosol concentrations, while Section 4 discusses the impact of SIP processes. Section 5 provides a summary.

2 Simulation framework

2.1 The Meso-NH model

In this study the 3D atmospheric mesoscale model Meso-NH (Lac et al., 2018) in version 5-7 is used to simulate different idealized thunderstorms. Meso-NH is the high-resolution limited area research model of the French community and enables performing simulations of idealized cases or real meteorological situations over complex terrain initialized and forced at the lateral boundaries from model outputs. The model has a complete set of physical parameterizations: convection, turbulence, microphysics, aerosols, chemistry, radiation, atmospheric electricity... In the present study, a focus is done on the explicit coupling between the quasi two-moment microphysical scheme LIMA (Vié et al., 2016) and the electrical scheme CELLS (Barthe et al., 2012).

2.1.1 Microphysical scheme

LIMA (Liquid Ice Multiple Aerosols, Vié et al., 2016) is a quasi two-moment microphysics scheme which includes five types of hydrometeors: cloud droplets, raindrops, pristine ice crystals, snow/aggregates, and graupel. ~~The microphysical scheme~~ It predicts the mass mixing ratio for all five categories of hydrometeors and the number concentrations only for cloud droplets, raindrops, and ice crystals. ~~This microphysics scheme~~ LIMA includes a representation of the aerosols as multi-lognormal distributions of aerosols acting as CCN or INP. Details on the LIMA microphysics scheme can be found in Vié et al. (2016).

Here we focus on the different ice crystal production processes: heterogeneous and homogeneous ice nucleation, and SIP processes. Heterogeneous ice nucleation is parameterized as proposed by Phillips et al. (2008, 2013). Activated INP are computed by integration of a reference activity spectrum which depends on supersaturation and temperature. Ice crystals can also form through homogeneous nucleation of cloud droplets at temperatures below -35°C , with the homogeneous cloud droplets freezing rate taken from Eadie (1971). Three SIP processes are available in LIMA. First, the Hallett-Mossop process (HM), often referred to as rime splintering (Hallett and Mossop, 1974) produces an ice splinter each time a graupel is rimed with 200 droplets having diameters between 12 and 25 μm (Beheng, 1987) as described in Vié et al. (2016). Then, the collisional ice break-up (CIBU) mechanism deals with the production of ice splinters during collisions between fragile snow/aggregate particles and large and dense graupel particles (Vardiman, 1978; Takahashi, 1975; Yano and Phillips, 2011). The parameterization implemented in LIMA implementation of this parameterization in LIMA is described by Hoarau et al. (2018). It depends on the impact velocity between the two particles. ~~It~~, and enables varying the number of ice fragments, which can also be randomized (Hoarau et al., 2018). Finally, the ~~third SIP process implemented in LIMA is the~~ raindrop shattering freezing (RDSF) process that was recently implemented in LIMA. It deals with ice splinters production during raindrop freezing. The number of fragments depends on the raindrop diameter (Lawson et al., 2015) and the probability of shattering has a Gaussian temperature dependency centered at 258 K, as introduced in Sullivan et al. (2018) on the basis of droplet levitation experiments (Leisner et al., 2014; Keinert et al., 2020). A general form of the equation describing the RDSF process can be written:

$$\frac{\partial n_i}{\partial t} = \alpha n_i n_r \quad (1)$$

where n_i and n_r are the particle size distribution of cloud ice and raindrops, respectively. An expression for α is:

$$\alpha = \mathcal{N}_{\text{RDSF}} V_r \frac{\pi}{4} D_r^2 \quad (2)$$

where V_r is the impact velocity of a raindrop of size D_r at the surface of the ice crystal. $\mathcal{N}_{\text{RDSF}}$ is the number of ice fragments per raindrop freezing and is parameterized as:

$$\mathcal{N}_{\text{RDSF}} = p_{sh}(T) \chi D_r^4 \quad (3)$$

p_{sh} is the shattering probability depending on temperature (T). According to Lawson (2015), χ is set to $2.5 \times 10^{13} \text{ m}^{-4}$ and $p_{sh}(T) = 0.2 N(258\text{K}, 5\text{K})$ where $N(258\text{K}, 5\text{K})$ is a normal distribution centered around 258 K and with a variance of 5 K. The maximum of the shattering probability was found to be 20 % based on laboratory experiments (Leisner et al., 2014).

2.1.2 The electrical scheme

The Cloud ELectrification and Lightning Scheme (CELLS) (Barthe et al., 2012) is implemented in Meso-NH and can simulate the electrification of clouds and their lightning activity. The scheme computes the evolution of the bulk charge carried by each type of hydrometeors and also takes into account free ions. Several non-inductive charge separation parameterizations are available which are all associated with collisions between a rimmed particle (graupel or snow/aggregates) and an ice particle growing mostly by deposition (ice crystal or snow/aggregates) (~~Takahashi, 1978; Saunders and Brooks, 1992; Saunders and Peck, 1998; ?~~) (Takahashi, 1978; Saunders et al., 1991; Saunders and Peck, 1998; Tsenova et al., 2013). Inductive charging resulting from collisions between graupel and droplets in a preexisting electric field is also considered following the approach of Ziegler et al. (1991). ~~The~~ While the cloud electrification scheme described in Barthe et al. (2012) was built upon the ICE3 one-moment bulk microphysics scheme (Pinty and Jabouille, 1998). ~~Recently, developments have been made,~~ recent developments in Meso-NH ~~so that now allow~~ the cloud electrification scheme ~~can also to~~ be used with the LIMA two-moments microphysics scheme. The electric field is computed at each time step following the procedure described in ~~Barthe and Pinty (2007a)~~ Barthe and Pinty (2007b) and is updated after charge neutralization by each lightning flash. Lightning flashes are triggered when the electric field exceeds a height-dependent threshold (Marshall et al., 1995). Flashes are described first as a bidirectional phase (vertical extension) and secondly, the branches spread horizontally according to a fractal law (see Barthe et al., 2012, for a full description of the lightning flash scheme).

2.2 Case studies and numerical set-up

Different environmental conditions may impact ice crystals formation pathways, and consequently cloud electrification. Therefore, numerical simulations of three contrasted idealized thunderstorms were performed. Figure 1 depicts the different cloud phases (warm, mixed, and cold) during the early electrification of the three simulated thunderstorms. The first case (hereinafter called WARM) is a tropical maritime thunderstorm. It has the warmest cloud base temperature (23°C) and is associated with the deepest warm phase (4 km). The second case (hereinafter called MID-WARM) is a continental case with a slightly warm cloud base (16°C). It has a 2.5 km deep warm phase, is less vertically developed (up to 10 km), and is associated with a shallow cold phase of about 1 km high. The last case (hereinafter called COLD) is a continental storm with a cold cloud base (10°C) and a very shallow warm phase (1.5 km).

All the simulations were performed with the same horizontal ($\Delta x = \Delta y = 500$ m) and vertical ~~grid spacing~~ ($\Delta z = 250$ m) grid spacing. The WARM case ~~was is the thunderstorm observed on the 18th of July 2002 during the CRYSTAL-FACE~~ (Cirrus Regional Study of Tropical Anvils and Cirrus Layers - Florida Area Cirrus Experiment; Leroy et al., 2009) experiment in southern Florida. Meso-NH was run for 1 hour with a 2.5 s time step over a domain of $256 \times 256 \times 72$ gridpoints. A warm bubble of 1.5°C was used to trigger convection. The MID-WARM case was simulated using the sounding from Klemp and Wilhelmson (1978) ~~-(see Fig. S1 in supplement)~~. A warm bubble of 1.5°C was also used to trigger convection. The simulation lasted 1 hour and was run with a time step of 3 s. A $200 \times 200 \times 60$ gridpoints domain was used. The ~~MID-WARM case is the~~ thunderstorm observed on the 18th of July 2002 during the CRYSTAL-FACE (Cirrus Regional Study of Tropical Anvils and Cirrus Layers

experiment in southern Florida. Meso-NH was run for 1 hour with a 2.5 time step over a domain of $256 \times 256 \times 72$ gridpoints.

The COLD case study is the 10 July 1996 thunderstorm observed during the STERAO (Stratospheric-Tropospheric Experiment: Radiation, Aerosols, and Ozone) experiment near the Wyoming-Nebraska-Colorado border. The initial sounding comes from Skamarock et al. (2000) (see Fig. S1 in supplement) and is applied on a domain of $270 \times 270 \times 72$ gridpoints. Unlike Skamarock et al. (2000), who used three warm bubbles to initiate this storm, here a single warm bubble was used. Indeed, the objective was not to reproduce the observed thunderstorm but to simulate a single cell storm with a cold cloud base to simplify the analysis. For each simulation, output files are available every 5 min. Microphysical budgets are calculated and integrated every 5 min over a 5 min period.

Here, the aerosol populations acting as CCN and INP are prescribed using a single mode for each, with a mean radius of 125 nm and 0.8 μm , respectively. The concentration of aerosols acting as CCN (N_{CCN}) is set to a constant value between the ground and 1000 m height, and it decreases exponentially up to 10,000 m, where it reaches the constant value of 0.01 cm^{-3} . The concentration of aerosols acting as INP (N_{INP}) is homogeneous and set to a constant value. In the MID-WARM and COLD simulations, continental aerosols (ammonium sulfate, kappa = 0.61) were used as CCN, while marine aerosols (sea salt, kappa = 1.28) were used in the WARM simulations. The type of aerosols used for INP remained consistent across all simulations and consisted of the same mixture of different species, including dust, black carbon, and organic particles.

In CELLS, the non-inductive charge separation is parameterized following Takahashi (1978). INP mode is composed of 61% of dust, 33% of black carbon and the 6% of organic matter in all simulations (Phillips et al., 2008; Vié et al., 2016).

The choice of the non inductive charging parameterization can impact model results, both in terms of charge structure and total number of flashes (Helsdon Jr. et al., 2001; Altaratz et al., 2005; Mansell et al., 2005; Barthe and Pinty, 2007a; Fierro et al., 2006; Ku. Both the parameterizations of Saunders and Peck (1998) and Takahashi (1978) have been widely used to simulate the electrical activity of thunderstorms. However, recent laboratory studies have shown strong similarities between the charge reversal line in Takahashi (1978) and the ones in Pereyra et al. (2000), Saunders et al. (2006) or Emersic and Saunders (2010), leading us to choose the parameterization of Takahashi (1978) for the non-inductive charge separation in this study. The inductive charging is also activated.

Concerning the lightning scheme, the fractal parameters are set to $\chi = 2.3$ and $L_\chi = 1000 \text{ m}$.

2.3 Sensitivity tests

Since ice crystal production depends on both primary and secondary ice production processes, their contributions are tested separately. A first series of tests is carried out on CCN and INP concentrations. Simulations were performed with N_{CCN} that can take five different values (500, 1000, 5000, 8000, and $10,000 \text{ cm}^{-3}$), and N_{INP} that can also take five different values (0.1, 1, 10, 100, and 1000 L^{-1}). For each fixed N_{INP} , N_{CCN} was varied over its five possible values resulting in a total of 25 simulations. A large range of values of N_{CCN} were used in this study as done in Mansell and Ziegler (2013) where was inspired by the literature. Mansell and Ziegler (2013) made N_{CCN} varied over 13 values between 50 and 8000 cm^{-3} , while Tan et al. (2017) used four different values between 100 and 3000 cm^{-3} . A large range of values for N_{INP} was also set according to observations and literature, while Yang et al. (2020) As for Yang et al. (2020), they used INP concentrations of 0.3;

0.8 and 1.3. In a summary of INP measurements separated based on air mass type, Kanji et al. (2017) showed that 300, 800 and 1300 L⁻¹. In the remainder of the paper, low N_{CCN} refers to 500 cm⁻³, medium to 1000 and 5000 cm⁻³ and high to 8000 and 10,000 cm⁻³. In the same way, low N_{INP} can reach several hundred particles per liter in dust air masses (Boose et al., 2016) at temperatures around -30 corresponds to 0.1, 1 and 10 L⁻¹, medium to 100 L⁻¹ and high to 1000 L⁻¹. In this first set of simulations, only the HM process as a SIP mechanism is activated. For decades, two-moment schemes include a parameterization of the HM process (e.g. Ferrier, 1994; Straka and Mansell, 2005; Seifert and Beheng, 2006; Vié et al., 2016), while the CIBU and RDSF mechanisms have been only recently included in microphysics schemes (Phillips et al., 2017a, 2018; Hoarau et al., 2018; Sullivan et al., 2018) with uncertainties remaining regarding the number of fragments produced by these processes (Grzegorzczuk et al., 2025b). Moreover, CIBU and RDSF can be activated or deactivated in LIMA at the user's discretion while HM is systematically activated. Therefore, it was decided to keep HM active in these first series of simulations.

Four additional simulations are performed for each storm to analyze the impact of SIP processes on cloud electrification and lightning activity. In this series of simulations, aerosol concentrations are kept constant with $N_{CCN} = 1000$ and $N_{INP} = 10$. First the CIBU process is activated in addition to the HM process. The number of fragments is randomly generated between 1 per 10 collisions and 100 per collision (Hoarau et al., 2018). Secondly, the RDSF process is activated in addition to the HM process. The maximum shattering probability is set to 20%. In the third test the HM process is disabled, resulting in a simulation in which no SIP process is considered (hereinafter referred to as NOSIP). In the last simulation, all three SIP processes are activated (hereinafter referred to as ALLSIP). In this series of simulations, aerosol concentrations representative of average aerosol conditions are used. Rose et al. (2021) have surveyed aerosol concentrations using the network of Global Atmosphere Watch (GAW) stations. Using particle number concentration in the range 100-500 nm as a proxy for potential CCN population, they showed that the potential CCN concentration ranges between a few hundreds to a few thousands particles cm⁻³ over the continents. Mansell and Ziegler (2013) and Sun et al. (2021) used values around 1000 cm⁻³ in their modeling studies. Regarding INP concentrations, Kanji et al. (2017) showed that most studies exhibit INP concentrations between 0.5 and 50 L⁻¹ at temperatures colder than -15°C. Therefore, $N_{CCN} = 1000$ cm⁻³ and $N_{INP} = 10$ L⁻¹ are used in all these simulations. The HM simulation where only the HM process is activated corresponds to the simulation with $N_{CCN} = 1000$ cm⁻³ and $N_{INP} = 10$ L⁻¹ from the first set of simulations. The structure of these three storms is illustrated for this particular setup in the Supplementary Material (Figure S2).

3 Aerosol impact on cloud electrification and lightning activity

3.1 Electrical activity

Figure 2a-c represents the total flash number for each N_{CCN} and N_{INP} pairing during 1 hour. The normalized total number of flashes per simulation is plotted in color; the total number of lightning flashes is normalized by its minimum and maximum values for each storm. The lightning activity shows a large variability between the three thunderstorms, and depending on N_{CCN} and N_{INP} . For the same fixed values of N_{CCN} and N_{INP} , the three idealized cases produce a different amount of flashes during 1 h. The MID-WARM case is the most electrically active storm with a total number of flashes ranging from 625 to 4461.

The WARM and COLD cases have a less intense lightning activity with the total number of flashes not exceeding 2800. In the remainder of this paper we will focus on the modification of the electrical activity and of the microphysics of each idealized case due to the sensitivity tests rather than on the differences between the three cases with the same aerosol concentration and SIP process conditions.

~~Whatever~~ For all values of N_{INP} , the total number of flashes is minimum for each storm when $N_{\text{CCN}} = 500 \text{ cm}^{-3}$. In general, the total number of lightning flashes tends to increase with N_{CCN} , as in the WARM case at low N_{INP} . However, in certain cases, threshold values of N_{CCN} can be observed, beyond which the total number of lightning flashes decreases. At low N_{INP} (0.1, 1 and 10), the threshold is at 5000 for the COLD case with a light decrease of the, the total number of flashes above this value. In the MID-WARM case, this threshold is rather around 8000 slowly decreases for $N_{\text{CCN}} > 5000 \text{ cm}^{-3}$. In the WARM case, there is no threshold observed in the simulations with low N_{INP} , and the total number of flashes increases monotonically with N_{CCN} . A in the COLD case. When $N_{\text{INP}} = 100 \text{ L}^{-1}$, this threshold value for N_{CCN} appears for the three storms when $N_{\text{INP}} = 100$: is observed at 5000 cm^{-3} for the MID-WARM and COLD cases, and 8000 cm^{-3} for the WARM case. For the highest high N_{INP} (1000), this threshold effect is still present for the MID-WARM and the COLD cases at a lower and higher N_{CCN} , respectively. However, in general, the electrical activity is less intense at high than at low or medium N_{INP} . When considering the increase of N_{INP} while keeping N_{CCN} constant, a threshold effect is also observed for the three storms (Fig. 2a-e). For N_{CCN} higher than 5000, the total lightning activity peaks when $N_{\text{INP}} = 100$. Overall, lightning activity shows an intensification when N_{CCN} increases up to a threshold which was observed in other numerical studies (Mansell and Ziegler, 2013; Tan et al., 2017). They observed a relatively low N_{CCN} threshold of 1000 above which lightning activity and non inductive charging decrease or remain steady. In the present study, it is shown that N_{CCN} threshold value depends on N_{INP} and the type of storms defined by the cloud base temperature.

The time of the first lightning flash for each simulation is plotted on Fig. 2d-f. The WARM and MID-WARM storms exhibit little variability compared to the COLD storm. The first flash is triggered between 18 and 27 min for the WARM storm, between 20 and 32 min for the MID-WARM storm, but between 15 and 43 min for the COLD storm. The general trend is an earlier triggering of the first flash when N_{CCN} or N_{INP} is increased. The earlier triggering time of the first flash when N_{CCN} or N_{INP} are increased could be associated with a higher non-inductive charging rate in these conditions (Fig. 3).

The variability in the time of the first lightning and the total number of flashes can be mostly explained by the non-inductive charging rate in the convective zone as represented in Fig. 3. A difference in the time the first flash depends on the intensity of the non-inductive charging rate, the depth over which it is active, and the start of cloud electrification (not shown). At high N_{INP} (1000) cloud electrification starts 5 earlier in each storms. An earlier An earlier triggering of the first flash can be due to an earlier cloud electrification, a more intense non-inductive charging rate or a deeper region where the non-inductive charging occurs favors the triggering of the first flash earlier in the cloud lifetime. At high N_{INP} , cloud electrification starts 5 min earlier in each storm (not shown). In the WARM case and for $N_{\text{INP}} = 1000 \text{ L}^{-1}$, the mean non-inductive charging rate is rather low and does not evolve too much when N_{CCN} is increased (Fig. 3a) which is consistent with the lightning activity in such N_{INP} and N_{CCN} conditions (Fig. 2a, first line). At lower N_{INP} , the non-inductive charging rate in the WARM case intensifies when N_{CCN} is increased. However high charging rate occurs on a restricted altitude range at $N_{\text{CCN}} = 10,000 \text{ cm}^{-3}$ which is inline with the

decrease of the total lightning flash number (Fig. 2a). It must be noted that the horizontally averaged non-inductive charging rate in the convective zone in the WARM storm does not show negative charging of the graupel. It leads to the formation of a strong positive layer of charge at low altitude with relatively low negative charges in the storm for low N_{CCN} regardless of N_{INP} (see Fig. S3 in Supplement). Increasing N_{CCN} results in higher average positive and negative charge density located at higher altitude, and in more complex charge structures.

A similar behavior is observed for the MID-WARM case in terms of non-inductive charging rate intensity when N_{INP} and N_{CCN} are varied (Fig. 3, middle column). ~~The MID-WARM simulation enables~~ However, the negative charging of graupel ~~particles for is enabled for low and medium N_{INP} less or equal to 100,~~ and for N_{CCN} higher than a threshold value. This threshold value decreases from $10,000 \text{ cm}^{-3}$ at $N_{INP} = 100 \text{ L}^{-1}$ to 1000 cm^{-3} for low N_{INP} values. Referring to the ~~Takahashi~~ Takahashi (1978)'s diagram, the negative charging of graupel between -10°C and -30°C occurs for cloud water content higher than $0.2\text{--}0.3 \text{ g m}^{-3}$ and lower than 4 g m^{-3} . Then, the negative charging of the graupel signs the presence of significant liquid water content at cold temperatures associated with the transport of a large number of small droplets by the updraft when N_{CCN} increases. The negative charge acquired by the graupel induces a negative layer shifted toward the cloud base (see Fig. S3 in supplement) potentially changing the cloud electrical structure when N_{CCN} and N_{INP} vary.

Concerning the COLD case, the relatively low variability in the total number of lightning flashes (ratio of 5.2 between the minimum and maximum total number of flashes) translates into a low variability in the intensity and altitude range of the non-inductive charging rate (Fig. 3, right column). ~~It must be noted that the~~ Negative charging of graupel occurs at high N_{INP} or high N_{CCN} values. According to the parameterization of Takahashi (1978), the negative charging of graupel occurs as soon as the temperature falls below -10°C meaning that the cloud water content exceeds 1 g m^{-3} at this altitude ~~according to the parameterization of Takahashi (1978). Negative charging of graupel occurs whatever N_{INP} but for high N_{CCN} values.~~

Therefore, in general, the total flash number can be mainly explained by the amount of charge exchanged in the convective region by the non-inductive mechanism.

It must be noted that the total number of flashes and the time of the first flash evolve the same way when N_{CCN} is increased for ~~low $N_{INP}=0.1, 1$ and 10 .~~ In the remainder of this study ~~the three lowest~~ only the simulation with N_{INP} are treated together $= 10 \text{ L}^{-1}$ is shown as a representative for low N_{INP} simulations since they have similar tendencies in their electrical and microphysical properties.

3.2 Microphysical structure of the storms

In order to explain the differences in the electrical activity among the three storms under different aerosol concentration configurations, the key factors contributing to cloud electrification are analyzed. In the parameterization of Takahashi (1978) used in this study, the cloud water content (CWC) and the temperature determine the sign and amount of charge acquired by graupel particles and ice crystals. Therefore, in the following, CWC, ice crystal number concentration and graupel mass are investigated in the convective zone during the initial stage of cloud electrification. The convective region is defined as the region where the maximum vertical velocity is higher than 5 m s^{-1} or the instantaneous precipitation rate is higher than 20 mm h^{-1} .

The initial stage of cloud electrification is defined as the first 10 min during which the absolute value of the non-inductive charging rate integrated over the volume of the convective region is greater than 0.1 C s^{-1} .

3.2.1 Cloud-water content

Figure 6 shows the mean vertical profiles of CWC in the convective region during the early stage of cloud electrification for the three storms and all the sensitivity studies on N_{CCN} and N_{INP} . CWC shows important variations with increasing N_{CCN} and N_{INP} in the three simulated thunderstorms. However, some general characteristics can be highlighted. In general, higher N_{CCN} leads to higher CWC at each altitude, except for $N_{INP} = 100$ in the COLD case in which the highest CWC is reached for $N_{CCN} = 8000$. Moreover, the maximum altitude where values of CWC higher than 0.01 can be found, tends to increase as N_{CCN} rises. It is admitted that higher N_{CCN} yields to higher number concentration of smaller droplets which tends to suppress collection and coalescence processes (Albrecht, 1989; Rosenfeld, 1999). These smaller cloud droplets can be transported at higher altitude where they are converted into ice crystals. The latent heat release is increased leading to stronger updrafts, increased upward transport of cloud droplets, and more CWC at higher altitudes (van den Heever et al., 2006; Sun et al., 2021).

The effect of N_{INP} on the mean vertical profile of CWC in the convective region is more variable. In the WARM case and for $N_{INP} \leq 100$ (Fig. 6g and d), similar mean CWC profiles are obtained whatever N_{CCN} . The maximum values of the mean CWC profiles range between 0.2 and 0.3, and are located at 3 altitude (~ 2 below the 0 isotherm). At the altitude of the 0 and the -10 isotherms, CWC does not exceed 0.2 and 0.08, respectively. In contrast, the mean CWC profiles show a different behavior at the highest N_{INP} (Fig. 6a). At low N_{CCN} , the maximum value of CWC is ~ 0.25 and is located at 2 altitude. Then CWC increases with N_{CCN} , and the altitude where it peaks is shifted upward. For $N_{CCN} = 10,000$, CWC reaches 0.55 at 5 altitude. However, at the altitude of the -10 isotherm, all curves converge to values between 0.05 and 0.1. Finally, mean CWC higher than 0.01 can be found up to 8 to 10 altitude. According to Takahashi's diagram, graupel could only gain a positive charge during non-inductive charging in these conditions (Fig. 3a, 3d and 3g).

In the MID-WARM case (middle column in Fig. 6), the mean CWC profile reaches its maximum between 3 and 4 altitude, i.e. around the 0 isotherm, whatever the altitude, N_{CCN} and N_{INP} . For $N_{INP} = 1000$, the highest mean values of CWC are observed (around 1.1), but CWC is almost null at temperature colder than -10. When N_{INP} decreases, the maximum value of the mean CWC decreases, but significant values of CWC can be found at higher altitudes extending the non-inductive charging zone up to the -20 and -30 isotherms when $N_{INP} = 100$ and $N_{INP} \leq 10$, respectively. According to Takahashi's diagram, due to relatively high CWC for temperatures colder than -10, negative charging of graupel occurs for $N_{INP} = 100$ and high N_{CCN} (Fig. 3e), and for $N_{INP} \leq 10$ and $N_{CCN} \geq 5000$ (Fig. 3e).

The COLD case (Fig. 6e) shows less variability than the two other storms. For high N_{CCN} values, and whatever N_{INP} , the mean CWC is between 0.1 and 0.2 at the altitude of the -30 isotherm. The COLD case is thus favorable for negative graupel charging at relatively low altitude, between the -10 and the -20 isotherms as also revealed by Fig. 3e, 3f and 3i.

While the effect of varying N_{CCN} for a fixed N_{INP} is mainly the same regardless the warm phase thickness, the effect of varying N_{INP} for a fixed N_{CCN} is less straightforward. It will be examined further in the following.

3.2.1 Ice crystal concentration

Ice crystal number concentration is ~~another factor essential~~ an essential factor for non-inductive charging. ~~It~~ it impacts the number of collisions and the amount of charge acquired by each particle. Figure 4 shows the mean vertical profiles of ice crystal number concentration. Most profiles show two main peaks which presence and amplitude depend on aerosol concentration and storm type. These peaks are generally found around the -40°C and -5°C isotherms. To better understand the variability of ice crystal number concentration relative to storm type and aerosol concentrations, the tendencies of the ice production processes are plotted on Fig. 5 for each simulation.

Firstly, we focus on the peak concentration of ice crystals located around -40°C . At such temperature, ice crystals can be produced by heterogeneous (first row in Fig. 5) or homogeneous (second row in Fig. 5) nucleation. Logically, the ~~heterogeneous nucleation has a clear dependence on N_{INP}~~ the production of ice crystals through heterogeneous nucleation (Fig. 5a-c) increases with N_{INP} . ~~However, the ice crystal number concentration in the upper part of the cloud while their production through homogeneous nucleation is favored by high N_{CCN} values~~ (Fig. 4) ~~does not increase with the heterogeneous nucleation rate. Increasing~~ 5d-f). Indeed, ~~when N_{CCN} results in more important homogeneous nucleation rates whatever N_{INP} . Indeed, increases,~~ a larger number of small cloud droplets are produced ~~and~~ transported in the updraft ~~when N_{CCN} increases. Part and part~~ of them are available for homogeneous freezing when reaching the -35°C isotherm. ~~At low N_{INP} , homogeneous Homogeneous~~ nucleation is the most effective at low N_{INP} (Fig. 5d-f). Then, increasing N_{CCN} from 500 to 5000 cm^{-3} results in a 3-4 order of magnitude increase in the ice crystal number concentration in the upper part of the cloud (Fig. 4g-i). Above 5000 cm^{-3} and ~~whatever regardless of N_{INP}~~ , the mean ice crystal number concentration in the upper part of the cloud is not significantly enhanced. On the contrary, at ~~very-high N_{INP} (1000)~~, a different behavior is observed between the MID-WARM and WARM cases, and the COLD case. In the upper part of the cloud, all curves representing different N_{CCN} values are almost merged for MID-WARM and WARM, while the curves for low ~~(500 and 1000)~~ and high ~~(8000 and 10,000)~~ high N_{CCN} are separated by 2 orders of magnitude next to the -40°C isotherm in the COLD case. Indeed, in the COLD case, the lack of efficiency of the warm-rain processes leading to smaller and more numerous droplets, ~~compensates for the Bergeron effect,~~ and allows a significant quantity of supercooled water to reach the -40°C isotherm (Fig. 6c) and freeze ~~(Fig. 5f)~~. On the contrary, for the WARM and MID-WARM cases, in such high N_{INP} , supercooled droplets are riming ice particles and are competing for water vapor with INP, leading to less droplets available for homogeneous nucleation of ice crystals (Phillips et al., 2007; van den Heever et al., 2006).

Lastly, the Hallett-Mossop process is responsible for the second peak of ice crystal number concentration close to the 0°C isotherm. For the WARM and COLD cases, the HM process rate is maximum for ~~high N_{INP} ($\geq 100\text{ L}^{-1}$) and high and~~ N_{CCN} ($\geq 5000\text{ cm}^{-3}$). For the MID-WARM case, the maximum values of the ice crystal production rate via the HM process are also obtained for $N_{\text{INP}} \geq 100\text{ L}^{-1}$, but for medium N_{CCN} values ~~(1000-5000)~~. These differences are the result of a combination of factors that can add up or cancel each other out depending on N_{CCN} , N_{INP} and the warm-phase thickness. The parameterization of the HM process in LIMA follows ~~Beheng (+1987)~~ Beheng (1987). Accordingly, the efficiency of this process increases with the number of cloud droplets with diameter in the range 12 to $26\text{ }\mu\text{m}$, and the graupel mass in the region where the temperature

is between -3°C and -8°C . Now, cloud droplet number concentration increases monotonically with N_{CCN} , while their size decreases (not shown). At low and high N_{CCN} , cloud droplets are therefore either too large or too small to be effective at rime splintering (Takahashi, 1984; Borys et al., 2003; Mansell and Ziegler, 2013). That is why the HM process is the most intense ~~for intermediate values of~~ not for the highest N_{CCN} but at 8000 and 1000 cm^{-3} in the WARM and MID-WARM cases, respectively. As for the impact of N_{INP} on the HM process, it is via the cloud water and graupel content, ~~the later being discussed in Seet. 3.2.3. that are discussed in the next sections.~~

3.2.2 Cloud water content

Figure 6 shows the mean vertical profiles of CWC in the convective region during the early stage of cloud electrification for the three storms and all the sensitivity studies on N_{CCN} and N_{INP} . CWC shows important variations with increasing N_{CCN} and N_{INP} in the three simulated thunderstorms. However, some general characteristics can be highlighted. In general, higher N_{CCN} leads to higher CWC at each altitude, except for $N_{\text{INP}} = 100 \text{ L}^{-1}$ in the COLD case in which the highest CWC is reached for $N_{\text{CCN}} = 8000 \text{ cm}^{-3}$. Moreover, the maximum altitude at which values of CWC higher than 0.01 g m^{-3} can be found, tends to increase as N_{CCN} rises. It is admitted that higher N_{CCN} yields to higher number concentration of smaller droplets which tends to suppress collection and coalescence processes (Albrecht, 1989; Rosenfeld, 1999). These smaller cloud droplets can be transported at higher altitude where they are converted into ice crystals. The latent heat release is increased leading to stronger updrafts, increased upward transport of cloud droplets, and more CWC at higher altitudes (van den Heever et al., 2006; Sun et al., 2021)

The effect of N_{INP} on the mean vertical profile of CWC in the convective region is more variable. In the WARM case and for $N_{\text{INP}} \leq 100 \text{ L}^{-1}$ (Fig. 6g and d), similar mean CWC profiles are obtained independently of N_{CCN} . At the altitude of the 0°C and the -15°C isotherms, CWC does not exceed 0.2 and 0.08 g m^{-3} , respectively. In contrast, at high N_{INP} (Fig. 6a), CWC increases with N_{CCN} , and the altitude where it peaks is shifted upward. However, at the altitude of the -15°C isotherm, all curves converge to values between 0.05 and 0.1 g m^{-3} . Finally, mean CWC higher than 0.01 g m^{-3} can be found up to 8 to 10 km altitude. According to Takahashi (1978)'s diagram, graupel could only gain a positive charge during non-inductive charging in these conditions (Fig. 3a, 3d and 3g).

In the MID-WARM case (middle column in Fig. 6), the mean CWC profile reaches its maximum between 3 and 4 km altitude, i.e. around the 0°C isotherm, regardless of altitude, N_{CCN} and N_{INP} . For $N_{\text{INP}} = 1000 \text{ L}^{-1}$, the highest mean values of CWC are observed ($\sim 1.1 \text{ g m}^{-3}$), but CWC is almost null at temperature colder than -15°C . When N_{INP} decreases, the maximum value of the mean CWC decreases, but significant values of CWC can be found at higher altitudes extending the non-inductive charging zone above -15°C up to 7km and 9km altitude at medium and low N_{INP} , respectively. According to the diagram of Takahashi (1978), due to relatively high CWC for temperatures colder than -10°C , negative charging of graupel occurs for medium N_{INP} and high N_{CCN} (Fig. 3e), and for low N_{INP} and $N_{\text{CCN}} \geq 5000 \text{ cm}^{-3}$ (Fig. 3e).

The COLD case (Fig. 6c) shows less variability than the two other storms. For high N_{CCN} values, and for any value of N_{INP} , the mean CWC is between 0.1 and 0.2 g m^{-3} at the altitude of the -30°C isotherm. The COLD case is thus favorable for

negative graupel charging at relatively low altitude, between the -10°C and the -20°C isotherms as also revealed by Fig. 3c, 3f and 3i.

While the effect of varying N_{CCN} for a fixed N_{INP} is mainly the same with an increase of CWC at higher N_{CCN} , regardless the warm phase thickness, the effect of varying N_{INP} for a fixed N_{CCN} is less straightforward.

3.2.3 Graupel mass

The total mass of graupel in the convective zone during cloud electrification and between 0°C and -40°C is displayed in Fig. 7. The three storms show different impact of N_{CCN} and N_{INP} variations on graupel mass. While in the WARM case, maximum values of graupel mass are achieved for $N_{\text{CCN}} \geq 8000 \text{ cm}^{-3}$ and $N_{\text{INP}} \leq 100 \text{ L}^{-1}$, in the MID-WARM case, they are obtained for $N_{\text{INP}} = 100 \text{ L}^{-1}$ whatever for any value of N_{CCN} . As for In the COLD case, both low N_{CCN} and N_{INP} values are conducive to large graupel mass. However, for each storm type, the ratio between the maximum and the minimum graupel mass is between 1.5 and 1.7. It suggests graupel mass is not a limiting ingredient for cloud electrification in these storms, but it can modulate the amplitude of the charge exchanged during the non-inductive process.

Graupel formation and growth are the result of many mixed-phase processes. Increasing N_{CCN} enhances CWC in the mixed-phase region of clouds by forming more cloud droplets (Sect. 3.2.2) that can be transported above the 0°C isotherm and contribute to the riming growth of graupel. However this trend is only observed in only the WARM case where shows an increase of the total mass of graupel increases with N_{CCN} whatever and independent of N_{INP} . The response of graupel mass to a variation in In the MID-WARM case, the graupel mass remains almost constant when N_{CCN} is different in the MID-WARM and COLD cases. In the COLD case, high graupel mass values are not correlated with high non-inductive charging rates (Fig. 3) and high total flash number (Fig. 2), confirming that graupel mass is not a limiting factor for lightning activity. In this storm varies for a fixed N_{INP} . In the COLD case, higher graupel mass is linked to larger riming rates of raindrops on graupel at low N_{CCN} . The cold cloud base of this storm prevents the growth of most of the raindrops to precipitation size, promoting the transport of smaller raindrops at sub-zero temperatures. In the MID-WARM case, the graupel mass remains almost constant when N_{CCN} varies for a fixed N_{INP} .

In all storms, the graupel mass decreases at high N_{INP} . The formation-Indeed, graupel formation is accelerated through the rapid formation of ice crystals by heterogeneous nucleation increases with when N_{INP} increases (Fig. 5c). Graupel formation is accelerated through the rapid formation of ice crystals that can aggregate and grow These crystals aggregate, then graupel mass is increased by riming, and raindrops that freeze raindrop freezing after collisions with ice crystals. The formation is accelerated but the intensity is weaker At high N_{INP} , graupel formation is thus accelerated, but its growth rate is limited, leading to a lower graupel mass at high N_{INP} .

3.3 The relationship between aerosols, microphysics and electrification

In general, increasing N_{CCN} and N_{INP} leads to an amplification of the lightning activity up to a threshold value due to an increased production of ice crystals. Previous lightning activity due to increased ice crystal production. In this study, the enhancement of lightning activity with increasing N_{CCN} varies between the three storms, with maximum enhancement factors

of 11, 7, and 4 in the WARM, MID-WARM, and COLD cases, respectively. These values are of the same order of magnitude as the ones in the literature. Sun et al. (2023) found that the total number of flashes was multiplied by 5 when N_{CCN} increased from 400 to 6,400 cm^{-3} in a simulated multicell storm developing in a high CAPE environment. Huang et al. (2025) reported a nearly 60-fold increase of the total lightning number in a simulated squall line when the aerosol concentration increased from 400 to 4,000 cm^{-3} . Observational studies based on observations and simulations have found similar results focusing on AOD and N_{CCN} , respectively. AOD and lightning strikes data report similar increases in lightning activity, with enhancement factors ranging from 1.6 to 9 (Thornton et al., 2017; Naccarato et al., 2003; Proestakis et al., 2016).

This study confirms that increasing aerosol concentration leads to an amplification of lightning activity, but only up to a threshold value. Previous numerical experiments found a N_{CCN} threshold around 2000-2,000 cm^{-3} (Mansell and Ziegler, 2013; Tan et al., 2017). In the present study, we further show that the N_{CCN} threshold lies in a large range of N_{CCN} threshold (1000-8000 values (1,000 - 8,000 cm^{-3}) is found depending on both the cloud-base temperature of the storm and depends on both warm-phase thickness and N_{INP} .

Increasing N_{INP} naturally results in higher heterogeneous nucleation rate. However, it is less efficient to produce large concentrations of ice crystals compared to homogeneous nucleation and HM process. The HM process is shown to depend upon N_{CCN} as already highlighted by Takahashi (1984), Mansell and Ziegler (2013) and Borys et al. (2003). But the relationship is less straightforward than in previous studies due to the combined effects of varying N_{INP} and warm-phase thickness. Increasing N_{INP} naturally results in higher heterogeneous nucleation rate. However, compared to homogeneous nucleation and HM process, it is less efficient to produce large concentrations of ice crystals.

The warm-phase thickness influences the warm-phase processes and subsequently the mixed-phase processes. A deeper warm phase favors raindrop formation and precipitation, reducing the supercooled water content in the mixed-phase region at cold temperatures which is essential for cloud electrification. On the contrary, a shallower warm-phase region does not provide an environment where cloud droplets can grow through collision-coalescence processes. In this type of storm these conditions, smaller raindrops and cloud droplets are more easily found at sub-zero temperatures, increasing the depth of the region where the non-inductive charging can occur.

In this study, the graupel mass in the convective region during the early stage of cloud electrification was marginally impacted by aerosol concentrations compared to CWC and ice crystal concentration.

Variations in aerosol concentrations modify both the amplitude and the sign of the charge exchanged during the non-inductive process, and thus the polarity of the cloud's charge structures. Numerous studies have shown that the choice of the non-inductive process parameterization can strongly influence model results, both in terms of charge structure and number of flashes (Helsdon Jr. et al., 2000). Therefore, the charge structures shown in this study would be different if the Saunders and Peck (1998) parameterization was used. However, the objective of this study is not to evaluate which parameterization of the non-inductive process is the best suited for storm modeling, but rather to isolate and explore the effect of ice production on cloud electrification.

4 Effect of secondary ice production on cloud electrification and lightning activity

4.1 Electrical activity

Figures 8 and 9 show the total number of flashes and the time of the first flash, and the total charge gained by graupel in the convective zone during cloud electrification, respectively, for all ~~the sensitivity tests dedicated to the effect of SIP processes~~SIP-related sensitivity tests. When no SIP process is activated (NOSIP), the total number of flashes is the lowest and the first flash is triggered the latest among all tests, for all three cases. This is due to a very low charging rate (≤ 1 pC m⁻³ s⁻¹) in a very small cloud depth (≤ 1.5 km).

Activating the HM process leads to a higher number of flashes, especially in the COLD case; ~~In this case,~~ the total number of flashes is doubled from 466 to 1009 between the NOSIP and the HM simulations. Additionally, the first flash is triggered 3 to 7 min earlier ~~when the HM process is activated. Figure 9 clearly shows an enhancement of the~~ and the non-inductive charging zone ~~when the HM process is active~~ is clearly enhanced (Fig. 9).

Activating the CIBU process in addition to the HM process (HM+CIBU) ~~considerably increases~~ multiplies the total number of flashes ~~for the three storms. The total number of flashes is multiplied by a factor by~~ ~ 25 for the WARM and COLD cases, and by ~ 8 for the MID-WARM case. In addition, the time of the first flash is further reduced compared to the HM simulations (between 2 and 6 min). This higher and earlier lightning activity is associated with a dramatic increase of the non-inductive charging rate up to 30 pC m⁻³ s⁻¹ ~~as shown in~~ (Fig. 9).

When the RDSF process is activated in addition to the HM process (HM+RDSF), the total number of flashes increases compared to the HM simulations, for the WARM and MID-WARM cases. This enhancement is 7 times higher in the WARM case than in the MID-WARM case. In ~~terms of time of first flash, the effect of RDSF is the same as the one of CIBU for the~~ WARM case, RDSF has a slightly higher impact than CIBU, whereas in the MID-WARM case RDSF has half the impact of CIBU. RDSF and CIBU share the same time of the first flash in these two storms (Fig. 8b). ~~However~~ In contrast, the RDSF process does not affect the electrical activity (Fig. 8a) and the non-inductive charging rate (Fig. 9) of the COLD storm.

Finally, when all SIP processes are ~~active~~ activated (ALLSIP), the total number of flashes is maximum and the time of the first flash is minimum for the three types of storm. The total number of flashes is multiplied by 75, 21 and 53 compared to the NOSIP simulations for the WARM, MID-WARM and COLD cases, respectively. In the WARM ~~case and MID-WARM cases,~~ the dramatic increase in the total number of flashes is largely due to the combined effect and significant impact of the RDSF and CIBU processes. ~~In the WARM and MID-WARM cases, both the CIBU and RDSF processes have a significant impact on the total number of flashes.~~ However, the effect of CIBU is almost 25 times, ~~two~~ 2 times, and similar to that of RDSF in terms of total lightning activity in the COLD, MID-WARM, and WARM cases, respectively. The first flash is triggered 10 min earlier compared to the NOSIP simulations. ~~The reasons for such differences are investigated in the following~~

In general, SIP processes intensify the average density of charge and can modify the charge structure of the cloud (see Fig. S4 in Supplement). While HM and CIBU processes impact the charge structure below 10 km altitude, the effect of CIBU is more visible above 10 km altitude, in all storms.

4.2 Microphysics

490 4.2.1 Ice crystal number concentration

Figure 10 shows the SIP tendencies summed on the convective region of each storm for each simulation while Fig. 11 displays the mean vertical profiles of ice crystal number concentration for each storm and each simulation.

The HM ~~mechanism~~ process has the lowest tendency among the three SIP, due to the restricted range of temperature in which it is active and the relative low number of splinters produced. However, this is enough to increase the ice crystal number
495 concentration by more than one order of magnitude between 5 and 7 km altitude in the three storms (Fig. 11) ~~in the three storms~~. The ice crystal number concentration is also increased between 8 and 12 km altitude when the HM process is active in the WARM and MID-WARM simulations. From Fig. 10a and 10b, this peak can be assigned to an increase of the homogeneous nucleation tendency. This is due to ~~a delayed cloud electrification in the NOSIP simulation. The~~ the conditions for cloud electrification that are met 5 min later in NOSIP, at a time when homogeneous nucleation is less active.

500 In the WARM case, the HM process tendency is ~~identical~~ similar for the two pairs of simulations HM and HM+CIBU ($6.5 \times 10^9 \text{ kg}^{-1} \text{ s}^{-1}$), and HM+RDSF and ALLSIP (7.1 and $7.2 \times 10^9 \text{ kg}^{-1} \text{ s}^{-1}$), meaning that RDSF has a positive impact on the HM process. The CIBU process is very efficient in producing ice crystals over the whole mixed and cold cloud depth, leading to an increase of ice crystal number concentration by around two orders of magnitude compared to the NOSIP simulation (green and blue lines in Fig. 11a). It peaks at 10 km altitude with value $\sim 300 \text{ L}^{-1}$. RDSF is the most efficient SIP
505 in this storm; it induces a maximum of 1000 L^{-1} at 15 km altitude (orange line in Fig. 11a). Despite being the most active at -15°C (7.5 km in the WARM case), the RDSF process results in high N_i throughout the whole mixed and cold cloud depth, as the CIBU process, due to vertical transport. In ~~a~~ such a deep warm cloud depth, cloud droplets can be efficiently converted into raindrops, providing a favorable environment for the RDSF process. When the three SIP processes are active (ALLSIP), they add up to produce mean ice crystal number concentration that reaches a maximum of 1500 L^{-1} .

510 In the MID-WARM case, the HM process increases the mean ice crystal number concentration by up to 3 orders of magnitude between 4 and 6 km altitude in the vicinity of its active temperature range, producing ice crystal number concentration up to 8 L^{-1} (black line in Fig. 11b). The RDSF process increases the ice crystal number concentration by a factor 10 around 6 km altitude which is inline with its parameterization. ~~In contrast, CIBU~~ CIBU also makes the ice crystal number concentration increase by up to a factor of 10, but over the whole mixed and cold cloud depth ~~by up to a factor of 10.~~ As in the WARM, case,
515 the ALLSIP simulation produces the ~~highest~~ highest mean ice crystal number concentration. Despite a stronger tendency for the RDSF process than for the CIBU process, the HM+RDSF simulation presents lower values of ice crystal concentration along the vertical profile. Indeed, the RDSF process produces a high amount of ice crystals at the early stage of the storm but becomes rapidly inactive. Grzegorzczuk et al. (2025a) found a similar evolution of the RDSF process which get surpassed by the HM mechanism when the storm starts to glaciate. Actually, RDSF needs a deep warm-phase cloud depth and a moderate updraft
520 which will help raindrops to grow and to be lifted up to the right temperature region, around -15°C (Sullivan et al., 2018). Interestingly, in the ALLSIP simulation, the RDSF process tendency is tripled compared to the HM+RDSF simulation 10. This

demonstrates a positive feedback from the CIBU process: the production of additional ice crystals increases the collisions with rain drops.

Figure 10c shows that the COLD case has a particular behavior compared to the WARM and MID-WARM cases (Fig. 10a and 10b, respectively). Due to its limited warm cloud depth (less than 1.5 km thick, Fig. 1c), there is little opportunity for warm rain to form (Gupta et al., 2023), and to further participate to ice multiplication through the RDSF process (Fig. 10c). Consequently the curves of the mean ice crystal number concentration are merged in Fig. 11c for the HM and HM+RDSF simulations. In contrast, the HM and CIBU processes ~~are very efficient in producing ice crystals. They~~ increase the mean ice crystal number concentration by up to a factor of 1000 in the temperature range in which they are active, i.e. between -3 and -8°C and in the mixed-phase region, respectively. In the HM+CIBU and ALLSIP simulations, the mean ice crystal number concentration reaches 500 L^{-1} at 11 km altitude.

4.2.2 Cloud water content

Figure 12 shows the mean vertical profiles of CWC during cloud electrification. In the WARM case, the NOSIP simulation produces the lowest CWC. It reaches a maximum of 0.08 g m^{-3} near the 0°C isotherm against a maximum of 0.15 g m^{-3} in all simulations where SIP processes are activated. As soon as one SIP process is activated, all mean vertical profiles of CWC are merged in the WARM and MID-WARM cases. In the MID-WARM case, in the altitude range between the 10°C and -10°C isotherms, CWC is higher in the NOSIP simulation than in all simulations where SIP processes are activated ~~near the 0 isotherm~~. At temperatures colder than -10°C , CWC exponentially decreases in the NOSIP simulation, while higher CWC are found at higher altitude when SIP processes are considered.

It is important to note that the beginning of the electrification period may be different in the different sensitivity studies. As SIP processes accelerate the formation of ice particles, cloud electrification starts ~~5 min~~ earlier as soon as one SIP process is activated compared to the NOSIP simulation, in the WARM and MID-WARM storms. When the mean vertical profile of CWC is computed at the same time period as the one when SIP processes are activated (not shown), CWC is lower when SIP processes are considered. This is in agreement with previous numerical studies of SIP impact (Zhao and Liu, 2022; Grzegorzczuk et al., 2025a). Indeed, SIP processes are sink of CWC through the riming of snow/aggregates and graupel, ~~and through the Bergeron effect~~. The COLD case does not show any impact of the SIP processes on the average CWC profile in the early cloud electrification stage. The SIP processes do not change the timing of cloud electrification onset in the COLD storm. As cloud electrification starts during the development stage of the cloud, SIP processes have not yet consumed CWC. Despite the presence of significant CWC in the mixed phase region above the 0°C isotherm in NOSIP simulations across all storms, the non-inductive charging process only occurs at high altitude (between 7.5 and 11 km), where ice crystals are available (Fig. 9 and Fig. 11).

4.2.3 Graupel mass

As in Sect. 3.2.3, mild-only slight changes are found in ~~graupel mass between the sensitivity tests in which SIP processes are activated or not~~ the graupel mass in the SIP series of simulations. In general, the total mass of graupel is higher when no SIP

555 process is activated. ~~In the WARM and MID-WARM cases, the~~ The total graupel mass decreases from 16×10^8 to 14×10^8 kg, and from 12×10^8 to 9.8×10^8 kg in the WARM and MID-WARM cases, respectively, as soon as the HM process is taken into account. In the COLD case, the total graupel mass only varies between 6.9×10^8 and 6.5×10^8 kg, with the maximum value for the NOSIP simulation. However, in this case study, only the inclusion of the CIBU process reduces the total mass of graupel.

560 SIP processes reduce graupel mass through two different pathways. The HM and RDSF processes directly consume graupel to form ice crystals while the CIBU parameterization of Hoarau et al. (2018) considers that splinters originate from breaking aggregates during collisions with graupel. The reduction of graupel mass when SIP processes are activated can also be attributed to the competition for CWC, which is shared between the riming of snow particles and the riming of graupel. SIP mechanisms produce numerous ice crystals, which can aggregate or grow into snow particles by water vapor deposition. As a result, cloud
565 droplets increasingly rime onto snow/aggregates at the expense of graupel riming growth.

4.2.4 The relationship between SIP processes, microphysics and electrification

In the NOSIP simulations, ~~the cloud electrification is weak leading to a late triggering of flashes and a low lightning activity in all storms. In these simulations,~~ only homogeneous and heterogeneous nucleation produce ice crystals resulting in low ice crystal number concentration at warm temperatures which limits the non-inductive charging rates. It can explain the late
570 triggering of flashes and the low lightning activity in all storms.

Activating SIP processes enhances the ice crystal number concentration and the lightning activity, with an impact ~~tenfold~~ 5
times greater than that of aerosol concentration. ~~However in terms of the number of flashes. This is lower than the 100-fold~~
flash rate increase deduced from Huang et al. (2025) when multiple SIP processes are activated, especially under high CCN
concentrations. Using the WRF model, Yang et al. (2024) also found that when SIP processes are taken into account in a
575 simulation of a cold-season thundersorm, collisions between graupel and ice crystals are enhanced leading to an increase of
the vertical electric field and flash rate.

Numerical studies consistently highlight the dominant role of SIP processes over primary ice production especially in
the mixed phase region (Huang et al., 2022; Grzegorzczuk et al., 2025a). However, SIP efficiency can vary with microphysical
conditions. For instance, Zhao and Liu (2022) found reduced SIP rates when using a stronger primary ice nucleation parameterization:
580 cloud glaciation is accelerated, and rain and graupel formation is reduced which inhibits SIP processes. In the present study,
SIP sensitivity was tested using only one set of N_{CCN} and N_{INP} , and the sensitivity to SIP parameterization has not been
explored. Prior work (e.g. Mansell and Ziegler, 2013) has shown that different HM parameterizations significantly influence
electrification.

In general, all SIP processes produce a high ice crystal number concentration at altitudes lower than those at which homogeneous
585 and heterogeneous nucleation occurs. This enables the cohabitation between CWC, high ice crystal number concentration and
graupel particles which results in more intense cloud electrification. However, each SIP mechanism has a different impact according to the cloud base temperature of the storm. ~~The~~ HM and CIBU processes enhance cloud electrification and lightning activity in every storm. Activating the HM process makes the first flash to be triggered approximately 5 min earlier. When

combined with CIBU or RDSF, it can advance the triggering of the first flash by up to 10 min. ~~All SIP processes produce high ice crystal number concentration at altitudes lower than the ones where homogeneous and heterogeneous nucleation occur. This enables the cohabitation between CWC, high ice crystal number concentration and graupel particles which results in more intense cloud electrification.~~ The RDSF process requires specific and challenging conditions to take place. In the COLD case, this process is not active due to the lack of raindrop formation. In the MID-WARM storm, the RDSF process can take place and produce a large number of ice crystals, but it becomes rapidly inactive. In contrast, in the WARM storm, the RDSF process is very efficient in producing secondary ice crystals and is active during the whole cloud electrification period thanks to a constant raindrops supply and efficient updraft.

SIP processes also impact the mean CWC vertical profile and the graupel mass, though to a lesser extent within the short window of cloud electrification defined in this study. The main differences observed on these two parameters are due to different cloud electrification onsets and to the enhanced production of snow/aggregate particles that can grow by riming of cloud droplets at the expense of graupel. However, as already shown in Sect. 3.2.3, graupel mass is not a limiting factor in cloud electrification and lightning activity ~~whatever the storm.~~ across the three storms.

The weak sensitivity of CWC to SIP processes may result from several factors. Data sampling during the electrification period limits the detection of differences, which occur more significantly during the storm's mature stage. Another possible explanation is the use of a saturation adjustment scheme in LIMA. This adjustment, applied after all other microphysical processes, forces the environment to reach a strict equilibrium at water saturation at the end of each time step. The use of saturation adjustment can overestimate condensate mass and enhance rain formation, reducing supercooled water (Khain et al., 2015; Zhang et al., 2020). In contrast, studies showing stronger CWC responses to SIP does not use saturation adjustment schemes (Phillips and Patade, 2022; Grzegorzewski et al., 2023). Additionally, in the version of LIMA used in this study, snow and graupel number concentrations are not prognostic, potentially accelerating their formation and depleting liquid and small ice species in comparison with a full two-moment version of LIMA (Taufour et al., 2024).

5 Conclusions

Three idealized thunderstorms that differed by their warm-phase cloud thickness were simulated in order to assess the influence of ice production processes on cloud electrification and lightning activity. This was done using the cloud-resolving model Meso-NH with the quasi two-moment microphysics scheme LIMA coupled with the explicit electrical scheme CELLS. A first set of simulations was performed by simultaneously varying the number concentration of aerosols acting as INP and CCN. A second set of simulations was conducted in which three different SIP processes were alternately active or deactivated. Our results indicate that both aerosol concentration and SIP processes alter the cloud microphysics and the subsequent electrical activity. Several effects can be observed: a delay in the onset of cloud electrification and in the triggering of the first lightning flash, as well as a change in the total number of flashes.

Sensitivity tests on aerosol concentration show that an increase in N_{CCN} and N_{INP} generally enhances lightning activity; ~~either monotonically or up to a specific threshold (N_{INP} : 100 ; N_{CCN} : 1000–8000). Beyond these thresholds, the total number~~

~~of flashes decreases.~~ Aerosol concentrations affect cloud electrification by modulating the vertical profiles of CWC and ice crystal number concentration, ~~while having a comparatively smaller impact on graupel mass.~~ A higher N_{CCN} leads to a greater CWC, expanding the mixed-phase region of the cloud, while a higher N_{INP} depletes CWC at high altitude, altering both the sign and magnitude of the charge exchanged during the non-inductive charging mechanism. N_{INP} favours an accelerated production of rimed particles as a result of a sequence of microphysical processes. Aerosols also control homogeneous nucleation, which is dominant at high N_{CCN} and low N_{INP} . ~~As for the HM process, it is maximum at intermediate or high N_{CCN} levels (1000–10,000), and enables the production of a large number of ice crystals directly in the mixed-phase region of the cloud. By Therefore,~~ by increasing the ice crystal number concentration, aerosol concentration controls the number of ice crystal-graupel collisions, thereby influencing the amount of charge exchanged at each grid point. ~~Moreover N_{INP} favours an accelerated production of rimed particles which is usually the outcome of a sequence of microphysical processes.~~ Despite a similar onset of cloud electrification, the triggering time of the first flash can differ according to the intensity of charge separation by the non-inductive process. ~~Previous observation-~~

However, the increase of lightning activity with aerosol concentration can be monotonic or up to a specific threshold (N_{INP} : 100 L⁻¹; N_{CCN} : 1000–8000 cm⁻³). The total number of flashes decreases beyond these thresholds. Previous observational and numerical modeling studies have also found an enhancement of lightning activity with high aerosol loading up to a threshold value (Altartatz et al., 2010; Shi et al., 2020; Mansell and Ziegler, 2013; Tan et al., 2017). Numerical modeling studies ~~have found N_{CCN} threshold value of identified a CCN threshold around 2000 cm⁻³ (Mansell and Ziegler, 2013; Tan et al., 2017) -This value~~ (Altartatz et al., 2010; Shi et al., 2020; Mansell and Ziegler, 2013; Tan et al., 2017) which is of the same order of magnitude as ~~that of~~ our study. Mansell and Ziegler (2013) attributed the decrease of lightning activity with N_{CCN} to the HM process, while Tan et al. (2017) hypothesized that vapor competition leads to a decrease in ice crystal size and mixing ratio. Most ~~simulation-numerical modeling~~ studies have focused only on N_{CCN} or N_{INP} . Our findings highlight a complex interaction between CCN and INP. ~~At a fixed N_{INP} , the N_{CCN} threshold value varies. Thus, both particles has, and both particles have~~ to be taken into account to understand the aerosol impact on cloud electrification and lightning activity.

Sensitivity tests on SIP processes (rime splintering, raindrop shattering by freezing, and collision ice breakup) demonstrated that they are essential to produce high ice crystal number concentration, especially at low altitudes where primary ice production does not occur. When no SIP process is activated and ~~whatever regardless of~~ the simulated storm, the ice crystal number concentration remains low, resulting in a weak cloud electrification and lightning activity. When they are active, the intensity of each process depends on the thickness of the cloud's warm-phase. A thick cloud's warm-phase region favors the growth of cloud droplets and their conversion to raindrops, enabling raindrop shattering freezing afterwards. On the contrary, a thinner cloud's warm-phase region creates an environment with fewer raindrops and more supercooled cloud droplets aloft conducive to the HM process. The CIBU process is active ~~whatever regardless of~~ the cloud base temperature. ~~Yang et al. (2024) studied the impact of ice multiplication on a cold-season thunderstorm with simulations. They found that when ice multiplication processes are taken into account collisions between graupel and ice crystals are enhanced leading to an increase of the vertical electric field and flash rate. However the activity of RDSF is more intense in their study compared to our results. Each SIP~~

~~process has a different impact~~ This differentiated effect of SIP processes on cloud electrification. ~~This~~ is consistent with Phillips and Patade (2022) results for a cold-base thunderstorm in which HM and RDSF are almost inactive.

Comparing the impact of aerosol and SIP processes on cloud microphysics and electrification, it is clear that both ice production pathways are essential for cloud electrification through the non-inductive charging mechanism. However, SIP processes
660 have a more important impact on the cloud electrification and the resulting lightning activity. While variations in the aerosol concentration can increase the total number of flashes by up to an order of magnitude, activating the SIP processes can multiply the total number of flashes by ~~500~~ 50 in some cases. It is also important to note that the relationship between aerosol concentration, SIP processes, and cloud electrification is complex and varies depending on the cloud base temperature. This study highlights the importance of taking into account the formation of ice crystals via SIPs, as this largely determines the conditions
665 required for the non-inductive mechanism to take effect. However, uncertainties arise from the parameterizations of the SIP processes in the model. For example, several studies have proposed a more complex parameterization of the CIBU process by including dependence on physical parameters (Phillips et al., 2017b; Grzegorzczak et al., 2025a). Additionally, there is still no consensus on the parameterization of the non-inductive process, and several existing parameterizations should be tested.

The next step will be to simulate a thunderstorm observed during the EXAEDRE (EXploiting new Atmospheric Electricity
670 Data for Research and the Environment) field campaign that took place in Corsica in 2018. This campaign offers many observations of cloud microphysics and electric activity, including data from operational weather radar, from a suite of airborne microphysics probes and the airborne 95 GHz Doppler cloud radar RASTA onboard the French Falcon research aircraft, and from the SAETTA network (Coquillat et al., 2019). It provides a robust database for comparison with the numerical simulations. This study will focus on SIP processes and will be crucial for improving our understanding of SIP processes and their
675 role in cloud electrification, as well as validating the findings presented in this study.

Code availability. Version 5-7 of Meso-NH is under the CeCILL-C license agreement and freely available at <http://mesonh.aero.obs-mip.fr/mesonh57>.

Author contributions. IV: Writing – original draft, Methodology, Investigation, Formal analysis, Visualization. CB: Writing - Review & Editing, Conceptualization, Software, Methodology, Investigation, Validation, Funding acquisition, Supervision

680 *Competing interests.* The contact author has declared that none of the authors has any competing interests.

Acknowledgements. This work supported by Agence Nationale de la Recherche (ANR) of France under grant ANR-21-CE01-0006 (ANR ICCARE). Computations were performed on the local computer cluster of Laboratoire d'Aérodynamique.

References

- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- Altaratz, O., Reisin, T., and Levin, Z.: Simulation of the electrification of winter thunderclouds using the three-dimensional Regional Atmospheric Modeling System (RAMS) model: Single cloud simulations, *J. Geophys. Res.: Atmos.*, 110, 2004JD005616, <https://doi.org/10.1029/2004JD005616>, 2005.
- Altaratz, O., Koren, I., Yair, Y., and Price, C.: Lightning response to smoke from Amazonian fires, *Geophys. Res. Lett.*, 37, 2010GL042679, <https://doi.org/10.1029/2010GL042679>, 2010.
- Baker, B., Baker, M. B., Jayaratne, E. R., Latham, J., and Saunders, C. P. R.: The Influence of Diffusional Growth Rates On the Charge Transfer Accompanying Rebounding Collisions Between Ice Crystals and Soft Hailstones, *Q. J. Roy. Meteor. Soc.*, 113, 1193–1215, <https://doi.org/10.1002/qj.49711347807>, 1987.
- Barthe, C. and Pinty, J.: Simulation of electrified storms with comparison of the charge structure and lightning efficiency, *J. Geophys. Res.-Atmos.*, 112, 2006JD008241, <https://doi.org/10.1029/2006JD008241>, 2007a.
- Barthe, C. and Pinty, J.-P.: Simulation of a supercellular storm using a three-dimensional mesoscale model with an explicit lightning flash scheme, *J. Geophys. Res.: Atmos.*, 112, <https://doi.org/10.1029/2006JD007484>, 2007b.
- Barthe, C., Chong, M., Pinty, J.-P., Bovalo, C., and Escobar, J.: CELLS v1.0: updated and parallelized version of an electrical scheme to simulate multiple electrified clouds and flashes over large domains, *Geosci. Model Dev.*, 5, 167–184, <https://doi.org/10.5194/gmd-5-167-2012>, 2012.
- Beheng, K. D.: Microphysical Properties of Glaciating Cumulus Clouds: Comparison of Measurements With A Numerical Simulation, *Q. J. Roy. Meteor. Soc.*, 113, 1377–1382, <https://doi.org/10.1002/qj.49711347815>, 1987.
- Boose, Y., Sierau, B., García, M. I., Rodríguez, S., Alastuey, A., Linke, C., Schnaiter, M., Kupiszewski, P., Kanji, Z. A., and Lohmann, U.: Ice nucleating particles in the Saharan Air Layer, *Atmos. Chem. Phys.*, 16, 9067–9087, <https://doi.org/10.5194/acp-16-9067-2016>, 2016.
- Borys, R. D., Lowenthal, D. H., Cohn, S. A., and Brown, W. O. J.: Mountaintop and radar measurements of anthropogenic aerosol effects on snow growth and snowfall rate, *Geophys. Res. Lett.*, 30, 2002GL016855, <https://doi.org/10.1029/2002GL016855>, 2003.
- Coquillat, S., Defer, E., de Guibert, P., Lambert, D., Pinty, J.-P., Pont, V., Prieur, S., Thomas, R. J., Krehbiel, P. R., and Rison, W.: SAETTA: high-resolution 3-D mapping of the total lightning activity in the Mediterranean Basin over Corsica, with a focus on a mesoscale convective system event, *Atmos. Meas. Tech.*, 12, 5765–5790, <https://doi.org/10.5194/amt-12-5765-2019>, 2019.
- Dayeh, M. A., Farahat, A., Ismail-Aldayeh, H., and Abuelgasim, A.: Effects of aerosols on lightning activity over the Arabian Peninsula, *Atmos. Res.*, 261, 105723, <https://doi.org/10.1016/j.atmosres.2021.105723>, 2021.
- Eadie, W. J.: A molecular theory of the homogeneous nucleation of ice from supercooled water, The University of Chicago, 1971.
- Emersic, C. and Saunders, C.: Further laboratory investigations into the Relative Diffusional Growth Rate theory of thunderstorm electrification, *Atmos. Res.*, 98, 327–340, <https://doi.org/10.1016/j.atmosres.2010.07.011>, 2010.
- Emersic, C. and Saunders, C.: The influence of supersaturation at low rime accretion rates on thunderstorm electrification from field-independent graupel-ice crystal collisions, *Atmos. Res.*, 242, 104962, <https://doi.org/10.1016/j.atmosres.2020.104962>, 2020.
- Ferrier, B. S.: A Double-Moment Multiple-Phase Four-Class Bulk Ice Scheme. Part I: Description, *J. Atmos. Sci.*, 51, 249 – 280, [https://doi.org/10.1175/1520-0469\(1994\)051<0249:ADMMPF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<0249:ADMMPF>2.0.CO;2), 1994.

- Field, P. R., Lawson, R. P., Brown, P. R. A., Lloyd, G., Westbrook, C., Moisseev, D., Miltenberger, A., Nenes, A., Blyth, A., Choularton, T., Connolly, P., Buehl, J., Crosier, J., Cui, Z., Dearden, C., DeMott, P., Flossmann, A., Heymsfield, A., Huang, Y., Kalesse, H., Kanji, Z. A., Korolev, A., Kirchgaessner, A., Lasher-Trapp, S., Leisner, T., McFarquhar, G., Phillips, V., Stith, J., and Sullivan, S.: Chapter 7. Secondary Ice Production - current state of the science and recommendations for the future, *Meteor. Mon.*, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0014.1>, 2016.
- Fierro, A. O., Gilmore, M. S., Mansell, E. R., Wicker, L. J., and Straka, J. M.: Electrification and Lightning in an Idealized Boundary-Crossing Supercell Simulation of 2 June 1995*, *Mon. Wea. Rev.*, 134, 3149–3172, <https://doi.org/10.1175/MWR3231.1>, 2006.
- Fuchs, B. R., Rutledge, S. A., Bruning, E. C., Pierce, J. R., Kodros, J. K., Lang, T. J., MacGorman, D. R., Krehbiel, P. R., and Rison, W.: Environmental controls on storm intensity and charge structure in multiple regions of the continental United States, *J. Geophys. Res.-Atmos.*, 120, 6575–6596, <https://doi.org/10.1002/2015JD023271>, 2015.
- Glassmeier, F., Arnold, L., Dietlicher, R., Paukert, M., and Lohmann, U.: A Modeling Study on the Sensitivities of Atmospheric Charge Separation According to the Relative Diffusional Growth Rate Theory to Nonspherical Hydrometeors and Cloud Microphysics, *J. Geophys. Res.-Atmos.*, 123, <https://doi.org/10.1029/2018JD028356>, 2018.
- Grzegorzczak, P., Wobrock, W., Canzi, A., Niquet, L., Tridon, F., and Planche, C.: Investigating secondary ice production in a deep convective cloud with a 3D bin microphysics model: Part II - Effects on the cloud formation and development, *Atmos. Res.*, 314, 107 797, <https://doi.org/10.1016/j.atmosres.2024.107797>, 2025a.
- Grzegorzczak, P., Wobrock, W., Canzi, A., Niquet, L., Tridon, F., and Planche, C.: Investigating secondary ice production in a deep convective cloud with a 3D bin microphysics model: Part I - Sensitivity study of microphysical processes representations, *Atmos. Res.*, 313, 107 774, <https://doi.org/https://doi.org/10.1016/j.atmosres.2024.107774>, 2025b.
- Gupta, A. K., Deshmukh, A., Waman, D., Patade, S., Jadav, A., Phillips, V. T. J., Bansemer, A., Martins, J. A., and Gonçalves, F. L. T.: The microphysics of the warm-rain and ice crystal processes of precipitation in simulated continental convective storms, *Commun. Earth Environ.*, 4, 226, <https://doi.org/10.1038/s43247-023-00884-5>, 2023.
- Hallett, J. and Mossop, S. C.: Production of secondary ice particles during the riming process, *Nature*, 249, 26–28, <https://doi.org/10.1038/249026a0>, 1974.
- Helsdon Jr., J. H., Wojcik, W. A., and Farley, R. D.: An examination of thunderstorm-charging mechanisms using a two-dimensional storm electrification model, *J. Geophys. Res.: Atmos.*, 106, 1165–1192, <https://doi.org/10.1029/2000JD900532>, 2001.
- Hoarau, T., Pinty, J.-P., and Barthe, C.: A representation of the collisional ice break-up process in the two-moment microphysics LIMA v1.0 scheme of Meso-NH, *Geosci. Model Dev.*, 11, 4269–4289, <https://doi.org/10.5194/gmd-11-4269-2018>, 2018.
- Huang, S., Yang, J., Li, J., Chen, Q., Zhang, Q., and Guo, F.: Impact of secondary ice production on thunderstorm electrification under different aerosol conditions, *Atmos. Chem. Phys.*, 25, 1831–1850, <https://doi.org/10.5194/acp-25-1831-2025>, 2025.
- Huang, Y., Wu, W., McFarquhar, G. M., Xue, M., Morrison, H., Milbrandt, J., Korolev, A. V., Hu, Y., Qu, Z., Wolde, M., Nguyen, C., Schwarzenboeck, A., and Heckman, I.: Microphysical processes producing high ice water contents (HIWCs) in tropical convective clouds during the HAIC-HIWC field campaign: dominant role of secondary ice production, *Atmospheric Chemistry and Physics*, 22, 2365–2384, <https://doi.org/10.5194/acp-22-2365-2022>, 2022.
- Jayaratne, E. R., Saunders, C. P. R., and Hallett, J.: Laboratory studies of the charging of soft-hail during ice crystal interactions, *Q. J. Roy. Meteor. Soc.*, 109, 609–630, <https://doi.org/10.1002/qj.49710946111>, 1983.
- Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Krämer, M.: Overview of Ice Nucleating Particles, *Meteor. Mon.*, 58, 1.1–1.33, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0006.1>, 2017.

- Keinert, A., Spannagel, D., Leisner, T., and Kiselev, A.: Secondary Ice Production upon Freezing of Freely Falling Drizzle Droplets, *J. Atmos. Sci.*, 77, 2959–2967, <https://doi.org/10.1175/JAS-D-20-0081.1>, 2020.
- Khain, A. P., Beheng, K. D., Heymsfield, A., Korolev, A., Krichak, S. O., Levin, Z., Pinsky, M., Phillips, V., Prabhakaran, T., Teller, A.,
760 van den Heever, S. C., and Yano, J.-I.: Representation of microphysical processes in cloud-resolving models: Spectral (bin) microphysics versus bulk parameterization, *Reviews of Geophysics*, 53, 247–322, <https://doi.org/https://doi.org/10.1002/2014RG000468>, 2015.
- Klemp, J. B. and Wilhelmson, R. B.: The Simulation of Three-Dimensional Convective Storm Dynamics, *J. Atmos. Sci.*, 35, 1070–1096, [https://doi.org/10.1175/1520-0469\(1978\)035<1070:TSOTDC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<1070:TSOTDC>2.0.CO;2), 1978.
- Korolev, A. and Leisner, T.: Review of experimental studies of secondary ice production, *Atmos. Chem. Phys.*, 20, 11 767–11 797,
765 <https://doi.org/10.5194/acp-20-11767-2020>, 2020.
- Kuhlman, K. M., Ziegler, C. L., Mansell, E. R., MacGorman, D. R., and Straka, J. M.: Numerically Simulated Electrification and Lightning of the 29 June 2000 STEPS Supercell Storm, *Mon. Wea. Rev.*, 134, 2734–2757, <https://doi.org/10.1175/MWR3217.1>, 2006.
- Lac, C., Chaboureaud, J.-P., Masson, V., Pinty, J.-P., Tulet, P., Escobar, J., Leriche, M., Barthe, C., Aouizerats, B., Augros, C., Aumond, P.,
770 Auguste, F., Bechtold, P., Berthet, S., Bielli, S., Bosseur, F., Caumont, O., Cohard, J.-M., Colin, J., Couvreur, F., Cuxart, J., Delautier, G., Dauhut, T., Ducrocq, V., Filippi, J.-B., Gazen, D., Geoffroy, O., Gheusi, F., Honnert, R., Lafore, J.-P., Lebeaupin Brossier, C., Libois, Q., Lunet, T., Mari, C., Maric, T., Mascart, P., Mogé, M., Molinié, G., Nuissier, O., Pantillon, F., Peyrillé, P., Pergaud, J., Perraud, E., Pianezze, J., Redelsperger, J.-L., Ricard, D., Richard, E., Riette, S., Rodier, Q., Schoetter, R., Seyfried, L., Stein, J., Suhre, K., Taufour, M., Thouron, O., Turner, S., Verrelle, A., Vié, B., Visentin, F., Vionnet, V., and Wautelet, P.: Overview of the Meso-NH model version 5.4 and its applications, *Geosci. Model Dev.*, 11, 1929–1969, <https://doi.org/10.5194/gmd-11-1929-2018>, 2018.
- 775 Lawson, R. P., Woods, S., and Morrison, H.: The Microphysics of Ice and Precipitation Development in Tropical Cumulus Clouds, *J. Atmos. Sci.*, 72, 2429–2445, <https://doi.org/10.1175/JAS-D-14-0274.1>, 2015.
- Leisner, T., Pander, T., Handmann, P., and Kiselev, A.: Secondary ice processes upon heterogeneous freezing of cloud droplets, 14th Conf. on Cloud Physics and Atmospheric Radiation, Amer. Meteor. Soc, Boston, <https://ams.confex.com/ams/14CLOUD14ATRAD/webprogram/Paper250221.html> (last access: 12 December 2024), 2014.
- 780 Leroy, D., Wobrock, W., and Flossmann, A. I.: The role of boundary layer aerosol particles for the development of deep convective clouds: A high-resolution 3D model with detailed (bin) microphysics applied to CRYSTAL-FACE, *Atmos. Res.*, 91, 62–78, <https://doi.org/10.1016/j.atmosres.2008.06.001>, 2009.
- Mansell, E. R. and Ziegler, C. L.: Aerosol Effects on Simulated Storm Electrification and Precipitation in a Two-Moment Bulk Microphysics Model, *J. Atmos. Sci.*, 70, 2032–2050, <https://doi.org/10.1175/JAS-D-12-0264.1>, 2013.
- 785 Mansell, E. R., MacGorman, D. R., Ziegler, C. L., and Straka, J. M.: Charge structure and lightning sensitivity in a simulated multicell thunderstorm, *J. Geophys. Res.: Atmos.*, 110, 2004JD005 287, <https://doi.org/10.1029/2004JD005287>, 2005.
- Marshall, T. C., McCarthy, M. P., and Rust, W. D.: Electric field magnitudes and lightning initiation in thunderstorms, *J. Geophys. Res.-Atmos.*, 100, 7097–7103, <https://doi.org/10.1029/95JD00020>, 1995.
- Naccarato, K. P., Pinto Jr., O., and Pinto, I. R. C. A.: Evidence of thermal and aerosol effects on the cloud-to-ground lightning density and
790 polarity over large urban areas of Southeastern Brazil, *Geophys. Res. Lett.*, 30, <https://doi.org/10.1029/2003GL017496>, 2003.
- Norville, K., Baker, M., and Latham, J.: A numerical study of thunderstorm electrification: Model development and case study, *J. Geophys. Res.: Atmos.*, 96, 7463–7481, <https://doi.org/10.1029/90JD02577>, 1991.
- Pereyra, R. G., Avila, E. E., Castellano, N. E., and Saunders, C. P. R.: A laboratory study of graupel charging, *J. Geophys. Res. Atmos.*, 105, 20 803–20 812, <https://doi.org/10.1029/2000JD900244>, 2000.

- 795 Phillips, V. T. J. and Patade, S.: Multiple Environmental Influences on the Lightning of Cold-Based Continental Convection. Part II: Sensitivity Tests for Its Charge Structure and Land–Ocean Contrast, *J. Atmos. Sci.*, 79, 263–300, <https://doi.org/10.1175/JAS-D-20-0234.1>, 2022.
- Phillips, V. T. J., Donner, L. J., and Garner, S. T.: Nucleation Processes in Deep Convection Simulated by a Cloud-System-Resolving Model with Double-Moment Bulk Microphysics, *J. Atmos. Sci.*, 64, 738–761, <https://doi.org/10.1175/JAS3869.1>, 2007.
- 800 Phillips, V. T. J., DeMott, P. J., and Andronache, C.: An Empirical Parameterization of Heterogeneous Ice Nucleation for Multiple Chemical Species of Aerosol, *J. Atmos. Sci.*, 65, 2757–2783, <https://doi.org/10.1175/2007JAS2546.1>, 2008.
- Phillips, V. T. J., Demott, P. J., Andronache, C., Pratt, K. A., Prather, K. A., Subramanian, R., and Twohy, C.: Improvements to an Empirical Parameterization of Heterogeneous Ice Nucleation and Its Comparison with Observations, *J. Atmos. Sci.*, 70, 378–409, <https://doi.org/10.1175/JAS-D-12-080.1>, 2013.
- 805 Phillips, V. T. J., Yano, J.-I., Formenton, M., Ilotoviz, E., Kanawade, V., Kudzotsa, I., Sun, J., Bansemer, A., Detwiler, A. G., Khain, A., and Tessendorf, S. A.: Ice Multiplication by Breakup in Ice–Ice Collisions. Part II: Numerical Simulations, *J. Atmos. Sci.*, 74, 2789–2811, <https://doi.org/10.1175/JAS-D-16-0223.1>, 2017a.
- Phillips, V. T. J., Yano, J.-I., and Khain, A.: Ice Multiplication by Breakup in Ice–Ice Collisions. Part I: Theoretical Formulation, *J. Atmos. Sci.*, 74, 1705–1719, <https://doi.org/10.1175/JAS-D-16-0224.1>, 2017b.
- 810 Phillips, V. T. J., Patade, S., Gutierrez, J., and Bansemer, A.: Secondary Ice Production by Fragmentation of Freezing Drops: Formulation and Theory, *J. Atmos. Sci.*, 75, 3031 – 3070, <https://doi.org/10.1175/JAS-D-17-0190.1>, 2018.
- Pinty, J. P. and Jabouille, P.: A mixed-phase cloud parameterization for use in mesoscale non hydrostatic model: simulations of a squall line and of orographic precipitations, in: *Proceedings of Conf. on Cloud Physics*, p. 217–220, Amer. Meteor. Soc Everett, WA, <https://worldcat.org/oclc/39920280>, 1998.
- 815 Proestakis, E., Kazadzis, S., Lagouvardos, K., Kotroni, V., and Kazantzidis, A.: Lightning activity and aerosols in the Mediterranean region, *Atmos. Res.*, 170, 66–75, <https://doi.org/10.1016/j.atmosres.2015.11.010>, 2016.
- Qu, Z., Korolev, A., Milbrandt, J. A., Heckman, I., Huang, Y., McFarquhar, G. M., Morrison, H., Wolde, M., and Nguyen, C.: The impacts of secondary ice production on microphysics and dynamics in tropical convection, *Atmos. Chem. Phys.*, 22, 12 287–12 310, <https://doi.org/10.5194/acp-22-12287-2022>, 2022.
- 820 Reynolds, S. E., Brook, M., and Gourley, M. F.: Thunderstorm charge separation, *J. Atmos. Sci.*, 14, 426 – 436, [https://doi.org/10.1175/1520-0469\(1957\)014<0426:TCS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1957)014<0426:TCS>2.0.CO;2), 1957.
- Rose, C., Collaud Coen, M., Andrews, E., Lin, Y., Bossert, I., Lund Myhre, C., Tuch, T., Wiedensohler, A., Fiebig, M., Aalto, P., Alastuey, A., Alonso-Blanco, E., Andrade, M., Artñano, B., Arsov, T., Baltensperger, U., Bastian, S., Bath, O., Beukes, J. P., Brem, B. T., Bukowiecki, N., Casquero-Vera, J. A., Conil, S., Eleftheriadis, K., Favez, O., Flentje, H., Gini, M. I., Gómez-Moreno, F. J., Gysel-Beer, M., Hallar, A. G., Kalapov, I., Kalivitis, N., Kasper-Giebl, A., Keywood, M., Kim, J. E., Kim, S.-W., Kristensson, A., Kulmala, M., Lihavainen, H., Lin, N.-H., Lyamani, H., Marinoni, A., Martins Dos Santos, S., Mayol-Bracero, O. L., Meinhardt, F., Merkel, M., Metzger, J.-M., Mihalopoulos, N., Ondracek, J., Pandolfi, M., Pérez, N., Petäjä, T., Petit, J.-E., Picard, D., Pichon, J.-M., Pont, V., Putaud, J.-P., Reisen, F., Sellegri, K., Sharma, S., Schauer, G., Sheridan, P., Sherman, J. P., Schwerin, A., Sohmer, R., Sorribas, M., Sun, J., Tulet, P., Vakkari, V., Van Zyl, P. G., Velarde, F., Villani, P., Vratolis, S., Wagner, Z., Wang, S.-H., Weinhold, K., Weller, R., Yela, M., Zdimal, V., and Laj, P.: Seasonality of the particle number concentration and size distribution: a global analysis retrieved from the network of Global Atmosphere Watch (GAW) near-surface observatories, *Atmos. Chem. Phys.*, 21, 17 185–17 223, <https://doi.org/10.5194/acp-21-17185-2021>, 2021.

- Rosenfeld, D.: TRMM observed first direct evidence of smoke from forest fires inhibiting rainfall, *Geophys. Res. Lett.*, 26, 3105–3108, <https://doi.org/10.1029/1999GL006066>, 1999.
- Rosenfeld, D., Lohmann, U., Raga, G. B., O'Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A., and Andreae, M. O.: Flood or Drought: How
835 Do Aerosols Affect Precipitation?, *Science*, 321, 1309–1313, <https://doi.org/10.1126/science.1160606>, 2008.
- Saunders, C. P. R. and Brooks, I. M.: The effects of high liquid water content on thunderstorm charging, *J. Geophys. Res-Atmos.*, 97, 14 671–14 676, <https://doi.org/10.1029/92JD01186>, 1992.
- Saunders, C. P. R. and Peck, S. L.: Laboratory studies of the influence of the rime accretion rate on charge transfer during crystal/graupel collisions, *J. Geophys. Res-Atmos.*, 103, 13 949–13 956, <https://doi.org/10.1029/97JD02644>, 1998.
- 840 Saunders, C. P. R., Keith, W. D., and Mitzeva, R. P.: The effect of liquid water on thunderstorm charging, *J. Geophys. Res.: Atmos.*, 96, 11 007–11 017, <https://doi.org/10.1029/91JD00970>, 1991.
- Saunders, C. P. R., Bax-Norman, H., Emersic, C., Avila, E. E., and Castellano, N. E.: Laboratory studies of the effect of cloud conditions on graupel/crystal charge transfer in thunderstorm electrification, *Q. J. R. Meteorol. Soc.*, 132, 2653–2673, <https://doi.org/10.1256/qj.05.218>, 2006.
- 845 Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part 1: Model description, *Meteor. Atmos. Phys.*, 92, 45–66, <https://doi.org/10.1007/s00703-005-0112-4>, 2006.
- Shi, Z., Wang, H., Tan, Y., Li, L., and Li, C.: Influence of aerosols on lightning activities in central eastern parts of China, *Atmos. Sci. Lett.*, 21, e957, <https://doi.org/10.1002/asl.957>, 2020.
- Skamarock, W. C., Powers, J. G., Barth, M., Dye, J. E., Matejka, T., Bartels, D., Baumann, K., Stith, J., Parrish, D. D., and Hubler, G.: Nu-
850 merical simulations of the July 10 Stratospheric-Tropospheric Experiment: Radiation, Aerosols, and Ozone/Deep Convection Experiment convective system: Kinematics and transport, *J. Geophys. Res-Atmos.*, 105, 19 973–19 990, <https://doi.org/10.1029/2000JD900179>, 2000.
- Straka, J. M. and Mansell, E. R.: A Bulk Microphysics Parameterization with Multiple Ice Precipitation Categories, *J. Appl. Meteor.*, 44, 445 – 466, <https://doi.org/10.1175/JAM2211.1>, 2005.
- Sullivan, S. C., Hoose, C., Kiselev, A., Leisner, T., and Nenes, A.: Initiation of secondary ice production in clouds, *Atmos. Chem. Phys.*, 18, 1593–1610, <https://doi.org/10.5194/acp-18-1593-2018>, 2018.
- 855 Sun, M., Liu, D., Qie, X., Mansell, E. R., Yair, Y., Fierro, A. O., Yuan, S., Chen, Z., and Wang, D.: Aerosol effects on electrification and lightning discharges in a multicell thunderstorm simulated by the WRF-ELEC model, *Atmos. Chem. Phys.*, 21, 14 141–14 158, <https://doi.org/10.5194/acp-21-14141-2021>, 2021.
- Sun, M., Qie, X., Mansell, E. R., Liu, D., Yair, Y., Fierro, A. O., Yuan, S., and Lu, J.: Aerosol Impacts on Storm Electrifica-
860 tion and Lightning Discharges Under Different Thermodynamic Environments, *J. Geophys. Res. Atmos.*, 128, e2022JD037 450, <https://doi.org/https://doi.org/10.1029/2022JD037450>, 2023.
- Takahashi, C.: Deformations of Frozen Water Drops and Their Frequencies, *J. Meteorol. Soc. Jpn. Ser. II*, 53, 402–411, https://doi.org/10.2151/jmsj1965.53.6_402, 1975.
- Takahashi, T.: Riming Electrification as a Charge Generation Mechanism in Thunderstorms, *J. Atmos. Sci.*, 35, 1536–1548, [https://doi.org/10.1175/1520-0469\(1978\)035<1536:REACG>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<1536:REACG>2.0.CO;2), 1978.
- 865 Takahashi, T.: Thunderstorm Electrification—A Numerical Study, *J. Atmos. Sci.*, 41, 2541–2558, [https://doi.org/10.1175/1520-0469\(1984\)041<2541:TENS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1984)041<2541:TENS>2.0.CO;2), 1984.
- Takahashi, T., Sugimoto, S., Kawano, T., and Suzuki, K.: Riming Electrification in Hokuriku Winter Clouds and Comparison with Laboratory Observations, *J. Atmos. Sci.*, 74, 431–447, <https://doi.org/10.1175/JAS-D-16-0154.1>, 2017.

- 870 Tan, Y., Shi, Z., Chen, Z., Peng, L., Yang, Y., Guo, X., and Chen, H.: A numerical study of aerosol effects on electrification of thunderstorms, *J. Atmos. Sol-Terr. Phys.*, 154, 236–247, <https://doi.org/10.1016/j.jastp.2015.11.006>, 2017.
- Taufour, M., Pinty, J.-P., Barthe, C., Vié, B., and Wang, C.: LIMA (v2.0): A full two-moment cloud microphysical scheme for the mesoscale non-hydrostatic model Meso-NH v5-6, *Geosci. Model Dev.*, 17, 8773–8798, <https://doi.org/10.5194/gmd-17-8773-2024>, 2024.
- Thornton, J. A., Virts, K. S., Holzworth, R. H., and Mitchell, T. P.: Lightning enhancement over major oceanic shipping lanes, *Geophys. Res. Lett.*, 44, 9102–9111, <https://doi.org/10.1002/2017GL074982>, 2017.
- 875 Tsenova, B., Barthe, C., Mitzeva, R., and Pinty, J.-P.: Impact of parameterizations of ice particle charging based on rime accretion rate and effective water content on simulated with MésóNH thunderstorm charge distributions, *Atmos. Res.*, 128, 85–97, <https://doi.org/10.1016/j.atmosres.2013.03.011>, 2013.
- van den Heever, S. C., Carrió, G. G., Cotton, W. R., DeMott, P. J., and Prenni, A. J.: Impacts of Nucleating Aerosol on Florida Storms. Part I: Mesoscale Simulations, *J. Atmos. Sci.*, 63, 1752–1775, <https://doi.org/10.1175/JAS3713.1>, 2006.
- 880 Vardiman, L.: The Generation of Secondary Ice Particles in Clouds by Crystal–Crystal Collision, *J. Atmos. Sci.*, 35, 2168–2180, [https://doi.org/10.1175/1520-0469\(1978\)035<2168:TGOSIP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2168:TGOSIP>2.0.CO;2), 1978.
- Vié, B., Pinty, J.-P., Berthet, S., and Leriche, M.: LIMA (v1.0): A quasi two-moment microphysical scheme driven by a multimodal population of cloud condensation and ice freezing nuclei, *Geosci. Model Dev.*, 9, 567–586, <https://doi.org/10.5194/gmd-9-567-2016>, 2016.
- 885 Wang, H., Tan, Y., Shi, Z., Yang, N., and Zheng, T.: Diurnal differences in the effect of aerosols on cloud-to-ground lightning in the Sichuan Basin, *Atmos. Chem. Phys.*, 23, 2843–2857, <https://doi.org/10.5194/acp-23-2843-2023>, 2023.
- Yang, J., Huang, S., Yang, T., Zhang, Q., Deng, Y., and Liu, Y.: Impact of ice multiplication on the cloud electrification of a cold-season thunderstorm: a numerical case study, *Atmos. Chem. Phys.*, 24, 5989–6010, <https://doi.org/10.5194/acp-24-5989-2024>, 2024.
- Yang, Y., Sun, J., Zhu, Y., and Zhang, T.: Examination of the impacts of ice nuclei aerosol particles on microphysics, precipitation and electrification in a 1.5D aerosol-cloud bin model, *J. Aerosol Sci.*, 140, 105 440, <https://doi.org/10.1016/j.jaerosci.2019.105440>, 2020.
- 890 Yano, J.-I. and Phillips, V. T. J.: Ice–Ice Collisions: An Ice Multiplication Process in Atmospheric Clouds, *J. Atmos. Sci.*, 68, 322–333, <https://doi.org/10.1175/2010JAS3607.1>, 2011.
- Zhang, Y., Fan, J., Li, Z., and Rosenfeld, D.: Impacts of cloud microphysics parameterizations on simulated aerosol–cloud interactions for deep convective clouds over Houston, *Atmospheric Chemistry and Physics*, 21, 2363–2381, <https://doi.org/10.5194/acp-21-2363-2021>, 2021.
- 895 Zhao, X. and Liu, X.: Primary and secondary ice production: interactions and their relative importance, *Atmos. Chem. Phys.*, 22, 2585–2600, <https://doi.org/10.5194/acp-22-2585-2022>, 2022.
- Ziegler, C. L., MacGorman, D. R., Dye, J. E., and Ray, P. S.: A model evaluation of noninductive graupel-ice charging in the early electrification of a mountain thunderstorm, *J. Geophys. Res-Atmos.*, 96, 12 833–12 855, <https://doi.org/10.1029/91JD01246>, 1991.




Figure 1. Thickness of the warm, mixed and cold-phase regions of the WARM (a), MID-WARM (b) and COLD (c) storms during the cloud electrification period.




Figure 2. Normalized total number of flashes in colors (first line, a-c) and normalized time of the first flash (colors) (second line, d-f) as a function of N_{CCN} and N_{INP} , for the WARM (left column), MID-WARM (middle column) and COLD (right column) simulations. The normalization is obtained by subtracting the minimum and dividing by the difference between the maximum and minimum. The text in each grid box corresponds to the total number of lightning flashes (first line) and the time of the first flash in minutes (second line) in the corresponding simulation.




Figure 3. Non-inductive charge separation rate on graupel and ice crystals averaged in the convective zone during cloud electrification as a function of N_{CCN} , for $N_{INP} = 1000 L^{-1}$ (top line), $N_{INP} = 100 L^{-1}$ (middle line) and low N_{INP} (bottom line) of the WARM (left column), MID-WARM (center column), and COLD (right column) simulations. A positive (negative) value corresponds to a positive (negative) charge gained by the graupel particle after collision with an ice crystal.

Mean vertical profiles of cloud water content (CWC, in-) in the convective region during cloud electrification as a function of N_{CCN} , for $N_{INP} = 1000$ (top line), $N_{INP} = 100$ (middle line) and low N_{INP} (bottom line) of the WARM (left column), MID-WARM (center column), and COLD (right column) simulations. The 0, -10, -20, -30, and -40 isotherms are plotted with black dashed lines:




Figure 4. Mean vertical profiles of ice crystal number concentration (L^{-1}) for $N_{INP} = 1000 L^{-1}$ (top line), $N_{INP} = 100 L^{-1}$ (middle line) and low N_{INP} (bottom line) in the convective region during cloud electrification of the WARM (left column), MID-WARM (center column), and COLD (right column) cases. In each panel, the blue, black, green, orange and pink curves correspond to the mean vertical profiles of ice crystal number concentration for $N_{CCN} = 500, 1000, 5000, 8000$ and $10,000 cm^{-3}$, respectively. The $0^{\circ}C$ and $-40^{\circ}C$ isotherms are plotted with black dashed lines.




Figure 5. Normalized tendencies of the three ice production processes summed on the vertical (colors) of the WARM (left column), MID-WARM (center column) and COLD (right column) simulations : heterogeneous nucleation (first line), homogeneous nucleation (middle line) and Hallett-Mossop process (bottom line). The text in each grid box corresponds to the ice production processes tendencies summed on the vertical ($\times 10^9 s^{-1}$) in the corresponding simulation.




Figure 6. Mean vertical profiles of cloud water content (CWC, in gm^{-3}) in the convective region during cloud electrification as a function of N_{CCN} , for $N_{\text{INP}} = 1000 \text{ L}^{-1}$ (top line), $N_{\text{INP}} = 100 \text{ L}^{-1}$ (middle line) and low N_{INP} (bottom line) of the WARM (left column), MID-WARM (center column), and COLD (right column) simulations. The 0°C , -10°C , -20°C , -30°C , and -40°C isotherms are plotted with black dashed lines.




Figure 7. Normalized total mass of graupel (colors) in the convective zone during cloud electrification between $0^{\circ}C$ and $-40^{\circ}C$ in the WARM (a), MID-WARM (b) and COLD (c) simulations as a function of N_{INP} and N_{CCN} . The text in each grid box corresponds to the total mass of graupel ($\times 10^8 kg$) in the corresponding simulation.




Figure 8. Normalized total number of flashes (a) and normalized time of the first flash (b), in colors, as a function of the SIP processes activated (NOSIP, HM, HM+CIBU, HM+RDSF, ALLSIP) and of the storm type (WARM, MID-WARM, COLD). The number in each grid box corresponds to (a) the total number of lightning flashes and (b) to the time of the first flash (min).




Figure 9. Non-inductive charge separation rate between graupel and ice crystals summed in the convective zone during cloud electrification as a function of the SIP processes activated (NOSIP, HM, HM+CIBU, HM+RDSF, ALLSIP) and of the storm type (WARM, MID-WARM, COLD). A positive (negative) value corresponds to a positive (negative) charge gained by the graupel particle after collision with an ice crystal.




Figure 10. Normalized ~~tendencies~~-ice production rate of the SIP processes summed on the vertical (colors) of the (a) WARM, (b) MID-WARM and (c) COLD cases for all sensitivity tests about SIP processes. The text in each grid box corresponds to the ice production processes tendencies summed on the vertical ($\times 10^9 \text{ s}^{-1}$) in the corresponding simulation.




Figure 11. Mean vertical profiles of ice crystal number concentration (L^{-1}) in the convective region during cloud electrification of the (a) WARM, (b) MID-WARM, and (c) COLD cases. In each panel, the blue, black, green, orange and pink curves correspond to the mean vertical profiles of ice crystal number concentration for the NOSIP, HM, HM+RDSF, HM+CIBU and ALLSIP tests, respectively. The 0 and -40°C isotherms are plotted with black dashed lines.




Figure 12. Mean vertical profiles of CWC (g.m^{-3}) in the convective region during cloud electrification of the WARM (1a), MID-WARM (b), and COLD (c) simulations. In each panel, the blue, black, green, orange and pink curves correspond to the mean vertical profiles of CWC for each SIP sensitivity tests. The 0°C , -10°C , -20°C , -30°C , and -40°C isotherms are plotted with black dashed lines.




Figure 13. Total mass of graupel in the convective zone between -40°C and 0°C in the WARM (a), MID-WARM (b) and COLD (c) simulation. The text in each grid box corresponds to the total mass of graupel ($\times 10^8$ kg) in the corresponding simulation.