

Plant community composition controls spatial variation in year-round methane fluxes in a boreal rich fen

Eeva Järvi-Laturi¹, Teemu Tahvanainen², Eero Koskinen³, Efrén López-Blanco^{4,5}, Juho Lämsä³, Hannu Marttila¹, Mikhail Mastepanov^{3,4}, Riku Paavola³, Maria Väisänen⁶, Torben R. Christensen^{1,4}

¹Water, Energy and Environmental Engineering Research Unit, University of Oulu, Oulu, 90570, Finland

²Department of Environmental and Biological Sciences, University of Eastern Finland, Joensuu, 80100, Finland

³Oulanka Research Station, University of Oulu, Kuusamo, 93900, Finland

⁴Department of Ecoscience – Arctic Ecosystem Ecology, Aarhus University, Roskilde, 4000, Denmark

⁵Department of Environment and minerals, Greenland Institute of Natural Resources, Nuuk, 3900, Greenland

⁶Ecology and Genetics Research Unit, University of Oulu, Oulu, 90570, Finland

Correspondence to: Eeva Järvi-Laturi (eeva.jarvi-laturi@oulu.fi)

Abstract Climate change is expected to impact the methane (CH₄) budget of boreal peatlands, highlighting the need to understand the factors that influence methane cycling, including plant community structure. In northern peatlands, the majority of methane is transported through plants, and the magnitude of this process is strongly linked to plant community composition. Therefore, detailed information about the role of plants regulating year-round methane fluxes is highly valuable. This paper explores the causes of spatial variability in plot-scale methane fluxes in a northern boreal rich fen. Methane fluxes were measured using the manual chamber technique in the context of fine-scale biomass variations in plant community compositions from 36 study plots over 232 days throughout a full year. The mean methane flux rates for snow-free and snow seasons were 2.55 and 0.21 mg CH₄/m²/h, respectively. We found a significant correlation between methane fluxes and a plant community type associated with the presence of sedge *Carex rostrata* during three studied periods: year-round, snow-free and snow season. *C. rostrata* was present at 13 out of 36 plots, and these 13 plots contributed 44–49 % of the total measured methane fluxes. The biomass of vascular plants, sedges, and *C. rostrata*, as well as the ratio of vascular plant to bryophyte biomass, also significantly correlated with methane fluxes during year-round and snow-free seasons. By identifying vegetation-driven emission hotspots, these results can enhance efforts to upscale emission predictions and improve ecosystem-scale methane modeling. Thus, our findings provide valuable insights for predicting realistic future changes in peatland methane emissions throughout the year.

1 Introduction

Northern peatlands are an intrinsic part of the global carbon cycle and currently, these peatlands store more than a third of all terrestrial carbon, act as strong sinks of carbon dioxide (CO₂) and are among the main natural terrestrial source of methane (CH₄) (Ramage et al., 2024; Schuur et al., 2022). Indeed, wetlands produce about 25 % of the total methane emissions that are

still considered an important source of uncertainty in the global methane budget (Saunois et al., 2024). The uncertainty arises from several factors: the relative contributions of methane emissions from tropical and northern wetlands, how these regions respond to rising temperatures, and the spatiotemporal patterns of the emissions (Christensen, 2024; Yuan et al., 2024).

Climate change is predicted to affect the hydrology of peatlands by increasing the water table depth (WTD) (Evans et al., 2021; Helbig et al., 2020; Swindles et al., 2019), which is one of the most well-known regulators of methane fluxes along with temperature and vegetation (Turetsky et al., 2014). According to several studies, an increased WTD would decrease methane fluxes (e.g. Pearson et al., 2015; Riutta et al., 2020) and increase the rate of decomposition and soil CO₂ emissions (Ma et al., 2022). These ecosystem-level processes are complex, though, as the increasing level of atmospheric CO₂ is predicted to enhance plant productivity (Forkel et al., 2016) and thereby the rate of root exudation (Nielsen et al., 2017) that via priming leads to greater methane emissions (Turner et al., 2020; Waldo et al., 2019). The expected rise in methane production could be balanced by increased methane oxidation in the topsoil, which is a probable response to enhanced microbial activity and oxygen availability caused by rising temperatures (Zhang et al. 2021). Warming climate may, however, also increase the areal cover of wet fens in the Arctic region due to permafrost thaw, which could potentially create new sources for methane release (e.g. Christensen et al., 2023; Grimes et al., 2024).

Vegetation type and responses to environmental changes are highly relevant for methane flux dynamics, as up to 90 % of ecosystem-level methane is transported through plants in northern peatlands (Ge et al., 2023; Korrensalo et al., 2022). The aerenchymatous tissues of certain vascular plant species allow methane to move from anoxic soil through the plant, thus avoiding the oxidation in oxic soil layers (Ge et al., 2023; Joabsson et al., 1999) and, indeed, plant species and their specific traits have been found to reliably predict methane flux rates (Korrensalo et al., 2022). *Sphagnum* mosses of wet environments can also host methanotrophic microbes and thus have a potential to oxidize methane and affect the magnitude of the total emissions (Larmola et al., 2010). Climate change is predicted to accelerate the natural vegetational succession in boreal rich fens towards *Sphagnum*-dominated plant communities even in stable hydrological conditions (Kolari et al., 2021), which could have major impacts on methane production and release. To improve our understanding and the predictions of future methane emissions, it is important to have more focus on the vegetation composition and the specific plant species controlling the magnitude of the fluxes (Riutta et al., 2020).

The relationship between plant community composition and methane fluxes remains an important topic of study (e.g. Lai et al., 2014; Riutta et al., 2007; Ström et al., 2015) and research on individual plant species has shown significant variation in the magnitude of flux rates and transport efficiencies (e.g. Bhullar et al., 2013; Koelbener et al., 2010; Korrensalo et al., 2022). However, year-round methane flux data from northern boreal rich fens at the plot scale are limited. This study aims to better understand the causes behind local spatial variability of methane fluxes year-round, and to provide a new perspective on assessing plant-mediated methane emissions. To this end, we focus on fine-scale variations in plant community composition based on species' biomass, using non-destructive in situ methods. We intend to answer the following questions: (1) Do methane flux variations correlate with plant community type at a study plot scale? (2) Does plant community composition correlate with methane flux variability alone or in combination with other environmental factors? We hypothesize that (1) the plant

community composition affects the methane flux and that (2) the flux is highest on study plots with largest biomass of vascular plants in absolute terms or in proportion to the biomass of bryophytes.

2 Materials and methods

2.1 Study site

70 This study was implemented in Puukkosuo, an open and slightly sloping calcareous fen located in the northern boreal zone at Oulanka National Park in Kuusamo, Northeast Finland (66.377299° N, 29.308062° E) (Fig. 1). The long-term (1992–2022) mean annual, January and July temperatures were 0.6, -13.0, and 15.3 °C, respectively, and the mean annual precipitation was 557.4 mm. The study period (19.10.2021–31.10.2022) was slightly warmer and drier than the long-term mean with temperatures 0.9, -11.4, and 16.5 °C, respectively, and with total annual precipitation of 528.8 mm. During the snow-free
75 season of 2022, the mean pH (n=179) measured from peat porewater at approximately 10 cm depth, was 7.0, ranging from 6.74 to 7.38. The deepest measured water table was 9.3 cm below the peat surface, while the highest was 7.0 cm above the surface (Fig. A1). The plot-scale variation in WTD during the snow-free season of 2022 was 3.8–9.1 cm with an average of 6.5 cm. The vegetation at the site is dominated by vascular plants typical of rich fens (*Carex* spp., *Trichophorum* spp., *Molinia caerulea*, *Potentilla erecta*, *Menyanthes trifoliata*), as well as brown mosses (*Scorpidium cossonii*, *Campylium stellatum*,
80 *Cinclidium stygiym*) and peat mosses (*Sphagnum* spp., mostly *S. warnstorffii*).

2.2 Experimental design

The study area was approximately one hectare in size and included 12 spatial blocks. Each block had three study plots resulting in 36 study plots. The plots were established in summer 2018, and the size of a plot was 2 m × 3.5 m which included a 0.5 m wide buffer zone. Wooden boardwalks, built to minimize stepping on the surface of the peat, led to the plots. Half of the study
85 plots (n=18) were located inside a fence, built in spring 2019, to exclude grazing by reindeer (Fig. 1). At the time of this study, the exclusion had lasted for 2–3 years. The location of the plots and the fence followed the hydrological gradient of the fen (Fig. 1). The study plots were also assigned to snow level manipulations that were started in January 2019. Within each block, one plot was an untreated control with ambient snow level, one a snow removal plot where the snow depth was maintained at 0.25 m throughout the snow season, and one plot a snow addition plot, where the snow from the removal plot was placed. The
90 snow treatments had no statistically significant effects on the methane fluxes (Fig. B6).




Figure 1. A map of Puukkosuo rich fen showing the location of boardwalks, experimental area and the fence excluding reindeer. The small inset shows the location of the study area, Oulanka, in NE Finland. Orthomosaic © Petra Korhonen 2024.

2.3 Methane flux measurements

95 For methane flux measurements, a round PVC collar (inner diameter 29.5 cm) was inserted approximately 5 cm into the ground in the rear end of each plot in September 2020. During 19.10.2021–31.10.2022, we measured methane fluxes ($\text{mg CH}_4/\text{m}^2/\text{h}$) over 232 individual days between 8 am and 6 pm, conducting one measurement per plot per day. On most days, only half of the plots were measured using randomized plot selection. We used manual, closed chamber technique (e.g. Christensen et al., 2000) with a portable LI-COR $\text{CH}_4/\text{CO}_2/\text{H}_2\text{O}$ Trace Gas Analyzer (LI-7810) and a transparent polycarbonate chamber (height 100 38 cm, diameter 29 cm) equipped with a small fan to circulate the air inside the chamber (Fig. 2a). For each measurement during snow-free season, the chamber was placed on the PVC collar for an airtight seal (Fig. 2b). Each measurement lasted for 5 minutes. During winter, when snow covered the collars and it was not possible to place the chamber directly on them, we measured the fluxes on top of the snowpack, also known as floating chamber technique, to avoid disturbance to the snow and methane diffusion (Björkman et al., 2010) and consequently to the experimental plots. In comparison to the snow-free 105 season, these measurements were taken at a slightly different spot closer to the boardwalks. Due to these limitations, snow season flux measurements should be examined with caution. Altogether, 4121 individual measurements were used in this study.

To calculate methane flux rates for each plot, we used a Python script that calculated the slope of methane concentration change during three centremost minutes of a five-minute measurement period and computed the flux in $\text{mg CH}_4/\text{m}^2/\text{h}$ using ambient air pressure and air temperature at the time of each measurement (linear regression model, e.g. Pirk 110

et al., 2016a). We accepted the measurements with an R^2 value ≥ 0.95 ($n = 3589$) and inspected all the rest ($n = 691$) individually, leaving out measurements showing very strong non-linearity or any other sign of failed measurement ($n = 159$). We examined the fluxes in three periods: 1) year-round (19.10.2021–31.10.2022), 2) snow-free (13.5.–26.10.2022), and 3) snow season (19.10.2021–12.5.2022 and 27.10.–31.10.2022). Snow-free season presented a period when we were able to measure fluxes on the collar (part of the experimental plot or the experimental site could be covered with snow). Snow covered season presented a season when we measured the fluxes on top of the snow, not on the collar. Annual accumulated flux (1.11.2021–31.10.2022) was estimated by calculating a 24-hour accumulated flux for each available datapoint by multiplying the hourly mean flux by 24. These daily flux values were then summed to obtain the annual total. The days which were missing a measurement were given the value from a previous measurement, assuming the fluxes did not vary remarkably diurnally or over the days.




Figure 2. A flux measurement carried out with (a) a portable LI-COR Trace Gas Analyzer and (b) a clear polycarbonate chamber placed on a round collar installed at each plot in 2020. Photos by Eeva Järvi-Laturi.

2.4 Plant community data

We studied the plant communities by identifying species and their abundance within each collar, collecting separate biomass samples from outside of the main research area, and using the mean biomass (BM) value of each species' samples to determine the community composition within each collar. Plant surveys and sample collection were done during 25.7.–12.8.2022.

To survey the vascular plant (VP) species within the collars, we counted each aboveground shoot individually and separated them into flowering and non-flowering categories, as the flowering shoots may hold higher biomass than the non-flowering ones. Altogether, 31 vascular plant species were identified from the study site. We collected separate biomass

samples of each identified species. The biomass sampling was randomized by first selecting an area where vegetation heights resembled the heights of the vegetation within the collars. Then, we randomly threw a marker and selected the first ten non-flowering individuals of target vascular plant species close to the marker. Additionally, we collected ten flowering individuals for those species, which were found flowering within the collars. Shoots were cut at the peat surface. For uncommon species (*Angelica sylvestris*, *Carex dioica*, *Carex panicea*, *Dactylorhiza* sp., *Drosera* sp., *Eriophorum angustifolium*, *Festuca ovina*, *Pinguicula* sp., *Saussurea alpina* and *Viola epipsila*), randomization could not be put into practice. For these, samples were collected from where they could be found.

For bryophytes, we visually estimated the percentage cover of each species within each collar. Altogether, 10 bryophyte species were identified from the study site. We took either three samples representing 5 % (diam. 6.6 cm) or one sample representing 1 % (diam. 2.95 cm) of the collar area with small plastic jars. The larger sample size was collected for species that could easily be found in pure patches (*Aulacomnium palustre*, *Campylium stellatum*, *Scorpidium cossonii*, *Sphagnum* spp. and *Tomentypnum nitens*), and the smaller sample size for the remaining, more scattered and sparse species (*Aneura pinguis*, *Cinclidium stygium*, *Fissidens adianthoides*, *Mesoptychia rutheana* and *Paludella squarrosa*). The sampling locations were selected so that the target bryophyte species could be found as “pure monoculture” as possible. We then cleaned the samples of other species and litter, removed the non-living parts, and included only the colorful or leafy parts to represent the aboveground biomass.

For BM estimation, we dried the samples (n=454) in a hot air circulation oven at 40 °C for minimum of two days and weighed them with a four decimal precision (Denver Instrument SI-234, Table A1). The dry weights were normalized either by shoot (vascular plant, g/shoot) or by cover percent (bryophyte, g/1 %). For each vascular plant species, we used the total number of shoots, separated by flowering and non-flowering categories, within a collar, multiplied by the mean dry mass per shoot to obtain total species BM for each collar. For each bryophyte species, we used the total percentage cover within a collar, multiplied by the mean dry mass per 1 % area to obtain total species BM for each collar. We also calculated a VP to bryophyte BM ratio for each collar using these BM values (Table A2). We acknowledge that studying the plant communities using non-destructive methods and a mean BM of the separate samples, instead of the actual BM of the plant communities within the collars, introduces a margin of error in the calculations.

2.5 Environmental variables

We used weekly measures of WTD from June to October from regular measuring points located approximately 1 m from the methane flux measurement points. At every plot, soil temperature was recorded with 10-min intervals at 5 cm depth (model p107 CS CR1000, Campbell Scientific Inc., Logan, UT, USA). In October 2023, we measured peat layer thickness at the rear edge of each study plot with a thin, metallic 300 cm long auger. To determine peat chemistry, we collected pore-water during frost-free periods. In the beginning of snow-free season 2022, rhizon samplers (Rhizosphere Research Products, The Netherlands) were installed (10 cm depth) for pore water sampling in the middle of the collar in each experimental plot. Pore water was sampled five times during the frost-free periods (31 May, 29 June, 22 July, 28 August, 29 September). Samples

165 were collected into evacuated opaque syringes over a period of 24–48 h, filtered (0.45 µm, sterile nylon, Sarstedt, Germany) and frozen (–18 °C). Thawed samples from all sampling campaigns were analyzed for pH (913 pH/DO Meter, Metrohm), dissolved organic (DOC) and inorganic carbon (DIC; Shimadzu DOC-VCX, Trios) and dissolved organic nitrogen (DON), ammonium (NH₄⁺) and nitrite + nitrate (NO₂⁻+NO₃⁻; AA500 Seal Analytical) and for DOC and DON reported as mg/l and for NH₄⁺ and NO₂⁻+NO₃⁻ as µg/l. In analyses, we used the mean value of each measured variable. Additionally, we estimated the
170 percentage cover of litter inside the collar while identifying the vegetation.

2.6 Data analyses

To analyse the vegetation, we divided the plant communities into three species combinations: 1) all species, 2) VPs, and 3) bryophytes. We analysed these combinations separately using plant BM estimates (see 2.4) by hierarchical cluster analysis using Sorensen (Bray–Curtis) distance measure and Flexible Beta group linkage method (McCune et al., 2002) with beta-value
175 of -0.25. Clusters with less than six samples (study plots) were discarded. To evaluate which species were statistically most connected to the different clusters, we carried out an indicator species analysis separately to all three species combinations. We tested the differences between the clusters with Tukey’s Honestly Significant Difference (HSD) test following an Analysis of Variance (aov function, R Core Team, 2024) to assess the significance of clusters in relation to methane fluxes in different time periods (snow-free, year-round and snow season). With clustered data, we performed detrended correspondence analysis
180 (DCA) to identify patterns in species composition and canonical correspondence analysis (CCA) to relate the species composition to environmental variables (McCune et al., 2002). We conducted the analyses separately for VPs and bryophytes to determine the main environmental factors characterizing the composition of these plant communities. Pairwise correlations between multiple BM and environmental variables were examined with a separate CCA, using snow-free season averages. The resulting Pearson correlation matrix (Table B2) provides insights into these relationships. Finally, we conducted local
185 regression models (LOESS, locally estimated scatterplot smoothing) to explore nonlinear trends between various variables and methane fluxes in relation to VP clusters.

Cluster, indicator species and correspondence analyses were implemented with PC-ORD version 7.09 (McCune and Mefford, 2018). Regression models and Tukey’s HSD tests were performed using RStudio version 2024.4.2.764 (Posit team, 2024). Both programmes were used for data visualization: PC-ORD for cluster dendrograms, boxplots, and ordination graphs,
190 RStudio for line graphs and scatterplots. Significance was defined as $p < 0.05$. We used the following R packages: readxl v1.4.3 (Wickham and Bryan, 2023), dplyr v1.1.4 (Wickham et al., 2023a), tidyr v1.3.1 (Wickham et al., 2024), ggplot2 v3.5.1 (Wickham, 2024), forcats v1.0.0 (Wickham, 2023), scales v1.3.0 (Wickham et al., 2023b), paletteer v1.3.0 (Hvitfeldt, 2021), ggnewscale v0.5.0 (Campitelli, 2024) and viridis v0.6.5 (Garnier et al., 2024). R scripts were created with the assistance of Microsoft 365 Copilot, an AI-powered productivity tool.

3.1 Methane fluxes

During the snow-free season, methane fluxes were the highest, ranging from 0.02 to 9.17 mg CH₄/m²/h (i.e., 0.48–220 mg CH₄/m²/d), with an overall mean of 2.55 mg CH₄/m²/h (i.e., 61 mg CH₄/m²/d). During this season, the mean of plot-scale methane fluxes ranged from 0.51 to 4.67 mg CH₄/m²/h, and daily coefficient of variation spanned from 38.9 % to 85.4 %. The highest individual fluxes per plot varied between 1.51 and 9.17 mg CH₄/m²/h. The lowest flux values of the snow-free period (0.02–1.18 mg CH₄/m²/h) were measured in May after a spring burst (1.4.–12.5.2022), and at the end of the season in late October (Fig. 3). The magnitude of the spring burst differed among the plots, with maximum individual fluxes ranging from 0.15 to 6.65 mg CH₄/m²/h. During the snow season, flux measurements fluctuated between 0 and 6.65 mg CH₄/m²/h (i.e., 0–160 mg CH₄/m²/d), with an overall mean of 0.21 mg CH₄/m²/h (i.e., 5 mg CH₄/m²/d). The mean of plot-scale fluxes ranged from 0.07 to 0.56 mg CH₄/m²/h, and daily coefficient of variation spanned from 39.3% to 300.4%. Further, the snow season fluxes accounted on average for 8.2% of the estimated annual accumulated flux, with values ranging from 2.3 % up to 21.3 % across the study plots (Table B3). On a year-round scale, the overall mean flux was 1.37 mg CH₄/m²/h, with mean fluxes varying between 0.29 and 2.52 mg CH₄/m²/h, and daily coefficient of variation ranging from 38.9 % to 300.4 %.




Figure 3. Year-round methane fluxes (mg CH₄/m²/h) at Puukkosuo with average soil temperature at 5 cm depth. The grey flux range shows the overall variation of the fluxes. Smoothed conditional means (LOESS method) and data points, representing individual plot-scale flux measurements, are segregated by vascular plant clusters defined in section 3.2. The vertical dashed lines mark the start and end of the snow-free season (13.5.–26.10.2022).

3.2 Plant community structures

The plots differed in their plant community structures, with wide variation in plant BM. VP BM per plot ranged from 62.1 g/m² to 486 g/m², while the bryophyte BM ranged from 65.1 g/m² to 269 g/m². The lowest total BM for an individual plot was

167 g/m², while the highest was 674 g/m². The percentage of VPs in the total BM varied between 21 % and 72 % (Table A2). The species producing the most BM, based on combined dry mass estimates from all plots, were sedges *Carex lasiocarpa* (122 g) and *C. rostrata* (84 g), and bryophytes *Scorpidium cossonii* (193 g), *Campylium stellatum* (113 g), and *Sphagnum* spp. (92 g) (Table A3).

The cluster and indicator species analyses for all species, VPs, and bryophytes all yielded three clusters (groups of plots) with significant ($p < 0.05$) indicator species statistically connected to the clusters (Figs. B1–B3). The cluster analyses indicated that bryophytes (*Sphagnum* spp., *S. cossonii*, and *C. stellatum*) showed the strongest connection to the clusters when analyses were done with all plant species or bryophytes (Figs. B1, B3). On the other hand, when analysing VPs alone, the clusters connected mostly with different sedges (*C. rostrata* with the first, *C. chordorrhiza* with the second, and *Trichophorum cespitosum*, *C. lasiocarpa*, and *Potentilla erecta* (a forb) with the third cluster). From here on, the clusters are referred to as *C. rostrata*-cluster, *C. chordorrhiza*-cluster, and *T. cespitosum*-cluster, respectively. The characteristics of each VP cluster, including their community structure and indicator species, were studied by comparing the indicator values of the species (Table B1). The results demonstrated that the plant communities differed between the clusters, as most species were abundant in only one or two clusters, and only three species (*Campylium stellatum*, *Vaccinium oxycoccos*, and *Scorpidium cossonii*) were common in all clusters (Table B1). We also found an association between *Sphagnum* mosses and *C. rostrata* in the plant communities: the *C. rostrata*-cluster had the highest abundance of *Sphagnum* mosses among the three vascular plant clusters (Table B1). The spatial division of the vascular plant communities was clear – the *C. rostrata*-community plots were all located in the upper part of the sloping fen, while the other two clusters were more widely distributed (Fig. B7).

3.3 Clusters' relation to methane fluxes

An Analysis of Variance (ANOVA) revealed that VP clusters correlated significantly with snow-free season ($F = 10.71$, $p < 0.001$) and year-round ($F = 10.92$, $p < 0.001$) methane fluxes but not with snow season fluxes ($F = 2.14$, $p > 0.05$). Tukey's HSD test indicated that *C. rostrata*-cluster, which had the highest fluxes especially during snow-free season (Fig. 3), differed significantly from *C. chordorrhiza*- and *T. cespitosum*-clusters in snow-free ($p < 0.01$ and $p < 0.001$, respectively) and year-round periods ($p < 0.01$ and $p < 0.001$, respectively) but not in snow season (Fig. 4a). There were no significant differences between *C. chordorrhiza*- and *T. cespitosum*-clusters in any of the periods (Fig. 4a–c). All species-clusters did not correlate significantly with methane fluxes in any period ($F = 1.51$, $p > 0.2$ for snow-free, $F = 1.45$, $p > 0.2$ for year-round, and $F = 0.57$, $p > 0.5$ for snow season, Fig. 4b). Similarly, the bryophyte clusters did not correlate significantly with methane fluxes ($F = 1.23$, $p > 0.3$ for snow-free, $F = 1.26$, $p > 0.2$ for year-round, and $F = 0.90$, $p > 0.4$ for snow season, Fig. 4c).




Figure 4. Methane fluxes in snow-free season (left panel), the whole year (middle panel), and snow season (right panel), as divided by cluster analyses of (a) vascular plants (VP), (b) all species (ALL) and (c) bryophytes (BRYO). Abbreviations for the clusters and the species with highest fidelity according to indicator species analysis: C.Ros = *C. rostrata*, C.Cho = *C. chordorrhiza*, T.Ces = *C. lasiocarpa*, *P. erecta*, and *T. cespitosum*, Sco = *S. cossonii*, Sph = *Sphagnum* spp. and Cam = *C. stellatum*. Asterisks above the bars denote significant differences between the clusters at the level $p < 0.01^{**}$ and $p < 0.001^{***}$.

3.4 Ordination analyses


Within VP communities, the *C. rostrata*-cluster diverged from the *C. chordorrhiza*- and *T. cespitosum*-clusters in both detrended correspondence analysis (DCA) and canonical correspondence analysis (CCA), while the *C. chordorrhiza*- and *T. cespitosum*-clusters diverged in DCA, but not in CCA (Fig. B4a–b). In bryophyte communities, the *Sphagnum*-cluster diverged from the *S. cossonii*- and *C. stellatum*-clusters in both DCA and CCA, while the *S. cossonii*- and *C. stellatum*-clusters diverged in DCA, but not in CCA (Fig. B5a–b). The main compositional gradients of VP and bryophyte communities displayed different correlation patterns with environmental variables. The first ordination axis of the VP communities correlated with snow-free and year-round methane fluxes ($r = 0.775$ and 0.782 in DCA, $r = -0.856$ and -0.866 in CCA, respectively, Fig. B4a–b). Additionally, the first ordination axis correlated with snow season fluxes ($r = 0.445$ in DCA, -0.402 in CCA). The correlation was also significant for the ratio of VP to bryophyte BM ($r = 0.562$ in DCA, -0.730 in CCA), peat layer depth ($r = 0.342$ in DCA and -0.580 in CCA) as well as the combined nitrate (NO_3^-) and nitrite (NO_2^-) concentration ($r = 0.529$ in DCA, -0.453 in

CCA). In contrast, the strongest compositional gradient of the bryophyte data correlated with WTD and pH in both DCA ($r = -0.606$ and 0.473 , respectively) and CCA ($r = -0.614$ and 0.408 , respectively) (Fig. B5a–b). The correlation between the bryophyte communities and methane was not significant in any of the periods ($r = 0.267$, 0.262 , and 0.132 in DCA, $r = 0.316$, 0.326 , and 0.130 in CCA in snow-free, year-round, and snow season, respectively).

3.5 BM, environmental variables and methane

In the Pearson correlation matrix (Table B2), the total BM of VPs correlated with the total BM of sedges ($r = 0.98$) and the total BM of *C. rostrata* ($r = 0.93$) indicating that sedges were the main functional group of VPs, and *C. rostrata* the main VP species, producing BM in our study site. The BM variables that had a significant pairwise correlation with methane fluxes in year-round and snow-free season were the total BM of VPs ($p < 0.001$), the total BM of sedges ($p < 0.001$), the total BM of *C. rostrata* ($p < 0.001$), and the ratio of VP to bryophyte BM ($p < 0.01$) (Fig. 5a–d). Significant correlations during the snow season were not discovered. Environmental variables that had a significant pairwise correlation with methane fluxes in snow-free and year-round periods were pH ($p < 0.05$ and $p \leq 0.01$, respectively, Fig. 5e) and combined concentration of NO_3^- and NO_2^- in peat pore water ($p < 0.05$ for both periods, Fig. 5f). There was no significant correlation between methane fluxes and WTD or soil temperature in any period. All correlation coefficients are listed in a correlation matrix in Table B2.

We found that when *C. rostrata* BM was 0 g, there was large variation in the magnitude of methane flux (Fig. 5c). Therefore, we additionally examined the clusters' relation to fluxes in plots without any *C. rostrata* shoots (23 plots out of 36), as well as the relationship between methane fluxes and environmental variables using linear regression models (lm function, R Core Team, 2024). This revealed that, in the absence of *C. rostrata*, pH and litter cover were the primary environmental drivers of snow-free season methane fluxes, showing significant relationships ($p < 0.001$ and $p = 0.01$, respectively) and Pearson correlation values of -0.66 and -0.29 , respectively.



285 **Figure 5. Scatterplots showing the relationships between methane (CH₄) flux (mg/m²/h) and environmental variables: a) total biomass (BM) of vascular plants ($p < 0.001$, $r = 0.54$), b) total BM of sedges ($p < 0.001$, $r = 0.57$), c) BM of *C. rostrata* ($p < 0.001$, $r = 0.60$), d) BM ratio of vascular plants and bryophytes ($p < 0.01$, $r = 0.45$), e) pH ($p < 0.05$, $r = -0.38$), and f) NO₃+NO₂ concentration ($p < 0.05$, $r = 0.34$) across different vascular plant clusters. Vascular plant clusters are defined in section 3.2. The smoothing lines were fitted using local polynomial regression (LOESS) with a span value of 1. Biomass values are in grams (g) of dry weight.**

4 Discussion

290 4.1 Methane flux variation over time and space

During the snow-free season, methane fluxes ranged from 0.48 to 220 mg CH₄/m²/d, with an overall mean of 61 mg CH₄/m²/d. Similar fluxes have been measured from a northern boreal rich fen (Jammet et al., 2017). We observed that snow season fluxes, with a range from 0–160 mg CH₄/m²/d and an overall mean of 5 mg CH₄/m²/d, accounted for 2.3–21.3 % of the estimated annual accumulated flux (Table B3). Although we measured fluxes on top of an undisturbed snowpack, our results align with a study from fens in central Finland, where 6–17 % of the annual methane release was observed during wintertime when measuring fluxes on peat surface after clearing the snow (Alm et al., 1999). We acknowledge that direct measurements from the ground surface would have improved our ability to connect winter fluxes with the studied plant communities. However, we do not expect significant differences in flux values obtained using a closed chamber on top of the snowpack, as

methane diffusion through the snowpack may be relatively unrestricted (Alm et al., 1999; Pirk et al., 2016b). Since winter may
300 account for up to 20 % of the annual methane flux in boreal fens (this study, Alm et al., 1999), any changes in wintertime
processes may impact future methane emissions from these regions. Therefore, our results highlight the importance of
including winter in methane flux studies, as this information may help reduce the current large uncertainties in the net carbon
balance. Furthermore, the reduced variability of fluxes and the lack of significant variation among vegetation clusters during
mid-winter (Figs. 3, 4) suggest that winter fluxes could be effectively captured even with lower but more optimized sampling
305 efforts, aiding in the estimation of annual emissions (Vargas and Le, 2023).

Flux rates during the snow-free season reached up to 9.17 mg CH₄/m²/h at certain study plots, while they never
exceeded 2 mg CH₄/m²/h in some others (Fig. 3). The high BM of vascular plants, and especially presence *C. rostrata*,
contributes to the spatial variability in fluxes: all plots with more than five flux measurements exceeding 6 mg CH₄/m²/h (n =
9) contained *C. rostrata* shoots but only three out of 19 plots that never recorded fluxes above 6 mg CH₄/m²/h contained *C.*
310 *rostrata*. However, *C. rostrata* biomass cannot be the sole factor of high methane fluxes, as one plot containing no *C. rostrata*
shoots also measured fluxes up to 9.05 mg CH₄/m²/h (see 4.3). Unlike the snow-free season, the high flux values during the
spring burst at the end of the snow season could not be explained by plant biomass or any other studied environmental variable.
Other reasons for the spatial variation in flux rates may include species-specific plant traits such as rooting characteristics (Ge
et al., 2023), ecohydrological aspects such as peat water holding capacity (Zhang et al., 2020), or microbial metabolic
315 interactions such as nutrient cycling (Kujala et al., 2024; Yavitt et al., 2012), which contribute to soil conditions, substrate
availability, and microbial activity. Interestingly, the methane fluxes from *C. rostrata*-community plots were the highest from
late July to late August (Fig. 3) when vegetation at our site remained predominantly green, even though fluxes from *C. rostrata*
shoots have been reported to be the highest when leaves are senescing (Ge et al., 2024). The period from late July to late
August may coincide with the peak development of permeable root surface area (Reid et al., 2015) which extent, in turn, is a
320 key factor influencing methane transport in plants (Henneberg et al., 2012). Therefore, the seasonal changes in methane flux
rates associated with *C. rostrata* may be controlled by the belowground parts of the plant (Ge et al., 2024).

4.2 Plant communities, biomass and methane

During the growing season, vegetation is known to correlate substantially with methane fluxes (e.g. Lai et al., 2014; Riutta et
al., 2007; Ström et al., 2015), but we found significant correlations year-round supporting our first hypothesis. Thus, our results
325 indicate that the influence of plant communities on methane flux dynamics is not limited to active growing season or plant
senescence, and that some species are likely to be more efficient at supporting methane production and transport, even under
the snowpack. Indeed, continuous production and spatial variation in methane flux rates has been observed throughout the cold
season, and the variation has been associated with differences in plant community composition and the adjoining differences
in substrate quantity and quality (Pirk et al., 2016b). However, the significant correlation between vegetation and methane
330 fluxes during the snow season identified by CCA, but not by linear regression models, suggests that the relationship between
vegetation and snow season methane fluxes is complex and non-linear.

Both the total vascular plant biomass and the ratio of vascular plant to bryophyte biomass correlated significantly with methane fluxes during snow-free and year-round periods, supporting our second hypothesis (Fig. 5a, d; Table B2). There was a strong correlation between the biomass of vascular plants, sedges and *C. rostrata* (Table B2). Additionally, the biomass of vascular plants, sedges and *C. rostrata* also correlated with methane fluxes (Fig. 5a–c; Table B2). These findings suggest, similar to earlier studies (e.g. Ge et al., 2023; Korrensalo et al., 2022), that plant functional type and species largely determine the magnitude of the fluxes. High amounts of vascular plant, sedge, and *C. rostrata* biomass likely enhance methane production and release by supplying labile organic carbon substrates for methanogenesis through deep root systems throughout the year (Alm et al., 1999; Joabsson et al., 1999, Saarinen, 1996). High flux rates from *C. rostrata* dominated plots (Fig. 7) may be due to the species' high methane transport rate (Ge et al., 2023), the high porosity of its roots (Ge et al., 2023), and a low capacity to oxidize methane into CO₂ in the rhizosphere (Ström et al., 2005). Additionally, the perennial nature and deep rooting traits of *C. rostrata* (Saarinen, 1996; 1998) could support methane production and transport during the cold season by providing substrates for microbial processes in deeper peat layers and a potential pathway from belowground to the atmosphere. Approximately 40 % of *C. rostrata* shoots at our study site overwinter green (Cunow et al., unpublished data), indicating the potential to transport gases also during wintertime. Moreover, the significant correlation between the ratio of vascular plant to bryophyte biomass and methane fluxes brings a new perspective to the discussion, as previous studies have mainly focused on finding differences between single plant species (e.g. Bhullar et al., 2013; Ge et al., 2023; Koelbener et al., 2010; Korrensalo et al., 2022) or their role in a plant community (e.g. Lai et al., 2014; Riutta et al., 2007; Ström et al., 2015). This biomass ratio could be obtained remotely (e.g., multispectral imaging, Wolff et al., 2023) and potentially used as a predictive parameter for modeling peatland methane fluxes.

A majority (75 %) of the highest methane fluxes observed in this study originated from *C. rostrata* dominated plots. However, the magnitude of the fluxes was not solely dictated by *C. rostrata* biomass; the fluxes increased rapidly in a non-linear manner and only slightly with higher biomass, being relatively high and stable in most *C. rostrata*-cluster plots (Fig. 5c). As a species with high gas transport efficiency, even a single *C. rostrata* shoot can transport a substantial volume of gases – potentially the same amount as a community with multiple shoots, where gas transport is distributed among many individuals (Koelbener et al., 2010; Korrensalo et al., 2022). This division of gas transport among multiple shoots could explain the observed saturation in methane flux rates: as the biomass of *C. rostrata* increases, the total flux remains consistent regardless of the number of shoots. In our analysis, an increase in biomass corresponds to a higher number of shoots, and therefore we cannot argue that the size of individual *C. rostrata* shoots did not impact the plant's methane transport efficiency.

4.3 Environmental factors and methane

We also asked if the spatial variability of methane fluxes could be explained by the combination of plant communities and environmental variables, and found that vegetation composition was the primary driver of this variation. However, in some cases, the role of an environmental variable seemed to depend on the presence of plants. Firstly, we analyzed snow-free season methane fluxes from plots without any *C. rostrata* shoots and found that pH and litter cover emerged as main drivers of

365 methane fluxes. However, as the significance of pH lowered when including the plots with *C. rostrata*, it is likely that the significant relationship between methane fluxes and pH in snow-free and year-round periods is explained through vegetation. Secondly, when *C. rostrata* was present, NO_3^- and NO_2^- levels correlated positively with methane fluxes during snow-free and year-round periods. In plots without *C. rostrata* shoots, this significant relationship disappeared, suggesting that the positive correlation is related to the presence of the species. The positive correlation was unexpected, as these compounds usually
370 inhibit methanogenesis (e.g. Knorr & Blodau, 2009). More precisely, higher levels of NO_3^- may promote methane oxidation, leading to lower fluxes (Song et al., 2022), but the impact of these electron acceptors remains debated (Zhang et al., 2021). Thus, our findings underscore the need for further investigation into the interaction between *C. rostrata* and NO_3^- and NO_2^- in driving methane fluxes.

Water table depth is commonly thought to regulate methane fluxes in peatlands (Lai, 2009; Turetsky et al., 2014) but
375 did not correlate with methane fluxes at our site in any period. Indeed, in fens with a stable water table, such as our study site (standard deviation of WTD 1.05–2.78 cm), the depth of the water table may not control the variability of the fluxes (e.g. Ge et al., 2023). Peat layer did not show a significant pairwise correlation with methane fluxes which supports the previous findings that most of methane is produced from the fresh root litter and root exudates instead of old, recalcitrant peat (Ström et al., 2012). However, peat layer thickness correlated significantly with the first ordination axis in CCA and had a significant
380 pairwise correlation with the biomass of vascular plants, sedges, and *C. rostrata*, as well as the biomass ratio of vascular plants and bryophytes (Table B2), all of which were proxies for higher methane fluxes. This suggest that peat layer composition and depth can also impact methane production through an indirect relationship. Lastly, methane fluxes did not correlate with peat temperature at 5 cm depth. Indeed, methane fluxes in boreal rich fens associate with deeper soil temperatures, which connect to water table position, rather than with surface temperatures influenced by air temperature (Olefeldt et al., 2017). All these
385 findings highlight that vegetation, rather than environmental factors, was the main driver of methane fluxes at our site.

Conclusions

Our year-round methane flux data showed significant spatial and temporal variations in flux magnitude in a northern boreal rich fen. Plant community composition, particularly the biomass of *C. rostrata*, effectively explained the plot-scale spatial variation of methane fluxes during snow free and year-round periods. Multivariate analysis also revealed that a vascular plant
390 cluster characterized by *C. rostrata* correlated with snow season methane fluxes. In addition, the total biomass of vascular plants and the ratio of vascular plant to bryophyte biomass also showed a significant positive relationship with methane fluxes in both year-round and snow-free seasons. On the other hand, the role of other environmental factors, such as pH and nitrogen, in driving methane fluxes was subtle and seemed to depend on vegetation. Our findings suggest that the biomass ratio of vascular plants and bryophytes could be used as a parameter for predicting methane emissions at study sites with similar
395 vegetation to ours. Importantly, these findings help to predict more accurately and realistically future changes in peatland methane emissions, which is essential for estimating the potential impacts of ongoing climate change.

Appendices

Appendix A with additional site information. Appendix B with additional figures and tables of the results.

Data availability

400 Full datasets used are available on request.

Author contribution

MV, RP, MM and TRC designed the experimental setup of the study site. EJJ, TT, TRC and JL designed the study. EJJ sampled the vegetation and performed the statistical analyses with assistance from JL, EK and TT. EJJ, JL and EK were responsible for data curation. EK and MM wrote the code for the flux calculations. MV designed and provided resources for
405 sampling the water chemistry. Resource management of the data collection were overseen by RP. Manual methane flux measurements, pore water sampling and water level measurements were conducted by numerous individuals, including EJJ and EK. EJJ prepared the manuscript with contributions from all co-authors.

Competing interests

The authors declare that they have no conflict of interest.

410 Acknowledgements

Maik Bischoff, Pyry Runko, Noémie Perrier-Malette, Valentin Krieger, Polina Saarinen and Johannes Väisänen, among others, are gratefully acknowledged for their help with the fieldwork. A special thanks goes to the Oulanka Research station staff for their continuous efforts. We also acknowledge that the help of AI was used when creating the R codes. EJJ was supported by Green-Digi-Basin project funded by Research Council of Finland (RCF, 347704) and DIWA-Digital Waters flagship PhD
415 pilot program funded by Ministry of Education (RCF, 359228). ELB, MM and TRC consider this study a contribution to GreenFeedBack (Greenhouse gas fluxes and earth system feedbacks) funded by the European Union's HORIZON research and innovation program under grant agreement No 101056921. Other funds during the project came from Maa- ja vesiteknikan tuki ry (to MV). The work is part of EcoClimate experimental platform at Oulanka Research Station.

References

420 Alm, J., Saarnio, S., Nykänen, H., Silvola, J., and Martikainen, P.J.: Winter CO₂, CH₄ and N₂O fluxes on some natural and drained boreal peatlands, *Biogeochemistry*, 44, 163–186, <https://doi.org/10.1007/bf00992977>, 1999.

- Bhullar, G. S., Irvani, M., Edwards, P. J., and Olde Venterink, H.: Methane transport and emissions from soil as affected by water table and vascular plants, *BMC Ecol*, 13, 32, <https://doi.org/10.1186/1472-6785-13-32>, 2013.
- 425 Björkman, M. P., Morgner, E., Cooper, E. J., Elberling, B., Klemetsson, L., and Björk, R. G.: Winter carbon dioxide effluxes from Arctic ecosystems: An overview and comparison of methodologies, *Global Biogeochemical Cycles*, 24, <https://doi.org/10.1029/2009GB003667>, 2010.
- 430 Brancaloni, L., Carbognani, M., Gerdol, R., Tomaselli, M., and Petraglia, A.: Refugial peatlands in the Northern Apennines. Vegetation-environment relationships and future perspectives, *Phytocoenologia*, 51, 275–298, <https://doi.org/10.1127/phyto/2022/0405>, 2022.
- Campitelli, E. ggnewscale: Multiple Fill and Colour Scales in 'ggplot2'. R package version 0.5.0, <https://CRAN.R-project.org/package=ggnewscale/>. 2024.
- 435 <https://CRAN.R-project.org/package=ggnewscale/>. 2024.
- Christensen, T. R.: Wetland emissions on the rise, *Nat. Clim. Chang.*, 14, 210–211, <https://doi.org/10.1038/s41558-024-01938-y>, 2024.
- 440 Christensen, T. R., Friberg, T., Sommerkorn, M., Kaplan, J., Illeris, L., Soegaard, H., Nordstroem, C., and Jonasson, S.: Trace gas exchange in a high-Arctic valley: 1. Variations in CO₂ and CH₄ Flux between tundra vegetation types, *Global Biogeochemical Cycles*, 14, 701–713, <https://doi.org/10.1029/1999GB001134>, 2000.
- Christensen, T. R., Scheller, J., Scheel, M., Rudd, D. A., Jackowicz-Korczynski, M., Mastepanov, M., and López-Blanco, E.: Rapid shift in greenhouse forcing of emerging arctic peatlands, *Sci Rep*, 13, 2828, <https://doi.org/10.1038/s41598-023-29859-4>, 2023.
- 445 Evans, C. D., Peacock, M., Baird, A. J., Artz, R. R. E., Burden, A., Callaghan, N., Chapman, P. J., Cooper, H. M., Coyle, M., Craig, E., Cumming, A., Dixon, S., Gauci, V., Grayson, R. P., Helfter, C., Heppell, C. M., Holden, J., Jones, D. L., Kaduk, J., Levy, P., Matthews, R., McNamara, N. P., Misselbrook, T., Oakley, S., Page, S. E., Rayment, M., Ridley, L. M., Stanley, K. M., Williamson, J. L., Worrall, F., and Morrison, R.: Overriding water table control on managed peatland greenhouse gas emissions, *Nature*, 593, 548–552, <https://doi.org/10.1038/s41586-021-03523-1>, 2021.

- 455 Forkel, M., Carvalhais, N., Rödenbeck, C., Keeling, R., Heimann, M., Thonicke, K., Zaehle, S., and Reichstein, M.: Enhanced seasonal CO₂ exchange caused by amplified plant productivity in northern ecosystems, *Science*, 351, 696–699, <https://doi.org/10.1126/science.aac4971>, 2016.
- 460 Garnier, S., Ross, N., Rudis, R., Camargo, A.P., Sciaini, M. and Scherer, C. *viridis*(Lite): Colorblind-Friendly Color Maps for R. *viridis* package version 0.6.5. <https://cran.r-project.org/web/packages/viridis/>. 2024.
- Ge, M., Korrensalo, A., Laiho, R., Lohila, A., Makiranta, P., Pihlatie, M., Tuittila, E.-S., Kohl, L., Putkinen, A., and Koskinen, M.: Plant phenology and species-specific traits control plant CH₄ emissions in a northern boreal fen, *New Phytologist*, 238, 1019–1032, <https://doi.org/10.1111/nph.18798>, 2023.
- 465 Ge, M., Korrensalo, A., Putkinen, A., Laiho, R., Kohl, L., Pihlatie, M., Lohila, A., Makiranta, P., Siljanen, H., Tuittila, E.-S., Wang, J., and Koskinen, M.: CH₄ transport in wetland plants under controlled environmental conditions – separating the impacts of phenology from environmental variables, *Plant Soil*, <https://doi.org/10.1007/s11104-024-06756-x>, 2024.
- 470 Grimes, M., Carrivick, J. L., Smith, M. W., and Comber, A. J.: Land cover changes across Greenland dominated by a doubling of vegetation in three decades, *Sci Rep*, 14, 3120, <https://doi.org/10.1038/s41598-024-52124-1>, 2024.
- 475 Helbig, M., Waddington, J. M., Alekseychik, P., Amiro, B. D., Aurela, M., Barr, A. G., Black, T. A., Blanken, P. D., Carey, S. K., Chen, J., Chi, J., Desai, A. R., Dunn, A., Euskirchen, E. S., Flanagan, L. B., Forbrich, I., Friborg, T., Grelle, A., Harder, S., Heliasz, M., Humphreys, E. R., Ikawa, H., Isabelle, P.-E., Iwata, H., Jassal, R., Korkiakoski, M., Kurbatova, J., Kutzbach, L., Lindroth, A., Löfvenius, M. O., Lohila, A., Mammarella, I., Marsh, P., Maximov, T., Melton, J. R., Moore, P. A., Nadeau, D. F., Nicholls, E. M., Nilsson, M. B., Ohta, T., Peichl, M., Petrone, R. M., Petrov, R., Prokushkin, A., Quinton, W. L., Reed, D. E., Roulet, N. T., Runkle, B. R. K., Sonnentag, O., Strachan, I. B., Taillardat, P., Tuittila, E.-S., Tuovinen, J.-P., Turner, J., Ueyama, M., Varlagin, A., Wilmking, M., Wofsy, S. C., and Zyrianov, V.: Increasing contribution of peatlands to boreal evapotranspiration in a warming climate, *Nat. Clim. Chang.*, 10, 555–560, <https://doi.org/10.1038/s41558-020-0763-7>, 2020.
- 480 Henneberg, A., Sorrell, B. K., and Brix, H.: Internal methane transport through uncus effusus: experimental manipulation of morphological barriers to test above- and below-ground diffusion limitation, *New Phytologist*, 196, 799–806, <https://doi.org/10.1111/j.1469-8137.2012.04303.x>, 2012.
- 485 Hvitfeldt, E. *paletteer: Comprehensive Collection of Color Palettes*. R package version 1.3.0, <https://github.com/EmilHvitfeldt/paletteer/>. 2021.

- Jammet, M., Dengel, S., Kettner, E., Parmentier, F.-J. W., Wik, M., Crill, P., and Friborg, T.: Year-round CH₄ and CO₂ flux dynamics in two contrasting freshwater ecosystems of the subarctic, *Biogeosciences*, 14, 5189–5216, <https://doi.org/10.5194/bg-14-5189-2017>, 2017.
- Jiménez-Alfaro, B., Aunina, L., Carbognani, M., Dítě, D., Fernández-Pascual, E., Garbolino, E., Hájek, O., Hájková, P., Ivchenko, T. G., Jandt, U., Jansen, F., Kolari, T. H. M., Pawlikowski, P., Pérez-Haase, A., Peterka, T., Petraglia, A., Plesková, Z., Tahvanainen, T., Tomaselli, M., and Hájek, M.: Habitat-based biodiversity responses to macroclimate and edaphic factors in European fen ecosystems, *Global Change Biology*, 29, 6756–6771, <https://doi.org/10.1111/gcb.16965>, 2023.
- Joabsson, A., Christensen, T.R., Wallén, B. Vascular plant controls on methane emissions from northern peatforming wetlands. *Trends Ecol. Evol.*, 14, 385–388, [https://doi.org/10.1016/S0169-5347\(99\)01649-3](https://doi.org/10.1016/S0169-5347(99)01649-3), 1999.
- Keith, D. A., Ferrer-Paris, J. R., Nicholson, E., Bishop, M. J., Polidoro, B. A., Ramirez-Llodra, E., Tozer, M. G., Nel, J. L., Mac Nally, R., Gregr, E. J., Watermeyer, K. E., Essl, F., Faber-Langendoen, D., Franklin, J., Lehmann, C. E. R., Etter, A., Roux, D. J., Stark, J. S., Rowland, J. A., Brummitt, N. A., Fernandez-Arcaya, U. C., Suthers, I. M., Wiser, S. K., Donohue, I., Jackson, L. J., Pennington, R. T., Iliffe, T. M., Gerovasileiou, V., Giller, P., Robson, B. J., Pettorelli, N., Andrade, A., Lindgaard, A., Tahvanainen, T., Terauds, A., Chadwick, M. A., Murray, N. J., Moat, J., Plischoff, P., Zager, I., and Kingsford, R. T.: A function-based typology for Earth's ecosystems, *Nature*, 610, 513–518, <https://doi.org/10.1038/s41586-022-05318-4>, 2022.
- Knorr, K.-H. and Blodau, C.: Impact of experimental drought and rewetting on redox transformations and methanogenesis in mesocosms of a northern fen soil, *Soil Biology and Biochemistry*, 41, 1187–1198, <https://doi.org/10.1016/j.soilbio.2009.02.030>, 2009.
- Koelbener, A., Ström, L., Edwards, P. J., and Olde Venterink, H.: Plant species from mesotrophic wetlands cause relatively high methane emissions from peat soil, *Plant Soil*, 326, 147–158, <https://doi.org/10.1007/s11104-009-9989-x>, 2010.
- Kolari, T. H. M., Korpelainen, P., Kumpula, T., and Tahvanainen, T.: Accelerated vegetation succession but no hydrological change in a boreal fen during 20 years of recent climate change, *Ecology and Evolution*, 11, 7602–7621, <https://doi.org/10.1002/ece3.7592>, 2021.
- Korrensalo, A., Mammarella, I., Alekseychik, P., Vesala, T., and Tuittila, E.-S.: Plant mediated methane efflux from a boreal peatland complex, *Plant Soil*, 471, 375–392, <https://doi.org/10.1007/s11104-021-05180-9>, 2022.

- Lai, D. Y. F., Moore, T. R., and Roulet, N. T.: Spatial and temporal variations of methane flux measured by autochambers in a temperate ombrotrophic peatland, *Journal of Geophysical Research: Biogeosciences*, 119, 864–880, <https://doi.org/10.1002/2013JG002410>, 2014.
- 525
- Laitinen, J., Nyberg, J., Kaakinen, E., Küttim, M., Muurinen, L., Ulvinen, T., Virtanen, R., and Tahvanainen, T.: Site types, species composition, species richness and ecological gradients of rich fens in the Oulanka region, north-boreal Finland, *Phytocoenologia*, 52, 1–27, <https://doi.org/10.1127/phyto/2023/0421>, 2024.
- 530
- Larmola, T., Tuittila, E.-S., Tiirola, M., Nykänen, H., Martikainen, P. J., Yrjälä, K., Tuomivirta, T., and Fritze, H.: The role of *Sphagnum* mosses in the methane cycling of a boreal mire, *Ecology*, 91, 2356–2365, 2010.
- Ma, L., Zhu, G., Chen, B., Zhang, K., Niu, S., Wang, J., Ciais, P., and Zuo, H.: A globally robust relationship between water table decline, subsidence rate, and carbon release from peatlands, *Commun Earth Environ*, 3, 1–14, <https://doi.org/10.1038/s43247-022-00590-8>, 2022.
- 535
- McCune, B., Grace, J. B., and Urban, D. L.: *Analysis of Ecological Communities*, MjM Software Design, Gleneden Beach, OR, 300 pp., ISBN 0-9721290-0-6, 2002.
- 540
- McCune, B. and Mefford, M.J. PC-ORD. Multivariate Analysis of Ecological Data. Version 7.09. 2018.
- Nielsen, C. S., Michelsen, A., Strobel, B. W., Wulff, K., Banyasz, I., and Elberling, B.: Correlations between substrate availability, dissolved CH₄, and CH₄ emissions in an arctic wetland subject to warming and plant removal, *Journal of Geophysical Research: Biogeosciences*, 122, 645–660, <https://doi.org/10.1002/2016JG003511>, 2017.
- 545
- Olefeldt, D., Euskirchen, E. S., Harden, J., Kane, E., McGuire, A. D., Waldrop, M. P., and Turetsky, M. R.: A decade of boreal rich fen greenhouse gas fluxes in response to natural and experimental water table variability, *Global Change Biology*, 23, 2428–2440, <https://doi.org/10.1111/gcb.13612>, 2017.
- 550
- Pearson, M., Penttilä, T., Harjunpää, L., Laiho, R., Laine, J., Sarjala, T., Silvan, K., and Silvan, N.: Effects of temperature rise and water-table-level drawdown on greenhouse gas fluxes of boreal sedge fens, 489–505, 2015.
- Pirk, N., Mastepanov, M., Parmentier, F.-J. W., Lund, M., Crill, P., and Christensen, T. R.: Calculations of automatic chamber flux measurements of methane and carbon dioxide using short time series of concentrations, *Biogeosciences*, 13, 903–912, <https://doi.org/10.5194/bg-13-903-2016>, 2016a.
- 555

Pirk, N., Tamstorf, M. P., Lund, M., Mastepanov, M., Pedersen, S. H., Mylius, M. R., Parmentier, F.-J. W., Christiansen, H. H., and Christensen, T. R.: Snowpack fluxes of methane and carbon dioxide from high Arctic tundra, *Journal of Geophysical Research: Biogeosciences*, 121, 2886–2900, <https://doi.org/10.1002/2016JG003486>, 2016b.

560

Raivonen, M., Smolander, S., Backman, L., Susiluoto, J., Aalto, T., Markkanen, T., Mäkelä, J., Rinne, J., Peltola, O., Aurela, M., Lohila, A., Tomasic, M., Li, X., Larmola, T., Juutinen, S., Tuittila, E.-S., Heimann, M., Sevanto, S., Kleinen, T., Brovkin, V., and Vesala, T.: HIMMELI v1.0: Helsinki Model of Methane build-up and emission for peatlands, *Geoscientific Model Development*, 10, 4665–4691, <https://doi.org/10.5194/gmd-10-4665-2017>, 2017.

565

Posit team. RStudio: Integrated Development Environment for R. Posit Software, PBC, Boston, MA. URL <http://www.posit.co/>. 2024.

570

R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>. 2024.

575

Ramage, J., Kuhn, M., Virkkala, A.-M., Voigt, C., Marushchak, M. E., Bastos, A., Biasi, C., Canadell, J. G., Ciais, P., López-Blanco, E., Natali, S. M., Olefeldt, D., Potter, S., Poulter, B., Rogers, B. M., Schuur, E. A. G., Treat, C., Turetsky, M. R., Watts, J., and Hugelius, G.: The Net GHG Balance and Budget of the Permafrost Region (2000–2020) From Ecosystem Flux Upscaling, *Global Biogeochemical Cycles*, 38, e2023GB007953, <https://doi.org/10.1029/2023GB007953>, 2024.

580

Reid, M. C., Pal, D. S., and Jaffé, P. R.: Dissolved gas dynamics in wetland soils: Root-mediated gas transfer kinetics determined via push-pull tracer tests, *Water Resources Research*, 51, 7343–7357, <https://doi.org/10.1002/2014WR016803>, 2015.

Riutta, T., Laine, J. ., Aurela, M., Rinne, J. ., Vesala, T., Laurila, T., Haapanala, S., Pihlatie, M., and Tuittila, E.-S.: Spatial variation in plant community functions regulates carbon gas dynamics in a boreal fen ecosystem, *Tellus B: Chemical and Physical Meteorology*, 59, 838, <https://doi.org/10.1111/j.1600-0889.2007.00302.x>, 2007.

585

Riutta, T., Korrensalo, A., Laine, A. M., Laine, J., and Tuittila, E.-S.: Interacting effects of vegetation components and water level on methane dynamics in a boreal fen, *Biogeosciences*, 17, 727–740, <https://doi.org/10.5194/bg-17-727-2020>, 2020.

Saarinén, T.: Biomass and production of two vascular plants in a boreal mesotrophic fen, *Can. J. Bot.*, 74, 934–938, <https://doi.org/10.1139/b96-116>, 1996.

Saarinen, T.: Demography of *Carex rostrata* in a boreal mesotrophic fen: Shoot dynamics and biomass development, *Ann. Bot. Fennici.*, 36, 203–209, ISSN 0003-3847, 1998.

Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P., Regnier, P., Canadell, J. G., Jackson, R. B., Patra, P. K.,
 595 Bousquet, P., Ciais, P., Dlugokencky, E. J., Lan, X., Allen, G. H., Bastviken, D., Beerling, D. J., Belikov, D. A., Blake, D. R.,
 Castaldi, S., Crippa, M., Deemer, B. R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L., Holgerson, M. A.,
 Hopcroft, P. O., Hugelius, G., Ito, A., Jain, A. K., Janardanan, R., Johnson, M. S., Kleinen, T., Krummel, P., Lauerwald, R.,
 Li, T., Liu, X., McDonald, K. C., Melton, J. R., Mühle, J., Müller, J., Murguia-Flores, F., Niwa, Y., Noce, S., Pan, S., Parker,
 R. J., Peng, C., Ramonet, M., Riley, W. J., Rocher-Ros, G., Rosentreter, J. A., Sasakawa, M., Segers, A., Smith, S. J., Stanley,
 600 E. H., Thanwerdas, J., Tian, H., Tsuruta, A., Tubiello, F. N., Weber, T. S., Van Der Werf, G., Worthy, D. E., Xi, Y., Yoshida,
 Y., Zhang, W., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: Global Methane Budget 2000–2020, [https://doi.org/10.5194/essd-](https://doi.org/10.5194/essd-2024-115)
[2024-115](https://doi.org/10.5194/essd-2024-115), 6 June 2024.

Schuur, E. A. G., Abbott, B. W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G., Grosse, G., Jones, M., Koven,
 605 C., Leshyk, V., Lawrence, D., Loranty, M. M., Mauritz, M., Olefeldt, D., Natali, S., Rodenhizer, H., Salmon, V., Schädel, C.,
 Strauss, J., Treat, C., and Turetsky, M.: Permafrost and Climate Change: Carbon Cycle Feedbacks From the Warming Arctic,
Annual Review of Environment and Resources, 47, 343–371, <https://doi.org/10.1146/annurev-environ-012220-011847>, 2022.

Singh, P., Těšitel, J., Plesková, Z., Peterka, T., Hájková, P., Dítě, D., Pawlikowski, P., Hájek, M., and Fraser, L.: The ratio
 610 between bryophyte functional groups impacts vascular plants in rich fens, *Applied Vegetation Science*, 22, 494–507,
<https://doi.org/10.1111/avsc.12454>, 2019.

Song, S., Wang, X., Wang, Y., Li, T., and Huang, J.: NO₃[−] is an important driver of nitrite-dependent anaerobic methane
 oxidation bacteria and CH₄ fluxes in the reservoir riparian zone, *Environ Sci Pollut Res*, 29, 16138–16151,
 615 <https://doi.org/10.1007/s11356-021-16914-8>, 2022.

Ström, L., Mastepanov, M., and Christensen, T. R.: Species-specific Effects of Vascular Plants on Carbon Turnover and
 Methane Emissions from Wetlands, *Biogeochemistry*, 75, 65–82, <https://doi.org/10.1007/s10533-004-6124-1>, 2005.

620 Ström, L., Tagesson, T., Mastepanov, M., and Christensen, T. R.: Presence of *Eriophorum scheuchzeri* enhances substrate
 availability and methane emission in an Arctic wetland, *Soil Biology and Biochemistry*, 45, 61–70,
<https://doi.org/10.1016/j.soilbio.2011.09.005>, 2012.

- Ström, L., Falk, J. M., Skov, K., Jackowicz-Korczynski, M., Mastepanov, M., Christensen, T. R., Lund, M., and Schmidt, N.
625 M.: Controls of spatial and temporal variability in CH₄ flux in a high arctic fen over three years, *Biogeochemistry*, 125, 21–
35, <https://doi.org/10.1007/s10533-015-0109-0>, 2015.
- Swindles, G. T., Morris, P. J., Mullan, D. J., Payne, R. J., Roland, T. P., Amesbury, M. J., Lamentowicz, M., Turner, T. E.,
Gallego-Sala, A., Sim, T., Barr, I. D., Blaauw, M., Blundell, A., Chambers, F. M., Charman, D. J., Feurdean, A., Galloway, J.
630 M., Galka, M., Green, S. M., Kajukalo, K., Karofeld, E., Korhola, A., Lamentowicz, L., Langdon, P., Marcisz, K., Mauquoy,
D., Mazei, Y. A., McKeown, M. M., Mitchell, E. A. D., Novenko, E., Plunkett, G., Roe, H. M., Schoning, K., Sillasoo, Ü.,
Tsyganov, A. N., Van Der Linden, M., Väliranta, M., and Warner, B.: Widespread drying of European peatlands in recent
centuries, *Nat. Geosci.*, 12, 922–928, <https://doi.org/10.1038/s41561-019-0462-z>, 2019.
- 635 Turetsky, M. R., Kotowska, A., Bubier, J., Dise, N. B., Crill, P., Hornibrook, E. R. C., Minkinen, K., Moore, T. R., Myers-
Smith, I. H., Nykänen, H., Olefeldt, D., Rinne, J., Saarnio, S., Shurpali, N., Tuittila, E.-S., Waddington, J. M., White, J. R.,
Wickland, K. P., and Wilking, M.: A synthesis of methane emissions from 71 northern, temperate, and subtropical wetlands,
Global Change Biology, 20, 2183–2197, <https://doi.org/10.1111/gcb.12580>, 2014.
- 640 Turner, J. C., Moorberg, C. J., Wong, A., Shea, K., Waldrop, M. P., Turetsky, M. R., and Neumann, R. B.: Getting to the Root
of Plant-Mediated Methane Emissions and Oxidation in a Thermokarst Bog, *Journal of Geophysical Research: Biogeosciences*,
125, e2020JG005825, <https://doi.org/10.1029/2020JG005825>, 2020.
- Vargas, R. and Le, V. H.: The paradox of assessing greenhouse gases from soils for nature-based solutions, *Biogeosciences*,
645 20, 15–26, <https://doi.org/10.5194/bg-20-15-2023>, 2023.
- Waldo, N. B., Hunt, B. K., Fadely, E. C., Moran, J. J., and Neumann, R. B.: Plant root exudates increase methane emissions
through direct and indirect pathways, *Biogeochemistry*, 145, 213–234, <https://doi.org/10.1007/s10533-019-00600-6>, 2019.
- 650 Wickham, H. *forcats: Tools for Working with Categorical Variables (Factors)*. R package version 1.0.0, [https://CRAN.R-
project.org/package=forcats/](https://CRAN.R-project.org/package=forcats/), 2023.
- Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. R package version 3.5.1, [https://cran.r-
project.org/web/packages/ggplot2/](https://cran.r-project.org/web/packages/ggplot2/), 2024.
- 655 Wickham, H. and Bryan, J. *readxl: Read Excel Files*. R package version 1.4.3, <https://CRAN.R-project.org/package=readxl/>,
2023.

- Wickham, H., François, R., Henry, L., Müller, K. and Vaughan, D. dplyr: A Grammar of Data Manipulation. R package version 1.1.4, <https://CRAN.R-project.org/package=dplyr/>, 2023a.
- Wickham, H., Pedersen, T. and Seidel, D. _scales: Scale Functions for Visualization. R package version 1.3.0, <https://CRAN.R-project.org/package=scales/>. 2023b.
- Wickham, H., Vaughan, D. and Girlich, M. tidyr: Tidy Messy Data. R package version 1.3.1, <https://CRAN.R-project.org/package=tidyr/>, 2024.
- Wolff, F., H. M. Kolari, T., Villoslada, M., Tahvanainen, T., Korpelainen, P., A. P. Zamboni, P., and Kumpula, T.: RGB vs. Multispectral imagery: Mapping aapa mire plant communities with UAVs, *Ecological Indicators*, 148, 110140, <https://doi.org/10.1016/j.ecolind.2023.110140>, 2023.
- Yavitt, J. B., Yashiro, E., Cadillo-Quiroz, H., and Zinder, S. H.: Methanogen diversity and community composition in peatlands of the central to northern Appalachian Mountain region, North America, *Biogeochemistry*, 109, 117–131, <https://doi.org/10.1007/s10533-011-9644-5>, 2012.
- Yuan, K., Li, F., McNicol, G., Chen, M., Hoyt, A., Knox, S., Riley, W. J., Jackson, R., and Zhu, Q.: Boreal–Arctic wetland methane emissions modulated by warming and vegetation activity, *Nat. Clim. Chang.*, 14, 282–288, <https://doi.org/10.1038/s41558-024-01933-3>, 2024.
- Zhang, H., Tuittila, E.-S., Korrensalo, A., Räsänen, A., Virtanen, T., Aurela, M., Penttilä, T., Laurila, T., Gerin, S., Lindholm, V., and Lohila, A.: Water flow controls the spatial variability of methane emissions in a northern valley fen ecosystem, *Biogeosciences*, 17, 6247–6270, <https://doi.org/10.5194/bg-17-6247-2020>, 2020.
- Zhang, H., Tuittila, E.-S., Korrensalo, A., Laine, A. M., Uljas, S., Welti, N., Kerttula, J., Maljanen, M., Elliott, D., Vesala, T., and Lohila, A.: Methane production and oxidation potentials along a fen-bog gradient from southern boreal to subarctic peatlands in Finland, *Global Change Biology*, 27, 4449–4464, <https://doi.org/10.1111/gcb.15740>, 2021.

Appendix A




Figure A1. The depth of the water table (WTD) at 36 study plots in Puukkosuo fen during June–October 2022. The average minimum value for WTD was -4.8 cm and the average maximum +1.8 cm from the peat surface.

5 Table A1. Distribution (Dist), mean (M), median (Mdn) and standard deviation (SD) of the heights (h) and dry biomass (BM) of vascular plant (VP) and bryophyte (Bryo) species identified from the 36 study plots (P) in July 2022. The number of VPs within the studied area were counted by the number of shoots (Shoots_P). BMs are reported only for samples (S) collected for BM estimations. Separate samples were collected for plants in different growth stage: flowering (FL) and non-flowering (NF) VP species, except for four species for which these were combined (Comb). The sample size (Sample) of bryophytes depended on the commonness of the species and it represented either 1 or 5 % of the studied area. For all Bryo species a BM for 1 % coverage of the studied area was calculated (BM 1%). Abbreviations for the plant species: *Andromeda polifolia* (AndPol), *Angelica sylvestris* (AngSyl), *Betula nana* (BetNan), *Carex chordorrhiza* (CarCho), *Carex dioica* (CarDio), *Carex flava* (CarFla), *Carex lasiocarpa* (CarLas), *Carex limosa* (CarLim), *Carex panicea* (CarPan), *Carex rostrata* (CarRos), *Dactylorhiza* species (DactSp.), *Drosera anglica* (DroAng), *Drosera rotundifolia* (DroRot), *Equisetum fluviatile* (EquFlu), *Equisetum variegatum* (EquVar), *Eriophorum angustifolium* (EriAng), *Eriophorum gracile* (EriGra), *Eriophorum latifolium* (EriLat), *Festuca ovina* (FesOvi), *Menyanthes trifoliata* (MenTri), *Molinia caerulea* (MolCae), *Pedicularis palustris* (PedPal), *Pinguicula* species (PingSp.), *Potentilla erecta* (PotEre), *Saussurea alpina* (SauAlp), *Selaginella selaginoides* (SelSel), *Tofieldia pusilla* (TofPus), *Trichophorum alpinum* (TriAlp), *Trichophorum cespitosum* (TriCes), *Vaccinium oxycoccos* (VacOxy), *Viola epipsila* (VioEpi), *Aneura pinguis* (AnePin), *Aulacomnium palustre* (AulPal), *Campylium stellatum* (CamSte), *Cinclidium stygium* (CinSty), *Fissidens adianthoides* (FisAdi), *Mesoptychia rutheana* (MesRut), *Paludella squarrosa* (PalSqu), *Scorpidium cossonii* (ScoCos), *Sphagnum warnstorffii* (SphWar) and *Tomentypnum nitens* (TomNit). Plot-scale vegetation data can be obtained from the correspondent.

10

15

20

VP species	Growth stage	Shoots_P (n)	h_Dist_P (cm)	h_Dist_S (cm)	h_M_P (cm)	h_M_S (cm)	h_Mdn_P (cm)	h_Mdn_S (cm)	SD_P (cm)	SD_S (cm)	BM_Dist (g)	BM_M (g)	BM_Mdn (g)	BM_SD (g)
AndPol	Comb	217	3–9	3–8	5.4	4.8	5	4	1.6	2	0.0190–0.0941	0.0499	0.0394	0.03
AngSyl	NF	3	15	9–21	15	14.9	15	15.5	0	3.8	0.0356–0.2916	0.1317	0.1251	0.09
BetNan	Comb	48	4–16	4–17	8.8	10.4	9	10	3.6	3.5	0.0228–0.6131	0.1646	0.1192	0.17
CarCho	FL	122	13–22	11–22	15.6	16.4	15	16.5	2.8	3.2	0.0250–0.0824	0.0431	0.0359	0.02
CarCho	NF	596	12–22	11–19	14	14.7	13	14	2.6	2.5	0.0165–0.0611	0.0344	0.0295	0.01
CarDio	FL	6	9–28	12–24	17.3	17.8	15	16.5	9.7	3.5	0.0110–0.0359	0.0222	0.0227	0.01
CarDio	NF	141	8–15	9–16	9.8	12.3	9	12	2.1	2.6	0.0073–0.0326	0.0146	0.0126	0.01
CarFla	FL	2	18–35	28–44	26.5	32.8	26.5	32	12	5.1	0.1524–0.3685	0.2204	0.1908	0.06
CarFla	NF	7	18–23	16–29	20.5	23.8	20.5	25	3.5	4.7	0.0421–0.1581	0.1072	0.1045	0.04
CarLas	FL	1	53	27–62	53	52	53	55.5	0	11.8	0.1367–0.4719	0.3274	0.3528	0.1
CarLas	NF	457	37–63	41–66	48.7	55.4	48	54	6.3	8.7	0.1493–0.5235	0.2667	0.2483	0.11
CarLim	FL	3	18–20	20–28	19	24.4	19	24	1.4	3	0.0431–0.1291	0.0932	0.0986	0.03
CarLim	NF	31	15–18	15–24	16.5	19.8	16.5	20	2.1	3.4	0.0306–0.1366	0.0306	0.1366	0.03
CarPan	FL	3	23–24	13–42	23.5	29.8	23.5	30	0.7	11	0.0199–0.1986	0.1001	0.0943	0.05
CarPan	NF	45	15–36	14–38	23.1	25	23	26.5	6.4	8.1	0.0334–0.3339	0.1343	0.118	0.09
CarRos	FL	11	45–56	49–60	51	54.7	53	54	5.6	3.8	0.8980–1.8110	1.3596	1.3199	0.31
CarRos	NF	50	25–57	49–60	46	54.7	52	54	11	3.8	0.8980–1.8110	1.3788	1.3135	0.31
DactSp.	FL	1	24	24	24	24	24	24	0	0	0.4342	0.4342	0.4342	0
DactSp.	NF	8									0.0126–0.0486	0.0245	0.0222	0.01
DroAng	Comb	6									0.0025–0.0565	0.0191	0.0161	0.02
DroRot	Comb	18									0.0049–0.1010	0.0202	0.0099	0.03
EquFlu	NF	120	10–36	15–41	24.2	27.5	24	27	7.05	6.5	0.0580–0.2401	0.1387	0.1349	0.05
EquVar	NF	12	6–10	9–16	8	11.7	8	11.5	2.8	1.9	0.0258–0.0529	0.0355	0.0336	0.01
EriAng	NF	4	20–26	26–47	23.7	34.8	25	34	3.2	5.6	0.1752–0.4143	0.2841	0.2794	0.07
EriGra	FL	1	25	23–40	25	29.9	25	30.5	0	5	0.0509–0.1255	0.0761	0.0736	0.2
EriGra	NF	2	12	8–13	12	10.7	12	10.5	0	1.6	0.0305–0.0613	0.0437	0.0388	0.01
EriLat	NF	18	14–22	13–26	17.8	19.8	17.5	19.5	2.8	3.9	0.0523–0.3484	0.1556	0.1067	0.1
FesOvi	NF	2	7	7	7	7	7	7	0	0	0.0287–0.0307	0.0298	0.0298	0.001
MenTri	FL	2	7–8	16–23	7.5	18.1	7.5	19.5	0.7	2.1	0.1573–0.3780	0.2538	0.2322	0.1
MenTri	NF	180	3–10	5–14	6.5	9.4	6	9.5	2.1	3.3	0.0173–0.2177	0.0942	0.0628	0.07
MolCae	FL	1	33	31–74	33	55.7	33	56.5	0	3.3	0.1985–0.4128	0.3198	0.3343	0.07

MolCae	NF	173	21–36	21–36	25.5	28.4	24	29	4.7	5.9	0.0289–0.1339	0.0675	0.0527	0.04
PedPal	FL	5	12–21	10–37	15.6	23.1	14	24.5	3.8	8.2	0.1104–1.4398	0.6056	0.3976	0.49
PedPal	NF	7	4–14	5–14	9	7.8	9	6	7.1	3	0.0091–0.0616	0.0267	0.0153	0.02
PingSp.	NF	3									0.0074–0.0680	0.0312	0.0284	0.02
PotEre	FL	3	18	15–31	18	21	18	20	0	4.5	0.0836–0.9890	0.3105	0.2053	0.29
PotEre	NF	26	10–25	9616	14	12.5	11	12	5.7	2.5	0.0335–0.1021	0.0674	0.0722	0.02
SauAlp	NF	3	13	5–22	13	12.8	13	13	0	4.7	0.0141–0.3586	0.1127	0.0806	0.1
SelSel	FL	31	4–6	2–9	4.6	5.4	4	5	0.9	2.2	0.0074–0.0310	0.017	0.0154	0.01
TofPus	FL	1	21	13–24	21	17.9	21	17	0	3.6	0.0182–0.1026	0.0564	0.0558	0.03
TofPus	NF	30									0.0065–0.0316	0.0191	0.0213	0.01
TriAlp	FL	71	13–25	15–24	18.4	20.2	17.5	21.5	3.5	3.2	0.0128–0.0332	0.022	0.0223	0.01
TriCes	FL	1828	6–27	15–27	19.4	20.5	20	20	5.3	3.2	0.0168–0.0379	0.0247	0.0245	0.01
VacOxy	NF	86	3–15	3–13	6.8	7.8	5	6.5	3.6	4	0.0057–0.0326	0.0175	0.0177	0.01
VioEpi	NF	4	3–5	4–8	4	6.3	4	6.5	1.4	1.4	0.0065–0.0378	0.0228	0.0200	0.01
Bryo species	Sample	BM_Dist (g)	BM_M (g)	BM_Mdn (g)	BM_SD (g)	BM 1% (g)								
AnePin	1 x 1 %	0.0126	0.0126	0.0126	0	0.0126								
AulPal	3 x 5 %	0.6562–1.2110	0.9212	0.8965	0.28	0.1842								
CamSte	3 x 5 %	0.3650–0.4697	0.4172	0.417	0.05	0.0834								
CinSty	1 x 1 %	0.0307	0.0307	0.0307	0	0.0307								
FisAdi	1 x 1 %	0.0691	0.0691	0.0691	0	0.0691								
MesRut	1 x 1 %	0.0264	0.0264	0.0264	0	0.0264								
PalSqu	1 x 1 %	0.0695	0.0695	0.0695	0	0.0695								
ScoCos	3 x 5 %	0.5623–1.0105	0.777	0.7582	0.22	0.1554								
SphWar	3 x 5 %	0.5028–1.1476	0.6669	0.8417	0.32	0.1334								
TomNit	3 x 5 %	0.5402–0.9949	0.7166	0.6148	0.24	0.1433								

Table A2. Total plot-scale dry mass weight biomasses (BM) of vascular plants (VP) and bryophytes (Bryo) in grams inside the collar (A=660.5 cm²) at each experimental plot (37–72), a VP to Bryo BM ratio (VP:Bryo ratio) and the portion of VP BM of total BM (VP of total BM).

Plot	BM VP	BM Bryo	VP:Bryo ratio	VP of total BM
37	17.8538	15.0746	1.18	54 %
38	22.0626	17.7820	1.24	55 %
39	6.2854	13.3400	0.47	32 %
40	7.9262	8.0765	0.98	50 %
41	6.4044	8.0765	0.79	44 %
42	16.8502	9.1706	1.84	65 %
43	10.6523	13.0900	0.81	45 %
44	9.7756	11.7626	0.83	45 %
45	8.9990	10.1400	0.89	47 %
46	10.4179	8.7790	1.19	54 %
47	14.3059	12.8400	1.11	53 %
48	7.5385	13.8365	0.54	35 %
49	6.6854	4.2710	1.57	61 %
50	32.1099	12.4200	2.59	72 %
51	15.2690	8.8400	1.73	63 %
52	11.7328	9.3945	1.25	56 %
53	9.5022	15.5400	0.61	38 %
54	9.1127	8.3400	1.09	52 %
55	9.9275	11.7626	0.84	46 %
56	8.2885	9.7800	0.85	46 %
57	10.3305	8.7000	1.19	54 %
58	4.1176	15.1273	0.27	21 %
59	5.3803	13.3400	0.40	29 %
60	6.7374	10.9706	0.61	38 %
61	8.9244	15.5400	0.57	36 %
62	8.2568	8.8896	0.93	48 %
63	11.0370	7.7520	1.42	59 %
64	10.2857	11.2200	0.92	48 %
65	13.5974	11.8010	1.15	54 %
66	8.2232	15.1800	0.54	35 %
67	5.9341	15.5400	0.38	28 %
68	7.2415	15.2906	0.47	32 %
69	8.8783	13.3400	0.67	40 %
70	11.2556	11.6765	0.96	49 %
71	7.4490	12.2400	0.61	38 %
72	9.0483	13.3400	0.68	40 %

Table A3. Combined dry mass weight estimation (g) of all the study plots for each species. Vascular plants on the top, bryophytes at the bottom of the list in alphabetical order. For abbreviations of the species, see Table A1.

Species	Mass (g)
CarLas	121.9426
CarRos	83.8956
TriCes	40.2116
CarCho	25.7606
MenTri	17.5578
MolCae	16.6548
EquFlu	16.644
AndPol	11.3273
BetNan	7.9008
TriAlp	6.502
CarPan	6.3438
PedPal	4.4261
PotEre	3.2905
EriLat	3.2676
CarDio	2.1918
VacOxy	1.505
CarLim	1.2282
CarFla	1.1912
EriAng	1.1364
TofPus	1.0885
SauAlp	1.0143
SelSel	0.663
DactSp.	0.6302
EquVar	0.426
AngSyl	0.3951
DroRot	0.3636
PingSp.	0.1872
EriGra	0.1635
DroLon	0.1146
VioEpi	0.0912
FesOvi	0.0596
ScoCos	192.8514
CamSte	112.5066
SphWar	91.6458
AulPal	18.2358
CinSty	2.0262
MesRut	1.584
PalSqu	1.529
FisAdi	1.1056
TomNit	0.7166
AnePin	0.063

Appendix B




Figure B1. Cluster dendrogram of all species-data. Indicator species by clusters: Sco = *Scorpidium cossonii*, Sph = *Sphagnum* spp. and *Paludella squarrosa* and Cam = *Campylium stellatum*.




Figure B2. Cluster dendrogram of vascular plant data. Indicator species by clusters: C.ros= *Carex rostrata*, C.cho= *Carex chordorrhiza* and T.ces = *Carex lasiocarpa*, *Potentilla erecta* and *Trichophorum cespitosum*.




Figure B3. Cluster dendrogram of bryophyte data. Indicator species by clusters: Sco = *Scorpidium cossonii*, Sph = *Sphagnum* spp. and *Paludella squarrosa* and Cam = *Campylium stellatum*.

15 Table B1. Indicator values (% of perfect indication, based on combining the values of relative abundance and relative frequency) of
all identified species (n = 40) of Puukkosuo segregated by the vascular plant clusters. The values for the clusters' significant indicator
species are bolded. Species with indicator value lower than 20 % were excluded from the table. Abbreviations for the species most
strongly connected to the clusters: C.ros = *C. rostrata*, C.cho = *C. chordorrhiza* and T. ces = *C. lasiocarpa*, *P. erecta*, and *T. cespitosum*.
Abbreviations for the species: CarRos = *Carex rostrata*, MenTri = *Menyanthes trifoliata*, TriAlp = *Trichophorum alpinum*, CarLim =
Carex limosa, CinSty = *Cinclidium stygium*, CarCho = *Carex chordorrhiza*, VacOxy = *Vaccinium oxycoccos*, BetNan = *Betula nana*,
AndPol = *Andromeda polifolia*, EriLat = *Eriophorum latifolium*, TriCes = *Trichophorum cespitosum*, PotEre = *Potentilla erecta*, CarLas
= *Carex lasiocarpa*, EquFlu = *Equisetum fluviatile*, MolCae = *Molinia caerulea*, TofPus = *Tofieldia pusilla*, PingSp. = *Pinguicula* sp.,
20 CarDio = *Carex dioica*, CamSte = *Campyllum stellatum*, ScoCos = *Scorpidium cossonii* and SphWar = *Sphagnum warnstorffii*.

		C.ros	C.cho	T.ces
C.ros				
	CarRos (p < 0.01)	95	0	1
	MenTri	38	16	18
	TriAlp	31	2	8
	CarLim	20	0	0
C.cho				
	CinSty (p = 0.01)	8	64	2
	CarCho (p = 0.02)	31	58	5
	VacOxy	9	46	11
	BetNan	26	39	3
	AndPol	31	36	25
	EriLat	7	31	2
T.ces				
	TriCes (p < 0.01)	7	13	65
	PotEre (p < 0.01)	0	0	55
	CarLas (p < 0.01)	36	11	51
	EquFlu	35	8	41
	MolCae	6	15	41
	TofPus	4	1	27
	PingSp.	0	0	25
	CarDio	0	18	23
Relative frequencies of common species				
	CamSte	70	67	75
	VacOxy	40	83	45
	ScoCos.	40	50	75
	SphWar	60	17	20




Figure B4. Ordination graphs of vascular plant data. (a) DCA with rare species downweighed (eigenvalue axis 1 = 0.463, axis 2 = 0.154) and (b) CCA (eigenvalue axis 1 = 0.423, axis 2 = 0.163). Plant community clusters: C.ros = *Carex rostrata*, C.cho = *Carex chondorrhiza* and T.ces = *Carex lasiocarpa*, *Potentilla erecta* and *Trichophorum cespitosum*. Timespans for methane fluxes: CH4 Year-round (19.10.2021–31.10.2022), CH4 Snow-free (13.5.–26.10.2022) and CH4 Snow (19.10.2021–12.5.2022, 27.–31.10.2022). NO₃+NO₂ is the combined concentration of nitrate (NO₃⁻) and nitrite (NO₂⁻), BMratio is the ratio of vascular plant and bryophyte biomasses, and Peat (m) is the peat layer thickness.




Figure B5. Ordination graphs of bryophyte data. (a) DCA with rare species downweighed (eigenvalue axis 1 = 0.950, axis 2 = 0.364) and (b) CCA (eigenvalue axis 1 = 0.711, axis 2 = 0.510). Abbreviations for plant community clusters: Sco = *Scorpidium cossonii*, Sph = *Sphagnum* spp. and *Paludella squarrosa* and Cam = *Campylium stellatum*. WTD stands for water table depth and CarRos for biomass of *C. rostrata*.

35


Table B2. Pearson's correlation matrix among the environmental and biomass variables. Significant correlations bolded, p-value ≤ 0.05 ($r \geq 0.33$) marked with one asterisk (*), p-value ≤ 0.01 ($r \geq 0.42$) with two asterisks () and p-value ≤ 0.001 ($r \geq 0.53$) with three asterisks (***). CH4_y = year-round methane flux, CH4_sf = snow-free season methane flux, CH4_s = snow season methane flux, WTD = water table depth, NO3+NO2 = nitrate and nitrite, NH4 = ammonium, TOTN = total nitrogen, TOC = total organic carbon, IC = inorganic carbon, BMratio = vascular plant to bryophyte biomass ratio, BMvp = vascular plant biomass, BMbryo = bryophyte biomass, BMsed = biomass of all sedges, BMsph = biomass of *Sphagnum* mosses, Peat = peat layer thickness, Litter = litter percentage, CarRos = biomass of *Carex rostrata*, SoilT_y = average year-round soil temperature at -5 cm, SoilT_sf = average snow-free season soil temperature at -5 cm and SoilT_s = average snow season soil temperature at -5 cm. WTD and porewater chemistry concentrations are values of the snow-free season.**

	CH4_y	CH4_sf	CH4_s	WTD	pH	NO3+NO2	NH4	TOTN	TOC	IC	BMratio	BMvp	BMbryo	BMsed	BMsph	Peat	Litter	C.Ros	SoilT_y	SoilT_sf	SoilT_s
CH4_y		0.99***	0.60***	-0.05	-0.41*	0.34*	0.17	0.21	0.24	-0.08	0.45**	0.54***	0.03	0.57***	0.16	0.09	0.12	0.61***	0.22	0.06	-0.29
CH4_sf	0.99***		0.55***	-0.06	-0.38*	0.34*	0.17	0.22	0.22	-0.08	0.45**	0.54***	0.05	0.57***	0.19	0.07	0.12	0.60***	0.17	0.06	-0.27
CH4_s	0.60***	0.55***									0.24	0.16	-0.19	0.21	0.11	0.17	0.09	0.20	0.23	0.24	-0.14
WTD	-0.05	-0.06			-0.48**	0.15	-0.18	0.10	0.07	0.13	-0.13	-0.17	-0.13	-0.15	-0.59***	-0.20	0.00	-0.23	0.19	0.21	-0.21
pH	-0.41*	-0.38*				-0.11	0.23	-0.21	-0.28	-0.05	-0.30	-0.26	0.16	-0.35	0.55***	0.12	-0.15	-0.22	-0.25	-0.07	0.36*
NO3+NO2	0.34*	0.34*		0.15	-0.11		0.20	-0.18	0.05	0.24	0.13	0.14	-0.07	0.14	-0.09	0.28	-0.03	0.24	0.07	0.03	-0.12
NH4	0.17	0.17		-0.18	0.23	0.20		-0.04	-0.07	0.03	-0.13	-0.10	0.13	-0.09	0.13	0.15	-0.34*	0.01	-0.01	-0.15	0.28
TOTN	0.21	0.22		0.10	-0.21	-0.18	-0.04		0.08	-0.14	0.36*	0.23	-0.16	0.24	-0.09	0.12	0.09	0.29	0.06	0.10	-0.19
TOC	0.24	0.22		0.07	-0.28	0.05	-0.07	0.08		0.41*	0.28	0.22	-0.10	0.19	-0.03	0.34*	-0.03	0.13	-0.01	-0.10	-0.09
IC	-0.08	-0.08		0.13	-0.05	0.24	0.03	-0.14	0.41*		0.09	0.20	0.17	0.15	-0.17	0.31	0.20	0.11	-0.13	-0.21	0.07
BMratio	0.45**	0.45**	0.24	-0.13	-0.30	0.13	-0.13	0.36*	0.28	0.09		0.81***	-0.46**	0.78***	-0.02	0.37*	0.46**	0.71***	-0.06	-0.14	-0.15
BMvp	0.53***	0.54***	0.16	-0.17	-0.26	0.14	-0.10	0.23	0.22	0.20	0.81***		0.11	0.98***	0.13	0.37*	0.59***	0.93***	0.03	-0.08	-0.09
BMbryo	0.03	0.05	-0.19	-0.13	0.16	-0.07	0.13	-0.16	-0.10	0.17	-0.46**	0.11		0.14	0.22	-0.03	0.06	0.19	0.03	0.06	0.16
Bmsedg	0.57***	0.57***	0.21	-0.15	-0.35*	0.14	-0.09	0.24	0.19	0.15	0.78***	0.98***	0.14		0.06	0.34*	0.62***	0.93***	0.01	-0.12	-0.14
				-																	
BMsph	0.16	0.19	0.11	0.59***	0.55***	-0.09	0.13	-0.09	-0.03	-0.17	-0.02	0.13	0.22	0.06		0.12	-0.02	0.16	-0.12	0.02	0.22
Peat	0.09	0.07	0.17	-0.20	0.12	0.28	0.15	0.12	0.34*	0.31	0.37*	0.37*	-0.03	0.34*	0.12		0.22	0.40*	-0.01	-0.06	0.09
				-																	
Litter	0.12	0.12	0.09	0.00	-0.15	-0.03	0.34*	0.09	-0.03	0.20	0.46**	0.59***	0.06	0.62***	-0.02	0.22		0.48**	-0.25	-0.23	-0.22
C.Ros	0.61***	0.60***	0.20	-0.23	-0.22	0.24	0.01	0.29	0.13	0.11	0.71***	0.93***	0.19	0.93***	0.16	0.40*	0.48**		0.11	-0.08	-0.13
SoilT_y	0.22	0.17	0.23	0.19	-0.25	0.07	-0.01	0.06	-0.01	-0.13	-0.06	0.03	0.03	0.01	-0.12	-0.01	-0.25	0.11		0.62***	0.03
SoilT_sf	0.06	0.06	0.24	0.21	-0.07	0.03	-0.15	0.10	-0.10	-0.21	-0.14	-0.08	0.06	-0.12	0.02	-0.06	-0.23	-0.08	0.62***		0.09
SoilT_s	-0.29	-0.27	-0.14	-0.21	0.36	-0.12	0.28	-0.19	-0.09	0.07	-0.15	-0.09	0.16	-0.14	0.22	0.09	-0.22	-0.13	0.03	0.09	

40

Table B3. Accumulated annual (1.11.2021–31.10.2022, AccAnnual) and snow season (1.11.2021–12.5.2022 and 27.10.–31.10.2022, AccSnow) methane flux rates in mg CH₄/m² and the portion of snow season fluxes of the annual flux (%SnowOfAnnual) calculated for each plot from the manual static chamber measurements. Ordered by the highest portion of annual flux.

Plot	AccAnnual (mg CH ₄ /m ²)	AccSnow (mg CH ₄ /m ²)	%SnowOfAnnual
41	6859.75	1462.88	21.33
64	10490.26	2050.96	19.55
71	3907.02	724.26	18.54
48	14210.27	2261.19	15.91
72	3901.88	605.24	15.51
46	15486.28	2307.50	14.90
49	8821.98	1256.22	14.24
67	2262.12	244.18	10.79
55	11231.20	1202.56	10.71
69	3623.34	375.92	10.37
62	6730.06	660.54	9.81
58	14024.16	1372.14	9.78
68	3992.76	342.10	8.57
52	12463.48	981.63	7.88
59	9800.62	771.15	7.87
57	9525.56	745.81	7.83
40	9417.67	725.66	7.71
70	4104.88	313.75	7.64
42	19409.26	1451.68	7.48
63	9436.37	678.56	7.19
38	15134.87	1037.47	6.85
39	16616.16	1113.47	6.70
56	14225.52	939.73	6.61
50	19267.61	1263.10	6.56
43	16525.27	1078.93	6.53
65	9711.21	620.80	6.39
54	6300.25	387.94	6.16
51	13298.34	750.18	5.64
47	19544.40	1064.65	5.45
61	9024.76	486.75	5.39
60	7824.93	414.35	5.30
37	16767.20	823.34	4.91
44	15123.26	640.19	4.23
53	8097.61	339.17	4.19
45	13906.22	419.71	3.02
66	11038.68	250.06	2.27
TOTAL	392105.20	32163.79	8.20



50

Figure B6. The relation of snow level manipulation experiment's treatments and methane flux in different time periods. There were no significant ($p < 0.05$) differences between the treatments in relation to the fluxes. Control plots represented the natural snow level, while in removal plots, the snow depth was maintained at 0.25 m by shovelling the snow and distributing it evenly onto the addition plots.



55

Figure B7. Location of the study plots at Puukkosuo fen segregated by vascular plant clusters. Indicator species most strongly connected to the vascular plant clusters: C.ros = *C. rostrata*, C.cho = *C. chordorrhiza* and T. ces = *C. lasiocarpa*, *P. erecta*, and *T. cespitosum*. Orthomosaic © Petra Korhonen 2024.