

Measurement report: Variations and environmental impacts of atmospheric N₂O₅ concentrations in urban Beijing during the 2022 Winter Olympics

Tiantian Zhang¹, Peng Zuo², Yi Chen², Tong Liu¹, Linghan Zeng², Weili Lin^{1,3*}, Chunxiang Ye^{2*}

¹NEAC Key Laboratory of Ecology and Environment in Minority Areas, College of Life and Environmental Sciences, Minzu University of China, Beijing, 100081, China

²State Key Joint Laboratory for Environmental Simulation and Pollution Control, College of Environmental Sciences and Engineering, Peking University, Beijing, 100871, China

³Institute of National Security, Minzu University of China, Beijing, 100081, China

Correspondence to: Weili Lin (linwl@muc.edu.cn) and Chunxiang Ye (c.ye@pku.edu.cn)

Abstract. The chemistry of nitrate radical (NO_3) and dinitrogen pentoxide (N_2O_5) plays a pivotal role in tropospheric nighttime chemistry. Given their close linkage to precursor variations, emission reduction during the 2022 Beijing Winter Olympics likely affected NO_3 and N_2O_5 behavior. In this study, we measured N_2O_5 , NO_2 , O_3 , etc. during and after the Olympics, and compared pollutant levels as well as the contributions of reaction pathways to the loss of NO_3 and N_2O_5 . Throughout the entire observation period, NO_3 production rate averaged 0.5 ± 0.4 ppbv h^{-1} , and the N_2O_5 mixing ratio could reach up to 875 pptv within 1 minute, indicating their active production. The relatively long nighttime N_2O_5 lifetime ($\tau_{\text{N}_2\text{O}_5}$), with an average of 11.9 ± 11.8 minutes, suggested a slow N_2O_5 loss rate during winter season. Despite low NO mixing ratio (below 3 ppbv), it dominated NO_3 loss (79.0%). VOCs oxidation contributed 0.2%, primarily driven by styrene. During the Olympics, emission reductions led to decreased NO and VOCs, which in turn reduced their reaction with NO_3 . The heterogeneous uptake of N_2O_5 , another key NO_3 loss pathway—accounted for 20.8% of NO_3 loss during the Olympics, but this contribution decreased to 10.6% after the Olympics. This uptake is crucial for nighttime NO_3 removal and would be essential for winter nitrate formation in urban Beijing. Our results highlight that under emission control scenarios, the relative importance of heterogeneous processes in nocturnal NO_3 cycling increases, providing new insights into how emission reduction measures shape nighttime oxidation processes in polluted urban environments.

1 Introduction

Nitrate radical (NO_3) and dinitrogen pentoxide (N_2O_5) play crucial roles in the nocturnal atmospheric chemical cycle, controlling the removal and conversion of nitrogen oxides (NO_x) and volatile organic compounds (VOCs). They significantly contribute to the formation of nitrate and secondary organic aerosols during the nighttime (Crutzen, 1979; Wayne et al., 1991). NO_3 primarily originates from the reaction of NO_2 with O_3 (R1), while it rapidly establishes a thermodynamic equilibrium (R2) with N_2O_5 . This tight coupling species are frequently considered simultaneously in atmospheric chemistry studies. During daytime, however, the rapid photolysis of NO_3 (R3) and its reaction with NO (R4) result in a short NO_3 lifetime (< 5 s). Consequently, the concentrations of NO_3 and N_2O_5 usually fall below the detection limit of analytical instruments during daylight hours.

The direct removal pathways of NO_3 include heterogeneous reactions on particulate matter surfaces and gas-phase reactions with NO (R4) and VOCs (R5), which can influence the atmospheric lifetimes of nighttime NO_x and VOCs (Ng et al., 2017; Wayne et al., 1991). NO_3 is also capable of reacting with alkenes in an addition reaction (R5) and subsequently with O_2 to form peroxy radicals (RO_2), which further generates organic nitric acid, one of the important sources of secondary organic aerosols (Fry and Sackinger, 2012; Hoyle et al., 2007; Pye et al., 2010). The removal pathways of N_2O_5 represent indirect removal pathways for NO_3 chemistry (R6), primarily involving reactions with water vapor and heterogeneous reactions on cloud droplets or particle surfaces (Brown and Stutz, 2012; Chang et al., 2011).





64 In recent years, anthropogenic emission control measures have played a pivotal role in improving air pollution
 65 in China with notable declines in NO_x emissions (Li et al., 2020; Zhang et al., 2016). However, NO_x emission
 66 intensity remains relatively high (Li et al., 2024). Reactive nitrogen-containing compounds have emerged as a
 67 prominent factor in China's complex air pollution scenario (Zhu et al., 2023). With advancements in
 68 measurement techniques, several research teams have explored the core processes of reactive nitrogen species
 69 in atmospheric pollution, particularly in regions with severe atmospheric pollution such as the North China
 70 Plain, the Yangtze River Delta, and the Pearl River Delta (Li et al., 2020; Tham et al., 2016; Wang et al., 2018,
 71 2020, 2024, 2013; Yun et al., 2018). While NO_3 reactivity is typically attenuated under low-temperature winter
 72 conditions, thereby restricting its oxidative capacity, multiple studies—including winter campaigns such as Yun
 73 et al., (2018) and Yan et al., (2023)—have demonstrated the significance of nocturnal NO_3 chemistry in cold
 74 seasons.

75 Emerging evidence further highlights the sensitivity of nocturnal NO_3 chemistry to emission controls: increases
 76 in nocturnal NO_3 production rates enhance nighttime oxidation capacity (Wang et al., 2023a), and urban
 77 nocturnal NO_3 concentrations may even experience “explosive” growth under stringent emission reduction
 78 policies (Wang et al., 2023b). In the Beijing region specifically, Yan et al., (2023) showed that the contribution
 79 of nocturnal nitrogen chemistry to winter haze has increased in recent years. Despite these advances, gaps
 80 remain in our understanding how short-term, large-scale emission control measures—such as those
 81 implemented during major events—influence the balance between $\text{NO}_3/\text{N}_2\text{O}_5$ production and loss, and their
 82 subsequent impacts on nighttime secondary pollution.

83 The 2022 Beijing Winter Olympics (BWO) provided a unique opportunity to address this gap. To ensure high
 84 air quality during the event, a series of strict emission reduction measures were implemented in Beijing and its
 85 surrounding areas, leading to a significant decline in $\text{PM}_{2.5}$ concentrations (average $24 \mu\text{g m}^{-3}$) compared to
 86 historical levels for the same period (Huang et al., 2024). Given the close linkage between $\text{NO}_3/\text{N}_2\text{O}_5$ and their
 87 precursors (NO_2 , O_3 , and VOCs), this emission cuts were expected to alter the behavior of NO_3 and N_2O_5 .

88 In this study, we conducted field observations of N_2O_5 , NO_2 , O_3 , NO , VOCs, and meteorological parameters in
 89 urban Beijing during (5–20 February 2022) and after (21 February–3 March 2022) the BWO. Our objectives

were to: (1) characterize the temporal variations of N_2O_5 and NO_3 in urban Beijing during winter; (2) quantify the contributions of different reaction pathways to NO_3 and N_2O_5 loss; and (3) assess how the BWO emission reduction measures modulated nocturnal $\text{NO}_3/\text{N}_2\text{O}_5$ chemistry. The findings provide critical insights into the response of nighttime reactive nitrogen chemistry to short-term emission controls, with implications for air quality management in polluted urban environments.

2 Methods

2.1 Site description

Field measurements were conducted from 5 February to 3 March 2022 at an urban site in Beijing. The observation location was situated on the rooftop of the NO. 1 Science Building at Peking University in Beijing (39.99° N , 116.31° E , 61.6 m asl). As shown in Figure 1, the location is proximal to the North Fourth Ring Road—one of Beijing’s major traffic arteries—and within 1 km of two primary traffic corridors (east-west along the North Fourth Ring Road and north-south along Zhongguancun Street). The surrounding area features mixed land use, including residential complexes (within 500 m) and low-intensity commercial facilities (within 1 km), with no large industrial sources within a 5 km radius. This setting makes the site representative of a typical urban mixed-use area impacted by fresh anthropogenic emissions (e.g., traffic-related NO_x and VOCs), consistent with previous characterizations of this location (Hu et al., 2023; Wang et al., 2017a; Yao et al., 2023).




Figure 1. Measurement site, surroundings, and wind rose (winter 2022). Base map adapted from <https://map.baidu.com/>); wind rose generated from on-site meteorological observations.

Sampling inlets were installed 1.5 meters above the rooftop, approximately 20 meters above ground level. Throughout the measurement period, the prevailing wind direction was predominantly from the northwest, with an average wind speed of $2.0 \pm 2.0 \text{ m s}^{-1}$.

2.2 Instrument setup

2.2.1 Measurement of NO₃ and N₂O₅

Ambient concentrations of NO₃ and N₂O₅ were determined utilizing an in-house-developed cavity ring-down spectroscopy (CRDS) analyzer, operating at a wavelength of 662 nm (Zhang et al., 2024). The analyzer employs a dual-channel design for simultaneous detection: Channel 1 directly measures NO₃ concentrations under ambient temperature conditions; Channel 2 maintained at 150 °C to thermally decompose N₂O₅ into NO₃, enabling quantification of the total concentration of [NO₃ + N₂O₅].

During this observation campaign, the NO₃-specific detection channel (Channel 1) malfunctioned due to the damage of the mirror, limiting measurements to the combined [NO₃ + N₂O₅] signal from Channel 2. NO₃ concentrations were subsequently derived using the thermodynamic equilibrium relationship between NO₃ and N₂O₅ (Eq. (1) in Section 2.3), with input parameters including measured N₂O₅, NO₂ concentrations, and ambient temperature. Under winter conditions (low temperature and relatively high NO₂ levels), the NO₃/N₂O₅ ratio is inherently low (~1%), ensuring that [NO₃ + N₂O₅] is dominated by N₂O₅ and derived NO₃ concentrations are reliable.

The limit of detection (LOD) of the CRDS system was calculated as 2.9 pptv, with a measurement uncertainty of ±13.7% (Zhang et al., 2024). To ensure measurement accuracy, regular calibrations were performed using a stable dynamic generation system for NO₃ and N₂O₅ standard gases (Zhang et al., 2026). The detailed technical specifications of the instrument are provided in Text S1 of the Supporting Information.

2.2.2 Measurement of other species

Concentrations of NO, NO₂, O₃, VOCs, and meteorological parameters were measured using commercial or well-validated analytical instruments, with details summarized in Table 1. NO and O₃ mixing ratios were measured using commercial instruments, specifically the Model 42i-Y and Model 49i from Thermo Fisher Scientific (USA). NO₂ mixing ratios were observed via a cavity-enhanced absorption spectroscopy (CEAS) (Zhou et al., 2022). Calibrations of these instruments are performed weekly using the standard gases of known concentrations, and the R² of the standard curve for each calibration is greater than 0.99. VOC concentrations were determined using a gas chromatograph equipped with mass spectrometry and flame ionization detectors (Wang et al., 2014). This system measures 99 VOC species with a time resolution of 1 hour, LOD range of 1–26 pptv, and accuracy of 0.8–6.1%. Quality control included weekly zero/span checks (using ultra-high-purity nitrogen and a multi-component VOC standard) and a post-campaign full calibration, which confirmed linearity (R² > 0.996) and negligible intercepts for all target VOCs. The photolysis rate constants (*j*-values) were obtained using a spectroradiometer (Metcon CCD-Spectrograph, Garmisch-Partenkirchen, Germany) (Bohn et al., 2008). This instrument quantifies two primary photolysis channels (NO₃ + *hν* → NO₂ + O(³P) and NO₃ + *hν* → NO + O₂), with total *j*(NO₃) calculated as the sum of the two channel-specific rate constants (*j*(NO₃)_{total} = *j*(NO₃)_M

+ $j(\text{NO}_3)_R$). Meteorological parameters, including wind direction, wind speed, temperature (T), and relative humidity (RH), were monitored utilizing a sensor meteorological measurement system (Metone, USA). The $\text{PM}_{2.5}$ concentration data were obtained from the Beijing Municipal Ecological and Environmental Monitoring Center (bjmemc.com.cn). Detailed information about these instruments is listed in Table 1.

Table 1 Species measured in the field work

Species	Time resolution (s)	Limit of Detection/working range	Methods	Accuracy (%)	References
O_3	60	1 ppbv (parts per billion by volume)	UV photometry	$\pm 1 \%$	-
NO_2	30	8 pptv	CEAS	$< 6 \%$	(Zhou et al., 2022)
NO	60	50 pptv	chemiluminescence	$\pm 1 \%$	-
N_2O_5	60	2.9 pptv	CRDS	$\pm 18 \%$	(Zhang et al., 2024)
VOCs	3600	1–26 pptv	GC-MS/FID	0.8 % – 6.1 %	(Wang et al., 2014)
T	60	-50–50°C	A three-element composite thermistor	$\pm 0.1^\circ\text{C}$	www.metone.com
RH	60	0–100 %	Thin film polymer capacitor	$\pm 0.2 \%$	www.metone.com

2.3 Calculation methods

When N_2O_5 concentrations are obtained, the concentration of NO_3 can be determined by dividing the concentration of N_2O_5 by the equilibrium constant K_{eq} and the concentration of NO_2 (Osthoff et al., 2006; Wang et al., 2017b), which is specified in Eq. (1).

$$[\text{NO}_3] = \frac{[\text{N}_2\text{O}_5]}{K_{\text{eq}}[\text{NO}_2]} \quad (1)$$

Here, K_{eq} represents the temperature-dependent equilibrium constant established when NO_3 attains steady-state equilibrium with N_2O_5 , and is given by $5.50 \times 10^{-27} \times \exp(10724/T)$, where T is the temperature in Kelvin (Wang et al., 2024).

The primary source of NO_3 and N_2O_5 is the chemical reaction of NO_2 with O_3 . Consequently, the concentrations of NO_2 and O_3 are key factors influencing the production rate of NO_3 ($P(\text{NO}_3)$). This production rate is mathematically represented by Eq. (2) (Brown et al., 2011). Assuming that the formation and loss processes of NO_3 and N_2O_5 are in a state of dynamic equilibrium, the lifetime of N_2O_5 , denoted as $\tau_{\text{N}_2\text{O}_5}$, can be expressed as the ratio of its concentration to the rate of NO_3 production, as determined by Eq. (3) (Brown and Stutz, 2012;

163 Lin et al., 2022; Wang et al., 2017a).

$$164 \quad P(\text{NO}_3) = k_{\text{NO}_2+\text{O}_3} \times [\text{NO}_2] \times [\text{O}_3] \quad (2)$$

$$165 \quad \tau_{\text{N}_2\text{O}_5} = \frac{[\text{N}_2\text{O}_5]}{P(\text{NO}_3)} = \frac{[\text{N}_2\text{O}_5]}{k_{\text{NO}_2+\text{O}_3} \cdot [\text{NO}_2] \cdot [\text{O}_3]} \quad (3)$$

166 The total reactivity of NO_3 (k_{NO_3}) represents the sum of all first-order loss processes for NO_3 , including
 167 photolysis, reaction with NO , reaction with VOCs, and indirect loss via N_2O_5 heterogeneous uptake. It is
 168 calculated using Eq. (4) (Wang et al., 2020): The nocturnal NO_3 loss rate, denoted as $L(\text{NO}_3)$, is calculated via
 169 Eq. (5).

$$170 \quad k_{\text{NO}_3} = j(\text{NO}_3) + k_{\text{NO}_3+\text{NO}} \cdot [\text{NO}] + k_{\text{NO}_3+\text{VOC}_i} \cdot [\text{VOC}_i] + k_{\text{N}_2\text{O}_5} \cdot K_{\text{eq}} \cdot [\text{NO}_2] \quad (4)$$

$$171 \quad L(\text{NO}_3) = \sum k_i [\text{VOC}_i] \cdot [\text{NO}_3] + k_{\text{NO}_3+\text{NO}} \cdot [\text{NO}] [\text{NO}_3] + k_{\text{N}_2\text{O}_5} \cdot [\text{N}_2\text{O}_5] \quad (5)$$

172 Here, $j(\text{NO}_3)$ denotes the photolysis rate constant for NO_3 decomposition. The rate coefficients $k_{\text{NO}_2+\text{O}_3}$ and
 173 $k_{\text{NO}_3+\text{NO}}$ correspond to the rate coefficients for reaction Eqs. (1) and (4), respectively, as referenced in Atkinson
 174 et al., (2004). The reactivity of NO_3 with VOCs ($k_{\text{NO}_3+\text{VOC}_i}$) is characterized by a first order loss rate constant,
 175 calculated as the product of the reaction rate constant k_i and the VOC concentrations $[\text{VOC}_i]$.

176 $k_{\text{N}_2\text{O}_5}$ represents the total first-order loss rate coefficient for the heterogeneous uptake of N_2O_5 at the aerosol
 177 surface, which is governed by the uptake coefficient $\gamma(\text{N}_2\text{O}_5)$, the aerosol surface area density S_a ($\mu\text{m}^2 \text{ cm}^{-3}$),
 178 and the mean molecular speed of N_2O_5 , c , as described in Eq. (6).

$$179 \quad k_{\text{N}_2\text{O}_5} = \frac{1}{4} c S_a \gamma(\text{N}_2\text{O}_5) \quad (6)$$

180 The mean molecular speed c is calculated as $\sqrt{8RT/(\pi M)}$, where R is the gas constant ($8.314 \text{ J mol}^{-1} \text{ K}^{-1}$), T is
 181 temperature (K), and M is the molar mass of N_2O_5 ($0.108 \text{ kg mol}^{-1}$). $\gamma(\text{N}_2\text{O}_5)$ is a critical parameter describing
 182 the efficiency of N_2O_5 uptake on aerosol surfaces, influenced by aerosol composition (e.g., organic coatings,
 183 nitrate/sulfate content) and meteorological conditions (RH, temperature) (Bertram et al., 2009; Tang et al., 2014;
 184 Yu et al., 2020). It was determined using the steady-state method (Brown et al., 2016; Lin et al., 2022; Lu et al.,
 185 2022), which relies on linear regression of $K_{\text{eq}}[\text{NO}_2]\tau(\text{N}_2\text{O}_5)^{-1}$ and $\frac{1}{4} c S_a K_{\text{eq}}[\text{NO}_2]$ (Eq. (7)).

$$186 \quad K_{\text{eq}}[\text{NO}_2] \tau(\text{N}_2\text{O}_5)^{-1} = \frac{1}{4} c S_a \gamma(\text{N}_2\text{O}_5) K_{\text{eq}}[\text{NO}_2] + k_{\text{NO}_3} \quad (7)$$

187 Here, the slope of the regression equals $\gamma(\text{N}_2\text{O}_5)$, and the intercept equals k_{NO_3} . To minimize interference from
 188 non-steady-state conditions, data fitting was data fitting was performed under the following constraints:

(1) Meteorological constraint: $RH < 70\%$ (avoiding excessive water vapor interference); (2) Chemical constraints: $NO < 1$ ppbv (suppressing $NO-NO_3$ titration), $[N_2O_5] > LOD$ (2.9 pptv, ensuring reliable equilibrium calculations); (3) Timing constraint: Data selected 2–3 hours post-sunset (when the $NO_3-N_2O_5$ equilibrium is most stable). Negative intercepts (physically implausible, arising from k_{NO_3}) were excluded, resulting in 23 valid data points for $\gamma(N_2O_5)$ calculation.

Aerosol surface area density (S_a)

Due to the unavailability of direct particle size distribution measurements, S_a was derived from $PM_{2.5}$ concentrations using an empirical formula validated for winter Beijing conditions (Zhang et al., 2022):

$$S_a = 60.03 \times [PM_{2.5}]^{0.62} \quad (8)$$

This formula exhibits a strong linear correlation ($R^2 = 0.82$) with $PM_{2.5}$ and is applicable for $PM_{2.5}$ concentrations $< 200 \mu g m^{-3}$ —consistent with the $PM_{2.5}$ range observed in this study (average: $24 \pm 21 \mu g m^{-3}$, maximum: $131 \mu g m^{-3}$).

3 Results

3.1 Measurements overview

Figure 2 illustrates time-series variations in the mixing ratios of N_2O_5 and associated trace gases, alongside meteorological parameters, captured during the 2022 Beijing Winter Olympics (BWO) at a temporal resolution of 1 minute. Valid data were systematically acquired over a 26-day span, from 5 February to 3 March. In alignment with the 2022 Beijing Winter Olympics timeline, the observation interval was bifurcated into two distinct periods: (1) the Olympic Games Period (OGP; spanning 5-20 February), and (2) the Post-Olympics Period (POP; extending from 21 February to 3 March). Comprehensive statistical metrics for each period are meticulously detailed in Table 2.




Figure 2. Time series for N_2O_5 , NO_3 , related trace gases, and meteorological data (T and RH) measured in Beijing from 5 February to 3 March 2022.

3.1.1 Meteorological conditions and $\text{PM}_{2.5}$

Meteorological conditions differed notably between the OGP and POP, driving variations in pollutant dispersion and chemical reactivity:

Temperature: Nocturnal temperatures during the OGP were predominantly below freezing (average: -1.4 ± 3.6 °C), while the POP saw a marked warming trend—nocturnal temperatures rose to 3.5 ± 3.5 °C (all-day average: 5.6 ± 3.9 °C, vs. -0.4 ± 3.9 °C in the OGP).

Relative Humidity (RH): The OGP exhibited higher RH (nocturnal average: $29 \pm 13\%$) compared to the POP (nocturnal average: $20 \pm 4\%$). A heavy snowfall event occurred during 13–14 February (OGP), coinciding with peak RH ($> 60\%$) and a transient increase in $\text{PM}_{2.5}$ concentration ($68 \mu\text{g m}^{-3}$).

$\text{PM}_{2.5}$: The overall average $\text{PM}_{2.5}$ concentration across the entire observation period was $24 \pm 21 \mu\text{g m}^{-3}$, consistent with the improved air quality during the BWO (Huang et al., 2024). While $\text{PM}_{2.5}$ levels were similar

between the OGP (nocturnal average: $26 \pm 2 \mu\text{g m}^{-3}$) and POP (nocturnal average: $23 \pm 2 \mu\text{g m}^{-3}$), the POP experienced a severe pollution episode with $\text{PM}_{2.5}$ peaking at $131 \mu\text{g m}^{-3}$ —likely driven by relaxed emissions and stagnant meteorological conditions.

The aerosol surface area density, a key parameter for N_2O_5 heterogeneous uptake calculations (Eq. (8)), was derived from $\text{PM}_{2.5}$ concentrations. The overall average S_a was $402 \pm 215 \mu\text{m}^2 \text{cm}^{-3}$, with values in the OGP (nocturnal average: $417 \pm 208 \mu\text{m}^2 \text{cm}^{-3}$) slightly higher than in the POP (nocturnal average: $385 \pm 205 \mu\text{m}^2 \text{cm}^{-3}$), reflecting the influence of $\text{PM}_{2.5}$ and RH-driven aerosol growth.

Table 2. Summary of observed parameters for the two periods (mean $\pm$ standard deviation).

Species	All time	OGP		POP	
		All day	Nighttime	All day	Nighttime
O_3 (ppbv)	28.6 ± 12.8	29.9 ± 9.5	27.4 ± 10.3	26.7 ± 10.6	19.8 ± 12.1
NO_2 (ppbv)	14.8 ± 11.5	12.6 ± 8.2	14.5 ± 9.3	18.2 ± 12.3	20.7 ± 13.1
NO (ppbv)	3.5 ± 7.2	1.9 ± 2.3	1.0 ± 1.2	5.7 ± 6.1	4.8 ± 6.0
N_2O_5 (pptv)	86.7 ± 116.5	87.3 ± 71.6	137.6 ± 112.7	62.1 ± 57.7	97.8 ± 90.3
NO_3 (pptv)	0.6 ± 0.7	0.4 ± 0.4	0.6 ± 0.6	0.3 ± 0.4	0.5 ± 0.6
Total VOCs (ppbv)	17.36 ± 10.10	15.67 ± 7.45	16.02 ± 7.74	19.72 ± 11.93	19.68 ± 12.17
$\text{PM}_{2.5}$ ($\mu\text{g m}^{-3}$)	24 ± 21	25 ± 2	26 ± 2	23 ± 3	23 ± 2
T ($^{\circ}\text{C}$)	2.1 ± 5.7	-0.4 ± 3.9	-1.4 ± 3.6	5.6 ± 3.9	3.5 ± 3.5
RH (%)	24 ± 12	27 ± 13	29 ± 13	19 ± 4	20 ± 4
$P(\text{NO}_3)$ (ppbv h^{-1})	0.5 ± 0.4	0.5 ± 0.2	0.5 ± 0.2	0.6 ± 0.4	0.5 ± 0.3
$\tau_{\text{N}_2\text{O}_5}$ (min)	11.9 ± 11.8	10.9 ± 17.0	17.0 ± 17.0	7.4 ± 4.4	11.6 ± 6.8

3.1.2 Precursor gases (NO , NO_2 , O_3)

Concentrations of NO_x ($\text{NO} + \text{NO}_2$) and O_3 —key precursors for $\text{NO}_3/\text{N}_2\text{O}_5$ production (Reaction R1)—exhibited distinct differences between the OGP and POP, directly reflecting the impact of emission controls:

NO: The OGP observed significantly lower NO concentrations compared to the POP—nocturnal NO averaged 1.0 ± 1.2 ppbv (OGP) vs. 4.8 ± 6.0 ppbv (POP). This contrast confirms BWO emission reductions effectively curbed primary NO emissions (e.g., traffic, industry), which further modulated O_3 and NO_2 dynamics.

NO_2 : Inverse to NO , NO_2 concentrations were higher in the POP. Nocturnal NO_2 averaged 20.7 ± 13.1 ppbv in the POP, compared to 14.5 ± 9.3 ppbv in the OGP. The increase in NO_2 during the POP is attributed to enhanced NO oxidation by O_3 , driven by higher post-Olympic NO emissions.

O₃: Nocturnal O₃ levels were substantially higher in the OGP (27.4 ± 10.3 ppbv) than in the POP (19.8 ± 12.1 ppbv), despite similar all-day averages (OGP: 29.9 ± 9.5 ppbv; POP: 26.7 ± 10.6 ppbv). The lower nocturnal O₃ in the POP results from intensified O₃ titration by elevated NO, a process that simultaneously increases NO₂ concentrations. Notably, the overall mean O₃ concentration (28.6 ± 12.8 ppbv) was lower than spring values in Beijing (Wang et al., 2018), consistent with reduced photochemical O₃ production in winter.

The average NO_x concentration (18.2 ± 16.6 ppbv) during the study was also substantially lower than autumn values at this site (typically > 30 ppbv) (Li et al., 2022; Wang et al., 2017a), further highlighting the effectiveness of winter emission controls.

3.1.3 N₂O₅ and NO₃ concentrations

N₂O₅ concentrations exhibited significant temporal variability throughout the observation period, with a mean daily value of 86.7 ± 116.5 pptv and a maximum of 874.9 pptv (00:15 LST on 18 February, OGP)—coinciding with moderate NO₂ (14.6 ppbv), O₃ (26.8 ppbv) and extremely low NO (0.4 ppbv) that minimized NO₃ titration.

NO₃ concentrations were derived from the NO₃-N₂O₅ equilibrium (Eq. (1)), with a mean value of 0.6 ± 0.7 pptv (maximum: 4.6 pptv). Nocturnal NO₃ levels followed the same trend as N₂O₅: higher in the OGP (0.6 ± 0.6 pptv) than in the POP (0.5 ± 0.6 pptv). This derived NO₃ concentration is notably lower than observations in Shanghai (16 ± 9 pptv) (Wang et al., 2013), likely due to Beijing's lower winter temperatures (which suppress NO₃ production) and higher NO emissions (which enhance NO₃ loss).

3.1.4 NO₃ production rate and N₂O₅ lifetime

The NO₃ production rate, calculated via Eq. (2) as the product of the NO₂ + O₃ reaction rate constant and precursor concentrations, averaged 0.5 ± 0.4 ppbv h⁻¹ across the entire observation period, with a maximum of 2.4 ppbv h⁻¹. This value aligns with winter observations in the Beijing area (0.4 ppbv h⁻¹) (Wang et al., 2021) and summer measurements at Mount Tai (0.45 ± 0.40 ppbv h⁻¹) (Wang et al., 2017c), but is lower than autumn values in Beijing (2.25 ± 2.02 ppbv h⁻¹) (Wang et al., 2017a) and summer measurements in Taizhou (1.2 ± 0.3 ppbv h⁻¹) (Li et al., 2020). The relatively low $P(\text{NO}_3)$ in this study is primarily driven by winter's low temperatures, which reduce the NO₂ + O₃ reaction rate constant: for example, at identical NO₂ (15 ppbv) and O₃ (30 ppbv) concentrations, increasing temperature from -1 °C to 5 °C raises the rate constant from 1.59×10^{-17} to 1.94×10^{-17} cm³ molecule⁻¹ s⁻¹, leading to a 19% increase in $P(\text{NO}_3)$ (from 0.70 to 0.83 ppbv h⁻¹).

The N₂O₅ lifetime, a key indicator of N₂O₅ removal efficiency (Eq. (3)), averaged 11.9 ± 11.8 minutes across the study—longer than summer values in Beijing (270 ± 240 seconds) (Wang et al., 2018) and rural Wangdu (77–172 seconds) (Tham et al., 2016), but shorter than observations in the Hong Kong boundary layer (Brown et al., 2016). This prolonged winter $\tau_{\text{N}_2\text{O}_5}$ suggests slower nocturnal NO₃/N₂O₅ loss, consistent with lower winter temperatures and reduced heterogeneous reactivity.

Notably, $\tau_{\text{N}_2\text{O}_5}$ differed significantly between the OGP and POP: the nocturnal mean lifetime was 17.0 ± 17.0 minutes in the OGP, compared to 11.6 ± 6.8 minutes in the POP—a ~5-minute reduction. This difference is primarily driven by variations in N_2O_5 concentrations (rather than $P(\text{NO}_3)$, which was similar between periods: OGP nocturnal $P(\text{NO}_3) = 0.5 \pm 0.2$ ppbv h^{-1} ; POP = 0.5 ± 0.3 ppbv h^{-1}).

3.2 Mean diurnal variations

Figure 3 displays the average diurnal variations in NO , NO_2 , N_2O_5 , NO_3 , O_3 mixing ratios, and $P(\text{NO}_3)$ during the study period. Specifically, panel (a) presents the daily mean patterns for the OGP, whereas panel (b) depicts those for the POP. Notable differences in concentration but similar diurnal trends were observed between the two periods.




Figure 3. Mean diurnal variations in NO , NO_2 , N_2O_5 , NO_3 , O_3 mixing ratios and $P(\text{NO}_3)$ during (a) the Olympic Games Period (OGP) and (b) the Post-Olympics Period (POP). Data represent hourly means with error bars indicating the standard deviation; non-physical values < 0 (for NO_3) have been excluded.

3.2.1 Diurnal cycles of precursor gases (NO , NO_2 , O_3)

The diurnal variations of NO , NO_2 , and O_3 exhibit strong interdependencies, consistent with well-documented atmospheric chemical processes (e.g., O_3 titration by NO , NO_2 photolysis) and anthropogenic emission patterns.

Nocturnal NO mixing ratios were substantially lower in the OGP than in the POP, with OGP nighttime values remaining below 2 ppbv (average: 1.0 ± 1.2 ppbv) and POP values frequently exceeding 4 ppbv (average: 4.8 ± 6.0 ppbv). Both periods showed two distinct daily peaks in NO , aligned with morning (06:00–08:00 LST)

and evening (18:00–20:00 LST) traffic rush hours—confirming traffic as the primary NO source. The morning peak was particularly pronounced in the POP, reaching 20.4 ppbv at 08:00 LST (vs. 6.4 ppbv at 06:00 LST in the OGP), directly attributable to relaxed post-Olympic traffic emission controls.

NO₂ concentrations displayed an inverse diurnal trend to O₃, with nocturnal levels consistently higher than daytime values (OGP: 14.5 ± 9.3 ppbv nighttime vs. 12.6 ± 8.2 ppbv all-day; POP: 20.7 ± 13.1 ppbv nighttime vs. 18.2 ± 12.3 ppbv all-day). This pattern arises from reduced daytime NO₂ photolysis ($\text{NO}_2 + h\nu \rightarrow \text{NO} + \text{O}$) and enhanced nocturnal NO oxidation. The POP saw higher NO₂ concentrations across all hours, with the nocturnal peak (24.3 ppbv at 21:00 LST) exceeding the OGP peak (15.9 ppbv at 21:00 LST) by ~35%—a result of elevated NO emissions driving more O₃-to-NO₂ conversion.

O₃ exhibited a classic mid-afternoon peak in both periods, rising gradually after sunrise (07:00 LST) as photochemical production intensified, and peaking between 15:00–16:00 LST (OGP: 39.9 ppbv; POP: 41.2 ppbv). Nocturnal O₃ levels, however, differed sharply: the OGP maintained higher nighttime O₃ (27.4 ± 10.3 ppbv) compared to the POP (19.8 ± 12.1 ppbv), with the POP O₃ concentration dropping to a minimum of 14.6 ppbv at 07:00 LST (vs. 22.2 ppbv in the OGP). This discrepancy stems from intensified O₃ titration by NO in the POP—higher NO concentrations rapidly consume O₃ via $\text{NO} + \text{O}_3 \rightarrow \text{NO}_2 + \text{O}_2$, reducing the O₃ pool available for NO₃ production (Reaction R1).

3.2.2 Diurnal cycles of NO₃ production rate

P(NO₃) exhibited a strong diurnal pattern tied to the availability of its precursors (NO₂ and O₃) and the suppression of daytime NO₃ loss.

In both periods, *P*(NO₃) peaked shortly after sunset (19:00 LST), when O₃ concentrations remained relatively high (OGP: 33.3 ppbv; POP: 30.6 ppbv) and NO emissions (and thus NO₃ titration) were still low. The OGP peak *P*(NO₃) was 0.74 ppbv h⁻¹, slightly lower than the POP peak (0.92 ppbv h⁻¹)—a difference driven by the POP's higher NO₂ concentrations (18.4 ppbv at 19:00 LST vs. 14.2 ppbv in the OGP) offsetting its lower O₃. Throughout the night, *P*(NO₃) gradually declined in both periods: by 04:00 LST, it dropped to 0.33 ppbv h⁻¹ (OGP) and 0.20 ppbv h⁻¹ (POP), mirroring the nocturnal decrease in O₃.

Notably, *P*(NO₃) showed a strong first-order exponential decay correlation with NO concentrations (Figure S2): when NO < 5 ppbv, *P*(NO₃) decreased sharply with increasing NO (from 1.2 ppbv h⁻¹ to 0.5 ppbv h⁻¹); when NO > 10 ppbv, *P*(NO₃) stabilized at < 0.3 ppbv h⁻¹. This confirms NO is a dominant inhibitor of NO₃ production—even small NO increases (e.g., POP rush hours) rapidly consume NO₃ as it forms.

3.2.3 Diurnal cycles of N₂O₅ and NO₃

N₂O₅ and NO₃ exhibited nearly identical diurnal patterns in both periods, reflecting their tight thermodynamic

equilibrium (Reaction R2) and shared dependence on precursor availability and loss pathways:

Post-sunset accumulation: Both species began accumulating rapidly after sunset (18:00 LST), as photolysis (a major daytime NO₃ loss pathway) ceased. They reached peak concentrations around 20:00 LST—shortly after the *P*(NO₃) peak—with OGP peaks (N₂O₅: 208.2 pptv; NO₃: 1.1 pptv) exceeding POP peaks (N₂O₅: 171.2 pptv; NO₃: 1.0 pptv) by ~22%. This difference is driven by the OGP’s lower NO concentrations, which reduce NO₃ titration (Reaction R4) and allow more N₂O₅ to form via equilibrium.

Nocturnal decline: After the 20:00 LST peak, N₂O₅ and NO₃ concentrations gradually decreased in the OGP, falling to near-detection limits by sunrise (07:00 LST). The POP, however, showed a steeper decline between 02:00–04:00 LST: N₂O₅ dropped from 94.1 pptv to 61.8 pptv, and NO₃ from 0.6 pptv to 0.3 pptv—attributed to elevated nocturnal NO emissions in the POP (peaking at 9.9 ppbv at 04:00 LST) that accelerate NO₃ loss.

Morning secondary peak (POP only): A notable secondary peak in N₂O₅ (112.1 pptv at 06:00 LST) and NO₃ (0.5 pptv at 06:00 LST) occurred in the POP, coinciding with a transient increase in O₃ (1.3 ppbv) and decrease in NO (2.9 ppbv) before morning rush hour. This peak is absent in the OGP, likely because lower POP emission controls led to a larger “pre-rush” O₃ pool that drives NO₃ production, while OGP NO emissions remained too low to support such a transient precursor balance.

The daily average variation trends of both N₂O₅ and NO₃ aligned with those reported for the Yangtze River Delta and North China regions (Li et al., 2020; Wang et al., 2022, 2017b). While the chemical conditions in this study bore similarities to those in summer Beijing, the meteorological conditions differed, notably characterized by higher relative humidity during the summer. The average nocturnal N₂O₅ concentration over the observation period was 113.7 ± 103.3 pptv, which was higher than that observed in the Changping area of Beijing (Wang et al., 2018), indicating that the loss process of NO₃ and N₂O₅ in Beijing during winter is more sluggish compared to that in the summer.

4 Discussion

4.1 Factoring influencing N₂O₅ lifetime

RH and *S_a* are well-recognized as pivotal factors regulating N₂O₅ lifetime in nocturnal atmospheric chemistry, as they directly modulate the efficiency of N₂O₅ heterogeneous uptake—the primary indirect loss pathway for NO₃ (Brown et al., 2017; Lin et al., 2022). The correlation between these parameters and $\tau_{\text{N}_2\text{O}_5}$ is presented in Figure 4.

4.1.1 Relationship between $\tau_{\text{N}_2\text{O}_5}$ and RH

As shown in Figure 4(a), the correlation between $\tau_{\text{N}_2\text{O}_5}$ and RH exhibits a humidity-dependent dual pattern. At RH < 35%, $\tau_{\text{N}_2\text{O}_5}$ increased with RH—slight humidity rises softened hydrophobic organic coatings on aerosols (from traffic VOC oxidation), thereby reducing N_2O_5 heterogeneous uptake. RH > 35%, the heterogeneous uptake rate of N_2O_5 increases, and $\tau_{\text{N}_2\text{O}_5}$ decreases with increasing RH.

RH < 35%: Counterintuitive $\tau_{\text{N}_2\text{O}_5}$ increases with rising RH. Minimal aerosol liquid water content drives hydrophobic organic components—primarily oxidation products of traffic-related anthropogenic VOCs (AVOCs, e.g., styrene, propylene)—to condense into dense, impermeable coatings on particle surfaces (Bertram et al., 2009; Folkers et al., 2003; McNeill et al., 2006; Tang et al., 2014). These coatings act as a diffusion barrier, preventing N_2O_5 from reaching reactive aqueous sites (e.g., nitrate/sulfate-rich droplets) and lowering the heterogeneous uptake coefficient $\gamma(\text{N}_2\text{O}_5)$ (Anttila et al., 2006; Yu et al., 2020). For example, at RH = 25%, $\tau_{\text{N}_2\text{O}_5}$ averaged 15.5 minutes, 43% longer than the 8.9 minutes observed at RH = 15%.

RH > 35%: $\tau_{\text{N}_2\text{O}_5}$ decreases with increasing RH (conventional trend). Above 35% RH, the relationship aligns with physical expectations: $\tau_{\text{N}_2\text{O}_5}$ declines as RH increases, driven by two synergistic effects. First, hygroscopic growth of aerosols (e.g., ammonium sulfate, sodium chloride) increases S_a , providing more reactive surface sites for N_2O_5 uptake. Second, elevated aerosol liquid water content accelerates N_2O_5 hydrolysis, which becomes the dominant N_2O_5 loss pathway (Brown et al., 2016; Chang et al., 2011). This effect is particularly pronounced when RH > 60%: during the snowfall events on 13–14 February 2022, $\tau_{\text{N}_2\text{O}_5}$ approached zero (0.8 ± 0.3 minutes), as high RH (> 85%) maximized both aerosol growth and hydrolysis efficiency.

Notably, scattered data points in the 30–40% RH range (e.g., $\tau_{\text{N}_2\text{O}_5}$ ranging from 2 to 33 minutes at RH = 35%) suggest interference from transient factors—such as sudden NO spikes or shifts in aerosol organic composition—which obscure the RH- $\tau_{\text{N}_2\text{O}_5}$ relationship. A more comprehensive discussion of these confounding factors requires consideration of the organic composition of the aerosol.




Figure 4. The relationship between $\tau_{\text{N}_2\text{O}_5}$ and S_a as well as RH during the observation period.

4.1.2 Relationship between $\tau_{\text{N}_2\text{O}_5}$ and S_a

Figure 4(b) depicts the dependence of $\tau_{\text{N}_2\text{O}_5}$ on S_a , reflecting the interplay between S_a and other regulating factors (e.g., RH, organic coatings).

$S_a < 325 \mu\text{m}^2 \text{cm}^{-3}$: $\tau_{\text{N}_2\text{O}_5}$ gradually increases with rising S_a . For low S_a values, $\tau_{\text{N}_2\text{O}_5}$ gradually rises from ~ 10 to 12 minutes as S_a increases. This non-monotonic pattern is driven by the co-occurrence of low S_a with extremely dry conditions (RH < 25% for 68% of data points in this S_a range).

$S_a = 500\text{--}1000 \mu\text{m}^2 \text{cm}^{-3}$: Robust negative correlation. Above a threshold S_a of $\sim 500 \mu\text{m}^2 \text{cm}^{-3}$, a clear negative correlation emerges: $\tau_{\text{N}_2\text{O}_5}$ decreases from ~ 12 to 6 minutes as S_a increases. This aligns with physical expectations, as higher S_a provides more reactive surface area for N_2O_5 heterogeneous uptake (Lin et al., 2022; Wang et al., 2020; Zhou et al., 2018).

Quantitative discrepancies between our results and summer studies (e.g., Zhou et al., 2018 reported $\tau_{\text{N}_2\text{O}_5} < 5$ minutes at $S_a > 600 \mu\text{m}^2 \text{cm}^{-3}$ in urban Beijing) are attributed to seasonal differences in aerosol liquid water content. Winter's lower RH (average: $24 \pm 12\%$) reduces aerosol liquid water content, lowering $\gamma(\text{N}_2\text{O}_5)$ and slowing N_2O_5 loss—even at high S_a . This highlights the need for season-specific parameterizations of S_a - $\tau_{\text{N}_2\text{O}_5}$ relationships in air quality models.

4.2 NO_3 and N_2O_5 loss pathways

To quantify the mechanisms governing NO_3 and N_2O_5 removal, we calculated total NO_3 reactivity via Eq. (4) and dissected the contributions of individual loss pathways. The reaction rate constants for the interaction between VOCs and the oxidizing agent NO_3 were obtained from the literature ((Atkinson and Arey, 2003; Brown et al., 2011) or extracted from the National Institute of Standards and Technology database (accessible via <http://webbook.nist.gov/chemistry/>). For certain VOC species where quantitative laboratory reaction rate constants were unavailable, these values were estimated based on the reaction rate constants of analogous species.

4.2.1 NO_3 Loss via VOC oxidation and N_2O_5 uptake

VOC-driven NO_3 loss is negligible in winter urban Beijing, but distinct patterns emerge in the concentrations and reactivity of different VOC categories. Detailed statistical data regarding VOC concentrations (e.g., styrene, isoprene, and other anthropogenic species) and their corresponding NO_3 reaction rates—are provided in Table S1 (auxiliary material). Time series plots of the concentrations of several highly reactive VOCs (Figure S3) show that the styrene concentration peaks at 86 pptv, while the isoprene concentration peaks at 96 pptv.

Comparative analysis reveals these high-VOC periods coincide with enhanced NO (e.g., NO spikes to 24.8 ppbv on February 24, POP), suggesting VOCs and NO share a common emission source (traffic exhaust)—consistent with the site’s proximity to urban traffic corridors (Section 2.1).

High-concentration AVOCs contribute minimally. The most abundant VOCs—ethane (3.8 ± 1.8 ppbv), propane (2.1 ± 1.3 ppbv), and acetone (1.4 ± 0.8 ppbv)—exhibit extremely low k_{NO_3} (e.g., $k(\text{NO}_3 + \text{propane}) = 9.49 \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 298 K (Atkinson and Arey, 2003)). As a result, their combined contribution to total VOC-driven NO_3 reactivity is $< 5\%$ ($0.04 \times 10^{-3} \text{ s}^{-1}$), emphasizing that high VOC concentration does not equate to strong NO_3 reactivity. When all AVOCs are considered, they dominate NO_3 reactivity ($\sim 70.4\%$ of total VOC-driven NO_3 loss), exceeding the contribution of biogenic VOCs (BVOCs, $\sim 29.6\%$) (Figure S4).

Reactive VOCs dominate VOC-driven NO_3 loss. Despite their low concentrations, styrene and isoprene account for $\sim 74\%$ of total VOC-driven NO_3 reactivity (Figure S4), due to their high k_{NO_3} .

Styrene: Average reactivity = $0.34 \times 10^{-3} \text{ s}^{-1}$ ($k = 1.5 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$), contributing $\sim 44\%$ of VOC reactivity. Styrene emissions in Beijing are primarily from vehicle exhaust, with minor contributions from evergreen plant emissions (Li et al., 2014).

Isoprene: Average reactivity = $0.25 \times 10^{-3} \text{ s}^{-1}$, contributing $\sim 30\%$ of VOC reactivity. Isoprene has dual sources: traffic exhaust (anthropogenic) and deciduous/evergreen plant emissions (biogenic), with biogenic sources dominating in winter (Cheng et al., 2018; Yuan et al., 2009).

Notably, biogenic VOCs (BVOCs) other than isoprene (e.g., limonene, α -pinene) were not detected, leading to potential underestimation of BVOC reactivity. For example, the rate constant for limonene ($\sim 1.6 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) is $\sim 20\times$ higher than isoprene’s, so including it could increase the total VOC reactivity.

$\gamma(\text{N}_2\text{O}_5)$ —a key parameter describing the efficiency of N_2O_5 uptake on aerosol surfaces—was calculated via Eq. (7) (steady-state method) with strict data selection, and the calculation results for S_a are presented in Figure S5. For most of the time, $\gamma(\text{N}_2\text{O}_5)$ ranged from 0.01 to 0.12 (average: 0.032 ± 0.049 ; Table S2), consistent with observations in urban Beijing (0.01–0.09) (Li et al., 2022; Wang et al., 2017a; Xia et al., 2021; Zhou et al., 2018) but higher than rural sites: Wangdu (Hebei), 0.006–0.034 (Tham et al., 2018); Hong Kong boundary layer, 0.014 ± 0.007 (Brown et al., 2016); Rural Germany, 0.028 ± 0.029 (Phillips et al., 2016). The elevated $\gamma(\text{N}_2\text{O}_5)$ in our study is attributed to urban aerosols’ higher water content and reactive composition (e.g., nitrate, sulfate, and organic acids), which enhance N_2O_5 hydrolysis efficiency (Bertram et al., 2009; Tang et al., 2014).

4.2.2 Temporal Variations in NO_3 Reactivity

To characterize how NO_3 loss dynamics respond to emission controls and transient environmental events, we analyzed the time series of total NO_3 reactivity—the sum of all first-order loss pathways, including reaction

with NO, heterogeneous uptake of N₂O₅, oxidation by VOCs, and photolysis (daytime only). This analysis is supported by Figure 5 (time series of total k_{NO_3} and key environmental drivers).




Figure 5. Time series variation of k_{NO_3} (reactions with NO and VOCs, heterogeneous uptake of N₂O₅ and photolysis of NO₃).

OGP: Total k_{NO_3} averaged 1.14 s⁻¹, with minimal day-to-day variability. A notable exception occurred on 13 February, when a heavy snowfall event (RH = 71%, PM_{2.5} = 61 µg m⁻³) triggered a transient spike in k_{NO_3} to 2.35 s⁻¹—nearly double the OGP average. This spike was driven by enhanced N₂O₅ heterogeneous uptake (Section 4.1.1), as high RH increased aerosol liquid water content and reactive surface sites.

POP: Total k_{NO_3} surged to 3.06 s⁻¹ (2.7× higher than OGP), driven by a 3.7× increase in NO reactivity (from 0.81 s⁻¹ to 3.00 s⁻¹; Table 3). The elevated NO reactivity aligns with post-Olympic NO emission rebound (Section 3.1.2), which intensified NO₃ titration (Reaction R4) and crowded out N₂O₅ uptake.

Figure 5 also captures the response of k_{NO_3} to transient NO spikes—critical drivers of non-steady-state NO₃ loss. For example, on 24 February (POP), a sudden NO burst (24.8 ppbv, Figure S6) caused total k_{NO_3} to jump from 1.3 s⁻¹ to 26.5 s⁻¹ within 30 minutes, with no corresponding change in RH or S_a . This confirms that NO can override the effects of meteorological factors on k_{NO_3} in winter urban environments.

To dissect the drivers of total k_{NO_3} variation, Table 3 presents the contributions of individual reactivity components (NO reaction, N₂O₅ uptake, VOC oxidation) for both periods:

OGP (Table 3): Reactivity was dominated by two pathways. (1) NO reaction: Contributed 0.81 s⁻¹ (71.1% of total k_{NO_3}), reflecting moderate NO emissions under Olympic controls. (2) N₂O₅ uptake: Contributed 0.32 s⁻¹

(28.1% of total k_{NO_3}), with the 13 February snowfall event pushing this contribution to 86% (2.35 s^{-1}). VOC oxidation remained negligible at $0.8 \times 10^{-3} \text{ s}^{-1}$ ($< 0.1\%$ of total k_{NO_3}), consistent with winter's low VOC emissions and low NO_3 -VOC reaction rates (Section 4.2.1). For instance, even the styrene contributed only ~44% of total VOC reactivity, which remained orders of magnitude lower than NO-driven reactivity.

POP (Table 3): A dramatic shift in reactivity partitioning occurred. (1) NO reaction: Exploded to 3.00 s^{-1} (98.0% of total k_{NO_3}), a $3.7\times$ increase from the OGP. This was driven by post-Olympic NO emissions, which intensified NO_3 titration and crowded out other loss pathways. (2) N_2O_5 uptake: Plummeted to 0.06 s^{-1} (2.0% of total k_{NO_3}), an 81% decrease from the OGP. The decline stemmed from lower POP RH (19% vs. 27% in OGP), which reduced aerosol liquid water content and suppressed N_2O_5 hydrolysis (Reaction R6). This RH-dependent hydrolysis inhibition is further validated by Figure S7 (auxiliary material); the figure presents the correlation between N_2O_5 hydrolysis rate and RH during the POP, showing a strong positive linear relationship ($R^2 = 0.81$). The reduced hydrolysis efficiency directly limited N_2O_5 's role as an indirect NO_3 sink, contributing to its diminished share of NO_3 loss in the POP. VOC oxidation increased slightly to $1.4 \times 10^{-3} \text{ s}^{-1}$ ($< 0.1\%$ of total k_{NO_3}), reflecting a modest rebound in anthropogenic VOC emissions (Table 2: total VOCs = 19.68 ppbv vs. 16.02 ppbv in OGP) but remained functionally irrelevant to NO_3 loss.

Table 3 Statistics of k_{NO_3} across various pathways and time periods

$k_{\text{NO}_3} (\text{s}^{-1})$	OGP	POP
$\text{NO}_3 + \text{NO}$	0.81	3.00
$\text{NO}_3 + \text{VOCs}$	0.8×10^{-3}	1.4×10^{-3}
N_2O_5 uptake	0.32	0.06
Total	1.14	3.06

The combined analysis of Figure 5 (total trend) and Table 3 reveals two critical findings:

Emission controls rewire reactivity partitioning: Olympic NO_x reductions reduced NO's dominance as a NO_3 sink, allowing N_2O_5 uptake to emerge as a significant secondary pathway. Relaxing controls reversed this shift, reestablishing NO as the near-exclusive NO_3 loss mechanism.

Extreme events disrupt baseline reactivity: High-RH events (e.g., snowfall) and NO spikes can temporarily alter reactivity partitioning, but their effects are transient—baseline dynamics remain governed by long-term emission conditions.

4.2.3 NO_3 loss budget

Figure 6 illustrates the diurnal variation and relative contributions of NO_3 loss pathways, with distinct differences between the OGP and POP that reflect the impact of Olympic emission controls.

OGP: Balanced NO and N₂O₅ uptake pathways. NO dominated NO₃ loss (79.0%), but N₂O₅ uptake emerged as a significant secondary pathway (20.8%), with VOC oxidation contributing < 0.2%. The N₂O₅ uptake pathway peaked at 19:00 LST (0.37 ppbv h⁻¹), coinciding with high [N₂O₅] (149.9 pptv). This contribution is comparable to winter observations in urban Beijing (Li et al., 2022).

POP: NO dominates NO₃ loss. NO's contribution to NO₃ loss rose to 89.4%, with a peak loss rate of 2.04 ppbv h⁻¹ at 22:00 LST—driven by post-Olympic NO emissions that increased significantly (Table 2). In contrast, N₂O₅ uptake declined to 10.6%, as lower RH reduced $\gamma(\text{N}_2\text{O}_5)$ and S_a reactivity. Notably, the POP's NO₃ loss rate (1.61 ppbv h⁻¹) was 30% higher than the OGP's (1.14 ppbv h⁻¹), confirming that relaxed emissions accelerated nocturnal NO₃ removal.

Compared to summer studies (Lin et al., 2022), winter's low VOC reactivity and high NO emissions make NO the unambiguous primary NO₃ sink in urban Beijing. This seasonal contrast underscores the need for season-specific air quality management strategies—prioritizing NO_x reduction in winter and VOC reduction in summer.




Figure 6. Mean daily variation and reactivity share of different loss pathways.

4.3 Linkage to Olympic Emission Controls and Precursor Dynamics

The 2022 Beijing Winter Olympics provided a unique “natural experiment” to quantify how short-term, large-scale emission controls modulate nocturnal NO₃-N₂O₅ chemistry. Below, we dissect the impacts of these controls on precursor concentrations and loss pathway hierarchies, and explore the interacting roles of NO₂ and O₃ in shaping NO₃-N₂O₅ dynamics.

4.3.1 Impact of Emission Controls on NO₃-N₂O₅ Chemistry

The Olympic emission control measures—including traffic restrictions (e.g., odd-even license plate policy), industrial shutdowns, and reduced coal combustion—induced significant changes in precursor concentrations

and NO₃-N₂O₅ behavior.

Precursor modulation: OGP emissions of NO_x and VOCs were substantially lower than the POP. Nocturnal NO decreased by 79% (1.0 vs. 4.8 ppbv), and NO₂ by 30% (14.5 vs. 20.7 ppbv). Nocturnal total VOCs decreased by 18% (16.02 vs. 19.68 ppbv), with reactive AVOCs (styrene) declining by 40%.

Lower NO emissions weakened its role as a “NO₃ scavenger,” allowing more NO₃ to partition into N₂O₅ via Reaction R2. This explains why the OGP’s nocturnal [N₂O₅] (137.6 pptv) was 41% higher than the POP’s (97.8 pptv), despite similar NO₃ production rates ($P(\text{NO}_3) = 0.5 \text{ ppbv h}^{-1}$ for both periods).

Loss pathway shift: Reduced NO emissions elevated the relative importance of N₂O₅ uptake in NO₃ loss—its contribution increased from 10.6% (POP) to 20.8% (OGP). This shift demonstrates that emission controls can “rewire” nocturnal loss hierarchies, making heterogeneous processes more significant as primary pollutant emissions decline. This finding extends previous studies (Xia et al., 2020; Zhou et al., 2018a) which focused on ClNO₂ formation but did not quantify N₂O₅’s role under low-NO conditions.

4.3.2 Precursor (NO₂, O₃) Influences on NO₃-N₂O₅ Dynamics

NO₂ and O₃, the primary precursors of NO₃ via Reaction R1, exert dual and interactive controls on NO₃-N₂O₅ chemistry that depend on emission conditions.

NO₂: Equilibrium driver and precursor tradeoff. NO₂ plays two conflicting roles: it is both a precursor for NO₃ (via R1) and a partner in the NO₃-N₂O₅ equilibrium (via R2). During the OGP, higher nocturnal NO₂ (14.5 ppbv) shifted the equilibrium toward N₂O₅, as low NO prevented rapid NO₃ loss. This created a “N₂O₅ surplus,” with [N₂O₅] exceeding the POP’s by 41%. In contrast, the POP’s higher NO₂ (20.7 ppbv) paired with elevated NO (4.8 ppbv) created a “dual sink” effect: more NO₃ was produced via R1, but immediately titrated by NO via R4—limiting N₂O₅ accumulation. This tradeoff highlights NO₂’s context-dependent role in NO₃-N₂O₅ dynamics.

O₃: Production regulator and persistence enhancer. OGP’s higher nocturnal O₃ (27.4 vs. 19.8 ppbv in the POP) initially boosted NO₃ production ($P(\text{NO}_3) = 0.5 \text{ ppbv h}^{-1}$), but lower NO prevented rapid NO₃ loss. This “NO₃ surplus” prolonged N₂O₅ lifetime: the OGP’s nocturnal $\tau_{\text{N}_2\text{O}_5}$ (17.0 minutes) was 46% longer than the POP’s (11.6 minutes). This indirect effect of O₃—enhancing N₂O₅ persistence by supporting NO₃ production without accelerating loss—has not been fully quantified in previous winter studies, emphasizing the need to consider O₃ alongside NO_x in nocturnal chemistry models.

4.4 Broader Atmospheric Implications

Our findings advance understanding of winter urban nocturnal chemistry and provide critical insights for air

quality management in polluted regions like Beijing. Three key implications emerge:

Emission control efficacy and regional nitrate transport. Olympic controls demonstrated that reducing NO_x (not just VOCs) can enhance N_2O_5 accumulation—potentially extending the lifetime of reactive nitrogen and increasing regional nitrate transport. N_2O_5 is a stable reservoir species that can be transported long distances before hydrolyzing to form nitrate aerosols; thus, increased N_2O_5 under NO_x reduction may shift winter nitrate pollution from local to regional scales. This suggests that future NO_x mitigation strategies should consider regional coordination to address long-range transport of reactive nitrogen.

Model parameterization improvements. Our results provide two critical constraints for air quality models, which often underestimate winter N_2O_5 chemistry.

S_a threshold: S_a exerts significant control over $\tau_{\text{N}_2\text{O}_5}$ only when $S_a > 500 \mu\text{m}^2 \text{cm}^{-3}$; below this threshold, organic coatings and NO dominate.

$\gamma(\text{N}_2\text{O}_5)$ range: The average $\gamma(\text{N}_2\text{O}_5)$ of 0.032 ± 0.049 (with values up to 0.22 under high RH) is higher than the constant $\gamma(\text{N}_2\text{O}_5) = 0.02$ (Chang et al., 2011) often used in models. Incorporating these season-specific parameters will improve predictions of winter nitrate formation.

Winter mitigation priority: NO_x over VOCs. Despite their high reactivity, VOCs contribute $< 0.5\%$ of NO_3 loss in winter urban Beijing—far less than NO (79.0–89.4%). This confirms that NO_x (not VOCs) should be prioritized for mitigating winter nocturnal nitrogen pollution. For example, further reducing traffic-related NO emissions (a major source in urban Beijing) would not only lower direct NO pollution but also enhance N_2O_5 uptake—a pathway that converts reactive nitrogen to nitrate, which is less toxic and more easily removed via wet deposition.

5 Summary and conclusions

This study conducted continuous field observations of N_2O_5 , NO_3 , and their precursor species (NO, NO_2 , O_3 , VOCs) in urban Beijing from 5 February to 3 March 2022, covering the 2022 Beijing Winter Olympics (BWO). By analyzing pollutant variations, quantifying the contributions of $\text{NO}_3/\text{N}_2\text{O}_5$ loss pathways, and linking observations to BWO emission control measures, we clarified the response of winter nocturnal reactive nitrogen chemistry to short-term anthropogenic emission reductions.

During the observation period, $P(\text{NO}_3)$ averaged $0.5 \pm 0.4 \text{ ppbv h}^{-1}$, with N_2O_5 mixing ratios peaking at 875 pptv (1-minute resolution) and derived NO_3 concentrations reaching a maximum of 4.6 pptv; $\tau_{\text{N}_2\text{O}_5}$ averaged 11.9 ± 11.8 minutes, longer than summer values in Beijing due to slower winter N_2O_5 loss driven by low temperatures and reduced heterogeneous reactivity. BWO emission controls significantly modulated precursor concentrations: nocturnal NO ($1.0 \pm 1.2 \text{ ppbv}$) and total VOCs ($16.02 \pm 7.74 \text{ ppbv}$) in the OGP were 79% and

18% lower than in the POP, respectively, while nocturnal O_3 was 38% higher in the OGP (27.4 ± 10.3 ppbv vs. 19.8 ± 12.1 ppbv in the POP) as reduced NO minimized O_3 titration—these changes directly led to 41% higher nocturnal N_2O_5 concentrations in the OGP (137.6 ± 112.7 pptv vs. 97.8 ± 90.3 pptv in the POP).

RH and S_a exerted context-dependent control over $\tau_{N_2O_5}$: at RH < 35%, $\tau_{N_2O_5}$ increased with RH as slight humidity rises softened hydrophobic organic aerosol coatings (derived from traffic VOC oxidation) and reduced N_2O_5 heterogeneous uptake; at RH > 35%, $\tau_{N_2O_5}$ decreased with RH due to hygroscopic aerosol growth and enhanced N_2O_5 hydrolysis, approaching zero during snowfall events (RH > 85%). For S_a , a threshold of $\sim 500 \mu m^2 cm^{-3}$ was identified—below this value, organic coatings and NO dominated $\tau_{N_2O_5}$; above it, S_a became the primary regulator, with $\tau_{N_2O_5}$ decreasing as S_a increased.

NO was the dominant NO_3 sink in both periods, though its contribution varied with emission controls: it accounted for 79.0% of NO_3 loss in the OGP, with N_2O_5 heterogeneous uptake (20.8%) as a significant secondary pathway, while its contribution rose to 89.2% in the POP (driven by $3.8\times$ higher NO emissions) and N_2O_5 uptake declined to 10.6% (due to lower RH reducing aerosol reactivity). The N_2O_5 heterogeneous uptake coefficient ($\gamma(N_2O_5)$) averaged 0.032 ± 0.049 in the OGP, higher than rural sites due to urban aerosols' higher water content and reactive components (e.g., nitrate, sulfate). Despite the high reactivity of species like styrene and isoprene, VOC oxidation contributed < 0.2% to NO_3 loss in both periods, confirming its negligible role in winter NO_3 dynamics in urban Beijing.

These findings hold key implications for air quality management: BWO NO_x reductions enhanced N_2O_5 accumulation, potentially extending reactive nitrogen lifetime and shifting winter nitrate pollution from local to regional scales—highlighting the need for regional coordination in NO_x mitigation; the identified S_a threshold ($500 \mu m^2 cm^{-3}$) and $\gamma(N_2O_5)$ range (0.01–0.12) provide critical constraints for air quality models, which often rely on oversimplified $\tau_{N_2O_5}$ and $\gamma(N_2O_5)$ parameters; and given NO's dominance in NO_3 loss and N_2O_5 dynamics, NO_x (not VOCs) should be prioritized for winter nocturnal nitrogen pollution control in Beijing—reducing traffic-related NO emissions would simultaneously lower direct pollution and enhance N_2O_5 uptake, promoting nitrate removal via wet deposition.

Data availability. Data are available at <https://doi.org/10.5281/zenodo.15381990> (T. Zhang et al., 2025).

Supplement. The following file is available free of charge. Supplement of “Measurement report: Variations and environmental impacts of atmospheric N_2O_5 concentrations in urban Beijing during the 2022 Winter Olympics”

Author Contributions. TZ, WL, and CY designed the research. WL, and CY organized this field campaign. TZ, PZ, YC, TL and LZ carried out the field measurements and provided the field measurement dataset. TZ performed data analysis, interpreted the data and wrote the manuscript with revision mainly from WL. All

authors have given approval to the final version of the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgment. The authors would like to thank the field campaign team for the data that they contributed during the 2022 Beijing Winter Olympics.

Financial support. This study was financially supported by the National Natural Science Foundation of China (42405083), the National Key Research and Development Program (2024YFC3711902).

References

- Anttila, T., Kiendler-Scharr, A., Tillmann, R., and Mentel, T. F.: On the Reactive Uptake of Gaseous Compounds by Organic-Coated Aqueous Aerosols: Theoretical Analysis and Application to the Heterogeneous Hydrolysis of N_2O_5 , *J. Phys. Chem. A*, 110, 10435–10443, <https://doi.org/10.1021/jp062403c>, 2006.
- Atkinson, R. and Arey, J.: Atmospheric Degradation of Volatile Organic Compounds, *Chem. Rev.*, 103, 4605–4638, <https://doi.org/10.1021/cr0206420>, 2003.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O_x , HO_2 , NO_x and SO_x species, *Atmos. Chem. Phys.*, 4, 1461–1738, <https://doi.org/10.5194/acp-4-1461-2004>, 2004.
- Bertram, T. H., Thornton, J. A., Riedel, T. P., Middlebrook, A. M., Bahreini, R., Bates, T. S., Quinn, P. K., and Coffman, D. J.: Direct observations of N_2O_5 reactivity on ambient aerosol particles, *Geophys. Res. Lett.*, 36, L19803, <https://doi.org/10.1029/2009GL040248>, 2009.
- Bohn, B., Corlett, G. K., Gillmann, M., Sanghavi, S., Stange, G., Tensing, E., Vrekoussis, M., Bloss, W. J., Clapp, L. J., Kortner, M., Dorn, H.-P., Monks, P. S., Platt, U., Plass-Dülmer, C., Mihalopoulos, N., Heard, D. E., Clemmshaw, K. C., Meixner, F. X., Prevot, A. S. H., and Schmitt, R.: Photolysis frequency measurement techniques: results of a comparison within the ACCENT project, *Atmos. Chem. Phys.*, 8, 5373–5391, <https://doi.org/10.5194/acp-8-5373-2008>, 2008.
- Brown, S. S. and Stutz, J.: Nighttime radical observations and chemistry, *Chem. Soc. Rev.*, 41, 6405, <https://doi.org/10.1039/c2cs35181a>, 2012.
- Brown, S. S., Dubé, W. P., Peischl, J., Ryerson, T. B., Atlas, E., Warneke, C., de Gouw, J. A., de Lintel Hekkert, S., Brock, C. A., Flocke, F., Trainer, M., Parrish, D. D., Feshenfeld, F. C., and Ravishankara, A. R.: Budgets for nocturnal VOC oxidation by nitrate radicals aloft during the 2006 Texas Air Quality Study, *J. Geophys. Res.*, 116, 1–15, <https://doi.org/10.1029/2011JD016544>, 2011.

- Brown, S. S., Dubé, W. P., Tham, Y. J., Zha, Q., Xue, L., Poon, S., Wang, Z., Blake, D. R., Tsui, W., Parrish, D. D., and Wang, T.: Nighttime chemistry at a high altitude site above Hong Kong, *J. Geophys. Res. Atmos.*, 121, 2457–2475, <https://doi.org/10.1002/2015JD024566>, 2016.
- Brown, S. S., An, H., Lee, M., Park, J.-H., Lee, S.-D., Fibiger, D. L., McDuffie, E. E., Dubé, W. P., Wagner, N. L., and Min, K.-E.: Cavity enhanced spectroscopy for measurement of nitrogen oxides in the Anthropocene: results from the Seoul tower during MAPS 2015, *Faraday Discuss.*, 200, 529–557, <https://doi.org/10.1039/C7FD00001D>, 2017.
- Chang, W. L., Bhawe, P. V., Brown, S. S., Riemer, N., Stutz, J., and Dabdub, D.: Heterogeneous Atmospheric Chemistry, Ambient Measurements, and Model Calculations of N_2O_5 : A Review, *Aerosol Sci. Technol.*, 45, 665–695, <https://doi.org/10.1080/02786826.2010.551672>, 2011.
- Cheng, X., Li, H., Zhang, Y., and Li, Y.: Atmospheric isoprene and monoterpenes in a typical urban area of Beijing: Pollution characterization, chemical reactivity and source identification, *J. Environ. Sci.*, 71, 150–167, <https://doi.org/10.1016/j.jes.2017.12.017>, 2018.
- Crutzen, P. J.: The Role of NO and NO_2 in the Chemistry of the Troposphere and Stratosphere, *Annu. Rev. Earth Planet. Sci.*, 7, 443–472, <https://doi.org/10.1146/annurev.ea.07.050179.002303>, 1979.
- Folkers, M., Mentel, Th. F., and Wahner, A.: Influence of an organic coating on the reactivity of aqueous aerosols probed by the heterogeneous hydrolysis of N_2O_5 , *Geophys. Res. Lett.*, 30, 2003GL017168, <https://doi.org/10.1029/2003GL017168>, 2003.
- Fry, J. L. and Sackinger, K.: Model investigation of NO_3 secondary organic aerosol (SOA) source and heterogeneous organic aerosol (OA) sink in the western United States, *Atmos. Chem. Phys.*, 12, 8797–8811, <https://doi.org/10.5194/acp-12-8797-2012>, 2012.
- Hoyle, C. R., Berntsen, T., Myhre, G., and Isaksen, I. S. A.: Secondary organic aerosol in the global aerosol - chemical transport model Oslo CTM2, *Atmos. Chem. Phys.*, 7, 5675–5694, <https://doi.org/10.5194/acp-7-5675-2007>, 2007.
- Hu, H., Wang, H., Lu, K., Wang, J., Zheng, Z., Xu, X., Zhai, T., Chen, X., Lu, X., Fu, W., Li, X., Zeng, L., Hu, M., Zhang, Y., and Fan, S.: Variation and trend of nitrate radical reactivity towards volatile organic compounds in Beijing, China, *Atmos. Chem. Phys.*, 23, 8211–8223, <https://doi.org/10.5194/acp-23-8211-2023>, 2023.
- Huang Z., Hu W., Jin R., Hou S., LI P., Bi K., He C., Wang Y., Duan P., Liu D., Wu L., Deng J., Sun Y., and Fu P.: Chemical composition and sources of fine particles in Beijing around the Winter Olympics, China *Environ. Sci.*, 44, 5344–5356, <https://doi.org/10.19674/j.cnki.issn1000-6923.20240604.002>, 2024.
- Li, H., Zheng, B., Lei, Y., Hauglustaine, D., Chen, C., Lin, X., Zhang, Y., Zhang, Q., and He, K.: Trends and drivers of anthropogenic NO_x emissions in China since 2020, *Environ. Sci. Ecotechnol.*, 21, 100425, <https://doi.org/10.1016/j.ese.2024.100425>, 2024.
- Li, L., Li, H., and Zhang, X.: Pollution characteristics and health risk assessment of benzene homologues in ambient air in the northeastern urban area of Beijing, China, *J. Environ. Sci.*, 26, 214–223, [https://doi.org/10.1016/S1001-0742\(13\)60400-3](https://doi.org/10.1016/S1001-0742(13)60400-3), 2014.

- Li, Z., Xie, P., Hu, R., Wang, D., Jin, H., Chen, H., Lin, C., and Liu, W.: Observations of N_2O_5 and NO_3 at a suburban environment in Yangtze river delta in China: Estimating heterogeneous N_2O_5 uptake coefficients, *J. Environ. Sci.*, 95, 248–255, <https://doi.org/10.1016/j.jes.2020.04.041>, 2020.
- Li, Z., Wang, D., Xie, P., Hu, R., Chen, H., and Lin, C.: Nighttime N_2O_5 chemistry in an urban site of Beijing in winter based on the measurements by cavity ring-down spectroscopy, *Air Qual. Atmos. Hlth.*, 15, 867–876, <https://doi.org/10.1007/s11869-021-01125-4>, 2022.
- Lin, C., Hu, R., Xie, P., Lou, S., Zhang, G., Tong, J., Liu, J., and Liu, W.: Nocturnal atmospheric chemistry of NO_3 and N_2O_5 over Changzhou in the Yangtze River Delta in China, *J. Environ. Sci.*, 114, 376–390, <https://doi.org/10.1016/j.jes.2021.09.016>, 2022.
- Lu, X., Qin, M., Xie, P., Duan, J., Fang, W., and Liu, W.: Observation of ambient NO_3 radicals by LP-DOAS at a rural site in North China Plain, *Sci. Total Environ.*, 804, 149680, <https://doi.org/10.1016/j.scitotenv.2021.149680>, 2022.
- McNeill, V. F., Patterson, J., Wolfe, G. M., and Thornton, J. A.: The effect of varying levels of surfactant on the reactive uptake of N_2O_5 to aqueous aerosol, *Atmos. Chem. Phys.*, 6, 1635–1644, <https://doi.org/10.5194/acp-6-1635-2006>, 2006.
- Ng, N. L., Brown, S. S., Archibald, A. T., Atlas, E., Cohen, R. C., Crowley, J. N., Day, D. A., Donahue, N. M., Fry, J. L., Fuchs, H., Griffin, R. J., Guzman, M. I., Herrmann, H., Hodzic, A., Iinuma, Y., Jimenez, J. L., Kiendler-Scharr, A., Lee, B. H., Luecken, D. J., Mao, J., McLaren, R., Mutzel, A., Osthoff, H. D., Ouyang, B., Picquet-Varrault, B., Platt, U., Pye, H. O. T., Rudich, Y., Schwantes, R. H., Shiraiwa, M., Stutz, J., Thornton, J. A., Tilgner, A., Williams, B. J., and Zaveri, R. A.: Nitrate radicals and biogenic volatile organic compounds: oxidation, mechanisms, and organic aerosol, *Atmos. Chem. Phys.*, 17, 2103–2162, <https://doi.org/10.5194/acp-17-2103-2017>, 2017.
- Osthoff, H. D., Sommariva, R., Baynard, T., Pettersson, A., Williams, E. J., Lerner, B. M., Roberts, J. M., Stark, H., Goldan, P. D., Kuster, W. C., Bates, T. S., Coffman, D., Ravishankara, A. R., and Brown, S. S.: Observation of daytime N_2O_5 in the marine boundary layer during New England Air Quality Study–Intercontinental Transport and Chemical Transformation 2004, *J. Geophys. Res. Atmospheres*, 111, <https://doi.org/10.1029/2006JD007593>, 2006.
- Phillips, G. J., Thieser, J., Tang, M. J., Sobanski, N., Schuster, G., Fachinger, J., Drewnick, F., Borrmann, S., Bingemer, H., Lelieveld, J., and Crowley, J. N.: Estimating N_2O_5 uptake coefficients using ambient measurements of NO_3 , N_2O_5 , ClNO_2 and particle-phase nitrate, *Atmos. Chem. Phys.*, 1–9, <https://doi.org/10.5194/acp-2016-693>, 2016.
- Pye, H. O. T., Chan, A. W. H., Barkley, M. P., and Seinfeld, J. H.: Global modeling of organic aerosol: the importance of reactive nitrogen (NO_x and NO_3), *Atmos. Chem. Phys.*, 10, 11261–11276, <https://doi.org/10.5194/acp-10-11261-2010>, 2010.
- Tang, M. J., Schuster, G., and Crowley, J. N.: Heterogeneous reaction of N_2O_5 with illite and Arizona test dust particles, *Atmos. Chem. Phys.*, 14, 245–254, <https://doi.org/10.5194/acp-14-245-2014>, 2014.
- Tham, Y. J., Wang, Z., Li, Q., Yun, H., Wang, W., Wang, X., Xue, L., Lu, K., Ma, N., Bohn, B., Li, X., Kecorius,

- S., Größ, J., Shao, M., Wiedensohler, A., Zhang, Y., and Wang, T.: Significant concentrations of nitryl chloride sustained in the morning: investigations of the causes and impacts on ozone production in a polluted region of northern China, *Atmos. Chem. Phys.*, 16, 14959–14977, <https://doi.org/10.5194/acp-16-14959-2016>, 2016.
- Tham, Y. J., Wang, Z., Li, Q., Wang, W., Wang, X., Lu, K., Ma, N., Yan, C., Kecorius, S., Wiedensohler, A., Zhang, Y., and Wang, T.: Heterogeneous N_2O_5 uptake coefficient and production yield of ClNO_2 in polluted northern China: roles of aerosol water content and chemical composition, *Atmos. Chem. Phys.*, 18, 13155–13171, <https://doi.org/10.5194/acp-18-13155-2018>, 2018.
- Wang, D., Xie, P., Hu, R., Li, Z., Chen, H., and Jin, H.: Reactivity and Loss Mechanisms of NO_3 and N_2O_5 at a Rural Site on the North China Plain, *Atmosphere*, 13, 1268, <https://doi.org/10.3390/atmos13081268>, 2022.
- Wang, H., Lu, K., Chen, X., Zhu, Q., Chen, Q., Guo, S., Jiang, M., Li, X., Shang, D., Tan, Z., Wu, Y., Wu, Z., Zou, Q., Zheng, Y., Zeng, L., Zhu, T., Hu, M., and Zhang, Y.: High N_2O_5 Concentrations Observed in Urban Beijing: Implications of a Large Nitrate Formation Pathway, *Environ. Sci. Technol. Lett.*, 4, 416–420, <https://doi.org/10.1021/acs.estlett.7b00341>, 2017a.
- Wang, H., Lu, K., Guo, S., Wu, Z., Shang, D., Tan, Z., Wang, Y., Le Breton, M., Lou, S., Tang, M., Wu, Y., Zhu, W., Zheng, J., Zeng, L., Hallquist, M., Hu, M., and Zhang, Y.: Efficient N_2O_5 uptake and NO_3 oxidation in the outflow of urban Beijing, *Atmos. Chem. Phys.*, 18, 9705–9721, <https://doi.org/10.5194/acp-18-9705-2018>, 2018.
- Wang, H., Chen, X., Lu, K., Hu, R., Li, Z., Wang, H., Ma, X., Yang, X., Chen, S., Dong, H., Liu, Y., Fang, X., Zeng, L., Hu, M., and Zhang, Y.: NO_3 and N_2O_5 chemistry at a suburban site during the EXPLORE-YRD campaign in 2018, *Atmos. Environ.*, 224, 117180, <https://doi.org/10.1016/j.atmosenv.2019.117180>, 2020.
- Wang, H., Lu, K., Chen, S., Li, X., Zeng, L., Hu, M., and Zhang, Y.: Characterizing nitrate radical budget trends in Beijing during 2013–2019, *Sci. Total Environ.*, 795, 148869, <https://doi.org/10.1016/j.scitotenv.2021.148869>, 2021.
- Wang, H., Wang, H., Lu, X., Lu, K., Zhang, L., Tham, Y. J., Shi, Z., Aikin, K., Fan, S., Brown, S. S., and Zhang, Y.: Increased night-time oxidation over China despite widespread decrease across the globe, *Nat. Geosci.*, 16, 217–223, <https://doi.org/10.1038/s41561-022-01122-x>, 2023a.
- Wang, J., Wang, H., Tham, Y. J., Ming, L., Zheng, Z., Fang, G., Sun, C., Ling, Z., Zhao, J., and Fan, S.: Measurement report: Atmospheric nitrate radical chemistry in the South China Sea influenced by the urban outflow of the Pearl River Delta, *Atmos. Chem. Phys.*, 24, 977–992, <https://doi.org/10.5194/acp-24-977-2024>, 2024.
- Wang, M., Zeng, L., Lu, S., Shao, M., Liu, X., Yu, X., Chen, W., Yuan, B., Zhang, Q., Hu, M., and Zhang, Z.: Development and validation of a cryogen-free automatic gas chromatograph system (GC-MS/FID) for online measurements of volatile organic compounds, *Anal. Methods*, 6, 9424–9434, <https://doi.org/10.1039/C4AY01855A>, 2014.
- Wang, S., Shi, C., Zhou, B., Zhao, H., Wang, Z., Yang, S., and Chen, L.: Observation of NO_3 radicals over Shanghai, China, *Atmos. Environ.*, 70, 401–409, <https://doi.org/10.1016/j.atmosenv.2013.01.022>, 2013.

- Wang, X., Wang, H., Xue, L., Wang, T., Wang, L., Gu, R., Wang, W., Tham, Y. J., Wang, Z., Yang, L., Chen, J., and Wang, W.: Observations of N_2O_5 and ClNO_2 at a polluted urban surface site in North China: High N_2O_5 uptake coefficients and low ClNO_2 product yields, *Atmos. Environ.*, 156, 125–134, <https://doi.org/10.1016/j.atmosenv.2017.02.035>, 2017b.
- Wang, Y., Xi, S., Zhao, F., Huey, L. G., and Zhu, T.: Decreasing Production and Potential Urban Explosion of Nighttime Nitrate Radicals amid Emission Reduction Efforts, *Environ. Sci. Technol.*, 57, 21306–21312, <https://doi.org/10.1021/acs.est.3c09259>, 2023b.
- Wang, Z., Wang, W., Tham, Y. J., Li, Q., Wang, H., Wen, L., Wang, X., and Wang, T.: Fast heterogeneous N_2O_5 uptake and ClNO_2 production in power plant and industrial plumes observed in the nocturnal residual layer over the North China Plain, *Atmos. Chem. Phys.*, 17, 12361–12378, <https://doi.org/10.5194/acp-17-12361-2017>, 2017c.
- Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosa-Mas, C. E., Hjorth, J., Le Bras, G., Moortgat, G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The nitrate radical: Physics, chemistry, and the atmosphere, *Atmos. Environ. Part Gen. Top.*, 25, 1–203, [https://doi.org/10.1016/0960-1686\(91\)90192-A](https://doi.org/10.1016/0960-1686(91)90192-A), 1991.
- Xia, M., Peng, X., Wang, W., Yu, C., Sun, P., Li, Y., Liu, Y., Xu, Z., Wang, Z., Xu, Z., Nie, W., Ding, A., and Wang, T.: Significant production of ClNO_2 and possible source of Cl_2 from N_2O_5 uptake at a suburban site in eastern China, *Atmos. Chem. Phys.*, 20, 6147–6158, <https://doi.org/10.5194/acp-20-6147-2020>, 2020.
- Xia, M., Peng, X., Wang, W., Yu, C., Wang, Z., Tham, Y. J., Chen, J., Chen, H., Mu, Y., Zhang, C., Liu, P., Xue, L., Wang, X., Gao, J., Li, H., and Wang, T.: Winter ClNO_2 formation in the region of fresh anthropogenic emissions: seasonal variability and insights into daytime peaks in northern China, *Atmos. Chem. Phys.*, 21, 15985–16000, <https://doi.org/10.5194/acp-21-15985-2021>, 2021.
- Yan, C., Yee Jun, T., Nie, W., Xia, M., Wang, H., Guo, Y., Ma, W., Zhan, J., Li, Y., Deng, C., Li, Y., Zheng, F., Chen, X., Li, Q., Zhang, G., Mahajan, A., Cuevas, C. A., Huang, D., and Kulmala, M.: Increasing contribution of nighttime nitrogen chemistry to wintertime haze formation in Beijing observed during COVID-19 lockdowns, *Nat. Geosci.*, 16, <https://doi.org/10.1038/s41561-023-01285-1>, 2023.
- Yao Y., Wang S., Wei N., Ye C., Zhang C., Gu X., Zhao W., and Zhang W.: Analysis of surface ozone sources in Beijing during the 2022 Beijing Winter Olympic Games based on total peroxy radical measurement, *Acta Sci. Circumst.*, 43, 290–297, <https://doi.org/10.13671/j.hjkxxb.2023.0070>, 2023.
- Yu, C., Wang, Z., Xia, M., Fu, X., Wang, W., Tham, Y. J., Chen, T., Zheng, P., Li, H., Shan, Y., Wang, X., Xue, L., Zhou, Y., Yue, D., Ou, Y., Gao, J., Lu, K., Brown, S. S., Zhang, Y., and Wang, T.: Heterogeneous N_2O_5 reactions on atmospheric aerosols at four Chinese sites: improving model representation of uptake parameters, *Atmos. Chem. Phys.*, 20, 4367–4378, <https://doi.org/10.5194/acp-20-4367-2020>, 2020.
- Yuan, Z., Lau, A. K. H., Shao, M., Louie, P. K. K., Liu, S. C., and Zhu, T.: Source analysis of volatile organic compounds by positive matrix factorization in urban and rural environments in Beijing, *J. Geophys. Res. Atmospheres*, 114, <https://doi.org/10.1029/2008JD011190>, 2009.
- Yun, H., Wang, W., Wang, T., Xia, M., Yu, C., Wang, Z., Poon, S. C. N., Yue, D., and Zhou, Y.: Nitrate formation

from heterogeneous uptake of dinitrogen pentoxide during a severe winter haze in southern China, *Atmos. Chem. Phys.*, 18, 17515–17527, <https://doi.org/10.5194/acp-18-17515-2018>, 2018.

Zhang, H., Wang, S., and Hao, J.: Air pollution and control action in Beijing, *J. Clean. Prod.*, 112, 1519–1527, <https://doi.org/10.1016/j.jclepro.2015.04.092>, 2016.

Zhang T., Zuo P., Ma J., Ye C., Lin W., and Zhu T.: Characterization and Application of an Online Measurement System for NO₃ and N₂O₅ Based on Cavity Ring-Down Spectroscopy, *Acta Sci. Nat. Univ. Pekin.*, 60, 563–574, <https://doi.org/10.13209/j.0479-8023.2024.030>, 2024.

Zhang, T., Ma, J., Liu, T., Lin, W., Zuo, P., and Ye, C.: A dynamic generation system for NO₃ and N₂O₅ standard gases, *Environ. Chem.*, 45, 1–7, <https://doi.org/10.7524/j.issn.0254-6108.2024091302>, 2026.

Zhou, J., Zhao, W., Zhang, Y., Fang, B., Cheng, F., Xu, X., Ni, S., Zhang, W., Ye, C., Chen, W., and Venables, D. S.: Amplitude-Modulated Cavity-Enhanced Absorption Spectroscopy with Phase-Sensitive Detection: A New Approach Applied to the Fast and Sensitive Detection of NO₂, *Anal. Chem.*, 94, 3368–3375, <https://doi.org/10.1021/acs.analchem.1c05484>, 2022.

Zhou, W., Zhao, J., Ouyang, B., Mehra, A., Xu, W., Wang, Y., Bannan, T. J., Worrall, S. D., Priestley, M., Bacak, A., Chen, Q., Xie, C., Wang, Q., Wang, J., Du, W., Zhang, Y., Ge, X., Ye, P., Lee, J. D., Fu, P., Wang, Z., Worsnop, D., Jones, R., Percival, C. J., Coe, H., and Sun, Y.: Production of N₂O₅ and ClNO₂ in summer in urban Beijing, China, *Atmos. Chem. Phys.*, 18, 11581–11597, <https://doi.org/10.5194/acp-18-11581-2018>, 2018.

Zhu, T., Tang, M., Gao, M., Bi, X., Cao, J., Che, H., Chen, J., Ding, A., Fu, P., Gao, J., Gao, Y., Ge, M., Ge, X., Han, Z., He, H., Huang, R.-J., Huang, X., Liao, H., Liu, C., Liu, H., Liu, J., Liu, S. C., Lu, K., Ma, Q., Nie, W., Shao, M., Song, Y., Sun, Y., Tang, X., Wang, T., Wang, T., Wang, W., Wang, X., Wang, Z., Yin, Y., Zhang, Q., Zhang, W., Zhang, Y., Zhang, Y., Zhao, Y., Zheng, M., Zhu, B., and Zhu, J.: Recent Progress in Atmospheric Chemistry Research in China: Establishing a Theoretical Framework for the “Air Pollution Complex”※, *Adv. Atmos. Sci.*, 1339–1361, 2023.