

Bacterial community composition changes independently of soil edaphic parameters following localized permafrost disturbance

Patrick Neuberger¹, Alireza Saidi-Mehrabad¹, Duane Froese², Brian Lanoil¹

5 ¹Department of Biological Sciences, University of Alberta, Edmonton, T6G 2E9, Canada.

²Department of Earth and Atmospheric Sciences, Edmonton, T6G 2E3, Canada.

Correspondence to: Brian Lanoil (lanoil@ualberta.ca)

Abstract. Microbial degradation of frozen organic carbon increases with permafrost thaw, resulting in greater fluxes of the greenhouse gases CO₂ and CH₄. To examine the effect of disturbance-induced permafrost thaw on microbial communities, we assessed the microbial diversity of soils near a gold mine where thaw was induced by stripping the vegetation and topsoil at Dominion Creek, Yukon, Canada. Bacterial metabarcoding and soil physicochemical parameters were assessed across this disturbance including surface samples and three cores which included active layer and permafrost horizons. Bacterial communities changed in the absence of physicochemical parameter shifts after only 6 weeks of thaw, with a high proportion of active layer indicator species becoming more abundant with permafrost thaw. Three distinct communities emerged: (1) undisturbed active layer, (2) lower active layer, disturbed active layer, and disturbed permafrost samples, and (3) intact permafrost. Community composition shifts correlated with pH, Zn and community cohesion. These results suggest that active layer communities rapidly colonized thawed permafrost at our sample site, combining with and replacing many resident permafrost taxa. Disturbances may induce a strong microbial community change in permafrost-affected soils before soil physicochemical parameter shifts.

1 Introduction

There is roughly 1,700 Gt of carbon in northern circumpolar soils, more than twice as much as currently in the atmosphere (Schuur et al., 2015; van Huissteden & Dolman, 2012). Although increased vegetation and greening of the Arctic may offset carbon output from permafrost thaw, the increased rate of forest fires (Chen et al., 2021) and thermokarst formation as a result of abrupt permafrost thaw (Turetsky et al., 2020) may already have shifted the Arctic from a net carbon sink to a net carbon source as carbon mineralization outpaces fixation (Schuur et al., 2020). Models suggest that thawed permafrost sediments will

release 41 PgC per 1°C of warming as greenhouse gases by the year 2100, representing one of the most significant biospheric sources of atmospheric carbon (IPCC, 2022). Newer models suggest that of this predicted 208 PgC per year, approximately 80 PgC will be sourced from abrupt thaw areas which cover only 5% of global permafrost-affected areas (Turetsky et al., 2020). However, there is significant uncertainty in the timing and mechanisms of these models. A significant source of uncertainty is how the microbial communities will respond to permafrost thaw.

Permafrost thaw acts as a disturbance to the environment, changing the ice and water content, leading to subsidence or mass wasting of overlying layers, and releasing preserved organic material (Kokelj & Jorgenson, 2013). Such disturbances can be the result of the rapidly rising temperatures in the Northern polar regions (i.e., polar amplification of climate warming) or industrial activity such as mining or road construction (Richter-Menge, et al., 2016). The extent and severity of these disturbances can vary from relatively small scale and minor (e.g. subsidence and ponding due to road building) to large scale and extreme (e.g. active layer detachment and mass wasting due to the formation of retrogressive thaw slumps) (Kokelj & Jorgenson, 2013). These disturbances strongly affect bacterial communities in the permafrost, leading to shifts in both the structure and function of bacterial communities residing in permafrost (Wu et al., 2018). However, our knowledge of such shifts is mostly limited to laboratory-based experiments, where there are no other sources of bacteria than the permafrost itself.

Bacterial community activity regulates the rate of carbon mineralization from permafrost, and so the characterization of these communities is paramount to modelling carbon flux from thawing permafrost soils (Monteux et al., 2018; McCalley et al., 2014). Although permafrost microbial communities are not as diverse as in the seasonally thawing

active layer, permafrost microbial diversity is substantial (Hultman et al., 2015). Within the last decade, an abundance of information has been collected regarding the microbial response to *in situ* permafrost thaw (Taş et al., 2014; Taş et al., 2018; Singleton et al., 2018; Woodcroft et al., 2018; Emerson et al., 2018; Hultman et al., 2015). Field experiments have shown contradicting results; either that (1) thaw shifts the permafrost microbial community to become similar to the overlying active layer (Taş et al., 2014; Monteux et al., 2018), or that (2) subsidence creates an altogether new microbial community not previously seen (Mondav et al., 2014; Woodcroft et al., 2018).

Soil microbial activity refers to the growth and metabolic processes undertaken by soil microbial communities (Nazir et al., 2014). Increased microbial activity in thawing permafrost has been observed through increases in respiration, methanogenesis, and denitrification (Hultman et al., 2015; Palmer et al., 2012). While specific taxa upregulate broad transcriptional activity (with increased total RNA production) during permafrost thaw, only a fraction of the total community participates in biogeochemical cycling (Coolen & Orsi, 2015). Despite low levels of activity under frozen conditions, the fraction of viable microorganisms in permafrost is highly variable: 18-63% (La Ferla et al., 2017; Hansen et al., 2007; Mackelprang et al., 2017); much lower than the 91.3% viability ratio found in thawed temperate soils (Janssen et al., 2002). It is unknown whether increased microbial activity with thaw is driven by an increase in microbial viability or greater activity of a small fraction of live microbes.

Dramatic changes are induced by permafrost disturbances, with long-lasting impacts on plant community composition, plant physiognomy, and soil chemistry in both modern and historic sites (Forbes et al., 2001). Soil horizon removal is the most severe of these disturbances, requiring 20-75 years to recover plant community function (Forbes et al., 2001).

As industrial activity becomes more commonplace in the subarctic, anthropogenic disturbances may disrupt plant communities and shift the source of this fluxed carbon to old, frozen, and recalcitrant deposits. Both natural and artificial disturbances can induce lake formation. In other permafrost affected environments outside of the Arctic, artificial disturbances – namely infrastructure development – are the most common causes of thermokarst lake formation in the Qinghai-Tibetan Plateau (Lin et al., 2016). The construction of oil wells, pipelines, gold mines, and other mineral extractions are also becoming more common in the Arctic (Elias, 2014). These industrial developments are conducive to the formation of thermokarst ponds across the Circumpolar North. It is unknown how the microbial community of active layer and permafrost will change in response to these direct industrial disturbances or if industrial disturbances differ from natural disturbances.

In this site-specific study, we assess the soil microbial community structure shifts and drivers at a locally disturbed site where soils and vegetation were stripped in preparation for mining at a permafrost-affected site near Dominion Creek, Yukon, Canada. The study site was disturbed by the installation of a drainage ditch a few weeks prior to our arrival at the site by a large excavator (Figure 1). As microbial diversity has been closely linked to edaphic parameters, we hypothesized that microbial community shifts in disturbed permafrost are dependent on edaphic parameter changes. In addition, we hypothesized that thaw would induce the proliferation of viable microbial cells, increasing microbial viability ratios with disturbance, due to the exposure to atmospheric temperatures for six weeks. This information helps clarify how permafrost communities restructure upon thaw.

3 Methods and Materials

3.1 Field Site and Sampling Procedure

To assess the impacts of disturbance and thaw on permafrost-affected soils, we sampled after a disturbance along a gradient from undisturbed to a thawed thermokarst pond (Figure 1). Surface samples and soil cores were collected in May 2016 at a gold mining site near Dominion Creek, Yukon Territory, Canada (NTS:116-B/3; DMS: 60°42'59.5"N, 138°33'0.14"W). Cores were taken with a light portable gas-powered permafrost drill (EDR-260, ECHO Inc., Lake Zurich, IL, USA), (Saidi-Mehrabad et al., 2020). Cores were collected using a system devised in Calmels et al. (2005) which takes individual ~40 cm cores which are then broken off from the surrounding permafrost and recovered. The study site was disturbed in preparation for placer gold mining a few weeks prior to our arrival at the site by a large excavator. The primary disturbance was the development of a drainage ditch (Figure 1c) which removed the upper active layer at Core Site 2, and all of the active layer at Core Site 3 (Figure 1b). Site 1 was located away from the disturbance site.




Figure 1: Dominion Creek core and surface sample locations. (a) Map of Dominion Creek area and sampling site within Yukon Territory (Source: USGS & The Government of Canada). The map of the Dominion Creek region was assembled in ArcGIS 10.6.1. (b) Cross-section of disturbance gradient soils and permafrost cores. The horizontal line represents the

surface soil profile as it had been disturbed from road construction preparation. The undulating line refers to the forming thermokarst pond. A scale depicts the depth of each permafrost sampling layer. The dashed blue line refers to the permafrost table. (c) The disturbance gradient soils including A (least disturbed), B-D (disturbed active layer), and E-F (disturbed permafrost). Disturbed permafrost samples are inset and were found 3 metres away at a shallower location of the thermokarst pond.

Quadruplicate surface samples (unfrozen soils) were taken at six locations including undisturbed permafrost-affected soils (site A; 474 cm from the thermokarst pond); disturbed active layer soils (sites B-D; 363 cm, 288 cm, and 213 cm from the thermokarst pond, respectively); and putatively disturbed permafrost sediments (sites E-F; 109 cm and 0 cm from the thermokarst pond, respectively). Organic layers and surface vegetation were removed with a spade before sampling. A separate spade was cleaned with 10% bleach and 70% ethanol before sampling at approximately 5 cm depth and stored in sterile Whirlpak® bags and homogenized by mixing both in the field and in the lab prior to chemical and biological analyses.

Soil cores, which were 1-3 metres in depth, were collected at three sites along the same thaw gradient (from Core 1 – Core 3). Core 1 was sampled in a nearby clearing analogous to the conditions at sample A (the “undisturbed” site). Core 2 was sampled 2.6 m from the thermokarst pond between surface sample sites C and D. The Core 2 site had all vegetation, the organic soil horizon, and the surface soil horizon of active layer removed; however, active layer soils were still present in Core 2. Core 3 was sampled directly adjacent to the thermokarst pond, with all soil layers above the permafrost table removed. The thaw depth was 14 cm, 34 cm, and 85 cm for Cores 1, 2, and 3, respectively (Figure S1). All

thawed soil above 15 cm was removed prior to coring. Each core subsection had a 10 cm diameter and ranged in length from 14 – 36 cm long, the total core depth and reached at least 1 m for every core as measured from the soil surface. Loose material on the surface of soil cores was removed by scraping with razor blades and discarded. Frozen soil cores were placed in clear polypropylene bags (Uline, Pleasant Prairie, WI, USA), transferred to coolers with ice packs in the field, transported frozen to the laboratory, and stored at -20°C until further analysis. Permafrost cores were collected frozen while surface samples were thawed at the time of collection. All samples were transported back to a freezer in Dawson City, frozen to ca. -20 ° C and transported frozen to the University of Alberta Permafrost Archives Laboratory.

Surface samples along a disturbance gradient were collected to assess the impact of thaw as a result of the disturbance, allowing for direct comparison to undisturbed frozen soil cores beneath them. In this study we consider the following as active layer samples: Core 1 - 30 cm, Core 2 - 30 cm, and Core 2 - 45 cm. Active layer soil was distinguished from permafrost by measuring gravimetric water content at 1 cm intervals along each core. A pronounced increase in water content was observed at the transition zone at 45 cm as shown in Figure S1: Core 1 - 45 cm, Core 1 75 cm, Core 1 95 cm, Core 2 75 cm, Core 2 95 cm, Core 3 75 cm, and Core 3 95 cm. Soil water accumulates at the bottom of the active layer, allowing for determination of the maximum active layer depth through the determination of gravimetric water content (Chang et al., 2024). Surface sample A was denoted as an undisturbed active layer soil, surface samples B-E were denoted as disturbed active layer soil, and sample F was denoted as a disturbed permafrost soil (Figure 1). Adjacent undisturbed areas were used as controls to assess the impact of disturbance on microbial community

composition. These soils were collected as nearby as possible to minimize the impact of soil spatial heterogeneity, however, some underlying spatial variation cannot be fully removed.

3.2 Soil Chemistry Subsampling and Analysis

Physicochemical analysis was performed as in Saidi-Mehrabad et al., (2020). Surface soils were subsampled as follows: 0.5 g for loss on ignition (LOI), 0.5 g for gravimetric water content; 0.5 g for pH; 5 g for nitrate analysis; 0.5 g for dissolved metal and TN/TC (Total Nitrogen/Total Carbon) analyses. Cores were longitudinally sectioned into ~1/3 (3 cm width) and ~2/3 (7 cm width) sections for chemical and biological analyses, respectively. An additional two longitudinal subsections were taken from the 1/3 portions (chemical analysis section) of the cores. Subsections were scraped with a razor blade to remove all thawing and contaminating materials. The first section was cut into 1 cm³ cubes and used for pH, SOM, and gravimetric water content throughout the entire profile of each core. Soil water content was determined by oven drying the samples at 105°C for 24 hours. Soil pH was determined using a soil to 0.01 M CaCl₂ ratio of 1:2 with an AB15 pH meter (Fisher Scientific, Waltham, MA, USA). Soil organic matter was measured using loss on ignition (LOI) procedures (Lim & Jackson 1982), using a Lindberg SB Muffle Furnace (Thermo Fisher Scientific Inc., USA). Subsamples from the second section were homogenized, and 0.5 g of material was taken for TN, TC and dissolved metals while the remainder was used for NO₃ and NH₄ measurements at the Natural Resources Analytical Laboratory (NRAL, Edmonton, AB, Canada). Total Nitrogen and TC were analyzed using standard dry combustion methods with a Costech Model EA 4010 Elemental Analyzer (Costech International Strumatzione, Florence, Italy, 2003) (Sparks et al., 1996). Extraction of NO₃ and NH₄ was performed using a 2M KCl solution and measured using a Diazo Coupling method with a SmartChem Discrete Wet Chemistry Analyzer, Model 47 200 (Westco Scientific, Brookfield, CT, USA) (Maynard et

al., 1993). Trace metal analysis was conducted after an HNO₃/HCl acid digest using Inductively Coupled Plasma-Optical Emission Spectroscopy (ICP-OES) which measured P, K, S, Mg, Ca, Fe, Cu, Mn, Zn, Na using standard protocols on a Thermo iCAP6300 Duo ICP-OES (ThermoFisher, Cambridge, United Kingdom) (Skoog et al., 2007).

3.3 Biological Subsampling and Contamination Detection

The 2/3 portions of the cores (“biological sections”) were cut into discrete 5 cm long sections and subsamples were taken at 15 cm, 30 cm, 45 cm, 75 cm, and 95 cm below the soil surface: Core 1 was subsampled at all horizons; Core 2 was subsampled at 30 cm, 45 cm, 75 cm, and 95 cm; Core 3 was subsampled at 75 cm, and 95 cm. Contamination detection was performed as in (Saidi-Mehrabad et al., 2020). To detect contamination, each subsample was painted with 10 mL of 1 × PBS solution containing *E. coli*:DH10B transformed with pBAD *mNeonGreen* at 8.3×10^9 cells/mL; for a total of 8.3×10^{10} cells/core in a 4°C walk-in refrigeration unit. All solutions (including 10% bleach and water) were kept at 4°C to maintain the integrity of the core. Any thawed soil was removed using an autoclaved and sterile razor blade. The remaining soil was reserved for DNA extraction. Following decontamination, the presence of PCR-amplifiable pBAD vector DNA would be interpreted as potential contamination. DNA extraction subsamples were prepared in a class 10,000 clean lab at the University of Alberta which has not been used previously for DNA amplification. To decontaminate cores, samples were: 1) immersed in 100% bleach solution, 2) washed with DNA-free water, and 3) scraped approximately 4-7 g of material using a razor blade; this cycle was repeated twice. Subsamples were then homogenized within a sterile Whirlpak® bag. The resulting decontaminated soils were then stored at -20°C until DNA extraction.

Genomic DNA was extracted from 0.5 g samples using the MOBio PowerSoil kit according

to the manufacturer's instructions. Replicate extractions were pooled before contamination checks and sequencing.

Conditions for pBAD PCRs are outlined in Saidi-Mehrabad et al., (2020). In brief, contamination was assessed using primers pBAD-forward and pBAD-reverse under standard PCR conditions with Platinum Taq DNA polymerase. No PCR products were observed in all DNA extractions, indicating no contamination. Blank extractions were performed on each of the three DNA extraction kits. To check for DNA contaminants, PCRs of blank exactions were performed using the primers 341F and 518R (Muyzer et al., 1993). No bands were observed following gel electrophoresis of blank extract PCR amplification.

3.4 16S rRNA Gene Amplicon Sequencing and Analysis

The microbial communities of soil samples were surveyed by small-subunit rRNA gene amplicon sequencing. DNA from the V4 region of the 16S rRNA gene was amplified with the primers 515F and 806R primers (Caporaso et al., 2012). Sequencing was performed commercially by Microbiome Insights (Microbiome Insights, Vancouver, BC, Canada), with an Illumina MiSeq using V2 chemistry (Illumina, San Diego, California, United States). Before sequence processing, a total of 552,252 reads were recovered from 18 samples including surface samples, core samples, as well as 5 blank extractions, negatives. The read count ranged from 10,525 to 47,389 reads per sample, and most samples had high coverage (Figure S3). Sequence processing was performed in USEARCH v10 (Edgar, 2010).

Overlapping reads were combined from forward and reverse reads with the following qualifiers: no unknown base pairs allowed (Ns), a maximum number of sequence mismatches of 10 nt in the alignment, a minimum merge length of 230 bp, a maximum merge length of 300 bp, and an ID cut-off of 80%. Of the 552,252 reads recovered, 515,916 reads were aligned into overlapping reads (93.4% of the total), at an average length of 253 bp, with an

240 expected size of 291 bp. Quality filtering with a maximum expected error cut-off of 1.0
resulted in 510,757 (99.0% of the total) reads passing (Edgar & Flyvbjerg, 2015).
Dereplication identified 148,294 unique sequences and 105,589 singletons (71.2% of the
unique reads) for the total dataset. OTU clustering at 97% using UPARSE identified a total of
5,964 OTUs as well as identifying and removing 12,508 chimeras while removing all
245 singletons and doubletons (Edgar, 2013). After sequence processing 386,419 reads remained.
Taxonomy was assigned using *de novo* picking as implemented in SINTAX using the RDP
v16 database at a bootstrap cut-off of 80 (Ribosomal Database Project) (Edgar, 2016).
Sequences were separated for bacterial (377,578 reads) and archaeal (8,841 reads) analysis.
Taxa identified as chloroplast/streptophyta using SINTAX were removed from the
250 downstream analysis and no mitochondrial reads were observed, leaving 361,034 reads in the
bacterial dataset (Edgar, 2016). Reads identified in extraction blank samples were removed
from analysis. A NEWICK formatted 16S rRNA phylogenetic tree was constructed using
USEARCH v10. For bacterial analysis, samples were rarefied to 10,000 reads.

3.5 Microbiological Diversity, Taxonomy and Assembly

255 α -diversity metrics, including species richness, Chao 1 richness estimation, Shannon
Diversity, Shannon Evenness, Simpson Diversity, Simpson Evenness, and Heip's Evenness
were calculated using Mothur v 1.39.5 (Schloss et al., 2009). A Kruskal-Wallis test followed
by posthoc testing with Kruskal-Wallis multiple comparisons as implemented in SigmaPlot
v13.0 tested for significant differences between samples. β -diversity was calculated
260 USEARCH v10 using Bray-Curtis, Jaccard presence/absence, and weighted UniFrac distance
metrics. Visualization of community dissimilarities and significance testing were carried out
using R v3.4.1, in the RStudio IDE with the *vegan* package v2.4-4 (Philip, 2003).
Hierarchical clustering was visualized with the dendextend package v1.5.2 (Galili, 2015).

Statistical differences in community composition between clusters were assessed using

265 permutational multivariate analysis of variance in the *vegan* package v2.4-4. Environmental parameters were fitted to the NMDS based on a significant Spearman correlation ($p < 0.05$) plotted using a total of 999 permutations. The direction of vector indicates the direction of change while the length is proportional to the correlation between the communities and variable. Differences in soil chemistry were visualized in a redundancy analysis (RDA).

270 Values used a Z-score standardization technique (“standardize”) to create a standard normal distribution of 0 with a mean standard deviation of 1. A backward selection of the model using ordiR2step in *vegan* was used to determine the most appropriate variables for the RDA which included: P (%), K (%), S (%), Mg (%), Ca (%), Fe (%), Cu (mg/kg), Mn (mg/kg), Zn (mg/kg), Na (mg/kg), NO₃ (mg/kg), NH₄ (mg/kg), TN (%), TC (%), SOM (%), pH, and

275 water content. Differences between permafrost – and permafrost layers – as well as active layer – and active layer – were determined using a PERMANOVA test in R. Phylum, class, and genus level tables were constructed using SINTAX (Edgar, 2016). A stacked bar chart of relative abundance for class-level taxonomic classification was constructed in MS Excel. Assessment and visualization of significant ($p < 0.05$) differential abundances were assessed

280 using analysis of the composition of microbiomes (ANCOM) (Mandal et al., 2015). Significance testing was followed by post-hoc analysis using a false discovery rate to correct for multiple comparisons. Only significantly different groups were visualized. Indicator OTUs for either permafrost cores or active layer cores were determined using the *indicspecies* package in R (De Caceres et al., 2016) using an $\alpha < 0.05$ and a minimum IndVal

285 of 0.8. Active layer OTUs were defined from Section 3.1, where active layer samples included Core 1 - 30 cm, Core 2 - 30 cm, and Core 2 - 45 cm while permafrost samples included Core 1 - 45 cm, Core 1 75 cm, Core 1 95 cm, Core 2 75 cm, Core 2 95 cm, Core 3

75 cm, and Core 3 95 cm. The presence of either permafrost indicator OTUs or active layer OTUs was then determined in surface soils using Mothur v 1.39.5 using the shared richness function.

3.6 Cell Enumeration

Permafrost, active layer, and surface soils (0.5 g) were diluted 1:100 (w/v) in 10 mM tetrasodium pyrophosphate (Fisher Scientific, Hampton, NH, USA). The resulting soil slurry was hand shaken for 15 minutes with 5 ml of 2 mm glass beads then sonicated at 42kHz for 2 × 40 s in an ultrasonic bath (Elma, Singen, Germany), and allowed to settle for 5 minutes. Slurry taken from the middle of the sample bottle was further diluted to a final 1:100 (or 1:500 for surface sample A) in 10 mM tetrasodium pyrophosphate (v/v) before pre-filtration through a 5 µm pore size polysulfone filter (Burlington, Massachusetts, United States) to remove larger particles. The resulting filtrate was then passed through a black polycarbonate 0.22 µm filter (Burlington, Massachusetts, United States). Bacterial cell counts were determined by epifluorescence microscopy using the LIVE/DEAD® BacLight™ staining solution with 6µM Syto 9 and 30 µM propidium iodide (Carlsbad, California, United States) (Boulos et al., 1999). Filters were stained in a 1:1 mixture (300µL stain solution: 300µL 0.85% NaCl) for 1 h in the dark at 4°C. Fifteen random fields of view were counted on a Leica DMRXA fluorescent microscope. The resulting live and dead cell counts were averaged for each sample, and corrected for by dry weight, with the sum of both live and dead representing the total cell counts.

Surface samples were sampled in triplicate (n = 3; A-F). Permafrost samples included Core 1 (45 – 50 cm, 75 – 80 cm, 95 – 100 cm), Core 2 (75 – 80 cm and 95 – 100 cm) and Core 3 (75 – 80 cm and 95 – 100 cm) (n = 7). Active layer samples included Core 1 (15 – 20 cm, 30 – 35 cm), and Core 2 (30 – 35 cm, 45 – 50 cm) (n = 4).

4 Results

4.1 Edaphic Parameter Analysis of Dominion Creek Soils

Edaphic parameters separated active layer soils from permafrost soils in soil cores in an RDA (Figure 2 and Supplementary Tables 1 and 2; $p = 0.035$). Disturbed soils did not form a separate grouping to the exclusion of permafrost and active layer core samples ($p = 0.084$). Of the surface soils, A-E grouped with active layer soils, while F grouped with permafrost soils (Figure 2; $p = 0.022$), indicating that while A-E likely originated from active layer soils, F likely originated as a permafrost soil. Surface soil samples of C and D were distinct from other soil samples (Figure 2); C and D had higher Mn, P, SOM, and water content than other samples (Table S1 & S2).




Figure 2: Thawed permafrost surface soils were similar to intact permafrost samples, while thawed active layer surface soils were similar to intact active layer samples. Redundancy analysis of soil physicochemical parameters in permafrost, active layer, and disturbance gradient soils ($R^2 = 0.9890434$). Soil chemical parameters are represented by arrows, and sites are distinguished by coloured circles. Darker circles represent greater disturbance across the disturbance gradient soils.

4.2 Viability is negatively impacted by disturbance

To assess the impact of disturbance on active layer and permafrost microbial cell abundance and viability, counts of live and dead cells were determined across surface samples and core samples. The proportion of viable cells (Table 1) ranged from 18-86%. Microbial cell counts ranged from 9.10×10^7 (in Core 1 95 cm) to 9.20×10^8 (in surface sample A) cells gdw^{-1} . Total microbial abundance in active layer cores was greater than in permafrost cores, with microbial abundance averages of $1.26 \times 10^8 \pm 6.30 \times 10^7$ cells gdw^{-1} in active layer cores and $2.30 \times 10^7 \pm 2.79 \times 10^7$ cells gdw^{-1} in permafrost cores (Table 1). The viability ratios of permafrost soils were not significantly different from active layer soils ($p = 0.07$). Surface soil cell abundance declined significantly in surface samples E and F ($\sim 1.8 \times 10^8$ cells gdw^{-1}) in comparison to other surface soils ($\sim 5.2 \times 10^8$ cells gdw^{-1}) (Table 1). Viability ratios did not change significantly with the degree of disturbance, although viability ratios in disturbed permafrost and active layer soils (B – F) were significantly lower than undisturbed active layer (A) (Table 1).

Table 1: Live-dead cell counts of viable and non-viable microbial cells across the disturbance (A-F) gradient, as compared to undisturbed permafrost and active layer core

samples. Counts are grouped (like letters are the same group) based on a one-way ANOVA (P<0.05) with a post-hoc Tukey HSD calculator. Error indicates standard deviation as calculated through propagation of error across each field of view as well as biological replicates.

	Viable (cells gdw ⁻¹)	Non-Viable (cells gdw ⁻¹)	Total (cells gdw ⁻¹)	Ratio (%)
A	$6.28 \times 10^8 \pm 1.27 \times 10^8$ (a)	$2.92 \times 10^8 \pm 1.46 \times 10^8$ (ab)	$9.20 \times 10^8 \pm 1.31 \times 10^8$ (a)	26.79 ± 10.14 (a)
B	$1.15 \times 10^8 \pm 4.86 \times 10^7$ (bc)	$1.68 \times 10^8 \pm 1.41 \times 10^8$ (bc)	$2.83 \times 10^8 \pm 1.87 \times 10^8$ (bcd)	18.13 ± 3.87 (ab)
C	$8.65 \times 10^7 \pm 2.46 \times 10^7$ (bc)	$2.78 \times 10^8 \pm 7.37 \times 10^7$ (ab)	$3.64 \times 10^8 \pm 8.48 \times 10^7$ (bc)	21.29 ± 13.59 (b)
D	$1.29 \times 10^8 \pm 4.53 \times 10^7$ (bc)	$3.95 \times 10^8 \pm 9.24 \times 10^7$ (a)	$5.24 \times 10^8 \pm 1.88 \times 10^7$ (b)	26.93 ± 4.98 (b)
E	$8.57 \times 10^7 \pm 8.09 \times 10^7$ (bc)	$1.18 \times 10^8 \pm 5.1 \times 10^7$ (bc)	$2.03 \times 10^8 \pm 1.15 \times 10^7$ (cd)	49.08 ± 11.86 (b)
F	$4.20 \times 10^7 \pm 6.76 \times 10^6$ (bc)	$1.16 \times 10^8 \pm 4.25 \times 10^7$ (bc)	$1.58 \times 10^8 \pm 4.84 \times 10^7$ (cd)	36.74 ± 7.76 (b)
Permafrost	$2.30 \times 10^7 \pm 2.79 \times 10^7$ (b)	$6.77 \times 10^7 \pm 2.86 \times 10^7$ (c)	$9.07 \times 10^7 \pm 4.11 \times 10^7$ (d)	25.43 ± 14.41 (b)
Active Layer	$1.26 \times 10^8 \pm 6.3 \times 10^7$ (c)	$1.84 \times 10^8 \pm 7.27 \times 10^7$ (bc)	$3.10 \times 10^8 \pm 1.14 \times 10^7$ (bc)	39.92 ± 9.21 (b)

4.3 Diversity of disturbed soil resembles active layer soils

Observed OTU richness across all samples ranged from a low of 387 OTUs in the deepest Core 1 permafrost sample to a high of 1738 OTUs in surface sample F, the disturbed permafrost soil (Table S3). Higher Shannon diversity, Shannon evenness, and OTU richness in active layer indicate that active layer communities harboured a more diverse microbial community at the OTU level than in permafrost (Figure 3 and Table S3). Lower evenness suggests the permafrost microbial communities were dominated by a small number of abundant OTUs.




Figure 3: Microbial diversity in permafrost responds to abrupt thaw. Box and whisker plots

of diversity metrics across active layer, permafrost, and disturbance gradient soils.

Disturbance gradient soils were combined for statistical testing. Letters signify statistically different groupings ($p < 0.05$) as assessed using one-way ANOVA with a post-hoc Tukey test. Surface samples were assessed as a single statistical grouping ($n = 6$). Permafrost samples were grouped across Core 1, Core 2, and Core 3, including depths 75 – 80 cm and 95 – 100 cm as well as 45 – 50 cm in Core 1 ($n = 7$). Active layer samples were grouped together across Core 1 (15 – 20 cm, 30 – 35 cm) and Core 2 (30 – 35 cm, 45 – 50 cm) ($n = 4$).

Surface soils were not significantly different from active layer soils in any α -diversity metric; both had higher richness, diversity, and evenness than permafrost soils (Figure 3 and Table S3). Microbial diversity of disturbance gradient soils most closely resembled undisturbed active layer soils and differed from permafrost.

4.4 Microbial community composition shifts with disturbance

There were three distinct bacterial community clusters in the studied soils: (1) upper active layer and undisturbed soils (surface sample A); (2) lower active layer and the disturbed soils (surface samples B-F); and (3) exclusively permafrost ($p < 0.05$) (Figure 4). These clusters were observed with different community distance metrics, including Jaccard presence/absence (Figure S4), OTU composition (Figure 4), and phylogenetic dissimilarity

(Figure 6). All disturbed soil communities clustered with lower active layer soil communities (including Core 1 30 – 35 cm, Core 2 30 – 35 cm, and Core 2 45 – 50 cm). Sample groups separated across NMDS1, while NMDS2 separated within-cluster community differences. Active layer and disturbed soils group together to the exclusion of permafrost soils (Figure 4).




Figure 4: Dissimilarity of microbial communities visualized by nonmetric multidimensional scaling (NMDS) of total microbial assemblages. Groupings of soil samples are based on clusters recovered from a hierarchical clustering algorithm using Bray-Curtis community distances. Ellipses encompass clusters determined by weighted average algorithms: (1) upper active layer and undisturbed soils, (2) lower active layer and disturbed soils, and (3) permafrost. Stress = 0.06.

Both Bray-Curtis and Jaccard presence/absence distances within clusters correlated significantly with differences in pH ($r^2 = 0.4316$ and 0.5000 , respectively, $p < 0.05$), while differences across clusters correlated with Zn concentration ($r^2 = 0.3609$ and 0.3461 , respectively, $p < 0.05$) (Figure 4).

In an attempt to describe additional community variation, cohesion metrics (where positive cohesion represents a situation in which both taxa become more abundant with association while negative cohesion where the presence of one taxon negatively impacts another taxon) were compared against Bray-Curtis dissimilarity. Positive cohesion ($r^2 = 0.6760$) and negative cohesion ($r^2 = 0.8069$) explained 29.0% of bacterial community variation and 40.2% of the community variation when combined with pH and Zn (Figure 4). Cohesion not only explained a greater fraction of bacterial community variation but also explained an additional 20.3% of variation beyond environmental parameter differences, as Zn and pH explained only 19.9% of the variation alone.

4.5 Persistence of permafrost indicator OTUs in disturbed permafrost soils

Indicator taxon analysis identified 94 active layer indicator OTUs (Table S4) and 60 bacterial permafrost indicator OTUs (Table S5). Despite disturbance and thaw, permafrost indicator OTUs persisted following permafrost thaw at relatively high proportions in sample F. The proportion of permafrost indicator OTUs present in surface samples increased in both disturbed active layer and disturbed permafrost soils, with less than 5% of permafrost indicator OTUs present in the undisturbed surface sample (sample A) increasing to 80% of bacterial permafrost indicator OTUs present in sample F (Figure 5a). By comparison, the presence of permafrost indicator OTUs ranged from 93-100% in permafrost soils.




Figure 5: Indicator species analysis and shared richness assessed across the disturbance gradient as compared to combined active layer and permafrost soils. (a) The proportion of indicator OTUs assigned to permafrost or active layer soils that are shared within disturbance gradient soils. Indicator OTUs were defined as having indicator values above 0.80. Each indicator group is calculated independently. The proportion represents the fraction of indicator OTUs from either active layer or permafrost that are present within each surface soil. (b) Shared richness of disturbance gradient soils compared against the combined assemblages of permafrost or active layer. “Uniques” were unique to the disturbance gradient soil, and “Boths” were found in both active layer samples and permafrost samples.

Across the surface soils, F shared the highest proportion of OTUs with permafrost soils (9.0%) and the lowest with active layer soils (26.8%) (Figure 5b). Likewise, the undisturbed soil shared the lowest proportion of OTUs with permafrost soils (0.1%), and the

highest with active layer soils (45.7%). Endemism was highest in soil F (29.0%), but lowest in intermediately disturbed soils C and D (19.1% and 22.3% respectively). Endemic OTUs were lowest in disturbed active layer but increased within disturbed permafrost. OTUs shared with permafrost soil communities increased in prominence with greater levels of disturbance. Thus, although the overall community composition in the most disturbed soils was quite similar to active layer soils, indicator bacterial OTUs of relict permafrost were still present. We interpret these findings to indicate that the permafrost microbial communities were still present but were being supplanted by a colonizing active layer community following thaw.

4.6 Community membership of disturbed soils resembles active layer soils

Phylogenetic distances of community composition (weighted UniFrac) indicated similar cluster structure as found in Bray-Curtis and Jaccard distances (Figure 6). Community dissimilarity was greatest in permafrost soils, indicating that permafrost soil microbial communities were more heterogeneous than those of disturbed active layer and undisturbed active layer soils. A total of 26 phyla, 66 classes, and 235 genera were assigned across all soils, with 32.4%, 46.1%, and 62.3% of all sequences unassigned at the phylum, class, and genus levels, respectively. At the phylum level, all bacterial communities combined were predominantly composed of Proteobacteria (21.1%), Actinobacteria (8.6%), Acidobacteria (7.9%), Bacteroidetes (7.9%), and Firmicutes (5.5%), with substantial proportions of Verrucomicrobia (4.2%), and Planctomycetes (4.0%) and above a 3% cut-off. Across class levels, Actinobacteria (8.5%), Alphaproteobacteria (6.6%), Deltaproteobacteria (4.2%), Planctomycetia (4.0%), Gammaproteobacteria (3.7%), Clostridia (3.3%), and Betaproteobacteria (3.2%) were most abundant and above a 3% cut-off (Figure 6). Bacilli, Clostridia, Anaerolineae, and Bacteroidia were significantly more abundant in permafrost soils than in active layer soils, while Acidobacteria group 4, Acidobacteria group 17, and

Nitrospira were all significantly more abundant in active layer soils than permafrost (Figure S2). Class level taxonomic composition across the surface soils was relatively static with some exceptions: Clostridia (A= 0.04%, B =0.529%, F= 1.96%), Bacteroidia (A=0.03%, B=0.49%, F=2.1%), and Acidobacteria group 7 (A=0.71%, B=1.81%, F= 3.19%) all increased with disturbance, while Planctomycetia decreased (A=2.85%, B=1.63%, F=0.775%). Bacteroidia and Clostridia were more abundant in disturbed and permafrost soils than in active layer soils.




Figure 6: Hierarchical clustering of cores and surface sample communities alongside class level relative abundances. Classes <1% in any sample was grouped into “other” alongside unassigned taxa. Samples which are not significantly different ($P < 0.05$) across weighted UniFrac distances have the same colour. Active layer samples are labelled in red, permafrost is labelled in blue, and disturbance gradient soils are labelled in black.

5 Discussion

5.1 Analogues of natural disturbances

Discrete naturally occurring physical disturbances to active layer and permafrost soils (such as retrogressive thaw slumps, thermokarst thaw ponds, and active layer detachment

slides) are becoming increasingly common. Removal of the active layer, as well as subsequent detachment slides of the newly forming active layer, results in a similar geographic formation as anthropogenic excavation. Indeed, anthropogenic disturbances can also result in active layer detachment slides and thermokarst formation (Lin et al., 2016). Our data indicate that the edaphic parameters measured, including a variety of micronutrients, did not shift as a result of permafrost thaw or disturbance. The exposure of preserved organic acids, such as fulvic acids and humic acids, likely decreased the pH in our disturbed permafrost soils. Both excavated sites and naturally disturbed sites undergo slow re-vegetation, and we predict that although microbial community replacement occurs rapidly, plant community succession may occur over decadal time scales in anthropogenically and naturally disturbed sites (Forbes et al., 2001; Lantz et al., 2009). With time, additional nutrients may be released into the newly formed surface soils from permafrost soils, including $\text{NO}_3\text{-N}$, $\text{NH}_4^+\text{-N}$, Na, and S found in disturbed permafrost samples in this study. Future studies should follow industrially disturbed sites over multiple years to investigate how evolving edaphic parameters change with increased mobilization.

5.2 Anthropogenic Disturbance does not reliably shift edaphic parameters

As water potential, nitrogen content, carbon content, and nutrient availability are amongst the most important drivers of microbial community composition in these soils, permafrost thaw is expected to lead to microbial community shifts. Anthropogenic disturbance alters soil pH, while thawing permafrost leaches micronutrients and macronutrients into downstream thaw waters (Forbes et al., 2011; Lantz et al., 2011). Organic matter in permafrost is generally more recalcitrant and holds higher concentrations of dissolved nitrogen than active layer soils above it (Keuper et al., 2012). The dissolved nitrogen preserved within permafrost can be rapidly released following permafrost thaw,

500 releasing nutrients into the active layer and downstream aquatic systems (Reyes & Lougheed, 2015). Thus, we expected dramatic changes in soil edaphic parameters following disturbance in the Dominion Creek system, including a lowered pH, increased mineral nitrogen, increased carbon content, and increased micronutrient abundance. However, the measured edaphic parameters did not shift within six weeks of thaw, although higher nitrogen contents found in 505 disturbed permafrost were similar to the high levels of nitrogen found in undisturbed permafrost. The lack of a shift may be due to the short timeline since thaw; environmental parameters may need more time to shift in these disturbed soils, particularly as many of our analyzed metals have a slow rate of change.

5.3 Permafrost bacterial communities respond to disturbance

510 Most experiments studying *in situ* permafrost thaw show either that microbial communities shift to resemble the active layer (Tas et al., 2014), or that new microbial communities develop (Mondav et al., 2017). Unlike *in situ* experiments, laboratory thaw experiments identify significant changes in microbial community structure, transcriptional activity, respiration, and genetic potential (Coolen & Orsi, 2015; Coolen et al., 2011; 515 Oelbermann et al., 2008; Mackelprang et al., 2011). Interestingly, we observe rapid microbial community shifts in disturbed permafrost soils in response to thaw. Previous incubation experiments identify microbial community changes in both active layer and permafrost soils with rapid thaw and suggest that the rapid succession of microbial communities is related to carbon recalcitrance (Coolen et al., 2011). However, while we observed a significant shift in 520 microbial composition in disturbed permafrost soils, we did not observe the same trend in disturbed active layer soils. Together, we suggest that microbial community dynamics with significant physical disturbance is likely to induce rapid microbial community shifts in permafrost soils, but not in active layer soils. This can be extended not only to anthropogenic

disturbances, but to thermokarst formation as well. During this *in situ* thaw experiment,
525 microbial community successional patterns were observed; however, interpretations of
function from taxonomic assignment must be made with caution, requiring further study into
the genetic potential and activity of the community. Thermokarst formation as found in this
study can induce anaerobic conditions, increasing acetogenic fermentation and
methanogenesis (Coolen & Orsi et al., 2015). Future study into methane cycling, such as
530 through the use of stable isotope probing of methanotrophic communities, may be
particularly beneficial to understanding functional shifts in anthropogenically disturbed
permafrost soils such as the ones studies here.

Indeed, longer-term field warming experiments show a deepening of the ice table,
allowing the colonization of plant roots into deeper soil horizons (Monteux et al., 2018).

535 Microbial communities in these thawed permafrost layers shifted to resemble those of the
active layer. The mechanism of this change is the colonization of active layer taxa into deeper
soil horizons. Our results support these findings, as permafrost microbial communities shifted
to resemble the community of the lower active layer. However, in the absence of plant roots,
we suggest the perturbation of soil horizons by anthropogenic activities allowed the
540 colonization of active layer microbial communities into thawed permafrost horizons. Physical
mixing may also contribute to these community shifts, which has previously been shown in
laboratory experiments (Doherty et al., 2025). These results are supported by the presence of
both active layer and permafrost indicator OTUs within disturbed permafrost soils, as a small
abundance of permafrost microbiota may have persisted after disturbance. We expect that
545 disturbances, such as thermokarst formation, in the Arctic and subarctic are likely to create
homogenous microbial communities similar to the active layer regardless of the soil source.
Rather than observing indigenous microbial communities shifting in abundance in response

to thaw, we observed dispersal of active layer microorganisms into exposed permafrost layers. Soil fauna, such as collembola, may be responsible for the transport of soil

microorganisms into thawing permafrost layers (Monteux et al., 2022). Therefore, we suggest that discrepancies between *in situ* thaw experiments and laboratory thaw experiments may be due in part to the lack of surrounding active layer disturbance in the *in situ* experiments.

5.4 Drivers of Microbial Community Composition are Biotic

Resident microbiota must adapt to numerous shifting environmental variables in response to permafrost thaw or they will be supplanted by active layer microbiota.

Physicochemical parameters are suggested to control the fate of microbial communities in response to permafrost thaw (Hayden et al, 2012; Rousk et al, 2012); however, rapid microbial community shifts have been observed under only seven days of warming, before any significant changes in chemistry or physical parameters in the system would be possible (Mackelprang et al 2011). Therefore, it is unknown if: (1) physicochemical characteristics control microbial communities following permafrost thaw, and so microbial communities in samples with similar environmental parameters should resemble each other (environmental filtering) or (2) permafrost thaw induces microbial community shifts, and so environmental and microbial groupings do not resemble one another (biotic filtering). Soil microbial

communities in thawing permafrost are generally driven by environmental factors, most significant of which is pH (Chen et al., 2017; Tripathi et al., 2018). We therefore expected that disturbance and thaw would result in rapid changes to pH, water content, micronutrient concentration, and exchangeable nitrogen content which would correlate to microbial community composition. These anticipated changes reflect documented biogeochemical responses to thaw (Scheel et al., 2022; Chen et al., 2017; Tripathi et al., 2018). However, in our study the variance in bacterial community composition was poorly explained by

physicochemical parameters. One exception within our study, the correlation of Zn to bacterial community composition, may be due to the site-specific mobilization of Zn during permafrost thaw or due to other underlying processes (Burn et al., 2025). Previous studies have shown permafrost and active layer prokaryotic communities are controlled by distinct mechanisms (Chen et al., 2017; Tripathi et al., 2018). However, we observed no such correlation between microbial composition and these parameters in our system. Our results therefore support hypothesis 2, that microbial community changes are induced by biotic filtering.

Interestingly, while the short 6-week time frame since disturbance in our study did not lead to a significant shift in environmental parameters, the microbial community profiles found in permafrost underwent substantial changes. We suggest that the indigenous microbial community of permafrost cannot respond to the opportunities presented by permafrost thaw (e.g. increased nutrient availability, increased temperatures, and the availability of new niches) as rapidly as the overlying active layer community. Instead of a “blooming” viable permafrost microbial community, active layer taxa appear to invade the disturbed permafrost environment. While taxon membership in disturbed soils appears to be partly of permafrost origin, as indicated by the presence of numerous permafrost indicator taxa in disturbed soils, this signal is not strong enough to affect overall community membership or composition. The disturbed permafrost microbial community converges with the active layer community, is driven by biotic processes, and likely occurs through the invasion of active layer taxa. Cohesion metrics support the hypothesis that biotic filtering was impacting microbial community composition with disturbance.

5.5 Taxonomic profiles resemble other regional studies across the Arctic

Taxonomic profiles in our study were similar to other periglacial soils and differential abundance analysis identified numerous taxa that may be ecologically significant across active layer and permafrost habitats (Malard & Pearce, 2018). Previous studies have indicated that complex hydrocarbons preserved in permafrost are both aerobically and anaerobically degraded into greenhouse gases, including CO₂ and CH₄, as well as volatile fatty acids. While physiology cannot be directly inferred from taxonomy, trends in the membership of these communities are consistent with these previous findings. Classes containing aerobic heterotrophic bacteria (e.g. class Bacilli and most class Actinobacteria), anaerobic heterotrophic bacteria (e.g. class Clostridia), and volatile fatty acid oxidizers (e.g. *Smithella propionica*) were all prominent across Dominion Creek soils and are common globally in active layer and permafrost soils (Jansson & Taş, 2014; Malard & Pearce, 2018; Metje & Frenzel, 2007). Planctomycetes phylum sequences were more abundant in active layer than in permafrost, and disturbance also decreased the relative abundance of Planctomycetes-related sequences in these surface soils, as has been seen in previous studies (Taş et al., 2014). Relatives of spore-forming bacteria, including Clostridia and Bacilli, were more abundant in permafrost soils than in active layer soils. Increased abundance of these putative spore formers is likely due to niche partitioning and survival strategies precluding spore formation (Johnson et al., 2007). The distinct life strategies between permafrost and active layer classes suggest that community functionality may also differ across permafrost, active layer, and disturbed soil communities, similar to previous findings (Hultman et al., 2015).

5.6 Viability and survivability in permafrost

The impact of permafrost thaw on microbial survivability is unknown. We posit that shifts in microbial community structure could be accomplished by growth of active bacteria

and/or death of poorly adapted bacteria. Viability did not differ between the active layer and permafrost soils at Dominion Creek; we propose this lack of difference is due to long-term adaptation to both seasonal thaw in the active layer (Schostag et al., 2015), as well as constant and perennial stressor adaptations in the permafrost horizons (Mackelprang et al., 2017). Our data suggests that abrupt thaw does not dramatically increase microbial cell abundance or viability of microorganisms in permafrost soils. However, previous studies have suggested increases in microbial activity, transcription, and genetic potential with permafrost thaw in the laboratory, and in the field (Coolen et al., 2011; Coolen & Orsi, 2015; Hultman et al., 2015; Mackelprang et al., 2011). Due to these increases in metabolic activity, transcriptional activity, and genetic potential under both rapid and long-term thaw, lower biomass and lower viability with discrete and substantial perturbation were unexpected. Future studies may investigate whether activity changes result from a small but viable community in thawing permafrost which becomes more abundant with thaw.

6 Conclusions

Six weeks of thaw following disturbance was sufficient to shift bacterial community composition, membership, and diversity in disturbed permafrost soils to become more similar to lower active layer soils at a single permafrost-affected site. These communities are driven primarily by the biotic filtering of an invading active layer microbial community in disturbed permafrost communities. Thaw does not increase the viability or abundance of bacteria, suggesting increasing activity, transcription, and genetic potential found in other thawing soils may be due to a more active, yet sparse, microbial community. This study should be expanded to determine if disturbance creates similar microbial community shifts in naturally occurring rapid permafrost thaw zones. Interpretation of activity measurements and characterization of important active microbial taxa may suggest future avenues of research to

develop our understanding of microbial succession in response to permafrost thaw induced by direct anthropogenic disturbance.

645 **Data Availability**

The data for the findings in this manuscript can be found at:

<https://dataview.ncbi.nlm.nih.gov/object/PRJNA999916>.

Author Contributions

Supervision and resources were provided by BL and DF; conceptualization was undertaken
650 by BL, DF, and ASM; methodology was designed and undertaken by ASM and PN; curation
of data, formal analysis, and manuscript writing was performed by PN; reviewing and editing
of the manuscript was performed by BL, DF, and ASM. All authors made substantial
contributions to the publication of this work.

Funding Disclosure

655 The research was made possible by NSERC Discovery Grants (BL and DF) and a NSERC
Northern Research Supplement to DF. A Yukon Scientists and Explorers License was
acquired from the Yukon Government for coring the permafrost samples.

Conflict of interest

The researchers declare no conflict of interest.

660 **Acknowledgements**

We would like to thank Ali Naeimi Nezamabad for designing the sampling location
map, as well as Sasiri Bandara, Casey Buchanan, and Joseph Young for assisting in the
coring of permafrost. We would also like to thank the University of Alberta (UANRA) and

Polar Knowledge Canada (NSTP) for student academic support. Finally, thank you to Dr.
665 Martin Sharp, Dr. Alberto Reyes, and Richard Mah for providing the facilities necessary for
this research. This work was supported by NSERC Discovery Grants to BL and DF, and a
Northern Research Supplement to DF.

References

- 670 Boulou, L., Prevost, M., Barbeau, B., Coallier, J., and Desjardins, R.: LIVE/DEAD BacLight : application of a new rapid staining method for direct enumeration of viable and total bacteria in drinking water, *J Microbiol Methods*, 37, 77-86, [https://doi.org/10.1016/s0167-7012\(99\)00048-2](https://doi.org/10.1016/s0167-7012(99)00048-2), 1999.
- 675 Burn, C., Bartsch, A., Chakraborty, E., Das, S., Frauenfelder, R., Gärtner-Roer, I., Gissnäs, K., Herring, T., Jones, B., Kokelj, S., Langer, M., Lathrop, E., Murton, J., Nielsen, D., Niu, F., Olson, C., O'Neill, H. ., Opfergelt, S., Overduin, P., Schaefer, K., Schuur, E. ., Skierszkan, E., Smith, S., Stuenzi, S., Tank, S., van der Sluijs, J., Vieira, G., Westermann, S., Wolfe, S. and Yarmak, E.: Developments in Permafrost Science and Engineering in Response to Climate Warming in Circumpolar and High Mountain Regions, 2019–2024, *Permafrost and Periglacial Processes*, 36, 167-188. <https://doi.org/10.1002/ppp.2261>, 2025.
- 680 Calmels, F., Gagnon, O., and Allard, M.: A portable earth-drill system for permafrost studies, *Permafrost and Periglacial Processes*, 16, 311-315, <https://doi.org/10.1002/ppp.529>, 2005.
- Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S. M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J. A., Smith, G., and Knight, R.: Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms, *ISME J*, 6, 1621-1624, <https://doi.org/10.1038/ismej.2012.8>, 2012.
- 685 Chang, T., Yi, Y., Jiang, H., Li, R., Lu, P., Liu, L., Wang, L., Wang, L., Zhao, L., Zwieback, S., and Zhao, J.: Unraveling the non-linear relationship between seasonal deformation and permafrost active layer thickness, *npj Climate and Atmospheric Science*, 7, 308, <https://doi.org/10.1038/s41612-024-00866-0>.
- 690 Chen, Y., Romps, D. M., Seeley, J. T., Veraverbeke, S., Riley, W. J., Mekonnen, Z. A., and Randerson, J. T.: Future increases in Arctic lightning and fire risk for permafrost carbon, *Nature Climate Change*, 11, 404+, <https://doi.org/10.1038/s41558-021-01011-y>, 2021.
- Chen, Y. L., Deng, Y., Ding, J. Z., Hu, H. W., Xu, T. L., Li, F., Yang, G. B., and Yang, Y. H.: Distinct microbial communities in the active and permafrost layers on the Tibetan Plateau, *Mol Ecol*, 26, 6608-6620, <https://doi.org/10.1111/mec.14396>, 2017.
- 695 Coolen, M. J. and Orsi, W. D.: The transcriptional response of microbial communities in thawing Alaskan permafrost soils, *Front Microbiol*, 6, 197, <https://doi.org/10.3389/fmicb.2015.00197>, 2015.
- 700 Coolen, M. J., van de Giessen, J., Zhu, E. Y., and Wuchter, C.: Bioavailability of soil organic matter and microbial community dynamics upon permafrost thaw, *Environ Microbiol*, 13, 2299-2314, <https://doi.org/10.1111/j.1462-2920.2011.02489.x>, 2011.
- De Cáceres, M., Sol, D., Lapiedra, O., and Legendre, P.: A framework for estimating niche metrics using the resemblance between qualitative resources, *Oikos*, 120, 1341-1350, <https://doi.org/10.1111/j.1600-0706.2011.19679.x>, 2011.
- 705 Dixon, P.: VEGAN, a package of R functions for community ecology, *Journal of Vegetation Science*, 14, 927-930, [https://doi.org/10.1658/1100-9233\(2003\)014\[0927:Vaporf\]2.0.Co;2](https://doi.org/10.1658/1100-9233(2003)014[0927:Vaporf]2.0.Co;2), 2003.

- 710 Doherty, S., Thurston, A., and Barbato, R.: Active layer and permafrost microbial community
coalescence increases soil activity and diversity in mixed communities compared to
permafrost alone, *Frontiers in Microbiology*, 16, 1579156, <https://doi.org/10.3389/fmicb.2025.1579156>, 2025
- Edgar, R. C.: Search and clustering orders of magnitude faster than BLAST, *Bioinformatics*,
26, 2460-2461, <https://doi.org/10.1093/bioinformatics/btq461>, 2010.
- 715 Edgar, R. C.: SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS
sequences, *bioRxiv*, 074161, <https://doi.org/10.1101/074161>, 2016.
- Edgar, R. C. and Flyvbjerg, H.: Error filtering, pair assembly and error correction for next-
generation sequencing reads, *Bioinformatics*, 31, 3476-3482,
<https://doi.org/10.1093/bioinformatics/btv401>, 2015.
- 720 Edgar, R. C. and Flyvbjerg, H.: Error filtering, pair assembly and error correction for next-
generation sequencing reads, *Bioinformatics*, 31, 3476-3482,
<https://doi.org/10.1093/bioinformatics/btv401>, 2015.
- Elias, S. A.: Conservation issues: Tundra ecosystems, in: Reference Module in Earth Systems
and Environmental Sciences, Elsevier, <https://doi.org/10.1016/b978-0-12-409548-9.09183-1>,
725 2014.
- Emerson, J. B., Roux, S., Brum, J. R., Bolduc, B., Woodcroft, B. J., Jang, H. B., Singleton, C.
M., Solden, L. M., Naas, A. E., Boyd, J. A., Hodgkins, S. B., Wilson, R. M., Trubl, G., Li, C.,
Frolking, S., Pope, P. B., Wrighton, K. C., Crill, P. M., Chanton, J. P., Saleska, S. R., Tyson,
G. W., Rich, V. I., and Sullivan, M. B.: Host-linked soil viral ecology along a permafrost
730 thaw gradient, *Nat Microbiol*, 3, 870-880, <https://doi.org/10.1038/s41564-018-0190-y>, 2018.
- Forbes, B. C., Ebersole, J. J., and Strandberg, B.: Anthropogenic disturbance and patch
dynamics in circumpolar arctic ecosystems, *Conservation biology*, 15, 954-969, 2001.
- Galili, T.: dendextend: an R package for visualizing, adjusting and comparing trees of
hierarchical clustering, *Bioinformatics*, 31, 3718-3720,
735 <https://doi.org/10.1093/bioinformatics/btv428>, 2015.
- Hansen, A. A., Herbert, R. A., Mikkelsen, K., Jensen, L. L., Kristoffersen, T., Tiedje, J. M.,
Lomstein, B. A., and Finster, K. W.: Viability, diversity and composition of the bacterial
community in a high Arctic permafrost soil from Spitsbergen, Northern Norway, *Environ
Microbiol*, 9, 2870-2884, <https://doi.org/10.1111/j.1462-2920.2007.01403.x>, 2007.
- 740 Hayden, H. L., Mele, P. M., Bougoure, D. S., Allan, C. Y., Norng, S., Piceno, Y. M., Brodie,
E. L., DeSantis, T. Z., Andersen, G. L., and Williams, A. L.: Changes in the microbial
community structure of bacteria, archaea and fungi in response to elevated CO₂ and warming
in an Australian native grassland soil, *Environmental microbiology*, 14, 3081-3096,
<https://doi.org/10.1111/j.1462-2920.2012.02855.x>, 2012.
- 745 Hultman, J., Waldrop, M. P., Mackelprang, R., David, M. M., McFarland, J., Blazewicz, S.
J., Harden, J., Turetsky, M. R., McGuire, A. D., Shah, M. B., VerBerkmoes, N. C., Lee, L.
H., Mavrommatis, K., and Jansson, J. K.: Multi-omics of permafrost, active layer and
thermokarst bog soil microbiomes, *Nature*, 521, 208-212,
<https://doi.org/10.1038/nature14238>, 2015.

- 750 Intergovernmental Panel on Climate, C.: *Climate Change 2022 – Impacts, Adaptation and Vulnerability*, Cambridge University Press, Cambridge, <https://doi.org/10.1017/9781009325844>, 2023.
- Janssen, P. H., Yates, P. S., Grinton, B. E., Taylor, P. M., and Sait, M.: Improved culturability of soil bacteria and isolation in pure culture of novel members of the divisions
- 755 Acidobacteria, Actinobacteria, Proteobacteria, and Verrucomicrobia, *Appl Environ Microbiol*, 68, 2391-2396, <https://doi.org/10.1128/AEM.68.5.2391-2396.2002>, 2002.
- Jansson, J. K. and Tas, N.: The microbial ecology of permafrost, *Nat Rev Microbiol*, 12, 414-425, <https://doi.org/10.1038/nrmicro3262>, 2014.
- Johnson, S. S., Hebsgaard, M. B., Christensen, T. R., Mastepanov, M., Nielsen, R., Munch,
- 760 K., Brand, T., Gilbert, M. T., Zuber, M. T., Bunce, M., Ronn, R., Gilichinsky, D., Froese, D., and Willerslev, E.: Ancient bacteria show evidence of DNA repair, *Proc Natl Acad Sci U S A*, 104, 14401-14405, <https://doi.org/10.1073/pnas.0706787104>, 2007.
- Keuper, F., van Bodegom, P. M., Dorrepaal, E., Weedon, J. T., van Hal, J., van Logtestijn, R. S. P., and Aerts, R.: A frozen feast: thawing permafrost increases plant-available nitrogen in
- 765 subarctic peatlands, *Global Change Biology*, 18, 1998-2007, <https://doi.org/10.1111/j.1365-2486.2012.02663.x>, 2012.
- Kokelj, S. V. and Jorgenson, M. T.: *Advances in Thermokarst Research, Permafrost and Periglacial Processes*, 24, 108-119, <https://doi.org/10.1002/ppp.1779>, 2013.
- La Ferla, R., Azzaro, M., Michaud, L., Caruso, G., Lo Giudice, A., Paranhos, R., Cabral, A. S., Conte, A., Cosenza, A., Maimone, G., Papale, M., Rappazzo, A. C., and Guglielmin, M.: Prokaryotic Abundance and Activity in Permafrost of the Northern Victoria Land and Upper
- 770 Victoria Valley (Antarctica), *Microb Ecol*, 74, 402-415, <https://doi.org/10.1007/s00248-017-0955-5>, 2017.
- Lantz, T. C., Kokelj, S. V., Gergel, S. E., and Henryz, G. H. R.: Relative impacts of
- 775 disturbance and temperature: persistent changes in microenvironment and vegetation in retrogressive thaw slumps, *Global Change Biology*, 15, 1664-1675, <https://doi.org/10.1111/j.1365-2486.2009.01917.x>, 2009.
- Lim, C. H. and Jackson, M. L.: Dissolution for total elemental analysis, *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties*, 9, 1-12, 1982.
- 780 Lin, Z. J., Luo, J., and Niu, F. J.: Development of a thermokarst lake and its thermal effects on permafrost over nearly 10 yr in the Beiluhe Basin, Qinghai-Tibet Plateau, *Geosphere*, 12, 632-643, <https://doi.org/10.1130/Ges01194.1>, 2016.
- Mackelprang, R., Burkert, A., Haw, M., Mahendrarajah, T., Conaway, C. H., Douglas, T. A., and Waldrop, M. P.: Microbial survival strategies in ancient permafrost: insights from
- 785 metagenomics, *ISME J*, 11, 2305-2318, <https://doi.org/10.1038/ismej.2017.93>, 2017.
- Mackelprang, R., Waldrop, M. P., DeAngelis, K. M., David, M. M., Chavarria, K. L., Blazewicz, S. J., Rubin, E. M., and Jansson, J. K.: Metagenomic analysis of a permafrost microbial community reveals a rapid response to thaw, *Nature*, 480, 368-371, <https://doi.org/10.1038/nature10576>, 2011.

- 790 Malard, L. A. and Pearce, D. A.: Microbial diversity and biogeography in Arctic soils, *Environ Microbiol Rep*, 10, 611-625, <https://doi.org/10.1111/1758-2229.12680>, 2018.
- Mandal, S., Van Treuren, W., White, R. A., Eggesbo, M., Knight, R., and Peddada, S. D.: Analysis of composition of microbiomes: a novel method for studying microbial composition, *Microb Ecol Health Dis*, 26, 27663, <https://doi.org/10.3402/mehd.v26.27663>, 795 2015.
- Maynard, D. G., Kalra Y. P., and Crumbaugh J. A.: Soil sampling and methods of analysis: Nitrate and exchangeable ammonium nitrogen, Lewis Publishers and CRC Press, 2007.
- McCalley, C. K., Woodcroft, B. J., Hodgkins, S. B., Wehr, R. A., Kim, E. H., Mondav, R., Crill, P. M., Chanton, J. P., Rich, V. I., Tyson, G. W., and Saleska, S. R.: Methane dynamics 800 regulated by microbial community response to permafrost thaw, *Nature*, 514, 478-481, <https://doi.org/10.1038/nature13798>, 2014.
- Metje, M. and Frenzel, P.: Methanogenesis and methanogenic pathways in a peat from subarctic permafrost, *Environ Microbiol*, 9, 954-964, <https://doi.org/10.1111/j.1462-2920.2006.01217.x>, 2007.
- 805 Mondav, R., McCalley, C. K., Hodgkins, S. B., Frolking, S., Saleska, S. R., Rich, V. I., Chanton, J. P., and Crill, P. M.: Microbial network, phylogenetic diversity and community membership in the active layer across a permafrost thaw gradient, *Environ Microbiol*, 19, 3201-3218, <https://doi.org/10.1111/1462-2920.13809>, 2017.
- Monteux, S., Weedon, J. T., Blume-Werry, G., Gavazov, K., Jassey, V. E. J., Johansson, M., 810 Keuper, F., Olid, C., and Dorrepaal, E.: Long-term in situ permafrost thaw effects on bacterial communities and potential aerobic respiration, *ISME J*, 12, 2129-2141, <https://doi.org/10.1038/s41396-018-0176-z>, 2018.
- Monteux, S., Mariën, J., and Krab, E.: Dispersal of bacteria and stimulation of permafrost decomposition by Collembola, *Biogeosciences*, 19, 4089-4105, <https://doi.org/10.5194/bg-19-4089-2022>, 815 2022.
- Muyzer, G., de Waal, E. C., and Uitterlinden, A. G.: Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA, *Appl Environ Microbiol*, 59, 695-700, <https://doi.org/10.1128/aem.59.3.695-700.1993>, 1993.
- 820 Nazir, A., Farooq, B., Farooq, M., Anjum, S., Farooq, U., Shameem, N., Egamberdieva, D., and Fazeli-Nasab, B.: Chapter 2 - soil microbial diversity and functions, *Microbiome drivers of ecosystem function, Annual Reviews: Academic Press*, 17-29. doi: 10.1016/B978-0-443-19121-3.00011-9, 2024.
- Oelbermann, M., English, M., and Schiff, S. L.: Evaluating carbon dynamics and microbial 825 activity in arctic soils under warmer temperatures, *Canadian Journal of Soil Science*, 88, 31-44, <https://doi.org/10.4141/Cjss07060>, 2008.
- Palmer, K., Biasi, C., and Horn, M. A.: Contrasting denitrifier communities relate to contrasting N₂O emission patterns from acidic peat soils in arctic tundra, *ISME J*, 6, 1058-1077, <https://doi.org/10.1038/ismej.2011.172>, 2012.

- 830 Reyes, F. R. and Lougheed, V. L.: Rapid nutrient release from permafrost thaw in arctic aquatic ecosystems, *Arctic Antarctic and Alpine Research*, 47, 35-48, <https://doi.org/10.1657/Aaar0013-099>, 2015.
- Richter-Menge, J. A.: Arctic report card 2010, DIANE Publishing, 2011.
- 835 Rousk, J., Frey, S. D., and Baath, E.: Temperature adaptation of bacterial communities in experimentally warmed forest soils, *Glob Chang Biol*, 18, 3252-3258, <https://doi.org/10.1111/j.1365-2486.2012.02764.x>, 2012.
- Saidi-Mehrabad, A., Neuberger, P., Cavaco, M., Froese, D., and Lanoil, B.: Optimization of subsampling, decontamination, and DNA extraction of difficult peat and silt permafrost samples, *Sci Rep*, 10, 14295, <https://doi.org/10.1038/s41598-020-71234-0>, 2020.
- 840 Saidi-Mehrabad, A., Neuberger, P., Hajihosseini, M., Froese, D., and Lanoil, B. D.: Permafrost Microbial Community Structure Changes Across the Pleistocene-Holocene Boundary, *Frontiers in Environmental Science*, 8, 548929, ARTN 133, <https://doi.org/10.3389/fenvs.2020.00133>, 2020.
- 845 Scheel, M., Zervas, A., Jacobsen, C.S., Christensen, T.R.: Microbial Community Changes in 26,500-Year-Old Thawing Permafrost, *Frontiers in Microbiology*, 13, 787146, <https://doi.org/10.3389/fmicb.2022.787146>, 2022.
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski, R. A., Oakley, B. B., Parks, D. H., Robinson, C. J., Sahl, J. W., Stres, B., Thallinger, G. G., Van Horn, D. J., and Weber, C. F.: Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities, *Appl Environ Microbiol*, 75, 7537-7541, <https://doi.org/10.1128/AEM.01541-09>, 2009.
- 850 Schostag, M., Stibal, M., Jacobsen, C. S., Baelum, J., Tas, N., Elberling, B., Jansson, J. K., Semenchuk, P., and Prieme, A.: Distinct summer and winter bacterial communities in the active layer of Svalbard permafrost revealed by DNA- and RNA-based analyses, *Front Microbiol*, 6, 399, <https://doi.org/10.3389/fmicb.2015.00399>, 2015.
- 860 Schuur, E. A. G., Bracho, R., Celis, G., Belshe, E. F., Ebert, C., Ledman, J., Mauritz, M., Pegoraro, E. F., Plaza, C., Rodenhizer, H., Romanovsky, V., Schädel, C., Schirokauer, D., Taylor, M., Vogel, J. G., and Webb, E. E.: Tundra Underlain By Thawing Permafrost Persistently Emits Carbon to the Atmosphere Over 15 Years of Measurements, *Journal of Geophysical Research-Biogeosciences*, 126, e2020JG006044, ARTN e2020JG006044, <https://doi.org/10.1029/2020JG006044>, 2021.
- 865 Schuur, E. A., McGuire, A. D., Schadel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171-179, <https://doi.org/10.1038/nature14338>, 2015.
- 870 Singleton, C. M., McCalley, C. K., Woodcroft, B. J., Boyd, J. A., Evans, P. N., Hodgkins, S. B., Chanton, J. P., Frolking, S., Crill, P. M., Saleska, S. R., Rich, V. I., and Tyson, G. W.: Methanotrophy across a natural permafrost thaw environment, *ISME J*, 12, 2544-2558, <https://doi.org/10.1038/s41396-018-0065-5>, 2018.

- Skoog, D. A., Holler, F. J., Crouch, S. R.: Principles of Instrumental Analysis, Chapter 10: Atomic Emission Spectrometry, Thomson Brooks/Cole, Belmont, 2007.
- 875 Sparks, D. L., Page, A. L., Helmke, P. A., and Loeppert, R. H.: Methods of soil analysis, part 3: Chemical methods, John Wiley & Sons, 2020.
- Tas, N., Prestat, E., McFarland, J. W., Wickland, K. P., Knight, R., Berhe, A. A., Jorgenson, T., Waldrop, M. P., and Jansson, J. K.: Impact of fire on active layer and permafrost microbial communities and metagenomes in an upland Alaskan boreal forest, *ISME J*, 8, 1904-1919, <https://doi.org/10.1038/ismej.2014.36>, 2014.
- 880 Tas, N., Prestat, E., Wang, S., Wu, Y., Ulrich, C., Kneafsey, T., Tringe, S. G., Torn, M. S., Hubbard, S. S., and Jansson, J. K.: Landscape topography structures the soil microbiome in arctic polygonal tundra, *Nat Commun*, 9, 777, <https://doi.org/10.1038/s41467-018-03089-z>, 2018.
- 885 Tripathi, B. M., Kim, M., Kim, Y., Byun, E., Yang, J. W., Ahn, J., and Lee, Y. K.: Variations in bacterial and archaeal communities along depth profiles of Alaskan soil cores, *Sci Rep*, 8, 504, <https://doi.org/10.1038/s41598-017-18777-x>, 2018.
- 890 Turetsky, M. R., Abbott, B. W., Jones, M. C., Anthony, K. W., Olefeldt, D., Schuur, E. A. G., Grosse, G., Kuhry, P., Hugelius, G., Koven, C., Lawrence, D. M., Gibson, C., Sannel, A. B. K., and McGuire, A. D.: Carbon release through abrupt permafrost thaw, *Nature Geoscience*, 13, 138-143, <https://doi.org/10.1038/s41561-019-0526-0>, 2020.
- Van Huissteden, J. and Dolman, A.: Soil carbon in the Arctic and the permafrost carbon feedback, *Current Opinion in Environmental Sustainability*, 4, 545-551, <https://doi.org/10.1016/j.cosust.2012.09.008>, 2012.
- 895 Woodcroft, B. J., Singleton, C. M., Boyd, J. A., Evans, P. N., Emerson, J. B., Zayed, A. A. F., Hoelzle, R. D., Lamberton, T. O., McCalley, C. K., Hodgkins, S. B., Wilson, R. M., Purvine, S. O., Nicora, C. D., Li, C., Frolking, S., Chanton, J. P., Crill, P. M., Saleska, S. R., Rich, V. I., and Tyson, G. W.: Genome-centric view of carbon processing in thawing permafrost, *Nature*, 560, 49-54, <https://doi.org/10.1038/s41586-018-0338-1>, 2018.
- 900 Wu, X. D., Zhao, L., Liu, G. M., Xu, H. Y., Zhang, X. L., and Ding, Y. J.: Effects of permafrost thaw-subsidence on soil bacterial communities in the southern Qinghai-Tibetan Plateau, *Applied Soil Ecology*, 128, 81-88, <https://doi.org/10.1016/j.apsoil.2018.04.007>, 2018.