



Vertical and seasonal variations in airborne endotoxins in a coastal megacity of North China: insights from 3-hydroxy fatty acids

Wenxin Zhang¹, Wei Hu^{1, 2*}, Mutong Niu¹, Quanfei Zhu³, Na An³, Qiang Zhang¹, Rui Jin¹, Xiaoli Fu¹,
Jian Hao⁴, Jianbo Yang⁴, Jingle Liu⁵, Jing Shi⁵, Suqin Han⁶, Junjun Deng¹, Libin Wu¹, Yuqi Feng³,
5 Kimitaka Kawamura⁷, Pingqing Fu^{1, 8*}

¹Institute of Surface-Earth System Science, School of Earth System Science, Tianjin University, Tianjin 300072, China

²Tianjin Bohai Rim Coastal Earth Critical Zone National Observation and Research Station, Tianjin University, Tianjin 300072, China

³School of Bioengineering and Health, Wuhan Textile University, Wuhan 430072, China

10 ⁴Tianjin Institute of Meteorological Science, Tianjin 300074, China

⁵Tianjin Meteorological Radar Research and Test Centre, Tianjin 300061, China

⁶Tianjin Environmental Meteorological Center, Tianjin 300074, China

⁷Chubu Institute for Advanced Studies, Chubu University, Kasugai, Aichi 487-8501, Japan

15 ⁸Tianjin Key Laboratory of Earth Critical Zone Science and Sustainable Development in Bohai Rim, Tianjin University, Tianjin 300072, China

Correspondence to: Wei Hu (huwei@tju.edu.cn), Pingqing Fu (fupingqing@tju.edu.cn)

Abstract. Endotoxins, integral components of Gram-negative bacteria, are released into atmosphere during bacterial fragmentation and pose health risks. This study investigated 3-hydroxy fatty acids (3-OH-FAs, C₈ – C₁₈) in inhalable particles (PM₁₀) from urban Tianjin, a coastal megacity in northern China, to estimate endotoxin levels utilizing UPLC-MS. Results
20 revealed seasonal and altitudinal variations in 3-OH-FAs and endotoxin levels. Total endotoxin concentrations averaged 21.5 ng m⁻³ at near ground (2 m) and 16.1 ng m⁻³ at a higher altitude (220 m), corresponding to total 3-OH-FAs (C₁₀ – C₁₈) concentrations of 2.75 ng m⁻³ and 2.01 ng m⁻³, respectively. Maximum endotoxin level (26.5 ng m⁻³) occurred near ground during winter, attributed to enhanced near-surface emissions. Bioactive endotoxins peaked at 12.4 ng m⁻³ near ground in winter, exceeding the exposure threshold, while averaging 8.12 ng m⁻³ in other seasons. Short-chain 3-OH-FAs (C₈ – C₁₃) exhibited
25 significant correlations with meteorological factors (e.g., temperature, humidity, and wind speed) at both altitudes, indicating regulation through microbial growth dynamics and photochemical processes. Long-chain homologues (C₁₄ – C₁₈) were affected by both meteorological conditions and particulate pollutants (e.g., OC, WSOC, K⁺, Ca²⁺), reflecting mixed influences from natural sources (e.g., soils) and anthropogenic activities (e.g., combustion). These findings advance understanding of
30 spatiotemporal variations in airborne endotoxins within complex urban environments, providing critical data for assessing health risks associated with particulate pollution and informing urban air quality management strategies.



1 Introduction

Inhalable particles with aerodynamic diameters less than 10 μm (PM_{10}) may originate from both natural and anthropogenic sources (Yin et al., 2022; Xue et al., 2024), and could further result in various respiratory and cardiovascular diseases (Makkonen et al., 2010; Mukherjee and Agrawal, 2017). Among the biological components of PM_{10} , endotoxins are of extreme importance due to strong chronic health effects (Mahapatra et al., 2018; Mueller-Anneling et al., 2004).

Endotoxin, also known as lipopolysaccharide (LPS), is an integral component of the outer membrane of Gram-negative bacteria (GNB) (Rylander, 2002). An endotoxin contains three regions, consisting of a core polysaccharide, a long-chain polysaccharide, and a nonpolar lipid named lipid A, with lipid A being the most associated with toxicity (Schneier et al., 2020; Rylander, 2002; Spaan et al., 2006). As a potent inflammagen, an endotoxin can cause fever, shaking chills, septic shock, toxic pneumonitis and respiratory symptoms, through inhalation, dermal and/or eye contact or ingestion (Rylander, 2002; Farokhi et al., 2018). The health risks of endotoxins to humans have been investigated (Farokhi et al., 2018; Laboha et al., 2023; Liebers et al., 2008). Their harmful effects depend on both the composition of the inhaled particles and the degree and duration of exposure (Liebers et al., 2020; Lundin and Checkoway, 2009). Data about the quantity and quality of LPS in environmental samples are critical to understanding the biological effects of environmental endotoxin exposure (Park et al., 2004).

Previous studies have investigated endotoxin levels in various environments, including occupational settings (Heederik and Douwes, 1997; Liebers et al., 2020; Liebers et al., 2006), indoor environments (Amin et al., 2023; Fonseca Gabriel et al., 2021; Phiri et al., 2023), and ambient aerosols and dust (Cheng et al., 2012; Hwang and Park, 2019; Lang-Yona et al., 2014; Hines et al., 2003; Park et al., 2000). Airborne endotoxin levels exhibited spatial and temporal variations, influenced by geographical location (Moretti et al., 2018), season (Makkonen et al., 2010; Hwang and Park, 2019), emission sources (Mueller-Anneling et al., 2004; Mahapatra et al., 2018), meteorological conditions (Guan et al., 2014; Rolph et al., 2018), and pollution levels (Guo et al., 2018). These variations might reveal the potential sources, atmospheric processes, and the survival mechanisms of airborne microbes (Hu et al., 2020a). However, the factors influencing endotoxin levels are not fully understood (Hwang and Park, 2019; Mahapatra et al., 2018). To gain further clarity, outdoor sampling coupled with simultaneous monitoring of meteorological and air quality parameters is essential (Amin et al., 2023). Additionally, the sources of endotoxins in ambient environment, whether from long-range transport or local emissions, have yet to be definitively identified.

Previous studies on endotoxins have primarily focused on occupational exposure and the health effects of excessive endotoxin exposure (Liebers et al., 2006; Amin et al., 2023). However, research on endotoxin levels in the ambient atmosphere remains limited. The vertical distribution, spatiotemporal patterns of airborne endotoxins in the urban boundary layer, and their possible influencing factors have yet to be reported.

Multiple factors including sample types (personal or stationary), sampling method (active or passive), extraction procedure, and storage conditions, can affect the determination of endotoxins (Liebers et al., 2020). One mainstream detection method of



endotoxins is the *Limulus* amoebocyte lysate (LAL) test (Rylander, 2002). This enzyme activation-based method provides direct information on bioactive endotoxins. Despite its sensitivity, the LAL test may still underestimate endotoxin levels, because many endotoxins, including bioactive ones, may remain concealed within the intact bacterial structure, making them unavailable to react with the *Limulus* enzymes (Mattsby-Baltzer et al., 1991). False-negative results may occur when endotoxins are masked by buffer constituents, product formulation, cell culture medium compositions, and surfactants (Schwarz et al., 2017; Schneier et al., 2020). Whilst, false-positive results can be caused by β -1,3-glucan, which might as well trigger the enzymatic coagulation of the blood (Cheng et al., 2012; Uhlig et al., 2016).

The alternative approach most commonly applied to detect endotoxins in environmental samples is to measure 3-hydroxy fatty acids (3-OH-FAs) with carbon chain lengths from 10 to 18 as endotoxin biomarkers utilizing chemical analytical methods such as gas chromatography-mass spectrometry (GC-MS) (Bikkina et al., 2021a; Cheng et al., 2012). In contrast to the LAL test, 3-OH-FAs measured by GC-MS do not reflect the biologically active endotoxin, but rather total concentration (Liebers et al., 2008). In addition, the sample pretreatment in GC-MS with strong alkaline hydrolysis may result in 3-OH-FAs turning into 3-unsaturated fatty acids while eliminating water during the derivatization process (trifluoroacetamide) (Binding et al., 2004; Wollenweber and Rietschel, 1990). Liquid chromatography-mass spectrometry (LC-MS) with its high sensitivity and selectivity, and broad adaptability, is regarded as a powerful analytical tool for small molecules (Zhang et al., 2019; Paba et al., 2019). Recently, ultra-high-performance liquid chromatography mass spectrometry (UHPLC-MS) combined with isotope labeling has been used to effectively separate, identify and quantify positional isomers of hydroxy fatty acids, demonstrating exceptional accuracy and precision in complex environmental matrices including aerosols (Niu et al., 2024; Zhu et al., 2020).

The coastal zone is a transition area between the ocean and the land, which is affected by natural continental processes (e.g., dust storms) and marine emissions, as well as intense human activities due to a large population. In addition, sea-land breeze is one of the most common circulations over coastal areas and has a great impact on the local meteorology and atmospheric environment (Xiao et al., 2023). These interactions make the atmospheric environment of coastal zones more complex, providing unique conditions for investigating the effects of local emissions and regional or long-distance transport on atmospheric environment. Meteorological tower-based studies complement ground-based observations (Yang et al., 2023), offering new insights into the formation, transformation, and transport of organic matter under different atmospheric environments (Lei et al., 2021). Comprehensive vertical measurements of the physical and chemical properties of aerosol particles in the lower boundary layer have been conducted to evaluate the roles of local emissions and regional and long-distance transport in air pollution (Fan et al., 2022; Li et al., 2022). However, the levels, sources, and health impacts of airborne endotoxins in coastal zones remain largely unknown.

Herein, diurnal PM₁₀ samples were collected at two altitudes using a 255-m meteorological tower in urban Tianjin, the largest coastal megacity in North China, enabling the investigation of the vertical distribution of airborne endotoxins. The UHPLC-



MS system, combined with isotope labeling, was applied for accurate determination of 3-OH-FAs ($C_8 - C_{18}$) in PM_{10} . The
95 amounts of endotoxins and GNB dry mass in PM_{10} were estimated using the concentrations of 3-OH-FAs ($C_{10} - C_{18}$) according
to previous studies (Bikkina et al., 2021a; Tyagi et al., 2015b; Tyagi et al., 2016). This study will provide: (1) the quantification
and distribution patterns of the bacterial marker 3-OH-FAs in ambient PM_{10} within the urban boundary layer in Tianjin; (2)
insights into the vertical, seasonal and diurnal variations in airborne endotoxins, along with estimates of the bioactive portion
of 3-OH-FAs and the mass loading of GNB for assessing the potential allergenic impact of airborne endotoxin levels, and (3)
100 the preliminary identification of influencing factors of endotoxin levels, in combination with key environmental parameters,
including meteorological and air quality data.

2 Materials and methods

2.1 Samples collection

Tianjin is a coastal megacity located in northeast part of the North China Plain (NCP), bordering the Bohai Sea in the east,
105 leaning against the Yanshan Mountain in the north (Fig. S1). It is 120 kilometers from Beijing, the capital of China, in the
northwest. The sampling was conducted by utilizing the 255-m meteorological tower at the Tianjin Atmospheric Boundary
Layer Observatory of the China Meteorological Administration, located at the southern area of urban Tianjin. Two
automatic high-volume aerosol samplers (Digitel DHA-80) were set at two floors in the meteorological tower, i.e., the near-
ground (2 m) and high altitude (220 m), respectively. PM_{10} samples were collected on pre-combusted (450°C for 6 hours)
110 quartz fiber filter (PALL, 150mm) during the autumn (18 – 24 September) and winter (8 – 14 December) of 2020, and the
spring (26 – 31 May) and summer (2 – 8 August) of 2021. After collection, the samples were stored at -20°C in a freezer until
analysis.

2.2 Samples pretreatment

The extraction and detection methods of 3-OH-FAs are available in our previous study (Niu et al., 2024). Briefly, aliquots of
115 aerosol samples (Φ 28 mm) were cut and placed in a centrifuge tube, followed by adding 4 mL of EtOAc, 1 min of vortex, and
15 min of sonicating at room temperature. Then, 4 mL of Milli-Q water was added, and the resulting solution was vortexed for
remixing. Centrifugation was performed for 5 min at 5000 rpm, and the supernatant was extracted afterward, and then dried
under nitrogen. Subsequently, 100 μ L of ACN was added to the dried centrifuge tube and vortexed for 2 min, then derivatization
reagents (10 μ L of 20 μ mol/mL CMPI, 20 μ L of 20 μ mol/mL TEA, and 20 μ L of 20 μ mol/mL DMED) were added and vortexed
120 for 2 min, and the mixture was incubated for 1 h at 40°C and 1500 rpm, then dried under nitrogen. Milli-Q water and internal
standards (*d*₇-DMED-labeled hydroxy fatty acid standard) (v/v, 1/9) were added to each dried centrifuge tube, vortexed to
redissolve, and then centrifuged for 5 min at 4°C and 12,000 rpm to extract the supernatant. Details of the chemicals, reagents



and method validation are described in Text S1 in the Supporting Information.

2.3 UHPLC-MS analysis

125 The UHPLC-MS system comprised an ACQUITY UPLC I-Class LC (Waters, Milford, MA, USA) coupled with an AB SCIEX
6500+ triple quadrupole MS (AB SCIEX, Framingham, MA, USA). Separation of the target compounds was achieved using
ACQUITY HPLC HSS T3 column (1.8 μm , 2.1 mm \times 50 mm, Thermo Fisher Scientific, CA, USA). The mobile phases
consisted of solvent A (0.1% formic acid in water) and solvent B (ACN), delivered at a flow rate of 0.4 mL/min. The gradient
program was set as follows: 0-3 min at 5% B, 3-15 min from 22% to 60% B, 15-17 min at 60% B, 17-24 min at 95% B, 24-
130 40 min at 5% B. Mass spectrometric analysis was performed in positive ion mode using electrospray ionization (ESI) and
operated in multiple reaction monitoring (MRM) mode. The ESI conditions were set as follows: curtain gas, 35 L min⁻¹;
collision gas, 8 L min⁻¹; ion spray voltage, 5500 V; temperature, 500 °C; ion source gas 1, 50 L min⁻¹; ion source gas 2, 45 L
min⁻¹. Data acquisition and quantification were conducted using SCIEX OS version 2.0.1 software. The mass concentrations
of 3-OH-FAs (C₈ – C₁₈) were quantified using internal standard method, employing *d*₄-DMED-labeled 3-OH-FA standards as
135 internal standards. The detailed analytical procedure is described in (Niu et al., 2024).

2.4 Endotoxin and GNB dry mass estimation

The amounts of endotoxins and GNB dry mass were estimated using the following formulas:

$$\text{Endotoxin}(ng\ m^{-3}) = \sum[3 - OH - FAs (C_{10} - C_{18})(nmol\ m^{-3}) \div 4] \times 8000 (g\ mol^{-1}) \quad (1)$$

140 where 4 means each lipid A carries 4 mol 3-OH-FAs (C₁₀ – C₁₈), while the multiplication factor 8000 represents the average
molecular weight of endotoxin (Laitinen et al., 2001).

$$\text{GNB dry mass}(mg\ m^{-3}) = \sum[3 - OH - FAs (C_{10} - C_{18})(nmol\ m^{-3})] \div 15(nmol\ mg^{-1}) \quad (2)$$

where 15 refers to the biomarker-to-microbial mass conversion factor of 15 nmol of 3-OH-FAs per mg dry cell weight (Lee et
al., 2004; Balkwill et al., 1988).

2.5 Other measurements

145 The concentrations of organic carbon (OC) and elemental carbon (EC) were determined using a Thermal/Optical Carbon
Analyzer (Model RT-4, Sunset Laboratory Inc., USA). Analytical errors were controlled within $\pm 10\%$ via a duplicate analysis
of each filter. The concentrations of water-soluble organic carbon (WSOC) were measured using Total Organic Carbon (TOC)
Analyzer (Shimadzu 5000A, JAPAN). Water-soluble inorganic anions and cations (Cl⁻, SO₄²⁻, NO₃⁻, Na⁺, NH₄⁺, K⁺, Mg²⁺,
Ca²⁺) were analyzed using an Ion Chromatography system (Thermo, USA). Gradient meteorological parameters were
150 continuously recorded by automatic weather stations installed on the 255 m high meteorological tower, while both gaseous
and particulate pollutants were collected simultaneously at the ground platform.



2.6 Concentration-weighted trajectory (CWT) analysis

Concentration-weighted trajectories (CWTs) were calculated per hour, and starting height of 220 m, based on the 3-day (72 h) backward trajectories of air masses with the HYSPLIT model in conjunction with the measured concentrations of 3-OH-FAs as follows:

$$CWT_{ij} = \frac{\sum_l C_l n_{ijl}}{\sum_l n_{ijl}} \quad (3)$$

where l is the number of the trajectory; C_l is the average concentration of 3-OH-FAs, and n_{ijl} is the number of trajectory endpoints that lie in the grid cell (i, j) . The CWT value indicates the potential distant sources impacting the sampling site. The analyses were performed for $1^\circ \times 1^\circ$ grids covering the area encompassed by $20\text{--}80^\circ\text{N}$ and $60\text{--}150^\circ\text{E}$. A weight function (W_{ij}) in Eq. (4) was applied in the CWT analysis to increase statistical stability, and N_{ij} is the number of trajectory endpoints that lie in the grid cell (i, j) .

$$W_{ij} = \begin{cases} 0.1, & N_{ij} < 2 \\ 0.4, & 2 \leq N_{ij} < 4 \\ 0.7, & 4 \leq N_{ij} < 8 \\ 1.0, & \text{elsewhere} \end{cases} \quad (4)$$

2.7 Statistical analysis

Spearman rank analysis was conducted to evaluate the correlation between airborne endotoxins and air pollutants. Data were analyzed using unpaired t-tests, two-tailed for single comparisons. Graphs were generated using RStudio software version 2023.09.1+494, R version 4.3.2 and Origin 2021. The distribution characteristics of 3-OH-FAs and airborne endotoxins were described in range (minimum-maximum) and arithmetic mean.

3 Results and discussion

3.1 Molecular distribution of 3-hydroxy fatty acids

The molecular distribution of 3-OH-FAs with carbon numbers from 8 to 18 ($C_8 - C_{18}$) is illustrated in Figure 1. In terms of diurnal variations, the mass concentrations of most 3-OH-FAs were higher at night than during the daytime (Fig. 1a). This feature, observed predominantly at ground level may be partly due to the lower boundary layer height at night, which contributes to the accumulation of pollutants (Li et al., 2017; Qiu et al., 2019).




Figure 1. The vertical, diurnal and seasonal variations in molecular distribution of 3-OH-FAs (C₈ – C₁₈). (a-b) The mean values (indicated by points) and standard deviations (error bars) of 3-OH-FAs concentrations. (c-d) The differences in 3-OH-FAs concentrations between 2 m and 220 m. The short lines indicate the mean values of the differences.

180

The molecular distribution of 3-OH-FAs also showed apparent seasonal variations (Fig. 1b). Even-carbon homologues (C₁₆, C₁₈) were predominant in all seasons, especially in winter, with average mass concentrations of 1335 pg m⁻³ (C₁₆) and 855.4 pg m⁻³ (C₁₈) at near ground, 882.6 pg m⁻³ (C₁₆) and 467.7 pg m⁻³ (C₁₈) at 220 m. C₁₁, C₁₃, and C₁₅ 3-OH-FAs were the predominant odd-carbon 3-OH-FAs, particularly in summer, with average mass concentrations of 367.4 pg m⁻³ (C₁₁), 412.7 pg m⁻³ (C₁₃), and 405.8 pg m⁻³ (C₁₅) at near ground, respectively. In terms of 3-OH-FAs with different carbon numbers, most 3-OH-FAs were more abundant at near ground, suggesting that these homologues were mainly influenced by near-surface emissions. In contrast, C₉ at 220 m remained higher than at 2 m in autumn and summer, regardless of diurnal changes (Fig. 1c, d). Other short-chain homologues (C₈, C₁₀, C₁₁) also presented higher concentrations at 220 m during summer, indicating that they were possibly contributed by regional transport or photochemical oxidation processes (Lei et al., 2021).

190

The result of concentration-weighted trajectory (CWT) analysis revealed that the mass concentration of 3-OH-FAs at 220 m was influenced by marine sources to varying extents in spring, autumn, and summer (Fig. 2). During autumn and winter, the major source areas were the Beijing-Tianjin-Hebei region, with transport pathways originating from the northwest of Tianjin, including Mongolia, Inner Mongolia and Beijing, before reaching the sampling site. In summer, predominant origin areas included Beijing-Tianjin-Hebei region, Heilongjiang province, and the Bohai Sea in the northeast. Photochemical oxidation of organic matter is prone to occur in summer due to high temperatures and intense solar radiation. The elevated concentrations

195



of short-chain 3-OH-FAs ($C_8 - C_{11}$) at 220 m may result from secondary transformation processes during long-range transport or photochemical oxidation of the local emissions. Bikkin et al. (2019) reported that photochemical oxidation of marine organic matter likely accounts for the predominance of odd-carbon and short-chain 3-OH-FAs in sea-spray aerosols.

200




Figure 2. Seasonal concentration-weighted trajectory (CWT) maps for 3-OH-FAs measured at the Atmospheric Boundary Layer Observatory of Tianjin. This analysis is based on air mass backward trajectories during observation periods.

205

3.2 Spatial-temporal patterns of airborne endotoxin levels

Total mass concentrations of endotoxins in PM_{10} from urban Tianjin during 2020-2021 were estimated based on the measured 3-OH-FAs ($C_{10} - C_{18}$) (Table 1, Fig. 3a). There was no obvious diurnal difference in total endotoxin levels (Fig. 3a). The endotoxin level was significantly higher ($p < 0.05$) at 2 m than at 220 m during each season (Fig. 4a), indicating that endotoxin emissions were mainly dominated by near-ground sources. It was suggested that the majority of airborne endotoxins may originate from soils (Brooks et al., 2006), as aliphatic 3-OH-FAs ($< C_{20}$) originate from soil microorganisms (Zelles, 1999; Bikkin et al., 2021a). 3-OH-FAs are also detected in marine ultrafiltered dissolved organic matter (Wakeham et al., 2003), groundwater sediments and estuarine sediments (Parker et al., 1982). Marine source could also be a dominant contributor to elevated endotoxin levels in coastal areas. Lang-Yona et al. (2014) observed high endotoxins content correlated with cyanobacteria at a coastal site of the eastern Mediterranean Sea, assumed that higher wind speeds with low pressure system led to increased sea spray and the consequent primary aerosol emission. Therefore, the endotoxins at 220 m in Tianjin might

215



be originated either from the vertical transport of ground emissions or long-range transport from marine areas.

In the near-ground (2 m) air, the average endotoxin mass concentrations were the highest (26.5 ng m⁻³) in winter, followed by 23.7 ng m⁻³ in summer, 20.5 ng m⁻³ in fall, and the lowest 14.0 ng m⁻³ in spring (Table 1, Fig. 4a). While at higher altitude (220 m), the average endotoxin level was the highest in summer (18.3 ng m⁻³), followed by winter (16.8 ng m⁻³) and fall (12.8 ng m⁻³). The average endotoxin ratio of 220 m/2 m during summer was 0.78, higher than those during autumn (0.65) and winter (0.67), suggesting an increased contribution of possible regional transport or photochemical oxidation processes during summer. Most airborne bacteria could not survive during long-range transport due to ultraviolet radiation, which further lead to photodegradation of proteinaceous matter and significant losses of microbial viability (Hu et al., 2021; Yin et al., 2021).

LPS is then released into the air during cell death (Petsch and Anspach, 2000), cause the rising in higher endotoxin level. Moreover, the ratio of 220 m/2 m was higher during the daytime (0.77) than nighttime (0.64). The average endotoxin mass concentration was slightly higher at night than during the daytime in spring, summer and autumn, but lower at night during winter (Fig. 4a). The increasing concentration of certain bacteria at night may also attribute to the diurnal changes in endotoxins. Hu et al. (2020b) observed richer operational taxonomic units per PM_{2.5} sample in Hangzhou at night during spring, with higher relative abundance of *Thermomicrobia*, a Gram-negative bacterial class, generally distributed away from anthropogenic regions. Abdel Hameed et al. (2009) found the diurnal variations in number concentration of airborne bacteria peaked in the evening at Helwan, Egypt.

Table 1. Seasonal variations in airborne endotoxin concentrations in PM₁₀ from urban Tianjin at different attitudes during 2020-2021.

Sampling site	Group	Endotoxin concentration (ng m ⁻³)			
		Average	Standard deviation	Range	
Meteorological Tower (220 m)	Season	Autumn (n=12)	12.8	2.93	7.12–18.2
		Winter (n=13)	16.8	8.67	4.48–33.5
		Spring (n=0)	/	/	/
		Summer (n=13)	18.3	5.28	12.1–32.7
	Day/Night	Day (n=18)	16.8	6.57	4.5–33.5
		Night (n=20)	15.5	6.52	4.86–32.7
	Total	SUM (n=38)	16.1	6.57	4.48–33.5
Near-ground (2 m)	Season	Autumn (n=12)	20.5	5.15	13.9–30.8
		Winter (n=13)	26.5	13.0	5.02–42.2
		Spring (n=11)	14.0	6.61	6.12–25.6
		Summer (n=13)	23.6	6.13	16.1–39.0
	Day/Night	Day (n=23)	19.7	9.77	5.02–41.8
		Night (n=26)	23.1	9.13	6.30–42.2
	Total	SUM (n=49)	21.5	9.59	5.02–42.2




Figure 3. Time series of concentrations of total endotoxins (a), particulate matter (b), and gaseous pollutants (c, d) during the sampling periods.

240 The relative contributions of different carbon-numbered 3-OH-FAs to total endotoxin mass concentrations in PM₁₀ of Tianjin presented seasonal and altitudinal variations (Fig. 4b). Overall, the even-carbon long-chain C₁₆ and C₁₈ 3-OH-FAs made up the largest proportion in all seasons, with higher relative contributions (46.5%) to total endotoxins at near-ground level than at 220 m (39.7%), suggesting their local emissions. The relative contribution of the even-carbon C₁₆ and C₁₈ homologues were the highest in winter (56.0% at near ground and 53.3% at 220 m), followed by autumn (49.1%, 38.6%), and the lowest in

245 summer (36.7%, 27.0%). This finding is consistent with those obtained in suburban Tokyo (Bikkina et al., 2021a), Chichijima island (Bikkina et al., 2021b), and Jeju island (Tyagi et al., 2017), in which 3-OH-FAs were also characterized by a strong even-carbon predominance. Hydroxy fatty acids exhibiting a strong even-carbon preference are usually derived from biological pathways (Bikkin et al., 2019) or indicative of biogenic sources (Tyagi et al., 2015a), such as the lipid fractions of microorganisms (bacteria, algae, fungi, etc.) and vascular plant surface waxes (Kawamura et al., 2003; Simoneit et al., 2004).

250




Figure 4. Concentrations of total endotoxins and relative contributions of 3-OH-FAs in urban aerosols from Tianjin.

(a) The vertical, diurnal and seasonal variations in mass concentrations of endotoxins. (b-c) Seasonal (b) and diurnal (c) variations in the relative contributions of 3-OH-FAs with different carbon number to total endotoxins at different altitudes.

255

However, in this study, the odd-carbon homologues (C₁₁, C₁₃ and C₁₅) accounts for 36.7% and 41.9% of total endotoxins at near ground and at 220 m, respectively. Their relative contribution was maximum in summer (45.0% at near ground and 51.6% at 220 m), followed by autumn (34.5% at near ground and 41.4% at 220 m), and the lowest in winter (30.2% at near ground and 32.5% at 220 m). Compared with remote islands dominated by natural emissions such as microorganisms and vegetation (Bikkina et al., 2021b; Tyagi et al., 2015a; Tyagi et al., 2017), the urban aerosols in Tianjin are largely affected by both anthropogenic emissions and secondary formation (Li et al., 2022). These odd-carbon 3-OH-FAs (C₁₁, C₁₃ and C₁₅) may originate from various sources, such as anthropogenic activities (e.g., fossil fuel combustion, biomass emissions), or from the photochemical oxidation of long-chain fatty acids or hydroxy fatty acids during atmospheric transport (Bikkin et al., 2019).

265

In addition, long-chain C₁₆, C₁₈ and odd-carbon homologues (C₁₁, C₁₃ and C₁₅) 3-OH-FAs exhibited similar variation trend at the two heights. Long-chain C₁₆, C₁₈ 3-OH-FAs contributed more at 2 m, while odd-carbon homologues (C₁₁, C₁₃ and C₁₅) 3-OH-FAs contributed more at 220 m, regardless of diurnal changes (Fig. 4c).



3.3 Comparison of the endotoxin level with literature results

The airborne endotoxin level at near-ground observed in Tianjin is compared with results in literatures (Fig. 5). Different sampling methods and endotoxin assay methodologies may affect the results of the endotoxin assessment (De Rooij et al., 2017). Herein only the endotoxin level in aerosol samples assessed by chemical analytical method with 3-OH-FAs as biomarkers are compared.




Figure 5. Comparisons of average endotoxin concentrations in aerosols with various size ranges during different seasons. All data are estimated from 3-OH-FAs measured as biomarkers by chemical analytical method. The error bars are the standard derivations. Note: 1, 1990-1993; 2, 2001-2003; 3, total endotoxins in this study (at ground); 4, bioactive endotoxins in this study (at ground); 5, rural region, Hongkong University of Science and Technology; 6, urban region, Hongkong Science Museum; 7, urban region, Hong Kong

During all seasons, the mean mass concentration of airborne endotoxins in Tianjin was much higher than that in remote islands such as Gosan (except for spring) (Tyagi et al., 2017), Chichijima (Bikkina et al., 2021b; Tyagi et al., 2015a), but close to the endotoxin levels measured in developed areas such as Tokyo (Bikkina et al., 2021a), and the Pearl River Delta in China (Cheng et al., 2012). These results imply that high endotoxin mass concentrations in urban areas may be caused by enhanced human activities (Lee et al., 2004). Guan et al. (2014) discovered a higher endotoxin level in a high-traffic urban setting than in a low-traffic residential area, and similarly Madsen (2006) reported that the endotoxin concentrations on congested streets were higher than in residential areas.

Different from higher endotoxin levels observed in winter and summer in Tianjin, the endotoxin level was much higher in spring at Gosan, Korea, which was probably due to the enhanced mobilization of mineral dust through high altitudinal long-range atmospheric transport (Tyagi et al., 2017). Dutch Expert Committee on Occupational Safety (DECOS), a Committee of



the Health Council of the Netherlands stated that exposure to endotoxin above 9 ng m^{-3} for 8 hours or more increases the risk of respiratory diseases (Decos, 2010). The endotoxin concentrations reported in this study and in previous studies exceeded this value, except for remote islands such as Chichijima (Fig. 5), raising an environmental health issue caused by airborne endotoxins.

295 It should be noted that the endotoxin concentration proposed by DECOS refers to the mass concentration of bioactive endotoxin rather than total mass concentration (Decos, 2010). In light of the multiple studies, bioactive endotoxin appears to be correlated with C_{10} , C_{14} , C_{16} 3-OH-FAs in coarse mode particles in the Pearl River Delta region (Cheng et al., 2012); C_{10} , C_{12} , C_{14} 3-OH-FAs in house dust (Saraf et al., 1997; Saraf et al., 1999; Park et al., 2004; Uhlig et al., 2016); C_{14} , C_{16} 3-OH-FAs in airplane seat dust (Hines et al., 2003); and C_{12} , C_{14} 3-OH-FAs in agricultural dust (Reynolds et al., 2005). In this regard, C_{10} , C_{12} , C_{14} ,
300 C_{16} 3-OH-FAs were selected to estimate the bioactive endotoxin potential in this study. Correspondingly, the estimated concentrations of bioactive endotoxins were substantially lower compared to total endotoxins (Fig. 5). The level of bioactive endotoxins at 2 m was 6.59 ng m^{-3} in spring, 8.68 ng m^{-3} in autumn, lower than the endotoxin exposure threshold (9 ng m^{-3}). While the level of bioactive endotoxins exceeded the threshold in summer (9.10 ng m^{-3}) and winter (12.4 ng m^{-3}), implying possible health risks of airborne endotoxin exposure.

305 3.4 Mass loading of airborne Gram-negative bacteria

Based on the empirical equations, the dry mass loading of airborne GNB was estimated from the endotoxin mass concentration. The mass concentrations of airborne GNB measured in this study and previous literatures measured in urban, marine and alpine aerosols are listed in Table 2. The dry mass concentration of airborne GNB showed obvious regional variations. The airborne GNB dry mass concentration in Tianjin was the highest in winter ($883 \pm 167 \text{ ng m}^{-3}$), followed by summer ($788 \pm 538 \text{ ng m}^{-3}$),
310 autumn ($684 \pm 465 \text{ ng m}^{-3}$), and the lowest in spring ($468 \pm 220 \text{ ng m}^{-3}$) at 2 m. These levels were close to those in the megacity Tokyo (Bikkina et al., 2021a), urban area in Hong Kong (HKSCM) (Lee et al., 2004) and spring Gosan (Tyagi et al., 2017). The airborne GNB loading in Tianjin was much higher than those measured in remote islands as Chichijima island, and at Mount Tai during the low biomass burning events.

Amato et al. (2007) observed that airborne GNB were more abundant in summer than winter in terms of aerobic cultivable
315 microorganisms presented in cloud water. The phenomenon was further explained by the preferential occurrence of microbial strains on vegetation is far more developed in summer than in winter and GNB are usually more resistant to ultraviolet (UV) damage. Heavy biomass burning activities may also result in higher GNB mass concentration. GNB mass concentration measured at Mount Tai, China during the high biomass burning events, were 5 times higher than that during the low biomass burning period (Tyagi et al., 2016). During biomass burning events (i.e., wildfires and prescribed fires), microbes can be
320 aerosolized from terrestrial sources into the atmosphere (Bonfantine et al., 2024; Kobziar et al., 2024). It was previously assumed that pyrogenic carbon or smoke produced by biomass burning provides temporary habitat for microbes aerosolized



from soils and vegetations (Kobziar and Thompson, 2020; Bonfantine et al., 2024), while particulate matter in smoke confers attenuation of ultraviolet-B (UVB) and UV-A radiation (Mims et al., 1997), further leading to higher microbial cell concentration (Moore et al., 2021). Thus, the high GNB dry mass loading in Tianjin, extremely during summer and winter, could be resulted from the contributions of specific strains of GNB and/or microbial emissions during biomass burning. There are few descriptions of the atmospheric GNB mass concentrations in Tianjin in the literatures, and the distribution of different GNB community, namely the species distribution and their effects on human health remain to be further investigated.

Table 2. Dry mass concentrations of GNB estimated in this study and previous studies.

Sampling (period)	site	Sample types	Time / Location	GNB (ng m ⁻³) Mean ± SD	References
Tianjin, China (2020-2021)		PM ₁₀	Autumn (2 m/220 m)	684±464/ 427±237	This study
			Winter (2 m/220 m)	883±167/ 560±149	
			Spring (2 m)	468±220	
			Summer (2 m/220 m)	788±538/ 611±402	
Hong Kong, China (2002-2003)	Kong,	PM _{2.5}	HKUST	182	(Lee et al., 2004)
			HKSCM	313	
		PM _{2.5-10}	HKUST	45	
			HKSCM	93	
		PM ₁₀	HKUST	228	
			HKSCM	408	
Mount Tai, China (2006)		TSP	H-BEES	390	(Tyagi et al., 2016)
			L-BEES	75	
Chichijima, Japan (1990-1993, 2001-2003)		TSP	Spring	66, 200	(Bikkina et al., 2021b)
			Summer	48, 100	
			Autumn	35, 58	
			Winter	50, 194	
Tokyo, Japan (1994)		TSP	May	313	(Bikkina et al., 2021a)
			September	636	
Gosan, Korea (2001-2002)		TSP	Spring	928	(Tyagi et al., 2017)
			Summer	118	
			Autumn	194	
			Winter	380	

Note: SD, standard derivation; HKUST, Hongkong University of Science and Technology; HKSCM, Hongkong Science Museum; H-BEES, high biomass-burning samples; L-BEES, low biomass-burning samples.



3.5 Influencing factors of airborne endotoxins

3.5.1 Effects of meteorological conditions and gaseous pollutants

As components of GNB cell membranes, endotoxins are released during bacterial growth or death, meaning their mass
335 concentration is determined not only by the abundance of GNB but also by factors such as GNB sources, meteorological
conditions, surface types, and gaseous pollutants (Guan et al., 2014; Rolph et al., 2018).

Previous studies have demonstrated that air temperature has a more predominant role than other environmental factors
(Traversi et al., 2010; Carty et al., 2003). Higher endotoxin level was found in winter and autumn in Korea, and a negative
association was observed between endotoxins and air temperature (Hwang and Park, 2019). Guan et al. (2014) observed high
340 endotoxin level at 4–10°C, and assumed this temperature range was suitable for bacterial growth. And the endotoxin
concentration was higher under very dry or wet conditions compared to relative humidity (RH) was 40–80%. Cheng et al.
(2012) suggested the differences in bacterial fauna and growth rates can cause different endotoxin level under similar climatic
conditions. In this study, during the winter period in Tianjin, the RH was the lowest (46%) compared to other seasons, while
wind speed reached the highest at 220 m (4.80 m s⁻¹) and the temperature was -7–5°C, which might be suitable for the survival
345 of specific GNB species, resulting the higher endotoxin level. In winter when air pollution is frequent, pollutants and other
elements originating from anthropogenic sources act as toxins to GNB, thereby increasing the concentration of endotoxins as
a result of cell death level (Guan et al., 2014; Mahapatra et al., 2018), providing a possible explanation for the highest endotoxin
level in winter. Moreover, meteorological conditions characterized by stable weather patterns, such as high atmospheric
pressure (Ormanova et al., 2020), reduced mixing layer height (MLH) and temperature inversions create unfavorable
350 conditions for pollutant dispersion (Li et al., 2022; Yao et al., 2022). These conditions can lead to the accumulation of pollutants,
exacerbating air pollution levels, particularly during the winter season in Tianjin.

Hierarchical cluster analysis classified the mass concentration of 3-OH-FAs (C₈ – C₁₈) into 3 groups: short-chain (C₈, C₉),
mid-chain (C₁₀ – C₁₃), and long-chain (C₁₄ – C₁₈) 3-OH-FAs. Each group likely shares similar sources or influencing factors.
Spearman correlation analysis (Fig. S2) and Mantel test (Fig. 6) were used to evaluate the relationships between 3-OH-FAs,
355 endotoxins, and environmental factors (meteorological parameters, carbonaceous fraction, anions and cations, air pollutants).
Among meteorological parameters, RH and temperature mainly accounted for observed variations in 3-OH-FAs concentrations
(Fig. 6) (Mantel's $p < 0.01$). Most short/mid-chain (C₈ – C₁₃) 3-OH-FAs are positively correlated with air temperature (TEM)
and relative humidity, and negatively correlated with wind speed (WS) at both ground level and 220 m (Fig. S2). Long-chain
(C₁₄ – C₁₈) 3-OH-FAs were negatively correlated with TEM and WS at near ground. These findings align with previous study
360 (Guan et al., 2014), which reported significant correlations between endotoxin levels and TEM and RH. Moderate TEM and
RH are considered conducive to bacterial growth (Allen et al., 2011; Guan et al., 2014). During summer, the average air
temperature (27.8 °C at 2 m, 25.7 °C at 220 m) and relative humidity (72% at 2 m, 81% at 220 m) were the highest, with



relatively intense solar radiation (156 W m^{-2}). These meteorological conditions were favorable for photochemical reactions to occur (Huo et al., 2024; Pavuluri et al., 2015), indicating alternative photochemical reaction pathways of generating airborne short/mid-chain ($\text{C}_8 - \text{C}_{13}$) 3-OH-FAs.




Figure 6. Mantel test of 3-OH-FAs with meteorological conditions and chemical components of PM_{10} .

Previous studies have shown that gaseous pollutants (O_3 , SO_2 , NO_2 , CO_2 , and CO) can affect bacterial diversity and richness (Yan et al., 2018; Qi et al., 2020; Dong et al., 2016), potentially leading to varying distributions of 3-OH-FAs and endotoxins. The endotoxins mass concentrations exhibited a similar trend with $\text{PM}_{2.5}$, PM_{10} , NO_2 and CO (Fig. 2). Spearman's correlation analysis revealed significant positive correlations ($p < 0.01$) between endotoxin mass concentrations, NO_2 , and CO near the ground (2 m) (Fig. S2, 7c, 7d), along with a significant negative correlation ($p < 0.05$) with O_3 (Fig. S2, 7b). These results are consistent with previous studies (Guan et al., 2014; Hwang and Park, 2019), suggesting that gaseous pollutants, including SO_2 , NO_x and O_3 , may kill GNB, thereby increasing endotoxin levels as result of cell death. Exposure to high concentration of SO_2 and NO_2 significantly increases the fragility and disruption of bioaerosols as pollen, and may possibly increase the incidence of allergic airway disease in sensitized individuals by facilitating the bioavailability of airborne bioaerosols (Ouyang et al., 2016). Atmospheric photodegradation of proteinaceous matter may occur through reactions with O_3 , NO_2 , and relative radicals, causing the damage of the main components of cell membranes (Green et al., 2013; Wang et al., 2019; Xu et al., 2020). Additionally, the combined concentration of O_3 and NO_2 , defined as the total oxidants in previous studies, has been used to estimate atmospheric oxidizing capacity (Hu et al., 2016; Hu et al., 2017). In this study, endotoxin mass concentrations presented a positive correlation with total oxidants in winter while a negative correlation in spring (Fig. 7e, f), suggesting



385 differing mechanisms of endotoxin release and presence between these two seasons. In winter, strong atmospheric oxidizing conditions ($O_x=68-98 \mu\text{g m}^{-3}$) likely induce oxidative stress, causing cell membrane fragmentation and subsequent endotoxin release (Hwang and Park, 2019; Mahapatra et al., 2018). While in spring with relative high total oxidants ($O_x=78-126 \mu\text{g m}^{-3}$), the community structure of bacteria may change, with certain bacteria adapting to the polluted atmosphere and survived the circumstances, and the species becoming more limited (Yin et al., 2021). No significant correlations were found between endotoxins and total oxidants in summer ($O_x=92-199 \mu\text{g m}^{-3}$) or autumn ($O_x=69-140 \mu\text{g m}^{-3}$).

390




Figure 7. Spearman's correlation analyses between mass concentrations of endotoxins and air pollutants. (a) PM_{2.5} and PM_{2.5-10}; (b) O₃; (c) NO₂; (d) CO; (e-f) Total oxidants [O₃+NO₂] in winter and spring.

395 **3.5.2 Effects of particulate pollutants**

The endotoxin levels generally increased significantly under high PM pollution, showing a strong correlation with PM₁₀ concentrations (Shen et al., 2019; Zhong et al., 2019). Previous studies demonstrated that 60% of biological active endotoxins (measured by LAL method) were associated with coarse-mode particles (PM_{2.5-10}) emitted from natural sources, including soil dust and vegetation. Meanwhile, 80% of total endotoxins (measured by GC-MS method) were associated with fine particles
400 (PM_{2.5}) originating from anthropogenic combustion processes (Cheng et al., 2012; Seinfeld et al., 1998). The endotoxin mass concentrations measured by LC-MS in this study showed a similar temporal trend to PM_{2.5} and PM₁₀ (Fig. 3a, 3b). Here, airborne endotoxin levels were significantly and positively correlated with PM_{2.5} ($p < 0.01$), while no significant correlation



with $PM_{2.5-10}$ was observed (Fig. 7a). This result indicates that total airborne endotoxins are most likely related to anthropogenic sources.

405 Spearman correlation analysis (Fig. S2) and Mantel test (Fig. 6) revealed that short-chain 3-OH-FAs ($C_{10} - C_{13}$) at 220 m were mainly associated with meteorological conditions, WSOC, and Mg^{2+} , reflecting their emissions from natural sources and secondary transformation during transport. At near ground level, all 3-OH-FAs showed significant correlations with OC, WSOC, K^+ , Ca^{2+} , Cl^- , SO_4^{2-} , NO_3^- , NH_4^+ , Mg^{2+} and temperature ($p < 0.05$). This finding suggests that the emissions of long-chain 3-OH-FAs are influenced by multiple factors, including natural sources such as soil and anthropogenic sources such as
410 combustion (Zhong et al., 2019). Composting facilities, intensive farming (e.g., cattle, swine and poultry), and wastewater treatment facilities are all potential anthropogenic sources of airborne endotoxins (Hu et al., 2020a; Rolph et al., 2018). According to Mahapatra et al. (2018), certain species of GNB may release more endotoxins when disrupted by anthropogenic contamination, or the source of higher pollution may themselves be significant sources of endotoxins.

415 **4 Conclusions**

This study establishes the first continuous time series of airborne endotoxins in Tianjin through comprehensive quantification of 3-OH-FAs ($C_{10} - C_{18}$) in PM_{10} . By resolving vertical, seasonal, and diurnal variability patterns, we demonstrate significant altitude-dependent gradients (2 m vs. 220 m) and distinct source-driven temporal dynamics. Key findings include: (1) Higher total endotoxin concentrations at ground level compared to higher altitudes, peaking in winter due to combined contributions
420 of near-ground emissions from both natural (e.g., soil microbial activity) and anthropogenic sources (e.g., biomass combustion); (2) Source differentiation via chain length specificity, with short- and mid-chain 3-OH-FAs ($C_8 - C_{13}$) associated with atmospheric processes (microbial emissions and photochemical reactions), while long-chain homologues ($C_{14} - C_{18}$) correlated with anthropogenic markers; (3) Total endotoxins are comparable to those in industrialized regions; the marked vertical attenuation emphasizes the localized nature of ground-level emission impacts. Bioactive endotoxin levels remained below
425 DECOS thresholds, except during winter maxima, indicating seasonally variable health risks. (4) The significant correlation between total oxidants ($O_3 + NO_2$) and endotoxin levels in winter and spring underscores atmospheric oxidative capacity as a critical modulator of endotoxin bioactivity.

Further research is essential to disentangle specific factors influencing endotoxin concentrations, explore the dynamics between total and bioactive endotoxins, and identify GNB species contributing to airborne endotoxins in Tianjin.

430

Data availability. The dataset for this paper is available upon request from the corresponding author (huwei@tju.edu.cn; fupingqing@tju.edu.cn).



Author contribution. PF and WH designed the entire study and the experiments. WZ visualized the data, and determined the structure of the article. WZ, MN, QZ, and NA executed experiments. WZ analyzed the data. WZ and WH wrote the draft
435 manuscript with input from all the authors. All the authors discussed, edited, and approved the final manuscript.

Competing interests. The authors declare that they have no conflict of interests.

Acknowledgments. This work was supported by the National Natural Science Foundation of China (Grant No. 42130513). We thank Yajie Wang and Donghuan Zhang for their help in sampling.

440 **References**

Abdel Hameed, A. A., Khoder, M. I., Yuosra, S., Osman, A. M., and Ghanem, S.: Diurnal distribution of airborne bacteria and fungi in the atmosphere of Helwan area, Egypt, *Sci. Total Environ.*, 407, 6217-6222, <https://doi.org/10.1016/j.scitotenv.2009.08.028>, 2009.

Allen, J., Bartlett, K., Graham, M., and Jackson, P.: Ambient concentrations of airborne endotoxin in two cities in the interior
445 of British Columbia, Canada, *J. Environ. Monit.*, 13, 631-640, <https://doi.org/10.1039/c0em00235f>, 2011.

Amato, P., Parazols, M., Sancelme, M., Laj, P., Mailhot, G., and Delort, A. M.: Microorganisms isolated from the water phase of tropospheric clouds at the Puy de Dome: major groups and growth abilities at low temperatures, *FEMS Microbiol. Ecol.*, 59, 242-254, <https://doi.org/10.1111/j.1574-6941.2006.00199.x>, 2007.

Amin, H., Santl-Temkiv, T., Cramer, C., Finster, K., Real, F. G., Gislason, T., Holm, M., Janson, C., Jogi, N. O., Jogi, R.,
450 Malinovski, A., Marshall, I. P. G., Modig, L., Norback, D., Shigdel, R., Sigsgaard, T., Svanes, C., Thorarinsdottir, H., Wouters, I. M., Schlunssen, V., and Bertelsen, R. J.: Indoor Airborne Microbiome and Endotoxin: Meteorological Events and Occupant Characteristics Are Important Determinants, *Environ. Sci. Technol.*, 57, 11750-11766, <https://doi.org/10.1021/acs.est.3c01616>, 2023.

Balkwill, D. L., Leach, F. R., Wilson, J. T., McNabb, J. F., and White, D. C.: Equivalence of microbial biomass measures based
455 on membrane lipid and cell wall components, adenosine triphosphate, and direct counts in subsurface aquifer sediments, *Microb. Ecol.*, 16, 73-84, <https://doi.org/10.1007/BF02097406>, 1988.

Bikkin, P., Kawamura, K., Bikkina, S., Kunwar, B., Tanaka, K., and Suzuki, K.: Hydroxy Fatty Acids in Remote Marine Aerosols over the Pacific Ocean: Impact of Biological Activity and Wind Speed, *ACS Earth Space Chem.*, 3, 366-379, <https://doi.org/10.1021/acsearthspacechem.8b00161>, 2019.

460 Bikkina, P., Kawamura, K., Bikkina, S., and Tanaka, N.: Hydroxy Fatty Acids in Rainwater and Aerosols from Suburban Tokyo in Central Japan: The Impact of Long-Range Transport of Soil Microbes and Plant Waxes, *ACS Earth Space Chem.*, 5, 257-267, <https://doi.org/10.1021/acsearthspacechem.0c00275>, 2021a.

Bikkina, P., Kawamura, K., Bikkina, S., and Yamaguchi, H.: Decadal Variations in Hydroxy Fatty Acids Over Chichijima



- Island in the North Pacific: Long-Term Seasonal Variability in Plant and Microbial Markers, *Journal of Geophysical Research-*
465 *Atmospheres*, 126, e2020JD033347, <https://doi.org/10.1029/2020JD033347>, 2021b.
- Binding, N., Jaschinski, S., Werlich, S., Bletz, S., and Witting, U.: Quantification of bacterial lipopolysaccharides (endotoxin) by GC-MS determination of 3-hydroxy fatty acids, *J. Environ. Monit.*, 6, 65-70, <https://doi.org/10.1039/b309237b>, 2004.
- Bonfantine, K., Vuono, D. C., Christner, B. C., Moore, R., Fox, S., Dean, T., Betancourt, D., Watts, A., and Kobziar, L. N.: Evidence for Wildland Fire Smoke Transport of Microbes From Terrestrial Sources to the Atmosphere and Back, *Journal of*
470 *Geophysical Research-Biogeosciences*, 129, e2024JG008236, <https://doi.org/10.1029/2024JG008236>, 2024.
- Brooks, J. P., Tanner, B. D., Gerba, C. P., and Pepper, I. L.: The measurement of aerosolized endotoxin from land application of Class B biosolids in Southeast Arizona, *Can. J. Microbiol.*, 52, 150-156, <https://doi.org/10.1139/w05-115>, 2006.
- Carty, C. L., Gehring, U., Cyrus, J., Bischof, W., and Heinrich, J.: Seasonal variability of endotoxin in ambient fine particulate matter, *J. Environ. Monit.*, 5, 953-958, <https://doi.org/10.1039/b308488d>, 2003.
- 475 Cheng, J. Y. W., Hui, E. L. C., and Lau, A. P. S.: Bioactive and total endotoxins in atmospheric aerosols in the Pearl River Delta region, China, *Atmos. Environ.*, 47, 3-11, <https://doi.org/10.1016/j.atmosenv.2011.11.055>, 2012.
- de Rooij, M. M., Heederik, D. J., Borlee, F., Hoek, G., and Wouters, I. M.: Spatial and temporal variation in endotoxin and PM₁₀ concentrations in ambient air in a livestock dense area, *Environ. Res.*, 153, 161-170, <https://doi.org/10.1016/j.envres.2016.12.004>, 2017.
- 480 DECOS: Endotoxins. Health-based recommended occupational exposure limit. , 2010.
- Dong, L., Qi, J., Shao, C., Zhong, X., Gao, D., Cao, W., Gao, J., Bai, R., Long, G., and Chu, C.: Concentration and size distribution of total airborne microbes in hazy and foggy weather, *Sci. Total Environ.*, 541, 1011-1018, <https://doi.org/10.1016/j.scitotenv.2015.10.001>, 2016.
- Fan, M. Y., Zhang, Y. L., Hong, Y. H., Lin, Y. C., Zhao, Z. Y., Cao, F., Sun, Y. L., Guo, H., and Fu, P. Q.: Vertical Differences
485 of Nitrate Sources in Urban Boundary Layer Based on Tower Measurements, *Environmental Science & Technology Letters*, 9, 906-912, <https://doi.org/10.1021/acs.estlett.2c00600>, 2022.
- Farokhi, A., Heederik, D., and Smit, L. A. M.: Respiratory health effects of exposure to low levels of airborne endotoxin - a systematic review, *Environ. Health*, 17, 14, <https://doi.org/10.1186/s12940-018-0360-7>, 2018.
- Fonseca Gabriel, M., Paciência, I., Felgueiras, F., Cavaleiro Rufo, J., Castro Mendes, F., Farraia, M., Mourão, Z., Moreira, A.,
490 and de Oliveira Fernandes, E.: Environmental quality in primary schools and related health effects in children. An overview of assessments conducted in the Northern Portugal, *Energy and Buildings*, 250, 111305, <https://doi.org/10.1016/j.enbuild.2021.111305>, 2021.
- Green, M. C., Fedorov, D. G., Kitaura, K., Francisco, J. S., and Slipchenko, L. V.: Open-shell pair interaction energy decomposition analysis (PIEDA): formulation and application to the hydrogen abstraction in tripeptides, *J. Chem. Phys.*, 138,



- 495 074111, <https://doi.org/10.1063/1.4790616>, 2013.
- Guan, T. J., Yao, M. S., Wang, J. X., Fang, Y. H., Hu, S. H., Wang, Y., Dutta, A., Yang, J. N., Wu, Y. S., Hu, M., and Zhu, T.: Airborne endotoxin in fine particulate matter in Beijing, *Atmos. Environ.*, 97, 35-42, <https://doi.org/10.1016/j.atmosenv.2014.08.005>, 2014.
- Guo, Z., Wang, Z., Qian, L., Zhao, Z., Zhang, C., Fu, Y., Li, J., Zhang, C., Lu, B., and Qian, J.: Biological and chemical
500 compositions of atmospheric particulate matter during hazardous haze days in Beijing, *Environ. Sci. Pollut. Res. Int.*, 25, 34540-34549, <https://doi.org/10.1007/s11356-018-3355-6>, 2018.
- Heederik, D. and Douwes, J.: Towards an occupational exposure limit for endotoxins, *Ann. Agric. Environ. Med.*, 4, 17-19, 1997.
- Hines, C. J., Waters, M. A., Larsson, L., Petersen, M. R., Saraf, A., and Milton, D. K.: Characterization of endotoxin and 3-
505 hydroxy fatty acid levels in air and settled dust from commercial aircraft cabins, *Indoor Air*, 13, 166-173, <https://doi.org/10.1034/j.1600-0668.2003.00175.x>, 2003.
- Hu, W., Hu, M., Hu, W. W., Zheng, J., Chen, C., Wu, Y. S., and Guo, S.: Seasonal variations in high time-resolved chemical compositions, sources, and evolution of atmospheric submicron aerosols in the megacity Beijing, *Atmospheric Chemistry and Physics*, 17, 9979-10000, <https://doi.org/10.5194/acp-17-9979-2017>, 2017.
- 510 Hu, W., Wang, Z. H., Huang, S., Ren, L. J., Yue, S. Y., Li, P., Xie, Q. R., Zhao, W. Y., Wei, L. F., Ren, H., Wu, L. B., Deng, J. J., and Fu, P. Q.: Biological Aerosol Particles in Polluted Regions, *Current Pollution Reports*, 6, 65-89, <https://doi.org/10.1007/s40726-020-00138-4>, 2020a.
- Hu, W., Liu, D. D., Su, S. H., Ren, L. J., Ren, H., Wei, L. F., Yue, S. Y., Xie, Q. R., Zhang, Z. M., Wang, Z. H., Yang, N., Wu, L. B., Deng, J. J., Qi, Y. L., and Fu, P. Q.: Photochemical Degradation of Organic Matter in the Atmosphere, *Advanced
515 Sustainable Systems*, 5, 2100027, <https://doi.org/10.1002/adsu.202100027>, 2021.
- Hu, W. W., Hu, M., Hu, W., Jimenez, J. L., Yuan, B., Chen, W. T., Wang, M., Wu, Y. S., Chen, C., Wang, Z. B., Peng, J. F., Zeng, L. M., and Shao, M.: Chemical composition, sources, and aging process of submicron aerosols in Beijing: Contrast between summer and winter, *Journal of Geophysical Research-Atmospheres*, 121, 1955-1977, <https://doi.org/10.1002/2015jd024020>, 2016.
- 520 Hu, Z., Liu, H., Zhang, H., Zhang, X., Zhou, M., Lou, L., Zheng, P., Xi, C., and Hu, B.: Temporal discrepancy of airborne total bacteria and pathogenic bacteria between day and night, *Environ. Res.*, 186, 109540, <https://doi.org/10.1016/j.envres.2020.109540>, 2020b.
- Huo, Y. X., Lyu, X., Yao, D. W., Zhou, B. N., Yuan, Q., Lee, S. C., and Guo, H.: Exploring the Formation of High Levels of Hydroxyl Dicarboxylic Acids at an Urban Background Site in South China, *Journal of Geophysical Research-Atmospheres*,
525 129, e2023JD040096, <https://doi.org/10.1029/2023JD040096>, 2024.



- Hwang, S. H. and Park, D. U.: Ambient Endotoxin and Chemical Pollutant (PM_{2.5}, PM₁₀, and O₃) Levels in South Korea, *Aerosol and Air Quality Research*, 19, 786-793, <https://doi.org/10.4209/aaqr.2018.06.0235>, 2019.
- Kawamura, K., Ishimura, Y., and Yamazaki, K.: Four years' observations of terrestrial lipid class compounds in marine aerosols from the western North Pacific, *Global Biogeochem. Cycles*, 17, 3-1-3-19, <https://doi.org/10.1029/2001gb001810>, 2003.
- 530 Kobziar, L. N. and Thompson, G. R., 3rd: Wildfire smoke, a potential infectious agent, *Science*, 370, 1408-1410, <https://doi.org/10.1126/science.abe8116>, 2020.
- Kobziar, L. N., Lampman, P., Tohidi, A., Kochanski, A. K., Cervantes, A., Hudak, A. T., McCarley, R., Gullett, B., Aurell, J., Moore, R., Vuono, D. C., Christner, B. C., Watts, A. C., Cronan, J., and Ottmar, R.: Bacterial Emission Factors: A Foundation for the Terrestrial-Atmospheric Modeling of Bacteria Aerosolized by Wildland Fires, *Environ. Sci. Technol.*, 58, 2413-2422, <https://doi.org/10.1021/acs.est.3c05142>, 2024.
- 535 Laboha, P., Sychrova, E., Brozman, O., Sovadinova, I., Blahova, L., Prokes, R., Ondracek, J., and Babica, P.: Cyanobacteria, cyanotoxins and lipopolysaccharides in aerosols from inland freshwater bodies and their effects on human bronchial cells, *Environ. Toxicol. Pharmacol.*, 98, 104073, <https://doi.org/10.1016/j.etap.2023.104073>, 2023.
- Laitinen, S., Kangas, J., Husman, K., and Susitaival, P.: Evaluation of exposure to airborne bacterial endotoxins and peptidoglycans in selected work environments, *Ann. Agric. Environ. Med.*, 8, 213-219, 2001.
- 540 Lang-Yona, N., Lehahn, Y., Herut, B., Burshtein, N., and Rudich, Y.: Marine aerosol as a possible source for endotoxins in coastal areas, *Sci. Total Environ.*, 499, 311-318, <https://doi.org/10.1016/j.scitotenv.2014.08.054>, 2014.
- Lee, A. K. Y., Chan, C. K., Fang, M., and Lau, A. P. S.: The 3-hydroxy fatty acids as biomarkers for quantification and characterization of endotoxins and Gram-negative bacteria in atmospheric aerosols in Hong Kong, *Atmos. Environ.*, 38, 6307-6317, <https://doi.org/10.1016/j.atmosenv.2004.08.013>, 2004.
- 545 Lei, L., Sun, Y., Ouyang, B., Qiu, Y., Xie, C., Tang, G., Zhou, W., He, Y., Wang, Q., Cheng, X., Fu, P., and Wang, Z.: Vertical Distributions of Primary and Secondary Aerosols in Urban Boundary Layer: Insights into Sources, Chemistry, and Interaction with Meteorology, *Environ. Sci. Technol.*, 55, 4542-4552, <https://doi.org/10.1021/acs.est.1c00479>, 2021.
- Li, Y., Du, A., Lei, L., Sun, J., Li, Z., Zhang, Z., Wang, Q., Tang, G., Song, S., Wang, Z., Wang, Z., and Sun, Y.: Vertically Resolved Aerosol Chemistry in the Low Boundary Layer of Beijing in Summer, *Environ. Sci. Technol.*, 56, 9312-9324, <https://doi.org/10.1021/acs.est.2c02861>, 2022.
- 550 Li, Z. Q., Guo, J. P., Ding, A. J., Liao, H., Liu, J. J., Sun, Y. L., Wang, T. J., Xue, H. W., Zhang, H. S., and Zhu, B.: Aerosol and boundary-layer interactions and impact on air quality, *National Science Review*, 4, 810-833, <https://doi.org/10.1093/nsr/nwx117>, 2017.
- 555 Liebers, V., Bruning, T., and Raulf-Heimsoth, M.: Occupational endotoxin-exposure and possible health effects on humans, *Am. J. Ind. Med.*, 49, 474-491, <https://doi.org/10.1002/ajim.20310>, 2006.



- Liebers, V., Bruning, T., and Raulf, M.: Occupational endotoxin exposure and health effects, *Arch. Toxicol.*, 94, 3629-3644, <https://doi.org/10.1007/s00204-020-02905-0>, 2020.
- Liebers, V., Raulf-Heimsoth, M., and Bruning, T.: Health effects due to endotoxin inhalation (review), *Arch. Toxicol.*, 82, 203-560 210, <https://doi.org/10.1007/s00204-008-0290-1>, 2008.
- Lundin, J. I. and Checkoway, H.: Endotoxin and cancer, *Environ. Health Perspect.*, 117, 1344-1350, <https://doi.org/10.1289/ehp.0800439>, 2009.
- Madsen, A. M.: Airborne endotoxin in different background environments and seasons, *Ann. Agric. Environ. Med.*, 13, 81-86, <https://doi.org/10.1016/j.ijheh.2010.03.001>, 2006.
- 565 Mahapatra, P. S., Jain, S., Shrestha, S., Senapati, S., and Puppala, S. P.: Ambient endotoxin in PM(10) and association with inflammatory activity, air pollutants, and meteorology, in Chitwan, Nepal, *Sci. Total Environ.*, 618, 1331-1342, <https://doi.org/10.1016/j.scitotenv.2017.09.249>, 2018.
- Makkonen, U., Hellen, H., Anttila, P., and Ferm, M.: Size distribution and chemical composition of airborne particles in south-eastern Finland during different seasons and wildfire episodes in 2006, *Sci. Total Environ.*, 408, 644-651, 570 <https://doi.org/10.1016/j.scitotenv.2009.10.050>, 2010.
- Mattsby-Baltzer, I., Lindgren, K., Lindholm, B., and Edebo, L.: Endotoxin shedding by enterobacteria: free and cell-bound endotoxin differ in Limulus activity, *Infect. Immun.*, 59, 689-695, <https://doi.org/10.1128/iai.59.2.689-695.1991>, 1991.
- Mims, F. M., 3rd, Holben, B. N., Eck, T. F., Montgomery, B. C., and Grant, W. B.: Smoky skies, mosquitoes, and disease, *Science*, 276, 1774-1775, <https://doi.org/doi:10.1126/science.276.5320.1773c>, 1997.
- 575 Moore, R. A., Bomar, C., Kobziar, L. N., and Christner, B. C.: Wildland fire as an atmospheric source of viable microbial aerosols and biological ice nucleating particles, *ISME J*, 15, 461-472, <https://doi.org/10.1038/s41396-020-00788-8>, 2021.
- Moretti, S., Smets, W., Oerlemans, E., Blust, R., and Lebeer, S.: The abundance of urban endotoxins as measured with an impinger-based sampling strategy, *Aerobiologia*, 34, 487-496, <https://doi.org/10.1007/s10453-018-9525-7>, 2018.
- Mueller-Anneling, L., Avol, E., Peters, J. M., and Thorne, P. S.: Ambient endotoxin concentrations in PM₁₀ from Southern 580 California, *Environ. Health Perspect.*, 112, 583-588, <https://doi.org/10.1289/ehp.6552>, 2004.
- Mukherjee, A. and Agrawal, M.: World air particulate matter: sources, distribution and health effects, *Environmental Chemistry Letters*, 15, 283-309, <https://doi.org/10.1007/s10311-017-0611-9>, 2017.
- Niu, M. T., An, N., Zhang, W. X., Hu, W., Fu, X. L., Kawamura, K., Feng, Y. Q., Zhu, Q. F., and Fu, P. Q.: Position-specific isomers of monohydroxy fatty acids in the land-atmosphere interface: identification and quantification, *Npj Climate and 585 Atmospheric Science*, 7, 75, <https://doi.org/10.1038/s41612-024-00621-5>, 2024.
- Ormanova, G., Karaca, F., and Kononova, N.: Analysis of the impacts of atmospheric circulation patterns on the regional air quality over the geographical center of the Eurasian continent, *Atmospheric Research*, 237, 104858,



<https://doi.org/10.1016/j.atmosres.2020.104858>, 2020.

Ouyang, Y., Xu, Z., Fan, E., Li, Y., and Zhang, L.: Effect of nitrogen dioxide and sulfur dioxide on viability and morphology
590 of oak pollen, *International Forum of Allergy & Rhinology*, 95-100,

Paba, E., Chiominto, A., Marcelloni, A. M., Tombolini, F., Paci, E., and Tranfo, G.: Endotoxin Analysis: Correlation Between
Biological and Chemical Methods, *Biomedical Journal of Scientific & Technical Research*, 19,
<https://doi.org/10.26717/BJSTR.2019.19.003339>, 2019.

Park, J. H., Szponar, B., Larsson, L., Gold, D. R., and Milton, D. K.: Characterization of lipopolysaccharides present in settled
595 house dust, *Appl. Environ. Microbiol.*, 70, 262-267, <https://doi.org/10.1128/AEM.70.1.262-267.2004>, 2004.

Park, J. H., Spiegelman, D. L., Burge, H. A., Gold, D. R., Chew, G. L., and Milton, D. K.: Longitudinal study of dust and
airborne endotoxin in the home, *Environ. Health Perspect.*, 108, 1023-1028, <https://doi.org/10.1289/ehp.001081023>, 2000.

Parker, J. H., Smith, G. A., Fredrickson, H. L., Vestal, J. R., and White, D. C.: Sensitive assay, based on hydroxy fatty acids
from lipopolysaccharide lipid A, for Gram-negative bacteria in sediments, *Appl. Environ. Microbiol.*, 44, 1170-1177,
600 <https://doi.org/10.1128/aem.44.5.1170-1177.1982>, 1982.

Pavuluri, C. M., Kawamura, K., Mihalopoulos, N., and Swaminathan, T.: Laboratory photochemical processing of aqueous
aerosols: formation and degradation of dicarboxylic acids, oxocarboxylic acids and α -dicarbonyls, *Atmospheric Chemistry and
Physics*, 15, 7999-8012, <https://doi.org/10.5194/acp-15-7999-2015>, 2015.

Petsch, D. and Anspach, F. B.: Endotoxin removal from protein solutions, *J. Biotechnol.*, 76, 97-119,
605 [https://doi.org/10.1016/s0168-1656\(99\)00185-6](https://doi.org/10.1016/s0168-1656(99)00185-6), 2000.

Phiri, Y. V. A., Wu, M. M., Chen, Y. H., Zou, M. L., Jiang, C. B., Wu, C. D., Huang, H. C., Lung, S. C. C., Chien, L. C., Lo, Y.
C., Lee, F. Y., and Chao, H. J.: Environmental determinants of household microbial and allergen levels in the Greater Taipei
Area, *Build. Environ.*, 230, 110003, <https://doi.org/10.1016/j.buildenv.2023.110003>, 2023.

Qi, Y., Li, Y., Xie, W., Lu, R., Mu, F., Bai, W., and Du, S.: Temporal-spatial variations of fungal composition in PM(2.5) and
610 source tracking of airborne fungi in mountainous and urban regions, *Sci. Total Environ.*, 708, 135027,
<https://doi.org/10.1016/j.scitotenv.2019.135027>, 2020.

Qiu, Y. M., Xie, Q. R., Wang, J. F., Xu, W. Q., Li, L. J., Wang, Q. Q., Zhao, J., Chen, Y. T., Chen, Y. F., Wu, Y. Z., Du, W.,
Zhou, W., Lee, J., Zhao, C. F., Ge, X. L., Fu, P. Q., Wang, Z. F., Worsnop, D. R., and Sun, Y. L.: Vertical Characterization and
Source Apportionment of Water-Soluble Organic Aerosol with High-resolution Aerosol Mass Spectrometry in Beijing, China,
615 *ACS Earth Space Chem.*, 3, 273-284, <https://doi.org/10.1021/acsearthspacechem.8b00155>, 2019.

Reynolds, S. J., Milton, D. K., Heederik, D., Thorne, P. S., Donham, K. J., Croteau, E. A., Kelly, K. M., Douwes, J., Lewis,
D., Whitmer, M., Connaughton, I., Koch, S., Malmberg, P., Larsson, B. M., Deddens, J., Saraf, A., and Larsson, L.:
Interlaboratory evaluation of endotoxin analyses in agricultural dusts--comparison of LAL assay and mass spectrometry, *J.*



- Environ. Monit., 7, 1371-1377, <https://doi.org/10.1039/b509256f>, 2005.
- 620 Rolph, C. A., Gwyther, C. L., Tyrrel, S. F., Nasir, Z. A., Drew, G. H., Jackson, S. K., Khera, S., Hayes, E. T., Williams, B., Bennett, A., Collins, S., Walsh, K., Kinnersley, R., and Gladding, T. L.: Sources of Airborne Endotoxins in Ambient Air and Exposure of Nearby Communities—A Review, *Atmosphere*, 9, 375, <https://doi.org/10.3390/atmos9100375>, 2018.
- Rylander, R.: Endotoxin in the environment--exposure and effects, *J. Endotoxin Res.*, 8, 241-252, <https://doi.org/10.1179/096805102125000452>, 2002.
- 625 Saraf, A., Larsson, L., Burge, H., and Milton, D.: Quantification of ergosterol and 3-hydroxy fatty acids in settled house dust by gas chromatography-mass spectrometry: comparison with fungal culture and determination of endotoxin by a *Limulus* amoebocyte lysate assay, *Appl. Environ. Microbiol.*, 63, 2554-2559, <https://doi.org/10.1128/aem.63.7.2554-2559.1997>, 1997.
- Saraf, A., Park, J. H., Milton, D. K., and Larsson, L.: Use of quadrupole GC-MS and ion trap GC-MS-MS for determining 3-hydroxy fatty acids in settled house dust: relation to endotoxin activity, *J. Environ. Monit.*, 1, 163-168, <https://doi.org/10.1039/a809019j>, 1999.
- 630 Schneiher, M., Razdan, S., Miller, A. M., Briceno, M. E., and Barua, S.: Current technologies to endotoxin detection and removal for biopharmaceutical purification, *Biotechnol. Bioeng.*, 117, 2588-2609, <https://doi.org/10.1002/bit.27362>, 2020.
- Schwarz, H., Gornicec, J., Neuper, T., Parigiani, M. A., Wallner, M., Duschl, A., and Horejs-Hoeck, J.: Biological Activity of Masked Endotoxin, *Sci. Rep.*, 7, 44750, <https://doi.org/10.1038/srep44750>, 2017.
- 635 Seinfeld, J. H., Pandis, S. N., and Noone, K. J.: *Atmospheric Chemistry and Physics*. , *Physics Today*, John Wiley & Sons, Inc, The United States of America. 1998.
- Shen, F., Niu, M., Zhou, F., Wu, Y., and Zhu, T.: Culturability, metabolic activity and composition of ambient bacterial aerosols in a surrogate lung fluid, *Sci. Total Environ.*, 690, 76-84, <https://doi.org/10.1016/j.scitotenv.2019.07.005>, 2019.
- Simoneit, B. R. T., Kobayashi, M., Mochida, M., Kawamura, K., Lee, M., Lim, H. J., Turpin, B. J., and Komazaki, Y.: 640 Composition and major sources of organic compounds of aerosol particulate matter sampled during the ACE-Asia campaign, *Journal of Geophysical Research-Atmospheres*, 109, D19S10, <https://doi.org/10.1029/2004jd004598>, 2004.
- Spaan, S., Wouters, I. M., Oosting, I., Doekes, G., and Heederik, D.: Exposure to inhalable dust and endotoxins in agricultural industries, *J. Environ. Monit.*, 8, 63-72, <https://doi.org/10.1039/b509838f>, 2006.
- Traversi, D., Alessandria, L., Schiliro, T., Chiado Piat, S., and Gilli, G.: Meteo-climatic conditions influence the contribution 645 of endotoxins to PM₁₀ in an urban polluted environment, *J. Environ. Monit.*, 12, 484-490, <https://doi.org/10.1039/b913314c>, 2010.
- Tyagi, P., Ishimura, Y., and Kawamura, K.: Hydroxy fatty acids in marine aerosols as microbial tracers: 4-year study on β - and ω -hydroxy fatty acids from remote Chichijima Island in the western North Pacific, *Atmos. Environ.*, 115, 89-100, <https://doi.org/10.1016/j.atmosenv.2015.05.038>, 2015a.



- 650 Tyagi, P., Yamamoto, S., and Kawamura, K.: Hydroxy fatty acids in fresh snow samples from northern Japan: long-range atmospheric transport of Gram-negative bacteria by Asian winter monsoon, *Biogeosciences*, 12, 7071-7080, <https://doi.org/10.5194/bg-12-7071-2015>, 2015b.
- Tyagi, P., Kawamura, K., Fu, P. Q., Bikkina, S., Kanaya, Y., and Wang, Z. F.: Impact of biomass burning on soil microorganisms and plant metabolites: A view from molecular distributions of atmospheric hydroxy fatty acids over Mount Tai, *Journal of Geophysical Research-Biogeosciences*, 121, 2684-2699, <https://doi.org/10.1002/2016jg003324>, 2016.
- 655 Tyagi, P., Kawamura, K., Kariya, T., Bikkina, S., Fu, P. Q., and Lee, M.: Tracing atmospheric transport of soil microorganisms and higher plant waxes in the East Asian outflow to the North Pacific Rim by using hydroxy fatty acids: Year-round observations at Gosan, Jeju Island, *Journal of Geophysical Research-Atmospheres*, 122, 4112-4131, <https://doi.org/10.1002/2016jd025496>, 2017.
- 660 Uhlig, S., Negard, M., Heldal, K. K., Straumfors, A., Madso, L., Bakke, B., and Eduard, W.: Profiling of 3-hydroxy fatty acids as environmental markers of endotoxin using liquid chromatography coupled to tandem mass spectrometry, *J. Chromatogr. A*, 1434, 119-126, <https://doi.org/10.1016/j.chroma.2016.01.038>, 2016.
- Wakeham, S. G., Pease, T. K., and Benner, R.: Hydroxy fatty acids in marine dissolved organic matter as indicators of bacterial membrane material, *Org. Geochem.*, 34, 857-868, [https://doi.org/10.1016/S0146-6380\(02\)00189-4](https://doi.org/10.1016/S0146-6380(02)00189-4), 2003.
- 665 Wang, S., Song, T., Shiraiwa, M., Song, J., Ren, H., Ren, L., Wei, L., Sun, Y., Zhang, Y., Fu, P., and Lai, S.: Occurrence of Aerosol Proteinaceous Matter in Urban Beijing: An Investigation on Composition, Sources, and Atmospheric Processes During the "APEC Blue" Period, *Environ. Sci. Technol.*, 53, 7380-7390, <https://doi.org/10.1021/acs.est.9b00726>, 2019.
- Wollenweber, H.-W. and Rietschel, E. T.: Analysis of lipopolysaccharide (lipid A) fatty acids, *J. Microbiol. Meth.*, 11, 195-211, [https://doi.org/10.1016/0167-7012\(90\)90056-c](https://doi.org/10.1016/0167-7012(90)90056-c), 1990.
- 670 Xiao, Y. T., Yang, J. B., Cui, L. H., Deng, J. J., Fu, P. Q., and Zhu, J. L.: Weakened Sea-Land Breeze in a Coastal Megacity Driven by Urbanization and Ocean Warming, *Earths Future*, 11, e2022EF003341, <https://doi.org/10.1029/2022EF003341>, 2023.
- Xu, Y., Xiao, H., Wu, D., and Long, C.: Abiotic and Biological Degradation of Atmospheric Proteinaceous Matter Can Contribute Significantly to Dissolved Amino Acids in Wet Deposition, *Environ. Sci. Technol.*, 54, 6551-6561, <https://doi.org/10.1021/acs.est.0c00421>, 2020.
- 675 Xue, T., Wang, R., Wang, M., Wang, Y., Tong, D., Meng, X., Huang, C., Ai, S., Li, F., Cao, J., Tong, M., Ni, X., Liu, H., Deng, J., Lu, H., Wan, W., Gong, J., Zhang, S., and Zhu, T.: Health benefits from the rapid reduction in ambient exposure to air pollutants after China's clean air actions: progress in efficacy and geographic equality, *Natl Sci Rev*, 11, nwad263, <https://doi.org/10.1093/nsr/nwad263>, 2024.
- 680 Yan, D., Zhang, T., Su, J., Zhao, L. L., Wang, H., Fang, X. M., Zhang, Y. Q., Liu, H. Y., and Yu, L. Y.: Structural Variation in



- the Bacterial Community Associated with Airborne Particulate Matter in Beijing, China, during Hazy and Nonhazy Days, *Appl. Environ. Microbiol.*, 84, e00004-00018, <https://doi.org/10.1128/AEM.00004-18>, 2018.
- Yang, T., Li, H., Xu, W., Song, Y., Xu, L., Wang, H., Wang, F., Sun, Y., Wang, Z., and Fu, P.: Strong Impacts of Regional Atmospheric Transport on the Vertical Distribution of Aerosol Ammonium over Beijing, *Environmental Science & Technology Letters*, 11, 29-34, <https://doi.org/10.1021/acs.estlett.3c00791>, 2023.
- 685
- Yao, Q., Ma, Z., Liu, J., Qiu, Y., Hao, T., Yao, L., Ding, J., Tang, Y., Cai, Z., and Han, S.: Effects of Meteorological Factors on the Vertical Distribution of Peroxyacetyl Nitrate in Autumn in Tianjin, *Aerosol and Air Quality Research*, 22, 220226, [10.4209/aaqr.220226](https://doi.org/10.4209/aaqr.220226), 2022.
- Yin, Y., Qi, J., Gong, J., and Gao, D.: Distribution of bacterial concentration and viability in atmospheric aerosols under various weather conditions in the coastal region of China, *Sci. Total Environ.*, 795, 148713, <https://doi.org/10.1016/j.scitotenv.2021.148713>, 2021.
- 690
- Yin, Z., Wan, Y., Zhang, Y., and Wang, H.: Why super sandstorm 2021 in North China?, *Natl Sci Rev*, 9, nwab165, <https://doi.org/10.1093/nsr/nwab165>, 2022.
- Zelles, L.: Fatty acid patterns of phospholipids and lipopolysaccharides in the characterisation of microbial communities in soil: a review, *Biol. Fertil. Soils*, 29, 111-129, <https://doi.org/10.1007/s003740050533>, 1999.
- 695
- Zhang, T. Y., Li, S., Zhu, Q. F., Wang, Q., Hussain, D., and Feng, Y. Q.: Derivatization for liquid chromatography-electrospray ionization-mass spectrometry analysis of small-molecular weight compounds, *Trac-Trends in Analytical Chemistry*, 119, 115608, <https://doi.org/10.1016/j.trac.2019.07.019>, 2019.
- Zhong, S., Zhang, L., Jiang, X., and Gao, P.: Comparison of chemical composition and airborne bacterial community structure in PM(2.5) during haze and non-haze days in the winter in Guilin, China, *Sci. Total Environ.*, 655, 202-210, <https://doi.org/10.1016/j.scitotenv.2018.11.268>, 2019.
- 700
- Zhu, Q. F., An, N., and Feng, Y. Q.: In-Depth Annotation Strategy of Saturated Hydroxy Fatty Acids Based on Their Chromatographic Retention Behaviors and MS Fragmentation Patterns, *Anal. Chem.*, 92, 14528-14535, <https://doi.org/10.1021/acs.analchem.0c02719>, 2020.

705