

We thank the two anonymous reviewers for their valuable comments and constructive suggestions on the manuscript. Below, we explain how the comments and suggestions are addressed and highlight revisions made in the revised manuscript. The reviewers' comments are in blue color. Our responses are in black, and our corresponding revisions in the manuscript are in red.

Reviewer #1

General Comments:

Particulate nitrate plays a critical role in aerosol radiative forcing and leads to very large uncertainty in climate projection. But very few global climate models have nitrate included, and only sparse studies evaluated the nitrate simulation performance. This work provides a comprehensive evaluation of the nitrate simulation in six global climate models against in-situ observations over North American, Australia, Japan and South Asia, Europe and Africa. This work not only evaluates the simulation of nitrate mass concentration at the surface level, but also for the vertical profile and gas-aerosol partitioning of nitrate. I think this work fits well with the scope of ACP. Although the scientific value is appreciated for this work, I would like to make a few suggestions that could help improve the presentation of this article for publishing in ACP.

Reply: We thank the reviewer for the encouraging comments.

Minor Comments:

1. IPCC AR5 2013 is referred. I recommend refer to the latest IPCC AR6 2021, to keep the research up to date.

Reply: Thank you for the suggestion. We did cite IPCC AR6 chapter 6 (i.e., Szopa et al., 2021) in the first paragraph for the important role that nitrate plays in the Earth's climate and air quality. Unfortunately, IPCC AR6 does not have an updated estimate of nitrate radiative forcing. In Figure 6.12 from AR6 chapter 6 (Szopa et al., 2021), nitrate is not included in the breakdown of individual contributions by aerosol species to the total ERF and the contribution of NO_x to the total ERF does not include nitrate either.

2. Line 75-77. This sentence is not clear. Please rephrase it.

Reply: We have rewritten two paragraphs in the introduction in response to a separate comment. The following revision addresses this comment:

“Previous studies found large relative differences (up to 150%) of nitrate concentrations at co-located desert and marine sites over the U.S. between the Clean Air Status and Trends Network (CASTNET) and the Interagency Monitoring of Protected Visual Environments (IMPROVE) (Ames and Malm, 2001; Sickles and Shadwick, 2008), which suggest significant fractions of coarse-mode nitrate ($\text{PM}_{>2.5}$, PM with diameter larger than $2.5\ \mu\text{m}$) related to heterogeneous reactions on dust and sea salt particles.”

3. Thanks for providing comprehensive review of previous studies in nitrate simulation, in the Introduction section. However, the presentation, or the writing style, would need some improvement. Rather than just listing what has been done previously (e.g., L80-90, and many other), but also better to summarize a key message you would like to express to audience.

Reply: We have rewritten the two paragraphs in the introduction section (L63-L89) to deliver clear key messages to readers:

“In the past decades, there are only a few regionally focused studies providing observational insights into the mass size distribution of nitrate aerosol, especially the significant contribution of coarse-mode nitrate. They found large relative differences (up to 150%) of nitrate concentrations at co-located desert and marine sites over the U.S. between the Clean Air Status and Trends Network (CASTNET) and the Interagency Monitoring of Protected Visual Environments (IMPROVE) (Ames and Malm, 2001; Sickles and Shadwick, 2008), which suggest significant fractions of coarse-mode nitrate ($\text{PM}_{>2.5}$, PM with diameter larger than $2.5\ \mu\text{m}$) related to heterogeneous reactions on dust and sea salt particles. Long-term measurements of $\text{PM}_{2.5}$ and PM_{10} (PM with diameter less than 2.5 and $10\ \mu\text{m}$, respectively) of nitrate surface concentrations at sites in Japan also show that $\text{PM}_{2.5-10}$ (PM with diameter between 2.5 and $10\ \mu\text{m}$) nitrate accounts for $\sim 50\%$ to $\sim 80\%$ of all PM_{10} nitrate (Khan et al., 2010; Itahashi et al., 2016; Uno et al., 2017; Pan et al., 2018). The same studies also showed that while $\text{PM}_{2.5}$ nitrate has a distinct winter-high-summer-low seasonal variation, the seasonal variation of $\text{PM}_{2.5-10}$ nitrate is relatively flat. Furthermore, dust induced $\text{PM}_{2.5-10}$ nitrate mainly contributes to $\text{PM}_{2.5-10}$ nitrate in spring, while sea salt induced $\text{PM}_{2.5-10}$ nitrate dominates in other seasons. Although recent studies attempted to compile surface observational data around globe for both fine and coarse aerosols (e.g., Mahowald et al., 2025), there remains a lack of observational constraints on the mass size distribution of nitrate aerosol, especially from a global view.

Previous regional modeling studies showed that including nitrate formation on coarse sea salt and dust particles through heterogenous reactions can significantly shift the mass size distribution of nitrate aerosol (e.g., Chen et al., 2020; Zhai et al., 2023), as it competes for

HNO₃ with formation of ammonium nitrate on fine particles. Chen et al. (2020) showed that heterogeneous reactions on sea salt shift mass size distribution of nitrate from fine to coarse mode, compared with an experiment turning off sea salt emission using WRF-Chem. The simulated mass size distribution of nitrate agrees well with measurements at Melpitz in Europe, where coarse-mode nitrate (PM_{>1.2}, PM with diameter larger than 1.2 μm) accounts for ~20% of total nitrate aerosol in marine air mass in September. Zhai et al. (2023), comparing simulated nitrate concentrations from GEOS-Chem with observations (PM₁ and PM₄, PM with diameter less than 1 and 4 μm, respectively) from the Korea-United States Air Quality (KORUS-AQ) campaign, found that including heterogeneous reactions on anthropogenic coarse particulate matter, mainly composed of anthropogenic dust, significantly reduces the overestimation of fine-mode nitrate in previous versions of GEOS-Chem.

Finally, only a few global modeling studies have evaluated the spatiotemporal distribution of fine-mode nitrate in GCMs against observations. There remains a lack of comprehensive analysis on the mass size distribution of nitrate aerosol (both fine and coarse mode nitrate) in GCMs. Mezuman et al. (2016) evaluated vertical profiles of PM₁ nitrate simulated in the GISS model against measurements from 14 aircraft campaigns and found systematic underestimation of nitrate over US and Europe. Bian et al. (2017) compared vertical profiles of fine-mode nitrate (PM with diameter less than 1 or 2.5 μm) simulated in AeroCom phase III models with measurements from the Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) campaign and found the models tend to underestimate fine-mode nitrate below 4 km. Some previous studies compared PM_{2.5} nitrate surface concentrations simulated in GCMs with IMRPOVE observations over the U.S. (e.g., Skeie et al., 2010; Bellouin et al., 2011; Hauglustaine et al., 2014; Zaveri et al., 2021; Lu et al., 2021). More recently, Tsimpidi et al. (2024) evaluated the simulated PM₁ and PM_{2.5} nitrate surface concentrations in the EMAC model against PM₁ data from field campaigns and PM_{2.5} data from regional networks, respectively, over the past 20 years. This comparison provides a robust basis for assessing model performance in capturing long-term trends and variability of nitrate across key regions in the northern hemisphere.”

4. “MOSAIC explicit treats the heterogeneous reactions on HNO₃ on dust and sea salt particles ...”. What about particles of other species? Are reactions treated independently for each size mode, or a bulk treatment for particles spanning all modes?

Reply: In the MAM4 scheme, aerosol species are assumed to be internally mixed within modes and externally mixed among modes. Reactions are treated independently for each size mode. We revised the description in section 2.1.1 to clarify it.

“In MAM4, aerosol species are assumed to be internally mixed within modes and externally mixed among modes. Wu et al. (2022) added nitrate (NO_3^-) and ammonium (NH_4^+) aerosol to the Aitken, accumulation, and coarse modes of MAM4. MOSAIC explicitly and independently treats the heterogeneous reactions of HNO_3 on particles containing dust (i.e., CaCO_3) and/or sea salt (i.e., NaCl) in the Aitken, accumulation, and coarse modes.”

5. Line-150. Why? Or you may want citation here to support your argument.

Reply: The equilibrium timescale for supermicron particles exceeds the typical GCM timestep, which is usually less than one hour. TEQMs does not account for the kinetic limitation of coarse particles and, therefore, may lead to an overestimation of nitrate in the coarse mode.

We have rewritten that part to address this comment:

“Previous studies showed that supermicron (coarse) nitrate particles take significantly longer, ranging from several hours to days, to reach equilibrium between the gas (HNO_3) and particles phase than submicron (fine) particles that equilibrate within minutes (Meng and Seinfeld, 1996; Fridlind and Jacobson, 2000). This equilibrium timescale for supermicron particles exceeds the typical GCM timestep, which is usually less than one hour. TEQMs assume instantaneous equilibrium between gas and particle phases with each timestep, which does not account for the kinetic limitation of coarse particles and therefore may lead to an overestimation of nitrate in the coarse mode.”

6. last line of page-6. Change to “with 1 year spin-up and the last 10-year results for analysis”.

Reply: Revised.

7. Line 184. Should be Table-2?

Reply: Corrected.

8. In the discussion, where possible, better to briefly engage some discussion about ammonia, sulfate, sea-salt and dust simulations. Since, this could help point towards clearer directions for improving nitrate simulation. Excessive ammonia after neutralizes sulfuric acid can then contribute to particulate nitrate.

Reply: We have added the following discussion in the conclusion:

“Most models have negative biases in all four regions. The model differences in gas-aerosol partitioning and large uncertainties in the simulated lifecycle of dust and sea salt contribute to the uncertainties in simulating mass size distribution of nitrate. E3SMv2, EMEP, and INCA slightly overestimate the annual mean surface nitrate $\text{PM}_{2.5}/\text{PM}_{10}$ ratio, but they still have negative biases for considerable number of sites.”

“The observed surface molar ratios at sites dominated by fine-mode nitrate show strong seasonal variations (summer-low-winter-high) with maximum values between 0.5 and 0.8 during winter and minimum values around 0.2 during summer. There are large spreads in the modeled surface molar ratios at the selected sites, which is caused by not only the model differences in gas-aerosol partitioning but also the large uncertainties in modeled HNO_3 or total nitrate ($\text{NO}_3^- + \text{HNO}_3$). Multiple processes such as gas phase chemistry (O_3 - NO_x - HO_x chemistry or N_2O_5 hydrolysis) and wet removal of HNO_3 can also greatly affect the abundance of HNO_3 and therefore change the molar ratio. In addition to those processes, sulfate aerosol formation as well as other processes can affect the abundance of free NH_3 available to react with HNO_3 and form particulate ammonium nitrate.”

“All models capture the seasonal variations to varying degrees but have larger spread in the modeled molar ratio in winter than in summer at the sites dominated by fine-mode nitrate. Differences in temperature, precipitation and sulfate concentrations, which favor nitrate formation in winter, and different dominant nitrate formation pathways contribute to the different model spread between winter and summer.”

Reviewer #2

General Comments:

This manuscript focuses on characterizing the ability of atmospheric models to capture the mass size distribution of nitrate aerosols, by comparing a set of models with relevant observations from ground-based networks and aircraft campaigns. Simulating nitrate aerosols remains challenging, and this work is a valuable contribution for the modeling community in the direction of better constraining the representation of nitrate. The paper is within the scope of ACP, well-structured and written, and I recommend publication in ACP. I would like to offer some recommendations to improve the clarity and presentation of this article for publication:

Reply: We thank the reviewer for the encouraging comments.

Major Comments:

Section: 2.1 Model description

1. The E3SMv2 and CESM2 models couple MOSAIC with MAM4, however, when reading, it is not clear which nitrate relevant processes are treated by MOSAIC and which by MAM4. Can you add more detail?

Reply: The implementation of MOSAIC in E3SMv2 and CESM2 replaces the default MAM4 treatment of gas-aerosol exchange between gases, including H_2SO_4 , HNO_3 , NH_3 , HCl , and a single lumped secondary organic aerosol (SOA) precursor, and aerosols.

To improve the clarity, we have now added the following description in Section 2.1.1 to give more details.

“Wu et al. (2022) implemented the Model for Simulating Aerosol Interactions and Chemistry (MOSAIC) module (Zaveri et al., 2008) in E3SMv2 and coupled it with the Model for Ozone and Related chemical Tracers gas chemistry (MOZART-4) (Emmons et al., 2010; Tilmes et al., 2015) and an enhanced version of the four-mode version of Modal Aerosol Module (MAM4) (Liu et al., 2016; Wang et al., 2020) following previous model development effort in CESM2 (Lu et al., 2021; Zaveri et al., 2021). MOSAIC is a comprehensive aerosol chemistry module which simulates the dynamic partitioning between semivolatile gases and particles of different sizes in an accurate but computationally efficient way. MOSAIC implemented in EAMv2 replaces the default MAM4 treatment of gas-aerosol exchange between gases, including H_2SO_4 , HNO_3 , NH_3 , HCl , and a single lumped secondary organic aerosol (SOA) precursor, and aerosols. The aqueous chemistry (i.e., occurring with cloud water) is also modified to include reactions of HNO_3 , NH_3 , and HCl .”

2. Is the size bins treatment in MOSAIC the same in E3SMv2 and CESM2, in terms of number and size range of bins for nitrate? I would suggest to explicitly add these information in the models description, since (mass) size distribution is a key focus of the paper. Also adding some additional information on how nitrate size distribution is treated in the other AeroCom models would be beneficial (maybe as a supplementary Table). This is important context also for the cutoff assumptions when comparing with observations.

Reply: The geometric standard deviations (σ_g) of particle sizes in the accumulation and coarse mode of MAM4 are different between CAM6 and EAMv2, which has significant impacts on the lifecycle of dust through dry deposition (Wu et al., 2020). The upper and lower bound of the number median diameter in the three modes are also different between CAM6 and EAMv2.

We have added details on the model treatment of aerosol size bins or modes containing nitrate to Table 1 for E3SMv2, CESM2, and the AeroCom models.

Table 1: Nitrate chemical mechanisms and physical properties.

Model	Gas-aerosol partitioning method for ammonium nitrate (fine and coarse)	Gas chemistry	DU/SS-nitrate treatment (fine and coarse)	Bins/modes for nitrate	Resolution
E3SMv2	DYN (MOSAIC) (Zaveri et al., 2008; Wu et al., 2022)	MOZART-4 (Emmons et al. 2010)	DYN	Aitken, accumulation, and coarse modes ($\sigma_g=1.6, 1.8$, and 1.8)	1° , 72
CESM2	DYN (MOSAIC) (Zaveri et al., 2008, 2021; Lu et al., 2021)	MOZART-TS1 (Emmons et al. 2020)	DYN	Aitken, accumulation, and coarse modes ($\sigma_g=1.6, 1.6$, and 1.2)	$1.25^\circ \times 0.9^\circ$, 56
EMAC	TEQM (ISORROPIA-II) (Fountoukis and Nenes, 2007)	MECCA (Sander et al., 2011)	TEQM	nucleation, Aitken, accumulation, and coarse modes	$2.8^\circ \times 2.8^\circ$, 31
EMEP	TEQM (MARS) (Saxena et al., 1986)	EmChem09 (Simpson et al., 2012)	first-order loss	fine and coarse modes ($D_p=0.33$ and $3.0 \mu\text{m}$; $\sigma_g=1.8$ and 1.6)	$0.5^\circ \times 0.5^\circ$, 20
GMI	TEQM (RPMARES) (Saxena et al., 1986)	Strahan et al., 2007	first-order loss	three bins ($D_p<0.1, 0.1-2.5, >2.5 \mu\text{m}$)	$2.5^\circ \times 2^\circ$, 72
INCA	TEQM (INCA) (Hauglustaine et al., 2004)	Hauglustaine et al., 2004; Folberth et al., 2006	first-order loss	fine and coarse modes	$1.9^\circ \times 3.75^\circ$, 39
OsloCTM2	TEQM (EQSAM_v03d) (Metzger and Leliveld, 2007)	Berntsen and Isaksen, 1997	TEQM (only SS)	fine ($D_p=0.1 \mu\text{m}$; $\sigma_g=2.0$) and coarse modes	$2.8^\circ \times 2.8^\circ$, 60

Note. DYN: dynamic mass transfer; DU: dust; SS: sea salt.

We have also added the following discussion:

“Nitrate and ammonium aerosol are explicitly simulated in the Aitken, accumulation, and coarse modes. However, as shown in Table 1, the geometric standard deviations (σ_g) in the accumulation and coarse mode of MAM4 are different between CAM6 and EAMv2, which has significant impacts on the lifecycle of dust through dry deposition (Wu et al., 2020). The upper and lower bound of the number median diameter in the three modes are also different between CAM6 and EAMv2.”

“AeroCom models have different treatments for the size distribution of nitrate, which affects the calculation of nitrate concentration at cut off size.”

Section 2.3 Observations

3. Ground based observations: are the measurements of nitrate assumed to be dry mass (RH=0%) or some level of hydration is taken into account in the measured mass? In particular, are these assumptions the same for CASTNET and IMPROVE, and if not, which source of error can this introduce in the PM_{2.5}/PM₁₀ mass ratio, given that nitrate is strongly hydrophilic?

Reply: The measurements of nitrate are reported as dry mass. We believe that the water vapor should have limited impact on the measurements of nitrate. Both CASTNET and IMPROVE use ion chromatography (IC) to analyze the extract from the Teflon and nylon filter, respectively, instead of measuring gravimetric mass of the material collected on the filters (before and after sampling). The material collected on the filters is extracted by extraction solution and subsequently analyzed for nitrate anions by a Dionex IC. We have discussed the main differences between CASTNET and IMPROVE sampling protocols and their impacts on biases in estimating PM_{2.5}/PM₁₀ ratios in section 2.3.

Aircraft observations: as acknowledged, not all the simulated periods in the models overlap with of aircraft campaigns deployment, thus comparison need to be treated cautiously. Can you briefly add which sources of uncertainty this might introduce and are they stronger at different heights (e.g. above/below PBL)?

Reply: The differences in the anthropogenic, biomass burning, and dust emissions between the simulation period and the observation period may contribute to the uncertainty of the comparison. Those differences in emissions may have large impact on

the simulated mass size distribution of nitrate in the middle to upper troposphere over remote oceans, regions sampled during the ATom campaigns.

We have added a few words to clarify this.

“The model biases in fine-mode nitrate and nitrate PM_1/PM_4 ratio, when compared to aircraft measurements, should be interpreted with caution due to discrepancies, particularly in the anthropogenic, biomass burning, and dust aerosol emissions, between the simulation period and the observation period. Those differences in emissions may have stronger impact on the simulated mass size distribution of nitrate in the middle to upper troposphere than in the lower troposphere over remote oceans (e.g., ATom), as sea salt is dominant in the marine boundary layer and fine sulfate/carbonaceous aerosols and coarse dust aerosols are dominant in the middle to upper troposphere (Thompson et al., 2021).”

Section 3 Results

5. Figures 4, 5, 9, 10: For added clarity in the vertical distribution plots, I suggest to plot observations on top of all the model lines, as done for the surface concentrations plots (e.g. fig 7).

Reply: Following the great suggestion, we have revised the figures to plot observations on top of all model lines to improve the clarity.




Figure 4: Vertical profiles of fine-mode nitrate concentrations ($\mu\text{g m}^{-3}$ in STP) from model simulations (colored lines) and PM_{10} nitrate concentrations from five aircraft campaigns: (a-c) ARCTAS, (d-e) DC3, (f-g) SEAC⁴RS, (h-i) WINTER, and (j-k) KORUS-AQ. Black dots denote mean values of observations with one standard deviation on each side marked in grey lines. The numbers in each panel are median concentrations.




Figure 5: Vertical profiles of fine-mode nitrate concentrations ($\mu\text{g m}^{-3}$ in STP) from model simulations (colored dots) and PM₁ nitrate concentrations from ATom 1-4 campaigns (with black dots for mean values and grey lines marking one standard deviation of observations). The numbers in each panel are median concentrations.




Figure 9: Vertical profiles of nitrate PM_1/PM_4 ratios from model simulations (colored dots) and five aircraft campaigns: (a-c) ARCTAS, (d-e) DC3, (f-g) SEAC⁴RS, (h-i) WINTER, and (j-k) KORUS-AQ (black dots).




Figure 10: Same as Figure 9 but for the eight regions (marked by boxes in Figure 2) during ATom 1-4 campaigns.

Section 4 Discussion and Conclusion

6. L481-517: This first part of the discussion just states the different results of the model observation comparison but doesn't explicitly links them to the uncertainty in the representation of underlying processes. E.g. L516: All models capture the seasonal variations to varying degrees but have larger spread in the modeled molar ratio in winter. Why? Which processes are causing the model spread particularly in winter? This part would benefit from an additional discussion on the potential sources of uncertainty in the modeled processes that may be driving the observed biases in the comparison.

Reply: Further data analysis has revealed large spreads in modeled HNO_3 surface concentrations. The large spreads in the modeled surface molar ratios at the selected sites is caused by not only the model differences in in gas-aerosol partitioning but also the large uncertainties in modeled HNO_3 or total nitrate ($\text{NO}_3^- + \text{HNO}_3$). Multiple processes such as gas phase chemistry ($\text{O}_3\text{-NO}_x\text{-HO}_x$ chemistry or N_2O_5 hydrolysis) and wet removal of HNO_3 can also greatly affect the abundance of HNO_3 and therefore change the molar ratio. In addition to those processes, sulfate aerosol formation as well as other processes can affect the abundance of free NH_3 available to react with HNO_3 and form

particulate ammonium nitrate.

Although the model spread of the simulated HNO_3 at these sites is large across all seasons, the colder temperatures, lower precipitation, and lower sulfate concentrations at the same locations in winter promote more favorable conditions for nitrate formation compared to summer. Also, as indicated by Fig. 8, the nitrate formation pathway in winter is mainly through thermodynamic interactions between HNO_3 and NH_3 in most models, while nitrate formation through heterogeneous reactions on coarse dust and sea salt also tends to have significant contribution in summer in many models.

We have added the following discussions in the conclusion:

“The observed surface molar ratios at sites dominated by fine-mode nitrate show strong seasonal variations (summer-low-winter-high) with maximum values between 0.5 and 0.8 during winter and minimum values around 0.2 during summer. There are large spreads in the modeled surface molar ratios at the selected sites, which is caused by not only the model differences in gas-aerosol partitioning but also the large uncertainties in modeled HNO_3 or total nitrate ($\text{NO}_3^- + \text{HNO}_3$). Multiple processes such as gas phase chemistry (O_3 - NO_x - HO_x chemistry or N_2O_5 hydrolysis) and wet removal of HNO_3 can also greatly affect the abundance of HNO_3 and therefore change the molar ratio. In addition to those processes, sulfate aerosol formation as well as other processes can affect the abundance of free NH_3 available to react with HNO_3 and form particulate ammonium nitrate. All models capture the seasonal variations to varying degrees but have larger spread in the modeled molar ratio in winter than in summer at the sites dominated by fine-mode nitrate. Differences in temperature, precipitation and sulfate concentrations which favor nitrate formation in winter and different dominant nitrate formation pathway contribute to the different model spread between winter and summer.”

We have also added some detailed discussions in the section 3.3:

“Figures 12 and 13 show the seasonal variations of modeled surface $\text{NO}_3^-/(\text{NO}_3^- + \text{HNO}_3)$ molar ratio in comparison with observations at the same selected sites in Figs. 7 and 8, respectively. In general, there are large spreads in the modeled surface molar ratios. The large spreads in modeled HNO_3 surface concentrations (Figs. S7 and S8) partially contribute to the large uncertainties in the modeled surface molar ratios, which can be further related to model differences in multiple processes, such as gas-aerosol partitioning between nitrate aerosol and gas phase HNO_3 , sulfate aerosol formation, gas phase chemistry (e.g., He et al., 2015), and wet removal of HNO_3 (e.g., Luo et al., 2019). Comparisons of nitrate surface concentration with CASTNET measurements are also provided in the supplement (Figs. S9 and S10).”

“These factors can contribute to the high molar ratios and weak seasonal variations at the

two sites. OsloCTM2 and OsloCTM3 have positive biases in the modeled surface molar ratios at the two marine sites, which is mainly due to their gas-aerosol partitioning methods (only TEQMs). However, EMAC has negative biases at the two sites, which is mainly due to its significant positive biases in HNO_3 .”

“All models capture the seasonal variations to varying degrees and have larger spread in the modeled molar ratio in winter, ranging from below 0.2 to above 0.8, than in summer. Although the model spread in the simulated HNO_3 at these sites is large across all seasons, the colder temperatures, lower precipitation, and lower sulfate concentrations at the same locations in winter promote more favorable conditions for nitrate formation compared to summer. Also, as indicated by Fig. 8, the nitrate formation pathway in winter is mainly through thermodynamic interactions between HNO_3 and NH_3 in most models, while nitrate formation through heterogeneous reactions on coarse dust and sea salt particles also tends to have a significant contribution in summer in many models.”

For conciseness, we have put the related figures in supplement.




Figure S7. Same as Figure 7 but for HNO_3 surface concentrations ($\mu\text{g m}^{-3}$).




Figure S8. Same as Figure 8 but for HNO₃ surface concentrations (μg m⁻³).




Figure S9. Same as Figure 7 but for nitrate surface concentrations ($\mu\text{g m}^{-3}$).




Figure S10. Same as Figure 8 but for nitrate surface concentrations ($\mu\text{g m}^{-3}$).

7. L521-523: the results suggest that coarse mode nitrate is better captured when representing heterogeneous formation on sea salt and dust as first order loss approximation or dynamic mass transfer. I know the models setup is very different among models, but can you derive any conclusion from the results in terms of which method seems to be better capturing fine mode nitrate (simple TEQM, more complex TEQM, dynamic mass transfer)?

Reply: It is difficult to derive any conclusion in terms of which gas-aerosol partitioning method (TEQM only, TEQM combined with first order loss approximation, and dynamic mass transfer) lead to smaller model biases in simulating fine-mode nitrate.

In Figure 3, OsloCTM2 and OsloCTM3, which use only TEQMs, have similar negative biases in the annual mean $\text{PM}_{2.5}$ nitrate surface concentration averaged across all sites as GMI and INCA, which use TEQMs with first order loss approximation. EMAC, which uses only TEQM but considers the kinetic limitation, has the smallest biases in the annual mean $\text{PM}_{2.5}$ nitrate surface concentration among all models. OsloCTM2 and OsloCTM3 have much larger negative biases than other models at the Australian, South African, and Japanese sites, which is consistent with that the two models have much larger negative

biases in $\text{PM}_{2.5}/\text{PM}_{10}$ at the South African and Japanese sites.

In Figures 4 and 5, OsloCTM2 and OsloCTM3 give much lower fine-mode nitrate than other models and have significant negative biases compared with all aircraft campaigns. However, some models (e.g., INCA and EMAC) have much larger biases (positive) than OsloCTM2 and OsloCTM3.

We have also revised the text in the conclusion section to add implications on simulating fine-mode nitrate:

“Most of the AeroCom models underestimate PM_1 nitrate concentrations below 600 hPa compared to the ARCTAS, DC3, SEAC⁴RS, and KORUS-AQ campaigns. OsloCTM2 and OsloCTM3, which both use only TEQMs for the formation of coarse-mode nitrate, have uniformly significant negative biases in fine-mode nitrate concentration. EMAC, which uses only TEQM but considers the kinetic limitation, has the smallest biases in the annual mean $\text{PM}_{2.5}$ nitrate surface concentration among all models.”

“Our analysis suggests that future studies and model development efforts should better represent heterogeneous reactions of nitrate formation on coarse dust and sea salt particles and use first-order loss approximation with TEQMs or dynamic mass transfer approach for gas-aerosol partitioning between nitrate aerosol and HNO_3 gas, as all models that use only TEQMs have larger biases than the other models. Using TEQMs only may also significantly underestimate fine-mode nitrate compared with the other two methods.”

Minor Comments:

1. L86-89: “Zhai et al (2023)...found that including anthropogenic coarse particulate matter mainly composed of anthropogenic dust, significantly reduces the overestimation of fine-mode nitrate in previous versions of GEOS-Chem” add what process was responsible for this decrease (less HNO_3 available for fine mode nitrate?)

Reply: Including nitrate formation through heterogeneous reactions on coarse dust particles competes for HNO_3 with formation of ammonium nitrate on fine particles and therefore improves the agreement with observations. We have rewritten this part to address this comment:

“Previous regional modeling studies show that including nitrate formation on coarse sea salt and dust particles through heterogenous reactions can significantly shift the mass size distribution of nitrate aerosol (e.g., Chen et al., 2020; Zhai et al., 2023), as it competes for HNO_3 with formation of ammonium nitrate on fine particles. Chen et al. (2020) showed

that heterogeneous reactions on sea salt shift mass size distribution of nitrate from fine to coarse mode compared with an experiment turning off sea salt emission using WRF-Chem. The simulated mass size distribution of nitrate agrees well with measurements at Melpitz in Europe, where coarse-mode nitrate ($PM_{>1.2}$, PM with diameter larger than 1.2 μm) accounts for ~20% of total nitrate aerosol in marine air mass in September. Zhai et al. (2023) compared simulated nitrate concentrations from GEOS-Chem with observations (PM_1 and PM_4 , PM with diameter less than 1 and 4 μm , respectively) from the Korea-United States Air Quality (KORUS-AQ) campaign and found that including heterogeneous reactions on anthropogenic coarse particulate matter, mainly composed of anthropogenic dust, significantly reduces the overestimation of fine-mode nitrate in previous versions of GEOS-Chem.”

2. Table 1: I would add for additional clarity which nitrate mode (fine/coarse) is treated in by the 'Gas-aerosol partitioning method'(fine only) and the 'DU/SS-nitrate treatment' (fine+coarse).

Reply: We have changed “Gas-aerosol partitioning method” to “Gas-aerosol partitioning method for ammonium nitrate (fine and coarse)” and “DU/SS-nitrate treatment” to “DU/SS-nitrate treatment (fine and coarse)” in Table 1 for additional clarity.

3. L22-225: for E3SMv2 and CESM2, are monthly output from 2014-only the one chosen for comparison to WINTER, KORUS-AQ and Atom? Please specify the sampling year of model output.

Reply: We have added a few words to clarify it.

“We use the corresponding monthly mean model results for the aircraft campaigns operated during the simulation period. Note that all aircraft campaigns except for ARCTAS are outside the model simulation year (2008) of AeroCom phase III models. For comparisons with WINTER, KORUS-AQ, and ATom campaigns that are not within the simulation period (2005-2014) of E3SMv2 and CESM2, we use the ten-year average of the corresponding month from the two models.”

4. Table 3: Is there a reason why you have chosen to report the data in terms of moles instead of mass? For most studies (like in Bian et al.2017) they are expressed in terms of mass (Tg, mass mixing ratio).

Reply: We are not reporting the data of nitrate and HNO_3 in mol. We only report the

tropospheric burden ratio of $\text{NO}_3^-/(\text{NO}_3^- + \text{HNO}_3)$ and the burden ratio of $\text{PM}_{2.5}/\text{PM}_{10}$ nitrate in mol mol^{-1} which is essentially unitless and can better represent the partitioning of between reactive nitrogen species. We report $\text{PM}_{2.5}$ nitrate, total nitrate, and tropospheric HNO_3 burden in mass of N (Tg N) which is widely used in literature and comparable to other studies (e.g., Hauglustaine et al., 2014; Zaveri et al., 2021).

References

He, J., Zhang, Y., Tilmes, S., Emmons, L., Lamarque, J.-F., Glotfelty, T., Hodzic, A., and Vitt, F.: CESM/CAM5 improvement and application: comparison and evaluation of updated CB05_GE and MOZART-4 gas-phase mechanisms and associated impacts on global air quality and climate, *Geosci. Model Dev.*, 8, 3999–4025, <https://doi.org/10.5194/gmd-8-3999-2015>, 2015.

Luo, G., Yu, F., and Schwab, J.: Revised treatment of wet scavenging processes dramatically improves GEOS-Chem 12.0.0 simulations of surface nitric acid, nitrate, and ammonium over the United States, *Geosci. Model Dev.*, 12, 3439–3447, <https://doi.org/10.5194/gmd-12-3439-2019>, 2019.

Mahowald, N. M., Li, L., Vira, J., Prank, M., Hamilton, D. S., Matsui, H., Miller, R. L., Lu, P. L., Akyuz, E., Meidan, D., Hess, P., Lihavainen, H., Wiedinmyer, C., Hand, J., Alaimo, M. G., Alves, C., Alastuey, A., Artaxo, P., Barreto, A., Barraza, F., Becagli, S., Calzolari, G., Chellam, S., Chen, Y., Chuang, P., Cohen, D. D., Colombi, C., Diapouli, E., Dongarra, G., Eleftheriadis, K., Engelbrecht, J., Galy-Lacaux, C., Gaston, C., Gomez, D., González Ramos, Y., Harrison, R. M., Heyes, C., Herut, B., Hopke, P., Hüglin, C., Kanakidou, M., Kertesz, Z., Klimont, Z., Kyllönen, K., Lambert, F., Liu, X., Losno, R., Lucarelli, F., Maenhaut, W., Marticorena, B., Martin, R. V., Mihalopoulos, N., Morera-Gómez, Y., Paytan, A., Prospero, J., Rodríguez, S., Smichowski, P., Varrica, D., Walsh, B., Weagle, C. L., and Zhao, X.: AERO-MAP: a data compilation and modeling approach to understand spatial variability in fine- and coarse-mode aerosol composition, *Atmos. Chem. Phys.*, 25, 4665–4702, <https://doi.org/10.5194/acp-25-4665-2025>, 2025.

Tsimpidi, A. P., Scholz, S. M. C., Milousis, A., Mihalopoulos, N., and Karydis, V. A.: Aerosol Composition Trends during 2000–2020: In depth insights from model predictions and multiple worldwide observation datasets, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-3590>, 2024.