

From column to surface: connecting the performance in simulating aerosol optical properties and PM_{2.5} concentrations in the NASA GEOSCCM

Caterina Mogno^{1,2}, Peter R. Colarco², Allison B. Collow^{1,3}, Sampa Das^{2,4}, Sarah A. Strode^{2,5}, Vanessa Valenti^{6,7}, Michael E. Manyin^{2,6}, Qing Liang², Luke Oman², Stephen D. Steenrod^{1,2}, K. Emma Knowland^{3,5,8}

¹ Goddard Earth Sciences Technology and Research II (GESTAR II), University of Maryland Baltimore County, Baltimore, Maryland, USA

² Atmospheric Chemistry and Dynamics Lab, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

³ Global Modeling and Assimilation Office, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

⁴ Earth System Science Interdisciplinary Center (ESSIC), University of Maryland, College Park, Maryland, USA

⁵ Goddard Earth Sciences Technology and Research II (GESTAR II), Morgan State University, Baltimore, Maryland, USA

⁶ Science Systems and Applications, Inc., Lanham, Maryland, USA

⁷ Computational and Information Sciences and Technology Office, NASA Goddard Space Flight Center Greenbelt, Maryland, USA

⁸ now at: NASA Headquarters, Washington, District of Columbia, USA

Correspondence to: Caterina Mogno (caterina.mogno@nasa.gov)

Abstract. Aerosols are a key climate forcer and harmful to human health at the surface. Accurately modeling aerosol optical properties, mass loading and their relationship is important for constraining aerosol-climate forcing and characterizing particulate matter pollution exposure. We investigate the drivers of uncertainties in the NASA Goddard Earth Observing System Chemistry Climate Model (GEOSCCM) in simulating aerosols by focusing on the link between aerosol optical properties and mass. We compare a GEOSCCM hindcast with long-term coincident observations including satellite AOD measurements, speciated PM_{2.5} datasets from observations-model data fusion, and ground-based measurements of aerosol mass and optical properties. We analyze regional trends and seasonal variations of AOD and PM_{2.5}, and surface aerosol properties, including relative humidity's role in hygroscopic enhancement. This work also presents the first extensive assessment of GEOSCCM's aerosol component with observational data. Our findings show that biases in PM_{2.5} components and relative humidity significantly impact simulated aerosol scattering at the surface, while scattering efficiency assumptions align with observations. This indicates that errors in simulated scattering relate more to simulated aerosol **speciated** mass and relative humidity than optical properties and size distribution assumptions in GEOSCCM. Our work highlights the importance of relative humidity biases on aerosol scattering enhancement for climate models where meteorology is not prescribed. Findings suggest improvements in GEOSCCM aerosols mass and optical properties could be achieved through updating emission inventories, especially over biomass burning regions, reducing nitrate biases, and improving relative humidity simulation.

35 **1 Introduction**

Aerosols are an important climate forcer and are harmful pollutants to human health. Aerosols impact climate directly through scattering and absorption of solar radiation (aerosol-radiation interaction) and indirectly by acting as nuclei for cloud droplet and ice particle formation (aerosol-cloud interactions) (IPCC, 2023). Uncertainties in aerosol properties, distributions, and processes are the main driver of uncertainty in the overall anthropogenic forcing of the climate system, with impacts of aerosol-radiation and aerosol-clouds interactions remaining one of the largest drivers of uncertainty in anthropogenic climate forcing (IPCC, 2023; Kahn et al., 2023; Li et al., 2022). Close to Earth's surface, exposure to ambient fine particulate matter, i.e. aerosols with diameter less than 2.5 μm ($\text{PM}_{2.5}$), is a leading contributor to the global burden of disease. It has been associated with short-term acute health effects and long-term chronic respiratory and cardiovascular diseases, accounting for more than 4 million premature deaths per year (Brauer et al., 2024; Fuller et al., 2022).

45 The ability of atmospheric models to correctly represent aerosols and their properties is thus fundamental for constraining aerosol climate forcing and projecting the impacts of changing emissions of aerosols and trace gases on the state of the climate, as well as for characterizing the distribution and changes in particulate matter pollution exposure.

The interaction of atmospheric aerosols with radiation is complex and depends on factors such as the aerosol mass, composition, size, shape, mixture, and hygroscopicity (IPCC, 2023). Current atmospheric models generally account for the aerosol-radiation interaction by pre-calculating aerosol extinction, scattering and absorption of radiation and their dependence on relative humidity through look-up optical tables (LUT), which are used to convert simulated aerosol mass to optical quantities. Recently, atmospheric models have been utilized in combination with satellite measurements of aerosol optical depth (AOD) to produce satellite-derived estimates of surface $\text{PM}_{2.5}$ (van Donkelaar et al., 2019, 2021; Hammer et al., 2020; Xu et al., 2015). The linking of AOD (an optical column measurement) to surface concentrations remains a key uncertainty in these derivations, because it is addressed by using model-based assumptions of the $\text{PM}_{2.5}$ /AOD ratio, which is subject to further assumptions in the model representation of aerosol processes, properties, vertical distribution and hygroscopic enhancement (Zhai et al., 2021). Inaccuracies in modeled $\text{PM}_{2.5}$ /AOD ratios can propagate into errors in satellite-derived surface $\text{PM}_{2.5}$ estimates, especially in regions with complex aerosol mixtures or vertical distribution (Zhu et al., 2024). In particular, recent aerosol model intercomparison and evaluation efforts highlight the need to investigate model biases by looking beyond standard aerosol global optical depth (AOD) evaluations and incorporating additional analysis such as the characterization of regional and seasonal variations, aerosol hygroscopic growth, and whether model biases in optical depth are linked to biases in aerosol mass or optical property assumptions (Gliß et al., 2021).

In this context, the use of simultaneous observations from various sources are essential to improve our representation of modeled aerosol-radiation interactions and of particulate matter. The expanding atmosphere observing system, which includes satellite, aircraft and ground-based platforms, can provide a wide range of long-term, complementary and coincident aerosol measurements. The simultaneous use of these observations can enhance our understanding and interpretation of observed and modeled aerosol burdens, the relationships between their optical, physical, and chemical properties (Bharath et al., 2024; Kahn

et al., 2023; Mortier et al., 2020). Specifically, co-located aerosol chemistry and size data, relative humidity, and light scattering measurements can be used to better constrain model aerosol optical property assumptions (Latimer and Martin 2019; Burgos et al. 2020).

In this work we investigate the drivers of uncertainties in the current NASA Goddard Earth Observing System Chemistry Climate Model (GEOSCCM) in simulating aerosols by focusing on the link between aerosol optical properties and mass. We accomplish this by comparing a GEOSCCM hindcast with a variety of long-term coincident observations of aerosol, including satellite measurements of AOD, datasets of speciated $\text{PM}_{2.5}$ from observation-model data fusion products, and ground-based measurements of aerosol mass and optical properties. We focus our analysis on regional trends and seasonal variations of AOD and $\text{PM}_{2.5}$, and on the properties of aerosols at the surface. In particular, we examine speciated $\text{PM}_{2.5}$ and the consistency of GEOSCCM $\text{PM}_{2.5}$ simulation with optical properties, including the role of relative humidity in hygroscopic scattering enhancement, using surface observations of co-located long-term speciated $\text{PM}_{2.5}$, scattering coefficient, and relative humidity. This is also the first time that the GEOSCCM aerosol component is extensively benchmarked against observations.

In Section 2 we introduce the GEOSCCM model, the GOCART aerosol module and the simulation analyzed in this study. Section 3 describes the aerosol observation datasets used for evaluating GEOSCCM. In Section 4 we present our results. Finally, Section 5 present the final discussion and conclusions.

2 Model Description

2.1 GEOSCCM and the GOCART Aerosol Module

GEOSCCM is a configuration of the NASA Goddard Earth Observing System (GEOS) global atmospheric general circulation model (AGCM; Molod et al. 2015) with coupled and radiatively interactive atmospheric chemistry and aerosols. GEOSCCM has been extensively evaluated for stratospheric ozone photochemistry and transport processes in various model intercomparison studies (Eyring T., 2010; Morgenstern et al., 2017), and has been used in a variety of studies to investigate trends in stratospheric chemistry and dynamics (Li et al., 2018; Oman et al., 2008, 2010), tropospheric ozone (Liu et al., 2022; Strode et al., 2017, 2019), and impacts of aerosols on atmospheric chemistry and climate (Aquila et al., 2014; Case et al., 2024; Rollins et al., 2017). For this study we use GEOSCCM version Icarus-3_2_MEM_22x.

The GEOS AGCM uses a finite volume dynamical core on a cubed sphere based on (Putman and Lin, 2007). The model has 72 vertical layers, with hybrid-sigma coordinates, transitioning from terrain-following at the surface to pressure levels at 180 hPa and a model top at ~ 80 km (0.01 hPa). The convective scheme is based on a modified version of the relaxed Arakawa-Schubert parametrization (Moorthi and Suarez, 1992), while the turbulence parameterization is based on (Lock et al., 2000; Louis, 1979). Radiative processes are described by (Chou, M. and Suarez, M. J., 2001) for longwave radiation and (Chou, M.

and Suarez, M. J., 1999) for shortwave radiation. Fluxes at the land/atmosphere interface are determined using the Catchment Land Surface Model (Koster et al., 2000), while the surface layer is determined as in (Helfand and Schubert, 1995).

The current GEOSCCM model couples the Global Modeling Initiative (GMI) chemistry mechanism (Nielsen et al. 2017; Strahan et al., Duncan, and Hoor 2007; Duncan et al. 2007; Douglass et al. 2004) with the Goddard Chemistry, Aerosol, Radiation, and Transport (GOCART) aerosol module (Chin et al. 2014, 2002; Colarco et al. 2010). The GMI module includes 120 species and over 400 reactions, combining tropospheric and stratospheric chemistry.

GOCART treats externally mixed dust, sea salt, sulfate, nitrate, and black and organic carbon aerosol species. Dust and sea salt are each partitioned into five non-interacting size bins and have surface windspeed-dependent source functions. Black and organic carbon aerosols are partitioned into hydrophobic and hydrophilic modes, with anthropogenic and biomass burning sources for both and a biogenic source for organic aerosol. A constant ratio of 1.8 is assumed between organic mass and organic carbon (OM:OC). GOCART utilizes a reduced-complexity secondary organic aerosol (SOA) mechanism that relates volatile organic carbon (VOC) emissions to CO emissions across anthropogenic, biofuel and biomass burning sectors, as established by (Kim et al., 2015). The generated SOA is allocated to hydrophilic components of organic carbon. Additionally, biogenic sources of SOA are from an online version of the Model of Emissions of Gases and Aerosols from Nature (MEGAN). Throughout the rest of the paper, we refer to model OA as the organic aerosol contributions from all primary and secondary sources (biomass, biogenic, biofuel, and anthropogenic). The sulfur mechanism in GOCART includes tracers for dimethyl sulfide (DMS), sulfur dioxide (SO₂), sulfate aerosol (SO₄), and methanesulfonic acid (MSA). DMS has surface wind-speed dependent emissions based on sea water DMS concentrations. SO₂ has direct emissions from biomass burning, volcanic degassing and eruptions, and anthropogenic sources from aircraft, ships, and land-based pollution sources. Sulfate has direct emissions from ships but is otherwise a product of oxidation of precursor DMS and SO₂ species following the mechanism in (Chin et al., 1996). The nitrate mechanism is described in Huisheng Bian et al. 2017 and tracks ammonia (NH₃), bulk ammonium aerosol, and three sizes of nitrate aerosol. There are anthropogenic and biomass burning sources of ammonia. Ammonium and fine-mode nitrate production are calculated in an equilibrium thermodynamic module that balances water, sulfur, and nitrogen species. Coarse mode nitrate is produced from heterogeneous reaction of nitric acid (HNO₃) on sea salt and dust surfaces. GOCART aerosol optical properties are pre-computed for each aerosol tracer and saved in look up tables (LUTs). Optical properties are primarily based on Mie calculations with most parameters as in Chin et al. 2002. Non-spherical dust optical properties are as in Colarco et al. 2014. In the version of GEOSCCM evaluated here, GOCART aerosols provide surfaces for heterogeneous chemistry and impact photolysis rates in GMI. GMI oxidants and nitric acid are one-way coupled to GOCART sulfate and nitrate algorithms.

2.2 The Ref-D1 Experiment

The simulation analyzed in this study is a global hindcast run at performed with c90 (~1°) horizontal resolution covering the period 1960 – 2018. The simulation follows the protocol for the Chemistry Climate Model Initiative

130 (<https://blogs.reading.ac.uk/ccmi/>) “Ref-D1” experiment (IGAC/SPARC, REFD1, 2022) and was submitted as input to the
2022 WMO Scientific Assessment of Ozone Depletion (Scientific Assessment of Ozone Depletion: 2022, 2025). The
simulation is performed with a free running atmosphere and prescribed sea surface temperature (SST) and sea ice
concentrations (SICs) from the global HadISST1 data set as monthly mean boundary conditions (Rayner et al., 2003). The
Ref-D1 simulation includes anthropogenic emissions provided for the Coupled Modelling Intercomparison project Phase 6
135 (CMIP6), which are from the Community Emissions Data System (Hoesly et al., 2018) for years 1960-2014, and from the
CMIP6 ScenarioMIP experiment (SSP2-4.5 scenario) for the years 2015-2018 (Gidden et al., 2019). Biomass burning
emissions for years 1960-2015 are also from CMIP6, which are derived from harmonizing different emissions inventories and
models as described in (van Marle et al., 2017). In this dataset satellite-derived biomass-burning emissions from the Global
Fire Emissions Database version 4 with small fires (GFED4.1s) are used from 1997 to 2015 (Randerson, J.T et al., 2018). For
140 the remainder years not covered by CMIP6 emissions (2016 to




Figure 1: Global spatial distribution of emissions and chemical production of aerosol species in our Ref-D1 experiment, averaged from 1960-2018. SO₄ includes direct emissions + production, NO₃ refers to production. OA includes all organic mass (emissions + SOA production).




Figure 2. Ref-D1 regional annual timeseries of emissions and production of aerosols by species for the period 1960-2018. ANTH=anthropogenic, BB=biomass burning, BIO=biogenic. SO₄ includes direct emissions + production but no emissions from volcanic eruptions, NO₃ refers to production. OA includes all organic mass (emissions + production).

2018), biomass burning emissions are also obtained from GFED4.1. Both anthropogenic and biomass burning emissions include seasonal (monthly) and interannual variability. Finally, for natural emissions sources, biogenic emissions are calculated online in the model using the Model of Emissions of Gases and Aerosols from Nature (MEGAN) (Guenther et al., 2006). Dust and sea salt sources are wind driven following (Ginoux et al., 2001) and (Gong, 2003; Jaeglé et al., 2011), respectively. Volcanic emissions (eruptive + outgassing) follow (Carn et al., 2017). The model also uses prescribed species for aerosol chemistry not calculated by GM1: open-ocean emission of NH₃ (Bouwman et al., 1997) and dimethyl sulfide (DMS), whose concentrations are prescribed with seasonally varying climatology and emissions are calculated based on wind speed (Lana et al., 2011; Liss and Merlivat, 1986).

Because the Ref-D1 is not driven by real meteorology, we focus our analysis on monthly and yearly timescales. For regional analysis over land, we divide the globe in 8 geographical regions: North America, South America, Northern Africa, Southern Africa, Europe, Siberia, Southern Asia, Australia. The geographical boundaries for the region are reported in Supplemental Figure 1. This regional aggregation follows the approach from Collow et al. (2024), which was designed to capture continental-scale trends and behaviours in aerosol distributions. In particular, South Asia and East Asia are grouped together because they are both dominated by anthropogenic aerosol sources, particularly from major populated regions in India and China. While we acknowledge that there are important sub-regional differences within this large domain, the continental-scale analysis allows us to examine broad patterns in model performance across regions with similar dominant source characteristics.

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto, Ligatures: None

Formatted: Font: (Default) Times New Roman, Not Italic, Underline, Font color: Dark Red, Ligatures: None

Formatted: Font: (Default) Times New Roman, Not Italic, Underline, Font color: Dark Red, Ligatures: None

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto, Ligatures: None

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Dark Red, Ligatures: None

Regional statistics have been calculated as area-weighted averages of model grid-boxes falling within the boundary for each region. The mathematical definition of the metrics used for evaluating the model against observation, namely normalized mean bias (NMB), root mean square error (RMSE) and Pearson's correlation coefficient (r), are reported in the Supplement S1.

Deleted: ¶ ... [1]

Deleted: ¶ ... [2]




Figure 3: Ref-D1 regional annual timeseries of (a) total column AOD with (cyan dashed) and without (solid blue) the contribution of volcanic eruptions and (b) surface PM_{2.5} for the period 1960-2018.

2.3 Modelled long-term time-series of emissions, AOD and PM_{2.5}

Figure 1 presents the spatial distribution of different aerosol species in Ref-D1, while Figure 2 shows their regional timeseries. Timeseries of emissions by region are reported in Supplemental Figure 2. Emissions from biomass burning (OA_BB, BC_BB) shows higher variability and magnitude starting in the late 1990s, reflecting the use of satellite observations in informing biomass burning estimates from that point forward.

Figure 3 presents regional annual timeseries in Ref-D1 for total AOD and surface PM_{2.5}, while Supplemental Figure 3 show the timeseries for speciated AOD and PM_{2.5}. Spikes in regional AOD are driven by major volcanic events (e.g. Agung 1963, El Chicon 1982, Pinatubo 1991, Supplemental Figure 3a). Regional changes and trends in emissions are reflected in the regional AOD and surface PM_{2.5} trends. The decline in anthropogenic SO₄ AOD since the 1980s drives a steady decline in total AOD over Europe, and to a lesser extent over North America, where total AOD has been slowly decreasing. Over Siberia the decline in SO₄ AOD is counterbalanced by the increase of OA AOD from biomass burning since the early 2000s. On the other hand, increase in SO₄ and NO₃ AOD drive the increase in total AOD in Southern Asia since the 1960s. Dust AOD is a relatively constant but important contribution in this region. Total AOD over Southern Asia peaked in the 2010s coincident with the peak in SO₄ and carbonaceous aerosol emissions (Figure 2). Southern Africa and South America also show an increase in total AOD, driven by increase in OA AOD from biomass burning. In these regions SO₄ AOD is also an important steady contributor to total AOD. Northern Africa and Australia AOD are dominated by natural component of dust and sea salt, resulting in an almost constant total AOD in the last decades.

Similar to total AOD, the declines in PM_{2.5} in Europe, North America and Siberia starting from the 1980s are driven mainly by the declines in anthropogenic SO₄ PM_{2.5}, which was more than halved in all the regions between the 1990s and late 2010s. Further decline in surface PM_{2.5} is driven by reduction in the NO₃ and NH₄ components in Europe and Siberia, and to a lesser extent North America, where NH₄ exhibit a slow decline while the NO₃ component remained stable. These declines are partially compensated in Siberia and North America by the increase in magnitude and variability of OA from biomass burning emissions (Figure 2). The OA component in PM_{2.5} is also the dominant component in Southern Africa and South America, driving the total increasing trends in total PM_{2.5} since the 1960s and the early 2000s respectively. As for AOD, PM_{2.5} in Southern Asia increased mainly due to the increase in SO₄, NO₃ and OA. This increase is on top of a generally constant contribution from fine dust which dominates PM_{2.5} in the beginning of the time series.

3 Observations and methodology used for model evaluation

Here we describe in detail each dataset and the methodology used in the evaluation of mass and optical properties of aerosols in the Ref-D1 run. Table 1 summarizes the observations used.

3.1 Satellite AOD from MODIS Neural Net Retrieval (NNR)

We compare the global and regional modelled total column AOD with observations of total column AOD at 550 nm retrieved by the Moderate Resolution Imaging Spectroradiometer (MODIS) onboard the Aqua spacecraft. The version of the dataset used for the comparison is based on the Neural Network Retrieval (NNR), a bias-corrected and quality-controlled retrieval of AOD based on MODIS reflectance observations suitable for use in aerosol data assimilation (Randles et al., 2017). The NNR approach retrieves AOD using cloud-screened and homogenized reflectances from the standard Dark Target (Levy et al., 2013) and Deep Blue (Hsu et al., 2013) retrieval algorithms that have been calibrated to co-located AOD measurements from the Aerosol Robotic Network (AERONET, see Section 3.2) using a neural network algorithm. In general, satellite retrievals of aerosols may be subject to uncertainties due to cloud contamination, inadequate representation of surface reflectance, and retrieval aerosol model assumptions (e.g., particle composition and size). The NNR retrieval is less susceptible to these uncertainties than traditional retrievals because it connects the observed radiances to directly observed AOD measurements and so does not explicitly invoke an aerosol model that assumes particle properties. Because our experiment is a free-running model and no data assimilation is invoked we are here comparing the statistical representation of aerosol interannual, seasonal, monthly, and regional variability in our model run to data rather than focusing on specific events.

We obtain NNR data global monthly means by averaging all the overpasses within each month, and regrid it from its native resolution ($0.25^\circ \times 0.3125^\circ$) to the model spatial resolution. For each month, the model outputs have been spatially sampled to match where monthly NNR data were available. This approach is appropriate given that the model is not run with real meteorology, and we are looking at long-term timeseries and climatological statistics. For the region of Siberia, we consider only months from May to October for each year, given the lack of data coverage in winter months in this region due to snow-covered surfaces not being retrieved over.

3.2 Ground-based AOD and Angstrom Exponent (AE) from AERONET

We compare the Ref-D1 experiment to the column spectral AOD observations from AERONET. AERONET is a federated global network of ground-based sun photometers that measure total column aerosols properties (Holben et al., 1998). We consider the cloud-screened quality assured (Version 3 Level 2) monthly AOD and Armstrong Exponent (AE) (Giles et al., 2019). AE conveys information about the spectral variability in AOD, which itself is an indicator of particle size, where large values of AE (>1 , indicating strong wavelength dependence) are generally associated with small particles (sub-micron) and small values (<1 , indicating more spectrally flat AOD) are more indicative of coarse particles. For our climatological analysis, we consider all AERONET sites which have at least 10 months for each of the 12 climatological months over the period 2000-2018, similarly to what was done by (Gliß et al., 2021). For sites without direct measurements of AOD at 550 nm, AOD available at 500 nm or 532 nm and 667 nm or 675 nm are used to obtain AOD at 550 nm. For comparison with modelled AE at 470-870 nm, we use observations of AE at 440-870 nm. We obtain a total of 46 sites matching the selection criteria, and

Deleted:

they are shown in Supplemental Figure 4 and listed in Supplemental Table 1. The model has been spatially sampled by co-locating the site coordinates with the nearest latitude-longitude model grid cell.

3.3 Surface PM_{2.5} from satellite-derived datasets

The modelled surface total PM_{2.5} are compared with satellite-derived PM_{2.5} model-data fusion datasets developed by the Atmospheric Composition Analysis Group at Washington University in St. Louis and available at <https://sites.wustl.edu/acag/datasets/surface-pm2-5/>. In these datasets, monthly high-resolution surface PM_{2.5} over land are estimated combining information from satellite derived AOD, a chemical transport model, and surface observations from ground-based monitoring networks. The AOD retrieved from multiple satellites is combined to produce a monthly best-estimate satellite-based AOD using a weighted average based on the level of agreement with AERONET. Simulations from the GEOS-Chem chemical transport model were used to establish the relationship between the total AOD and surface PM_{2.5} and applied to the satellite AOD best estimate to produce a geophysical estimates of surface PM_{2.5}. Subsequent statistical fusion incorporated additional information and corrections from PM_{2.5} ground-based measurements.

We use the dataset developed in van Donkelaar et al. 2021 at $0.1^\circ \times 0.1^\circ$ resolution (dataset version V5.GL.03) to compare GEOSCCM simulated global and regional surface total PM_{2.5}. Monthly speciated PM_{2.5} mass concentrations at $0.01^\circ \times 0.01^\circ$ are provided for North America in separate dataset (van Donkelaar et al., 2019), which we use to compare the modeled total PM_{2.5} and components over the contiguous United States (CONUS). We use percentage PM_{2.5} composition (dataset version V4.NA.02) applied to total PM_{2.5} mass (dataset version V4.NA.03), to ensure mass closure as recommended by the data provider.

The main uncertainties related to these satellite-derived hybrid PM_{2.5} estimates are linked to AOD retrieval limitations, AOD to PM_{2.5} assumed relationship in the GEOS-Chem model, sparse ground monitoring in certain areas, and subgrid-scale features. Monthly per-pixel uncertainties range from ~20% in populous Asian regions to 30-50% in North America but decrease significantly when aggregating data, making these PM_{2.5} estimates reliable for regional-scale assessments despite pixel-level limitations.

To be compared with the model, the hybrid datasets have been spatially regridded to match the Ref-D1 spatial resolution. We also exclude areas where the hybrid dataset has no data in the monthly means at high latitudes (latitude > 68°).

3.4 Speciated PM_{2.5} and scattering coefficient from IMPROVE

We compare simulated speciated surface PM_{2.5} concentrations and surface aerosol scattering coefficient with long-term observations over the US from the Interagency Monitoring of Protected Visual Environments Network mainly located in US national parks (Jenny L. Hand, et al., 2023). The IMPROVE network provides measurements of total PM_{2.5} and of speciated

285 components (dust obtained from elemental species data, sea salt, OA, BC, nitrate, and sulfate). Filter samples are collected for
24h consecutive every 3 days. Collected PM mass is subsequently analyzed at controlled laboratory “dry” RH (30%-40%)
conditions. For our climatological analysis of PM_{2.5}, we calculate monthly means from all IMPROVE sites that had at least
80% of measurements data with ‘valid’ status flag over the period 2000-2018 for total PM_{2.5} and all its components (>1850
data points per station). We obtained a total of 101 sites matching the selection criteria, and they are shown in Supplemental
Figure 5 and listed in Supplemental Table 2. We sampled the model output by co-locating the site coordinates with the nearest
290 latitude-longitude model grid cell. From averaging modelled and observed PM_{2.5} and individual components across all the
selected sites, we obtain average modelled and observed monthly speciated PM_{2.5} across the US, which we use for our surface
PM_{2.5} comparison. For the comparison of total PM_{2.5} we apply at a post-processing stage a growth factor correction with relative
humidity at RH=35% as defined in the GEOS documentation (Collow A., 2023) to match the IMPROVE PM_{2.5} measurement
conditions.

295 Additionally, we use long-term co-located measurements of speciated PM_{2.5} and scattering coefficient to investigate the
relationship between surface PM_{2.5} and surface aerosol optical properties. Scattering coefficient at 550 nm together with
ambient RH is measured at a subset of IMPROVE sites through open-air integrating nephelometers. Measurements are reported
hourly. We filter hourly nephelometer measurements with RH <=95%, with scattering coefficient < 5000 m⁻¹ and rate of
change less than 50 Mm⁻¹h⁻¹ to exclude the interference of meteorology (e.g. fog episodes), following the procedure used in
300 (Latimer and Martin, 2019). Subsequently, we consider only days with at least 12 valid hourly measurements to obtain average
daily values. We then consider stations with co-located scattering and PM_{2.5} measurements and subset only days where both
valid PM_{2.5} and scattering are measured. We select months with at least 6 valid daily measurements out of a max of 11 (PM
are measurements are done every 3 calendar days). This selection procedure results in 6 sites with long-term measurements of
both speciated PM_{2.5} and scattering coefficient and RH, presented in Supplemental Figure 6 and listed in Supplemental Table
305 3. An important source of uncertainty in our model-observations comparison with the IMPROVE data (as well as the
AERONET data) stems from the representativeness error when comparing point measurements with model grid cells (1°×1°).
Focusing our our comparisons on seasonal and monthly statistics derived from long-term observations can help mitigate this
sampling mismatch. Additionally, IMPROVE measurements can be subjected to sampling artifacts, particularly volatilization
losses during particle collection and handling. Ammonium nitrate, for instance, can evaporate during filter sampling, leading
310 to underestimation of nitrate concentrations and total PM_{2.5} mass (J.L. Hand, 2023; Ward et al., 2025).

Formatted: Justified, Indent: First line: 0.25", Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto, Ligatures: None

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto, Ligatures: None

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto, Ligatures: None

Deleted: ¶ ... [3]

3.5 Reconstruction of surface scattering from model diagnostics and IMPROVE observations

We use the long-term co-located measurements of speciated PM_{2.5} and scattering coefficient to investigate the relationship
between surface PM_{2.5} concentrations and aerosol optical properties. Total surface scattering coefficient of an aerosol
315 population σ_{scat} [m⁻¹] is a function of mass concentration, size distribution, optical properties, and local relative humidity for

each aerosol component in the population. In GOCART, the total scattering coefficient σ_{sca} of an aerosol population of k species is calculated assuming externally mixed aerosol as:

$$\sigma_{\text{sca}} = \sum_{n=1}^k (b_{\text{sca},n}(\text{LUT}_n, \text{RH}) * m_n) \quad 1$$

where $b_{\text{sca},n}$ is the mass scattering efficiency for each aerosol species n , which depends on the model LUTs that contain the assumed information on aerosol optical properties and size distribution and their dependence on RH. m_n is the dry mass concentration for each aerosol species n . The k species considered in GEOSCCM Ref-D1 in the calculation are organic carbon, black carbon, nitrate, sulfate, dust and sea salt.

Following from the approach developed in (Latimer and Martin, 2019), we compare the observed σ_{sca} from IMPROVE with the reconstructed scattering $\sigma_{\text{sca,RE}}$. We reconstruct the scattering coefficient in Equation 1 using GEOSCCM LUTs for 3 different combinations of modeled and observed RH and $\text{PM}_{2.5}$ composition to isolate the importance of modeled LUT, RH and $\text{PM}_{2.5}$ composition respectively in simulating σ_{sca} (Section 4.4). As indicated in Section 3.5, the IMPROVE optical measurements are at ambient RH, while $\text{PM}_{2.5}$ and PM_{10} measurements are at a controlled, low RH comparable to dry conditions.

One limitation of our approach is that we use fine mode speciated mass to reconstruct total (fine+coarse) observed scattering coefficient, because of the lack of IMPROVE observations on coarse mode. This could lead to underestimating the reconstructed scattering. On the other hand, nephelometers are affected by a truncation error, which is a systematic error increasing as the particle size increases, thus scattering from the coarse mode can be underestimated (Molenar J. V., 1997). To reduce the uncertainty in determining the scattering coefficient for both the observed and reconstructed scattering, we restrict our analysis to conditions dominated by fine-mode aerosols i.e. when IMPROVE observed $\text{PM}_{2.5}/\text{PM}_{10} > 0.5$.

Table 1: summary table of aerosols observations used for comparison with the GEOSCCM Ref-D1 simulation.

observation	platform	time coverage	spatial coverage	source	reference
Aerosol Optical Depth	satellite	2003-2018	global	MODIS Aqua (NNR retrieval)	(Randles et al., 2017)
Aerosol Optical Depth	ground-based	2000-2018	global	AERONET	(Holben et al., 1998)
Angstrom Exponent	ground-based	2000-2018	global	AERONET	(Holben et al., 1998)
Surface total $\text{PM}_{2.5}$ mass	data-model fusion	1998-2018	global	Washington U. St. Louis	(van Donkelaar et al., 2021)
Speciated surface $\text{PM}_{2.5}$ mass	data-model fusion	2000-2016	US	Washington U. St. Louis	(van Donkelaar et al., 2019)
Speciated surface $\text{PM}_{2.5}$ mass	ground-based	2001-2018	US	IMPROVE network	(Jenny L. Hand, et al., 2023)
Surface aerosol scattering coefficient	ground-based	2001-2018	US	IMPROVE network	(Jenny L. Hand, et al., 2023)

4 Results

4.1 Global column AOD

345 Figure 4 shows the comparison of regional mean monthly timeseries and mean annual cycle of AOD between Ref-D1 and
MODIS NNR retrievals. The global climatological monthly spatial distribution is reported in Supplemental Figure 7.
GEOSCCM reproduces month-to-month variability of AOD over land for all regions ($0.6 < r < 0.9$) except in Europe and
Siberia ($r \approx 0.3$). However, there are strong regionally and seasonally varying biases evident. GEOS CCM simulated AOD is
350 biased low by 38%-58% in biomass burning regions of Southern Africa, North America, South America, Siberia during peak
fire months (Figure 4b). For South America and Southern Africa, simulated AOD is underestimated throughout the entire year.
Over Northern Africa, the model underestimates AOD during the spring season and overestimates the seasonal summertime
and early autumn peak in magnitude (within ~30%) and shows a delay of 1-2 months in the peak of AOD compared to
observations, with bias mainly concentrated over the South Sahara steppe and woodlands (Supplemental Figure 7). In Southern
Asia the model shows a good agreement with observations in all seasons except winter, when AOD is overestimated (+26%).
355 A persistent positive bias is found over Eastern China almost throughout the year (Supplemental Figure 7). Over Europe the
model has a high bias in AOD in all seasons (between +36% and +64%) except summer, when the AOD is underestimated (-
13%). Finally, over Australia, the model tends to overestimate AOD in summer (+35%) and slightly underestimate AOD over
winter (-13%), but the overall seasonal variability is well reproduced.

We find consistent results with the comparison with satellite retrieved AOD when comparing the modeled total column
360 AOD with observations at selected AERONET sites (Supplemental Figure 8). GEOSCCM AOD is on average underestimated
in biomass burning influenced regions and overestimated over Europe, and GEOSCCM overestimates AOD at sites over North
America during all seasons except summer (i.e., peak biomass burning season). However, at sites over Asia in India and China
the model underestimates AOD compared to AERONET, which is somewhat at odds with the results obtained with the
comparison with satellited retrieved AOD, for which modeled AOD was overestimated. Total column AE tends to be globally
365 underestimated in the model compared to AERONET observations (Supplemental Figure 9), suggesting that the model
simulates either too much coarse aerosol or not enough smaller aerosols, as also found on average in the AeroCom multi-
model evaluation (Gliß et al., 2021). An exception are Asian sites and central Brazil, where AE is overestimated. Additional
climatological comparison and statistics of AOD and AE for each selected AERONET site are presented in Supplemental
Figure 10, 11 and Supplemental Table 4.






Figure 4: Comparison of GEOSCCM Ref-D1 and MODIS Aqua (a) regional annual timeseries and (b) regional seasonal cycle of AOD for 2003-2018. For GEOSCCM, the AOD seasonal cycle is reported as speciated. Siberia seasonal cycle is from May to October only.




375 **Figure 5: Comparison of GEOSCCM Ref-D1 (red) and satellite derived (black) PM_{2.5} for the period 1998 – 2018.**
 Shown are (a) regional annual timeseries and (b) regional annual cycle. Note the y-axes do not necessarily start at zero.

4.2 Global surface total PM_{2.5} concentrations


380 Comparisons of the GEOSCCM Ref-D1 simulated PM_{2.5} concentrations with satellite-derived PM_{2.5} show similarities with the comparison with total column AOD. Figure 5 shows the regional mean monthly timeseries and mean annual cycle comparison for surface PM_{2.5} concentrations. The global climatological monthly spatial distribution is reported in Supplemental Figure 12. The model captures the month-to-month variability in regional PM_{2.5} compared to the satellite-

385 derived $PM_{2.5}$ in all regions ($r > 0.5$), especially in Southern Asia, North America, and Siberia ($0.73 < r < 0.84$). However, the modeled $PM_{2.5}$ concentrations are biased low in all regions throughout the seasonal cycle, except for Northern Africa, which has a small positive bias, and Australia, which has a mixed bias.

Similar to AOD, the model underestimates $PM_{2.5}$ in biomass burning regions, but does overall reproduce the seasonal cycle, except for Southern Africa, where the model predicts the peak season to occur one month earlier than compared to observations. During peak fire season, highest biases are found in South America and Southern Africa ($\sim 70\%$) and Siberia ($\sim 60\%$), while the model performs better in North America ($\sim 10\%$). Over Northern Africa, the model overestimates $PM_{2.5}$ by around 20% in all seasons except spring. For Southern Asia, the model underestimates $PM_{2.5}$, especially during wintertime (-20%), while in Europe the bias is largest in the summertime (-37%). Model bias is largest in Australia during the southern hemisphere winter (20%). Notably, over Europe there is a much better correlation of the climatological modeled $PM_{2.5}$ with observations ($r = 0.8$) than for AOD ($r = 0.33$), suggesting some disconnect between the modeled mass at the surface and the aerosols optical properties at the column level and or some uncertainties related to the transport in the model.



Formatted: Centered



Deleted:

Figure 6. Comparison of speciated $PM_{2.5}$ over CONUS between GEOSCCM (REFD1, red) and satellite-derived $PM_{2.5}$ (black) at $RH=35\%$. (a) Annual timeseries and (b) mean seasonal cycle for the period 2000-2003.

4.3 PM_{2.5} composition over CONUS

We use both satellite-derived and ground-based observations of speciated PM_{2.5} over CONUS to investigate in more detail the GEOSCCM ability to represent aerosol composition at the surface. Figure 6 shows the comparison of speciated surface PM_{2.5} concentration timeseries and seasonal cycle between GEOSCCM and satellite-derived data. The spatial distribution comparison is reported in Supplemental Figure 13. Figure 7 shows the comparison of speciated surface PM_{2.5} concentrations timeseries and spatial distribution between GEOSCCM and ground-based observations from the IMPROVE network.

GEOSCCM has a similar decreasing trend in total PM_{2.5} as in the satellite-derived observations (Figure 6), although it has a positive bias during wintertime. The performance for individual species varies. GEOSCCM Ref-D1 reproduces the decreasing trend in SO₄ both for ground-based observations and satellite-derived SO₄, and to a lesser extent the decreasing trend in NH₄ (available for satellite-derived data only). However, the model is not able to fully capture the SO₄ and NH₄ seasonal cycle and is positively biased especially in winter months. The comparison with both satellite-derived data and ground-based observations shows an average monthly positive bias in natural aerosols of sea salt (Normalized Mean Bias NMB $\approx$ 2 and NMB $\approx$ 12 respectively) and dust (NMB $\approx$ 0.3 and NMB $\approx$ 0.1 respectively). This is most pronounced in the summer, and may be linked to excessive inland intrusion of dust and sea salt mainly from the Atlantic Ocean, and an overestimation of local dust sources from the Mojave and possibly the Chihuahuan Deserts (Figure 7b and Supplemental Figure 13). In addition the model doesn't account for land-use changes that affect dust emissions over time, while observations capture these evolving conditions. This likely explains the increasing bias between modeled and observed dust concentrations especially in the final years of the study period. The model shows a high positive bias in nitrate during the winter when compared both to ground-based observations (NMB = 3.61 and satellite-derived data NMB = 1.14), driving an overall positive bias in total PM_{2.5} in the winter, which is also affected by model-overestimated SO₄ and NH₄. On the other hand, GEOSCCM is skillful in reproducing OA interannual variability ($r = 0.9$) and capture better magnitude compared to the ground-based observations (NMB $\approx$ -0.12) than to the satellite derived estimates (NMB $\approx$ -0.43). For BC, the model has also a smaller magnitude bias against the ground-based observations than for satellite-derived ones (NMB = 0.03 vs NMB = -0.67 respectively) but is not able to capture the month-to-month variability in either ($r < \sim 0.2$). Despite the described biases, the model captures the spatial distribution of observations for most PM_{2.5} components, except for sea-salt and dust as already noted and for nitrate (Figure 7b).

The differences between the comparison of GEOSCCM with the satellite derived PM_{2.5} and the ground-based monitors may arise from the fact that IMPROVE ground-based sites are representative of background concentrations, being sites located in national parks across the US, while the satellite-derived PM_{2.5} are representative of different chemical environments at higher resolution (background, rural, urban). This is evident for example in the performance of BC. The model underestimation of BC when compared with satellite derived component but not when compared with the IMPROVE ground-based stations, which might indicate that the model is missing some anthropogenic BC in the emission inventory used.

Deleted: This is most pronounced in the summer, and may be linked to excessive inland intrusion of dust and sea salt mainly from the Atlantic Ocean (Figure 7b and Supplemental Figure 13).

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, Not Italic, Font color: Auto

Deleted: , which has a smaller bias in the comparison with observations from the IMPROVE sites than the satellite-derived ones....

Formatted: Font: (Default) Times New Roman, Font color: Auto, Ligatures: None

Formatted: Font: (Default) Times New Roman, Underline, Font color: Dark Red, Ligatures: None

Formatted: Font: (Default) Times New Roman, Font color: Auto, Ligatures: None

In addition, this comparison could be sensitive to different assumptions in the AOD and PM_{2.5} relationship in GEOSCCM versus the one assumed in the GEOS-Chem model which is used to obtain the satellite-derived PM_{2.5}, especially in areas where the incorporated additional information and corrections from PM_{2.5} ground-based measurements is sparse.





Figure 7. (a): Ref-D1 modelled (red, Ref-D1) vs observed (black, IMPROVE) monthly mean timeseries of speciated PM_{2.5}, averaged over the observation sites throughout CONUS. Lines represent the median, while shading

455

encompasses the 25th -75th percentiles. (b) Modelled vs observed (circles, IMPROVE) yearly average of surface PM_{2.5} and composition over CONUS (2001-2018). **Total PM_{2.5} are compared at RH=35%, while individual species are measured and simulated at RH=0% (dry mass).**

Formatted: Subscript



460

Figure 8. Seasonal cycle of (a) surface scattering coefficient and (b) total PM_{2.5} (RH=35%) at selected IMPROVE sites. Black lines represent IMPROVE observations, solid lines are Ref-D1 and dashed lines are Ref-D1 with the nitrate component removed.

465

4.4 Connecting surface PM_{2.5} mass and scattering coefficient

470

Differences in the comparisons of AOD and PM_{2.5} to observations prompt us to investigate the relationship between optical properties and aerosol mass in the model. For this, we use co-located IMPROVE measurements of speciated PM_{2.5} and the scattering coefficient at selected sites, as described in Section 3.5. We find that the model has a better performance in reproducing surface total PM_{2.5} than scattering coefficient for the selected sites (Figure 8, Supplemental Table 5). Although the bias in the scattering coefficient and PM_{2.5} varies across sites, the model is not able to capture the seasonal variability of the scattering coefficient at all sites ($-0.2 < r < 0.2$), while it does better for the seasonality of PM_{2.5} ($-0.25 < r < 0.59$). The performance of the model improves significantly both for the scattering coefficient and the total PM_{2.5} when removing the

Deleted: Table 2: statistics referring to Figure 8. station station

... [4]

Deleted: Table 2

nitrate component, which we found is largely overestimated in the model in winter over CONUS (Section 4.3), although it is still better for PM_{2.5} ($0.24 < r < 0.84$) than for the scattering coefficient ($-0.18 < r < 0.6$).

We reconstruct the scattering coefficient ($\sigma_{\text{sca,RE}}$) for different combinations of modeled and observed RH, PM_{2.5} and LUT (Section 3.6) and compare it with the observed scattering coefficient (σ_{sca}) to help understand the role of each of these factors in the disconnect between the model performance for the scattering coefficient and PM_{2.5}. For each species n these combinations are: 1) [$LUT_{\text{mod}}, RH_{\text{obs}}, m_{\text{obs}}$] to isolate the impact of GEOSCCM assumed optical tables; 2) [$LUT_{\text{mod}}, RH_{\text{mod}}, m_{\text{obs}}$] to isolate the impact of simulated relative humidity; and 3) [$LUT_{\text{mod}}, RH_{\text{obs}}, m_{\text{mod}}$] to isolate the impact of simulated PM_{2.5} composition.

Figure 9 shows the comparison between the reconstructed and observed scattering coefficient. We find that the $\sigma_{\text{sca,RE}}$ using the model LUT with the observed RH and PM_{2.5} (Figure 9, blue line) is in good agreement with the observed σ_{sca} with generally the lowest NMB and RMSE and highest correlation for this case, indicating that the combination of optical properties and size distribution assumptions in the LUT are generally compatible with observations. By contrast, the reconstructed $\sigma_{\text{sca,RE}}$ that uses the model LUT, observed PM_{2.5} and modeled RH (Figure 9, green line) exhibits a consistent negative bias compared to observations for all sites. This is expected since the model-simulated RH is generally lower than observed at all sites (Figure 10b) and the scattering coefficient increases as RH increases. The results are more ambiguous for $\sigma_{\text{sca,RE}}$ using the model LUT, observed RH, and modeled PM_{2.5} (Figure 9, orange line) suggesting that the modeled composition plays a more complicated role in affecting the simulated scattering coefficient. At the selected sites, beside the high bias in nitrate, GEOSCCM Ref-D1 is not able to fully capture the observed PM_{2.5} seasonal composition (Figure 10a), with biases for each component in line with what is found for the comparison of speciated PM_{2.5} averaged over CONUS in Section 4.3. However, when removing the nitrate component and reconstructing the scattering, the performance of the model improves (Supplemental Figure 14, Supplemental Table Q).

Deleted: 5





Figure 9. Reconstructed scattering for 3 different combinations of modelled and observed RH, PM_{2.5} and LUT as described in Section 3.6. (a) Seasonal cycle of reconstructed scattering at the selected sites. Observations are reported with the black line. (b) Scatter plot of the reconstructed scattering with all monthly data.

Table 3: metrics for the reconstructed scattering $s_{sca,RE}$ compared to IMPROVE scattering observations for the three combination of model and observations considered: 1) $[LUT_{mod}, RH_{obs}, m_{obs}]$, 2) $[LUT_{mod}, RH_{mod}, m_{obs}]$, 3) $[LUT_{mod}, RH_{obs}, m_{mod}]$.

station	NMB 1	NMB 2	NMB 3	RMSE 1 [Mm-1]	RMSE 2 [Mm-1]	RMSE 3 [Mm-1]	r 1	r 2	r 3
BIBE1	0.12	-0.48	-0.16	5.18	13.52	9.21	0.84	0.64	0.06
GRCA2	0.14	-0.37	3.25	5.73	6.73	46.71	0.67	0.58	0.31
GRSM1	0.00	-0.68	0.1	13.97	54.85	40.28	0.92	0.86	0.16
MACA1	0.08	-0.62	0.1	12.09	41.45	23.08	0.9	0.86	0.48
MORA1	0.02	-0.62	-0.22	9.85	20.02	15.13	0.71	0.62	0.24
SHEN1	0.13	-0.63	0.1	11.55	38.51	31.6	0.95	0.9	0.07

515



520

Figure 10. Comparison between Ref-D1 and observations for the seasonal cycle of (a) surface PM_{2.5} components (a) and (b) surface RH at selected IMPROVE sites. Individual species are measured and compared at RH=0% (dry mass).




Figure 11. (a) Comparison of the observed and reconstructed scattering coefficient for different RH ranges and different dominant component in the $PM_{2.5}$ (fraction > 0.5) using daily IMPROVE observations of speciated $PM_{2.5}$, RH and scattering coefficient. (b) Same as (a) but with reconstructed scattering with updated LUT for OC.

We further analyze the role of model assumptions of aerosol optical properties by sorting the reconstructed scattering coefficient for different RH range and dominant $PM_{2.5}$ component, as shown in Figure 11. Overall, where the dominant component is OA, dust, or SO_4 , the distribution of the reconstructed scattering coefficient agrees well with observations for all different RH ranges, suggesting that the model assumptions of aerosol optics for each of these species is adequate. However, it is possible to further improve the model representation of aerosol properties. For example, the reconstructed scattering coefficient derived using an updated set of optical properties developed by Das et al. (2024) for biomass burning organic aerosol, is in better agreement with observed scattering coefficients compared to using default optics, especially in regions where OA is the dominant component (Figure 11b).

Deleted: D

5 Discussion

In this study we assessed GEOSCCM capabilities and limitations in simulating aerosols by focusing on the link between aerosol optical properties and mass, examining key factors that contribute to uncertainties in simulated long-term variability, regional and seasonal variations of AOD and $PM_{2.5}$, and the consistency of GEOSCCM $PM_{2.5}$ simulation with optical properties, including the role of simulated RH on hygroscopic scattering enhancement. This work also presents the first extensive comparison between the GEOSCCM aerosol component and observational data.

Our results show that the GEOCCM reproduces month-to-month variability and long-term trends of total column AOD and surface $PM_{2.5}$ concentrations over most land regions. However, GEOSCCM shows regional biases in AOD and $PM_{2.5}$ that show similar global patterns. In particular, GEOSCCM underestimates both AOD and $PM_{2.5}$ in biomass burning regions and overestimates dust over the Sahara Desert. Performance for speciated $PM_{2.5}$ varies. Specifically, the GEOSCCM model overestimates secondary inorganic $PM_{2.5}$ concentrations across the continental United States, with nitrate showing the most significant overestimation. Additionally, the model produces excessive values for sea-salt in coastal regions and dust during summer months. The model performs better for carbonaceous species (OA and BC). Biases in the magnitude of individual $PM_{2.5}$ components also have a significant impact in the simulated aerosol scattering, as does the model-simulated relative humidity, which is generally lower than observed, while we find that the scattering efficiency assumptions in GEOSCCM are consistent with observations.

One cause of aerosol loading biases in biomass burning regions can be related to biases in the emission inventory used in the GEOSCCM Ref-D1 simulation, which is based on fire emissions estimations from burned-area estimates (Section 2.2). Burned-area driven fire inventories have been shown to give lower overall biomass burning aerosol emissions and result in low AOD over biomass burning regions in GEOS compared to emissions derived from fire radiative power, which are by design constrained by observed satellite AOD, as well as use different set of emissions factors (Pan et al., 2020). This is further confirmed by comparing the simulated AOD by GEOSCCM with the one simulated with a recent benchmark run of GEOS, GOCART-2G run, (Collow et al., 2024) but using a fire radiative power-based inventory (QFEDv2.5 (Darmenov, A. and da Silva, A., 2015)). The GOCART-2G run estimates higher AOD over biomass burning region than GEOSCCM and is closer to observations especially during peak fire season, since by design it is specifically tuned to produce a reasonable observed AOD. While increasing the emissions of biomass burning would improve GEOSCCM performance in simulating AOD and $PM_{2.5}$ mass over relevant regions, there might be additional reasons that contribute to the low bias, such as not utilizing a plume-rise parametrization (Ke et al. 2021), incorrect lifetime from incorrect precipitation simulation, and particle size-distribution assumptions that affect the assumed mass scattering efficiencies (Zhong et al., 2022).

The underestimation of surface $PM_{2.5}$ despite reasonable AOD agreement in non-biomass burning regions likely stems from multiple factors. Emission inventory deficiencies, particularly for anthropogenic sources, may contribute to systematic regional biases. Additionally, the $PM_{2.5}$ -AOD relationship depends heavily on aerosol vertical distribution (Zhu et al. 2024), and previous studies show that global models tend to simulate excessive aerosol transport to the free troposphere at the expense

Deleted: and Conclusions

Deleted: , except Europe

Formatted: Subscript

Formatted: Subscript

of near-surface concentrations (Koffi et al. 2016; Yu et al. 2010). This vertical redistribution may contribute to explain why column-integrated AOD appears reasonable while surface $PM_{2.5}$ remains underestimated.

Formatted: Subscript

For the overestimation in GEOSCCM of natural aerosols of dust and sea salt there might also be multiple factors at play. GEOSCCM dust summer peaks might be higher than observations because the current model does not include seasonal vegetation cover changes that likely suppress dust emissions, especially over Southern Saharan steppe and woodlands in summer. While both for sea salt and dust, the model coarse resolution might lead to more inland intrusion of these species due to higher numerical diffusion over coastal regions as found for the regional $PM_{2.5}$ composition analysis over CONUS. Additionally, the internal resolution-dependent dust emission scaling factor used may have been inappropriate. Finally, the free running meteorology in the GEOSCCM configuration can introduce differences in the long-range transport dynamics and removal processes compared to a run using replayed or assimilated meteorology. As shown in Supplemental Figure 15 GEOSCCM is relatively more efficient at transporting dust from the Sahara across the North Atlantic than the GEOS GOCART2G run performed with the MERRA-2 assimilated meteorology (Collow et al. 2024).

Analysis of the speciated $PM_{2.5}$ over CONUS highlights that GEOSCCM overall can reproduce long-term trends of total and speciated $PM_{2.5}$ and their spatial distribution features, but there are biases in individual components. Besides the biases in biomass burning and natural aerosols already discussed, GEOSCCM shows a positive bias in secondary inorganic components (NO_3 , SO_4 NH_4), especially during wintertime. The nitrate aerosol component shows the greatest positive bias, which also drives a high positive bias in total $PM_{2.5}$ during winter. One main reason for the bias in nitrate can be related to potential problems in nitric acid (HNO_3) mass closure in the coupled GMI-GOCART scheme in GEOSCCM. As described in Section 2.2, nitric acid chemistry is computed in GMI and then passed as an input to GOCART for nitrate chemistry computation. However, nitric acid loss due to nitrate formation is not fed back to GMI, and likely results in an overestimation of total HNO_3 available. Indeed, when using a configuration with decoupled chemistry-aerosols using scaled HNO_3 fields as input to GOCART, the GEOS model has a better performance for nitrate (Collow et al., 2024). In addition, the positive bias in natural aerosols provides more surface for heterogeneous production of nitrate onto sea salt and dust particles, which is the main pathway for nitrate aerosol formation (Bian et al., 2017), leading to potential overestimation of nitrate formed through heterogeneous chemistry.

The analysis of co-located scattering coefficient, speciated $PM_{2.5}$ and reconstructed scattering coefficient shed lights over the uncertainties driving the link between aerosol mass and aerosol optical properties in current GEOSCCM. We found that current optics LUT assumptions used in the model are overall compatible with observations. However, aerosol optical properties can be further improved, as shown for the biomass burning emitted OA example. Extending the assessment of GEOSCCM assumptions of aerosol optical properties beyond the CONUS will be also desirable, in order to increase the confidence in optics LUT on a global scale spanning different physio-chemical environments. The main uncertainties in the aerosol mass-optical properties relationship are driven by model-simulated relative humidity and model simulated $PM_{2.5}$ composition, which play an important role in determining biases in simulated surface scattering coefficient. In particular, model-simulated RH is generally lower than observed over CONUS, driving a constant negative bias in simulated scattering

605 coefficient, that could be linked to deficiencies in our methodology in deriving surface RH accurately enough with free-running meteorology. It has also been shown that model calculated AOD is sensitive to the spatial and temporal resolution of atmospheric relative humidity, due to the highly non-linear relationship between RH and the aerosol mass extinction efficiency (Bian et al., 2009). The bias introduced by the modeled aerosol composition has a more mixed effect in affecting the simulated scattering coefficient (both positive and negative biases), which can be more complex to disentangle. Nevertheless, improving the model's ability to correctly simulate aerosol composition would improve the simulated scattering coefficient, as shown in the experiment with nitrate removed.

6 Conclusions

615 In this work we investigated the drivers of uncertainties in the current NASA Goddard Earth Observing System Chemistry Climate Model (GEOSCCM) in simulating aerosols by focusing on the link between aerosol optical properties and mass. In conclusion, this study demonstrates that the primary drivers of simulated aerosol scattering uncertainty in GEOSCCM stem from biases in PM_{2.5} component concentrations and the simulated relative humidity field rather than from model assumptions regarding optical properties or size distributions, which are in good agreement with observations.

620 These findings serve as a benchmark for GEOSCCM aerosol improvements, with the following priorities to reduce uncertainties in the aerosols mass, optical properties and their relationship: 1) aerosol mass loading and optical properties can both be improved in GEOSCCM, possibly by using a fire radiative power based inventory fire emissions which will reduced seasonal biases in biomass burning regions; 2) biases in PM_{2.5} and surface scattering coefficient can be reduced by reducing biases in nitrate mass loading component, which require further investigation in the coupled GMI-GOCART module and in the nitrate heterogeneous production onto natural aerosols surface; 3) and accurate model representation of RH could substantially improve the representation of the aerosol scattering coefficient. For free-running meteorology models like GEOSCCM, increased spatial resolution could improve convective representation and relative humidity fields, leading to better aerosol extinction simulations (Bian et al., 2009), though computational costs must be balanced against increased accuracy. Additionally, incorporating land-atmosphere coupling could better represent surface processes affecting atmospheric RH through enhanced soil-vegetation-atmosphere feedbacks (Santanello et al., 2018). In addition, further systematic assessment of the model's aerosol vertical profile representation against lidar observations will be important and complementary to this work for understanding the aerosol mass-optical relationship, as well as the connection between surface concentrations and column-integrated AOD retrievals.

635 The challenges identified in GEOSCCM's aerosol representation reflect issues that are present more broadly in the context of aerosol modeling. Our identified RH biases reflect a persistent challenge across Earth system models (Dunn et al. 2017). In particular, our work highlights the implication of biases in the simulated relative humidity on the aerosol scattering coefficient, which is of importance for climate models where the meteorology is free running and not prescribed based on reanalysis as in chemistry transport models. This is in addition to the diversity in how models represent aerosol light scattering enhancement with humidity, emphasizing that hygroscopicity plays an important role in simulated aerosol extinction (Latimer and Martin

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, Pattern: Clear

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, Pattern: Clear

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Auto, Pattern: Clear

Formatted: Subscript

2019; Burgos et al. 2020). Assumptions about aerosol size distributions and incorrect mass extinction coefficients can also
640 significantly impact scattering calculations (Latimer and Martin 2019; Zhong et al. 2022).

Our work reinforces the importance of synergistic use of multiple co-located and coincident observations to effectively
investigate and constrain modeled aerosols properties in climate models. By simultaneously assessing aerosol AOD, surface
PM_{2.5} mass concentration and composition, and surface scattering coefficient, it was possible to identify key sources of
uncertainties in GEOSCCM's current aerosol representation and in the link between aerosol optical properties and mass. This
645 co-incident and co-located observational constraint approach provides a framework for model improvement that extends
beyond traditional independent parameters evaluation, helping to clarify the interconnected properties and processes affecting
aerosol representation in atmospheric models, and improve their subsequent applications.

Code Availability

GEOS, including the GEOSCCM configuration, is a publicly available Earth System model with source code
650 at <https://github.com/GEOS-ESM>.

Data Availability

655 All observational data used are from publicly available datasets.
MODIS Level 2 reflectances are available from https://doi.org/10.5067/MODIS/MOD04_L2.006 (Levy, R. et al., 2015a) for
Terra and https://doi.org/10.5067/MODIS/MYD04_L2.006 (Levy, R. et al., 2015b) for Aqua.
AERONET observations can be downloaded at https://aeronet.gsfc.nasa.gov/cgi-bin/webtool_aod_v3.
IMPROVE network data can be downloaded from the Federal Land Manager Environmental Database at
660 <http://views.cira.colostate.edu/fed/DataWizard/Default.aspx>.
Satellite-derived PM_{2.5} datasets are available at <https://sites.wustl.edu/acag/datasets/surface-pm2-5/>.
GOCART optical tables are available at <https://portal.nccs.nasa.gov/datashare/iesa/aerosol/AerosolOptics/>.
(Last access for all the data May 7, 2025).

Author Contribution

665 CM, PRC, conceived the study. PRC, QL, LH, KEN, VV, SAS, SS, MM participated in the set-up and run of the RefD1
experiment. CM led the software, data curation and analysis with contribution from PRC and ABC. CM was responsible for
original draft preparation, and PRC, ABC, SAS, SD and QL contributed to review and editing.

Competing Interests

The authors declare no competing interests.

670 **Acknowledgements**

Simulations were performed at the NASA Center for Climate Simulation (NCCS).

IMPROVE is a collaborative association of state, tribal, and federal agencies, and international partners. US Environmental Protection Agency is the primary funding source, with contracting and research support from the National Park Service. The
675 Air Quality Group at the University of California, Davis is the central analytical laboratory, with ion analysis provided by Research Triangle Institute, and carbon analysis provided by Desert Research Institute.

We thank the AERONET (PI(s) and Co-I(s)) and their staff for establishing and maintaining the 46 sites used in this investigation.

680 **Financial Statement**

This work is funded by the NASA Modeling, Analysis, and Prediction (MAP) program (PM: David Considine) under the Chemistry-Climate Modeling (CCM) project.

685

Bibliography

Aquila, V., Garfinkel, C. I., Newman, P. a., Oman, L. d., and Waugh, D. w.: Modifications of the quasi-biennial oscillation by a geoengineering perturbation of the stratospheric aerosol layer, *Geophys. Res. Lett.*, 41, 1738–1744, <https://doi.org/10.1002/2013GL058818>, 2014.

690 Bharath, J., Kumar, T. V. L., Salgueiro, V., Costa, M. J., and Mall, R. K.: Aerosol characteristics in CMIP6 models’ global simulations and their evaluation with the satellite measurements, *Int. J. Climatol.*, 44, 217–236, <https://doi.org/10.1002/joc.8324>, 2024.

Bian, H., Chin, M., Rodriguez, J. M., Yu, H., Penner, J. E., and Strahan, S.: Sensitivity of aerosol optical thickness and aerosol direct radiative effect to relative humidity, *Atmospheric Chem. Phys.*, 9, 2375–2386, <https://doi.org/10.5194/acp-9-2375-2009>,
695 2009.

Bian, H., Chin, M., Hauglustaine, D. A., Schulz, M., Myhre, G., Bauer, S. E., Lund, M. T., Karydis, V. A., Kucsera, T. L., Pan, X., Pozzer, A., Skeie, R. B., Steenrod, S. D., Sudo, K., Tsigaridis, K., Tsimpidi, A. P., and Tsyro, S. G.: Investigation of global particulate nitrate from the AeroCom phase III experiment, *Atmospheric Chem. Phys.*, 17, 12911–12940, <https://doi.org/10.5194/acp-17-12911-2017>, 2017.

- 700 Bouwman, A. F., Lee, D. S., Asman, W. a. H., Dentener, F. J., Van Der Hoek, K. W., and Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, *Glob. Biogeochem. Cycles*, 11, 561–587, <https://doi.org/10.1029/97GB02266>, 1997.
- Brauer, M., Roth, G. A., Aravkin, A. Y., Zheng, P., Abate, K. H., Abate, Y. H., Abbafati, C., Abbasgholizadeh, R., Abbasi, M. A., Abbasian, M., Abbasifard, M., Abbasi-Kangevari, M., ElHafeez, S. A., Abd-Elsalam, S., Abdi, P., Abdollahi, M., Abdoun, M., Abdulah, D. M., Abdullahi, A., Abebe, M., Abedi, A., Abedi, A., Abegaz, T. M., Zuñiga, R. A. A., Abiodun, O., Abiso, T. L., Aboagye, R. G., Abolhassani, H., Abouzid, M., Aboye, G. B., Abreu, L. G., Abualruz, H., Abubakar, B., Abu-Gharbieh, E., Abukhadajah, H. J. J., Aburuz, S., Abu-Zaid, A., Adane, M. M., Addo, I. Y., Addolorato, G., Adedoyin, R. A., Adekanmbi, V., Aden, B., Adetunji, J. B., Adeyeoluwa, T. E., Adha, R., Adibi, A., Adnani, Q. E. S., Adzigbli, L. A., Afolabi, A. A., Afolabi, R. F., Afshin, A., Afyouni, S., Afzal, M. S., Afzal, S., Agampodi, S. B., Agbozo, F., Aghamiri, S., Agodi, A., Agrawal, A., Agyemang-Duah, W., Ahinkorah, B. O., Ahmad, A., Ahmad, D., Ahmad, F., Ahmad, N., Ahmad, S., Ahmad, T., Ahmed, A., Ahmed, A., Ahmed, A., Ahmed, L. A., Ahmed, M. B., Ahmed, S., Ahmed, S. A., Ajami, M., Akalu, G. T., Akara, E. M., Akbarialiabad, H., Akhlaghi, S., Akinosoglou, K., Akinyemiju, T., Akkaif, M. A., Akkala, S., Akombi-Inyang, B., Awaity, S. A., Hasan, S. M. A., Alahdab, F., AL-Ahdal, T. M. A., Alalalmech, S. O., Alalwan, T. A., Al-Aly, Z., Alam, K., Alam, N., Alanezi, F. M., Alanzi, T. M., Albakri, A., AlBataineh, M. T., Aldhaleei, W. A., et al.: Global burden and strength of evidence for 88 risk factors in 204 countries and 811 subnational locations, 1990–2021: a systematic analysis for the Global Burden of Disease Study 2021, *The Lancet*, 403, 2162–2203, [https://doi.org/10.1016/S0140-6736\(24\)00933-4](https://doi.org/10.1016/S0140-6736(24)00933-4), 2024.
- 715 Carn, S. A., Fioletov, V. E., McLinden, C. A., Li, C., and Krotkov, N. A.: A decade of global volcanic SO₂ emissions measured from space, *Sci. Rep.*, 7, 44095, <https://doi.org/10.1038/srep44095>, 2017.
- Case, P. A., Colarco, P. R., Toon, O. B., and Newman, P. A.: Simulating the Volcanic Sulfate Aerosols From the 1991 Eruption of Cerro Hudson and Their Impact on the 1991 Ozone Hole, *Geophys. Res. Lett.*, 51, e2023GL106619, <https://doi.org/10.1029/2023GL106619>, 2024.
- 720 Chin, M., Jacob, D. J., Gardner, G. M., Foreman-Fowler, M. S., Spiro, P. A., and Savoie, D. L.: A global three-dimensional model of tropospheric sulfate, *J. Geophys. Res. Atmospheres*, 101, 18667–18690, <https://doi.org/10.1029/96JD01221>, 1996.
- Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: Tropospheric Aerosol Optical Thickness from the GOCART Model and Comparisons with Satellite and Sun Photometer Measurements, 2002.
- 725 Chin, M., Diehl, T., Tan, Q., Prospero, J. M., Kahn, R. A., Remer, L. A., Yu, H., Sayer, A. M., Bian, H., Geogdzhayev, I. V., Holben, B. N., Howell, S. G., Huebert, B. J., Hsu, N. C., Kim, D., Kucsera, T. L., Levy, R. C., Mishchenko, M. I., Pan, X., Quinn, P. K., Schuster, G. L., Streets, D. G., Strode, S. A., Torres, O., and Zhao, X.-P.: Multi-decadal aerosol variations from 1980 to 2009: a perspective from observations and a global model, *Atmospheric Chem. Phys.*, 14, 3657–3690, <https://doi.org/10.5194/acp-14-3657-2014>, 2014.
- 730 Chou, M. and Suarez, M. J.: A solar radiation parameterization for atmospheric studies, Technical Report Series on Global Modeling and Data Assimilation, available at: <http://gmao.gsfc.nasa.gov/pubs/docs/Chou136.pdf>, 1999.
- Chou, M. and Suarez, M. J.: An efficient thermal infrared radiation parameterization for use in general circulation models, Technical Report Series on Global Modeling and Data Assimilation, available at: <https://gs6101-gmao.gsfc.nasa.gov/pubs/docs/Chou137.pdf>, 2001.
- 735 Colarco, P., da Silva, A., Chin, M., and Diehl, T.: Online simulations of global aerosol distributions in the NASA GEOS-4 model and comparisons to satellite and ground-based aerosol optical depth, *J. Geophys. Res. Atmospheres*, 115, <https://doi.org/10.1029/2009JD012820>, 2010.

- 740 Collow, A. B., Colarco, P. R., da Silva, A. M., Buchard, V., Bian, H., Chin, M., Das, S., Govindaraju, R., Kim, D., and Aquila, V.: Benchmarking GOCART-2G in the Goddard Earth Observing System (GEOS), *Geosci. Model Dev.*, 17, 1443–1468, <https://doi.org/10.5194/gmd-17-1443-2024>, 2024.
- Collow A., B., V. ., Chin, M. ., Colarco, P. ., Darnenov, A. ., and da Silva, A.: Supplemental Documentation for GEOS Aerosol Products, GMAO Office Note No. 22 (Version 1.0), 2023.
- 745 Darnenov, A. and da Silva, A.: The quick fire emissions dataset (QFED)– Documentation of versions 2.1, 2.2 and 2.4. NASA/TM-2015-104606, Vol. 38, NASA Global Modeling and Assimilation Office, 183 pp., <https://gmao.gsfc.nasa.gov/pubs/docs/Darnenov796.pdf>, 2015.
- van Donkelaar, A., Martin, R. V., Li, C., and Burnett, R. T.: Regional Estimates of Chemical Composition of Fine Particulate Matter Using a Combined Geoscience-Statistical Method with Information from Satellites, Models, and Monitors, *Environ. Sci. Technol.*, 53, 2595–2611, <https://doi.org/10.1021/acs.est.8b06392>, 2019.
- 750 van Donkelaar, A., Hammer, M. S., Bindle, L., Brauer, M., Brook, J. R., Garay, M. J., Hsu, N. C., Kalashnikova, O. V., Kahn, R. A., Lee, C., Levy, R. C., Lyapustin, A., Sayer, A. M., and Martin, R. V.: Monthly Global Estimates of Fine Particulate Matter and Their Uncertainty, *Environ. Sci. Technol.*, 55, 15287–15300, <https://doi.org/10.1021/acs.est.1c05309>, 2021.
- 755 Duncan, B. N., Logan, J. A., Bey, I., Megretskaya, I. A., Yantosca, R. M., Novelli, P. C., Jones, N. B., and Rinsland, C. P.: Global budget of CO, 1988–1997: Source estimates and validation with a global model, *J. Geophys. Res. Atmospheres*, 112, <https://doi.org/10.1029/2007JD008459>, 2007.
- Eyring T., Shepherd and D. Waugh (Eds.): SPARC CCMVal Report on the Evaluation of Chemistry-Climate Models. V. SPARC Report No. 5, WCRP-30/2010, WMO/TD – No. 40, available at www.sparc-climate.org/publications/sparc-reports/, 2010.
- 760 Fuller, R., Landrigan, P. J., Balakrishnan, K., Bathan, G., Bose-O'Reilly, S., Brauer, M., Caravanos, J., Chiles, T., Cohen, A., Corra, L., Cropper, M., Ferraro, G., Hanna, J., Hanrahan, D., Hu, H., Hunter, D., Janata, G., Kupka, R., Lanphear, B., Lichtveld, M., Martin, K., Mustapha, A., Sanchez-Triana, E., Sandilya, K., Schaeffli, L., Shaw, J., Seddon, J., Suk, W., Téllez-Rojo, M. M., and Yan, C.: Pollution and health: a progress update, *Lancet Planet. Health*, 6, e535–e547, [https://doi.org/10.1016/S2542-5196\(22\)00090-0](https://doi.org/10.1016/S2542-5196(22)00090-0), 2022.
- 765 Gidden, M. J., Riahi, K., Smith, S. J., Fujimori, S., Luderer, G., Kriegler, E., van Vuuren, D. P., van den Berg, M., Feng, L., Klein, D., Calvin, K., Doelman, J. C., Frank, S., Fricko, O., Harmsen, M., Hasegawa, T., Havlik, P., Hilaire, J., Hoesly, R., Horing, J., Popp, A., Stehfest, E., and Takahashi, K.: Global emissions pathways under different socioeconomic scenarios for use in CMIP6: a dataset of harmonized emissions trajectories through the end of the century, *Geosci. Model Dev.*, 12, 1443–1475, <https://doi.org/10.5194/gmd-12-1443-2019>, 2019.
- 770 Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korokin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 database – automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements, *Atmospheric Meas. Tech.*, 12, 169–209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.
- 775 Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S.-J.: Sources and distributions of dust aerosols simulated with the GOCART model, *J. Geophys. Res. Atmospheres*, 106, 20255–20273, <https://doi.org/10.1029/2000JD000053>, 2001.

- 780 Gliß, J., Mortier, A., Schulz, M., Andrews, E., Balkanski, Y., Bauer, S. E., Benedictow, A. M. K., Bian, H., Checa-Garcia, R., Chin, M., Ginoux, P., Griesfeller, J. J., Heckel, A., Kipling, Z., Kirkevåg, A., Kokkola, H., Laj, P., Le Sager, P., Lund, M. T., Lund Myhre, C., Matsui, H., Myhre, G., Neubauer, D., van Noije, T., North, P., Olivieri, D. J. L., Rémy, S., Sogacheva, L., Takemura, T., Tsigaridis, K., and Tsyro, S. G.: AeroCom phase III multi-model evaluation of the aerosol life cycle and optical properties using ground- and space-based remote sensing as well as surface in situ observations, *Atmospheric Chem. Phys.*, 21, 87–128, <https://doi.org/10.5194/acp-21-87-2021>, 2021.
- 785 Gong, S. L.: A parameterization of sea-salt aerosol source function for sub- and super-micron particles, *Glob. Biogeochem. Cycles*, 17, <https://doi.org/10.1029/2003GB002079>, 2003.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chem. Phys.*, 6, 3181–3210, <https://doi.org/10.5194/acp-6-3181-2006>, 2006.
- 790 Hammer, M. S., van Donkelaar, A., Li, C., Lyapustin, A., Sayer, A. M., Hsu, N. C., Levy, R. C., Garay, M. J., Kalashnikova, O. V., Kahn, R. A., Brauer, M., Apte, J. S., Henze, D. K., Zhang, L., Zhang, Q., Ford, B., Pierce, J. R., and Martin, R. V.: Global Estimates and Long-Term Trends of Fine Particulate Matter Concentrations (1998–2018), *Environ. Sci. Technol.*, 54, 7879–7890, <https://doi.org/10.1021/acs.est.0c01764>, 2020.
- Helfand, H. M. and Schubert, S. D.: Climatology of the Simulated Great Plains Low-Level Jet and Its Contribution to the Continental Moisture Budget of the United States, 1995.
- 795 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.
- 800 Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET—A Federated Instrument Network and Data Archive for Aerosol Characterization, *Remote Sens. Environ.*, 66, 1–16, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.
- Hsu, N. C., Jeong, M.-J., Bettenhausen, C., Sayer, A. M., Hansell, R., Seftor, C. S., Huang, J., and Tsay, S.-C.: Enhanced Deep Blue aerosol retrieval algorithm: The second generation, *J. Geophys. Res. Atmospheres*, 118, 9296–9315, <https://doi.org/10.1002/jgrd.50712>, 2013.
- 805 IGAC/SPARC, REF1: Proposed Chemistry-Climate Model Initiative Simulations in support of the 2022 WMO/UNEP Scientific Assessment of Ozone Depletion, Accessible at: https://blogs.reading.ac.uk/ccmi/files/2021/04/CCMI-2022_Scenarios_proposal_20210422.pdf, 2022.
- 810 IPCC: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)] Geneva, Switzerland, pp. 35–115, doi: 10.59327/IPCC/AR6-9789291691647., 2023.
- Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B., and Lin, J.-T.: Global distribution of sea salt aerosols: new constraints from in situ and remote sensing observations, *Atmospheric Chem. Phys.*, 11, 3137–3157, <https://doi.org/10.5194/acp-11-3137-2011>, 2011.
- Jenny L. Hand, et al.: Spatial and Seasonal Patterns and Temporal Variability of Haze and its Constituents in the United States Report VI, Interagency Monitoring of the protected Visual Environment (IMPROVE), 2023.

J.L. Hand: IMPROVE DATA USER GUIDE 2023 (VERSION 2), 2023.

- 820 Kahn, R. A., Andrews, E., Brock, C. A., Chin, M., Feingold, G., Gettelman, A., Levy, R. C., Murphy, D. M., Nenes, A., Pierce, J. R., Popp, T., Redemann, J., Sayer, A. M., da Silva, A. M., Sogacheva, L., and Stier, P.: Reducing Aerosol Forcing Uncertainty by Combining Models With Satellite and Within-The-Atmosphere Observations: A Three-Way Street, *Rev. Geophys.*, 61, e2022RG000796, <https://doi.org/10.1029/2022RG000796>, 2023.
- 825 Kim, P. S., Jacob, D. J., Fisher, J. A., Travis, K., Yu, K., Zhu, L., Yantosca, R. M., Sulprizio, M. P., Jimenez, J. L., Campuzano-Jost, P., Froyd, K. D., Liao, J., Hair, J. W., Fenn, M. A., Butler, C. F., Wagner, N. L., Gordon, T. D., Welti, A., Wennberg, P. O., Crounse, J. D., St. Clair, J. M., Teng, A. P., Millet, D. B., Schwarz, J. P., Markovic, M. Z., and Perring, A. E.: Sources, seasonality, and trends of southeast US aerosol: an integrated analysis of surface, aircraft, and satellite observations with the GEOS-Chem chemical transport model, *Atmospheric Chem. Phys.*, 15, 10411–10433, <https://doi.org/10.5194/acp-15-10411-2015>, 2015.
- Koster, R. D., Suarez, M. J., Ducharne, A., Stieglitz, M., and Kumar, P.: A catchment-based approach to modeling land surface processes in a general circulation model: 1. Model structure, *J. Geophys. Res. Atmospheres*, 105, 24809–24822, <https://doi.org/10.1029/2000JD900327>, 2000.
- 830 Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E., and Liss, P. S.: An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, *Glob. Biogeochem. Cycles*, 25, <https://doi.org/10.1029/2010GB003850>, 2011.
- Latimer, R. N. C. and Martin, R. V.: Interpretation of measured aerosol mass scattering efficiency over North America using a chemical transport model, *Atmospheric Chem. Phys.*, 19, 2635–2653, <https://doi.org/10.5194/acp-19-2635-2019>, 2019.
- 835 Levy, R., Hsu, C., and et al.: MODIS Atmosphere L2 Aerosol Product, NASA MODIS Adaptive Processing System, Goddard Space Flight Center [data set], USA, https://doi.org/10.5067/MODIS/MOD04_L2.006, 2015a.
- Levy, R., Hsu, C., and et al.: MODIS Atmosphere L2 Aerosol Product, NASA MODIS Adaptive Processing System, Goddard Space Flight Center [data set], USA, https://doi.org/10.5067/MODIS/MYD04_L2.006, 2015b.
- 840 Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F., and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, *Atmospheric Meas. Tech.*, 6, 2989–3034, <https://doi.org/10.5194/amt-6-2989-2013>, 2013.
- Li, F., Newman, P., Pawson, S., and Perlwitz, J.: Effects of Greenhouse Gas Increase and Stratospheric Ozone Depletion on Stratospheric Mean Age of Air in 1960–2010, *J. Geophys. Res. Atmospheres*, 123, 2098–2110, <https://doi.org/10.1002/2017JD027562>, 2018.
- 845 Li, J., Carlson, B. E., Yung, Y. L., Lv, D., Hansen, J., Penner, J. E., Liao, H., Ramaswamy, V., Kahn, R. A., Zhang, P., Dubovik, O., Ding, A., Lacis, A. A., Zhang, L., and Dong, Y.: Scattering and absorbing aerosols in the climate system, *Nat. Rev. Earth Environ.*, 3, 363–379, <https://doi.org/10.1038/s43017-022-00296-7>, 2022.
- 850 Liss, P. S. and Merlivat, L.: Air-Sea Gas Exchange Rates: Introduction and Synthesis, in: *The Role of Air-Sea Exchange in Geochemical Cycling*, edited by: Buat-Ménard, P., Springer Netherlands, Dordrecht, 113–127, https://doi.org/10.1007/978-94-009-4738-2_5, 1986.
- Liu, J., Strode, S. A., Liang, Q., Oman, L. D., Colarco, P. R., Fleming, E. L., Manyin, M. E., Douglass, A. R., Ziemke, J. R., Lamsal, L. N., and Li, C.: Change in Tropospheric Ozone in the Recent Decades and Its Contribution to Global Total Ozone, *J. Geophys. Res. Atmospheres*, 127, e2022JD037170, <https://doi.org/10.1029/2022JD037170>, 2022.

- 855 Lock, A. P., Brown, A. R., Bush, M. R., Martin, G. M., and Smith, R. N. B.: A New Boundary Layer Mixing Scheme. Part I: Scheme Description and Single-Column Model Tests, 2000.
- Louis, J.-F.: A parametric model of vertical eddy fluxes in the atmosphere, *Bound.-Layer Meteorol.*, 17, 187–202, <https://doi.org/10.1007/BF00117978>, 1979.
- van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A.-L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015), *Geosci. Model Dev.*, 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.
- 860 Molener J. V.: Analysis of the real world performance of the Optec NGN-2 ambient nephelometer. in: *Visual Air Quality: Aerosols and Global Radiation Balance*, pp. 243-265, 1997.
- 865 Molod, A., Takacs, L., Suarez, M., and Bacmeister, J.: Development of the GEOS-5 atmospheric general circulation model: evolution from MERRA to MERRA2, *Geosci. Model Dev.*, 8, 1339–1356, <https://doi.org/10.5194/gmd-8-1339-2015>, 2015.
- Moorthi, S. and Suarez, M. J.: Relaxed Arakawa-Schubert. A Parameterization of Moist Convection for General Circulation Models, 1992.
- Morgenstern, O., Hegglin, M. I., Rozanov, E., O’Connor, F. M., Abraham, N. L., Akiyoshi, H., Archibald, A. T., Bekki, S., Butchart, N., Chipperfield, M. P., Deushi, M., Dhomse, S. S., Garcia, R. R., Hardiman, S. C., Horowitz, L. W., Jöckel, P., Josse, B., Kinnison, D., Lin, M., Mancini, E., Manyin, M. E., Marchand, M., Marécal, V., Michou, M., Oman, L. D., Pitari, G., Plummer, D. A., Revell, L. E., Saint-Martin, D., Schofield, R., Stenke, A., Stone, K., Sudo, K., Tanaka, T. Y., Tilmes, S., Yamashita, Y., Yoshida, K., and Zeng, G.: Review of the global models used within phase 1 of the Chemistry–Climate Model Initiative (CCMI), *Geosci. Model Dev.*, 10, 639–671, <https://doi.org/10.5194/gmd-10-639-2017>, 2017.
- 870 Mortier, A., Glib, J., Schulz, M., Aas, W., Andrews, E., Bian, H., Chin, M., Ginoux, P., Hand, J., Holben, B., Zhang, H., Kipling, Z., Kirkevåg, A., Laj, P., Lurton, T., Myhre, G., Neubauer, D., Olivié, D., von Salzen, K., Skeie, R. B., Takemura, T., and Tilmes, S.: Evaluation of climate model aerosol trends with ground-based observations over the last 2 decades – an AeroCom and CMIP6 analysis, *Atmospheric Chem. Phys.*, 20, 13355–13378, <https://doi.org/10.5194/acp-20-13355-2020>, 2020.
- 875 Nielsen, J. E., Pawson, S., Molod, A., Auer, B., da Silva, A. M., Douglass, A. R., Duncan, B., Liang, Q., Manyin, M., Oman, L. D., Putman, W., Strahan, S. E., and Wargan, K.: Chemical Mechanisms and Their Applications in the Goddard Earth Observing System (GEOS) Earth System Model, *J. Adv. Model. Earth Syst.*, 9, 3019–3044, <https://doi.org/10.1002/2017MS001011>, 2017.
- 880 Oman, L., Waugh, D. W., Pawson, S., Stolarski, R. S., and Nielsen, J. E.: Understanding the Changes of Stratospheric Water Vapor in Coupled Chemistry–Climate Model Simulations, <https://doi.org/10.1175/2008JAS2696.1>, 2008.
- 885 Oman, L. D., Waugh, D. W., Kawa, S. R., Stolarski, R. S., Douglass, A. R., and Newman, P. A.: Mechanisms and feedback causing changes in upper stratospheric ozone in the 21st century, *J. Geophys. Res. Atmospheres*, 115, <https://doi.org/10.1029/2009JD012397>, 2010.
- Pan, X., Ichoku, C., Chin, M., Bian, H., Darmanov, A., Colarco, P., Ellison, L., Kucsera, T., da Silva, A., Wang, J., Oda, T., and Cui, G.: Six global biomass burning emission datasets: intercomparison and application in one global aerosol model, *Atmospheric Chem. Phys.*, 20, 969–994, <https://doi.org/10.5194/acp-20-969-2020>, 2020.
- 890

- Putman, W. M. and Lin, S.-J.: Finite-volume transport on various cubed-sphere grids, *J. Comput. Phys.*, 227, 55–78, <https://doi.org/10.1016/j.jcp.2007.07.022>, 2007.
- Randerson, J.T, G.R. van der Werf, L. Giglio, G.J. Collatz, and P.S. Kasibhatla: Global Fire Emissions Database, Version 4.1 (GFEDv4). ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1293>, 2018.
- 895 Randles, C. A., Silva, A. M. da, Buchard, V., Colarco, P. R., Darmenov, A., Govindaraju, R., Smirnov, A., Holben, B., Ferrare, R., Hair, J., Shinozuka, Y., and Flynn, C. J.: The MERRA-2 Aerosol Reanalysis, 1980 Onward. Part I: System Description and Data Assimilation Evaluation, <https://doi.org/10.1175/JCLI-D-16-0609.1>, 2017.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *J. Geophys. Res. Atmospheres*, 108, <https://doi.org/10.1029/2002JD002670>, 2003.
- 900 Rollins, A. W., Thornberry, T. D., Watts, L. A., Yu, P., Rosenlof, K. H., Mills, M., Baumann, E., Giorgetta, F. R., Bui, T. V., Höpfner, M., Walker, K. A., Boone, C., Bernath, P. F., Colarco, P. R., Newman, P. A., Fahey, D. W., and Gao, R. S.: The role of sulfur dioxide in stratospheric aerosol formation evaluated by using in situ measurements in the tropical lower stratosphere, *Geophys. Res. Lett.*, 44, 4280–4286, <https://doi.org/10.1002/2017GL072754>, 2017.
- 905 Santanello, J. A., Dirmeyer, P. A., Ferguson, C. R., Findell, K. L., Tawfik, A. B., Berg, A., Ek, M., Gentine, P., Guillod, B. P., Heerwaarden, C. van, Roundy, J., and Wulfmeyer, V.: Land–Atmosphere Interactions: The LoCo Perspective, <https://doi.org/10.1175/BAMS-D-17-0001.1>, 2018.
- Strahan, S. E., Duncan, B. N., and Hoor, P.: Observationally derived transport diagnostics for the lowermost stratosphere and their application to the GMI chemistry and transport model, *Atmospheric Chem. Phys.*, 7, 2435–2445, <https://doi.org/10.5194/acp-7-2435-2007>, 2007.
- 910 Strobe, S. A., Douglass, A. R., Ziemke, J. R., Manyin, M., Nielsen, J. E., and Oman, L. D.: A Model and Satellite-Based Analysis of the Tropospheric Ozone Distribution in Clear Versus Convectively Cloudy Conditions, *J. Geophys. Res. Atmospheres*, 122, 11,948–11,960, <https://doi.org/10.1002/2017JD027015>, 2017.
- Strobe, S. A., Ziemke, J. R., Oman, L. D., Lamsal, L. N., Olsen, M. A., and Liu, J.: Global changes in the diurnal cycle of surface ozone, *Atmos. Environ.*, 199, 323–333, <https://doi.org/10.1016/j.atmosenv.2018.11.028>, 2019.
- 915 Ward, R. X., Baliaka, H. D., Schulze, B. C., Kerr, G. H., Crounse, J. D., Hasheminassab, S., Bahreini, R., Dillner, A. M., Russell, A., Ng, N. L., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Poorly quantified trends in ammonium nitrate remain critical to understand future urban aerosol control strategies, *Sci. Adv.*, 11, eadt8957, <https://doi.org/10.1126/sciadv.adt8957>, 2025.
- 920 Scientific Assessment of Ozone Depletion: 2022: <https://csl.noaa.gov/assessments/ozone/2022/>, last access: 28 March 2025.
- Xu, J.-W., Martin, R. V., van Donkelaar, A., Kim, J., Choi, M., Zhang, Q., Geng, G., Liu, Y., Ma, Z., Huang, L., Wang, Y., Chen, H., Che, H., Lin, P., and Lin, N.: Estimating ground-level PM_{2.5} in eastern China using aerosol optical depth determined from the GOCI satellite instrument, *Atmospheric Chem. Phys.*, 15, 13133–13144, <https://doi.org/10.5194/acp-15-13133-2015>, 2015.
- 925 Zhai, S., Jacob, D. J., Brewer, J. F., Li, K., Moch, J. M., Kim, J., Lee, S., Lim, H., Lee, H. C., Kuk, S. K., Park, R. J., Jeong, J. I., Wang, X., Liu, P., Luo, G., Yu, F., Meng, J., Martin, R. V., Travis, K. R., Hair, J. W., Anderson, B. E., Dibb, J. E., Jimenez, J. L., Campuzano-Jost, P., Nault, B. A., Woo, J.-H., Kim, Y., Zhang, Q., and Liao, H.: Relating geostationary satellite measurements of aerosol optical depth (AOD) over East Asia to fine particulate matter (PM_{2.5}): insights from the KORUS-AQ

930 aircraft campaign and GEOS-Chem model simulations, *Atmospheric Chem. Phys.*, 21, 16775–16791,
<https://doi.org/10.5194/acp-21-16775-2021>, 2021.

Zhong, Q., Schutgens, N., van der Werf, G. R., van Noije, T., Bauer, S. E., Tsigaridis, K., Mielonen, T., Checa-Garcia, R., Neubauer, D., Kipling, Z., Kirkevåg, A., Olivie, D. J. L., Kokkola, H., Matsui, H., Ginoux, P., Takemura, T., Le Sager, P., Rémy, S., Bian, H., and Chin, M.: Using modelled relationships and satellite observations to attribute modelled aerosol biases over biomass burning regions, *Nat. Commun.*, 13, 5914, <https://doi.org/10.1038/s41467-022-33680-4>, 2022.

935 Zhu, H., Martin, R. V., van Donkelaar, A., Hammer, M. S., Li, C., Meng, J., Oxford, C. R., Liu, X., Li, Y., Zhang, D., Singh, I., and Lyapustin, A.: Importance of aerosol composition and aerosol vertical profiles in global spatial variation in the relationship between $PM_{2.5}$ and aerosol optical depth, *Atmospheric Chem. Phys.*, 24, 11565–11584, <https://doi.org/10.5194/acp-24-11565-2024>, 2024.

940

Page 7: [1] Deleted Mogno, Caterina (GSFC-614.0)[UNIVERSITY OF MARYLAND BALTIMORE CO] 11/20/25
11:48:00 AM

Page 7: [2] Deleted Mogno, Caterina (GSFC-614.0)[UNIVERSITY OF MARYLAND BALTIMORE CO] 11/6/25 3:27:00
PM

Page 11: [3] Deleted Mogno, Caterina (GSFC-614.0)[UNIVERSITY OF MARYLAND BALTIMORE CO] 11/6/25
4:16:00 PM

Page 19: [4] Deleted Mogno, Caterina (GSFC-614.0)[UNIVERSITY OF MARYLAND BALTIMORE CO] 11/4/25
12:24:00 PM