

Deposition velocity concept does not apply to fluxes of ambient aerosols

Rostislav Kouznetsov¹, Mikhail Sofiev¹, Andreas Uppstu¹, and Risto Hänninen¹

¹Finnish Meteorological Institute, Helsinki, Finland

Correspondence: Rostislav Kouznetsov (Rostislav.Kouznetsov@fmi.fi)

Abstract. The process of dry deposition in chemistry-transport models is usually implemented assuming a proportionality between the deposition flux and the corresponding concentration of a tracer at some reference height. The coefficient of proportionality, called deposition velocity, V_d , is to be parameterized and validated experimentally. ~~We analyse~~ Insufficient attention to the proportionality requirement leads to large discrepancies between field and wind-tunnel measurements of V_d of aerosols with aerodynamic diameters ranging from approximately ~~to~~ 0.1 μm to 2 μm . In seemingly similar conditions, the deposition velocities reported in different experiments may differ by up to two orders of magnitude, with field measurements showing much higher values than experiments performed in controlled environments with known particle properties. We demonstrate that the bulk of the discrepancy can be explained by fast chemical reactions and a particle-to-gas conversion in the immediate vicinity of the surface. ~~By applying~~ With the chemistry-transport model SILAM ~~equipped~~ equipped with gas-particle partitioning for ammonium nitrate, we demonstrate that in ~~the~~ the presence of even small amounts of ammonium nitrate, the vertical flux of total aerosol mass is not controlled by particle deposition but rather by aerosol-gas partitioning in the vicinity of the surface. ~~While there are many other non-conservative~~ Presence of other semi-volatile components in ambient aerosols ~~apart from ammonium nitrate, we demonstrate that the abundance of~~ may further complicate the case, but ammonium nitrate alone is sufficient to ~~render typical ambient aerosol~~ turn "ambient aerosol" into a non-conservative substance. Under these conditions, the deposition flux is not proportional to the concentration, and the concept of deposition velocity as a proportionality coefficient between concentration and deposition flux falls apart. By simulating a renowned field experiment with the SILAM model, we are able to reproduce the magnitudes and temporal behaviors of ambient particle fluxes using the deposition parameterization derived from wind-tunnel studies. Combining these simulations with a set of computational experiments, we suggested guidelines for accounting for the relevant processes in regional atmospheric composition models.

20 1 Introduction

Dry deposition is one of the key mechanisms for removal of gaseous and particulate species from the atmosphere. The measure of dry deposition is the flux F of a substance, expressed as amount (e.g. mass or number) deposited per unit area per unit time. In many applications, the flux is *assumed* to be proportional to the substance concentration C at some reference height above the surface:

$$25 \quad F = V_d(\text{substance, flow, surface}) \cdot C, \tag{1}$$

where V_d is a proportionality coefficient of a velocity dimension, called the deposition velocity. The assumption implies that V_d can be a function of flow, surface, ~~and~~ substance properties, and/or reference height, but not a function of C .

All dry deposition schemes for particles used in atmospheric dispersion models that we are aware of can be expressed in the form of Eq. (??), (e.g. ??????). Eq. (??) follows from the basic law of mass conservation (?) if one assumes: (1) a
30 quasi-steady-state process, (2) horizontal homogeneity at a sufficient scale, (3) absence of sources and sinks of the substance between the reference height and the surface, and (4) zero concentration *at* the surface. The assumptions (1)-(3) taken together are equivalent to the constant-flux assumption, i.e. the flux is constant along the vertical between the reference height and the surface (?). The assumption (4) implies that particles, once reach the surface, get captured regardless of the presence of other particles. If any of these assumptions is violated, the flux proportionality to the concentration at the reference height does not
35 hold, and the concept of deposition velocity ~~becomes~~ is inapplicable.

Along with the definition given by Eq. (??), a wider definition of deposition velocity has been adopted within the ~~flux measurement community (???????)~~ measurement community and some modelling studies (???????).

The deposition velocity $V_{d,a}$ of a substance is defined as the vertical flux F of the substance at a reference height above the surface, normalized with its concentration C at the same height, i.e.

40
$$V_{d,a} = F/C. \tag{2}$$

$V_{d,a}$ is thus a feature of each individual measurement, and computed from the two observed quantities. Moreover, its independence from the concentration postulated in the Eq. (??) is not declared. In the following, we will use the term "apparent deposition velocity" for the one defined by Eq. (??) to distinguish it from the definition given by Eq. (??). These two definitions are equivalent only if all four assumptions behind Eq. (??) hold.

45 Both definitions appeared together for the first time probably in the review of ?, however no explicit distinction between them has been ~~done there~~ made. An extensive review of ? adopts definition (?), then mentions the importance of negligibility of storage and phase changes below the observation heights. ? used definition similar to (??) and explicitly mentioned the requirement of absence of chemical reactions, but in the next sentence V_d was calculated using the average deposition velocity as defined in (?). A recent review of ? uses the definition in form of (??) and mentions that V_d is independent from ambient
50 concentration, but without explicitly specifying that this independence is based on *assumptions* and not guaranteed for arbitrary particles. A few experimental studies (??) clearly mention the requirement of proportionality between the flux and concentration therefore stick to Eq. (??). A substantial number of experimental studies (???????) address deposition velocity without giving an explicit definition, and proportionality between flux and concentration was probably assumed in some of them, but obviously not considered in some others.

55 Many studies have been conducted to infer deposition velocities for different airborne substances in gaseous and aerosol forms under both laboratory and field conditions. They can be classified into two categories: those where the deposition flux is calculated from the amount deposited *onto the surface* (direct method applying Eq. ??), and those where the flux is inferred *above the surface* from gradient or eddy-covariance measurements (indirect method). The direct method is more laborious and requires a well-defined and easily identifiable substance, and it is therefore only applicable in laboratory conditions. ~~While the~~

60 The indirect method can be applied also outdoors, but ensuring that assumptions (1)-(4) hold for the ambient aerosol mixture is non-trivial.

Literature values for deposition velocities of aerosols in the accumulation-mode size range, i.e. from 0.1 μm up to a few micrometers, exhibit a large scatter, which has been puzzling researchers ~~already since several~~ for decades (?). Studies inferring deposition fluxes from direct measurements (????) show that typical values of accumulation mode aerosol deposition velocities
65 onto a smooth surface are of the order of 0.01 cm s^{-1} . These studies were performed in wind tunnels for both smooth solid and water surfaces, and for surfaces with small roughness elements. Deposition velocities of the same order of magnitude were also found in the study by ?, who used a surrogate surface to collect material deposited on a natural lake. On the other hand, field studies that apply flux or gradient methods above water surfaces have reported two orders of magnitude higher deposition velocities of about 1 cm s^{-1} (?). Moreover, observations over snow or low vegetation show a scatter of about two orders
70 of magnitude. While some of them (??) agree with wind-tunnel measurements, others (??) indicate 1–2 orders of magnitude higher velocities.

Indirect measurements of particle deposition onto high vegetation have yielded deposition velocities of 1 cm s^{-1} , with little dependence on the particle size within a range of ~~to~~ 0.2 μm to 5 μm , as summarized by (?). These deposition velocities are two orders of magnitude higher than those predicted by mechanistic models (?). To the best of our knowledge, there are no direct
75 measurements of V_d onto high vegetation.

Many of the original studies considered by ? report strong temporal variability and even a change of direction of the particle fluxes over high vegetation. The reported fraction of upward particle fluxes over forest ecosystems range from 30% to 60% (??????). For this reason many of these studies avoid the term "deposition velocity" in favour of "transfer velocity", while still calling the downward particle flux "deposition".

80 Several mechanisms causing such discrepancies have been suggested to explain the upward fluxes. ? and ? pointed out that vertical fluxes of ambient aerosols can be substantially affected by NH_4NO_3 formation and decomposition due to the $\text{NH}_3 + \text{HNO}_3 \leftrightarrow \text{NH}_4\text{NO}_3$ equilibrium. Among other mechanisms breaking the particle mass conservation are interactions between aerosol particles (coagulation, agglomeration, fragmentation) or aerosol particles and the carrying gas (mass transfer of water vapour, condensation or evaporation of semi-volatile compounds), etc. (?). If any of these mechanisms impacts the
85 vertical flux, it is no longer proportional to the concentration, rendering Eq. (??) inapplicable. This discrepancy has been recognised in many deposition flux studies (e.g. ?). The role of ammonium nitrate ~~in particle flux formation~~ formation in modifying particle fluxes has been studied in details by ?. However, particle concentrations dominated by non-volatile ammonium sulfate were considered sufficient justification to neglect gas-particle partitioning (?). Therefore, apparent deposition velocities reported in field studies have been used to develop parametrisations for particle deposition ~~used in atmospheric transport in~~
90 atmospheric composition models (????). As a result, the obtained dry deposition schemes predict 1 – 2 orders of magnitude higher deposition fluxes than mechanistic models based on "first-principles" (?).

The goal of the present study is to explore the discrepancy between different experimental studies of V_d . We show that the discrepancy does not originate from the natural variability of the particle-surface interaction, but rather from the inconsistency between the deposition velocity defined by Eq. (??) and the apparent deposition velocity defined by Eq. (??). While

95 we primarily focus on the deposition of NH_4NO_3 , our findings should be valid also for other non-inert aerosols. We use a chemistry-transport model SILAM that applies a first-principles based deposition scheme and accounts explicitly for relevant aerosol processes to simulate fluxes and depositions observed in one of the measurement campaigns. This allows us to directly identify the processes responsible for observed particle fluxes (Sec. ??). Then, by using a single-column setup of the model for a system undergoing the reaction $\text{NH}_3 + \text{HNO}_3 \leftrightarrow \text{NH}_4\text{NO}_3$, we demonstrate the difference between the particle deposition process and the processes responsible for particle fluxes above the surface (Sec. ??). Finally, we estimate the amount of in-air NH_4NO_3 needed to break the linear relationship between flux and concentration, and suggest an approach to bridge the gap between observed apparent deposition velocities and deposition parametrisations based on Eq. (??).

2 Methods

To demonstrate the ability of a chemistry-transport model to reproduce the observed range of apparent deposition velocities, we use the Eulerian chemistry transport model SILAM (System for Integrated modelLing of Atmospheric coMposition, <http://silam.fmi.fi>, accessed 26.04.2025). The model features mass-conservative transport schemes (??), and a mechanistic particle deposition scheme derived from first principles (?). It also applies a scheme for secondary inorganic aerosol formation and gas-particle equilibrium that is capable of bi-directional gas-particle partitioning of ammonium nitrate with ammonia and nitric acid (?). NH_4NO_3 is formed when NH_3 and HNO_3 are in abundance, and it decomposes back to the gaseous constituents when the product of their partial pressures drops below a temperature-dependent threshold (?). This simplified approach is consistent with external-mixing representation of aerosols in the model, but ignores the process of bidirectional exchange of NH_3 and HNO_3 gases with (usually aqueous) mixed aerosols that include other than NH_4NO_3 compounds, such as chlorides and sulphates. Such a simplification leads to somewhat overstated gas-phase concentrations in the model, but should not affect the qualitative results of this study. Besides that SILAM does not consider secondary emission of once-deposited NH_3 , and is thus not expected to reproduce the upward flux of NH_3 or NH_4NO_3 -at the surface. The emissions of NH_3 are treated independently of deposition.

The model has been extensively used and evaluated in numerous applications of air quality (?????), atmospheric composition (??) and emergency response (??). Along with regular air-quality evaluation, various setups of the model have been evaluated for depositions and aerosol composition (??).

120 For the present study, we have performed two simulations with SILAM. The first one was a simulation of a field campaign by ? at the Speulder forest site in the Netherlands on June 29-30, 1993. The simulations included regular atmospheric pollutants, with the model setup being similar to that of Copernicus Atmosphere Monitoring Service (CAMS) operational forecasts of SILAM <https://regional.atmosphere.copernicus.eu/> (accessed 26.04.2025). The simulation was driven by ECMWF ERA5 meteorology and nested into global SILAM simulations for the corresponding period. The European model setup generally followed the CAMS configuration (?)-as of 2013, with the MACC anthropogenic emission inventory (?) for 2007. The global simulations closely followed the corresponding operational setup of (?). From the results of the regional run, we extracted con-

centrations and depositions of $\text{PM}_{2.5}$, NH_3 , HNO_3 , NH_4NO_3 for the site location and the time period of the measurements performed by ?, allowing a direct comparison of modelled and measured time series.

130 The second simulation applied an idealized single-column setup to explore the difference between the particle fluxes at the reference height and the actual deposition fluxes at the surface. We simulated the 1D case with the disabled regular transport and wet deposition processes, thus keeping only gas-particle partitioning, vertical diffusion, settling of particles, and dry deposition. The simulation had a vertical resolution of 1 m close to the surface, to enable explicit evaluation of profiles of vertical fluxes and actual deposition. Since the SILAM deposition model for smooth and water surfaces (?) has no tuning parameters, the simulations were performed over water picking a location in Northern Sea (56N,0E). The simulation was driven with the hourly
135 meteorological fields of ERA5 (vertical profiles of wind, temperature and humidity) for 72 hours starting from 2005-06-01. The meteorological fields were used to derive parameters for dry deposition, vertical turbulent diffusion, and gas-particle equilibrium. The domain extended up to 250 m above the surface, at which height a zero-flux boundary condition was enforced.

~~The In both simulations NH_4NO_3 aerosol was simulated as represented with a single bin with of dry diameter. Local. For the Speulder case, 0.3 μm dry diameter was used to match the observed size range, while for the second case 0.7 μm dry diameter was used, as in the operational SILAM air quality simulations.~~
140 The local equilibrium between NH_3 , HNO_3 , and NH_4NO_3 was enforced at every each model time step. To separate diffusion- and chemistry-driven fluxes, we introduced a gaseous chemically-inert substance with diffusion characteristics equivalent to those of SO_2 . For such a gas, the constant-flux assumption is fulfilled in 1D simulations. Then the aerodynamic resistance between two adjacent layers could be evaluated from the deposition flux of the pseudo- SO_2 and the difference in its

145 concentrations between the levels. Having the aerodynamic resistance, one can evaluate the diffusion-driven vertical flux of all species from gradients of their concentrations at the corresponding levels.

Since the model uses a process split, it is important to make sure that the order of the processes and the time step of the simulations does not affect the results. To confirm it, we have performed a series of the simulations with time steps from 5 seconds to 10 minutes. ~~The impact of the time step on the concentrations of the species of interest becomes noticeable at time~~
150 ~~steps longer than one minute and different thicknesses of the vertical levels.~~

The gas-particle equilibrium in SILAM is established instantly. To investigate the effect of this assumption, additional simulations were performed with the equilibrium relaxation time varying from zero up to 10,000 seconds.

3 ~~Simulating~~ Simulation of particle fluxes under real-life conditions

The particle deposition scheme in SILAM (?) predicts deposition velocities for sub-micron particles of a fraction of a millimeter
155 per second. At the same time, other schemes fitted with the data of field ~~experments~~ experiments (???) predict much higher deposition velocities, ~~~~~




Figure 1. Observed apparent-"surface" deposition velocity " V_{ds} " and simulated apparent deposition velocity, concentrations and deposition fluxes of the key species in the campaign of ?. V_{ds} in the original paper stands for the apparent deposition velocity compensated for aerodynamic resistance. For our purpose the difference between V_{ds} and $V_{d,a}$ is negligible. Panel a: a sketch of the timeseries of aerosol fluxes observed by ?, Fig. 1 there with an overlay of the apparent deposition velocity derived from SILAM, panel b: SILAM-predicted concentrations of NH_4NO_3 and total $\text{PM}_{2.5}$; panel c: SILAM-predicted concentrations of HNO_3 and NH_3 ; panel d: SILAM-predicted dry deposition fluxes of HNO_3 and NH_3 .

$\sim 1 \text{ cm s}^{-1}$. Consider a simulation of one of the experimental datasets used to justify high deposition velocities. Figure ??a shows a sketch¹ of the particle-fluxes time series observed by ? above a coniferous forest at the Speulder site in the Netherlands on June 29–30, 1993. ~~Other panels~~ The quantity V_{ds} is so-called surface deposition velocity which is the apparent deposition velocity at the measurement location compensated for aerodynamic resistance between the measurement height and the roughness height. For the SILAM simulation the difference between V_{ds} and $V_{d,a}$ for the particles is negligible due to slow particle deposition in the model. Other panels in Fig. ?? show the concentrations and deposition fluxes of the SILAM simulation for the same time period and location. The model concentrations are those for the near-surface model layer of 25 m thick.

The simulated time series of $\text{PM}_{2.5}$ and NH_4NO_3 indicate that during the considered period, concentrations of NH_3 and HNO_3 were high enough to ~~form~~ allow for particulate NH_4NO_3 to be present. Since the deposition rates of both gases are much higher than that of NH_4NO_3 , the downward particle flux was largely dominated by decomposition of NH_4NO_3 near the surface and the subsequent deposition of NH_3 and HNO_3 . This very process was hypothesised to be the reason for the observed fluxes by ?. The periods of observed strong aerosol "deposition" fluxes closely correspond to periods when the concentrations of NH_4NO_3 and both gases were high in the model. Under these conditions, the modelled deposition flux is controlled by intensively depositing HNO_3 and NH_3 . The resulting near-surface deficit of the gases shifts the gas-particle fractionation and causes disintegration of NH_4NO_3 , which, in turn, creates a net downward flux of the aerosol well-comparable-with-that-is-close to the observed fluxes.

For a comparison, we have plotted an apparent deposition velocity $V_{d,a}$ deduced from the simulation (Figure ??a) as if the total particle flux was fully controlled by the deposition of HNO_3 . This apparent deposition velocity can be deduced from the concentration of $C_{\text{PM}_{2.5}}$ and the ~~dry-deposition~~ dry deposition flux D_{HNO_3} at the surface reported by the model.

$$V_{d,a} = \begin{cases} 0, & \text{if } C_{\text{NH}_4\text{NO}_3} < 1 \mu\text{g m}^{-3}; \\ \frac{M_{\text{NH}_4\text{NO}_3}}{M_{\text{HNO}_3}} \frac{D_{\text{HNO}_3}}{C_{\text{PM}_{2.5}}}, & \text{otherwise.} \end{cases} \quad (3)$$

The ratio of molar masses M is needed to convert the deposition mass flux of HNO_3 to the equivalent mass flux of NH_4NO_3 .

We have chosen a concentration threshold for ~~the~~ NH_4NO_3 of $1 \mu\text{g m}^{-3}$ as a criterion of abundance of particulate phase. Without this threshold, the division of small values caused excessive noise in the time series. The observed ~~triple-peak~~ triple-peak pattern and the magnitude of the simulated $V_{d,a}$ timeseries is well reproduced by the model. The imperfect timing of the peaks can be attributed to the uncertainties in the model, imperfect emission inventories, limited representativeness of point-wise observations and 0.1° grid cell, as well as differences between the ERA5 meteorology and the actual weather conditions at the campaign site.




Figure 2. The temporal evolution of vertical profiles of concentrations of ammonium NH_4NO_3 (a), NH_3 (b), and HNO_3 (c) in the model experiment, and corresponding time series of the deposition fluxes at the surface (d), the downward fluxes at 2 m above the surface (e), and apparent deposition velocity at 2m (f).

The temporal evolution of a system where NH_3 , HNO_3 , and NH_4NO_3 interact to form an equilibrium has been studied in a single-column SILAM simulation. The simulation was initialised ~~at~~ with a constant molar mixing ratio of NH_4NO_3 corresponding to a concentration of $5 \mu\text{g m}^{-3}$. Fig. ?? shows the evolution of the vertical profiles of the ~~mixing-ratios~~ concentrations for the three compounds, as well as their deposition rates and vertical fluxes at 2m. The apparent deposition velocities of the
 190 species, i.e. their vertical fluxes normalised with the corresponding concentrations, are plotted in Fig. ??f.

As seen in Fig. ??, the initial NH_4NO_3 immediately partly decomposes to form an altitude-dependent equilibrium with the gases NH_3 and HNO_3 . The deposition fluxes of the gases are mostly controlled by their respective concentrations in the lowest layer, and gradually decrease as the surface layer becomes depleted. The deposition flux of sub-micron NH_4NO_3 is several orders of magnitude smaller than that of the gases. While the equilibrium allows for the existence of NH_4NO_3 at the
 195 domain bottom, there is a substantial downward aerosol flux replenishing the decomposed amount in the lowest model layer. A ~~remarkable~~ minor upward flux of NH_3 (panel e) is due to its somewhat slower deposition rate compared to HNO_3 in the simulation. As a result, the disintegration of NH_4NO_3 (controlled by the gas-phase removal rate) produces more NH_3 than becomes deposited. ~~The excess~~, so the excess NH_3 diffuses upwards.

The apparent deposition velocity, which would be reported by flux measurements at 2 m above the surface, is about 1 cm s^{-1}
 200 for most of the period, matching the one observed in many field campaigns used by ?. However, the simulations show that the apparent deposition velocity originates from the chemical decomposition of NH_4NO_3 and has no relation to the process of deposition of fine particles.

~~Another~~ A remarkable feature of the simulated particle flux is that it is not proportional to the corresponding concentration. Indeed, once the NH_4NO_3 concentration drops (panels a-c), the apparent deposition velocity of NH_4NO_3 increases and,
 205 theoretically speaking, approaches infinity at the moment when the NH_4NO_3 concentration tends toward zero (panel f). In reality, infinite apparent deposition velocities are not found in outdoor experiments due to the presence of other aerosols and finite gas-particle conversion rates.

As long as NH_4NO_3 is present in the lowest layer, the fluxes of both gases are also heavily affected by the aerosol decomposition, which replenishes the gases while they are deposited. When NH_4NO_3 vanishes, so does the supporting mechanism that
 210 kept the gaseous concentrations stable, and the downward fluxes of both gases become proportional to their concentrations. Therefore, the concept of deposition velocity is not only inapplicable to NH_4NO_3 fluxes, but also to the fluxes of NH_3 and HNO_3 ~~fluxes~~, as long as NH_4NO_3 is present.

5 Discussion

In this section, we consider implications of ~~the existence of non-conservative aerosols on the~~ ambient semi-volatile aerosols on
 215 particle fluxes and apparent deposition velocities.

¹The original paper is copyrighted by the Elsevier Group, which refused to give a permission to use their intellectual property in the openly-distributable publication. Readers are encouraged to reach out to the paper and compare our results to the original figure.

5.1 Idealised systems with gas-particle equilibrium




Figure 3. The process of ammonium nitrate deposition onto the surface. The thickness of the arrows qualitatively represents the magnitude of the flux. NH_4NO_3 flux to the surface appears only if the concentrations of gases at the surface allow for NH_4NO_3 existence

The deposition process described above is shown in Fig. ???. The downward flux of particles is controlled by gas-phase deposition at the surface rather than particle deposition. The resulting near-surface deficit of the gases is replenished via particle-to-gas conversion, which, in turn, creates a deficit of aerosols, leading to a net downward flux of ammonium nitrate.

220 Quite evidently, the presence of any semi-volatile substance with a high deposition rate of the gas phase will lead to the same result: the aerosol flux at a reference height will be controlled by gaseous deposition and particle-to-gas conversion.

While it is clear that gas-particle conversion breaks the proportionality between the fluxes and concentrations for both phases, the question arises: What concentration of a ~~non-conservative aerosol~~ semi-volatile species is needed to dominate the net particle downward flux? To answer this question, let us consider a mixture of NH_4NO_3 and an inert sub-micron aerosol
 225 with a deposition velocity of ~~$V_p = 0.001 \text{ m/s}$~~ and at $V_p = 0.1 \text{ cm s}^{-1}$ and a concentration of C_p . The presence of ammonium nitrate implies that the concentrations of NH_3 and HNO_3 are sufficiently high for its formation, i.e. several $\mu\text{g m}^{-3}$, depending on the temperature. For such a system, the net downward particulate flux at the reference height will be a sum of the fluxes of the inert aerosol and ammonium nitrate:

$$F_{\text{aer}} = F_{\text{NH}_4\text{NO}_3} + F_p = \max(C_{\text{HNO}_3, \text{sat}} V_{\text{HNO}_3}, C_{\text{NH}_3, \text{sat}} V_{\text{NH}_3}) + C_p V_p \quad (4)$$

230 The apparent aerosol deposition velocity can be computed as a ratio of the downward flux and aerosol concentration. Assuming that NH_3 is deposited slower than HNO_3 , one obtains that

$$V_{\text{aer}, a} = \frac{F_{\text{aer}}}{C_{\text{aer}}} = \frac{C_{\text{HNO}_3, \text{sat}} V_{\text{HNO}_3} + C_p V_p}{C_{\text{NH}_4\text{NO}_3} + C_p} \quad (5)$$

If the ammonium nitrate concentration $C_{\text{NH}_4\text{NO}_3}$ is small compared to C_p , the apparent deposition velocity for the total aerosols becomes simply:

$$235 \quad V_{\text{aer,a}} = V_p + V_{\text{HNO}_3} \frac{C_{\text{HNO}_3,\text{sat}}}{C_p}, \quad (6)$$

and the requirement of the small disturbance of the deposition velocity reads as:

$$C_p \gg C_{\text{HNO}_3,\text{sat}} \frac{V_{\text{HNO}_3}}{V_p} \quad (7)$$

This condition does not depend on $C_{\text{NH}_4\text{NO}_3}$ (as long as it is small enough) and, since $V_{\text{HNO}_3} = 0.1 \text{ m/s}$ and $V_p = 0.001 \text{ m/s}$ $V_{\text{HNO}_3} = 10 \text{ cm s}^{-1}$ and $V_p = 0.1 \text{ cm s}^{-1}$, leads to a requirement for the concentration of the inert aerosol $C_p \gg 100 \mu\text{g/m}^3$ $C_p \gg 100 \mu\text{g m}^{-3}$. Such concentrations of inert aerosols are rare even in highly polluted locations. Therefore, in a vast majority of real-life cases, even a minor presence of ammonium nitrate in the air guarantees that the apparent deposition velocity of particles is heavily affected by the particle-to-gas conversion near the ground.

5.2 Effect of finite gas-particle equilibration kinetics

The above calculations were performed assuming instant gas-particle conversion. The actual conversion rates depend on the features of the semi-volatile substance(s) and atmospheric conditions. Equilibration times have been considered by ? for inorganic aerosols and by ? for organic aerosols. The main mechanism for such equilibration is condensation/evaporation of material to/from the particle surface. The rate of this process is controlled by the diffusivity of the gases and by the availability of particle surface. The latter is a function of particle size and concentration.

? concluded that for the considered cases of ambient environment, except for very low relative humidity conditions when particles stayed dry, the conversion of nitric acid to nitrate was completed within 1000 seconds. This time scale is probably relevant for most of ambient conditions, since there is strong evidence for ammonium-nitrate particles not crystallizing at down to a 10 – 20% relative humidity (?), while in the simulations of ? deliquescence occurred at 50%.

For the validity of instantaneous gas-particle partitioning of organic compounds, ? conclude that it is clearly established “for relatively high volatility compounds partitioning into liquid particles, but ... breaks down for partitioning of low and semi-volatile compounds into liquid and semi-solid particles”. For liquid particles the equilibration time scale is also quite short and approaches 1000 s only for particle sizes closer to 1 μm at PM concentrations below a few $\mu\text{g m}^{-3}$.

Using a 1000-s equilibration time as a conservative estimate, one can conclude that the vertical flux of particles with true deposition velocity of 0.1 cm s^{-1} is affected by the gas-particle conversion at a reference height of -,and-detailed-1 m, and explicit calculations are needed to account for it reproduce it.

To evaluate the effect of the resolution and particle kinetics on the simulated particle fluxes and depositions, we implemented a finite relaxation time for gas-particle partitioning into SILAM. The idealized case simulations of Sec ?? were subsequently repeated with several prescribed relaxation times. Sensitivity simulations were performed for the fine-scale vertical layers, used in Sec ??, and for the coarser vertical layers typically used for the regional air quality simulations: four layers of 25 m, 50 m, 100 m and 200 m thick. The vertical fluxes were evaluated at the interface between the lowermost levels of the coarse vertical and the closest interface of the fine-resolution one (25 m above the surface).

The sensitivity simulations were performed with identical initial conditions and meteorological forcing for both verticals and three gas-particle equilibration times of $\tau = 0$ s, 100 s and 10000 s. For each case, the model time step was varying to explore the effect of the time split between the gas-particle transformation and diffusion-deposition processes. Time steps of 12 s, 30 s, 1 min, 2 min, 5 min, 10 min, and 30 min were tested for all setups. For each simulated case, plots similar to the one in Fig. ?? were produced (see Supplementary Materials).

In all cases, the removal of ammonium and nitrate was driven by gas-phase deposition onto the surface, and the corresponding rates were quite insensitive to the vertical resolution, as long as the model time step was sufficiently short: despite the implicit diffusion implementation, a finer vertical resolution required a shorter time step.

Large upward fluxes of the gas-phase compounds visible in the beginning of the simulations with fast equilibration while both phases co-exist, were artificial. The flux arises from the difference in equilibrium gas-phase concentrations different levels due to different temperatures and humidity provided by the meteorological model. This effect was not seen for 2 m fluxes (Fig. ??), since the layers above and below the flux sampling point had the same temperature and humidity.

Long equilibration time substantially reduces the gas-phase deposition at the very beginning of the simulations due to lack of the gas phase to deposit. One can also notice some 20% lower peak deposition rates for $\tau = 10000$ s compared to the instant equilibration case, due to a slower conversion of particles to depositing gases. At $\tau = 100$ s the difference in fluxes and depositions between the instant and non-instant conversion vanishes.

The particle fluxes and apparent deposition velocities at the reference height appear to be quite sensitive to the relaxation time. For shorter relaxation times, gas-particle conversion occurs closer to the surface, so that the deposition of gases is maintained by the downward flux of particles, leading to quite high apparent deposition velocities. The vertical fluxes of particles at the reference height deviate from corresponding surface fluxes even with an unrealistically long relaxation time of several hours. The high sensitivity of the particle fluxes and apparent deposition velocities on the gas-particle conversion rates confirm that these parameters cannot provide a reliable constraint for a deposition parameterization in atmospheric models.

Considering the low sensitivity of deposition fluxes and reference height concentrations of both gases and aerosols to the relaxation time, one can conclude that for air quality and deposition assessment, the instant gas-particle conversion is a valid assumption. There is no need to account for the kinetics of the process unless one is interested in resolving the profiles of gas and particle fluxes separately.

5.3 Intermittency, sign and scaling of particle fluxes

The intermittency of ambient particle fluxes can be clearly seen in the observational time series obtained with fast-response instruments, such as those reported by ? and many others (e.g. ????). One of plausible explanations is that the particle flux is shifting back and forth between the ~~gas-phase and particle-phase driven a regimes due to gas- and particle- phases driven by~~ changes in ammonia and nitrate concentrations ~~, or changes in or by changing~~ ambient temperature and humidity ~~altering,~~ which shift the gas-particle equilibrium point.

In many cases, the observed fluxes of particles change their ~~sign. The upward flux of particles signs manifesting the upward diffusion of aerosols. This~~ can occur due to particle formation ~~from the gas phase in the vicinity of the surface or if near~~




Figure 4. Hourly downwards particle fluxes vs concentrations multiplied with friction velocity, simulated for the Speulder site in the Netherlands for the month of June 1993, same simulation as in Fig. ?? Different markers used for the cases with and without NH_4NO_3 particles.

300 the surface if, for instance, ammonia evaporates from the surface at a sufficient rate, enriching the near-surface layer with ammonium-nitrate particles soil prompting the formation of ammonium nitrate (?).

So far, we have considered externally mixed particles, i.e., or a conservative aerosol. In the presence of internally-mixed particles, that have internally mixed particles with a relatively stable core covered with ammonium nitrate staying in gas-particle equilibrium, opposite-sign fluxes of different particle sizes can occur, the flux direction may depend on the particle sizes. Larger particles enriched with ammonium nitrate would then follow the surface flux of ammonia. If Indeed, if the gas concentration drops below its saturation limit level, the ammonium nitrate shells evaporate, resulting in a shrinkage of the particle size and the formation of an opposite-sign flux of smaller particles. This mechanism was explored by ?, explains the rather strong anti-correlation of particle fluxes of different sizes reported by ? and ?.

305

Several field studies report a linear scaling of particle fluxes with friction velocity u_* , i. e. with the apparent deposition velocity being linear to it(????). This scaling is consistent with the assumption that conclusion that the particle fluxes are controlled by fast-depositing the fast depositing gases, whose deposition scales linearly with the turbulent diffusion intensity in the surface layer.

310

The driving role of the gas-phase deposition is also consistent with the behavior of the apparent deposition velocity, normalized with u_* , as a function of atmospheric stratification. The minimum of the deposition velocity at a slightly-stable stratification has been observed by ? and by (??), with the shape of the function agreeing with the behavior of the stability function for scalar fluxes in the stratified surface layer.

315

5.4 Is the flux proportional to the concentration?

The vast majority of experimental studies on particle fluxes report the fluxes in terms of the apparent deposition velocity, without explicit verification of proportionality between the flux and the concentration. While the proportionality rather naturally holds for well-controlled wind-tunnel studies, it is not guaranteed for ambient particles in outdoor experiments.

320




Figure 5. Mean fraction of NH_4NO_3 in $\text{PM}_{2.5}$ for the year of 2024 (a), and fraction of hourly average values with NH_4NO_3 exceeding $0.1 \mu\text{g m}^{-3}$, according to SILAM forecasts made for the CAMS2-40 regional service

To illustrate the non-trivial nature of this requirement, consider a scatter plot of $\text{PM}_{2.5}$ fluxes at a reference height vs the corresponding $\text{PM}_{2.5}$ concentrations (Fig ??), simulated by SILAM for the month of the (?) study (~~considered in Sec. ??~~). We use different markers for the cases with and without presence of NH_4NO_3 component in $\text{PM}_{2.5}$. As a reference, we plotted a line corresponding to the dimensionless deposition velocity suggested by ? for neutral and stable stratification, i.e.

$$325 \quad E_d \equiv v_d/u_* = 0.002. \quad (8)$$

The points in the plot form two clearly distinguishable clusters corresponding to the two aforementioned regimes ~~of the flux~~. The lower cluster corresponds to the flux of the particles in the absence of ammonium nitrate, rendering them inert in the model. ~~The cluster indicates~~ This cluster manifests a high correlation between the flux and the concentration. The residual scatter is caused by the different size distributions and water contents of the $\text{PM}_{2.5}$ components, as well as the varying meteorological conditions, ~~in accordance to the SILAM deposition scheme (?).~~ The upper cluster corresponds to the regime of gas-phase controlled particle flux, ~~exhibiting~~ much higher fluxes, a two orders of magnitude larger scatter of the values, and a much less pronounced dependency on the concentration.

We have not found any study with such a scatter plot ~~of for~~ the observed fluxes, or another way of examination of the proportionality between the flux and the concentration. However, time series ~~that can be found~~ in many studies (e.g ??), suggest indicate that the correlation between fluxes and concentrations is rather low in ~~the~~ outdoor studies.

5.5 ~~Oecurrence~~ Abundance of ammonium-nitrate

To estimate the frequency of ~~the occurrence of the near-surface ammonium nitrate~~ nonzero concentrations of ammonium nitrate in the surface layer, we analyzed ~~the~~ SILAM operational air quality forecasts for 2024 ~~performed within the framework of the~~ Copernicus CAMS2-40 regional operational service (?). The service consists of 11 European ~~air-quality~~ air quality models that
340 produce daily forecasts of atmospheric pollution at a spatial resolution of $0.1^\circ \times 0.1^\circ$ and ~~at~~ a temporal resolution of one hour. The forecasts have been extensively evaluated against multiple types of observational data ~~Copernicus publishes quarterly reports on the performance of each model~~ (<https://atmosphere.copernicus.eu/regional-services>, accessed 15.4.2025), ~~and has an extensive tool for online model evaluation~~ (<https://regional-evaluation.atmosphere.copernicus.eu/>, accessed 15.4.2025). Apart from the regulated air pollutants, the model performance has been evaluated for a wide range of atmospheric trace gases
345 and aerosols. The service provides the ~~by far~~ most reliable forecasts ~~for of~~ atmospheric pollution in Europe, and SILAM is among the ~~best-performing~~ best performing models of the ensemble.

For the analysis, we ~~chose the first day of the SILAM forecasts over the whole year and~~ extracted the near-surface concentrations of $\text{PM}_{2.5}$ and NH_4NO_3 ~~Figure ??a shows the mean relative contribution of ammonium nitrate to the~~ and evaluated the mean contribution of ammonium nitrate to total $\text{PM}_{2.5}$ ~~over the year~~ (Figure ??a). One can see that for most of the European
350 territory, the fraction of ~~ammonium nitrate~~ ammonium nitrate is within the range of 10–30%, whereas the mere presence of ammonium nitrate ~~aerosol~~ is an indicator of the gas-particle equilibrium strongly affecting the deposition fluxes. The frequency of such occurrences was estimated from the fraction of the hourly NH_4NO_3 concentrations exceeding $0.1 \mu\text{g m}^{-3}$ (Figure ??b). For ~~the most part~~ most of the European territory, NH_4NO_3 aerosol is present in the air ~~near the surface~~ for more than 30% of time. The particle fluxes over the Speulder forest site in the Netherlands ~~studied by ?~~, are impacted by NH_4NO_3 for more
355 than 50% of the time, at the Hyytiälä site in central Finland (?????) — for more than 20% of the time, at the Waldstein site in Bavaria (?) for over 50% of the time, and at the Yatir forest research station in Israel (?) for ~~about 10 to 20~~ 10-20% of the time.

6 Recommendations

Based on the above findings and considerations one can formulate recommendations on a consistent way to simulate particles and semi-volatile compounds and their depositions in chemistry-transport models. For such simulations, a model should
360 have implementation of the gas-particle partitioning and deposition processes for both phases. An instant-equilibrium model for partitioning provides a good approximation if one is interested in bulk depositions of ammonium and/or nitrates. For applications analyzing profiles of gaseous and particulate fluxes, more sophisticated schemes with finite conversion rates may be preferable.

A dry deposition scheme of the form (??) should not use values of "effective deposition velocity" or apparent deposition
365 velocity obtained for poorly-known particles. Application of such values would heavily over-deposit inert particles, while being applied to semi-volatile substances it would enforce the false linear relationship between corresponding fluxes and concentrations.

370 Parameterizations of V_d should be calibrated and evaluated with the data on particles, for which the deposition velocity concept is applicable, i.e., inert aerosols not affected by chemical or phase-transition processes. Such parameterizations can be derived from wind tunnel studies with well-identified and controlled particle features. One can also use mechanistic considerations to obtain the V_d dependencies. Such schemes have a very limited number of tuning parameters and agree well with the V_d values observed for passive particles (??).

375 Ambient observations of aerosol concentrations and fluxes, while not measuring the deposition fluxes themselves, can provide invaluable material for model evaluation. One should just use the full chemical and physical transformation schemes, as well as corresponding emission data, while simulating these cases.

Sensitivity studies reveal only a weak dependence of the deposition flux calculations on vertical resolution near the surface and the model time step. However, the first model layer of 10-20 m thickness can be suggested as a good compromise between simulation costs and accuracy. The model time step will, for the vast majority of cases, be decided by a relaxation time of this layer.

380 As demonstrated above, such a setup, while relying on seemingly very low deposition rates of aerosols, can successfully reproduce the observed fluxes of particles under ambient conditions. Moreover, together with gas-particle partitioning, this approach reproduces observed intermittency and bidirectional fluxes of particles above natural surfaces.

7 Conclusions

We identified ~~a terminological inconsistency in the definition of the deposition velocity~~. For the modeling community an inconsistency in using the concept of deposition velocity in the literature. For modeling applications, the concept of deposition velocity implies a proportionality between the deposition flux and the atmospheric concentration, ~~and all parameterizations of dry deposition velocity rely on this assumption~~ where V_d is a proportionality coefficient. On the other hand, many experimental studies define the deposition velocity as ~~just~~ a ratio of observed downward flux and concentration, ~~formally calculated from the observed variables and being~~ specific for each measurement ("apparent deposition velocity" $V_{d,a}$). We have not found any experimental study where the proportionality between the observed fluxes and the concentrations was explicitly examined across the campaign. ~~Time~~ Conversely, the time series available from some studies suggest that the correlation between fluxes and concentrations is rather low.

There are several mechanisms that break the proportionality. We considered one of them, the evaporation of particles between the measurement height and the surface due to the particle-to-gas transition of ammonium nitrate. SILAM model simulations of ~~both an~~ idealized and real-life cases demonstrated that the vertical flux of particles is frequently often not proportional to the particle concentration at a reference height. ~~The proportionality is broken by the sink of particles between the measurement height and the surface, due to gas-particle transformation of ammonium nitrate.~~

The SILAM model, with the particle deposition scheme based on first principles and explicit gas-particle equilibration for ammonium nitrate, is capable of reproducing observed fluxes of ambient aerosols that ~~corresponds to velocity of~~. However, the correspond to apparent deposition velocity of 1 cm s^{-1} , usually reported by outdoor studies. At the same time, the SILAM

aerosol deposition velocity ~~in the model is of the order of~~. Moreover, ~~the model is about~~ 0.01 cm s^{-1} . ~~The model also~~ reproduces the intermittent nature and temporal evolution of the observed particle fluxes.

We have shown that ~~very low concentration even very small amount~~ of ammonium nitrate is sufficient to break the flux conservation ~~, as because~~ it is broken rather by the presence of sufficiently high concentrations of nitric acid and ammonia, which trigger the gas-particle conversion. ~~Data on the abundance of ammonium nitrate over Europe from the CAMS2-40 European service suggest~~ Copernicus AQ forecasts for Europe show that ammonium nitrate ~~conversion occurs over Europe is present in the air~~ for about half of the time:-

~~, which implies that at least a fraction of outdoor particle-flux measurements are affected by gas-particle conversion.~~ In addition to ammonium nitrate, the presence of any semi-volatile substance may break the particle mass conservation near the surface ~~following the same mechanism~~.

In laboratory experiments, the nature of the aerosol is usually well known, so the assumption of particle conservation ~~holds. In natural conditions, the~~ can be ensured. ~~In ambient conditions, this~~ assumption should be explicitly tested by evaluating whether the particle concentrations and corresponding fluxes are proportional to each other ~~for each specific set of conditions~~.

The presence of gas-particle conversion is sufficient to explain ~~the~~ 100-fold differences between the apparent deposition velocities obtained in different experiments for similar surfaces and aerosol sizes.

The concept of deposition velocity (in a sense of Eq. ??) ~~is a very powerful tool to~~ can reduce the complexity of dry deposition schemes in models. However, ~~it should be accompanied with phase conversion calculations to make the scheme applicable~~ under ambient conditions, ~~the presence of where~~ semi-volatile species ~~breaks the aerosol~~ break the aerosol mass conservation, and ~~where~~ the observed vertical fluxes cannot be directly used to infer the dry deposition velocity ~~for models~~. Nevertheless, ~~well documented well documented~~ observations of ambient fluxes provide invaluable means for ~~the~~ assessment of chemistry-transport models as a whole, when all relevant processes are considered, ~~including gas-particle conversion near the surface~~.

Code and data availability. The code of SILAM model that can be used to reproduce the results of the current study is available from GitHub <https://github.com/fmidev/silam-model> under the GPLv3 public license, which in particular allows for reusing the code in other models. The specific revision used for this paper has been archived at Zenodo <https://doi.org/10.5281/ZENODO.14608973>, (access 06.06.2025), (?). The implementation of the particle dry deposition scheme (?) can be found from `depositions.silam.mod.f90` file, and implementation of ammonium-nitrate partitioning in `aerosol_dynamics_simple.f90`.

Author contributions. RK performed the numerical simulations and evaluations, wrote the initial text and prepared the figures. MS contributed to the case conceptualization, participated in writing and editing of the manuscript. AU and RH participated in the model development and editing the manuscript.

Competing interests. The authors declare that no competing interests are present.

Disclaimer. The paper represents the authors' personal opinions and views, which might or might not agree with the positions of their organizations.

Acknowledgements. The study was ~~supported by the~~ funded by the taxpayers of the European Union via Horizon Europe-funded ~~CAMAERA~~ project (No-projects CAMAERA (grant No. 101134927) and RI-URBANS (grant No. 101036245) and by the taxpayers of Finland via the Research Council of Finland project VFSP-WASE (grant No 359421), which we gratefully acknowledge.