

Replies to Referees

Liisa Juusola^{1,2}, Ilkka Virtanen³, Spencer Mark Hatch⁴, Heikki Vanhamäki³, Maxime Grandin², Noora Partamies⁵, Urs Ganse¹, Ilja Honkonen², Abiyot Workayehu¹, Antti Kero⁶, and Minna Palmroth^{1,2}

¹Department of Physics, University of Helsinki, Helsinki, Finland

²Finnish Meteorological Institute, Helsinki, Finland

³Space Physics and Astronomy Unit, University of Oulu, Oulu, Finland

⁴Department of Physics and Technology, University of Bergen, Bergen, Norway

⁵Department of Arctic Geophysics, University Centre in Svalbard (UNIS), Longyearbyen, Norway

⁶Sodankylä Geophysical Observatory, University of Oulu, Sodankylä, Finland

Reply to Referee #1

We thank Referee #1 for constructive comments and agree with almost all of them. Please, see below for our point-by-point replies. The original review is written in *black* and our replies in blue. The line numbers given refer to the original submitted manuscript, unless otherwise stated.

5

This paper presents a thorough and mathematically rigorous analysis of the relation between ground magnetic perturbations and ionospheric conductances measured by the EISCAT radars. The statistics are carefully performed, and the separation of the equivalent currents into divergence-free and curl-free parts is a significant improvement over prior similar studies.

10 *There are several drawbacks to the model that should be addressed in more detail in the discussion section as noted in the attached annotated file. Nevertheless, the model described here can be very useful for the study of individual events localized over suitably spaced ground magnetometers.*

Overall, this paper represents a significant contribution that will facilitate future studies and improved insights on auroral electrodynamic processes. I consider the paper publishable after the comments in the attached file are addressed.

The comments extracted from the file:

15 *Line 26: difficult 'to' obtain*

We have corrected the typo.

Line 96: What was the intrinsic range resolution of the radar measurements?

Range resolution of the fitted plasma parameters depends on altitude and may vary between radar modes. The resolution is typically about 3 km in the lower E region (below 110 km altitude) and about 20 km at 330 km altitude.

20 We have added this explanation to the text.

Line 96: This is rather arbitrary. What percentage of the profiles were eliminated by this test, and how would that percentage change with different thresholds. Is the test done only in the altitude range up to 330 km? Does the presence of satellite echoes above this altitude affect the electron densities measured at other altitudes?

25 The limit was chosen to be close to, but clearly above, the largest density one may realistically expect in the F region. The test was done only for the interpolated data up to 330 km altitude. Also satellites above 330 km altitude may affect




Figure 1. Fraction of rejected points as function of the electron density threshold.

the densities measured below 330 km, because satellite echoes from above may be aliased below 330 km in the radar signal processing.

The fraction of profiles rejected based on this criterion was 0.066% for the UHF and 0.035% for the ESR. Fraction of rejected points as function of the threshold is shown in the attached Figure 1. The deep decline at thresholds smaller than $3 \cdot 10^{12} \text{ m}^{-3}$ is believed to be from largest real electron densities, while $5 \cdot 10^{12} \text{ m}^{-3}$ is close to beginning of a plateau probably caused by satellite echoes and other outliers.

We have added this explanation (without the figure) to the text.

Line 260: Why are there so many conductance values below that given by the Moen and Brekke model? Weren't they also based on EISCAT measurements?

Yes, the Moen and Brekke (1993) model is based on EISCAT measurements, but only on nine days of data (cf. line 56). Most likely the limited amount of data causes the model to be less than ideal for describing the larger data set we now have. This observation was part of the motivation for the current study (cf. line 69).

Line 409: The

We have corrected the typo.

40 Line 416: The authors should consider comparing the field-aligned current determinations from those derived from AMPERE. This would help highlight the importance of spatial scales in the conductances and electrodynamic modeling results.

Thank you for a very good suggestion, we have added AMPERE data to Fig. 13.

Line 436: There is no discussion of the effects of a neutral wind on the calculations.

45 Thank you for pointing this out. We have added the following discussion:

“Neutral wind v_n perpendicular to the ionospheric background magnetic field B_0 generally needs to be considered in ionospheric electrodynamics (e.g., Hatch et al., 2024). For example, in a coordinate system where the neutral wind is not zero, Ohm’s law (Eq. 3) becomes

$$\mathbf{J} = \Sigma_P(\mathbf{E} + \mathbf{v}_n \times \mathbf{B}_0) + \Sigma_H(\mathbf{E} + \mathbf{v}_n \times \mathbf{B}_0) \times \hat{\mathbf{e}}_r. \quad (1)$$

50 Neutral wind is not required to estimate the electron density N_e from EISCAT observations and, thus, to estimate the conductivities with Eqs 6 and 7. The Ebec model is parametrized with electric current, which is independent on the frame of reference determined by the neutral wind. Thus, neutral wind is not expected to introduce errors to the conductances given by Ebec, and the electric field estimated from equivalent current density and Ebec conductances is given in the frame of reference where the neutral wind is zero.”

55 Line 459: While artificially introducing gradients improves the agreement with Swarm, this represents an ad hoc solution that cannot be used in the absence of satellite measurements.

This is true, and for this reason we have included the demonstration as part of the discussion only.

Line 477: Some discussion should be added of the recent models that correlate conductances with field-aligned currents, thus ensuring the gradients are self-consistent with the parallel currents. The difference in spatial resolution between the EISCAT measurements and the equivalent currents derived from magnetometers is a significant drawback to the model presented here.

60 Thank you for the suggestion, we have added the following discussion and a new Figure 15:

“Recent models by Robinson et al. (2020) and Wang and Zou (2022) correlate conductances from the Poker Flat Incoherent Scattering Radar (PFISR) with field-aligned currents from AMPERE and Swarm, respectively, thus ensuring the conductances and their gradients are self-consistent with the parallel currents. The latitudinal scale of AMPERE field-aligned currents is about 3° (Waters et al., 2020) and field-aligned current variations with a latitudinal scale of 150 km were used by Wang and Zou (2022). Thus, although the difference in spatial resolution between the EISCAT measurements and the equivalent currents derived from IMAGE magnetometers is a significant drawback to the Ebec model, the models parametrized by field-aligned currents also have similar limitations.

70 Figure 15 shows Σ_P and Σ_H as given by the conductance models of Robinson et al. (2020) and Wang and Zou (2022) when applied to the AMPS field-aligned current density (Fig. 10d). Note that due to the large differences in conductance amplitudes, the color scales are not the same as those in Fig. 9b–e. Apart from the amplitudes, the most obvious difference in the conductances given by Robinson et al. (2020) and Wang and Zou (2022) compared to SWIPE and Ebec is the latitudinally wide double oval structure caused by parametrizing by both upward and downward field-aligned current density. The coefficients of the COMPASS model by Wang and Zou (2022) are defined for three-hour MLT sectors, resulting in the discontinuous conductance distributions that could be smoothed, e.g., by interpolation.”

75

Line 486: The authors should note that the overall agreement with Ahn et al. suggests that these relations are not dependent on the geographic locations of the magnetometer measurements.

Do you mean that this indicates that the internal contribution to magnetometer measurements is not significant?

The internal part is typically some tens of percent of the total horizontal variation magnetic field amplitude, and directed such that it strengthens the external horizontal field. However, this depends much on geographic location. For example, it is possible that the internal contribution to the magnetometer data used by Ahn et al. (1998) may have been particularly small. Without a more thorough investigation, we do not think that an overall agreement between the Ahn et al. (1998) and our model indicates that the internal contribution to magnetometer measurements in general can be considered insignificant.

Line 497: The authors should discuss whether this method can help model the extreme magnetic perturbations associated with GICs. Is it possible these GIC events are associated with large EDF values?

Indeed, this is the case as discussed in Juusola et al. (2025). We have added:

“Especially, the possibility of extracting E_{DF} on the ground surface (Juusola et al., 2025) allows detailed studies of extreme magnetic perturbation events associated with GICs.”

Figure 8: The vertical scale should be adjusted to better show the spread in conductance values.

We have adjusted the scale.

In addition, we have updated the discussion paper reference Juusola et al. (2024) to the final version Juusola et al. (2025), and corrected a few typos.

Reply to Referee #2

We thank Referee #2 for constructive comments and agree with all of them. Please, see below for our point-by-point replies. The original review is written in *black* and our replies in blue.

The paper introduced a new empirical model which can predict conductance based on divergence-free part of the horizontal current density. They built the model using EISCAT observations. They further introduced a method to further solve the field-aligned currents and potential. They validated the model against other empirical models and SWARM observations. Even though the validation result is not perfect, the method itself is novel. If more details about the methodology can be added, I will suggest publishing it.

Minor suggestions:

Section 3.2: More introduction to AMPS and SWIPE in the first paragraph would be helpful. For example, what is the input and output of these models. For example, is E_{DF} assumed to be zero in AMPS?

We have added: “Both AMPS and SWIPE are parametrized with magnetic latitude, MLT, dipole tilt angle, solar wind speed, $F_{10.7}$, and Interplanetary Magnetic Field (IMF) $B_{GSM,y}$ and $B_{GSM,z}$. AMPS provides the DF and CF part of the horizontal ionospheric current density and the field-aligned current density. SWIPE provides the ionospheric Hall and Pedersen conductances and horizontal potential electric field and corresponding potential. E_{DF} is assumed to be zero.”

Introduce the name of the model at the beginning. Usage of AMPS-EISCAT is a bit confusing.

Thank you for the suggestion, we have changed this accordingly.

Font size in Figure 11 and 12 is too small. How are the internal and external E_{DF} separated?

We have increased the font size.

115

We have added the following explanations in section 2.3 to clarify how the internal and external E_{DF} separated:

Line 173 in the old version / line 180 in the new vesion: “Using two DF SECS layers, one at 1 m depth and the other at 90 km altitude, will separate the external and internal contributions to the magnetic field variations.”

120

Line 180 in the old version / line 188 in the new version: “The DF SECS amplitudes of the telluric equivalent current sheet at 1 m depth can be used to calculate the internal contribution to E_{DF} and the DF SECS amplitudes of the ionospheric equivalent current sheet at 90 km altitude can be used to calculate the external contribution to E_{DF} . The total E_{DF} is then obtained as a sum of the internal and external contributions.”

In addition, we have updated the discussion paper reference Juusola et al. (2024) to the final version Juusola et al. (2025), and corrected a few typos.

References

- 125 Ahn, B.-H., Richmond, A. D., Kamide, Y., Kroehl, H. W., Emery, B. A., de la Beaujardière, O., and Akasofu, S.-I.: An ionospheric conductance model based on ground magnetic disturbance data, *J. Geophys. Res.*, 103(A7), 14 769–14 780, <https://doi.org/https://doi.org/10.1029/97JA03088>, 1998.
- Hatch, S. M., Vanhamäki, H., Laundal, K. M., Reistad, J. P., Burchill, J. K., Lomidze, L., Knudsen, D. J., Madelaire, M., and Tesfaw, H.: Does high-latitude ionospheric electrodynamics exhibit hemispheric mirror symmetry?, *Ann. Geophys.*, 42, 229–253, <https://doi.org/https://doi.org/10.5194/angeo-42-229-2024>, 2024.
- 130 Juusola, L., Vanhamäki, H., Marshalko, E., Kruglyakov, M., and Viljanen, A.: Estimation of the 3-D geoelectric field at the Earth’s surface using Spherical Elementary Current Systems, *EGUsphere*, [preprint], <https://doi.org/https://doi.org/https://doi.org/10.5194/egusphere-2024-2831>, 2024.
- Juusola, L., Vanhamäki, H., Marshalko, E., Kruglyakov, M., and Viljanen, A.: Estimation of the 3-D geoelectric field at the Earth’s surface using Spherical Elementary Current Systems, *Ann. Geophys.*, 43, 271–301, <https://doi.org/https://doi.org/10.5194/angeo-43-271-2025>, 2025.
- 135 Moen, J. and Brekke, A.: The solar flux influence on quiet time conductances in the auroral ionosphere, *Geophysical Research Letters*, 20, 971–974, <https://doi.org/https://doi.org/10.1029/92GL02109>, 1993.
- Robinson, R. M., Kaeppler, S. R., Zanetti, L., Anderson, B., Vines, S. K., Korth, H., and Fitzmaurice, A.: Statistical relations between auroral electrical conductances and field-aligned currents at high latitudes, *Journal of Geophysical Research: Space Physics*, 125, e2020JA028 008, <https://doi.org/https://doi.org/10.1029/2020JA028008>, 2020.
- 140 Wang, Z. and Zou, S.: COMPASS: A new CONductance Model based on PFISR And SWARM Satellite observations, *Space Weather*, 20, e2021SW002 958, <https://doi.org/https://doi.org/10.1029/2021SW002958>, 2022.
- Waters, C. L., Anderson, B. J., Green, D. L., Korth, H., Barnes, R. J., and Vanhamäki, H.: Science Data Products for AMPERE, in: *Ionospheric Multi-Spacecraft Analysis Tools*, pp. 5–33, ISSI Scientific Report Series 17, https://doi.org/https://doi.org/10.1007/978-3-030-26732-2_7, 2020.
- 145