

1 Modelled dynamics of floating and grounded icebergs, with 2 application to the Amundsen Sea

3 Yavor Kostov¹, Paul R. Holland¹, Kelly A. Hogan¹, James A. Smith¹, Nicolas C. Jourdain², Pierre
4 Mathiot², Anna Olivé Abelló², Andrew H. Fleming¹, Andrew J. S. Meijers¹.

5 ¹British Antarctic Survey, Cambridge, UK

6 ²Univ. Grenoble Alpes/CNRS/IRD/G-INP/INRAE, Institut des Géosciences de l'Environnement, Grenoble, France

7 *Correspondence to:* Yavor Kostov (yastov@bas.ac.uk)

8 **Abstract.** Icebergs that ground on the submarine Bear Ridge in the Amundsen Sea are known to block the drift of sea ice,
9 playing a crucial role in maintaining shelf sea ocean conditions. This important iceberg—sea ice—ocean interaction is
10 commonly observed around the Antarctic shelf seas. To better represent the drift, grounding, and ungrounding of icebergs in
11 the vicinity of such seabed ridges, we introduce new dynamics into the iceberg component of the Nucleus for European
12 Modelling of the Ocean (NEMO) ocean general circulation model. We implement a physically-motivated grounding scheme
13 with parameter choices guided by observations from the Amundsen Sea. When the bergs are grounded, they now experience
14 bottom sediment resistance, bedrock friction, and an iceberg acceleration due to gravity acting down topographic slopes. We
15 also improve the representation of ocean turbulent drag and ocean pressure gradients, both for freely-floating and grounded
16 icebergs, by incorporating the depth-dependence of these forces. We examine the diverse set of forces acting on simulated
17 icebergs in the Amundsen Sea, and compare our simulations with iceberg observations near Bear Ridge. The new iceberg
18 physics pave the way for future studies to explore the existence of possible feedback mechanisms between iceberg grounding,
19 changing sea ice and ocean conditions, and iceberg calving from the ice shelves.

20 1 Introduction

21 Icebergs play a major role in the redistribution of freshwater in the polar oceans, contributing roughly half of the freshwater
22 discharge from Antarctica and Greenland (Cenedese and Straneo, 2022; Davison et al., 2023). Compared to basal melt of ice
23 shelves, which causes localized freshening, icebergs tend to release meltwater farther offshore (Merino et al., 2016; Fox-
24 Kemper et al, 2019). At the same time, there is diversity in the freshwater contribution among icebergs of different dimensions;
25 large tabular bergs may become grounded in shallow shelf seas shortly after they calve and hence release melt water closer to
26 the shore (Cenedese and Straneo, 2022; Olivé Abelló et al., 2025). Nevertheless, the largest icebergs have the longest life-
27 span, allowing for a sustained contribution of meltwater (Cenedese and Straneo, 2022). Stern et al. (2016) and Fox-Kemper
28 et al. (2019) point out that modelling the behaviour of the largest icebergs is particularly challenging, especially when they are
29 represented as point particles. This modelling approach does capture the integrated effect of ocean and atmospheric properties
30 over the entire breadth and depth of large icebergs (Stern et al., 2017). Martin and Adcroft (2010), as well as Merino et al.
31 (2016) partially mitigate this issue in NEMO by imposing that icebergs respond to spatially averaged ambient properties, and
32 that iceberg keels are aware of bottom topography.

33 Icebergs of all sizes are important not only for the thermodynamics of the polar oceans, but also for marine biology
34 and ecology. For example, iceberg debris is a significant source of silica, whose supply to the ocean enhances the biological
35 carbon pump by promoting the growth of diatom phytoplankton populations (Hawkings et al., 2017). Cefareli et al. (2016)
36 identify diatom species attached to the submerged parts of “iceberg walls,” providing further evidence for the biological
37 relevance of icebergs. Tarling et al. (2024) argue that icebergs impact the distribution of phytoplankton and particulate matter
38 via changes in the salinity-driven stratification, while Lucas et al. (2024) suggest that iceberg melting allows for deep nutrient-
39 rich water to penetrate layers shallower than the stratification maximum. The latter promotes primary production and hence,

40 the carbon pump (Smith et al., 2007; Cenedese and Streaneo, 2022; Lucas et al., 2024). Last but not least, icebergs also
41 represent one of the largest sources of iron in the polar ocean (Duprat et al., 2016), while at the same time they are also very
42 heterogeneous and variable in their iron content (Hopwood et al., 2019).


43 As they drift, icebergs may run into topographic obstacles or plough into seafloor sediment, a phenomenon that has
44 been the subject of scientific inquiry as far back as the 19th century, when Charles Darwin published a short communication
45 on iceberg interaction with the ocean bottom (Darwin 1855). More recently, a number of studies on iceberg scouring have had
46 a focus on protecting underwater infrastructure (e.g., Chari 1979; Chari et al., 1980; and Lopez et al., 1981). Other literature
47 explores past iceberg scours in order to infer paleoclimatic changes in the North Atlantic (Hill and Condron, 2014) and the
48 Southern Ocean (Starr et al., 2021).

49 Topographic blocking by grounded icebergs can also significantly modify regional ocean conditions (Nakayama et
50 al., 2014; Stern et al., 2015; St-Laurent et al., 2015; St-Laurent et al., 2024), in shelf seas such as the Amundsen Sea
51 Embayment, where there are hundreds of icebergs present at any time and almost 90% of them are smaller than 2 km² (Mazur
52 et al., 2019). Bett et al. (2020) suggest the importance of iceberg grounding along Bear Ridge in the Amundsen Sea, Antarctica
53 (Figure 1; Mazur et al., 2019) to the oceanography of the region. In particular, Bett et al. (2020) point out that icebergs grounded
54 on Bear Ridge form a physical barrier to westward sea ice transport, and in their study, this is the dominant mechanism creating
55 a dipole in the sea-ice concentration (Figure 1). Icebergs blocking the supply of drifting sea ice creates the Amundsen Sea
56 Polynya to the west of Bear Ridge, which has a strong cooling effect on the water column and is important to the biological
57 productivity of the region (e.g., Arrigo et al., 2012; Person et al., 2019; St-Laurent et al., 2019). To the east of Bear Ridge,
58 icebergs blocking the export of sea ice suppresses polynya activity, reducing the local formation of Winter Water and thereby
59 promoting the intrusion of warmer modified Circumpolar Deep Water (mCDW; Bett et al., 2020). The drift and grounding of
60 icebergs in the Amundsen Sea is shown in the Supplementary Movie in Appendix A (Kostov et al., 2025a, an update of the
61 supplementary movie in Bett et al., 2020), which contains Copernicus Sentinel-1 SAR (synthetic aperture radar) images taken
62 over the time period 2017-2024. The impact on sea ice in the region is also clearly visible.

63 Crucially, these dynamics offer the possibility of a local positive feedback loop. The warm mCDW on the shelf in the
64 Amundsen Sea supports rapid melting of the surrounding ice shelves (Jacobs et al., 2012; Dutrieux et al., 2014). Changes in the
65 supply of warm CDW has led to rapid thinning and acceleration of these ice shelves and their tributary ice streams (Mouginot
66 et al., 2016; Shepherd et al., 2019; Naughten et al., 2022). This has caused an overall increase in iceberg calving, punctuated
67 by several large ice shelf collapse events that released large numbers of icebergs (Miles et al., 2020; Joughin et al., 2021,
68 Davison et al., 2023). Changes in the supply of icebergs to Bear Ridge (and the thickness of those bergs) could affect the
69 characteristics of the grounded “iceberg wall,” influencing sea ice blocking, hence CDW temperatures, and ultimately ice shelf
70 melting and calving (Bett et al., 2020). The sign and strength of this feedback mechanism and its future response to projected
71 climate change are unknown yet have important implications for the sea-level contribution from the West Antarctic Ice Sheet.
72 Therefore, the representation of iceberg grounding in models is essential not only for simulating the trends and variability of
73 the Amundsen Sea polynya but also for projecting major components of future sea-level rise.

74 These important roles of icebergs in the climate system necessitate their proper representation in ocean models (Fox-
75 Kemper et al., 2019). Some of the earliest efforts in modelling iceberg trajectories are attributed to Mountain (1980) and Smith
76 and Banke (1983), while many of the currently used iceberg models can be traced back to Bigg et al. (1997) and Gladstone et
77 al. (2001). Martin and Adcroft (2010) introduce icebergs in a fully coupled climate model, and Marsh et al. (2015) pioneer the
78 first version of the iceberg module in the state-of-the-art Nucleus for European Modelling of the Ocean (NEMO-v4.2.0) ocean
79 model (Madec, G. and NEMO-Team, 2016; Madec, G. and NEMO Systems Team, 2022). Bigg et al. (2018) use NEMO to
80 model iceberg hazard, while Merino et al. (2016) further develop the representation of icebergs in NEMO and, more
81 specifically, simulate the impact of depth-varying ocean properties on iceberg dynamics and thermodynamics.

82 In this study, we introduce a new physically and observationally motivated iceberg grounding capability in NEMO.
 83 This is particularly designed and tuned for the Amundsen Sea, given the importance of icebergs grounding on Bear Ridge, but
 84 the underlying physics should be generally applicable to other areas. Sequences of satellite images from the Amundsen Sea
 85 region reveal complex patterns of iceberg motion near Bear Ridge with multiannual episodes of iceberg grounding (see the
 86 Supplementary Movie). Seafloor records of iceberg scours collected from multibeam bathymetry sounding in the vicinity of
 87 Bear Ridge also reflect the dynamics that arise from the combination of multiple forces acting on a grounded berg. A
 88 compilation of sediment density and shear strength data from representative sediment cores in the Amundsen Sea point to the
 89 important role that the so-called ‘silt resistance’ (Chari, 1979; hereafter referred to as ‘sediment resistance’) may play for the
 90 deceleration and arrest of grounded icebergs. These various observations, along with theoretical analysis of the underlying
 91 physical principles behind iceberg motion, inform our improvements to the NEMO iceberg module. This paper first develops
 92 the physical principles controlling the grounding of icebergs in NEMO, before applying these new physics in a regional model
 93 of the Amundsen Sea.



94
 95 **Figure 1:** Left: Amundsen Sea and Bear Ridge topography [m] based on the BedMachine Antarctica dataset (Morlighem et al., 2020)
 96 and re-gridded on the model configuration grid. Gray contours are spaced 80 m apart. Thin solid black contours delimit the bounds
 97 of boxes a and b in Figure 2, indicating the sites of observed iceberg scours in Section 2.1 with magenta labels matching boxes a and
 98 b in Figure 2. Dark red crosses mark the locations of sediment samples analysed in Section 2.2. The thick purple line shows the
 99 ice shelf edge, with major ice shelves labelled: Dotson, Crosson, Thwaites, and Pine Island Glacier (PIG). Thick solid (thin dashed)
 100 black lines show the boundaries of the boxes where icebergs are released in the SHORT (LONG) simulations. Right: Copernicus
 101 Sentinel-1 SAR (synthetic aperture radar) image from 12 February 2022 over Bear Ridge with white shades indicating the presence
 102 of icebergs and sea-ice.

103 In Section 2 we present and discuss observations of iceberg scouring and sediment properties in the Amundsen Sea.
 104 In Section 3, we describe the theoretical and modelling advances for iceberg dynamics and grounding and their implementation
 105 in NEMO. In Section 4 we test our modelling developments in an Amundsen Sea regional configuration of NEMO and we
 106 describe the force balance of freely floating vs grounded icebergs. In Section 5 we summarise the results of our observationally-
 107 motivated numerical simulations with the updated NEMO model and then conclude.

109 **2.1 Observed iceberg scours**

110 Iceberg scours (also called ploughmarks) are widespread in two Amundsen Sea regions. First, they are extensive across the
 111 middle—outer continental shelf where water depths shoal sufficiently to have allowed icebergs to touch the seabed (~ 700
 112 m), and second, they occur on the shallow banks between the landward-deepening glacial troughs on the continental shelf
 113 (Figure 1). Scour populations are dominated by single-keeled v- or u-shaped forms with relatively small widths and incision
 114 depths, although a few multi-keeled forms are present. A study of $>10,800$ scours in Pine Island Trough (East) by Wise et al.
 115 (2017) returned average single-keel scour widths and depths of 115 m and 2.8 m, respectively (ranges were 12–449 m and 0.1–
 116 22 m). Further west, on the banks around the Dotson-Getz glacial trough, typical scour widths and depths are 100–250 m and
 117 4–10 m, respectively (Graham et al., 2009).

118 Most of the scours in deeper water (> 700 m) on the middle-outer shelf are likely to be relict forms incised during the
 119 last deglaciation, given the keel depths of modern iceberg populations and bathymetric impediments on the inner shelf
 120 (Dowdeswell & Bamber, 2007; Wise et al., 2017). In contrast, satellite imagery from the last decade shows clear accumulations
 121 of modern icebergs on the shallow banks to the west of calving glaciers in the eastern Amundsen Sea (see Supplementary
 122 Movie in Appendix A). After they are calved, the icebergs are moved westwards by prevailing winds and ocean currents and
 123 run aground wherever water depths shoal significantly. Bear Ridge, a N/NW-trending remnant of crystalline basement (Gohl
 124 et al., 2013) with water depths of 150–450 m, exhibits one of the most persistent collections of grounded icebergs today on its
 125 eastern margin, with icebergs of all sizes running aground there (Mazur et al., 2019). Here, we manually map more than 60
 126 scours from existing multibeam-bathymetric data on the eastern flank and central part of Bear Ridge as a representative
 127 population of modern scours. We describe their morphology qualitatively with the aim of characterising modern iceberg
 128 grounding events, and then we provide metrics on their dimensions to calibrate the modelled scouring in Section 4.

129 Multibeam-bathymetric data from 17 research cruises with tracks across Bear Ridge were gridded with 30-m square
 130 grid cells and visualised in ArcGIS Pro 3.3.1 and QPS Fledermaus 8.6.1 for analysis. More than 60 iceberg scours were
 131 mapped, visualised with hillshading, and their cross-sectional widths and depths measured. Overall metrics are presented in
 132 Table 1.

133 Iceberg scours on Bear Ridge exhibit a variety of forms (Figure 2). Many incisions consist of single v- to u-shaped
 134 curvilinear depressions with shallow berms on either side; incision depths range from 2.5–15 m, widths are 90–360 m, and
 135 berm heights are typically 1–3 m (Figure 2c). The orientation of these scours is typically E–W or SE–NW (average orientation
 136 is 0.94°) and is thus consistent with iceberg transport from the Pine Island and Thwaites glacier fronts with the coastal current.
 137 However, the cross sections of some scours can change direction and vary along their length from a single v-shaped to wider
 138 u- or w-shaped, the latter indicating that the geometry of the iceberg base has varied as the iceberg was dragged through the
 139 sediment (consistent with either keel break-off or iceberg rotation). Tabular iceberg keels also have v-shaped protrusions along
 140 the rough surface of their bases. Occasionally, a large tabular iceberg can even plough multiple parallel scours. Some scours
 141 also terminate abruptly in rounded depressions, encircled by a shallow berm. Such iceberg grounding pits or iceberg plough
 142 ridges indicate where seafloor scouring terminated (e.g., Jakobsson et al., 2011). Most often there are no other incisions on the
 143 seafloor that cross-cut the plough ridges, suggesting that the icebergs either lifted-off from the seafloor once enough melting
 144 had occurred or perhaps capsized (rotated vertically due to an unstable height:width ratio; e.g., Bass, 1980 and Ruffman, 2005)
 145 allowing them to float away freely without incising the seafloor further.

146 Detailed inspection of the scours reveals that some have either sinuous edges over length scales of a few hundred
 147 metres, variable depths along the centre of the scour, again at length scales of hundreds of metres, or exhibit both morphologies
 148 together. We interpret this variability as being formed by icebergs that have ploughed the seafloor and “wobbled”, either from
 149 side to side or are close to flotation and have bobbed up and down on the seafloor under the influence of tides, currents and/or

winds, or a combination of these factors (cf. Barnes and Lien, 1988; Lien et al., 1989). A particularly intriguing seafloor morphology of crossed u-shaped scours that appear to be linked together are observed on the southern part of Bear Ridge in water depths of ~ 275 m (Figure 2a). We suggest that these scours were formed by flat-bottomed icebergs that incised the seafloor as they were transported on to a shallower part of the ridge, where they halted. Then, under the influence of local currents, winds or tides, the bergs were dragged through the sediment in a perpendicular direction to their original pathway. Alternatively, two different icebergs may have ploughed scours perpendicular to each other, thus generating a crossed shape. Like the iceberg plough ridges, these scours have sharply defined and uninterrupted edges and berms suggesting that when the icebergs did eventually move off the seafloor it was due to melting or capsizing and no further marks on the seafloor were made.

Table 1: Bear Ridge iceberg scour metrics

Av. depth (m)	Depth range (m)	Av. width (m)	Width range (m)	Av. orientation
7	2.5-14	189	90-357	0.94°




Figure 2: Examples of iceberg scour types from Bear Ridge (boxes a and b located as shown on the map in Figure 1). The red arrows show likely iceberg transport paths westwards from the eastern Amundsen Sea and across Bear Ridge; blue arrows show iceberg plough ridges (berms); black arrows show the perpendicular direction of iceberg scouring in the crossed forms. Panel c) presents a schematic of a typical scour cross-section found on Bear Ridge.

2.2 Sediment density and shear strength


Saturated density and shear strength determine the sediment resistance forces acting on grounded icebergs. Data from representative sediment cores recovered from the Amundsen Sea (Figure 3; Smith et al., 2011; Clark et al., 2024) provide a range of values for both the saturated density and shear strength. Whole core saturated density was determined using a GEOTEK multi-sensor core logger following standard methods (Niessen et al., 2007) while shear strength was measured with a shear vane (Smith et al., 2011; Clark et al., 2024).

173 Sediment thicknesses across different polar regions (and hence the maximum scour depth) differ markedly between

174 areas of glacially eroded bedrock that are completely stripped of sediment and thick sequences of more deformable
175 glaciomarine muds and subglacial tills. The composition of post-glacial seafloor (<15 m) sediments in the Amundsen Sea is
176 variable, but typically consists of three broad lithological units. The uppermost unit, directly at the seafloor, consists of soft,
177 water-rich muds with variable biogenic content. These sediments were deposited in an open or seasonally sea-ice covered
178 glaciomarine environment distal (>25 km) from the ice sheet grounding line (Smith and Hogan, 2024) and are characterised
179 by low shear strength and density values (<5 kPa, <1.6 × 10³ kg/m³). Beneath these sediments, coarser-grained stratified to
180 structureless, sandy to gravelly, terrigenous sediments occur. These ‘transitional’ sediments were deposited at or close to (<25
181 km) to the ice sheet grounding line and lie directly above subglacial sediments below. The subglacial sediments in the basal
182 unit consist of stiffer/denser, and largely structureless diamictons deposited subglacially during the Last Glacial Maximum,
183 either as ‘soft’ deformation till (>5-20 kPa, 1.8-2.2 × 10³ kg/m³) or ‘stiff’ compacted tills (>20 kPa, >2.0 × 10³ kg/m³).
184 Thicknesses of each unit vary, but postglacial glaciomarine mud tends to be thicker on the inner shelf, particularly where
185 sedimentation is focussed in bathymetric troughs or depressions. Conversely, on shallow banks such as former pinning points
186 (see Hogan et al., 2021), post-glacial sediments tend to be much thinner (<5 m), so that stiffer sediments and (crystalline)
187 bedrock occur much closer to the seafloor. Figure 3 presents samples from all of the three units characterized by different
188 saturated density and shear strength.

189 Compared to other sectors of the Amundsen Sea, comparatively few sediment cores have been recovered from Bear

190 Ridge due to a combination of perennial sea-ice cover, and a general scientific focus on the glacial troughs. The seafloor drill
191 rig MARUM-MeBo70 was used at three locations on the western flank of Bear Ridge (Gohl et al., 2017), revealing that
192 indurated mudstones, likely of Eocene-Miocene age, crop out very close to the seafloor. Crucially, in areas such as Bear Ridge
193 where post-glacial sediments are thin and crystalline basement/sedimentary strata crop out close to the seafloor iceberg
194 ploughing depth will be retarded via solid body and static friction.



195

196 **Figure 3: Observed sediment properties for selected cores from the Amundsen Sea: the uppermost sediments consist of soft, water-**
197 **rich muds (solid black box); coarser-grained stratified to structureless diamictons and ‘soft’ deformation tills (solid blue box); stiffer**
198 **tills and/or denser subglacial diamictons (dashed black box). The soft/stiff till boundary follows Reinardy et al. (2009). Values outside**
199 **the boxes reflect other sediment types such as iceberg diamictons, glaciogenic debris flows etc.**

200 **3 Modelling iceberg dynamics and grounding**

201 We improve the capability to simulate iceberg drift, grounding, and ungrounding by building up a theoretical framework that
202 defines the relevant forces at play. We consider drivers of iceberg motion – such as surface winds and ocean pressure gradients
203 – as well as forces that dissipate mechanical energy via the effects of bottom sediment resistance and solid bedrock friction.

204 **3.1 Forces acting on freely floating icebergs and their representation in NEMO**

205 In order to fully understand the behaviour of grounded icebergs, we first explore their dynamics while they are in free flotation.
 206 Floating icebergs are driven by the combined effect of relative winds (on the iceberg's upper surface and sides of the iceberg
 207 freeboard), sea-ice drag, ocean waves, and ocean pressure gradients and relative ocean currents integrated over the depth of
 208 the submerged keel and along the iceberg's basal surface (Merino et al., 2016). Icebergs are also subject to the Coriolis force
 209 (Marsh et al., 2015; Cenedese and Straneo, 2022). The effect of tides (if represented) is indirect: tides alter the sea surface
 210 height gradients, and hence, the water column thickness, ocean pressure gradients, and ocean drag. The acceleration of a freely-
 211 floating iceberg as it moves (Bigg et al, 1996; Condron and Hill, 2021; Marsh et al., 2018) can be expressed as:

$$212 \frac{d\vec{u}}{dt} = \vec{a}_{\text{pressure gradient}} + \vec{a}_{\text{Coriolis}} + \vec{a}_{\text{ocean drag}} + \vec{a}_{\text{atmos. drag}} + \vec{a}_{\text{sea-ice drag}} + \vec{a}_{\text{waves}} \quad (1)$$

213 where terms on the right-hand side denote accelerations due to the ocean pressure gradient, the Coriolis force, ocean drag,
 214 atmospheric drag on icebergs, sea-ice drag, and wave drag (see the supplementary Table D1 for a list of all symbols used).
 215 Importantly, grounded icebergs are also decelerated by an additional term, \vec{a}_{ground} , which represents the sum of all dissipative
 216 forces acting on an iceberg. We assume that there is no continuous field of interacting icebergs, and so $\nabla \vec{u} = 0$ because icebergs
 217 are treated as isolated rigid, solid bodies (in NEMO they are in fact represented as moving point particles when it comes to
 218 advection).

219 The direct acceleration that the atmosphere and ocean flows impose on icebergs can be understood in terms of the
 220 turbulent drag that the fluid exerts on the berg. This turbulent drag is assumed to be a quadratic function of the speed of the
 221 iceberg relative to the fluid:

$$222 \vec{a}_{\text{drag}} = -C_{\text{drag}} |\vec{u} - \vec{u}_{\text{fluid}}| (\vec{u} - \vec{u}_{\text{fluid}}) \quad (2)$$

223 where C_{drag} represents an effective form drag (Wagner et al., 2022) coefficient that is different for air, seawater and sea-ice,
 224 and has units of an inverse length scale related to the horizontal extent of the iceberg (Martin and Adcroft, 2010). The effective
 225 drag coefficient takes into account the efficiency of momentum transfer between the fluid and the iceberg, the density ratio
 226 between the fluid and the iceberg, and the iceberg area exposed to the drag. The typical order of magnitude of this parameter
 227 when applied to ocean—iceberg drag is $10^{-4} [\text{m}^{-1}]$ for thick tabular icebergs in the Amundsen Sea. Analogously to a turbulent
 228 drag, the sea-ice pack also exerts a drag on icebergs, especially in regions with a high sea-ice concentration (Marson et al.,
 229 2024). While the quadratic formulation of equation (2) is well established for high Reynolds numbers in the ocean and
 230 atmosphere, its application to sea-ice is used for simplicity, but little is known about its validity.

231 When considering ocean drag, it is important to account for the vertical shear in the upper ocean, which can
 232 significantly impact iceberg motion (FitzMaurice et al., 2016). The existing NEMO algorithms use the depth-averaged ocean
 233 velocity to estimate the ocean drag term:

$$234 \text{Old } \vec{a}_{\text{ocean drag}} = \frac{C_{\text{drag}}}{D_{\text{keel}}} \left| \vec{u} - \int_{D_{\text{keel}}}^0 \vec{u}_{\text{ocean}}(z) dz \right| \left(\vec{u} - \int_{D_{\text{keel}}}^0 \vec{u}_{\text{ocean}}(z) dz \right), \quad (3)$$

235 but the nonlinearity of the drag formulation means that equation (3) is inaccurate. In the update that we propose here, we
 236 consider the ocean currents at each depth level and average the resulting drag over the iceberg keel depth to arrive at an
 237 improved representation of ocean drag:

$$238 \vec{a}_{\text{ocean drag}} = \frac{C_{\text{drag}}}{D_{\text{keel}}} \int_{D_{\text{keel}}}^0 |\vec{u} - \vec{u}_{\text{ocean}}(z)| (\vec{u} - \vec{u}_{\text{ocean}}(z)) dz, \quad (4)$$

239 where D_{keel} is the keel depth.

240 Another significant source of acceleration for icebergs is the horizontal gradient in ocean pressure at different depth
 241 levels. Gradients in sea-surface height (SSH) form a large component of the horizontal pressure gradients, but density
 242 variations also modify pressure gradients within the water column. This latter effect was not represented in the existing iceberg
 243 dynamics in NEMO, and is introduced here. For example, in the x-direction, at each depth level z , we have a pressure gradient
 244 component:

245
$$\frac{\partial(\frac{P(z)}{\rho_0})}{\partial x} = gh(z) \frac{\partial(\frac{\langle \rho'(z) \rangle}{\rho_0})}{\partial x} + g \left(\frac{\langle \rho'(z) \rangle}{\rho_0} \right) \frac{\partial h(z)}{\partial x} + g \frac{\partial h(z)}{\partial x} , \quad (5)$$

246 where $\langle \rho'(z) \rangle = \frac{1}{z} \int_0^z (\rho(z) - \rho_0) dz'$ is the density anomaly, relative to the reference Boussinesq density ρ_0 , averaged over
 247 the overlying water column, g is the magnitude of the acceleration due to gravity, and $h(z)$ is the water-column thickness
 248 above level z . Under the Boussinesq approximation, the contribution of the second term on the RHS may be neglected. In
 249 addition to large-scale dynamical features, tides also contribute to SSH gradients, while SSH anomalies are considered a major
 250 driver of iceberg motion, especially for weakly-grounded icebergs (Brown et al., 2017).

251 In addition to drag and pressure forces, icebergs furthermore respond to the Coriolis force due to the planetary rotation,
 252 which gives rise to an acceleration at a right angle relative to the velocity vector:

253
$$\vec{a}_{Coriolis} = -f \hat{k} \times \vec{u} , \quad (6)$$

254 where f is the Coriolis parameter.

255 3.2 Sediment resistance

256 Grounded icebergs are subject to additional sources of acceleration and deceleration such as sediment resistance and solid-
 257 body friction, which have so far not been represented in the NEMO iceberg module and are introduced here for the first time.
 258 Here we make a distinction between several different cases: icebergs that plough only into the sediment and remain in motion
 259 or halt; icebergs whose base reaches the solid basement i.e., crystalline bedrock but remain in motion; and icebergs whose
 260 motion comes to a stop when grounded in the sediment and/or on the basement. Here we first describe the resistance
 261 experienced by a moving berg whose keel ploughs into the sediment. Guided by the most typical observed scour shapes in the
 262 Amundsen Sea, we assume that each iceberg ploughs a single v-shaped trench approximately 8-m deep.

263 There is a specialized body of literature discussing the interaction of icebergs with seafloor sediments. Empirical
 264 observations and theoretical considerations suggest that sediment resistance can be decomposed into three different
 265 mechanisms. An iceberg ploughing into sediments creates a wedge of sediment, composed of fracture plates (Chari et al.,
 266 1975; Chari et al., 1980). The weight of this wedge pushes back on the part of the iceberg's frontal face that has ploughed into
 267 the sediment with a force (in units of N):

268
$$\frac{\gamma'(H+D)^2 W}{2} \approx \frac{\gamma'D^2 W}{2} \quad (7)$$

269 where γ' is the submerged unit weight of the bottom sediment (in units of $N m^{-3}$), W is the width of the scour, D is the depth
 270 of the scour, and $H \ll D$ is the height of the excavated sediment above the level of the surrounding seafloor (Chari et al., 1975;
 271 Chari et al., 1980). The submerged unit weight γ' [$N m^{-3}$] in turn can be expressed as a function of the saturated sediment
 272 density ρ_{sat} [kg/m^3] and the density of water ρ_{water} [kg/m^3]:

273
$$\gamma' = g(\rho_{sat} - \rho_{water}) . \quad (8)$$

274 Furthermore, there are shear stresses acting along the surfaces of the fracture plates within the wedge of sediment
 275 excavated by the iceberg:

276
$$Shear \text{ within wedge} = 2\tau DW \quad (9)$$

277 where τ is the shear strength of the sediments in units of Pa (Chari, 1975; Chari et al., 1980). Lastly, as the excavated wedge
 278 is pushed forward along the path of the iceberg scour, the sides of the wedge experience lateral friction against the undisturbed
 279 surrounding sediments:

280
$$Lateral \text{ friction} = \frac{\sqrt{2}}{2} \tau D^2 . \quad (10)$$

281 This gives rise to a three-term expression for the maximum possible sediment resistance, which acts in the direction
 282 opposite to the iceberg drift if it is in motion, or opposite to the net driving force if the iceberg is static:


283
$$\vec{a}_{sediment \ max} = - \left[\frac{\gamma'D^2 W}{2} + 2\tau DW + \frac{\sqrt{2}}{2} \tau D^2 \right] \cdot \begin{cases} \hat{u}/M_{iceberg} , & |\vec{u}_0| > 0 \\ \hat{a}_{net \ drivers}/M_{iceberg} , & |\vec{u}_0| = 0 \end{cases} \quad (11)$$

284 where $M_{iceberg}$ is the iceberg mass, $\hat{a}_{net\ drivers}$ indicates a unit vector in the direction of the net driving force (the sum of all
 285 nondissipative forces that drive iceberg motion), \hat{u} indicates a unit vector along the *projected* direction of future motion, which
 286 depends both on the present velocity and on the net driving force (see below), and \vec{u}_0 is the present velocity of the iceberg. In
 287 the model implementation (Section 3.4), the sediment resistance is not allowed to drive a net acceleration but only to decelerate
 288 iceberg motion or to oppose the forces trying to set an iceberg in motion, hence why the above is formulated as a maximum
 289 force.

290 The latter can be 10s to 100s of meters thick, particularly in bathymetric depressions such as glacial troughs.
 291 Following our Amundsen Sea example, the observed scours (Section 2.1) and compiled sediment properties (Section 2.2)
 292 inform our decision to assume the existence of a uniform 8-m sediment layer with a saturated density of $1.8 \times 10^3 \text{ kg/m}^3$ and
 293 shear strength of 6 kPa. Beneath this single uniform sediment layer we assume the existence of a basement. Thus, iceberg
 294 ploughing is allowed to occur up to a maximum of 8-m depth while it remains in motion, but then other forces come into play,
 295 described next.

296 3.3 Gravity, solid-body friction, and static friction

297 When a moving grounded iceberg ploughs through all the sediment and reaches a sloping basement, it can move up the
 298 basement slope if the berg has enough momentum and/or sufficient sources of sustained acceleration (Chari 1975). As the berg
 299 moves up the slope, it is lifted upwards, its freeboard increases, and as a result, the buoyancy force of the ocean no longer
 300 balances the weight of the iceberg and a force directed into the basement supports the remainder (Figure 4). This gives rise to
 301 a gravitational force, which converts the kinetic energy of the iceberg motion into potential energy upon grounding, and also
 302 causes a solid body friction between the berg and the basement.



303
 304 **Figure 4: Vertical force budget.** An idealized schematic of an iceberg trajectory (green arrow) going up a topographic slope. The
 305 angle of the trajectory (solid green arrow) relative to its projection on the horizontal x-y plane (dashed grey arrow) is denoted α .
 306 The local maximum slope of the topography is denoted β . The vector normal to the topography is indicated with a brown line.
 307 Schematic angles not shown to scale with respect to the relevant topographic slopes.

308 Lifting the iceberg out of floatation means that a portion of the iceberg's weight is no longer supported by the
 309 buoyancy of ice relative to seawater. The vertical displacement yields a deficit of $\rho_{water} H_{freeboard\ anom.}$ in the mass of
 310 seawater displaced by the berg per unit area, and a corresponding net downwards gravitational force. Dividing this force by
 311 the berg mass per unit area, we arrive at an acceleration whose projection down the sloping topography is

312
$$\text{Projection of gravitational acceleration} = \frac{\rho_{water}}{\rho_{ice}} \frac{H_{freeboard\ anom.}}{H_{Total}} \bar{g} \sin \beta. \quad (12)$$

313 where the above takes into account the density difference between the iceberg $\rho_{ice} = 875 \text{ kg/m}^3$ – including firn air trapped
 314 in the ice (Veldhuijsen et al., 2023) – and the ambient sea water $\rho_{water} = 1025 \text{ kg/m}^3$, and \bar{g} is the gravitational acceleration
 315 vector, and the topography has a local maximum slope β (Figure 4). It is the horizontal components of the force balance that

316 are of interest here. We can furthermore express the projection of the gravitational force directed straight down the slope as a
 317 horizontal acceleration term:

$$318 \quad \vec{a}_{\text{gravity}} = \frac{\rho_{\text{water}} H_{\text{freeboard anom.}}}{\rho_{\text{ice}} H_{\text{Total}}} g (\sin \beta \cos \beta) \hat{s} \quad (13)$$

319 where \hat{s} is a unit vector in the x-y horizontal plane aligned with the direction of the maximum topographic slope at the location
 320 of the iceberg, and g is the magnitude of the gravitational acceleration.

321 There is another gap in the literature when it comes to the solid-body friction between the base of the iceberg and the
 322 basement beneath any sediments. One may, however, draw a qualitative analogy to the friction between glaciers and rock,
 323 whose representation in models can also be reduced to a solid-body (Coulomb type) sliding law (Fowler et al., 2010), which
 324 is independent of the speed in certain regimes. This term in the force balance is briefly discussed in Chari's dissertation (Chari,
 325 1975). Generally, Coulomb friction is assumed to be proportional to the normal confining force at the interface between the
 326 two solid objects, in our case the seafloor basement and the base of the iceberg. When the iceberg is in contact with basement,
 327 this normal force is a projection of the net vertical imbalance between gravity and buoyancy. If an iceberg moves up a slope
 328 at an angle α relative to the horizontal plane, the magnitude of the Coulomb friction in the sloping plane can be expressed as

$$329 \quad \text{Coulomb friction} = \frac{\rho_{\text{water}} H_{\text{freeboard anom.}}}{\rho_{\text{ice}} H_{\text{Total}}} g \mu \cos \beta \cos \alpha \quad (14)$$

330 where μ is a dimensionless coefficient of solid body friction, whose value for the interaction between icebergs and the seafloor
 331 basement is not known. Equation (14) describes a deceleration term acting against the horizontal direction of motion \hat{u} (Figure
 332 4) in the case of moving icebergs. The kinetic friction has a vector orientation opposing the iceberg velocity:

$$333 \quad \vec{a}_{\text{solid}} = - \frac{\rho_{\text{water}} H_{\text{freeboard anom.}}}{\rho_{\text{ice}} H_{\text{Total}}} g \mu \cos \beta \cos \alpha \hat{u}. \quad (15)$$

334 In the model implementation (Section 3.4), the magnitude of the kinetic friction vector \vec{a}_{solid} is capped, so that it
 335 never drives a net acceleration. The value of the non-dimensional coefficient μ cannot be constrained directly and could only
 336 be inferred from the resulting iceberg grounding behaviour. In this study, we test a range of values including $\mu = 0$, (which
 337 corresponds to no Coulomb friction but only sediment resistance and gravity), $\mu = 0.002$, and $\mu = 0.2$, but we use $\mu = 0.002$
 338 in simulations with a realistic Amundsen Sea model configuration (See Appendix B for a discussion of parameter sensitivity).
 339 Our results are also robust with respect to the choice of value for the Coulomb coefficients μ and μ_{stat} (Appendix B). We
 340 find that varying the magnitude of μ does not affect the episodes of motionless grounding in numerical simulations (Figure B1
 341 in Appendix B). In contrast, the static episodes marked appear to be more influenced by the sediment resistance and the
 342 processes that thin the icebergs.

343 We note that the coefficient of Coulomb friction μ is usually assumed to be different in the case of moving objects
 344 (kinetic friction) versus objects at rest (static friction, μ_{stat}). Usually, the coefficient of static friction μ_{stat} is greater than or
 345 equal to the value of the kinetic friction coefficient for the same type of surfaces under similar conditions. Another noteworthy
 346 difference between the kinetic and static solid body friction is related to the orientation of the frictional force vector. In the
 347 case of static friction, there is no motion, so friction does not act in a direction opposite to the velocity vector, which is zero.
 348 Instead, static Coulomb friction and sediment resistance jointly act in a direction opposing the net sum of all acceleration terms
 349 that try to set an object into motion. Therefore, in the case of grounded icebergs, the maximum possible decelerating force per
 350 unit mass due to static friction is:

$$351 \quad \vec{a}_{\text{static-solid}} = - \frac{\rho_{\text{water}} H_{\text{freeboard anom.}}}{\rho_{\text{ice}} H_{\text{Total}}} g \mu_{\text{stat}} \cos \beta \cos \alpha \widehat{a}_{\text{net}} \quad (16)$$

352 which is similar to the expression for kinetic friction, but here, we orient the vector against the sign of the net acceleration
 353 $\vec{a}_{\text{net drivers}}$, which includes fluid and ice drag, the Coriolis force, and gravitational accelerations but none of the dissipative
 354 accelerations, and is aligned with the unit vector \widehat{a}_{net} . In our study we impose a static coefficient μ_{stat} that is 100 times larger
 355 than the corresponding kinetic coefficient μ , which is an arbitrary choice tested below.

356 Last but not least, an iceberg moving along solid-basement topography with non-constant slopes experiences changes
357 in motion due to the surface geometry. For example, if an iceberg moves up a steeper slope, some of its horizontal momentum
358 is reoriented in the vertical direction, a “curvature term” $\vec{a}_{curvature}$ discussed in Appendix B. In total, then, the new iceberg
359 grounding model is made up of 4 terms, each of which applies under different conditions:

360 $\vec{a}_{ground} = \vec{a}_{sediment} + \vec{a}_{gravity} + \vec{a}_{solia} + \vec{a}_{curvature},$ (17)

361 and when an iceberg is grounded the above terms (if present) are added to the full iceberg momentum budget.

362 **3.3 Representation of iceberg melting and capsizing in NEMO and their role in ungrounding**

363 Melting plays an important role in the ungrounding of icebergs from bottom topography. NEMO breaks down the iceberg
364 melting into three processes: basal melting, lateral melting, and wave erosion (Martin and Adcroft, 2010). Wave erosion is a
365 function of the relative wind velocity and the ambient ocean temperature. Basal melting is represented as a function of
366 temperature and velocity relative to the ocean, while the melting due to buoyant convection along the side walls is assumed to
367 depend only on temperature (Martin and Adcroft, 2010). Wave erosion and buoyant convection decrease the horizontal width
368 and length of a simulated iceberg, while basal melting reduces the vertical thickness, which is relevant to ungrounding.
369 However, the freshwater released by all three simulated processes, including basal melting, is injected at the NEMO ocean
370 surface.

371 For icebergs that float or follow the old grounding schemes, the rate of basal melting is proportional to the relative
372 velocity between the iceberg and the ocean current at the height of the iceberg base. However, we allow grounded icebergs to
373 plough into the sediment. In this case, for the purpose of calculating basal melting, we compute the relative velocity at the
374 deepest ocean level above the sediment.

375 When the iceberg’s geometry becomes unstable and its centre of mass shifts, the iceberg can spontaneously roll over
376 and capsize (Wagner et al., 2017 OM). The existing NEMO code uses a ratio between the iceberg’s longer horizontal length
377 and its keel depth as a criterion for stability against rolling (Weeks and Mellor, 1978). However, (Wagner et al., 2017 OM)
378 point out that this approach used in models is unphysical. The appropriate indicator of vulnerability to capsizing is the ratio
379 between the horizontal width and the full thickness (Wagner et al., 2017 OM).

380 Olivé Abelló et al. (2025), a companion study, updates the rolling criteria in NEMO by appropriately considering the
381 ratio between the iceberg width and thickness. Olivé Abelló et al. (2025) build up on the approach used by MacAyeal et al.
382 (2003) and Bigg et al. (1997 and 2018), while correcting errors highlighted by Wagner et al. (2017, OM).

383 **3.4 Implementation of the new iceberg dynamics algorithms in NEMO**

384 We implement the relevant iceberg dynamics and grounding theory as an update (Kostov et al., 2025b) to the existing
385 representation of icebergs in NEMO. The iceberg model in NEMO operates within the ocean domain. However, iceberg motion
386 is not confined to the resolved ocean grid, and instead bergs are dynamically represented as Lagrangian mass particles that
387 propagate across and within ocean grid cells. For the purposes of calculating melt rates, as well as turbulent drag, each iceberg
388 is associated with horizontal and vertical dimensions: length, width, and thickness. The iceberg motion is calculated using a
389 4th-order Runge-Kutta scheme (Marsh et al., 2015; Merino et al., 2016). Acceleration, velocity, local SSH slope, and horizontal
390 position are calculated at each stage of the Runge-Kutta scheme and the final estimate for a given iceberg timestep is a weighted
391 sum of the result from each of the four Runge-Kutta stages.

392 As the icebergs move within the resolved grid cells, ocean properties such as velocity, temperature, SSH, and
393 bathymetry are horizontally interpolated onto the sub-gridscale position of the bergs. We calculate vertically-averaged ocean
394 drag terms that we update and apply as acceleration terms within each stage of the Runge-Kutta scheme for iceberg dynamics.

395 At each Runge-Kutta stage, we also apply horizontal pressure gradients interpolated onto iceberg positions. The pre-existing
 396 code included only the gradient components due to SSH anomalies. Here we also account for the pressure gradient terms that
 397 arise due to water density anomalies, exactly as they are calculated and applied in the dynamics of the liquid ocean domain.
 398 Our calculation of the pressure gradient terms is therefore compatible with the multiple vertical coordinate systems that are
 399 available in NEMO, e.g, fixed z-level or z^* -coordinates that stretch and contract matching the SSH anomalies. In our
 400 implementation, icebergs do not feel the component of the ocean pressure gradient due to the iceshelf load over iceshelf
 401 cavities, as icebergs themselves cannot enter the cavities. However, icebergs are often found near the iceshelf edge even long
 402 after they have calved. That is why, in algorithm, we mask the ice-shelves, such that icebergs in their vicinity can approach
 403 the shelf edge without being exposed to unphysical pressure gradients.

404 The NEMO model has some existing, crude representations of iceberg grounding (Olivé Abelló et al., 2025). At each
 405 of the four Runge-Kutta stages, these routines consider criteria under which iceberg velocity or components of the iceberg
 406 velocity are set to zero as topographic obstacles are encountered. The grounding routines use the projected forward motion of
 407 the iceberg and therefore have an implicit element. The pre-existing algorithms allow for two options: 1) icebergs come to a
 408 complete stop when their keel reaches a topographic obstacle; or 2) the component of an iceberg's velocity vector normal to a
 409 topographic obstacle is set to zero, while the tangential component is preserved.

410 In our new implementation, we also apply grounding routines at each stage in the Runge-Kutta scheme. However, we
 411 represent the interaction between icebergs and the seafloor in terms of acceleration terms due to gravity, solid-body friction,
 412 and sediment resistance (Section 4). These acceleration vectors are projected along the x and y-axes (zonal and meridional) of
 413 iceberg motion, which are aligned with the ocean model grid (Figure 4). The acceleration/deceleration terms that we introduce
 414 depend on the local slope of the topography and on the angle at which an iceberg moves relative to the horizontal plane (which
 415 may be different from the maximum topographic gradient, Figure 4). In order to calculate these angles for each Lagrangian
 416 iceberg particle, we track and store interpolated bottom topography along an iceberg trajectory and calculate the corresponding
 417 slope angles.

418 In addition, we apply different solid-body friction coefficients depending on whether the grounded iceberg is in
 419 motion or at rest. For numerical purposes, we do not set a zero speed as the threshold for assuming no motion. Instead, when
 420 the speed drops below a small magnitude ($5 \times 10^{-5} \text{ m/s}$), we set it to zero and assume a regime of static friction. This way
 421 we avoid situations where we see numerical stick and slip of icebergs alternating between no motion and brief periods of very
 422 slow motion.

423 While turbulent drag, geometric deflection, and gravity can act in any direction to accelerate or decelerate a berg, the
 424 dissipative sediment resistance and solid-body friction terms only act to decelerate motion. As described above, we do not
 425 allow the two dissipative terms, solid friction and sediment resistance, to drive motion but only to bring an iceberg to a stop
 426 and/or keep it at rest. Therefore, we impose that

$$427 \frac{d\vec{u}}{dt} = -\frac{\vec{u}_0}{\Delta t} \text{ for } |\vec{a}_{\text{solid}} + \vec{a}_{\text{sediment}}| \geq \left| \vec{a}_{\text{net drivers}} + \frac{\vec{u}_0}{\Delta t} \right|, \quad (18)$$

428 where Δt is the model timestep and $\vec{a}_{\text{net drivers}}$ is the net sum of all acceleration terms that do not dissipate mechanical energy.
 429 Even if the condition in (18) is achieved at only one of the four Runge-Kutta stages of the velocity calculation, the zeroing of
 430 an x- or y-velocity component is applied to the output of all four stages. Furthermore, if setting the velocity component to zero
 431 brings the total iceberg speed below $5 \times 10^{-5} \text{ m/s}$, we force the iceberg to stop.

432 The fact that iceberg motion is computed using a separate numerical scheme within the NEMO ocean domain allows
 433 us to apply temporal substepping within the iceberg code while keeping the same time-step size for the ocean and sea-ice. We
 434 thus define a parameter that sets the ratio of the ocean timestep to the shorter iceberg timestep. We find that allowing a shorter
 435 iceberg timestep is numerically important when modelling the shape and evolution of iceberg scours because forces such as
 436 gravity along the sloping topography may act on much shorter timescales compared to other sources of acceleration. (See
 437 Appendix B, where the numerical relevance of sub-stepping is demonstrated in an idealized experiment with gravity but no

438 dissipative forces along a sloping bottom). At the same time, when modelling iceberg residence times and locations on the
439 resolved ocean grid, substepping becomes less critical (Appendix B).

440 In summary, the implementation of our updated algorithms affects the dynamics of both freely floating and grounded
441 icebergs through the following set of changes: 1) we use a vertical average of the full profile of non-linear ocean drag along
442 the keel; 2) we make the ocean pressure gradients on icebergs a function of both SSH and density; 3) we allow icebergs to
443 experience sediment resistance when grounding; 4) we allow icebergs to experience Coulomb friction and gravitational
444 acceleration when the keels reach the solid basement.

445 **3.5 Amundsen Sea NEMO test cases**

446 We test our updated iceberg dynamics in a regional Amundsen Sea configuration of the Nucleus for European
447 Modelling of the Ocean (NEMO-v4.2.0, Madec et al. 2022). This includes the SI3 sea-ice component (Vancoppenolle et al.,
448 2023), a representation of ocean–ice-shelf interactions (Mathiot et al., 2017), and a Lagrangian iceberg module (Marsh et al.,
449 2015; Merino et al., 2016).

450 The extent of the regional domain, the horizontal resolutions (0.25°), and the topography are the same as in Caillet et
451 al. (2023), except that we do not impose a “wall of icebergs” over Bear Ridge to block sea ice. We use 121 vertical levels as
452 in Mathiot and Jourdain (2023), with a thickness increasing from 1 m at the surface to 20–30 m between 100 and 1000 m
453 depth. The cell thicknesses are slightly compressed or stretched to follow the SSH variations (Z^* coordinates), and cells
454 adjacent to the sea bed and ice shelf draft are described as partial steps (Adcroft et al., 1997).

455 The initial state and lateral boundary conditions are from 5-day mean outputs of the global simulation of Mathiot and
456 Jourdain (2023). Tides are neither generated nor prescribed at the boundaries, which is expected to have a limited impact in
457 the Amundsen Sea (Jourdain et al., 2019). Tides are relatively weak in the Amundsen Sea relative to other Antarctic sectors
458 (Jourdain et al., 2019). The surface boundary conditions are calculated through the bulk formula of Large and Yeager (2008)
459 from the JRA55-do atmospheric reanalysis (Tsujino et al., 2018).

460 We release icebergs in January of the first year of a simulation, i.e., 1979. Our aim with these simulations is to
461 demonstrate the new iceberg grounding physics. We are interested in a realistic historical setting, but in this study, we do not
462 aim to explore a broad range of long-term natural variability in the Amundsen Sea and its impact on icebergs. All the icebergs
463 in the Amundsen Sea simulations are released east of Bear Ridge and then allowed to drift, ground, and unground according
464 to the modelled ocean, atmosphere, sea ice and iceberg conditions. We conduct two sets of tests, each with a different iceberg
465 population. In the first set, labelled THICK hereafter, all test icebergs are tabular, with the same mass, 3.9×10^{11} kg, uniform
466 density of 875 kg/m^3 , and thickness of 395 m. In the second set of tests, labelled MEDIUM, all icebergs are 250 m thick, each
467 with mass of 3.8×10^{10} kg. However, MEDIUM icebergs are not thick enough to interact with the topography along Bear
468 Ridge, and hence, they do not feature in our grounding analysis. We analyse the force balance of MEDIUM icebergs in free
469 flotation.


470 In this study we explore the results of two types of simulations, where each type of simulation is performed with each
471 type of iceberg thickness. In the short simulations, labelled ‘SHORT’ hereafter, we initially seed a total of 497 identical test
472 icebergs one in each grid cell ($\sim 1/4^\circ$ by $\sim 1/4^\circ$) in the region east of Bear Ridge enclosed by 109.9° W, 106.2° W, 74.3° S, and
473 72.0° S (Figure 1). While it is not realistic to expect hundreds of thick tabular icebergs to simultaneously occupy this region,
474 our icebergs do not interact with each other or block sea-ice (while sea-ice does impact icebergs). In our SHORT simulations,
475 even when icebergs melt, we do not allow any iceberg meltwater to be injected in the ocean, so as not to perturb the background
476 conditions with each set of 497 icebergs. We run this simulation with a large ensemble of icebergs but only over the course of
477 one full year spanning all seasons. Each of the grid boxes that we seed is a possible location where an iceberg may be found,
478 and hence, we explore the behaviour of thick tabular icebergs when exposed to the local conditions. This large population

479 allows us to analyze the statistics of iceberg dynamics while focusing specifically on the thick tabular icebergs that are most
480 likely to exhibit a long residence time when grounding on Bear Ridge. We aim to explore the differences in the dynamical
481 behaviour of icebergs of different sizes by comparing the THICK and MEDIUM populations. In order to preserve the large
482 thicknesses and horizontal areas in the THICK iceberg population, we do not allow the latter to melt during a SHORT 1-year
483 simulation.

484 We also run longer simulations labelled ‘LONG’, which are extended farther in time over 4 years to enable a
485 comparison between modelled and observed iceberg scours. We seed a much smaller subset of 14 identical icebergs in a
486 narrower box, bounded by 109.9° W, 109.2° W, 73.9° S, and 73.5° S (Figure 1), on the eastern flank of Bear Ridge. We explore
487 the simulated scours left by the THICK icebergs whose motion is arrested on Bear Ridge. In simulation LONG, we do allow
488 THICK icebergs to melt, as that affects their motion and residence times along the bottom: in our experiments with tabular
489 icebergs, melting (Section 3.3) is a mechanism that can allow icebergs to unground and resume free flotation. The released
490 meltwater is injected in the ocean and added to the precipitation at the surface.

491 **4 Results: Force balance and behaviour of thick tabular icebergs in the Amundsen Sea configuration**

492 In our Amundsen Sea numerical simulations, we observe the behaviour of thick tabular icebergs that are less typical
493 of the Arctic but often found near Antarctica and whose large volume and mass plays an important role in their dynamics
494 (Wagner et al., 2017). We are interested in the forces that act on these large icebergs in free flotation, upon grounding, and
495 upon ungrounding. Although our main focus is the behaviour of grounded icebergs, we also aim to understand the forces acting
496 on freely floating icebergs before they may ground and after they unground. In this analysis, we also test the impact of the new
497 iceberg dynamics and grounding algorithms that we have implemented in NEMO in the important Amundsen Sea regional
498 context.




501 **Figure 5: Climatology (1979 through 1982) of the Amundsen Sea configuration. a) potential density [kg/m³] referenced to the surface,
502 vertically averaged over the top 310 m with shallower regions masked; b) sea-surface height [m]; c) wind stress vectors [N/m²]; d)
503 barotropic streamfunction contours [0.1 Sv intervals]. The bottom depth [m] is superimposed in c and d. The depth range in c
504 chosen to highlight the location of the shallow Bear Ridge and a wider depth range in d to highlight the alignment of streamlines
505 and topography. The ice shelf edge is marked by the thick gray line.**

500

501
502
503
504
505

506




508
509 **Figure 6: Contributions of atmospheric drag (a), the oceanic pressure gradients (b), the Coriolis force (c), the ocean drag (d), the net
510 acceleration (e), and the average of a-d across all test icebergs and all timesteps (f) to the momentum budget of simulated freely
511 floating THICK icebergs in the Amundsen Sea in the SHORT simulation. Directions are denoted as F (forward, along the iceberg's
512 direction of motion), B (backward, against the direction of motion), L (to the left of the direction of motion), R (to the right). The
513 full length of the radial bars indicates the probability density function of acceleration in a given direction. For each panel in a-e, the
514 sum of all directions is 100%, but the bars along some of the directions are relatively small. For each bar in a-e, the range of
515 acceleration magnitudes is indicated by the colours, and the probability of each magnitude class is indicated by the radial distance.
516 Colour sections in a-e indicate the frequency distribution of the accelerations in each magnitude class. Note that the radial axis is
517 different between panels a-e, while the colour scale is common to panels a-e. Panel f has a separate legend and a different colour
518 scheme. The first 100 timesteps of the simulation, when the iceberg motion is initialized from rest, are omitted. Software written by
519 Daniel Pereira was used to create the wind rose (<https://www.mathworks.com/matlabcentral/fileexchange/47248-wind-rose>, last access: November 2025, MATLAB Central File Exchange)**

520

521

522



523

524

Figure 7: Same as Figure 6 but for freely floating MEDIUM icebergs in the Amundsen Sea in the SHORT simulation.

525 **4.1 Background conditions and force balance of freely-floating icebergs**

526 In order to understand the trajectories and behaviour of icebergs in the eastern Amundsen Sea, we first explore the wider
 527 climatic conditions, including the oceanic and atmospheric climatology and variability in the region. There is a relatively lower
 528 water column density to the east of the ridge compared to the water column over the ridge (Figure 5a), but this is compensated
 529 by a positive zonal SSH gradient between 112°W and 110°W (Figure 5b), which in combination with the Coriolis force would
 530 drive icebergs southward near the Ridge (Figure 5a). Overall, there is very little ocean transport across Bear Ridge although
 531 the prevailing winds are oriented north-westward across the ridge (Figure 5c,d).

532 Using model output from the large ensemble of freely-floating icebergs in Amundsen Sea Simulation SHORT, we
 533 compute statistics showing how often a given force acts along a particular direction relative to the iceberg motion (the length
 534 and orientation of the bars in the Figure 6 rose plots for THICK icebergs and Figure 7 for MEDIUM icebergs). We furthermore
 535 consider how often (the length of the individual color segments in Figures 6 and 7) a given force has a magnitude that falls
 536 within a particular range (the color scheme indicates the magnitude range). We point out that some forces represented in the
 537 iceberg model play a less prominent role compared to the main drivers. For example, the acceleration due to wave radiation is
 538 much smaller than $1.5 \times 10^{-6} \text{ m s}^{-2}$ and so is neglected from the figures, while the acceleration due to sea-ice drag is always
 539 smaller than the other terms (see Figure C1 in Appendix C).

540 We first consider the dynamics of freely-floating THICK icebergs in the period before they encounter the Bear Ridge
 541 or the shelf-edge, where they may ground or come to a stop. This helps us obtain a more comprehensive understanding of
 542 iceberg behaviour in the region. We also omit the first 100 timesteps of the simulation, when iceberg motion is initialized from
 543 rest. For freely-floating THICK icebergs, the most common situation by far is that the atmospheric form drag, the Coriolis
 544 force acting on icebergs, and the ocean form drag are nearly in a three-way balance (Figure 6) with very little net acceleration.
 545 The Coriolis force is oriented solely to the left, perpendicular to the direction of iceberg motion, while the acceleration due to
 546 wind stress and ocean pressure gradients is mostly oriented to the right of the direction of motion. On average, the
 547 corresponding forces on THICK icebergs are nearly orthogonal to the direction of motion and therefore do very little work on
 548 the freely floating large icebergs. This result is not peculiar to tabular icebergs in the Eastern Amundsen Sea but is more
 549 fundamental and agrees with the theoretical arguments of Wagner et al. (2017).

550 In our results, the (updated) ocean drag on freely floating icebergs plays a less prominent role compared to other
 551 forces and mostly opposes the forward motion of the bergs. Compare the small positive projection of the atmospheric drag
 552 along the forward direction of motion (Figure 6a and 7a) and the small deceleration due to ocean drag (Figure 6d and 7d). This
 553 result for large THICK icebergs may be expected on theoretical grounds. Here we briefly revisit the Wagner et al. (2017)
 554 arguments about the motion of large icebergs from a new perspective and with an attempt to recover the same result in fewer
 555 steps. We also specifically avoid making the assumption that the ocean's water column is in geostrophic balance, which is not
 556 necessarily applicable when modelling iceberg dynamics (e.g., Mountain, 1980). Unlike Wagner et al. (2017), we attempt to
 557 account for a realistic Ekman drift in our theoretical analysis and show that the same arguments remain valid.

558 We first consider the decomposition of the iceberg velocity \vec{u} into a component that matches exactly the ocean's
 559 horizontal velocity \vec{u}_{ocean} and a relative mismatch $\Delta\vec{u}$. The evolution, in an Eulerian sense, of the local ocean velocity over
 560 the upper ocean layer can be approximately expressed as:

$$561 \frac{\partial \vec{u}_{ocean}}{\partial t} \approx -f\hat{k} \times \vec{u}_{ocean} + (\text{Pressure gradient terms}) + (\text{Ekman ageostrophic acceleration due to wind stress}) \quad (19)$$

562 where we have assumed that the main source of ageostrophic transport in the Ekman layer (of thickness δ) comes from the
 563 tendency for Ekman drift due to surface wind stress $\vec{\tau}_{atm-ocean}$ (see Bigg et al., 1996). Under equilibrium conditions, the
 564 geostrophic component of $f\hat{k} \times \vec{u}_{ocean}$ balances the pressure gradient terms and the ageostrophic component of $f\hat{k} \times \vec{u}_{ocean}$
 565 vertically averaged over the Ekman layer balances the wind-stress acceleration.

566 While the ocean water column contains an Ekman spiral, the icebergs do not, and this is a potential source of mismatch
 567 between the velocity of icebergs and the surrounding ocean. In our study, we focus on thick icebergs, whose keel depth exceeds
 568 the depth of the Ekman layer. If we consider the relative iceberg-to-ocean velocity $\Delta\vec{u}$ averaged over the keel depth, we can
 569 express the Ekman acceleration term vertically integrated over the Ekman layer and then averaged over the full keel depth of
 570 the icebergs as

571 $|\text{Ekman acceleration}| = \left| \frac{\tau_{atm-ocean}}{\rho_0 D_{keel}} \right| \text{ over the keel depth,} \quad (20)$

572 such that the vertically-averaged Ekman component of \vec{u}_{ocean} is $\frac{\tau_{atm-ocean}}{f \rho_0 D_{keel}}$. The Ekman acceleration averaged over the depth
 573 of the Ekman layer is oriented along the wind direction such that the average Ekman drift $\frac{\tau_{atm-ocean}}{f \rho_0 D_{keel}}$ is to the left of the wind
 574 direction in the Southern Hemisphere.

575 The ocean pressure gradients exert the same force on the submerged portion of the iceberg as they would on an
 576 equivalent water parcel. Therefore, we can express the evolution of the iceberg-to-ocean velocity mismatch $\Delta\vec{u}$ as arising from
 577 a set of ageostrophic and dissipative processes. We can ignore the small terms due to sea-ice drag and wave drag. Furthermore,
 578 assuming that the horizontal gradients of \vec{u}_{ocean} are small along the iceberg trajectory, we can focus on the Eulerian evolution
 579 of $\Delta\vec{u}$:

$$580 \frac{\partial(\Delta\vec{u})}{\partial t} = \frac{\partial\vec{u}}{\partial t} - \frac{\partial\vec{u}_{ocean}}{\partial t} \approx \\ 581 \approx -f\hat{k} \times \Delta\vec{u} - C_{drag-ocean}\Delta\vec{u}|\Delta\vec{u}| - C_{drag-atm}(\vec{u} - \vec{u}_{atmos})|\vec{u} - \vec{u}_{atmos}| \\ 582 - \frac{\tau_{atm-ocean}}{\rho_0 D_{keel}} \quad (21)$$

583 Under the additional assumption that typically $|\vec{u}| \ll |\vec{u}_{atmos}|$, the above becomes

$$584 \frac{\partial(\Delta\vec{u})}{\partial t} \approx -f\hat{k} \times \Delta\vec{u} - C_{drag-ocean}\Delta\vec{u}|\Delta\vec{u}| + C_{drag-atm}\vec{u}_{atmos}|\vec{u}_{atmos}| - \frac{\tau_{atm-ocean}}{\rho_0 D_{keel}} \quad (22)$$

585 We assume that the main source of the mismatch $\Delta\vec{u}$, if present, can be attributed to the different atmospheric drag felt by
 586 icebergs relative to the surrounding water column, which experiences Ekman drift. Notice that the last two RHS terms depend
 587 primarily on winds as an external source of acceleration, while the first two RHS terms depend on the relative iceberg-to-ocean
 588 velocity $\Delta\vec{u}$. When winds input momentum into iceberg motion, and into the surrounding water column, this creates a
 589 differential acceleration between the iceberg and the ocean. If equation (22) is not in balance, the iceberg's relative speed $\Delta\vec{u}$
 590 increases, and motion gets deflected to the left by the Coriolis force until the first two terms on the RHS balance the last two,
 591 wind-dominated terms. In a steady state, the Coriolis force and ice-ocean drag balance the wind-dominated last two terms in
 592 equation 22.

593 The question is whether the Coriolis term (the first on the RHS) is sufficiently large to balance the excess atmospheric
 594 drag on icebergs (last two terms), or whether the atmospheric drag on icebergs and the Ekman acceleration on the water column
 595 are balanced by a relative drag between the ocean and the iceberg (second RHS term). With $\tau_{atm-ocean}$ that is typically
 596 $\sim 3 \times 10^{-1} N m^{-2}$ in our simulated region of interest, for THICK icebergs, the Ekman acceleration is

$$597 |\text{Ekman acceleration}| = \left| \frac{\tau_{atm-ocean}}{\rho_0 D_{keel}} \right| \sim \frac{\sim 3 \times 10^{-1} N m^{-2}}{(1025 \text{ kg m}^{-3})(\sim 350 \text{ m})} \sim 10^{-8} \text{ to } 10^{-7} \text{ m s}^{-2} \quad (23)$$

598 We point out that, following equation (2), both the acceleration due to wind drag and the acceleration due to ocean drag scale
 599 proportionally to their corresponding effective coefficient of form drag C_{drag} . The latter has units of a characteristic inverse
 600 length scale describing the iceberg $C_{drag} \propto 1/L_{iceberg}$ (see equation 2). In contrast, the Coriolis component of acceleration is
 601 not proportional to the iceberg's characteristic lengthscale. Therefore, the ratios between the groups of acceleration terms are:

$$\begin{aligned}
& \frac{\left| C_{drag-atm} \vec{u}_{atmos} \right|}{\left| f \hat{k} \times \Delta \vec{u} \right|} \propto \frac{C_{drag-atm} |\vec{u}_{atmos}|^2}{f |\Delta \vec{u}|} \propto \frac{(|\vec{u}_{atmos}|^2)}{L_{iceberg} f |\Delta \vec{u}|} \\
& \frac{|Ekman|}{\left| f \hat{k} \times \Delta \vec{u} \right|} \propto \frac{\tau_{atm-ocean}}{\rho_0 D_{keel} f |\Delta \vec{u}|} \\
& \frac{|f \hat{k} \times \Delta \vec{u}|}{c_{drag-ocean} \Delta \vec{u} |\Delta \vec{u}|} \propto \frac{f / |\Delta \vec{u}|}{c_{drag-ocean}} \propto L_{iceberg} f / |\Delta \vec{u}| \tag{24}
\end{aligned}$$

However, these effective coefficients for the atmospheric and oceanic form drag (Wagner et al., 2022) on icebergs are dependent not only on the iceberg dimensions ($\sim L_{iceberg}$), but are also proportional to the density of the respective ambient fluid: water or air. Hence, the coefficient for ocean drag is much larger than the coefficient of atmospheric drag, while the characteristic lengthscales $\frac{1}{c_{drag-ocean}} < \frac{1}{c_{drag-atm}}$. For example, in our LONG simulation with THICK icebergs, direct output from the model shows that $\frac{1}{c_{drag-ocean}} \sim 10^4$ m, while the atmospheric drag coefficient $\frac{1}{c_{drag-atm}}$ is on the order of 10^7 m, consistent with a 1:1000 density ratio between air and water. In our region, the southeastern Amundsen Sea, $f \sim 1.4 \times 10^{-4} \text{ s}^{-1}$, and the surface wind speed $|\vec{u}_{atmos}|$ is typically on the order of ~ 1 to 10 ms^{-1} .

The scaling of terms (24) implies that if ocean drag were to match to the Coriolis force, the speed of icebergs relative to the ambient currents would have to reach unrealistically high values $|\Delta \vec{u}| \sim \frac{|f|}{c_{drag-ocean}} \sim 1 \text{ ms}^{-1}$. On the other hand, the set of proportionalities (24) implies that a relative iceberg-ocean speed

$$|\Delta \vec{u}| \sim \frac{|\vec{u}_{atmos}|^2}{1000} \sim 1 \text{ cm s}^{-1} \tag{25}$$

is enough for an appropriately oriented Coriolis force to balance the direct atmospheric drag on large tabular icebergs (order 10^{-6} m s^{-2}) and also the smaller but nonnegligible Ekman term (up to order 10^{-7} m s^{-2}). Assuming that this balance holds, the ocean drag that results from $\Delta \vec{u}$ can be estimated to be

$$C_{drag-ocean} \Delta \vec{u} |\Delta \vec{u}| \sim 10^{-8} \text{ to } 10^{-7} \text{ m s}^{-2} \tag{26}$$

consistent with our findings (Figure 6). This suggests that for large tabular icebergs, the fictitious Coriolis force is sufficient *in magnitude* to balance the excess atmospheric drag while the relative iceberg-ocean drag remains small (Figure 6) in agreement with our results for THICK icebergs. Notice that the iceberg-ocean drag is similar in magnitude to the impact of Ekman drift. Therefore, we are able to recover the force balance of large THICK icebergs even in the presence of small but nonnegligible ageostrophic Ekman flow.

In addition, the scaling proportionalities (24) explain the orientation of the force vectors needed to achieve balanced motion. Only the Coriolis force is large enough to balance the excess atmospheric drag on icebergs relative to the water column. However, for that to happen, the iceberg's motion relative to the ocean has to be oriented at 90° to the left of the atmospheric drag, so that the Coriolis force can oppose the atmospheric drag. In this case, ocean drag would act against the direction of motion relative to the ocean. In the case of freely-floating THICK icebergs, the icebergs' motion is on average at an angle of 63° relative to the wind direction, which means that the Coriolis force balances almost 80% of the acceleration due to wind-stress (Figure 6f). The remaining 20% are balanced by the relatively smaller ocean drag, which is indeed oriented against the direction of motion (Figure 6f), as predicted on theoretical grounds.

The Coriolis force forms part of the dominant balance and steers motion in a direction nearly perpendicular to the driving force due to winds (Wagner et al., 2017). As a result, most of the time, the orthogonal wind stress force does not do significant work along the pathway of motion of large tabular icebergs. The important Coriolis effect also projects onto a small net acceleration to the left of the iceberg direction of motion (Figure 6), which differs from the Wagner et al. (2017) assumption of zero net acceleration but still does very little work along the iceberg pathway.


643 We can contrast the above results for THICK icebergs with our simulated MEDIUM icebergs, where the iceberg-
644 ocean relative speed and the associated ocean drag play a more prominent role (Compare Figure 7 for MEDIUM and 6 for
645 THICK icebergs). The ratio between ocean drag and the Coriolis force is 2/7 for THICK icebergs (Figure 6), while it is closer
646 to 1 for MEDIUM ones (Figure 7). At the onset of the SHORT simulations, the horizontal and vertical length scales of
647 MEDIUM icebergs are 2.6 and 1.6 times smaller, respectively, compared to THICK ones. Consistently, we see that $\frac{1}{c_{drag-atm}}$
648 at the beginning of the simulations is 2.5 times smaller for MEDIUM icebergs relative to the wider THICK ones. MEDIUM
649 icebergs experience further lateral and basal melting which causes additional reduction of their size relative to THICK icebergs.

650 Naturally, the Coriolis acceleration cannot change orientation and always remains directed to the left of the motion
651 even as the balance of forces on MEDIUM icebergs is shifted. At the same time, for our MEDIUM icebergs, the atmospheric
652 drag deviates from the orthogonality to the iceberg trajectory and projects more strongly along the axis of motion (Figure 7).
653 Thus, atmospheric drag seems to drive a more pronounced forward motion for MEDIUM icebergs than for THICK ones during
654 the course of the simulation. This in turn is matched by the enhanced ocean drag deceleration.

655 Both for THICK and MEDIUM freely-floating icebergs, the pressure gradient force is oriented mostly to the right of
656 the direction of motion, but with larger and more noticeable deviations for MEDIUM icebergs (Compare Figure 6 and Figure
657 7). We can invoke a different line of reasoning to explain the orientation of the pressure gradient force. We expect on the
658 aforementioned theoretical grounds (scaling arguments 24, and equations 25-26) that THICK large tabular floating icebergs
659 move with velocities close to the ambient ocean currents (see also Wagner et al., 2017) in terms of both magnitude and
660 direction. In our study, this notion is confirmed by the smaller ocean drag (Figure 6d), which suggests a small relative velocity
661 between icebergs and the ambient water. This implies that the relationship between ocean pressure gradients and iceberg
662 motion reflects the underlying relationship between ocean flow and pressure gradients. The consistent orientation of the ocean
663 pressure gradient force nearly orthogonal to the direction of motion of THICK tabular icebergs (interpret Figure 6b) can be
664 interpreted as an indication that geostrophic transport is a dominant component of the ambient ocean flow in the region of our
665 simulated icebergs, when vertically averaging over the THICK keel depth. As in the case of wind-induced acceleration, the
666 ocean pressure gradient force is mostly perpendicular to the iceberg trajectory and therefore does little work on the iceberg.

667 MEDIUM icebergs are released in the same region as THICK ones and over the same simulated historical period. So
668 we expect that the ocean flow has a similarly important geostrophic component. However, in the case of MEDIUM icebergs
669 (see Figure 7), we have a larger superimposed iceberg-to-ocean relative velocity. Therefore, the net direction of iceberg motion
670 force is less clearly orthogonal to the ocean pressure gradient force. The second major difference between MEDIUM and
671 THICK icebergs is that for MEDIUM ones, a larger fraction of their shorter keel is within the Ekman layer, and hence the
672 ocean drag on MEDIUM icebergs is more strongly influenced by the ageostrophic Ekman spiral in the upper ocean. This
673 contributes to further deviations of the iceberg motion away from the direction of the water column's predominantly
674 geostrophic flow, which otherwise would have been purely orthogonal to the iceberg's trajectory. The resulting nonzero
675 projection of the ocean pressure gradient force onto the MEDIUM iceberg trajectories means that this force does work on the
676 icebergs.

677 We furthermore explore the background conditions that give rise to the ocean pressure gradient force. The pre-existing
678 NEMO iceberg module represented the pressure gradients as a function of SSH alone. We analyse our new simulations with
679 the updated algorithms. Our results show that under realistic historical initial and boundary conditions, horizontal density
680 gradients in the ocean exert a noticeable impact on horizontal spatial variability in pressure (Figure 7 and Figure 8) and hence
681 iceberg accelerations in the Amundsen Sea. For MEDIUM ones, horizontal density gradients contribute a smaller component
682 of the full pressure gradient force. This decomposition of pressure, however, is averaged over multiple icebergs and timesteps.
683 The orientation and dominance of the density and SSH gradients varies across timesteps of the simulation and individual
684 icebergs.



686
687 **Figure 8: a) same as Figure 6f but for the contributions of SSH and density gradients to the acceleration due to pressure gradients**
688 **689 acting on THICK freely floating icebergs in the box where they were launched (Fig. 1), away from shallow topography and ice- shelves and b) as in a but for freely floating MEDIUM icebergs.**


690

691 **4.2 Force balance of grounded icebergs**


692 In the real Amundsen Sea, a fraction of the freely floating icebergs end up grounding along Bear Ridge. We now
693 explore the force balance of grounded icebergs. In order to better understand the iceberg-bathymetry interactions, we first run
694 a short NONDISSIPATIVE simulation without sediment resistance or solid-body friction along the bottom. In that case,
695 icebergs whose keels reach the bottom topography are only subject to the force of gravity that pushes them down the slope. If
696 the westward driving accelerations up the eastern flank of Bear Ridge are not large enough to overcome gravity, the icebergs
697 slide eastward down the slope until they resume free flotation. The Coriolis force curves the trajectories of these eastward-
698 moving icebergs northward and then westward, once again pushing them up the slope. This gives rise to a continuous repetitive
699 motion of icebergs up and down the eastern slope of Bear Ridge but also a net northward motion of the bergs parallel to the
700 Ridge (Figure 9). The shape traced out by the iceberg trajectory looks like a stretched-out spiral. Interestingly, such northward
701 displacement of icebergs is indeed observed along Bear Ridge, although not in this unrealistic continuous fashion but instead
702 with intermittent iceberg grounding (see Supplementary Movie).

703

704



705
 706 Figure 9: Simulated motion of an iceberg along Bear Ridge in an Amundsen Sea configuration of NEMO without sediment resistance
 707 or solid-body friction along the bottom. The topography, with contours 4 m apart in elevation, and the superimposed iceberg
 708 trajectory are shown in the Mollweide projection. The right panel shows the enlarged iceberg trajectory. Most of the iceberg
 709 trajectory corresponds to a state of free flotation, and the iceberg's vertical displacement along the topographic slope is only on the
 710 order of centimeters.




711
 712 Figure 10: Maps showing 14 iceberg trajectories at 20 minute time-intervals (different shades of purple corresponding to individual
 713 icebergs) over the four-year LONG simulation with THICK icebergs, where the colorbar indicates the seafloor depth along Bear
 714 Ridge highlighting regions shallower than 400 m. All icebergs in this test are launched east of 109.9° W and north of 74° S.

715 The second scenario we explore is that along Bear Ridge we impose a uniform 8-m deep layer of sediment with
 716 properties observed in the Amundsen Sea, as described in Section 2. We explore this scenario in our Amundsen Sea Simulation

717 LONG. In the LONG simulation with 14 THICK icebergs, three are advected southward and avoid Bear Ridge (Figure 10). A
 718 total of 11 icebergs ground on the eastern side of Bear Ridge. In contrast, the keels of MEDIUM icebergs do not extend deep
 719 enough to ground along the bottom.

720 In contrast to our NONDISSIPATIVE simulation, in experiment LONG, we have THICK icebergs grounded on Bear
 721 Ridge that are subject to sediment resistance. The latter counteracts forces that may otherwise accelerate the icebergs.
 722 Occasionally, icebergs also come into contact with the basement and experience Coulomb friction. In this scenario, we do not
 723 see a clear northward displacement of the icebergs as they ground and unground. Instead, they come to a stop quite soon after
 724 their keels start ploughing scours into the seafloor sediment. The ploughed scours are mostly oriented perpendicular to Bear
 725 Ridge and along the zonal direction (Figure 11). The icebergs grounded in the seafloor sediment melt in place, which thins
 726 them. In addition, simulated icebergs sporadically receive a strong enough push by the ocean and/or the atmosphere, which
 727 allows them to resume their motion briefly and intermittently and to plough scours deeper into the soft sediment (Figure 11b).
 728 These ploughing episodes are marked by sporadic short-lived thrusts of acceleration into the sediment (Figure 11) while, most
 729 of the time, the icebergs actually remain static when they are embedded in the sediment. This combination of processes creates
 730 scours whose depth relative to the surrounding ocean bottom is not uniform or changing monotonically along the scour length
 731 but is marked by vertical undulations (Figure 11a), which is reminiscent of the complexity of the observed scours (Figure 2).




732
 733 **Figure 11: Simulated scours along Bear Ridge from the LONG simulation with THICK icebergs. Panel a shows scour depth [m] and**
 734 **b shows the temporal evolution of the scours [days of the simulation].**

735 During the episodes of kinetic grounding, the dominant forces are the sediment resistance, which decelerates the
 736 icebergs, and atmospheric drag, which tries to push them forward (Figure 12). The ocean drag also makes a smaller relative
 737 contribution towards slowing down kinetically grounded icebergs that plough sediment scours. In the case of kinetic grounding,
 738 the force due to ocean drag is much weaker than the pressure gradient force and the Coriolis force. Moreover, the ocean drag
 739 does not exhibit a clear prevailing orientation against the direction of motion (Figure 12), and the ocean pressure gradients are
 740 not purely orthogonal to the iceberg trajectories. Instead of pointing mostly to the right, as is the case for freely-floating
 741 icebergs, the pressure gradient force on kinetically grounded icebergs can point in any of multiple directions. This can be
 742 compared with the very clear decelerating effect of ocean drag on freely floating icebergs (Figure 6d). The difference in the
 743 relative orientation of the sea-ice drag is noticeable, as well (Figure C1 in Appendix C). When the icebergs are grounded, sea-
 744 ice drag acts to push them forward rather than decelerate them or act in a direction orthogonal to their trajectory (Figure C1).
 745 Even more importantly, kinetically grounded icebergs experience a large forward push by atmospheric drag. However, the net
 746 acceleration, which is also positive, is smaller (Figure 12) due to the large dissipative forces (Figure 13 b,c) where sediment
 747 resistance plays a major role (Figure 13). The latter means that while in motion, these kinetically grounded icebergs moving
 748 within the sediment and along the bottom topography are rarely in force balance. Note that here we are considering the net

749 acceleration as a weighted average across all four Runge Kutta stages of a given model timestep but before the ‘stop’ flag
750 described in Section 3.4 is applied. Therefore, the metric in Figure 12e does not include this additional deceleration imposed
751 on icebergs as soon as the condition in equation 21 is fulfilled at any stage of the solver.

752 We furthermore explore the relative contribution of different physical mechanisms to the sediment resistance force
753 and the dominant terms that affect its magnitude (Figure 14). We consider icebergs with an assumed keel width (and hence
754 scour width) $W=90\text{m}$. We see that for scour depths D shallower than 3 m, the shear strength term that is proportional to D
755 dominates over the term due to the weight of the ploughed sediment in accordance with equation 11. In the deeper scours, the
756 density of the ploughed sediment determines to a large extent the total resistance experienced by a grounded iceberg. However,
757 the leading density and shear terms remain comparable for scour depths of 8 m. These results indicate that knowledge of both
758 the sediment density and the shear strength are necessary for an accurate estimate of the sediment resistance experienced by
759 grounding icebergs. On the other hand, the third component of equation (11), which is independent of keel width, remains
760 negligible for all scour depths when the scour width is $W=90\text{m}$. The scour width (which matches the keel width) must be
761 comparable to the scour depth for all three sediment resistance terms to be characterized by the same order of magnitude, and
762 this is not seen in the observations (Figure 2). However, such narrow scours are not resolvable in our observational dataset
763 (Table 1) and could occur in some settings.



765
766 **Figure 12:** As Figure 6 and 7 and with the same colorscale but for the case of kinetic grounding of THICK icebergs. The net term in
767 (e) is dominated by the dissipative forces shown in Figure 13. Occasionally, the orientation of the Coriolis force is not strictly to the
768 left of the iceberg motion for numerical reasons: the force is calculated using the projected directions of future motion at each Runge-
769 Kutta stage of the iceberg dynamics algorithm.

770 When icebergs plough all the way to the basement, they experience Coulomb friction and down-slope gravitational
771 acceleration (Figure 13). These additional forces bring icebergs to a halt within one day of contact with the basement, and the
772 icebergs are no longer kinetically grounded.

773 We draw a distinction between the term “kinetic grounding” which may still allow for motion along the bottom and
774 a static regime which implies no motion of the grounded iceberg. Icebergs do come to a complete stop even before their keels
775 plough deep enough to reach the solid basement beneath the sediment. In that case, sediment resistance balances the net
776 potential drivers of iceberg motion.




777 When the iceberg keels plough deep into the sediment and eventually reach the basement beneath, they start to feel
778 the effect of Coulomb friction and gravity. Our chosen kinetic Coulomb coefficient value of $\mu = 0.002$ is large enough to
779 bring icebergs rapidly to a halt once they reach the basement along Bear Ridge. In our simulation, we find that when the
780 icebergs reach the basement, they come to a complete stop within the same day and enter a static regime. In this regime, to a
781 first order, static Coulomb friction ($\mu_{stat} = 0.2$) balances the gravitational force that tries to pull the icebergs down the slope
782 (Figure 15). When the icebergs are statically grounded on the solid basement, Coulomb friction dominates, while sediment
783 resistance and other forces play a less prominent role (compare the orders of magnitude in Figures 13 and 15).

784 During the first year of the LONG simulation with THICK icebergs, the curvature term $\vec{a}_{curvature}$ in each horizontal
785 direction only has a nonnegligible magnitude ($> 1 \times 10^{-6} \text{ m s}^{-2}$) over less than 23 hours. During the times when the
786 geometric curvature term is greater than $1 \times 10^{-6} \text{ m s}^{-2}$ in magnitude, solid-body friction and gravitational forces dominate.
787 We also explore a range of values for the Coulomb coefficient μ , and they also give qualitatively similar results (Appendix B).

788 4.3 Comparison with observations

789 We next consider simulated iceberg residence times along Bear Ridge and qualitatively compare them with observations of
790 the real Amundsen Sea. This allows us to test the fidelity of our new algorithms and the validity of the assumptions we have
791 made regarding sediment and scour properties along Bear Ridge. We furthermore compare our new results with the output of
792 a pre-existing iceberg grounding algorithm in NEMO.

793
794
795
796
797
798
799
800



801

802


803

804

805

806


Figure 13: As Figure 12 but for the gravitational acceleration and the deceleration due to dissipative forces in the case of kinetic grounding of THICK icebergs: (a) gravitational acceleration present in a small fraction of the iceberg timestep samples (b), the Coulomb solid-body frictional deceleration (c), deceleration due to kinetic sediment resistance. Occasionally, the orientation of the deceleration due to sediment resistance force is not strictly against the iceberg motion for numerical reasons: the force is calculated using the projected directions of future motion at each Runge-Kutta stage of the iceberg dynamics algorithm.



807
808 **Figure 14: Components of the sediment resistance force [N] as a function of the scour depth D [m] for icebergs with keel width**
809 **W=90m and sediment properties typical of the Amundsen Sea. The individual terms and their sum refer to equations 6**
810 **through 11.**

811 In particular, we consider a pre-existing very simple iceberg grounding scheme in which icebergs whose keels reach
812 topography are moved back to their previous floating position and have their horizontal velocity set to zero (Olivé Abelló et
813 al., 2025 and Figure 16a). We apply this algorithm in the SIMPLE GROUNDING simulation with THICK icebergs, and we
814 find that bergs along Bear Ridge quickly leave the region and float away or are moved back to their previous floating position
815 and have their horizontal velocity set to zero, consistent with the simulations of Olivé Abelló et al. (2025). If the icebergs do
816 ‘ground’, that grounding is often near the ice shelf fronts. If our pressure gradient algorithm masks the ice shelves, as described
817 in section 3.4, then we see that we avoid trapping as many icebergs near the ice shelf edges (Figure 16b). In that case most
818 icebergs that we launch go around the northern side of the shallow Bear Ridge and quickly end up in the western Amundsen
819 Sea, within a month. Five icebergs circumvent the shallowest part of Bear Ridge by going around its southern end. However,
820 the SIMPLE GROUNDING scheme still remains fundamentally unable to keep the icebergs grounded along Bear Ridge for
821 long enough (Figure 16b).

822 In contrast, with our new updated grounding algorithm, THICK icebergs remain trapped along Bear Ridge for years
823 (Figure 16c). Crucially, assuming a deformable sediment layer above the basement is what allows the icebergs to remain
824 grounded along Bear Ridge (Figure 16 and Appendix B). This stands in contrast to the rapid northward export of icebergs in
825 the scenario without seafloor sediments or solid-body friction (compare Figures 9 against 10, 11 and 16). While icebergs in
826 the real Amundsen Sea do propagate northward, they also remain grounded for long periods of time suggesting that frictional
827 forces are an important consideration and should be retained in the model. Our results are in qualitative agreement with the
828 observations suggesting that thick icebergs may be trapped on Bear Ridge for months or years.




829

830 **Figure 15:** As Figure 12 and 13 but in the case of static grounding of THICK icebergs on solid basement with (a) showing the net
 831 productive force per unit mass excluding gravity, (b) the gravitational force per unit mass, (c) the sediment resistance per unit mass
 832 and (d) static Coulomb friction in the case of static grounding of THICK icebergs. The scale is non-linear and shows the order of
 833 magnitude. Note that unlike Figures 12-13, the directions are here the cardinal points as the icebergs do not move. In order to show
 834 meaningful geographical directions, here we include only icebergs grounded on Bear Ridge and nowhere else. In the case of
 835 grounding along the basement, the icebergs in the simulation always come to rest and enter a static regime within a day. The static
 836 Coulomb frictional force per unit mass shown here is the maximum achievable magnitude, but the effective friction does not exceed
 837 the sum of the other sources of acceleration, which are dominated by gravity.

838

839



840
841 **Figure 16: Locations of THICK icebergs during the course of the (a) pre-existing SIMPLE GROUNDING; (b) the pre-existing
842 SIMPLE GROUNDING scheme and with masked pressure gradient forces along the iceshelf edge; and (c) LONG
843 dissipative simulation with the new grounding scheme. The topography shallower than 560 m is superimposed with gray
844 contours, 20 m apart. The blue line denotes the ice shelf fronts, where many icebergs come to a stop under the pre-existing SIMPLE GROUNDING scheme.
845 Iceberg locations 4 years after launch are shown for the simulation with new grounding scheme only.**

846 While the transition between static and kinematic grounding and scouring is determined by basal melt rates and winds,
847 the full ungrounding of icebergs is a much slower process. In the real ocean, grounded icebergs may capsize if their horizontal
848 dimensions shrink faster than the decrease in their vertical thickness and their new geometric orientation becomes unstable. In
849 order to explore the potential for capsizing, we have extended our four-year LONG simulation with THICK icebergs up to 20
850 years by applying a repeat cycle of the same four-year surface boundary conditions. We see that in our extended LONG
851 simulation, the rate of lateral melting indeed exceeds the rate of basal melting, with a potentially unrealistically large
852 contribution of wave erosion to the former. One may expect that this causes widespread capsizing of grounded icebergs.
853 However, the pre-existing NEMO capsizing criterion incorrectly compares horizontal length rather than horizontal width
854 against vertical thickness. As a result, the excessive lateral melting of individual grounded icebergs eventually reduces the
855 horizontal width to zero within 17 years and completely destroys the iceberg before the horizontal length decreases enough for
856 the iceberg to capsize (not shown). Olivé Abelló et al. (2025) correct the unphysical capsizing criterion in NEMO, a change
857 that will be implemented in new model configurations. On the other hand, eliminating the relevant bias in the horizontal wave
858 erosion remains an important outstanding issue left to future studies.

859 **5 Summary and conclusions**

860 We have introduced a new and improved representation of iceberg dynamics and iceberg grounding capability in NEMO.
861 These updates to the model are verified using idealized test configurations before being applied to an eddy-permitting regional
862 simulation of the Amundsen Sea forced with historical boundary conditions. We note that in this initial study our specific
863 intention is to develop the new model physics, and so we delay a full multi-year realistic simulation of iceberg calving and
864 grounding to a subsequent study. A companion paper presents an improved iceberg thickness distribution in NEMO based on
865 the thickness of the Antarctic ice-shelves from which icebergs calve (Olivé Abelló et al., 2025). The companion paper also
866 introduces an objective definition of iceberg size classes implemented in NEMO (Olivé Abelló et al., 2025). Future work will
867 combine these two approaches. Once we are in possession of a full realistic simulation, a much fuller comparison to observed
868 iceberg grounding can take place, with additional tuning of the iceberg grounding parameters. This would require the
869 application of advanced techniques for tracking iceberg grounding episodes from satellite imagery, perhaps involving artificial
870 intelligence (AI) techniques. A full comparison of simulated and observed iceberg scours would also be extremely powerful
871 in constraining the model. In addition, future radar sounding observations may reveal the distribution of iceberg keel shapes.

872 The Amundsen Sea experiments suggest that in this region the ocean pressure gradients acting on icebergs are not
873 determined solely by spatial variability in SSH. Instead, our results highlight the important direct contribution of horizontal
874 gradients in the ocean density, which are separate from the contribution of density through steric SSH variability. In addition,
875 we show that large freely drifting tabular icebergs often enter a dynamical regime characterized as a balance between surface
876 winds and the Coriolis force. In contrast, smaller icebergs experience stronger turbulent ocean drag. Overall, freely floating
877 tabular icebergs in the Amundsen Sea are not subject to strong net acceleration along or against their direction of motion, but
878 tend to get deflected to the left by Coriolis force. This is consistent with the theoretical arguments presented by Wagner et al.
879 (2017) who suggest a similar force balance for large tabular icebergs when the ocean is in a purely geostrophic regime. In our
880 analysis, we derive more general scaling arguments that describe the momentum budget of large and medium icebergs even in
881 the presence of ageostrophic Ekman flow.

882 Our results for the force balance of freely floating icebergs also highlight the potential dynamical importance of
883 iceberg fragmentation. If large tabular icebergs get fragmented, the smaller fragments will get advected in a different direction
884 compared to a parent iceberg. Under identical oceanic and atmospheric conditions, large and small icebergs originating from
885 the same place drift towards and melt in different geographical locations. England et al. (2022) highlight this effect by directly
886 imposing the Wagner et al. (2017) analytical result on simulated icebergs. Furthermore, our findings for the dependence of the
887 turbulent iceberg-ocean drag on iceberg size have implications for the rate of iceberg subsurface melting. Jenkins et al. (2010)
888 and Davis et al. (2023) point out that the rate of ice shelf basal melting depends on the velocity beneath the shelf. By analogy,
889 the rate of iceberg subsurface melting, especially for large tabular icebergs, exhibits a similar dependence on the horizontal
890 length scale through the latter's impact on the relative velocity. Note that the formulas in NEMO explicitly represent the direct
891 effect of iceberg lengthscale on basal melting through a -0.2 power law, while the potentially large impact of horizontal size
892 via changes in relative velocity is an emergent phenomenon. This once again points to the importance of the fragmentation of
893 large icebergs into smaller ones, a first order process, which is not currently represented in NEMO but is implemented in other
894 models such as GFDL (Huth et al., 2022).

895 Our updated algorithms show the force balance of icebergs interacting with the ocean bottom. The new grounding
896 representation in NEMO allows us to reproduce realistic bottom scours and grounding along Bear Ridge in the Amundsen Sea.
897 We compare our simulations against observations in the region. Grounding in deformable sediments is marked by alternating
898 periods of slowdown and arrest followed by episodic acceleration that thrusts the icebergs deeper into the sediment (Figure
899 11). The sediment resistance force is characterized by comparable contributions due to the density and the shear strength of
900 the deformable layer. Consistent with observations, the presence of a sediment layer in our simulations allows thick icebergs
901 to remain trapped along Bear Ridge on a timescale of years (Figure 16c) despite their susceptibility to basal and lateral melting.

902 The long residence time of icebergs along the Ridge and their eventual northward motion affect the distribution of iceberg
903 meltwater (Bett et al., 2020) but also, more importantly, create the Amundsen Sea Polynya and the dipole in the distribution
904 of sea ice growth on either side of Bear Ridge (Figure 1). In contrast, in the absence of sediment resistance and solid-body
905 friction with the basement, icebergs would continuously move northward along the eastern slope of Bear Ridge.

906 The model algorithms in this study will allow us to explore the existence of possible feedback mechanisms between
907 iceberg grounding along Bear Ridge in the Amundsen Sea and the calving of new icebergs from the shelf under changing
908 climate conditions, as suggested by Bett et al. (2020). Ultimately, these processes may be more accurately represented
909 circumpolarly, with consequent improvements in regional ocean modelling, particularly over the continental shelves. This is
910 a region that models struggle to accurately recreate, notably the interactions between melt and sea ice distribution, polynyas,
911 dense shelf water formation and wider ocean vertical circulation (e.g. Aguiar et al., 2024). The strength of this mechanism
912 and its future response to climate change have major implications for the freshwater budget of the Amundsen Sea, for the
913 stability of ice shelves, and for the Antarctic contribution global sea level rise. Therefore, the proper representation of Antarctic
914 iceberg grounding and residence times in models is an important prerequisite for developing reliable future climate projections.

915

916 **Appendix A. Supplementary Movie**

917 We enclose a hyperlink (<https://doi.org/10.5446/70447>) to a supplementary movie composed of Copernicus Sentinel-1 SAR
918 (synthetic aperture radar) images taken over the time period 2017-2024, an extension of the movie provided in Bett et al.
919 (2020). The movie highlights the locations of grounded icebergs of various sizes along the shallow Bear Ridge. Many of the
920 still images in the movie also show the dipole in sea-ice concentration with larger values to the east of the Bear Ridge.



922

923 **Supplementary Movie A. Available at the TIB AV Portal (Kostov et al., 2025a) : <https://doi.org/10.5446/70447>**

924

925

926

927

928 **Appendix B. Effects of topographic curvature, substepping parameters, and the Coulomb coefficient of solid-body
929 friction**

930 We explore the parameter space of possible values for the unconstrained Coulomb coefficient of solid-body friction μ across
931 different shorter repetitions of the Amundsen Sea Simulation B that extend for more than 350 days. In each case, we vary the
932 coefficient of kinetic friction, while the coefficient of static friction is always set to be 100 times larger than the kinetic one.

933 We find that varying the value of the kinetic μ and static coefficient μ_{stat} by different orders of magnitude does not
934 affect the statistics for episodes of motionless grounding (Figure B1). We infer two conclusions from this result. First, our
935 results are robust with respect to the unconstrained value of the Coulomb coefficient for solid body friction. Second, the
936 duration of episodes of grounding along the bottom seem to be governed by sediment resistance and melting processes that
937 thin the icebergs.


938 We furthermore test the importance of the “curvature term” from equation 17. When an iceberg moves upwards from
939 a gentler slope to an even steeper slope, some of its horizontal momentum is reoriented in the vertical direction. This gives rise
940 to an apparent horizontal deceleration term $\vec{a}_{curvature}$ that is purely geometric, independent of gravity and buoyancy, unrelated
941 to turbulent drag, and unrelated to dissipative forces such as sediment resistance or solid body friction. For example, if the
942 topographic slope β changes at a rate $\partial\beta/\partial x$ along the pathway of a grounded iceberg moving in the x-direction, then u , the
943 x-component of the iceberg’s horizontal velocity also changes as:

$$944 \vec{a}_{curvature} \cdot \hat{x} = -|u^2 \tan \beta| \frac{\partial \beta}{\partial x} \hat{x} \quad (B1)$$


945 and analogously for a component $\vec{a}_{curvature} \cdot \hat{y}$ along the y-direction. The term $\vec{a}_{curvature}$ represents the direct impact of
946 seafloor basement geometry on horizontal velocity and can be understood through dimensional analysis. For example, in
947 equation (B1), the factor u^2 has dimensions of length squared over time squared $\left[\frac{L^2}{T^2}\right]$, while the horizontal spatial derivative
948 $\frac{\partial}{\partial x}$ has dimensions of inverse length $[L^{-1}]$. Therefore, $\vec{a}_{curvature}$ is an acceleration term with dimensions of length over squared
949 time $\left[\frac{L}{T^2}\right]$. Qualitatively, the tangent $\tan \beta$ of the topographic slope β tells us how changes in the slope along the horizontal
950 direction re-orient the velocity vector along the vertical direction. For instance, when the slope gets steeper, the iceberg’s
951 velocity vector is deflected more towards the vertical direction. Numerical tests demonstrate that the term $\vec{a}_{curvature}$ becomes
952 non-negligible compared to other acceleration terms when icebergs slide along realistic topography. However, this happens
953 only when the angle of the interpolated topography changes along the iceberg’s pathway. The curvature factor is non-negligible
954 if we focus on the small-scale shape and evolution of iceberg bottom scours, but does not play a first-order role in the overall
955 residence times of grounded icebergs on the scale of the ocean domain resolution as revealed by the tests in Section 4. We also
956 see that the effect of topographic curvature does not contribute to qualitative differences in the shape and evolution of iceberg
957 scours (Figure B1).

958 We also show that in the presence of a sediment layer above the crystalline basement, the time-stepping of the
959 interaction with the shape of the topography becomes less important. Using or not using temporal sub-stepping does not play
960 a first order role when icebergs ground in the sediment above the crystalline base (Figure B1).

961 We do find that having no Coulomb friction (Figure B1) or having neither friction, nor sediment resistance (Figure 9
962 in the main text) are extreme cases where the absence of processes makes a noticeable difference. If there is no solid-body
963 friction but there is sediment resistance, icebergs plough swirls and half-loops or exhibit a repetitive back and forth motion
964 along the slope (Figure B1). If no dissipative force acts on a grounded iceberg, then its trajectory can trace out a spiral motion
965 (Figure 9 in the main text). Interestingly, iceberg scours in the shape of swirls and spirals are indeed observed along the ocean
966 bottom in the real world, but they are traditionally attributed to the effect of tides (See Figure 2 and Section 2). In our model
967 configurations, we do not represent tidal effects, yet we observe such features and point out the important role of the Coriolis
968 force for driving oscillatory iceberg motion.



971 **Figure B1:** As in Figure 11 but in simulations with a range of kinetic Coulomb coefficient values μ , sub-stepping choices, and
 972 representations of the topographic curvature effect.



976
 977 **Figure B2:** As in Figure B1 but in simulations with zero Coulomb friction and 8-m sediment thickness. Each row represents an
 978 individual iceberg. The left column shows scour depths, and the right column shows their evolution in time during the LONG
 979 simulation with THICK icebergs.


980 We also probe the impact of temporal substepping. In the absence of sediment resistance forces, simulating the effect
 981 of gravity on grounded icebergs does require shorter timesteps and may necessitate substepping for iceberg dynamics. Here
 982 we present an important test of the iceberg substepping and its impact on gravitational acceleration/deceleration. In this test,
 983 we release a 97-m thick iceberg in an idealized rectangular basin that is not in a rotating frame of reference (the Coriolis force
 984 is set to zero). All wind/ocean/sea ice drag terms are set to zero. Iceberg melting is switched off, and so is solid-body friction.
 985 The sediment depth along the bottom is set to zero, eliminating any sediment resistance. Under these conditions, the only term
 986 remaining in the momentum balance is the gravitational disequilibrium term $\vec{a}_{gravity}$. The iceberg is initialized with south-
 987 eastward velocity above a sloping bottom whose depth decreases by 10m per 1 km in the southward direction, going from

988 100m to 80m. As the iceberg moves up the bottom slope and out of free flotation, its kinetic energy is not dissipated but rather
989 converted to potential energy. The gravity term should then convert this back into kinetic energy, accelerating the berg back
990 down the slope, and this descent should exactly mirror the berg's ascent up the slope, like an idealized perfectly un-damped
991 pendulum.


992 We first test a case where the iceberg timestep equals the ocean timestep. The iceberg travels south-eastward up the
993 slope, decelerating under gravity, and then reverses direction to slide north-eastward, accelerated by the same term. However,
994 when the iceberg starts sliding back down the slope, its descent is not symmetric with respect to its initial ascent, despite the
995 absence of external driving or dissipative forces (Figure B3); there is an asymmetry in the iceberg's meridional velocity going
996 up the slope and coming back down (Figure B3).

997 In our numerical tests, we found that this error is caused by an inadequate time-resolution of the iceberg deceleration
998 and acceleration by the gravitational acceleration, which can be much larger than other sources in realistic cases. When we
999 introduce temporal substepping within the iceberg module (four iceberg steps per one ocean step), we see a better symmetry
1000 between the iceberg's ascent and descent along the slope (Figure B3). This indicates an improved conservation of mechanical
1001 energy which is converted between kinetic and potential energy and this was found to be important in realistic cases.

1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020



1021



1022

1023 **Figure B3: Asymmetry between the ascent and descent (a); and deceleration and acceleration (b) of an iceberg along a frictionless**
 1024 **slope with and without 1075 sub-stepping.**

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037


1038

1039

1040

1041 **Appendix C. Sea-ice drag on freely-floating and grounded icebergs**

1042 We calculate the average magnitude and orientation of the sea-ice drag for the case of freely-floating THICK and
1043 MEDIUM icebergs, as well as, kinetically grounded THICK icebergs. In the case of statically grounded icebergs, the solid-
1044 body friction and gravity per unit mass are four orders of magnitude larger than the sea-ice contribution, and we do not consider
1045 that case here. Figure C1 highlights the different orientation of the sea-ice drag for freely-floating icebergs and grounded ones:..
1046 the drag projects onto the forward direction of motion of kinetically grounded icebergs.
1047



1048
1049 **Figure C1: As Figure 6f and 7f in the main text but for sea-ice drag on freely-floating THICK icebergs (dark blue), freely-floating**
1050 **MEDIUM icebergs (cyan), and kinetically-grounded THICK icebergs (purple).**

1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066

D1. Table of Symbols

Symbol	Unit	Description
$\vec{a}_{ocean\ drag}$	ms^{-2}	Acceleration due to ocean drag
$\vec{a}_{atmos.drag}$	ms^{-2}	Acceleration due to atm. drag
$\vec{a}_{pressure\ gradient}$	ms^{-2}	Acceleration due to horizontal oceanic pressure gradient forces
$\vec{a}_{Coriolis}$	ms^{-2}	Acceleration due to Coriolis
$\vec{a}_{sea-ice\ drag}$	ms^{-2}	Acceleration due to sea-ice drag
\vec{a}_{waves}	ms^{-2}	Acceleration due to ocean waves
$\vec{a}_{sediment}$	ms^{-2}	Deceleration due to sediment resistance
$\vec{a}_{gravity}$	ms^{-2}	Acceleration due to gravity
\vec{a}_{ground}	ms^{-2}	Deceleration due to all forces active only on grounded icebergs
\vec{a}_{solid}	ms^{-2}	Deceleration due to solid-body friction
$\vec{a}_{static-solid}$	ms^{-2}	Deceleration due to solid-body friction in a static regime
$\vec{a}_{curvature}$	ms^{-2}	Effective horizontal acceleration due to topographic curvature
\vec{a}_{net}	ms^{-2}	Net acceleration
$\vec{a}_{net\ drivers}$	ms^{-2}	Acceleration due to all nondissipative forces
C_{drag}	m^{-1}	Drag coefficient parameter
$C_{drag-ocn}$	m^{-1}	Ocean drag coefficient parameter
$C_{drag-atm}$	m^{-1}	Atmospheric drag coefficient over icebergs
$C_{drag-atm-to-ocean}$	m^{-1}	Atmospheric drag coefficient over ocean
D	m	Scour depth
D_{keel}	m	Keel depth
f	s^{-1}	Coriolis parameter
g	ms^{-2}	Magnitude of the acceleration due to gravity
\vec{g}	ms^{-2}	Acceleration due to gravity
H_{Total}	m	Iceberg thickness
$H_{freeboard\ anom.}$	m	Iceberg freeboard anomaly (out of buoyancy-gravity equilibrium)
\vec{k}		Vertical direction
$L_{iceberg}$	m	Characteristic iceberg lengthscale
$M_{iceberg}$	kg	Iceberg mass
\hat{n}		Unit vector normal to the topographic plane

P	Pa	Ocean pressure
\hat{s}		Unit vector in the direction of the topographic slope
Δt	s	Timestep length
$\Delta \vec{u}$	ms^{-1}	Difference between the iceberg velocity and the velocity of the ambient ocean
\vec{u}_{fluid}	ms^{-1}	Velocity of ambient fluid
$\vec{u}_{atmos.}$	ms^{-1}	Atmospheric velocity
\hat{u}	ms^{-1}	Unit vector in the direction of the iceberg motion
W	m	Scour width
x, y, z		Horizontal dimensions
α	rad	Angle of the trajectory relative to the horizontal plane
β	rad	Maximum local angle of the topographic slope
γ'	Nm^{-3}	Submerged unit weight
δ	m	Thickness of the Ekman layer
μ		Coefficient of Coulomb friction
μ_{static}		Coefficient of static Coulomb friction
ρ_0	$kg\ m^{-3}$	Reference density of ocean water
ρ_{sat}	$kg\ m^{-3}$	Saturated density of sediment
ρ_{water}	$kg\ m^{-3}$	Water density
ρ_{ice}	$kg\ m^{-3}$	Iceberg density
τ	$Pa\ (Nm^{-2})Pa$	Shear strength resistance of sediment
$\tau_{atm-ocean}$	$Pa\ (Nm^{-2})$	Magnitude of the wind stress at the ocean surface

1069

1070 **Code availability**

1071 The updated code for the NEMO iceberg module is publicly available at the following repository
 1072 <https://doi.org/10.5281/zenodo.15484879>

1073

1074 **Acknowledgements**

1075 The multibeam-bathymetric data were compiled from the NCEI (National Centers for Environmental Information; USA) and
 1076 the NERC PDC (Polar Data Centre; UK) data centres. We thank the masters and crew of all ships involved in data collection;
 1077 we thank Frank Nitsche for his help collating the bathymetric data. This research was supported by Ocean Cryosphere
 1078 Exchanges in ANtarctica: Impacts on Climate and the Earth system, OCEAN ICE, which is funded by the European Union,
 1079 Horizon Europe Funding Programme for research and innovation under grant agreement Nr. 101060452, 10.3030/101060452.
 1080 OCEAN ICE Internal contribution number 26. This work was funded by UK Research and Innovation (UKRI) under the UK

1081 government's Horizon Europe funding Guarantee 10048443. A. Olivé Abello, N. Jourdain and P. Mathiot received funding
1082 from Agence Nationale de la Recherche - France 2030 as part of the PEPR TRACCS programme under grant numbers ANR-
1083 22-EXTR-0008 (IMPRESSION-ESM) and ANR-22-EXTR-0010 (ISCLIM).
1084

1085 **Author contribution**

1086 YK developed new model algorithms in coordination with AOA's companion manuscript, conducted numerical simulations,
1087 and analysed output. KAH, JAS, and AHF provided observationally-based data and images. YK, PRH, KAH, JAS, NCJ, PM,
1088 AOA, AHF, and AJSM contributed content and commented on the manuscript.

1089 **Competing interests**

1090 NCJ is a member of the TC editorial board. The authors declare that they have no other conflict of interest.
1091

1092 **References**

1093 Aguiar, W., Lee, S.-K., Lopez, H., Dong, S., Seroussi, H., Jones, D. C., & Morrison, A. K. (2023). Antarctic Bottom Water
1094 sensitivity to spatio-temporal variations in Antarctic meltwater fluxes. *Geophysical Research Letters*, 50, e2022GL101595.
1095 <https://doi.org/10.1029/2022GL101595>

1096
1097 Arrigo, K.R., Lowry, K.E., and van Dijken, G.L. (2012) Annual changes in sea ice and phytoplankton in polynyas of the
1098 Amundsen Sea, Antarctica, *Deep Sea Res. Part II: Topical Studies in Oceanography*, Vol. 71–76, 5–15,
1099 <https://doi.org/10.1016/j.dsr2.2012.03.006>.

1100
1101 Asay-Davis, X. S., Cornford, S. L., Durand, G., Galton-Fenzi, B. K., Gladstone, R. M., Gudmundsson, G. H., Hattermann, T.,
1102 Holland, D. M., Holland, D., Holland, P. R., Martin, D. F., Mathiot, P., Pattyn, F., and Seroussi, H.: Experimental design for
1103 three interrelated marine ice sheet and ocean model intercomparison projects: MISMIP v. 3 (MISMIP +), ISOMIP v. 2
1104 (ISOMIP +) and MISOMIP v. 1 (MISOMIP1), *Geosci. Model Dev.*, 9, 2471–2497, <https://doi.org/10.5194/gmd-9-2471-2016>,
1105 2016.

1106
1107 Barnes, P. W., and R. Lien (1988), Icebergs rework shelf sediments to 500-m off Antarctica, *Geology*, 16(12), 1130–1133.

1108
1109 Bass DW. Stability of Icebergs. *Annals of Glaciology*. 1980; 1:43–47. doi:10.3189/S0260305500016943

1110
1111 Bett, D. T., et al. (2020) The impact of the Amundsen Sea freshwater balance on ocean melting of the West Antarctic Ice
1112 Sheet. *JGR: Oceans*, <https://doi.org/10.1029/2020JC016305>

1113
1114 Bigg GR, et al. 1996. Prediction of iceberg trajectories for the North Atlantic and Arctic oceans. *Geophys. Res. Lett.*
1115 23:243587–90
1116

1117
1118 Bigg, G.R., et al. (1997) Modelling the dynamics and thermodynamics of icebergs, *Cold Regions Science and Technology*,
Vol. 26, 2, 113–135, [https://doi.org/10.1016/S0165-232X\(97\)00012-8](https://doi.org/10.1016/S0165-232X(97)00012-8)

1119 Bigg, G.R., Cropper, T.E., O'Neill, C.K. *et al.* A model for assessing iceberg hazard. *Nat Hazards* **92**, 1113–1136 (2018).
1120 <https://doi.org/10.1007/s11069-018-3243-x>

1121 Christopher S. Brown, Andrew M.W. Newton, Mads Huuse, Francis Buckley, Iceberg scours, pits, and pockmarks in the North
1122 Falkland Basin, *Marine Geology*, Volume 386, 2017, Pages 140-152, <https://doi.org/10.1016/j.margeo.2017.03.001>.

1123

1124 Caillet, Justine & Jourdain, Nicolas & Mathiot, Pierre & Hellmer, Hartmut & Mouginot, Jérémie. (2022). Drivers and
1125 Reversibility of Abrupt Ocean State Transitions in the Amundsen Sea, Antarctica. *Journal of Geophysical Research: Oceans*.
1126 128. 10.1029/2022JC018929.

1127

1128 Cefarelli, Adrián & Ferrario, Martha & Vernet, Maria. (2016). Diatoms (Bacillariophyceae) associated with free-drifting
1129 Antarctic icebergs: taxonomy and distribution. *Polar Biology*. 39. 10.1007/s00300-015-1791-z.

1130

1131 Cenedese, C. and Straneo, F. Icebergs Melting. *Annu. Rev. Fluid Mech.* 2023. 55:377–402

1132

1133 Chari, T. R. (1975) Some geotechnical aspects of iceberg grounding. Doctoral (PhD) thesis, Memorial University of
1134 Newfoundland

1135

1136 Chari, T.R. et al. (1980) Environmental factors affecting iceberg scour estimates, *Cold Regions Sci. and Tech.*, DOI:
1137 [https://doi.org/10.1016/0165-232X\(80\)90050-6](https://doi.org/10.1016/0165-232X(80)90050-6).

1138

1139 R.W. Clark, J.S. Wellner, C. Hillenbrand, R.L. Totten, J.A. Smith, L.E. Miller, R.D. Larter, K.A. Hogan, A.G.C. Graham,
1140 F.O. Nitsche, A.A. Lehrmann, A.P. Lepp, J.D. Kirkham, V.T. Fitzgerald, G. Garcia-Barrera, W. Ehrmann, and L. Wacker,
1141 (2024) Synchronous retreat of Thwaites and Pine Island glaciers in response to external forcings in the presatellite era, *Proc.
1142 Natl. Acad. Sci. U.S.A.* 121 (11) e2211711120, <https://doi.org/10.1073/pnas.2211711120>.

1143 Condron, A., Hill, J.C. Timing of iceberg scours and massive ice-rafting events in the subtropical North Atlantic. *Nat
1144 Commun* 12, 3668 (2021). <https://doi.org/10.1038/s41467-021-23924-0>

1145

1146 Davis, P.E.D., Nicholls, K.W., Holland, D.M. *et al.* Suppressed basal melting in the eastern Thwaites Glacier grounding
1147 zone. *Nature* **614**, 479–485 (2023). <https://doi.org/10.1038/s41586-022-05586-0>

1148

1149 Davison BJ, Hogg AE, Gourmelen N, Jakob L, Wuite J, Nagler T, Greene CA, Andreasen J, Engdahl ME. (2023) Annual mass
1150 budget of Antarctic ice shelves from 1997 to 2021. *Sci Adv.* Oct 13; 9(41):eadi0186. doi: 10.1126/sciadv.adi0186

1151

1152 Darwin, C. R. 1855. On the power of icebergs to make rectilinear uniformly-directed grooves across a submarine undulatory
1153 surface. *The London, Edinburgh and Dublin Philosophical Magazine* 10(64), 96–98.
1154 <https://doi.org/10.1080/14786445508641938>

1155

1156 Dowdeswell, J. A., and Bamber J. L. (2007), Keel depths of modern Antarctic icebergs and implications for sea-floor scouring
1157 in the geological record, *Marine Geology*, 243, 120-131.

1158

1159 Duprat, L., Bigg, G. & Wilton, D. (2016) Enhanced Southern Ocean marine productivity due to fertilization by giant
1160 icebergs. *Nature Geosci* **9**, 219–221. <https://doi.org/10.1038/ngeo2633>

1161 Pierre Dutrieux *et al.*, Strong Sensitivity of Pine Island Ice-Shelf Melting to Climatic Variability. *Science* **343**, 174–178 (2014).
1162 DOI:[10.1126/science.1244341](https://doi.org/10.1126/science.1244341)

1163 England, M.R. et al. ,Modeling the breakup of tabular icebergs. *Sci. Adv.* **6**, eabd1273(2020). DOI:[10.1126/sciadv.abd1273](https://doi.org/10.1126/sciadv.abd1273)

1164 FitzMaurice, A., F. Straneo, C. Cenedese, and M. Andres (2016), Effect of a sheared flow on iceberg motion and
1165 melting, *Geophys. Res. Lett.*, **43**, 12,520–12,527, doi:[10.1002/2016GL071602](https://doi.org/10.1002/2016GL071602).

1166 Fowler AC. Weertman, Lliboutry and the development of sliding theory. *Journal of Glaciology*. 2010;56(200):965-972.
1167 doi:[10.3189/002214311796406112](https://doi.org/10.3189/002214311796406112)

1168

1169 Fox-Kemper B, Adcroft A, Böning CW, Chassignet EP, Curchitser E, Danabasoglu G, Eden C, England MH, Gerdes R,
1170 Greatbatch RJ, Griffies SM, Hallberg RW, Hanert E, Heimbach P, Hewitt HT, Hill CN, Komuro Y, Legg S, Le Sommer J,
1171 Masina S, Marsland SJ, Penny SG, Qiao F, Ringler TD, Treguier AM, Tsujino H, Uotila P and Yeager SG (2019) Challenges
1172 and Prospects in Ocean Circulation Models. *Front. Mar. Sci.* **6**:65. doi: [10.3389/fmars.2019.00006](https://doi.org/10.3389/fmars.2019.00006)

1173

1174 Gladstone, Rupert M.; Bigg, Grant R.; Nicholls, Keith W. 2001 Iceberg trajectory modeling and meltwater injection in the
1175 Southern Ocean. *Journal of Geophysical Research*, **106** (C9). 19,903-19,915. <https://doi.org/10.1029/2000JC000347>

1176

1177 Graham, A.G.C., Larter, R.D., Gohl, K., Hillenbrand, C.-D., Smith, J.A., and Kuhn, G. (2009), Bedform signature of a West
1178 Antarctic palaeo-ice stream reveals a multi-temporal record of flow and substrate control, *Quaternary Science Reviews*, **28**,
1179 2774-2793.

1180

1181 Hawking, J. R., Wadham, J. L., Benning, L. G., Hendry, K. R., Tranter, M., Tedstone, A., Nienow, P., & Raiswell, R. (2017).
1182 Ice sheets as a missing source of silica to the polar oceans. *Nature Communications*, **8**, Article 14198.
1183 <https://doi.org/10.1038/ncomms14198>

1184

1185 Hill, J., Condron, A. Subtropical iceberg scours and meltwater routing in the deglacial western North Atlantic. *Nature
1186 Geosci* **7**, 806–810 (2014). <https://doi.org/10.1038/ngeo2267>

1187

1188 Hopwood MJ, Carroll D, Höfer J, Achterberg EP, Meire L, Le Moigne FAC, Bach LT, Eich C, Sutherland DA, González HE.
1189 Highly variable iron content modulates iceberg-ocean fertilisation and potential carbon export. *Nat Commun.* 2019 Nov
1190 20;10(1):5261. doi: [10.1038/s41467-019-13231-0](https://doi.org/10.1038/s41467-019-13231-0).

1191

1192 Huth, A., Adcroft, A., & Sergienko, O. (2022). Parameterizing tabular-iceberg decay in an ocean model. *Journal of Advances
1193 in Modeling Earth Systems*, **14**, e2021MS002869. <https://doi.org/10.1029/2021MS002869>

1194

1195

1196 Jacobs, S., A. Jenkins, H. Hellmer, C. Giulivi, F. Nitsche, B. Huber, and R. Guerrero. 2012. The Amundsen Sea and the
1197 Antarctic Ice Sheet. *Oceanography* **25**(3):154–163, <https://doi.org/10.5670/oceanog.2012.90>.

1198
1199 Jakobsson, M., et al. (2011), Geological record of ice shelf break-up and grounding line retreat, Pine Island Bay, West
1200 Antarctica, *Geology*, 39(7), 691–694.
1201
1202 Jenkins, A., K. W. Nicholls, and H. F. J. Corr, 2010: Observation and Parameterization of Ablation at the Base of Ronne Ice
1203 Shelf, Antarctica. *J. Phys. Oceanogr.*, 40, 2298–2312, <https://doi.org/10.1175/2010JPO4317.1>.
1204
1205 Jourdain, N. C., et al. 2019. Simulating or prescribing the influence of tides on the Amundsen Sea ice shelves, *Ocean
1206 Modelling*, Vol. 133, 44-55, <https://doi.org/10.1016/j.ocemod.2018.11.001>.
1207
1208 Kostov, Yavor; Holland, Paul R.; Hogan, Kelly A.; Smith, James A.; Jourdain, Nicolas C; Mathiot, Pierre; Olivé Abelló, Anna;
1209 Fleming, Andrew H.; Meijers, Andrew J. S.: Modelled dynamics of floating and grounded icebergs, with application to the
1210 Amundsen Sea. Supplementary Movie. Copernicus Publications, 2025a. <https://doi.org/10.5446/70447>
1211 Kostov et al. 2025b. Updates to the NEMO iceberg dynamics and grounding (Version 2). Zenodo.
1212 <https://doi.org/10.5281/zenodo.15484879>
1213 Ian Joughin *et al.*, Ice-shelf retreat drives recent Pine Island Glacier speedup. *Sci. Adv.* 7, eabg3080 (2021).
1214 DOI:[10.1126/sciadv.abg3080](https://doi.org/10.1126/sciadv.abg3080)
1215 Lemieux, J.-F., F. Dupont, P. Blain, F. Roy, G. C. Smith, and G. M. Flato (2016), Improving the simulation of landfast ice by
1216 combining tensile strength and a parameterization for grounded ridges, *J. Geophys. Res. Oceans*, 121, 7354–7368,
1217 doi:10.1002/2016JC012006.
1218 Lien, R., A. Solheim, A. Elverhoi, and K. Rokoengen (1989), Iceberg scouring and sea bed morphology on the eastern Weddell
1219 Sea shelf, Antarctica, *Polar Research*, 7(1), 43–57.
1220
1221 Lopez, R. et al., Hydrodynamic Effects On Iceberg Gouging. *Cold Regions Science and Technology*, 4 (1981) 55-61 55
1222
1223 Lucas, N., Brearley, J., Katherine Hendry et al. Giant icebergs increase mixing and stratification in upper-ocean layers, 10
1224 June 2024, PREPRINT (Version 1) available at Research Square [<https://doi.org/10.21203/rs.3.rs-4425629/v1>]
1225
1226 MacAyeal, D.R. , Scambos, T.A. , Hulbe, C.L. , Fahnestock, M.A. , 2003. Catastrophic ice-shelf break-up by an ice-shelf-
1227 fragment-capsize mechanism. *J. Glaciol.* 49 (164), 22–36.
1228
1229 Madec, G., & NEMO-Team. (2016). Note du Pôle de modélisation de l’Institut Pierre-Simon Laplace No 27. In *NEMO ocean
1230 engine, version 3.6 stable (Technical Report)*. France: IPSL.
1231
1232 Madec, G., & NEMO System Team. (2022). NEMO ocean engine. *Zenodo*. <https://doi.org/10.5281/zenodo.6334656>
1233
1234 Marsh, R.; Ivchenko, V.O.; Skliris, N.; Alderson, S.; Bigg, G.R.; Madec, G.; Blaker, A.T. ; Aksenov, Y. ; Sinha, B.; Coward,
1235 A.C. ; Le Sommer, J.; Merino, N.; Zalesny, V.B.. 2015 NEMO–ICB (v1.0): interactive icebergs in the NEMO ocean model
1236 globally configured at eddy-permitting resolution. *Geoscientific Model Development*, 8 (5). 1547-1562.
1237 <https://doi.org/10.5194/gmd-8-1547-2015>

1238
1239 Marsh, R., Bigg, G., Zhao, Y. et al. Prospects for seasonal forecasting of iceberg distributions in the North Atlantic. *Nat*
1240 *Hazards* 91, 447–471 (2018). <https://doi.org/10.1007/s11069-017-3136-4>

1241
1242 Marson, J. M., Myers, P. G., Garbo, A., Copland, L., & Mueller, D. (2024). Sea ice-driven iceberg drift in Baffin Bay.
1243 *Journal of Geophysical Research: Oceans*, 129, e2023JC020697. <https://doi.org/10.1029/2023JC020697>

1244
1245 Martin, T. and Adcroft, A. (2010) Parameterizing the fresh-water flux from land ice to ocean with interactive icebergs in a
1246 coupled climate model, *Ocean Modelling*, Vol. 34, Issues 3–4, 111-124, <https://doi.org/10.1016/j.ocemod.2010.05.001>

1247
1248 Mazur, A. K., Wahlin, A. K., & Kalén, O. (2019). The life cycle of small-to medium-sized icebergs in the Amundsen Sea
1249 Embayment. *Polar Research*, 38.

1250
1251 Merino, N., Julien Le Sommer, Gael Durand, Nicolas C. Jourdain, Gurvan Madec, Pierre Mathiot, Jean Tournadre, Antarctic
1252 icebergs melt over the Southern Ocean: Climatology and impact on sea ice, *Ocean Modelling*, Volume 104, 2016, Pages 99–
1253 110, <https://doi.org/10.1016/j.ocemod.2016.05.001>.

1254
1255 Miles BWJ, Stokes CR, Jenkins A, Jordan JR, Jamieson SSR, Gudmundsson GH. Intermittent structural weakening and
1256 acceleration of the Thwaites Glacier Tongue between 2000 and 2018. *Journal of Glaciology*. 2020;66(257):485-495.
1257 doi:10.1017/jog.2020.20

1258
1259 Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., et al. (2020). Deep glacial troughs and
1260 stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet. *Nature Geoscience*, 13(2), 132–
1261 137. <https://doi.org/10.1038/s41561-019-0510-8>

1262
1263 Mouginot, J., E. Rignot, and B. Scheuchl (2014), Sustained increase in ice discharge from the Amundsen Sea Embayment,
1264 West Antarctica, from 1973 to 2013, *Geophys. Res. Lett.*, 41, 1576–1584, doi:[10.1002/2013GL059069](https://doi.org/10.1002/2013GL059069).

1265
1266 Mountain, D.G. (1980) On predicting iceberg drift, *Cold Regions Science and Technology*, Vol. 1, 3–4, 1980, pp. 273-282,
1267 [https://doi.org/10.1016/0165-232X\(80\)90055-5](https://doi.org/10.1016/0165-232X(80)90055-5).

1268
1269 Naughten, K. A., Holland, P. R., Dutrieux, P., Kimura, S., Bett, D. T., & Jenkins, A. (2022). Simulated twentieth-century
1270 ocean warming in the Amundsen Sea, West Antarctica. *Geophysical Research Letters*, 49,
1271 e2021GL094566. <https://doi.org/10.1029/2021GL094566>

1272
1273 Nakayama, Y. , Timmermann, R. , Schröder, M. and Hellmer, H. (2014): On the difficulty of modeling Circumpolar Deep
1274 Water intrusions onto the Amundsen Sea continental shelf , *Ocean Modelling*, 84 , pp. 26-34 . doi:
1275 10.1016/j.ocemod.2014.09.007

1276
1277 Olivé Abelló, A., Mathiot, P., Jourdain, N. C., Kostov, Y., Holland, P. R., Gascoin, S., & Rousset, C. (2025). Iceberg
1278 grounding enhances the release of freshwater on the Antarctic continental shelf. *Journal of Geophysical Research: Oceans*, 130, e2025JC022857. <https://doi.org/10.1029/2025JC022857>

1280 Person, R., Aumont, O., Madec, G., Vancoppenolle, M., Bopp, L., and Merino, N.: Sensitivity of ocean biogeochemistry to
1281 the iron supply from the Antarctic Ice Sheet explored with a biogeochemical model, *Biogeosciences*, 16, 3583–3603,
1282 <https://doi.org/10.5194/bg-16-3583-2019>, 2019.

1283

1284 Reinardy, B. T.I., C. J. Pudsey, C.-D. Hillenbrand, T. Murray, J. Evans (2009) Contrasting sources for glacial and interglacial
1285 shelf sediments used to interpret changing ice flow directions in the Larsen Basin, Northern Antarctic Peninsula, *Marine
1286 Geology*, Vol. 266, 1–4, pp 156-171, <https://doi.org/10.1016/j.margeo.2009.08.003>.

1287

1288 Ruffman, A. (2005) The First Known Attempts Off Eastern Canada 'to Tow, or Cant' Icebergs: a seed to the understanding of
1289 iceberg scour processes at the seafloor. CMOS Bulletin SCMO. Vol. 33, 99-123 <http://cmosarchives.ca/Bulletin/b3304.pdf>

1290

1291 Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., et al. (2019). Trends in Antarctic Ice Sheet
elevation and mass. *Geophysical Research Letters*, 46, 8174–8183. <https://doi.org/10.1029/2019GL082182>

1292

1293 Smith, S.D., E. G. Banke (1983) The influence of winds, currents and towing forces on the drift of icebergs, *Cold Regions
Science and Technology*, Vol. 6, 3, pp. 241-255, [https://doi.org/10.1016/0165-232X\(83\)90045-9](https://doi.org/10.1016/0165-232X(83)90045-9).

1294

1295 Smith, J.A., Claus-Dieter Hillenbrand, Gerhard Kuhn, Robert D. Larter, Alastair G.C. Graham, Werner Ehrmann, Steven G.
1296 Moreton, Matthias Forwick, Deglacial history of the West Antarctic Ice Sheet in the western Amundsen Sea Embayment,
Quaternary Science Reviews, Volume 30, Issues 5-6, 2011, Pages 488-505, <https://doi.org/10.1016/j.quascirev.2010.11.020>

1297

1298 Starr, A., Hall, I.R., Barker, S. et al. Antarctic icebergs reorganize ocean circulation during Pleistocene glacials. *Nature* **589**,
236–241 (2021). <https://doi.org/10.1038/s41586-020-03094-7>

1299

1300 Stern, A. A., et al. (2015), Wind-driven upwelling around grounded tabular icebergs, *J. Geophys. Res. Oceans*, 120, 5820–
5835, doi:[10.1002/2015JC010805](https://doi.org/10.1002/2015JC010805).

1301

1302 Stern, A. A., A. Adcroft, and O. Sergienko (2016), The effects of Antarctic iceberg calving-size distribution in a global climate
model, *J. Geophys. Res. Oceans*, 121, 5773–5788, doi:[10.1002/2016JC011835](https://doi.org/10.1002/2016JC011835).

1303

1304 Stern, A. A., A. Adcroft, O. Sergienko, and G. Marques (2017), Modeling tabular icebergs submerged in the ocean, *J. Adv.
Model. Earth Syst.*, 9, 1948–1972, doi:[10.1002/2017MS001002](https://doi.org/10.1002/2017MS001002).

1305

1306 St-Laurent, P., J. M. Klinck, and M. S. Dinniman (2015), Impact of local winter cooling on the melt of Pine Island Glacier,
1307 Antarctica, *J. Geophys. Res. Oceans*, 120, 6718–6732, doi:[10.1002/2015JC010709](https://doi.org/10.1002/2015JC010709)

1308

1309 St-Laurent, P., Yager, P. L., Sherrell, R. M., Oliver, H., Dinniman, M. S., & Stammerjohn, S. E. (2019). Modeling the
1310 seasonal cycle of iron and carbon fluxes in the Amundsen Sea Polynya, Antarctica. *Journal Geophysical Research:
1311 Oceans*, 124, 1544–1565. <https://doi.org/10.1029/2018JC014773>

1312

1313 St-Laurent, P., Stammerjohn, S. E., & Maksym, T. (2024). Response of onshore oceanic heat supply to yearly changes in the
1314 Amundsen Sea icescape (Antarctica). *Journal of Geophysical Research: Oceans*, 129,
1315 e2023JC020467. <https://doi.org/10.1029/2023JC020467>

1316

1317 Tarling, Geraint A. ; Thorpe, Sally E. ; Henley, Sian F.; Burson, Amanda ; Liszka, Cecilia M. ; Manno, Clara ; Lucas, Natasha
1318 S. ; Ward, Freyja; Hendry, Katharine R. ; Woodward, E. Malcolm S.; Wootton, Marianne; Abrahamsen, E. Povl. 2024 Collapse
1319 of a giant iceberg in a dynamic Southern Ocean marine ecosystem: in situ observations of A-68A at South Georgia. *Progress*
1320 *in Oceanography*, 226, 103297. 17, pp. <https://doi.org/10.1016/j.pocean.2024.103297>

1321

1322 Tsujino, H., et al. (2018) JRA-55 based surface dataset for driving ocean–sea-ice models (JRA55-do), *Ocean Modelling*, Vol.
1323 130, 79-139, <https://doi.org/10.1016/j.ocemod.2018.07.002>.

1324

1325 Veldhuijsen, S. B. M., van de Berg, W. J., Brils, M., Kuipers Munneke, P., and van den Broeke, M. R.: Characteristics of the
1326 1979–2020 Antarctic firn layer simulated with IMAU-FDM v1.2A, *The Cryosphere*, 17, 1675–1696,
1327 <https://doi.org/10.5194/tc-17-1675-2023>, 2023.

1328

1329 Wagner, T. J. W., R. W. Dell, and I. Eisenman, 2017: An Analytical Model of Iceberg Drift. *J. Phys. Oceanogr.*, 47, 1605–
1330 1616, <https://doi.org/10.1175/JPO-D-16-0262.1>.

1331

1332 Wagner, T. J. W., I. Eisenman, A. M. Ceroli, and N. C. Constantinou, 2022: How Winds and Ocean Currents Influence the
1333 Drift of Floating Objects. *J. Phys. Oceanogr.*, 52, 907–916, <https://doi.org/10.1175/JPO-D-20-0275.1>.

1334

1335 Weeks, W.F. , Mellor, M. , 1978. Some Elements of Iceberg Technology. Technical report. Cold Regions Research and
1336 Engineering Laboratory . March

1337

1338 Wise, M.G., Dowdeswell, J.A., Jakobsson, M.J., and Larter, R.D. (2017), Evidence of marine ice-cliff instability in Pine Island
1339 Bay from iceberg-keel plough marks, *Nature*, 550(7677), 506-510.

1340