



New quantitative measurements and spectroscopic line parameters of ammonia in the 685-1250 cm⁻¹ spectral region for atmospheric remote sensing

Daniel J. L. Coxon^{1,2}, Jeremy. J. Harrison^{1,2}, D. Chris Benner³, V. Malathy Devi³

¹School of Physics and Astronomy, Space Park Leicester, University of Leicester, Leicester, LE4 5SP, United Kingdom

²National Centre for Earth Observation, UK

³Department of Physics, The College of William and Mary, Williamsburg, 23187-8795, United States

Correspondence to: Daniel J. L. Coxon (djlcl1@leicester.ac.uk), Jeremy J. Harrison (jh592@leicester.ac.uk)

Abstract.

Ammonia (NH₃) is a toxic pollutant, generally linked to agricultural emissions, and plays a major role in the formation of fine aerosols which have a significant and detrimental effect on human health. NH₃ is one of the key pollutants that can be monitored by satellite instruments orbiting the Earth, including the Infrared Atmospheric Sounding Interferometer (IASI) and the Cross-track Infrared Sounder (CrIS). The interpretation of these measured atmospheric spectra requires accurate radiative transfer modelling, which relies on the quality of the input spectroscopic line parameters.

In this work we present new high quality high-resolution infrared spectra of self- and air-broadened NH₃ at 296 K using a Bruker IFS 125HR spectrometer and a 24.45 cm pathlength sample cell with silver chloride windows. Using a multispectrum fitting approach, we then determine new spectroscopic line parameters over the range 685 cm⁻¹ to 1250 cm⁻¹ for the NH₃ 0100 00 0 s ← 0000 00 0 a and 0100 00 0 s ← 0000 00 0 s transitions associated with the ν_2 mode; the Q branches of these transitions are the strongest NH₃ features observed in atmospheric spectra. Our analysis utilises the Voigt lineshape, with speed-dependent Voigt and Rosenkranz line mixing for the strongest lines. To date this is the most complete experimental and multispectrum analysis of air-broadened NH₃ over this spectral region. Our derived spectroscopic line parameters reproduce the new measurements substantially better than line parameters from the HITRAN 2020 database, which were derived from a mixture of *ab initio* calculations and previous laboratory measurements. We have revised values for parameters such as line intensities and air-broadened Lorentz halfwidths, in some cases by almost 10%. We have substantially lowered the uncertainties of key parameters, such as line intensities. In addition to the measured speed dependence and Rosenkranz line mixing parameters, which we believe are the first reported for the ν_2 band of NH₃ in air, we also determine a range of parameters for the ν_2 band that are not currently in HITRAN, for example self- and air-pressure-induced shifts. We expect these new parameters to provide a more accurate basis for incorporation into atmospheric radiative transfer models to measure NH₃ concentrations from satellite.



1 Introduction

Ammonia (NH_3) is found throughout the universe, from the interstellar medium (Ho et al., 1983) to the atmospheres of exoplanets, e.g. GJ 504 b (Mâlin et al., 2025). Closer to home, it can be found throughout the Solar System, e.g. in the atmospheres of the gas giants Jupiter and Saturn (Irwin et al., 2025), and also in the Earth's atmosphere where it is the most abundant alkaline gas. On Earth, it is emitted through a range of anthropogenic sources (Sutton et al., 2008; Behara et al., 2013; Wyer et al., 2022) of which agriculture, mostly accounted for by livestock and nitrogen-based fertilisers, is responsible for over 80% of the global total emissions (van Damme et al., 2021). Atmospheric NH_3 is responsible for a variety of negative environmental and health effects. It contributes to the eutrophication and acidification of aquatic ecosystems, causing adverse effects on a variety of different flora (Sutton et al., 2009; García-Gómez et al., 2014). Additionally, NH_3 serves as a significant precursor to fine particulate matter, $\text{PM}_{2.5}$ (Brunekreef et al., 2015; Megaritis et al., 2013; Thakrar et al. 2020). It is estimated that atmospheric NH_3 accounts for 30% and 50% of the total production of $\text{PM}_{2.5}$ in the US and Europe, respectively (Erisman et al., 2004; Behara et al., 2010; Bauer et al., 2016; Han et al., 2020). $\text{PM}_{2.5}$ contributes to visibility degradation (Ting et al., 2022; Yang et al., 2022) and, when inhaled, can cause a number of diseases, including chronic obstructive pulmonary disorder (COPD) and lung cancer (Apte et al., 2018; Lelieveld et al., 2015; Yu et al., 2000).

Atmospheric NH_3 can be monitored near the Earth's surface using a variety of in situ methods such as wet chemistry or spectroscopic techniques (Twigg et al., 2022). Increasingly over the last few decades, it has been possible to monitor NH_3 by infrared spectrometers onboard satellites, such as the Infrared Atmospheric Sounding Interferometer (IASI) (Clarisse et al., 2009; van Damme et al., 2014; van Damme et al., 2018; Clarisse et al., 2019), the Cross-track Infrared Sounder (CrIS) (Shephard et al., 2015), the Atmospheric InfraRed Sounder (AIRS) (Warner et al., 2016; Warner et al., 2017), and the Tropospheric Emission Spectrometer (TES) (Shephard et al., 2011). Among the most prominent NH_3 features in the infrared used for remote sensing are the two Q branches of the ν_2 ammonia band at ~ 10.4 and $10.7 \mu\text{m}$. Accurate retrievals of NH_3 concentrations from the measurements recorded by the various instruments mentioned above depend upon our ability to model atmospheric radiative transfer, which in turn depends on the underlying NH_3 spectroscopic line parameters. It is therefore crucially important that the line parameters for the ν_2 NH_3 band are both accurately and precisely defined.

In 2013, a full re-analysis of $^{14}\text{NH}_3$ spectroscopy was conducted (Down et al., 2013). In particular, line positions and intensities for the ν_2 band were revised. This work led to the most recent significant spectroscopic update for NH_3 in the HITRAN 2012 (Rothman et al., 2013) and GEISA 2015 (Jacquinet-Husson et al., 2016) databases. A study was also conducted to determine self- and air-broadened Lorentz halfwidths at half-maximum (HWHM) for the ν_2 $^{14}\text{NH}_3$ band for $J \leq 9$ (Nemtchinov et al., 2004); these results were included in the HITRAN 2012 database. In the case of $^{14}\text{NH}_3$ lines where $J > 9$, the self-broadened halfwidths at each K level are given arbitrary values of 0.5, and the air-broadened halfwidths at each K level ($K \leq 9$) are assigned the corresponding values for the $J = 9$ levels, except for $K > 9$ levels for which arbitrary values are assigned. High



65 resolution spectra of $^{15}\text{NH}_3$ have been recorded (Canè et al., 2019; Canè et al., 2020), and line positions derived from these studies being included in the HITRAN 2020 database (Gordon et al., 2021) with line intensities derived through *ab initio* calculations (Yurchenko et al., 2015). Currently, no lines of other minor isotopologues of NH_3 , such as NH_2D , are included in the HITRAN database.

70 In this work, we label vibrational states by $\nu_1\nu_2\nu_3\nu_4l_3l_4li$, where ν_n represents the quantum number of the n th vibrational mode. Unlike the non-degenerate modes ν_1 and ν_2 , ν_3 and ν_4 are doubly degenerate and give rise to the vibrational angular momentum labels l_3 and l_4 , respectively, which can take the values $-\nu$, $-\nu + 2$, ..., $\nu - 2$, ν . These two angular momenta couple to produce the total vibrational angular momentum, l . The ν_2 mode is associated with the inversional umbrella motion between the two equivalent equilibrium structures, and is considered a classic textbook example of quantum mechanical
75 tunnelling. As a result of this large amplitude vibration, each energy level of ammonia splits into a symmetric and antisymmetric level. For this reason, vibrational states additionally require an inversion label, i , denoted by s or a depending on whether the state is symmetric or antisymmetric on inversion through the planar configuration. The rotational levels are characterised by J , the total angular momentum quantum number, and K , its component along the molecular axis, which can take the values $0, 1, 2, \dots, J$.

80 In this work, we derive a set of new line parameters for the ν_2 NH_3 band, including positions, intensities, self- and air-broadened halfwidths and, for the first time, a complete set of self- and air-pressure-induced shifts, and, where possible, speed dependence and Rosenkranz line mixing parameters. Although this work has focused primarily on deriving new line parameters for the $0100\ 00\ 0\ s \leftarrow 0000\ 00\ 0\ a$ and $0100\ 00\ 0\ a \leftarrow 0000\ 00\ 0\ s$ transitions (excitation from the ground state to
85 $\nu_2 = 1$) of $^{14}\text{NH}_3$, which are of particular importance for remote sensing, we have also derived line parameters within the $685\text{--}1250\ \text{cm}^{-1}$ spectral window for the same transitions of $^{15}\text{NH}_3$, alongside a variety of other less intense $^{14}\text{NH}_3$ lines in the ν_4 , $2\nu_2$ (first overtone), and $\nu_2 + \nu_4$ (combination) bands. The line parameters for these weaker bands are more fully detailed in the Supplementary Information.

90 2 Experimental details

A new laboratory for the purposes of producing spectroscopic data for remote sensing of trace gases in the Earth's atmosphere has recently been set up at Space Park Leicester. Dubbed the SPectroscopy for ENvironmental SENSing Research (SPENSER) facility, the principal instrument is a Bruker IFS 125HR spectrometer. In this work, we carried out measurements of pure and air-broadened NH_3 samples at 296 K with a focus on the spectral window $685\text{--}1250\ \text{cm}^{-1}$. A glass sample cell, situated inside
95 the sample compartment of the spectrometer, was configured with high IR transmission AgCl windows. Its pathlength was determined as 24.45 cm using high resolution measurements of the $0002 \leftarrow 0000$ band of N_2O and high accuracy line intensities



from a previous study at the Physikalisch-Technische Bundesanstalt (Werwein et al., 2017). The pressure inside the absorption cell and gas line was monitored by four calibrated Inficon SKY CDG-45D manometers, with different full-scale pressures ranging from 0.1 to 1000 Torr. The temperature of the sample was maintained at a constant value of 296 K using a F32-HE Julabo circulator that flowed water around the outer jacket of the cell; the temperature was measured by four calibrated PRTs connected to an Isotech milliK precision thermometer and millisKanner channel expander. A schematic of the experimental setup is shown in Fig. 1. The spectrometer settings and additional measurement information are listed in Table 1.




Figure 1. Schematic representation of the experimental setup.

Sample gas mixtures were made from high purity gases, namely NH_3 (BOC; 99.98 % purity) and dry synthetic air (primary reference material (PRM) from the National Physical Laboratory (NPL); 0.95 % Ar, 20.95 % O_2 in N_2 , relative uncertainties Ar 1.0 %, O_2 0.2 %). For the pure NH_3 measurements, a spectral resolution of 0.003 cm^{-1} was used, defined as the Bruker instrument resolution of 0.9/MOPD; MOPD = maximum optical path difference. For the air-broadened measurements, the spectral resolution was reduced according to the linewidth, with a lowest resolution of 0.02 cm^{-1} used where the total pressure approached 1000 hPa. Table 2 lists the pressures and resolutions selected for the measurements, which were optimised for the Earth's atmospheric conditions. For the analysis, NH_3 spectra were ratioed against appropriate empty cell spectra (taken at 0.03 cm^{-1} spectral resolution) to produce transmittance spectra. These spectra were unapodised except for the highest resolution measurements ($< 0.005 \text{ cm}^{-1}$, equivalent to MOPD $> 180 \text{ cm}$) for which the following apodisation function (a_p) was applied (Devi et al., 2003):



$$\left[1 - \left(\frac{p}{\text{MOPD}}\right)^2\right]^2, \quad (1)$$

where p is the path difference, and a_p is defined as zero when $p > \text{MOPD}$. Finally, additional N_2O spectra were recorded for the purposes of calibrating the wavenumber scales of the NH_3 spectra. This is discussed in more detail in Section 3.

120

Table 1. Spectrometer settings and measurement information for the pure and air-broadened NH_3 spectra.

Spectrometer	Bruker IFS 125HR
Light source	MIR globar
Beam splitter	KBr
Detector	MCT D316
Focal length of collimating lens	418 mm
Aperture diameter	1.7 mm
MOPD	45-300 cm
Spectral resolution	0.02-0.003 cm^{-1}
Apodisation	Eq. 1 for $\text{MOPD} > 180$ cm
Spectral window	685-1250 cm^{-1}
Temperature	296 K
Cell path length	0.2445 m
Cell windows	AgCl
Signal-to-noise (SNR)	750-2500
NH_3	BOC, 99.98 % purity
Dry synthetic air	NPL PRM, 0.95 % Ar, 20.95 % O_2 in N_2 , relative uncertainties Ar 1.0 %, O_2 0.2 %
Wavenumber calibration	N_2O

Table 2. Experimental conditions for the measured NH_3 spectra.

NH_3 amount fraction	Total pressure (hPa)	Spectral resolution (cm^{-1})
1	0.0171	0.0030
1	0.121	0.0030
1	0.320	0.0030
1	0.579	0.0030
1	0.995	0.0030



0.003740	9.94	0.0030
0.002160	19.91	0.0040
0.002450	29.88	0.0045
0.001810	50.21	0.0060
0.001755	73.63	0.0080
0.002135	99.54	0.0100
0.001485	199.2	0.0150
0.001355	302.7	0.0150
0.001350	498.8	0.0200
0.001260	749.5	0.0200
0.001320	978.3	0.0200

125

3 Analysis and data retrievals

The pure and air-broadened ammonia spectra taken in this study were analysed using a nonlinear least-squares multispectrum fitting software package known as Labfit (Benner et al., 1995), which has an extensive history in deriving non-Voigt line parameters for remote sensing, e.g. (Benner et al. 2016). In this work, line parameters were determined for both Voigt and speed-dependent Voigt lineshapes with Rosenkranz first order line mixing (Rosenkranz et al., 1988). The normalised Voigt lineshape with Rosenkranz line mixing is defined in Labfit (Benner et al., 1995; Letchworth et al., 2007) as:

$$I = \sqrt{\frac{\ln(2)}{\pi}} \frac{1}{\Gamma_D} [K(x, y) + YL(x, y)], \quad (2)$$

where Y is the Rosenkranz line mixing parameter,

$$K(x, y) = \frac{y}{\pi} \int_{-\infty}^{\infty} \frac{\exp(-t^2)}{(x-t)^2 + y^2} dt, \quad (3)$$

and

$$L(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{(x-t)\exp(-t^2)}{(x-t)^2 + y^2} dt. \quad (4)$$

These are simply related to the complex probability function, W , by

$$W(z) = K(x, y) + iL(x, y) = \frac{i}{\pi} \int_{-\infty}^{\infty} \frac{\exp(-t^2)}{x+iy-t} dt, \quad (5)$$

140 where x and y are given by



$$x = \sqrt{\ln(2)}(v - v_0 - \Delta)/\Gamma_D, \quad (6)$$

and

$$y = \sqrt{\ln(2)}\Gamma_0/\Gamma_D. \quad (7)$$

In the above equations, $v - v_0$ is the wavenumber detuning relative to the unperturbed line centre v_0 , Γ_0 is the Lorentz (collisional) halfwidth at half maximum, Γ_D the Doppler halfwidth at half maximum, and Δ the pressure-induced shift of the line centre (so that $v - v_0 - \Delta$ represents the wavenumber detuning relative to the pressure-shifted line centre, $v_0 + \Delta$).

When the speed dependence of collisions is accounted for, the Voigt lineshape is modified into the speed-dependent Voigt profile. With the inclusion of Rosenkranz line mixing, the mathematical form of this profile is defined in Labfit (Benner et al., 1995; Ciurylo et al., 1998) by rewriting the Lorentz halfwidth as a function of v , defined as the ratio of the speed of an absorbing molecule to its most probable speed, and integrating over the full range of velocities. This leads to K and L being redefined as:

$$K(x, y, a_w) = \frac{2}{\pi} \int_{-\infty}^{\infty} v e^{-v^2} \arctan \left[\frac{x+v}{y+a_w y(v^2-3/2)} \right] dv, \quad (8)$$

and

$$L(x, y, a_w) = \frac{1}{\pi} \int_{-\infty}^{\infty} v e^{-v^2} \ln \left\{ 1 + \left[\frac{x+v}{y+a_w y(v^2-3/2)} \right]^2 \right\} dv, \quad (9)$$

where

$$a_w = \frac{\Gamma_2}{\Gamma_0}, \quad (10)$$

and Γ_2 is the (quadratic) speed-dependence parameter for the Lorentz halfwidth. In this work, we do not consider the speed-dependence of the pressure-induced shifts. These shifts are expected to be small, and difficult to separate from the effects of line mixing.

The parameters Γ_0 , Δ , and Y , introduced in Eqs. 2-10, are pressure dependent and can be expanded as

$$\Gamma_0 = p[\gamma_{air}^0(1-x) + \gamma_{self}^0 x], \quad (11)$$

$$\Delta = p[\delta_{air}^0(1-x) + \delta_{self}^0 x], \quad (12)$$

$$Y = pY_{air}^0. \quad (13)$$

In the above expressions, γ^0 , δ^0 , and Y^0 are defined at $p_{ref} = 1$ atm, p is the pressure in atmospheres, and x is the amount fraction of NH_3 in the sample. As all measurements in this work were taken at 296 K, we have not considered temperature dependence terms in Eqs. 2-13. Also note that for each line in the spectrum, the current version of Labfit can only fit one speed dependence parameter (a_w) and one Rosenkranz line-mixing parameter (Y^0) in a solution, making no distinction between self



and foreign contributions; however, the NH_3 amount fractions in the high-pressure measurements are very small (less than
 170 ~ 0.004) so that the self contributions to the air-broadened spectra are smaller than the experimental uncertainty. The definitions
 of the retrieved line parameters are summarised in Table 3.

In addition to line parameters, Labfit is able to correct during the analysis for minor imperfections in the spectra associated
 with the field of view or residual phase errors. The software can also fit spectral baselines using Chebyshev polynomials, and
 175 the zero level of individual spectra, and all spectra can be weighted in accordance with their individual signal-to-noise ratios
 (SNRs).

Table 3. Line parameters derived in this study.

Symbol	Description	Units
ν_0	Line centre position at zero pressure	cm^{-1}
S	Line intensity at 296 K and 1 atm	$\text{cm}^{-1}/(\text{molecule cm}^{-2})$
γ_{self}^0	Self-broadened Lorentz halfwidth at 296 K and 1 atm	$\text{cm}^{-1}/\text{atm}$
γ_{air}^0	Air-broadened Lorentz halfwidth at 296 K and 1 atm	$\text{cm}^{-1}/\text{atm}$
δ_{self}^0	Self-pressure-induced line shift at 296 K and 1 atm	$\text{cm}^{-1}/\text{atm}$
δ_{air}^0	Air-pressure-induced line shift at 296 K and 1 atm	$\text{cm}^{-1}/\text{atm}$
$a_w(\text{air})$	Ratio of the speed dependence of the air-broadened Lorentz halfwidth to the air-broadened Lorentz halfwidth at 296 K	Unitless
Y_{air}^0	First-order (Rosenkranz) line mixing parameter in air at 296 K and 1 atm	$1/\text{atm}$

180 Using Labfit (Benner et al., 1995) it is possible to relate line positions via the corresponding upper and lower state rovibrational
 energies, and parameterise them in terms of the band centre and the rotational constants B , D , H , etc. By constraining line
 positions to these quantum mechanical expressions, fewer parameters are floated and correlations between parameters are
 reduced. This method has the advantage that blended lines can easily be separated using appropriate constraints. In Labfit,
 however, these position constraints are optimised for linear molecules such as carbon dioxide. Fitting spectra of ammonia, a
 185 symmetric rotor whose energy levels require an additional quantum number, K , adds an additional level of complexity to the
 problem that is too cumbersome without substantial modifications to the programming code. Therefore, for this work all line
 parameters, including positions, were solved on a line-by-line basis. This does present challenges, for example, when fitting
 two lines of similar intensity that are blended even in the lowest pressure spectra. In such cases, some parameters of one or
 both of the lines need to be fixed, as outlined in Section 4. In all cases, preference is given to fitting lines from the ν_2 band.



190 The multispectrum fitting technique requires that all spectra are consistently calibrated prior to the spectral fitting. Lines
present in the measured spectra originating from water absorption inside the spectrometer (not the cell) were used to provide
a relative wavenumber calibration between spectra. The absolute calibration of the wavenumber scales of these water lines
was determined by measuring a low-pressure spectrum of N₂O at room temperature and calibrating against NIST heterodyne
calibration tables (Maki et al., 1991).


195

The NH₃ multispectrum fit was initialized using parameters from HITRAN 2020 (Gordon et al., 2021), and proceeded by
fitting all pure sample spectra before adding the air-broadened spectra to the solution and floating selected parameters for all
spectra simultaneously. Initial fits were carried out over the entire 685-1250 cm⁻¹ region, however there was evidence from
the saturated lines in the pure sample spectra that the zero level was not constant across this wide fitted interval; the current
200 version of Labfit can only apply a single zero level offset correction to each spectrum. To counter this situation as much as
possible, the spectra were split into three regions (685-900 cm⁻¹, 885-1015 cm⁻¹ and 1000-1250 cm⁻¹), allowing three different
offsets to be applied to each spectrum as required.

The approach taken here was to fit all absorption features within the fitted interval, and adjust positions, intensities, Lorentz
205 halfwidths, pressure-induced shifts, Rosenkranz line mixing parameters, and quadratic speed dependence for each line
separately. Line mixing and speed dependence parameters were included in the fit as judged by inspection of the fit residuals
and the derived uncertainties determined by the fit.

4 Discussion of results and comparison with HITRAN2020

210 The final multispectrum fit for all 16 NH₃ spectra (self- and air-broadened) over the entire 685-1250 cm⁻¹ range is plotted in
Fig. 2. The top panel shows the superimposed weighted spectral fit residuals (observed minus calculated), with the bottom
panel showing the observed spectra. The vast majority of the residuals, including the entire Q branch region, are within ±0.5%.
The only exceptions are a small number of saturated features in the R branch region of the pure NH₃ spectra, which are slightly
greater. These small but persistent residuals are predominantly associated with the lowest pressure measurements, and are
215 likely caused by small variations in the zero level for these spectra that are not adequately captured by Labfit (Benner et al.,
1995).



220 **Figure 2: Overlay of all the transmittance spectra across the entire measured spectral region, 685-1250 cm^{-1} , alongside their weighted fit residuals, defined as the difference between the observed and calculated spectra.**

The principal focus of this work is the ν_2 band, however there are a number of other weaker bands present in the spectral region of interest. Table 4 lists the vibrational assignments of these bands along with the rotational lines and isotopologues that were included in the fit. In total, parameters were fit for 1374 lines with intensities above a threshold of $1 \times 10^{-23} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$. For the weakest of these lines, only positions and intensities were determined; for the strongest, the whole suite of parameters listed in Table 3 were determined. In general, self- and air-broadened Lorentz halfwidths were fit for lines with intensities above $4 \times 10^{-22} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ and self- and air-pressure-induced shifts for lines with intensities above $1.5 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$. Selected lines of intensity $> 7.5 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ were fit for one or both of speed dependence and Rosenkranz line mixing parameters. The total number of each type of line parameter determined for each of the bands listed in Table 4 is given in Table 5.



235 **Table 4. Main NH₃ bands analysed in this study within the 685-1250 cm⁻¹ spectral region. Some bands were only partially observed in the spectral window, for which only R branch lines were fit. The number ranges in columns 5 to 7 denote the range of analysed J'' values for the observed transitions. However, a number of lines associated with K sub-levels are not included in the fit if the relevant lines are either outside the measured spectral window or have an intensity below the fitting threshold of 1 x 10⁻²³ cm⁻¹ / (molecule cm⁻²) at 296 K.**

Upper vib		Lower vib		P branch	Q branch	R branch	Isotopologue
Full notation	Short notation	Full notation	Short notation	range in J''	range in J''	range in J''	
0100 00 0 s	$\nu_2(s)$	0000 00 0 a	0(a)	2-13	1-19	0-16	¹⁴ N ¹ H ₃
0100 00 0 a	$\nu_2(a)$	0000 00 0 s	0(s)	1-14	1-19	1-14	¹⁴ N ¹ H ₃
0200 00 0 s	$2\nu_2(s)$	0100 00 0 a	$\nu_2(a)$	0	0	2-12	¹⁴ N ¹ H ₃
0200 00 0 a	$2\nu_2(a)$	0100 00 0 s	$\nu_2(s)$	1-11	1-13	1-11	¹⁴ N ¹ H ₃
0100 00 0 s	$\nu_2(s)$	0000 00 0 a	0(a)	2-10	1-13	0-11	¹⁵ N ¹ H ₃
0100 00 0 a	$\nu_2(a)$	0000 00 0 s	0(s)	1-10	1-12	1-11	¹⁵ N ¹ H ₃
0001 01 1 a	$\nu_4(a)$	0100 00 0 a	$\nu_2(a)$	0	6-8	1-11	¹⁴ N ¹ H ₃
0101 01 1 s	$\nu_2 + \nu_4(s)$	0001 01 1 a	$\nu_4(a)$	2-5	1-9	1-6	¹⁴ N ¹ H ₃
0101 01 1 a	$\nu_2 + \nu_4(a)$	0001 01 1 s	$\nu_4(s)$	2-5	1-9	1-6	¹⁴ N ¹ H ₃

240

Table 5. The number of each type of fitted line parameter for the main NH₃ bands analysed in this study.

	Number of floated parameters									Total # of lines
	$\nu_2(s) \leftarrow 0(a)$ (¹⁴ NH ₃)	$\nu_2(a) \leftarrow 0(s)$ (¹⁴ NH ₃)	$2\nu_2(s) \leftarrow \nu_2(a)$	$2\nu_2(a) \leftarrow \nu_2(s)$	$\nu_2(s) \leftarrow 0(a)$ (¹⁵ NH ₃)	$\nu_2(a) \leftarrow 0(s)$ (¹⁵ NH ₃)	$\nu_4(a) \leftarrow \nu_2(a)$	$\nu_2 + \nu_4(s) \leftarrow \nu_4(a)$	$\nu_2 + \nu_4(a) \leftarrow \nu_4(s)$	
ν_0	313	319	63	174	145	141	82	58	53	1374
S	313	315	63	174	145	141	82	58	53	1370
γ_{air}^0	219	220	30	82	0	4	5	0	2	569
γ_{self}^0	225	223	30	82	0	4	5	3	2	578
δ_{air}^0	189	142	3	13	0	0	0	0	0	350
δ_{self}^0	132	135	5	13	0	0	0	0	0	289
γ^0	46	42	0	0	0	0	0	0	0	89
a_w	41	27	0	0	0	0	0	0	0	68

245 Tables 6 and 7 list the newly derived line parameters for lines with an intensity $> 1 \times 10^{-21}$ cm⁻¹ / (molecule cm⁻²) at 296 K for the 0100 00 0 s $\leftarrow$ 0000 00 0 a and the 0100 00 0 a $\leftarrow$ 0000 00 0 s transitions. This intensity cut-off was selected because in all but two cases, it was the minimum required to enable floating of the pressure-induced line shift parameters. Throughout this work we will often refer to these transitions using the shorthand notation $\nu_2(s) \leftarrow 0(a)$ and $\nu_2(a) \leftarrow 0(s)$, respectively. Tables



in the Supplementary Information include line parameters for the lines with lower intensity, line parameters for the $^{15}\text{NH}_3$ isotopologue, and the other transitions listed in Table 4.

Figure 3 shows the spectral region covering the $\nu_2(a) \leftarrow 0(s)$ Q branch, with spectral residuals that result from our fit (panel a) compared to those obtained using the values from HITRAN 2020 (panel b). A significant improvement over the fit residuals in 3(a) are obtained in 3(b) as a result of using the line parameters determined from our multispectrum fit.




Figure 3. Overlay of all the transmittance spectra across the $\nu_2(a) \leftarrow 0(s)$ Q branch, 955-970 cm^{-1} , alongside their weighted residuals using (a) the new line parameters determined in the present study, and (b) the line parameters from the HITRAN 2020 database.

Table 6. Measured parameters at 296 K for lines (labelled ${}^{AK}AJ(J'', K'')_{J''}$) with spectral intensity $> 1 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ for the transition $\nu_2(s) \leftarrow 0(a)$. The numbers in parentheses are 1σ uncertainties in the units of the last digit listed. Missing entries indicate a lack of convergence for a given parameter, which was set to zero in the final solution. An asterisk (*) indicates that the line parameter was fixed to its value in HITRAN2020. A table of line parameters with spectral intensity $< 1 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ for this transition is provided in the Supplementary Information.

Line	ν_0 cm^{-1}	$S \times 10^{19}$ $\text{cm}^{-1} / (\text{mol. cm}^{-2})$	γ_{air}^0 $\text{cm}^{-1}/\text{atm}$	γ_{self}^0 $\text{cm}^{-1}/\text{atm}$	δ_{air}^0 $\text{cm}^{-1}/\text{atm}$	δ_{self}^0 $\text{cm}^{-1}/\text{atm}$	Y^0 $1/\text{atm}$	a_w
${}^Q P(12,9)_a$	691.023944(10)	0.0154(1)	0.0774(30)	0.448(19)				
${}^Q P(11,10)_a$	700.965472(12)	0.0131(1)	0.0911(45)	0.547(23)				
${}^Q P(12,6)_a$	702.940985(13)	0.0114(1)	0.0566(26)	0.315(23)				
${}^Q P(11,9)_a$	707.046787(5)	0.0352(1)	0.0885(16)	0.515(9)				
${}^Q P(11,8)_a$	712.087270(9)	0.0182(1)	0.0967(35)	0.466(16)				
${}^Q P(11,7)_a$	716.242510(9)	0.0175(1)	0.0830(29)	0.405(17)				
${}^Q P(11,6)_a$	719.636566(5)	0.0320(1)	0.0722(13)	0.337(9)				
${}^Q P(11,5)_a$	722.366957(10)	0.0144(1)	0.0687(27)	0.252(18)				



$^Q P(10,9)_a$	723.270108(4)	0.0554(1)	0.0874(10)	0.545(7)	0.00817(99)		
$^Q P(11,4)_a$	724.509302(10)	0.0139(1)	0.0559(21)	0.277(19)			
$^Q P(11,3)_a$	726.127029(6)	0.0256(1)	0.0501(10)	0.213(11)			
$^Q P(11,2)_a$	727.243373(11)	0.0122(1)	0.0483(19)	0.137(20)			
$^Q P(11,1)_a$	727.901739(11)	0.0120(1)	0.0451(18)	0.133(20)			
$^Q P(10,8)_a$	728.574279(10)	0.0383(2)	0.0803(26)	0.535(14)			
$^Q P(10,7)_a$	732.943438(5)	0.0398(1)	0.0806(12)	0.417(8)			
$^Q P(10,6)_a$	736.508920(3)	0.0796(1)	0.0777(6)	0.402(5)	0.00590(58)		
$^Q P(10,5)_a$	739.374623(5)	0.0382(1)	0.0736(11)	0.357(8)			
$^Q P(10,4)_a$	741.621288(5)	0.0363(1)	0.0719(12)	0.327(9)			
$^Q P(10,3)_a$	743.306420(3)	0.0680(1)	0.0606(5)	0.248(5)	0.01031(48)		
$^Q P(10,2)_a$	744.486107(5)	0.0331(1)	0.0547(9)	0.218(9)			
$^Q P(10,1)_a$	745.177922(5)	0.0320(1)	0.0538(9)	0.193(9)			
$^Q P(9,8)_a$	745.292189(10)	0.0543(1)	0.0900(11)	0.548(7)	0.00330(109)	-0.050(14)	
$^Q P(10,0)_a$	745.419764(3)	0.0635(1)	0.0466(4)	0.167(5)	0.01180(36)		
$^Q P(9,7)_a$	749.862933(3)	0.0770(1)	0.0862(7)	0.503(5)	0.00775(70)		
$^Q P(9,6)_a$	753.590428(5)	0.1682(2)	0.0836(3)	0.454(3)	0.00547(31)	-0.021(7)	
$^Q P(9,5)_a$	756.584160(7)	0.0860(1)	0.0806(6)	0.409(5)	0.00554(57)	-0.044(9)	
$^Q P(9,4)_a$	758.929555(7)	0.0840(1)	0.0725(5)	0.355(5)	0.00812(50)	-0.031(9)	
$^Q P(9,3)_a$	760.693882(2)	0.1613(2)	0.0638(2)	0.315(3)	0.00964(22)		
$^Q P(9,2)_a$	761.918038(3)	0.0793(1)	0.0611(4)	0.254(4)	0.01120(42)		
$^Q P(9,1)_a$	762.640495(3)	0.0782(1)	0.0547(4)	0.218(4)	0.01104(36)		
$^Q P(8,7)_a$	767.037327(3)	0.1006(1)	0.0899(6)	0.552(5)			
$^Q P(8,6)_a$	770.913750(3)	0.2886(4)	0.0871(2)	0.508(2)	0.00300(20)	-0.031(5)	
$^Q P(8,5)_a$	774.025275(5)	0.1656(2)	0.0840(3)	0.452(3)	0.00679(32)	-0.021(7)	
$^Q P(8,4)_a$	776.461608(5)	0.1707(2)	0.0792(3)	0.412(3)	0.00675(29)	-0.034(6)	
$^Q P(8,3)_a$	778.289916(3)	0.3389(4)	0.0724(2)	0.362(2)	0.00838(13)	-0.035(5)	
$^Q P(8,2)_a$	779.563955(4)	0.1702(2)	0.0679(3)	0.308(3)	0.00992(26)	-0.021(6)	
$^Q P(8,1)_a$	780.314084(2)	0.1683(2)	0.0626(2)	0.276(3)	0.01004(21)		
$^Q P(8,0)_a$	780.567647(3)	0.3325(4)	0.0597(1)	0.269(2)	0.01092(10)	-0.014(5)	
$^Q P(7,6)_a$	788.510315(3)	0.3416(4)	0.0904(2)	0.561(2)	0.00060(18)	-0.024(5)	
$^Q P(7,5)_a$	791.726747(4)	0.2576(3)	0.0886(3)	0.508(2)	0.00290(23)	-0.024(6)	
$^Q P(7,4)_a$	794.244042(3)	0.2977(4)	0.0861(2)	0.450(2)	0.00373(19)	-0.030(5)	
$^Q P(7,3)_a$	796.134168(3)	0.6272(5)	0.0786(1)	0.400(1)	0.00563(9)	-0.030(5)	
$^Q P(7,2)_a$	797.447704(2)	0.3244(4)	0.0749(2)	0.351(2)	0.00704(14)		
$^Q P(7,1)_a$	798.222246(3)	0.3254(4)	0.0694(2)	0.310(2)	0.00744(13)	-0.016(5)	
$^Q P(6,5)_a$	809.714703(4)	0.2810(4)	0.0932(3)	0.556(2)		-0.038(6)	0.0121(26)
$^Q P(6,4)_a$	812.301057(3)	0.4267(5)	0.0905(2)	0.498(2)	0.00108(19)	-0.044(5)	0.0093(20)
$^Q P(6,3)_a$	814.241454(2)	1.0007(6)	0.0858(1)	0.436(1)	0.00303(8)	-0.039(4)	0.0084(9)



${}^Q P(6,2)_a$	815.591048(3)	0.5385(5)	0.0811(1)	0.381(1)	0.00358(13)	-0.026(5)	0.0136(16)	
${}^Q P(6,1)_a$	816.386214(3)	0.5591(6)	0.0769(1)	0.338(1)	0.00502(13)	-0.020(5)	-0.0093(19)	
${}^Q P(6,0)_a$	816.650989(1)	1.1273(7)	0.0747(1)	0.322(1)	0.00597(7)		0.0047(9)	
${}^Q P(5,4)_a$	830.653100(3)	0.4227(5)	0.0942(2)	0.551(2)	-0.00136(20)	-0.045(5)	0.0049(21)	
${}^Q P(5,3)_a$	832.634814(2)	1.3161(8)	0.0933(2)	0.473(1)	0.00121(6)	-0.028(4)	0.0026(1)	0.124(5)
${}^Q P(5,2)_a$	834.012176(2)	0.7814(6)	0.0874(1)	0.414(1)	0.00276(8)	-0.021(4)		
${}^Q P(5,1)_a$	834.823783(1)	0.8500(7)	0.0846(2)	0.347(1)	0.00331(7)			0.106(9)
${}^Q P(4,3)_a$	851.326964(1)	1.1990(4)	0.0964(1)	0.530(1)	-0.00115(5)	0.039(3)	0.0034(5)	
${}^Q P(4,2)_a$	852.724762(1)	0.9344(4)	0.0934(1)	0.438(1)	0.00163(6)	0.056(3)	0.0058(7)	
${}^Q P(4,1)_a$	853.548208(1)	1.1072(6)	0.0902(1)	0.365(1)	0.00269(4)	0.086(3)		0.111(5)
${}^Q P(4,0)_a$	853.817828(1)	2.2584(7)	0.0883(1)	0.336(1)	0.00413(3)	0.116(3)	0.0053(1)	0.109(2)
${}^Q P(3,2)_a$	871.736811(2)	0.7845(4)	0.0980(1)	0.484(1)	-0.00167(8)	0.021(3)	0.0077(8)	
${}^Q P(3,1)_a$	872.567121(1)	1.1918(5)	0.0957(1)	0.444(1)	0.00135(4)	0.072(3)	0.0033(1)	0.096(4)
${}^Q P(2,1)_a$	891.881924(1)	0.9172(6)	0.0987(2)	0.399(1)	-0.00106(6)	-0.009(3)		0.094(6)
${}^Q P(2,0)_a$	892.156656(1)	2.4022(7)	0.0956(1)	0.328(1)	0.00040(3)	-0.011(3)	0.0064(1)	0.117(2)
${}^Q Q(15,15)_a$	900.472748(4)	0.0285(1)	0.0930(14)	0.567(8)	0.00606(136)			
${}^Q Q(14,14)_a$	904.800019(4)	0.0332(1)	0.0921(12)	0.562(7)	0.01131(115)			
${}^Q Q(13,13)_a$	908.767389(2)	0.0721(1)	0.0950(6)	0.587(4)	0.01068(57)			
${}^Q Q(12,12)_a$	912.386259(1)	0.2901(2)	0.0935(2)	0.627(2)	0.00912(14)			
${}^Q Q(14,13)_a$	913.290967(7)	0.0171(1)	0.0906(22)	0.576(12)				
${}^Q Q(11,11)_a$	915.667346(1)	0.2766(2)	0.0937(2)	0.621(2)	0.00805(15)			
${}^Q Q(13,12)_a$	916.428023(5)	0.0756(1)	0.0890(5)	0.546(3)	0.00945(49)	0.033(7)		
${}^Q Q(10,10)_a$	918.620597(2)	0.4889(3)	0.0931(1)	0.627(1)	0.00852(9)	0.029(3)		
${}^Q Q(12,11)_a$	919.266284(2)	0.0794(1)	0.0914(5)	0.557(3)	0.00802(48)			
${}^Q Q(14,12)_a$	920.290102(6)	0.0181(1)	0.0888(20)	0.542(11)	0.00812(201)			
${}^Q Q(9,9)_a$	921.255116(1)	1.6239(5)	0.0946(1)	0.627(1)	0.00791(3)	0.041(3)	0.0043(1)	0.119(3)
${}^Q Q(11,10)_a$	921.811877(3)	0.1537(1)	0.0897(3)	0.555(2)	0.00795(25)	0.024(4)		
${}^Q Q(13,11)_a$	922.752591(5)	0.0208(1)	0.0869(17)	0.526(10)	0.00883(170)			
${}^Q Q(8,8)_a$	923.579066(1)	1.2556(5)	0.0948(1)	0.624(1)	0.00747(4)	0.044(3)	0.0015(1)	0.106(3)
${}^Q Q(10,9)_a$	924.070189(2)	0.5492(3)	0.0886(1)	0.579(1)	0.00740(9)	0.035(3)	0.0141(12)	
${}^Q Q(12,10)_a$	924.949970(3)	0.0438(1)	0.0859(8)	0.492(5)	0.01065(80)			
${}^Q Q(7,7)_a$	925.599645(1)	1.7950(6)	0.0957(1)	0.620(1)	0.00656(3)	0.052(3)	0.0037(1)	0.124(3)
${}^Q Q(9,8)_a$	926.045700(2)	0.4628(3)	0.0891(1)	0.572(1)	0.00747(9)	0.037(3)		
${}^Q Q(11,9)_a$	926.884570(3)	0.1732(1)	0.0839(2)	0.504(2)	0.00814(21)	0.019(4)		
${}^Q Q(6,6)_a$	927.323065(1)	4.7285(8)	0.0962(1)	0.618(1)	0.00582(2)	0.063(3)	0.0054(1)	0.122(1)
${}^Q Q(8,7)_a$	927.741994(1)	0.7174(4)	0.0896(1)	0.570(1)	0.00667(6)	0.042(3)		
${}^Q Q(13,10)_a$	927.977270(9)	0.0112(1)	0.0669(23)	0.405(17)				
${}^Q Q(10,8)_a$	928.557879(3)	0.1599(1)	0.0860(3)	0.503(2)	0.00659(25)	0.022(4)		
${}^Q Q(5,5)_a$	928.754524(1)	2.8482(7)	0.0976(1)	0.617(1)	0.00470(3)	0.054(3)	0.0084(1)	0.126(2)



$^Q Q(7,6)_a$	929.161739(1)	2.0515(7)	0.0925(1)	0.559(1)	0.00543(3)	0.047(2)	0.0061(1)	0.129(2)
$^Q Q(12,9)_a$	929.642370(3)	0.0498(1)	0.0809(7)	0.469(5)	0.00526(78)			
$^Q Q(4,4)_a$	929.898150(1)	3.0751(7)	0.0989(1)	0.613(1)	0.00369(3)	0.058(3)	0.0032(1)	0.127(2)
$^Q Q(9,7)_a$	929.970487(2)	0.2683(2)	0.0844(2)	0.514(2)	0.00626(21)	0.024(3)	0.0636(36)	
$^Q Q(6,5)_a$	930.306629(1)	1.3243(5)	0.0919(1)	0.565(1)	0.00476(4)	0.053(3)		
$^Q Q(3,3)_a$	930.757072(1)	5.8232(10)	0.0999(1)	0.605(1)	0.00284(2)	0.056(3)	0.0087(1)	0.126(1)
$^Q Q(11,8)_a$	931.062166(3)	0.0497(1)	0.0871(10)	0.447(5)				
$^Q Q(8,6)_a$	931.122043(1)	0.8195(5)	0.0846(1)	0.515(1)	0.00623(7)	0.038(3)		
$^Q Q(5,4)_a$	931.177372(1)	1.5284(7)	0.0949(1)	0.545(1)	0.00374(5)	0.049(3)		0.111(4)
$^Q Q(2,2)_a$	931.333294(1)	2.2597(8)	0.1018(1)	0.580(1)	0.00091(3)	0.043(3)	0.0234(1)	0.135(2)
$^Q Q(1,1)_a$	931.627798(1)	1.1633(6)	0.1017(2)	0.526(1)	0.00153(6)	0.052(3)		0.126(5)
$^Q Q(4,3)_a$	931.773587(1)	3.0229(8)	0.0963(1)	0.520(1)	0.00326(3)	0.061(3)	0.0051(1)	0.126(2)
$^Q Q(7,5)_a$	932.011239(2)	0.5746(4)	0.0866(1)	0.513(1)	0.00513(9)	0.037(3)		
$^Q Q(3,2)_a$	932.094067(1)	1.1647(6)	0.0958(1)	0.486(1)	0.00296(6)	0.050(3)		
$^Q Q(2,1)_a$	932.136195(2)	0.5311(4)	0.0972(1)	0.464(1)	0.00242(12)	0.074(3)		
$^Q Q(10,7)_a$	932.234806(2)	0.0915(1)	0.0793(4)	0.443(3)	0.01021(40)			
$^Q Q(13,9)_a$	932.292613(8)	0.0131(1)	0.1044*	0.500*	0.01341(417)			
$^Q Q(6,4)_a$	932.635735(1)	0.7171(4)	0.0887(1)	0.491(1)	0.00442(6)	0.040(3)		0.072(7)
$^Q Q(3,1)_a$	932.881241(2)	0.2789(2)	0.0930(2)	0.369(1)	0.00252(18)	0.071(3)		
$^Q Q(5,3)_a$	932.992346(1)	1.5059(6)	0.0912(1)	0.469(1)	0.00271(4)	0.044(2)	0.0115(1)	0.125(3)
$^Q Q(4,2)_a$	933.075863(2)	0.6078(4)	0.0908(1)	0.439(1)	0.00288(9)	0.050(3)		
$^Q Q(9,6)_a$	933.157404(2)	0.3071(2)	0.0798(1)	0.454(1)	0.00619(13)	0.033(3)		
$^Q Q(12,8)_a$	933.508227(11)	0.0144(1)	0.0724(18)	0.436(18)				
$^Q Q(8,5)_a$	933.826075(2)	0.2336(2)	0.0796(2)	0.448(2)	0.00647(19)	0.030(4)		
$^Q Q(4,1)_a$	933.842554(3)	0.1460(1)	0.0892(4)	0.345(2)	0.00512(34)	0.053(4)		
$^Q Q(7,4)_a$	934.235784(2)	0.3118(2)	0.0797(2)	0.433(2)	0.00797(28)	0.026(3)	-0.501(39)	
$^Q Q(5,2)_a$	934.252181(2)	0.3050(2)	0.0830(2)	0.403(1)	0.00183(26)	0.034(3)	0.541(40)	
$^Q Q(6,3)_a$	934.379647(1)	0.6989(4)	0.0827(1)	0.438(1)	0.00589(6)	0.043(3)		
$^Q Q(11,7)_a$	934.486087(4)	0.0286(1)	0.0741(11)	0.378(7)	0.00961(108)			
$^Q Q(5,1)_a$	934.994052(5)	0.0730(1)	0.0827(5)	0.313(3)	0.00509(46)	0.033(7)		
$^Q Q(10,6)_a$	935.221193(2)	0.1054(1)	0.0744(3)	0.389(3)	0.00714(28)			
$^Q Q(6,2)_a$	935.591689(3)	0.1444(1)	0.0788(2)	0.363(2)	0.00660(23)	0.034(4)		
$^Q Q(9,5)_a$	935.707391(4)	0.0875(1)	0.0748(4)	0.386(3)	0.00678(35)	0.034(6)		
$^Q Q(7,3)_a$	935.903810(2)	0.3072(2)	0.0754(1)	0.385(1)	0.00415(13)	0.031(3)		
$^Q Q(8,4)_a$	935.937325(3)	0.1272(1)	0.0746(3)	0.376(2)	0.00607(32)	0.024(5)	0.0493(47)	
$^Q Q(6,1)_a$	936.304884(3)	0.0343(1)	0.0752(9)	0.302(6)	0.00756(84)			
$^Q Q(7,2)_a$	937.059332(6)	0.0630(1)	0.0708(4)	0.307(4)	0.00770(43)	0.031(7)		
$^Q Q(11,6)_a$	937.268250(3)	0.0329(1)	0.0713(8)	0.339(6)	0.00577(82)			
$^Q Q(8,3)_a$	937.515514(1)	0.1253(1)	0.0705(2)	0.331(2)	0.00949(22)			



${}^Q Q(10,5)_a$	937.611525(4)	0.0297(1)	0.0693(10)	0.315(7)	0.00719(95)			
${}^Q Q(9,4)_a$	937.698851(3)	0.0475(1)	0.0694(7)	0.325(5)	0.00937(65)			
${}^Q Q(7,1)_a$	937.740407(7)	0.0151(1)	0.0674(19)	0.279(12)	0.00959(189)			
${}^Q Q(8,2)_a$	938.617525(4)	0.0255(1)	0.0622(9)	0.268(7)	0.00595(89)			
${}^Q Q(9,3)_a$	939.198142(3)	0.0464(1)	0.0634(5)	0.272(4)	0.00586(49)			
${}^Q Q(10,4)_a$	939.478940(6)	0.0160(1)	0.0594(13)	0.264(11)	0.01281(143)			
${}^Q Q(9,2)_a$	940.227526(11)	0.0100(1)	0.0598*	0.500*				
${}^Q Q(10,3)_a$	940.865857(6)	0.0158(1)	0.0596(13)	0.230(11)	0.01128(130)			
${}^Q R(0,0)_a$	951.776277(1)	1.7200(6)	0.1043(1)	0.357(1)	0.00453(3)	0.186(2)	-0.0012(1)	0.162(2)
${}^Q R(1,1)_a$	971.882067(1)	1.2105(5)	0.1022(1)	0.555(1)	0.00306(4)	0.066(3)		0.108(4)
${}^Q R(2,2)_a$	991.690541(1)	1.1878(5)	0.1018(1)	0.590(1)	0.00248(5)	0.047(3)		0.110(4)
${}^Q R(2,1)_a$	992.450303(1)	1.8033(7)	0.0966(1)	0.398(1)	0.00373(3)	0.080(2)	-0.0024(1)	0.133(3)
${}^Q R(2,0)_a$	992.698752(1)	3.8725(8)	0.0929(1)	0.301(1)	0.00522(2)	0.127(2)	0.0063(1)	0.118(2)
${}^Q R(3,3)_a$	1011.203704(1)	2.0782(6)	0.1002(1)	0.601(1)	0.00302(3)	0.066(3)	0.0013(1)	0.103(3)
${}^Q R(3,2)_a$	1012.445179(1)	1.6183(6)	0.0962(1)	0.473(1)	0.00388(3)	0.091(3)	0.0037(1)	0.119(3)
${}^Q R(3,1)_a$	1013.175596(1)	1.9152(6)	0.0907(1)	0.355(1)	0.00504(3)	0.103(3)	0.0014(1)	0.118(2)
${}^Q R(4,4)_a$	1030.422430(2)	0.8371(4)	0.0980(1)	0.617(1)	0.00368(6)	0.051(3)		
${}^Q R(4,3)_a$	1032.131113(1)	2.6106(6)	0.0953(1)	0.517(1)	0.00450(2)	0.093(3)	0.0045(1)	0.114(2)
${}^Q R(4,2)_a$	1033.315854(1)	1.5550(6)	0.0898(1)	0.426(1)	0.00494(3)	0.099(3)	0.0081(1)	0.119(3)
${}^Q R(4,1)_a$	1034.012794(1)	1.6845(6)	0.0847(1)	0.346(1)	0.00639(3)	0.104(3)	-0.0089(1)	0.118(3)
${}^Q R(4,0)_a$	1034.244907(1)	3.4515(8)	0.0818(1)	0.313(1)	0.00785(2)	0.128(2)	0.0045(1)	0.119(2)
${}^Q R(5,5)_a$	1049.346438(2)	0.6316(3)	0.0973(1)	0.623(1)	0.00411(7)	0.069(3)		
${}^Q R(5,4)_a$	1051.512043(1)	0.9667(5)	0.0936(1)	0.542(1)	0.00488(5)	0.084(3)		0.086(5)
${}^Q R(5,3)_a$	1053.130522(1)	2.2900(6)	0.0882(1)	0.464(1)	0.00565(2)	0.105(3)	0.0022(1)	0.119(2)
${}^Q R(5,2)_a$	1054.252803(1)	1.2316(5)	0.0827(1)	0.392(1)	0.00621(3)	0.098(3)	0.0083(1)	0.122(3)
${}^Q R(5,1)_a$	1054.912663(1)	1.2743(5)	0.0770(1)	0.330(1)	0.00741(3)	0.105(3)	-0.0039(1)	0.123(3)
${}^Q R(6,6)_a$	1067.974489(1)	0.8950(5)	0.0982(1)	0.622(1)	0.00493(6)	0.066(3)		0.104(5)
${}^Q R(6,5)_a$	1070.591112(2)	0.6670(3)	0.0917(1)	0.558(1)	0.00554(7)	0.084(3)		
${}^Q R(6,4)_a$	1072.627458(2)	0.7717(4)	0.0855(1)	0.489(1)	0.00580(5)	0.082(3)		
${}^Q R(6,3)_a$	1074.149268(1)	1.6381(5)	0.0807(1)	0.415(1)	0.00624(3)	0.099(3)	0.0091(1)	0.131(2)
${}^Q R(6,2)_a$	1075.203269(1)	0.8398(4)	0.0749(1)	0.356(1)	0.00756(4)	0.094(3)		0.143(4)
${}^Q R(6,1)_a$	1075.822975(1)	0.8463(4)	0.0671(1)	0.302(1)	0.00755(4)	0.092(3)		
${}^Q R(6,0)_a$	1076.033148(1)	1.6881(5)	0.0640(1)	0.285(1)	0.00940(2)	0.105(3)		
${}^Q R(7,7)_a$	1086.304290(2)	0.2974(2)	0.0963(2)	0.618(2)	0.00574(15)	0.048(4)		
${}^Q R(7,6)_a$	1089.370025(2)	0.8582(4)	0.0903(1)	0.561(1)	0.00633(5)	0.083(3)		
${}^Q R(7,5)_a$	1091.812026(2)	0.4849(3)	0.0845(1)	0.491(1)	0.00601(8)	0.055(3)		
${}^Q R(7,4)_a$	1093.711470(2)	0.5018(3)	0.0777(1)	0.430(1)	0.00649(7)	0.081(3)		
${}^Q R(7,3)_a$	1095.129363(1)	1.0079(4)	0.0708(1)	0.370(1)	0.00711(3)	0.079(3)		0.119(4)
${}^Q R(7,2)_a$	1096.112845(2)	0.5010(3)	0.0648(1)	0.312(1)	0.00815(5)	0.089(3)		



$^QR(7,1)_a$	1096.689838(2)	0.4958(3)	0.0586(1)	0.264(1)	0.00763(5)	0.087(3)	-0.0907(36)
$^QR(8,8)_a$	1104.332585(3)	0.1864(1)	0.0964(2)	0.606(2)	0.00725(23)	0.037(5)	
$^QR(8,7)_a$	1107.849515(3)	0.2614(2)	0.0904(2)	0.551(2)	0.00655(15)	0.052(4)	
$^QR(8,6)_a$	1110.689002(2)	0.5662(3)	0.0831(1)	0.506(1)	0.00677(7)	0.075(3)	
$^QR(8,5)_a$	1112.949309(2)	0.2877(2)	0.0775(1)	0.440(2)	0.00649(11)	0.051(4)	
$^QR(8,4)_a$	1114.706573(2)	0.2816(2)	0.0706(1)	0.383(2)	0.00629(10)	0.056(4)	
$^QR(8,3)_a$	1116.019697(2)	0.5411(3)	0.0641(1)	0.327(1)	0.00749(5)	0.079(3)	
$^QR(8,2)_a$	1116.926964(2)	0.2663(2)	0.0573(1)	0.267(3)	0.00773(8)	0.046(4)	
$^QR(8,1)_a$	1117.459089(2)	0.2616(2)	0.0517(1)	0.220(1)	0.00676(7)	0.069(4)	
$^QR(8,0)_a$	1117.647823(2)	0.5153(3)	0.0493(1)	0.210(1)	0.00809(4)	0.082(3)	
$^QR(9,9)_a$	1122.055163(3)	0.2177(1)	0.0933(3)	0.612(2)	0.01045(32)	0.050(4)	
$^QR(9,8)_a$	1126.029250(4)	0.1489(1)	0.0907(3)	0.555(3)	0.00770(26)	0.026(5)	
$^QR(9,7)_a$	1129.261825(4)	0.1580(1)	0.0842(2)	0.501(2)	0.00739(22)	0.042(5)	
$^QR(9,6)_a$	1131.869636(2)	0.3067(2)	0.0769(1)	0.451(2)	0.00689(11)	0.055(4)	
$^QR(9,5)_a$	1133.944268(4)	0.1470(1)	0.0715(2)	0.383(2)	0.00621(19)	0.033(5)	
$^QR(9,4)_a$	1135.556420(4)	0.1388(1)	0.0643(2)	0.329(2)	0.00654(17)	0.035(5)	-0.0157(44)
$^QR(9,3)_a$	1136.757508(3)	0.2612(2)	0.0542(1)	0.280(2)	0.00684(8)	0.064(4)	
$^QR(9,2)_a$	1137.592299(4)	0.1271(1)	0.0524(2)	0.222(2)	0.00751(14)	0.040(5)	
$^QR(9,1)_a$	1138.077579(4)	0.1232(1)	0.0465(1)	0.185(2)	0.00641(12)	0.064(5)	
$^QR(10,10)_a$	1139.466977(3)	0.0606(1)	0.0961(7)	0.611(6)	0.00657(69)		
$^QR(10,9)_a$	1143.907945(4)	0.1583(1)	0.0903(3)	0.558(3)	0.00762(25)	0.036(5)	
$^QR(10,8)_a$	1147.532764(2)	0.0814(1)	0.0852(4)	0.493(4)	0.00769(43)		
$^QR(10,7)_a$	1150.478293(6)	0.0777(1)	0.0790(4)	0.449(4)	0.00733(41)	0.038(8)	
$^QR(10,6)_a$	1152.852822(4)	0.1422(1)	0.0726(2)	0.401(3)	0.00703(20)	0.038(5)	
$^QR(10,5)_a$	1154.740824(7)	0.0654(1)	0.0654(4)	0.337(4)	0.00683(37)	0.026(9)	
$^QR(10,4)_a$	1156.207390(3)	0.0607(1)	0.0584(4)	0.291(4)	0.00640(34)		
$^QR(11,11)_a$	1156.561951(16)	0.0309(3)	0.0933(13)	0.540(13)	0.00748(129)		
$^QR(10,3)_a$	1157.304589(4)	0.1131(1)	0.0547(2)	0.238(3)	0.00752(17)	0.033(6)	
$^QR(10,2)_a$	1158.058166(3)	0.0540(1)	0.0476(3)	0.199(5)	0.00693(29)		
$^QR(10,1)_a$	1158.494324(3)	0.0526(1)	0.0434(3)	0.179(5)	0.00569(26)		
$^QR(10,0)_a$	1158.667119(4)	0.1037(1)	0.0414(1)	0.159(3)	0.00691(17)	0.038(6)	
$^QR(11,10)_a$	1161.483434(4)	0.0398(1)	0.0898(10)	0.565(7)	0.00624(95)		0.051(15)
$^QR(11,9)_a$	1165.502964(2)	0.0787(1)	0.0849(5)	0.503(4)	0.00909(45)		
$^QR(11,8)_a$	1168.779793(12)	0.0364(1)	0.0775(9)	0.458(7)	0.00843(84)		
$^QR(11,7)_a$	1171.440325(4)	0.0323(1)	0.0719(9)	0.392(8)	0.00920(85)		
$^QR(12,12)_a$	1173.333522(5)	0.0307(1)	0.0962(14)	0.633(9)	0.01201(136)		
$^QR(11,6)_a$	1173.583646(8)	0.0576(1)	0.0665(4)	0.361(5)	0.00708(43)	0.039(10)	
$^QR(11,5)_a$	1175.287053(5)	0.0260(1)	0.0603(8)	0.304(9)	0.00529(82)		
$^QR(11,4)_a$	1176.609701(5)	0.0237(1)	0.0515(7)	0.260(10)	0.00513(73)		



${}^Q R(11,3)_a$	1177.591221(3)	0.0436(1)	0.0438(3)	0.212(6)	0.00668(32)			
${}^Q R(11,2)_a$	1178.277610(6)	0.0208(1)	0.0433(7)	0.192(11)	0.00680(65)			
${}^Q R(11,1)_a$	1178.660802(6)	0.0200(1)	0.0394(6)	0.147(11)	0.00383(60)			
${}^Q R(12,11)_a$	1178.752575(7)	0.0187(1)	0.0998(24)	0.522(14)				
${}^Q R(12,10)_a$	1183.172650(8)	0.0181(1)	0.0864(20)	0.539(14)				
${}^Q R(12,9)_a$	1186.777525(5)	0.0319(1)	0.0817(11)	0.459(8)	0.00798(104)			
${}^Q R(12,8)_a$	1189.713243(10)	0.0138(1)	0.0798(23)	0.419(18)				
${}^Q R(12,7)_a$	1192.094983(11)	0.0119(1)	0.0666(21)	0.372(20)	0.00787(206)			
${}^Q R(12,6)_a$	1194.012659(6)	0.0209(1)	0.0610(11)	0.357(12)	0.00591(104)			
${}^Q R(13,12)_a$	1195.711523(9)	0.0166(1)	0.0892(23)	0.563(16)	0.01004(229)			
${}^Q R(12,3)_a$	1197.607283(8)	0.0148(1)	0.0489(11)	0.148(15)	0.00653(107)			
${}^Q R(12,0)_a$	1198.708581(27)	0.0138(1)	0.0384(8)	0.128(16)	0.00532(82)	0.144(33)		
${}^Q R(13,9)_a$	1207.676748(12)	0.0111(1)	0.0816(30)	0.415(22)				

265 **Table 7. Measured parameters at 296 K for lines (labelled ${}^{AK} \Delta J(J'', K'')_{l''}$) with spectral intensity $> 1 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ for the transition $\nu_2(a) \leftarrow 0(s)$. The numbers in parentheses are 1σ uncertainties in the units of the last digit listed. Missing entries indicate a lack of convergence for a given parameter, which was set to zero in the final solution. An asterisk (*) indicates that the line parameter was fixed to its value in HITRAN2020. A table of line parameters with spectral intensity $< 1 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ for this transition is provided in the Supplementary Information.**

Line	ν_0 cm^{-1}	$S \times 10^{19}$ $\text{cm}^{-1} / (\text{mol. cm}^{-2})$	γ_{air}^0 $\text{cm}^{-1}/\text{atm}$	γ_{self}^0 $\text{cm}^{-1}/\text{atm}$	δ_{air}^0 $\text{cm}^{-1}/\text{atm}$	δ_{self}^0 $\text{cm}^{-1}/\text{atm}$	Y^0 $1/\text{atm}$	a_w
${}^Q P(12,9)_s$	724.061132(11)	0.0151(1)	0.0755(29)	0.461(19)				
${}^Q P(12,6)_s$	726.929270(13)	0.0111(1)	0.0535(25)	0.297(24)				
${}^Q P(11,10)_s$	743.072899(13)	0.0127(1)	0.0887(44)	0.480(23)				
${}^Q P(11,9)_s$	743.957997(5)	0.0347(1)	0.0917(17)	0.527(10)				
${}^Q P(11,8)_s$	744.850693(9)	0.0183(1)	0.0870(30)	0.483(17)				
${}^Q P(11,7)_s$	745.705453(9)	0.0171(1)	0.0817(29)	0.416(17)				
${}^Q P(11,6)_s$	746.489888(5)	0.0314(1)	0.0666(12)	0.300(10)				
${}^Q P(11,5)_s$	747.181003(10)	0.0148(1)	0.0581(21)	0.343(19)				
${}^Q P(11,4)_s$	747.762549(11)	0.0136(1)	0.0618(25)	0.257(20)				
${}^Q P(11,3)_s$	748.226456(6)	0.0256(1)	0.0575(12)	0.236(11)				
${}^Q P(11,2)_s$	748.569636(11)	0.0124(1)	0.0443(17)	0.200(21)				
${}^Q P(11,1)_s$	748.767023(12)	0.0119(1)	0.0399(16)	0.151(21)				
${}^Q P(11,0)_s$	748.834248(6)	0.0230(1)	0.0420(9)	0.099(11)				
${}^Q P(10,9)_s$	764.077311(4)	0.0541(1)	0.0952(12)	0.564(7)				
${}^Q P(10,8)_s$	764.823394(5)	0.0371(1)	0.0881(15)	0.506(9)				
${}^Q P(10,7)_s$	765.561045(5)	0.0402(1)	0.0823(13)	0.451(8)				
${}^Q P(10,6)_s$	766.252403(3)	0.0784(1)	0.0746(6)	0.378(5)				
${}^Q P(10,5)_s$	766.870430(5)	0.0372(1)	0.0698(11)	0.334(8)				
${}^Q P(10,4)_s$	767.395851(5)	0.0351(1)	0.0618(10)	0.265(9)				



$^Q P(10,3)_s$	767.808735(8)	0.0669(1)	0.0532(4)	0.239(5)	-0.00169(41)	-0.055(11)		
$^Q P(10,2)_s$	768.128157(5)	0.0327(1)	0.0500(8)	0.200(9)				
$^Q P(10,1)_s$	768.311240(5)	0.0319(1)	0.0462(7)	0.181(9)				
$^Q P(9,8)_s$	784.980367(4)	0.0530(1)	0.0935(15)	0.552(7)				
$^Q P(9,7)_s$	785.596210(3)	0.0756(1)	0.0876(8)	0.510(5)				
$^Q P(9,6)_s$	786.190565(5)	0.1646(2)	0.0823(3)	0.454(3)	-0.00217(31)	-0.031(7)		
$^Q P(9,5)_s$	786.732205(3)	0.0836(1)	0.0748(5)	0.384(5)				
$^Q P(9,4)_s$	787.198874(7)	0.0817(1)	0.0687(5)	0.345(5)		-0.035(10)		
$^Q P(9,3)_s$	787.576067(2)	0.1582(2)	0.0625(2)	0.294(3)				
$^Q P(9,2)_s$	787.854562(3)	0.0782(1)	0.0546(4)	0.246(5)				
$^Q P(9,1)_s$	788.021371(7)	0.0764(1)	0.0507(4)	0.190(4)	-0.00820(38)	-0.067(10)		
$^Q P(9,0)_s$	788.077590(5)	0.1512(2)	0.0485(2)	0.187(3)	-0.00654(20)	-0.039(7)		
$^Q P(8,7)_s$	805.778912(3)	0.0975(1)	0.0953(7)	0.560(5)				
$^Q P(8,6)_s$	806.274178(2)	0.2796(3)	0.0877(2)	0.521(2)	-0.00206(21)			
$^Q P(8,5)_s$	806.737669(2)	0.1598(2)	0.0825(3)	0.452(3)				
$^Q P(8,4)_s$	807.144066(5)	0.1668(2)	0.0759(3)	0.388(3)		-0.024(7)		
$^Q P(8,3)_s$	807.471744(3)	0.3301(4)	0.0670(2)	0.347(2)	-0.00279(12)	-0.036(5)		
$^Q P(8,2)_s$	807.722260(5)	0.1664(2)	0.0620(2)	0.275(3)	-0.00345(31)	-0.020(6)	0.0573(59)	
$^Q P(8,1)_s$	807.871553(5)	0.1642(2)	0.0572(2)	0.242(3)	-0.00679(27)	-0.043(6)	-0.0530(55)	
$^Q P(7,6)_s$	826.470100(2)	0.3332(4)	0.0955(2)	0.576(2)	-0.00132(26)		-0.0078(28)	
$^Q P(7,5)_s$	826.855255(2)	0.2509(3)	0.0895(3)	0.520(3)	-0.00156(24)			
$^Q P(7,4)_s$	827.201102(2)	0.2908(4)	0.0829(2)	0.447(2)	-0.00198(26)		0.0174(35)	
$^Q P(7,3)_s$	827.487784(3)	0.6109(6)	0.0760(1)	0.387(1)	-0.00253(12)	-0.020(5)	0.0336(18)	
$^Q P(7,2)_s$	827.702055(3)	0.3119(4)	0.0693(2)	0.336(2)	-0.00308(23)	-0.022(5)	0.0160(47)	
$^Q P(7,1)_s$	827.833356(3)	0.3168(4)	0.0644(2)	0.289(2)	-0.00585(21)	-0.045(5)	-0.0994(43)	
$^Q P(7,0)_s$	827.877707(2)	0.6302(6)	0.0613(1)	0.277(1)	-0.00764(8)	-0.060(4)		
$^Q P(6,5)_s$	847.051290(2)	0.2723(3)	0.0971(3)	0.570(3)	-0.00114(33)		0.0096(37)	
$^Q P(6,4)_s$	847.337605(2)	0.4129(5)	0.0906(2)	0.511(2)	-0.00096(16)			
$^Q P(6,3)_s$	847.578087(2)	0.9696(8)	0.0842(1)	0.442(1)	-0.00184(7)	-0.028(4)		
$^Q P(6,2)_s$	847.762950(3)	0.5232(6)	0.0781(2)	0.377(1)	-0.00296(17)	-0.025(5)	0.0329(35)	
$^Q P(6,1)_s$	847.876314(3)	0.5429(6)	0.0735(1)	0.325(1)	-0.00499(14)	-0.042(5)	-0.0407(31)	
$^Q P(5,4)_s$	867.520135(2)	0.4134(3)	0.0985(1)	0.551(1)	0.00049(16)	0.056(3)	0.0125(21)	
$^Q P(5,3)_s$	867.719609(1)	1.2761(7)	0.0934(1)	0.477(1)	-0.00014(5)	0.051(3)	0.0110(1)	0.113(4)
$^Q P(5,2)_s$	867.872920(2)	0.7584(5)	0.0864(1)	0.404(1)	-0.00062(7)	0.044(3)		
$^Q P(5,1)_s$	867.968907(2)	0.8178(6)	0.0813(1)	0.343(1)	-0.00265(9)	0.019(3)	-0.0357(13)	
$^Q P(5,0)_s$	868.001571(1)	1.6857(8)	0.0802(1)	0.315(1)	-0.00447(4)			0.114(4)
$^Q P(4,3)_s$	887.876851(1)	1.1339(6)	0.0983(1)	0.527(1)	0.00188(8)	0.071(3)	-0.0134(14)	
$^Q P(4,2)_s$	887.999812(2)	0.9049(5)	0.0934(1)	0.440(1)	0.00050(9)	0.046(3)	0.0154(18)	
$^Q P(4,1)_s$	888.079400(1)	1.0670(5)	0.0885(1)	0.362(1)	-0.00158(6)	0.013(3)		



${}^Q P(3,2)_s$	908.112353(1)	0.7564(5)	0.0987(1)	0.477(1)	0.00220(9)	0.063(3)	-0.0077(10)	
${}^Q P(3,1)_s$	908.176885(1)	1.1511(8)	0.0962(2)	0.365(1)				0.115(6)
${}^Q P(3,0)_s$	908.199144(1)	2.5424(9)	0.0935(1)	0.330(0)	-0.00160(3)	-0.019(2)	-0.0013(1)	0.121(3)
${}^Q P(2,1)_s$	928.232054(1)	0.8893(5)	0.1025(2)	0.486(1)	0.00120(6)	0.017(3)		0.118(5)
${}^Q P(1,0)_s$	948.232082(1)	1.4413(5)	0.1034(1)	0.358(1)	0.00441(4)	0.183(2)	0.0046(1)	0.156(3)
${}^Q Q(15,15)_s$	952.431082(4)	0.0274(1)	0.0931(15)	0.604(8)	-0.00719(143)			
${}^Q Q(14,14)_s$	954.346928(4)	0.0314(1)	0.0965(13)	0.576(7)				
${}^Q Q(14,13)_s$	954.578178(7)	0.0162(1)	0.0853(22)	0.530(14)				
${}^Q Q(14,12)_s$	955.055778(7)	0.0173(1)	0.0863(20)	0.517(12)	-0.01028(202)			
${}^Q Q(13,13)_s$	956.150988(6)	0.0688(1)	0.0933(6)	0.595(4)	-0.00610(58)	0.061(7)		
${}^Q Q(13,12)_s$	956.396724(5)	0.0724(1)	0.0886(5)	0.546(5)	-0.00444(54)	0.033(7)		
${}^Q Q(13,11)_s$	956.841807(6)	0.0195(1)	0.0819(17)	0.479(11)				
${}^Q Q(13,10)_s$	957.398965(10)	0.0112(1)	0.0906*	0.500*				
${}^Q Q(12,12)_s$	957.838971(2)	0.2762(2)	0.0913(2)	0.626(2)	-0.00326(18)	0.077(3)	-0.0347(22)	
${}^Q Q(13,9)_s$	958.005751(8)	0.0123(1)	0.0677(22)	0.376(16)				
${}^Q Q(12,11)_s$	958.088723(5)	0.0759(1)	0.0863(5)	0.547(4)	-0.00453(51)	0.055(7)		
${}^Q Q(12,10)_s$	958.496699(3)	0.0423(1)	0.0808(8)	0.518(5)	-0.00677(77)			
${}^Q Q(12,9)_s$	958.989336(3)	0.0478(1)	0.0785(7)	0.457(5)	-0.00440(65)			
${}^Q Q(11,11)_s$	959.407100(2)	0.2642(2)	0.0916(2)	0.626(2)	-0.00404(20)	0.067(3)	-0.0246(29)	
${}^Q Q(12,8)_s$	959.513850(8)	0.0140(1)	0.0732(24)	0.409(14)				
${}^Q Q(11,10)_s$	959.652317(3)	0.1475(1)	0.0884(3)	0.562(2)	-0.00329(34)	0.043(4)	-0.0203(51)	
${}^Q Q(11,9)_s$	960.019880(3)	0.1667(1)	0.0823(3)	0.510(2)	-0.00302(23)	0.047(4)		
${}^Q Q(11,8)_s$	960.448221(7)	0.0482(1)	0.0751(6)	0.452(5)	-0.00432(64)	0.056(9)		
${}^Q Q(10,10)_s$	960.852157(2)	0.4678(3)	0.0909(1)	0.637(1)	-0.00426(10)	0.066(3)		
${}^Q Q(11,7)_s$	960.893028(4)	0.0281(1)	0.0734(12)	0.427(8)				
${}^Q Q(10,9)_s$	961.085933(2)	0.5250(3)	0.0859(1)	0.580(1)	-0.00360(8)	0.066(3)		
${}^Q Q(11,6)_s$	961.322387(25)	0.0294(9)	0.0702(9)	0.336(12)				
${}^Q Q(10,8)_s$	961.411013(3)	0.1530(1)	0.0834(2)	0.497(2)	-0.00227(30)	0.046(4)	-0.0321(44)	
${}^Q Q(10,7)_s$	961.776163(4)	0.0894(1)	0.0788(4)	0.448(3)	-0.00258(39)	0.039(6)		
${}^Q Q(10,6)_s$	962.144371(4)	0.1023(1)	0.0721(4)	0.401(3)		0.029(5)		
${}^Q Q(9,9)_s$	962.171436(1)	1.5504(6)	0.0919(1)	0.632(1)	-0.00351(4)	0.090(3)	-0.0136(1)	0.090(3)
${}^Q Q(9,8)_s$	962.388336(2)	0.4436(3)	0.0866(1)	0.576(1)	-0.00394(9)	0.063(3)		
${}^Q Q(10,5)_s$	962.488979(4)	0.0292(1)	0.0650(9)	0.339(7)				
${}^Q Q(9,7)_s$	962.669827(2)	0.2586(2)	0.0825(2)	0.507(2)	-0.00286(14)	0.050(3)		
${}^Q Q(10,4)_s$	962.790703(6)	0.0157(1)	0.0592(16)	0.252(12)				
${}^Q Q(9,6)_s$	962.973614(2)	0.2959(2)	0.0771(1)	0.455(1)	-0.00207(15)	0.050(3)	-0.0047(24)	
${}^Q Q(10,3)_s$	963.057363(6)	0.0158(1)	0.0529(13)	0.310(12)				
${}^Q Q(9,5)_s$	963.269115(5)	0.0851(1)	0.0734(4)	0.386(3)		0.031(6)		
${}^Q Q(8,8)_s$	963.362693(1)	1.2016(6)	0.0931(1)	0.633(1)	-0.00309(4)	0.079(3)	-0.0142(1)	0.101(4)



$^Q Q(9,4)_s$	963.534215(3)	0.0461(1)	0.0645(7)	0.326(5)				
$^Q Q(8,7)_s$	963.558533(1)	0.6891(4)	0.0876(1)	0.573(1)	-0.00298(7)	0.073(3)		
$^Q Q(9,3)_s$	963.736449(3)	0.0452(1)	0.0610(6)	0.252(5)				
$^Q Q(8,6)_s$	963.796216(1)	0.7923(4)	0.0830(1)	0.509(1)	-0.00231(6)	0.065(3)		0.061(6)
$^Q Q(8,5)_s$	964.041207(2)	0.2262(2)	0.0785(2)	0.452(2)	-0.00256(17)	0.037(4)		
$^Q Q(8,4)_s$	964.268488(3)	0.1234(1)	0.0763(3)	0.377(2)	-0.00130(28)	0.021(5)		
$^Q Q(7,7)_s$	964.424034(1)	1.7179(7)	0.0942(1)	0.629(1)	-0.00170(4)	0.094(3)	-0.0338(1)	0.118(3)
$^Q Q(8,3)_s$	964.467014(3)	0.1218(1)	0.0668(3)	0.327(2)	-0.00230(30)	0.019(5)		
$^Q Q(7,6)_s$	964.595752(1)	1.9825(7)	0.0901(1)	0.562(1)	-0.00198(3)	0.080(3)		0.103(3)
$^Q Q(8,2)_s$	964.609589(5)	0.0250(1)	0.0595(15)	0.268(9)				
$^Q Q(7,5)_s$	964.790231(2)	0.5592(3)	0.0838(1)	0.501(1)	-0.00174(8)	0.063(3)		
$^Q Q(7,4)_s$	964.979620(2)	0.3033(2)	0.0790(1)	0.434(1)	-0.00275(12)	0.053(3)		
$^Q Q(7,3)_s$	965.137736(2)	0.2983(2)	0.0738(1)	0.381(1)	-0.00218(12)	0.040(3)		
$^Q Q(7,2)_s$	965.272750(2)	0.0615(1)	0.0719(7)	0.315(4)				
$^Q Q(6,6)_s$	965.353930(1)	4.5603(10)	0.0961(1)	0.623(1)	-0.00178(2)	0.097(3)	-0.0086(1)	0.128(2)
$^Q Q(6,5)_s$	965.499444(1)	1.2850(7)	0.0927(1)	0.561(1)	-0.00105(5)	0.082(3)	0.0018(1)	0.122(4)
$^Q Q(6,4)_s$	965.652036(1)	0.6898(4)	0.0858(1)	0.496(1)	-0.00120(7)	0.066(3)		
$^Q Q(6,3)_s$	965.791313(1)	0.6793(4)	0.0810(1)	0.426(1)	-0.00297(7)	0.051(3)		
$^Q Q(6,2)_s$	965.899176(3)	0.1397(1)	0.0756(3)	0.351(2)	-0.00500(27)	0.028(4)		
$^Q Q(6,1)_s$	965.967690(3)	0.0333(1)	0.0695(10)	0.315(6)	-0.00625(97)			
$^Q Q(5,5)_s$	966.151154(1)	2.7467(9)	0.0979(1)	0.617(1)	-0.00069(3)	0.102(3)	-0.0150(1)	0.132(2)
$^Q Q(5,4)_s$	966.269252(1)	1.4774(7)	0.0953(1)	0.549(1)	-0.00147(5)	0.071(3)	0.0458(1)	0.137(3)
$^Q Q(5,3)_s$	966.379938(1)	1.4551(7)	0.0911(1)	0.474(1)	-0.00016(4)	0.063(2)		0.145(4)
$^Q Q(5,2)_s$	966.473673(2)	0.2954(2)	0.0842(2)	0.405(1)	-0.00123(19)	0.049(3)	-0.0563(41)	
$^Q Q(5,1)_s$	966.532416(2)	0.0708(1)	0.0782(6)	0.345(3)				
$^Q Q(4,4)_s$	966.814749(1)	2.9580(10)	0.0998(1)	0.614(1)		0.099(3)	0.0014(1)	0.132(2)
$^Q Q(4,3)_s$	966.905162(1)	2.9139(10)	0.0966(1)	0.526(1)	0.00057(3)	0.089(3)	0.0066(1)	0.134(2)
$^Q Q(4,2)_s$	966.981017(2)	0.5873(4)	0.0901(1)	0.446(1)	-0.00052(11)	0.053(3)		
$^Q Q(4,1)_s$	967.030884(3)	0.1415(1)	0.0871(4)	0.349(2)	-0.00269(36)	0.020(4)	-0.1625(82)	
$^Q Q(3,3)_s$	967.346381(1)	5.4605(13)	0.1002(1)	0.597(1)	0.00075(3)	0.101(3)	0.0066(1)	0.120(2)
$^Q Q(3,2)_s$	967.406845(1)	1.1240(6)	0.0960(1)	0.486(1)	0.00095(9)	0.067(3)	0.0135(37)	
$^Q Q(3,1)_s$	967.449136(2)	0.2682(2)	0.0929(3)	0.369(1)	0.00183(28)	0.017(3)	-0.122(14)	
$^Q Q(2,2)_s$	967.738455(1)	2.1603(8)	0.1030(1)	0.575(1)	0.00071(5)	0.078(3)	0.0375(1)	0.130(3)
$^Q Q(2,1)_s$	967.774785(1)	0.5086(4)	0.0974(2)	0.400(1)	0.00091(15)		-0.1654(27)	
$^Q Q(1,1)_s$	967.997801(1)	1.1096(5)	0.1021(1)	0.592(1)	0.00233(7)	0.043(3)		
$^Q R(1,1)_s$	1007.540521(2)	1.1401(12)	0.0984(2)	0.528(2)	0.00209(15)	0.101(7)		
$^Q R(1,0)_s$	1007.547125(1)	3.0649(14)	0.0979(1)	0.291(1)	0.00033(6)	0.063(7)		0.143(2)
$^Q R(2,2)_s$	1027.032962(1)	1.1380(7)	0.0986(1)	0.582(1)	0.00382(9)	0.108(3)		
$^Q R(2,1)_s$	1027.047044(1)	1.7326(9)	0.0956(1)	0.383(1)	0.00129(6)	0.066(3)		0.143(3)



$^QR(3,3)_s$	1046.374706(1)	1.9941(11)	0.0968(1)	0.606(1)	0.00406(8)	0.110(4)		
$^QR(3,2)_s$	1046.388053(1)	1.5660(14)	0.0928(3)	0.467(2)	0.00259(11)	0.097(4)		0.105(8)
$^QR(3,1)_s$	1046.400598(3)	1.8092(28)	0.0848(2)	0.379(4)		0.171(12)		
$^QR(3,0)_s$	1046.405551(1)	3.9377(31)	0.0872(1)	0.299(1)	-0.00058(6)			0.176(3)
$^QR(4,4)_s$	1065.563836(2)	0.7653*	0.0863(5)	0.504(9)	-0.00588(68)		-0.2000(52)	
$^QR(4,3)_s$	1065.565532(2)	2.3830*	0.0884(2)	0.601(4)	0.00482(24)	0.063(9)	0.0544(16)	
$^QR(4,2)_s$	1065.581753(1)	1.5936(11)	0.0949(2)	0.380(1)	0.00117(9)	0.143(3)		0.202(5)
$^QR(4,1)_s$	1065.594368(1)	1.6653(11)	0.0853(2)	0.334(1)	-0.00061(7)	0.117(3)		0.205(4)
$^QR(5,4)_s$	1084.583656(2)	0.9251(7)	0.0883(2)	0.546(3)	0.00119(14)	0.116(6)		
$^QR(5,3)_s$	1084.593124(2)	2.2158(15)	0.0862(2)	0.469(4)		0.222(7)		0.186(4)
$^QR(5,5)_s$	1084.599299(3)	0.6087(8)	0.0841(3)	0.555(9)	0.00431(29)	0.199(10)		
$^QR(5,2)_s$	1084.609882(1)	1.1923(9)	0.0745(1)	0.382(2)		0.098(4)		
$^QR(5,1)_s$	1084.623733(2)	1.2026(16)	0.0709(2)	0.346(3)	-0.00836(15)	0.147(9)		
$^QR(5,0)_s$	1084.629060(2)	2.5156(18)	0.0685(1)	0.310(1)	-0.00467(7)	0.042(7)		
$^QR(6,4)_s$	1103.430522(4)	0.7339(17)	0.0811(3)	0.511(6)	0.00299(24)	0.146(8)		
$^QR(6,5)_s$	1103.434373(6)	0.6421(16)	0.0923(4)	0.445(18)	-0.00287(38)	0.042(15)		
$^QR(6,3)_s$	1103.441239(2)	1.5817(9)	0.0751(1)	0.430(2)	-0.00123(9)	0.073(7)		
$^QR(6,2)_s$	1103.469829(2)	0.8048(6)	0.0679(1)	0.355(2)	-0.00303(11)	0.084(4)		
$^QR(6,6)_s$	1103.479582(2)	0.8463(7)	0.0889(2)	0.550(4)	-0.00232(16)			
$^QR(6,1)_s$	1103.485776(2)	0.8197(7)	0.0624(1)	0.335(2)	-0.00610(10)	0.037(6)		
$^QR(7,5)_s$	1122.093743(2)	0.4677(3)	0.0785(2)	0.474(2)	-0.00047(16)	0.074(4)		
$^QR(7,4)_s$	1122.104028(2)	0.4834(4)	0.0727(2)	0.433(2)	0.00182(16)	0.071(4)		
$^QR(7,6)_s$	1122.117766(2)	0.8160(5)	0.0823(1)	0.560(1)	0.00277(11)	0.121(3)		
$^QR(7,3)_s$	1122.132977(2)	0.9699(6)	0.0668(1)	0.364(1)	0.00038(7)	0.116(3)		
$^QR(7,2)_s$	1122.160044(2)	0.4847(3)	0.0567(1)	0.314(1)	-0.00308(11)	0.049(3)		
$^QR(7,1)_s$	1122.178400(2)	0.4735(4)	0.0537(1)	0.253(2)	-0.01636(13)	0.023(5)		
$^QR(7,0)_s$	1122.185302(2)	0.9551(6)	0.0505(1)	0.263(1)	-0.00861(6)	0.046(5)		
$^QR(7,7)_s$	1122.203657(3)	0.2841(2)	0.0860(3)	0.614(2)	-0.00501(25)	0.082(4)		
$^QR(8,5)_s$	1140.578024(6)	0.2689*	0.0769(5)	0.413(8)		0.075(13)		
$^QR(8,6)_s$	1140.579298(3)	0.5305*	0.0750(4)	0.520(5)	-0.00270(12)	0.098(8)		
$^QR(8,4)_s$	1140.603793(3)	0.2731(2)	0.0670(2)	0.373(2)	-0.00176(19)	0.062(4)		
$^QR(8,3)_s$	1140.627329(3)	0.5227(5)	0.0584(1)	0.327(2)	-0.00386(12)	0.104(5)		
$^QR(8,7)_s$	1140.632137(4)	0.2517(4)	0.0846(4)	0.555(4)	0.00151(39)	0.082(6)		
$^QR(8,2)_s$	1140.679206(3)	0.2569(2)	0.0509(1)	0.269(2)	-0.00206(12)	0.043(4)		
$^QR(8,1)_s$	1140.699717(3)	0.2515(2)	0.0466(1)	0.234(2)	-0.01067(10)	0.012(4)		
$^QR(8,8)_s$	1140.770681(3)	0.1776(1)	0.0935(3)	0.610(2)	-0.00482(28)	0.078(5)		
$^QR(9,6)_s$	1158.865536(2)	0.2946(2)	0.0742(2)	0.449(2)	-0.00121(15)	0.078(4)		
$^QR(9,7)_s$	1158.884906(11)	0.1486(10)	0.0741(6)	0.488(6)		0.066(8)		
$^QR(9,5)_s$	1158.887592(11)	0.1411(10)	0.0729(7)	0.396(6)		0.080(9)		



$^QR(9,4)_s$	1158.928983(4)	0.1343(1)	0.0619(2)	0.333(3)		0.056(5)		
$^QR(9,8)_s$	1158.975947(4)	0.1408(1)	0.0739(4)	0.523(4)	-0.00139(39)	0.046(6)		
$^QR(9,3)_s$	1158.985042(3)	0.2527(2)	0.0539(1)	0.306(2)	0.00334(14)	0.055(4)		
$^QR(9,2)_s$	1159.025845(4)	0.1224(1)	0.0412(2)	0.246(3)	-0.00512(19)	0.019(5)		
$^QR(9,1)_s$	1159.047365(4)	0.1188(1)	0.0386(2)	0.177(3)	-0.01819(20)	-0.015(6)		
$^QR(9,0)_s$	1159.056104(1)	0.2346(2)	0.0384(1)	0.208(2)	-0.00906(9)			
$^QR(9,9)_s$	1159.180082(3)	0.2057(1)	0.0931(2)	0.616(2)	-0.00300(22)	0.072(5)		
$^QR(10,7)_s$	1176.963675(6)	0.0744(1)	0.0757(6)	0.443(4)		0.070(8)		
$^QR(10,6)_s$	1176.976760(4)	0.1366(1)	0.0712(3)	0.400(3)	-0.00169(31)	0.050(5)		
$^QR(10,8)_s$	1177.008560(6)	0.0781(1)	0.0796(7)	0.496(4)	-0.00297(70)	0.048(8)		
$^QR(10,5)_s$	1177.021239(7)	0.0633(1)	0.0635(6)	0.357(5)	-0.00218(60)	0.039(9)		
$^QR(10,4)_s$	1177.077707(7)	0.0586(1)	0.0573(5)	0.295(5)	-0.00148(50)	0.042(9)		
$^QR(10,3)_s$	1177.110798(4)	0.1096(1)	0.0473(2)	0.251(3)	-0.00306(20)	0.030(6)		
$^QR(10,9)_s$	1177.147801(4)	0.1501(1)	0.0832(3)	0.559(3)	-0.00321(32)	0.088(5)		
$^QR(10,2)_s$	1177.197873(3)	0.0522(1)	0.0411(4)	0.194(5)				
$^QR(10,1)_s$	1177.218002(3)	0.0507(1)	0.0362(3)	0.157(5)	-0.00967(27)			
$^QR(10,10)_s$	1177.431563(8)	0.0569(1)	0.0947(7)	0.603(5)	0.00249(74)	0.089(11)		
$^QR(11,7)_s$	1194.868068(67)	0.0343(21)	0.0619(18)	0.383(24)		0.134(30)		
$^QR(11,8)_s$	1194.869935(62)	0.0321(21)	0.0969(38)	0.501(27)	-0.00800(181)			
$^QR(11,6)_s$	1194.911206(8)	0.0560(1)	0.0652(6)	0.376(5)		0.049(10)		
$^QR(11,9)_s$	1194.948087(6)	0.0755(1)	0.0810(5)	0.520(4)	-0.00310(64)	0.047(8)		
$^QR(11,5)_s$	1194.976337(5)	0.0251(1)	0.0618(12)	0.299(10)				
$^QR(11,4)_s$	1195.046529(6)	0.0228(1)	0.0565(10)	0.240(10)				
$^QR(11,3)_s$	1195.133708(10)	0.0426(1)	0.0473(5)	0.243(6)	0.00377(42)	0.040(12)		
$^QR(11,10)_s$	1195.146565(4)	0.0377(1)	0.0769(12)	0.552(8)				
$^QR(11,2)_s$	1195.193054(6)	0.0203(1)	0.0342(8)	0.246(12)	-0.00418(81)			
$^QR(11,1)_s$	1195.206980(6)	0.0198(1)	0.0320(8)	0.187(13)	-0.01644(84)			
$^QR(11,0)_s$	1195.216938(4)	0.0385(1)	0.0331(4)	0.145(7)	-0.00848(38)			
$^QR(11,11)_s$	1195.525249(5)	0.0295(1)	0.0947(19)	0.610(10)	0.00468(149)			
$^QR(12,8)_s$	1212.558281(10)	0.0136(1)	0.0720(29)	0.463(18)				
$^QR(12,9)_s$	1212.581165(14)	0.0308(1)	0.0777(14)	0.487(9)	-0.00515(146)	0.074(18)		
$^QR(12,7)_s$	1212.595263(11)	0.0114(1)	0.0709(37)	0.331(21)				
$^QR(12,6)_s$	1212.665106(7)	0.0203(1)	0.0588(12)	0.350(12)	-0.00556(129)			
$^QR(12,10)_s$	1212.701556(8)	0.0172(1)	0.0796(23)	0.500(15)				
$^QR(12,3)_s$	1212.863473(8)	0.0150(1)	0.0409(9)	0.195(15)				
$^QR(12,11)_s$	1212.971122(8)	0.0179(1)	0.0828(20)	0.587(15)				
$^QR(12,12)_s$	1213.461314(15)	0.0287(1)	0.0854(12)	0.606(10)	-0.00389(123)	0.073(19)		
$^QR(13,9)_s$	1230.044062(13)	0.0109(1)	0.0674(24)	0.527(24)				
$^QR(13,12)_s$	1230.620500(68)	0.0147(7)	0.0716(21)	0.531(31)	-0.01945(234)	0.208(38)		



270

In order to analyse our measured dataset, we produced a number of plots of the individual line parameters, comparing our new line parameters for the NH_3 ν_2 band to those reported in the HITRAN 2020 database (Gordon et al., 2021). HITRAN is chosen here for the primary comparison because its NH_3 linelist forms the basis for the majority of remote sensing datasets of NH_3 in the Earth's atmosphere. Unless explicitly stated otherwise, the plots shown in the following sections display comparisons of data for the Q branch of the $\nu_2(a) \leftarrow 0(s)$ transition. Analogous comparison plots for the P and R branch for this transition, along with the P, Q and R branches of the $\nu_2(s) \leftarrow 0(a)$ transition, can be found in the Supplementary Information. As long as sufficient parameters were available, similar comparison plots are shown in the Supplementary Information for the other, less intense bands found in the measured spectral window. In every case, the uncertainties in the plots represent Labfit internal uncertainties at three standard deviations (3σ) for additional clarity. All uncertainties in tables are listed at 1σ .

280 **4.1 Uncertainties of the retrieved parameters**

The line parameter uncertainties directly provided by Labfit (Benner et al. 1995) correspond to Type A uncertainties (Joint Committee for Guides in Metrology, 2008), and are determined from the noise level of the spectrum, the spectral residuals, and the derived parameters in the solution. Labfit does not report Type B uncertainties (Joint Committee for Guides in Metrology, 2008) of the line parameters arising from uncertainties in quantities such as the sample conditions (i.e., temperature, pressure, pathlength), the partition sums, issues with the instrumental lineshape of the spectrometer, and uncertainties arising from a poor choice of lineshape function, so we calculate these Type B contributions separately. The overall uncertainty can be found by adding the Type A and Type B contributions in quadrature and taking the square root.

285

Table 8. Type B uncertainties for the measured NH_3 line parameters.

Spectral line parameter	Pressure (%)	Pathlength (%)	N_2O wavenumber calibration (cm^{-1})	Total Type B uncertainty
ν_0	-	-	0.00005	0.00005 cm^{-1}
S	0.65	0.40	-	0.76 %
γ_{self}^0	0.65	0.40	-	0.76 %
γ_{air}^0	0.09	0.40	-	0.41 %
δ_{self}^0	0.65	-	-	0.65 %
δ_{air}^0	0.09	-	-	0.09 %
$a_w(air)$	0.09	0.40	-	0.41 %
γ_{air}^0	0.09	0.40	-	0.41 %

290



Type B contributions are outlined in Table 8 and detailed below. The assumption we made in this study is that 1σ measurement uncertainties can be used to represent the maximum Type B uncertainty contributions to the derived line parameters. There are a number of contributions that are very small and so are not included in this table. For example, as the majority of parameters are temperature-dependent, there will be a very small uncertainty due to possible deviation of the true temperature from 296 K. An uncertainty of 0.05 K will contribute no more than 0.02 % to the Type B budget. Additionally, the stated NH_3 purity was 99.98%, so we can assume that the uncertainty on the absorber amount is no more than 0.02 %.

The NIST N_2O lines (Maki et al., 1991) used for the wavenumber calibration in the present analysis have stated uncertainties of $\sim 0.00005 \text{ cm}^{-1}$, which we can equate to a Type B uncertainty in the retrieved line positions. The other line parameters have contributions to their Type B uncertainty from uncertainties in pathlength and absorber pressure. The pathlength contribution is determined from the calibration process, namely 0.1% uncertainty in N_2O pressures, 0.24% uncertainty in N_2O line intensities (Devi et al., 2003), and 0.3 % resulting from the spectral fit.

Uncertainties in the pressures depend on which of the manometer pressure gauges were used for the pressure measurements and their calibrations; all were factory calibrated using reference gauges traceable to the PTB standard. The line intensity, self-broadened halfwidth, and self-pressure-induced shift parameters are most sensitive to the pure NH_3 pressure measurements, which are estimated to have $\sim 0.65\%$ uncertainty. The air-broadened halfwidths, air-pressure-induced shifts, speed dependence of the air-broadened halfwidth and line mixing parameters are most sensitive to the partial pressure of air. For the range 200–1000 mbar pressure of air used in our experiments we have assumed a representative estimation of 0.09% uncertainty. Given the small amount fraction of NH_3 in air (less than ~ 0.004), the contribution of NH_3 partial pressure uncertainties is minimal. Also note that the pressure-induced shifts do not depend on cell pathlength, and represent wavenumber differences so that the absolute wavenumber calibration has minimal contribution. Lastly, it should be noted that Rosenkranz line mixing used in the present analysis is only an approximation, and therefore the uncertainty contributions from its use cannot easily be quantified. The total uncertainties in our line parameters compare favourably to those for the corresponding line parameters in HITRAN 2020. For the strongest lines, uncertainties in line positions are comparable, in the range $0.00001 - 0.0001 \text{ cm}^{-1}$. For the strongest lines in the Q branches of the ν_2 band, the total uncertainties of our line intensities, and self- and air-broadened Lorentz halfwidths are of the order of 1 %, a significant improvement over those reported in HITRAN 2020: 10 – 20 %, 2 – 5 %, and 2 – 5 %, respectively.

4.2 Line positions

Newly determined line positions (ν_0) have been compared with those from the HITRAN 2020 database, whose positions are based upon the analysis of Down et al. (Down et al., 2013), by plotting the position differences in Fig. 4(a). For the $\nu_2(a) \leftarrow 0(s)$ Q branch lines up to $J = 6$, our line positions are higher by up to 0.00005 cm^{-1} , comparable to the uncertainty (~ 0.00005



cm⁻¹) of the N₂O lines from the NIST heterodyne wavenumber calibration tables (Maki et al., 1991) used in the wavenumber calibration. For lines with $J > 6$, our line positions tend to be lower than those reported in HITRAN 2020, although these lines are weaker with lower SNR in our overall analysis, and therefore the differences show more scatter. Table 9 reports that the average position difference between the new values and those from HITRAN is positive for the P, Q and R branches of both the $\nu_2(a) \leftarrow 0(s)$ and $\nu_2(s) \leftarrow 0(a)$ transitions, with the P and R branches displaying a greater shift in position than the Q branches.

4.3 Line intensities

While Figure 4(a) shows differences between our measured line positions and those from HITRAN 2020 (Gordon et al., 2021) for the NH₃ $\nu_2(a) \leftarrow 0(s)$ Q branch lines, Figure 4(b) shows a plot of the percentage deviations of our measured spectral line intensities (S) against those from the HITRAN 2020 database (Gordon et al., 2021), again with intensities based on the work of Down et al. (Down et al., 2013). The scatter in percentage deviation as J increases is a consequence of the relative decrease in SNR for these high- J lines. Overall, our line intensities are slightly higher than those in HITRAN 2020. Considering only lines with $S > 1 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ at 296 K, we report an average increase in S of 3.4% for lines in the two $\nu_2 \leftarrow 0$ Q branches. Similar increases of 3-7% are observed for the P and R branches; see Table 9 for further information.




Figure 4. (a) Differences between the measured line positions and those from HITRAN 2020 (Gordon et al., 2021) for the NH₃ $\nu_2(a) \leftarrow 0(s)$ Q branch lines, and (b) percentage deviations in intensity between the measured line intensities and those from HITRAN 2020 for the NH₃ $\nu_2(a) \leftarrow 0(s)$ Q branch lines. For both cases, only lines with line intensity $> 1 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ are plotted, and the error bars are 3σ .



Table 9. Average absolute position differences, and percentage deviations of intensity and Lorentz halfwidths between parameters derived in this work and those in HITRAN 2020 (Gordon et al., 2021); the percentage deviations are calculated as $100(\text{Labfit}/\text{HITRAN} - 1)$. The values were determined from lines with intensity $S > 1 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ at 296 K. Halfwidth deviations were only determined for those lines with distinct halfwidth parameters for the analogous lines in HITRAN 2020, as opposed to an arbitrary fill value discussed in Section 4.4.

Transition		No. of lines above S threshold	Average absolute position difference (10^{-5} cm^{-1})	Average S deviation (%)	Average γ_{self}^0 deviation (%)	Average γ_{air}^0 deviation (%)
$\nu_2(s) \leftarrow 0(a)$	P	62	7.52	4.7	3.7	-2.5
	Q	70	4.14	3.6	-3.6	-6.5
	R	83	9.66	4.5	7.7	-1.7
$\nu_2(a) \leftarrow 0(s)$	P	63	7.73	6.6	1.4	-4.6
	Q	69	3.55	3.2	-2.0	-8.2
	R	82*	8.24	2.8	6.2	-7.7

*For the $\nu_2(a) \leftarrow 0(s)$ R branch, four lines (two pairs) were blended together. In order to fit the other line parameters, their intensities were fixed to the values from HITRAN 2020.

4.4 Self- and air-broadened Lorentz halfwidths

Figures 5(a) and 6(a) show the present measured self- (γ_{self}^0) and air- (γ_{air}^0) broadened halfwidths for all lines with intensity $> 1 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ at 296 K compared to the corresponding values from HITRAN 2020, which were derived by Nemtchinov et al. using a polynomial function of m and K , apparently only from measurements in the P and R branches of the ν_2 band (Nemtchinov et al., 2004). Our measured values are found to show the same general trend, with both γ_{self}^0 and γ_{air}^0 increasing with K for the same J value. However, there are notable differences between the magnitudes of our halfwidths and those from HITRAN. Average percentage deviations of our measured parameters from HITRAN for the $\nu_2(a) \leftarrow 0(s)$ Q branch are plotted in Figs. 5(b) and 6(b), with averages for other branches listed in Table 9 for both γ_{self}^0 and γ_{air}^0 . We observe significant deviations from the Nemtchinov-derived values, 4-10% for γ_{self}^0 and 2-8% for γ_{air}^0 . Analysis of the individual branches shows that the γ_{self}^0 are 2-4% lower than the Nemtchinov values in the Q branches but 1-4% and 6-8% higher for the P and R branches, respectively. The γ_{air}^0 percentage deviations demonstrate a consistent pattern of overestimation by HITRAN 2020; our values are 2-5%, 6-8%, and 2-8% lower for the P, Q, and R branches, respectively. In addition, we report values of the halfwidth parameters for $J'' > 8$ for the first time. Previously, in HITRAN 2020, γ_{self}^0 values for $J'' > 8$ were set to 0.5, while γ_{air}^0 values for $J'' > 8$ were based on the widths reported for lower J values, using an extrapolation of the width parameters determined by Nemtchinov et al. (Nemtchinov et al., 2004).




Figure 5. Plots of (a) self-broadened halfwidth, γ_{self}^0 , and (b) percentage deviations from HITRAN of γ_{self}^0 plotted against K for each J value of the $\nu_2(a) \leftarrow 0(s)$ Q branch. The Labfit values are included with uncertainties of 3σ . For $J > 8$, HITRAN values for the self-broadened halfwidth were all set to 0.5, so percentage deviations have not been calculated for these cases.




Figure 6. Plots of (a) air-broadened halfwidth, γ_{air}^0 , and (b) percentage deviations from HITRAN of γ_{air}^0 plotted against K for each J value of the $\nu_2(a) \leftarrow 0(s)$ Q branch. The Labfit values are included with uncertainties of 3σ . For $J > 9$, HITRAN values for the air-broadened halfwidth were set to the same value as their corresponding K value at $J = 9$ (with values of $K > 9$ set to the value for $K = 9$), so percentage deviations have not been calculated for these cases.



4.5 Self- and air-pressure-induced shifts

Figure 7 reports the self- (δ_{self}^0) and air- (δ_{air}^0) pressure-induced shifts for lines with an intensity $> 1 \times 10^{-21} \text{ cm}^{-1} / (\text{molecule cm}^{-2})$ at 296 K. Since the HITRAN 2020 database does not list any values for δ_{self}^0 or δ_{air}^0 for NH_3 , we report values for these parameters for the complete ν_2 band for the first time. Due to the optimisation of our experimental measurements for atmospheric conditions, i.e. the smaller range of NH_3 pressures than air pressures, we observe larger errors in our determination of δ_{self}^0 compared with δ_{air}^0 .




Figure 7. Plots of (a) self-pressure-induced shifts, δ_{self}^0 , and (b) air-pressure-induced shifts, δ_{air}^0 , against K for each J value of the $\nu_2(a) \leftarrow 0(s)$ Q branch. Measured values are plotted with uncertainties of 3σ . The HITRAN 2020 database does not include values for these parameters.

According to Fig. 7(a), the values of our measured δ_{self}^0 parameters for the $\nu_2(a) \leftarrow 0(s)$ transition tend to decrease with increasing J, but increase with increasing K. A plot of δ_{self}^0 against J'' for only $K = 3$ levels is shown in Fig. 8(a), and further indicates that the decrease in δ_{self}^0 with increasing J holds for all three branches of the two $\nu_2 \leftarrow 0$ transitions, with the exception of a single line in the R branch. Figure 8(b) indicates a tendency for δ_{air}^0 values associated with the $\nu_2(a) \leftarrow 0(s)$ Q branch to decrease with both increasing J and K (apart from two more R branch lines). However, the plot of δ_{air}^0 against J'' for $K = 3$ in Fig. 8(b) suggests that the trend of δ_{air}^0 with J is dependent on the symmetry of the transition, unlike for δ_{self}^0 . For P, Q, R branches of the $\nu_2(s) \leftarrow 0(a)$ transition, δ_{air}^0 increases with J'' , but for P, Q, R branches of the $\nu_2(a) \leftarrow 0(s)$ transition, δ_{air}^0 decreases with J'' . There are not enough fitted δ_{self}^0 and δ_{air}^0 values for the other transitions to provide a more detailed analysis of these J-dependent observations.




Figure 8. Plots of (a) self-pressure-induced shifts, δ_{self}^0 , and (b) air-pressure-induced shifts, δ_{air}^0 , against J'' for $K = 3$, for the P, Q and R branches of the $\nu_2(s) \leftarrow 0(a)$ and $\nu_2(a) \leftarrow 0(s)$ transitions. The present measured values are given with uncertainties of 3σ . The HITRAN 2020 database does not include values for these parameters in this band.

Although values of δ_{self}^0 are not currently included in the HITRAN or GEISA databases, we were able to compare our values to previous data in the literature. Aroui et al. (2009) and Guinet et al. (2011) recorded measurements of several pure NH_3 spectra at wavenumbers $> 1000 \text{ cm}^{-1}$ and at spectral resolutions of 0.0038 cm^{-1} and 0.005 cm^{-1} respectively. They determined δ_{self}^0 values for several lines in the $\nu_2(s) \leftarrow 0(a)$ R branch. Figure 9(a) shows a comparison plot of δ_{self}^0 for our work against both studies, for values where $J = 9$. We predict higher δ_{self}^0 values, an average increase of $0.0333 \text{ cm}^{-1}/\text{atm}$ from Aroui et al. and $0.0570 \text{ cm}^{-1}/\text{atm}$ from Guinet et al. The data from the Guinet paper does however obey the same trend of δ_{self}^0 decreasing with K as found in our data.

With regards to the P and Q branches, Clar et al. (1988) and Ibrahimi et al. (1999) recorded measurements of several lines for both $\nu_2(s) \leftarrow 0(a)$ and $\nu_2(a) \leftarrow 0(s)$ transitions using tuneable diode laser spectrometers. Figure 9(b) shows a comparison plot of δ_{self}^0 values against the data from Clar et al. for $\nu_2(a) \leftarrow 0(s)$, where $J = 7$. We observe an average increase of $0.0532 \text{ cm}^{-1}/\text{atm}$ from the work of Clar, and the same general trend of increasing δ_{self}^0 with increasing K for the Q branch. Figure 9(c) and 9(d) show comparison plots of δ_{self}^0 against the values from Ibrahimi et al. for $\nu_2(s) \leftarrow 0(a)$ and $\nu_2(a) \leftarrow 0(s)$ respectively. We observe an average increase of $0.0545 \text{ cm}^{-1}/\text{atm}$ from the work of Ibrahimi, and observe the same general trends of δ_{self}^0 with J and K .

We found no studies reporting δ_{air}^0 values, and thus cannot make any comparisons to our values.





Figure 9. Comparison plots of self-pressure-induced shifts, δ_{self}^0 , from our work and other papers, against K , for (a) the R branch of the $\nu_2(s) \leftarrow 0(a)$ transition, where $J = 9$; (b) the Q branch of the $\nu_2(a) \leftarrow 0(s)$ transition, where $J = 7$; (c) the P, Q and R branches of the $\nu_2(s) \leftarrow 0(a)$ transition; (d) the P and Q branches of the $\nu_2(a) \leftarrow 0(s)$ transition. Unlike other plots, uncertainties are plotted here as 1σ to better compare with the previous studies. The HITRAN 2020 database does not include values for these parameters in this band.

4.6 Rosenkranz line mixing and speed dependence

Figure 10(a) shows the Rosenkranz line mixing parameters, Y_{air}^0 , plotted against the quantum number K for lines of the $\nu_2(a) \leftarrow 0(s)$ Q branch where values were determinable. The largest values of Y_{air}^0 for this transition occur for $K = 1$ levels. The lines corresponding with the parameters in Fig. 10(a) span a wavenumber range of 1 cm^{-1} , and there is significant overlap between the lines when air-broadened at pressures approaching 1000 hPa. Line mixing parameters are almost exclusively fit for the Q branches, which are more congested than the P and R branches.



445 **Figure 10. Plots of (a) Rosenkranz line mixing parameter, Y^0_{air} , and (b) speed dependence of the halfwidth parameter, $a_w(air)$, against K for each J value of the $\nu_2(a) \leftarrow 0(s)$ Q branch. The present measured values are shown with uncertainties of 3σ . The HITRAN 2020 database does not include values for these line parameters of NH_3 .**

Figure 10(b) provides a plot of the speed dependence of the width parameter, $a_w(air)$, where this parameter was determinable, against K for lines in the $\nu_2(a) \leftarrow 0(s)$ Q branch. The results show a decrease in a_w as J and K increase, although this trend is not observed in the Q branch for the $\nu_2(s) \leftarrow 0(a)$, where a_w remains constant across all values of J and K . We report an average value for a_w of 0.131 for $\nu_2(a) \leftarrow 0(s)$ and 0.118 for $\nu_2(s) \leftarrow 0(a)$.

As an illustration of the benefit in floating both Y^0_{air} and $a_w(air)$ parameters in the multispectrum fit, the plots in Fig. 11 show the final spectral fit residuals for both ν_2 Q branches, compared with residuals calculated from the same fit but with all Y^0_{air} and $a_w(air)$ values set to zero. It is clear that the inclusion of each parameter in the multispectrum fit improves the residuals, bringing them closer to zero.




Figure 11. Comparison plots over the spectral region containing the ν_2 Q branches (900-980 cm^{-1}) of the weighted fit residuals for the measured highest-pressure spectrum (total pressure 978.3 hPa): (a) with (black) and without (red) the inclusion of Rosenkranz line mixing; (b) with (black) and without (green) the inclusion of speed dependence; (c) shows the measured laboratory transmittance spectrum.

5 Conclusions

We have carried out the most complete experimental and multispectrum analysis of air-broadened NH_3 over the 675 cm^{-1} to 1250 cm^{-1} spectral region to date. This spectral region covers the most intense NH_3 features observed in atmospheric spectra, the $0100\ 00\ 0\ s \leftarrow 0000\ 00\ 0\ a$ and $0100\ 00\ 0\ a \leftarrow 0000\ 00\ 0\ s$ transitions of the ν_2 band. Our analysis utilises the Voigt and



speed-dependent Voigt lineshapes (depending on the SNR of the lines), with Rosenkranz line mixing for congested Q-branches. In all, we derive line positions, line intensities, self- and air-broadened Lorentz halfwidths, self- and air-pressure-induced line shifts, the speed dependence of the air-broadened Lorentz halfwidths, and first-order (Rosenkranz) line mixing parameters. Our derived spectroscopic line parameters reproduce the new laboratory measurements substantially better than those from the HITRAN 2020 database. We have revised values for parameters such as line intensities and air-broadened Lorentz halfwidths, in some cases by almost 10 %. For the strongest lines, the total uncertainties of key parameters such as line intensities, and self- and air-broadened Lorentz halfwidths have been considerably improved to $\sim 1\%$. In addition to the measured speed dependence and Rosenkranz line mixing parameters, which we believe are the first reported for the ν_2 band of NH_3 in air, we also determine a range of parameters for the ν_2 band that are not currently in HITRAN, for example self- and air-pressure-induced shifts. We expect these new parameters to provide a more accurate basis for incorporation into atmospheric radiative transfer models to measure NH_3 concentrations from satellite.

6 Acknowledgements

This study was funded as part of the UK Research and Innovation Natural Environment Research Council's support of the National Centre for Earth Observation, grant numbers NE/R016518/1 and NE/Y006216/1. We thank Manfred Birk at Deutsches Zentrum für Luft- und Raumfahrt (DLR) for providing us with the spectroscopic sample cell, although this was later rebuilt due to a breakage.

References

- Ho, P.T.P. and Townes, C.H.: Interstellar Ammonia, *Ann. Rev. Astron. Astrophys.*, 21, 239-270, doi:10.1146/annurev.aa.21.090183.001323, 1983.
- Mâlin, M. Boccaletti, A., Perrot, C., Baudoz, P., Rouan, D., Lagage, P.-O., Waters, R., Güdel, M., Henning, T., Vandenbussche, B., Absil, O., Barrado, D., Charnay, B., Choquet, E., Cossou, C., Danielski, C., Decin, L., Glauser, A.M., Pye, J., Olofsson, G., Glasse, A., Patapis, P., Royer, P., Scheithauer, S., Serabyn, E., Tremblin, P., Whiteford, N., van Dishoeck, E. F., Ostlin, G., Ray, T. P. and Wright, G.: First unambiguous detection of ammonia in the atmosphere of a planetary mass companion with JWST/MIRI coronagraphs, *Astron. Astrophys.*, 693, A315, doi:10.1051/0004-6361/202452695, 2025.
- Irwin, P. G. J., Hill, S. M., Fletcher, L. N., Alexander, C., Rogers, J. H.: Clouds and Ammonia in the Atmospheres of Jupiter and Saturn Determined From a Band-Depth Analysis of VLT/MUSE Observations, *J. Geophys. Res. Planets*, 130, e2024JE008622, doi:10.1029/2024JE008622, 2025.



- 500 Sutton, M. A., Erisman, J. W., Dentener, F., Möller, D.: Ammonia in the environment: from ancient times to the present, *Environ. Pollut.*, 156, 583-604, doi:10.1016/j.envpol.2008.03.013, 2008.
- Behara, S. N., Sharma, M., Aneja, V. P., Balasubramanian, R.: Ammonia in the atmosphere: a review on emission sources, atmospheric chemistry and deposition on terrestrial bodies, *Environ. Sci. Pollut. Control Ser.*, 20, 8092-8131, doi:10.1007/s11356-013-2051-9, 2013.
- 505 Wyer, K. E., Kelleghan, D. B., Blanes-Vidal, V., Schauburger, G., Curran, T. P.: Ammonia emissions from agriculture and their contribution to fine particulate matter: A review of implications for human health, *J. Environ. Manage.*, 323, 116285, doi:10.1016/j.jenvman.2022.116285, 2022.
- Van Damme, M., Clarisse, L., Franco, B., Sutton, M. A., Erisman, J. W., Kruit, R. W., van Zanten, M., Whitburn, S., Hadji-
510 Lazaro, J., Hurtmans, D., Clerbaux, C., Coheur, P.-F.: Global, regional and national trends of atmospheric ammonia derived from a decadal (2008–2018) satellite record, *Environ. Res. Lett.*, 16, 055017, doi:10.1088/1748-9326/abd5e0, 2021.
- Sutton, M. A., Reis, S., Baker, S. M. H., *Atmospheric Ammonia: Detecting Emission Changes and Environmental Impacts*, Springer Netherlands, 2009.
- 515 García-Gómez, H., Garrido, J. L., Vivanco, M. G., Lassaletta, L., Rábago, I., Àvila, A., Tsyro, S., Sánchez, G., González Ortiz, A., González-Fernández, I., Alonso, R.: Nitrogen deposition in Spain: modeled patterns and threatened habitats within the Natura 2000 network, *Sci. Total Environ.*, 485–486, 450-460, doi:10.1016/j.scitotenv.2014.03.112, 2014.
- 520 Brunekreef, B., Harrison, R. M., Künzli, N., Querol, X., Sutton, M. A., Heederik, D. J. J., Sigsgaard, T.: Reducing the health effect of particles from agriculture, *Lancet Respir. Med.*, 3, 831-832, doi:10.1016/S2213-2600(15)00413-0, 2015.
- Megaritis, A. G., Fountoukis, C., Charalampidis, P. E., Pilinis, C., Pandis, S. N.: Response of fine particulate matter concentrations to changes of emissions and temperature in Europe, *Atmos. Chem. Phys.*, 13, 3423-3443, doi:10.5194/acp-13-
525 3423-2013, 2013.
- Thakrar, S. K., Balasubramanian, S., Adams, P. J., Azevedo, I. M. L., Muller, N. Z., Pandis, S. N., Polasky, S., Arden Pope III, C. Robinson, A. L., Apte, J. S., Tessum, C. W., Marshall, J. D., Hill, J. D.: Reducing Mortality from Air Pollution in the United States by Targeting Specific Emission Sources, *Environ. Sci. Technol. Lett.*, 7, 639-645,
530 doi:10.1021/acs.estlett.0c00424, 2020.



Erismann, J. W., Schaap, M.: The need for ammonia abatement with respect to secondary PM reductions in Europe, *Environ. Pollut.*, 129, 159-163, doi:10.1016/j.envpol.2003.08.042, 2004.

535 Behera, S., Sharma, M.: Investigating the potential role of ammonia in ion chemistry of fine particulate matter formation for an urban environment, *Sci. Total Environ.*, 408, 3569-3575, doi:10.1016/j.scitotenv.2010.04.017, 2010.

Bauer, S. E., Tsigaridis, K., Miller, R.: Significant atmospheric aerosol pollution caused by world food cultivation, *Geophys. Res. Lett.*, 43, 5394-5400, doi:10.1002/2016GL068354, 2016.

540

Han, X., Zhu, L., Liu, M., Song, Y., Zhang, M.: Numerical analysis of agricultural emissions impacts on PM_{2.5} in China using a high-resolution ammonia emission inventory, *Atmos. Chem. Phys.*, 20, 9979-9996, doi:10.5194/acp-20-9979-2020, 2020.

545 Ting, Y. C., Young, L. H., Lin, T. H., Tsay, S. C., Chang, K. E., Hsiao, T. C.: Quantifying the impacts of PM_{2.5} constituents and relative humidity on visibility impairment in a suburban area of eastern Asia using long-term in-situ measurements, *Sci. Total Environ.*, 818, 151759, doi:10.1016/j.scitotenv.2021.151759, 2022.

Yang, Y., Zhang, Z., Yang, Y., Wang, Z., Chen, Y., He, H.: Impact of high PM_{2.5} nitrate on visibility in a medium size city of Pearl River Delta, *Atmos Pollut. Res.*, 13, 101592, doi:10.1016/j.apr.2022.101592, 2022.

550

Apte, J. S., Brauer, M., Cohen, A. J., Ezzati, M. Arden Pope III, C.: Ambient PM_{2.5} Reduces Global and Regional Life Expectancy, *Environ. Sci. Technol. Lett.*, 5, 546-551, doi:10.1021/acs.estlett.8b00360, 2018.

555 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367-371, doi:10.1038/nature15371, 2015.

Yu, O., Sheppard, L., Lumley, T., Koenig, J. Q., Shapiro, G. G.: Effects of ambient air pollution on symptoms of asthma in Seattle-area children enrolled in the CAMP study, *Environ. Health Perspect.*, 108, 1209-1214, doi:10.1289/ehp.001081209, 2000.

560

Twigg, M. M. Berkhout, A. J. C., Cowan, N. et al.: Intercomparison of in situ measurements of ambient NH₃: instrument performance and application under field conditions, *Atmos. Meas. Tech.*, 15, 6755-6787, doi:10.5194/amt-15-6755-2022, 2022.



- 565 Clarisse, L., Clerbaux, C., Dentener, F., Hurtmans, D., Coheur, P.-F.: Global ammonia distribution derived from infrared satellite observations, *Nat. Geosci.*, 2, 479-483, doi:10.1038/ngeo551, 2009.
- Van Damme, M., Clarisse, L., Heald, C. L., Hurtmans, D., Ngadi, Y., Clerbaux, C., Dolman, A. J., Erisman, J. W., Coheur, P.-F.: Global distributions, time series and error characterization of atmospheric ammonia (NH₃) from IASI satellite
570 observations, *Atmos. Chem. Phys.*, 14, 2905-2922, doi:10.5194/acp-14-2905-2014, 2014.
- Van Damme, M., Clarisse, L., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Clerbaux, C., Coheur, P.-F.: Industrial and agricultural ammonia point sources exposed, *Nature*, 564, 99-103, doi:10.1038/s41586-018-0747-1, 2018.
- 575 Clarisse, L., Van Damme, M., Gardner, W., Coheur, P.-F., Clerbaux, C., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D.: Atmospheric ammonia (NH₃) emanations from Lake Natron's saline mudflats, *Sci. Rep.*, 9, 4441, doi:10.1038/s41598-019-39935-3, 2019.
- Shephard, M. W., Cady-Pereira, K. E.: Cross-track Infrared Sounder (CrIS) satellite observations of tropospheric ammonia,
580 *Atmos. Meas. Tech.*, 8, 1323-1336, doi:10.5194/amt-8-1323-2015, 2015.
- Warner, J. X., Wei, Z., Strow, L. L., Dickerson, R. R., Nowak, J. B.: The global tropospheric ammonia distribution as seen in the 13-year AIRS measurement record, *Atmos. Chem. Phys.*, 16, 5467-5479, doi:10.5194/acp-16-5467-2016, 2016.
- 585 Warner, J. X., Dickerson, R. R., Wei, Z., Strow, L. L., Wang, Y., Liang, Q.: Increased atmospheric ammonia over the world's major agricultural areas detected from space, *Geophys. Res. Lett.*, 44, 2875-2884, doi:10.1002/2016GL072305, 2017.
- Shephard, M. W., Cady-Pereira, K. E., Luo, M., Henze, D. K., Pinder, R. W., Walker, J. T., Rinsland, C. P., Bash, J. O., Zhu, L., Payne, V. H., Clarisse, L.: TES ammonia retrieval strategy and global observations of the spatial and seasonal variability
590 of ammonia, *Atmos. Chem. Phys.*, 11, 10743-10763, doi:10.5194/acp-11-10743-2011, 2011.
- Down, M. J., Hill, C., Yurchenko, S. N., Tennyson, J., Brown, L. R., Kleiner, I.: Re-analysis of ammonia spectra: Updating the HITRAN 14NH₃ database, *J. Quant. Spectrosc. Radiat. Transf.*, 130, 260-272, doi:10.1016/j.jqsrt.2013.05.027, 2013
- 595 Rothman, L. S., Gordon, I. E., Babikov, Y. et al.: The HITRAN2012 molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transf.*, 130, 4-50, doi:10.1016/j.jqsrt.2013.07.002, 2013.



- Jacquinet-Husson, N., Scott, N. A., Armante, R. et al.: The 2015 edition of the GEISA spectroscopic database, *J. Mol. Spectrosc.*, 327, 31-72, doi:10.1016/j.jms.2016.06.007, 2016.
- Nemtchinov, V., Sung, K., Varanasi, P.: Measurements of line intensities and half-widths in the 10- μ m bands of 14NH₃, *J. Quant. Spectrosc. Radiat. Transf.*, 83, 243-265, doi:10.1016/S0022-4073(02)00354-0. 2004.
- Canè, E., Di Lonardo, G., Fusina, L., Tamassia, F., Predoi-Cross, A.: The $v_2 = 1, 2$ and $v_4 = 1$ bending states of 15NH₃ and their analysis at experimental accuracy, *J. Chem. Phys.*, 150, 194301, doi:10.1063/1.5088751, 2019.
- Canè, E., Di Lonardo, G., Fusina, L., Tamassia, F., Predoi-Cross, A.: Spectroscopic characterization of the $v_2 = 3$ and $v_2 = v_4 = 1$ states for 15NH₃ from high resolution infrared spectra, *J. Quant. Spectrosc. Radiat. Transf.*, 250, 106987, doi:10.1016/j.jqsrt.2020.106987, 2020.
- Gordon, I. E., Rothman, L. S., Hargreaves, R. J. et al.: The HITRAN2020 molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transf.*, 277, 107949, doi:10.1016/j.jqsrt.2021.107949, 2021.
- Yurchenko, S. N.: A theoretical room-temperature line list for 15NH₃, *J. Quant. Spectrosc. Radiat. Transf.*, 152, 28-36, doi:10.1016/j.jqsrt.2014.10.023, 2015.
- Werwein, V., Brunzendorf, J., Li, G., Serdyukov, A., Werhahn, O., Ebert, V.: High-resolution Fourier transform measurements of line strengths in the 0002-0000 main isotopologue band of nitrous oxide, *Appl. Opt.*, 56, E99-E105, doi:10.1364/AO.56.000E99, 2017.
- Devi, V. M., Benner, D. C., Smith, M. A. H., Brown, L. R., Dulick, M.: Absolute intensity measurements of the laser bands near 10 μ m, *J. Quant. Spectrosc. Radiat. Transf.*, 76, 393-410, doi:10.1016/S0022-4073(02)00067-5, 2003.
- Benner, D. C., Rinsland, C. P., Devi, V. M., Smith, M. H., Atkins, D.: A multispectrum nonlinear least squares fitting technique, *J. Quant. Spectrosc. Radiat. Transf.*, 6, 705-721, doi:10.1016/0022-4073(95)00015-D, 1995.
- Benner, D. C., Devi, V. M., Sung, K., Brown, L. R., Miller, C. E., Payne, V. H., Drouin, B. J., Yu, S., Crawford, T. J., Mantz, A. W., Smith, M. H., Gamache, R. R.: Line parameters including temperature dependences of air- and self-broadened line shapes of 12C16O₂: 2.06- μ m region, *J. Mol. Spectrosc.*, 326, 21-47, doi:10.1016/j.jms.2016.02.012, 2016.



- Rosenkranz, P. W., Interference coefficients for overlapping oxygen lines in air, *J. Quant. Spectrosc. Radiat. Transf.*, 39, 287-297, doi:10.1016/0022-4073(88)90004-0, 1988.
- 635 Letchworth, K. L. and Benner, D. C.: Rapid and accurate calculation of the Voigt function, *J. Quant. Spectrosc. Radiat. Transf.*, 107, 173-192, doi:10.1016/j.jqsrt.2007.01.052, 2007.
- Ciurylo, R., Shapes of pressure- and Doppler-broadened spectral lines in the core and near wings, *Phys. Rev. A*, 58, 1029, doi:10.1103/PhysRevA.58.1029, 1998.
- 640 Joint Committee for Guides in Metrology (JCGM), 100:2008 Evaluation of measurement data - Guide to the expression of uncertainty in measurement, doi:10.59161/JCGM100-2008E, 2008.
- Maki, A. G., Wells, J. S.: Wavenumber calibration tables from heterodyne frequency measurements. NIST Special Publication 821, National Institute of Standard and Technology, Gaithersburg, 1991.
- 645 Aroui, H., Laribi, H., Orphal, J., Chelin, P.: Self-broadening, self-shift and self-mixing in the ν_2 , $2\nu_2$ and ν_4 bands of NH_3 , *J. Quant. Spectrosc. Radiat. Transf.*, 110, 2037-2059, doi:10.1016/j.jqsrt.2009.05.002, 2009.
- 650 Guinet, M., Jeseck, P., Mondelain, D., Pepin, I., Janssen, C., Camy-Peyret, C., Mandin, J. Y.: Absolute measurements of intensities, positions and self-broadening coefficients of R branch transitions in the ν_2 band of ammonia, *J. Quant. Spectrosc. Radiat. Transf.*, 112, 1950-1960, doi:10.1016/j.jqsrt.2011.03.015, 2011.
- Clar, H.-J., Schieder, R., Winnewisser, G., Yamada, K. M.T., Pressure broadening and lineshifts in the ν_2 band of NH_3 , *J. Mol. Struct.*, 190, 447-456, doi:10.1016/0022-2860(88)80303-X, 1988.
- 655 Ibrahimi, M., Babay, A., Lemoine, B., Rohart, F., Pressure-Induced Frequency Lineshifts in the ν_2 Band of Ammonia: An Experimental Test of the Rydberg–Ritz Principle, *J. Mol. Spectrosc.*, 193, 277-284, doi:10.1006/jmsp.1998.7749, 1999.