

# Growing role of secondary organic aerosol in the North China Plain from 2014 to 2024

Chunshui Lin<sup>1</sup>, Ru-Jin Huang<sup>1,2\*</sup>, Jing Duan<sup>1</sup>, Jing Qu<sup>1</sup>, Jiahua Liu<sup>1</sup>, Yi Liu<sup>1</sup>, Yan Luo<sup>1</sup>, Wei Huang<sup>1</sup>, Wei Xu<sup>3</sup>, Yanan Zhan<sup>1,2</sup>, Zhitao Liu<sup>4,5</sup>, Sihao Liu<sup>4,5</sup>, Qingshuang Zhang<sup>6</sup>, Quan Liu<sup>7</sup>,  
5 Zirui Liu<sup>8</sup>, Shengrong Lou<sup>9,10</sup>, Huinan Yang<sup>10</sup>, Dan Dan Huang<sup>9</sup>, Cheng Huang<sup>9,11</sup>, and Hongli Wang<sup>9</sup>

<sup>1</sup>State Key Laboratory of Loess Science, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710061, China

<sup>2</sup>Institute of Global Environmental Change, School of Human Settlements and Civil Engineering, Xi'an Jiaotong University, Xi'an 710049, China

<sup>3</sup>Center for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, China

<sup>4</sup>Key Laboratory of Meteorology and Ecological Environment of Hebei Province, Shijiazhuang 050021, China

<sup>5</sup>Weather Modification Office of Hebei Province, Shijiazhuang 052100, China

<sup>6</sup>Shandong Province Dezhou Ecological Environment Monitoring Center, Dezhou 200235, China

<sup>7</sup>State Key Laboratory of Severe Weather & Key Laboratory of Atmospheric Chemistry of CMA, Chinese Academy of Meteorological Sciences, Beijing 100081, China

<sup>8</sup>State Key Laboratory of Atmospheric Environment and Extreme Meteorology, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

<sup>9</sup>Key Laboratory of Formation and Prevention of Urban Air Pollution Complex, Ministry of Ecology and Environment, Shanghai Academy of Environmental Sciences, Shanghai 200233, China

<sup>10</sup>School of Energy and Power Engineering, University of Shanghai for Science and Technology, Shanghai 200093, China

<sup>11</sup>Shanghai Environmental Monitoring Center, Shanghai 200235, China

25 *Correspondence to:* Ru-Jin Huang (rujin.huang@ieecas.cn)

**Abstract.** Since the Clean Air Act was implemented in 2013, China has witnessed a reduction of over 50% in the annual average concentration of fine particulate matter (PM<sub>2.5</sub>). Despite these emission cuts, the formation mechanism of secondary organic aerosols (SOA), a crucial constituent of PM<sub>2.5</sub>, remains inadequately understood. In this study, we performed a model-assisted analysis of field sampling data  
30 collected in Shijiazhuang. The results show that, compared to 2014, the contribution of SOA to the total organics (from 27% in 2014 to 87% to in 2024) exceeded that of primary organic aerosol (POA) during the winter haze in 2024. Although the model underestimated the measured SOA levels, incorporating the transformation of transported POA into SOA under high relative humidity (RH) conditions helped  
35 bridge the gap between model predictions and on-site measurements. The increase in SOA contribution occurred amidst large emission reductions, which accounted for 70% of the decline in POA levels, while meteorological factors contributed an additional 10%. Increased contribution of SOA was also found in

other North China Plains areas, which underscores the pressing necessity for coordinated regional initiatives to effectively mitigate SOA levels across the NCP, thereby tackling the transboundary nature of air pollution.

## 40 **1 Introduction**

Ambient PM<sub>2.5</sub> pollution is a critical environmental issue in China, contributing to millions of premature deaths annually (Zhao et al., 2018; Zhang et al., 2021; Chen et al., 2024). Since the implementation of the Clean Air Act in 2013, there have been large reductions in PM<sub>2.5</sub> concentrations nationwide (Zhang et al., 2019; Geng et al., 2021). However, improvements have slowed in recent years, e.g., from 2018 to 2020, compared to the initial surge  
45 of progress between 2013 and 2017 (Geng et al., 2024). The North China Plain (NCP), one of the country's most polluted areas, continues to experience severe haze episodes with elevated PM<sub>2.5</sub> levels, especially during winter months (Huang et al., 2014; Daellenbach et al., 2024). This persistent pollution usually stems from a complex interplay of local emission sources, transboundary pollutant transport, secondary formation, specific topographical features, and unfavorable meteorological conditions (Chang et al., 2022; Nie et al., 2022).

50 Organic aerosols (OA) represent a large portion (20-90%) of PM<sub>2.5</sub> in the atmosphere, arising from both primary and secondary processes (Jimenez et al., 2009; Lin et al., 2018; Huang et al., 2023; Chen et al., 2024). Recent research has increasingly focused on the budgets, formation pathways, and photochemical evolution of SOA due to their impacts on climate and human health (Chen et al., 2024; Daellenbach et al., 2024). SOA formation involves precursor VOCs released from both natural sources,  
55 such as forests and wildfires, and anthropogenic sources, including biomass burning and vehicular emissions (Chang et al., 2022). Despite this understanding, many SOA precursors remain unidentified, leading to uncertainties in global SOA budget estimations, as numerous climate models do not account for these unknowns (Shrivastava et al., 2017; Chang et al., 2022). Therefore, a better understanding of SOA formation, particularly in the context of emission reductions, is essential for evaluating their future  
60 implications for climate and human health (Chen et al., 2024).

In the capital city of Beijing, a recent study indicates that SOA primarily originates from solid-fuel combustion and aqueous processes involving multiphase chemistry with aromatic compounds (Daellenbach et al., 2024), with the most severe haze events linked to the transport from south of Beijing. Kuang et al. (2020) explored the effects of gas-phase and aqueous-phase photochemical processes on  
65 SOA formation, revealing that photochemical aqueous-phase processes contributed largely to daytime oxygenated organic aerosol (OOA) formation during winter in a sampling site near Beijing. While much

of the existing research has focused on Beijing and nearby areas, studies in Shijiazhuang (SJZ) are relatively scarce (Huang et al., 2019; Lin et al., 2020). As the capital of Hebei Province and one of the most polluted cities in China, particularly during the winter heating season, Shijiazhuang presents a critical context for further investigation(Huang et al., 2019; Lin et al., 2020). Huang et al. (2019) utilized a high-time resolution Aerosol Chemical Speciation Monitor (ACSM) and found that in wintertime Shijiazhuang 2014, submicron organic aerosol was primarily sourced from direct emissions without extensive atmospheric aging, indicating opportunities for improving air quality through emission reductions. Moreover, during pollution events with high relative humidity (RH), SOA prevailed due to intensified aqueous-phase chemistry highlighting the critical role of meteorological factors in PM pollution in this heavily polluted city (Huang et al., 2019).

In this study, we deployed a Soot Particle (SP) Long Time of Flight (LToF) Aerosol Mass Spectrometer (AMS) (Onasch et al., 2012) in Shijiazhuang from January to March 2024. We conducted a model-assisted analysis of variations in OA sources and formation, comparing data to 2014 levels(Huang et al., 2019). Our objectives were: (1) to elucidate the chemical mechanisms underlying SOA formation in light of emission reductions, and (2) to evaluate the relative contribution of emission reductions versus meteorological factors by comparing measurements and simulations from 2014 and 2024.

## 2 Method

### 2.1 Field Sampling

Shijiazhuang (SJZ), the capital of Hebei Province, is located approximately 270 km southwest of Beijing (Huang et al., 2019; Lin et al., 2020). In 2024, an SP-LToF-AMS (laser off mode) was deployed from January to March in the Luan Cheng district of southern SJZ. The sampling site was selected as a suburban background location, without strong nearby emission sources.

The design of the SP-LToF-AMS is detailed in Onasch et al. (2012), and its setup followed our previous studies (Lin et al., 2021, 2022). Briefly, ambient air was drawn through a stainless-steel tube via a PM<sub>2.5</sub> cyclone at a consistent flow rate of 5 L min<sup>-1</sup>. After being dried with a Nafion dryer (MD-700-24S, Perma Pure, Inc.), the sampled aerosol was focused into a particle beam using a PM<sub>2.5</sub> aerodynamic lens, then vaporized by a thermal standard vaporizer at approximately 600 °C and ionized through electron ionization (70 eV) to generate positive fragments. The resulting ion fragments were detected and analyzed by the LToF mass spectrometer.

Ionization efficiency (IE) and relative ionization efficiency (RIE) were calibrated using pure NH<sub>4</sub>NO<sub>3</sub> and (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> in the 300-350 nm range. Data acquired from the SP-LToF AMS were analyzed using SQUIRREL (version 1.61D)

and PIKA (1.21D) software, coded in Igor Pro 6.37 (WaveMetrics). To identify OA sources, positive matrix factorization (PMF) combined with multilinear engine (ME-2) analysis was performed on the high-mass spectra of OA. The data and error matrices for OA mass spectra in the m/z range of 12-120 were preprocessed, removing ions with a signal-to-noise ratio (S/N) < 0.2 and down-weighting those with 0.2 < S/N < 3 by a factor of 2 (Lin et al., 2018; Lin et al., 2022; Duan et al., 2024). An unconstrained PMF analysis was then conducted to explore a range of solutions and determine the potential number and types of OA factors. Subsequently, an ME-2 approach with a-value constraints was employed to obtain the most interpretable profiles and environmentally meaningful results. As a result, the time series of AMS + BC is strongly correlated with total PM<sub>2.5</sub> with slope of the linear fit close to 1, and the OA explained by PMF closely aligns with the measured OA (Fig. S1).

## 2.2 Model Analysis

The Community Multiscale Air Quality (CMAQ) model (version 5.4) (Pye et al., 2023) was employed to simulate ambient variations in emission reductions and the meteorological impacts on ambient aerosol on a regional scale in January 2024, compared to January 2014. The CMAQ model was configured with the aero7 aerosol module and the CB6r5 gas-phase mechanism. To generate the meteorological fields required for the CMAQ simulations, the Weather Research and Forecasting (WRF) model (version 4.4) was utilized with the National Centers for Environmental Prediction (NCEP) Final (FNL) dataset.

The emission inventory for 2014 was derived from the Multiresolution Emission Inventory for China (MEIC) (<http://meicmodel.org.cn>; last access: May 1, 2025) (Geng et al., 2024). Since the emission inventories are not up-to-date for 2024, the MEIC data were downscaled from the 2020 MEIC version. A scaling factor of 0.5 was applied to enable more accurate comparisons between model outputs and 2024 measurements (Fig. S2). The modeling period covered January 11 to 31 for both years, coinciding with the availability of ACSM/AMS data in January. The first three days were designated as a spin-up phase to stabilize the model, ensuring that subsequent data accurately reflected steady-state atmospheric conditions. Model performance metrics—including Mean Bias (MB), Normalized Mean Bias (NMB), Root Mean Square Error (RMSE), and correlation coefficient (R)—indicated good agreement between modeled and measured PM<sub>2.5</sub> levels (Fig. S2).

High relative humidity (RH) increases the availability of aerosol liquid water (ALW), thereby enhancing the uptake of both organic and inorganic gases. These gases are oxidized in aerosol water to form secondary aerosols (Zheng et al., 2015; Mcfall et al., 2020). High RH also promotes the uptake of oxidants, including hydroxyl (OH) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), which can generate OH radicals via Fenton chemistry (Nguyen et al., 2013). These oxidants oxidize POA and form SOA. To characterize the RH-mediated conversion of POA to SOA, we introduce a simplified surrogate equation for SOA formation:



In this equation, atmospheric water vapor ( $\text{H}_2\text{O}$ ) is linked to RH calculations. Higher RH (i.e., increased  $\text{H}_2\text{O}$ ) facilitates the apparent conversion of more POA into SOA. The conversion rate was tested until a closure was achieved between measured and simulated SOA. As a result, the model SOA and POA tracked the observed ones with a moderate correlation coefficient (Fig. S3). The diurnal pattern of SOA was simulated relatively well in 2024, whereas the model missed the midday peak of SOA in 2014, suggesting that daytime SOA formation was more important in 2014 than in 2024 (Fig. S4). Different scenarios were designed (Table S1). This simplified calculation serves as an empirical approach to account for the underestimated SOA formed during the aging of POA. Further research is needed to determine the effective uptake coefficients of precursor VOCs and oxidants, as well as the dynamics of POA aging.

### 3 Results

#### 3.1 Increased Secondary Aerosol Contribution in 2024 vs 2014

Figure 1a shows the variations of the major components of NR-PM (i.e., organics, sulfate, nitrate, ammonium, and chloride) at SJZ in 2014, while Figure 1b presents the data for 2024, categorizing non-refractory PM into deciles, ranging from the least polluted 10th percentile to the most polluted 10th percentile. As NR-PM increases from the lowest 10th percentile to the highest 10th percentile, organics remain the largest component, with mean concentrations rising from  $16.8 \mu\text{g m}^{-3}$  to  $206.0 \mu\text{g m}^{-3}$  in 2014, and from  $2.9 \mu\text{g m}^{-3}$  to  $45.4 \mu\text{g m}^{-3}$  in 2024. In 2014, sulfate was the second largest component, with concentrations rising from  $4.0 \mu\text{g m}^{-3}$  to  $90.3 \mu\text{g m}^{-3}$ . In 2024, however, sulfate became less prominent than nitrate, with mean concentrations increasing from  $1.3 \mu\text{g m}^{-3}$  to  $21.7 \mu\text{g m}^{-3}$ . In 2024, nitrate emerged as the second dominant component, rising from  $1.0 \mu\text{g m}^{-3}$  to  $33.6 \mu\text{g m}^{-3}$ , although its concentrations were still slightly lower than in 2014, which saw an increase from  $3.6 \mu\text{g m}^{-3}$  to  $43.5 \mu\text{g m}^{-3}$ .

The dominance of organic aerosol is linked to direct emissions from various combustion sources, such as biomass burning and coal combustion, as well as the formation of SOA. Figure 1c shows that non-cooking primary organic aerosol (ncPOA)—the sum of biomass burning organic aerosol (BBOA), coal combustion organic aerosol (CCOA), and hydrocarbon-like organic aerosol (HOA)—increased from  $7.1 \mu\text{g m}^{-3}$  to  $115.0 \mu\text{g m}^{-3}$  in 2014, while oxygenated organic aerosol (OOA, a surrogate for SOA) rose from  $5.2 \mu\text{g m}^{-3}$  to  $44.1 \mu\text{g m}^{-3}$ . In contrast, in 2024, OOA became more dominant than ncPOA, with OOA increasing from  $3.0 \mu\text{g m}^{-3}$  to  $47.5 \mu\text{g m}^{-3}$  (Fig. 1d), while ncPOA rose from less than  $1 \mu\text{g m}^{-3}$  to  $6.7 \mu\text{g m}^{-3}$ . This shift in dominance of OOA, a surrogate of SOA, underscores the increasing role of secondary aerosol formation processes in affecting air quality and public health, highlighting the need for enhanced monitoring, adaptive policy measures to control emissions, and interdisciplinary research efforts to tackle the emerging challenges posed by organic aerosols. Moreover, changes in PM components are known to drive differences in aerosol liquid water content (Kuang et al., 2020), and such changes may also alter the formation pathways of SOA.

In contrast to organics, the reduction in inorganics was relatively straightforward. Specifically, the decrease in sulfate is linked to the reduction of its precursor, SO<sub>2</sub>, and its conversion from SO<sub>2</sub> to sulfate. During the sampling period in 2014, the measured SO<sub>2</sub> concentration was 150 ± 84.5 μg m<sup>-3</sup> (± one standard deviation), while in 2024, it dropped to 6.5 ± 7.2 μg m<sup>-3</sup>. This represents an average reduction of 95.6% in SO<sub>2</sub> concentrations from 2014 to 165 2024. For measured sulfate, the average reduction was less pronounced than that of its precursor, at 82%. This indicates higher conversion efficiency from SO<sub>2</sub> to sulfate in 2024, resulting in a higher sulfur oxidation ratio (F<sub>SO<sub>4</sub></sub>) of 0.42 compared to 0.15 in 2014. It's important to note that the observed F<sub>SO<sub>4</sub></sub> may represent an upper limit, as regional transport and background sulfate levels were not factored into the calculation. In contrast to sulfate, nitrate experienced a smaller reduction, resulting in an increased nitrate fraction in 2024 compared to 2014. This change 170 is partly linked to NO<sub>x</sub> emissions, which have seen less reduction than SO<sub>2</sub> (Fig. S5). The increase in oxidizing capacity may convert more NO<sub>x</sub> into nitrate in the presence of abundant ammonia in recent years (Zang et al., 2022). Ammonium (NH<sub>4</sub>) is associated with ammonia emissions, which are converted to ammonium in the presence of sulfate and nitrate. Given the abundant ammonia emissions over the years (Liu et al., 2019), ammonium concentrations were influenced more by sulfate and nitrate than by ammonia itself. Chloride is connected to 175 biomass burning and coal combustion. Therefore, the reduction in ncPOA likely reflects a decrease in the corresponding emission strength, leading to lower chloride emissions.

On average, the measured NR-PM concentrations showed a reduction of 75% over the sampling period from 2014 to 2024, with NR-PM<sub>1</sub> averaging 188.0 ± 111.0 μg m<sup>-3</sup> in 2014 and NR-PM<sub>2.5</sub> averaging 46.85 ± 35.8 μg m<sup>-3</sup> in 2024. Considering the differences in size cuts (i.e., PM<sub>1</sub> from ACSM in 2014 and PM<sub>2.5</sub> from AMS in 2024) and the 180 sampling sites between the two campaigns (Huang et al., 2019), a direct comparison of the two campaigns may introduce limitations. However, given the strong time-series correlation in bulk PM<sub>2.5</sub> concentrations across cities in the NCP (as discussed in Section 3.4), this comparison can still reflect an overall trend.

Comparing Figures 1a and 1b, we observed that all components were reduced to varying extents across different pollution levels, from the cleanest 10<sup>th</sup> percentile to the most polluted 10<sup>th</sup> percentile. On average, organic 185 compounds were reduced by 79% in 2024 compared to 2014. During the most polluted 10% of the sampling period, organics saw a similar reduction of 78% compared to 2014. Sulfate experienced an average reduction of 82%, with a 76% reduction during the most polluted 10%. Ammonium was reduced by an average of 72.1%, with a 79% reduction in the most polluted period. Chloride also saw an average reduction of 76%, though during the most polluted period, the reduction was only 67%. In contrast, nitrate experienced the least reduction, with an average 190 decrease of 50% and only a 23% reduction during the most polluted period. This indicates that all major components saw similar reductions in 2024, except for nitrate, which became more important in driving pollution levels compared to 2014. Previously dominated by direct emissions (as indicated by high levels of ncPOA in Fig. 1c), organics are now increasingly influenced by secondary production. On average, ncPOA was reduced by 93% in 2024, while SOA was reduced by 45%.

### 3.2 Formation and Transport of Secondary Organic Aerosol

During the polluted periods, the shift in dominance from ncPOA to SOA in 2024 compared to 2014 indicates that secondary formation processes are becoming more important than primary emissions (Fig. 1). Plotting SOA against relative humidity (RH) reveals that as RH increases, SOA levels also rise (Fig. 2). The mean SOA in different RH brackets increased from 15.6  $\mu\text{g m}^{-3}$  to 27.9  $\mu\text{g m}^{-3}$  in 2024, while in 2014, it rose from 21.9  $\mu\text{g m}^{-3}$  to 28.9  $\mu\text{g m}^{-3}$  as RH increased from 35% to 95% (Fig. 2). The positive correlation between SOA and RH suggests that RH modulates aqueous phase reactions, facilitating SOA formation, consistent with previous studies indicating enhanced SOA production at high RH (Huang et al., 2019; Lin et al., 2020).

In contrast to SOA, ncPOA reached a plateau at high RH, with mean concentrations ranging from 58.4 to 60.9  $\mu\text{g m}^{-3}$  at RH levels above 55% in 2014, and from 4.4 to 5.0  $\mu\text{g m}^{-3}$  in 2024. While the continuous increase in SOA at high RH suggests enhanced SOA formation through heterogeneous reactions, the plateau in ncPOA concentrations at high RH may indicate either that ncPOA emissions ceased at high RH (which is unlikely) or that ncPOA was converted to SOA through aging processes. This conversion likely requires an oxidizing environment, and the increasing ozone ( $\text{O}_3$ ) concentrations observed in recent winters may reflect a more oxidizing atmosphere (Fig. S6). Consequently, more ncPOA is likely converted to SOA during transport. The aging of POA into SOA aligns with findings from winter case studies in Beijing, suggesting that ncPOA aging may partly explain the recent increase in SOA levels during the case study in Beijing (Wang et al., 2021). Additionally, the oxidation of semi-volatile and intermediate volatile organic compounds (SVOCs and IVOCs) may have been enhanced due to increased oxidation capacity (Huang et al., 2025).

In 2014, ncPOA was relatively well simulated (Fig. 3a). However, in 2024, ncPOA concentrations were overestimated by more than 100% (Fig. 3b). In contrast, SOA levels were underestimated by 40-60%. The base run considered the default SOA formation mechanism, which includes ncPOA evaporation and subsequent aging driven by hydroxyl (OH) radicals. The underestimation of SOA can be partly attributed to insufficient consideration of the conversion of ncPOA at high RH levels, which may not necessarily be driven by OH radicals. This factor has likely become more important in recent years due to a more oxidizing environment. The overestimation of ncPOA in 2024 suggests that a greater fraction of ncPOA should be converted to SOA.

In this study, emissions from SJZ set to zero for January 2014 and 2024 to study the local vs regional transport of ncPOA, which can be converted into SOA during transport. The differences between simulations with and without SJZ emissions were treated as the emissions impact. CMAQ simulations indicate that, on average, 39% of ncPOA in 2014 was from regional transport, while 61% originated from local emissions in SJZ. In 2024, 35% was from local emissions and 65% from regional transport. However, during pollution episodes ( $\text{PM}_{2.5} > 75 \mu\text{g m}^{-3}$ ), approximately 50% of ncPOA was sourced from regional transport in both 2014 and 2024 (Fig. S7). Given the large

contribution from regional ncPOA, accounting for the aging of transported ncPOA into SOA could help resolve these model-measurement discrepancies.

230 By adjusting the model with RH-related parameters, we can improve SOA simulations and narrow the gap between simulated and observed SOA levels (see Methods). After these adjustments, SOA concentrations in NCP were elevated to over  $20 \mu\text{g m}^{-3}$  in 2014 (Fig. 3c). In 2024, SOA levels were reduced by about 50%, which is a smaller reduction compared to the more than 90% decrease observed in ncPOA (Fig. 3b and 3d). In 2024, SOA constituted approximately 60% of the total OA for the adjusted model (Table S2) and 80% of the total OA for the observed OA  
235 (Fig. 3b), highlighting its dominance over POA. As a result (Fig. 3), the aging of POA under high RH, which leads to the formation of SOA, contributed to 30-31% of the total SOA in 2014-2024. Additionally, SOA generated from the oxidation of intermediate and semi volatile compounds comprised 23-37% of the total. Aromatic precursor oxidation accounted for 8-11% of the SOA, while SOA resulting from aqueous phase chemistry involving aldehydes contributed 23-34%. Note that biogenic SOA was not considered in the CMAQ configuration. However, given a  
240 reduced emission of biogenic VOCs in winter, SOA formation potential from biogenic SOA was less important in winter than in summer (Dou et al., 2025), and may have a minor impact on the overall contribution of the anthropogenic sources.

### 3.3 Rising contribution of SOA to the total OA Amidst Emission Reduction

As discussed in Section 3.1, total OA were reduced by an average of 79%. However, different types of organic  
245 aerosols were reduced to varying extents, with ncPOA decreasing by 93% and OOA by 45% from 2014 to 2024. POA is directly emitted from emission sources. By utilizing the emission inventory while keeping meteorological parameters constant, we can simulate how emissions reductions contributed to this decrease in 2024 compared to 2014 (Fig. 4; Table S1).

Figure 4a shows the mean POA concentrations using the 2014 emission inventory alongside the 2024  
250 meteorological fields, while Figure 4b presents mean POA concentrations with the 2024 emission inventory and the 2024 meteorological fields. The differences between these two scenarios highlight the impact of emissions reduction (Figure 4c). In the 2014 emissions scenario, POA concentrations were elevated across the NCP and the Fenwei Plain (FWP), with average concentrations around  $50 \mu\text{g m}^{-3}$ . Notably, POA concentrations in Shijiazhuang (SJZ), Beijing (BJ), Tianjin (TJ), Baoding (BD), Xingtai (XT), Handan (HD), and Xi'an (XA) were comparable,  
255 indicating widespread regional air pollution affecting multiple cities with hundreds of millions of residents.

In contrast, the 2024 emissions scenario showed a substantial reduction in POA concentrations, decreasing by up to 70%, with mean concentrations ranging from 1 to  $10 \mu\text{g m}^{-3}$  in the North China Plain (NCP) and Fenwei Plain (FWP). This reduction indicates that emission decrease accounts for approximately 30 to  $40 \mu\text{g m}^{-3}$  of the decline in POA levels in these regions, representing about 70% of the overall reduction. Thus, the modeled decrease in  
260 emission strength explains the majority of the observed reduction.

Figure S8 shows the impact of meteorological variations on modeled POA concentrations in January 2024 compared to 2014, accounting for about a 10% reduction attributed to these variations. In Figure S8, we compared mean POA concentrations using 2014 meteorological conditions alongside 2024 emissions with those under 2024 meteorological conditions and the same emissions. The differences between these two scenarios highlight the influence of meteorological factors. The results indicate that mean POA concentrations under the 2014 meteorological conditions were higher than those under the 2024 conditions. Overall, the emission reduction explained approximately 70% of POA in the NCP, with meteorological conditions contributing to an additional reduction of around 10%. The remaining unexplained reduction of 10-20% in POA is likely due to the conversion of POA to SOA, as discussed in Section 2.2.

### 270 3.4 Comparable Reduction in other Cities

Starting in 2013, China initiated two phases of clean air actions: Phase 1 (2013–2017) and Phase 2 (2018–2020). The years 2021-2024 are considered Phase 3, following these initial phases. Figure 5 illustrates the evolution of PM<sub>2.5</sub> during the heating periods across Phases 1, 2, and 3 for the seven cities. From Phase 1 to Phase 2, reductions were greater in the six cities of the NCP, with mean concentrations decreasing from 58-106  $\mu\text{g m}^{-3}$  in Phase 1 to 38-73  $\mu\text{g m}^{-3}$  in Phase 2. However, from Phase 2 to Phase 3, the reduction was less pronounced, with some cities experiencing a plateau or even a slight rebound, resulting in PM<sub>2.5</sub> concentrations around 30-52  $\mu\text{g m}^{-3}$ . An overall reduction in regional PM<sub>2.5</sub> has been observed for major cities in the NCP and FWP over the last decade (Fig. 5). This trend is consistent with the modeling results that compare 2024 to 2014 (Fig. 3), highlighting a regional decrease in PM<sub>2.5</sub> concentrations attributed to emission reductions over the past decade.

280 Xi'an (XA), the largest city in the FWP, exhibited different PM<sub>2.5</sub> variations compared to cities in the NCP. At the end of Phase 1, PM<sub>2.5</sub> concentrations increased slightly from 58 to 74  $\mu\text{g m}^{-3}$ , followed by a reduction in Phase 2 (from 62 to 51  $\mu\text{g m}^{-3}$ ), and a rebound during Phase 3 (from 43 to 50  $\mu\text{g m}^{-3}$ ). The differences in PM<sub>2.5</sub> variations between XA and other NCP cities are linked to emissions patterns in the Guanzhong Basin, which is more isolated from NCP emissions. Figure 5a illustrates the wind fields during haze periods at ground level. The wind patterns reveal anticyclonic flow at low speeds in NCP, while the winds in FWP were more stagnant, indicating that emissions were confined within the basin rather than being influenced by external sources. This observation aligns with previous studies in Xi'an that often saw higher pollution levels than in NCP (Zhong et al., 2020; Lin et al., 2023), whereas pollution in NCP has been primarily associated with southerly winds transporting emissions from nearby sources (Daellenbach et al., 2024). Figure 5c demonstrates that from Phase 1 to Phase 3, the correlation between PM<sub>2.5</sub> time series in NCP cities remain similarly elevated if not slightly increased. For example, the correlation coefficients (r) between BJ and TJ, BD, and SJZ were 0.65, 0.58, and 0.63 during Phase 1, 0.63, 0.60, and 0.57 during Phase 2, and 0.70, 0.69, and 0.63 in Phase 3. The correlation between SJZ and XT and HD was 0.80 and 0.74 in Phase 1, 0.83 and 0.72 in Phase 2, and remained 0.83 and 0.74 in Phase 3. In contrast, the correlation with

XA and other cities remained low, ranging from 0.39 to 0.56 in Phase 1, 0.28 to 0.56 in Phase 2, and 0.22 to 0.50  
295 in Phase 3. Thus, the high correlations among PM<sub>2.5</sub> levels in NCP cities highlight the increased importance of  
regional transport, while the lower correlations with XA indicate isolated emission sources in the FWP.

Removing emissions from SJZ alone was insufficient to reduce PM<sub>2.5</sub> levels to safe standards of less than 25 µg  
m<sup>-3</sup> (Fig. S9). This is largely due to the fact that approximately half of the PM<sub>2.5</sub> in SJZ was transported from outside  
the city (Fig. S9). However, the transport of pollutants appears similar in both 2014 and 2024. During pollution  
300 episodes (PM<sub>2.5</sub> > 75 µg m<sup>-3</sup>), with around 56-60% of PM<sub>2.5</sub> transported from regional sources and the remaining  
40-44% attributed to local emissions (Fig. S9). During relatively clean periods (PM<sub>2.5</sub> < 25 µg m<sup>-3</sup>), approximately  
46-51% of PM<sub>2.5</sub> originates from local sources, while 49-54% is derived from regional transport, highlighting the  
need for a coordinated regional effort to further reduce PM<sub>2.5</sub> levels across the NCP.

Figure S10 shows the contributions of five key sectors (in µg m<sup>-3</sup>) to the total PM<sub>2.5</sub> in January 2014 and 2024:  
305 power generation, industry (including industrial boilers, cement production, iron production, and other processes),  
residential burning (covering domestic biomass and fossil fuel burning), transportation (both on-road and off-road  
vehicles), and agriculture. In both years, residential domestic burning was the primary contributor to PM<sub>2.5</sub>,  
accounting for approximately 51% of PM<sub>2.5</sub> in 2014 and 48% in 2024. The industrial sector followed as the second-  
largest contributor, representing about 34% of PM<sub>2.5</sub> in 2014 and 33% in 2024. Combined, traffic, power generation,  
310 and agriculture accounted for 15% of total PM<sub>2.5</sub> in 2014, with a similar level of 18% in 2024. When comparing  
contributions between 2024 and 2014, the domestic sector experienced a large reduction, decreasing from 68.5  
µg m<sup>-3</sup> in 2014 to 22.4 µg m<sup>-3</sup> in 2024—a reduction of 67%. The industrial sector also saw a notable decrease of  
65%, from 44.7 µg m<sup>-3</sup> in 2014 to 15.6 µg m<sup>-3</sup> in 2024. Other sectors contributed minimally to PM<sub>2.5</sub>, but the  
reductions in these areas exhibited greater uncertainty, as indicated by larger error bars. The dominant contribution  
315 of residential and industrial emissions aligns with modelling results from other regions in other BTH area (Chang  
et al., 2019). This highlights the importance of targeting residential burning and industry in future mitigation  
strategies. Given the similar temporal and spatial distributions of sub-sectors within each major sector, it is feasible  
to estimate contributions from more detailed sectors based on specific emission rates. For instance, in SJZ,  
domestic biomass burning accounted for 57% of the primary PM<sub>2.5</sub> emissions, with the remaining 43% from fossil  
320 fuel combustion. Domestic burning SO<sub>2</sub>, a precursor of sulfate, was predominantly from domestic fossil fuel burning.  
In the industrial sector, industrial boilers, cement production, and steel manufacturing were responsible for 82% of  
the primary PM<sub>2.5</sub> emissions, with the rest coming from other industrial activities. Additionally, pollutants from  
domestic sources are primarily emitted at ground level, resulting in a larger impact compared to emissions from  
elevated industrial sources.

## 325 **4 Conclusion**

In conclusion, organics remain the most important component of non-refractory PM (NR-PM) in 2024. Emission reductions account for 70% of the decrease in POA compared to 2014, with meteorological conditions contributing an additional 10%. Haze pollution in 2024 was characterized by high RH, with nitrate levels exceeding those of sulfate and SOA levels surpassing POA levels—a reversal of the situation in 2014, in the context of increasing oxidizing capacity as summarized in Fig. 6. Therefore, further reductions in VOCs and NO<sub>x</sub> could lead to additional decreases in overall aerosol pollution levels. The transport of pollutants appears similar in both 2014 and 2024 during pollution episodes, with around 80% of PM<sub>2.5</sub> transporting from external sources and the remaining 20% attributing to local emissions, highlighting the need for coordinated regional efforts to further reduce PM<sub>2.5</sub> levels across the NCP. The positive correlation between SOA and RH suggests that RH modulates aqueous phase reactions, facilitating SOA formation in both 2014 and 2024. Accounting for the aging of transported ncPOA into SOA could help reconcile these model-measurement discrepancies. This factor is likely becoming increasingly important given the rising levels of oxidants.

### **Data availability**

All data used in the study are available from the corresponding author upon request.

### 340 **Code availability**

The code for CMAQ model is available at <https://github.com/USEPA/CMAQ>.

### **Supporting Information**

Supplementary Figures (Fig S1-S10) and Table (Table S1 and S2)

### **Corresponding author**

345 Ru-Jin Huang ([rujin.huang@ieecas.cn](mailto:rujin.huang@ieecas.cn))

### **Author Contributions**

C.L. and R.J.H. Designed the study, C.L. wrote the main manuscript text, C.L. prepared all figures. C.L., J.D., J.Q., J.L., Yi L., Yan L., W.H., W.X., and Y.Z, conducted the field sampling, S.L., Q.Z, Q.L., Z.L., D.D.H., C.H., and H.W

provided guidance on the research design and contributed to the interpretation of results, R.J.H. coordinated the  
350 project. All authors read and approved the final version of the manuscript.

### Competing interests

The authors declare that they have no conflicting interests.

### ACKNOWLEDGMENT

This work was supported by the National Natural Science Foundation of China (NSFC) under Grant Nos. 42525301,  
355 42277092, 42430708, and 42107126, the Key Research Program of Frontier Sciences from the Chinese Academy  
of Sciences (ZDBS-LY-DQC001), the Shaanxi Innovation Team for Science and Technology (2024RS-CXTD-40),  
the Shaanxi Innovation Capability Support Plan - Youth Science and Technology Star Project (2024ZC-KJXX-055),  
and the Natural Science Basic Research Program of Shaanxi Province (2023-JC-QN-0319).

### References

- 360 Chang, X., Wang, S., Zhao, B., Xing, J., Liu, X., Wei, L., Song, Y., Wu, W., Cai, S., Zheng, H., Ding, D., and Zheng,  
M.: Contributions of inter-city and regional transport to PM<sub>2.5</sub> concentrations in the Beijing-Tianjin-Hebei region  
and its implications on regional joint air pollution control, *Sci. Total Environ.*, 660, 1191-1200, 2019.
- Chang, X., Zhao, B., Zheng, H., Wang, S., Cai, S., Guo, F., Gui, P., Huang, G., Wu, D., Han, L., Xing, J., Man, H.,  
365 Hu, R., Liang, C., Xu, Q., Qiu, X., Ding, D., Liu, K., Han, R., Robinson, A. L., and Donahue, N. M.: Full-volatility  
emission framework corrects missing and underestimated secondary organic aerosol sources, *One Earth*, 5, 403-  
412, 2022.
- Chen, Q., Miao, R., Geng, G., Shrivastava, M., Dao, X., Xu, B., Sun, J., Zhang, X., Liu, M., Tang, G., Tang, Q., Hu,  
H., Huang, R.-J., Wang, H., Zheng, Y., Qin, Y., Guo, S., Hu, M., and Zhu, T.: Widespread 2013-2020 decreases and  
reduction challenges of organic aerosol in China, *Nat. Commun.*, 15, 4465, 2024.
- 370 Daellenbach, K. R., Cai, J., Hakala, S., Dada, L., Yan, C., Du, W., Yao, L., Zheng, F., Ma, J., Ungeheuer, F., Vogel,  
A. L., Stolzenburg, D., Hao, Y., Liu, Y., Bianchi, F., Uzu, G., Jaffrezo, J.-L., Worsnop, D. R., Donahue, N. M., and  
Kulmala, M.: Substantial contribution of transported emissions to organic aerosol in Beijing, *Nat. Geosci.*, 17, 747-  
754, 2024.
- Dou, J., Liu, T., Ge, D., Zhang, Y., Yin, J., Wang, L., Liu, H., Li, D., Niu, G., Chen, L., Wang, J., Qi, X., Nie, W., Chi,  
375 X., Huang, X., and Ding, A.: In-situ secondary organic aerosol formation from ambient air in suburban Eastern  
China: Substantially distinct characteristics between summer and winter, *Atmos. Environ.*, 356, 121295, 2025.
- Duan, J., Huang, R.-J., Wang, Y., Xu, W., Zhong, H., Lin, C., Huang, W., Gu, Y., Ovadnevaite, J., Ceburnis, D., and  
O'Dowd, C.: Measurement report: Size-resolved secondary organic aerosol formation modulated by aerosol water  
uptake in wintertime haze, *Atmos. Chem. Phys.*, 24, 7687-7698, 2024.
- 380 Geng, G., Zheng, Y., Zhang, Q., Xue, T., Zhao, H., Tong, D., Zheng, B., Li, M., Liu, F., Hong, C., He, K., and Davis,  
S. J.: Drivers of PM<sub>2.5</sub> air pollution deaths in China 2002–2017, *Nat. Geosci.*, 14, 645-650, 2021.
- Geng, G., Liu, Y., Liu, Y., Liu, S., Cheng, J., Yan, L., Wu, N., Hu, H., Tong, D., Zheng, B., Yin, Z., He, K., and Zhang,  
Q.: Efficacy of China's clean air actions to tackle PM<sub>2.5</sub> pollution between 2013 and 2020, *Nat. Geosci.*, 17, 987-  
994, 2024.
- 385 Huang, R.-J., Wang, Y., Cao, J. J., Lin, C., Duan, J., Chen, Q., Li, Y. J., Gu, Y., Yan, J., Xu, W., Fröhlich, R.,

- Canonaco, F., Bozzetti, C., Ovadnevaite, J., Ceburnis, D., Canagaratna, M., Jayne, J., Worsnop, D. R., El-Haddad, I., Prévôt, A. S., and O'Dowd, C.: Primary emissions versus secondary formation of fine particulate matter in the most polluted city (Shijiazhuang) in North China, *Atmos. Chem. Phys.*, 19, 2283-2298, 2019.
- 390 Huang, R.-J., Li, Y. J., Chen, Q., Zhang, Y., Lin, C., Chan, C. K., Yu, J. Z., de Gouw, J., Tong, S., Jiang, J., Wang, W., Ding, X., Wang, X., Ge, M., Zhou, W., Worsnop, D., Boy, M., Bilde, M., Dusek, U., Carlton, A. G., Hoffmann, T., McNeill, V. F., and Glasius, M.: Secondary organic aerosol in urban China: A distinct chemical regime for air pollution studies, *Science*, 389, eadq2840, 2025.
- 395 Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. E., and Prévôt, A. S. H.: High secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218-222, 2014.
- 400 Huang, X., Liu, Z., Ge, Y., Li, Q., Wang, X., Fu, H., Zhu, J., Zhou, B., Wang, L., George, C., Wang, Y., Wang, X., Su, J., Xue, L., Yu, S., Mellouki, A., and Chen, J.: Aerosol high water contents favor sulfate and secondary organic aerosol formation from fossil fuel combustion emissions, *npj Clim. Atmos. Sci.*, 6, 173, 2023.
- 405 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., E., Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525-1529, 2009.
- 410 Kuang, Y., He, Y., Xu, W., Yuan, B., Zhang, G., Ma, Z., Wu, C., Wang, C., Wang, S., Zhang, S., Tao, J., Ma, N., Su, H., Cheng, Y., Shao, M., and Sun, Y.: Photochemical Aqueous-Phase Reactions Induce Rapid Daytime Formation of Oxygenated Organic Aerosol on the North China Plain, *Environ. Sci. Technol.*, 54, 3849-3860, 2020.
- 415 Lin, C., Huang, R.-J., Duan, J., Zhong, H., and Xu, W.: Primary and Secondary Organic Nitrate in Northwest China: A Case Study, *Environ. Sci. Technol. Lett.*, 8, 947-953, 2021.
- Lin, C., Huang, R.-J., Duan, J., Zhong, H., and Xu, W.: Polycyclic aromatic hydrocarbons from cooking emissions, *Sci. Total Environ.*, 818, 151700, 2022.
- Lin, C., Huang, R.-J., Zhong, H., Duan, J., Wang, Z., Huang, W., and Xu, W.: Elucidating ozone and PM<sub>2.5</sub> pollution in the Fenwei Plain reveals the co-benefits of controlling precursor gas emissions in winter haze, *Atmos. Chem. Phys.*, 23, 3595-3607, 2023.
- 420 Lin, C., Huang, R.-J., Ceburnis, D., Buckley, P., Preissler, J., Wenger, J., Rinaldi, M., Facchini, M. C., O'Dowd, C., and Ovadnevaite, J.: Extreme air pollution from residential solid fuel burning, *Nat. Sustain.*, 1, 512-517, 2018.
- Lin, C., Huang, R.-J., Xu, W., Duan, J., Zheng, Y., Chen, Q., Hu, W., Li, Y., Ni, H., Wu, Y., Zhang, R., Cao, J., and O'Dowd, C.: Comprehensive Source Apportionment of Submicron Aerosol in Shijiazhuang, China: Secondary Aerosol Formation and Holiday Effects, *ACS Earth Space Chem.*, 4, 947-957, 2020.
- 425 Liu, M., Huang, X., Song, Y., Tang, J., Cao, J., Zhang, X., Zhang, Q., Wang, S., Xu, T., Kang, L., Cai, X., Zhang, H., Yang, F., Wang, H., Yu, J. Z., Lau, A. K. H., He, L., Huang, X., Duan, L., Ding, A., Xue, L., Gao, J., Liu, B., and Zhu, T.: Ammonia emission control in China would mitigate haze pollution and nitrogen deposition, but worsen acid rain, *Proc. Natl. Acad. Sci. U.S.A.*, 116, 7760-7765, 2019.
- 430 McFall, A. S., Johnson, A. W., and Anastasio, C.: Air-Water Partitioning of Biomass-Burning Phenols and the Effects of Temperature and Salinity, *Environ. Sci. Technol.*, 54, 3823-3830, 2020.
- Nguyen, T. B., Coggon, M. M., Flagan, R. C., and Seinfeld, J. H.: Reactive Uptake and Photo-Fenton Oxidation of Glycolaldehyde in Aerosol Liquid Water, *Environ. Sci. Technol.*, 47, 4307-4316, 2013.
- 435 Nie, W., Yan, C., Huang, D. D., Wang, Z., Liu, Y., Qiao, X., Guo, Y., Tian, L., Zheng, P., Xu, Z., Li, Y., Xu, Z., Qi, X., Sun, P., Wang, J., Zheng, F., Li, X., Yin, R., Dallenbach, K. R., Bianchi, F., Petäjä, T., Zhang, Y., Wang, M., Schervish, M., Wang, S., Qiao, L., Wang, Q., Zhou, M., Wang, H., Yu, C., Yao, D., Guo, H., Ye, P., Lee, S., Li, Y. J., Liu, Y., Chi,

- X., Kerminen, V.-M., Ehn, M., Donahue, N. M., Wang, T., Huang, C., Kulmala, M., Worsnop, D., Jiang, J., and Ding, A.: Secondary organic aerosol formed by condensing anthropogenic vapours over China's megacities, *Nat. Geosci.*, 15, 255-261, 2022.
- 440 Onasch, T. B., Trimborn, A., Fortner, E. C., Jayne, J. T., Kok, G. L., Williams, L. R., Davidovits, P., and Worsnop, D. R.: Soot Particle Aerosol Mass Spectrometer: Development, Validation, and Initial Application, *Aerosol Sci. Technol.*, 46, 804-817, 2012.
- Pye, H. O. T., Place, B. K., Murphy, B. N., Seltzer, K. M., D'Ambro, E. L., Allen, C., Piletic, I. R., Farrell, S., Schwantes, R. H., Coggon, M. M., Saunders, E., Xu, L., Sarwar, G., Hutzell, W. T., Foley, K. M., Pouliot, G., Bash, J., and Stockwell, W. R.: Linking gas, particulate, and toxic endpoints to air emissions in the Community Regional Atmospheric Chemistry Multiphase Mechanism (CRACMM), *Atmos. Chem. Phys.*, 23, 5043-5099, 2023.
- 445 Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., and Ng, N. L.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Reviews of Geophysics*, 55, 509-559, 2017.
- Wang, J., Ye, J., Zhang, Q., Zhao, J., Wu, Y., Li, J., Liu, D., Li, W., Zhang, Y., Wu, C., Xie, C., Qin, Y., Lei, Y., Huang, X., Guo, J., Liu, P., Fu, P., Li, Y., Lee, H. C., Choi, H., Zhang, J., Liao, H., Chen, M., Sun, Y., Ge, X., Martin, S. T., and Jacob, D. J.: Aqueous production of secondary organic aerosol from fossil-fuel emissions in winter Beijing haze, *Proc. Natl. Acad. Sci. U.S.A.*, 118, e2022179118, 2021.
- 450 Zang, H., Zhao, Y., Huo, J., Zhao, Q., Fu, Q., Duan, Y., Shao, J., Huang, C., An, J., Xue, L., Li, Z., Li, C., and Xiao, H.: High atmospheric oxidation capacity drives wintertime nitrate pollution in the eastern Yangtze River Delta of China, *Atmos. Chem. Phys.*, 22, 4355-4374, 2022.
- 455 Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu, W., Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L., Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu, B., Liu, Z., Yang, F., He, K., and Hao, J.: Drivers of improved PM<sub>2.5</sub> air quality in China from 2013 to 2017, *Proc. Natl. Acad. Sci. U.S.A.*, 116, 24463-24469, 2019.
- 460 Zhang, W., Yun, X., Meng, W., Xu, H., Zhong, Q., Yu, X., Ren, Y. a., Shen, H., Chen, Y., Shen, G., Ma, J., Cheng, H., Li, B., Liu, J., Wang, X., and Tao, S.: Urban residential energy switching in China between 1980 and 2014 prevents 2.2 million premature deaths, *One Earth*, 4, 1602-1613, 2021.
- Zhao, B., Zheng, H., Wang, S., Smith, K. R., Lu, X., Aunan, K., Gu, Y., Wang, Y., Ding, D., Xing, J., Fu, X., Yang, X., Liou, K.-N., and Hao, J.: Change in household fuels dominates the decrease in PM<sub>2.5</sub> exposure and premature mortality in China in 2005-2013, *Proc. Natl. Acad. Sci. U.S.A.*, 115, 12401-12406, 2018.
- 465 Zheng, B., Zhang, Q., Zhang, Y., He, K. B., Wang, K., Zheng, G. J., Duan, F. K., Ma, Y. L., and Kimoto, T.: Heterogeneous chemistry: a mechanism missing in current models to explain secondary inorganic aerosol formation during the January 2013 haze episode in North China, *Atmos. Chem. Phys.*, 15, 2031-2049, 2015.
- Zhong, H., Huang, R.-J., Duan, J., Lin, C., Gu, Y., Wang, Y., Li, Y., Zheng, Y., Chen, Q., Chen, Y., Dai, W., Ni, H., Chang, Y., Worsnop, D. R., Xu, W., Ovadnevaite, J., Ceburnis, D., and O'Dowd, C. D.: Seasonal variations in the sources of organic aerosol in Xi'an, Northwest China: The importance of biomass burning and secondary formation, *Sci. Total Environ.*, 737, 139666, 2020.

475






Figure 1. Aerosol Components and Organic Aerosol Subtypes. Aerosol components in (a) 2014 and (b) 2024 as a function of non-refractory PM deciles, ranging from the least polluted 10th percentile to the most polluted. Organic sources in (c) 2014 and (d) 2024, including non-cooking primary organic aerosol (ncPOA) and secondary organic aerosol, as a function of NR-PM deciles.

480




485 **Figure 2. Organic Aerosol Subtypes vs RH. Variation of non-cooking primary organic aerosol (ncPOA) and secondary organic aerosol (SOA) across different relative humidity (RH) brackets in 2014 (a) and (b) and in 2024 (c) and (d). The error bars represent one standard deviation.**

490



495 **Figure 3. SOA Underestimation in the Model.** (a) the base run (Base), and the adjusted (Adjust) CMAQ model for SJZ in 2014, along with (b) the corresponding data for 2024 as a comparison to the observation (Obs) from PMF result. (c) Distribution of the adjusted SOA in 2014 and (d) in 2024 during the pollution period. Error bars indicate one standard deviation.

500



**Figure 4. Emission Reduction Explains the Reduction in POA. CMAQ simulated monthly mean POA concentrations in (a) scenario using MEIC 2014 + Meteorology 2024; (b) base run with MEIC 2024 + Meteorology 2024. The difference (in  $\mu\text{g m}^{-3}$ ) due to emission is shown in (c) (equivalent to  $b - a$ ) and in percentages (equivalent to  $c/a$ ).**




Figure 5. Regional reduction in PM. The wind field at ground level (a); box plot of  $PM_{2.5}$  concentrations during the heating period (November 15 to March 15) for the cities of Beijing, Tianjin, Shijiazhuang, Xingtai, Handan, and Xi'an. The years 2013-2017 are considered Phase 1 of the Clean Air Action Plan, 2018-2020 as Phase 2 of the three-year action plan to win the blue sky war, and 2021-2024 as Phase 3. Only three full years of measurements are included in (b). The correlation coefficient ( $r$ ) between the seven cities for winter across the three phases is also shown.




Figure 6. Conceptual schematic illustrating the variations in chemical components (shift from sulfate to nitrate) and aerosol sources (shift from POA to SOA) from 2014 to 2024 against a backdrop of emission reductions.