

Effectiveness of Emission Controls on Atmospheric Oxidation and Air Pollutant Concentrations: Uncertainties due to Chemical Mechanisms and Inventories

Mingjie Kang^{1,2}, Hongliang Zhang³, Qi Ying^{4,a}

¹ School of Applied Meteorology, Nanjing University of Information Science and Technology. Nanjing 210044, China.

² Atmospheric Environment Center, Joint Laboratory for International Cooperation on Climate and Environmental Change, Ministry of Education, Nanjing University of Information Science and Technology. Nanjing 210044, China.

³ Department of Environmental Science and Engineering, Fudan University, Shanghai 200433, China.

⁴ Zachry Department of Civil and Environmental Engineering, Texas A&M University, College Station, Texas 77843-3136, USA.

^a Currently at the Division of Environment and Sustainability, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China.

Correspondence to: Qi Ying (qying@ust.edu)(qying@ust.hk)

Abstract. In this study, three photochemical mechanisms of varying complexity from the Statewide Air Pollution Research Center (SAPRC) family and two widely used anthropogenic emission inventories are employed to quantify the discrepancies in the predicted effectiveness of nitrogen oxides (NO_x) and volatile organic compound (VOC) emission controls on ozone (O_3), secondary inorganic aerosols (SIA), and hydroxyl (OH) and nitrate (NO_3) radicals using the Community Multiscale Air Quality (CMAQ) model. For maximum daily average 8-hour O_3 (O_3 -8h), relative reductions predicted using different emission inventory and mechanism combinations are consistent for up to 80% NO_x or VOC reductions, with maximum differences of approximately 15%. For ~~secondary inorganic aerosols (SIA)~~₂, while the predicted relative changes in their daily average concentrations due to NO_x reductions are quite similar, very large differences of up to 30% occur for VOC reductions. Sometimes even the direction of change (i.e., increase or decrease) is different. For the oxidants OH and NO_3 radicals, the uncertainties in the relative changes due to emission changes are even larger among different inventory-mechanism combinations, sometimes by as much as 200%. Our results suggest that while the O_3 -8h responses to emission changes are not sensitive to the choice of chemical mechanism and emission inventories, using a single model and mechanism to evaluate the effectiveness of emission controls on SIA and atmospheric oxidation capacity may have large errors. For these species, the evaluation of the control strategies may require an ensemble approach with multiple inventories and mechanisms.

1. Introduction

Tropospheric ozone (O₃) pollution remains a major global concern. O₃ plays a critical role in atmospheric chemistry and is an important target for air quality improvement (Lu et al., 2018; Lyu et al., 2023; Real et al., 2024) because high levels of surface O₃ negatively affect human health, agricultural crop yields and plant growth (Du et al., 2024; Feng et al., 2022; Ghude et al., 2016; Lu et al., 2020; West et al., 2006). Surface O₃ is mainly formed by the photochemical reactions of NO_x and volatile organic compounds (VOCs) emitted from anthropogenic and biogenic sources (Finlayson-Pitts and Pitts Jr, 1999; Seinfeld and Pandis, 2016). The development of effective O₃ control strategies is hampered by the considerable spatial and temporal variability of surface O₃ concentrations and their non-linear relationship with emissions and meteorological conditions.

Three-dimensional chemical transport models (CTMs) are a valuable tool for developing effective air pollution control strategies. They can provide spatial and temporal information on O₃, particulate matter, and toxic air pollutants by numerically solving the mathematical equations describing the emission, reaction, transport, and deposition of primary and secondary atmospheric pollutants (Byun and Schere, 2006; Russell, 1997). The veracity of modeling outcomes is contingent upon the gas-phase chemical mechanisms and emission inventories, among other variables. Reliance on a single chemical mechanism and emission inventory may result in substantial uncertainty in modelled pollutant concentrations. A common approach to reduce uncertainty in air quality model predictions is to use an ensemble of simulations with different emission inventories and chemical mechanisms (Hu et al., 2017a).

Photochemical mechanism is one of the core components of all CTMs. While the representation of inorganic chemistry is generally similar across mechanisms, the representation of atmospheric organic chemistry differs significantly in terms of the number of explicit model species, the lumping schemes, and the radical chemistry, leading to variations in the model predictions of O₃, PM_{2.5}, air toxics, and some important radical species (Griffith et al., 2016; Kim et al., 2009). Furthermore, the responses of the predictions to changes in emissions may also differ depending on the photochemical mechanism employed, which may impact the assessment of emission control strategies. In practical applications, it is of the utmost importance to strike a balance between mechanism complexity and computational efficiency. For long-term modeling studies of criteria pollutants and the evaluation of numerous emission control strategies, condensed mechanisms may prove to be the optimal choice. More detailed mechanisms are appropriate for a broader range of applications, such as

investigations into specific reaction products that are not explicitly represented in condensed mechanisms. However, they are more demanding on computational resources.

In this context, several mechanism comparison studies have been conducted. Differences in predicted O₃ levels using various photochemical mechanisms have been reported (Yu et al., 2010; Venecek et al., 2018). In addition to directly comparing the model predictions of O₃ concentrations, comparative analyses were also undertaken to examine the similarities and differences between these mechanisms in predicting O₃ changes in response to changes in precursor emissions. For example, Li et al. (2012) and Kang et al. (2022) compared several mechanisms from the SAPRC family, including the standard versions of SAPRC-99, SAPRC-07, SAPRC-11, and a highly condensed version of SAPRC-07. Their results showed that, despite discrepancies in the predictions for O₃, key radicals such as OH and HO₂, and oxidation products such as HNO₃, H₂O₂, NO₂, PAN, and HCHO, the relative changes in O₃ due to changes in NO_x and VOC emissions were almost identical.

On the other hand, the accuracy of model predictions is also significantly affected by uncertainties in anthropogenic emission inventories (Hu et al., 2017a; Kang et al., 2022; Placet et al., 2000), which primarily arise from uncertainties and variability in activity levels (e.g., industrial production or energy consumption) and emission factors (Akimoto et al., 2006; Lei et al., 2011; Streets et al., 2003). For example, when local speciation profiles are not available for the generation of mechanism-specific VOC emissions from the emission rate of non-methane hydrocarbons, average speciation profiles from the SPECIATE database developed by the US EPA were often adopted (Bray et al., 2019; Li et al., 2014; Streets et al., 2003; Wu and Xie, 2017). However, the emission factors from the SPECIATE profiles are predominantly representative of the characteristics of local emissions, as a consequence of the disparities in emission standards and control technologies among diverse geographical regions (Sha et al., 2021). This introduces uncertainties into the emissions and the predicted pollutant concentrations.

To date, several emission inventories covering China have been developed, such as the Multi-resolution Emission Inventory for China (MEIC), the Regional Emission inventory in ASia (REAS), and the Emission Database for Global Atmospheric Research (EDGAR). These inventories have been successfully applied in chemical transport modeling to investigate the concentration and spatial distribution of O₃ and other pollutants (Hu et al., 2017b; Kang et al., 2021; Li et al., 2019, 2018; Saikawa et al., 2017; Xue et al., 2020; Yamaji et al., 2008). Hu et al. (2017a) reported inconsistencies in emission inventories in predicting O₃ and PM_{2.5} using the Weather Research and Forecasting/Community Multiscale Air Quality (WRF/CMAQ)

model system. Ma et al. (2004) identified variations in NO_x and VOC emissions among different inventories as the main factors influencing modeled O₃ concentrations. Our previous study, which employed the condensed SAPRC-07 mechanism and two anthropogenic emission inventories (MEIC and REAS), also observed relatively larger differences in O₃ predictions between inventories in megacities of Beijing and Shanghai, especially on days with elevated O₃ levels (Kang et al., 2022).

While several studies have explored the differences in model predictions due to varying chemical mechanisms or emission inventories, a comprehensive analysis of the influence of diverse combinations of these two factors on the sensitivity of O₃ and other air pollutants to emission changes in China has yet to be undertaken. In this study, we address this gap by applying the CMAQ model, integrated with three photochemical mechanisms and two widely used emission inventories, to quantify the effects of different combinations of mechanisms and inventories on the predictions of maximum daily average 8-hour ozone (O₃-8h) and other secondary pollutants in different regions of China. In addition, the impacts on atmospheric oxidation capacity and key gaseous pollutants are investigated. To gain insight into the variations in pollutant sensitivity, a series of incremental emission reduction scenarios were used, thereby enabling the quantification of the influence of different mechanism and inventory combinations on the response of O₃ and related pollutants. The findings of this study can assist policymakers in the development of more effective and adaptive pollution control strategies.

2. Materials and methods

2.1 The CS07, SAPRC-11 and SAPRC-18 mechanisms

The SAPRC mechanism is a widely used photochemical mechanism that represents complex atmospheric reactions in computationally tractable forms. Instead of tracking the oxidation of individual precursor organic compounds and their reaction products explicitly, the SAPRC is a lumped-molecule chemical mechanism that groups structurally similar VOC species (e.g., alkanes, akenes, and aromatics) into several groups of lumped model species. Some important species, such as isoprene and formaldehyde, are represented explicitly. The reactions of each lumped model species with common oxidants (OH, NO₃ and O₃) are derived based on the reactions of individual species within that group, which are generated automatically using a mechanism generator (Carter et al., 2025). One of the complexities in representing VOC reactions is the intermediate oxidation products and radical species. In the SAPRC mechanism, the radicals such as peroxy radicals and intermediate products such as organic nitrates are

represented by a group of common species to reduce the number of reactions and model species. A brief comparison of the three SAPRC mechanisms is summarized in Table S1.

Three different chemical mechanisms from the SAPRC mechanism family were used in this study, i.e., the condensed SAPRC-07 (CS07) (Carter, 2010), the standard SAPRC-11 (S11) (Carter and Heo, 2013), and the standard SAPRC-18 (S18) (Carter, 2020). These mechanisms were selected to represent different levels of detail in gas-phase reactions in a regional chemical transport model. The CS07 was derived from the widely used SAPRC-07 mechanism and has a high condensation level similar to that of the Carbon Bond mechanism, which is also widely used. The S11 is an updated version of the SAPRC-07 mechanism, with significant revisions made to the aromatic chemistry. The S11 mechanism employed in this study is identical to that utilized in our previous study (Kang et al., 2021). For a detailed description of CS07 and S11 regarding O₃ source apportionment and emission sensitivity, please refer to Kang et al. (2022). The S18 mechanism represents a complete update of the SAPRC mechanism since SAPRC-07. S18 incorporates a greater number of model species, a more explicit representation of peroxy radical chemistry, and a lumping scheme that is more suitable for predicting secondary organic aerosol (SOA) formation. Due to these modifications, S18 is more extensive than S11 in terms of both the number of species and chemical reactions. Although it has been successfully applied in photochemical box models (Jiang et al., 2020; Li et al., 2022a, b), it has not yet been implemented in 3D regional CTMs.

2.2. Anthropogenic emission inventories

The present study compares two widely used anthropogenic emission inventories, MEIC (<http://www.meicmodel.org>) and REAS 3.1 (<https://www.nies.go.jp/REAS/>), to investigate O₃ pollution in China. The emission data from these inventories were processed using an in-house emission processor. Detailed VOC speciation profiles selected from the US EPA-developed SPECIATE database were processed using the speciation profile processor from W.P.L. Carter (2015) to generate profiles for the CS07, SAPRC-11, and SAPRC-18 mechanisms, which are used to estimate emissions of CMAQ-ready VOCs. The MEIC emission inventory includes only emission estimates in China, whereas the REAS emission inventory has complete spatial coverage for Asian countries. In the MEIC simulation, emissions from other countries are supplemented using data from the REAS inventory.

2.3 Model application

The CS07, S11, and S18 mechanisms were incorporated into the CMAQ model (version 5.0.2) to evaluate the differences of mechanisms and inventories in predicting O₃-8h, OH and NO₃ radicals, secondary inorganic aerosols, and reactive VOC species (HCHO) in July 2017 in China. July is widely used in modeling studies to represent conditions of a typical summer month (Kang et al., 2021). The model domain, which covers China and surrounding areas in eastern and southeastern Asia at a 36 km × 36 km horizontal resolution, along with the locations of cities mentioned in the manuscript, is illustrated in [Figure S1](#).

The simulations included model runs with the following combinations: S11 mechanism with MEIC inventory (S11-MEIC), S11 with REAS (S11-REAS), CS07 with MEIC (CS07-MEIC), CS07 with REAS (CS07-REAS), and S18 with REAS (S18-REAS). The MEIC only has five emission sectors, making it difficult to re-speciate the emissions for the S18 mechanism. Specifically, solvent utilization emissions are not represented in the public version of MEIC as a separate sector (Wang et al., 2018). Since solvent utilization accounts for a significant fraction of VOC emissions in urban areas and has very different emission characteristics than fuel combustion sources, re-speciating the five-sector MEIC VOC emissions will be inaccurate. It should be noted that while the MEIC emission inventory is for 2017, the most recent year in the REAS inventory is 2015, which is the one used in the current study.

The July anthropogenic emission inventories from MEIC and REAS were processed using an in-house emission processor, with updated speciation profiles employed to generate CMAQ-ready VOC emissions (Kang et al., 2022). The speciation profiles for different chemical mechanisms were derived from the same detailed speciation profiles extracted from the SPECIATE database. Comprehensive overviews of MEIC and REAS inventories were provided by Kang et al. (2022). Comparisons of major species, including NO_x (NO+NO₂), SO₂, ethene (ETHENE), formaldehyde (HCHO), higher olefins (OLE) (comprising OLE1 and OLE2, which are lumped alkene species with propylene and trans-2-pentene as representative compounds), isoprene (ISOPRENE), and monoterpenes (TRP1), are shown in [Tables S2-S3](#) and [Figures S2-S3](#) for municipalities and provinces in July 2017. Note that OLE and TRP1 emissions are ~~for~~from the S11 mechanisms. Emissions for CS07 are similar but some of the species in OLE and TRP1 are explicit species in S18. There are notable spatial differences in the weekday emissions of HCHO, ETHENE, OLE, SO₂, and NO_x between MEIC and REAS. Specifically, REAS exhibits higher HCHO emissions in locations like Beijing, Tianjin, Henan, Shanghai, and Guangzhou than MEIC ([Figure S2](#)). In the south of Henan, ethene emissions are

significantly higher in REAS relative to MEIC. The emissions of OLE in REAS are lower than those in MEIC in Beijing, Tianjin, Shanghai, Guangzhou, Chengdu, and Chongqing. SO₂ emissions in MEIC are generally lower than those in REAS, except for Shanghai and Guangzhou. NO_x emissions differ significantly between MEIC and REAS. In eastern China, including cities like Shanghai and Guangzhou, NO_x emissions from REAS are typically lower than those from MEIC, although some areas demonstrate a notable increase. These discrepancies inevitably affect the accuracy of air pollutant predictions, underscoring the necessity for a comprehensive assessment of emission inventories in the development of effective pollution control ~~polices~~policies.

Biogenic emissions were generated using the Model of Emissions of Gases and Aerosols from Nature (MEGAN) version 2.10, which has been observed to emit higher levels of isoprene and monoterpenes in comparison to anthropogenic sources (Figure S3). Open burning emissions were produced using FINN inventory from the National Center for Atmospheric Research (NCAR) (Wiedinmyer et al., 2011). Sea salt and windblown dust emissions were simulated online using the CMAQ model. Initial and boundary conditions for the model simulation were generated using CMAQ default profiles. The initial three days of the simulation serve as a spin-up and are excluded from subsequent analyses.

Meteorological inputs were generated using the Weather Research and Forecasting (WRF) model version 4.2 for the 36×36 km domain with 44 vertical layers. The initial and boundary conditions for WRF were derived from the global reanalysis data FNL (available at <https://rda.ucar.edu/datasets/ds083.2/>). Further details on the configuration of WRF model can be found in the work of Kang et al. (2022). WRF-derived meteorological parameters, including temperature and relative humidity at a height of 2 m above the surface and wind speed and direction at 10 m, have been validated against observational data from the National Climatic Data Center (NCDC), demonstrating good performance (Kang et al., 2021).

2.4 Sensitivity of O₃ and related pollutants to emission controls across different mechanisms and inventories

A large number of sensitivity simulations with systematic reductions in NO_x and VOC emissions were conducted to explore variations in the sensitivity of O₃-8h and related pollutants to emission reductions across different mechanisms and inventories. Three sets of simulations were performed for each mechanism/inventory combination considered in this study (see section 2.3). In the first set of simulations, NO_x emissions were reduced by 20, 40,

50, 60, and 80%, while the emissions of VOCs were maintained at their base-case level. In the second set of simulations, VOC emissions were reduced by 20, 40, 50, 60, and 80%, while NO_x emissions remained constant. In the third set of simulations, both NO_x and VOC emissions were reduced by 20, 40, 50, 60, and 80%.

3. Results and discussion

3.1 Model evaluation of O₃ and PM_{2.5} predictions across various mechanisms and inventories

Observations of O₃-8h and PM_{2.5} at a large number of surface monitoring stations nationwide in July 2017 were obtained from the publication website of the China National Environmental Monitoring Center (<http://www.cnemc.cn>). The model performance statistics were evaluated separately for different regions: the North China Plain (NCP), Yangtze River Delta (YRD), Central China (Center), Pearl River Delta (PRD), and Sichuan Basin (SCB). Specifically, NCP includes Beijing, Tianjin, and some cities in Hebei and Shandong provinces; Center includes cities in the provinces of Henan, Hubei, Hunan, and Jiangxi; YRD includes Shanghai and some cities in the provinces of Anhui, Jiangsu, and Zhejiang; PRD includes Shenzhen and some cities in Guangdong province; and SCB includes Chongqing and some cities in Sichuan province. As shown in Figures S4-S5, the model performance for O₃-8h and PM_{2.5} predicted by different mechanisms and emission inventories in major regions of China exhibits large variations, due to differences in climate, topography, and emission sources. Overall, the average values of mean normalized bias (MNB) and mean normalized error (MNE) for O₃-8h predictions across all combinations of mechanisms and inventories are generally within the ~~model performance criteria suggested by the US EPA~~recommended model performance criteria (MNB \leq \pm 0.15 and MNE \leq 0.3) (Emery et al., 2017) in most regions, except for underprediction in the PRD. Similarly, the model shows good performance for PM_{2.5} in most areas, except for the PRD region, where the mean fractional biases (MFB) for PM_{2.5} using S18-REAS, CS07-REAS, and S11-REAS are slightly outside the recommended range (Boylan and Russell, 2006). The underestimation of PM_{2.5} predictions using REAS in the PRD, as shown in Figure S5, is likely related to biases in this inventory specific to this region.

3.2 Spatial variations in predictions of O₃ and related pollutants by different mechanisms and inventories

Figure 1 shows the spatial distribution of monthly averaged O₃-8h concentrations predicted by S11-MEIC in July, along with the absolute differences between O₃-8h predictions from S11-

MEIC and those from other mechanisms and inventories. Based on S11-MEIC, high O₃-8h levels exceeding 80 ppb occurred in eastern China, especially in Beijing, Tianjin, Hebei, northern Henan, SCB, and Shanghai. Extremely high O₃-8h levels above 100 ppb are also observed over the Bohai Bay and the Yellow Sea, likely due to regional transport of polluted air from the continent, reduced NO_x titration with O₃, and lower O₃ dry deposition velocities over the ocean- (Luhar et al., 2017; Silva and Heald, 2018). Similar spatial distributions of O₃-8h concentrations are also found in the simulations using other mechanisms and inventories (Figure S6). In comparison, O₃-8h concentrations predicted by S11-REAS are generally 2–7 ppb higher than those by S11-MEIC in most parts of China, especially in the central region, Zhejiang, and Fujian provinces, but about 2–7 ppb lower in Beijing, Shanghai, Chengdu, and the PRD region. S18-REAS predicted significantly lower O₃-8h levels, with reductions greater than 7 ppb in regions such as Beijing, Tianjin, Hebei, Shanghai, SCB, PRD, and other developed areas such as Zhengzhou and Hefei.

In contrast, CS07-MEIC predicted lower O₃-8h levels overall, with apparent reductions of ~ 6–7 ppb in the SCB, central China, and coastal areas near Shanghai. Similarly, CS07-REAS also showed generally lower O₃-8h concentrations compared to S11-MEIC, particularly in Chengdu, Luoyang, Shanghai, Guangzhou, Ningbo, Hefei, Nanchang, and the Yellow Sea (~7 ppb lower or more), although some locations exhibit higher O₃-8h levels than those predicted by S11-MEIC. These discrepancies highlight that the accuracy of O₃ predictions is sensitive to variations in photochemical mechanisms and emission inventories. This underscores the importance of adopting and comparing multiple mechanisms and inventories when developing region-specific pollution control policies.

The difference in Figure 1(c) for other countries is mainly due to the difference between S11 and S18. Predicted O₃ concentrations by S18 are lower than those by S11, especially in urban areas (e.g., in Ulaanbaatar in Mongolia). This is consistent with the box-model simulation results reported by Carter (Carter, 2020). In Figure 1(a), the difference of the monthly O₃ in other countries is negligible when the same chemical mechanism is used, which further confirms our conclusion.

In Figures 1(b), O₃ concentrations from S11-REAS are significantly lower than those from S11-MEIC. This is because O₃ formation sensitivity regime over the yellow sea is likely NO_x-limited as the VOC-sensitive urban plume is advected to the marine environment (Vermeuel et al., 2019). Furthermore, satellite observed HCHO/NO_x columns over the Yellow Sea is greater than 6, clearly indicating a NO_x-limit regime (Kang et al., 2021). Since the NO_x emissions in the upwind regions in the REAS inventory are significantly lower than

these in the MEIC inventory, this leads to reduced O₃ formation in the S11-REAS results compared to the S11-MEIC results.

Figure 2 illustrates the differences in spatial distribution of monthly averaged secondary inorganic aerosol (SIA) concentrations modeled with different mechanisms and inventories. According to S11-MEIC, high SIA concentrations are mainly concentrated in NCP and SCB, exceeding 15 µg m⁻³. High SIA concentrations are also found in Bohai Bay, likely due to long-range transport of polluted air from land sources. Compared to S11-MEIC, S11-REAS predicts higher SIA concentrations in most regions, with increases of > 1 µg m⁻³ in most areas, especially in NCP, Henan, and SCB (higher by about 6 µg m⁻³ or even more). Nationwide, SIA concentrations predicted by S18-REAS are generally higher than those predicted by S11-MEIC, with increases of up to 6 µg m⁻³ or more in the NCP, Central China, SCB, Bohai Bay, and the Yellow Sea. CS07-MEIC shows similar SIA levels to S11-MEIC, with slightly lower concentrations of ~1 µg m⁻³ in SCB and Bohai Bay. The spatial differences in SIA predictions between CS07-REAS and S11-MEIC are similar to those between S11-REAS and S11-MEIC, suggesting that SIA concentrations from CS07-REAS and S11-REAS are comparable.

3.3 Impacts of mechanisms and inventories on predicted atmospheric oxidation capacity

Atmospheric oxidation capacity (AOC), which governs the removal rate of primary pollutants and the production of secondary pollutants (Elshorbany et al., 2009; Prinn, 2003), is primarily controlled by the hydroxyl (OH) and nitrate (NO₃) radicals in the atmosphere (Geyer et al., 2001; Liu et al., 2022). The spatial differences in OH and NO₃ predictions for different mechanisms and inventories are illustrated in Figures 3-4.

S11-MEIC predicts high OH concentrations exceeding 3.7×10^6 molecules cm⁻³ (0.15 ppt) mainly in northern China, including NCP, Inner Mongolia, and some western sites. Compared to S11-MEIC, S11-REAS predicts lower OH concentrations in NCP, Jiangsu, Shanghai, PRD, SCB, and other urban nuclei but higher concentrations in rural areas. In contrast, S18-REAS predicts higher OH concentrations than S11-MEIC in most regions, except in some northern locations. CS07-MEIC produces elevated OH concentrations in northwestern China, Chengdu, Chongqing, Shanghai, Hebei, Shandong, and northern Henan, but lower OH levels in other regions. Similar to S11-REAS, CS07-REAS shows much higher OH levels in northwestern China but lower OH levels in eastern regions than S11-MEIC. The variability in OH levels can significantly affect the formation of O₃ and other gaseous and particulate pollutants.

The spatial distribution of NO₃ concentrations modeled by S11-MEIC exhibits high values in Xinjiang, Inner Mongolia, NCP, and Bohai Bay, with NO₃ concentrations reaching up to 20 ppt. Obvious differences in NO₃ predictions between S11-REAS and S11-MEIC are primarily observed in Xinjiang and Inner Mongolia, with smaller differences in other regions. S18-REAS generally predicted higher NO₃ concentrations than S11-MEIC, especially in Xinjiang, Inner Mongolia, Bohai Bay, Yellow Sea, and Taiwan Strait, by ~6 ppt or more. In contrast, CS07-MEIC consistently predicts lower NO₃ levels than S11-MEIC. CS07-REAS predicted lower NO₃ concentrations in the eastern regions but higher concentrations in parts of Xinjiang compared to S11-MEIC.

3.4 Impacts of mechanisms and inventories on HCHO prediction

Spatial variations in AOC, as reflected in OH and NO₃ predictions from different mechanisms and inventories, imply that photochemical formation and loss rates for air toxics may also vary depending on the mechanism and inventory used. To explore how these variations affect the modeling results for gaseous pollutants, we examined HCHO, one of the most important gaseous air toxics with both primary and secondary sources. It is also a significant contributor to O₃ formation and OH production, thus playing a crucial role in tropospheric photochemistry (Wang et al., 2017; Zhang et al., 2013).

~~HCHO concentrations predicted by S11-MEIC~~ Monthly average surface HCHO concentrations from S11-MEIC are similar to the surface observations made in China (Zhang et al., 2021). HCHO concentrations are generally high in the SCB and eastern China, areas with significant biogenic VOC emissions (Kang et al., 2023), particularly in Chengdu, Shanghai, and Changsha, where HCHO concentrations reach or exceed 7 ppb (Figure 5). (Kang et al., 2023a), particularly in Chengdu, Shanghai, and Changsha, where HCHO concentrations reach or exceed 7 ppb (Figure 5). This suggests that a significant fraction of the HCHO is due to secondary formation. HCHO levels predicted by S11-REAS are similar to those predicted by S11-MEIC, with only minor differences of less than 1 ppb. In contrast, S18-REAS predicts significantly lower HCHO levels than S11-MEIC, with differences > 1 ppb, especially in SCB, NCP, PRD, YRD, and central China. CS07-MEIC and CS07-REAS predict higher HCHO levels than S11-MEIC, especially in SCB and eastern China, with differences exceeding 1 ppb. These results indicate that HCHO predictions are more strongly influenced by photochemical mechanisms than by uncertainties in emission inventories, due to its secondary formation.

3.5 Impacts of mechanisms and inventories on the sensitivity of O₃ and related pollutants to emission controls

Figures 6-10 and S6-S15 display the fractional changes in predictions of O₃ and related pollutants due to NO_x and VOC reductions using various mechanisms and inventories.

3.5.1 Impacts of mechanisms and inventories on O₃ sensitivity to emission controls

For all mechanisms and inventories, O₃-8h concentrations consistently decrease with reductions in NO_x, VOCs, or both in all five cities in July. The efficiency of emission controls improves with increasingly higher emission reductions of NO_x, VOCs. These trends are consistent with the findings of Kang et al. (2021), suggesting that the mechanisms and inventories do not affect the trend of change in O₃, but do affect the magnitude of the change. These results also show that in July, the sensitivity of O₃ formation to NO_x and VOCs is in transition or NO_x-limited regimes. In Beijing, when NO_x emissions are reduced, CS07-MEIC and CS07-REAS show the largest O₃-8h reductions compared to other mechanisms and inventories, while S18-REAS exhibits the smallest changes, especially with larger NO_x reductions. In Shanghai, the largest O₃-8h reductions due to NO_x controls are observed with CS07-REAS, followed by S11-REAS, while S18-REAS again shows the smallest changes. In Changsha, there are no significant differences among mechanisms and inventories in the sensitivity tests. In Shenzhen, S11-REAS shows the greatest decreases in O₃-8h, followed by CS07-REAS and S18-REAS, while S11-MEIC and CS07-MEIC show the smallest changes. Except for S18-REAS, predictions from other mechanisms and inventories are similar in Chongqing.

When only VOC emissions are reduced in Beijing, S11-MEIC predicts the largest O₃-8h reductions, while CS07-REAS predicts the smallest changes. Similar patterns are found in Shanghai and Changsha, with larger differences among different mechanisms and inventories in Shanghai. In Shenzhen, S11-MEIC also exhibits the largest O₃-8h decreases, followed by CS07-MEIC, while CS07-REAS, S11-REAS, and S18-REAS show smaller changes. In Chongqing, S18-REAS predicts the largest changes and CS07-MEIC the smallest, although the differences among these scenarios are not substantial. When both NO_x and VOC emissions are reduced, the predicted change rates of O₃-8h do not vary much across different mechanisms and emission inventories for all five cities. Additionally, the sensitivity tests for Shenzhen, as presented in Figure 6, suggest that differences in emission inventories may have a greater impact on emission control outcomes than differences in chemical mechanisms. O₃ in Shenzhen is lower than other cities included in Figure 6 and most O₃ there is background O₃

on typical days (Kang et al., 2023b). Thus, it is less responsive to emission reductions than other cities.

Figure S6 illustrates the national-scale relative changes in O₃-8h due to NO_x emission reductions evaluated using different mechanisms and inventories. The results are consistent with those shown in Figure 6, with the greatest O₃-8h reductions occurring in the SCB, Central, and YRD. For all mechanisms and inventories, increasingly higher NO_x reductions generally lead to increasingly lower O₃ in July, with relative changes in O₃-8h ranging from approximately 5% to 60% as NO_x reductions increase from 20% to 80%. In comparison, the O₃-8h reductions predicted by S18-REAS are less pronounced than other mechanisms and inventories in most areas. Similar to NO_x reductions, higher VOC reductions typically result in larger O₃ reductions in July, with significant reductions observed in NCP, Chengdu, PRD, Shanghai, and Bohai Bay. The relative changes in O₃-8h increase from 5% to 40% as VOC reductions increase from 20% to 80% in these areas (Figure S7). The comparison between Figure S6 and Figure S7 suggests that NO_x reduction tends to be more effective in controlling O₃ pollution in non-VOC-limited regions than VOC reduction, given the same level of emission reduction.

3.5.2 Impacts of mechanisms and inventories on SIA sensitivity to emission controls

Reducing NO_x emissions typically leads to decreasing SIA levels across the five cities, with the efficiency of reductions improving as NO_x reduction increases. As illustrated in Figure 7, the predicted changes in SIA are quite similar for different mechanisms and inventories. However, VOC controls do not always effectively reduce SIA concentrations, likely attributable to an increase in NO₃ radicals, as shown in Figure 9. In Beijing, changes in SIA in response to VOC reductions vary with mechanisms and inventories. For S11-MEIC and CS07-MEIC, SIA concentration first increases and then decreases with decreasing VOCs. S18-REAS shows a general decrease in SIA concentrations with decreasing VOCs. In contrast, CS07-REAS and S11-REAS predict an increase in SIA concentrations with VOC reductions, with the rate of SIA growth initially increasing and then decreasing as VOC emission decreases. In Shanghai, VOC reductions generally increase SIA concentrations across all mechanisms and inventories, especially for CS07-REAS and S11-REAS, which show that SIA concentrations consistently increase as VOCs are progressively reduced. In Changsha, SIA concentrations also consistently increase with increasing VOC reductions for all mechanisms and inventories. In Shenzhen, SIA concentrations increase with VOC reductions; however, the changes are more pronounced for S11-REAS compared to the less significant variations observed for CS07-

MEIC. Except for S18-REAS, Chongqing shows an increase in SIA concentrations with VOC reductions, particularly for CS07-REAS. Overall, simultaneous reductions in NO_x and VOCs decrease SIA levels in most cities, similar to the effects of NO_x controls alone.

Figure S8 shows the relative changes in SIA due to NO_x controls on a national scale, using different mechanisms and inventories. The stepwise reductions in NO_x emissions lead to an overall decrease in SIA concentrations, with the most pronounced reductions observed in the NCP, SCB, Jiangsu, Bohai Bay, and Yellow Sea. The decreases in SIA increase from ~5% to 60% as NO_x reduction increases from 20% to 80%. On the contrary, VOC reduction causes an overall increase in SIA concentrations (Figure S9), indicating that VOC controls are less effective than NO_x controls in reducing SIA levels in July. Notably, the largest increases in SIA are seen in SCB, Central, and YRD, where SIA increases grow from ~6% to 30% as VOC reduction increases from 20% to 80%. In comparison, S18-REAS shows relatively smaller changes in SIA, with some areas in NCP showing obvious decreases in SIA concentrations.

3.5.3 Impacts of mechanisms and inventories on OH and NO₃ sensitivity

As shown in Figures 8-9, the effects of NO_x reductions on OH concentrations vary depending on the mechanism and inventory used. In most regions, OH production decreases with reducing NO_x emissions, likely due to the decrease in O₃ concentrations, as O₃ photolysis in the presence of water vapor is a significant source of atmospheric OH (Seinfeld and Pandis, 2016). Except for Shenzhen, CS07-MEIC and CS07-REAS predict the largest decreases in OH due to NO_x control in other cities, while S18-REAS predicts the smallest OH change rates. In Shenzhen, OH levels predicted by S11-MEIC and CS07-MEIC initially increase and then decrease as NO_x emissions decrease, while those predicted by S18-REAS, CS07-REAS, and S11-REAS exhibit a consistent decrease in OH levels. When only VOCs are reduced, the changes in OH vary across different mechanisms and inventories. In Beijing, except for S18-REAS, OH concentration initially increases and then decreases with decreasing VOC emissions. In Shanghai, OH concentrations from CS07-REAS and S11-REAS basically increase with decreasing VOCs, while those from CS07-MEIC and S11-MEIC show an initial increase followed by a decrease. S18-REAS predicts a general decrease of OH with decreasing VOCs in Beijing and Shanghai. In Changsha, OH concentrations consistently increase with reduced VOCs across all mechanisms and inventories, with CS07-REAS showing a rapid increase of up to 75%. In Shenzhen, OH levels from CS07-REAS, S11-REAS, and S18-REAS increase with decreasing VOCs, while those from S11-MEIC and CS07-MEIC decrease. In Chongqing, all mechanisms and inventories except S18-REAS predict an increase in OH concentrations

with decreasing VOCs, with CS07-MEIC showing a significant increase of up to 75%. When both NO_x and VOC emissions are reduced, OH concentrations generally increase for all mechanisms and inventories, although the relative changes are relatively small ($< 25\%$).

Figures S10-S11 display the spatial distribution of OH changes due to incremental emission controls across different mechanisms and inventories. There is a nationwide decrease in OH concentrations due to NO_x reductions, with more pronounced decreases in central and eastern China, where OH levels drop by $\sim 20\%$ to 80% as NO_x reduction increases from 20% to 80% . ~~Instead, VOC reductions generally lead to increased OH concentrations across most of China, significantly in some regions with large vegetation cover (Kang et al., 2023), such as central and southeastern China.~~ Instead, VOC reductions generally lead to increased OH concentrations across most of China, significantly in some regions with large vegetation cover (Kang et al., 2023a), such as central and southeastern China. In these areas, OH levels increase by $20\sim 200\%$ in response to stepwise VOC reductions. This increase occurs because high atmospheric VOC concentrations in rural and vegetation-rich areas react with OH radicals to produce RO_2 and HO_2 , depleting OH, thus reducing VOCs in these areas increases OH levels. However, in some urban centers, such as Beijing and Shanghai, changes in OH levels due to VOC controls depend on the extent of reduction and the choice of mechanism and inventory.

Similar to OH, NO_3 levels predicted by different mechanisms and inventories basically decrease with decreasing NO_x emissions, likely related to the decline of O_3 concentrations since NO_3 radicals are predominantly formed by the reaction of NO_2 with O_3 (Geyer et al., 2001). In general, the change rates of NO_3 due to NO_x reductions are higher for CS07-MEIC and CS07-REAS, while lower for S18-REAS. When VOC emissions are reduced, NO_3 production increases, except for S18-REAS. In Changsha, a particularly dramatic increase of over 200% and 150% is observed for CS07-REAS and CS07-MEIC, respectively, when VOC emissions are cut by 80% . This increase in NO_3 radicals could be attributed to a decline in the rapid reaction of NO_3 with unsaturated hydrocarbons. However, for S18-REAS, NO_3 concentrations decrease with decreasing VOCs in Beijing, Shanghai, and Chongqing. Unlike OH, simultaneous reductions in NO_x and VOCs result in lower NO_3 levels across all mechanisms and inventories.

Figures S12-13 illustrate the spatial variation in NO_3 changes due to systematic emission reductions for all mechanisms and inventories. NO_3 radicals show a decreasing trend, particularly in eastern China, where reductions in NO_3 levels vary from 40% to 100% . Except for S18-REAS, other mechanisms and inventories generally predict a nationwide increase in NO_3 radicals with reduced VOCs, particularly in central and southeastern China, where NO_3

levels increase by ~ 30 – 300% or even more with increasing VOC reductions. S18-REAS predicts a decrease in NO₃ levels with reduced VOCs in some megacities such as Beijing, Shanghai, Chengdu, and Chongqing, while NO₃ levels increase elsewhere.

3.5.4 Impacts of mechanisms and inventories on the sensitivity of secondary gaseous pollutants to emission controls

As shown in Figure 10, HCHO concentrations decrease with decreasing NO_x emissions, likely due to decreased AOC, as evidenced by decreased OH and NO₃ levels. This indicates that secondary formation from the oxidation of atmospheric VOCs is the dominant source of HCHO, in accordance with previous studies (Wang et al., 2017; Yang et al., 2019; Zhang et al., 2013). In addition, the most significant decreases in HCHO levels due to NO_x reductions are observed for CS07-REAS in all five cities, particularly in Changsha and Chongqing, where HCHO levels drop by about 40%. When VOCs are reduced alone, HCHO levels decrease linearly across all mechanisms and inventories, a trend similar to the simultaneous reduction of NO_x and VOCs. Variations in HCHO changes across different mechanisms and inventories suggest the need to evaluate different mechanisms and inventories when formulating regional emission control policies for carbonyl pollution.

Figures S14- S15 illustrate the spatial distribution of relative changes in HCHO due to NO_x and VOC controls using different mechanisms and inventories. VOC and NO_x controls lead to reductions in HCHO concentration, especially in eastern China. However, VOC controls generally ~~leads~~lead to larger reductions in HCHO concentrations (~10–80%) than NO_x controls (~ 4 – 40%), as shown in Figures S14-15 and 10. This suggests that VOC controls are also essential and more effective for reducing secondary gaseous organic pollutants in the atmosphere.

4. Conclusions

This study utilized the CMAQ model to evaluate the impacts of different mechanisms and inventories on the prediction of O₃ and other air pollutants. It also examined how these mechanisms and inventories affect the sensitivity of O₃ and related species to emission reductions. For ~~maximum daily average 8-hour O₃-(O₃-8h)₂~~, relative reductions predicted using different emission inventory and mechanism combinations are consistent for up to 80% NO_x or VOC reductions, with maximum differences of approximately 15%. For ~~secondary inorganic aerosols-(SIA)₂~~, while the predicted relative changes in their daily average concentrations due to NO_x reductions are quite similar, very large differences of up to 30%

occur for VOC reductions. Sometimes even the direction of change (i.e., increase or decrease) is different. For the oxidants OH and NO₃ radicals, the uncertainties in the relative changes due to emission changes are even larger among different inventory-mechanism combinations, sometimes by as much as 200%. Our results suggest that while the O₃-8h responses to emission changes are not sensitive to the choice of chemical mechanism and emission inventories, using a single model and mechanism to evaluate the effectiveness of emission controls on SIA and atmospheric oxidation capacity may have large errors. For these species, the evaluation of the control strategies may require an ensemble approach with multiple inventories and mechanisms.

Data availability. The dataset for this paper is available upon request from the corresponding author (qying@ust.hk).

Author contributions. QY designed the study. QY and MJK developed the CMAQ model. MJK conducted the simulations, analyzed the data, and wrote the manuscript. All coauthors contributed to the discussion and revision of the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgments

This work was partly supported by the National Natural Science Foundation of China (No. 42307142).

References

- Akimoto, H., Ohara, T., Kurokawa, J., and Horii, N.: Verification of energy consumption in China during 1996–2003 by using satellite observational data, *Atmospheric Environment*, 40, 7663–7667, <https://doi.org/10.1016/j.atmosenv.2006.07.052>, 2006.
- Boylan, J. W. and Russell, A. G.: PM and light extinction model performance metrics, goals, and criteria for three-dimensional air quality models, *Atmospheric Environment*, 40, 4946–4959, <https://doi.org/10.1016/j.atmosenv.2005.09.087>, 2006.
- Bray, C. D., Strum, M., Simon, H., Riddick, L., Kosusko, M., Menetrez, M., Hays, M. D., and Rao, V.: An assessment of important SPECIATE profiles in the EPA emissions modeling platform and current data gaps, *Atmospheric Environment*, 207, 93–104, <https://doi.org/10.1016/j.atmosenv.2019.03.013>, 2019.
- Byun, D. and Schere, K. L.: Review of the Governing Equations, Computational Algorithms, and Other Components of the Models-3 Community Multiscale Air Quality (CMAQ) Modeling System, *Applied Mechanics Reviews*, 59, 51–77, <https://doi.org/10.1115/1.2128636>, 2006.
- Carter, W. P. L.: Development of a condensed SAPRC-07 chemical mechanism, *Atmospheric Environment*, 44, 5336–5345, <https://doi.org/10.1016/j.atmosenv.2010.01.024>, 2010.
- Carter, W. P. L.: Development of a database for chemical mechanism assignments for volatile organic emissions, *Journal of the Air & Waste Management Association*, 65, 1171–1184, <https://doi.org/10.1080/10962247.2015.1013646>, 2015.
- Carter, W. P. L.: Documentation of the SAPRC-18 mechanism, Report to the California Air Resources Board, 2020.
- Carter, W. P. L. and Heo, G.: Development of revised SAPRC aromatics mechanisms, *Atmos. Environ.*, 77, 404–414, <https://doi.org/10.1016/j.atmosenv.2013.05.021>, 2013.
- Carter, W. P. L., Jiang, J., Orlando, J. J., and Barsanti, K. C.: Derivation of atmospheric reaction mechanisms for volatile organic compounds by the SAPRC mechanism generation system (MechGen), *Atmospheric Chemistry and Physics*, 25, 199–242, <https://doi.org/10.5194/acp-25-199-2025>, 2025.
- Du, H., Yan, M., Liu, X., Zhong, Y., Ban, J., Lu, K., and Li, T.: Exposure to Concurrent Heatwaves and Ozone Pollution and Associations with Mortality Risk: A Nationwide Study in China, *Environmental Health Perspectives*, 132, 047012, <https://doi.org/10.1289/EHP13790>, 2024.
- Elshorbany, Y. F., Kurtenbach, R., Wiesen, P., Lissi, E., Rubio, M., Villena, G., Gramsch, E., Rickard, A. R., Pilling, M. J., and Kleffmann, J.: Oxidation capacity of the city air of Santiago, Chile, *Atmospheric Chemistry and Physics*, 9, 2257–2273, <https://doi.org/10.5194/acp-9-2257-2009>, 2009.

595 [Emery, C., Liu, Z., Russell, A. G., Odman, M. T., Yarwood, G., and Kumar, N.:
596 Recommendations on statistics and benchmarks to assess photochemical model
597 performance, Journal of the Air & Waste Management Association, 67, 582–598,
598 <https://doi.org/10.1080/10962247.2016.1265027>, 2017.](#)

599 Feng, Z., Xu, Y., Kobayashi, K., Dai, L., Zhang, T., Agathokleous, E., Calatayud, V., Paoletti, E.,
600 Mukherjee, A., Agrawal, M., Park, R. J., Oak, Y. J., and Yue, X.: Ozone pollution threatens the
601 production of major staple crops in East Asia, *Nat Food*, 3, 47–56,
602 <https://doi.org/10.1038/s43016-021-00422-6>, 2022.

603 Finlayson-Pitts, B. J. and Pitts Jr, J. N.: Chemistry of the upper and lower atmosphere:
604 theory, experiments, and applications, Elsevier, 1999.

605 Geyer, A., Alicke, B., Konrad, S., Schmitz, T., Stutz, J., and Platt, U.: Chemistry and oxidation
606 capacity of the nitrate radical in the continental boundary layer near Berlin, *Journal of*
607 *Geophysical Research: Atmospheres*, 106, 8013–8025,
608 <https://doi.org/10.1029/2000JD900681>, 2001.

609 Ghude, S. D., Chate, D. M., Jena, C., Beig, G., Kumar, R., Barth, M. C., Pfister, G. G., Fadnavis,
610 S., and Pithani, P.: Premature mortality in India due to PM_{2.5} and ozone exposure,
611 *Geophysical Research Letters*, 43, 4650–4658, <https://doi.org/10.1002/2016GL068949>,
612 2016.

613 Griffith, S. M., Hansen, R. F., Dusanter, S., Michoud, V., Gilman, J. B., Kuster, W. C., Veres, P.
614 R., Graus, M., de Gouw, J. A., Roberts, J., Young, C., Washenfelter, R., Brown, S. S., Thalman,
615 R., Waxman, E., Volkamer, R., Tsai, C., Stutz, J., Flynn, J. H., Grossberg, N., Lefer, B., Alvarez,
616 S. L., Rappenglueck, B., Mielke, L. H., Osthoff, H. D., and Stevens, P. S.: Measurements of
617 hydroxyl and hydroperoxy radicals during CalNex-LA: Model comparisons and radical
618 budgets, *Journal of Geophysical Research: Atmospheres*, 121, 4211–4232,
619 <https://doi.org/10.1002/2015JD024358>, 2016.

620 Hu, J., Li, X., Huang, L., Ying, Q., Zhang, Q., Zhao, B., Wang, S., and Zhang, H.: Ensemble
621 prediction of air quality using the WRF/CMAQ model system for health effect studies in
622 China, *Atmospheric Chemistry and Physics*, 17, 13103–13118, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-17-13103-2017)
623 [17-13103-2017](https://doi.org/10.5194/acp-17-13103-2017), 2017a.

624 Hu, J., Huang, L., Chen, M., Liao, H., Zhang, H., Wang, S., Zhang, Q., and Ying, Q.: Premature
625 Mortality Attributable to Particulate Matter in China: Source Contributions and Responses
626 to Reductions, *Environ. Sci. Technol.*, 51, 9950–9959,
627 <https://doi.org/10.1021/acs.est.7b03193>, 2017b.

628 Jiang, J., Carter, W. P. L., Cocker, D. R. I., and Barsanti, K. C.: Development and Evaluation of
629 a Detailed Mechanism for Gas-Phase Atmospheric Reactions of Furans, *ACS Earth Space*
630 *Chem.*, 4, 1254–1268, <https://doi.org/10.1021/acsearthspacechem.0c00058>, 2020.

631 Kang, M., Zhang, J., Zhang, H., and Ying, Q.: On the Relevancy of Observed Ozone Increase
632 during COVID-19 Lockdown to Summertime Ozone and PM_{2.5} Control Policies in China,
633 *Environ. Sci. Technol. Lett.*, 8, 289–294, <https://doi.org/10.1021/acs.estlett.1c00036>, 2021.

634 Kang, M., Hu, J., Zhang, H., and Ying, Q.: Evaluation of a highly condensed SAPRC chemical
 635 mechanism and two emission inventories for ozone source apportionment and emission
 636 control strategy assessments in China, *Science of The Total Environment*, 813, 151922,
 637 <https://doi.org/10.1016/j.scitotenv.2021.151922>, 2022.

638 Kang, M., Zhang, H., and Ying, Q.: Enhanced summertime background ozone by
 639 anthropogenic emissions – Implications on ozone control policy and health risk assessment,
 640 *Atmospheric Environment*, 314, 120116, <https://doi.org/10.1016/j.atmosenv.2023.120116>,
 641 [2023](https://doi.org/10.1016/j.atmosenv.2023.120116)[2023a](https://doi.org/10.1016/j.atmosenv.2023.120116).

642 [Kang, M., Zhang, H., and Ying, Q.: Enhanced summertime background ozone by](https://doi.org/10.1016/j.atmosenv.2023.120116)
 643 [anthropogenic emissions – Implications on ozone control policy and health risk assessment,](https://doi.org/10.1016/j.atmosenv.2023.120116)
 644 [Atmospheric Environment, 120116, https://doi.org/10.1016/j.atmosenv.2023.120116,](https://doi.org/10.1016/j.atmosenv.2023.120116)
 645 [2023b](https://doi.org/10.1016/j.atmosenv.2023.120116).

646 Kim, Y., Sartelet, K., and Seigneur, C.: Comparison of two gas-phase chemical kinetic
 647 mechanisms of ozone formation over Europe, *J Atmos Chem*, 62, 89–119,
 648 <https://doi.org/10.1007/s10874-009-9142-5>, 2009.

649 Lei, Y., Zhang, Q., He, K. B., and Streets, D. G.: Primary anthropogenic aerosol emission
 650 trends for China, 1990–2005, *Atmospheric Chemistry and Physics*, 11, 931–954,
 651 <https://doi.org/10.5194/acp-11-931-2011>, 2011.

652 Li, J., Zhang, H., and Ying, Q.: Comparison of the SAPRC07 and SAPRC99 photochemical
 653 mechanisms during a high ozone episode in Texas: Differences in concentrations, OH budget
 654 and relative response factors, *Atmospheric Environment*, 54, 25–35,
 655 <https://doi.org/10.1016/j.atmosenv.2012.02.034>, 2012.

656 Li, K., Jacob, D. J., Liao, H., Shen, L., Zhang, Q., and Bates, K. H.: Anthropogenic drivers of
 657 2013–2017 trends in summer surface ozone in China, *Proc Natl Acad Sci USA*, 116, 422–427,
 658 <https://doi.org/10.1073/pnas.1812168116>, 2019.

659 Li, K., Zhang, X., Zhao, B., Bloss, W. J., Lin, C., White, S., Yu, H., Chen, L., Geng, C., Yang, W.,
 660 Azzi, M., George, C., and Bai, Z.: Suppression of anthropogenic secondary organic aerosol
 661 formation by isoprene, *npj Clim Atmos Sci*, 5, 1–9, [https://doi.org/10.1038/s41612-022-](https://doi.org/10.1038/s41612-022-00233-x)
 662 [00233-x](https://doi.org/10.1038/s41612-022-00233-x), 2022a.

663 Li, M., Zhang, Q., Streets, D. G., He, K. B., Cheng, Y. F., Emmons, L. K., Huo, H., Kang, S. C., Lu,
 664 Z., Shao, M., Su, H., Yu, X., and Zhang, Y.: Mapping Asian anthropogenic emissions of non-
 665 methane volatile organic compounds to multiple chemical mechanisms, *Atmospheric*
 666 *Chemistry and Physics*, 14, 5617–5638, <https://doi.org/10.5194/acp-14-5617-2014>, 2014.

667 Li, N., He, Q., Greenberg, J., Guenther, A., Li, J., Cao, J., Wang, J., Liao, H., Wang, Q., and
 668 Zhang, Q.: Impacts of biogenic and anthropogenic emissions on summertime ozone
 669 formation in the Guanzhong Basin, China, *Atmospheric Chemistry and Physics*, 18, 7489–
 670 7507, <https://doi.org/10.5194/acp-18-7489-2018>, 2018.

671 Li, Q., Jiang, J., Afreh, I. K., Barsanti, K. C., and Cocker III, D. R.: Secondary organic aerosol
 672 formation from camphene oxidation: measurements and modeling, *Atmospheric Chemistry*
 673 *and Physics*, 22, 3131–3147, <https://doi.org/10.5194/acp-22-3131-2022>, 2022b.

674 Liu, T., Hong, Y., Li, M., Xu, L., Chen, J., Bian, Y., Yang, C., Dan, Y., Zhang, Y., Xue, L., Zhao, M.,
 675 Huang, Z., and Wang, H.: Atmospheric oxidation capacity and ozone pollution mechanism in
 676 a coastal city of southeastern China: analysis of a typical photochemical episode by an
 677 observation-based model, *Atmospheric Chemistry and Physics*, 22, 2173–2190,
 678 <https://doi.org/10.5194/acp-22-2173-2022>, 2022.

679 Lu, X., Hong, J., Zhang, L., Cooper, O. R., Schultz, M. G., Xu, X., Wang, T., Gao, M., Zhao, Y.,
 680 and Zhang, Y.: Severe surface ozone pollution in China: A global perspective, *Environmental*
 681 *Science & Technology Letters*, 5, 487–494, 2018.

682 Lu, X., Zhang, L., Wang, X., Gao, M., Li, K., Zhang, Y., Yue, X., and Zhang, Y.: Rapid Increases in
 683 Warm-Season Surface Ozone and Resulting Health Impact in China Since 2013, *Environ. Sci.*
 684 *Technol. Lett.*, 7, 240–247, <https://doi.org/10.1021/acs.estlett.0c00171>, 2020.

685 Luhar, A. K., Galbally, I. E., Woodhouse, M. T., and Thatcher, M.: An improved
 686 parameterisation of ozone dry deposition to the ocean and its impact in a global climate–
 687 chemistry model, *Atmos. Chem. Phys.*, 17, 3749–3767, [https://doi.org/10.5194/acp-17-](https://doi.org/10.5194/acp-17-3749-2017)
 688 3749-2017, 2017.

689 Lyu, X., Li, K., Guo, H., Morawska, L., Zhou, B., Zeren, Y., Jiang, F., Chen, C., Goldstein, A. H.,
 690 Xu, X., Wang, T., Lu, X., Zhu, T., Querol, X., Chatani, S., Latif, M. T., Schuch, D., Sinha, V.,
 691 Kumar, P., Mullins, B., Seguel, R., Shao, M., Xue, L., Wang, N., Chen, J., Gao, J., Chai, F.,
 692 Simpson, I., Sinha, B., and Blake, D. R.: A synergistic ozone-climate control to address
 693 emerging ozone pollution challenges, *One Earth*, 6, 964–977,
 694 <https://doi.org/10.1016/j.oneear.2023.07.004>, 2023.

695 Ma, J. and van Aardenne, J. A.: Impact of different emission inventories on simulated
 696 tropospheric ozone over China: a regional chemical transport model evaluation,
 697 *Atmospheric Chemistry and Physics*, 4, 877–887, <https://doi.org/10.5194/acp-4-877-2004>,
 698 2004.

699 Placet, M., Mann, C. O., Gilbert, R. O., and Niefer, M. J.: Emissions of ozone precursors from
 700 stationary sources:: a critical review, *Atmospheric Environment*, 34, 2183–2204,
 701 [https://doi.org/10.1016/S1352-2310\(99\)00464-1](https://doi.org/10.1016/S1352-2310(99)00464-1), 2000.

702 Prinn, R. G.: The Cleansing Capacity of the Atmosphere, *Annual Review of Environment and*
 703 *Resources*, 28, 29–57, <https://doi.org/10.1146/annurev.energy.28.011503.163425>, 2003.

704 Real, E., Megaritis, A., Colette, A., Valastro, G., and Messina, P.: Atlas of ozone chemical
 705 regimes in Europe, *Atmospheric Environment*, 320, 120323,
 706 <https://doi.org/10.1016/j.atmosenv.2023.120323>, 2024.

707 Russell, A.: REGIONAL PHOTOCHEMICAL AIR QUALITY MODELING: Model Formulations,
 708 History, and State of the Science, *Annual Review of Environment and Resources*, 22, 537–
 709 588, <https://doi.org/10.1146/annurev.energy.22.1.537>, 1997.

710 Saikawa, E., Kim, H., Zhong, M., Avramov, A., Zhao, Y., Janssens-Maenhout, G., Kurokawa, J.,
 711 Klimont, Z., Wagner, F., Naik, V., Horowitz, L. W., and Zhang, Q.: Comparison of emissions
 712 inventories of anthropogenic air pollutants and greenhouse gases in China, *Atmospheric*
 713 *Chemistry and Physics*, 17, 6393–6421, <https://doi.org/10.5194/acp-17-6393-2017>, 2017.

714 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to*
 715 *Climate Change*, John Wiley & Sons, 1152 pp., 2016.

716 Sha, Q., Zhu, M., Huang, H., Wang, Y., Huang, Z., Zhang, X., Tang, M., Lu, M., Chen, C., Shi, B.,
 717 Chen, Z., Wu, L., Zhong, Z., Li, C., Xu, Y., Yu, F., Jia, G., Liao, S., Cui, X., Liu, J., and Zheng, J.: A
 718 newly integrated dataset of volatile organic compounds (VOCs) source profiles and
 719 implications for the future development of VOCs profiles in China, *Science of The Total*
 720 *Environment*, 793, 148348, <https://doi.org/10.1016/j.scitotenv.2021.148348>, 2021.

721 [Silva, S. J. and Heald, C. L.: Investigating Dry Deposition of Ozone to Vegetation, *Journal of*](#)
 722 [*Geophysical Research: Atmospheres*, 123, 559–573,](#)
 723 [<https://doi.org/10.1002/2017JD027278>, 2018.](#)

724 Streets, D. G., Bond, T. C., Carmichael, G. R., Fernandes, S. D., Fu, Q., He, D., Klimont, Z.,
 725 Nelson, S. M., Tsai, N. Y., Wang, M. Q., Woo, J.-H., and Yarber, K. F.: An inventory of gaseous
 726 and primary aerosol emissions in Asia in the year 2000, *Journal of Geophysical Research:*
 727 *Atmospheres*, 108, <https://doi.org/10.1029/2002JD003093>, 2003.

728 Wang, C., Huang, X.-F., Han, Y., Zhu, B., and He, L.-Y.: Sources and Potential Photochemical
 729 Roles of Formaldehyde in an Urban Atmosphere in South China, *Journal of Geophysical*
 730 *Research: Atmospheres*, 122, 11,934–11,947, <https://doi.org/10.1002/2017JD027266>, 2017.

731 [Wang, P., Ying, Q., Zhang, H., Hu, J., Lin, Y., and Mao, H.: Source apportionment of](#)
 732 [secondary organic aerosol in China using a regional source-oriented chemical transport](#)
 733 [model and two emission inventories, *Environmental Pollution*, 237, 756–766,](#)
 734 [<https://doi.org/10.1016/j.envpol.2017.10.122>, 2018.](#)

735 West, J. J., Fiore, A. M., Horowitz, L. W., and Mauzerall, D. L.: Global health benefits of
 736 mitigating ozone pollution with methane emission controls, *Proceedings of the National*
 737 *Academy of Sciences of the United States of America*, 103, 3988,
 738 <https://doi.org/10.1073/pnas.0600201103>, 2006.

739 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and
 740 Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate
 741 the emissions from open burning, *Geoscientific Model Development*, 4, 625–641,
 742 <https://doi.org/10.5194/gmd-4-625-2011>, 2011.

743 Wu, R. and Xie, S.: Spatial Distribution of Ozone Formation in China Derived from Emissions
 744 of Speciated Volatile Organic Compounds, *Environ. Sci. Technol.*, 51, 2574–2583,
 745 <https://doi.org/10.1021/acs.est.6b03634>, 2017.

746 Xue, T., Zheng, Y., Geng, G., Xiao, Q., Meng, X., Wang, M., Li, X., Wu, N., Zhang, Q., and Zhu,
 747 T.: Estimating Spatiotemporal Variation in Ambient Ozone Exposure during 2013–2017

748 Using a Data-Fusion Model, *Environ. Sci. Technol.*, 54, 14877–14888,
 749 <https://doi.org/10.1021/acs.est.0c03098>, 2020.

750 Yamaji, K., Ohara, T., Uno, I., Kurokawa, J., Pochanart, P., and Akimoto, H.: Future prediction
 751 of surface ozone over east Asia using models-3 community multiscale air quality modeling
 752 system and regional emission inventory in Asia, *Journal of Geophysical Research:*
 753 *Atmospheres*, 113, 2008.

754 Yang, Z., Cheng, H. R., Wang, Z. W., Peng, J., Zhu, J. X., Lyu, X. P., and Guo, H.: Chemical
 755 characteristics of atmospheric carbonyl compounds and source identification of
 756 formaldehyde in Wuhan, Central China, *Atmospheric Research*, 228, 95–106,
 757 <https://doi.org/10.1016/j.atmosres.2019.05.020>, 2019.

758 Zhang, H., Li, J., Ying, Q., Guven, B. B., and Olaguer, E. P.: Source apportionment of
 759 formaldehyde during TexAQS 2006 using a source-oriented chemical transport model,
 760 *Journal of Geophysical Research: Atmospheres*, 118, 1525–1535,
 761 <https://doi.org/10.1002/jgrd.50197>, 2013.

762 [Zhang, K., Duan, Y., Huo, J., Huang, L., Wang, Y., Fu, Q., Wang, Y., and Li, L.: Formation](#)
 763 [mechanism of HCHO pollution in the suburban Yangtze River Delta region, China: A box](#)
 764 [model study and policy implementations, *Atmospheric Environment*, 267, 118755,](#)
 765 [<https://doi.org/10.1016/j.atmosenv.2021.118755>, 2021.](#)

766

767




Figure 1. Predicted monthly averages of MDA8 O₃ (O₃-8h) concentrations in July 2017 using (a) the S11 mechanism and MEIC emission inventory (base case), and the differences between the base case and cases using alternative photochemical mechanisms and emission inventories (alternative case – base case): (b) S11 and REAS, (c) S18 and REAS, (d) CS07 and MEIC, and (e) CS07 and REAS inventories. Units are ppb.





Figure 2. Predicted monthly averages of secondary inorganic aerosol (the sum of nitrate, sulfate and ammonium ion, SIA) concentrations in July 2017 using (a) the S11 mechanism and MEIC emission inventory (base case), and the differences between base case and cases using other photochemical mechanisms and emission inventories (alternative case – base case): (b) S11 and REAS, (c) S18 and REAS, (d) CS07 and MEIC, and (e) CS07 and REAS. Units are $\mu\text{g m}^{-3}$.



786

787

788 **Figure 3.** Predicted monthly averages of OH radical concentrations in July 2017 using (a) the
 789 S11 mechanism and MEIC emission inventory (base case), and the differences between the
 790 base case and cases using other photochemical mechanisms and emission inventories
 791 (alternative case – base case): (b) S11 and REAS, (c) S18 and REAS, (d) CS07 and MEIC, and
 792 (e) CS07 and REAS. Units are ppt. ($0.1 \text{ ppt} \sim 2.46 \times 10^6 \text{ molec cm}^{-3}$)




Figure 4. Predicted monthly averages of NO_3 radical concentrations in July 2017 using (a) the S11 mechanism and MEIC emission inventory (base case), and the differences between the base case and cases using alternative photochemical mechanisms and emission inventories (alternative case – base case): (b) S11 and REAS, (c) S18 and REAS, (d) CS07 and MEIC, and (e) CS07 and REAS. Units are in ppt. ($0.1 \text{ ppt} \sim 2.46 \times 10^6 \text{ molec cm}^{-3}$)




Figure 5. Predicted monthly averages of HCHO concentrations in July 2017 using (a) the S11 mechanism and MEIC emission inventory (base case), and the differences between the base case and cases using alternative photochemical mechanisms and emission inventories (alternative case – base case): (b) S11 and REAS, (c) S18 and REAS, (d) CS07 and MEIC, and (e) CS07 and REAS. Units are ppb.




Figure 6. Predicted changes of monthly average MDA8 O₃ (O₃-8h) concentrations in July 2017 due to reductions of NO_x only (first row), VOCs only (second row), and NO_x and VOCs (third row) using different photochemical mechanisms and emission inventories. The black line represents the average change across all mechanisms and inventories at different levels of emission reductions.




Figure 7. Predicted changes of monthly average secondary inorganic aerosol (nitrate, sulfate, and ammonium, SIA) concentrations in July 2017 due to reductions of NO_x only (first row), VOCs only (second row), and NO_x and VOCs (third row) using different photochemical mechanisms and emission inventories. The black line represents the average change across all mechanisms and inventories at different levels of emission reductions.




Figure 8. Predicted changes of monthly average OH radical concentrations in July 2017 due to reductions of NO_x only (first row), VOCs only (second row), and NO_x and VOCs (third row) using different photochemical mechanisms and emission inventories. The black line represents the average change across all mechanisms and inventories at different levels of emission reductions.




Figure 9. Predicted changes of monthly average NO_3 radical concentrations in July 2017 due to reductions of NO_x only (first row), VOCs only (second row), and NO_x and VOCs (third row) using different photochemical mechanisms and emission inventories. The black line represents the average change across all mechanisms and inventories at different levels of emission reductions.




Figure 10. Predicted changes of monthly average HCHO concentrations in July 2017 due to reductions of NO_x only (first row), VOCs only (second row), and NO_x and VOCs (third row) using different photochemical mechanisms and emission inventories. The black line represents the average change across all mechanisms and inventories at different levels of emission reductions.