

Assessment of the differences in European CH₄ emission estimates from three TROPOMI products

Aurélien Sicsik-Paré¹, Audrey Fortems-Cheiney^{1,*}, Isabelle Pison¹, Grégoire Broquet¹, Alvin Opler¹, Elise Potier^{1,*}, Adrien Martinez¹, Oliver Schneising², Michael Buchwitz², Joannes D. Maasakkers³, Tobias Borsdorff³, and Antoine Berchet¹

¹Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France

*now in Science Partners, Quai de Jemmapes, 75010 Paris, France

²Institute of Environmental Physics (IUP), University of Bremen FB1, Bremen, Germany

³SRON Netherlands Institute for Space Research, 2333 CA Leiden, The Netherlands

Correspondence: Aurélien Sicsik-Paré (aurelien.sicsik-pare@lsce.ipsl.fr)

Abstract. Satellite observations from the Sentinel-5P TROPOMI instrument, combined with inverse modeling, provide a valuable resource for quantifying regional methane (CH₄) emissions. This study compares the 2019 European emissions estimated from variational inversions assimilating three TROPOMI products of dry-column methane mole fractions (XCH₄). The SRON (v2.4, operational product), BLENDED (v1.0), and WFMD (v1.8) products are retrieved from distinct algorithms.

- 5 These retrievals differ in coverage, error characterization, and XCH₄ spatial distribution. Machine learning predictions of XCH₄ differences point to aerosol scattering and albedo sensitivity as the largest contributors to the differences. The derived 2019 European CH₄ emission budgets show an increase of +2% for SRON, and a decrease of -1%, -33% and -9% relative to the prior, respectively, for BLENDED, WFMD and surface-based inversions. The range of budget estimates reveals inconsistencies in the total emissions derived from the inversions. At the national and sub-national scale, spatial emission patterns are similar
- 10 for the non-independent SRON and BLENDED but differ substantially from WFMD. No inversion provides a systematically closer match to the spatial distribution of emissions derived from independent surface observations. Evaluation at surface stations shows that the residual between the observations and the posterior concentrations is reduced for 37%, 53% and 47% of the stations, respectively, for SRON, BLENDED and WFMD. Observing System Simulation Experiments (OSSEs) are used to disentangle the drivers of differences between the posterior emissions. Results show that aligning coverage and individual
- 15 observation errors increases the consistency between emission estimates. Residual differences in the OSSE posterior emissions can be attributed to differences in averaging kernels and prior profiles.

1 Introduction

The global emission pathways to limit global warming below the international objective of 1.5°C (IPCC, 2021) include significant reductions of methane (CH₄) emissions. Methane is the second most important anthropogenic greenhouse gas (GHG) after carbon dioxide (CO₂). Despite its relatively short atmospheric lifetime of 9.1 ± 0.9 years (Szopa et al., 2023), it has a strong radiative efficiency (Forster et al., 2021). The accounting of emissions is required to assess the current regulatory policies and to provide a robust reference for projections. For that, inverse modeling of CH₄ emissions combines atmospheric observations and transport simulations. This top-down (TD) approach complements bottom-up (BU) emission reporting by using independent information provided by atmospheric mixing ratios to reduce uncertainties on the emissions.

In the last decade, satellite data of CH₄ total column dry air mixing ratios (XCH₄) (Monteil et al., 2013; Cressot et al., 2014; Alexe et al., 2015; Bergamaschi et al., 2015) have provided wider spatial coverage than the surface data but with a double challenge: the estimation of the XCH₄ retrieval from the raw spectroscopic measurement, and the estimation of emissions based on the assimilation of retrievals. Satellite instruments can be divided into area flux mappers, designed to observe emissions at the global or regional scale, and point source imagers (Jacob et al., 2022). Point source imagers are fine-pixel instruments designed to quantify the point source emissions from the observation of the CH₄ plume. They are used to build datasets of local emitters (Schuit et al., 2023), available in public interactive tools such as the CAMS Methane Hotspot Explorer app and the Methane Alert and Response System (United Nations Environment Programme, 2025). Area flux mappers of methane include SCIAMACHY, launched on-board ENVISAT in 2002, the first instrument to provide global methane observations (Bovensmann et al., 1999); TANSO-FTS, launched in 2009 on-board the Greenhouse Gases Observing Satellite (GOSAT), that provides relatively accurate but sparse XCH₄ observations (Parker et al., 2020); IASI, on-board the MetOp satellites, a thermal-infrared (TIR) interferometer (Dils et al., 2024). The TROPOspheric Monitoring Instrument, also known as TROPOMI (Veefkind et al., 2012), launched in October 2017 on-board the satellite Sentinel-5P, now provides XCH₄ with a nadir resolution of $5.5 \text{ km} \times 7 \text{ km}$ and daily global coverage (Hu et al., 2016; Hasekamp et al., 2022). Its high resolution (relative to other instruments) together with its high coverage make it possible to quantify both point source and regional/global CH₄ emissions. TROPOMI retrievals have been successfully used to detect large releases from oil and gas facilities (Schneising et al., 2020; Zhang et al., 2020; Lauvaux et al., 2022; Veefkind et al., 2023), as well as coal mines and landfills (Schuit et al., 2023) and large persistent source regions (Vanselow et al., 2024). They have also been used to quantify national and sectoral emissions, in global (Qu et al., 2021; Yu et al., 2023; East et al., 2025) and regional inversions for the US (Shen et al., 2022; Nesser et al., 2024), the Middle East and North Africa (Chen et al., 2023), East Asia (Chen et al., 2022; Liang et al., 2023) and South America (Nathan et al., 2023; Hancock et al., 2024). Small sources account in aggregate to a large proportion of total emissions at the national scale (Williams et al., 2025): bridging the gap between the local emission estimates and aggregated national budgets is a current challenge (Santaren et al., 2021; Naus et al., 2023).

The TROPOMI XCH₄ total columns are retrieved from SWIR radiance measurements along with the averaging kernels (AKs), that assess the sensitivity of the retrievals to the different atmospheric layers. This derivation is a major challenge: the retrieved columns and the vertical sensitivity indeed strongly depend on the chosen algorithm and on different factors of the

radiative transfer (e.g., clouds, the assumed prior profile of the CH₄ dry air mixing ratio, surface albedo and aerosols). The systematic analysis of the TROPOMI XCH₄ data has pointed to biases linked to albedo and scattering due to the presence of aerosols (Barré et al., 2021).

In this context, several products of TROPOMI XCH₄ retrievals have been developed with different algorithms, and they are iteratively updated. The SRON Netherlands Institute for Space Research provides the operational Copernicus product (European Space Agency, 2021), which has been updated with the optimized settings of the research product (Lorente et al., 2021, 2023). It uses the full-physics algorithm RemoTeC and simultaneously retrieves XCH₄, surface albedo and atmospheric scattering properties. A destriping algorithm, based on moving median smoothing in the across-track directions and flight directions, has been developed for new XCH₄ data (Borsdorff et al., 2024). The BLENDED TROPOMI+GOSAT product (Balasus et al., 2023) is a corrected version of the SRON product. It relies on a machine learning model, trained to predict the differences between TROPOMI and GOSAT co-located retrievals. The modeled correction was applied to the whole SRON dataset. The scientific WFMD product from University of Bremen is independent of the other two and is based on the algorithm Weighting Function Modified Differential Optical Absorption Spectroscopy (WFMD-DOAS) (Schneising et al., 2019, 2023).

Previous inter-comparisons only include the SRON and WFMD products (T. Hilbig et al., 2023). A comparison of observations at high latitudes (Lindqvist et al., 2024) revealed higher XCH₄ for WFMD in comparison to SRON, as well as a persistent seasonal bias for the SRON (high values in spring, low values in autumn). The assimilation of older versions of SRON and WFMD products at high latitudes showed similarities but also clear differences in the posterior emissions, in terms of spatial and temporal distributions (Tsuruta et al., 2023). Despite the advancements in XCH₄ retrievals in recent updates and the additional product BLENDED, the literature on systematic comparisons between the products and between the estimated emissions remains limited.

This study aims at addressing these limitations by providing a comparison of CH₄ emissions estimated from the assimilation of the SRON, BLENDED and WFMD products in regional inversions. The objective is to assess the consistency of emission estimates, with respect to the aim of building emission inventories at the pixel, country and continental scale. This benchmark requires the characterization of the (in)consistencies in the observation coverage, observation errors and spatial biases of XCH₄ between the three datasets. The study covers the year 2019 and focuses on Europe, where an extensive network of surface stations provide independent CH₄ measurements that are used to support the comparison.

First, we compare the three products over Europe in 2019. We investigate the role of atmospheric, instrumental and algorithmic variables in driving the XCH₄ differences between products. Using a machine learning model, feature contributions to the prediction of the differences are quantified, following the methodology of Balasus et al. (2023).

Subsequently, we perform variational inversions to estimate the CH₄ fluxes at a $0.5^\circ \times 0.5^\circ$ spatial resolution and weekly temporal resolution by assimilating each product independently. We conduct Observing System Simulation Experiments (OSSEs) with synthetic pseudo-observations and perturbed prior fluxes. The OSSEs assess the ability of the inversion system to refine emission estimates and its sensitivity to parameters such as observation density, error characteristics, and inter-product differences. Scenarios focusing on specific changes provide insights into the drivers of differences in the posterior fluxes. We also investigate the role of the optimization of the boundary conditions and the stratospheric concentrations, but do not fur-

ther address the question of model errors and their potential correlations with retrieval errors. Finally, we compare European CH_4 emission estimates in 2019 from the three TROPOMI products and ground-based observations, at the pixel, country, and regional resolution, to assess the consistency of the results at these various scales. For this study, we use the recent inverse modeling platform Community Inversion Framework (CIF Berchet et al., 2021), coupled with the regional Eulerian atmospheric chemistry-transport model (CTM) CHIMERE (Menut et al., 2013; Mailler et al., 2017) and its adjoint code (Fortems-Cheiney et al., 2021). CHIMERE has already been used to model transport of GHGs at the regional scale, especially CO_2 (Broquet et al., 2011; Santaren et al., 2021) and CH_4 (Pison et al., 2018). The methane observations, the configuration of the CHIMERE CTM and the methodology for the variational inversions and OSSEs are described in Section 2. The results, particularly those from the inversions and the OSSEs, are detailed in Section 3.

The purpose of our atmospheric inversions is to correct prior estimates of the CH₄ emissions, to improve the fit between the simulations of CH₄ atmospheric mixing ratios by a CTM and the observations. In this section, we detail the inputs and components of the CIF-CHIMERE inversion system. Section 2.1 provides an overview and a comparison of the coverage, distribution and errors of the three TROPOMI products. It also introduces the other CH₄ observations used in this study. 100 Section 2.2 describes the prior emission estimates, and Section 2.3 details the configuration of the CHIMERE CTM. The methodology for the inversions and OSSEs is outlined in Section 2.4.

2.1 CH₄ observations

2.1.1 TROPOMI satellite products

TROPOMI is on-board Sentinel-5 Precursor (S5P) since 2017. The radiances measured in the shortwave infrared range (SWIR, 105 2314 – 2382 nm) are used to derive methane total columns. The pixel size at nadir was about 7.2×7.2 km² before 2019/08/06. It was upgraded to 5.6×7.2 km² after a change in the instrument settings on this date. The swath is about 2600 km on ground, allowing global daily coverage. The accuracy requirements for XCH₄ total columns are 1% bias and 1% random error, as defined in the S5P Calibration and Validation Plan (2017). In this study, the assimilated methane retrievals consist of three level 2 (L2) products, described in the following sections. Table 1 shows the number of observations available in the domain 110 in 2019 for each of these products.

Product	Obs count	Of which ocean
SRON	4,731,034	448,721 (9.5%)
BLENDED	4,731,022	448,720 (9.5%)
WFMD	7,627,595	283,722 (3.7%)
In common	3,432,335	106,775 (3.1%)
GOSAT	12,841	790 (6.2%)

Table 1. Number of available observations over the domain in 2019, for the three TROPOMI products SRON, BLENDED and WFMD used in this study, the "common" dataset (composed of the observations shared across the three TROPOMI products), and GOSAT.

SRON

The operational TROPOMI product (v2.4), hereafter called "SRON", is developed by the SRON Netherlands Institute for Space Research. It includes the iterative improvements of the research product (Lorente et al., 2021, 2022), in particular better 115 accounting for surface reflectance (Lorente et al., 2023). In this study, we use the reprocessed albedo bias-corrected version and apply the recommended filtering criteria (Apituley et al., 2024) to ensure high-quality data (quality flag superior to 0.5). This product is retrieved from TROPOMI measurements with the RemoTeC full-physics algorithm, that retrieves both the atmospheric methane mixing ratio and the physical scattering properties of the atmosphere. A detailed description of the algorithm is given in the Algorithm Theoretical Baseline Document (Hasekamp et al., 2022). A destriping procedure (Borsdorff

et al., 2024) is applied to new XCH₄ data from 2024/09/07 (v2.07), but older orbits have not been reprocessed. This empirical approach consists in removing the CH₄ background by a median smoothing in the cross-track direction, and then computing a per orbit stripe value as a median in the flight direction, which is used for correction (Borsdorff et al., 2019). However, this processing is not applied to the 2019 observations used in this study.

Blended GOSAT+TROPOMI product

The BLENDED product (Balasus et al., 2023) is a corrected version of SRON: a machine learning (ML) model has been trained to predict the differences between GOSAT and TROPOMI-SRON (v2.4) observations, based on SRON data features. The GOSAT XCH₄ product uses a proxy retrieval which is less sensitive to surface and atmospheric artifacts. It is described in further detail in Section 2.1.2. The global mean bias versus TCCON data (9.2 ppb) is subtracted from all GOSAT observations (Parker et al., 2020). The correction is modeled on co-located observations, then applied to the whole TROPOMI-SRON record. This method aims at taking advantage of the density of TROPOMI measurements, while mitigating the effect of known biases associated to surface albedo, coarse aerosol particles and striping, thanks to the GOSAT proxy approach. We keep only the highest quality data (quality flag of 1) and filter for coastal scenes as recommended by Balasus et al. (2023). This quality filter is theoretically stricter than the one recommended for the SRON product, but in practice it eliminates only 12 observations over the domain in 2019. It can thus be considered that both SRON and BLENDED products share the same observation sampling (see Table 1). In the BLENDED product, the observation errors, averaging kernels and prior profiles are directly taken from the SRON product (only the XCH₄ values differ between the two products).

WFMD

The WFMD scientific product v1.8 (Schneising et al., 2019, 2023), hereafter called "WFMD", is based on the Weighting Functions Modified Differential Optical Absorption Spectroscopy (WFMD-DOAS). Quality filtering is based on a random forest (RF) classifier: data with a quality flag of 0 is selected. Post-processing includes systematic bias correction (eg. due to albedo) based on a RF regressor, as well as a destriping filter based on combined wavelet–Fourier filtering.

Common observations

To separate the effect of varying coverage between products from the effect of the differences in the XCH₄ distributions, we extract a dataset of observations shared across all three products, so that they have identical spatio-temporal sampling. This "common" dataset is used for the analysis of the inter-product differences (Section 3.1) and for OSSEs (Sections 2.4.3, 3.4).

2.1.2 Other CH₄ observations

In Section 2.1.3, we compare TROPOMI observations with retrievals from the University of Leicester GOSAT Proxy XCH₄ product, v9.0 (Parker et al., 2020). The GOSAT satellite was launched in 2009. Methane estimates are based on the CO₂ proxy method, which takes advantage of CO₂ absorption in the measured 1.65 μm band, with a finer spectral resolution than TROPOMI. This approach significantly reduces the error in the retrieval, since it cancels aerosol and surface artifact

contributions that similarly affect XCO₂ and XCH₄ measurements. Only highest-quality data (quality flag of 0) is considered, yielding 12,841 observations over the domain in 2019. The global mean bias versus TCCON data is removed from the GOSAT observations shown in the following, subtracting 9.2 ppb from all retrievals, to be consistent with the dataset used by Balasus et al. (2023) to derive the BLENDED observations (see Section 2.1.1).

To evaluate the results of the inversions, we use independent in-situ measurements of CH₄ mixing ratios from flasks and continuous sampling sites. The 19 available surface sites for 2019 are listed in Table A1. The data are compiled from the ICOS atmospheric network (<https://www.icos-cp.eu>), the World Data Centre for Greenhouse Gases (WDCGG, <https://gaw.kishou.go.jp>), the NOAA ESRL discrete sampling network (<https://www.esrl.noaa.gov/gmd/>) and the EBAS data base. Data from sites with continuous measurements are averaged to hourly values. The evaluation consists in comparing the emissions derived from the satellite-based inversions to an inversion assimilating only these surface measurements (Section 3.3). We also compare the posterior simulated concentrations at the locations of the surface stations to these independent measurements (Section 3.5).

2.1.3 Comparison of the TROPOMI XCH₄ coverage, distribution and errors

Coverage - The inter-comparison of TROPOMI satellite retrievals reveals differences in coverage over Europe in 2019. The density of observations is a critical parameter for ensuring robust constraints on emissions during the inversion. Yet, substantial differences exist between SRON/BLENDED and WFMD, as shown in Table 1. WFMD exhibits 63% more observations in comparison to SRON and BLENDED over Europe in 2019. This higher observation count is attributable to the different filtering approaches: as mentioned in Section 2.1.1, WFMD employs machine learning techniques to identify scenes that are poorly characterized by the retrieval algorithm (Schneising, 2023); quality control criteria for SRON and BLENDED rely on threshold-based filters for cloud fraction, solar zenith angle and other scene description variables (Landgraf et al., 2024).

Spatial patterns of coverage differences are illustrated in Figure 1b. The WFMD dataset demonstrates a higher density of observations in arid regions (e.g., over Spain and Turkey). Moreover, it includes a few observations in mountainous areas (Alps, Norway), whereas SRON and BLENDED filter out all retrievals in these regions.

The temporal coverage shows consistent variations for the three products (Figure 1a): observation density is high between June and October and decreases during winter. An anomalously low density is noticeable for all products in May 2019. It is due to an increase of the Fractional Cloud Cover (CFC) over Europe during this month, as indicated by the CLARA-A3 record (Karlsson et al., 2023). Seasonal spatial variations are consistent among products, with an absence of observations at high latitudes ($\gtrsim 60^\circ$) during winter months (see Figure S1 in the Supplement).

Comparison of TROPOMI observations - The temporal and spatial XCH₄ distributions of the three products are shown in Figure 2. Average observed XCH₄ in 2019 is respectively 1850, 1843 and 1856 ppb for SRON, BLENDED and WFMD (Figure 3). Part of the differences between the distributions can be explained by the differences in spatial coverage between SRON/BLENDED and WFMD. When considering only the common observations across the three datasets, the average XCH₄ are slightly closer but still differ (1851, 1844 and 1856 ppb respectively), and the analysis of space-time distributions remains similar.




Figure 1. Number of observations per month (panel **a**) and per 0.5° longitude $\times$ 0.5° latitude cell 2019 (panel **b**) for the three TROPOMI products SRON, BLENDED and WFMD used in this study. The covered period includes the change in pixel size (7.2 km to 5.6 km) in the along track direction, starting on the 6th of August 2019.

The temporal variations of XCH_4 throughout the year show overall similar shapes (Figure 2a). Average concentrations decrease slightly from January to May, reaching a minimum in April/May, before rising during late summer and autumn, peaking in November. The difference between BLENDED and WFMD monthly averaged values is steady over the year (-11.7 ± 2.1 ppb). The differences between SRON and BLENDED (6.8 ± 2.3 ppb), as well as SRON and WFMD (-5.0 ± 2.5 ppb) depict more temporal variations: SRON observations tend to be closer to GOSAT and BLENDED in winter, but closer to WFMD in summer. Only BLENDED aligns rather well with GOSAT, due to its correction of the TROPOMI-GOSAT bias.

Overall, the spatial patterns of the distributions of CH_4 concentrations are consistent, yet with local discrepancies, e.g., in Scandinavia (higher XCH_4 than SRON for WFMD and lower for BLENDED, as seen in Figure 2b). BLENDED shows lower XCH_4 over the domain, but also consistent spatial gradients with SRON and WFMD (Figures 2b, S2).

To understand the drivers of the differences in XCH_4 distributions, we adopt the method outlined by Balasus et al. (2023). A machine learning (ML) model is trained to predict the difference in XCH_4 based on the retrieval variables. The contribution of individual features to the prediction is assessed to estimate the importance of each variable. The results and analysis of the biases are presented in Section 3.1. Detailed study of the differences of XCH_4 relative to TCCON is available in Appendix C.




Figure 2. Average XCH_4 observation per month (Panel **a**) and per $0.5^\circ \times 0.5^\circ$ cell in 2019 (Panel **b**) for the three TROPOMI products.

200 **Observation error** - The retrieval error (in terms of uncertainty) attributed to individual observations can impact the emissions from inversions by changing their spatial and temporal variations (Lu et al., 2025b), as well as call into question both very high and very low emission values (Zheng et al., 2026). The definition of this error differs between the SRON/BLENDED and WFMD products. For SRON (thus also for BLENDED, for which only XCH₄ values are corrected), the reported error is defined as the standard deviation of the retrieval noise, which follows from the error covariance matrix in the retrieval inversion
 205 procedure. This error describes the effect of the noise in the measured radiances on the retrieval (Lorente et al., 2021). For WFMD, the error is propagated in the retrieval algorithm from the noise in the measured spectra. In both cases, the characterized error σ only includes the effect of noise in the measured radiance, however the unknown noise components related to atmospheric conditions or instrumental features are not accounted for.

To avoid underestimating the observation uncertainty, the products have different error corrections, based on the validation
 210 of the satellite XCH₄ with respect to total columns observations of the ground-based TCCON stations. SRON and BLENDED provides the error σ as described before and suggest to multiply it with a factor 2 to reflect the scatter of errors in the TCCON validation (Landgraf et al., 2024). We apply this recommended correction. For WFMD, the error σ is linearly rescaled, based on a regression of the scatter relative to TCCON observations (Equation 1). The scaled error $\hat{\sigma}$ is directly provided in the product and described in the Algorithm Theoretical Baseline Document (Schneising, 2023).

$$215 \quad \hat{\sigma} = 4/3(\sigma + 5\text{ppb}) \quad (1)$$

Due to this difference in error definition between SRON/BLENDED and WFMD, the errors of SRON and BLENDED (3.9 ppb in average) are smaller by a factor of 3 than those of WFMD (12.2 ppb) (Figure 3b). If we try to apply the linear transformation of Equation 1 to SRON/BLENDED errors, the resulting error distribution aligns more closely with WFMD: the average scaled error is 11.9 ppb for SRON and BLENDED (Figure 3c). Therefore, the difference in rescaling the error based
 220 on the scatter relative to TCCON is the main source of difference in the final uncertainty provided in the products. The impact of this difference on the results of inversions is further evaluated in Section 3.4.

Vertical parameters - As illustrated in Figure 4 (only the common observations are considered here), the column averaging kernels (AKs) and prior profiles exhibit similar shapes, characteristic of SWIR retrievals. The per level relative difference between SRON/BLENDED and WFMD vertical profiles is below 5% for all levels. The WFMD AKs seem to be less sensitive
 225 to the layers close to the surface in comparison to SRON/BLENDED, but more sensitive to the stratosphere (pressures inferior to 200 hPa). Differences up to approximately 200 ppb occur between the prior profiles of SRON/BLENDED and WFMD, in particular for the layers close to the surface: the WFMD prior profiles are scaled to have uniform surface values of 1850 ppb, while SRON/BLENDED prior mixing ratios at the surface are between 1900 and 2000 ppb. The scaling of prior profiles for WFMD also explains the lower variability of prior mixing ratios in comparison to SRON/BLENDED, as evidenced by the
 230 narrower horizontal deviations in the right panel of Figure 4. These profiles are consistent with the ones presented in Lindqvist et al. (2024), which also highlight a systemically higher prior profile for SRON in comparison to the one of WFMD. The differences in the profiles lead to differences in the simulated XCH₄ distributions, as shown in Section 3.2.




Figure 3. Distributions of observations and observation errors (both in ppb) for the three TROPOMI products. For the error, panel (b) shows the raw errors, while panel (c) shows the same histogram after error scaling for the SRON and BLENDED products. SRON and BLENDED errors are exactly similar, as BLENDED errors are directly retrieved from the SRON product.




Figure 4. Column averaging kernels (left, unitless) and prior profiles (right, in ppb) averaged between the common observations of the TROPOMI products (dataset described in Section 2.1.1). The zoomed windows show the lower levels. SRON and BLENDED have exactly the same profile, since the only difference in the co-located dataset is the XCH₄ value. Horizontal lines are CHIMERE pressure levels (plain lines) and CAMS pressure levels in the stratosphere (dashed lines, for pressures lower than 200 hPa), for a surface pressure of 1000 hPa.

2.2 Prior estimates of the CH₄ emissions

Prior methane emissions are compiled from several BU inventories. Anthropogenic emissions are from EDGARv8.0 (Crippa et al., 2023). The biomass burning fluxes are taken from GFEDv-4.1s (Randerson et al., 2018). Natural fluxes consist of an ensemble of datasets provided by the Global Methane Budget protocol for inversions (GCP-CH₄, Saunois et al., 2020). Fluxes from wetlands (peatlands, inundated and mineral soils) are from the JSBACH-HIMMELI model (Raivonen et al., 2017). The

geological emissions are a climatology based on Etiope et al. (2019) and scaled down to a global total of 15 TgCH₄ yr⁻¹ in accordance with the maximum suggested by Petrenko et al. (2017). The emissions due to termites are a climatology based on the estimate of S. Castaldi, from GCP-CH₄ (Saunois et al., 2020). Finally, the ocean fluxes are a climatology based on Weber et al. (2019). All these datasets have been interpolated at the 0.5°×0.5° horizontal resolution of the CTM grid. The map of total emissions in 2019 is shown in Figure 5. Anthropogenic emissions contribute to about 72% of the total emission budget (25.2 TgCH₄ in 2019) for the countries listed in Table B1.




Figure 5. Prior CH₄ emissions ($\text{kg.m}^{-2}.\text{h}^{-1}$) in Europe in 2019, in log-scale. Black triangles are the locations of the surface stations listed in Table A1.

2.3 Configuration of the CHIMERE CTM for the simulation of CH₄ concentrations

The CTM used for inversion is the regional chemistry-transport model CHIMERE (Menut et al., 2013; Mailler et al., 2017) and its adjoint code (Fortems-Cheiney et al., 2021). The targeted domain spans from 15°W to 35°E and 32°N to 74°N. This domain has already been used for the intercomparison of inversions EUROCOM (Monteil et al., 2020). The grid used for discretizing emissions, meteorological data and other inputs and for running the CTM covers this domain at a 0.5° longitude × 0.5° latitude resolution. Vertically, the domain where transport is simulated by CHIMERE extends from the surface to 200 hPa (with 17 sigma-pressure levels). Above this pressure (approximately the tropopause) and up to 0.1 hPa, CH₄ concentration fields are taken from the CAMS reanalysis simulations (based on the Integrated Forecasting System (IFS) system of the European Centre for Medium-Range Weather Forecasts (ECMWF)) (Agustí-Panareda et al., 2023). Lateral boundary and initial conditions of methane mixing ratios are taken from the same CAMS product.

CHIMERE requires a set of meteorological variables, taken from the ECMWF IFS operational forecast (every three hours) retrieved at $0.25^\circ \times 0.25^\circ$ and interpolated onto the model's grid. The size of the domain makes it possible to neglect the chemistry of CH_4 because its oxidation by hydroxyl radicals leads to a lifetime from 8 to 10 years (Saunois et al., 2020), whereas the ventilation time of the domain is of the order of 10 days (Nygård et al., 2023).

2.4 Variational inversions in the CIF-CHIMERE inversion system

2.4.1 Principle of Bayesian variational inversion

In the following, we use notations according to the convention defined by Ide et al. (1997) and Rayner et al. (2019). The 4D-Var inversion adjusts the control vector $\mathbf{x}$, to improve the fit between observed satellite data and their simulated equivalents, by minimizing the cost function J :

$$J(\mathbf{x}) = \frac{1}{2}(\mathbf{x} - \mathbf{x}^b)^T \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}^b) + \frac{1}{2}(\mathcal{H}(\mathbf{x}) - \mathbf{y}^0)^T \mathbf{R}^{-1}(\mathcal{H}(\mathbf{x}) - \mathbf{y}^0) \quad (2)$$

where $\mathbf{x}^b$ and $\mathbf{y}^0$ are respectively the vector of prior information and the vector of observations, $\mathcal{H}$ the observation operator, $\mathbf{B}$ and $\mathbf{R}$ the covariance matrices of the control vector and observation errors. The latest combines errors in both the observation data (measurement or processing errors) and the observation operator (model error, representativity of the gridded model compared to point measurements, aggregation errors).

2.4.2 Variational inversion procedure

The control vector $\mathbf{x}$ contains CH_4 emissions, at a $0.5^\circ \times 0.5^\circ$ horizontal resolution and a weekly temporal resolution (similar to Petrescu et al. (2024) and Szénási et al. (2021)). It also contains the 4D CH_4 background field used to impose the initial, lateral and top boundary conditions as well as the concentrations in the stratosphere (above 200 hPa) at the native pixel resolution of the corresponding CAMS product ($3^\circ \times 2^\circ$) and 2-day temporal resolution. Indeed, the assimilated data is used by the inversion to retrieve information on emission fluxes but also on the background.

The error covariance matrix $\mathbf{B}$ is built by blocks, the errors on the emissions and the background being considered independent. For the CH_4 emission components, the relative error standard deviation (diagonal elements of $\mathbf{B}$) are set at 100%, similarly as Ioannidis et al. (2025). Spatial and temporal correlations are built with an e-folding decrease with correlation lengths of 150 km on land and 200 km on sea, and 2 weeks through time. For CH_4 background concentrations, we assume a 2% relative error standard deviation for diagonal elements. The non-diagonal elements account for spatial and temporal covariances, using a similar e-folding decrease with a spatial correlation length of 200 km and a temporal correlation length of 14 days.

The matrix $\mathbf{R}$ is diagonal: we consider no correlation of the errors from one observation to another. The diagonal elements are the errors associated with the individual retrievals, as described in Section 2.1.3.

For the inversions, additional filters are applied to TROPOMI observations (Section 2.1.1) to avoid large differences between observed and simulated XCH_4 caused by the model limitations. The CTM cannot capture subpixel pressure variations: to mitigate the impact of subpixel topographic variability, we remove observations with surface pressure deviating by more than 3σ from the mean pressure of the data aggregated at the scale of the CTM pixel. This filter eliminates 1.4% of the observations for SRON and BLENDED, 1.0% for WFMD. Data for which the difference between the observation and the simulation is more than 100 ppb are also filtered out. It impacts very few observations, removing respectively 420 observations for SRON/BLENDED and 934 for WFMD, i.e 0.009% and 0.012% of the data. The spatial distributions of the observations removed by these two filters are similar across the three products (Figure D1).

In this study, we perform four 4D-Var inversions assimilating real observations, using the CIF. The CIF is a modular inverse modeling platform developed as a python library (Berchet et al., 2021), designed in the framework of European and international projects. It can drive various data assimilation schemes and can be coupled to various CTMs. Here, the cost function J is minimized using the quasi-newtonian M1QN3 algorithm (Gilbert and Lemaréchal, 1989). The inversion is stopped when the gradient norm reduction exceeds 95% compared to its initial value. The first three inversions assimilate separately each one of the TROPOMI products. An additional inversion assimilating surface observations (stations listed in Table A1) is performed for evaluation. These four inversions are listed in Table 2.

ID	Observation type	Observation product
Inv-SRON	Satellite	SRON
Inv-BLD	Satellite	BLENDED
Inv-WFMD	Satellite	WFMD
Inv-Surface	Surface	Hourly data/flasks from the stations listed in Table A1

Table 2. Synthesis of the inversions performed in this study. Results are presented in Section 3.3.

2.4.3 Observing System Simulation Experiments

We also explore the capability of the inversion set-up to derive robust emission estimates based on the assimilation of TROPOMI products in a set of Observing System Simulation Experiments (OSSEs). The OSSE approach (Lahoz and Schneider, 2014) makes it possible to evaluate the sensitivity of the inversions to the observation coverage, the observation errors and the differences between products, comparing the results of multiple OSSE scenarios. The structure of an OSSE is described in Figure 6. The general principle consists of two main parts (Brasseur and Jacob, 2017): first, the sampling of "true" synthetic observations, given a true state defined by the prior emissions and background concentrations. Then, a Monte-Carlo ensemble of inversions with perturbed priors is performed, to evaluate the capability of the observing system to recover the true state of the emission and background estimates.

In more details, a forward run of the CHIMERE CTM produces a "true" 4D concentration field, based on the prior emissions and the prior background concentrations (called $\mathbf{x}^t$). Pseudo-observations $\mathbf{y}^0$ are sampled from this field following the specifications (date, location, averaging kernels...) of the observation dataset.




Figure 6. OSSE structure within the CIF. "True" pseudo-observations $\mathbf{y}^0$ are synthesized from the prior emissions $\mathbf{x}^t$ and then assimilated in an ensemble of inversions using perturbed prior fluxes $\mathbf{x}_i^b$. See <http://community-inversion.eu>.

310 The pseudo-observations are assimilated in an ensemble of $n = 4$ inversions using randomly perturbed priors: for each inversion member i , the priors are initially perturbed (called $\mathbf{x}_i^b$) according to the error statistics provided in $\mathbf{B}$ (Section 2.4.2):

$$\forall i \in [1, n], \mathbf{x}_i^b = \mathbf{x}^t + \boldsymbol{\eta}_i, \quad \boldsymbol{\eta}_i \sim \mathcal{N}(0, \mathbf{B}) \quad (3)$$

The metrics used to assess the capability of the system to improve the emission estimates (i.e. to make them closer to the truth) is the relative increment r . It compares the distance to the true initial fluxes $\mathbf{x}^t$ of the perturbed prior $\mathbf{x}_i^b$ and the posterior fluxes

315 $\mathbf{x}_i^a$, averaged over all the members of the ensemble:

$$r = \frac{1}{n} \sum_i r_i = \frac{1}{n} \sum_i \left(\left| \frac{\mathbf{x}_i^a - \mathbf{x}^t}{\mathbf{x}_i^b - \mathbf{x}^t} \right| - 1 \right) \quad (4)$$

The closer r is to -100% , the more the inversion improves the fluxes. Negative values of r correspond to posterior emissions that are closer to the truth than the prior (thus "better" emissions) while positive values correspond to posterior emissions further from the truth.

320 We define 6 OSSE scenarios, listed in Table 3, using different pseudo-observation datasets. To ensure a robust comparison of the scenarios, the seed of the random perturbations is kept constant. It is important to note that since SRON and BLENDED share the same sampling and parameters (errors, vertical profiles), the pseudo-observation datasets are identical. In the following, "SB" scenarios stand for the two products. The 6 scenarios are designed to evaluate the sensitivity of the inversions to multiple parameters:

- 325 – 3 OSSEs are performed as reference scenarios using the pseudo-observations of the TROPOMI products: "*Ref-SB*" for SRON and BLENDED, "*Ref-WFMD*" for WFMD; a third reference OSSE ("*Ref-Surf*") is performed using the pseudo-observations of the surface stations.
- To evaluate the sensitivity to the observation density, 2 OSSEs are carried out assimilating only the pseudo-observations corresponding to the common observations (as described in Section 2.1.1). The pseudo-observation datasets differ only
 330 by the observation errors and vertical profiles (prior profiles and averaging kernels). These scenarios are referred to as "*Common-SB*" and "*Common-WFMD*".
- To evaluate the sensitivity to observation errors, one scenario is defined using the SRON/BLENDED pseudo-observations dataset, rescaling the errors with Equation 1. It is called "*Err-SB*".

To evaluate the impact of the differences between products, we carry out two more inversions assimilating pseudo-observations.
 335 The method differs from previous OSSEs: in this case, synthetic observations are sampled and biased with the difference between products:

- The difference WFMD-SRON (computed over the common observations) is averaged over $0.5^\circ \times 0.5^\circ$ pixels and 1-hour-long periods, and added to the corresponding SRON pseudo-observations. The same correction is applied to WFMD pseudo-observations with respect to SRON-WFMD difference. To evaluate the impact of the inter-product difference
 340 on the posterior increments, each "corrected" data set is assimilated (priors are not perturbed), the inversions are called "*Diff-SRON*" and "*Diff-WFMD*".

ID	Product	Obs subset	Error correction	Difference correction
Ref-SB	SRON/BLENDED	All obs	No	No
Ref-WFMD	WFMD	All obs	No	No
Ref-Surf	Surface	All obs	No	No
Common-SB	SRON/BLENDED	Common obs	No	No
Common-WFMD	WFMD	Common obs	No	No
Err-SB	SRON/BLENDED	Common obs	Yes	No
Diff-SRON	SRON/BLENDED	Common obs	No	Yes (w.r.t. WFMD)
Diff-WFMD	WFMD	Common obs	No	Yes (w.r.t. SRON)

Table 3. Synthesis of all the OSSEs performed in this study. "Common obs" refers to the restriction to observations common to the 3 products, as described in Section 2.1.3. The "error correction" corresponds to the application of Equation 1 to the errors of SRON/BLENDED.

3 Results

3.1 Drivers of the differences in the XCH_4 observed distributions

To better understand the inter-product differences related to atmospheric, instrumental and algorithmic variables, we adopt the method of Balasus et al. (2023): we train a ML model to predict the ΔXCH_4 differences between observed values, for each pairwise combination of products, using retrieval parameters as input features. Ten features are included: surface altitude and roughness, SWIR surface albedo, fluorescence, the aerosol size parameter, the SWIR aerosol optical thickness (AOT), the a priori XCH_4 total column and the across-track pixel index, which are retrieved from the SRON product, and the observation errors of the two products for which ΔXCH_4 is predicted. The across-track pixel index provides information about the relative position of the pixel in the swath of the satellite, therefore it is related to the so-called "striping" effect — a systematic artifact observed in TROPOMI CO, $\text{H}_2\text{O}/\text{HDO}$, and XCH_4 products (Borsdorff et al., 2019). This effect consists in differences of XCH_4 measurements across consecutive parallel strips. The 3,432,335 observations shared across the three products (dataset described in Section 2.1.1) are randomly split into a training dataset (90% of the observations) and a validation dataset (10% of the observations). We search for the best-performing model between three ML algorithms that rely on decision trees: Random Forest, XGBoost and Light Gradient-Boosting Machine (LightGBM). Random Forest builds an ensemble of decision trees by combining bootstrap sampling and feature randomization, and produces predictions through averaging across the forest (Breiman, 2001). XGBoost and LightGBM apply gradient-boosted decision trees, a method where sequentially trained decision trees minimize a loss function by correcting the residual errors of the previous trees, thereby iteratively enhancing accuracy. XGBoost employs advanced regularization (L1 and L2 penalties) and parallelized tree construction (Chen and Guestrin, 2016). In contrast, LightGBM adopts a histogram-based learning approach and exclusive feature bundling (Ke et al., 2017).

In all cases, LightGBM outperforms the other models, while Random Forest systematically shows the lowest performances. Table 4 presents the validation results for the LightGBM model, which was ultimately chosen for further analyses. The predictive performances are consistent with the results of Balasus et al. (2023) for the prediction of the XCH_4 differences between TROPOMI and GOSAT, in terms of RMSE (Root Mean Square Error) and correlation (RMSE= 12.4 ppb, $R^2 = 0.53$).

Beyond its predictive performances, the trained model provides insights into the most important features — that is, the features that contribute most to the prediction of ΔXCH_4 . We conduct a study using SHapley Additive exPlanations (SHAP) on the predictions of the model, following Balasus et al. (2023). This approach partitions an individual prediction into contributions attributed to each feature, quantified as SHAP values (in ppb). The sum of all SHAP values for a given prediction corresponds to the difference between that prediction and the mean prediction across the entire dataset. The input features can then be ranked using the average absolute SHAP values, for each product pair comparison (Figure 7). Our findings align closely with those of Balasus et al. (2023): the most impacting features on the differences between satellite products are the across-track pixel index (thus striping patterns), aerosols and SWIR albedo. Calculating the mean ratio between the absolute SHAP value and the sum of the absolute SHAP values of all the features, the contributions are between 20 and 29% of the difference for aerosols, 13 to 19% for striping patterns, and 13 to 14% for albedo. The observation errors also seem to impact the prediction. Features related to aerosols are the SWIR aerosol optical thickness (AOT) and the aerosol size parameter, which are strongly anti-correlated

($R^2 = -0.82$). Since the SHAP analysis does not fully resolve correlations between variables, we aggregated the contributions of these two features into a single combined SHAP value, hereafter called "aerosols", to preserve interpretability.

Dataset 1	Dataset 2	RMSE (ppb)	Correlation (R^2)
SRON	WFMD	8.1	0.58
SRON	BLENDED	4.4	0.86
WFMD	BLENDED	8.5	0.44

Table 4. Validation results of predicted ΔXCH_4 for each combination of products, with the best-performing model (Light GBM).




Figure 7. Contributions of the individual features to the ΔXCH_4 predictions between pairs of TROPOMI products. The 9 most impacting predictors are ranked in order of importance. The contribution of a feature is defined as the average of absolute SHAP values, with the green bar showing the interquartile range.

We focus on the impact of albedo, aerosols and striping patterns on the XCH_4 differences between products: Figure 8 illustrates the variations in observed XCH_4 as functions of SWIR albedo, aerosol size parameter, and across-track pixel index.

380 The albedo and aerosol parameters are retrieved with the SRON algorithm: it is important to mention that the accuracy of these retrievals is not clearly known.

– **Albedo** - XCH_4 are influenced by surface reflectance, particularly in scenes with low SWIR albedo (e.g., snow, water bodies) or high SWIR albedo (sandy areas). In all the products, the albedo dependence is corrected based on a polynomial fit to the surface reflectance spectrum in the $2.3 \mu\text{m}$ spectral range. The polynomial degree was increased from 2 to 3 in recent updates of the products (Lorente et al., 2023; Schneising et al., 2023). Despite these adjustments, the products show different variations with SWIR albedo, as illustrated in Figure 8a. BLENDED and WFMD have a consistent low dependency on the albedo in the range of values between 0.05 and 0.5, where the density of observations is high. However, there is an offset of approximately 10 ppb, which decreases at high albedo values. SRON XCH_4 observations show more variations: they are close to those of WFMD at low albedo values, but lower than the other two products for high albedos.

– **Aerosols** - The effect of aerosols on methane retrievals is well-known: scattering by aerosol particles modifies the light path and induces errors that can compromise the accuracy of the retrieval (Butz et al., 2012). It depends on aerosol amount, type and size distribution. The latter is characterized by the aerosol size parameter α , which is the negative

exponent of the power law of the aerosol size distribution $n(r) \propto r^{-\alpha}$, with r the particle radius: the higher α is, the more $n(r)$ is shifted towards small particle sizes. It is simultaneously inferred and corrected during the XCH₄ retrieval process of SRON (Hasekamp et al., 2022), yet it still strongly impacts XCH₄ observations. TROPOMI SRON data is biased low relative to GOSAT for low values of α , indicating large particles (Balasus et al., 2023), as confirmed by Figure 8b. The correction applied in the BLENDED product makes its sensitivity to α closer to the one of WFMD, in comparison to SRON. Still, the three products show different behaviors and seem to be consistent only for $\alpha > 4.4$, which corresponds to a small part of the observations.

- **Striping patterns** - Without dedicated destriping, stripes in the flight direction are visible in the TROPOMI XCH₄ data, likely due to variations in the offsets and gains of detector pixels. While the across-track pixel index is included in the calibration process of the WFMD retrieval, residual vertical stripes persist in the data (Schneising et al., 2023). The latest WFMD product (v1.8) mitigates this effect using a wavelet–Fourier decomposition and filtering (Schneising et al., 2023). Indeed, Figure 8c) depicts the higher sensitivity of the SRON product to this effect. The correction with respect to GOSAT applied in the BLENDED product yields slightly lower sensitivity regarding the striping effect (Figure 8c). A recent method based on a double moving median smoothing along both the flight and cross-track directions resulted in promising enhancements in striping correction for the operational SRON product. It is implemented into newer versions of the product (Borsdorff et al., 2024), but older orbits have not been reprocessed. Future uses of the SRON destriped data should improve the consistency of XCH₄ observations with the other products.




Figure 8. Observed XCH₄ total column vs. SWIR albedo, aerosol size parameter and across-track pixel index. These three parameters are retrieved with SRON algorithm. The histogram depicts the density of common observations with respect to the corresponding variable.

3.2 Comparison between observed and simulated CH₄ total columns

The differences in the simulated XCH₄ between the three TROPOMI products are linked to the differences in observation coverage and in the vertical profiles that are necessary to compute their simulated equivalents, as described in Section 2.1.1.

The mean biases (MBs) of the difference $\Delta XCH_4^{os} = XCH_4^{obs} - XCH_4^{sim}$ with the prior inputs are approximately 6.7, -0.4 and 4.4 ppb and the RMSEs 18.2, 13.6 and 15.4 ppb respectively for SRON, BLENDED and WFMD (top row of Figure 9). BLENDED demonstrates both a smaller bias and a lower RMSE between observations and simulations, in comparison to SRON and WFMD.

When the analysis is restricted to the subset of matching observations across the three datasets, the RMSE decreases for all products, to respective values of 17.4, 13.1 and 13.0 ppb. This improvement indicates that the application of combined filters, which more rigorously select high-quality data, reduces the gap between observed and simulated concentrations.




Figure 9. Annual average difference between TROPOMI observations and their CHIMERE simulated equivalents, using prior emissions (first row) and posterior emissions (second row). MB is the mean bias and RMSE the root mean square error, in ppb. Only the cells with at least 50 observations in 2019 are shown.

The spatial distributions of the differences between observed and simulated concentrations ΔXCH_4^{os} (first row of Figure 9) reveal common patterns but also notable discrepancies across the three TROPOMI products. For all products, simulated XCH_4 are overall lower than observations over land. SRON has the highest difference between observed and simulated XCH_4 over land, especially in Western Europe. Over the sea, the simulations behave very differently across products: SRON shows a positive difference, while it is negative for BLENDED and close to 0 for WFMD. The difference between SRON and BLENDED is consistent with the systematic downward correction over the ocean depicted in Balasus et al. (2023).

Seasonal variations in the ΔXCH_4^{os} are generally consistent across the three products (Figure D3). ΔXCH_4^{os} are high during the first half of 2019 and decrease in the second half of the year; the RMSE decreases by more than 20% from winter (i.e. Jan-

uary to March) to summer (i.e July to September) for all products, though this overall reduction masks spatial heterogeneities. Scandinavia stands out as a region with marked discrepancies between observations and simulations: for SRON, $\Delta\text{XCH}_4^{\text{os}}$ reaches high positive values from April to June and is negative between October and December (Figure D3). The seasonal variations are consistent across the three products in this region, though with lower amplitude for BLENDED and WFMD. The differences in behaviour among products in regions like North Africa and Scandinavia highlight the influence of surface albedo and aerosols on observations. This effect predominantly affects observed concentrations, as simulated equivalents exhibit limited dependence on the parameters identified in Section 3.1 (not shown).

Overall, simulated concentrations are underestimated compared to observations over land across all TROPOMI products, with BLENDED providing the closest match. The residual (difference between the observed XCH_4 and the posterior simulations) is reduced after the inversion: the mean biases are decreased from the prior to the posterior simulation by approximately 93%, 95% and 99% for SRON, BLENDED and WFMD respectively (bottom row of Figure 9). Similarly, the RMSE decreases by 37%, 36%, and 28% for these products.

3.3 TROPOMI-derived CH_4 emissions : spatial and temporal distributions, annual budgets

Inversions are performed using the method detailed in Section 2.4.2. The three inversions assimilating each TROPOMI product, as well as the additional inversion using surface data for evaluation, are listed in Table 2. Figure 10 shows the corrections to the prior CH_4 emissions from these inversions, referred to as increments, both at the grid-cell and national resolutions.

At the grid-cell resolution, the increments differ among the inversions (Figure 10a). These differences result in large differences in the total CH_4 emission budgets for 2019, for the regions and the corresponding countries listed in Table B1. While the prior total emissions of 25.2 TgCH_4/yr are increased by 2% for the SRON (25.7 Tg/yr) inversion, they are reduced by respectively 1%, 9% and 33% for the BLENDED (25.0 Tg/yr), surface-based (23.0 Tg/yr) and WFMD (16.9 Tg/yr) inversions. The detailed comparison of prior and posterior emissions per country is shown in Table S1. While some regions exhibit consistent corrections across all three TROPOMI inversions (e.g. in Northern Africa, Italy or Romania), the magnitude of increments is generally larger in the WFMD-based inversion in comparison to SRON and BLENDED, consistently with the OSSEs of Section 3.4. Aggregation at the national scale mitigates some of these inconsistencies but reveals high differences in countries such as the Netherlands, Germany, France, and Poland (Figure 10a). SRON and BLENDED provide similar corrections to the prior CH_4 emissions in most countries, except Poland and Turkey, but they differ from WFMD particularly in Western and Central Europe (Figure S4, in the Supplement).

It is important to note that the surface-based inversion is influenced by the spatial distribution of the stations: regions with dense station coverage, such as Western Europe (8 stations), are well constrained, whereas regions with sparse coverage, like the Baltic states, Spain and Portugal show little correction of the prior fluxes. Scandinavia is poorly constrained in all satellite inversions, whereas small increments appear in Northern Finland in the surface-based inversion due to the presence of the Pallas station. Despite these limitations, the comparison of satellite-based inversions with the surface-based inversion shows some consistency: surface-based increments agree well with WFMD corrections over the Paris area and Italy, and with SRON and BLENDED corrections over the Netherlands and Switzerland. However, the surface-based inversion is not consistently

closer to one satellite-based inversion than to another, as indicated by the R^2 correlation coefficients between surface-based increments and satellite-based increments (0.24 for SRON, 0.34 for BLENDED, and 0.37 for WFMD).




Figure 10. (a) Spatial distributions of the increments to the prior CH₄ emissions from the inversions, in Gg/yr at the grid-cell resolution (first row) and in Tg/yr at the national resolution (second row). (b) Total emissions for the regions described in Table B1, in Tg/yr. (c) Monthly variations of the CH₄ emissions in the domain in 2019, in Tg/month. The shaded areas show the standard deviation within each month.

465 The temporal variations of CH₄ emissions is shown in Figure 10c. Satellite-based inversions exhibit similar temporal patterns, including high emissions during December and January, a peak in April or May, and lower emissions over the summer months. Despite these similarities, WFMD-based emissions are systematically lower than those from SRON and BLENDED, contributing to the lower annual total mentioned earlier. The monthly corrections to prior CH₄ emissions differ between satellite-based and surface-based inversions. For example, the peak of emissions during spring detected in the satellite in-

versions is not found in the surface-based inversion (Figure 10c). This peak is due to positive corrections in Western and Central Europe across all satellite-based inversions. This period coincides with a slight decline in observed XCH_4 (Figure 2a), and an increase (for the month of April) in the lateral boundary conditions (Figure D4). Yet, the origin of this emission peak has not been clarified.

Both TROPOMI-based and surface-based inversions capture the summer emission minimum, which contrasts with the prior emissions, where the minimum occurs in autumn and winter. This unexpected seasonal cycle, already identified in Pison et al. (2021), seems to be related to CHIMERE transport modeling since it did not occur when using other models like FLEXPART. It is likely due to a misrepresentation of the atmospheric transport or poor representation of the seasonality of boundary conditions (Pison et al., 2021). Other transport model errors (miscalculation of the tropopause height, mesoscale transport errors) can impact the calculation of simulated XCH_4 (Saad et al., 2014), thus biasing the results of the inversions. However, a dedicated study would go beyond the scope of this article, and we do not address the impact of model errors and how they interact with the retrieval biases.

The nearly identical emission increments for SRON and BLENDED (Figure 10) show that corrections for albedo and aerosol related biases in BLENDED do not lead to any significant differences in the derived emissions. Indeed, the logic for BLENDED is to solve these biases of the SRON XCH_4 retrievals using GOSAT retrievals. Therefore, this suggests that potential albedo and aerosol related biases (depicted in Section 3.1) are not significantly impacting the derived CH_4 emissions in Europe, within the scope of our study.

Figures S5 and S6 (in the Supplement) show scatter plots of posterior emissions against prior emissions, at the pixel and weekly scale (Figure S5) and aggregated at the monthly and national scale (Figure S6). The previously mentioned differences between the distributions of posterior emissions are clearly observed at the pixel scale. The variability decreases when aggregating the fluxes, yet remains substantial across the posteriors of the four inversions. It highlights the limits of the current use of the TROPOMI products to build national CH_4 budgets and attribute emissions.

To complete the analysis of posterior emission differences, a further comparison of the increments in the background components of the control vector (as defined in Section 2.4.2) reveals that, while the increments exhibit some similar patterns, they overall differ for the three TROPOMI products. The average background total columns are increased in average by 3.5, 1.4 and 2.1 ppb for respectively SRON, BLENDED and WFMD (Figure D5). The increments in the stratosphere share common patterns of positive increments over the Mediterranean basin, but with differing magnitudes: they are stronger for SRON and WFMD than for BLENDED. The lateral boundary increments are mainly positive over the western limit for SRON, BLENDED and the surface-based inversion, with the higher values for SRON. WFMD shows a different pattern for the increments of the lateral boundary conditions, with an increase of the background at the southern border of the domain. The time variations of the increments are overall consistent between products, only the magnitude changes across the inversions (Figure D4).

For BLENDED, the close agreement between observed and simulated XCH_4 leads to a lower magnitude of both flux and background increments in comparison to other inversions. SRON has higher background increments in average than WFMD, consistently with the higher difference of observed and simulated XCH_4 showed in Figure 9. However, for SRON the emissions are not pulled down as strongly as for WFMD. For this latter product, the background and flux increments do not seem to be

505 anti-correlated, which could have been expected as the differences between the observations and the prior simulations were not that high and the fluxes strongly decreased through the inversion. Therefore, the strong negative increments on the *Inv-WFMD* fluxes result from a complex balance between the local gradients of the increments on the background and on the fluxes: the system could have difficulty separating both when using the WFMD observations. The differences between products detailed in Sections 2.1.3, 3.1 and 3.2 could push the system towards different splits between background and emission optimization.

510 3.4 Separation of the causes of differences through OSSEs

Previous sections highlighted the differences in the products, in the observed and simulated XCH₄ distributions and in the results of inversions. To deepen the understanding of the differences of inversion outputs, we compare here the capacity of the system to improve emission estimates through the inversion, for the three products. As described in Section 2.4.3, OSSEs are a relevant tool to evaluate this capability, to perform sensitivity tests on the inversion procedure and to discriminate between the causes of the differences in the fluxes derived from various inversions. Following the method and notations of Section 2.4.3, the "prior" refers to the perturbed prior x^b , while the "truth" corresponds to the unperturbed prior x^t . The performances of an OSSE are assessed using the relative increment, as defined in Equation 4: the closer to -100% r is, the closer the posterior fluxes are to the truth (in comparison to the prior). This metric is calculated only over land to avoid averaging effects caused by weak fluxes over the sea. Table 5 shows the relative increment for the first six OSSE configurations (described in Table 3), as well as the RMSE and R² coefficient between the perturbed prior fluxes (averaged over all the members) and the true fluxes, and between the posterior and the true fluxes.

3.4.1 Reference scenarios

The *Ref-SB* and *Ref-WFMD* OSSEs demonstrate a consistent capacity to constrain emission estimates. Despite the relatively low values of r (in comparison to the aim of -100%), we focus on the relative differences of r between scenarios. The temporal (Figure 11a) and spatial (Figure 11b) variations of the relative increments are consistent across the three products. Monthly variations smooth the noisier weekly variations in Figure 11a, but relative differences between scenarios are conserved. WFMD presents the highest performance, with a mean relative increment of -5.7%, in comparison to -3.5% for SRON/BLENDED, and a larger reduction in RMSE (Table 5). This product thus presents a higher constraint capacity in comparison to SRON and BLENDED.

530 The prior temporal variations of emissions remain largely preserved through the assimilation of pseudo-observations. Consequently, the posterior time variations align closely with those of the unperturbed prior. However, the relative increment varies over the year (Figure 11a). These variations are partially influenced by the observational density, with lower r (i.e. closer to -100%) in summer, coinciding with an abundance of high-quality measurements, and higher (i.e. further from -100%) values in January and December, when coverage decreases due to cloud filtering. A small $|r|$ is also seen in May 2019, likely caused by the low number of observations (Figure 1b).

Regions with sparse observational coverage generally exhibit worse (i.e. closer to 0) relative increments. In particular, the system fails to improve emission estimates in Scandinavia across all three configurations (Figure 11b). This limitation is

ID	Prior-truth		Posterior-truth		Relative increment
	RMSE ($\text{g.m}^{-2}.\text{yr}^{-1}$)	R^2	RMSE ($\text{g.m}^{-2}.\text{yr}^{-1}$)	R^2	
Ref-SB	2.94	0.89	2.82	0.89	-3.5%
Ref-WFMD	2.94	0.89	2.74	0.90	-5.7%
Ref-Surf	2.94	0.89	2.71	0.90	-7.3%
Common-SB	2.94	0.89	2.82	0.89	-3.5%
Common-WFMD	2.94	0.89	2.77	0.90	-5.1%
Err-SB	2.94	0.89	2.81	0.89	-3.8%

Table 5. Evaluation results of 6 OSSE configurations. "Prior" refers to the perturbed prior fluxes, "Truth" to the unperturbed prior fluxes and "Posterior" to the optimized fluxes after the inversion. All the fluxes (thus the RMSE) are in units of $\text{gCH}_4.\text{m}^{-2}.\text{yr}^{-1}$. The prior and posterior have been averaged over all samples of the OSSE. The relative increment is calculated only over land. *Diff-SRON* and *Diff-WFMD* are not included since prior emissions were not perturbed. Scenario IDs refer to Table 3.

directly related to the sparse observation density in this region (Figure 1b), with nearly no observations available during winter. Due to a sparse observation density, the system also performs poorly in the United Kingdom (Figure 11b), contributing to the poor mean relative increment over Western Europe.

The system manages to make the posterior emissions closer to the truth in areas with high fluxes (northern Italy, Benelux, Romania) while struggling in areas with weaker signals, like Latvia and southern France (Figure 11b). This is linked to the prior error covariance **B** being proportional to the emissions (see Section 2.4.1), which limits the ability of the inversion to recover diffuse sources with low signal-to-noise ratios, as previously noted by Yu et al. (2021).

These reference OSSE scenarios demonstrate the capability of all TROPOMI products to bring emission posteriors closer to the truth compared to the perturbed priors. The *Ref-Surf* scenario provides a better overall enhancement, with an average relative increment of -7.31%. However, this better result on average masks a high spatial heterogeneity: the relative increments are indeed very good in Western Europe and Central Europe where a number of stations are located, but fail to provide enhancements in regions with no stations, such as Spain or Romania (Figure 11b). As expected, the wider coverage of satellite observations provides the advantage of constraining the emissions on a wider area, even if there are no surface stations.

The OSSEs can also be used to estimate the uncertainty reduction (independent of the control vector) for the inversions, computing the ratio between the standard deviation of priors and the one of posterior emissions across the ensemble samples. For the total budgets, it is estimated to 78% reduction for SRON and BLENDED, 74% for WFMD and 51% for surface stations. However, we do not use them for evaluating the uncertainties of our budget estimates in Section 3.3, because of 1) the small size of the ensemble (4 samples) and 2) the lack of proper uncertainty estimation for the prior (not provided with the emission inventories). The estimation of the uncertainty requires a deeper analysis of the propagation of errors through the system: observation errors (Zheng et al., 2026) as well as model errors due to the prior emissions and the transport (Lu et al., 2025a).

3.4.2 Drivers of the differences of increments

Observation density




Figure 11. (a) Average monthly relative increment (%) over 2019 and (b) maps of relative increment (%), for the different OSSE scenarios. These scenarios are described in Table 3. The lower the relative increment, the more posterior fluxes are closer to the truth (Equation 4). The regions are described in Table B1.

560 The relationship between observation density and constraint potential is explored through the *Common-SB* and *Common-WFMD* OSSEs. As expected, the relative increments are slightly closer to zero as observation density decreases (Table 5). The average difference in comparison to the reference scenario is very low (less than $0.01 \text{ gCH}_4.\text{m}^{-2}.\text{yr}^{-1}$ in RMSE) for SRON/BLENDED, with 73% of the observations kept in *Common-SB*, but slightly higher ($0.03 \text{ gCH}_4.\text{m}^{-2}.\text{yr}^{-1}$ in RMSE) for WFMD because the difference in observation density is larger (45% of observations kept). The time series of *Ref* and *Common* scenarios in Figure 11a illustrate that the relative increment is overall slightly deteriorated, yet not systematically, as the number of observations decreases. The observation density is thus a driver of the potential for constraining emissions through the inversion. However it only partially explains the differences of performance between the products, since *Common-WFMD* results in lower relative increments than *Common-SB*: even with the same number of observations, WFMD seems to show a better capability to bring emission closer to the truth. The key differences between these scenarios lie in the uncertainties associated with the measurements and the vertical parameters (such as pressure levels, AKs, and prior profiles) used to compute the satellite equivalents.

Error rescaling

575 The linear rescaling of the error associated with individual observations, described in Equation 1, increases these individual errors in the dataset of pseudo-observations. The comparison of *Err-SB* scenario to *Common-SB* shows a slight improvement of the emissions for all months (Figure 11a) and all regions (Figure 11b), when rescaling these errors. The spatial distributions are similar, as shown in Figure 11b, but with a slightly larger amplitude for *Err-SB*.

The *Err-SB* scenario is the closest SRON/BLENDED scenario to WFMD ones in terms of relative increment (Figure 11). The *Err-SB* and *Common-WFMD*, which share similar features in terms of coverage and errors, have the highest consistency 580 between SRON/BLENDED and WFMD OSSEs, highlighting these features as drivers of the differences in inversion results. The remaining differences are due to the vertical parameters (pressure levels, averaging kernels, and prior profiles).

Rescaling the observation error results in an enhancement of the emissions. This counterintuitive effect is likely due to overfitting and complex indirect effects within the inversion: because of the large number of observations and the low individual observation errors, the system could struggle to overfit the observations, thus degrading the performances. At the 0.5° resolution, several observations constrain each component of the control vector, leading to overfitting of some observations if the errors are small. This effect would be mitigated in the *Err-SB* scenario, leading to better (i.e. more negative) r . Still, this effect is limited (r varies by a few 0.1%) in comparison to the differences of r between products.

SRON-WFMD XCH_4 difference

590 The scenarios *Diff-SRON* and *Diff-WFMD* provide insights into how the bias between products impacts the retrieved flux distribution. In this section, we do not focus anymore on the relative increment (fluxes are not perturbed in the *Diff* scenarios), but on the increments derived from the assimilation of the biased observations.

As expected, the opposite biases result in overall opposite increments. The assimilation of biased observations leads to an increase in total emissions for SRON (from 25.2 to 25.3 Tg/yr) and a decrease for WFMD (from 25.2 to 24.3 Tg/yr), consis-

595 tently with the average positive difference between WFMD and SRON. However, these small differences in total emissions are the sum of large spatial variations, as shown in Figure 12b. The maps highlight the contribution of the WFMD-SRON difference to the SRON increments, with negative increments in Western Europe (UK, Ireland, eastern Spain), southern Italy and Austria/Czech Republic, particularly over mountainous regions like the Pyrenees and the High Tatras in Central Europe. Positive contributions are observed across most of Eastern Europe, Benelux/Germany, central Italy and the Alps. While these
600 contributions should be closely related to the difference of XCH_4 columns, shown in Figure 12a, no clear correlation emerges between the bias in the concentrations and the corrections in the emissions: the average SRON-WFMD difference is generally negative and smooth, in contrast to the localized, strong increments in the emissions.




Figure 12. Spatial averages of (a) the XCH_4 difference SRON-WFMD (in ppb), (b) the increments of the *Diff-SRON* and *Diff-WFMD* OSSEs (in Gg/yr) and (c) of the ratio between the increments of *Diff-SRON* by the difference of increments between inversions *Inv-WFMD* and *Inv-SRON* (in %). For the last map (c), pixels with ratios above 150% (corresponding to very close increments in *Inv-WFMD* and *Inv-SRON*) were filtered out.

Yet, these two OSSE scenarios help clarify some of the patterns observed in inversions with real data. The increments of *Diff-SRON* (biased by the WFMD-SRON difference) and the difference of increments between inversions *Inv-WFMD* and *Inv-SRON*
605 described in Section 3.3 are compared in Figure 12c. The ratio between these two quantities show that the differences in the inversion increments can be partially explained by the difference SRON-WFMD over the sea, in Western Europe or in Romania. However, blue areas in Central Europe and Eastern Europe show opposite variations, meaning that the OSSE increments are not sufficient to explain the increments in the inversions with real data. This suggests that while XCH_4 differences provide some explanation, they do not fully account for the inversion outputs. In inversions, the CAMS background is also optimized
610 and transport is not assumed perfect, contrary to OSSEs: it makes the tracing of the main factors impacting the increments on fluxes more complex.

3.5 Evaluation against independent surface measurements

Finally, we evaluate the TROPOMI-based inversions described in Section 3.3 with independent surface data. To evaluate the consistency of satellite-based and surface-based perspectives, we compare the posterior simulated concentrations to independent methane observations from the surface stations listed in Table A1. The mean biases, RMSEs and correlation coefficients
615 are summarized in Table 6.

Considering all surface measurements, the average difference between XCH₄ observations and the prior simulations is 2.0 ppb. It is closer to 0 than the average difference between XCH₄ observations and the posterior simulations for SRON and WFMD (respectively -9.9 and 9.1 ppb). For these two simulations, the RMSE is also higher than for the prior simulation.
 620 However, the absolute mean bias and RMSE are closer to 0 for BLENDED posterior simulations, in comparison to the prior: this product is more consistent with surface station measurements than the other two. In *Inv-Surface*, the differences are decreased for almost all stations, as expected.

The overall statistics mask a heterogeneous distribution of differences across individual stations. The corrections to the prior CH₄ emissions derived from satellite-based inversions improve the fit with surface measurements for about half of the stations:
 625 37% for SRON, 53% for BLENDED, 47% for WFMD (Figure 13). For SRON and BLENDED posterior simulations, the difference is deteriorated at most stations with simulated equivalents generally higher than the observations, especially over the UK, Ireland and France (MHD, RGL, TAC, SAC, OPE, Figure D2). It is consistent with the positive increments in these regions (Figure 10a). The bias approaches zero for only 3 stations: LUT, IPR, HPB (Figure 13). For the WFMD posterior simulation, the simulated equivalents are lower than the prior for almost all stations, due to negative flux increments in the
 630 inversions (Figure 10a). This adjustment improves the fit to surface measurements at stations mostly in Western Europe or Italy (e.g., MHD, LMP, CMN, TRN and TAC), but deteriorates the fit at most other stations (Figure 13).

These results highlight the gap between satellite-based and surface-based inversions: fitting satellite methane observations does not systematically improve the fit of the simulated CH₄ mixing ratios to in-situ measurements. Aligning estimates from satellite-based and surface-based inversions is crucial for the accurate evaluation of inferred emissions, as well as for ensuring consistency and reliability in methane flux estimates derived from different observational frameworks. This future work
 635 could take advantage of both in-situ CH₄ measurements and ground-based remote sensing observations from TCCON (see Appendix C) or COCCON.

Flux inputs	MB (ppb)	RMSE (ppb)	R ²
Prior	2.0	34.7	0.74
Posterior SRON	-9.9	36.5	0.74
Posterior BLENDED	-1.1	34.2	0.76
Posterior WFMD	9.1	36.7	0.71
Posterior Surface	1.3	22.3	0.91

Table 6. Comparison of the mean bias (MB), RMSE and R² between independent surface measurements and the simulated concentrations using respectively the prior emissions and the posterior emissions from the four inversions. The high RMSE (in comparison to the MB) highlights the variability of observation/simulation comparison for surface measurements.

4 Conclusions

The assimilation of TROPOMI CH₄ total columns into an inverse modeling system is a powerful tool for quantifying methane
 640 emissions (Jacob et al., 2022), as they provide complementary information to bottom-up inventories and surface measurements.




Figure 13. Differences between independent surface measurements and the simulations (in ppb) using the prior emissions and the posterior emissions from the SRON, BLENDED and WFMD and Surface-based inversions, for the stations described in Table A1.

In this study, we compare the emissions estimated from the inversions of three TROPOMI products. Their consistency is essential for the comparability of the subsequent analyses based on these products.

The retrievals are sensitive to a range of instrumental and atmospheric variables. A machine learning model is employed to assess the importance of features in predicting differences between the satellite products, showing that the main drivers of these differences are aerosols (20-29% of the predicted difference), striping patterns (13-19%), and extreme albedo values (13-14%). The effects of these variables on the fluxes derived from inversions should be the object of further investigation. Recent and ongoing developments (e.g., reprocessing of destriped orbits for SRON, improved aerosol event filtering for WFMD, enhanced cloud filtering for both products) are expected in new product updates and should improve the quality of the products.

Our findings demonstrate that assimilating the three TROPOMI products into regional inversions results in distinct posterior CH_4 emission estimates. Our top-down European emissions (countries listed in Table B1) are evaluated to 25.7 Tg/yr for SRON, 25.0 Tg/yr for BLENDED and 16.9 Tg/yr for WFMD in 2019. The range of estimated total and country-scale budgets has to be put into perspective within the framework of emission reporting to the United Nations Framework Convention on Climate Change (UNFCCC). At the monthly and national scales, the consistency between products remains insufficient for reliable budget estimates. Our study shows a good agreement between the non-independent SRON and BLENDED. Since BLENDED is a post-processed version of SRON that corrects the albedo and aerosol related XCH_4 biases with GOSAT observations, it suggests that these biases have relatively low impact on the differences of posterior emissions between TROPOMI products. The comparison with an inversion assimilating surface station data (23.0 Tg/yr) does not conclusively indicate which TROPOMI product yields posterior emissions most consistent with surface-based estimates: the choice of product depends on the specific goals of the study, each having its own strengths, weaknesses, and sensitivities. Better characterization of the uncertainties is required to statistically test the consistency of posterior emissions and to complete the comparison. If OSSEs

provide an estimation of the uncertainty reduction, 4D-Var inversions do not give direct access to the posterior uncertainty on the CH₄ emissions.

OSSEs further highlight the role of both observation density and errors on the capability of the inversion system to enhance the emissions estimates. The *Diff* scenarios pinpoint how the SRON-WFMD XCH₄ differences explain specific spatial patterns of the increments. The OSSEs further underscore the limitations of the inversion system and its complex dynamics. The optimization process involves a delicate balance between increments on emissions and on the background. The relative corrections on emissions and background differ across TROPOMI-based inversions, thus influencing the derived emission budgets. Future work is necessary to include corrections of the biases related to the boundary conditions (Nesser et al., 2025), and to account for other model errors that have not been investigated in this study. Furthermore, standardized observation error definitions are required. Specifically, we recommend to rescale the observation errors for SRON and BLENDED: instead of the multiplication of errors by a factor 2, a linear regression similar to Equation 1 should be derived for each product, based on a regression of the scatter relative to TCCON observations, as described for the WFMD product (Schneising, 2023).

In this study, we chose not to correct the XCH₄ products. Global inversions with albedo- and aerosol-corrected products, as well as the new destriping procedure for SRON, would provide a quantification of the impact of these parameters on the posterior emissions. To extend the evaluation of TROPOMI-based inversions against surface data, we also recommend deeper comparisons with local studies, such as coal mining emissions in Poland (Tu et al., 2022), oil and gas production emissions in Romania (Maazallahi et al., 2025; Kuhlmann et al., 2025) and in Algeria (Naus et al., 2023). The comparisons were not included in this study to focus on the relative comparison of satellite products at regional scale. Bridging the gap between inversions using satellite and surface observations, as well as between local and regional/global studies is challenging but essential for the validation of emission estimates (Santaren et al., 2021; Naus et al., 2023). The progress towards consistent CH₄ emission budgets at national and sub-national scales is crucial for validating the effectiveness of European mitigation strategies and for monitoring the reductions of the CH₄ emissions in line with the Global Methane Pledge's 2030 reduction targets.

Data availability. TROPOMI CH₄ product (v2.4) can be found here: <https://s5phub.copernicus.eu/dhus/#/home>, last access May 2021 (Landgraf et al., 2024). The WFMD methane data can be accessed via http://www.iup.uni-bremen.de/carbon_ghg/products/tropomi_wfmd/, last access November 2022 (Schneising et al., 2023). Blended TROPOMI+GOSAT Satellite Data Product for Atmospheric Methane was accessed from <https://registry.opendata.aws/blended-tropomi-gosat-methane>, last access May 2024 (Balasus et al., 2023).

Code and data availability. The CHIMERE code is available here: www.lmd.polytechnique.fr/chimere/ (Menut et al., 2013; Mailler et al., 2017). The CIF inversion system is available at: <http://community-inversion.eu/> (Berchet et al., 2021).

690 *Author contributions.* AB, AFC, IP and ASP contributed to the study conceptualization. AFC conducted the data collection with contribution of ASP, AM and AO; IP and ASP run the simulations. ASP conducted the analyses with contributions of AB, AFC, IP, EP, AO and GB. OS, MB, JDM and TB provided guidance on the TROPOMI data and discussed results. ASP wrote the article with input from all authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We thank the reviewers for taking the time and effort necessary to review the manuscript. Their thoughtful comments helped us improve the quality of the manuscript. A large part of the development and analysis were conducted in the frame of the H2020 VERIFY project, funded by the European Commission Horizon 2020 research and innovation programme (agreement no. 776810). This project has received funding from the European Union's Horizon Europe research and innovation programme (agreement no. 101081395). This work was also supported by the CNES (Centre National d'Etudes Spatiales), in the frame of the TOSCA ARGOS project. The last part of the analysis was conducted in the frame of the ESA initiative SMART-CH4 (Satellite Monitoring of Atmospheric Methane, contract no. 4000142730/23/I-NS), which is part of the EC-ESA Joint Earth System Science Initiative. This work was granted access to the HPC resources of TGCC under the allocations A0180102201 made by GENCI. We thank the data providers of TROPOMI products: SRON, the Atmospheric Chemistry Modeling Group at Harvard University, and University of Bremen. We also acknowledge the principal investigators of surface stations and TCCON sites for the data used for evaluation in this work, as well as the Japanese Aerospace Exploration Agency, the National Institute for Environmental Studies, and the Ministry of Environment for the GOSAT data.

705 University of Bremen acknowledges funding from the European Space Agency via project GHG-CCI+ (contract no. 4000126450/19/I-NB) and from the Bundesministerium für Bildung und Forschung within its project ITMS (grant no. 01 LK2103A). The TROPOMI/WFMD retrievals were performed on HPC facilities funded by the Deutsche Forschungsgemeinschaft (grant nos. INST 144/379-1 FUGG and INST 144/493-1 FUGG).

Finally, we wish to thank J. Bruna (LSCE) and his team for computer support and the use of the OBELIX computing facility at LSCE.

Table A1. Surface stations used for the evaluation, with their coordinates.

ID	Station	Country	Lat (°)	Lon (°)	Altitude [m a.s.l.]
Mountain					
CMN	Monte Cimone	Italy	44.17	10.68	2165
HPB	Hohenpeissenberg	Germany	47.80	11.02	934
JFJ	Jungfrauoch	Switzerland	46.55	7.99	3570
KAS	Kasprowy Wierch	Poland	49.23	19.98	1989
OXX	Ochsenkopf	Germany	50.03	11.81	1112
PUY	Puy de Dôme	France	45.77	2.97	1465
ZSF	Zugspitze	Germany	47.42	10.98	2666
Coastal					
IPR	Ispra	Italy	45.81	8.64	210
LMP	Lampedusa	Italy	35.52	12.63	45
LUT	Lutjewad	Netherlands	53.40	6.35	1
MHD	Mace Head	Ireland	53.33	-9.90	8
RGL	Ridge Hill	UK	52.00	-2.50	204
TAC	Tacolneston	UK	52.52	1.14	56
Other					
HEI	Heidelberg	Germany	49.42	8.67	116
HUN	Hegyhátsál	Hungary	46.96	16.65	248
OPE	Obs. pérenne de l’environnement	France	48.56	5.50	390
PAL	Pallas	Finland	67.97	24.12	565
SAC	Saclay	France	48.72	2.14	160
TRN	Trainou	France	47.96	2.11	131

B List of sub-continental regions

Table B1. European regions used in this study.

Region	Countries
Western Europe	Belgium, France, Ireland, Luxembourg, Netherlands, United Kingdom
Central Europe	Austria, Croatia, Czech Republic, Estonia, Germany, Hungary, Latvia, Lithuania, Poland, Slovakia, Slovenia, Switzerland
Southern Europe	Italy, Portugal, Spain
Northern Europe	Denmark, Finland, Norway, Sweden
South-Eastern Europe	Albania, Bosnia-Herzegovina, Bulgaria, Cyprus, Greece, Macedonia, Moldova, Montenegro, Romania, Serbia

C Comparison to TCCON observations

In addition to the comparison of the XCH_4 distributions of the TROPOMI products, we compare the TROPOMI-TCCON co-located observations for the seven TCCON stations listed in Table C1. The Total Carbon Column Observing Network (TCCON) is a network of ground-based stations equipped with similar high-resolution spectrometers (Bruker IFS) and using a common retrieval algorithm to ensure comparability of the measurements. The network consists of 28 operational sites, of which 7 are in the domain of this study. It is available at <https://tccodata.org/>. We use the last update of GGG2020 (Laughner et al., 2024). Previous studies have compared one or two TROPOMI products to TCCON observations (T. Hilbig et al., 2023; Balasus et al., 2023; Borsdorff et al., 2024; Lindqvist et al., 2024), but none of them have directly compared the 3 products all together.

ID	Station	Country	Lat (°)	Lon (°)	Altitude [m a.s.l.]
BRE	Bremen	Germany	53.1	8.85	30
GAR	Garmisch	Germany	47.48	11.06	745
KRL	Karlsruhe	Germany	49.1	8.44	110
NIC	Nicosia	Cyprus	35.14	33.38	185
ORL	Orléans	France	47.96	2.11	130
PAR	Paris	France	48.85	2.36	60
SOD	Sodankylä	Finland	67.37	26.63	188

Table C1. TCCON stations used for the evaluation.

To compare observed CH_4 total columns from TROPOMI and TCCON datasets, we consider the co-located observations that are within 1 h and 100 km of each other, with a maximum surface elevation difference of 250 m. For co-located observations, it is required to adjust the columns for the differences of vertical sensitivities and prior XCH_4 profiles used in the retrievals, using the TCCON profile as the common prior profile. Following Apituley et al. (2024), Schneising (2022) and Balasus et al. (2023), the vertical profiles of TCCON (51 levels) are interpolated on the TROPOMI layers l (20 layers for WFMD, 12 for SRON and BLENDED). The adjusted TROPOMI XCH_4 total column $\hat{y}_{adj}$ is thus, with $\mathbf{y}_{a,TC}$ and $\mathbf{y}_{a,TR}$ the TCCON and TROPOMI prior profiles, $\hat{y}$ the TROPOMI XCH_4 total column and $\mathbf{a}$ the column averaging kernel:

$$\hat{y}_{adj} = \hat{y} + \sum_l h^l (1 - a^l) (y_{a,TC}^l - y_{a,TR}^l) \quad (\text{C1})$$

The results of the comparison for 2019 are presented in Table C2 and Figure C1. WFMD tends to overestimate methane concentrations, SRON has the lowest mean of the daily averaged difference but higher deviations and lower correlation in comparison to the other products. BLENDED observations align more closely with TCCON in terms of R^2 and RMSE, with a negative offset that is rather uniform across the stations. The average for individual stations are consistent with Balasus et al. (2023) and T. Hilbig et al. (2023) for SRON and BLENDED. However, they differ from the results of Borsdorff et al. (2024) and Lindqvist et al. (2024) for WFMD and from the results of T. Hilbig et al. (2023) for SRON. Overall, the values of the differences between TROPOMI and TCCON XCH_4 fall in the range [-25,+25] ppb. Moreover, the relative accuracy (standard

735 deviation of the mean local offsets relative to TCCON at the individual sites) of TROPOMI products shown in Table C2 are below the 10 ppb threshold deemed suitable for regional inversions by Buchwitz (2015). BLENDED and WFMD have lower relative accuracies (3.1 and 3.3 ppb) than SRON (4.8 ppb).

Product	Mean (ppb)	R ²	RMSE (ppb)	Relative accuracy (ppb)
SRON	0.9	0.48	16.4	4.8
BLENDED	-2.6	0.62	11.1	3.1
WFMD	7.5	0.54	13.7	3.3

Table C2. Mean and RMSE of the daily averaged difference TROPOMI-TCCON, as well as the correlation (R^2) and the relative accuracy of the TROPOMI products relative to TCCON, over the co-located observations at the TCCON stations in 2019. The relative accuracy is the standard deviation of the mean local offsets relative to TCCON at the individual sites.

Analysis of individual stations reveals similar patterns for those located in Western Europe (Bremen, Karlsruhe, Orléans and Paris). For these stations, SRON and WFMD show comparable distributions (with WFMD values slightly higher), while
740 BLENDED systematically produces lower median values, consistently with the comparison of XCH₄ distributions detailed in Section 3.1. A similar pattern can be seen in Nicosia, except for SRON higher values. All the products have a similar positive offset in Garmisch in comparison to other Western Europe stations. For this station, located in the Northern Alps, the offset is likely due to a bias associated with albedo or difference in the altitude of the ground pixel of the satellite and the station.




Figure C1. Median and quartiles of the daily averaged differences between TROPOMI and TCCON XCH₄ (ppb) for each station selected for the evaluation, in 2019.

Figure S7 in the Supplement shows a seasonally-resolved version of Figure C1, and Figure S8 shows the time series of
745 the TROPOMI-TCCON difference for key sites. Seasonal distributions and time series indicate a low seasonal dependency in the differences, apart from Sodankylä. In this high-latitude station in Finland, the bias in 2019 is positive during spring, and negative in autumn, consistent with the findings of Lindqvist et al. (2024). This seasonal variation is only present in 2019. Due to this temporal variations and to the limited number of co-located observations at this latitude, the deviations of the

TROPOMI-TCCON differences are amplified, especially for SRON which has the largest seasonal variations. These results
750 highlight the challenges of using TROPOMI at high latitudes, where coverage is sparse, and uncertainties are large.

D Additional figures




Figure D1. Count of observations that have been filtered in the post-processing of the TROPOMI products, as described in Section 2.4.2




Figure D2. Evaluation map of the satellite-based and surface-based inversions: green (resp. red) circles are the surface stations for which the posterior simulated concentrations are in average closer (resp. further away) to the observations than the prior ones.




Figure D3. Seasonal average difference between TROPOMI observed concentration and CHIMERE simulated equivalent. MB is the mean bias and RMSE the root mean square error. Units are ppb. JFM, AMJ, JAS and OND are acronyms referring to the months of each season.




Figure D4. Time series of monthly averaged increments of the components of the background, for the 4 inversions presented in Section 2.4.2. The first panel shows the time series of averaged total columns, the second panel those with only the pixels used as lateral boundary conditions, and the third panel shows the time series of average stratosphere column (for pressures > 200 hPa).




Figure D5. Increments of the components of the background, for the 4 inversions presented in Section 2.4.2. The components of the background are the same as in Figure D4: averaged total columns (first row), pixels used as lateral boundary conditions (second row), and average stratosphere column for pressures > 200 hPa (third row).

References

- Agustí-Panareda, A., Barré, J., Massart, S., Inness, A., Aben, I., Ades, M., Baier, B. C., Balsamo, G., Borsdorff, T., Bousserez, N., Boussetta, S., Buchwitz, M., Cantarello, L., Crevoisier, C., Engelen, R., Eskes, H., Flemming, J., Garrigues, S., Hasekamp, O., Huijnen, V., Jones, L., Kipling, Z., Langerock, B., McNorton, J., Meilhac, N., Noël, S., Parrington, M., Peuch, V.-H., Ramonet, M., Razinger, M., Reuter, M., Ribas, R., Suttie, M., Sweeney, C., Tarniewicz, J., and Wu, L.: Technical note: The CAMS greenhouse gas reanalysis from 2003 to 2020, *Atmospheric Chemistry and Physics*, 23, 3829–3859, <https://doi.org/10.5194/acp-23-3829-2023>, 2023.
- Alexe, M., Bergamaschi, P., Segers, A., Detmers, R., Butz, A., Hasekamp, O., Guerlet, S., Parker, R., Boesch, H., Frankenberg, C., Scheepmaker, R. A., Dlugokencky, E., Sweeney, C., Wofsy, S. C., and Kort, E. A.: Inverse modelling of CH₄ emissions for 2010–2011 using different satellite retrieval products from GOSAT and SCIAMACHY, *Atmospheric Chemistry and Physics*, 15, 113–133, <https://doi.org/10.5194/acp-15-113-2015>, 2015.
- Apituley, A., Pedergnana, M., Sneep, M., Veeffkind, J., Loyola, D., Hasekamp, O., A., L., Borsdorff, T., Martinez Velarte, M., and Mandal, S.: Sentinel-5 precursor/TROPOMI Level 2 Product User Manual Methane, Tech. rep., https://sentiwiki.copernicus.eu/_attachments/1673595/SRON-S5P-LEV2-MA-001-Sentinel-5PLevel2ProductUserManualMethane2024-2.8.0.pdf?inst-v=f449a58f-77a5-411d-952f-30467b0f995b, 2024.
- Balagus, N., Jacob, D. J., Lorente, A., Maasakkers, J. D., Parker, R. J., Boesch, H., Chen, Z., Kelp, M. M., Nesser, H., and Varon, D. J.: A blended TROPOMI+GOSAT satellite data product for atmospheric methane using machine learning to correct retrieval biases, *Atmospheric Measurement Techniques*, 16, 3787–3807, <https://doi.org/10.5194/amt-16-3787-2023>, 2023.
- Barré, J., Aben, I., Agustí-Panareda, A., Balsamo, G., Bousserez, N., Dueben, P., Engelen, R., Inness, A., Lorente, A., McNorton, J., Peuch, V.-H., Radnoti, G., and Ribas, R.: Systematic detection of local CH₄ anomalies by combining satellite measurements with high-resolution forecasts, *Atmospheric Chemistry and Physics*, 21, 5117–5136, <https://doi.org/10.5194/acp-21-5117-2021>, 2021.
- Berchet, A., Sollum, E., Thompson, R. L., Pison, I., Thanwerdas, J., Broquet, G., Chevallier, F., Aalto, T., Berchet, A., Bergamaschi, P., Brunner, D., Engelen, R., Fortems-Cheiney, A., Gerbig, C., Groot Zwaftink, C. D., Haussaire, J.-M., Henne, S., Houweling, S., Karstens, U., Kutsch, W. L., Lujikx, I. T., Monteil, G., Palmer, P. I., van Peet, J. C. A., Peters, W., Peylin, P., Potier, E., Rödenbeck, C., Saunio, M., Scholze, M., Tsuruta, A., and Zhao, Y.: The Community Inversion Framework v1.0: a unified system for atmospheric inversion studies, *Geoscientific Model Development*, 14, 5331–5354, <https://doi.org/10.5194/gmd-14-5331-2021>, 2021.
- Bergamaschi, P., Corazza, M., Karstens, U., Athanassiadou, M., Thompson, R. L., Pison, I., Manning, A. J., Bousquet, P., Segers, A., Vermeulen, A. T., Janssens-Maenhout, G., Schmidt, M., Ramonet, M., Meinhardt, F., Aalto, T., Haszpra, L., Moncrieff, J., Popa, M. E., Lowry, D., Steinbacher, M., Jordan, A., O'Doherty, S., Piacentino, S., and Dlugokencky, E.: Top-down estimates of European CH₄ and N₂O emissions based on four different inverse models, *Atmospheric Chemistry and Physics*, 15, 715–736, <https://doi.org/10.5194/acp-15-715-2015>, 2015.
- Borsdorff, T., aan de Brugh, J., Schneider, A., Lorente, A., Birk, M., Wagner, G., Kivi, R., Hase, F., Feist, D. G., Sussmann, R., Rettinger, M., Wunch, D., Warneke, T., and Landgraf, J.: Improving the TROPOMI CO data product: update of the spectroscopic database and destripping of single orbits, *Atmospheric Measurement Techniques*, 12, 5443–5455, <https://doi.org/10.5194/amt-12-5443-2019>, publisher: Copernicus GmbH, 2019.
- Borsdorff, T., Martinez-Velarte, M. C., Sneep, M., ter Linden, M., and Landgraf, J.: Random Forest Classifier for Cloud Clearing of the Operational TROPOMI XCH₄ Product, *Remote Sensing*, 16, <https://doi.org/10.3390/rs16071208>, 2024.

- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: Mission Objectives and Measurement Modes, *Journal of the Atmospheric Sciences*, 56, 127–150, [https://doi.org/10.1175/1520-0469\(1999\)056<0127:SMOAMM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<0127:SMOAMM>2.0.CO;2), 1999.
- Brasseur, G. and Jacob, D.: *Modeling of Atmospheric Chemistry*, Cambridge University Press, ISBN 978-1-108-21095-9, 2017.
- Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.
- Broquet, G., Chevallier, F., Rayner, P., Aulagnier, C., Pison, I., Ramonet, M., Schmidt, M., Vermeulen, A. T., and Ciais, P.: A European summertime CO₂ biogenic flux inversion at mesoscale from continuous in situ mixing ratio measurements, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD016202>, 2011.
- Buchwitz, M.: The Greenhouse Gas Climate Change Initiative (GHG-CCI): Comparison and quality assessment of near-surface-sensitive satellite-derived CO₂ and CH₄ global data sets, 2015.
- Butz, A., Galli, A., Hasekamp, O., Landgraf, J., Tol, P., and Aben, I.: TROPOMI aboard Sentinel-5 Precursor: Prospective performance of CH₄ retrievals for aerosol and cirrus loaded atmospheres, *Remote Sensing of Environment*, 120, 267–276, <https://doi.org/10.1016/j.rse.2011.05.030>, 2012.
- Chen, T. and Guestrin, C.: XGBoost: A Scalable Tree Boosting System, in: *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, KDD '16*, pp. 785–794, Association for Computing Machinery, ISBN 978-1-4503-4232-2, <https://doi.org/10.1145/2939672.2939785>, 2016.
- Chen, Z., Jacob, D. J., Nesser, H., Sulprizio, M. P., Lorente, A., Varon, D. J., Lu, X., Shen, L., Qu, Z., Penn, E., and Yu, X.: Methane emissions from China: a high-resolution inversion of TROPOMI satellite observations, *Atmospheric Chemistry and Physics*, 22, 10 809–10 826, <https://doi.org/10.5194/acp-22-10809-2022>, 2022.
- Chen, Z., Jacob, D. J., Gautam, R., Omara, M., Stavins, R. N., Stowe, R. C., Nesser, H. O., Sulprizio, M. P., Lorente, A., Varon, D. J., Lu, X., Shen, L., Qu, Z., Pendergrass, D. C., and Hancock, S.: Satellite quantification of methane emissions and oil/gas methane intensities from individual countries in the Middle East and North Africa: implications for climate action, <https://doi.org/10.5194/egusphere-2022-1504>, 2023.
- Cressot, C., Chevallier, F., Bousquet, P., Crevoisier, C., Dlugokencky, E. J., Fortems-Cheiney, A., Frankenberg, C., Parker, R., Pison, I., Scheepmaker, R. A., Montzka, S. A., Krummel, P. B., Steele, L. P., and Langenfelds, R. L.: On the consistency between global and regional methane emissions inferred from SCIAMACHY, TANSO-FTS, IASI and surface measurements, *Atmospheric Chemistry and Physics*, 14, 577–592, <https://doi.org/10.5194/acp-14-577-2014>, 2014.
- Crippa, M., Guizzardi, D., Schaaf, E., Monforti-Ferrario, F., Quadrelli, R., Risquez Martin, A., Rossi, S., Vignati, E., Muntean, M., Brandao De Melo, J., Oom, D., Pagani, F., Banja, M., Taghavi-Moharamli, P., Köykkä, J., Grassi, G., Branco, A., and San-Miguel, J.: GHG emissions of all world countries – 2023, Publications Office of the European Union, <https://doi.org/doi/10.2760/953322>, 2023.
- Dils, B., Zhou, M., Camy-Peyret, C., De Mazière, M., Kangah, Y., Langerock, B., Prunet, P., Serio, C., Siddans, R., and Kerridge, B.: Independent Validation of IASI/MetOp-A LMD and RAL CH₄ Products Using CAMS Model, in Situ Profiles, and Ground-Based FTIR Measurements, *Atmospheric Measurement Techniques*, 17, 5491–5524, <https://doi.org/10.5194/amt-17-5491-2024>, 2024.
- East, J. D., Jacob, D. J., Jervis, D., Balasus, N., Estrada, L. A., Hancock, S. E., Sulprizio, M. P., Thomas, J., Wang, X., Chen, Z., Varon, D. J., and Worden, J. R.: Worldwide Inference of National Methane Emissions by Inversion of Satellite Observations with UNFCCC Prior Estimates, *Nature Communications*, 16, 11 004, <https://doi.org/10.1038/s41467-025-67122-8>, 2025.
- Etiopie, G., Ciotoli, G., Schwietzke, S., and Schoell, M.: Gridded maps of geological methane emissions and their isotopic signature, *Earth System Science Data*, 11, 1–22, <https://doi.org/10.5194/essd-11-1-2019>, 2019.

- European Space Agency: Sentinel-5P Calibration and Validation Plan for the Operational Phase, Tech. Rep. ESA-EOPG-CSCOP-PL-0073, <https://sentinels.copernicus.eu/documents/247904/2474724/Sentinel-5P-Calibration-and-Validation-Plan.pdf>, 2017.
- European Space Agency: Copernicus Sentinel-5P (processed by ESA), TROPOMI Level 2 Methane Total Column products. Version 02., 2021.
- 830 Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H.: Chapter 7: The Earth's energy budget, climate feedbacks, and climate sensitivity, <https://doi.org/10.25455/wgtn.16869671>, 2021.
- Fortems-Cheiney, A., Pison, I., Broquet, G., Dufour, G., Berchet, A., Potier, E., Coman, A., Siour, G., and Costantino, L.: Variational regional inverse modeling of reactive species emissions with PYVAR-CHIMERE-v2019, *Geoscientific Model Development*, 14, 2939–2957, <https://doi.org/10.5194/gmd-14-2939-2021>, 2021.
- 835 Gilbert, J. and Lemaréchal, C.: Some numerical experiments with variable storage quasi Newton algorithms, *Math. Program.*, 45, 407–435, <https://doi.org/10.1007/BF01589113>, 1989.
- Hancock, S. E., Jacob, D., Chen, Z., Nesser, H., Davitt, A., Varon, D. J., Sulprizio, M. P., Balasus, N., Estrada, L. A., East, J. D., Penn, E., Randles, C. A., Worden, J., Aben, I., Parker, R. J., and Maasakkers, J. D.: Satellite quantification of methane emissions from South American countries: A high-resolution inversion of TROPOMI and GOSAT observations, *EGUsphere*, pp. 1–33, <https://doi.org/10.5194/egusphere-2024-1763>, publisher: Copernicus GmbH, 2024.
- 840 Hasekamp, O., Lorente, A., Hu, H., Butz, A., aan de Brugh, J., and Landgraf, J.: Algorithm Theoretical Basis Document for Sentinel-5 Precursor Methane Retrieval, SRON Netherlands Institute for Space Research, Tech. rep., <https://sentinels.copernicus.eu/documents/247904/2476257/Sentinel-5P-TROPOMI-ATBD-Methane-retrieval.pdf>, 2022.
- 845 Hu, H., Hasekamp, O., Butz, A., Galli, A., Landgraf, J., Aan de Brugh, J., Borsdorff, T., Scheepmaker, R., and Aben, I.: The operational methane retrieval algorithm for TROPOMI, *Atmospheric Measurement Techniques*, 9, 5423–5440, <https://doi.org/10.5194/amt-9-5423-2016>, 2016.
- Ide, K., Ghil, M., and Lorenc, A.: Unified Notation for Data Assimilation: Operational, Sequential and Variational, *Journal of the Meteorological Society of Japan*, 75, 181–189, 1997.
- 850 Ioannidis, E., Meesters, A., Steiner, M., Brunner, D., Reum, F., Pison, I., Berchet, A., Thompson, R., Sollum, E., Koch, F.-T., Gerbig, C., Wang, F., Maksyutov, S., Tsuruta, A., Tenkanen, M., Aalto, T., Monteil, G., Lin, H., Ren, G., Scholze, M., and Houweling, S.: An Inter-Comparison of Inverse Models for Estimating European CH₄ Emissions, *Earth System Science Data Discussions*, pp. 1–42, <https://doi.org/10.5194/essd-2025-235>, 2025.
- IPCC: Climate Change 2021 - The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Tech. rep., IPCC, ISBN 978-1-107-41532-4, <https://www.ipcc.ch/report/ar6/wg1/#FullReport/>, 2021.
- 855 Jacob, D. J., Varon, D. J., Cusworth, D. H., Dennison, P. E., Frankenberg, C., Gautam, R., Guanter, L., Kelley, J., McKeever, J., Ott, L. E., Poulter, B., Qu, Z., Thorpe, A. K., Worden, J. R., and Duren, R. M.: Quantifying Methane Emissions from the Global Scale down to Point Sources Using Satellite Observations of Atmospheric Methane, *Atmospheric Chemistry and Physics*, 22, 9617–9646, <https://doi.org/10.5194/acp-22-9617-2022>, 2022.
- 860 Karlsson, K.-G., Riihelä, A., Trentmann, J., Stengel, M., Solodovnik, I., Meirink, J. F., Devasthale, A., Jääskeläinen, E., Kallio-Myers, V., Eliasson, S., Benas, N., Johansson, E., Stein, D., Finkensieper, S., Håkansson, N., Akkermans, T., Clerbaux, N., Selbach, N.,

- Marc, S., and Hollmann, R.: CLARA-A3: CM SAF cLoud, Albedo and surface RAdiation dataset from AVHRR data - Edition 3, https://doi.org/10.5676/EUM_SAF_CM/CLARA_AVHRR/V003, 2023.
- 865 Ke, G., Meng, Q., Finley, T., Wang, T., Chen, W., Ma, W., Ye, Q., and Liu, T.-Y.: LightGBM: a highly efficient gradient boosting decision tree, in: Proceedings of the 31st International Conference on Neural Information Processing Systems, NIPS'17, p. 3149–3157, Curran Associates Inc., ISBN 9781510860964, 2017.
- Kuhlmann, G., Stavropoulou, F., Schwietzke, S., Zavala-Araiza, D., Thorpe, A., Hueni, A., Emmenegger, L., Calcan, A., Röckmann, T., and Brunner, D.: Evidence of Successful Methane Mitigation in One of Europe's Most Important Oil Production Region, *Atmospheric*
870 *Chemistry and Physics*, 25, 5371–5385, <https://doi.org/10.5194/acp-25-5371-2025>, 2025.
- Lahoz, W. A. and Schneider, P.: Data assimilation: making sense of Earth Observation, *Frontiers in Environmental Science*, 2, <https://doi.org/10.3389/fenvs.2014.00016>, publisher: Frontiers, 2014.
- Landgraf, J., Lorente, A., Borsdorff, T., Mandal, S., Langerock, B., and Sha, M.: ATM-MPC Mission Performance Cluster Methane [L2_CH4___] Readme, Tech. rep., [https://sentiwiki.copernicus.eu/__attachments/1673595/](https://sentiwiki.copernicus.eu/__attachments/1673595/S5P-MPC-SRON-PRF-CH4-Sentinel-5PMethaneProductReadmeFile2024-2.7.pdf)
875 [S5P-MPC-SRON-PRF-CH4-Sentinel-5PMethaneProductReadmeFile2024-2.7.pdf](https://sentiwiki.copernicus.eu/__attachments/1673595/S5P-MPC-SRON-PRF-CH4-Sentinel-5PMethaneProductReadmeFile2024-2.7.pdf), 2024.
- Laughner, J. L., Toon, G. C., Mendonca, J., Petri, C., Roche, S., Wunch, D., Blavier, J.-F., Griffith, D. W. T., Heikkinen, P., Keeling, R. F., Kiel, M., Kivi, R., Roehl, C. M., Stephens, B. B., Baier, B. C., Chen, H., Choi, Y., Deutscher, N. M., DiGangi, J. P., Gross, J., Herkommer, B., Jeseck, P., Laemmel, T., Lan, X., McGee, E., McKain, K., Miller, J., Morino, I., Notholt, J., Ohyama, H., Pollard, D. F., Rettinger, M., Riris, H., Rousogenous, C., Sha, M. K., Shiomi, K., Strong, K., Sussmann, R., Té, Y., Velazco, V. A., Wofsy, S. C., Zhou, M., and
880 Wennberg, P. O.: The Total Carbon Column Observing Network's GGG2020 data version, *Earth System Science Data*, 16, 2197–2260, <https://doi.org/10.5194/essd-16-2197-2024>, 2024.
- Lauvaux, T., Giron, C., Mazzolini, M., d'Aspremont, A., Duren, R., Cusworth, D., Shindell, D., and Ciais, P.: Global assessment of oil and gas methane ultra-emitters, *Science*, 375, 557–561, <https://doi.org/10.1126/science.abj4351>, 2022.
- Liang, R., Zhang, Y., Chen, W., Zhang, P., Liu, J., Chen, C., Mao, H., Shen, G., Qu, Z., Chen, Z., Zhou, M., Wang, P., Parker, R. J.,
885 Boesch, H., Lorente, A., Maasackers, J. D., and Aben, I.: East Asian methane emissions inferred from high-resolution inversions of GOSAT and TROPOMI observations: a comparative and evaluative analysis, *Atmospheric Chemistry and Physics*, 23, 8039–8057, <https://doi.org/10.5194/acp-23-8039-2023>, publisher: Copernicus GmbH, 2023.
- Lindqvist, H., Kivimäki, E., Häkkinen, T., Tsuruta, A., Schneising, O., Buchwitz, M., Lorente, A., Martinez Velarte, M., Borsdorff, T., Alberti, C., Backman, L., Buschmann, M., Chen, H., Dubravica, D., Hase, F., Heikkinen, P., Karppinen, T., Kivi, R., McGee, E., Notholt, J.,
890 Rautiainen, K., Roche, S., Simpson, W., Strong, K., Tu, Q., Wunch, D., Aalto, T., and Tamminen, J.: Evaluation of Sentinel-5P TROPOMI Methane Observations at Northern High Latitudes, *Remote Sensing*, 16, 2979, <https://doi.org/10.3390/rs16162979>, 2024.
- Lorente, A., Borsdorff, T., Butz, A., Hasekamp, O., aan de Brugh, J., Schneider, A., Wu, L., Hase, F., Kivi, R., Wunch, D., Pollard, D. F., Shiomi, K., Deutscher, N. M., Velazco, V. A., Roehl, C. M., Wennberg, P. O., Warneke, T., and Landgraf, J.: Methane retrieved from TROPOMI: improvement of the data product and validation of the first 2 years of measurements, *Atmospheric Measurement Techniques*,
895 14, 665–684, <https://doi.org/10.5194/amt-14-665-2021>, 2021.
- Lorente, A., Borsdorff, T., Martinez-Velarte, M. C., Butz, A., Hasekamp, O. P., Wu, L., and Landgraf, J.: Evaluation of the methane full-physics retrieval applied to TROPOMI ocean sun glint measurements, *Atmospheric Measurement Techniques*, 15, 6585–6603, <https://doi.org/10.5194/amt-15-6585-2022>, publisher: Copernicus GmbH, 2022.
- Lorente, A., Borsdorff, T., Martinez-Velarte, M. C., and Landgraf, J.: Accounting for surface reflectance spectral features in TROPOMI
900 methane retrievals, *Atmospheric Measurement Techniques Discussions*, 2023, 1–15, <https://doi.org/10.5194/amt-2022-255>, 2023.

- Lu, L., Cohen, J. B., Qin, K., Li, X., and He, Q.: Identifying Missing Sources and Reducing NO_x Emissions Uncertainty over China Using Daily Satellite Data and a Mass-Conserving Method, *Atmospheric Chemistry and Physics*, 25, 2291–2309, <https://doi.org/10.5194/acp-25-2291-2025>, 2025a.
- Lu, L., Cohen, J. B., Qin, K., Tiwari, P., Hu, W., Gao, H., and Zheng, B.: New Perspective on Using Observational Uncertainty to Improve Reliability of NO_x Emissions Over Northern China, *IEEE Transactions on Geoscience and Remote Sensing*, 63, 1–15, <https://doi.org/10.1109/TGRS.2025.3620116>, 2025b.
- Maazallahi, H., Stavropoulou, F., Sutanto, S. J., Steiner, M., Brunner, D., Mertens, M., Jöckel, P., Visschedijk, A., Denier van der Gon, H., Dellaert, S., Velandia Salinas, N., Schwietzke, S., Zavala-Araiza, D., Ghemulet, S., Pana, A., Ardelean, M., Corbu, M., Calcan, A., Conley, S. A., Smith, M. L., and Röckmann, T.: Airborne in Situ Quantification of Methane Emissions from Oil and Gas Production in Romania, *Atmospheric Chemistry and Physics*, 25, 1497–1511, <https://doi.org/10.5194/acp-25-1497-2025>, 2025.
- Mailler, S., Menut, L., Khvorostyanov, D., Valari, M., Couvidat, F., Siour, G., Turquety, S., Briant, R., Tuccella, P., Bessagnet, B., Colette, A., Létinois, L., Markakis, K., and Meleux, F.: CHIMERE-2017: from urban to hemispheric chemistry-transport modeling, *Geoscientific Model Development*, 10, 2397–2423, <https://doi.org/10.5194/gmd-10-2397-2017>, 2017.
- Menut, L., Bessagnet, B., Khvorostyanov, D., Beekmann, M., Blond, N., Colette, A., Coll, I., Curci, G., Foret, G., Hodzic, A., Mailler, S., Meleux, F., Monge, J.-L., Pison, I., Siour, G., Turquety, S., Valari, M., Vautard, R., and Vivanco, M. G.: CHIMERE 2013: a model for regional atmospheric composition modelling, *Geoscientific Model Development*, 6, 981–1028, <https://doi.org/10.5194/gmd-6-981-2013>, 2013.
- Monteil, G., Houweling, S., Butz, A., Guerlet, S., Schepers, D., Hasekamp, O., Frankenberg, C., Scheepmaker, R., Aben, I., and Röckmann, T.: Comparison of CH₄ inversions based on 15 months of GOSAT and SCIAMACHY observations, *Journal of Geophysical Research: Atmospheres*, 118, 11,807–11,823, <https://doi.org/https://doi.org/10.1002/2013JD019760>, 2013.
- Monteil, G., Broquet, G., Scholze, M., Lang, M., Karstens, U., Gerbig, C., Koch, F.-T., Smith, N. E., Thompson, R. L., Luijkx, I. T., White, E., Meesters, A., Ciais, P., Ganesan, A. L., Manning, A., Mischurow, M., Peters, W., Peylin, P., Tarniewicz, J., Rigby, M., Rödenbeck, C., Vermeulen, A., and Walton, E. M.: The regional European atmospheric transport inversion comparison, EUROCOM: first results on European-wide terrestrial carbon fluxes for the period 2006–2015, *Atmospheric Chemistry and Physics*, 20, 12 063–12 091, <https://doi.org/10.5194/acp-20-12063-2020>, 2020.
- Nathan, B., Maasakkers, J. D., Naus, S., Gautam, R., Omara, M., Varon, D. J., Sulprizio, M. P., Lorente, A., Borsdorff, T., Parker, R. J., and Aben, I.: Assessing methane emissions from collapsing Venezuelan oil production using TROPOMI, <https://doi.org/10.5194/egusphere-2023-2887>, 2023.
- Naus, S., Maasakkers, J. D., Gautam, R., Omara, M., Stikker, R., Veenstra, A. K., Nathan, B., Irakulis-Loitxate, I., Guanter, L., Pandey, S., Girard, M., Lorente, A., Borsdorff, T., and Aben, I.: Assessing the Relative Importance of Satellite-Detected Methane Superemitters in Quantifying Total Emissions for Oil and Gas Production Areas in Algeria, *Environmental Science & Technology*, 57, 19 545–19 556, <https://doi.org/10.1021/acs.est.3c04746>, 2023.
- Nesser, H., Jacob, D. J., Maasakkers, J. D., Lorente, A., Chen, Z., Lu, X., Shen, L., Qu, Z., Sulprizio, M. P., Winter, M., Ma, S., Bloom, A. A., Worden, J. R., Stavins, R. N., and Randles, C. A.: High-resolution US methane emissions inferred from an inversion of 2019 TROPOMI satellite data: contributions from individual states, urban areas, and landfills, *Atmospheric Chemistry and Physics*, 24, 5069–5091, <https://doi.org/10.5194/acp-24-5069-2024>, publisher: Copernicus GmbH, 2024.

- Nesser, H., Bowman, K. W., Thill, M. D., Varon, D. J., Randles, C. A., Tewari, A., Cardoso-Saldaña, F. J., Reidy, E., Maasackers, J. D., and Jacob, D. J.: Predicting and Correcting the Influence of Boundary Conditions in Regional Inverse Analyses, *Geoscientific Model Development*, 18, 9279–9291, <https://doi.org/10.5194/gmd-18-9279-2025>, 2025.
- 940 Nygård, T., Papritz, L., Naakka, T., and Vihma, T.: Cold Wintertime Air Masses over Europe: Where Do They Come from and How Do They Form?, *Weather and Climate Dynamics*, 4, 943–961, <https://doi.org/10.5194/wcd-4-943-2023>, 2023.
- Parker, R. J., Webb, A., Boesch, H., Somkuti, P., Barrio Guillo, R., Di Noia, A., Kalaitzi, N., Anand, J. S., Bergamaschi, P., Chevallier, F., Palmer, P. I., Feng, L., Deutscher, N. M., Feist, D. G., Griffith, D. W. T., Hase, F., Kivi, R., Morino, I., Notholt, J., Oh, Y.-S., Ohyama, H., Petri, C., Pollard, D. F., Roehl, C., Sha, M. K., Shiomu, K., Strong, K., Sussmann, R., Té, Y., Velasco, V. A., Warneke,
- 945 T., Wennberg, P. O., and Wunch, D.: A decade of GOSAT Proxy satellite CH₄ observations, *Earth System Science Data*, 12, 3383–3412, <https://doi.org/10.5194/essd-12-3383-2020>, 2020.
- Petrenko, V., Smith, A., and Schaefer, H. e. a.: Minimal geological methane emissions during the Younger Dryas–Preboreal abrupt warming event, *Nature*, 548, 443–446, <https://doi.org/https://doi.org/10.1038/nature23316>, 2017.
- Petrescu, A. M. R., Peters, G. P., Engelen, R., Houweling, S., Brunner, D., Tsuruta, A., Matthews, B., Patra, P. K., Belikov, D., Thompson,
- 950 R. L., Höglund-Isaksson, L., Zhang, W., Segers, A. J., Etiope, G., Ciotoli, G., Peylin, P., Chevallier, F., Aalto, T., Andrew, R. M., Bastviken, D., Berchet, A., Broquet, G., Conchedda, G., Dellaert, S. N. C., Denier van der Gon, H., Gütschow, J., Haussaire, J.-M., Lauerwald, R., Markkanen, T., van Peet, J. C. A., Pison, I., Regnier, P., Solum, E., Scholze, M., Tenkanen, M., Tubiello, F. N., van der Werf, G. R., and Worden, J. R.: Comparison of Observation- and Inventory-Based Methane Emissions for Eight Large Global Emitters, *Earth System Science Data*, 16, 4325–4350, <https://doi.org/10.5194/essd-16-4325-2024>, 2024.
- 955 Pison, I., Berchet, A., Saunio, M., Bousquet, P., Broquet, G., Conil, S., Delmotte, M., Ganesan, A., Laurent, O., Martin, D., O’Doherty, S., Ramonet, M., Spain, T. G., Vermeulen, A., and Yver Kwok, C.: How a European network may help with estimating methane emissions on the French national scale, *Atmospheric Chemistry and Physics*, 18, 3779–3798, <https://doi.org/10.5194/acp-18-3779-2018>, 2018.
- Pison, I., Thompson, R., Sollum, E., Kouyaté, M., Berchet, A., and Haussaire, J.-M.: Methane and nitrous oxide fluxes from the CIF, Tech. rep., https://verify.lsce.ipsl.fr/images/PublicDeliverables/VERIFY_D410_Methane_and_nitrous_oxide_fluxes_from_the_CIF_v1.pdf, 2021.
- 960 Qu, Z., Jacob, D. J., Shen, L., Lu, X., Zhang, Y., Scarpelli, T. R., Nesser, H., Sulprizio, M. P., Maasackers, J. D., Bloom, A. A., Worden, J. R., Parker, R. J., and Delgado, A. L.: Global distribution of methane emissions: a comparative inverse analysis of observations from the TROPOMI and GOSAT satellite instruments, *Atmospheric Chemistry and Physics*, 21, 14 159–14 175, <https://doi.org/10.5194/acp-21-14159-2021>, 2021.
- 965 Raivonen, M., Smolander, S., Backman, L., Susiluoto, J., Aalto, T., Markkanen, T., Mäkelä, J., Rinne, J., Peltola, O., Aurela, M., Lohila, A., Tomasic, M., Li, X., Larmola, T., Juutinen, S., Tuittila, E.-S., Heimann, M., Sevanto, S., Kleinen, T., Brovkin, V., and Vesala, T.: HIMMELI v1.0: Helsinki Model of METHane buiLd-up and emIssion for peatlands, *Geoscientific Model Development*, 10, 4665–4691, <https://doi.org/10.5194/gmd-10-4665-2017>, 2017.
- Randerson, J., van der Werf, G., Giglio, L., Collatz, G., and Kasibhatla, P.: Global Fire Emissions Database, Version 4.1 (GFEDv4), Tech.
- 970 rep., <https://doi.org/10.3334/ORNLDAAAC/1293>, 2018.
- Rayner, P. J., Michalak, A. M., and Chevallier, F.: Fundamentals of data assimilation applied to biogeochemistry, *Atmospheric Chemistry and Physics*, 19, 13 911–13 932, <https://doi.org/10.5194/acp-19-13911-2019>, 2019.

- Saad, K. M., Wunch, D., Toon, G. C., Bernath, P., Boone, C., Connor, B., Deutscher, N. M., Griffith, D. W. T., Kivi, R., Notholt, J., Roehl, C., Schneider, M., Sherlock, V., and Wennberg, P. O.: Derivation of Tropospheric Methane from TCCON CH₄ and HF Total Column Observations, *Atmospheric Measurement Techniques*, 7, 2907–2918, <https://doi.org/10.5194/amt-7-2907-2014>, 2014.
- 975 Santaren, D., Broquet, G., Bréon, F.-M., Chevallier, F., Siméoni, D., Zheng, B., and Ciais, P.: A local- to national-scale inverse modeling system to assess the potential of spaceborne CO₂ measurements for the monitoring of anthropogenic emissions, *Atmospheric Measurement Techniques*, 14, 403–433, <https://doi.org/10.5194/amt-14-403-2021>, publisher: Copernicus GmbH, 2021.
- Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth System Science Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.
- 985 990 Schneising, O.: Product User Guide (PUG) TROPOMI WFM-DOAS (TROPOMI/WFMD) XCH₄, Tech. rep., https://www.iup.uni-bremen.de/carbon_ghg/products/tropomi_wfmd/data/v18/pug_wfmd.pdf, 2022.
- Schneising, O.: Algorithm Theoretical Basis Document (ATBD) - TROPOMI WFM-DOAS XCH₄, Tech. rep., https://climate.esa.int/media/documents/ATBDv3_GHG-CCI_CH4_S5P_WFMD.pdf, 2023.
- Schneising, O., Buchwitz, M., Reuter, M., Bovensmann, H., Burrows, J. P., Borsdorff, T., Deutscher, N. M., Feist, D. G., Griffith, D. W. T., Hase, F., Hermans, C., Iraci, L. T., Kivi, R., Landgraf, J., Morino, I., Notholt, J., Petri, C., Pollard, D. F., Roche, S., Shiomi, K., Strong, K., Sussmann, R., Velasco, V. A., Warneke, T., and Wunch, D.: A scientific algorithm to simultaneously retrieve carbon monoxide and methane from TROPOMI onboard Sentinel-5 Precursor, *Atmospheric Measurement Techniques*, 12, 6771–6802, <https://doi.org/10.5194/amt-12-6771-2019>, 2019.
- 995 1000 Schneising, O., Buchwitz, M., Reuter, M., Vanselow, S., Bovensmann, H., and Burrows, J. P.: Remote sensing of methane leakage from natural gas and petroleum systems revisited, *Atmospheric Chemistry and Physics*, 20, 9169–9182, <https://doi.org/10.5194/acp-20-9169-2020>, publisher: Copernicus GmbH, 2020.
- Schneising, O., Buchwitz, M., Hachmeister, J., Vanselow, S., Reuter, M., Buschmann, M., Bovensmann, H., and Burrows, J. P.: Advances in retrieving XCH₄ and XCO from Sentinel-5 Precursor: improvements in the scientific TROPOMI/WFMD algorithm, *Atmospheric Measurement Techniques*, 16, 669–694, <https://doi.org/10.5194/amt-16-669-2023>, 2023.
- 1005 Schuit, B. J., Maasackers, J. D., Bijl, P., Mahapatra, G., Van Den Berg, A.-W., Pandey, S., Lorente, A., Borsdorff, T., Houweling, S., Varon, D. J., McKeever, J., Jervis, D., Girard, M., Irakulis-Loitxate, I., Gorroño, J., Guanter, L., Cusworth, D. H., and Aben, I.: Automated detection and monitoring of methane super-emitters using satellite data, *Atmospheric Chemistry and Physics*, 23, 9071–9098, <https://doi.org/10.5194/acp-23-9071-2023>, 2023.
- Shen, L., Gautam, R., Omara, M., Zavala-Araiza, D., Maasackers, J. D., Scarpelli, T. R., Lorente, A., Lyon, D., Sheng, J., Varon, D. J., Nesser, H., Qu, Z., Lu, X., Sulprizio, M. P., Hamburg, S. P., and Jacob, D. J.: Satellite quantification of oil and natural gas methane
- 1010

- emissions in the US and Canada including contributions from individual basins, *Atmospheric Chemistry and Physics*, 22, 11 203–11 215, <https://doi.org/10.5194/acp-22-11203-2022>, 2022.
- 1015 Szopa, S., Naik, V., Adhikary, B., Artaxo, P., Bernsten, T., Collins, W., Fuzzi, S., Gallardo, L., Kiendler-Scharr, A., Klimont, Z., Liao, H., Unger, N., and Zanis, P.: *Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 1 edn., <https://doi.org/10.1017/9781009157896>, 2023.
- Szénási, B., Berchet, A., Broquet, G., Segers, A., van der Gon, H. D., Krol, M., Hullegie, J. J., Kiesow, A., Günther, D., Petrescu, A. M. R., Saunois, M., Bousquet, P., and Pison, I.: A pragmatic protocol for characterising errors in atmospheric inversions of methane emissions over Europe, *Tellus B: Chemical and Physical Meteorology*, 73, 1–23, <https://doi.org/10.1080/16000889.2021.1914989>, 2021.
- 1020 T. Hilbig, T., Schneising, O., and Bösch, H.: Report on CO₂ and CH₄ Satellite Datasets: TROPOMI XCH₄ Comparison, Tech. rep., https://folk.nilu.no/~rthompson/eyeclima_reports/EYECLIMA_D1.1_v1.pdf, 2023.
- Tsuruta, A., Kivimäki, E., Lindqvist, H., Karppinen, T., Backman, L., Hakkarainen, J., Schneising, O., Buchwitz, M., Lan, X., Kivi, R., Chen, H., Buschmann, M., Herkommer, B., Notholt, J., Roehl, C., Té, Y., Wunch, D., Tamminen, J., and Aalto, T.: CH₄ Fluxes Derived from Assimilation of TROPOMI XCH₄ in CarbonTracker Europe-CH₄: Evaluation of Seasonality and Spatial Distribution in the Northern High Latitudes, *Remote Sensing*, 15, <https://doi.org/10.3390/rs15061620>, 2023.
- 1025 Tu, Q., Schneider, M., Hase, F., Khosrawi, F., Ertl, B., Necki, J., Dubravica, D., Diekmann, C. J., Blumenstock, T., and Fang, D.: Quantifying CH₄ Emissions in Hard Coal Mines from TROPOMI and IASI Observations Using the Wind-Assigned Anomaly Method, *Atmospheric Chemistry and Physics*, 22, 9747–9765, <https://doi.org/10.5194/acp-22-9747-2022>, 2022.
- United Nations Environment Programme: *An Eye on Methane 2025: From Measurement to Momentum - Data Is Driving Action - Now the Pace Must Match the Promise*, Tech. rep., ISBN 978-92-807-4236-7, <https://doi.org/10.59117/20.500.11822/48664>, 2025.
- 1030 Vanselow, S., Schneising, O., Buchwitz, M., Reuter, M., Bovensmann, H., Boesch, H., and Burrows, J. P.: Automated Detection of Regions with Persistently Enhanced Methane Concentrations Using Sentinel-5 Precursor Satellite Data, *Atmospheric Chemistry and Physics*, 24, 10 441–10 473, <https://doi.org/10.5194/acp-24-10441-2024>, 2024.
- Veefkind, J., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H., de Haan, J., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sensing of Environment*, 120, 70–83, <https://doi.org/https://doi.org/10.1016/j.rse.2011.09.027>, the Sentinel Missions - New Opportunities for Science, 2012.
- 1035 Veefkind, J. P., Serrano-Calvo, R., de Gouw, J., Dix, B., Schneising, O., Buchwitz, M., Barré, J., van der A, R. J., Liu, M., and Levelt, P. F.: Widespread Frequent Methane Emissions From the Oil and Gas Industry in the Permian Basin, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037 479, <https://doi.org/10.1029/2022JD037479>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022JD037479>, 2023.
- 1040 Weber, T., Wiseman, N., and Kock, A.: Global ocean methane emissions dominated by shallow coastal waters, *Nat Commun*, 10, <https://doi.org/https://doi.org/10.1038/s41467-019-12541-7>, 2019.
- 1045 Williams, J. P., Omara, M., Himmelberger, A., Zavala-Araiza, D., MacKay, K., Benmergui, J., Sargent, M., Wofsy, S. C., Hamburg, S. P., and Gautam, R.: Small Emission Sources in Aggregate Disproportionately Account for a Large Majority of Total Methane Emissions from the US Oil and Gas Sector, *Atmospheric Chemistry and Physics*, 25, 1513–1532, <https://doi.org/10.5194/acp-25-1513-2025>, 2025.

- Yu, X., Millet, D. B., and Henze, D. K.: How well can inverse analyses of high-resolution satellite data resolve heterogeneous methane fluxes? Observing system simulation experiments with the GEOS-Chem adjoint model (v35), *Geoscientific Model Development*, 14, 7775–7793, <https://doi.org/10.5194/gmd-14-7775-2021>, 2021.
- Yu, X., Millet, D. B., Henze, D. K., Turner, A. J., Delgado, A. L., Bloom, A. A., and Sheng, J.: A high-resolution satellite-based map of global methane emissions reveals missing wetland, fossil fuel, and monsoon sources, *Atmospheric Chemistry and Physics*, 23, 3325–3346, <https://doi.org/10.5194/acp-23-3325-2023>, 2023.
- Zhang, Y., Gautam, R., Pandey, S., Omara, M., Maasakkers, J. D., Sadavarte, P., Lyon, D., Nesser, H., Sulprizio, M. P., Varon, D. J., Zhang, R., Houweling, S., Zavala-Araiza, D., Alvarez, R. A., Lorente, A., Hamburg, S. P., Aben, I., and Jacob, D. J.: Quantifying methane emissions from the largest oil-producing basin in the United States from space, *Science Advances*, 6, eaaz5120, <https://doi.org/10.1126/sciadv.aaz5120>, publisher: American Association for the Advancement of Science, 2020.
- Zheng, B., Cohen, J. B., Lu, L., Hu, W., Tiwari, P., Lolli, S., Garzelli, A., Su, H., and Qin, K.: How Can We Trust TROPOMI Based Methane Emissions Estimation: Calculating Emissions over Unidentified Source Regions, *Atmospheric Chemistry and Physics*, 26, 1931–1946, <https://doi.org/10.5194/acp-26-1931-2026>, 2026.