



Interannual variability of summertime cross-isobath exchanges in the northern South China Sea: ENSO and riverine influences

5 **Yunping Song¹, Yuxin Lin¹, Peng Zhan¹, Zhiqiang Liu^{1,2}, Zhongya Cai^{3,4}**

¹Department of Ocean Science and Engineering, Southern University of Science and Technology, Shenzhen, China

²Center for Complex Flows and Soft Matter Research, Southern University of Science and Technology, Shenzhen, China

10 ³State Key Laboratory of Internet of Things for Smart City, Department of Ocean Science and Technology, University of Macau, Macau, China

⁴Center for Ocean Research in Hong Kong and Macau, Hong Kong, China

Corresponding to: Zhiqiang Liu (liuzq@sustech.edu.cn); Zhongya Cai (zyc@um.edu.mo)

15



Abstract. This study investigates the interannual variability of summer shelf circulation in the Northern South China Sea (NSCS) from 2000 to 2022, combining long-term observations and high-resolution simulations. We elucidate the responses of NSCS shelf circulation to ENSO and Pearl River Estuary (PRE) freshwater runoff, revealing distinct spatial and mechanistic signatures. During El Niño years, a pronounced sea level anomaly dipole forms between the central and southern South China Sea, intensifying northward geostrophic currents in the southern basin and modulating Kuroshio intrusion. Simultaneously, an amplified PRE plume extends eastward to the 100 m isobath, markedly reducing nearshore salinity. Analysis of depth-integrated vorticity equations indicates that the pressure gradient force—driven by the joint effect of baroclinicity and bottom relief (JEBAR) and bottom pressure gradients—governs NSCS circulation variability. In coastal regions, cross-isobath velocity anomalies are primarily controlled by bottom stress curl and nonlinear vorticity advection, whereas JEBAR dominates offshore dynamics beyond the 100 m isobath. During El Niño summers, bottom density anomalies generate positive cross-isobath velocity anomalies through JEBAR, partially offset by negative anomalies from altered vertical stratification, sustaining a meandering shelf current. These results highlight the interplay of regional and remote forcings, advancing understanding of NSCS hydrographic dynamics.



1 Introduction

35 The Northern South China Sea (NSCS), particularly the shelf region offshore of the Pearl River Estuary (PRE), exhibits a complex circulation system shaped by monsoonal forcing, Kuroshio intrusion, and intricate topography (Fang et al., 1998; Xue et al., 2004; Nan et al., 2015). During summer, prevailing southwesterly monsoon winds drive intense coastal upwelling, significantly modulating the hydrography of the NSCS shelf. In addition to wind-driven Ekman transport, cross-isobath exchanges
40 are strongly influenced by topographic effects, as the NSCS shelf is characterized by nearly shore-parallel isobaths (Liu et al., 2020).

The Luzon Strait, to the east of the NSCS shelf and between Taiwan Island and Philippine (Figure 1), serves as the primary conduit for water exchange between the NSCS and the Pacific Ocean (Zhu et al., 2019). The joint effect of baroclinicity and bottom relief (JEBAR) plays an important role
45 comparable to wind stress in steering shelf and slope circulation, significantly influencing Kuroshio path variability near the Taiwan Strait (Oey et al., 2010, 2014). Recent studies also highlight JEBAR's critical role in cross-shelf transport across diverse shelf seas. For instance, Lin & Gan (2024) demonstrated that in the NSCS, the depth-averaged along-isobath pressure gradient is primarily driven by JEBAR, where baroclinicity dominates over topographic gradients. Tidal amplification of JEBAR
50 further enhances the westward pressure gradient force from the outer to inner shelves east of Hainan. Similarly, Renkl et al. (2025) identified JEBAR as the dominant factor in mean dynamic topography differences on the Scotian Shelf, Canada, where it balances offshore wind-driven Ekman transport with onshore flow, sustaining coastal upwelling. Closer to the PRE, the shelf circulation takes the form of a coastal current, with the freshwater plume displaying pronounced seasonal variability. During summer,
55 this plume extends eastward under the influence of southwesterly winds, modulating stratification and nearshore dynamics (Chen et al., 2001).

Previous studies of NSCS shelf circulation have primarily examined seasonal patterns and their wind-driven dynamics (Hu, 2000). Short-term summer variability in the shelf current has also been explored: Geng et al. (2024) demonstrated that tides modulate sea surface height, influencing shelf
60 pressure gradients, while Liu et al. (2020) showed that cross-isobath exchanges during upwelling and downwelling winds are sensitive to along-shelf pressure gradients shaped by complex bathymetry. The intensity of this upwelling exhibits notable interannual variability, often linked to the El Niño–Southern



Oscillation (ENSO) (Shu et al., 2018). For example, during the summer of 1998—an El Niño year—enhanced alongshore wind stress substantially intensified coastal upwelling (Jing et al., 2011).

65 Despite these advances, the interannual variability of summer shelf circulation remains poorly constrained. While expanded observational datasets, satellite products, and high-resolution modeling (Hong & Wang, 2008; Shu et al., 2011; Zu et al., 2020) have refined our understanding of NSCS dynamics, the role of interannual forcing—particularly the combined effects of ENSO, PRE runoff, and regional current meandering—has yet to be systematically quantified.

70 This study investigates the interannual variability of NSCS shelf circulation during summer from 2000 to 2022, using long-term observations and numerical modeling. We examine how ENSO modulates regional atmospheric and oceanic forcing, how the PRE runoff controls freshwater plume behavior, and how vorticity-related processes—including JEBAR, bottom stress curl, and nonlinear vorticity advection—govern cross-isobath exchanges and meandering shelf currents. By integrating
75 these processes, our study provides new insight into how regional and remote forcings interact to shape NSCS shelf circulation, with implications for hydrographic structure and ecosystem variability.

2 Methods and observations

To investigate the interannual variability of summer shelf circulation in the NSCS, we employed a high-resolution regional ocean model based on the Regional Ocean Modeling System (ROMS;
80 Shchepetkin & McWilliams, 2005). The model domain spans 105°–121° E and 16°–26° N, covering the NSCS shelf and adjacent deep waters. The model uses a curvilinear orthogonal Arakawa-C grid, with horizontal resolution ranging from 1 to 3 km across 800×400 grid points, enabling fine-scale representation of coastal and shelf processes. Vertically, the water column is discretized into 60 terrain-following *s*-coordinate layers, with enhanced resolution near the surface and bottom to better
85 resolve boundary layer dynamics (Song & Haidvogel, 1994). Turbulent mixing is parameterized using the Mellor–Yamada 2.5 turbulence closure scheme, widely validated for coastal applications (Mellor & Yamada, 1982). Horizontal and vertical momentum advection are calculated using a third-order upstream and a fourth-order centered scheme, respectively, minimizing numerical diffusion while preserving accuracy. Transport of temperature and salinity employs the multidimensional
90 positive-definite advection transport algorithm (MPDATA; Smolarkiewicz, 1984).



Open boundary conditions incorporate both tidal and subtidal forcing (Liu and Gan, 2020). Tidal elevation and velocity fields are constructed from nine principal constituents—semidiurnal (M_2 , S_2 , K_2 , N_2), diurnal (K_1 , O_1 , P_1 , Q_1), and the nonlinear M_4 component arising from M_2 interactions over the shallow PRE shelf (Mao et al., 2004). Harmonic constants are derived from satellite-based sea level anomaly (SLA) data processed by the Oregon Tide Inverse Software (Egbert & Erofeeva, 2002), following the approach of Zu et al. (2008). Subtidal elevation, currents, and hydrographic properties are obtained from the global Hybrid Coordinate Ocean Model (HYCOM), providing consistent large-scale boundary conditions. River runoff data for the PRE are obtained from the Chinese Ministry of Water Resources, capturing both seasonal and interannual fluctuations. Atmospheric forcing fields—including wind stress, heat fluxes, and surface pressure—are sourced from the fifth-generation ECMWF Reanalysis (ERA5), which offers high temporal and spatial resolution suitable for analyzing ENSO-related variability.




Figure 1. Overview map of the northern South China Sea. Shelf currents (colored arrows) and geostrophic currents (pink arrows) represent summer averages from 2000 to 2022. The upper panel shows the corresponding climatological wind field. Red markers indicate the observational sites used for model validation in Fig. 2. The upper left sub-graph shows the location of the northern South China Sea.



110 The model simulation spans the period 1994–2022, with analysis focused on summer months
(June–August) from 2000 to 2022. The initial six-year period (1994–1999) is excluded to allow for
model spin-up and dynamic equilibration. Model performance has been rigorously validated in
previous NSCS studies, demonstrating skill in reproducing sea level, currents, temperature, and salinity
(Liu et al., 2020; Deng et al., 2022; Song et al., 2024). Summer SST anomalies (2000–2022) at
115 monitoring stations (Fig. 1) show strong agreement with observations ($CC=0.74$, $RMSE=0.65^\circ\text{C}$; Fig.
2a). Notably, SSS anomalies achieve higher accuracy ($CC=0.83$, $RMSE=1.25\text{ psu}$; Fig. 2b),
particularly in capturing the PRE plume dynamics at interannual scales. However higher observed
variability may reflect unresolved sub-grid processes (e.g., localized rainfall or small-scale eddies),
suggesting that the smaller scaled variabilities in salinity could be further detailed. The model also
120 skillfully reproduces SLA variability from the EU Copernicus Marine Service (CMEMS) seaward of
the 200-m isobath ($CC=0.72$, $RMSE=2.27\text{ cm}$; Fig. 2c), effectively resolving ENSO-related and
regionally forced variabilities. These validation metrics demonstrate that the model reliably captures
key physical processes governing shelf circulation, cross-isobath transport, and vorticity dynamics in
the NSCS, providing a robust foundation for the analysis presented in this study.

125




Figure 2. Temporal variations of observed and simulated anomalies during summer (2000–2022): (a) sea surface




temperature (SST, °C) at site SM20; (b) sea surface salinity (SSS, psu) at site MM4; (c) sea level anomaly (SLA; cm) averaged seaward of the 200 m isobath within the model domain.

130

3 Results

3.1 Interannual variability of dynamic factors influencing NSCS shelf circulation in summer

To assess the interannual variability of summer shelf circulation and associated hydrographic properties in the NSCS, we analyzed a suite of dynamic factors encompassing atmospheric and oceanic processes, including evaporation minus precipitation (E-P), air temperature, wind stress, SST, SLA, and SSS. To extract coherent spatiotemporal patterns, we applied multivariate empirical orthogonal function (MVEOF) analysis using the eofs Python library (Dawson, 2016), which computes EOFs through eigendecomposition of the joint covariance matrix of the input variables. The covariance matrix was computed from the anomaly fields (time-mean removed) and normalized by $N-1$, where N is the number of temporal samples. This normalization provides an unbiased estimate of the population covariance. This approach identifies dominant modes of variability and their linkages to large-scale climate drivers, particularly ENSO as a key modulator of regional interannual variability.





145 **Figure 3.** Leading mode (Mode 1) of summer-averaged climatic and oceanic variables from multivariate EOF (MVEOF): (a) evaporation minus precipitation ($\text{kg m}^{-2} \text{s}^{-1}$); (b) air temperature at 2 m height ($^{\circ}\text{C}$); (c) sea surface temperature ($^{\circ}\text{C}$); (d) sea surface wind stress curl (N m^{-3}) and wind speed (m s^{-1}); and (e) sea level anomaly (SLA; cm) with geostrophic currents (cm s^{-1}), all representing the spatial pattern of MVEOF Mode 1; (f) sea surface salinity (SSS; psu) spatial mode from univariate EOF; (g) principal component time series of MVEOF Mode 1 (MVPC1, black line) with Niño 3.4 index (colored bars), and (h) principal component time series of SSS EOF Mode 1 (PC1, black line) with superimposed PRE summer runoff. Anomalies are relative to summer climatology, and vectors are scaled for clarity.

The leading mode (Mode 1) and its corresponding multivariate principal component (MVPC1) are shown in Fig. 3, highlighting spatial patterns and their temporal evolution. Mode 1 explains 60.60% of the total variance, with MVPC1 exhibiting a correlation coefficient of 0.62 with the Niño 3.4 index. The spatial structures associated with Mode 1 reveal contrasting responses during years with positive and negative MVPC1 phases, which, in general, correspond to El Niño and La Niña years, respectively. In positive MVPC1 years, a weakened Walker circulation displaces atmospheric pressure centers and trade-wind zones southward, while negative phases strengthen the circulation and shift them northward. This pattern during the positive MVPC1 years establishes positive wind stress curl in the South China Sea (SCS) basin, and a high-pressure anomaly over the southern SCS and generates a prominent SLA dipole across the SCS, featuring positive SLA anomalies south of 10°N and negative anomalies north of 10°N (Fig. 3e). Simultaneously, under the control of the anomalous atmospheric high-pressure (low-pressure), the clear (cloudy) sky during positive (negative) MVPC1 phases enhances (suppresses) evaporation and raises (lowers) air temperatures in the southern SCS. On the contrary, the air temperature and SST decrease (increase) in the mid-basin of the SCS during positive (negative) MVPC1 years.

Over the NSCS shelf, E-P (Fig. 3a) and air temperature (Fig. 3b) tend to decrease (increase) during positive (negative) MVPC1 years, leading to cooler (warmer) SST seaward of the 200 m isobath (Fig. 3c). In contrast, regions landward of this isobath experience the opposite trend—warmer (cooler) SST in positive (negative) phases—due to substantial freshwater input from the PRE and intensified rainfall over the Southern Chinese mainland. During El Niño summers (positive MVPC1 phases), northeasterly wind stress anomalies (Fig. 3d) develop over the NSCS, acting against the prevailing southwesterly monsoon and thereby suppressing coastal upwelling and contributing to SST increases nearshore. At the same time, the Kuroshio intrusion through the Luzon Strait intensifies to strengthen the intrusion of lower SST waters during the positive phase (Fig. 3e). The strengthened geostrophic currents in the



southern SCS (Wang et al., 2020) are linked to anticyclonic (cyclonic) wind stress curl anomalies in the central and western SCS basin and offshore Vietnam (Fig. 3e).

180 In contrast, interannual SSS variability in the NSCS is primarily governed by PRE runoff rather than directly linked to ENSO (Fig. 3f and h). A separated EOF analysis of SSS shows that Mode 1 accounts for 60.55% of the total variance, with PC1 closely tracking observed PRE runoff (CC = 0.83; Fig. 3h). PC1 of SSS EOF analysis (Fig. 3h) corresponds to PRE runoff variability, with positive phases reflecting large-runoff years (Table 2) and negative phases low-runoff years. During
185 large-runoff summers, SSS decreases markedly as the freshwater plume extends eastward to approximately the 100 m isobath near 115° E (Fig. 3f). This indicates that regions inside the 100 m isobath are more sensitive to riverine input, as the plume readily spreads across the shelf. These results emphasize the dominant role of riverine freshwater in modulating nearshore salinity and highlight the need to consider local riverine forcing alongside broader climatic influences.

190 The strong correlations between MVPC1 and the Niño 3.4 index (Fig. 3g), and between PC1 and PRE runoff (Fig. 3h), motivated analyses of shelf currents anomalies for the years listed in Tables 1 and 2, which respectively represent the conditions in the positive and negative MVPC1 and PC1 years. However, the relationship between summer PRE runoff and ENSO remains controversial. For instance, Gu et al. (2017) and Ouyang et al. (2014) reported increased precipitation and streamflow in the Pearl
195 River basin during La Niña compared to El Niño periods, whereas Wang et al. (2018) and Xu et al. (2007) noted higher precipitation in South China during El Niño and lower precipitation during La Niña. To isolate the riverine plume's influence on shelf currents, we selected large-runoff years from El Niño periods and low-runoff years from La Niña periods, denoted by bold labels in Table 1 and summarized in Table 2, to disentangle runoff effects. Notably, shelf circulation in El Niño/La Niña
200 years with low/large runoff (non-bold years in Table 1) typically shows an opposite pattern to large-runoff El Niño years (bold years in Table 1), and these cases were excluded from subsequent analyses.

Table 1. The El Niño (positive MVPC1) and La Niña (negative MVPC1) years

	El Niño	La Niña
Years	2000, 2001 , 2002 , 2004, 2005, 2006 , 2009 , 2011, 2012 , 2014 , 2015, 2018, 2019	2003 , 2007 , 2008, 2010 , 2013 , 2016, 2017, 2020 , 2021 , 2022

205



Table 2. The large runoff (positive PC1) and low runoff (negative PC1) years

	Large runoff	Low runoff
Years	2001, 2002, 2006, 2009, 2012, 2014, 2019	2003, 2007, 2010, 2013, 2020, 2021

3.2 Climatic characteristics of NSCS shelf circulation in summer

This section investigates the climatological characteristics (2000 – 2022) of summer NSCS shelf circulation, where the mean circulation is dominated by southwesterly monsoon winds (Fig. 1). These winds drive prominent upwelling currents in coastal waters shallower than 100 m (Fig. 4a). Conversely, offshore regions beyond the shelf break exhibit southwestward transport, primarily due to Kuroshio intrusion through the Luzon Strait, consistent with prior studies (e.g., Li et al., 2018). Interannual wind anomalies further modulate this circulation: during positive MVPC1 years (El Niño conditions; see Sect. 3.1), northeastward wind stress anomalies (Fig. 3e), reflecting ENSO's climatic influence on NSCS circulation.

To systematically characterize interannual anomalies in shelf currents and hydrographic properties, we adopted a two-stage regression approach in the following contents. First, we extracted anomalies referring to the climatological conditions in shelf currents and hydrographic properties during the analyzed years, then performed linear least-squares regression of these anomalies against their corresponding MVPC1/PC1 indices (Fig. 3g, 3h). This quantified each variable's sensitivity to the principal components (MVPC1/PC1) through slope coefficients (units of anomaly per PC unit). Second, we averaged the regressed anomalies across positive MVPC1/PC phase years (Tables 1/2) to emphasize their persistent spatial patterns. This two-stage regression approach combines the quantitative rigor of regression (isolating MVPC1/PC1-related variability) with the signal enhancement of composite analysis (highlighting recurrent features), while controlling for noise associated with short-term distortions.

To explore the cross-isobath water exchanges associated with the upwelling circulation, we decomposed the depth-averaged velocity field into along-isobath (positive values indicate flow with shallower waters on the right) and cross-isobath (positive values indicate flow



toward shallower waters) components. The summer-averaged cross-isobath velocity indicates net onshore transport across most of the NSCS shelf (Fig. 4b), except where meandering isobaths cause topographic steering to locally alter flow direction. During El Niño summers, interactions between northeastward wind stress anomalies and shelf topography generate a distinct southwestward meandering current anomaly across the central NSCS shelf (Figs. 4c, d). This pattern highlights the nonlinear interplay between wind forcing and bathymetric gradients in driving interannual shelf circulation variability.

240




Figure 4. Depth-averaged circulation and associated anomalies in the northern South China Sea: (a) mean velocity vectors (cm s^{-1}) and (b) mean cross-isobath velocity (cm s^{-1}) averaged over summers from 2000 to 2022, where positive cross-isobath velocity indicates flow from deeper to shallower waters; (c) regression map of velocity vector anomalies (cm s^{-1}), and (d) regression map of cross-isobath velocity anomalies (cm s^{-1}) during positive MVPC1 years (Table 1). Shaded areas in (c) and (d) denote regions where the 90% confidence level is not met, based on a two-tailed t-test using the estimated standard deviation and sample degrees of freedom. The two-stage regression approach is detailed in the section 3.2. Positive values in (c) and (d) indicate flow toward shallower waters.

250

To further assess water mass characteristics under varying climatic conditions, we conducted a hydrographic analysis along three cross-shelf transects (marked in pink in Fig. 4b),



with corresponding temperature and salinity sections shown in Fig. 5 during El Niño or large
runoff years. During positive MVPC1 years—coinciding with weakened monsoonal
255 forcing—all three transects exhibit elevated temperatures across the shelf, while colder,
Kuroshio-originated waters occupy the continental slope (Fig. 5a–c). Along Transect A, east
of Hainan Island, anticyclonic wind stress anomalies enhance the surface expansion of the
warm PRE plume, reinforcing the coastal current. In contrast, Transect C, located further east
and exposed to cyclonic anomalies, displays suppressed vertical mixing and elevated surface
260 temperatures, indicative of a regime that is unfavorable for upwelling circulations.

The influence of buoyant freshwater input becomes especially pronounced during years
with large PRE runoff. Under prevailing southwesterly winds, fresher water spreads widely
across the shelf, most prominently at Transect C (Fig. 5d–f). Concurrently, saltier Kuroshio
waters are observed over the slope, particularly at Transect A, contributing to enhanced
265 stratification. This establishes a stratified system driven by deeper saline inflows and
wind-induced divergence. In addition, the east-west contrast of this system manifests that the
water exchanges in the water column are prohibited over the eastern shelf due to surface
freshening, while upwelling-favorable conditions persist over the western shelf.




Figure 5. Regression maps of hydrographic anomalies along three cross-shelf transects (locations shown in Fig. 4b): (a–c) temperature anomaly profiles (°C) during positive MVPC1 years (Table 1) at Transects A, B, and C, respectively; (d–f) salinity anomaly profiles (psu) during positive PC1 years (Table 2) at the same transects. The two-stage regression approach is detailed in the section 3.2.



275 4 Mechanisms altering the interannual variability of cross-isobath transport

To investigate the mechanism altering the interannual variability of cross-isobath transport, we adopt the depth-averaged vorticity equation under conditions of low Rossby and Ekman numbers, appropriate for the interannual timescale considered, as outlined in Liu et al. (2020) as:

$$280 \quad PGF_{y*} = JEBAR + PGF_{y*}^b \quad (1)$$

$$PGF_{y*}^b = \frac{1}{H_{x*}} \overline{\nabla} \times \left(\frac{\tau_b}{\rho_0} \right) + \frac{1}{H_{x*}} \overline{\nabla} \times \left(-\frac{\tau_s}{\rho_0} \right) + \frac{1}{H_{x*}} J(\psi, \xi) + \left(\frac{\|\overline{\mathbf{v}}\|^2}{2} \right)_{y*} \quad (2)$$

In these equations, subscript (x^*) and (y^*) symbolize the cross-isobath and along-isobath directions, J is the Jacobi operator ψ is the transport streamfunction, and $\overline{\mathbf{v}}(\overline{u}, \overline{v})$ denotes the depth-averaged velocity, respectively. According to Mertz & Wright (1992), the pressure gradient force (PGF_{y*}) in the water column can be decomposed into two terms: JEBAR and bottom-layer pressure gradient force in the water column (PGF_{y*}^b). The PGF_{y*}^b can be further decomposed into bottom stress curl (BSC), surface stress curl (SSC), the nonlinear advection of relative vorticity (RVA), and the gradient of momentum flux (GMF) terms (Liu and Gan, 2015). All the composite terms above are normalized by the Coriolis parameter to further assess their contribution to the cross-isobath velocity anomaly, and the results are presented in Fig. 6. The relatively negligible contribution from the GMF is excluded from the following discussion.

The cross-isobath velocity anomaly associated with JEBAR (Fig. 6a) predominantly governs transport from the vicinity of the 100 m isobath to the SCS basin, whereas the influence of PGF_{y*}^b (Fig. 6b) is largely concentrated over the shelf within the 100 m isobath, as also shown in Liu et al. (2020). More specifically, the spatial distribution of water exchange induced by the nonlinear advection of relative vorticity (RVA; Fig. 6c) forms the principal structure of the PGF_{y*}^b contribution across the shelf, particularly between the 30 m and 100 m isobaths. For instance, over the shelf east of Hainan Island, the strong resemblance between Fig. 6b and 6c highlights RVA as the dominant contributor in this region. The BSC term (Fig. 6d) is notably influenced by complex bathymetry, such as near the Taiwan Shoal, where it generates offshore flow near the 50 m isobath and pronounced onshore transport within the 30 m isobath during positive MVPC1 years. Overall, the meandering cross-isobath velocity anomaly (Fig. 4d) results from the combined influence of JEBAR, which controls offshore



exchange beyond the 100 m isobath, and PGF_{y*}^b , which plays a central role in shaping hydrodynamics of the coastal seas.

305




Figure 6. Regression maps of horizontal cross-isobath velocity anomalies (cm s^{-1}) during positive MVPC1 years (Table 1), attributed to different dynamic components: (a) joint effect of baroclinicity and bottom relief (JEBAR); (b) bottom pressure gradient force (PGF_{y*}^b); (c) nonlinear relative vorticity advection (RVA); and (d) bottom stress curl (BSC). The two-stage regression approach is detailed in the section 3.2. Positive values in the figures indicate flow toward shallower waters.

310

Given that the 100 m isobath marks a critical boundary separating shelf dynamics from those of the deeper offshore region, the circulation patterns along this isobath are important to reveal the underlying mechanisms of cross-isobath exchange. During positive MVPC1 years, the regressed cross-isobath velocity anomaly over the 100 m isobath (Fig. 7a) reveals a vertically coherent offshore transport from the surface to the bottom, with notable intensification near the widened shelf around 115°E and over the shelf east of 118°E and west of 113°E . Despite the overall coherence of the velocity structure, the hydrographic properties associated with these currents show distinct characteristics. East of 115°E over the 100 m isobath, the flow is predominantly accompanied by warmer, buoyant freshwater discharged from the PRE (Fig. 7b), while west of 115°E , colder

320



Kuroshio-intruded water dominates the offshore transport, particularly during El Niño summers, as also previously shown in Fig. 3. From a salinity perspective, the PRE plume exerts a significantly stronger influence to the east of 115° E, especially in the surface and middle layers (Fig. 7c). This spatial contrast persists even during years of elevated freshwater runoffs, particularly under positive PC1 conditions. These findings highlight the essential role of the PRE plume in modulating the salinity structure of the NSCS, with its influence being most pronounced on the eastern shelf where it interacts with offshore currents and contributes to broader shelf-basin exchange processes through modulating, for example, the JEBAR.




Figure 7. Regression maps of hydrographic anomalies along the 100 m isobath: (a) cross-isobath velocity anomaly (cm s^{-1}) and (b) temperature anomaly profiles ($^{\circ}\text{C}$) during positive MVPC1 years (Table 1); (c) salinity anomaly profile (psu) during positive PC1 years (Table 2). The two-stage regression approach is detailed in the section 3.2.

Based on the previously identified interannual characteristics of hydrographic properties and drawing from Mertz & Wright (1992), we extend the analysis of the JEBAR in Equation 3:

$$\text{JEBAR} = J \left(\frac{g}{\rho_0} \int_{-H}^0 z \rho dz, H^{-1} \right) \quad (3)$$

Equation 3 can be explicitly rewritten in a Cartesian coordinate:

$$\text{JEBAR} = \left(\frac{g}{\rho_0} \int_{-H}^0 z \rho dz \right)_x \left(\frac{1}{H} \right)_y - \left(\frac{1}{H} \right)_x \left(\frac{g}{\rho_0} \int_{-H}^0 z \rho dz \right)_y \quad (4)$$

Given that the water depth H generally remains constant—since variations in the SLA are much



smaller than the topographic depth—and that the slope current streams in the NSCS show minimal
345 spatial displacements between positive and negative MVPC1 and PC1 years, we focus our analysis on
further understanding the zonal and meridional gradients of the baroclinic component: $\frac{g}{\rho_0} \int_{-H}^0 z \rho dz$,
expressed by:

$$\frac{g}{\rho_0} \int_{-H}^0 z \rho dz = -\frac{g}{\rho_0} \int_{-H}^0 \frac{z^2}{2} \frac{\partial \rho}{\partial z} dz - \frac{g}{\rho_0} \frac{H^2}{2} \rho_b \quad (5)$$

The first term on the right-hand side of Equation 5 quantifies the strength of density stratification
350 in the water column over the shelf, while the second term represents the density anomaly of the bottom
seawater. Our primary focus is on the interannual variability of the cross-isobath velocities, which are
shown in Fig. 5. To further investigate the anomaly of the baroclinic component, we compute it as
follows:

$$\frac{g}{\rho_0} \int_{-H}^0 z(\rho - \bar{\rho}) dz = \overbrace{-\frac{g}{\rho_0} \int_{-H}^0 \frac{z^2}{2} \frac{\partial(\rho - \bar{\rho})}{\partial z} dz}^{N'} + \overbrace{-\frac{g}{\rho_0} \frac{H^2}{2} (\rho - \bar{\rho})_b}^{P'} \quad (6)$$

355 The spatial distribution pattern of $\left(\frac{g}{\rho_0} \int_{-H}^0 z \rho dz\right)'$ aligns closely with the meandering pathways of
the shelf current along the 100 m isobath, as shown in Fig. 4d, particularly during positive PC1 years
under both strong (Fig. 8a) and low runoff conditions (Fig. 8d). The horizontal distributions of each
term in Equation 6 during positive MVPC1 years are displayed in Fig. 8, with corresponding patterns
for negative MVPC1 years shown in Fig. A1. It is evident that the P' (Fig. 8c and f) plays a dominant
360 role in shaping the pattern of $\left(\frac{g}{\rho_0} \int_{-H}^0 z \rho dz\right)'$. However, the negative values near 114°E, primarily
sourced from N' (Fig. 8b and e), are critical for cross-shelf water transport. Notably, the strength of the
term $\left(\frac{g}{\rho_0} \int_{-H}^0 z \rho dz\right)'$ is strong or weak is highly correlated with P' , the key determinant of which is the
bottom density anomaly. In contrast, the variation in N' , which is primarily influenced by the intensity
of density stratification, appears less pronounced during both positive and negative PC1 years,
365 especially near the 100 m isobath. However, in negative MVPC1 years, this term exhibits more
significant differences, as shown in Fig. A1a and A1c. Consequently, the negative values near the 100
m isobath are more prominent when runoff from the Pearl River Estuary (PRE) is relatively low, as
seen in Fig. A1.




Figure 8. Regression maps of horizontal components in the JEBAR term during summer and positive MVPC1 years, under different runoff conditions: (a–c) correspond to positive PC1 years, and (d–f) to negative PC1 years.

Panels show (a, d) the full baroclinic gradient term ($\frac{g}{\rho_0} \int_{-H}^0 z(\rho - \bar{\rho}) dz$), (b, e) the contribution from vertical

density stratification ($-\frac{g}{\rho_0} \int_{-H}^0 \frac{z^2}{2} \frac{\partial(\rho - \bar{\rho})}{\partial z} dz$), and (c, f) the contribution from bottom density anomaly ($-\frac{g}{\rho_0} \frac{H^2}{2} (\rho -$

$\bar{\rho})_b$). The MVPC1 time series is shown in Figure 3g, with corresponding positive-phase years listed in Table 1, while the PC1 time series is shown in Figure 3h with its positive-phase years listed in Table 2. The two-stage regression approach is detailed in the section 3.2.

Having identified the determining factor responsible for the interannual variability of cross-isobath water transport over the 100 m isobath, we now focus on examining the density and buoyancy frequency anomaly profiles during positive MVPC1 years to elucidate the underlying formation mechanisms. The scenarios during negative MVPC1 years are presented in Fig. A2. Longitudinal profiles at Transect B for different conditions are shown in Fig. A3. These profiles reveal that the density anomaly patterns west of 115° E (the widened shelf) during both positive and negative PC1 years exhibit a similar structure (Fig. 9a and c). A prominent feature is the exceptionally high core of density anomaly between 112° and 113° E, strongly influenced by the southward flow of Kuroshio water. In contrast, the eastern side of the widened shelf presents a markedly different situation. Due to the substantial influx of buoyant water from the PRE, surface density decreases during positive PC1 years (Fig. 9a), while bottom density remains largely unchanged (as shown in Fig. A2a and A2b). In essence, the term P' over the 100 m isobath shows little variation regardless of whether runoff is high or low, supporting the distribution patterns observed in Fig. 8c and 8f.




Figure 9. Regression maps of hydrographic structure anomalies along the 100 m isobath during summer and MVPC1 years (Table 1), under varying runoff conditions: (a, b) correspond to positive PC1 years, and (c, d) to negative PC1 years (Table 2). Panels show (a, c) density anomaly profiles (kg m^{-3}) and (b, d) buoyancy frequency squared anomaly profiles (s^{-2}). The two-stage regression approach is detailed in the section 3.2.

The vertical profile of the buoyancy frequency anomaly during both positive and negative PC1 years further reveals distinctions between the western and eastern parts of the shelf. During years with large runoff, the density stratification in the surface and middle layers is enhanced across almost the entire 100 m isobath (Fig. 9b). In contrast, during years with low runoff, stratification is only strengthened on the western side of the widened shelf (Fig. 9d), and its intensity is lower compared to positive PC1 years. Furthermore, surface density stratification during low runoff years is noticeably weaker than in the mean state (as shown in Fig. A2c and A2d). In summary, during positive MVPC1 years, the vertical integration of the buoyancy frequency anomaly over the 100 m isobath is more pronounced in years of large runoff from the PRE than during low runoff periods. This explains the increased presence of negative values over the 100 m isobath in Fig. 8d compared to Fig. 8a, which in turn directly influences the interannual variability of the spatial distribution of JEBAR.

5 Conclusions

This study investigates the interannual variability of shelf circulation in the NSCS during the summer months from 2000 to 2022, utilizing long-term observational and numerical simulation data. The findings indicate that this variability is strongly influenced by ENSO-driven climate dynamics and



415 river runoff from the PRE. During El Niño years, the warm pool of the tropical Northwest Pacific shifts eastward, leading to a cooler SST and a pronounced westward Kuroshio intrusion through the Luzon Strait in the eastern SCS. This shift induces a cyclonic wind stress curl anomaly in the NSCS and an anticyclonic wind stress curl anomaly in the southern region, resulting in a distinct dipole pattern in the SLA anomaly across the central and southern SCS basin adjacent to the NSCS. Additionally, the northward geostrophic current is intensified in the southern SCS during El Niño years. When a warmer and more extensive Pearl River plume is generated, likely due to increased terrestrial precipitation, the buoyant freshwater exerts a stronger influence on the coastal waters east of the estuary, extending to the vicinity of the 100 m isobath.

The depth-integrated vorticity equations in the cross-isobath direction are adapted to describe the mechanisms shaping the circulation structure in the NSCS. Building on previous findings that the along-isobath pressure gradient force, composed of JEBAR and PGF_{y*}^b , represents the dominant driver of the NSCS shelf circulation, this study further clarifies that the baroclinic and river plume anomalies shape the spatial distribution of cross-isobath velocity anomalies in the coastal waters, while JEBAR primarily governs the region extending from the 100 m isobath to the open ocean. Through further analysis of JEBAR structure, it is found that the bottom density anomaly contributes to positive cross-isobath velocity anomalies, while the density stratification anomaly generates negative values during El Niño years, which collectively influence the characteristic meandering pattern of the southwestward shelf current.

Although this study categorizes years based on ENSO phases without considering phase-dependent dynamics (e.g., developing vs. decaying events), several important aspects remain for future research. First, the phase-locking effects of ENSO could alter seasonal circulation patterns. Second, inter-basin interactions—particularly with the Indian Ocean Dipole (IOD)—deserve further exploration. A positive IOD (characterized by a cool western and warm eastern Indian Ocean) frequently coincides with El Niño, potentially amplifying atmospheric teleconnections that modulate SCS wind patterns and Kuroshio intrusion. Conversely, La Niña conditions, which often occur in conjunction with a negative IOD, may enhance southwesterly monsoon winds, strengthening coastal upwelling and altering freshwater plume dispersal patterns.



445

Appendix A




Figure A1. Regression maps of horizontal components in the JEBAR term during summer and negative MVPC1 years, under different runoff conditions: (a, b) correspond to positive PC1 years, and (c, d) to negative PC1 years.

450 Panels show (a, c) the full baroclinic gradient term ($\frac{g}{\rho_0} \int_{-H}^0 z(\rho - \bar{\rho}) dz$) and (b, d) the contribution from vertical density stratification ($-\frac{g}{\rho_0} \int_{-H}^0 \frac{z^2}{2} \frac{\partial(\rho - \bar{\rho})}{\partial z} dz$). The MVPC1 time series is shown in Figure 3g, with corresponding positive-phase years listed in Table 1, while the PC1 time series is shown in Figure 3h with its positive-phase years listed in Table 2. The two-stage regression approach is detailed in the section 3.2.

455






Figure A2. Regression maps of hydrographic anomaly profiles along the 100 m isobath during summer and negative MVPC1 years (Table 1), under different runoff conditions: (a, c) correspond to positive PC1 years, and (b, d) to negative PC1 years (Table 2). Panels show (a, b) density anomalies (kg m^{-3}) and (c, d) buoyancy frequency squared anomalies (s^{-2}). The two-stage regression approach is detailed in the section 3.2.




Figure A3. Regression maps of hydrographic anomaly profiles at Transect B under different climate and runoff conditions. Panels (a–d) show density anomalies (kg m^{-3}), and panels (e–h) show buoyancy frequency squared anomalies (s^{-2}). Results correspond to: (a, b, e, f) positive MVPC1 years (Table 1) and (c, d, g, h) negative MVPC1 years; with (a, c, e, g) representing positive PC1 years and (b, d, f, h) negative PC1 years (Table 2). The two-stage regression approach is detailed in the section 3.2.

Data availability.

The timeseries of long - term hydrographic data observed in the surface and bottom layers of Hong Kong waters is described in Deng et al. (2022) and is publicly accessible at <https://cd.epic.epd.gov.hk/EPICRIVER/marine/?lang=en>. The atmospheric forcing data utilized in this study can be obtained from the ECMWF ERA5 dataset (Hersbach et al., 2023), available at <https://doi.org/10.24381/cds.fl7050d7>. The sea surface temperature data are described in Maturi et al. (2017) and openly accessible through NOAA (2024) at https://coralreefwatch.noaa.gov/product/5km/index_5km_sst.php. Gridded sea level anomaly (SLA) and geostrophic velocity data are openly accessible through the EU Copernicus Marine Service (CMEMS), and the relevant DOI link is <https://doi.org/10.48670/moi-00148>.

Author contributions.

SYP: Conceptualization, methodology, formal analysis, investigation, writing – original draft, visualization.



485 LYX: Conceptualization, methodology, resources, data curation, writing – review and editing.

ZP: Software, validation, data analysis, writing – review and editing.

LZQ: Supervision, funding acquisition, project administration, writing – review and editing.

CZY: Investigation, methodology, writing – review and editing, project administration.

All authors have read and approved the final manuscript.

490

Competing interests.

The contact authors have declared that none of the authors has any competing interests.

Acknowledgement.

495 This project was supported by National Natural Science Foundation of China (42276004, 42376024, 42450181), Science and Technology Development Fund, Macau SAR (File/Project no. 001/2024/SKL, 0040/2023/R1A1), Science, Technology and Innovation Commission of Shenzhen Municipality (JCYJ20240813094213018). The work described in this paper was substantially supported by a grant from the Research Grants Council of the Hong Kong Special Administrative Region, China 500 (AoE/P-601/23-N). CORE is a joint research centre for ocean research between Laoshan Laboratory and HKUST. This work was performed in part at the SICCC, which is supported by the SKL-IOTSC, University of Macau.



505 **References**

- Chen, C. T. A., Wang, S. L., Wang, B. J., and Pai, S. C.: Nutrient budgets for the South China Sea basin, *Marine Chemistry*, 75, 281–300, doi:10.1016/s0304-4203(01)00041-x, 2001.
- Chen, G. and Tam, C.-Y.: Different impacts of two kinds of Pacific Ocean warming on tropical cyclone frequency over the western North Pacific, *Geophysical Research Letters*, 37, doi:10.1029/2009gl041708, 2010.
- Dawson, A.: EOFs: A library for EOF analysis of meteorological, oceanographic, and climate data. *Journal of Open Research Software*, 4(1), 14. doi:10.5334/jors.122, 2016.
- Deng, Y., Liu, Z., Zu, T., Hu, J., Gan, J., Lin, Y., Li, Z., Quan, Q., and Cai, Z.: Climatic Controls on the Interannual Variability of Shelf Circulation in the Northern South China Sea, *Journal of Geophysical Research-Oceans*, 127, doi:10.1029/2022jc018419, 2022.
- Egbert, G. D. and Erofeeva, S. Y.: Efficient inverse Modeling of barotropic ocean tides, *Journal of Atmospheric and Oceanic Technology*, 19, 183 – 204, doi:10.1175/1520-0426(2002)019<0183:EIMOBO>2.0.CO;2, 2002.
- Fang, G., Fang, W., Fang, Y., and Wang, K.: A survey of studies on the South China Sea upper ocean circulation, *Acta Oceanogr Taiwan*, 1998.
- Geng, B., Shu, Y., Hu, J., Li, S., Liu, N., Xiu, P., and Wang, D.: Influences of tides on shelf circulation in the northeastern South China Sea during summer, *Journal of Marine Systems*, 243, doi:10.1016/j.jmarsys.2024.103964, 2024.
- Gu, X., Zhang, Q., Singh, V. P., and Shi, P.: Hydrological response to large-scale climate variability across the Pearl River basin, China: Spatiotemporal patterns and sensitivity, *Global and Planetary Change*, 149, 1–13, doi:10.1016/j.gloplacha.2016.12.016, 2017.
- Hong, B. and Wang, D.: Sensitivity Study of the Seasonal Mean Circulation in the Northern South China Sea, *Advances in Atmospheric Sciences*, 25, 824 – 840, doi:10.1007/s00376-008-0824-8, 2008.
- Jing, Z., Qi, Y., and Du, Y.: Upwelling in the continental shelf of northern South China Sea associated with 1997-1998 El Nino, *Journal of Geophysical Research-Oceans*, 116, doi:10.1029/2010jc006598, 2011.
- Lin, S. and Gan, J.: Dynamics of Tidal Effects on Coastal Upwelling Circulation Over Variable Shelves in the Northern South China Sea, *Journal of Geophysical Research-Oceans*, 129, doi:10.1029/2024jc021193, 2024.
- Li, D., Zhou, M., Zhang, Z., Zhong, Y., Zhu, Y., Yang, C., Xu, M., Xu, D., and Hu, Z.: Intrusions of Kuroshio and Shelf Waters on Northern Slope of South China Sea in Summer 2015, *Journal of Ocean University of China*, 17, 477–486, doi:10.1007/s11802-018-3384-2, 2018.
- Liu, Z. and Gan, J.: Upwelling induced by the frictional stress curl and vertical squeezing of the vortex tube over a submerged valley in the East China Sea, *Journal of Geophysical Research-Oceans*, 120, 2571–2587, doi:10.1002/2015jc010715, 2015.
- Liu, Z. and Gan, J.: A modeling study of estuarine-shelf circulation using a composite tidal and subtidal open boundary condition, *Ocean Modelling*, 147, doi:10.1016/j.ocemod.2019.101563, 2020.
- Liu, Z., Zu, T., and Gan, J.: Dynamics of cross-shelf water exchanges off Pearl River Estuary in



- 545 summer, *Progress in Oceanography*, 189, 10.1016/j.pocean.2020.102465, 2020.
- Mao, Q. W., Shi, P., Yin, K. D., Gan, J. P., and Qi, Y. Q.: Tides and tidal currents in the Pearl River estuary, *Continental Shelf Research*, 24, 1797–1808, doi:10.1016/j.csr.2004.06.008, 2004.
- Mellor, G. L. and Yamada, T.: Development of a turbulence closure model for geophysical fluid problems, *Reviews of Geophysics*, 20, 851–875, doi:10.1029/RG020i004p00851, 1982.
- 550 Mertz, G. and Wright, D. G.: Interpretations of the JEBAR Term, *Journal of Physical Oceanography*, 22, 301–305, doi:10.1175/1520-0485(1992)022<0301:Jotjt>2.0.Co;2, 1992.
- Nan, F., Xue, H., and Yu, F.: Kuroshio intrusion into the South China Sea: A review, *Progress in Oceanography*, 137, 314–333, doi:10.1016/j.pocean.2014.05.012, 2015.
- Oey, L.-Y., Hsin, Y.-C., and Wu, C.-R.: Why does the Kuroshio northeast of Taiwan shift shelfward in 555 winter?, *Ocean Dynamics*, 60, 413–426, doi:10.1007/s10236-009-0259-5, 2010.
- Oey, L.-Y., Chang, Y. L., Lin, Y. C., Chang, M. C., Varlamov, S., and Miyazawa, Y.: Cross Flows in the Taiwan Strait in Winter, *Journal of Physical Oceanography*, 44, 801–817, doi:10.1175/jpo-d-13-0128.1, 2014.
- Ouyang, R., Liu, W., Fu, G., Liu, C., Hu, L., and Wang, H.: Linkages between ENSO/PDO signals and 560 precipitation, streamflow in China during the last 100 years, *Hydrology and Earth System Sciences*, 18, 3651–3661, doi:10.5194/hess-18-3651-2014, 2014.
- Renkl, C., Oliver, E. C. J., and Thompson, K. R.: The alongshore tilt of mean dynamic topography and its implications for model validation and ocean monitoring, *Ocean Science*, 21, 181–198, doi:10.5194/os-21-181-2025, 2025.
- 565 Shchepetkin, A. F. and McWilliams, J. C.: The regional oceanic modeling system (ROMS): a split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean Modelling*, 9, 347–404, doi:10.1016/j.ocemod.2004.08.002, 2005.
- Shu, Y., Zhu, J., Wang, D., and Xiao, X.: Assimilating remote sensing and in situ observations into a coastal model of northern South China Sea using ensemble Kalman filter, *Continental Shelf 570 Research*, 31, S24–S36, doi:10.1016/j.csr.2011.01.017, 2011.
- Shu, Y., Wang, D., Feng, M., Geng, B., Chen, J., Yao, J., Xie, Q., and Liu, Q.-Y.: The Contribution of Local Wind and Ocean Circulation to the Interannual Variability in Coastal Upwelling Intensity in the Northern South China Sea, *Journal of Geophysical Research-Oceans*, 123, 6766–6778, doi:10.1029/2018jc014223, 2018.
- 575 Smolarkiewicz, P. K.: A fully multidimensional positive definite advection transport algorithm with small implicit diffusion, *Journal of Computational Physics*, 54, 325–362, doi:10.1016/0021-9991(84)90121-9, 1984.
- Song, Y. and Haidvogel, D.: A semi-implicit ocean circulation model using a generalized topography-following coordinate system, *Journal of Computational Physics*, 115, 228–244, 580 doi:10.1006/jcph.1994.1189, 1994.
- Song, Y., Lin, Y., Quan, Q., Zu, T., Cai, Z., Zhan, P., and Liu, Z.: ENSO-Modulated Variability in Winter Shelf Circulation of the Northern South China Sea, *Journal of Geophysical Research-Oceans*, 129, doi:10.1029/2024jc021577, 2024.
- Wang, L., Li, Q., Mao, X.-z., Bi, H., and Yin, P.: Interannual sea level variability in the Pearl River 585 Estuary and its response to El Niño-Southern Oscillation, *Global and Planetary Change*, 162, 163–174, doi:10.1016/j.gloplacha.2018.01.007, 2018.
- Wang, Q., Zeng, L., Shu, Y., Liu, Q., Zu, T., Li, J., Chen, J., He, Y., and Wang, D.: Interannual variability of South China Sea winter circulation: response to Luzon Strait transport and El Niño



- wind, *Climate Dynamics*, 54, 1145–1159, doi:10.1007/s00382-019-05050-2, 2020.
- 590 Xu, Z. X., Li, J. Y., Takeuchi, K., and Ishidaira, H.: Long-term trend of precipitation in China and its
association with the El Nino-southern oscillation, *Hydrological Processes*, 21, 61–71,
doi:10.1002/hyp.6180, 2007.
- Xue, H., Chai, F., Pettigrew, N., Xu, D., Shi, M., and Xu, J.: Kuroshio intrusion and the circulation in
the South China Sea, *Journal of Geophysical Research: Oceans*, 109, doi:10.1029/2002JC001724,
595 2004.
- Zhu, Y., Sun, J., Wang, Y., Li, S., Xu, T., Wei, Z., and Qu, T.: Overview of the multi-layer circulation in
the South China Sea, *Progress in Oceanography*, 175, 171–182, doi:10.1016/j.pocean.2019.04.001,
2019.
- Zu, T., Gan, H., and Erofeeva, S. Y.: Numerical study of the tide and tidal dynamics in the South China
600 Sea, *Deep-Sea Research Part I-Oceanographic Research Papers*, 55, 137–154,
doi:10.1016/j.dsr.2007.10.007, 2008.
- Zu, T., Wang, D., Wang, Q., Li, M., Wei, J., Geng, B., He, Y., and Chen, J.: A revisit of the interannual
variation of the South China Sea upper layer circulation in summer: correlation between the
eastward jet and northward branch, *Climate Dynamics*, 54, 457–471,
605 doi:10.1007/s00382-019-05007-5, 2020.