

Supporting Information for "Linear trends of temperature, salinity, and oxygen in the North Pacific based on GOBAI-O₂ over 2 decades and their controlling factors"

Authors:

Miho Ishizu^{1, 2}, Tomomichi Ogata³

¹Center for Climate Physics, Institute for Basic Science, Busan 46241, Republic of Korea

²Pusan National University, Tonghappiggyegwan Bldg 2 Busandaehak-ro, 63 beon-gil, Geumjeong-gu, Busan 46241, Republic of Korea

³Japan Agency for Marine-Earth Science and Technology, Environmental Variability Prediction and Application Research Group, Yokohama Institute for Earth Sciences, 3173-25 Showa-machi, Kanagawa-ku, Yokohama 236-0001, Japan

Contents of this file:

Figure S1. Uncertainty estimates of dissolved oxygen concentration ($\mu\text{mol/kg}$) along the isopycnal surface of 25.0 (a), 26.0 (b), and 26.8 σ_0 (c). The uncertainty data is based on GOBAI-O₂ data [Sharp et al. 2023].

Figure S2. Coefficient of determination for linear regression lines along the isopycnal surface of 25.0 (a), 26.0 (b), and 26.8 σ_0 (c).




Figure S1. Uncertainty estimates of dissolved oxygen concentration ($\mu\text{mol/kg}$) along the isopycnal surface of 25.0 (a), 26.0 (b), and $26.8\sigma_\theta$ (c). The uncertainty data is based on GOBAI-O₂ data [Sharp et al. 2023].




Figure S2. Coefficient of determination for linear regression lines along the isopycnal surface of 25.0 (a), 26.0 (b), and $26.8\sigma_\theta$ (c).