

# Long-term Effects of Drainage and Rewetting on the Degradation and Preservation of Peat Organic Matter in ~~Warm~~ Sub-Tropical Climate

Guy Sapir<sup>1,2,3</sup>, Alon Angert<sup>1</sup>, Yoav Oved Rosenberg<sup>2</sup>, Rotem Golan<sup>3</sup>.

5 <sup>1</sup> The Institute of Earth Sciences, The Hebrew University of Jerusalem, Israel. <sup>2</sup> Geological Survey of Israel, Jerusalem, Israel.

<sup>3</sup> Institute of Soil, Water and Environmental Sciences. Volcani Institute, Agricultural Research Organization, Israel.

Correspondence to: [Alon Angert \(angert@huji.ac.il\)](mailto:angert@huji.ac.il) [Rotem Golan \(rotemg@volcani.agri.gov.il\)](mailto:rotemg@volcani.agri.gov.il)

**Abstract.** Peatlands cover about 3% of the earth's land surface, while storing about 20% of the total global soil organic carbon. These carbon stocks are largely at risk, as many peatlands have deteriorated since the Industrial  
10 Revolution due to conversion to agricultural land by drainage. Globally, peatland drainage is responsible for over 3.5% of anthropogenic greenhouse gas emissions. About Approximately 75% of these emissions originate from warm climate regions. Mitigation of these emissions can be achieved by rewetting degraded peatlands. This study focuses on a warm sub-tropical climate peatland that has been cultivated for the past ~70 years (Hula Valley, Israel). The historic marsh was drained in 1957 for agricultural use and underwent a hydrological restoration project  
15 for elevating and stabilizing groundwater table since 1994. This land management history resulted in a sedimentary peat column that can be divided into three distinct sub-sections: drained, rewetted and pristine peat. This setting enables studying the drainage and rewetting effects on soil organic matter (SOM) degradation and preservation under warm climates. For this purpose, five sediment cores, 4 m long each, were excavated from cropland located over the historic marsh area. Locations were chosen to match previous studies on this site. Each soil profile was  
20 characterized using Rock-Eval® thermal analysis of the organic matter, and short-term soil aerobic respiration experiments. Integration of these results with historic SOM content data and with SOM modelling was used to explore the long-term process and rate of degradation. We found that the mean SOM content in the top one meter of the soil profile declined from 68 ±4 ~~wt.~~% to 21 ±2 ~~wt.~~% by weight over the past 66 years, excluding the compaction effect. In comparison to the drained section, the rewetted and pristine sub-sections has have a mean  
25 SOM of 33 ±2 ~~wt.~~% and 64 ±2 ~~wt.~~%, respectively. A peak in pyrite concentration beneath the recent water table-level, was observed in most profiles, indicating anaerobic conditions and sulfur recycling. Rock-Eval® thermal analysis demonstrated that during decomposition, the residual SOM became more oxidized and contained a lower proportion of thermally labile SOM, with a significant difference found between drained and rewetted peat. These results imply that the raising of the water table (~30 years ago) effectively helped preserving organic matter

30 compared to the drained section. Long-term SOM field data were integrated and studied using an SOM decomposition model and by incorporating respiration fluxes. The resulting trends highlighted that the first few decades of exposure are highly significant for the fate of the carbon stock, leading to substantial CO<sub>2</sub> emissions. These emissions were lower by 60–85% after 70 years. Furthermore, our results suggest that currently, approximately 13-21 wt.% of SOM persists as resistant organic matter in the degraded peat.

## 35 1 Introduction

Peatlands cover about 3% of the earth's land surface and estimated to store between 500-700 Pg of carbon (C), which are about 20% of the global total soil organic stock (Page et al., 2016). Peatland has been exploited intensively by humans since the Industrial Revolution. At present, human activity of draining and mining affects approximately 10% of the remaining global peatlands, transforming these ecosystems from long-term carbon sinks  
40 into significant sources of greenhouse gases (GHG's) (Leifeld & Menichetti, 2018). This transformation promotes carbon loss, primarily through microbial respiration and dissolved organic carbon (DOC) leaching (Hiraishi et al., 2013). Degraded peatlands currently contain 80.8 Pg of carbon and 2.3 Pg of nitrogen (Leifeld & Menichetti, 2018). Without further protective actions, drained peatlands will release most of this stock as greenhouse gases into the atmosphere.

45 The primary factor that contributes to the preservation of peat organic matter (OM) is anaerobic conditions, which predominantly prevail in water-saturated conditions (Moore ~~et al.~~, 1989). In such environments, where the SOM supply rate exceeds the decomposition rate, a net carbon sink can evolve (Moore, 1989). Generally, organic matter is added to the surface of the  ~~mire-marsh~~ (peat-forming habitat) mainly from net primary production, and its accumulation is countered by decomposition processes. Decomposition in the top layer occurs under aerobic  
50 conditions, followed by deeper burial in the saturated horizon where oxygen is limited (Moore, 1989; Joosten, 2002). There, under anaerobic conditions, decomposition occurs at rates that are approximately a thousand times lower than the upper aerobic layer (Clymo et al., 1998; Kleinen et al., 2012). ~~These~~ low decomposition rates result in a carbon sink with a turnover time of several ~~thousands of thousand~~ years if ~~it~~ remains undisturbed (Yu et al., 2010). Lowering the water table level (WTL) results in oxidation of the peat column, stimulating aerobic microbial  
55 decomposition of the SOM (Scanlon et al., 2000; Joosten, 2002), which releases intense CO<sub>2</sub> and N<sub>2</sub>O fluxes into the atmosphere (Page et al., 2016). Draining peatlands by lowering the WTL to convert them into agricultural lands has been a widespread practice for centuries and continues today. Numerous studies, e.g., Hooijer et al. (2010) and Moore ~~et al.~~ & Knowles (1989), have demonstrated a clear correlation between low WTL and high CO<sub>2</sub> emissions

due to peat soil respiration. Currently, more than 90% of global drained peatlands are used for agriculture (Tubiello et al., 2016). This practice occurs across most climate zones, with the largest emissions rates from drained peatlands ~~are~~ currently observed in Southeast Asia (Hooijer et al., 2010; Tubiello et al., 2016). Emissions rates due to microbial respiration are strongly controlled by temperature, and therefore, by climate zones. Drained peatlands in tropical regions have the highest annual emissions per area ( $\sim 70\text{-ton CO}_2 \text{ ha}^{-1} \text{ yr}^{-1}$ ), which is four and two times higher than in boreal and warm temperate climate zones, respectively (Tubiello et al., 2016; Hiraishi et al., 2013). Global emissions stemming from drained peatlands are estimated to be  $\sim 1.9 \text{ Pg CO}_2\text{-eq yr}^{-1}$  (Leifeld & Menichetti, 2018), accounting for  $\sim 3.6\%$  of global anthropogenic  $\text{CO}_2\text{-eq}$  emissions, with  $\sim 77\%$  of those emissions coming from the tropical climate zone (Leifeld & Menichetti, 2018).

Peatland restoration, primarily through rewetting drained peat, can reduce global  $\text{CO}_2$  emissions by  $0.5\text{-}1.3 \text{ Pg CO}_2\text{-eq yr}^{-1}$  (Leifeld ~~and~~ & Menichetti, 2018; Gunther et al., 2020), and account for 30% of global mitigation solutions for forests and other land ecosystems (Roe et al., 2021). Successful rewetting often involves blocking drainage systems, stabilizing water levels, and re-establishing vegetation adapted to wet conditions (Convention on Wetlands, 2021). While drainage leads to high  $\text{CO}_2$  emissions, maintaining a high WTL can result in increased methane emissions. However, studies showed that raising the WTL to near the surface results in a net reduction of  $\text{CO}_2$  emissions that outweighs methane emissions (Gunther et al., 2020; Wilson et al., 2016; Nyberg et al., 2022). A global meta-analysis found that keeping the groundwater level close to the surface can reduce global GHG's emissions from wetlands by about 90% by the end of the 21st century (Zou et al., 2022). Despite many studies highlighting the importance of rewetting for climate mitigation, there is scarce data on organic matter preservation in rewetted tropical peatlands (Nabuurs et al., 2023; Wilson et al., 2016). Available data on rewetted sites are limited ~~for~~ to several to ten years (Wilson et al., 2016), thus limiting the evaluation of rewetting effect on SOC preservation and GHG's emissions. Moreover, information about the time-frame since drainage, during which rewetting ~~is~~ remains effective, is crucial for selecting sites for management and preservation actions.

Peatlands deposits are rich in organic matter, characterized by about ~~50-wt.%~~ by weight of SOM (Joosten, 2002). While drainage leads to SOM loss and mineralization, studies have shown that recalcitrant material in peatlands may influence decomposition rates after drainage (Leifeld et al., 2012). Leifeld et al. (2012) showed that, in drained peatlands, high respiration rates were correlated with high concentrations of O-alkyl-C (a functional group of labile compounds). Könönen et al. (2016) found that tropical peat drainage leads to an increase in recalcitrant compounds concentrations ~~such as lignin~~. Phenol oxidase is known as a key factor in accelerating the decomposition of recalcitrant SOM, such as lignin, after drainage (Freeman et al., 2004). However,  $\text{Fe(II)}^{2+}$  oxidation may inhibit phenol oxidase activity and further enhancing Fe(III) and lignin complexation, as suggested by the 'iron gate'

90 mechanism (Wang et al., 2017). These studies ~~attest-suggest~~ that the reactivity for decomposition depends on the different fractions and compounds of the SOM in peat. However, the fate of these recalcitrant compounds after prolonged peat degradation and mineralization ~~is-yet-unknown~~remains unclear.

Recent evidence in mineral soils studies suggests that stabilization by recalcitrant material, as proposed in the 'selective preservation' hypothesis, is not common and relevant only at the beginning of SOM decomposition and  
95 in rich SOM soils (Rowley et al., 2017). Besides the limitations of oxygen availability due to water-saturation, and extreme temperature constraints on bacterial and enzymatic activity (Rowley et al., 2017), the most significant preservation mechanisms in mineral soils are SOM aggregation and mineral sorption (Lützow et al., 2006; Kleber et al., 2015). SOM aggregation is a physical process by which SOM is separated from decomposers by ~~the occluded~~  
occlusion of organic and inorganic particles within aggregates with cementing agents (Six et al., 2002).  
100 Aggregation is known to be promoted both by microbial activity (Chenu ~~and~~ Cosentino, 2011) and inorganic components in the soil matrix (Kleber et al. 2015). The sorption mechanism involves a wide range of reactions between organic compounds and soil minerals. Various clay minerals, oxides, and cations exhibit a high affinity for bonding with organic matter, leading to stabilization (Kleber et al., 2015; Rowley et al., 2017). Recent studies indicate that microbial residue, known as necromass, and depolymerized organic compounds produced by microbes  
105 are highly correlated with mineral-bonded OM, more than complex recalcitrant plant-derived material (Liang et al., 2017; Manzoni & Cotrufo ~~Franceesea~~, 2024).

Pristine peatlands lack these mechanisms (i.e., aggregation and sorption) due to their low mineral content (Leifeld et al., 2012; Mirabito & Chambers, 2023). However, these preservation mechanisms may become relevant after prolonged mineralization of peat soils. ~~Limited-k~~Knowledge exists-remains limited regarding about SOM pools and stabilization mechanisms  
110 in degraded peat soils. ~~Thus,~~This making-makes it challenging to accurately characterize long-term peat degradation and select suitable models for describing peat SOM loss over time.

~~To the best of our knowledge~~ Moreover, despite the disproportionate importance of peatlands to global carbon stocks compared to mineral soils, to the best of our knowledge, long-term peat degradation data and models remain scarce ,no long-term data or models on peat decomposition are available in the literature, ~~although many studies~~  
~~have been conducted on mineral soils.~~  
115 For instance, multiple long-term Bare Fallow experiments, spanning several decades, have been conducted on mineral soils in cold temperate climate zones (Barre et al., 2010). These experiments are especially valuable for identifying stable SOM, as they exclude organic matter inputs to the soil,  
a condition resembling drained peat.

Here we study a peatland site located in the Hula Valley, Israel, which has several important characteristics: (1) The site is located in a warm, subtropical climate; (2) The peat drainage occurred nearly 70 years ago, initiating a peat degradation period; (3) ~~RA~~rewetting of the bottom part of the degraded peat column, occurred three decades ago, ~~and leaving~~ the upper peat column drained until present time; and (4) Numerous historical studies were conducted in this site, enabling long-term (66 years) tracking of SOM stock. This research ~~hat had~~ two main objectives: First, to evaluate the long-term effect of rewetting on SOM preservation after nearly four decades of drainage in a warm subtropical climate. Second, to examine the decomposition rate of nearly 70 years ~~of long-term~~ degradation of drained peat and to explore the fate of peat SOM after prolonged mineralization period. To achieve this, we integrate Rock-Eval analysis, short-term respiration experiments, historical geochemical data, and a SOM decomposition model to investigate the long-term effects of drainage and rewetting on SOM degradation and preservation in the Hula Valley peatland.

## 130 2 Study site

The Hula Valley, Israel (33.1°N, 35.6°E), spans 175 km<sup>2</sup>, of which 28 km<sup>2</sup> consists of drained marshland that accumulated peat, now used mostly as cropland. The basin sustained lacustrine environments for the past 4 million years (Bein, 1986), with the last Hula Marsh prevailing from 2.5 to 4 thousand years ago until its drainage in 1957 (Cowgill, 1969; Bein, 1986). The dominant species found in the accumulated peat soil of the historic marsh was Cyperus Papyrus (Bein ~~and~~ Horowitz, 1986), typically occurring at characteristic water table of 0.5-2 m (Jones, 1940; Dimentman et al., 1992). The peat has accumulated in the top section of the sediment, reaching up to 12 meters of peat at its central area (Bein, 1967). This central area is also known as the 'deep peat' area which spans 12 km<sup>2</sup> (Marish, 1986). In 1957, the Hula Marsh and Lake were drained, primarily for agricultural land use and of reduction of water loss through evaporation. Consequently, the habitat was destroyed, and the exposed peat underwent aerobic conditions, accelerating SOM oxidation in the following years. This led to spontaneous peat fires, erosion, soil loss, subsidence, and nutrient leaching into downstream waterways (Hambright ~~and~~ Zohary, 1998; Litaor et al., 2011). In 1994, a major agricultural peat land preservation project was initiated, in which partial rewetting was implemented to halt soil degradation and fires by means of raising and stabilizing the WTL and frequent irrigation of crops (Hambright and Zohary, 1998). As part of the protection efforts, a hydrological barrier was constructed, and a local pond for water was created. Since then, the water table ~~is~~ has been managed at approximately -100 cm (Tsipris et al., ~~2021~~2022). Hence, the Hula's deep peat area can be divided vertically into 3 sections: (1) The deep pristine peat section (that was not drained) (2) mid-section of peat drained for 37 years

(1957-1994) and then rewetted for the last 30 years (1994-present) and (3) the top peat which was drained over the last 66 years (1957- present).

150

~~The site, classified as located within the Mediterranean sub-tropical classification region, with~~ characterized by ambient air ~~annual~~ mean annual temperature ~~of~~  $\sim 20^{\circ}\text{C}$  (Marish, 1986; online data MIGAL, 2023), ~~which classifies the area as a tropical soil respiration zone (Bond-Lamberty et al., 2024).~~ Soil average daily temperature measured at -20 cm depth was  $\sim 30^{\circ}\text{C}$  in summer months (Jun-Oct 2023) and  $\sim 15^{\circ}\text{C}$  in winter months (Dec-Mar 2023), with annual mean of  $\sim 22^{\circ}\text{C}$  (online data MIGAL, 2023). The annual mean precipitation was 610 mm (1987-97; Tsipris and Meron, 1998), and the peat area is used for cropland which is regularly irrigated during summer months preventing the upper peat from drying. WTL are routinely measured in 19 observation wells and 4 operational drainage channels throughout the Hula Valley, as documented in MIGAL's multi-annual reports over the past 20 years (Tsipris et al., ~~2021~~2022). However, assessing WTL in the drainage period between 1957 and 1994, is more challenging due to limited data availability. To address this, we relied on historical studies. Yaari-Cohen et al. (1971b) provides WTL data from core samples collected in December 1970, specifying the WTL for each core (Table 1). A report by Marish (1986) presents a spatial WTL map for winter 1985, indicating areas where the WTL is above or below -120 cm, and notes that no significant changes in WTL have occurred since Yaari et al. (1971b) (Table 1). Therefore, we adopted Yaari-Cohen et al. (1971b) WTL data to represent the period from post-drainage (1957) until the rewetting in 1994. Further, we choose to follow Yaari-Cohen et al. (1971b) soil profile's locations for our study and soil core samples collection for comparison. Table 1 presents the complete dataset of WTL variations from 1957 to 2021 in our examined profiles.

165

170

In this research, we focused on the area north of the Agmon pond, where the WTL is managed between the eastern and western canals, bordered by underground hydrological barrier (emphasized by an orange outline; Fig. 1). Focusing in this selection targeted areas with substantial peat thickness, enabled investigation of both drainage and rewetting effects, and ensured the availability of historical WTL records and SOM data.

Table 1. ~~water~~ Water table levels in cm. Negative values (-) for indicate depth below ground surface

Profile	Yaari- <u>Cohen</u> et al. (1971b)	Marish (1986)	Monitoring (2001-2021) mean by annual ( $\pm$ STDV)
A*	-80	Above -120	-101 $\pm$ 18*
C	-220	Below -120	-226 $\pm$ 17
D*	-120	Above -120	-101 $\pm$ 18*

E*	-165	Below -120	-101 ± 18*
F	-200	Below -120	-95 ± 10

---

\* Based on data collected from observation well number 13. Locations marked in Figure 1.


---

175

180

### 2.22.1 Sample collection

Soil samples were collected in January-April of 2023 from the 'Deep Peat' area (Fig. 1). Five different locations were chosen based on the sampling locations of Yaari-Cohen et al. (1971a) designated there as Crossline B. For coordinates, see the Supplementary Material (1.8). To ensure that the sampling will included the pristine peat, each core was excavated to a depth of approximately -400-e m (much deeper than the WTL preceding the drainage; Table 1). We used a manual gauge auger (Eijkelkamp<sup>®</sup>) of either 1 or 0.5 m long cores and of 5 and 2 cm diameter, respectively. The soil sampling from the cores was based on two collection schemes: (1) sampling of discrete 2--5 5 gf samples, collected at 10 cm intervals, into pre-weighed 50 ml tubes. These samples were mainly used for



**Figure 1:** Study site satellite map – Hula valley, Israel (coordinates of Israel TM Grid). Polygons of the historic marsh boundaries in dark red, deep peat area in black dashed line (by-after Marish 1986), WTL observation wells in white circles, core locations in light blue, WTL management canals zoneunderground hydrological barrier in orange. Regional map to the right in WGS 1984 coordinates. Image made by ArcGIS software and is based on ESRI-provided data.

190 RockEval analysis, (2) sampling of homogeneous soil sections, based on lithology and historic WTL, at 30-60 cm intervals to cotton aired bags. These samples were mainly used for respiration incubation experiments.

### 3 Materials and Methods

#### 3.1 Rock-Eval ~~Analysis-analysis~~

195 ~~Rock-Eval pyrolysis is a widely accepted method for the A~~ analysis of ~~the peat and organic~~ soil samples for SOM characterization ~~and soil organic carbon (SOC) estimation (e.g., Disnar et al., 2008; Marchand et al., 2008; Sebag et al., 2016; Brown et al., 2023).~~ In this study, ~~was carried out with~~ the newest Rock-Eval 7S analyzer ~~was used~~ (RE-7S, Vinci Technologies). 107 samples of peat from all profiles were freeze-dried, ground, and homogenized for analysis. About 15-30 mg of ~~the~~ ground sample were taken for analysis. The analytical protocol included two  
200 phases: (1) pyrolysis in an inert N<sub>2</sub> atmosphere starting at a temperature of 200 °C, reaching 650 °C; (2) combustion using air, starting at a temperature of 300°C and rising to 850 or 1200°C when sulfates were also quantified (see below). ~~Both steps~~ The heating rate ~~for both steps is was~~ 25 °C·min<sup>-1</sup>, and combustion had a final ~~hold~~ time of 10 min at maximum temperature before cooling down. The heating cycle is based on Behar et al. (2001) and modified for soils based on Sebag et al. (2016). The gradual heating of the RE-7S promotes a sequential decomposition of  
205 organic and inorganic phases according to their thermal stability. Continuous monitoring of the generated products is carried via four different detectors: (1) a Flame Ionization detector (FID) for hydrocarbons (HC), (2) two Infrared (IR) detectors for CO and CO<sub>2</sub> and (3) Ultra violet (UV) detector for SO<sub>2</sub>.

Calibration of C was done using the IFP 160000 standard (Behar et al., 2001). As carbonates are also decomposed during a RE analysis, an empirical separation between organic and inorganic C was previously determined for  
210 rocks and mineral soils (Behar et al., 2001). Usually, the organic CO, and CO<sub>2</sub> at the pyrolysis phase is determined up to 400°C, and for the oxidation phase, at a local minimum above 650°C. The mineral carbon (MINC) is the remaining fraction of the CO<sub>2</sub> and CO curves, calculated between 400-650°C in the pyrolysis phase and approximately between 650-850°C in the oxidation phase. To validate the usefulness of this discrimination, we also applied a loss-on-ignition (LOI) procedure to remove SOM and to identify the correct separation between soil  
215 organic carbon (SOC) and mineral carbon (MINC) (see Supplementary Material 1.5 ~~and Fig. S4~~). SOC is calculated by summing the hydrocarbon signal (HC), CO, and CO<sub>2</sub> curves from both the pyrolysis and oxidation phases. The HC peak, representing carbon in a reduced state, is used to calculate the hydrogen index (HI) of the SOC mg HC g<sup>-1</sup> SOC<sup>-1</sup>. The HC signal was also integrated in 4 parts representing different thermal stability regimes according

to Sebag et al. (2016): A<sub>1</sub> (200-340°C), A<sub>2</sub> (340-400°C), A<sub>3</sub> (400-460°C), A<sub>4</sub> (460-650°C), which are used to  
220 calculate two indices: The R-index (RI) calculated as (A<sub>3</sub>+A<sub>4</sub>)/A<sub>1-4</sub> is indicative of the relatively thermal stable  
fraction of the HC, while the I-index ( $\log_{10} [(A_1+A_2)/A_3]$ ) is indicative for the thermal labile fraction of HC (Sebag  
et al., 2016). ~~Similarly to the HI, the~~ The oxygen index in its revised version-(OI[RE6]; mg CO<sub>2</sub> g<sup>-1</sup> SOC<sup>-1</sup>), represents  
the oxygen -emitted as CO<sub>2</sub> and CO in the pyrolysis phase (200-400°C), corresponding to oxygen content in the  
OM (Saenger et al. 2013; Cécillon et al., 2018; Delahaie et al., 2023).- The TpkS2 °C is the temperature at the  
225 maximum HC signal.

Calibration of sulfur signal in both the pyrolysis and combustion phases was done using pure organic S compound  
(Di-benzothiophene) and pure pyrite, following Cohen-Sadon et al. (2022). For real samples that contain multiple  
S fractions, the organic and pyrite fractions are separated during the pyrolysis phase and can be distinguished by a  
local minimum at approximately 500°C (Cohen-Sadon et al., 2022). However, during the oxidation phase, the  
230 residual organic and pyrite S are co-eluted at around 400°C, requiring an empirical relation to deconvolute (Cohen  
Sadon et al., 2022; Aboussou, 2018). Moreover, some of the pyritic S is retained during pyrolysis at the presence  
of carbonates and is released at higher temperature (~700-900°C). Sulfate sulfur measurement requires high  
temperatures and is released during combustion above 900-1000°C (Aboussou, 2018). In this contribution the  
cutoff between retained S and sulfate signals was set manually to the local minimum between them, at temperature  
235 of ~1000°C.

### 3.2 Soil incubation

Soil samples were oven-dried at 60°C overnight and then stored in the lab. Each profile consists of 6 samples, each  
sample was gently ground, if it contained large aggregates, and sieved to < 2 mm. From each sample, four splits  
of 1.5 g portion of soil were weighed into 6 ml vessels, then distilled water was added to 60% of its water-holding  
240 capacity. Water holding capacity was determined separately for degraded and pristine samples separately. This was  
done by saturating 15 g<sup>±</sup> of soil with 50 ml distilled water and measuring the drained volume of water, using  
Whatman 40 filter (8µm). All samples were preincubated for bacterial acclimation for five days at the designated  
temperature. Respiration experiments were conducted at 22°C and 33°C, which represent the annual average soil  
temperature and its maximal summer temperature in Hula valley, respectively. The experiments were aerobic and  
245 conducted in duplicates under two conditions: (1) for seven days at 22°C, and (2) for two days at 33°C. During the  
incubations, gas samples were collected for CO<sub>2</sub> and O<sub>2</sub> concentration analyses. Samples were taken using glass  
flasks (~3.7 mL or ~13 mL, depending on incubation duration) equipped with Louwer™ O-ring high-vacuum  
valves. Measurements were taken at halfway (t=1) and at the end (t=2) of the experiment to validate a linear change

in gas concentration during incubation. The O<sub>2</sub> [ $\mu\text{mol}$ ] and CO<sub>2</sub> [ $\mu\text{mol}$ ] of the air samples were measured in the  
 250 laboratory by a closed system (The “Hampadah”; Hilman and Angert, 2016). The system is based on two analyzers:  
 an infra-red gas analyzer (IRGA) for CO<sub>2</sub> measurement (LI 840A LI-COR; Lincoln, NE, USA) and a fuel-cell-  
 based analyzer (FC-10; Sable Systems International, Las Vegas, NV, USA) for measuring O<sub>2</sub>, and is fully  
 automated. Each sample was incubated and measured in a known volume, to derive the percentage and  
 concentration of CO<sub>2</sub> and O<sub>2</sub>. Fluxes rates [ $\mu\text{mol}\cdot\text{day}^{-1}\cdot\text{g}_{\text{SOC}}^{-1}$ ], and reaction constant ( $k$  [ $\text{yr}^{-1}$ ]) of CO<sub>2</sub> and O<sub>2</sub> were  
 255 calculated for both 22°C and 33°C, as shown in Equations 1 and 2, respectively. The apparent respiratory quotient  
 (ARQ), as shown in Equation 3, represents the ratio of CO<sub>2</sub> and O<sub>2</sub> fluxes (Angert et al., 2015). This ratio provides  
 information on the respiratory substrate stoichiometry as well as the underlying biotic and abiotic processes  
 (Hilman et al., 2022). Temperature sensitivity of organic matter decomposition, expressed as Q<sub>10</sub> (Eq. 4), is a metric  
 for assessing the vulnerability of organic matter to temperature increases (Hilasvuori et al., 2013). Q<sub>10</sub> was  
 260 calculated using the rate constant  $k$ , derived from respiration experiments ~~conducted at 22°C and 33°C, which~~  
~~represent the annual average soil temperature and its maximal summer temperature in Hula valley, respectively.~~

$$(Eq. 1) \text{ Flux} = \frac{[CO_2 (t=2)] [\mu\text{mol}] - [CO_2 \text{ ambient}] [\mu\text{mol}]}{\text{daytime} * \text{gr}_{\text{SOC}}}$$

$$(Eq. 2) k \text{ constant } [\text{yr}^{-1}] = \text{Flux} * 12 \frac{\text{g C}}{\text{mol}} * \frac{1 \text{ gr}}{10^6 \mu\text{gr}} * \frac{365 \text{ days}}{1 \text{ yr}}$$

$$(Eq. 2) k \text{ constant} = \text{Flux} * 12 \frac{\text{g C}}{\text{mol}} * \frac{1 \text{ g}}{10^6 \mu\text{g}} * \frac{365 \text{ days}}{1 \text{ yr}}$$

$$(Eq. 3) \text{ ARQ} = \frac{[CO_2 (t=2)] - [CO_2 \text{ ambient}]}{[O_2 (t=2)] - [O_2 \text{ ambient}]} \quad (Eq. 4) Q_{10} = \frac{k(33^\circ\text{C})^{33-22}}{k(22^\circ\text{C})^{10}}$$

270

### 3.3 Statistical analysis

Statistical analyses were performed using JMP software (JMP®, JMP Pro 17, SAS Institute Inc., Cary, NC, USA).  
 To determine the difference between the peat column sections, we used one-way analysis of variance (ANOVA)  
 and the Tukey-Kramer test. For unequal variances, we used a Welch's test and nonparametric comparisons using  
 275 Steel-Dwass test. Significant differences were determined at  $P < 0.05$ .

The geochemical and respiration data were statistically analyzed based on three peat profile sections: drained,  
 rewetted, and pristine, defined according to their respective historic and current WTL's (Table 1). The drained

section included samples from depths of -30 cm to -100 cm, except for profile C, in which it extended to -180 cm. The upper boundary at -30 cm was set to minimize the influence of agricultural practices such as soil tillage and root bias. The rewetted section, available only in profiles E and F, ranged from -110 cm to -160 cm in profile E and -110 cm to -190 cm in profile F. A rewetted section is absent from the other cores as because the WTL's post-drainage and post-rewetting were very similar (Fig. 2, purple and light blue shaded areas). The pristine section extended from -200 cm to -400 cm in profiles C, E, and F, while in profiles A and D, it began at a depth of -150 cm. Samples containing more than 30% carbonate content were excluded from the statistical analysis to avoid incorporating lake mineral horizons, which were primarily found at the bottom of the profile.

### 3.4 Historical data of SOM

Since its drainage in 1957, the 'deep peat' area of the Hula Valley (marked in Fig. 1) has been studied in numerous research projects, some of which have dealt specifically with SOM loss over time, such as the report by ~~the~~ Marish (1986) and Litaor et al., (2011). To assess the changes in peat soil organic matter (SOM) subjected to 66 years of drainage, we focused on the drained section identified in previous studies with multiple soil profiles (Schallinger et al., 1970; Yaari-Cohen et al., 1971 ~~a&b~~; Marish, 1986, Litaor et al., 2003). We constrained our data compilation to the boundaries of the 'deep peat' area to ensure consistency and avoid significant lithological variations, as well as the impact of soil fires, which mainly occurred south of this region. ~~For depth selections, we set the upper boundary at -30 cm to minimize the influence of agricultural practices such as soil tillage and root bias.~~ To focus on the upper, drained peat, the upper boundary was set at -30 cm (to minimize the influence of agriculture practices) and the lower boundary was established at -100 cm, which reflects the prevailing WTL in the region, given the spatial heterogeneity of the WTL. To convert the SOC results to SOM data, as reported by previous studies, we measured ~~gravimetrically the SOM by removing SOM through combustion (using the SOM gravimetrically by removing SOM through combustion (using the~~ LOI<sub>400</sub> procedure; Nelson and Sommers, 1996) ~~to~~ on a set of samples. We then correlated the SOM to SOC and applied a correction factor. Details of this correction are provided in the supplementary materials (1.5).

### 3.5 Fitted SOM Model

The modelling of organic matter loss through decomposition is a well-studied topic (Manzoni and Porporato, 2009), with various models proposed to represent these processes (Manzoni et al., 2012). Here we chose a model that comprises of two organic matter compartments (pools), as described by equation 5. These two pools are: (1) a labile

SOM pool decomposing exponentially with reaction constant  $k$ , and (2) a persistent SOM pool ( $SOM_p$ ), ~~which with is inert, with no decomposition or formation rates that are significantly lower than the observed duration of degradation for the labile pool.~~  $SOM_i$  represents the initial SOM value of the model,  $P$  denotes the fraction of persistent SOM out of  $SOM_i$ ,  $t$  denotes time in years, and  $k$  serves as the reaction constant for SOM decomposition. The unknown parameters,  $k$  and  $SOM_p$ , are optimized to minimize the root mean square deviation (RMSD; Eq. 6), which quantifies the mean difference between model predictions and observed data. Where  $SOM_{data}$  represents field data from various studies,  $SOM_{model}$  denotes the simulated values, and  $n$  is the number of observations. The initial SOM ( $SOM_i$ ) value is based on Ravikovitch 1948 (1945) work. Historical data and this current study (five data points,  $n=5$ ) are used as the observed data ( $SOM_{data}$ ) for RMSD calculation. The model simulates the long-term decline in organic matter following drainage from 1957 to 2023 (present study).

$$(Eq. 5) \text{SOM}(t)_{[wt.\%]} = (SOM_i * (1 - P)) * e^{-kt} + (SOM_p) , \quad P = \frac{SOM_p_{[wt.\%]}}{SOM_i_{[wt.\%]}}$$


$$(Eq. 6) \text{RMSD} = \sqrt{\frac{\sum_y^n (SOM_{data} - SOM_{model})^2}{n}} , n = 5$$

320

## 4 Results and Discussion

### 4.1 SOC oxidation and mineralization

Previous research on Hula peat has highlighted a decline in the SOM content within the drained section, attributed to aerobic conditions (Marish, 1986; Litaor et al., 2011). Based on these findings, we hypothesize a continued reduction in SOM content in the upper drained section and anticipate potential evidence of SOM preservation in the rewetted section, which has remained unexplored to date. Additionally, the pristine section, which remained under anaerobic conditions, is expected to exhibit minimal variations in SOM content. Figure 2 shows the measured SOC profiles wt. [% SOC] from west to east, along with their corresponding WTL as detailed in Table 1. Overall, SOC content in all profiles is lower in the drained and rewetted sections relative to the pristine section. The profiles can be categorized into two main types: (1) profiles with similar post-drainage and post-rewetting WTLs, resulting in only drained and pristine sections – profiles A, D and C, and (2) profiles with different historical WTLs, creating a rewetted section between the drained and pristine sections – (profiles E and F). In most profiles, there is a clear upward decline in SOC corresponding to the post-drainage WTL (purple bars, Fig. 2), indicating mineralization and SOC loss due to the lowering of the WTL. For instance, profiles A and C (type 1) had relatively constant but



**Figure 2:** SOC profiles (from west to east) of peat cores with respective WTLs - post drainage (purple – after Yaari-Cohen et al. 1971b) and post rewetting (light blue – MIGAL's Monitoring data).

335 different WTLs over 66 years of drainage. In both profiles, the depth in which SOC declines above it, aligns with the respective WTLs.

The trends in SOM content are further supported by additional organic geochemical data, such as the OI and RI indices. For example, a detailed composite plot of profile F (type 2) identifies three distinct sections (Fig. 3),  
340 corresponding to drained, rewetted, and pristine conditions, separated by the historical WTL. Alongside the declining SOC trend, a negative relationship exists between SOC and both OI and RI. ~~This indicating suggests~~ that as peat degradation progresses, ~~the~~ residual SOC becomes more oxidized, and ~~exhibits an increase in~~ the proportion of the thermally stable fraction ~~increases~~. Similar coupling of the historical WTL and the trends of the organic geochemical proxies ~~were was~~ found for the rest of the cores, and their composite plots are given in the  
345 Supplementary Material (1.4).

Sulfur ~~species speciations~~, obtained using the RE-7S, ~~were was~~ distinguished as detailed in Supplementary Material 1.9. In the drained section, a decline in both organic sulfur and pyrite is evident, while sulfate accumulates, ~~which reflecting reflects~~ aerobic conditions (Fig. 3 and Supplementary Material 1.4). In contrast, the pristine and rewetted sections show low sulfate levels ~~but accompanied by high elevated~~ pyrite concentrations, ~~with~~ ~~The rewetted section~~  
350 ~~exhibits~~ a maximum pyrite abundance ~~in the rewetted section, clearly indicating that rewetting successfully re-established reducing conditions~~. Assuming that pyrite was ~~fully largely~~ oxidized in the rewetted section during its ~~37-year~~ drainage period (~~37 years~~), this peak ~~likely is interpreted represents as~~ secondary pyrite formation through sulfate reduction under anaerobic conditions following rewetting. ~~Moreover, The the~~ peak in pyrite concentration beneath the recent WTL was observed in most profiles (supplementary material 1.4), suggesting a sulfur recycling  
355 mechanism in the drained and rewetted peat sections. ~~Specifically, soluble sulfate appears to be constantly washed from the drained section and accumulates under anoxic conditions, in its reduced forms, within the rewetted section~~. Ultimately, pyrite serves as an indicator ~~for of~~ both sulfate reduction in the sediment and ~~the~~ recent WTL regimes. The corresponding respiration fluxes reveal two key points (Fig. 3 and supplementary material 1.4): (1) respiration fluxes (normalized to SOC) in the drained and pristine sections are similar despite differences in SOC content and  
360 chemical properties (as indicated by OI~~[RE6]~~ and RI); and (2) respiration fluxes in the rewetted section are twice as high as in the other two sections. These findings are explored further in the next section.

pH levels remain mostly neutral but drop to 5.9 in profile F and as low as 3.5 in other profiles. Litaor et al. (2011) also found pH levels of 5-3.8 at a depth of 50 cm in carbonate ~~free~~ horizons, which they interpreted to result from pyrite oxidation, and nitrification.




Figure 3: Detailed composite plot for profile F7, including the historical water table levels.


365

#### 4.3.4.2 Average SOC characteristics of drained, rewetted and pristine peat

In the current section we present a statistical compilation of all the soil profiles data. The analyses are made for the predefined three peat column sections: drained, rewetted, and pristine. Figure 4 present the average thermal analysis indices for each section, including standard error and the number of available data points for each section. Overview of the results and statistical tests for sections comparison are presented in supplementary Tables S1 and S2. Generally, the data show a clear trend with depth for each of the thermal indices. Despite the expected natural variation that characterize such sedimentary basin, it was found that most of the thermal indices show significant difference ( $p < 0.05$ ) between all the three sections defined. These include the SOC, TpkS2, RI, and I-index. The only and one exception is the OI[RE6] values, which show no significant difference between that the drained section OI values is not significantly different from and the rewetted section). The mean SOC values for the drained, rewetted, and pristine sections are  $13 \pm 1$ ,  $20 \pm 2$  and  $35 \pm 1$  wt.%, respectively. Thus, it is evident indicating that 66 years of drainage have led to extensive degradation that resulted in, reducing


380 the lowering of SOC in the drained section to approximately one-third of its original pristine value, ~~(without~~  
~~considering~~ not accounting for compaction processes). The rewetted section shows higher SOC content than the  
drained section, suggesting that degradation ceased or decreased following rewetting. The OI [RE6] shows an  
opposite reverse pattern to the SOC, with a gradual decrease in its mean value from the drained to the rewetted and  
pristine sections (~~195-157 ±75~~, ~~160-133 ±11-8~~ and ~~100-88 ±7-5~~ mg CO<sub>2</sub> g<sup>-1</sup> SOC<sup>-1</sup>, respectively). This pattern can  
indicate a partial oxidation of the SOM due to drainage and suggests a slowdown in oxidation due to rewetting.  
385 ~~This Additional support is evident observation is further supported~~ by other SOM thermal indices (Fig.4; correlation  
in supplementary material Fig.S6). For example, Mmean TpkS2 values for the drained, rewetted, and pristine  
sections are 460±5, 431±7 and 378±5 °C, showing the increase of HC peak temperature along with degradation  
and SOC decline. The mean RI values for the drained, rewetted, and pristine sections are 0.67±0.01, 0.59±0.02,  
and 0.49±0.01, respectively, while the mean I-index values are -0.03±0.01, 0.1±0.02 and 0.26±0.01, respectively  
390 (Table S1). While the RI and I-index values of the pristine section represent organic soil, those of the rewetted and  
drained sections fall into the range typical for organic-mineral and mineral soil horizons (Sebag et al., 2016). The  
RI values of the drained and pristine peatlands further align with the findings of Brown et al. (2023), who reported  
higher RI values for disturbed peat sites compared to pristine peat sites, all measured from the top 50 cm peat layer  
(duration since drainage was not specified). The higher thermal stability of the OM observed in the degraded peat  
395 sections is presumably indicative of preferential utilization of the thermally labile fraction indicates that  
while during microbial respiration occurred, the organic matter being utilized is mostly the thermally labile  
fraction; †This relationship between available OM for decomposition and thermal stability is further explored in  
subsection 4.7 below.

Together, these results demonstrate that with decomposition the residual SOM becomes more oxidized and contains  
400 a lower proportion of thermally labile compounds. The increase in thermal stability could be due to a relative  
increase of litter recalcitrant material (Könönen et al., 2016; Barré et al., 2016), that the residual SOM is minerals  
associated (Saenger et al., ~~2015~~2014), or both. Moreover, the rewetted section exhibits a higher degree of SOM  
preservation compared to the drained section, highlighting the impact of rewetting on reducing degradation.



**Figure 4:** Rock Eval analysis results. mean value and standard error of all the profiles from three sections [cm]: drained [30-100] (n=25), rewetted [110-170] (n=10) and pristine [200-400] (n=26). (a) SOC content, (b) R-index, (c) Oxygen index [RE6] and (d)

405




**Figure 5:** Respiration experiment results - mean value and standard error of all the profiles from three sections [cm]: drained [30-100] (n=8), rewetted [110-170] (n=4) and pristine [200-400] (n=12). (a) CO<sub>2</sub> and O<sub>2</sub> fluxes at 22°C experiments. (b) CO<sub>2</sub> and O<sub>2</sub> fluxes at 33°C experiments. (c) ARQ -CO<sub>2</sub> fluxes divided by O<sub>2</sub> fluxes derived from 22°C and 33°C experiments. (d) Q10- derived from CO<sub>2</sub> and O<sub>2</sub> k constant.HC peak temperature - TpkS2.

**Figure 4:** Rock Eval analysis results—mean value and standard error of all the profiles from three sections [cm]: drained [30–100] (n=25), rewetted [110–170] (n=10) and pristine [200–400] (n=26). (a) SOC content, (b) R index, (c) Oxygen index [RE6] and (d) HC peak temperature  $T_{pkS2}$ .

**Figure 5:** Respiration experiment results—mean value and standard error of all the profiles from three sections [cm]: drained [30–100] (n=8), rewetted [110–170] (n=4) and pristine [200–400] (n=12). (a)  $CO_2$  and  $O_2$  fluxes at 22°C experiments. (b)  $CO_2$  and  $O_2$  fluxes at 33°C experiments. (c) ARQ— $CO_2$  fluxes divided by  $O_2$  fluxes derived from 22°C and 33°C experiments. (d)  $Q_{10}$ —derived from  $CO_2$  and  $O_2$  k constant.

410

The experimental respiration results were also statistically analysed in each of the defined sections for all the profiles collectively. Figure 5 displays the mean  $CO_2$  and  $O_2$  fluxes per gram of SOC at 22°C and 33°C, and their derivatives indices—ARQ and  $Q_{10}$ . Mean fluxes of  $CO_2$  of the drained, rewetted, and pristine sections at 22°C



**Figure 6:** Mean soil organic matter (SOM) content of Hula peat since the drainage, based on data from this study (triangles) and historical studies (Purple circles). All from depths of 30–100 cm.

were  $9 \pm 2$ ,  $16 \pm 2$  and  $9 \pm 1$   $\mu\text{mol CO}_2 \text{ g}^{-1} \text{ SOC}^{-1} \text{ day}^{-1}$ , respectively (Fig. 5a), showing a similar flux for the drained and pristine sections, and approximately a double flux in the rewetted section. This pattern is also evident for the  $\text{O}_2$  fluxes at  $22^\circ\text{C}$  ( $-16 \pm 3$ ,  $-37 \pm 4$  and  $-20 \pm 2$   $\mu\text{mol O}_2 \text{ g}^{-1} \text{ SOC}^{-1} \text{ day}^{-1}$ ) for the drained, rewetted and pristine sections, respectively; Fig. 5a). The same pattern is evident in the  $33^\circ\text{C}$  experiments in both  $\text{CO}_2$  and  $\text{O}_2$  (Fig. 5b), reinforcing the reliability of this finding. It should be noted that our incubation experiments were done at 60% of the soil water holding capacity, which is a standard approach aimed at measuring respiration at close to optimal moisture conditions (Rey et al., 2005). Of course, the soil moisture in the field is variable, and the upper soil is often drier or wetter than this 60% value, while the lower sections are constantly saturated. Hence, the rates measured during these incubation experiments are an estimate of the potential respiration rates and the SOM short-term stability. For example, the value measured for the pristine or the re-flooded section gives an estimate for the rate of carbon loss if these sections were to be drained (and kept moist).

We found similar fluxes in the drained and pristine sections despite the distinctly their different SOC, RI and OI values (Fig. 4). A similar observation was made by Bader et al. (2017, 2018), which also found that the degraded peat horizon in cold temperate sites exhibited similar flux per SOC as deeper, less disturbed peat. Moreover, in our study, the rewetted section which has intermediate SOC, RI and OI values (compared with the pristine and drained sections), had the highest respiration fluxes. These results indicate that SOC content and RE-7S proxies (RI, OI, and I-index) do not directly reflect SOM bioavailability for short-term bacterial respiration. The high  $\text{CO}_2$  fluxes that were measured during the incubation of rewetted peat at 60% water holding capacity can be related to the exposure of flooded soil to oxygen, which may cause a temporary increase in SOC oxidation. However, this does not explain the reason for the higher respiration rates in the rewetted peat versus the pristine peat, as the latter was also exposed to oxygen during incubation. Thus, it seems that the drying and re-wetting of this section caused destabilization of the SOM, potentially through the release of occluded and mineral-protected SOM (e.g. iron-bound), or by changes in enzyme activity. The exact mechanism warrants future investigation.

The first order reaction constant  $k$  (Eq. 2) was calculated for aerobic SOC decomposition at  $22^\circ\text{C}$ , resulting in  $-0.037$ ,  $-0.087$  and  $-0.033 \text{ yr}^{-1}$ , for the drained, rewetted and pristine sections, respectively. Temperature sensitivity, as indicated by the mean  $Q_{10}$  values, for  $\text{O}_2$  fluxes was  $2.9 \pm 0.4$  for the drained section and  $2.5 \pm 0.4$  for the pristine and rewetted sections, with no significant difference between the sections. Mean  $Q_{10}$  values from  $\text{CO}_2$  of across all sections were relatively similar ( $2.3 \pm 0.3$ ), with no statistically significant difference (Table S2). Bader et al. (2017, 2018) found that the mean  $Q_{10}$  (derived from  $\text{CO}_2$  flux) for drained peatlands in cold temperate sites was  $2.5 \pm 0.1$ , which resembles our findings despite the different climate.

The ARQ results also show no significant difference between the sections, with mean values at 22°C of 0.6 ±0.05  
445 , 0.43 ±0.07 and 0.45 ±0.04, for the drained, rewetted and pristine sections, respectively. In aerobic conditions the  
expected ARQ value can be indicative of the oxidation state of the decomposed organic carbon compounds (Hilman  
et al., 2022). Values of ARQ of 1.0 are attribute to sugars which are more labile, while lower values are attribute  
to compounds such as lipids (0.73) or proteins (0.77), which can derive from necromass and more recalcitrant  
material (Masiello et al., 2008; Hicks Pries et al., 2020). However, the pristine and rewetted ARQ results show  
450 lower values that ~~cannot be solely are not~~ attributed to ~~standard organic any~~ compounds. These low values thus  
may indicate oxygen consumption, which is not related to CO<sub>2</sub> production, such as partial oxidation of SOM or  
oxidation of reduced chemical species (Hilman et al., 2022), such as Fe<sup>2+</sup> and sulfides, which are prevalent in such  
peat soils and thus may indicate oxygen consumption by partial oxidation of SOM or oxidation of reduced chemical  
species such as Fe<sup>2+</sup> or sulfides (Markel., et al 1998).

455

#### **4.54.3 Long-term SOM decomposition in the Hula's drained peat**

The Hula peat exhibits lithological variations both spatially and with depth, resulting from the natural habitat's  
dependence on varying water regimes and plant diversity. Additionally, as mentioned above, a decline in SOM of  
Hula's peat since its drainage was reported (Marish, 1986; Litaor et al., 2011). Here, we evaluate long-term (7  
460 decades) decomposition of peat in the field, by compiling literature data focusing on the top drained section. To  
this end, studies that included comprehensive data of spatial distribution and sampling depths together with  
evaluations of SOM of the drained section (30-100 cm) were compiled and ~~analyzed~~analysed. The SOC data from  
this study were converted to SOM using correlations factors for the Hula peat soils determined here as 1.6 for  
drained samples and 1.8 for pristine samples (Supplementary Fig. S3). The full data set is summarized in Table 2  
465 and illustrated in Fig. 6, alongside our SOM analysis.

During the 66 years of drainage the mean SOM content of the top layer significantly decreased from 68±4% to  
21±2% (Fig. 6). This decline was continuous, following an exponential decay pattern, with the most substantial  
loss of organic matter occurring during the first three decades, after which the rate of SOM decomposition slowed.  
Notably, the initial SOM value reported by Ravikovitch ~~in (19481945)~~ aligns closely with the SOM data of the  
470 pristine section in our study (2023), supporting the ~~concept-claim~~ of preservation under saturated conditions (Fig.  
6). Additionally, the SOM value in the rewetted section also conforms to this timeline, suggesting that it has  
remained mostly unchanged since 1994. These agreements are illustrated by the grey arrows in Fig. 6.

Table 2. Changes in soil organic matter content in the Hula deep peat area since the drainage.

Year	Study	Mean SOM% $\pm$ SE	n	Spatial n	Depth [cm]	LOI Method
1948 <sup>a</sup>	Ravikovitch <a href="#">1948</a> <a href="#">1945</a>	68 $\pm$ 4	4	3	0-100	Not reported
1970	Schallinger et al., 1970	48 $\pm$ 2	30	10	25-100	Not reported
1971	Yaari-Cohen et al., 1971	47 $\pm$ 3	10	9	40-110	Method 1
1986	Marish, 1986	33 $\pm$ 2	9	4	40-100	Not reported
2000	Litaor et al., 2003	31 $\pm$ 4	27	27	50	Method 2
2023	This Study	21 $\pm$ 2	20	5	30-100	Method 3
2023	This Study	33 $\pm$ 2	8	2	110-170	Method 3
2023	This Study	64 $\pm$ 2	23	5	200-400	Method 3

**Method 1:** LOI at 850°C and carbonate correction following acid digestion. **Method 2:** Acid digestion and LOI at 850°C.

**Method 3:** This study - SOC analysis with RE-7S and LOI at 400°C conversion (Supplementary 1.5).

#### 4.64.4 The two-pool SOM model

475 The decay pattern in Fig. 6 shows an exponential decline, a behavior similar to long-term Bare Fallow experiments of mineral soils (Barre et al., 2010). To better understand the long-term decomposition process of Hula's peat, we explored the observed SOM data with a model. A simple one-box model proved inadequate for capturing the decomposition process of SOM in mineral soils (Barré et al., 2010; Manzoni et al., 2012; Clivot et al., 2019). The long-term data of Manzoni et al. (2012), based on the Rothamsted Bare Fallow experiment, serves as a good analog

480 to Hula's peat SOM for two reasons: (1) it involves decomposition without significant additional organic matter inputs, and (2) it follows an exponential decomposition pattern over time. Manzoni et al. (2012) suggest that a two-pool model effectively captures long-term soil respiration dynamics, making further complexity unnecessary. Comparable models were used by Barré et al. (2010) and Cécillon et al. (2021), in which one pool undergoes exponential decay while the other remains stable. Other models account for plant input flux (e.g., the AMG model;

485 Clivot et al., 2019) and leaching of dissolved organic carbon as a pathway for SOM decomposition (e.g., Page et al., 2016, Rixen et al., 2016). Here we assume that plant input to the deep peat (i.e., below 30 cm) is negligible and focus on total SOM loss rather than specific pathways (i.e., respiration vs. DOC leaching). In addition, although seasonal soil moisture and temperature variations influence SOM decomposition rates, their effects were not addressed explicitly in this analysis. Ultimately, we adopted a two-pool model, where a persistent pool of SOM is

490 initially present and assumed to have a negligible decay rate, while a labile-intermediate pool is consumed with a single reaction constant (Eq. 5).


#### 4.74.5 The SOM Model results

As discussed above, the drained and pristine sections of Hula peat have similar respiration fluxes per SOC-unit SOC, both for the CO<sub>2</sub> efflux and the O<sub>2</sub> influx (Fig.5). Similarly, Bader et al. (2017,2018) also found, in their study of peat in temperate-cold climate, that respiration rates per SOC unit in drained and undisturbed peat are comparable. Thus, we suggest postulate that both the pristine and drained peat sections can be characterized by the same reaction constant (see Eq. 5), despite their different organic characteristics. Practically, this similarity allows for a fixed reaction constant ( $k$ ) to be applied to the labile SOM pool throughout the entire model duration. Also, the similarity of the present-day pristine SOM value to those reported for the peat in the historic marsh of the Hula by Ravikovitch (1948,1945) (i.e., pre-drainage) allows us to assume a value of  $68 \pm 4$  wt.% for the initial SOM<sub>i</sub> (i.e., pre-drainage).

Hence, our two-pool model includes two unknown variables: SOM<sub>p</sub> and  $k$ . With these two degrees of freedom, the model yields a range of possible solutions. The model was run with multiple SOM<sub>p</sub> and  $k$  values, and the RMSD was calculated for each run (Eq. 6). Plotting SOM<sub>p</sub> versus  $k$  and RMSD for all the runs (Fig. S1) reveals that the two unknown variables are related and identifies a single local minimum for the solutions RMSD. In Fig. 7, we illustrate the model results as an envelope of those possible solutions (grey area). This envelope represents model solutions with the lowest RMSD values, defined by a maximum RMSD value of 2.2 wt.% of SOM, which ensures that all observed data fit within their standard error. The resulting model envelope ranges for  $k$  and SOM<sub>p</sub> are -0.03 to -0.042 yr<sup>-1</sup> and 13–21 wt.%, respectively. To further constrain our model results, we incorporated the reaction constant ( $k$ ) measured by the CO<sub>2</sub> fluxes in the short-term respiration experiments at 22°C (which is the mean annual temperature) into the model (Fig. 5a).- This reduces the model solution to a single degree of freedom, enabling a direct comparison with the two degrees of freedom model solution. Since the experimental  $k$  constant is calculated for the entire SOC content, it must be corrected if SOM<sub>p</sub>>0 according to the labile SOM [ $k_{\text{corrected}} = k_{\text{experimental}} / (1-P)$ ]. Assuming SOM<sub>p</sub> of 17 wt.% (the median result of the two degrees of freedom model run); the experimental  $k_{\text{corrected}}$  constant is  $-0.044 \pm 0.005$  yr<sup>-1</sup> (Fig. 7, green shade). Additionally, we run the model assuming no SOM<sub>p</sub> (SOM<sub>p</sub>=0) and as a result the experimental respiration  $k$  constant had no correction factor ( $k=-0.033 \pm 0.004$  yr<sup>-1</sup>; Fig. 7, blue line).

As Fig. 7 shows, using the experiment  $k$  as an input, and assuming that there is no stable fraction of SOM (SOM<sub>p</sub>=0), results in a predicted decomposition (blue line) which is much faster compared with the field data. This model run predicts a SOM content in 2023 of approximately 8%, which is inconsistent with the observed value of 21%. In contrast, the model run which used the experimental  $k$  and a SOM<sub>p</sub> value of 17% (green envelope), effectively

explains the field data and aligns closely with the RMSD envelope of the two degrees of freedom run (grey envelope). Therefore, the short-term respiration experiments can be used to bridge the gap with long-term (70- years) process of peat degradation, at least at this site. In summary, we suggest that peat degradation is a process lasting several decades under a warm [sub-tropical](#) climate, during which a fraction of persistent organic matter remains in the soil, even after long-term degradation given the prevailing land-use practices. Given the degradation duration of the labile SOM was of the order of 70 years, we estimate that the persistent SOM should be stable for several centuries given similar environmental conditions.



**Figure 7:** Mean soil organic matter content of the upper 1 meter of Hula peat since drainage in 1957, represented by markers. The fitted model result is shown by the grey envelope of possible solutions with RMSD lower than 2.2% SOM. The horizontal grey area and dashed line represent the persistent SOM pool range and median, respectively, predicted by the model. The experimental respiration decay result is represented by the green area (width is indicating the standard error), assuming 17% SOM<sub>p</sub>. The blue line shows the result of the experimental k with zero SOM<sub>p</sub>.

#### 530 [4.84.6](#) The model limitations


It is important to note some limitations: (1) [our](#) analysis assumes the drained section has not been significantly affected by the partial rewetting initiated in 1994. While this section technically remains drained after the rewetting

project, the increase in water content could influence soil respiration and potentially alter decomposition rates, (either enhancing or reducing them). (2) The respiration experiment might overestimate the actual rate since it is measured in optimal water holding capacity (60%), which does not typically occur in natural settings. We suggest that this might be the reason for the experimental  $k$  plot to project somewhat more intense decomposition compared to the observed data (Fig. 7, green shade). (3) Another potential limitation is the possibility of enhanced decomposition in the drained section due to the priming effect caused by root exudates from crops (Fontaine et al., 2004). However, Linkosalmi et al. (2012) found no evidence of a priming effect in their study of drained peatlands, suggesting that the introduction of labile carbon substrates, such as root exudates, did not significantly accelerate the decomposition of older, more persistent organic matter. Furthermore, in the model the top 30 cm of the profile werewas excluded, minimizing this potential effect. (4) It is acknowledged that additional SOM inputs and outputs that are not accounted for in the model may contribute to the overall trend. These may be root carbon contributions or DOC leaching for example. While the model, which is based on observed data, effectively captures the overall SOM losses and gains, this remains a limitation when directly comparing its results to respiration fluxes. Nevertheless, as pointed out above, we find the model in its current form - both satisfactory and informative.

#### 4.94.7 Persistent SOM in the Hula peat

While our model analysis suggests the presence of original persistent SOM, and is further confirms-supported it-by including the short-term experiments' decomposition rate constant, direct evidence for persistent SOM in the field-scale is still lacking. Recent studies have attempted to quantify persistent SOC using Rock-Eval analysis (Saenger et al., 20152014; Cécillon et al., 2021). Here, we discuss this knowledge regarding SOC persistence in the context of relation to our findings. Such persistent SOM could have been an original fraction of the peat-~~or~~, formed during the oxidation process, or resulted from a combination of both. Furthermore, the mechanism behind its persistence in the field (e.g., mineral association or other processes) in drained peat remains unknown.

In addition to our SOM model, the correlation between mean SOC and RI values across sections highlights the increase in thermal stability of the drained section compared to the pristine section. Figure 8a presents pyrograms of the Rock-Eval HC signal generated during the pyrolysis ramping. It shows that pristine samples (blue curves) exhibit two dominant HC peaks, corresponding to thermally labile and stable fractions (Sebag et al., 2016). In drained samples, these peaks persist but are diminished, with the high-temperature peak deceasedecreasing to a lesser extentless reduced. This suggests that during drainage, as bacterial-microbial activity accelerates organic matter decomposition, the thermally labile fraction (up to 400°C) is preferentially consumed, leaving behind residual thermally stable SOC, which contributing-contributes to higher RI values in drained samples.



**Figure 8:** (a) FID signal curves of hydrocarbons for pristine, rewetted, and drained samples from profile F, with colour scheme by section type and depth. The dashed line separates labile and stable hydrocarbons, following the method used for RI (Sebag et al., 2016). (b) RI values for the same samples, using the same colour scheme.

The mechanism and evaluation of persistent SOC ~~has~~ have been widely studied and debated (Cécillon et al., 2021). Barré et al. (2016), Gregorich et al. (2015), and Saenger et al. (20152014) demonstrate that while persistent SOM has diverse chemical compositions, it generally exhibits low energy content and hydrogen depletion, making it more refractory at high temperatures. This suggests a relation between biological availability of OM and thermal stability. This which aligns with our findings (Fig. 8).

There are generally three SOC pools that can be identified in soils: labile, particulate, and mineral-associated SOC, a classification widely accepted in soil science (Blume et al., 2016). Saenger et al. (20152014) proposed a thermal fractionation of those SOC pools using RE analysis, ~~identifying three SOC pools based on the pyrolysis HC curve: labile, particulate, and mineral associated SOC, a classification widely accepted in soil science.~~


Our thermal analysis partially aligns with thisese findingsmethod. The stable SOC HC peak (above 450°C), attributed by Saenger et al. (20152014) to mineral-associated SOC, appears in all sections. In pristine samples, it is as high as the labile SOC peak. However, mineral association with SOC is expected to be very low in pristine peat samples (Mirabito and Chambers, 2023). This demonstrates that attributing HC peaks to particulate and mineral-associated SOC presents a challenge in Hula's peat.

~~Additional~~An additional ~~new~~ approach to estimating persistent SOC in soils using Rock-Eval was proposed recently by Cécillon et al. (2021) through the PARTY<sub>SOC</sub> model (v2.0). The results indicate that, unlike the thermal indices we used above, the predicted stable SOM by the PARTY<sub>SOC</sub> model showed only a weak negative correlation with SOC content ( $R^2 = 0.25$ ; supplementary material 1.6). The PARTY<sub>SOC</sub> model estimates of the stable fraction ~~was were~~ multiplied by the SOC content, yielding a mean persistent SOC values of  $13.2 \pm 0.8$  wt.% in the pristine section and  $6.1 \pm 0.5$  wt.% in the drained section. However, persistent SOC is not expected to decline so fast during 66 years of peat decomposition, since it is defined as resilient for more than a century (Cécillon et al., 2021), which suggesting suggests a discrepancy in these results. Those discrepancies may ~~rise~~ arise as the PARTY<sub>SOC</sub> model was fitted to soils that are not as organic-rich as peat soils (Cécillon et al., 2021). Indeed, our findings show that the estimation of the ~~PARTY<sub>soe</sub>~~ PARTY<sub>SOC</sub>v2.0 model or pool separation by Saenger et al. (20152014) is inadequate inapplicable for the Hula's peat. ~~However~~ Nevertheless, while RI does not ~~deliberate allow for direct~~ quantify quantification of persistent SOC, it could serve as a practical and effective tool for analyzing and comparing degraded peat SOC, both in research contexts and for peatland management.

#### 4.104.8 Estimation of annual CO<sub>2</sub> emissions

Respiration models and global annual emission estimates typically incorporate factors such as peat SOM content, bulk density (BD), and WTL. However, these models often neglect the critical factor of time since drainage, which significantly influences emission rates (Hooijer et al., 2010; Tubiello et al., 2016). To emphasize the impact of drainage duration on carbon fluxes, we used a dual approach involving two distinct time frames and two independent flux estimation methods. These time frames represent the change in fluxes in the decades following peat drainage: (1) ~~compared~~ Comparing experimental respiration data from pristine and drained sections of the Hula's peat (sub-section 4.1). (2) Historical trends estimating the annual flux based on SOM decrease with time (sub-section 4.3, figure 7). This dual approach allows the comparison between the first decade following the exposure of the pristine peat by the onset of drainage, and the second scenario, representing the degraded peat after nearly 70 years of drainage. ~~representing the onset of drainage and nearly 70 years of degradation, respectively.~~ The average annual carbon emissions flux from the pristine and drained peat sections of the Hula at 22°C, derived from the short-term experiments (Flux per g<sub>f</sub> soil; Eq.1), are  $12 \pm 1$  and  $4.7 \pm 0.9$  mg C g<sub>f</sub>soil<sup>-1</sup> yr<sup>-1</sup>, respectively. An independent flux estimation was carried based on the two-pool box model. ~~Additionally~~ In this case, the average annual carbon loss rates ~~are~~ derived from the observation-based long-term model (Fig. 7, grey area), and was calculated as (SOC<sub>initial</sub>

- SOC (10yr)/10yr, where SOC is given in  $\text{mg C g}_{\text{soil}}^{-1}$ . Based on these calculations, the estimated fluxes are  $9 \pm 1$  and  $1.4 \pm 0.2 \text{ mg C g}_{\text{soil}}^{-1} \text{ yr}^{-1}$  for the 1<sup>st</sup> and (current) 7<sup>th</sup> decades, respectively. These two independent methods demonstrate the drastic difference in CO<sub>2</sub> fluxes between early and advanced stages of peat degradation. The comparison between these fluxes is presented in Fig. 9.



**Figure 9:** Average annual fluxes of CO<sub>2</sub> from peat soil, derived from short-term experiment (green) and long-term field model (black).

Next, these fluxes were converted to emissions estimates. This was done by incorporating additional data regarding the peat's bulk density, its shrinkage properties, and assuming a characteristic peat column depth of 100 cm as a baseline for calculation (supplementary material 1.10). This calculation was carried to represent the two timeframes mentioned above. For the first decade following the drainage, we assumed pristine peat exposure, incorporated data experimental flux data of pristine peat at 22°C (mean annual average), bulk density of  $0.22 \pm 0.07 \text{ g cm}^{-3}$  (Dansberg & Neuman, 1977), yielded an estimated emission of  $94 \pm 31 \text{ tons CO}_2 \text{ ha}^{-1} \text{ yr}^{-1}$ . This is in good agreement with the IPCC and global models estimated CO<sub>2</sub> emissions from drained tropical peatlands at  $73 \pm 14 \text{ tons CO}_2 \text{ ha}^{-1} \text{ yr}^{-1}$  (Hiraishi et al., 2013; Tubiello et al., 2016). This similarity may be attributed to the Hula valley's mean annual temperature ( $\sim 20^\circ\text{C}$ ), which falls within the classification of tropical soil respiration zone (Bond-Lamberty et al., 2024).

625 ~~For the 7<sup>th</sup> decade, representing the degraded peat layer after 66 years of drainage, we used experimental flux of degraded peat at 22°C, increased bulk density of  $0.52 \pm 0.04 \text{ g cm}^{-3}$  (Orlov-Levin and Meron, 2005), and assumed  $71 \pm 7\%$  soil shrinkage (after Yaari-Cohen et al., 1971b). This yielded an estimated emission of  $26 \pm 11 \text{ tons CO}_2 \text{ ha}^{-1} \text{ yr}^{-1}$ . Thus, the expected emissions from the upper degraded peat have decrease to  $\sim 30\%$  of the original estimated value.~~

630 ~~The IPCC and global models estimate CO<sub>2</sub> emissions from drained tropical peatlands at  $73 \pm 14 \text{ tons CO}_2 \text{ ha}^{-1} \text{ yr}^{-1}$  (Hiraishi et al., 2013; Tubiello et al., 2016). This sharp decline in ~~indicates that most CO<sub>2</sub> emissions, which occurs within the first decades following drainage, implying~~ a narrow window for effective restoration. The degradation trajectory observed in the Hula Valley may therefore be relevant to other peatlands in ~~subtropical and warm climatic regions subjected to long-term drainage.~~~~

## 5 Conclusions

Our observations show that active land management that includes partial rewetting of ~~the~~ once-drained peat column significantly mitigates CO<sub>2</sub> emissions and organic carbon loss, demonstrating its effectiveness even after prolonged drainage in a warm sub-tropical climate. The respiration results highlight the vulnerability of the rewetted peat horizon to re-exposure to air, showing the potential of enhanced fluxes, twice as high relative to the drained and pristine sections (per SOC). We also found a peak in pyrite concentration beneath the recent WTL, and sulfate accumulation in the drained section, which indicates sulfur recycling and re-establishment of anaerobic conditions. Short-term respiration experiments reveal that the drained and pristine sections share similar first-order reaction rates ( $k$ ) for SOM decomposition, and that these experiments successfully describe the observed long-term SOM decomposition in the ~~warm climate~~ Hula's peat. Model analysis, supported by those experimental results, indicates that approximately 13–21 ~~wt.~~% of the  $\sim 68$  ~~wt.~~% initial SOM persists as a stable fraction in the drained peat, emphasizing the role of the persistent organic pool in stabilizing carbon stocks. However, the mechanism of persistence SOM formation in degraded peatlands requires further investigation. The use of a thermal analysis (Rock-Eval) to characterize the OM properties, coupled with short-term aerobic respiration experiments, is shown here to be useful in bridging the gap to the field scale, to the modelling of long-term decomposition of peatlands, and to estimating their CO<sub>2</sub> emissions. The timeline of drained peat SOM decomposition shows that the first few decades are crucial, as SOM declines rapidly, resulting in high CO<sub>2</sub> emissions. Future peatland restoration projects, especially in sub-tropical and warm climates, can use this site as an analogue to make more informed decisions ~~for~~ theregarding priorities for sites rewetting priorities.

645

650

## 6 Acknowledgments

We would like to express our gratitude to the Hula Valley Monitoring Program, operated by ~~the~~MIGAL Institute, Tel-Hai College, and the Israel Water Survey, for their valuable collaboration and the availability of their data  
655 online. We also thank ~~to~~ [Dr. Oren Reichman and Prof. Michael Litaor](#) ~~and Oren Reichman~~ for providing access to their field data and helpful information. ~~We would like to~~ Our acknowledge-acknowledgments to Tal Weiner, Roe Katzir and Itay Eyal for their assistance ~~in our~~ during field and laboratory work, and the helpful comments of two anonymous reviewers. This research was partly supported by the Israel Science Foundation grant #300/23, and the Chief Scientist of the Ministry of Agriculture and Food Security, Israel, grant #20-03-0081.

660

Code availability

The code for the peat SOM Model can be accessed in this DOI link: <https://doi.org/10.5281/zenodo.15635973> .

## 7 Data availability

665 Data of Rock-Eval 7 analysis and respiration experiments can be accessed in this DOI link:

<https://doi.org/10.5281/zenodo.15635973> .

## 8 Competing interests

The authors declare that they have no conflict of interest.

## 9 Sample availability

670 Samples are stored in the Geological survey of Israel and can be accessed if needed, please reach the author for more information.

## 10 Author contribution

675 The Investigation was done by GS. RG, AA, and YR conceptualized the research, were in charge on the methodology, and supervised GS. GS wrote the original draft, and AA, YR and RG reviewed and edited it.

## #11 Disclaimer

OpenAI was used solely to improve the grammar and phrasing of the text. The authors declare that no other artificial intelligence or machine learning tools were employed in this study.

## #12 References

680

Aboussou, A. New Rock-Eval method for Pyritic and Organic Sulphur quantification: Application to study Organic Matter preservation in Jurassic sediments [(Doctoral dissertation, Sorbonne Université)], <https://doi.org/10.13140/RG.2.2.27845.68324>, 2018.

685 Angert, A., Yakir, D., Rodeghiero, M., Preisler, Y., Davidson, E. A., and Weiner, T. Using O<sub>2</sub> to study the relationships between soil CO<sub>2</sub> efflux and soil respiration. *Biogeosciences*, 12(7), 2089., <https://doi.org/10.5194/bg-12-2089-2015>, 2015.

Bader, C., Müller, M., Schulin, R., and Leifeld, J. Peat decomposability in managed organic soils in relation to land use, organic matter composition and temperature Copernicus GmbH., <https://doi.org/10.5194/bg-15-703-2018>, 2018.

690 Barré, P., Eglin, T., Christensen, B. T., Ciais, P., Houot, S., Kätterer, T., . . . Chenu, C. Quantifying and isolating stable soil organic carbon using long-term bare fallow experiments. *Biogeosciences*, 7(11), 3839., <https://doi.org/10.5194/bg-7-3839-2010>, 2010.

Barré, P., Plante, A. F., Cécillon, L., Lutfalla, S., Baudin, F., Bernard, S., . . . Chenu, C. The energetic and chemical signatures of persistent soil organic matter. *Biogeochemistry*, 130(1-2), 1., <https://doi.org/10.1007/s10533-016-0246-0>, 2016.

695 Behar, F., Beaumont, V. D. E. B., and Penteadó, H. D. B. Rock-Eval 6 technology: performances and developments. *Oil and Gas Science and Technology*, 56(2), 111-134., <https://doi.org/10.2516/ogst:2001013>, 2001.

Bein, A. Early evolution and transformation of organic matter in the active continental Jordan Rift Valley. *Organic geochemistry*, 10(4-6), 751-757., [https://doi.org/10.1016/S0146-6380\(86\)80012-2](https://doi.org/10.1016/S0146-6380(86)80012-2), 1986.

Bein, A. Hula valley report by the Geological Survey of Israel. [(in Hebrew)]., 1967.

700 Bein, A., & Horowitz, A. Papyrus-A a historic newcomer to the Hula Valley Israel?. *Review of palaeobotany and palynology*, 47(1-2), 89-95. [https://doi.org/10.1016/0034-6667\(86\)90007-2](https://doi.org/10.1016/0034-6667(86)90007-2), 1986.

Blume, H. P., Brümmer, G. W., Fleige, H., Horn, R., Kandeler, E., Kögel-Knabner, I., ... and Wilke, B. M. Soil organic matter. *Scheffer/Schachtschabel Soil Science*, 55-86., [https://doi.org/10.1007/978-3-662-49880-4\\_3](https://doi.org/10.1007/978-3-662-49880-4_3), 2016.

705 Bond-lamberty, B., Ballantyne, A., Berryman, E., Fluet-chouinard, E., Jian, J., Morris, K. A., . . . Vargas, R. Twenty years of progress, challenges, and opportunities in measuring and understanding soil respiration American Geophysical Union (AGU)., <https://doi.org/10.1029/2023jg007637>, 2024.

- Brown C., Boyd D.S., Sjögersten S., and Vane C.H.. Detecting tropical peatland degradation: Combining remote sensing and organic geochemistry. *PLoS ONE* 18(3): e0280187. <https://doi.org/10.1371/journal.pone.0280187>, 2023.
- Cécillon, L., Baudin, F., Chenu, C., Christensen, B. T., Franko, U., Houot, S., ... and Barré, P. Partitioning soil organic carbon into its centennially stable and active fractions with machine-learning models based on Rock-Eval® thermal analysis (PARTY SOC v2. 0 and PARTY SOC v2. 0 EU). *Geoscientific Model Development*, 14(6), 3879-3898., <https://doi.org/10.5194/gmd-14-3879-2021>, 2021.
- Cécillon, L., Baudin, F., Chenu, C., Houot, S., Jolivet, R., Kätterer, T., Lutfalla, S., Macdonald, A., van Oort, F., Plante, A. F., Savignac, F., Soucémarianadin, L. N., and Barré, P.: A model based on Rock-Eval thermal analysis to quantify the size of the centennially persistent organic carbon pool in temperate soils, *Biogeosciences*, 15, 2835–2849, <https://doi.org/10.5194/bg-15-2835-2018>, 2018.
- Chenu, C., and Cosentino, D. Microbial regulation of soil structural dynamics. In *The architecture and biology of soils: life in inner space* (pp. 37-70). Wallingford UK: CABI., <https://doi.org/10.1079/9781845935320.0037>, 2011.
- Clivot, H., Mouny, J. C., Duparque, A., Dinh, J. L., Denoroy, P., Houot, S., ... and Mary, B. Modeling soil organic carbon evolution in long-term arable experiments with AMG model. *Environmental modelling and software*, 118, 99-113., <https://doi.org/10.1016/j.envsoft.2019.04.004>, 2019.
- Clymo, R. S., Turunen, J., and Tolonen, K. Carbon accumulation in peatland. *Oikos*, 368-388., <https://doi.org/10.2307/3547086>, 1998.
- Cohen-Sadon, H., Amrani, A., Feinstein, S., and Rosenberg, Y. O. A new empirical approach for rapid quantification of organic and pyritic sulfur in sedimentary rocks using the Rock-Eval 7S. *Organic Geochemistry*, 166, 104350., <https://doi.org/10.1016/j.orggeochem.2022.104350>, 2022.
- Convention on Wetlands. Global guidelines for peatland rewetting and restoration. Ramsar technical report no. 11. Gland, Switzerland: Secretariat of the Convention on Wetlands., 2021.
- Cowgill, U. M. The waters of Merom: A study of Lake Huleh. I. Introduction and general stratigraphy of a 54 m core. *Archiv Fur Hydrobiologie*, Vol 66, No (3), P 249-272, 1969. 9 FIG, 1 TAB, 28 REF., <https://doi.org/10.1127/archiv-hydrobiol/66/1969/249>, 1969.
- Dasberg, S. and Neuman, S.P. Peat hydrology in the Hula Basin, Israel: I. Properties of peat. *J. Hydrol.*, 32: 219--239, 1977.
- Delahaie, A. A., Barré, P., Baudin, F., Arrouays, D., Bispo, A., Boulonne, L., Chenu, C., Jolivet, C., Martin, M. P., Ratié, C., Saby, N. P. A., Savignac, F., and Cécillon, L.: Elemental stoichiometry and Rock-Eval® thermal stability of organic matter in French topsoils, *SOIL*, 9, 209–229, <https://doi.org/10.5194/soil-9-209-2023>, 2023.
- Dimentman Ch., Bromley H.J., and Por F.D.. Lake Hula, Reconstruction of the Fauna and Hydrobiology of a Lost Lake. *The Israel Academy of Science and Humanities*, Jerusalem, 170 pp, 1992.
- Disnar, J.R., Jacob J., Morched-Issa M., Lottier N., and Arnaud F. Assessment of peat quality by molecular and bulk geochemical analysis: Application to the Holocene record of the Chautagne marsh (Haute Savoie, France). *Chemical geology*, 254(1-2):101-12, 2008.

- 740 Fontaine, S., Bardoux, G., Abbadie, L., and Mariotti, A. Carbon input to soil may decrease soil carbon content. *Ecology letters*, 7(4), 314-320., <https://doi.org/10.1111/j.1461-0248.2004.00579.x>, 2004.
- Freeman, C., Ostle, N. J., Fenner, N. and Kang, H. A regulatory role for phenol oxidase during decomposition in peatlands. *Soil Biol. Biochem.* 36, 1663--1667., <https://doi.org/10.1016/j.soilbio.2004.07.012>, 2004.
- Gregorich, E. G., Gillespie, A. W., Beare, M. H., Curtin, D., Sanei, H., & Yanni, S. F. Evaluating biodegradability of soil  
745 organic matter by its thermal stability and chemical composition. *Soil Biology and Biochemistry*, 91, 182-191, 2015.
- Günther, A., Barthelmes, A., Huth, V., Joosten, H., Jurasinski, G., Koebisch, F., & Couwenberg, J. Prompt rewetting of drained peatlands reduces climate warming despite methane emissions. *Nature communications*, 11(1), 1644, 2020.
- Hambright, K. D., and Zohary, T. Lakes Hula and Agmon: destruction and creation of wetland ecosystems in northern Israel. *Wetlands Ecology and Management*, 6, 83-89., <https://doi.org/10.1023/A:1008494705994>, 1998.
- 750 Hicks Pries, C., Angert, A., Castanha, C., Hilman, B., and Torn, M. S. Using respiration quotients to track changing sources of soil respiration seasonally and with experimental warming. *Biogeosciences*, 17(12), 3045., <https://doi.org/10.5194/bg-17-3045-2020>, 2020.
- Hilasvuori, E., Akujärvi, A., Fritze, H., Karhu, K., Laiho, R., Mäkiranta, P., . . . Liski, J. Temperature sensitivity of decomposition in a peat profile. *Soil Biology and Biochemistry*, 67, 47., <https://doi.org/10.1016/j.soilbio.2013.08.009>, 2013.
- 755 Hilman, B., and Angert, A. Measuring the ratio of CO<sub>2</sub>efflux to O<sub>2</sub>influx in tree stem respiration. *Tree Physiology*, <https://doi.org/10.1093/treephys/tpw057>, 2016.
- Hilman, B., Weiner, T., Haran, T., Masiello, C. A., Gao, X., and Angert, A. The apparent respiratory quotient of soils and tree stems and the processes that control it *American Geophysical Union (AGU)*., <https://doi.org/10.1029/2021jg006676>, 2022.
- Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M., and Troxler, T. G. 2013 supplement to the  
760 2006 IPCC guidelines for national greenhouse gas inventories: Wetlands. IPCC, Switzerland., 2014.
- Hooijer, A., Page, S., Canadell, J. G., Silvius, M., Kwadijk, J., Wösten, H., and Jauhiainen, J. Current and future CO<sub>2</sub> emissions from drained peatlands in southeast Asia. *Biogeosciences*, 7(5), 1505--1514., <https://doi.org/10.5194/bg-7-1505-2010>, 2010.
- Jones, R. F.. Report of the Percy Sladen Expedition to Lake Huleh: A Contribution to the Study of the Fresh Waters of Palestine: The Plant Ecology of the District. *Journal of Ecology*, 28(2), 357–376. <https://doi.org/10.2307/2256234>, 1940.
- 765 Joosten, H., and Clarke, D. Wise use of mires and peatlands. *International mire conservation group and international peat society*, 304., 2002.
- Kleber, M., Eusterhues, K., Keiluweit, M., Mikutta, C., Mikutta, R., & Nico, P. S.. Mineral–organic associations: formation, properties, and relevance in soil environments. *Advances in agronomy*, 130, 1-140. 2015.

- 770 Kleber, M., Eusterhues, K., Keiluweit, M., Mikutta, C., Mikutta, R., and Nico, P. S. Chapter One---Mineral--Organic associations: Formation, properties, and relevance in soil environments. *Advances in Agronomy*, 130, 1., <https://doi.org/10.1016/bs.agron.2014.10.005>, 2015.
- Kleinen, T., Brovkin, V., and Schuldt, R. J. A dynamic model of wetland extent and peat accumulation: results for the Holocene. *Biogeosciences*, 9(1), 235-248., <https://doi.org/10.5194/bg-9-235-2012>, 2012.
- 775 Könönen, M., Jauhiainen, J. R., Laiho, R. P., Spetz, P. K., Kusin, K. S., Limin, S. H., and Vasander, H. Land use increases the recalcitrance of tropical peat. *Wetlands Ecology and Management*, 24(6), 717., <https://doi.org/10.1007/s11273-016-9498-7>, 2016.
- Leifeld, J., and Menichetti, L. The underappreciated potential of peatlands in global climate change mitigation strategies. *Nature Communications*, 9(1), <https://doi.org/10.1038/s41467-018-03406-6>, 2018.
- 780 Leifeld, J., Steffens, M., and Galego-sala, A. Sensitivity of peatland carbon loss to organic matter quality American Geophysical Union (AGU)., <https://doi.org/10.1029/2012gl051856>, 2012.
- Liang, C., Schimel, J. P., and Jastrow, J. D. The importance of anabolism in microbial control over soil carbon storage. *Nature Microbiology*, 2(8), <https://doi.org/10.1038/nmicrobiol.2017.105>, 2017.
- 785 Linkosalmi, M., Biasi, C., Pumpanen, J., Heinonsalo, J., Lindén, A., Minkkinen, K., ... and Lohila, A. Can the contrasting carbon balances of forestry-drained peatlands be explained by different priming effects?. In *International Peat Congress* (pp. Article-No). International Peat Society., 2012.
- Litaor, M. I., Reichmann, O., and Shenker, M. Genesis, classification and human modification of peat and mineral-organic soils, Hula Valley, Israel. *Mires and Peat*, 9., <https://doi.org/10.19189/MaP.2011.OMB.116>, 2011.
- 790 Litaor, M. I., Reichmann, O., Belzer, M., Auerswald, K., Nishri, A., and Shenker, M. Spatial analysis of phosphorus sorption capacity in a semiarid altered wetland. *Journal of Environmental Quality*, 32(1), 335-343., <https://doi.org/10.2134/jeq2003.3350>, 2003.
- Lützow, M. v., Kögel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., and Flessa, H. Stabilization of organic matter in temperate soils: Mechanisms and their relevance under different soil conditions - a review. *European Journal of Soil Science*, 57(4), 426--445., <https://doi.org/10.1111/j.1365-2389.2006.00809.x>, 2006.
- 795 Manzoni, S. and Cotrufo, M. F. Mechanisms of soil organic carbon and nitrogen stabilization in mineral-associated organic matter – insights from modeling in phase space, *Biogeosciences*, 21, 4077–4098, <https://doi.org/10.5194/bg-21-4077-2024>, 2024.
- Manzoni, S., and Porporato, A. Soil carbon and nitrogen mineralization: Theory and models across scales. *Soil Biology and Biochemistry*, 41(7), 1355-1379., <https://doi.org/10.1016/j.soilbio.2009.02.031>, 2009.
- 800 Manzoni, S., Piñeiro, G., Jackson, R. B., Jobbágy, E. G., Kim, J. H., and Porporato, A. Analytical models of soil and litter decomposition: Solutions for mass loss and time-dependent decay rates Elsevier BV., <https://doi.org/10.1016/j.soilbio.2012.02.029>, 2012.

- Marchand C., Lallier-Vergès E., Disnar J.R., and Kéravis D. Organic carbon sources and transformations in mangrove sediments: a Rock-Eval pyrolysis approach. *Organic Geochemistry*,39(4):408-21, 2008.
- 805 Marish S. The Hula: A soil survey report (Report, in Hebrew). Israel Ministry of Agriculture / Water Commissioner, Department of Soil Conservation and Drainage, Lake Kinneret Authority., 1986.
- Markel, D., Sass, E., Lazar, B., and Bein, A. Biogeochemical evolution of a sulfur-iron rich aquatic system in a reflooded wetland environment (Lake Agmon, northern Israel). *Wetlands Ecology and Management*, 6, 103-120., <https://doi.org/10.1023/A:1008414423821>, 1998.
- 810 Masiello, C. A., Gallagher, M. E., Randerson, J. T., Deco, R. M., and Chadwick, O. A. Evaluating two experimental approaches for measuring ecosystem carbon oxidation state and oxidative ratio. *Journal of Geophysical Research: Biogeosciences*, 113(G3), <https://doi.org/10.1029/2007jg000534>, 2008.
- MIGAL - (Ministry of Galilee Research Institute). Data Reports reports by stations; Hava 1 station. Mop-Zafon website. ., 2023.
- 815 Mirabito, A. J., and Chambers, L. G. Quantifying mineral-associated organic matter in wetlands as an indicator of the degree of soil carbon protection Elsevier BV., <https://doi.org/10.1016/j.geoderma.2023.116327>, 2023.
- Moore, P. D. The ecology of peat-forming processes: A review Elsevier BV., [https://doi.org/10.1016/0166-5162\(89\)90048-7](https://doi.org/10.1016/0166-5162(89)90048-7), 1989.. *International Journal of Coal Geology*, 12(1-4), 89-103. [https://doi.org/10.1016/0166-5162\(89\)90048-7](https://doi.org/10.1016/0166-5162(89)90048-7), 1989.
- Moore, T. R., and Knowles, R. The influence of water table levels on methane and carbon dioxide emissions from peatland soils. *Canadian Journal of Soil Science*, 69(1), 33-38., <https://doi.org/10.4141/cjss89-004>, 1989.
- Nabuurs, G., Mrabet, R., Abu, A., Bustamante, M., Clark, H., Aoki, L., . . . Emmet-Booth, J. P. Agriculture, forestry and other land uses (AFOLU) Cambridge University Press., <https://doi.org/10.1017/9781009157926.009>, 2023.
- Nelson, D. W., & Sommers, L. E. Total carbon, organic carbon, and organic matter. *Methods of soil analysis: Part 3 Chemical methods*, 5, 961-1010, 1996.
- 825 Nyberg, M., Black, T. A., Ketler, R., Lee, S., . . . , Johnson, M., Merkens, M., . . . Knox, S. H. Impacts of active versus passive Re-Wetting on the carbon balance of a previously drained bog American Geophysical Union (AGU)., <https://doi.org/10.1029/2022jg006881>, 2022.
- Orlov-Levin, V., and Meron, M. Soil salinization in Hula Valley crops. Report [in Hebrew]. Israel Ministry of Agriculture., 2005.
- 830 Page, S. E., and Baird, A. J. Peatlands and global change: Response and resilience *Annual Reviews*., <https://doi.org/10.1146/annurev-environ-110615-085520>, 2016.
- Ravikovitch, S. The Peat soils and enriched organic matter soils of the Hula Valley. *Agric. Rep*, 1, 23-54., 1945.
- Rey, A., Petsikos, C., Jarvis, P. G., & Grace, J. Effect of temperature and moisture on rates of carbon mineralization in a Mediterranean oak forest soil under controlled and field conditions. *European Journal of Soil Science*, 56(5), 589-599, 2005.

- 835 Rixen, T., Baum, A., Wit, F., and Samiaji, J. Carbon leaching from tropical peat soils and consequences for carbon balances. *Frontiers in Earth Science*, 4, <https://doi.org/10.3389/feart.2016.00074>, 2016.
- Roe, S., Streck, C., Beach, R., Busch, J., Chapman, M., Daioglou, V., . . . Lawrence, D. Land-based measures to mitigate climate change: Potential and feasibility by country Wiley., <https://doi.org/10.1111/gcb.15873>, 2021.
- Rowley, M. C., Grand, S., and Verrecchia, É P. Calcium-mediated stabilisation of soil organic carbon Springer Science and  
840 Business Media LLC., <https://doi.org/10.1007/s10533-017-0410-1>, 2017.
- Saenger, A., Cécillon, L., Poulencard, J., Bureau, F., De Daniéli, S., Gonzalez, J., and Brun, J. Surveying the carbon pools of mountain soils: A comparison of physical fractionation and rock-aval pyrolysis. *Geoderma*, 241-242, 279., <https://doi.org/10.1016/j.geoderma.2014.12.001>, 2014.
- Saenger, A., Cécillon, L., Sebag, D., & Brun, J. J. Soil organic carbon quantity, chemistry and thermal stability in a  
845 mountainous landscape: A Rock-Eval pyrolysis survey. *Organic Geochemistry*, 54, 101-114., <https://doi.org/10.1016/j.orggeochem.2012.10.008>, 2013.
- Scanlon, D., and Moore, T. Carbon dioxide production from peatland soil profiles: the influence of temperature, oxic/anoxic conditions and substrate. *Soil Science*, 165(2), 153-160., <https://doi.org/10.1097/00010694-200002000-00007>, 2000.
- Schallinger, K.M. Investigations of the Agricultural Problems Concerning Peat Soils of the Hula Valley. Report number 673,  
850 Volcani Institute for Agricultural Research, Bet Dagan, Israel, 1-33 [in Hebrew], 1970.
- Sebag, D., Verrecchia, E. P., Cécillon, L., Adatte, T., Albrecht, R., Aubert, M., . . . Trombino, L. Dynamics of soil organic matter based on new rock-aval indices. *Geoderma*, 284, 185., <https://doi.org/10.1016/j.geoderma.2016.08.025>, 2016.
- Six, J., Conant, R. T., Paul, E. A., and Paustian, K. Stabilization mechanisms of soil organic matter: implications for C-saturation of soils. *Plant and soil*, 241, 155-176., <https://doi.org/10.1023/A:1016125726789>, 2002.
- 855 Tsipris, J., and Meron, M. Climatic and hydrological aspects of the Hula restoration project. *Wetlands Ecology and Management*, 6(2), 91-101., <https://doi.org/10.1023/A:1008424518122>, 1998.
- Tsipris, J., and Meron, M., Orlov-Levin, V. Meteorological and hydrological regime of the Hula valley in 2008-2021. In: Pearlson O., Klein D., and Kaplan D. (Eds.), *MIGAL -- Galilee Research Institution; Hula monitoring project report*. MIGAL -- Galilee Research Institution [in Hebrew], 2022.
- 860 Tubiello, F. N., Biancalani, R., Salvatore, M., Rossi, S., and Conchedda, G. A worldwide assessment of greenhouse gas emissions from drained organic soils MDPI AG., <https://doi.org/10.3390/su8040371>, 2016.
- Wang, Y., Wang, H., He, J. S., and Feng, X. Iron-mediated soil carbon response to water-table decline in an alpine wetland. *Nature communications*, 8(1), 15972., <https://doi.org/10.1038/ncomms15972>, 2017.
- Wilson, D., Environment, D. B., Change, C., Murdiyarso, D., Page, S. E., Blain, D., . . . Tuittila, E. -. Greenhouse gas emission  
865 factors associated with rewetting of organic soils., <https://doi.org/10.19189/MaP.2016.OMB.222>, 2016.
- Yaari-Cohen, G., Paldi, H., Moses, G. and Harpaz, A. A survey of the peat soils in the Hula- valley, Israel, part I. Ministry of agriculture, department of soil conservation and drainage, Kyriat Shemona, israel, 112 pp. [in Hebrew]. (1971a).

Yaari-Cohen, G., Paldi, C., Moses, G. and Harpaz, A. A survey of the peat soils in the Hula- valley, Israel, part II (Appendices). Ministry of agriculture, department of soil conservation and drainage, Kyriat Shemona, israel, 78 pp. [in Hebrew]. (1971b).

870 Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J. Global peatland dynamics since the last glacial maximum. *Geophysical Research Letters*, 37(13), <https://doi.org/10.1029/2010gl043584>, 2010.

Zou, J., Ziegler, A. D., Chen, D., Mcnicol, G., Ciais, P., Jiang, X., . . . Zeng, Z. Rewetting global wetlands effectively reduces major greenhouse gas emissions. *Nature Geoscience*, 15(8), 627., <https://doi.org/10.1038/s41561-022-00989-0>, 2022.