

East Greenland Ice Sheet retreat history from Scoresby Sund and Storstrømmen Glacier during the last deglaciation

Jacob T.H. Anderson¹, Nicolás E. Young¹, Allie Balter-Kennedy¹, Karlee K. Prince², Caleb K. Walcott-George^{2,3}, Brandon L. Graham⁴, Joanna Charton¹, Jason P. Briner², and Joerg M. Schaefer¹

¹Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA

²Department of Earth Sciences, University at Buffalo, Buffalo, NY, USA

³University of Kentucky, Department of Earth and Environmental Sciences, Lexington, KY, USA

⁴United States Geological Survey, Florence Bascom Geoscience Center, Reston, VA, USA

Correspondence to: Jacob Anderson (janderson@ldeo.columbia.edu)

Abstract

The lack of geological constraints on past ice-sheet change in marine-based sectors of the Greenland Ice Sheet (GrIS) following the Last Glacial Maximum limits our ability to assess (1) the drivers of ice-sheet change, and (2) the performance of ice-sheet models that are benchmarked against the paleo-record of GrIS change. Here, we provide new in situ ¹⁰Be surface exposure chronologies of ice-sheet margin retreat from the outer Scoresby Sund and Storstrømmen Glacier regions in eastern and northeastern Greenland, respectively. Ice retreated from Rathbone Island, east of Scoresby Sund, by ~14.1 ka, recording some of the earliest documentations of terrestrial deglaciation in Greenland. The mouth of Scoresby Sund deglaciated by ~13.2 ka, and retreated at an average rate of ~43 m/yr between 13.2 ka and 9.7 ka. Storstrømmen Glacier retreated from the outer coast to within ~3 km of the modern ice margin between ~12.7 ka and 8.6 ka at an average rate of ~28 m/yr. Retreat then slowed or reached a stillstand as ice retreated ~3 km between ~8.6 ka to the modern ice margin at ~8.0 ka. These retreat rates are consistent with late glacial and Holocene estimates for marine-terminating outlet glaciers across East Greenland, and comparable to modern retreat rates observed at the largest ice streams in northeastern, and northwestern Greenland.


1 Introduction

The rate of Greenland Ice Sheet (GrIS) mass loss and contribution to sea-level rise has increased over recent decades (Khan et al., 2022; The IMBIE Team, 2020). Projections under varying warming scenarios suggest that GrIS contributions to global mean sea level by 2100 range from -0.02–0.15 m to 0.05–0.23 m under SSP1-2.6 and SSP5-8.5 scenarios, respectively (IPCC, 2021). To place these contemporary short-term ice sheet dynamics in context, we examine retreat patterns of the GrIS since the Last Glacial Maximum (LGM; 26.5 – 19 ka; Clark et al., 2009).

Previous research has provided robust constraints on the timing and pattern of deglaciation in southwestern Greenland (e.g., Briner et al., 2020; Larsen et al., 2014; Lesnek et al., 2020; Levy et al., 2012; Young et al., 2011, 2020, 2021). Relying partly on these detailed constraints, Briner et al. (2020) suggested rates of southwestern GrIS mass loss, and perhaps the entire GrIS, over the twenty-first century are predicted to exceed rates experienced anytime in the last 12,000 years. Comparable geologic records from eastern (e.g., Larsen et al., 2020; 2022; Levy et al., 2016; Roberts et al., 2024) and fewer studies from northern (e.g., Kelly & Bennike, 1992; O'Regan et al., 2021) sectors of Greenland, have been used to constrain the timing and rates of ice sheet margin retreat during the last deglaciation. In some regions across East and North Greenland, however, data gaps hinder our ability to reconstruct the pattern and timing of GrIS retreat, and assess the accuracy of ice sheet models. This is particularly important in northeast Greenland, where major marine-terminating ice streams drain large portions of the GrIS. For example, the Northeast Greenland Ice Stream (NEGIS) is one of the largest drainage systems in Greenland, accounting for ~16% of the total GrIS discharge through three major marine-terminating outlet glaciers: Nioghalvfjærdsfjord Glacier (or 79N Glacier), Zachariae Isstrøm, and Storstrømmen Glacier (Hvidberg et al., 2020). Collectively, these glaciers contain 1.4 m of sea level equivalent (An et al., 2021; Rignot et al., 2022).

During the LGM, the East GrIS extended onto the continental shelf edge, with significant contributions of ice from expanded marine terminating ice streams such as NEGIS and in the Scoresby Sund region (Fig. 1). Marine geological data and geophysical surveys identifying glacial lineations and recessional moraines have been used to reconstruct patterns of ice sheet retreat along the continental shelf during the last deglaciation (Arndt et al., 2015, 2017; Evans et al., 2009; López-Quirós et al., 2024; Ó Cofaigh et al., 2004, 2025; Stein et al., 1996). On the northeast Greenland shelf, radiocarbon dates from marine sediments show that ice sheet retreat was underway on the shelf edge by 21.6 cal ka BP at Norske Trough, and by 19 cal ka BP at Westwind Trough (Ó Cofaigh et al., 2025). Ice retreated to the mid-shelf before 18.6 cal ka BP (Ó Cofaigh et al., 2025), and that ice retreated from the inner continental shelf east of NEGIS between 13.4 – 12.5 cal ka BP, which was primarily driven by an influx of relatively warm Atlantic water (Davies et al., 2022; Hansen et al., 2022). The NEGIS retreated from the outer coast to the present ice margin between 11.7 and ~9.0 ka (Larsen et al., 2018; Roberts et al., 2024).

Despite recent advances understanding the retreat history from the outer coastline to the modern ice margin of marine-terminating glaciers along eastern Greenland (e.g., Håkansson et al., 2007a; Larsen et al., 2020, 2022; Levy et al., 2016a), there are still major gaps in some regions (i.e., a lack of consistent ages or an absence of dated material). To record ice margin change, glacial landforms (such as erratic boulders and glacially polished bedrock) can be dated via in situ cosmogenic nuclides produced in rock surfaces (e.g. Ivy-Ochs and Briner, 2014; Lal, 1991; Nishiizumi et al., 1989). Here, we build on previous work studying the retreat history of marine-terminating glaciers in East Greenland, and present in situ cosmogenic ^{10}Be exposure ages from (i) the outer coast of Scoresby Sund, which improve age constraints at the mouth of the largest fjord system in Greenland, and (ii) near the modern ice margin of Storstrømmen Glacier, where few existing ages constrain East GrIS retreat. We integrate these data to calculate rates of ice margin retreat, and reconstruct a detailed East GrIS retreat pattern that we connect to potential mechanisms driving these GrIS changes.



75 **Figure 1.** Study area of East Greenland. The maximum LGM extent is marked by the black dashed line,
derived from Leger et al. (2024 and references therein). The Milne Land stade moraines are marked by the
blue line (Funder et al., 2011; Larsen et al., 2022). The ice margin at ~11 ka and 9 ka is marked by the red
and green dashed lines, derived from Larsen et al. (2020). Previously reported ^{10}Be ages from outer coast,
intermediate and ice margin sites (Dyke et al., 2014; Håkansson et al., 2007a; Hughes et al., 2012; Larsen
80 et al., 2022; Levy et al., 2016). Radiocarbon ages are listed in Table 1. The black rectangles show the
locations of Scoresby Sund (Fig. 3) and Dove Bugt region (Fig. 4). Basemap derived from IBCAO
bathymetry (Jakobsson et al., 2024).



Figure 2. Representative boulder sample sites from outer Scoresby Sund (a, b, c), and the unnamed island near the terminus of Storstrømmen Glacier (d, e, f). (a) Sample 22-GRO-101, Uunarteq at the northern mouth of Scoresby Sund. (b) Sample 22GRO-131, Kap Brewster (Kangikajjip Appalia) at the southern mouth of Scoresby Sund. (c) Sample 22GRO-142, Rathbone Island, ~3 km to the east of the main coastline of Liverpool Land. (d) Sample 23DMH-29, from the prominent M1 moraine on the island at the terminus

of Storstrømmen Glacier. (e) Sample 23DMH-33, adjacent to the historical moraine and modern ice margin.
90 (f) Sample 23DMH-20, outboard of the moraine.

2 Study sites and geological setting

The eastern margin of Greenland, from 79 N Glacier in the north, to Scoresby Sund in the south, comprises an extensive system of marine-terminating outlet glaciers, ice-free valleys, and fjords at lower elevations
95 that transect blockfield-covered plateaus at higher elevations (Skov et al., 2020). Retreat of the ice margin along East Greenland has been constrained by marine and terrestrial radiocarbon ages (Table 1) and surface exposure ages (Fig. 1). Along the eastern margin of Greenland, the ice-free landscapes comprise moraines, erratic boulders and polished bedrock surfaces that can be directly dated via cosmogenic nuclide exposure dating (e.g., Larsen et al., 2022; Levy et al., 2016; Skov et al., 2020). Grounded ice between Peary Land,
100 79 N Glacier, and Zachariae Isstrøm retreated westward out of the Greenland Sea and reached the present-day outer coastline by ~11.4 ka (Larsen et al., 2020). Between Storstrømmen Glacier and Kong Oscar Fjord, the GrIS receded to the outer coast between ~12.8 ka and 11.5 ka (Håkansson et al., 2007b; Larsen et al., 2022; Skov et al., 2020). The ice sheet subsequently retreated from the outer coast toward intermediate sites on the northeastern margin between ~11 – 9 ka (Larsen et al., 2018, 2022; Skov et al., 2020). The 79N
105 Glacier experienced slow early Holocene ice margin retreat from the outer coast to the present ice margin (Larsen et al., 2018; Bennike and Weidick, 2001). However, retreat rates from the inner fjord suggest rapid deglaciation between 10 and 8.5 ka (Roberts et al., 2024). Ice reached close to the modern ice margin by ~9 ka at Lambert Land and had retreated to Borgfjorden by ~8.9 ka in the Dove Bugt region, and Vandrepasset by ~8.6 ka in Bessel Fjord (Larsen et al., 2018, 2022).

110 A combination of raised marine beaches and terrestrial geological evidence has provided retreat chronologies of the East GrIS from the outer coast to the present-day ice margin (Fig. 1, Leger et al., 2024). Geomorphological mapping and dating glacial landforms and deposits using terrestrial cosmogenic nuclides (TCN; in situ ¹⁰Be) and radiocarbon ages have provided direct constraints of ice margin retreat (e.g. Håkansson et al., 2007a; Larsen et al., 2022; Levy et al., 2016). Our study sites focus on the glacial
115 geology of two marine embayments: Scoresby Sund and Dove Bugt (Figs. 1, 2).

2.1 Scoresby Sund

Scoresby Sund (Kangertittivaq) is a large (13,700 km²) fjord system situated between Jameson Land and Liverpool Land in the north, King Christian IX Land in the south, and Milne Land, and the Renland

120 Peninsula in the west (Fig. 3). The fjord system extends ~350 km inland from the outer coast (Dowdeswell et al., 1994a), with the outer fjord being ~120 km long, up to 50 km wide, and up to 650 m deep. Scoresby Sund is deepest in the south and gently rises to the north towards Jameson Land (Håkansson et al., 2007a). At the fjord mouth is a large submarine moraine, Kap Brewster moraine (Fig. 3b; Dowdeswell et al., 1994; Funder & Hansen, 1996).

125 Our study focuses on three presently ice-free sites on the outer coast of Scoresby Sund: Uunarteq on the northern mouth, Kap Brewster (Kangikajiip Appalia) on the southern mouth, and Rathbone Island, ~3 km to the east of the eastern coastline of Liverpool Land. At Uunarteq on the northern cape of Scoresby Sund, the local bedrock comprises Paleoproterozoic orthogneiss and granite (Kokfelt et al., 2023). Erratic boulders and cobbles of orthogneiss form a drift sheet over the local bedrock (Fig. 2a). At Kap Brewster on

130 the southern cape of Scoresby Sund, the local bedrock comprises Paleogene basalts of the North Atlantic Igneous Province (Wager, 1947). The eastern margin of Kap Brewster is characterized by steep basalt cliffs that extend over 200 m asl. Erratic sandstone and orthogneiss boulders and cobbles (>230 m asl) form a drift draped over the basaltic plateau above the cliffs (Fig 2b; Håkansson et al., 2007a). At Rathbone Island, a thin drift of orthogneiss boulders and cobbles to at least 120 m asl overlie Paleozoic granodiorite and

135 quartz diorite bedrock (Kokfelt et al., 2023; Fig 2c).

Surface exposure ^{10}Be ages from the mouth of Scoresby Sund suggest the ice margin retreated west of Kap Brewster by ~14.5 ka (Håkansson et al., 2007a). In Milne Land, ~150 km west of Kap Brewster, ^{10}Be exposure ages from the Milne Land stade moraines suggest ice retreated to southeastern Milne Land by ~11.4 ka (Levy et al., 2016). Notably, the Milne Land stade moraines have been mapped from Milne Land

140 to Germania Land (Fig. 1), and ^{10}Be and radiocarbon ages constrain both local mountain glacier and GrIS retreat during the Younger Dryas (12.9 – 11.7 ka; Funder et al., 2011; Kelly et al., 2008; 2025; Levy et al., 2016). In northwestern Scoresby Sund, the Milne Land stade moraines grade into raised marine terraces (Funder, 1978), and radiocarbon dates from Kjove Land, suggested the outer Milne Land stade moraines were deposited just before ~12.8 cal ka BP (Hall et al., 2008; Kelly et al. 2025). Additionally, radiocarbon

145 dated shell samples from Schuchert Dal reveal the Milne Land stade moraines were deposited during the early Younger Dryas or late Allerød (Hall et al., 2010). The surface exposure ages in Scoresby Sund generally agree with radiocarbon ages from marine sediment cores (Dowdeswell et al., 1994; Marienfeld, 1991), bivalves (Funder, 1978, 1990; Hall et al., 2008), and plant macrofossils (Bennike et al., 1999; Hansen, 2001; Ingólfsson et al., 1994) which record ice retreat through the main trough of Scoresby Sund

150 between 11.7 and 7.9 cal ka BP (Fig. 3).




Figure 3. Scoresby Sund with new measured ^{10}Be ages of boulders (red circles) with red sample names, and recalculated ^{10}Be ages of Håkansson et al. (2007a) with blue sample names at Kap Brewster. All ^{10}Be ages shown in kiloyears with 1σ internal uncertainties. Mean ^{10}Be ages from Uunarteq and Kap Brewster are shown in bold. Mean ^{10}Be age ($n = 8$; green circle) from Milne Land (Levy et al., 2016). Radiocarbon ages (orange circles) are listed in Table 1. ^{10}Be age outliers shown in italics. The grey dashed line denotes the flowline and ^{10}Be and radiocarbon ages used to calculate average retreat rates. The maximum LGM extent is marked by the black dashed line, derived from Leger et al. (2024 and references therein).

2.2 Storstrømmen Isstrom

Storstrømmen Isstrom is a major NEGIS outlet, and includes Kofoed-Hansen Bræ which drains to the northeast, and Storstrømmen Glacier which drains to the southeast (Fig. 4). The drainage basin (58,176 km²) of Storstrømmen Glacier contains enough ice to raise global sea level by 27 cm (Rignot et al., 2022). The southeastern lobe of Storstrømmen Glacier is a 23 km wide marine-terminating outlet glacier that drains into Borgfjorden of the Dove Bugt Embayment (Fig. 4). Storstrømmen is a surge glacier, with an estimated recurrence cycle of 50 – 70 years (Mouginot et al., 2018; Reeh et al., 1994). The terminus retreated 15 km between 1913 and 1950, remained stable between 1950 and 1978, and readvanced 8 km between 1978 and 1984 during a surge (Reeh et al., 1994). Grounding line retreat of 1.1 km from 2017 – 2021 has been attributed to increased surface ablation (Rignot et al., 2022). Recent observations suggest a possible surge onset projected to occur between 2027 and 2040 (Andersen et al., 2025).

At the terminus, Storstrømmen Glacier splits into two lobes that flow around an unnamed island where we collected samples for surface exposure dating. The western lobe merges with L. Bistrup Bræ Glacier, and the eastern lobe flows directly into Borgfjorden. The unnamed island acts as a pinning point, stabilizing the present-day glacier (Reeh et al., 1994). The local bedrock of the island comprises basement Paleoproterozoic orthogneiss (Kokfelt et al., 2023) and is overlain by moraines and glacially transported erratics (Fig 2d,e,f). At the southern end of the island, moraine deposits grade into a raised marine terrace at the local marine limit (28 m asl). This observation is consistent with the 28 m marine limit at Hellefjord in western Germania Land (Landvik, 1994; Weidick et al., 1996). Radiocarbon ages from marine bivalves in Neoglacial moraines indicate that Storstrømmen Glacier terminated inland from its present-day position between 5.4 – 1.2 cal ka BP, when a ~100 km long sound formed between Kofoed-Hansen Bræ and Storstrømmen (Weidick et al., 1996). The ice margin of the unnamed island is surrounded by fresh (little weathered) “historical moraine” ridges and mark a late Neoglacial readvance that culminated ~1850 (Weidick et al., 1996).

Table 1. Radiocarbon ages from eastern Greenland

Sample number	Latitude	Longitude	Location	Material dated	Radiocarbon age (¹⁴ C yr BP)	Radiocarbon age uncertainty (1σ)	Calibrated age (cal yr. BP, median)	Calibrated age range (cal yr. BP, 2σ)	Reference
Northeast Greenland shelf									
ETH-106030	80.037	-8.923	DA17-NG-ST03-039G	Mixed planktonic foraminifera	12070	60	13291*	13202-13584*	Hansen et al. (2022)
ETH-1113888	78.501	-16.28	DA17-NG-ST08-092G	Mixed benthic foraminifera	10200	190	12473*	11268-12595*	Davies et al. (2022)
AW12795.1.1	77.497	-18.14	PS100/270	Mixed benthic foraminifera	9437	104	10138	9670 - 10568	Syring et al. (2020)
121063.1.1	77.127	-10.68	DA17-NG-ST12-135G	Mixed benthic foraminifera	14450	220	16684	16080 - 17285	Lopez-Quiros et al. (2024)
Lambert Land									
442909	79.107	-19.75	Zachariae Isstrom	Marine bivalve	7595	55	7905	7624 - 8184	Bennike and Weidick (2001)
Clavering Ø									
1-9659	74.350	-21.87	Clavering Ø	Marine bivalve	8195	125	8148	7050 - 9396	Weidick (1978)
Kejser Franz Fjord									
MM/G3	73.550	-25.12	Kejser Franz Joseph Fjord	Marine bivalve	8020	100	8078	7179 - 9086	Wagner and Melles (2002)
60282.1.1	73.675167	-24.18	HH13-012	Mixed benthic foram	7470	130	7471	6518 - 8376	Olsen et al. (2022)
4491.1.1	73.474667	-24.61	PS2633	Foraminifera	8376	84	8452	7550 - 9447	Olsen et al. (2022)
PS2630	73.158	-18.07	PS2630	Foraminifera	13560	130	15544	15113 - 15989	Evans et al. (2002)
Kong Oscar Fjord									
1-9104	73.350	-26.47	Kong Oscar Fjord	Marine bivalve	7760	115	7985	7654 - 8329	Weidick (1977)
1-9658	73.167	-27.42	Kong Oscar Fjord	Marine bivalve	7160	125	7428	7067 - 7779	Weidick (1978)
Lu-1070	72.717	-26.83	Kong Oscar Fjord	Marine bivalve	7530	75	7698	7150 - 8297	Håkansson (1976)
Scoresby Sund									
UIC-7428	70.595	-21.54	Scoresby Sund	Plant macrofossil	8900	50	10026	9887 - 10197	Cremer et al. (2001)
AAR-199	70.338	-23.71	PS1715-1	Foraminifera	10340	150	11250	10674 - 11819	Mariénfeld (1991)

AAR-200	70.483	-24.67	PS1719-1	Foraminifera	10480	190	11645	11058 - 12392	Marienfeld (1991)
AAR-202	70.708	-25.00	PS1728-1	Foraminifera	10760	120	12023	11503 - 12486	Marienfeld (1991)
AAR-201	70.928	-24.98	PS1727-1	Foraminifera	9080	650	9589	7952 - 11272	Marienfeld (1991); Dowdeswell et al. (1994)
K-1461	71.167	-25.33	Scoresby Sund	Marine bivalve	9040	140	9650	9275 - 10108	Funder (1990)
K-3109	70.500	-23.45	Scoresby Sund	Marine bivalve	10560	145	11726	11199 - 12318	Funder (1990)
AAR-636	70.533	-23.63	Scoresby Sund	Marine bivalve	9170	140	9764	9398 - 10183	Gulliksen et al. (1991)
I-9492	70.500	-26.18	Scoresby Sund	Marine bivalve	9925	140	10882	10395 - 11310	Funder (1978)
AAR-3368	70.387	-23.87	Scoresby Sund	Plant macrofossils	9780	160	11186	10695 - 11743	Hansen (2001)
100380, 100381	70.843	-24.01	Scoresby Sund	Plant macrofossils	9665	55	11058	10780 - 11207	Bennike et al. (1999)
AAR-2540	70.850	-24.02	Scoresby Sund	Plant macrofossils	10130	130	11729	11257 - 12177	Bennike et al. (1999)
AAR-3369	70.850	-24.07	Scoresby Sund	Plant macrofossils	9440	65	10681	10501 - 11069	Hansen (2001)
AAR-3802	70.850	-24.15	Scoresby Sund	Marine bivalve	9985	60	10915	10581 - 11203	Hansen (2001)
AAR-2571	70.985	-24.00	Scoresby Sund	Plant macrofossils	7900	100	8751	8464 - 9006	Böcher and Bennike (1996)
AAR-1829	70.959444	-24.11361	Scoresby Sund	Marine bivalve	10140	100	11126	10696 - 11557	Funder and Hansen (1996)
AAR-2541	70.950	-24.12	Scoresby Sund	Plant macrofossils	9910	130	11411	10885 - 11873	Hansen (2001)
AAR-644	70.914	-24.21	Scoresby Sund	Marine bivalve	9670	180	10486	9930 - 11085	Ingolfsson et al. (1994)
87516	71.567	-23.97	Scoresby Sund	Plant macrofossils	7000	390	7848	7003 - 8601	Böcher and Bennike (1996); Bennike et al. (1999)
K-1915	71.350	-24.83	Scoresby Sund	Marine bivalve	10300	120	11423	11018 - 11902	Funder (1978)
I-5421	70.950	-28.15	Scoresby Sund	Marine bivalve	7540	130	7863	7560 - 8182	Funder (1978)
Kangerlussuaq Trough									
AA-4026	66.203	-30.66	10A	Foraminifera	13585	110	15628	15259 - 15995	Williams (1993)
AA-6848	66.764	-30.84	PO-175/15	Foraminifera	14845	190	17229	16717 - 17791	Williams et al. (1995)

AA-11584	68.116	-31.43	91-K11A	Foraminifera	9975	100	10942	10585 - 11239	Andrews et al. (1997)
CAMS-32047	68.100	-29.35	JM96-1207	Foraminifera	9800	60	10731	10495 - 11037	Jennings et al. (2002)
AA-29205	67.300	-30.97	JM96- 1214/2GC	Benthic foraminifera	11,380	80	12786	12601 - 13009	Smith and Licht (2000); Jennings et al. (2006)
AA-32045	67.047	-30.86	JM96- 1215/2GC	Benthic and planktic foraminifera	12900	50	14693	14326 - 15010	Smith and Licht (2000); Jennings et al. (2006)
AA-23221	65.963	-30.63	JM96- 1216/2-GC	Benthic foraminifera	14550	150	16862	16423 - 17296	Smith and Licht (2000); Jennings et al. (2008)
AA-43116	68.093	-27.84	MD99-2317	Bryozoans	11950	110	13405	13122 - 13717	Jennings et al. (2006)
AA-4666	67.410	-31.07	3	Foraminifera	9375	70	10133	9818 - 10424	Williams (1993); Mienert et al. (1992)

*Modelled median age derived from age-depth model (Davies et al., 2022; Hansen et al., 2022).




Figure 4. (A) Dove Bugt region showing outer coast (purple), intermediate (green) and ice margin (blue) exposure ages from Larsen et al. (2022). The outer coast mean ages are 12.8 ka (n = 1), 12.7 ka (n = 3),

12.6 ka (n = 3). The intermediate mean ages are 10.8 ka (n = 3), 10.6 ka (n = 4). The ice margin mean ages are 9.1 ka (n = 3), 8.9 ka (n = 3), 8.6 ka (n = 6). Black inset shows the location of Storstrømmen Glacier terminus in aerial photo (B) and in Fig. 5. The grey dashed line denotes the flowline used to calculate average retreat rates. The mean ^{10}Be ages used to calculate average retreat rates are 12.7 ka, 10.8 ka, and 8.9 ka. (B) View looking south at the Storstrømmen Glacier terminus showing the historical moraine (blue dashed line) and modern ice margin.

3 Methods

3.1 Field methods

We collected samples from the mouth of Scoresby Sund and Rathbone Island during the boreal summer of 2022, and from the island at the terminus of Storstrømmen Glacier during the boreal summer of 2023. We sampled 13 boulders from Scoresby Sund (Fig. 3), and 15 boulders and one bedrock surface from Storstrømmen (Fig. 5) for ^{10}Be analysis (Table 2). We targeted large (>0.5 m diameter) boulders perched on bedrock or on well-developed moraines. The bedrock sample (23DMH-CR1-SURFACE) was collected from a surface with visible signs of glacial abrasion (e.g. glacial polish, striae). Four of the samples from Kap Brewster at the mouth of Scoresby Sund were collected from friable white sandstone boulders displaying exfoliation and weathering pits. We avoided sampling surfaces near the exfoliation or weathering pits. Two erratic boulder samples (22GRO-126 and 22GRO-127) from Kap Brewster were collected from the same boulders (KB-1 and KB-2) as previously sampled by Håkansson et al. (2007a).

3.2 Analytical methods

Whole rock samples from Scoresby Sund were crushed, sieved and processed to quartz purification at the University at Buffalo. Whole rock samples from Storstrømmen Glacier were crushed, sieved and processed to quartz purification at Lamont-Doherty Earth Observatory. All samples were then processed at Lamont-Doherty Earth Observatory via conventional ^{10}Be extraction methods (Schaefer et al., 2009). $^{10}\text{Be}/^9\text{Be}$ ratios were measured at the Center for Accelerator Mass Spectrometry at Lawrence Livermore National Laboratory, and normalized to the 07KNSTD standard of Nishiizumi et al. (2007) with a nominal ratio of 2.85×10^{-12} . Blank corrections to measured $^9\text{Be}/^{10}\text{Be}$ ratios amounted to <2%, and process blanks range from 6200 – 11,855 ^{10}Be atoms (Table S2). Final errors in ^{10}Be concentrations are obtained by quadrature addition of the final accelerator mass spectrometry analytical error, and a 1% error in Be spike concentration.

We calculated our new surface exposure ages and recalculated previously reported ages using version 3 of the CRONUS-Earth calculator (<http://hess.ess.washington.edu/>, last access: February 2025; Balco et al.,

2008) using the Lm scaling scheme (Lal, 1991; Stone, 2000) and the Arctic production rate (Young, et al., 2013a; Table S1). We note that using the St scaling scheme results in almost identical ages for our samples, while ages derived using the LSDn scaling scheme differ by less than 120 years, and thus do not affect our interpretations. Four samples are quartz sandstone boulders collected from Kap Brewster (22GRO-126, 22GRO-127, 22GRO-128, 22GRO-131) and we use a density of 2.4 g cm^{-3} . All other samples are crystalline boulders or bedrock (e.g., orthogneiss, granodiorite) and we use a density of 2.65 g cm^{-3} . Unless otherwise noted, all radiocarbon ages (reported as: cal ka BP) mentioned in the text are recalculated using CALIB 8.2 (Stuiver & Reimer, 1993); <http://calib.org/calib/calib.html>, last access: April 2025), and the IntCal20 calibration curve (Reimer et al., 2020) for terrestrial samples, or the Marine20 calibration curve (Heaton et al., 2020) for marine samples (Table 1, Table S3). For marine samples, marine reservoir corrections were calculated using the Marine Reservoir Correction Database (<http://calib.org/marine/>, last access: April 2025)(Reimer & Reimer, 2001). Each site uses an average ΔR value based on water depth, distance to coast, and where possible species that are most representative of the study site, as suggested in Pearce et al. (2023).

Table 2 Cosmogenic ^{10}Be concentrations and apparent exposure ages.

Sample name	Latitude (DD)	Longitude (DD)	Elevation (masl)	Sample thickness (cm)	Topographic shielding	^{10}Be conc. (10^4 atoms g^{-1}) ^a	^{10}Be age (ka) ^{b,c,d}
Rathbone Island							
22GRO-139	70.66612	-21.3985	118	3.00	1	7.66 $\pm$ 0.18	16.4 $\pm$ 0.4 (0.6)
22GRO-140	70.66674	-21.3991	119	3.00	1	8.25 $\pm$ 0.20	17.7 $\pm$ 0.4 (0.6)
22GRO-142	70.66599	-21.404	97	2.5	1	6.45 $\pm$ 0.16	14.1 $\pm$ 0.3 (0.5)
Most likely:							
Unarteq							
22GRO-100	70.44798	-21.8879	99	3.00	0.998	6.19 $\pm$ 0.15	13.6 $\pm$ 0.3 (0.5)
22GRO-101	70.44798	-21.8879	99	3.00	0.998	6.04 $\pm$ 0.17	13.2 $\pm$ 0.4 (0.5)
22GRO-102	70.44791	-21.8876	94	3.00	0.998	5.56 $\pm$ 0.23	12.3 $\pm$ 0.5 (0.6)
22GRO-103	70.44786	-21.888	91	3.00	0.994	5.64 $\pm$ 0.14	12.5 $\pm$ 0.3 (0.4)
22GRO-104	70.44768	-21.888	103	3.00	0.998	12.20 $\pm$ 0.26	26.4 $\pm$ 0.6 (0.9)
Kap Brewster (Kangikajip Appalia)							
22GRO-126	70.151	-22.0754	225	3.00	1	5.99 $\pm$ 0.17	11.4 $\pm$ 0.3 (0.4)
22GRO-127	70.15112	-22.076	231	3.00	1	6.97 $\pm$ 0.18	13.2 $\pm$ 0.3 (0.5)
22GRO-128	70.15112	-22.0761	233	3.00	1	7.18 $\pm$ 0.16	13.6 $\pm$ 0.3 (0.5)
22GRO-131	70.15077	-22.0766	227	3.00	1	7.33 $\pm$ 0.17	13.9 $\pm$ 0.3 (0.5)
22GRO-134	70.15005	-22.0756	230	3.00	1	7.03 $\pm$ 0.15	13.4 $\pm$ 0.3 (0.5)
Weighted mean (Unarteq and Kap Brewster):							
13.2 $\pm$ 0.7 (n = 8)							
Storstrømmen Glacier							
Outboard							
23DMH-CR1-SURFACE	76.87522	-22.3197	166	1.65	1	5.634 $\pm$ 0.15	11.5 $\pm$ 0.3 (0.4)
23DMH-18	76.87489	-22.3197	161	2.79	1	5.93 $\pm$ 0.16	12.3 $\pm$ 0.3 (0.5)
23DMH-20	76.87555	-22.3165	164	2.19	1	4.25 $\pm$ 0.15	8.7 $\pm$ 0.3 (0.4)
23DMH-21	76.87409	-22.3181	163	1.85	1	4.33 $\pm$ 0.09	8.9 $\pm$ 0.2 (0.3)
Weighted mean:							
8.8 $\pm$ 0.3 (n = 2)							
M1 moraine							
23DMH-22	76.85698	-22.277	114	1.08	1	4.48 $\pm$ 0.11	9.6 $\pm$ 0.2 (0.4)
23DMH-23	76.85442	-22.281	92	3.53	1	3.82 $\pm$ 0.19	8.6 $\pm$ 0.4 (0.5)

23DMH-24	76.85425	-22.2807	91	3.37	1	3.72 ± 0.08	8.4 ± 0.2 (0.3)
23DMH-25	76.85428	-22.2824	91	2.54	1	3.84 ± 0.12	8.6 ± 0.3 (0.3)
23DMH-29	76.8687	-22.3197	152	3.67	1	4.10 ± 0.11	8.6 ± 0.2 (0.3)
23DMH-30	76.86914	-22.3229	151	2.41	1	4.19 ± 0.11	8.7 ± 0.2 (0.3)
23DMH-31	76.87427	-22.3491	124	3.07	1	3.92 ± 0.09	8.5 ± 0.2 (0.3)
Weighted mean:							
Inboard							
23DMH-26	76.84808	-22.2882	38	2.39	1	3.32 ± 0.13	7.9 ± 0.3 (0.4)
23DMH-27	76.84876	-22.2989	90	2.40	1	3.70 ± 0.09	8.3 ± 0.2 (0.3)
23DMH-28	76.8511	-22.3069	109	1.07	1	3.80 ± 0.09	8.2 ± 0.2 (0.3)
23DMH-32	76.88953	-22.3271	127	2.01	1	3.80 ± 0.10	8.1 ± 0.2 (0.3)
23DMH-33	76.89405	-22.3273	125	2.22	1	3.72 ± 0.09	8.0 ± 0.2 (0.3)
Weighted mean:							
8.1 ± 0.3 (n = 5)							

^a Samples normalized to the 07KNSTD standard of Nishiizumi et al. (2007).

^b Exposure ages calculated using the CRONUS-Earth calculator (<https://hess.ess.washington.edu/>, last access: February 2025), using the Lm scaling scheme (Lal, 1991; Stone, 2000) and the Arctic production rate (Young, et al., 2013a).

^c Both internal and external uncertainties shown at the 1 σ level. Internal uncertainties are analytical uncertainties only, and external uncertainties (given in parentheses) are absolute uncertainties and include production rate and scaling errors.

^d The weighted mean for each site includes the propagation of a 3.7% production rate uncertainty (Young et al., 2013a). Outliers in italics are excluded from the weighted mean age.

4 Results

4.1 Scoresby Sund

The 13 erratic boulder samples from the outer coast of the Scoresby Sund region produce ^{10}Be ages that range from 11.4 ± 0.3 – 26.4 ± 0.6 ka (Figs. 3,6). On Rathbone Island, east of Scoresby Sund, three erratic boulders were sampled between 97 and 118 m elevation and produce exposure ages of 14.1 ± 0.3 ka (22GRO-142), 16.4 ± 0.4 ka (22GRO-139), and 17.7 ± 0.4 ka (22GRO-140). At Uunarteq located at the northern mouth of Scoresby Sund, five erratic boulders sampled between 91 and 103 m elevation produce ^{10}Be ages that range from 12.3 ± 0.5 – 26.4 ± 0.6 ka. At Kap Brewster (Kangikajik) located at the southern mouth of Scoresby Sund, five erratic boulders sampled between 225 and 233 m elevation produce ^{10}Be ages that range from 11.4 ± 0.3 to 13.9 ± 0.3 ka. The ^{10}Be ages from samples 22GRO-126 and 22GRO-127 are 11.4 ± 0.3 ka, and 13.2 ± 0.6 ka, respectively; these samples were collected from the same boulders as samples KB-1 and KB-2 from Håkansson et al. (2007a), which yield recalculated ages of 18.3 ± 2.9 and 17.2 ± 2.2 ka. The large difference in ages is outside the 1σ error range, and may be attributed to analytical errors. After excluding one older outlier (26.4 ± 0.6 ka) from Uunarteq, and a younger outlier from Kap Brewster (11.4 ± 0.3 ka), the mean age from the two sites at the mouth of Scoresby Sund is 13.2 ± 0.7 ka ($n=8$; Table 2).

4.2 Storstrømmen Glacier

The 15 erratic boulder samples from the unnamed island at the Storstrømmen Glacier terminus produce ^{10}Be ages that range from 7.9 ± 0.3 – 12.3 ± 0.3 ka (Figs. 5,6). Excluding two outliers (23DMH-18, 23DMH-22) the mean age of the boulders on the island is 8.4 ± 0.4 ka ($n = 13$; Table 2). Of these, seven samples were collected from boulders on a prominent moraine (M1 moraine) deposited by the western lobe that extends ~5 km through the island, and produce ^{10}Be ages of 9.6 ± 0.2 ka, 8.6 ± 0.4 ka, 8.4 ± 0.2 ka, 8.6 ± 0.3 ka, 8.6 ± 0.2 ka, 8.7 ± 0.2 ka, and 8.5 ± 0.2 ka. The mean age for the M1 moraine is 8.6 ± 0.3 ka ($n = 6$), excluding sample 23DMH-22 (9.6 ± 0.2 ka), which was identified as an outlier (Fig. 6). ^{10}Be ages from three boulders ~500 m north and outboard of the moraine are 8.7 ± 0.3 , 8.9 ± 0.2 , and 12.3 ± 0.3 ka. The 12.3 ± 0.3 ka sample (23DMH-18) outboard of the M1 moraine was identified as an outlier as it falls outside of the 2σ range of the mean age of the boulders on the island. The one bedrock surface, ~500 m north of the moraine and in close proximity to 23DMH-18, produced a ^{10}Be age of 11.5 ± 0.3 ka (23DMH-CR1-SURFACE). The bedrock surface likely has inherited ^{10}Be and was not used to determine deglacial history.

330 On the southeastern side of the island, ^{10}Be ages from three boulders close to the raised marine terrace are 7.9 ± 0.3 , 8.3 ± 0.2 , and 8.2 ± 0.2 ka, and are inboard of the M1 moraine. On the northern margin of the island, ^{10}Be ages from two boulders adjacent to the historical moraine and modern ice margin are 8.1 ± 0.2 ka (23DMH-32) and 8.0 ± 0.2 ka (23DMH-33). These boulders were deposited by the eastern lobe.




Figure 5. Map of Storstrømmen Glacier terminus with extent of Little Ice Age moraine (blue dashes) and raised marine terrace limit (green dashes). The M1 moraine (magenta) deposited by the western lobe, grades into the marine terrace at 28 m elevation. Measured ^{10}Be ages of boulders (red circles) and bedrock (blue square) are shown in kiloyears (ka) with 1σ internal uncertainties. ^{10}Be age outliers shown in italics. Satellite image from WorldView-2 (© 2021, Maxar).




Figure 6. Camel plots (normal kernel density function) for ^{10}Be exposure ages Uunarteq, Kap Brewster and Rathbone Island near the outer coast of Scoresby Sund, and samples from the M1 moraine, and inboard and outboard of the M1 moraine at the terminus of the Storstrømmen Glacier. Colored camel plots show the relative probability for all ^{10}Be ages at each site.

5 Discussion

5.1 Glacier evolution in Scoresby Sund

The ^{10}Be exposure ages from Rathbone Island provide new minimum constraints on the retreat of the GrIS off the coast of Liverpool Land at 14.1 ± 0.3 ka, and record one of the earliest terrestrial deglaciation events in Greenland. Our mean ^{10}Be age of 13.2 ± 0.7 ka ($n = 8$) from the north and south sides of the mouth of Scoresby Sund is consistent with the minimum ^{10}Be exposure age from Kap Brewster reported by Håkansson et al. (2007a) of 14.5 ± 1.7 ka, and provides a minimum constraint on ice margin retreat past the mouth of Scoresby Sund (Fig. 6). Based on our age constraints at the mouth of Scoresby Sund, we infer the depositional age of the Kap Brewster submarine moraine to correspond with our mean age of ~ 13.2 ka (Fig. 3b).

At sites (≥ 300 km) north of Scoresby Sund, the outer coast deglaciated between 12.8 ± 0.6 and 11.5 ± 0.2 ka (Larsen et al., 2022). This retreat chronology is consistent with mountain glacier retreat between ~ 12.8 and 11.7 ka from Milne Land, Kjove Land, Gurreholm Dal and Schuchert Dal derived from ^{10}Be and radiocarbon ages (Hall et al., 2008; 2010; Kelly et al., 2008; 2025; Levy et al., 2016). We suggest the earlier onset of deglaciation at the outer coast of the Scoresby Sund region may be related to the closer proximity of our dated sites to the edge of the continental shelf than other parts of East Greenland. This interpretation is further supported by deglaciation chronologies before 13.4 cal ka BP, and 13.3 cal ka BP on the northeast Greenland shelf (Davies et al., 2022; Hansen et al., 2022).

The timing of retreat at Rathbone Island at ~ 14.1 ka, to the mouth of Scoresby Sund at ~ 13.2 ka, coincides with the Bølling-Allerød warm period (14.7 – 12.9 ka, Rasmussen et al., 2006), suggesting widespread retreat of the GrIS across the eastern continental shelf. Retreat from across the northeast Greenland shelf between 16.6 and 13.4 cal ka BP (Davies et al., 2022; López-Quirós et al., 2024), coincides broadly with Meltwater Pulse 1A, a period of rapid sea level rise between 14.6 and 14.3 ka (Deschamps et al., 2012; Hanebuth et al., 2000). Alongside our ^{10}Be ages from Rathbone Island, we suggest that ice margin retreat across the East Greenland shelf may have been influenced, in part, by enhanced sensitivity to abrupt global sea-level rise at this time. The retreat of ice in the Scoresby Sund region was likely attributed to the influx of warm Atlantic water that flowed onto the East Greenland continental shelf, perhaps related to the Bølling-Allerød warm period (Hansen et al., 2022). The inflow of Atlantic water beneath fresher and colder polar surface water has been inferred by high abundances of *Cassidulina neoteretis* (Foraminifera) in marine sediment cores, a species typically found in Atlantic sourced subsurface waters (Cage et al., 2021; Consolaro et al., 2018). In southeast Greenland, the high abundance of *C. neoteretis* in shelf cores also suggests the advection of Atlantic water into the Kangerlussuaq Trough drove basal melt and retreat during the Bølling-Allerød and the Younger Dryas (Jennings et al., 2006).

Our retreat chronology, and previously published ^{10}Be and radiocarbon ages in Scoresby Sund, suggest ice in the main fjord basin of Scoresby Sund retreated from the mouth of Scoresby Sund at ~ 13.2 ka to Renland

by 9.7 cal ka BP (Funder, 1978), and using the flowline in Fig. 3, we calculate an average retreat rate of ~43 m/yr. In southern Scoresby Sund, ice retreated ~150 km from the mouth to the southern end of Milne Land by ~11.4 ka (Levy et al., 2016) at an average retreat rate of ~83 m/yr.

5.2 Glacier evolution in the Dove Bugt region

Our mean ^{10}Be age of 8.6 ± 0.3 ka ($n = 6$) from the prominent M1 moraine at Storstrømmen Glacier record the conclusion of a stillstand or advance near the modern ice margin (Fig. 6). This age is consistent with the ^{10}Be retreat chronology at Bræ Øerne of 8.9 ± 0.2 ka (Larsen et al., 2022), ~14 km south of Storstrømmen Glacier (Fig. 4). We suggest Storstrømmen Glacier comprised two expanded ice lobes around the eastern and western margins of the unnamed island. As the glacier retreated, boulders and moraines were deposited between ~8.6 ka and 8.0 ka (Figs. 2,5). The western ice lobe deposited the prominent M1 moraine in the middle of the island and all the boulders inboard (west) of the moraine. As the glacier retreated the western lobe wrapped around the southern end of the island depositing the M2 moraine. Two lakes ~1.5 km south of sample 23DMH-31 on the M2 moraine are dammed at their southern end, and we interpret these to be moraine dammed lakes that formed as the western lobe retreated (Fig. 5). The eastern lobe ice likely retreated beyond the historical moraine, and present ice margin at ~8 ka.

The 28 m marine limits at the terminus of Storstrømmen (Weidick et al., 1996) is dated to an estimated age of ~8 cal ka BP with marine bivalves (Landvik, 1994; Fig. 7). Our exposure ages from the unnamed island are consistent with the radiocarbon ages of the raised marine sediments. Also, our ice margin retreat history is in agreement with deglaciation of inner Kejser Franz Joseph Fjord and Nordfjord, ~360 km south of Storstrømmen, constrained by radiocarbon ages of 8.5 and 7.5 cal ka BP, respectively (Olsen et al., 2022).

Our ^{10}Be ages from the M1 moraine at ~8.6 ka to near the modern ice margin ~8 ka suggests an ice margin stillstand or slowdown of ice retreat, broadly consistent with the abrupt 8.2 ka cooling event (Alley et al., 1997). The formation of the M1 moraine (8.6 ± 0.3 ka) could be related to a period of ice margin stagnation in response to local cooling or localized dynamics unrelated to climate at this time. In West Greenland, surface exposure dating of moraine systems record the ice margin responded directly to the 9.3 and 8.2 ka cooling events (Young et al., 2013b). These observations support regional cooling across Greenland during this interval. However, considering present-day Storstrømmen is a surge glacier (Andersen et al., 2025; Mouginot et al., 2018; Reeh et al., 1994), an alternative explanation is that the moraine records past surge dynamics, similar to modern-day processes. Another alternative explanation is that the ice margin was pinned to the island ~8.6 ka, slowing ice retreat during this time, independent of any climate event.

The response of the GrIS to climate variability is strongly influenced by ice flux (Young, et al., 2013b). In low ice flux glaciers, such as Storstrømmen, mass balance change driven by regional cooling (i.e., reduced summer ablation), may occur slowly resulting in a subtle response or stillstand. In contrast, the response to high flux glaciers, such as Jakobshavn Isbræ in West Greenland, to such cooling may occur relatively quickly resulting in an ice sheet advance (Young, et al., 2013b). For example, at Jakobshavn Isbræ, a high flux environment was observed between $\sim 8.5 - 8.1$ ka, directly depositing moraines during the 8.2 ka event (Young et al., 2013b). Our observations in northeast Greenland, are consistent with patterns observed in West Greenland (Young, et al., 2013b; Young et al., 2011), suggesting that marine-terminating glaciers show high sensitivity to abrupt warming and cooling on centennial timescales. In contrast, land-terminating glaciers in Greenland are less sensitive to abrupt climate change (Long et al., 2006; Young et al., 2011).

We estimate a minimum average retreat rate of ~ 28 m/yr for the outer coast near Store Koldewey at ~ 12.7 ka (Larsen et al., 2022) to within ~ 3 km of the modern ice margin by ~ 8.6 ka. The average retreat rate of Storstrømmen during the last deglaciation is an order of magnitude lower than the observed grounding line retreat of 1.1 km at a rate of $250 - 380$ m/yr between 2017 and 2021, attributed to surface ablation (Rignot et al., 2022). However, the modern retreat follows a surge event between 1978 and 1984 (Reeh et al., 1994), and present surge behavior may be related glacier instability. Therefore, direct comparisons between Holocene and modern retreat rates at Storstrømmen require caution. Additionally, differences in temporal resolution, the drainage system, and in external forcing such as ablation rates and ocean temperatures must be considered when comparing retreat rates across different timescales.




Figure 7. Ice margin retreat from Scoresby Sund and the Dove Bugt region. $\delta^{18}\text{O}$ record of the North Greenland Ice core Project (NGRIP members, 2004). Marine limit curve for the Dove Bugt region derived from Weidick et al. (1996). Normalized glacier length includes published radiocarbon ages from Scoresby Sund (Dowdeswell et al., 1994; Funder, 1978; Marienfeld, 1991), and published terrestrial cosmogenic nuclides (TCN) ^{10}Be ages from the Dove Bugt embayment (Larsen et al. 2022).

5.3 Rates of marine-terminating glacier retreat

As ice retreated westward out of the Greenland Sea, minimum average retreat rates of ice streams from the middle of the continental shelf to the outer coast of northeast Greenland were 96 m/yr from Foster Bugt to

the mouth of Kejser Franz Joseph Fjord (Olsen et al., 2022). At the Store Koldewey Trough, located east of Store Koldewey Island and the Dove Bugt embayment, retreat rates ranged between 80 – 400 m/yr (Olsen et al., 2020). In contrast, our minimum average retreat rates observed at Scoresby Sund (~43 m/yr) and Storstrømmen Glacier (~28 m/yr) are consistent with broader estimates for marine-terminating glaciers in East Greenland fjords, which range between 10 and 80 m/yr (Dyke et al., 2014; Hughes et al., 2012).

We summarize the late glacial and Holocene rates of retreat in East Greenland in Figure 8. For comparison, retreat rates include 33 m/yr at Nordfjord, and 35 m/yr at Kejser Franz Joseph Fjord (Olsen et al., 2022); ~10 m/yr at Kangerlussuaq Fjord (Dyke et al., 2014); and ~80 m/yr at Helheim Glacier (Hughes et al., 2012). Rapid retreat rates of 79N Glacier within Nioghalvfjærdsfjorden after 10 ka range between 248.5 m/yr and 613 m/yr record some of the highest retreat rates in both Greenland and Antarctica (Roberts et al., 2024). Our retreat rates at Scoresby Sund (~43 m/yr) are comparable to retreat rates for the entire central East Greenland region (35 ± 23 m/yr), while our retreat rates at Dove Bugt (~28 m/yr) are lower than for the entire Northeast Greenland region (44 ± 10 m/yr) that were estimated using transects across isochrones (Leger et al., 2024). In West Greenland, Young et al. (2011) estimated an average retreat rate of ~100 m/yr for Jakobshavn Isbræ between 8 – 7.5 ka. The thickest and fastest moving parts of Greenland's largest present-day ice streams correspond to the contemporary submarine melt rates at 79N (50 – 60 m/yr) in northeastern Greenland, and Petermann Glacier (40 – 50 m/yr) and Ryder Glacier (50 m/yr) in northwestern Greenland (Wilson et al., 2017). These modern rates are comparable to late glacial and Holocene retreat rates observed in East Greenland (Fig. 8).

Fast flowing glaciers that lack topographic pinning points that enhance ice stability (i.e., islands, upward sloping beds) are particularly susceptible to rapid retreat (Mouginot et al., 2015). Modern ice flow rates at Jakobshavn (11–17 km/yr; Joughin et al., 2014), Helheim (5–11 km/yr; Bevan et al., 2012) and 79N (1.4 km/yr; Vijay et al., 2019), all reflect fast flowing outlet glaciers of the GrIS with large volumes of ice discharge. By comparison, Storstrømmen Glacier has flow speeds of 0.19 km/yr (Vijay et al., 2019), an order of magnitude slower than the fastest flowing glaciers of Greenland. This observation is also reflected in late glacial and Holocene retreat patterns in East Greenland fjords. With the exception of 79N Glacier, retreat rates observed within the East Greenland fjords during the late glacial and Holocene were generally lower than those observed from the middle shelf to the outer coast. This pattern suggests a shift from large ice streams to topographically controlled retreat within fjord systems.

Numerical simulations from southwestern Greenland suggested ice calving within fjord systems significantly impact enhanced ice discharge at the terminus of outlet glaciers during the early Holocene (Cuzzone et al., 2022). Therefore, future ice sheet modeling, benchmarked against geologic constraints of

past East GrIS change (e.g., Briner et al., 2020) could greatly improve estimations of ice mass loss, rates or retreat, and associated processes from marine-terminating outlet glaciers across Greenland.




Figure 8. Rates of ice margin retreat (m/yr) for marine-terminating outlet glaciers in East Greenland during the late glacial and Holocene, compared with modern subglacial melt rates from 79N Glacier, Ryder Glacier and Petermann Glacier (Wilson et al., 2017).

6 Conclusion

We report new ¹⁰Be ages from Scoresby Sund on the outer coast in central east Greenland, and from near the modern ice margin of Storstrømmen Glacier in northeast Greenland. These new exposure ages provide direct constraints of the late glacial and early Holocene timing of ice retreat westward out of the Greenland Sea, within the present-day coastline, and within the modern ice margin. Our key findings include:

- Ice retreated from Rathbone Island, east of Scoresby Sund, by ~14.1 ka, recording one of the earliest known terrestrial deglaciation events in Greenland. Our ¹⁰Be ages support the idea that ice margin

485 retreat across the East Greenland shelf was perhaps in response to enhanced sensitivity of abrupt
global sea-level rise.

- Ice retreated from the mouth of Scoresby Sund at ~13.2 ka during the Bølling-Allerød warm period,
perhaps related to an influx of warm Atlantic water that flowed onto the East Greenland continental
shelf (Hansen et al., 2022).
- Storstrømmen Glacier reached ~3 km from the modern ice margin at ~8.6 ka and reached the
490 modern margin ~8 ka. The ice margin slowed or reached a stillstand between 8.6 – 8.0 ka. In low
ice flux environments, such as Storstrømmen, mass balance change is driven by regional cooling
and may occur slowly resulting in a subtle response or stillstand. These observations suggest that
marine-terminating sectors of the eastern GrIS may be highly sensitive to abrupt warming and
495 cooling events on centennial timescales.
- Average rates of marine-terminating glacier retreat at Scoresby Sund were ~43 m/yr between ~13.2
ka and 9.7 ka. Average rates of marine-terminating glacier retreat in the Dove Bugt region were
~28 m/yr between ~12.7 ka and ~8.6 ka. The rates of late glacial and Holocene retreat at Scoresby
Sund, and Storstrømmen Glacier are within previously reported estimates of marine-terminating
500 outlet glacier retreat across eastern Greenland fjords (10 – 80 m/yr; Dyke et al., 2014; Hughes et
al., 2012). These retreat rates are also comparable to modern retreat rates observed at the largest
ice streams in northeastern (50 – 60 m/yr), and northwestern Greenland (40 – 50 m/yr; Wilson et
al., 2017).
- The exposure ages from both regions of East Greenland provide robust boundary conditions for ice
505 sheet models, improving the accuracy of simulating past GrIS change. These data refine our
understanding of the timing, rates and mechanisms of ice sheet retreat during the last deglaciation,
and provide important context for understanding contemporary rates of ice mass loss from the GrIS.

510 **Data availability.** All data described in the paper are included in the Supplementary files.

Author contributions

NEY, JPB and JMS designed the study. NEY, AB-K, KKP, CKW-G, BG, JPB and JMS conducted
fieldwork and sample collection. JTHA and KKP processed the samples and JTHA conducted sample
analysis and initial interpretations. JTHA prepared the manuscript with contributions from all authors.

515 **Competing interests**

This information product has been peer reviewed and approved for publication as a preprint by the U.S. Geological Survey.

Acknowledgements

520 We thank Volcano Heli for field support; Liza Wilson for assistance in the field; and Roseanne Schwartz, Jean Hanley, Kylie Seward, and Maya Lasker for laboratory assistance. Alan Hidy and Tyler Anderson are thanked for AMS measurements at Lawrence Livermore National Laboratory. The USGS internal reviewers are thanked for their feedback. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

525 Financial support

The research was supported by the National Science Foundation Office of Polar Programs (Awards #2105908 to NEY and JMS; #2106971 JPB).

References

- 530 Alley, R. B., Mayewski, P. A., Sowers, T., Stuiver, M., Taylor, K. C., & Clark, P. U. (1997). Holocene climatic instability: A prominent, widespread event 8200 yr ago. *Geology*, 25(6), 483. [https://doi.org/10.1130/0091-7613\(1997\)025<0483:HCIAPW>2.3.CO;2](https://doi.org/10.1130/0091-7613(1997)025<0483:HCIAPW>2.3.CO;2)
- An, L., Rignot, E., Wood, M., Willis, J. K., Mouginot, erémie, Khan, S. A., & analyzed data, S. (2021). *Ocean melting of the Zachariae Isstrøm and Nioghalvfjerdsfjorden glaciers, northeast Greenland*. <https://doi.org/10.1073/pnas.2015483118/-/DCSupplemental.y>
- 535 Andersen, J. K., Meyer, R. P., Huiban, F. S., Dømggaard, M. L., Millan, R., & Bjørk, A. A. (2025). Brief communication: Storstrømmen Glacier, northeastern Greenland, primed for end-of-decade surge. *Cryosphere*, 19(4), 1717–1724. <https://doi.org/10.5194/tc-19-1717-2025>
- Arndt, J. E., Jokat, W., & Dorschel, B. (2017). The last glaciation and deglaciation of the Northeast Greenland continental shelf revealed by hydro-acoustic data. *Quaternary Science Reviews*, 160, 45–56. <https://doi.org/10.1016/j.quascirev.2017.01.018>
- 540 Arndt, J. E., Jokat, W., Dorschel, B., Myklebust, R., Dowdeswell, J. A., & Evans, J. (2015). A new bathymetry of the Northeast Greenland continental shelf: Constraints on glacial and other processes. *Geochemistry, Geophysics, Geosystems*, 16(10), 3733–3753. <https://doi.org/10.1002/2015GC005931>
- 545 Balco, G., Stone, J. O., Lifton, N. A., & Dunai, T. J. (2008). A complete and easily accessible means of calculating surface exposure ages or erosion rates from ¹⁰Be and ²⁶Al measurements. *Quaternary Geochronology*, 3(3), 174–195. <https://doi.org/https://doi.org/10.1016/j.quageo.2007.12.001>

- 550 Bennike, O., Bjorck, S., Bocher, J., Hansen, L., Heinemeier, J., & Wohlfarth, B. (1999). Early Holocene Plant and Animal Remains from North-East Greenland. In *Source: Journal of Biogeography* (Vol. 26, Issue 3).
- Bevan, S. L., Luckman, A. J., & Murray, T. (2012). Glacier dynamics over the last quarter of a century at Helheim, Kangerdlugssuaq and 14 other major Greenland outlet glaciers. *Cryosphere*, 6(5), 923–937. <https://doi.org/10.5194/tc-6-923-2012>
- 555 Briner, J. P., Cuzzone, J. K., Badgley, J. A., Young, N. E., Steig, E. J., Morlighem, M., Schlegel, N. J., Hakim, G. J., Schaefer, J. M., Johnson, J. V., Lesnek, A. J., Thomas, E. K., Allan, E., Bennike, O., Cluett, A. A., Csatho, B., de Vernal, A., Downs, J., Larour, E., & Nowicki, S. (2020). Rate of mass loss from the Greenland Ice Sheet will exceed Holocene values this century. *Nature*, 586(7827), 70–74. <https://doi.org/10.1038/s41586-020-2742-6>
- 560 Clark, P. U., Dyke, A. S., Shakun, J. D., Carlson, A. E., Clark, J., Wohlfarth, B., Mitrovica, J. X., Hostetler, S. W., & McCabe, A. M. (2009). The Last Glacial Maximum. *Science*, 325(5941), 710–714. <https://doi.org/10.1126/science.1172873>
- 565 Cuzzone, J. K., Young, N. E., Morlighem, M., Briner, J. P., & Schlegel, N. J. (2022). Simulating the Holocene deglaciation across a marine-terminating portion of southwestern Greenland in response to marine and atmospheric forcings. *Cryosphere*, 16(6), 2355–2372. <https://doi.org/10.5194/tc-16-2355-2022>
- 570 Davies, J., Mathiasen, A. M., Kristiansen, K., Hansen, K. E., Wacker, L., Alstrup, A. K. O., Munk, O. L., Pearce, C., & Seidenkrantz, M. S. (2022). Linkages between ocean circulation and the Northeast Greenland Ice Stream in the Early Holocene. *Quaternary Science Reviews*, 286. <https://doi.org/10.1016/j.quascirev.2022.107530>
- Denton, G. H., Alley, R. B., Comer, G. C., & Broecker, W. S. (2005). The role of seasonality in abrupt climate change. *Quaternary Science Reviews*, 24(10–11), 1159–1182. <https://doi.org/10.1016/j.quascirev.2004.12.002>
- 575 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., Henderson, G. M., Okuno, J., & Yokoyama, Y. (2012). Ice-sheet collapse and sea-level rise at the Bølling warming 14,600 years ago. *Nature*, 483(7391), 559–564. <https://doi.org/10.1038/nature10902>
- Dowdeswell, J. A., Uenzelmann-Neben, G., Whittington, R. J., & Marienfeld, P. (1994). The Late Quaternary sedimentary record in Scoresby Sund, East Greenland. *Boreas*, 23(4), 294–310. <https://doi.org/10.1111/j.1502-3885.1994.tb00602.x>
- 580 Dowdeswell, J. A., Whittington, R. J., & Marienfeld, P. (1994). The origin of massive diamicton facies by iceberg rafting and scouring, Scoresby Sund, East Greenland. *Sedimentology*, 41(1), 21–35. <https://doi.org/10.1111/j.1365-3091.1994.tb01390.x>
- 585 Dyke, L. M., Hughes, A. L. C., Murray, T., Hiemstra, J. F., Andresen, C. S., & Rodés, Á. (2014). Evidence for the asynchronous retreat of large outlet glaciers in southeast Greenland at the end of the last glaciation. *Quaternary Science Reviews*, 99, 244–259. <https://doi.org/10.1016/j.quascirev.2014.06.001>
- Evans, J., Cofaigh, C. Ó., Dowdeswell, J. A., & Wadhams, P. (2009). Marine geophysical evidence for former expansion and flow of the Greenland Ice Sheet across the north-east Greenland continental shelf. *Journal of Quaternary Science*, 24(3), 279–293. <https://doi.org/10.1002/jqs.1231>

- 590 Funder, S. (1978). *Holocene stratigraphy and vegetation history in the Scoresby Sund area, East Greenland Seven plates in pocket 1978*.
- Funder, S., & Hansen, L. (1996). The Greenland ice sheet - a model for its culmination and decay during and after the last glacial maximum. *Bulletin of the Geological Society of Denmark*, 42, 137–152. <https://doi.org/10.37570/bgsd-1995-42-12>
- 595 Funder, S., Kjeldsen, K. K., Kjær, K. H., & O Cofaigh, C. (2011). The Greenland Ice Sheet During the Past 300,000 Years: A Review. In *Developments in Quaternary Science* (Vol. 15). <https://doi.org/10.1016/B978-0-444-53447-7.00050-7>
- Funder, S. V. (1990). *Scoresby Sund, sheet 12, Descriptive text to Quaternary map of Greenland 1:500.000*. .
- 600 Håkansson, L., Briner, J., Alexanderson, H., Aldahan, A., & Possnert, G. (2007a). 10Be ages from central east Greenland constrain the extent of the Greenland ice sheet during the Last Glacial Maximum. *Quaternary Science Reviews*, 26(19–21), 2316–2321. <https://doi.org/10.1016/j.quascirev.2007.08.001>
- 605 Håkansson, L., Graf, A., Strasky, S., Ivy-ochs, S., Kubik, P.W., Hjort, C., and Schlüchter, C. (2007b). Cosmogenic 10Be-ages from the Store Koldewey island, NE Greenland: *Geografiska Annaler. Series A. Physical Geography*, v. 89, no. 3, p. 195–202, <https://doi.org/10.1111/j.1468-0459.2007.00318.x>.
- Hall, B., Baroni, C., Denton, G., Kelly, M. A., & Lowell, T. (2008). Relative sea-level change, Kjøve Land, Scoresby Sund, East Greenland: Implications for seasonality in Younger Dryas time. *Quaternary Science Reviews*, 27(25–26), 2283–2291. <https://doi.org/10.1016/j.quascirev.2008.08.001>
- 610 Hall, B. L., Baroni, C., & Denton, G. H. (2010). Relative sea-level changes, Schuchert Dal, East Greenland, with implications for ice extent in late-glacial and Holocene times. *Quaternary Science Reviews*, 29(25–26), 3370–3378. <https://doi.org/10.1016/j.quascirev.2010.03.013>
- 615 Hanebuth, T., Stattegger, K., & Grootes, P. M. (2000). Rapid Flooding of the Sunda Shelf: A Late-Glacial Sea-Level Record. *Science*, 12. <https://doi.org/10.1126/science.288.5468.1033>
- Hansen, K. E., Lorenzen, J., Davies, J., Wacker, L., Pearce, C., & Seidenkrantz, M. S. (2022). Deglacial to Mid Holocene environmental conditions on the northeastern Greenland shelf, western Fram Strait. *Quaternary Science Reviews*, 293. <https://doi.org/10.1016/j.quascirev.2022.107704>
- 620 Hansen, L. (2001). *Landscape and Coast Development of a Lowland Fjord Margin following Deglaciation, East Greenland*. <https://doi.org/https://www.jstor.org/stable/521544>
- Heaton, T. J., Köhler, P., Butzin, M., Bard, E., Reimer, R. W., Austin, W. E. N., Bronk Ramsey, C., Grootes, P. M., Hughen, K. A., Kromer, B., Reimer, P. J., Adkins, J., Burke, A., Cook, M. S., Olsen, J., & Skinner, L. C. (2020). Marine20 - The Marine Radiocarbon Age Calibration Curve (0-55,000 cal BP). *Radiocarbon*, 62(4), 779–820. <https://doi.org/10.1017/RDC.2020.68>
- 625 Hughes, A. L. C., Rainsley, E., Murray, T., Fogwill, C. J., Schnabel, C., & Xu, S. (2012). Rapid response of Helheim Glacier, southeast Greenland, to early Holocene climate warming. *Geology*, 40(5), 427–430. <https://doi.org/10.1130/G32730.1>

- 630 Hvidberg, C. S., Grinsted, A., Dahl-Jensen, D., Khan, S. A., Kusk, A., Andersen, J. K., Neckel, N.,
Solgaard, A., Karlsson, N. B., Kjar, H. A., & Vallelonga, P. (2020). Surface velocity of the
Northeast Greenland Ice Stream (NEGIS): Assessment of interior velocities derived from satellite
data by GPS. *Cryosphere*, 14(10), 3487–3502. <https://doi.org/10.5194/tc-14-3487-2020>
- 635 Ingólfsson, Ó., Lyså, A., Funder, S., Möller, P., & Björck, S. (1994). Late Quaternary glacial history of
the central west coast of Jameson Land, East Greenland. *Boreas*, 23(4), 447–458.
<https://doi.org/10.1111/j.1502-3885.1994.tb00612.x>
- IPCC. (2021). *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to
the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge
University Press. <https://doi.org/10.1017/9781009157896>
- 640 Ivy-Ochs, S., Briner, J. P. (2014). Dating Disappearing Ice with Cosmogenic Nuclides. *Elements*, 10 (5)
351-356 [doi:10.2113/gselements.10.5.351](https://doi.org/10.2113/gselements.10.5.351)
- Jakobsson, M., Mohammad, R., Karlsson, M., Salas-Romero, S., Vacek, F., Heinze, F., Bringensparr, C.,
Castro, C. F., Johnson, P., Kinney, J., Cardigos, S., Bogonko, M., Accettella, D., Ambblas, D., An,
L., Bohan, A., Brandt, A., Bünz, S., Canals, M., ... Mayer, L. (2024). The International Bathymetric
Chart of the Arctic Ocean Version 5.0. *Scientific Data*, 11(1), 1420. [https://doi.org/10.1038/s41597-](https://doi.org/10.1038/s41597-024-04278-w)
645 024-04278-w
- Jennings, A. E., Hald, M., Smith, M., & Andrews, J. T. (2006). Freshwater forcing from the Greenland
Ice Sheet during the Younger Dryas: Evidence from southeastern Greenland shelf cores. *Quaternary
Science Reviews*, 25(3–4), 282–298. <https://doi.org/10.1016/j.quascirev.2005.04.006>
- 650 Joughin, I., Smith, B. E., Shean, D. E., & Floricioiu, D. (2014). Brief communication: Further summer
speedup of jakobshavn isbræ. *Cryosphere*, 8(1), 209–214. <https://doi.org/10.5194/tc-8-209-2014>
- Kelly, M.A., Lowell, T.V., Hall, B.L., Levy, L.B., Smith, C.A., Salamido, K., Schwartz, R. and Howley,
J.A. (2025). Relative summer temperature changes from glacial fluctuations in the Scoresby Sund
region, Central East Greenland, during late-glacial time. *Quaternary Science Reviews*, 367,
p.109531.
- 655 Kelly, M. A., Lowell, T. V., Hall, B. L., Schaefer, J. M., Finkel, R. C., Goehring, B. M., Alley, R. B., &
Denton, G. H. (2008). A 10Be chronology of lateglacial and Holocene mountain glaciation in the
Scoresby Sund region, east Greenland: implications for seasonality during lateglacial time.
Quaternary Science Reviews, 27(25–26), 2273–2282.
<https://doi.org/10.1016/j.quascirev.2008.08.004>
- 660 Kelly, M., & Bennike, O. (1992). Quaternary geology of western and central North Greenland. *Rapport
Grønlands Geologiske Undersøgelse*, 153, 1–34. <https://doi.org/10.34194/rapgggu.v153.8164>
- Khan, S. A., Colgan, W., Neumann, T. A., van den Broeke, M. R., Brunt, K. M., Noël, B., Bamber, J. L.,
Hassan, J., & Björk, A. A. (2022). Accelerating Ice Loss From Peripheral Glaciers in North
Greenland. *Geophysical Research Letters*, 49(12). <https://doi.org/10.1029/2022GL098915>
- 665 Kokfelt, T. F., Willerslev, E., Bjerager, M., Heijboer, T., Keulen, N., Larsen, L. M., Pedersen, C. B.,
Pedersen, M., Weng, W. L., Walentin, K. T., Søndersholm, M., & Svennevig, K. (2023). *Digital
1:500 000 scale geological map of Greenland, version 2.0*.
<https://doi.org/10.22008/FK2/FWX5ET>

- 670 Kuijpers, A., Troelstra, S. R., Prins, M. A., Linthout, K., Akhmetzhanov, A., Bouryak, S., Bachmann, M.
F., Lassen, S., Rasmussen, S., & Jensen, J. B. (2003). Late Quaternary sedimentary processes and
ocean circulation changes at the Southeast Greenland margin. *Marine Geology*, 195(1–4), 109–129.
[https://doi.org/10.1016/S0025-3227\(02\)00684-9](https://doi.org/10.1016/S0025-3227(02)00684-9)
- Lal, D. (1991). Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion
models. *Earth and Planetary Science Letters*, 104(2), 424–439.
- 675 Landvik, J. Y. (1994). The last glaciation of Germania Land and adjacent areas, northeast Greenland.
Journal of Quaternary Science, 9(1), 81–92. <https://doi.org/10.1002/jqs.3390090108>
- Larsen, N. K., Funder, S., Kjær, K. H., Kjeldsen, K. K., Knudsen, M. F., & Linge, H. (2014). Rapid early
Holocene ice retreat in West Greenland. *Quaternary Science Reviews*, 92, 310–323.
<https://doi.org/10.1016/j.quascirev.2013.05.027>
- 680 Larsen, N. K., Levy, L. B., Carlson, A. E., Buizert, C., Olsen, J., Strunk, A., Bjørk, A. A., & Skov, D. S.
(2018). Instability of the Northeast Greenland Ice Stream over the last 45,000 years. *Nature
Communications*, 9(1). <https://doi.org/10.1038/s41467-018-04312-7>
- Larsen, N. K., Søndergaard, A. S., Levy, L. B., Olsen, J., Strunk, A., Bjørk, A. A., & Skov, D. (2020).
Contrasting modes of deglaciation between fjords and inter-fjord areas in eastern North Greenland.
685 *Boreas*, 49(4), 903–917. <https://doi.org/10.1111/bor.12475>
- Larsen, N. K., Søndergaard, A. S., Levy, L. B., Strunk, A., Skov, D. S., Bjørk, A., Khan, S. A., & Olsen,
J. (2022). Late glacial and Holocene glaciation history of North and Northeast Greenland. *Arctic,
Antarctic, and Alpine Research*, 54(1), 294–313. <https://doi.org/10.1080/15230430.2022.2094607>
- Leger, T. P. M., Clark, C. D., Huynh, C., Jones, S., Ely, J. C., Bradley, S. L., Diemont, C., & Hughes, A.
690 L. C. (2024). A Greenland-wide empirical reconstruction of paleo ice sheet retreat informed by ice
extent markers: PaleoGrIS version 1.0. *Climate of the Past*, 20(3), 701–755.
<https://doi.org/10.5194/cp-20-701-2024>
- Lesnek, A. J., Briner, J. P., Young, N. E., & Cuzzone, J. K. (2020). Maximum Southwest Greenland Ice
Sheet Recession in the Early Holocene. *Geophysical Research Letters*, 47(1).
695 <https://doi.org/10.1029/2019GL083164>
- Levy, L. B., Kelly, M. A., Howley, J. A., & Virginia, R. A. (2012). Age of the Ørkendalen moraines,
Kangerlussuaq, Greenland: Constraints on the extent of the southwestern margin of the Greenland
Ice Sheet during the Holocene. *Quaternary Science Reviews*, 52, 1–5.
<https://doi.org/10.1016/j.quascirev.2012.07.021>
- 700 Levy, L. B., Kelly, M. A., Lowell, T. V., Hall, B. L., Howley, J. A., & Smith, C. A. (2016a). Coeval
fluctuations of the Greenland ice sheet and a local glacier, central East Greenland, during late glacial
and early Holocene time. *Geophysical Research Letters*, 43(4), 1623–1631.
<https://doi.org/10.1002/2015GL067108>
- 705 Levy, L. B., Kelly, M. A., Lowell, T. V., Hall, B. L., Howley, J. A., & Smith, C. A. (2016b). Coeval
fluctuations of the Greenland ice sheet and a local glacier, central East Greenland, during late glacial
and early Holocene time. *Geophysical Research Letters*, 43(4), 1623–1631.
<https://doi.org/10.1002/2015GL067108>

- 710 Levy, L. B., Larsen, N. K., Knudsen, M. F., Egholm, D. L., Bjørk, A. A., Kjeldsen, K. K., Kelly, M. A.,
Howley, J. A., Olsen, J., Tikhomirov, D., Zimmerman, S. R. H., & Kjær, K. H. (2020). Multi-phased
deglaciation of south and southeast Greenland controlled by climate and topographic setting.
Quaternary Science Reviews, 242. <https://doi.org/10.1016/j.quascirev.2020.106454>
- Long, A. J., Roberts, D. H., & Dawson, S. (2006). Early Holocene history of the west Greenland Ice
Sheet and the GH-8.2 event. *Quaternary Science Reviews*, 25(9–10), 904–922.
<https://doi.org/10.1016/j.quascirev.2005.07.002>
- 715 López-Quirós, A., Junna, T., Davies, J., Andresen, K. J., Nielsen, T., Haghipour, N., Wacker, L., Olsen
Alstrup, A. K., Munk, O. L., Rasmussen, T. L., Pearce, C., & Seidenkrantz, M. S. (2024). Retreat
patterns and dynamics of the former Norske Trough ice stream (NE Greenland): An integrated
geomorphological and sedimentological approach. *Quaternary Science Reviews*, 325.
<https://doi.org/10.1016/j.quascirev.2023.108477>
- 720 Marienfeld, P. (1991). *Evolution of sedimentation in Scoresby Sund, East-Greenland during the Holocene*
[PhD thesis, University of Bremen].
- Mouginot, J., Bjørk, A. A., Millan, R., Scheuchl, B., & Rignot, E. (2018). Insights on the Surge Behavior
of Storstrømmen and L. Bistrup Bræ, Northeast Greenland, Over the Last Century. *Geophysical
Research Letters*, 45(20), 11,197–11,205. <https://doi.org/10.1029/2018GL079052>
- 725 Mouginot, J., Rignot, E., Scheuchl, B., Fenty, I., Khazendar, A., Morlighem, M., Buzzi, A., & Paden, J.
(2015). Fast retreat of Zachariæ Isstrøm, northeast Greenland. *Science*, 350(6266), 1357–1361.
<https://doi.org/10.1126/science.aac7111>
- NGRIP members. (2004). High-resolution record of Northern Hemisphere climate extending into the last
interglacial period. *Nature*, 431(7005), 147–151. <https://doi.org/10.1038/nature02805>
- 730 Nishiizumi, K., Winterer, E.L., Kohl, C.P., Klein, J., Middleton, R., Lal, D. and Arnold, J.R., (1989).
Cosmic ray production rates of ^{10}Be and ^{26}Al in quartz from glacially polished rocks. *Journal of
Geophysical Research: Solid Earth*, 94(B12), pp.17907-17915.
- Nishiizumi, K., Imamura, M., Caffee, M. W., Southon, J. R., Finkel, R. C., & McAninch, J. (2007).
Absolute calibration of ^{10}Be AMS standards. *Nuclear Instruments and Methods in Physics
Research Section B: Beam Interactions with Materials and Atoms*, 258(2), 403–413.
735 <https://doi.org/10.1016/j.nimb.2007.01.297>
- Ó Cofaigh, C., Dowdeswell, J. A., Evans, J., Kenyon, N. H., Taylor, J., Mienert, J., & Wilken, M. (2004).
Timing and significance of glacially influenced mass-wasting in the submarine channels of the
Greenland Basin. *Marine Geology*, 207(1–4), 39–54. <https://doi.org/10.1016/j.margeo.2004.02.009>
- 740 Ó Cofaigh, C., Lloyd, J. M., Callard, S. L., Gebhardt, C., Streuff, K. T., Dorschel, B., Smith, J. A., Lane,
T. P., Jamieson, S. S. R., Kanzow, T., & Roberts, D. H. (2025). Shelf-edge glaciation offshore of
northeast Greenland during the last glacial maximum and timing of initial ice-sheet retreat.
Quaternary Science Reviews, 359, 109326. <https://doi.org/10.1016/j.quascirev.2025.109326>
- 745 Olsen, I. L., Laberg, J. S., Forwick, M., Rydningen, T. A., & Husum, K. (2022). Late Weichselian and
Holocene behavior of the Greenland Ice Sheet in the Kejser Franz Josef Fjord system, NE
Greenland. *Quaternary Science Reviews*, 284. <https://doi.org/10.1016/j.quascirev.2022.107504>

- Olsen, I. L., Rydningen, T. A., Forwick, M., Laberg, J. S., & Husum, K. (2020). Last glacial ice sheet dynamics offshore NE Greenland - A case study from Store Koldewey Trough. *Cryosphere*, 14(12), 4475–4494. <https://doi.org/10.5194/tc-14-4475-2020>
- 750 O'Regan, M., Cronin, T. M., Reilly, B., Alstrup, A. K. O., Gemery, L., Golub, A., Mayer, L. A., Morlighem, M., Moros, M., Munk, O. L., Nilsson, J., Pearce, C., Detlef, H., Stranne, C., Vermassen, F., West, G., & Jakobsson, M. (2021). The Holocene dynamics of Ryder Glacier and ice tongue in north Greenland. *The Cryosphere*, 15(8), 4073–4097. <https://doi.org/10.5194/tc-15-4073-2021>
- 755 Pearce, C., Özdemir, K. S., Forchhammer Mathiasen, R., Detlef, H., & Olsen, J. (2023). The marine reservoir age of Greenland coastal waters. *Geochronology*, 5(2), 451–465. <https://doi.org/10.5194/gchron-5-451-2023>
- 760 Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard-Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., & Ruth, U. (2006). A new Greenland ice core chronology for the last glacial termination. *Journal of Geophysical Research Atmospheres*, 111(6). <https://doi.org/10.1029/2005JD006079>
- Reeh, N., Bøggild, C. E., & Oerter, H. (1994). *Surge of Storstrømmen, a large outlet glacier from the Inland lee of North-East Greenland*. <https://doi.org/https://doi.org/10.34194/rapggu.v162.8263>
- 765 Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., ... Talamo, S. (2020). The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon*, 62(4), 725–757. <https://doi.org/10.1017/RDC.2020.41>
- 770 Reimer, P. J., & Reimer, R. W. (2001). A marine reservoir correction database and on-line interface. *Radiocarbon*, 43(2 PART I), 461–463. <https://doi.org/10.1017/s0033822200038339>
- Rignot, E., Bjork, A., Chauche, N., & Klauke, I. (2022). Storstrømmen and L. Bistrup Bræ, North Greenland, Protected From Warm Atlantic Ocean Waters. *Geophysical Research Letters*, 49(5). <https://doi.org/10.1029/2021GL097320>
- 775 Roberts, D. H., Lane, T. P., Jones, R. S., Bentley, M. J., Darvill, C. M., Rodes, A., Smith, J. A., Jamieson, S. S. R., Rea, B. R., Fabel, D., Gheorghiu, D., Davidson, A., Cofaigh, C., Lloyd, J. M., Callard, S. L., & Humbert, A. (2024). The deglacial history of 79N glacier and the Northeast Greenland Ice Stream. *Quaternary Science Reviews*, 336. <https://doi.org/10.1016/j.quascirev.2024.108770>
- 780 Schaefer, J. M., Denton, G. H., Kaplan, M., Putnam, A., Finkel, R. C., Barrell, D. J. A., Andersen, B. G., Schwartz, R., Mackintosh, A., Chinn, T., & Schlüchter, C. (2009). *High-Frequency Holocene Glacier Fluctuations in New Zealand Differ from the Northern Signature*. <https://www.science.org>
- Skov, D. S., Andersen, J. L., Olsen, J., Jacobsen, B. H., Knudsen, M. F., Jansen, J. D., Larsen, N. K., & Egholm, D. L. (2020). Constraints from cosmogenic nuclides on the glaciation and erosion history of Dove Bugt, northeast Greenland. *Bulletin of the Geological Society of America*, 132(11–12), 2282–2294. <https://doi.org/10.1130/b35410.1>
- 785 Stein, R., Nam, S., Grobe H., & Hubberton, H. (1996). *Late Quaternary glacial history and short-term ice-rafted debris fluctuations along the East Greenland continental margin* (J. A. W. E. N. B. H. &

J. A. E. Andrews, Ed.; Vol. 111). Geological Society of London, Special Publication.
<https://doi.org/https://doi.org/10.1144/GSL.SP.1996.111.01.09>

- 790 Stone, J. O. (2000). Air pressure and cosmogenic isotope production. *Journal of Geophysical Research: Solid Earth* (1978–2012), 105(B10), 23723–23753.
- Stuiver, M., & Reimer, P. J. (1993). Extended 14C data base and revised CALIB 3.0 14C age calibration program. *Radiocarbon*, 35(1), 215–230. <https://doi.org/10.1017/S0033822200013904>
- The IMBIE Team. (2020). Mass balance of the Greenland Ice Sheet from 1992 to 2018. *Nature*, 579(7798), 233–239. <https://doi.org/10.1038/s41586-019-1855-2>
- 795 Vijay, S., Khan, S. A., Kusk, A., Solgaard, A. M., Moon, T., & Bjørk, A. A. (2019). Resolving Seasonal Ice Velocity of 45 Greenlandic Glaciers With Very High Temporal Details. *Geophysical Research Letters*, 46(3), 1485–1495. <https://doi.org/10.1029/2018GL081503>
- Wager, L. R. (1947). Geological investigations in East Greenland. Part IV: The Stratigraphy and Tectonics of Knud Rasmussens Land and the Kangerdlugssuqq Region. *Meddelelser Om Grønland*, 3(134), 64.
- 800 Weidick, A., Andreassen, C., Oerter, H., & Reeh, N. (1996). Neoglacial Glacier Changes around Storstrommen, North-East Greenland. In *Polarforschung* (Vol. 64, Issue 3).
- Wilson, N., Straneo, F., & Heimbach, P. (2017). Satellite-derived submarine melt rates and mass balance (2011–2015) for Greenland’s largest remaining ice tongues. *Cryosphere*, 11(6), 2773–2782.
- 805 <https://doi.org/10.5194/tc-11-2773-2017>
- Young, N. E., Briner, J. P., Miller, G. H., Lesnek, A. J., Crump, S. E., Pendleton, S. L., Schwartz, R., & Schaefer, J. M. (2021). Pulsebeat of early Holocene glaciation in Baffin Bay from high-resolution beryllium-10 moraine chronologies. In *Quaternary Science Reviews* (Vol. 270). Elsevier Ltd. <https://doi.org/10.1016/j.quascirev.2021.107179>
- 810 Young, N. E., Briner, J. P., Miller, G. H., Lesnek, A. J., Crump, S. E., Thomas, E. K., Pendleton, S. L., Cuzzone, J., Lamp, J., Zimmerman, S., Caffee, M., & Schaefer, J. M. (2020). Deglaciation of the Greenland and Laurentide ice sheets interrupted by glacier advance during abrupt coolings. *Quaternary Science Reviews*, 229. <https://doi.org/10.1016/j.quascirev.2019.106091>
- 815 Young, N. E., Briner, J. P., Rood, D. H., Finkel, R. C., Corbett, L. B., & Bierman, P. R. (2013). Age of the Fjord Stade moraines in the Disko Bugt region, western Greenland, and the 9.3 and 8.2 ka cooling events. *Quaternary Science Reviews*, 60, 76–90. <https://doi.org/10.1016/j.quascirev.2012.09.028>
- 820 Young, N. E., Briner, J. P., Stewart, H. A. M., Axford, Y., Csatho, B., Rood, D. H., & Finkel, R. C. (2011). Response of Jakobshavn Isbræ, Greenland, to Holocene climate change. *Geology*, 39(2), 131–134. <https://doi.org/10.1130/G31399.1>
- Young, N. E., Schaefer, J. M., Briner, J. P., & Goehring, B. M. (2013). A 10Be production-rate calibration for the Arctic. *Journal of Quaternary Science*, 28(5), 515–526. <https://doi.org/10.1002/jqs.2642>