

~~Spring melting season~~ Melt period methane emissions in northern high latitude wetlands are governed by the length of the ~~season~~ period and presence of permafrost

Sara Hyvärinen¹, Maria K. Tenkanen¹, Aki Tsuruta¹, Anttoni Erkkilä¹, Kimmo Rautiainen¹, Hermanni Aaltonen¹, Motoki Sasakawa², and Tuula Aalto¹

¹Finnish Meteorological Institute, 00101 Helsinki, Finland

²National Institute for Environmental Studies, Ibaraki, Japan

Correspondence: Sara Hyvärinen (sara.hyvarinen@fmi.fi)

Abstract. Northern high latitude wetlands are significant sources of methane, with emissions driven by seasonal soil freezing and thawing. To better understand the seasonality of northern high latitude methane emissions, we defined the ~~spring melting season~~ melt period occurring in spring time using the remote sensing Soil Moisture and Ocean Salinity Freeze/Thaw data from 2011–2021. To estimate methane emissions in the northern high latitudes, we used the atmospheric inverse model CarbonTracker Europe-CH₄. The ~~melting season~~ melt period was defined for three permafrost zones and for a seasonally frozen non-permafrost region using two approaches: region-based, which considered climatological conditions of permafrost regions, and grid-based, which defines the ~~melting season~~ melt period at a finer 1° × 1° scale.

The length and timing of the ~~melting season~~ melt period varied significantly depending on the approach. The ~~melting season~~ melt period generally occurred between March and June and was influenced by ~~the~~ air temperature, with a negative correlation between the length and the mean temperature of the melt period. The longest ~~melting season~~ melt period was in the non-permafrost zone and the shortest varied between the two methods. The ~~spring melting season~~ melt period emissions were on average 1.83 Tg with the region-based approach and 0.45 Tg with the grid-based approach, the non-permafrost zone having the largest share of the ~~spring emissions~~. ~~The emissions~~ emissions. They were largely dependent on the season's length. Year-to-year variation was modest, within 15 % (region-based) and 23 % (grid-based) of average emissions, and there was also no trend during the study period. Our dual-method approach allows for robust comparison with both large-scale regional studies and localized site-level research.

1 Introduction

Northern high latitude wetlands are an important and dynamic part of the climate system (Hugelius et al., 2020). They are a large source of methane (CH₄), which is the second most important anthropogenic greenhouse gas causing climate change after carbon dioxide (CO₂) ~~,-having~~ (Forster et al., 2021). It has a 29.8 times stronger global warming potential ~~in over a~~ 100- year timescale than CO₂ without considering climate feedbacks (Forster et al., 2021). A large portion of the total soil carbon is stored in northern wetlands and the underlying permafrost (Scharlemann et al., 2014; Hugelius et al., 2020) ~~but due~~ , containing

~ 80 % (415 ± 150 Pg C (Hugelius et al., 2020)) of the total global peatland carbon, with nearly half being permafrost affected peatland (Hugelius et al., 2020; Scharlemann et al., 2014). Due to climate change and Arctic amplification, thawing permafrost could affect ~~the this~~ carbon stock (Schuur et al., 2015; Knoblauch et al., 2018; Voigt et al., 2019; Turetsky et al., 2020). Even ~~if the permafrost does not fully thaw, the deepening of the active layer — the top layer of soil that thaws in summer and freezes in winter — can still release a significant amount of carbon.~~ Permafrost ~~though permafrost~~ thaw will likely lead to soil drying and increased drainage, ~~potentially accelerating which potentially accelerates~~ organic matter decomposition and CO₂ emissions while suppressing CH₄ emissions (Lawrence et al., 2015) ~~but,~~ the effect of the released CH₄ might be as large due to its stronger global warming potential (Schuur and Abbott, 2011). ~~According to Neumann et al. (2019), increasing~~ Increasing rainfall and warming soils could increase near-term global warming, and the total annual boreal ~~methane~~ CH₄ emissions could rise 4 Tg per year (Neumann et al., 2019). The study by Poulter et al. (2017) using a biochemical model concluded that boreal wetland CH₄ emissions have already increased by 1.2 Tg yr⁻¹ between 2000–2012. ~~Even if the permafrost does not fully thaw, the deepening of the active layer – the top layer of soil that thaws in summer and freezes in winter – can still release a significant amount of carbon.~~

In addition to permafrost thaw, Arctic amplification is expected to significantly impact snowmelt and seasonal soil thawing, including the active layer, in northern high latitude wetlands. There are still a lot of open questions related to the future of methane emissions in the northern high latitudes due to uncertainties in the wetness of the area and possible feedback loops. For instance, de Vrese et al. (2023) demonstrated that atmospheric feedbacks resulting from ~~the~~ increasingly dry Arctic may offset the effects of growing wetland extent and result in comparable CH₄ emissions as in the case where the Arctic would remain wet. The changing Arctic hydrology affects the ~~spring melting season~~ melt period emissions as well (Rawlins and Karmalkar, 2024). The hydrological conditions during the ~~spring melting season~~ melt period in the northern high latitudes are especially uncertain due to the unreliability in the amount of snow as well as the melting and evaporation of snow. This lack of certainty brings unpredictability to the ~~melting season~~ melt period methane emissions.

CH₄ emissions from the northern high latitude wetlands are low during winter when the soil is frozen and high during summer after the soil has thawed (Aselmann and Crutzen, 1989; Rinne et al., 2018), also depending on the soil temperature and hydrological factors (Rychlik, 2009; Zhu et al., 2013; Turetsky et al., 2008). During the spring thawing, the emissions increase rapidly. Typically, the CH₄ storage is low during the ~~spring~~ thawing after the dormant cold season, and the CH₄ is released gradually as the soil melts (Raz-Yaseef et al., 2017). However, previous studies have reported large bursts of CH₄ from the wetlands during the spring thawing (Jin et al., 1999; Tokida et al., 2007; Song et al., 2012; Raz-Yaseef et al., 2017), though the reports are scarce with only a few measurement sites and just a few years of flux measurements from before the thawing starts (e.g. winter flux data) (Raz-Yaseef et al., 2017). The springtime bursts of CH₄ have been linked to rain-on-snow events which enhance soil cracking (Raz-Yaseef et al., 2017). However, the frequency and impact of these pulses to spring methane emissions are still highly uncertain. Our research did not focus on these springtime bursts, but that does not negate their existence. We focused on large scale methane emissions in the northern high latitudes.

Previous studies have mostly focused on methane emissions during the growing season while the other seasons, such as the autumn freezing period, winter season and the spring thaw season, has received less attention (e.g. Vourlitis et al., 1993; Sachs

et al., 2008; Zona et al., 2009; Parmentier et al., 2011). The regional studies focusing on the spring season, have often defined the timing of the ~~spring thaw~~ whole spring season simply as certain months, e.g. March, April and May (e.g. Castro-Morales et al., 2018; Ito et al., 2023) or in a case of site-level studies, using site-specific soil temperature measurements (e.g. Zona et al., 2016; Bao et al., 2020; Tagesson et al., 2012; Raz-Yaseef et al., 2017). Using soil temperature to define the soil thawing would be ideal, but ~~a~~ reliable soil temperature data is available only from point-wise measurements. Thus, to be able to study ~~spring season~~ the melt period on a regional scale, we need to use a proxy for the soil thawing. In this study, we define the ~~spring melting season~~ melt period in the spring time using the remote-sensed Soil Moisture and Ocean Salinity (SMOS) soil Freeze/Thaw (F/T) data (~~Rautiainen et al., 2016~~) (Rautiainen et al., 2025). The SMOS F/T data provides daily information of the freezing and thawing state of the soil in the northern high latitudes at ~~the a~~ resolution of 25 km. Using the SMOS F/T data gives us a dynamic picture of the soil thawing including the active layer of the permafrost during the spring and enables us to define the ~~melting season~~ melt period for the whole northern high latitude ~~regions~~ region instead of using a static definition (e.g. specific months). This allows us to focus on the CH₄ emissions during the actual thawing of the soil instead of the whole spring season with inverse modeling. The SMOS F/T data have been used successfully to define summer thaw, autumn freezing, and winter cold seasons in the northern high latitude wetlands (Tenkanen et al., 2021; Erkkilä et al., 2023). In this paper, the SMOS F/T data is used for the first time to define the ~~spring melting season~~ melt period.

In addition to in-situ measurement based studies, the spring CH₄ emissions ~~has~~ have been studied with process-based models (Ito et al., 2023; Castro-Morales et al., 2018). Process-based models estimate methane fluxes by simulating physical, chemical, and biological processes. Another ~~type of modeling approach to estimate fluxes~~ is inverse modeling. Inverse models are statistical approaches which can be used to inform and "re-evaluate" the process-based estimates and decrease the uncertainties in CH₄ emissions using atmospheric CH₄ measurements, so that the differences between simulations and measurements is minimized (Wittig, 2023).

The aim of this study is to estimate the ~~spring melting season and its~~ melt period occurring during the spring season and the corresponding CH₄ emissions in the northern high latitude permafrost and wetland regions. We use the SMOS F/T data to define the ~~spring melting season~~ melt period and determine methane emissions with the global atmospheric inverse model Carbon Tracker Europe-CH₄ (CTE-CH₄) (Tsuruta et al., 2017) at high spatial resolution of 1° latitude × 1° longitude. The years studied are 2011–2021 due to availability of the SMOS data. Previously, CTE-CH₄ has been used to study northern high latitude wetland emissions for various seasons (Tenkanen et al., 2021; Erkkilä et al., 2023), but this is the first time with a focus on ~~spring melting season. The spring melting season~~ the melt period. The melt period is defined for four different permafrost zones (sporadic, discontinuous and continuous permafrost and a seasonally frozen zone non-permafrost) with a region-based approach, as well as individually for each 1° × 1° grid cell ~~to determine the melting season~~ with a grid-based approach. For comparison between the two methods to define the melt period, we divided the grid-based melt period into the same four permafrost zones. This allows us to estimate emissions both in climatological and local scales.

90 2 Materials and methods

2.1 SMOS F/T soil state estimates

To define the ~~melting season~~ melt period of the northern high latitude wetlands, we used the European Space Agency's (ESA) SMOS Soil Freeze and Thaw State product (version 3.0) (Rautiainen et al., 2016; European Space Agency, 2023). ~~It provides daily information on the soil state in the northern latitudes based on observations from the ESA SMOS satellite (Kerr et al., 2010). The SMOS payload instrument Microwave Imaging Radiometer with Aperture Synthesis measures the natural microwave radiation emitted by the Earth's surface at L-band ($f \approx 1.4$ GHz) (Kerr et al., 2010). The intensity of the natural microwave radiation is related to~~ The product is derived from observations of the ESA Soil Moisture and Ocean Salinity (SMOS) satellite (Kerr et al., 2010) and provides daily categorical information on near-surface freeze-thaw state at 25 km spatial resolution. The main observational input consists of Centre Aval de Traitement des Données SMOS (CATDS) daily gridded Level-3 brightness temperature data (Al Bitar et al., 2017). Brightness temperature depends on both the physical temperature of the target and its emissivity; it is quantified by the brightness temperature. Detection of the soil F/T state with SMOS data is based on the high contrast in permittivity between free surface and its microwave emissivity. At L-band frequencies, the strong contrast in dielectric properties between liquid water and ice at L-band (Rautiainen et al., 2016). Soil containing liquid water has a higher permittivity and therefore lower emissivity than frozen soil. This causes the brightness temperature of thawed soil to be lower than that of frozen soil. L-band measurements are especially useful for soil observations due to their relatively long wavelength because it is less affected by vegetation cover and cloud layers compared to higher frequencies. Additionally, the deeper penetration depth of L-band microwaves enables observations not only from the surface skin but also from the top 0–5 cm enables sensitivity to near-surface soil freeze-thaw transitions, with partial penetration through vegetation and the upper soil layer. Importantly, L-band refers to low frequency (1–2 GHz) passive microwave observations are not dependent on sunlight, which is important for determining the melting period in the northern high latitudes since the melting season can start already during the time when sunlight is limited due to the polar night (Rautiainen et al., 2025).

The SMOS F/T soil state detection algorithm utilizes Centre Aval de Traitement des Données SMOS daily gridded level 3 brightness temperature data (Al Bitar et al., 2017) as input data. Additionally, two ancillary data sets were used: European Centre for Medium-Range Weather Forecast (ECMWF) ERA5 2-m air temperature (Hersbach et al., 2020) and National Snow and Ice Data Center Interactive Multisensor Snow and Ice Mapping System snow cover (U.S. National Ice Center, 2004). ~~The F/T detection algorithm is threshold-based: observations are compared to empirically defined frozen and thawed soil references and categorized as frozen~~ L3FT retrieval was developed by the Finnish Meteorological Institute in collaboration with GAMMA Remote Sensing, and the full algorithm description and validation are given in Rautiainen et al. (2025). Briefly, the algorithm computes the normalized polarization ratio (NPR) from horizontally and vertically polarized SMOS brightness temperatures, applies filtering and temporal smoothing to the NPR time series, and rescales NPR using grid cell-specific frozen and thawed reference states. These reference states are estimated empirically from multi-year NPR time series using ancillary air temperature and snow information (Rautiainen et al., 2025). The resulting scaled NPR is classified into three soil-state categories using fixed thresholds: thawed (< 50 %), partially frozen, or thawed soil based on these thresholds (50–70 %), and

frozen (> 70 %). Of the three categories, the ~~thawing state of the soil~~ thawed soil state is used in this study to define the melting season. ~~Two distinct soil state products are provided: one using ascending orbit observations (6 AM) and the other using descending orbit observations (6 PM). In this study, we used only the ascending orbit data, as they are less affected by radio-frequency melt period.~~

Although the SMOS freeze-thaw product is available beyond 2021, we restricted the analysis to the period 2011–2021 to ensure temporal consistency and robust data quality across the study domain. ~~In the later part of the SMOS mission, increasing radio-frequency interference (RFI) over the Eurasian continent and were better for the purpose of our study since the melting snow during daytime affect the descending orbit observations in the evening.~~ has affected observations over Europe and parts of Eurasia. SMOS, as an L-band aperture-synthesis radiometer, is particularly sensitive to RFI, which reduces observation density and degrades retrieval reliability (Oliva et al., 2016; Rautiainen et al., 2025). The SMOS freeze-thaw product is provided separately for ascending (approximately 06:00 local time) and descending (approximately 18:00 local time) satellite overpasses, with operational RFI monitoring indicating orbit-dependent differences in data availability over Europe, the Middle East, and Asia (Uranga et al., 2022). To maximize data quality and spatial coverage, we therefore restricted the analysis to ascending-orbit SMOS observations within the 2011–2021 period.

The SMOS ~~FL3~~ freeze-thaw dataset used in this study is publicly available through the SMOS online dissemination service, with persistent access provided via the dataset DOI (<https://doi.org/10.57780/sm1-fbf89e0>) ~~(ESA) and from FMI (SMOS). The data are available from July 2010 until the current day with a latency of a couple of days. Its resolution is 25 km × 25 km. To combine with inverse model results, the resolution of the data was changed to 1° × 1° resolution by selecting the pixels whose centre was inside the 1° × 1° grid cell and calculating the fraction of pixels that were defined as thaw~~ (European Space Agency, 2023).

2.2 CarbonTracker Europe-CH₄

The methane fluxes were estimated with the CarbonTracker Europe-CH₄ (CTE-CH₄) inverse model (Tsuruta et al., 2017). In its assimilation phase, a Bayesian cost function was minimized:

$$J = (\mathbf{x} - \mathbf{x}^b)^T \mathbf{P}^{-1} (\mathbf{x} - \mathbf{x}^b) + (\mathbf{y} - H(\mathbf{x}))^T \mathbf{R}^{-1} (\mathbf{y} - H(\mathbf{x})), \quad (1)$$

where \mathbf{x} is a state vector which contains a set of scaling factors that multiply the prior CH₄ surface fluxes, which are meant to be optimized, and \mathbf{x}^b is the prior state vector (see Section 2.2.3). \mathbf{P} is the state vector error covariance matrix. \mathbf{y} is a vector consisting of the atmospheric methane observations (see Section 2.2.2) and \mathbf{R} is the error covariance matrix of the observations \mathbf{y} . H is an observation operator, which is an atmospheric transport model TM5 in this study (see Section 2.2.1).

The model used the ensemble Kalman filter (EnKF) (Evensen, 2003; Peters et al., 2005) data assimilation scheme within the CarbonTracker Data Assimilation Shell (CTDAS) (van der Laan-Luijkx et al., 2017) with an ensemble size of 500 and a 5-week lag to optimize the fluxes (Peters et al., 2005; Tsuruta et al., 2017). The anthropogenic and natural fluxes were optimized separately but simultaneously in a weekly temporal resolution. The fluxes in the high northern latitudes were optimized at the spatial resolution of 1° × 1°, and regionally elsewhere. The spatial correlation followed an exponential decay model (Peters

et al., 2005) with correlation lengths of 100 km for $1^\circ \times 1^\circ$ grid-based domains, 500 km for other land domains, and 900 km for oceanic domains. Anthropogenic and natural CH_4 fluxes were assumed to be uncorrelated, as were land and ocean domains.

2.2.1 TM5 chemistry model

160 TM5 is an atmospheric chemistry transport model (Krol et al., 2005). Here, it was used as the observation operator to transform the methane fluxes to atmospheric mole fractions. In this study, its global horizontal resolution was $4^\circ \times 6^\circ$ with an intermediate zoom region of $2^\circ \times 3^\circ$ (14°N – 82°N , 36°W – 54°E) framing a $1^\circ \times 1^\circ$ zoom over Europe (24°N – 74°N , 21°W – 45°E). The model used preprocessed meteorological data from ECMWF ERA5 reanalysis data with a 3-hour resolution (Hersbach et al., 2020). The vertical domain was divided into 25 hybrid sigma pressure levels from the surface to the upper atmosphere. The
165 chemical loss of CH_4 in the atmosphere to the sinks of OH, was based on monthly precalculated values by Houweling et al. (2014), and Cl and $\text{O}(^1\text{D})$ sinks were based on the atmospheric chemistry general circulation model ECHAM5/MESSy1 (Jöckel et al., 2006; Kangasaho et al., 2022). The variability of the atmospheric sinks between different years was not considered, but varied monthly, and the sinks were not optimized.

2.2.2 Observations

170 In addition to the observations from the ObsPack v4.0 (Schuldt et al., 2021; Masarie et al., 2014), observations from two stations in Finland (Kumpula, Sodankylä) (Tsuruta et al., 2019) and from nine stations in Siberia (Sasakawa et al., 2010, 2025) were used. All stations are listed in the Appendix A1 and the location can be seen in Fig A1. Globally, 183 stations had observations between ~~2011–2021~~2011–2021, with some stations having two or three institutions contributing. The data included weekly discrete air samples and hourly continuous measurements of CH_4 , and the data was filtered according to the institutions’
175 quality flags. Only data points that represented well-mixed conditions were included, which means that daily averages were calculated from 12 to 4 pm local time, except for high mountain sites where averages are calculated from 0 to 4 am local time, following Tsuruta et al. (2017).

Observational uncertainties, or "model–data mismatches", were estimated for each site based on site-specific factors, measurement accuracy, and the capability of TM5 to simulate atmospheric CH_4 mole fractions (Bruhwiler et al., 2014; Tsuruta
180 et al., 2019). These discrepancies arose from TM5’s resolution and transport errors, with e.g. better performance at remote marine sites compared to those affected by strong local emissions. Sites were categorized, for example, as marine boundary layer (4.5 ppb), terrestrial (25 ppb), mixed marine and terrestrial (15 ppb), and strong local influence (30 ppb). Uncertainties ranged from 4.5 to 75 ppb.

2.2.3 Prior fluxes

185 The prior anthropogenic emissions were taken from the Emission Database for Global Atmospheric Research (EDGAR v6.0) (Monforti Ferrario et al., 2021). The emissions from LPX-Bern DYPTOP v1.4 (Lienert and Joos, 2018) were used as the natural biospheric prior emissions. Methane emissions from other sources were: Weber et al. (2019) for ocean, the Global

fire emission database (GFED v4.1s) (van der Werf et al., 2017; Randerson et al., 2017) for biomass burning, and VISIT (Ito and Inatomi, 2012) (Ito and Inatomi, 2012; Tsuruta et al., 2019) for termites. ~~Other fluxes, excluding biospheric~~ Biospheric and anthropogenic fluxes ~~;~~ were optimized globally. Other fluxes were not optimized. We only analyze the optimized biospheric fluxes in this study. The biospheric prior LPX-Bern DYPTOP represents ecosystem area fractions across several land-cover types: (i) peatlands suitable for peat growth as defined by DYPTOP (Stocker et al., 2014); (ii) rice paddies coinciding with croplands and the presence of rice paddies (Spahni et al., 2011); (iii) inundated wetlands other than peatlands or rice paddies; (iv) wet mineral soils, which are not wetlands, peatlands, rice paddies, permanent freshwater bodies, or ice/sea water covered areas, but are occasionally wet; and (v) dry mineral soils, which are areas identical to wet mineral soils but are "dry" in general. Of these categories, we did not include rice paddies in our prior as they are not present in the northern high latitude region. Soilsink was included in the biospheric prior.

For both the anthropogenic and biospheric fluxes, we used the prior uncertainty of 80 % for terrestrial fluxes and 20 % for oceanic fluxes, assuming uncorrelated uncertainties, following previous studies (e.g., Tsuruta et al., 2017; Bruhwiler et al., 2014).

~~Extents of the four permafrost zones in $1^\circ \times 1^\circ$ resolution.~~

2.3 Permafrost map

The permafrost extent (v3.0) is one of the variables belonging to ESA's Climate Change Initiative (CCI) (Bartsch et al., 2020; Obu et al., 2019). The temporal resolution of the data is one year and the spatial resolution is 926.63 m (~~(Obu et al., 2021)~~) (Obu et al., 2021). Using the permafrost and SMOS F/T soil state data, we split the northern latitudes into four zones depending on how much of the exposed land surface is underlain by permafrost: 90–100 % for the continuous permafrost, 50–90 % for the discontinuous permafrost, 14–50 % for the sporadic zone and the non-permafrost zone mainly comprising areas that were frozen seasonally based on the SMOS F/T data with some potential permafrost spots, since the CCI permafrost data had ~~not no~~ values lower than 14 % (Fig.1). Areas where no SMOS F/T data was available, were excluded from this study. This masking excluded areas such as southern regions below 40° N where almost no permafrost is located, as well as oceans, most of Greenland, and some areas close to the Great Lakes in the USA.

The permafrost data was only available until 2018. Average permafrost zones were calculated from the years 2011–2018. The four permafrost zones had only minor changes between the different years. The ~~average~~ permafrost percentage was calculated ~~for each $1^\circ \times 1^\circ$ grid cell after the resolution was changed to at a $1^\circ \times 1^\circ$ from the resolution by averaging the original 1 km resolution by taking mean for each grid data within each grid cell.~~ The average areas were used in this study to define the four permafrost zones for all the years from 2011–2021. We are aware that a newer version of the permafrost data (v4.0) including the years 2019–2021 has been published but it was not used in this study (Westermann et al., 2024).

2.4 ERA5 2 m temperature data

ERA5 is the latest global reanalysis from the ECMWF available from 1959 to the present day (ECMWF; Hersbach et al., 2020). It provides hourly estimates of multiple land, oceanic and atmospheric climate variables. Here, the ERA5 2 m air temperature




Figure 1. Extents of the four permafrost zones in 1° × 1° resolution.

reanalysis data was used to study the relationship between the air temperature and CTE-CH₄ emissions during the **melting season**melt period. The ERA5 temperature data was additionally used as an auxiliary dataset in the SMOS F/T data and as input data in the CTE-CH₄.

2.5 Defining the **melting season**melt period and its methane emissions

225 2.5.1 Definition of the **melting season**melt period

In this study, the **melting season**melt period was defined as a season in the northern latitudes in spring, when the soil in a specific region turned from frozen to thawed based on the "thawed" state in the SMOS F/T data (see Section 2.1). The word "**melting**melt" is used here instead of "**thawing**thaw" because the SMOS F/T data can indicate the melting of the snow instead of the soil, especially at the beginning of the **melting season**melt period. This is because the microwave radiation signal from wet snow resembles the signal from thawed soil (Rautiainen et al., 2016). However, methane emissions are possible even at the very beginning of the **melting season**melt period because the air temperature rises above zero and melted or rain water can trickle into the soil and wintertime methane reserves are released (Hargreaves et al., 2001; Rinne et al., 2007; Song et al., 2012; Raz-Yaseef et al., . It is thus justified to start the **melting season**melt period from the melting of the snow. The boundaries used in this study were similar to the ones used by Erkkilä et al. (2023) to define different seasons in the northern high latitude wetlands, but instead
 235 of frozen or partially frozen state of the SMOS F/T data, we used the thawed soil state to define the melt period. The resolution

of the SMOS F/T data was changed from 25 km × 25 km to 1° × 1° by calculating the fraction of the thawed 25 km × 25 km pixels whose center was inside the 1° × 1° grid cell.

The ~~melting-season-melt period~~ was defined for four permafrost zones: non-permafrost, sporadic, discontinuous and continuous permafrost (region-based approach), and separately for each 1° × 1° grid cell (grid-based approach) in these zones. 240 The region-based approach gives information about the permafrost areas constrained by their specific climatological conditions while the grid-based ~~melting-season-melt period~~ was studied to illustrate the local variation in the melting of the soil.

With the region-based approach, the ~~melting-season-melt period~~ was set to start 1) when the mean thawing fraction of a permafrost zone had surpassed the minimum thawing fraction of that year by 0.1 ($\text{thaw}(\%) \geq \text{thaw}_{\text{min,year}}(\%) + 10\%$), and 2) ~~after the day when the last day in the spring, when the zone reaches its~~ minimum annual mean thawing fraction ~~was reached,~~ 245 ~~was surpassed before mid July.~~ During some years, the freezing of the soil continued past the turn of the year, which meant that the ~~first (1)~~ boundary was reached before the maximum freezing of the soil. This meant that ~~an additional-the additional~~ (2) condition for the beginning of the ~~melting-season-melt period~~ had to be defined. With the ~~second (2)~~ condition, the ~~melting season-melt period~~ could be separated from the autumn freezing period. In regions with permafrost, the first (1) condition was surpassed later during the spring than the second (2) condition, but in the non-permafrost zone the second (2) condition was 250 needed.

The season ended when the mean thawing fraction of the whole zone had surpassed 0.8 of the maximum thawing fraction of all years which was 1 for all zones ($\text{thaw}(\%)/\text{thaw}_{\text{max,all}}(\%) \geq 80\%$). The fraction of 0.8 was chosen because there was not as much variation in the thawing fraction that ~~late in the spring close to summer~~ indicating a stable thaw state. The day when the ~~melting-season-melt period~~ ended was included in the ~~melting-season. The size of the grid-cells in the north-south~~ 255 ~~direction was taken into account when defining the melting season. The thawing fraction-melt period. The melt period~~ in each grid cell was defined from the amount of thawed pixels in a grid cell. ~~The amount of pixels in a grid cell diminished towards,~~ ~~emphasizing more the small grid cells in~~ the north. The mean thawing fraction of all ~~grid-cells-grid cells~~ in a permafrost zone was then calculated. ~~The selected thresholds were chosen to define a robust, zone-mean transition period rather than the exact timing of soil thaw at individual grid cells. In early spring, the SMOS F/T signal can respond to wet snow in addition to thawed~~ 260 ~~soil, as liquid water in the snowpack produces a microwave signature similar to that of thawed soil (Rautiainen et al., 2025). Consequently, thawing fractions very close to 0% or 100% are more sensitive to short-term fluctuations in the microwave signal. Defining the season boundaries away from these extremes ensures that the melt period reflects a sustained, large-scale transition.~~

In the grid-based approach, the start of the ~~melting-season-melt period~~ was defined differently. The season started when 1) 265 one SMOS F/T pixel (25 km × 25 km) in the 1° × 1° grid cell had melted ~~and 2) after the day when the grid cell reaches its minimum annual thawing fraction before the end of July. The second condition was needed for the same reason as in the region-based approach.~~ There were a maximum of 18 SMOS F/T pixels and a minimum of 1 pixel in each 1° × 1° grid cell depending on data availability and geographic location. For example, in a grid cell with 18 pixels, the season started when the thawing fraction was 1/18, meaning that 5.6 % of the area of that grid cell has melted. The season ended when the thawing 270 fraction in the grid cell had surpassed 0.8 of the maximum thawing fraction of that year. To separate the spring ~~melting-season~~

melt period from the autumn freezing season, the melting-season-melt period was defined to start and end before the 212th day of the year. ~~There were grid-cells where there was missing data on some days. The missing data (end of July).~~ Missing data in the grid cells was replaced by interpolating linearly between the previous and the coming day with an existing value. If none of the pixels melted in a grid cell, or if the thawing fraction ~~during the spring never became~~ never was below 0.8 of the maximum thawing fraction of that year before the end of July, meaning that the grid cell did not freeze, the grid cell did not have a melting-season-melt period. In some of the ~~grid-cells~~ grid cells, the condition for the end of the melting-season-melt period was never surpassed before the 212th day of the year even though the season had a beginning. For those ~~grid-cells~~ grid cells, the end of the melting-season-melt period was defined as a day when the soil had thawed the maximum amount during the spring-season-melt period. However, this only happened in 2 ~~grid-cells~~ grid cells in 2017. Additionally ~~there were a couple of grid-cells where the minimum thawing fraction during the spring time was reached, in a few grid cells, the minimum thaw fraction occurred~~ after the maximum was already had already been reached. In ~~those grid-cells these cases, the onset of~~ the season was defined to begin before the maximum was reached occur prior to the maximum. Excluding the first three years, fewer than 1 % of grid cells in the study area did not have a melting-season-melt period annually. In the first three years, this percentage ranged from 6.5 % in 2011 to 1.1 % in 2013.

285 2.5.2 Calculating melting-season-melt period methane emissions

After defining the melting-season-melt period, the methane emissions were calculated from the CTE-CH₄ inverse model biospheric posterior emission estimates. The methane emissions were optimized weekly, and to calculate the daily emissions, the model data was interpolated linearly from one optimized weekly value to the next weekly value. From these daily values, the posterior methane emissions were analysed during the melting-season-melt period. Similarly to the melting-season-melt period,
290 the regional emissions were calculated separately for the four permafrost zones: non-permafrost, sporadic, discontinuous and continuous permafrost zones.

The total melting-season-melt period CH₄ emissions in a region or grid cell in teragrams of methane (Tg CH₄ per region per season, hereafter denoted simply as Tg) were calculated from the daily methane emission estimate. The daily values of all grid cells were added together in each permafrost zone during the melting-season-melt period. The average emissions were
295 calculated in the unit of $\mu\text{gCH}_4 \text{ m}^{-2}\text{s}^{-1}$ by dividing the total emission average daily emissions during the melt period in a permafrost zone by the area of the specific permafrost zone and the length of the melting season. The emissions were studied against the land area of the permafrost zones instead of the actual area of wetlands or permafrost in each zone. This was done because the exact area of wetlands is uncertain (~~Saunois et al., 2020~~)(Saunois et al., 2025), and using the estimated wetland area extent would have added another source of uncertainty.

300 As the melting-season-melt period was defined separately for each $1^\circ \times 1^\circ$ grid cell, the melting-season-melt period emissions were calculated separately as well. The sum of the emissions per grid cell per melting-season-melt period was calculated in each grid from the first day of the melting-season-melt period to the last. To calculate average of the methane emissions in the unit of $\mu\text{gCH}_4 \text{ m}^{-2}\text{s}^{-1}$, the emissions in each grid average daily emissions during the melt period in a grid cell were divided

by the respective area of the grid cell ~~and the length of its melting season~~. To compare with the other method, we divided the
305 grid-based emissions to the four permafrost regions as well.

3 Results

3.1 Length of the ~~melting season~~ melt period

The average length of the ~~melting season~~ melt period over all the permafrost zones was approximately four times longer when
the region-based approach (45 days) was used than when the grid-based approach was used (10 days) (Fig. 2). Using the grid-
310 based approach, the average length of the ~~melting season~~ melt period was the longest in the southernmost zone, non-permafrost
(12 days), and the shortest in the northernmost zone, continuous permafrost (7 days) (Fig. 2 ~~and~~, Fig. 3 and Fig. 4). For the
region-based ~~melting season~~ melt period, there was no as clear gradient in the north–south direction as there was in the grid-
based mean lengths. However, the longest ~~melting season~~ melt period was typically still in the southernmost permafrost zone,
the non-permafrost (57 days), except for 2011, when the longest season was in the continuous permafrost zone (54 days), and
315 in 2018, when it was the longest in the sporadic zone (57 days). On average, the region-based season was the shortest in the
sporadic and discontinuous zones (40 days), which also had smaller areas than the other two zones. The years with the longest
and shortest ~~melting seasons~~ melt periods differed between the two methods and grid-based and region-based lengths of the
~~melting season~~ melt period also did not strongly correlate (Fig. A2).

Most of the of grid-based ~~melting seasons~~ melt periods lasted only a few days (Fig. 4). However, some grid cells had a very
320 long ~~melting season~~ melt period, some as long as the region-based ~~melting season~~ melt period. For example, the West Coast
of Canada and the USA had large areas with a longer ~~melting season~~ melt period (Fig. 3). Other areas with ~~noticeably longer~~
~~melting season~~ a noticeably longer melt period include the coast of Norway, Iceland, East Coast of the USA, Mongolia, the
southern parts of Russia and Northern China. ~~The shortest melting season~~ The shortest melt period in a grid cell was one day
in all the zones and the longest season was found in the sporadic zone and it lasted 212 days, the length of the the maximum
325 duration of the ~~melting season~~ melt period based on our method. It was clearly an outlier, since 95 % of grid cells in the
sporadic zone had a ~~melting season~~ melt period season shorter than 31 days. In addition, 95th percentiles for other zones were
40 days in the non-permafrost zone, 26 days in the discontinuous permafrost zones, and 20 days in the continuous permafrost
zone. From Fig. 4 we can see the gradient in the north–south direction of the longest (97,5 percentile) grid-based melt periods,
with the longest melt periods on average in the southernmost zone, non-permafrost (Fig. 4 a).

330 With the region-based approach, the length of the ~~melting season~~ melt period had a larger inter-annual variation (7–17 days
depending on the permafrost zone) than the average length of the grid-based season (Fig. 2), which might be linked to the inter-
annual variations of ~~weather~~ climate in the permafrost zones, as the region-based method focuses more on the climatological
averages. Hence, the grid-based method ~~might be~~ is better at detecting the actual melting of the snow and soil on a local level,
because it detects the faster changes in a specific ~~grid-cell~~ grid cell. An example of inter-annual variation of weather affecting
335 the region-based length of the ~~melting season~~ melt period is in 2011, when in the continuous permafrost zone, the average
ERA5 2 m mean air temperature was the lowest and the ~~melting season~~ melt period was the longest. In contrast, in 2017 and

2019, the 2 m mean temperature was higher and ~~melting-season-melt period~~ shorter in the continuous permafrost zone. In 2013, the spring ~~melting-season-melt period~~ was the shortest in the sporadic and discontinuous permafrost zones. According to the ERA5 2 m air temperature data, the ~~melting-season-melt period~~ mean temperature was higher in these two zones than
340 during other years of our study period. In 2015, the mean temperature on these two zones was higher during the spring ~~melting-season-melt period~~ as well. This suggest that the ~~melting-season-melt period~~ was shorter and warmer after the colder weather surpassed. In 2017–2019, the sporadic zone had its longest ~~melting-season-melt period~~ while the mean temperatures were the lowest. This all indicates that the ~~melting-season-melt period~~ timing and length were linked to the mean temperature of the zone.

345 The longer ~~melting-season-melt period~~ in the southern regions could have been caused by an early onset of the ~~melting-season-melt period~~, followed by variation between negative and positive temperatures, which would have slowed down the melting. The early onset of the ~~melting-season-melt period~~ and the consecutive variation between temperatures would have made the mean temperature lower and the ~~melting-season-melt period~~ longer. To confirm this, the relationship between the length of the ~~melting-season-melt period~~ and mean temperature was studied (Fig. 5). A negative correlation was found between
350 the two variables, which was the strongest in the discontinuous permafrost zone (region-based $p = 0.008$) and sporadic zone (grid-based $p = 0.005$). With the region-based approach, the correlation between the variables was significant ($p < 0.05$) for all the zones and there was a more prominent negative correlation between the length and the starting day of the ~~melting-season-melt period~~ in all the permafrost zones ($p < 0.001$ for all the zones, Fig. A3). This indicates that the later the season started, the shorter the ~~melting-season-melt period~~ was, at least in the larger permafrost zone scale. Additionally, there was a
355 positive correlation between the region-based starting day of the ~~melting-season-melt period~~ and the mean temperature of the ~~melting-season-melt period~~ (sporadic: $p < 0.001$, continuous: $p < 0.01$, and non-permafrost and discontinuous: $p < 0.05$). This all indicates that the ~~melting-season-melt period~~ started earlier, when the mean temperature was lower, on a larger scale.

With the grid-based approach there was not as strong a correlation between the mean starting day of the ~~melting-season-melt period~~ and the mean temperature or mean length of the ~~melting-season-melt period~~ ($p \geq 0.1$ for most of the permafrost
360 zones). Additionally, the ~~melting-season-melt period~~ mean temperature was higher with the grid-based approach than with the region-based approach (Fig. 5). ~~Looking at the individual grid cells could have given a different output.~~ The mean values of the length and temperature-, ~~temperature and starting day~~ of the grid-based ~~melting-season-melt period~~ ~~might not have been the best to describe the relationship between the length of the melting season and the temperature of the melting season-melt period may be insufficient to describe their relationship~~ because the variation between different grid cells is not seen.

365 3.1.1 Start and end days of the ~~melting-season-melt period~~

With the region-based ~~melting-season-melt period~~, the start and end of the season varied from year to year in each permafrost zone (Tab. A2). The range of variation of the starting days was 21 days in the non-permafrost and sporadic zones, 15 days in the discontinuous permafrost zone, and 18 days in the continuous permafrost zone. The range of variation of the ending days was shorter, only 6 days in the non-permafrost zone, 14 days in the sporadic zone, and 8 days in the discontinuous and continuous
370 permafrost zones. Between different ~~grid-cells-grid cells~~, the start and end of the ~~melting-season-melt period~~ varied more within




Figure 2. The length of the melting-season-melt period defined with the two methods in the northern high latitude permafrost zones. The dotted lines represent the grid-based mean length of the melting-season-melt period. The solid lines represent the region-based melting-season-melt period. The size of grid-cells in the north-south direction was taken into account when defining the melting season with both methods melting-season-melt period.

a year than the averages from year to year. However, even inside one grid-cell-grid cell, there was more inter-annual variation that was not seen from the average (Fig. A4 b and c). On average, the region-based melting-season-melt period started earlier and ended later than the grid-based melting-season-melt period. However, in some grid-cells, the melting-season-grid cells, the melt period started much earlier and ended later than the region-based season, as expected since at least 10 % of the grid-cells-grid cells had to be thawed for the region-based melting-season-melt period to start and 80 % had to be melted for it to end.

With both methods, the season typically started and ended the earliest in the southernmost regions and the latest in the northernmost regions, with the season starting later in some of the southernmost grid cells (mostly mountainous regions). In the southernmost regions, the season started earliest in January, but the average start of the season-Using the grid-based approach, the melt period in the southernmost regions began as early as January. However, the average onset occurred in April in the non-permafrost zone was in April. In and in May in the continuous permafrost zone, although the earliest onset in the season started earliest in February but the average start of the melting season was in May. In the non-permafrost zone, the season-latter was already observed in February. Similarly, the melt period ended earliest in January and on average in April. In in the non-permafrost zone and in March in the continuous permafrost zone. On average, the season ended earliest in March and on average in June. With in April in the non-permafrost zone and in June in the continuous permafrost zone. Using




Figure 3. Grid-based (figures a,b and c) and region-based (figures d,e and f) average length, start day, and end day of the melting season-melt period in the northern high latitudes, averaged over 2011–2021. Notice the different color-range in the melting season-melt period figures on the left side (a and d). In the region-based figures, only the illustrated length, start and end days are ticked on the color-bar. The color range in the start and end day figures are the same in the grid- and region-based figures.

385 the region-based approach in the, the melt period in the non-permafrost region, the melting season started in the middle of March and ended in the beginning of zone extended from mid-March to early May. In the sporadic region, the melting season started in the middle of April and ended in the middle of May permafrost zone, it lasted from mid-April to mid-May. In the discontinuous permafrost region, the melting season started zone, the season began in the second half of April and ended at the




Figure 4. The length of the melting-season-melt period defined with the two methods in the northern high latitude zones: a) non-permafrost, b) sporadic, c) discontinuous and d) continuous permafrost. The green-violin plots represent the grid-based length distribution in the four zones. The medians as well as the min and 97,5 percentile max values for each year are shown as well with the green lines. Grid cells where the length of the melting-season-melt period was longer than the 97,5 percentile have-been-masked-out-were-not-included. The blue lines represent the region-based length of the season.

end-of-in-late May. In the continuous permafrost region, the melting-season-started-in-the-beginning-of-May-and-ended-in-the
 390 middle-of-June-zone, the melt period occurred from early May to mid-June. In the Eurasian continent, the melting-season-melt
period started earlier in the west and later in the east. The regions with the most permafrost are located in the eastern part of
 Eurasia which means that the regions with less permafrost started to melt first. In the American continent, an east–west thawing
 gradient is not apparent. However, the melting-season-melt period started and ended the latest where the most permafrost is
 located. Our region-based melt period started on average 10-20 days earlier in all the zones compared to the start of the thaw
 395 season defined by Erkkilä et al. (2023). With our grid-based approach the melt period started on average 1-10 days later than
the thaw season.




Figure 5. Figure a depicts the region-based relationship between the length of the melting-season-melt period and the mean temperature of the melting-season-melt period. Figure b depicts the grid-based relationship between the mean length and the mean temperature of the melting-season-melt period. The scatter plot color gradient represents the different years with 2011 being the lightest color and 2021 being the darkest. R^2 and p in the legends are the coefficient of determination and p -values of the slopes from linear regression fit, indicating statistical significance of the coefficient of determination.

3.1.2 Melting-season-Melt period in the Hudson Bay lowlands and the Western Siberian lowlands

The melting-season-melt period was additionally studied in the Hudson Bay lowlands (land area in 50–60° N latitudes and 75–96° W longitudes) and the Western Siberian lowlands (land area in 52–74° N latitudes and 60–94.5° E longitudes). Hudson Bay lowlands and Western Siberian lowlands are some of the largest methane emitting wetlands in the northern high latitudes (Pickett-Heaps et al., 2011; Umezawa et al., 2012). They consist of the four permafrost zones, except the Hudson Bay lowlands where there is no continuous permafrost. On average, the grid-based melting-season-melt period was slightly shorter in the Western Siberian lowlands (~9 days) than in the Hudson Bay lowlands (~10 days). The season started and ended a few days earlier on average in the Western Siberian lowlands than in the Hudson Bay lowlands. The region-based melting-season-melt period length was not defined separately for these wetland regions, but the original region-based lengths in the four permafrost zones were used.

Overall, in the Hudson Bay lowlands, there was a negative correlation between the mean temperature and mean length of the melting-season-melt period ($p < 0.05$ in all zones) with the grid-based approach. In the Western Siberian lowlands, there was also a negative correlation in all of the permafrost zones but it was statistically significant only in the non-permafrost and sporadic zones ($p < 0.05$). This means that the season was shorter when the mean temperature was higher at least with the grid-based approach.

With the region-based approach, there was not as clear negative correlation between the length of the ~~melting-season-melt period~~ and the mean temperature of the ~~melting-season-melt period~~ especially in the Western Siberian lowlands ($p < 0.05$ only in the sporadic zone). In the Hudson Bay lowlands, there was a statistically significant negative correlation in the sporadic and discontinuous zones ($p < 0.05$). If the ~~melting-season-melt period~~ was defined with the region-based approach separately in the Hudson Bay and the Western Siberian lowlands areas, then the correlations might have been more significant.

3.2 ~~Melting-season-Melt period~~ methane emissions

~~The biospheric CH₄ emissions were analyzed with both the region-based and grid-based approach. From this point onward, we will focus exclusively on biospheric emissions, regardless of whether this is explicitly mentioned. Biospheric emissions refer to optimized fluxes for which prior estimates were derived from ecosystem model simulations of wetland emissions as well as terrestrial sink fluxes (see Section 2.2.3). The average annual biospheric posterior region-based melting-season-melt period CH₄ emissions in the whole northern high latitude region in our study were 1.83 ± 0.27 Tg (Table A3), where \pm is the maximum difference from the mean here and hereafter, and grid-based annual melting-season emissions were 0.45. Grid-based annual biospheric posterior melt period emissions were 0.451 ± 0.10 Tg (Table A4). The year-to-year variation was modest, within 15% of the average emissions with the region-based approach and 23 % of the average emissions with the grid-based approach. The grid-based emissions were much smaller than the region-based emissions in all the zones due to shorter melting seasons and a shorter melt period on average (see Section 3.1). The melting-season-posterior melt period methane emissions were the largest in the southernmost zone, non-permafrost, which was also the largest zone, and smallest in the discontinuous permafrost zone (Fig. 3.2.1).~~

To study which variables were linked to ~~melting-season-melt period~~ methane emissions, the relationship between the length of the ~~melting-season and the melt period and the posterior~~ methane emissions was studied (Fig. 87). With both methods, the correlation between the length of the ~~melting-season-melt period~~ and the total ~~melting-season-melt period~~ emissions was positive in all the permafrost zones, meaning that the melting-season had higher emissions when the season was longer. The positive correlation can be seen most clearly (Fig. 7 a and c), indicating higher emissions during longer seasons. This relationship was most pronounced in the non-permafrost zone. The correlations were statistically significant in all zones for both methods ($p < 0.01$) in all the zones with both methods).

Additionally, the ~~posterior~~ emission rate, as the units of $\mu\text{g}(\text{CH}_4) \text{ m}^{-2} \text{ s}^{-1}$, was calculated for the different zones. The region-based mean emission rate was defined by dividing the total emissions with the area of the zone and the length of the melting-season. The grid-based mean emission rate was defined as the average emission rate between the different grid-cells (see Section 2.5.2). The correlation between the rate of emissions and the length of the ~~melting-season-melt period~~ was not as clear as between the total ~~melting-season-melt period~~ emissions and length, especially with the grid-based approach ($p > 0.05$ in all zones) (Fig. 7 d). There was a small negative correlation in the sporadic and discontinuous zones and positive ones in the other two zones. With the region-based approach (Fig. 8), there was a stronger negative correlation in the sporadic zone between emission rate and length of the season ($p < 0.01$) (Fig. 7 b). In other zones, the correlation between emission rate and length was insignificant ($p > 0.05$). The negative correlation in the sporadic zone could exist because of the negative correlation

between the length and temperature of the ~~melting-season~~melt period, and therefore the temperature would correlate positively with the emission rate. This relationship was studied in the grid-based Fig. A5, where grid cells with a negative emission rate were masked out, but no correlation was found in any of the zones. The region-based relationship between the variables was very similar. This indicates that the total emissions grew when the season was longer but there was no clear indication that the
450 emission rate would be stronger or weaker when the season is longer.

Average posterior grid-based emissions in the Hudson Bay lowlands were 0.03 ± 0.01 Tg and in the Western Siberian lowlands they were 0.11 ± 0.04 Tg (Table A5). The corresponding region-based emissions in the Hudson Bay lowlands were 0.10 ± 0.03 Tg and 0.47 ± 0.14 Tg in the Western Siberian lowlands. ~~When the contribution of different permafrost zones~~
An analysis of permafrost zone contributions in the Hudson Bay ~~lowlands~~Lowlands and the Western Siberian ~~lowlands were~~
455 ~~studied, most of the~~Lowlands shows that, in most years, grid-based emissions ~~in these areas came from~~were dominated by the non-permafrost zone~~during most of the years~~. In 2012 and 2016 in the Hudson Bay lowlands, most emissions ~~came~~were from the sporadic zone, instead. With the region-based approach, most emissions ~~came~~were from the non-permafrost zone every year in both Hudson Bay lowlands and Western Siberian lowlands. The non-permafrost zone was the largest permafrost zone in both regions.

460 The years with the highest emissions in the entire northern latitude ~~regions were 2011, 2014–2017, and 2019~~ (Tables A3, A4). ~~With~~region were 2014 with the region-based approach, ~~the year with the largest emissions was 2014 and and 2017~~ with the grid-based approach ~~; it was 2017~~. (Tables A3, A4). Maps of some of the highest emitting ~~melting seasons~~melt periods in the Hudson Bay lowlands and Western Siberian lowlands are shown in Figures ~~7–8~~ and A6. ~~We can see that some of the grid cells~~
Some of the grid cells with the highest emissions were located ~~inside these two lowlands~~within these two lowland regions.
465 However, the years with highest emissions in the Hudson Bay lowlands and Western Siberian lowlands differed from the years with highest emissions of the whole northern latitude regions. On the other hand, the years with the highest emissions in the Hudson Bay lowlands and Western Siberian lowlands matched between the two methods. The highest grid-based emissions in the Hudson Bay lowlands were found in 2012 and 2020, and region-based in 2012 and 2014. In the Western Siberian lowlands, the highest emissions were in 2014 and 2015 with both approaches. The highest emissions of the Western Siberian
470 lowlands align better with the overall highest emitting years, likely due to the larger area of the Western Siberian lowlands to the Hudson Bay lowlands. However, there ~~were not major differences~~is no major difference in the emissions between the years in both regions. The average grid-based ~~melting-season~~melt period length varied between the permafrost zones during the years with the most emissions in the Hudson Bay lowlands and the Western Siberian lowlands. ~~Usually the season was longer than average at least in~~
During those years, at least one of the four ~~regions in both lowlands during those years~~zones
475 in each lowland experienced a longer-than-average season. In the Hudson Bay lowlands during 2012 and 2020, the ~~melting season~~melt period was usually longer in a region where the mean temperature of the ~~melting-season~~melt period was also colder than average. Same was seen in the Western Siberian lowlands in 2014.

The region-based ~~melting-season~~melt period was not longer than the average in any of the zones in 2012, but in 2014, it was longer than the average in all the zones, except in the sporadic zone. In 2015, it was only longer than average in the non-
480 permafrost zone. However, the region-based ~~melting-season~~melt period was defined from the whole permafrost zone instead


of only the area within Hudson Bay lowlands and Western Siberian lowlands, which might explain why the ~~melting seasons were melt period was~~ not exceptionally long during the years with highest emissions in these two regions. Both regions were also mostly covered by the non-permafrost zone, which explains why the emissions were generally larger when the ~~melting season-melt period~~ was longer in the non-permafrost zone.

485 3.2.1 Comparison to prior emissions

The total prior emissions during the melt period were 1.81 ± 0.29 Tg with the region based approach and 0.447 ± 0.11 Tg with the grid-based approach. This means that the inversion increased the total melt period emissions from the prior to the posterior. In the permafrost zones the inversion increased emissions from the prior in areas with less permafrost (Fig. 3.2.1 a) and decreased in areas with more permafrost (Fig. 3.2.1 d). This was more evident with the region-based approach and
490 was seen the most distinctly in the non-permafrost and continuous permafrost zones. In the non-permafrost zone the posterior was almost always larger than the prior with both methods, despite already having the highest emissions. In the continuous permafrost zone the posterior emissions were always smaller than the prior emissions with both methods.

To see where the differences between prior and posterior emissions were most noticeable, we plotted a map of the total difference of the grid-based emissions during the melt period (Fig A7) including the same years as in Fig. 8 and an average
495 of all the years. The region-based figure was very similar to the grid-based, thereby only the grid-based figure is shown in the Appendix. Grid cells with no prior emissions during the melt period were masked out in the figure, though there were only a few. During these years, we can see spatial variation throughout the years. The most notable differences appear in the Western Siberian Lowlands and the Hudson Bay Lowlands region. The average region-based biospheric prior CH₄ emissions during the melt period in the Western Siberian lowlands were 0.45 ± 0.11 Tg and in the Hudson Bay lowlands they were 0.09 ± 0.026
500 Tg. The corresponding grid-based emissions in the Western Siberian lowlands were 0.11 ± 0.03 Tg and in the Hudson Bay lowlands 0.03 ± 0.01 Tg. The change from the prior to posterior is percentage wise larger in the Hudson Bay lowlands and Western Siberian lowlands than it is in the total northern high latitude region used in our study. In the total northern high latitude region, the posterior is about 1 % larger than the prior with both methods. In the Hudson Bay lowlands the posterior is about 6 % larger than the prior with both methods and in the Western Siberian lowlands it is about 3 % (grid-based) to 4 %
505 (region-based) larger. In the four permafrost zones, the posterior emissions were 0.4 % (discontinuous permafrost) to 3 % larger (non-permafrost) or 6 % smaller (continuous permafrost) than the prior on average with the region-based approach. With the grid-based approach the posterior emissions were about 2 % larger (non-permafrost and sporadic zones) or 2 % (discontinuous) to 8.6 % (continuous permafrost) smaller than the prior on average.

We further examined the relationship between the difference in posterior and prior emissions and the length of the melt
510 period (Fig. A8). Using the region-based approach, a significant positive correlation was found in the continuous permafrost zone between the difference in emission rate and melt-period length ($p < 0.05$; Fig. A8 b). In contrast, the grid-based approach revealed a negative correlation in the sporadic permafrost zone between melt-period length and both the emission difference ($p < 0.05$) and the emission rate difference ($p < 0.01$). Although differences between posterior and prior emissions were also observed in the other zones, no statistically significant correlations were detected between posterior-prior emission differences



Grid-based emission maps of the melting season in the years that had some of The darker color represents the highest grid-based melt period emissions in the Hudson Bay lowlands and Western Siberian lowlands. Hudson Bay lowlands have been outlined with red borders, and Western Siberian lowlands with blue borders lighter colors the region-based emissions.

Grid-based emission maps of the melting season in the years that had some of The darker color represents the highest grid-based melt period emissions in the Hudson Bay lowlands and Western Siberian lowlands. Hudson Bay lowlands have been outlined with red borders, and Western Siberian lowlands with blue borders lighter colors the region-based emissions.

Figure 6. Total prior and posterior emissions in the four permafrost zones: a) non-permafrost, b) sporadic, c) discontinuous and d) continuous permafrost. The region-based prior emissions are coloured-colored in blue-purple and the grid-based posterior emissions in green-blue.

Grid-based emission maps of the melting season in the years that had some of The darker color represents the highest grid-based melt period emissions in the Hudson Bay lowlands and Western Siberian lowlands. Hudson Bay lowlands have been outlined with red borders, and Western Siberian lowlands with blue borders lighter colors the region-based emissions.

515 (or emission rate differences) and melt-period length. Overall, these results suggest that the inversion only marginally altered the relationship between emissions and melt-period duration.




Figure 7. Figures a and b depict the region-based relationship between the total emissions (a) or the emission rate (b) and the length of the season. Figures c and d depict the grid-based relationship between the total emissions (c) or the mean emission rate (d) and the mean length of the [melting season](#)[melt period](#). Notice the different value range on the y-axis in a and c. The scatter plot color gradient represents the different years with 2011 being the lightest color and 2021 being the darkest. R^2 and p in the legends are the coefficient of determination and p -values of the slopes from linear regression fit, indicating statistical significance of the coefficient of determination.




Figure 8. Grid-based emission maps of the melt period in the years that had some of the highest emissions in the Hudson Bay lowlands and Western Siberian lowlands. Hudson Bay lowlands have been outlined with red borders, and Western Siberian lowlands with blue borders.

4 Discussion

4.1 The ~~melting season~~melt period

4.1.1 Definition of the ~~melting season~~melt period and the SMOS F/T algorithm

520 The ~~spring melting season~~melt period occurring during springtime was defined for the northern high latitude wetlands for the first time based on the SMOS F/T data. ~~Because~~As the SMOS F/T algorithm cannot distinguish between thawed soil and wet snow, it likely ~~first detects the melting of snow instead of thawing of the soil in springtime. Thus, the defined melting seasons probably already started when the snow began to melt and before the soil thawing and made the season longer than the actual soil thawing lasted. However, methane emissions can start already during snow melting and cracking of the soil, when~~
525 ~~liquid water can seep to the soil (Rinne et al., 2007; Raz-Yaseef et al., 2017) and wintertime methane reserves are released. Hence~~extended the season beyond the duration of true soil thawing. E.g. Tenkanen et al. (2021) found that the SMOS F/T data showed later soil freezing but earlier soil thawing than a process-based ecosystem model. However, it is justified to define the ~~melting season~~melt period to start from the melting of the snow (see Section 2.5.1). In addition, including either snow depth data from in-situ measurements or fractional snow cover data from satellites could have helped our algorithm to detect the
530 starting of the soil thawing.

The amount of SMOS F/T pixels in a $1^\circ \times 1^\circ$ grid cell made the definition of ~~melting season~~melt period spatially inconsistent. The number of pixels decreased towards the north, requiring a higher fraction of thawed soil in northern grid cells before the ~~melting season~~melt period began. However, with our definition, the absolute area, which had to be thawed before the season could start, was similar across the study area, though it also meant that the estimated thawing state of the soil was
535 more uncertain if there were missing SMOS F/T datapoints. The difference in the amount of pixels in a grid cell might have introduced systematic biases, causing longer ~~melting seasons~~melt periods in south and shorter in north.

With both methods, the ~~melting season~~melt period generally started the earliest in the southern regions. However, there were some mountainous regions in the south, for example, the mountains in Mongolia, and the Rocky Mountains in the western coast of North America (Fig. 3), where the season started and ended later, causing long ~~melting seasons~~melt periods. This is
540 likely due to higher altitude and lower temperatures. The SMOS F/T data is not as reliable in the mountainous regions as it is elsewhere, mainly because there is less soil substance, especially at higher elevations. Since the SMOS satellite measures the soil freezing through the permittivity difference between ice and water, it does not measure the soil freezing correctly in areas with less absorbent soil. Thus, the melting of snow or ice on solid rock does not significantly affect what is measured. A sufficient difference between thawed and frozen states requires soil that has absorbed water. Additionally, the topography
545 affects the remote sensing measurements: the measurement angle varies a lot in the mountains, which means that with different measurement angles, the measured thawing state of the soil also changes.

4.1.2 Region-based and grid-based ~~melting season~~melt period

The region-based ~~melting season~~melt period was on average about 4 times as long as the grid-based ~~melting season~~melt period. The region-based approach ~~might does~~ not represent the local ~~melting season~~melt period accurately because of the substantial
550 area of the permafrost zones and the difference between the start and end days inside one zone was large. In other words, the ~~melting season~~melt period in the southern grid cells might have already ended, while soil was still frozen in northern parts of the zone. However, with this method, the thawing fraction varied less during the ~~melting season~~melt period, and the beginning and end of the ~~melting season~~melt period were more consistent, making it easier to compare the different permafrost zones' ~~melting seasons~~melt periods and methane emissions with each other ~~–~~

555 ~~Each of the four permafrost zones consisted of areas in the North American and the Eurasian continents. Especially the region-based melting season could have been more precise if it was defined separately for the two continents due to their different year-to-year variability of climates on a larger climatological scale.~~

The grid-based ~~melting season~~melt period was on average almost twice as long in the southernmost zone: non-permafrost (12 days) compared to the northernmost permafrost zone: continuous permafrost (7 days). This suggests that the season was
560 shorter when there was more permafrost and the further north the region was, except for the mountainous permafrost regions in the south (Fig. 3). In a small area, the soil ~~probably melts~~melts almost simultaneously leading to a shorter ~~melting season~~melt period. Additionally, ~~the season could not start because the grid-based period could not begin~~ before the minimum ~~thawing value was surpassed for the last time in the spring. This meant that for~~thaw fraction was exceeded for the final time in spring, some grid cells ~~– there might have been variation in the thawing value before this date, and this variation could~~may have
565 ~~experienced fluctuations in thaw fraction prior to this date. This variability could therefore~~ have been included ~~in the melting season~~within the defined melt period. However, with this method the ~~spring melting season~~melt period was separated from the autumn freezing season if it continued past the turn of the year.

~~Each of the four permafrost zones consisted of areas in the North American and the Eurasian continents. Especially the region-based melt period could have been more precise if it was defined separately for the two continents due to their different
570 year-to-year variability of climates.~~

4.1.3 Timing and length of the ~~melting season~~melt period

Different methods used to study the timing of snowmelt and spring thawing season have produced varying results. According to the definitions of the Finnish Meteorological Institute (FMI) based on temperature data, spring starts approximately in the middle of March in southern Finland and in April in northern Finland (Ilmatieteen laitos, 2025) and lasts for two to four weeks
575 during which snow melts and the growing season starts. Our estimated ~~melting seasons~~melt periods in Finland, which is located mostly in the non-permafrost area, coincide with this: on average, the ~~melting season~~melt period started in March–April using both the grid-based and the region-based methods.

According to regional studies in northern Alaska, underlain by continuous permafrost, the spring thawing season in the tundra region was approximately 20 days long (Zona et al., 2016; Bao et al., 2020), which is shorter than the region-based

580 ~~melting-season-melt period~~ defined here (45 days on average, Table A2) and longer than the average grid-based ~~melting-season~~
~~melt period~~ (10 days). With our grid-based method, the average length of the ~~melting-season-melt period~~ during our study
period in the ~~grid-cells-grid cells~~ closest to the coordinates of the measurement stations used by Zona et al. (2016) and Bao
et al. (2020) varied between 4-19 days. Overall, in many ~~grid-cells-grid cells~~ in the whole northern high latitude domain, the
length of the ~~melting-season-melt period~~ was ~20 days. Both Zona et al. (2016) and Bao et al. (2020) used soil temperature data
585 to define the thawing of the soil. We did not use in-situ soil temperature data in this study as in-situ measurements of frozen
soil are rare, and thus not suitable for our larger region study. Other option would have been to use reanalysis soil temperature
data such as ERA5, which has shown higher skills than other products and a significant improvement over its predecessor (Li
et al., 2020). However, it is not well-suited for permafrost research (Cao et al., 2020). Thus, SMOS F/T data is a good proxy
for the soil thawing.

590 ~~In the study by Tenkanen et al. (2021), the SMOS F/T data showed later soil freezing but earlier soil thawing than a~~
~~process-based ecosystem model, resulting in a shorter winter season. The region-based season instead was longer. This might~~
~~be due to the~~ The SMOS F/T data starting the ~~melting-season-melt period~~ already from the melting of the snow instead of the
soil ~~is likely the reason for the longer region-based melt period~~. According to our prior LPX-Bern DYPTOP v1.4 (Fig. A7A9),
the soil temperature rose to 0 °C during the region-based ~~melting-season-melt period~~ on all permafrost zones. This means that
595 the SMOS F/T ~~melting-season-melt period~~ started earlier than the process-based ecosystem model ~~melting-seasonmelt period~~.
However, the prior had a monthly temporal resolution compared to the daily resolution of the SMOS F/T data, and so the soil
temperature rising above 0 °C in the middle of the ~~melting-season-melt period~~ is a good indicator for the correct timing of our
~~melting-season-melt period~~.

According to the prior emissions used in this study (LPX-Bern DYPTOP v1.4), peat emissions were relatively high during
600 the ~~spring-melting-season-than-melt period compared to~~ emissions from inundation for all other permafrost zones but continuous
permafrost (Fig. A7A9). However, even in the continuous permafrost zone, the peat emissions are higher than ~~those from~~
inundation in April, and are rising during the ~~melting-seasonmelt period~~. At the beginning of spring, extensive snow melt
inundation causes a large part of methane emissions, and after the end of the ~~melting-seasonmelt period~~, the methane emissions
are dominated by peat emissions. Even though ~~methane emissions from peat dominated over emissions caused by inundation~~
605 for a large part of the year, there were evident peaks in inundation emissions in every permafrost zone during and/or right after
the ~~melting-seasonmelt period~~. This indicates that the ~~melting-season-melt period~~ timing we defined is reasonable.

4.2 Methane emissions during the ~~melting-seasonmelt period~~

The way we defined the ~~melting-season-also-melt periods~~ affected the estimated ~~melting-season-melt period~~ methane emissions.
The emissions were optimized weekly, which were interpolated linearly to daily values from one optimized weekly value to
610 the next weekly value before the calculation of the ~~melting-season-melt period~~ emissions. This introduced a new source of
uncertainty and affected the estimated ~~melting-season-melt period~~ methane emissions, especially when using the grid-based
approach, as the length of the grid-based season was much shorter and often less than a week (between 51% ~~of the grid-cells~~
in the non-permafrost zone and 70% in the continuous permafrost zone).

In the Hudson Bay lowlands and Western Siberian lowlands, as well as in the four permafrost zones, the methane emissions were distributed unevenly (Fig. 78). This ~~might~~ may be explained by the distribution of wetlands ~~in~~ within the lowlands, as wetlands ~~typically have generally exhibit~~ higher methane emissions than ~~, e.g., other land-cover types, such as~~ forests, which might even be a sink of methane. The years with ~~some of the~~ highest emissions were distributed evenly across the study period (Fig. 7), ~~which means~~ 8, indicating that there is no ~~direct indication of increasing melting season~~ clear evidence of increasing melt period emissions.

620 4.2.1 Methane bursts during the ~~spring melting season~~ melt period

Multiple previous studies have shown higher emission rates to our results with many of them showing large bursts of CH₄ from wetlands during the spring thawing season (Jin et al., 1999; Tokida et al., 2007; Song et al., 2012; Tagesson et al., 2012; Mastepanov et al., 2013; Zona et al., 2016; Raz-Yaseef et al., 2017; Bao et al., 2020). These events typically last for a short time, from few hours to a few days (Song et al., 2012). Such emission bursts could have caused large emissions even during a shorter ~~melting season~~ melt period, causing the relationship between the length and the emissions of the ~~melting season~~ melt period to be non-linear. However, we found a positive correlation between the total ~~melting season~~ melt period methane emissions and length, which was observed with both the grid-based and the region-based methods in permafrost zones and Western Siberian and Hudson Bay lowlands. This indicates that the bursts were not large enough to be detected with our model resolution (1° × 1°) and weekly temporal optimization of the fluxes. The emission rates were also estimated from the whole area of the 1° × 1° ~~grid-cell~~ grid cell instead of the wetland area, which made the emission rates smaller compared to the local studies. Another reason for the positive correlation between the values could be that in our study, we focused ~~to study on~~ emissions within larger permafrost zones. In individual ~~grid-cells~~ grid cells, some emission peaks could have been detected but that was not explicitly studied here. Above mentioned studies reporting large CH₄ bursts during the spring thaw used field measurements to define the emission rate which have ~~a sensitivity for a~~ sensitivity for smaller areas compared to our inverse model. Emission rates defined from the eddy covariance measurement measure the local methane bursts rather than the average rates from a larger permafrost zone.

Using a process-based model Castro-Morales et al. (2018) showed an emission rate closer to what we found in this study. The atmospheric inverse model used here used atmospheric CH₄ measurements to inform and "re-evaluate" the process-based estimates. In most years, our inversion showed an increase from the prior emission estimates which were based on the process model LPX-Bern DYPTOP. This means that there might have been short-lived emission bursts during the ~~spring time~~ melt period, which the process model was not able to produce due to the poor spatial and temporal resolution or missing processes, or that the process model emission estimates were overall too low during ~~the spring season~~ this period.

With our inverse model, the grid-based emission rates were higher in some grid cells (Fig. A5), and on average, the grid-based emission rates were higher than the region-based (Fig. 87). The grid-based ~~melting season likely represent~~ melt period likely represents local emission bursts during soil thawing better than the region-based approach. ~~Inside one~~ Within a single permafrost region, ~~the fluxes vary a lot, as the whole area is not~~ fluxes vary widely because the area does not consist entirely of permafrost or wetlands. This lowers the ~~region-based~~ CH₄ emission rate. The local measurement studies might be more

comparable to the ~~spring melting season melt period~~ emissions in the Hudson Bay lowlands and Western Siberian lowlands in grid cells, where the total area of the cell is mostly wetland. However, the focus of this study was to estimate the total ~~melting season emissions during the springtime melt period emissions~~ rather than occasional hotspots.

4.2.2 Magnitude of the ~~spring melting season melt period~~ methane emissions

According to Saunio et al. (2025), the annual global methane emissions estimated from atmospheric inversions were 575 [553–586] Tg CH₄ yr⁻¹, in 2010–2019. Of these emissions, 165 [145–214] Tg CH₄ yr⁻¹ were from wetlands. The ~~mean-average~~ global biospheric emissions using the CTE-CH₄ data from this study for the years 2011–2021 were 124 Tg CH₄ yr⁻¹, including also soil sink (LPX-Bern DYPTOP prior soil sink 33 Tg CH₄ yr⁻¹). In the northern high latitudes (defined in the chapter 2.3), the average annual methane emission were 23 Tg CH₄ yr⁻¹. The region-based ~~spring melting season melt period posterior~~ emissions were 1.83 Tg (8.1% of the annual northern high latitude emissions) and grid-based were 0.45 Tg (2% of the annual northern high latitude emissions), only a small portion of the annual total global emissions. The length of the ~~melting season melt period~~ with the grid-based method was 3%, and with the region-based method 12%, of the total length of a year.

Methane emissions during other seasons and the high northern latitudes have been studied previously. According to Tenkanen et al. (2021), the winter cold season (November to April) emissions were 3.3 Tg in the northern high latitudes (above 50° N), which included partly emissions from both the autumn freezing and spring thaw seasons. The autumn freezing season methane emissions from the same model with a different setup were 1 Tg (Tenkanen, 2019); ~~which means that the region-based spring melting season emissions were a bit larger than the autumn emissions, and grid-based were smaller.~~ Erkkilä et al. (2023) estimated the autumn freezing, winter cold season and summer thaw season emissions in the northern high latitudes with the same inverse model but a different setup to ours. They found emissions to be 1.2 Tg in winter, 0.73 Tg in the freezing period, and 16.2 Tg in the thaw (summer) period. ~~Both~~ In both studies, winter and freezing season emissions were smaller than our region-based ~~spring melt period~~ emissions but larger than the grid-based emissions.

According to a study which used upscaled flux measurements, the ~~spring~~-methane emissions during the melting of the soil in wetlands in a northern high latitude permafrost region were 0.5–0.97 Tg in 2011 (Song et al., 2012). In a study by Ito et al. (2023), the spring season (March–May) methane emissions were calculated with multiple process-based ecosystem ~~model~~ models for the northern wetlands (>45°), and their mean value was 0.80 ± 1.89 Tg CH₄ yr⁻¹ (3.07 ± 9.61 % of the annual emissions), where the error is the variation between the maximum and minimum model result. These ~~values estimates~~ are closer to the grid-based emissions estimated in this study ~~compared to the region-based emissions~~.

Defined from the inverse model used in this study, the mean annual emissions in the Hudson Bay lowlands and Western Siberian lowlands were the size of 2.9 Tg and 5.0 Tg, respectively. This is close to the values defined in other studies (Thompson et al., 2017; Peltola et al., 2019; Tenkanen et al., 2021). The average region-based ~~melting season melt period~~ emission in the Hudson Bay lowlands and the Western Siberian lowlands were the size of 0.1 Tg and 0.47 Tg, respectively. The grid-based emission were smaller with the magnitude of 0.03 Tg and 0.11 Tg, respectively. This means that together they produced approximately 31% of the total ~~melting season melt period~~ emissions from the northern high latitude wetlands with both methods. This is ~~quite~~ a significant part of the total ~~melting season melt period~~ emissions when their areas only represent a

portion of the total study region (Hudson Bay lowlands is 2.5% and Western Siberian lowlands is 9.6% of the total study region land area). The region-based melt period emissions were 7% and grid-based emissions were 1.7% of the annual emissions from the Hudson Bay lowlands and the Western Siberian lowlands. The increase in emissions from prior to posterior estimates was also more pronounced in these two regions than across the northern high-latitude domain as a whole. This may be attributed to the greater extent of continuous wetlands in these regions and/or the higher density of the observational network (Fig. A1), which likely influenced the prior estimates. Additionally, there are large areas in the high northern latitudes, where the difference between the posterior and prior emissions was close to zero during the melt period (Fig. A7). In grid cells where prior emissions were zero, posterior emissions during the melt period were also zero. This likely contributed to the minimal differences observed between prior and posterior estimates in those areas. However, the emissions in the Western Siberian lowlands could have been overestimated, because the NIES observation sites ~~used~~, which we did not adjust before using the measurements in the inversion, had a different calibration scale (3.0 to 5.5 ppb higher) than WMO CH4 X2004A scale as in the ObsPack (Sasakawa et al., 2025). The NIES observations are now included in the newer version of ObsPack, with scale corrected. ~~In conclusion, the Western Siberian lowlands and Hudson Bay lowlands regions produced a large portion of the total melting season emissions. A higher density of observations in the northern high latitudes but these emissions represented only a small portion of the total annual emissions from these regions could have improved our inverse method estimates.~~

5 Summary and conclusion

The ~~spring melting season~~ melt period was defined for the northern high latitude wetlands with the SMOS F/T soil thawing data separately for four permafrost zones: seasonally frozen non-permafrost, sporadic, discontinuous and continuous permafrost. The ~~melting season~~ melt period was defined separately with a region-based and a grid-based approach. The region-based method is comparable to studies with monthly emission estimates, as it focuses more on the climatological differences between large regions, while the grid-based method is more comparable to local studies. The region-based ~~season~~ period started and ended the earliest in the ~~most~~ southernmost permafrost zone and latest in the northernmost zone. With the grid-based approach, there was more variation in the start and end days of the season in the north-south direction mostly because of the mountainous regions. The region-based ~~melting season~~ melt period lengths varied more from ~~year-to-year~~ year to year than the grid-based seasons. ~~With both methods, the~~ with he non-permafrost zone ~~had~~ having the longest season on average with both methods. The average grid-based ~~melting season~~ melt period was shortest in the continuous permafrost zone and the region-based was shortest in the sporadic and discontinuous zones. We found that the length of the ~~melting season~~ melt period was dependent on the average 2 m temperature of the ~~melting season~~ melt period, with a longer season having a lower mean temperature. This could have been caused by the early onset of melting, followed by variation between temperatures, making the mean temperature of longer seasons lower. We found the SMOS F/T data to be useful in estimating ~~spring melting season~~ the melt period, and for example, using it to inform process-based models to account for soil freeze/thaw state could lead to better constrained methane emission estimates.

715 In the four permafrost zones, the ~~melting-season-melt period~~ methane emissions were the largest in the non-permafrost zone and the smallest in the discontinuous zone with both methods. The total region-based ~~melting-season-melt period~~ emissions for the four zones were 1.83 ± 0.27 Tg CH₄ per season and grid-based emissions were 0.45 ± 0.10 Tg CH₄ per season. These emissions represented 8.1 % and 2 % of the total annual northern high latitude emissions according to our inverse model results (23 Tg). We found that the total ~~melting-season-melt period~~ emissions depended on the length of the ~~melting-season-melt period~~, with more methane emitted during longer seasons.

720 We additionally studied the emissions in the Hudson Bay lowlands and Western Siberian lowlands, as they are the largest wetlands in the northern high latitudes. Their ~~melting-season-melt period~~ methane emissions were a significant portion of the total ~~melting-season-melt period~~ emissions (31% with both methods and both lowlands combined) but only a small part of the total annual northern high latitude emissions (2.5% with the region-based approach and 0.6% with the grid-based approach). The Western Siberian lowlands had higher emissions than the Hudson Bay lowlands ~~with both methods, likely due to its' larger~~
725 ~~size, even though it had a slightly shorter and colder grid-based melting-season-melt period on average than the Hudson Bay lowlands~~~~but with a lower average mean temperature with the grid-based approach.~~

The total ~~melting-season-melt period~~ emissions in the northern high latitudes were only a small portion of the total annual emissions. However, ~~the melting-season emissions could grow in the future due to future~~ climate change and ~~the thawing of permafrost associated permafrost thawing could amplify melt period emissions, resulting in elevated CH₄ bursts. Our results~~
730 ~~showed that the melt period was the longest with highest emissions in the non-permafrost region where the average temperature is higher throughout the year.~~ On the other hand, our results showed that ~~a shorter melting season when the mean temperature was higher on average during the melt period, the period was shorter and~~ had smaller total emissions~~and that a shorter season had a higher mean temperature on average.~~. Increasing temperatures could ~~instead~~ lead to shorter ~~melting seasons and lower melting-season-methane emissions but also a longer thaw season. Additionally~~ melt periods and consequently lower melt period
735 ~~methane emissions. Subsequently, a longer summer thaw season could lead to higher annual emissions. However,~~ it is still unclear how ~~climate change~~ feedback loops will affect emissions and ~~the thawing of the permafrost shoulder season lengths in the northern high latitudes, as permafrost thawing and soil drying could lead to higher CO₂ emissions instead.~~ To get a better look at the ~~spring melting-season-melt period~~ emissions, the results from several inversions could be compared to each other. In addition, going further to the level of individual wetlands and comparing their flux measurements ~~in the springtime during~~
740 ~~the melting~~ could bring more clarity to true wetland emissions.

Data availability. Article data will be published in a public data repository. Until that time, data is available from the main author (Sara Hyvärinen).

Appendix A

Table A1. List of surface observation sites used in inversions. Observation Uncertainty (Obs. Unc.) is used to define diagonal values in the observation covariance matrix. The data type is categorized into two measurements (discrete (D) and continuous (C)).

site code	Site Name	Site Country	Contributor	Site Longitude [°E]	Site Latitude [°N]	Altitude* [m a.s.l.]	Obs. Unc. [ppb]	Data Type D/C	Dates min. [year/month]	Dates max. [year/month]
ABP	Arembepe, Bahia	Brazil	NOAA	-38.17	-12.76	6.00	4.5	D	2006/10	2010/01
ABT	Abbotsford, British Columbia	Canada	EC	-122.34	49.01	93.00	30.0	C	2014/03	2020/12
ALT	Alert, Nunavut	Canada	NOAA	-62.51	82.45	190.00	15.0	D	2000/01	2021/10
ALT	Alert, Nunavut	Canada	EC	-62.51	82.45	195.00	15.0	C	2000/01	2020/12
AMY	Anmyeon-do	Republic of Korea	NOAA	126.33	36.54	87.00	30.0	D	2013/12	2021/11
ARA	Arcturus	Australia	CSIRO	148.47	-23.86	185.00	15.0	D	2010/05	2013/10
ASC	Ascension Island	United Kingdom	NOAA	-14.40	-7.97	90.00	15.0	D	2000/01	2021/11
ASK	Assekrem	Algeria	NOAA	5.63	23.26	2715.00	25.0	D	2000/01	2019/11
AZR	Terceira Island, Azores	Portugal	NOAA	-27.36	38.76	24.00	15.0	D	2000/01	2021/09
AZV	Azovo	Russian Federation	NIES	73.03	54.71	190.00	30.0	C	2009/10	2018/12
BAR	Baranova	Russian Federation	FMI	101.62	79.28	30.00	4.5	C	2015/11	2021/02
BCK	Behchoko, Northwest Territories	Canada	EC	-115.92	62.80	220.00	15.0	C	2010/10	2020/12
BHD	Baring Head Station	New Zealand	NOAA	174.87	-41.41	90.00	4.5	D	2002/03	2021/10
BIK	Bialystok	Poland	MPI-BGC	23.01	53.23	483.00	25.0	C	2005/07	2014/06
BIR	Birkenes	Norway	NILU	8.25	58.39	225.00	25.0	C	2014/01	2014/01
BKT	Bukit Kototabang	Indonesia	NOAA	100.31	-0.20	875.00	75.0	D	2004/01	2021/11
BLK	Baker Lake, Nunavut	Canada	EC	-96.01	64.33	61.00	15.0	C	2017/07	2019/11
BME	St. Davids Head, Bermuda	United Kingdom	NOAA	-64.65	32.37	17.00	15.0	D	2000/01	2010/01
BMW	Tudor Hill, Bermuda	United Kingdom	NOAA	-64.88	32.26	33.00	15.0	D	2000/01	2021/12
BRA	Bratt's Lake Saskatchewan	Canada	EC	-104.71	50.20	630.00	75.0	C	2009/10	2020/12
BRW	Barrow Atmospheric Baseline Observatory	United States	NOAA	-156.61	71.32	27.46	15.0	C	2000/01	2021/12
BRW	Barrow Atmospheric Baseline Observatory	United States	NOAA	-156.58	71.32	16.00	15.0	D	2000/01	2021/11
BRZ	Berezorechka	Russian Federation	NIES	84.33	56.15	248.00	75.0	C	2008/05	2018/12
BSD	Bilsdale	United Kingdom	UNIVBRIS	-1.15	54.36	628.00	30.0	C	2014/01	2019/12
CBA	Cold Bay, Alaska	United States	NOAA	-162.71	55.21	25.00	15.0	D	2000/01	2021/11
CBY	Cambridge Bay, Nunavut Territory	Canada	EC	-105.06	69.13	47.00	15.0	C	2012/12	2020/12
CFA	Cape Ferguson	Australia	CSIRO	147.06	-19.28	5.00	25.0	D	2000/01	2021/04
CGO	Cape Grim, Tasmania	Australia	NOAA	144.68	-40.68	164.00	4.5	D	2000/01	2021/10
CGO	Cape Grim	Australia	CSIRO	144.68	-40.68	94.00	15.0	C	2012/07	2021/07
CGR	Charles Point, Darwin	Australia	CSIRO	12.65	37.67	9.00	25.0	C	2015/04	2018/12
CHL	Churchill, Manitoba	Canada	EC	-93.82	58.74	89.00	15.0	C	2011/10	2020/12
CHM	Chibougamau, Quebec	Canada	EC	-74.34	49.69	423.00	30.0	C	2007/08	2011/04
CHR	Christmas Island	Republic of Kiribati	NOAA	-157.15	1.70	5.00	15.0	D	2000/01	2020/01
CPS	Chapais, Quebec	Canada	EC	-74.98	49.82	431.00	15.0	C	2011/12	2020/12
CPT	Cape Point	South Africa	NOAA	18.49	-34.35	260.00	25.0	D	2010/02	2021/10
CRI	Cape Rama	India	CSIRO	73.83	15.08	66.00	75.0	D	2000/01	2013/01
CRV	Carbon in Arctic Reservoirs Vulnerability Expe...	United States	NOAA	-147.60	64.99	643.13	15.0	C	2011/10	2021/12
CRZ	Crozet Island	France	NOAA	51.85	-46.43	202.00	4.5	D	2000/01	2021/01
CUR	Monte Curcio	Italy	IIA	16.42	39.32	1801.00	15.0	C	2014/12	2017/12
CYA	Casey Station, Antarctica	Australia	CSIRO	110.52	-66.28	55.00	4.5	D	2000/01	2021/01
DEM	Demyanskoe	Russian Federation	NIES	70.87	59.79	155.00	30.0	C	2005/09	2018/12
DRP	Drake Passage	NOAA	NOAA	-61.68	-59.07	10.00	4.5	D	2006/03	2021/05
DSI	Dongsha Island	Taiwan	NOAA	116.73	20.70	8.00	15.0	D	2010/03	2021/10
DVV	Danville, Virginia	United States	PSU	-79.44	36.71	492.00	15.0	C	2016/07	2017/12
EGB	Egbert, Ontario	Canada	EC	-79.78	44.23	276.00	25.0	C	2005/03	2020/12
EIC	Easter Island	Chile	NOAA	-109.45	-27.13	72.00	4.5	D	2000/01	2019/11
ENA	Eastern North Atlantic, Graciosa, Azores	Portugal	LBNL-ARM	-28.03	39.09	40.48	25.0	C	2015/07	2019/12

ESP	Estevan Point, British Columbia	Canada	EC	-126.54	49.38	47.00	25.0	C	2009/03	2020/12
EST	Esther, Alberta	Canada	EC	-110.21	51.67	757.00	30.0	C	2010/01	2020/11
ETL	East Trout Lake, Saskatchewan	Canada	EC	-104.99	54.35	598.00	30.0	C	2005/08	2020/10
FNE	Fort Nelson, British Columbia	Canada	EC	-122.57	58.84	376.00	30.0	C	2014/07	2020/12
FSD	Fraserdale	Canada	EC	-81.57	49.88	250.00	30.0	C	2000/01	2020/05
GAT	Gartow	Germany	ICOS-ATC,HPB	11.44	53.07	411.00	25.0	C	2016/05	2021/12
GCI	Millerville, AL	United States	PSU	-85.89	33.18	428.00	25.0	C	2017/10	2018/05
GMI	Mariana Islands	Guam	NOAA	144.66	13.39	8.00	15.0	D	2000/01	2021/09
GPA	Gunn Point	Australia	CSIRO	131.04	-12.25	37.00	75.0	D	2010/08	2021/02
HBA	Halley Station, Antarctica	United Kingdom	NOAA	-26.21	-75.61	35.00	4.5	D	2000/01	2021/02
HEI	Heidelberg	Germany	IUP	8.68	49.42	143.00	30.0	C	2005/01	2014/09
HNP	Hanlan's Point, Ontario	Canada	EC	-79.39	43.61	97.00	25.0	C	2014/06	2020/12
HPB	Hohepeissenberg	Germany	ICOS-ATC,HPB	11.02	47.80	1065.00	25.0	C	2015/09	2021/12
HSU	Humboldt State University	United States	NOAA	-124.44	41.57	7.60	30.0	D	2008/05	2017/05
HTM	Hyltemossa	Sweden	ICOS-ATC,LUND-CEC	13.42	56.10	265.00	25.0	C	2017/04	2021/12
ICE	Storhofdi, Vestmannaeyjar	Iceland	NOAA	-20.29	63.40	121.70	15.0	D	2000/01	2021/11
IGR	Igrim	Russian Federation	NIES	64.41	63.19	89.00	30.0	C	2005/04	2013/07
INU	Inuvik, Northwest Territories	Canada	EC	-133.53	68.32	123.00	15.0	C	2012/02	2020/12
IPR	Ispra	Italy	ICOS-ATC,JRC	8.64	45.81	310.00	30.0	C	2007/10	2021/12
IZO	Izana, Tenerife, Canary Islands	Spain	NOAA	-16.48	28.30	2377.90	25.0	D	2000/01	2021/10
JFJ	Jungfrauoch	Switzerland	ICOS-ATC,HFSJG	7.99	46.55	3585.00	15.0	C	2005/02	2021/12
KEY	Key Biscayne, Florida	United States	NOAA	-80.20	25.67	6.00	25.0	D	2000/02	2021/12
KIT	Karlsruhe	Germany	ICOS-ATC,HPB	8.42	49.09	310.00	30.0	C	2016/12	2021/12
KJN	Kjolnes	Norway	UEXE, MPI-BGC	29.23	70.85	20.00	15.0	C	2013/10	2018/08
KMP	Kumpula	Finland	FMI	24.96	60.20	53.00	30.0	C	2010/01	2021/12
KRE	Kresin u Pacova	Czech Republic	ICOS	15.08	49.57	784.00	25.0	C	2017/04	2021/12
KRS	Karasevoe	Russian Federation	NIES	82.42	58.25	156.00	30.0	C	2004/09	2018/12
KUM	Cape Kumukahi, Hawaii	United States	NOAA	-155.01	19.51	3.00	15.0	D	2000/01	2021/12
LEF	Park Falls, Wisconsin	United States	NOAA	-90.27	45.95	868.00	30.0	C	2010/09	2021/12
LEF	Park Falls, Wisconsin	United States	NOAA	-90.26	45.95	868.00	30.0	D	2000/01	2021/12
LIN	Lindenberg	Germany	ICOS-ATC,HPB	14.12	52.17	171.00	30.0	C	2015/10	2021/12
LLB	Lac La Biche, Alberta	Canada	NOAA	-112.45	54.95	546.10	30.0	D	2008/01	2013/02
LLB	Lac La Biche, Alberta	Canada	EC	-112.47	54.95	590.00	30.0	C	2007/04	2020/12
LLN	Lulin	Taiwan	NOAA	120.86	23.47	2867.00	25.0	D	2006/08	2021/11
LMP	Lampedusa	Italy	ICOS-ATC,ENEA	12.63	35.52	53.00	25.0	C	2020/01	2021/12
LMT	Lamezia Terme	Italy	ISAC	16.23	38.88	14.00	30.0	C	2015/01	2016/12
LPO	Ile Grande	France	LSCE	-3.58	48.80	20.00	15.0	D	2005/01	2013/08
LUT	Lutjewad	Netherlands	ICOS-ATC,RUG	6.35	53.40	61.00	25.0	C	2006/05	2021/12
MAA	Mawson, Antarctica	Australia	CSIRO	62.87	-67.62	32.00	4.5	D	2000/02	2021/02
MEX	High Altitude Global Climate Observation Center	Mexico	NOAA	-97.31	18.98	4469.00	15.0	D	2009/01	2021/09
MID	Sand Island, Midway	United States	NOAA	-177.38	28.21	8.00	15.0	D	2000/01	2021/11
MKN	Mt. Kenya	Kenya	NOAA	37.30	-0.06	3649.00	25.0	D	2003/12	2011/06
MLO	Mauna Loa, Hawaii	United States	NOAA	-155.58	19.54	3437.00	15.0	C	2000/01	2021/12
MLO	Mauna Loa, Hawaii	United States	NOAA	-155.58	19.54	3402.00	15.0	D	2000/01	2021/12
MNM	Minamitorishima	Japan	JMA	153.98	24.29	27.10	15.0	C	2000/01	2020/12
MQA	Macquarie Island	Australia	CSIRO	158.97	-54.48	13.00	4.5	D	2000/01	2021/02
MRC	Marcellus Pennsylvania	United States	PSU	-76.42	41.47	652.00	75.0	C	2015/05	2018/12
NAT	Farol De Mae Luiza Lighthouse	Brazil	NOAA	-35.19	-5.51	20.00	15.0	D	2010/09	2020/03

NGL	Neuglobsow	Germany	UBA	13.03	53.14	62.00	75.0	C	2005/01	2013/12
NMB	Gobabeb	Namibia	NOAA	15.01	-23.58	461.00	25.0	D	2000/01	2021/09
NOR	Norunda	Sweden	ICOS-ATC,LUND-CEC	17.48	60.09	146.00	15.0	C	2017/04	2021/12
NOY	Noyabrsk	Russian Federation	NIES	75.78	63.43	188.00	30.0	C	2005/10	2018/12
NWR	Niwot Ridge, Colorado	United States	NOAA	-105.57	40.05	3526.00	15.0	D	2000/01	2021/11
OPE	Observatoire perenne de l'environnement	France	ICOS-ATC,LSCE	5.50	48.56	510.00	30.0	C	2011/07	2021/12
OTA	Otway Basin	Australia	CSIRO	142.82	-38.52	50.00	30.0	D	2005/09	2014/08
OXK	Ochsenkopf	Germany	ICOS-ATC,HPB	11.81	50.03	1185.00	30.0	C	2006/06	2021/12
PAL	Pallas-Sammaltunturi, GAW Station	Finland	ICOS-ATC,FMI	24.12	67.97	577.00	15.0	C	2005/01	2021/12
PDM	Pic du Midi	France	LSCE	0.14	42.94	2887.00	15.0	D	2005/02	2018/02
POC	Pacific Ocean	Pacific Ocean	NOAA	-130.75	0.12	20.00	15.0	D	2000/01	2017/07
PSA	Palmer Station, Antarctica	United States	NOAA	-64.05	-64.77	15.00	4.5	D	2000/01	2021/05
PTA	Point Arena, California	United States	NOAA	-123.74	38.95	22.00	25.0	D	2000/01	2011/05
PUI	Puijo	Finland	ICOS-ATC, UEF	27.66	62.91	84.00	30.0	C	2011/11	2020/12
PUY	Puy de Dome	France	ICOS-ATC,LSCE	2.97	45.77	1475.00	15.0	C	2010/07	2021/12
RPB	Ragged Point	Barbados	NOAA	-59.43	13.16	20.00	15.0	D	2000/01	2021/11
RUN	La Réunion	France	ICOS-ATC,LSCE	55.38	-21.08	2160.00	15.0	C	2018/05	2021/12
RYO	Ryori	Japan	JMA	141.82	39.03	280.00	15.0	C	2000/01	2020/12
SAC	Saclay	France	ICOS-ATC,CEA	2.14	48.72	260.00	75.0	C	2017/05	2021/12
SCT	Beech Island, South Carolina	United States	NOAA	-81.83	33.41	420.20	75.0	C	2015/08	2021/12
SDZ	Shangdianzi	China	NOAA	117.12	40.65	298.00	15.0	D	2009/09	2015/09
SEY	Mahe Island	Seychelles	NOAA	55.53	-4.68	7.00	15.0	D	2000/01	2021/08
SGP	Southern Great Plains, Oklahoma	United States	NOAA	-97.50	36.62	339.00	75.0	D	2002/04	2021/12
SGP	Southern Great Plains, Oklahoma	United States	LBNL-ARM	-97.49	36.61	374.00	75.0	C	2011/01	2020/12
SHM	Shemya Island, Alaska	United States	NOAA	174.08	52.72	28.00	25.0	D	2000/01	2021/07
SMO	Tutuila	American Samoa	NOAA	-170.56	-14.23	60.30	15.0	D	2000/01	2021/11
SMR	Hyytiala	Finland	ICOS-ATC,UHEL5	24.29	61.85	306.00	25.0	C	2016/12	2021/12
SNB	Sonnblick	Austria	EAA	47.05	12.96	3111.00	15.0	C	2012/04	2018/12
SOD	Sodankylä	Finland	FMI	26.64	67.36	227.00	25.0	C	2012/01	2021/12
SPO	South Pole, Antarctica	United States	NOAA	-24.80	-89.96	2821.30	4.5	D	2000/01	2021/10
STE	Steinkimmen	Germany	ICOS-ATC,HPB	8.46	53.04	281.00	75.0	C	2019/07	2021/12
SUM	Summit	Greenland	NOAA	-38.42	72.60	3214.54	15.0	D	2000/08	2021/07
SVB	Svartberget	Sweden	ICOS-ATC,SLU	19.77	64.26	419.00	25.0	C	2017/06	2021/12
SYO	Syowa Station, Antarctica	Japan	NOAA	39.59	-69.00	16.00	4.5	D	2000/01	2020/12
TAC	Tacolneston	United Kingdom	NOAA	1.14	52.52	236.00	25.0	D	2014/06	2016/01
TAP	Tae-ahn Peninsula	Republic of Korea	NOAA	126.13	36.73	21.00	75.0	D	2000/01	2021/10
THD	Trinidad Head, California	United States	NOAA	-124.15	41.05	112.00	25.0	D	2002/04	2017/06
TIK	Hydrometeorological Observatory of Tiksi	Russia	NOAA	128.89	71.60	29.00	15.0	D	2011/08	2018/09
TIK	Tiksi	Russian Federation	FMI	128.89	71.60	29.00	15.0	C	2010/09	2020/12
TOH	Torfhaus	Germany	ICOS-ATC,HPB	10.53	51.81	948.00	25.0	C	2017/12	2021/12
TPD	Turkey Point, Ontario	Canada	EC	-80.56	42.64	266.00	25.0	C	2012/11	2020/12
TRN	Trainou	France	ICOS-ATC,LSCE	2.11	47.96	311.00	25.0	C	2006/08	2021/12
USH	Ushuaia	Argentina	NOAA	-68.31	-54.85	32.00	4.5	D	2000/01	2021/06
UTA	Wendover, Utah	United States	NOAA	-113.72	39.90	1332.00	25.0	D	2000/01	2021/12
UTO	Uto	Finland	FMI	21.37	59.78	65.00	25.0	C	2012/03	2018/03
UTO	Uto	Finland	ICOS-ATC,FMI	21.37	59.78	65.00	25.0	C	2018/03	2021/12
UUM	Ulaan Uul	Mongolia	NOAA	111.10	44.45	1012.00	25.0	D	2000/01	2020/10
VGN	Vaganovo	Russian Federation	NIES	62.32	54.50	277.00	30.0	C	2008/06	2018/12

VKV	Voekovo	Russian Federation	MGO	30.70	59.95	76.00	25.0	C	2008/05	2014/12
WIS	Weizmann Institute of Science at the Arava Ins...	Israel	NOAA	35.06	29.96	482.00	25.0	D	2000/01	2021/11
WLG	Mt. Waliguan	Peoples Republic of China	NOAA	100.90	36.27	3890.00	15.0	D	2000/01	2021/09
WPC	Western Pacific Cruise	Western Pacifi	NOAA	143.70	0.13	10.00	15.0	D	2004/12	2013/06
WSA	Sable Island, Nova Scotia	Canada	EC	-60.01	43.93	8.00	25.0	C	2003/06	2020/03
YAK	Yakutsk	Russian Federation	NIES	129.36	62.09	344.00	30.0	C	2007/09	2013/12
YON	Yonagunijima	Japan	JMA	123.01	24.47	50.00	30.0	C	2000/01	2020/12
ZEP	Ny-Alesund, Svalbard	Norway and Sweden	ICOS-ATC,NILU	11.89	78.91	489.00	15.0	C	2012/04	2021/12
ZOT	Zotino	Russian Federation	MPI-BGC	89.21	60.48	415.00	25.0	C	2009/05	2016/12
ZOT	Zotino	Russian Federation	MPI-BGC	89.21	60.48	415.00	15.0	D	2006/10	2013/06

*Sampling heights from which atmospheric CH₄ is sampled in TM5. **Observations used in this study between 2010 and 2021.




Figure A1. The locations of the atmospheric CH₄ measurement sites and the type of measurement used (continuous, discrete or both) in the inversions. The areas optimised regionally are shown with blues and reds, and the grey colour shows the area optimised at 1° latitude × 1° longitude resolution.




Figure A2. Relationship between grid-based and region-based melting seasons/melt periods in 2011-2021. R^2 and p in the legends are the coefficient of determination and p -values of the slopes from linear regression fit, indicating statistical significance of the coefficient of determination.






Figure A3. Relationship between region-based melt period and its' starting day (a) and grid-based melt period and its' average starting day (b) melting seasons in 2011-2021. R^2 and p in the legends are the coefficient of determination and p -values of the slopes from linear regression fit, indicating statistical significance of the coefficient of determination.



Table A2. Region-based melting-season-melt period start and end days, as well as the length of the melting-season-melt period in days for the four permafrost zones: non-permafrost, sporadic, discontinuous and continuous permafrost defined for each zone. The start and end days are represented as a number of days from the beginning of the year, with number one being the first day of the year (day-of-year).

	non-permafrost			sporadic			discontinuous			continuous permafrost		
	Start	End	Length	Start	End	Length	Start	End	Length	Start	End	Length
2011	80	130	51	105	138	34	109	149	41	114	167	54
2012	76	129	54	106	136	31	110	150	41	126	163	38
2013	74	130	57	113	140	28	117	148	32	121	168	48
2014	72	130	59	100	138	39	102	154	53	125	170	46
2015	71	129	59	105	134	30	114	148	35	127	167	41
2016	61	126	66	97	139	43	114	147	34	126	165	40
2017	75	132	58	93	142	50	115	150	36	132	171	40
2018	82	131	50	92	148	57	113	155	43	126	168	43
2019	77	127	51	94	137	44	114	150	37	128	165	38
2020	70	130	61	100	139	40	103	152	50	124	169	46
2021	69	129	61	99	137	39	113	150	38	125	168	44
mean	73	129	57	100	139	40	111	150	40	125	167	43

a) Melt period [max-min]



b) Start date [max-min]



c) End date [max-min]





Figure A4. Grid-based difference between the maximum and minimum length (a), start day (b) and end day (c) of the melting-season-melt period during the study period (2011-2021).

Table A3. Region-based melting-season-posterior melt period methane emissions in the four permafrost zones; non-permafrost, sporadic, discontinuous, and continuous permafrost, as well as the total sum of the emissions.

	non-permafrost (Gg)	sporadic (Gg)	discontinuous (Gg)	continuous permafrost (Gg)	total emissions (Gg)
2011	1293.7	205.1	116.9	315.5	1931.1
2012	1014.6	206.2	127.6	209.3	1557.7
2013	1121.9	209.3	99.5	277.0	1707.7
2014	1248.4	294.4	183.4	327.9	2054.1
2015	1275.5	210.3	150.6	249.1	1885.6
2016	1378.8	271.8	132.2	201.2	1984.0
2017	1268.7	266.2	121.8	254.3	1911.1
2018	1102.7	302.4	162.8	261.1	1829.0
2019	1063.5	231.5	124.5	161.1	1580.5
2020	1216.6	250.3	144.6	253.7	1865.2
2021	1195.7	230.2	129.5	271.9	1827.2
mean	1198.2	243.4	135.8	252.9	1830.3

Table A4. Grid-based melting-season-posterior melt period emissions in the four permafrost zones; non-permafrost, sporadic, discontinuous, and continuous permafrost, as well as the total sum of the emissions.

	non-permafrost (Gg)	sporadic (Gg)	discontinuous (Gg)	continuous permafrost (Gg)	total emissions (Gg)
2011	257.2	63.2	36.9	52.3	409.5
2012	260.8	77.2	27.0	32.2	397.2
2013	257.1	62.3	21.5	36.6	377.5
2014	304.1	84.8	37.0	49.6	475.5
2015	333.2	70.6	34.8	35.9	474.5
2016	356.3	75.9	37.0	38.6	507.8
2017	402.5	75.8	32.3	45.7	556.3
2018	310.0	81.0	32.3	42.9	466.1
2019	350.2	77.0	34.6	28.0	489.8
2020	281.7	64.7	35.2	27.5	409.1
2021	255.3	74.5	36.9	35.4	402.0
mean	306.2	73.4	33.2	38.6	451.4




Figure A5. Grid-based melting-season-posterior melt period methane emissions and the mean temperature in the four permafrost zones; a) non-permafrost, b) sporadic, c) discontinuous, and d) continuous permafrost. Grid-cells-grid cells with negative emission rate were masked out. R^2 and p in the legends are the coefficient of determination and p -values of the slopes from linear regression fit, indicating statistical significance of the coefficient of determination.

Table A5. ~~Melting season~~ Melt period posterior methane emissions in the Hudson Bay lowlands and Western Siberian lowlands.

	region-based [Gg]			grid-based [Gg]		
	HBL	WSL	all	HBL	WSL	all
2011	84.4	514.0	598.4	24.7	73.3	97.9
2012	115.7	393.6	509.2	44.3	81.7	125.9
2013	91.9	407.6	499.5	21.0	89.1	110.1
2014	116.9	533.1	650.0	41.2	137.8	179.0
2015	70.7	609.8	680.5	17.4	142.4	159.8
2016	81.6	514.6	596.3	22.4	126.2	148.5
2017	101.6	532.6	634.2	30.4	123.2	153.5
2018	111.6	420.6	532.2	39.4	113.4	152.9
2019	84.5	334.2	418.7	28.8	119.3	148.1
2020	114.2	461.5	575.7	45.7	89.7	135.4
2021	98.0	446.3	544.3	29.1	108.7	137.8
mean	97.4	469.8	567.2	31.3	109.5	140.8




Figure A6. Region-based [posterior](#) emission maps of the [melting season-melt period](#) in the years that had some of the highest emissions in the Hudson Bay lowlands and Western Siberian lowlands. Hudson Bay lowlands have been outlined with red borders, and Western Siberian lowlands with blue borders.




Figure A7. Average monthly emissions from peat Difference between prior and inundation for posterior emissions of the melt period in the years 2011-2019, from that had some of the prior LPX-Bern-DYPTOP v1.4 highest emissions in the Hudson Bay lowlands and Western Siberian lowlands (a,b,c,d,e and f) as well as the mean difference of all years of this study (g). The average region-based melting season is showed Hudson Bay lowlands and Western Siberian lowlands have been outlined with gray shading black.




Figure A8. Figures a and b depict the region-based relationship between the difference of prior and posterior total emissions (a) or the emission rate (b) and the length of the season. Figures c and d depict the grid-based relationship between the difference of prior and posterior total emissions (c) or the mean emission rate (d) and the mean length of the melt period. The scatter plot color gradient represents the different years with 2011 being the lightest color and 2021 being the darkest. R^2 and p in the legends are the coefficient of determination and p -values of the slopes from linear regression fit, indicating statistical significance of the coefficient of determination.




Figure A9. Average monthly emissions from peat, inundation and wetsoil for the years 2011-2019, from the prior LPX-Bern DYPTOP v1.4 emissions. Average temperature is indicated by a black line. The average region-based melt period is showed with gray shading.

Author contributions. S.H., M.T. and T.A. participated in the design of the study. T.A. supervised the project. M.T and T.A. offered advice of
745 the analysis of the results. M.T. performed the model runs with the CTE-CH₄ and A.T. helped in setting up the model runs and interpretation
of the results. S.H. did the data-analysis and prepared the visualizations, as well as wrote the original manuscript with the help of M.T. and
T.A.. K.R. provided and helped to interpret the SMOS F/T soil state data. A.E. postprocessed the SMOS F/T data for the analysis and helped
to interpret the results of the SMOS F/T data. M.S. provided CH₄ mole fraction measurements from sites in Western Siberian lowlands.
750 H.A. provided CH₄ mole fraction measurements from Kumpula and Sodankylä sites. All authors have read and commented and approved
the published version of the manuscript.

Competing interests. The authors declare no conflict of interest. Also, the funders had no role in the design of the study; in the collection,
analysis, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Acknowledgements. We thank the team behind the LPX-Bern DYPTOP v1.4 for providing the CH₄ emission estimates. The authors would like to thank the ICOS and ICOS-Finland PIs for providing the data on CH₄ mole fractions. We thank the Finnish Meteorological Institute (PAL, UTO, KMP, SOD), University of Eastern Finland (PUI) and University of Helsinki (SMR) for providing the methane data in Finland. We are grateful for CSIRO Oceans and Atmosphere, Climate Science Centre (CSIRO), Environment and Climate Change Canada (ECCC), the Hungarian Meteorological Service (HMS), the Institute for Atmospheric Sciences and Climate (ISAC), the Institute on Atmospheric Pollution of the National Research Council (IIA), the Institute of Environmental Physics, University of Heidelberg (IUP), Laboratoire des Sciences du Climat et de l'Environnement (LSCE), Lawrence Berkeley National Laboratory (LBNL-ARM), the Environment Division Global Environment and Marine Department Japan Meteorological Agency (JMA), the Main Geophysical Observatory (MGO), the Max Planck Institute for Biogeochemistry (MPIBG), National Institute for Environmental Studies (NIES), Norwegian Institute for Air Research (NILU), National Oceanic and Atmospheric Administration Earth System Research Laboratories (NOAA ESRL), the Pennsylvania State University (PSU), Swedish University of Agricultural Sciences (SLU), the Swiss Federal Laboratories for Materials Science and Technology (EMPA), Umweltbundesamt Germany/Federal Environmental Agency (UBA), Umweltbundesamt Austria/Environment Agency Austria (EAA) as the data provider for Sonnblick, University of Bristol (UNIVBRIS), University of Exeter (Univ. Exeter), and University of Urbino (UNIURB) for performing high-quality CH₄ measurements at global sites and making them available through the Global Atmosphere Watch - World Data Centre for Greenhouse Gases (GAW-WDCGG) and personal communications.

Financial support. We thank the European Space Agency ESRIN Contract No: 4000124500/18/I-EF (SMOS F/T Service) 2:44, 4000125046/18/I-NB (MethEO), 4000137895/22/I-AG MethaneCAMP, AO/1-10901/21/I-DT AMPAC-Net, FIRI - ICOS Finland (345531), ICOS-ERIC (281250), and 351311 (GHGSUPER), 337552 and 359196 (Flagships ACCC and FAME), ESA AO/1-11844/23/I-NS SMART-CH₄ and the Research Council of Finland (364034) for financial support.

References

- Al Bitar, A., Mialon, A., Kerr, Y. H., Cabot, F., Richaume, P., Jacquette, E., Quesney, A., Mahmoodi, A., Tarot, S., Parrens, M., Al-Yaari, A., Pellarin, T., Rodriguez-Fernandez, N., and Wigneron, J.-P.: The global SMOS Level 3 daily soil moisture and brightness temperature maps, *Earth System Science Data*, 9, 293–315, <https://doi.org/10.5194/essd-9-293-2017>, 2017.
- Aselmann, I. and Crutzen, P. J.: Global distribution of natural freshwater wetlands and rice paddies, their net primary productivity, seasonality and possible methane emissions, *Journal of Atmospheric Chemistry*, 8, 307–358, <https://doi.org/10.1007/BF00052709>, 1989.
- Bao, T., Xu, X., Jia, G., Billesbach, D. P., and Sullivan, R. C.: Much stronger tundra methane emissions during autumn freeze than spring thaw, *Glob Change Biology*, 27, 376–387, <https://doi.org/10.1111/gcb.15421>, 2020.
- Bartsch, A., Westermann, S., Strozzi, T., Wiesmann, A., and Kroisleitner, C.: Permafrost cci. D1.2 Product Specifications Document (PSD), Tech. rep., ESA, (version 3.0), 2020.
- Bruhwyler, L., Dlugokencky, E., Masarie, K., Ishizawa, M., Andrews, A., Miller, J., Sweeney, C., Tans, P., and Worthy, D.: CarbonTracker-CH₄: an assimilation system for estimating emissions of atmospheric methane, *Atmospheric Chemistry and Physics*, 14, 8269–8293, <https://doi.org/10.5194/acp-14-8269-2014>, 2014.
- Cao, B., Gruber, S., Zheng, D., and Li, X.: The ERA5-Land soil temperature bias in permafrost regions, *The Cryosphere*, 14, 2581–2595, <https://doi.org/10.5194/tc-14-2581-2020>, 2020.
- Castro-Morales, K., Kleinen, T., Kaiser, S., Zaehle, S., Kittler, F., Kwon, M. J., Beer, C., and Göckede, M.: Year-round simulated methane emissions from a permafrost ecosystem in Northeast Siberia, *Biogeosciences*, 15, 2691–2722, <https://doi.org/10.5194/bg-15-2691-2018>, 2018.
- de Vrese, P., Beckebanze, L., Galera, L. d. A., Holl, D., Kleinen, T., Kutzbach, L., Rehder, Z., and Brovkin, V.: Sensitivity of Arctic CH₄ emissions to landscape wetness diminished by atmospheric feedbacks, *Nature Climate Change*, 13, 832–839, <https://doi.org/10.1038/s41558-023-01715-3>, 2023.
- ECMWF: ECMWF Reanalysis v5 (ERA5), <https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-v5>, last access: 11 Jun 2025.
- Erkkilä, A., Tenkanen, M. K., Tsuruta, A., Rautiainen, K., and Aalto, T.: Environmental and Seasonal Variability of High Latitude Methane Emissions Based on Earth Observation Data and Atmospheric Inverse Modelling, *Remote Sensing*, 15, <https://doi.org/10.3390/rs15245719>, 2023.
- ESA: ESA SMOS Dissemination server, <https://smos-diss.eo.esa.int/>, last access: 9 December 2024.
- European Space Agency: SMOS Soil Freeze and Thaw State, Version 300, <https://doi.org/10.57780/sm1-fbf89e0>, accessed on: March 4, 2026, 2023.
- Evensen, G.: The Ensemble Kalman Filter: theoretical formulation and practical implementation, *Ocean Dynamics*, 53, 343–367, <https://doi.org/10.1007/s10236-003-0036-9>, 2003.
- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H.: The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity, vol. 18, p. 923–1054, Cambridge University Press, <https://doi.org/10.1017/9781009157896.009>, 2021.
- Hargreaves, K. J., Fowler, D., Pitcairn, C. E. R., and Aurela, M.: Annual methane emission from Finnish mires estimated from eddy covariance campaign measurements, *Theoretical and Applied Climatology*, 70, 203–213, <https://doi.org/10.1007/s007040170015>, 2001.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren,

- P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J.,
810 Hólm, E., Janisková, M., Keeley, S., Lalouaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Vil-
laume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049,
<https://doi.org/10.1002/qj.3803>, 2020.
- Houweling, S., Krol, M., Bergamaschi, P., Frankenberg, C., Dlugokencky, E. J., Morino, I., Notholt, J., Sherlock, V., Wunch, D., Beck,
V., Gerbig, C., Chen, H., Kort, E. A., Röckmann, T., and Aben, I.: A multi-year methane inversion using SCIAMACHY, accounting
815 for systematic errors using TCCON measurements, *Atmospheric Chemistry and Physics*, 14, 3991–4012, <https://doi.org/10.5194/acp-14-3991-2014>, 2014.
- Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M., Siewert,
M. B., Treat, C., Turetsky, M., Voigt, C., and Yu, Z.: Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw,
Proceedings of the National Academy of Sciences, 117, 20438–20446, <https://doi.org/10.1073/pnas.1916387117>, 2020.
- 820 Ilmatieteen laitos: Kevätsään tilastot, <https://www.ilmatieteenlaitos.fi/kevattilastot>, last access: 23 August 2023, 2025.
- Ito, A. and Inatomi, M.: Use of a process-based model for assessing the methane budgets of global terrestrial ecosystems and evaluation of
uncertainty., *Biogeosciences*, 9, 759–773, <https://doi.org/10.5194/bg-9-759-2012>, 2012.
- Ito, A., Li, T., Qin, Z., Melton, J. R., Tian, H., Kleinen, T., Zhang, W., Zhang, Z., Joos, F., Ciais, P., Hopcroft, P. O., Beerling, D. J.,
Liu, X., Zhuang, Q., Zhu, Q., Peng, C., Chang, K.-Y., Fluet-Chouinard, E., McNicol, G., Patra, P., Poulter, B., Sitch, S., Riley, W.,
825 and Zhu, Q.: Cold-Season Methane Fluxes Simulated by GCP-CH₄ Models, *Geophysical Research Letters*, 50, e2023GL103037,
<https://doi.org/10.1029/2023GL103037>, e2023GL103037 2023GL103037, Supporting Information, 2023.
- Jin, H., Wu, J., Cheng, G., Nakano, T., and Sun, G.: Methane emissions from wetlands on the Qinghai-Tibet Plateau, *Chinese Science
Bulletin*, 44, 2282–2286, <https://doi.org/10.1007/BF02885940>, 1999.
- Jöckel, P., Tost, H., Pozzer, A., Brühl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M. G., Sander, R., Steil, B.,
830 Stiller, G., Tanarhte, M., Taraborrelli, D., van Aardenne, J., and Lelieveld, J.: The atmospheric chemistry general circulation model
ECHAM5/MESSy1: consistent simulation of ozone from the surface to the mesosphere, *Atmospheric Chemistry and Physics*, 6, 5067–
5104, <https://doi.org/10.5194/acp-6-5067-2006>, 2006.
- Kangasaho, V., Tsuruta, A., Backman, L., Mäkinen, P., Houweling, S., Segers, A., Krol, M., Dlugokencky, E. J., Michel, S., White, J. W. C.,
and Aalto, T.: The Role of Emission Sources and Atmospheric Sink in the Seasonal Cycle of CH₄ and $\delta^{13}\text{-CH}_4$: Analysis Based on the
835 Atmospheric Chemistry Transport Model TM5, *Atmosphere*, 13, <https://doi.org/10.3390/atmos13060888>, 2022.
- Kerr, Y. H., Waldteufel, P., Wigneron, J.-P., Delwart, S., Cabot, F., Boutin, J., Escorihuela, M.-J., Font, J., Reul, N., Gruhier, C., Juglea, S. E.,
Drinkwater, M. R., Hahne, A., Martín-Neira, M., and Mecklenburg, S.: The SMOS Mission: New Tool for Monitoring Key Elements of the
Global Water Cycle, *Proceedings of the IEEE*, 98, 666–687, <https://doi.org/10.1109/JPROC.2010.2043032>, 2010.
- Knoblauch, C., Beer, C., Liebner, S., Grigoriev, M., and Pfeiffer, E.: Methane production as key to the greenhouse gas budget of thawing
840 permafrost., *Nature Climate Change*, 8, 309–312, <https://doi.org/10.1038/s41558-018-0095-z>, 2018.
- Krol, M., Houweling, S., Bregman, B., van den Broek, M., Segers, A., van Velthoven, P., Peters, W., Dentener, F., and Bergamaschi, P.:
The two-way nested global chemistry-transport zoom model TM5: algorithm and applications, *Atmospheric Chemistry and Physics*, 5,
417–432, <https://doi.org/10.5194/acp-5-417-2005>, 2005.
- Lawrence, D. M., Koven, C. D., Swenson, S. C., Riley, W. J., and Slater, A. G.: Permafrost thaw and resulting soil moisture changes
845 regulate projected high-latitude CO₂ and CH₄ emissions, *Environmental Research Letters*, 10, 094011, <https://doi.org/10.1088/1748-9326/10/9/094011>, 2015.

- Li, M., Wu, P., and Ma, Z.: A comprehensive evaluation of soil moisture and soil temperature from third-generation atmospheric and land reanalysis data sets, *International Journal of Climatology*, 40, 5744–5766, <https://doi.org/10.1002/joc.6549>, 2020.
- Lienert, S. and Joos, F.: A Bayesian ensemble data assimilation to constrain model parameters and land-use carbon emissions, *Biogeosciences*, 15, 2909–2930, <https://doi.org/10.5194/bg-15-2909-2018>, 2018.
- 850 Masarie, K. A., Peters, W., Jacobson, A. R., and Tans, P. P.: ObsPack: a framework for the preparation, delivery, and attribution of atmospheric greenhouse gas measurements, *Earth System Science Data*, 6, 375–384, <https://doi.org/10.5194/essd-6-375-2014>, 2014.
- Mastepanov, M., Sigsgaard, C., Tagesson, T., Ström, L., Tamstorf, M. P., Lund, M., and Christensen, T. R.: Revisiting factors controlling methane emissions from high-Arctic tundra, *Biogeosciences*, 10, 5139–5158, <https://doi.org/10.5194/bg-10-5139-2013>, 2013.
- 855 Monforti Ferrario, F., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Lo Vullo, E., Solazzo, E., Olivier, J., and Vignati, E.: EDGAR v6.0 Greenhouse Gas Emissions., European Commission, Joint Research Centre (JRC) [Dataset], PID: <http://data.europa.eu/89h/97a67d67-c62e-4826-b873-9d972c4f670b>, 2021.
- Neumann, R. B., Moorberg, C. J., Lundquist, J. D., Turner, J. C., Waldrop, M. P., McFarland, J. W., Euskirchen, E. S., Edgar, C. W., and Turetsky, M. R.: Warming effects of spring rainfall increase methane emissions from thawing permafrost., *Geophysical Research Letters*, 46, 1393–1401, <https://doi.org/10.1029/2018GL081274>, 2019.
- 860 Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H. H., Dashtseren, A., Delaloye, R., Elberling, B., Eitzelmüller, B., Kholodov, A., Khomutov, A., Kääh, A., Leibman, M. O., Lewkowicz, A. G., Panda, S. K., Romanovsky, V., Way, R. G., Westergaard-Nielsen, A., Wu, T., Yamkhin, J., and Zou, D.: Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale, *Earth-Science Reviews*, 193, 299–316, <https://doi.org/10.1016/j.earscirev.2019.04.023>, 2019.
- 865 Obu, J., Westermann, S., Barboux, C., Bartsch, A., Delaloye, R., Grosse, G., Heim, B., Hugelius, G., Irrgang, A., and Kääh, A.: ESA permafrost climate change initiative (Permafrost_cci): Permafrost extent for the Northern Hemisphere, v3. 0, The Centre for Environmental Data Analysis, RAL Space, 2021.
- Oliva, R., Daganzo, E., Richaume, P., Kerr, Y., Cabot, F., Soldo, Y., Anterrieu, E., Reul, N., Gutierrez, A., Barbosa, J., and Lopes, G.: Status of Radio Frequency Interference (RFI) in the 1400-1427 MHz passive band based on six years of SMOS mission, *Remote Sensing of Environment*, 180, 64–75, <https://doi.org/10.1016/j.rse.2016.01.013>, 2016.
- 870 Parmentier, F. J. W., van Huissteden, J., van der Molen, M. K., Schaepman-Strub, G., Karsanaev, S. A., Maximov, T. C., and Dolman, A. J.: Spatial and temporal dynamics in eddy covariance observations of methane fluxes at a tundra site in northeastern Siberia, *Journal of Geophysical Research: Biogeosciences*, 116, <https://doi.org/10.1029/2010JG001637>, 2011.
- Peltola, O., Vesala, T., Gao, Y., Rätty, O., Alekseychik, P., Aurela, M., Chojnicki, B., Desai, A. R., Dolman, A. J., Euskirchen, E. S., Friborg, T., Göckede, M., Helbig, M., Humphreys, E., Jackson, R. B., Jocher, G., Joos, F., Klatt, J., Knox, S. H., Kowalska, N., Kutzbach, L., Lienert, S., Lohila, A., Mammarella, I., Nadeau, D. F., Nilsson, M. B., Oechel, W. C., Peichl, M., Pypker, T., Quinton, W., Rinne, J., Sachs, T., Samson, M., Schmid, H. P., Sonntag, O., Wille, C., Zona, D., and Aalto, T.: Monthly gridded data product of northern wetland methane emissions based on upscaling eddy covariance observations, *Earth System Science Data*, 11, 1263–1289, <https://doi.org/10.5194/essd-11-1263-2019>, 2019.
- 875 Peters, W., Miller, J. B., Whitaker, J., Denning, A. S., Hirsch, A., Krol, M. C., Zupanski, D., Bruhwiler, L., and Tans, P. P.: An ensemble data assimilation system to estimate CO₂ surface fluxes from atmospheric trace gas observations, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/10.1029/2005JD006157>, 2005.

- Pickett-Heaps, C. A., Jacob, D. J., Wecht, K. J., Kort, E. A., Wofsy, S. C., Diskin, G. S., Worthy, D. E. J., Kaplan, J. O., Bey, I., and Drevet, J.: Magnitude and seasonality of wetland methane emissions from the Hudson Bay Lowlands (Canada), *Atmospheric Chemistry and Physics*, 11, 3773–3779, <https://doi.org/10.5194/acp-11-3773-2011>, 2011.
- 885 Poulter, B., Bousquet, P., Canadell, J. G., Ciais, P., Peregon, A., Saunio, M., Arora, V. K., Beerling, D. J., Brovkin, V., Jones, C. D., Joos, F., Gedney, N., Ito, A., Kleinen, T., Koven, C. D., McDonald, K., Melton, J. R., Peng, C., Peng, S., Prigent, C., Schroeder, R., Riley, W. J., Saito, M., Spahni, R., Tian, H., Taylor, L., Viovy, N., Wilton, D., Wiltshire, A., Xu, X., Zhang, B., Zhang, Z., and Zhu, Q.: Global wetland contribution to 2000–2012 atmospheric methane growth rate dynamics, *Environmental Research Letters*, 12, 094 013, <https://doi.org/10.1088/1748-9326/aa8391>, 2017.
- 890 Randerson, J., Van Der Werf, G., Giglio, L., Collaz, G., and Kasibhatla, P.: Global Fire Emissions Database, Version 4.1 (GFEDv4), <https://doi.org/10.3334/ORNLDAAC/1293>, 2017.
- Rautiainen, K., Parkkinen, T., Lemmetyinen, J., Schwank, M., Wiesmann, A., Ikonen, J., Derksen, C., Davydov, S., Davydova, A., Boike, J., Langer, M., Drusch, M., and Pulliainen, J.: SMOS prototype algorithm for detecting autumn soil freezing, *Remote Sensing of Environment*, 180, 346–360, <https://doi.org/10.1016/j.rse.2016.01.012>, special Issue: ESA’s Soil Moisture and Ocean Salinity Mission - Achievements and Applications, 2016.
- 895 Rautiainen, K., Holmberg, M., Cohen, J., Mialon, A., Schwank, M., Lemmetyinen, J., de la Fuente, A., and Kerr, Y.: An operational SMOS soil freeze–thaw product, *Earth System Science Data*, 17, 5337–5353, <https://doi.org/10.5194/essd-17-5337-2025>, 2025.
- Rawlins, M. A. and Karmalkar, A. V.: Regime shifts in Arctic terrestrial hydrology manifested from impacts of climate warming, *The Cryosphere*, 18, 1033–1052, <https://doi.org/10.5194/tc-18-1033-2024>, 2024.
- 900 Raz-Yaseef, N., Torn, M. S., Wu, Y., Billesbach, D. P., Liljedahl, A. K., Kneafsey, T. J., Romanovsky, V. E., Cook, D. R., and Wullschleger, S. D.: Large CO₂ and CH₄ emissions from polygonal tundra during spring thaw in northern Alaska, *Geophysical Research Letters*, 44, 504–513, <https://doi.org/10.1002/2016GL071220>, 2017.
- Rinne, J., Riutta, T., Pihlatie, M., Aurela, M., Haapanala, S., Tuovinen, J.-P., Tuittila, E.-S., and Vesala, T.: Annual cycle of methane emission from a boreal fen measured by the eddy covariance technique, *Tellus B: Chemical and Physical Meteorology*, 59, 449–457, <https://doi.org/10.1111/j.1600-0889.2007.00261.x>, 2007.
- 905 Rinne, J., Tuittila, E.-S., Peltola, O., Li, X., Raivonen, M., Alekseychik, P., Haapanala, S., Pihlatie, M., Aurela, M., Mammarella, I., and Vesala, T.: Temporal Variation of Ecosystem Scale Methane Emission From a Boreal Fen in Relation to Temperature, Water Table Position, and Carbon Dioxide Fluxes, *Global Biogeochemical Cycles*, 32, 1087–1106, <https://doi.org/10.1029/2017GB005747>, 2018.
- 910 Rychlik, S.: Relating interannual variability of atmospheric CH₄ growth rate to large-scale CH₄ emissions from northern wetlands, Master’s thesis, Department of Physical Geography and Ecosystem Analysis, Lund University, 2009.
- Sachs, T., Wille, C., Boike, J., and Kutzbach, L.: Environmental controls on ecosystem-scale CH₄ emission from polygonal tundra in the Lena River Delta, Siberia, *Journal of Geophysical Research: Biogeosciences*, 113, <https://doi.org/10.1029/2007JG000505>, 2008.
- Sasakawa, M., Shimoyama, K., Machida, T., Tsuda, N., Suto, H., Arshinov, M., Davydov, D., Fofonov, A., Krasnov, O., Saeki, T., Koyama, Y., and Maksyutov, S.: Continuous measurements of methane from a tower network over Siberia, *Tellus B: Chemical and Physical Meteorology*, <https://doi.org/10.1111/j.1600-0889.2010.00494.x>, 2010.
- 915 Sasakawa, M., Tsuda, N., Machida, T., Arshinov, M., Davydov, D., Fofonov, A., and Belan, B.: Revised methodology for CO₂ and CH₄ measurements at remote sites using a working standard-gas-saving system, *Atmospheric Measurement Techniques*, 18, 1717–1730, <https://doi.org/10.5194/amt-18-1717-2025>, 2025.

- 920 Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, 925 J. R., Morino, I., Müller, J., Murguía-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth System Science Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 930 2020.
- Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P. A., Regnier, P., Canadell, J. G., Jackson, R. B., Patra, P. K., Bousquet, P., Ciais, P., Dlugokencky, E. J., Lan, X., Allen, G. H., Bastviken, D., Beerling, D. J., Belikov, D. A., Blake, D. R., Castaldi, S., Crippa, M., Deemer, B. R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L., Holgerson, M. A., Hopcroft, P. O., Hugelius, G., Ito, A., Jain, A. K., Janardanan, R., Johnson, M. S., Kleinen, T., Krummel, P. B., Lauerwald, R., Li, T., Liu, X., McDonald, K. C., Melton, J. R., 935 Mühle, J., Müller, J., Murguía-Flores, F., Niwa, Y., Noce, S., Pan, S., Parker, R. J., Peng, C., Ramonet, M., Riley, W. J., Rocher-Ros, G., Rosentreter, J. A., Sasakawa, M., Segers, A., Smith, S. J., Stanley, E. H., Thanwerdas, J., Tian, H., Tsuruta, A., Tubiello, F. N., Weber, T. S., van der Werf, G. R., Worthy, D. E. J., Xi, Y., Yoshida, Y., Zhang, W., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: Global Methane Budget 2000–2020, *Earth System Science Data*, 17, 1873–1958, <https://doi.org/10.5194/essd-17-1873-2025>, 2025.
- Scharlemann, J. P., Tanner, E. V., Hiederer, R., and Kapos, V.: Global soil carbon: understanding and managing the largest terrestrial carbon 940 pool, *Carbon Management*, 5, 81–91, <https://doi.org/10.4155/cmt.13.77>, 2014.
- Schuldts, K., Aalto, T., Andrews, A., Aoki, S., Arduini, J., Baier, B., Bergamaschi, P., Biermann, T., Biraud, S., Boenisch, H., and et al.: Multi-Laboratory Compilation of Atmospheric Methane Data for the Period 1983–2020, Tech. rep., NOAA Earth System Research Laboratory, Global Monitoring Laboratory, Silver Spring, MD, USA, 2021.
- Schuur, E. and Abbott, B.: High risk of permafrost thaw, *Nature*, 480, 32–33, <https://doi.org/10.1038/480032a>, 2011.
- 945 Schuur, E., McGuire, A., Schädel, C., Grosse, G., Harden, J., Hayes, D., Hugelius, G., Koven, C., Kuhry, P., Lawrence, D., Natali, S., Olefeldt, D., Romanovsky, V., Schaefer, K., Turetsky, M., Treat, C., and Vonk, J.: Climate change and the permafrost carbon feedback., *Science Advances*, 520, 171–179, <https://doi.org/10.1038/nature14338>, 2015.
- SMOS: SMOS Level 3 Soil Freeze/Thaw Service, <http://nsdc.fmi.fi/services/SMOSService/>, last access: 5 March 2024.
- Song, C., Xu, X., Sun, X., Tian, H., Sun, L., Miao, Y., Wang, X., and Guo, Y.: Large methane emission upon spring thaw from natural 950 wetlands in the northern permafrost region, *Environmental Research Letters*, 7, 034 009, <https://doi.org/10.1088/1748-9326/7/3/034009>, 2012.
- Spahni, R., Wania, R., Neef, L., van Weele, M., Pison, I., Bousquet, P., Frankenberg, C., Foster, P. N., Joos, F., Prentice, I. C., and van Velthoven, P.: Constraining global methane emissions and uptake by ecosystems, *Biogeosciences*, 8, 1643–1665, <https://doi.org/10.5194/bg-8-1643-2011>, 2011.
- 955 Stocker, B. D., Spahni, R., and Joos, F.: DYPTOP: a cost-efficient TOPMODEL implementation to simulate sub-grid spatio-temporal dynamics of global wetlands and peatlands, *Geoscientific Model Development*, 7, 3089–3110, <https://doi.org/10.5194/gmd-7-3089-2014>, 2014.

- 960 Tagesson, T., Migliavacca, M., Mastepanov, M., Sigsgaard, C., Tamstorf, M. P., Lund, M. N., Falk, J. M., Lindroth, A., Christensen, T. R., and Ström, L.: Land-atmosphere exchange of methane from soil thawing to soil freezing in a high-arctic wet tundra ecosystem, *Global Change Biology*, 18, 1928–1940, <https://doi.org/10.1111/j.1365-2486.2012.02647.x>, 2012.
- Tenkanen, M. K.: Inversiomalli CarbonTracker Europe - CH₄:n arviot pohjoisten soiden metaaniemissioista maan routaantumisen aikana, Master's thesis, Turun yliopisto, Fysiikan ja tähtitieteen laitos, 2019.
- 965 Tenkanen, M. K., Tsuruta, A., Rautiainen, K., Kangasaho, V., Ellul, R., and Aalto, T.: Utilizing Earth Observations of Soil Freeze/Thaw Data and Atmospheric Concentrations to Estimate Cold Season Methane Emissions in the Northern High Latitudes, *Remote Sensing*, 13, <https://doi.org/10.3390/rs13245059>, 2021.
- Thompson, R. L., Sasakawa, M., Machida, T., Aalto, T., Worthy, D., Lavric, J. V., Lund Myhre, C., and Stohl, A.: Methane fluxes in the high northern latitudes for 2005–2013 estimated using a Bayesian atmospheric inversion, *Atmospheric Chemistry and Physics*, 17, 3553–3572, <https://doi.org/10.5194/acp-17-3553-2017>, 2017.
- 970 Tokida, T., Mizoguchi, M., Miyazaki, T., Kagemoto, A., Nagata, O., and Hatano, R.: Episodic release of methane bubbles from peatland during spring thaw, *Chemosphere*, 70, 165–171, <https://doi.org/10.1016/j.chemosphere.2007.06.042>, 2007.
- Tsuruta, A., Aalto, T., Backman, L., Hakkarainen, J., van der Laan-Luijkx, I. T., Krol, M. C., Spahni, R., Houweling, S., Laine, M., Dlugokencky, E., Gomez-Pelaez, A. J., van der Schoot, M., Langenfelds, R., Ellul, R., Arduini, J., Apadula, F., Gerbig, C., Feist, D. G., Kivi, R., Yoshida, Y., and Peters, W.: Global methane emission estimates for 2000–2012 from CarbonTracker Europe-CH₄ v1.0, *Geoscientific Model Development*, 10, 1261–1289, <https://doi.org/10.5194/gmd-10-1261-2017>, 2017.
- 975 Tsuruta, A., Aalto, T., Backman, L., Krol, M. C., Peters, W., Lienert, S., Joos, F., Miller, P. A., Zhang, W., Laurila, T., Hatakka, J., Leskinen, A., Lehtinen, K. E. J., Peltola, O., Vesala, T., Levula, J., Dlugokencky, E., Heimann, M., Kozlova, E., Aurela, M., Lohila, A., Kauhaniemi, M., and Gomez-Pelaez, A. J.: Methane budget estimates in Finland from the CarbonTracker Europe-CH₄ data assimilation system, *Tellus B: Chemical and Physical Meteorology*, 71, 1565 030, <https://doi.org/10.1080/16000889.2018.1565030>, 2019.
- 980 Turetsky, M., Abbott, B., Jones, M., Anthony, K., Olefeldt, D., Schuur, E., Grosse, G., Kuhry, P., Hugelius, G., Koven, C., Lawrence, D., Gibson, C., Sannel, A., and McGuire, A.: Carbon release through abrupt permafrost thaw., *Nature Geoscience*, 13, 138–143, <https://doi.org/10.1038/s41561-019-0526-0>, 2020.
- Turetsky, M. R., Treat, C. C., Waldrop, M. P., Waddington, J. M., Harden, J. W., and McGuire, A. D.: Short-term response of methane fluxes and methanogen activity to water table and soil warming manipulations in an Alaskan peatland, *Journal of Geophysical Research: Biogeosciences*, 113, <https://doi.org/10.1029/2007JG000496>, 2008.
- 985 Umezawa, T., Machida, T., Aoki, S., and Nakazawa, T.: Contributions of natural and anthropogenic sources to atmospheric methane variations over western Siberia estimated from its carbon and hydrogen isotopes, *Global Biogeochemical Cycles*, 26, <https://doi.org/https://doi.org/10.1029/2011GB004232>, 2012.
- Uranga, E., Llorente, , González, J., de la Fuente, A., Oliva, R., Soldo, Y., and Jorge, F.: SMOS ESA RFI Monitoring and Information Tool: Lessons Learned, *Remote Sensing*, 14, <https://doi.org/10.3390/rs14215387>, 2022.
- 990 U.S. National Ice Center: IMS Daily Northern Hemisphere Snow and Ice Analysis at 1 km, 4 km, and 24 km Resolutions, Version 1, <https://doi.org/10.7265/N52R3PMC>, 2004.
- van der Laan-Luijkx, I. T., van der Velde, I. R., van der Veen, E., Tsuruta, A., Stanislawski, K., Babenhauerheide, A., Zhang, H. F., Liu, Y., He, W., Chen, H., Masarie, K. A., Krol, M. C., and Peters, W.: The CarbonTracker Data Assimilation Shell (CTDAS) v1.0: implementation and global carbon balance 2001–2015, *Geoscientific Model Development*, 10, 2785–2800, <https://doi.org/10.5194/gmd-10-2785-2017>, 995 2017.

- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth System Science Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- 1000 Voigt, C., Marushchak, M., Mastepanov, M., Lamprecht, R., Christensen, T., Dorodnikov, M., Jackowicz-Korczyński, M., Lindgren, A., Lohila, A., Nykänen, H., Oinonen, M., Oksanen, T., Palonen, V., Treat, C., Martikainen, P., and Biasi, C.: Ecosystem carbon response of an Arctic peatland to simulated permafrost thaw., *Global change biology*, 25, 1746–1764, <https://doi.org/10.1111/gcb.14574>, 2019.
- Vourlitis, G. L., Oechel, W. C., Hastings, S. J., and Jenkins, M. A.: A System for Measuring in situ CO₂ and CH₄ Flux in Unmanaged Ecosystems: An Arctic Example, *Functional Ecology*, 7, 369–379, <http://www.jstor.org/stable/2390217>, 1993.
- 1005 Weber, T., Wiseman, N. A., and Kock, A.: Global ocean methane emissions dominated by shallow coastal waters, *Nature Communications*, 10, 4584, <https://doi.org/10.1038/s41467-019-12541-7>, 2019.
- Westermann, S., Barboux, C., Bartsch, A., Delaloye, R., Grosse, G., Heim, B., Hugelius, G., Irrgang, A., Kääb, A., Matthes, H., Nitze, I., Pellet, C., Seifert, F., Strozzi, T., Wegmüller, U., Wiczorek, M., and Wiesmann, A.: ESA Permafrost Climate Change Initiative (Permafrost_cci): Permafrost extent for the Northern Hemisphere, v4.0, <https://doi.org/10.5285/93444bc1c4364a59869e004bf9bfd94a>, 4 April 2024, 2024.
- 1010 Wittig, S.: Estimating methane sources and sinks in the Arctic using atmospheric data assimilation, Theses, Université Paris-Saclay, <https://theses.hal.science/tel-03997135>, 2023.
- Zhu, X., Zhuang, Q., Qin, Z., Glagolev, M., and Song, L.: Estimating wetland methane emissions from the northern high latitudes from 1990 to 2009 using artificial neural networks, *Global Biogeochemical Cycles*, 27, 592–604, <https://doi.org/10.1002/gbc.20052>, 2013.
- 1015 Zona, D., Oechel, W. C., Kochendorfer, J., Paw U, K. T., Salyuk, A. N., Olivas, P. C., Oberbauer, S. F., and Lipson, D. A.: Methane fluxes during the initiation of a large-scale water table manipulation experiment in the Alaskan Arctic tundra, *Global Biogeochemical Cycles*, 23, <https://doi.org/10.1029/2009GB003487>, 2009.
- Zona, D., Gioli, B., Commane, R., Lindaas, J., Wofsy, S. C., Miller, C. E., Dinardo, S. J., Dengel, S., Sweeney, C., Karion, A., Chang, R. Y.-W., Henderson, J. M., Murphy, P. C., Goodrich, J. P., Moreaux, V., Liljedahl, A., Watts, J. D., Kimball, J. S., Lipson, D. A., and Oechel, W. C.: Cold season emissions dominate the Arctic tundra methane budget, *Proceedings of the National Academy of Sciences*, 113, 40–45, <https://doi.org/10.1073/pnas.1516017113>, 2016.
- 1020