

Predicting and correcting the influence of boundary conditions in regional inverse analyses

Hannah Nesser^{1,2}, Kevin W. Bowman¹, Matthew D. Thill¹, Daniel J. Varon², Cynthia Randles^{3*}, Ashutosh Tewari^{3*}, Felipe J. Cardoso-Saldaña³, Emily Reidy³, Joannes D. Maasackers⁴, Daniel J. Jacob²

¹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

²Harvard University School of Engineering and Applied Sciences, Cambridge, MA, USA

³ExxonMobil Technology and Engineering Company, Annandale, NJ, USA

⁴SRON Space Research Organization Netherlands, Leiden, the Netherlands

*Now at Scepter, Inc., San Francisco, CA, USA

*Now at Amazon Supply Chain Optimization Technologies, Seattle, WA, USA

Correspondence to: Hannah Nesser (hannah.o.nesser@jpl.nasa.gov)

Abstract. Regional inverse analyses of atmospheric trace gas observations quantify gridded two-dimensional surface fluxes by fitting the observations to simulated concentrations from a transport model, usually by Bayesian optimization regularized by a gridded prior flux estimate. Regional inversions rely on the specification of background concentrations given by the boundary conditions (BCs) at the edges of the inversion domain, but biases in the BCs propagate to biases in the optimized fluxes. We develop a theoretical framework to explain how errors in the BCs influence the optimized fluxes as a function of the prior and observing system error statistics and of model transport. We derive a preview metric to estimate the BC-induced errors before conducting an inversion to support domain specification and a diagnostic metric to accurately quantify these errors after solving the inversion. We compare two methods to correct BC biases as part of an inversion, either directly by optimizing BC concentrations (boundary method) or indirectly by expanding the domain and correcting grid cell fluxes outside the region of interest (buffer method). We demonstrate that the boundary method is generally more accurate, physically grounded, and computationally tractable.

1 Introduction

Regional inversions of observed atmospheric concentrations of long-lived trace gases quantify surface fluxes by fitting simulated concentrations from a transport model to the observations assuming boundary conditions (BCs) at the edge of the inversion domain. Such analyses can improve knowledge of fluxes and their trends on local to continental scales at high spatiotemporal resolution while avoiding the need to accurately quantify fluxes and concentrations globally (Sargent et al., 2021; Nesser et al., 2024; Byrne et al., 2024). However, BCs are often uncertain and biases in the BCs propagate to the

Deleted: chemical

Deleted: (CTM)

Deleted: s

Deleted: CTM

Deleted: domain

Deleted:

Deleted: trace gas

Deleted: chemical

Deleted: (CTM)

Deleted: background concentrations given by

Deleted: concentrations

inferred fluxes. Here we examine the problem of how BC biases affect regional inversions of greenhouse gas fluxes and develop a framework to predict and correct the influence of these biases.

45 BCs can be provided by coarse-resolution global simulations (Göckede et al., 2010; Wecht et al., 2014), by **statistical or meteorological analysis** of trace gas observations (Lauvaux et al., 2016; Balashov et al., 2020), or by combining simulated and observed concentrations (Sargent et al., 2018; Estrada et al., 2025). The information used to constrain BCs is often spatiotemporally sparse and may be biased. Even small BC biases can **cause** large biases in the inferred fluxes (Göckede et al., 2010; Lauvaux et al., 2012; Karion et al., 2021). BC biases are particularly critical for **regional** inverse analyses of long-
50 lived gases such as carbon dioxide and methane where concentrations and variability can depend significantly on inflow.

Regional inversions generally infer gridded surface fluxes (the state vector) by minimizing a Bayesian cost function that accounts for the error statistics of the observing system (including the observations and **transport model**) and of the prior flux estimate used to regularize the solution. BC biases may be corrected as part of the inversion by optimizing BC concentrations as part of the state vector (Lauvaux et al., 2012; Wecht et al., 2014), by allowing buffer grid cells at the edge
55 of the domain to absorb BC biases (Shen et al., 2021; Varon et al., 2022), or by combining these approaches (Estrada et al., 2025).

We present here a theoretical framework to predict, diagnose, and correct the influence of BC biases on the posterior gridded
60 surface fluxes generated by a regional inversion (Sect. 2). We consider an inert trace gas with no sources or sinks within the inversion domain other than the surface fluxes. We demonstrate this framework with increasingly complex one- and two-dimensional simulation experiments that invert pseudo-observations of a long-lived trace gas generated using known fluxes and BCs (Sects. 3 and 4, respectively). Throughout, we compare different methods to correct BC biases within inversions and **present** metrics that estimate the effect of BC biases.

65 2 Analytical solution for the effect of boundary condition errors

We describe the analytical inverse solution, which we use to derive an exact solution (diagnostic) for the sensitivity of the posterior fluxes to a BC bias (Sect. 2.1). We apply this diagnostic to a simple one-dimensional transport model to determine how these errors depend on **constant** inversion parameters (Sect. 2.2). We then **generalize the results using a two-box model applicable to two-dimensional inversions with variable inversion parameters. We use this two-box model to derive a preview**
70 **metric that predicts the sensitivity of posterior fluxes to BC biases for a given inversion configuration, allowing the user to improve inversion parameters as needed before solving the inversion** (Sect. 2.3).

Deleted: The

Deleted: used to define background concentrations

Deleted: at the domain edge

Deleted: result in

Deleted: within the domain of interest

Deleted: concentrations

Deleted:

Deleted: CTM

Deleted: with potentially unphysical flux corrections

Deleted: Buffer grid cells increase the inversion's domain and, therefore, its computational cost...

Deleted: demonstrate

Deleted: Quantifying

Deleted: the

Deleted: derive a simplified version of

Deleted: the

Deleted: diagnostic (

Deleted:)

Deleted: to

Deleted: the

Deleted: before conducting the inversion

Deleted: to support domain specification

2.1 Diagnostic equation

95 Given an n -dimensional state vector of gridded fluxes \mathbf{x} and an m -dimensional vector of observations \mathbf{y} , both with normally distributed errors, the optimal flux estimate \mathbf{x} is obtained by minimizing a Bayesian cost function

$$J(\mathbf{x}) = (\mathbf{x} - \mathbf{x}_A)^T \mathbf{S}_A^{-1} (\mathbf{x} - \mathbf{x}_A) + (\mathbf{y} - F(\mathbf{x}))^T \mathbf{S}_0^{-1} (\mathbf{y} - F(\mathbf{x})), \quad (1)$$

100 where \mathbf{x}_A and \mathbf{S}_A are the prior flux estimate and error covariance matrix, respectively, \mathbf{S}_0 is the observing system error covariance matrix representing uncertainties in the observations and the transport model, and F is the transport model (Brasseur and Jacob, 2017). We assume as is standard that the transport model initial conditions are given by a sufficiently long spin-up simulation driven by the BCs so that the initial conditions are consistent with the BCs. If the transport model is linear, then $F(\mathbf{x}) = \mathbf{K}\mathbf{x} + \mathbf{c}$ where $\mathbf{K} = \partial F / \partial \mathbf{x}$ is the Jacobian matrix. The vector \mathbf{c} represents the model background defined by the transport of the BCs to the same spatiotemporal locations as the observations. If the BC concentrations are optimized as part of the inversion, information about the background is instead contained in the state vector and in the columns of the Jacobian matrix so that $\mathbf{c} = \mathbf{0}$.

We can write the analytical solution for the cost function minimum that yields the optimal (posterior) flux estimate \mathbf{x} as

$$\mathbf{x} = \mathbf{x}_A + \mathbf{G}(\mathbf{y} - (\mathbf{K}\mathbf{x}_A + \mathbf{c})) \quad (2)$$

where

$$\mathbf{G} = \frac{\partial \mathbf{x}}{\partial \mathbf{y}} = (\mathbf{K}^T \mathbf{S}_0^{-1} \mathbf{K} + \mathbf{S}_A^{-1})^{-1} \mathbf{K}^T \mathbf{S}_0^{-1} \quad (3)$$

is the gain matrix that represents the sensitivity of the posterior fluxes to the observations. We can separate the model-observation difference into the contributions from the errors in the prior fluxes relative to the true fluxes (\mathbf{x}_T) and the errors in the observing system ($\boldsymbol{\varepsilon} = \mathbf{y} - (\mathbf{K}\mathbf{x}_T + \mathbf{c})$) so that

$$\mathbf{x} = \mathbf{x}_A + \mathbf{A}(\mathbf{x}_T - \mathbf{x}_A) + \mathbf{G}\boldsymbol{\varepsilon} \quad (4)$$

125 where $\mathbf{A} = \partial \mathbf{x} / \partial \mathbf{x} = \mathbf{G}\mathbf{K}$ is the averaging kernel matrix, a measure of inversion information content that gives the sensitivity of the posterior fluxes to the true fluxes. The trace of the averaging kernel matrix gives the degrees of freedom for signal (DOFS), the number of independent pieces of information quantified by the inversion (Rodgers, 2000).

Deleted: CTM

Deleted: CTM

Deleted: unbiased compared to

Deleted: CTM


130

The posterior error induced by a BC error (ϵ_C) is derived by comparing the posterior fluxes produced with the true BC (c_T) to an inversion with the BC error ($c = c_T + \epsilon_C$):

$$\Delta x = -G\epsilon_C. \quad (5)$$

135

We define Eq. (5) as the diagnostic that estimates the effect of BC bias on the posterior fluxes given an explicitly constructed Jacobian matrix and assumed BC error statistics, which can be estimated from observed variability of background concentrations. BC-induced posterior errors are controlled by the gain matrix, which is a function of the observing system errors, prior flux errors, and transport as represented by the Jacobian matrix.



140

Figure 1: One-dimensional and two-box models of a passive trace gas to quantify the influence of BCs on inverse analyses. The one-dimensional model (a) simulates the concentrations of an inert trace gas over n grid cells of length L given a prescribed BC c , fluxes $x = [x_1, x_2, \dots, x_n]^T$, and advection with wind speed U . The two-box model (c) generalizes an inverse analysis optimizing fluxes over a two-dimensional grid (b) where all grid cells have a representative grid cell length L given the wind direction. Each grid cell in the domain is reduced to a two-box model composed of a cluster of the j upwind grid cells (index j_0) and the grid cell itself (index j). In all models, the inversion is solved with the averaged observation y_i over each grid cell $i = \{1, \dots, n\}$ with corresponding error standard deviation $\sigma_{0,i}$. The inversion is regularized by the prior flux $x_{A,i}$ in each grid cell with corresponding prior error standard deviation $\sigma_{A,i}$.

145

Deleted: BC

Deleted: assumed BC error statistics and

Deleted: .

Deleted: (Heald et al., 2004)

Deleted: ¶

We compare the posterior fluxes generated by an inversion with the true BC and no correction method to the posterior fluxes generated by an inversion with a perturbed BC and an error correction method. The increase in the state vector dimension resulting from the error correction method results in the addition of rows to the gain matrix (Eq. (3)). We define the reduced gain matrix including only the rows corresponding to the state vector elements within the domain of interest as G' , which has the same dimension as the gain matrix for the inversion with no correction method (G). The matrix and vector elements associated with the non-domain state vector elements (BC concentrations or buffer grid cells) are denoted with a positive superscript (*). The difference between the posterior fluxes is then:¶

We compare the posterior fluxes generated by an inversion with the true BC and no correction method to the posterior fluxes generated by an inversion with a perturbed BC and an error correction method. The increase in the state vector dimension resulting from the error correction method results in the addition of rows to the gain matrix (Eq. (3)). We define the reduced gain matrix including only the rows corresponding to the state vector elements within the domain of interest as G' , which has the same dimension as the gain matrix for the inversion with no correction method (G). The matrix and vector elements associated with the non-domain state vector elements (BC concentrations or buffer grid cells) are denoted with a positive superscript (*). The difference between the posterior fluxes is then:¶

$$\Delta x = (A' - A)(x_T -$$

where $A' = G'K$ and $A^{*+} = G^*K^*$ are the averaging kernel matrices for the corrected inversion associated with the domain and non-domain state vector elements, respectively. The first two terms of Eq. (6) represent the change in information content resulting from the correction method while the third term represents the change in the influence of the BC bias. The true - prior flux difference ($x_T - x_A$) is unknown but can be approximated by the prior flux standard deviations.¶

Deleted: upstream

2.2 One-dimensional model

195 To better understand how observing system errors, prior flux errors, and transport determine the influence of BC biases on
 200 posterior flux estimates (Eq. (5)), we consider a one-dimensional horizontal transport model with constant wind speed U .
 Figure 1 (panel a) depicts this model for n grid cells of length L . The Jacobian matrix for this transport model is derived
 through steady-state mass balance (Supplement S1) to be lower diagonal with constant values given by


$$\tau = \alpha \tau' \quad (6)$$

205 where α converts between the flux units and the observation units and τ' is the grid cell residence time calculated as $\tau' =$
 L/U . The trace gas sources in the model are the BC \mathbf{c} and fluxes $\mathbf{x} = [x_1, x_2, \dots, x_n]^T$. We assume that the m observations
 are uniformly distributed in space and time with constant uncorrelated error variance σ_0^2 . We solve the inversion (Eq. (2))
 using the averaged observations in each grid cell so that the observing system error covariance matrix \mathbf{S}_0 is diagonal with
 210 constant error variances σ_0^2/m_g where $m_g = m/n$ is the number of observations in each grid cell. We assume constant prior
 fluxes x_A and diagonal prior error covariance matrix \mathbf{S}_A with constant error variances σ_A^2 .

We define the dimensionless, domain-average information ratio R of the prior error variances in concentration units to the
 observing system error variances

$$R = m_g \left(\frac{\tau \sigma_A}{\sigma_0} \right)^2. \quad (7)$$


215 The information ratio increases with decreasing observing system error standard deviation, increasing prior error standard
 deviation, and increasing residence time so that the observations are more sensitive to the fluxes relative to the BC. Large
 values of the information ratio ($R \gg 1$) represent the case where the prior errors are larger than the observing system errors
 so that the posterior fluxes are strongly constrained by the observations (observation-rich). Small values ($R \ll 1$) correspond
 to inversions that are limited by the number or uncertainty of the observations (observation-limited), including the common
 case of spatially heterogeneous observations. Consider an illustrative inversion of a methane-like trace gas with $\tau = 1.4$ h
 (corresponding to $U = 5$ m s⁻¹ and $L = 25$ km), $\sigma_A = 12.5$ ppb h⁻¹ (corresponding to 50% uncertainty on relatively large
 220 prior emissions of 25 ppb h⁻¹), and $\sigma_0 = 10$ ppb. In this case, an average of $m_g = 200$ observations per grid cell are needed
 to achieve $R = 1$. The information ratio for an inversion can be increased by increasing the inversion duration to include
 more observations or by coarsening the resolution of the gridded fluxes optimized by the inversion.



225 **Figure 2:** BC-induced error on posterior surface fluxes as a function of the information ratio R (Eq. (7)) for a one-dimensional
 atmospheric trace gas inversion over ten grid cells with constant parameters. The inversions assume $\tau = 1.4$ h corresponding to
 $U = 5 \text{ m s}^{-1}$ and $L = 25$. The solid lines (b) represent the error induced by a 10 ppb BC bias relative to the prior emissions of 25
 ppb h^{-1} in each grid cell (thin lines) and the total error integrated across the full domain (thick black line). The dashed line
 230 corresponds to an information ratio of 1, while the dotted lines show representative values for which the limiting cases described in
 Eq. (8) hold. The distribution of the BC-induced errors across the domain for these representative values is also shown (a).

The effect of a constant BC bias ε_c on the posterior fluxes is calculated as a function of the information ratio with the
 diagnostic (Eq. (5)). We derive (Supplement S2) the BC-induced error in the limiting cases of inversions that are
 observation-limited ($R \ll 1$) or observation-rich ($R \gg 1$):

$$\Delta x_j = \begin{cases} -(\tau^{-1}\varepsilon_c)(n-j+1)R, & R \ll 1 \\ -(\tau^{-1}\varepsilon_c)R^{-j+1}, & R \gg 1 \end{cases} \quad j = \{1, \dots, n\}. \quad (8)$$



Deleted:

Deleted:

Deleted: 8

Deleted: small and large

Deleted: 9

Figure 2 shows the BC-induced error in these limiting cases (panel a) and for intermediate values of the information ratio ~~calculated using Eq. (5)~~ (panel b) for an illustrative inversion with $n = 10$, $\epsilon_c = 10$ ppb, and $\tau = 1.4$ h corresponding to $U = 5$ m s⁻¹ and $L = 25$ km. ~~In all cases, BC-induced errors decrease across the domain with the rate of decay bounded by the limiting cases in Eq. (8).~~ In the observation-limited case, the BC-induced errors decrease linearly with the distance from the upwind boundary. The total BC-induced error is relatively small because of the strong constraint from the prior estimate. As the information ratio increases from $R \ll 1$, BC-induced errors increase because of the decreased prior constraint. The errors increase the most in the most upwind grid cells. As the information ratio approaches one, the total error converges to the total flux needed to explain the error in the BC ($\tau^{-1}\epsilon_c$). Further increases in the observational constraint decrease the length scale over which the BC bias influences the posterior solution but not the total BC-induced error. In the observation-rich case, the BC-induced errors decay geometrically as the distance from the upwind boundary increases so that the BC-induced errors are limited to the most upwind grid cell.

We also ~~consider the effect of inverse methods to decrease BC biases~~ within the constant parameter, one-dimensional inverse model. We compare the effect of optimizing the BC concentrations (boundary method) or fluxes outside of the domain of interest (buffer method). The boundary method optimizes one or more terms corresponding to BC concentrations along with the domain fluxes (Lauvaux et al., 2012; Wecht et al., 2014; Hancock et al., 2025). The buffer ~~method~~ expands the inversion domain to optimize fluxes in outlying grid cells, which are allowed to vary unphysically to absorb BC biases and are ~~excluded from the final analysis~~ (Shen et al., 2021; Varon et al., 2022). Both approaches increase the state vector dimension, thereby altering the Jacobian matrix and prior error covariance matrix. The buffer ~~method~~ also applies large prior error standard deviations to the buffer grid cells, often by aggregating the buffer grid cells into large clusters.

We ~~derive the conditions under which the boundary and buffer methods are equivalent for~~ an inversion of a single grid cell. For the boundary method, we use a BC prior error standard deviation equal to the constant BC bias ϵ_c and update the Jacobian matrix to include the sensitivity of the grid cell average observation to the BC. For the buffer ~~method~~, we add a buffer grid cell and scale its prior error standard deviation by a factor of p . We assume observations are equally distributed over both grid cells. We solve both inversions. The posterior fluxes are equivalent when

$$p = \frac{\epsilon_c}{\tau\sigma_A\sqrt{R+2}}. \quad (9)$$

Assuming steady state and knowledge of the domain wind speed, the buffer scale factor p can be chosen so that ~~buffer and boundary methods~~ perform ~~equivalently~~. The scale factor goes to infinity for very small values of the information ratio (observation limited) and to zero for very large values (observation rich). The dependence of the scale factor on residence time implies that the buffer ~~method~~ is more vulnerable to varying wind speeds than the boundary method.

Deleted: (

Deleted: the

Deleted: from the prior estimate

Deleted: integrated over the domain

Deleted: compare

Deleted: the boundary and buffer methods for correcting BC biases...

Deleted: the framework of

Deleted: Equation (5) does not account for inverse methods used to correct for BC biases. ...

Deleted: approach

Deleted: therefore

Deleted: approach

Deleted: to minimize the increase in the state vector dimension

Deleted: consider for this purpose

Deleted: approach

Deleted: .

Deleted: s

Deleted: to the boundary method

Deleted: approach

Deleted: †

2.3 Preview for boundary condition errors

Equation (5) can estimate the effect of BC-induced errors on the posterior solution once model transport is characterized by the construction of the Jacobian matrix. We estimate BC-induced errors to inform the choice of inversion domain before building the Jacobian matrix by applying Eq. (5) to a simple steady-state two-box model that generalizes a gridded flux inversion. Figure 1 (panels b and c) depicts the model. The domain has BC c and constant wind speed U . All grid cells have representative grid cell length L given the wind direction. We consider a grid cell (with index j) that is j grid cells from the domain boundary and define the upwind grid cell (with index j_0) as the aggregate of the upwind j grid cells. We conduct an inversion of the average observations over these two grid cells $\mathbf{y} = [y_{j_0} \ y_j]^T$ regularized by prior fluxes $\mathbf{x}_A = [x_{A,j_0} \ x_{A,j}]^T$. We define the prior error covariance matrix as

$$\mathbf{S}_A = \begin{bmatrix} \sigma_{A,j_0}^2 & 0 \\ 0 & \sigma_{A,j}^2 \end{bmatrix} \quad (10)$$

and use an equivalent structure for the observing system error covariance matrix. We assume the observing system error standard deviations account for the reduction in error resulting from averaging the observations. The Jacobian matrix is derived by steady-state mass balance (Supplement S1) to be a function of τ (Eq. (6)) given by

$$\mathbf{K} = \tau \begin{bmatrix} j & 0 \\ j & 1 \end{bmatrix}. \quad (11)$$

We solve for the effect of a constant BC bias $\boldsymbol{\varepsilon}_c = [\varepsilon_c \ \varepsilon_c]^T$ on the posterior flux in the grid cell of interest Δx_j by applying the diagnostic (Eq. (5)) to compute $\Delta \mathbf{x} = [\Delta x_{j-1} \ \Delta x_j]^T$. This defines the preview

$$\Delta x_j = - \frac{(\tau^{-1} \varepsilon_c) R_j}{1 + R_j + \beta R_{j_0} + R_j R_{j_0}} \quad (12)$$

where R_{j_0} and R_j refer to the information ratio (Eq. (7)) in the upwind and selected grid cell, respectively, with $\tau_{j_0} = j\tau$, and

$$\beta = \frac{m_{g,j}}{m_{g,j_0}} + 1. \quad (13)$$

Here, m_{g,j_0} and $m_{g,j}$ refer to the number of observations in the upwind and selected grid cell, respectively, so that β represents the relative observation density across the grid cells. For grid cells abutting the boundary, all upwind values are set to 0 so that the BC-induced error approaches $\tau^{-1} \varepsilon_c$ when the grid cell information ratio is large and 0 when the grid cell

Deleted: 7

Deleted: β

Deleted: 3

Deleted: W

Deleted: and R_{j_0} and R_j refer to the information ratio (Eq. (8)) in the upwind and selected grid cell, respectively, with $\tau_{j_0} = j\tau$.

Deleted:

information ratio is small. As distance from the boundary increases, the BC-induced errors on average decrease due to the accumulation of upwind fluxes and observations, which increases the upwind information ratio R_{j0} . Variability in the prior fluxes and observation density as quantified by the information ratio R_j is super-imposed on this decay.

Deleted: so that the BC-induced error approaches $\tau^{-1}\epsilon_C$ so that $\Delta x_j = -\frac{(\tau^{-1}\epsilon_C)R_j(1+R_j)^{-1}}{(\tau^{-1}\epsilon_C)R_j(1+R_j)^{-1}}$ corresponding to an inversion solved only for the grid cell of interest excluding any upwind components...

By analogy to our one-dimensional model, we consider the observation-limited and observation-rich limiting cases assuming constant prior fluxes, prior flux standard deviations, observation density, and observing system errors. In this case, $R_{j0} = j^4 R_j$ due to the dependence on the number of grid cells j of the upstream observation count ($m_{g,j0} = jm_g$), grid cell residence time ($\tau_{j0} = j\tau$), and prior flux uncertainty (assuming the uncertainties are uncorrelated, $\sigma_{A,j0} = \sigma_A/\sqrt{j}$). The resulting BC-induced errors

Deleted: $\sigma_{A,j0}^2$

Deleted: assuming the observing system errors are uncorrelated

$$\Delta x_j = \begin{cases} -(\tau^{-1}\epsilon_C)R, & R \ll 1 \\ -(\tau^{-1}\epsilon_C)j^{-4}R^{-1}, & R \gg 1 \end{cases} \quad (14)$$

match the general form of the one-dimensional approximation (Eq. (8)), but with the geometric decay in the observation-controlled limiting case approximated as quartic decay.

Deleted: Δc

Deleted: Δc

Deleted: 5

Deleted: 5

Deleted: 9

Deleted: quintic

Deleted: , which results from the dependence of the upstream prior error standard deviation, observing system error standard deviation, and residence time on the grid cell index j

3 One-dimensional numerical solution

We demonstrate the preview, diagnostic, and correction methods using illustrative, numerical inversions of a one-dimensional model for the horizontal transport of an inert trace gas (Sect. 2.2; Nesser, 2025). Table S1 summarizes the model parameters and Table 1 the inversion parameters, which we select to simulate realistic inversions. We use these demonstration inversions to illustrate the effect of constant BC biases in inversions with constant and varying wind speeds (Sect. 3.1). We then consider periodic BC biases (Sect. 3.2). We finally generalize the results by varying the parameter choice to understand the variables controlling BC-induced errors (Sect. 3.3).

Deleted:

Deleted: and varying

Deleted: then

Deleted: the behavior of the

Deleted: 2

Deleted: and variable

3.1 Constant boundary condition perturbations

We quantify the sensitivity of posterior fluxes to BC biases by perturbing the true BC in our one-dimensional model, solving the inversion, and comparing the posterior fluxes to those generated by an inversion solved with the true BC (Nesser, 2025). The inversions apply no other sources of bias (including transport errors) to isolate the effect of BC biases. The domain-average information ratio $R = 0.2$ for these demonstration inversions reflects the low DOFS achieved (5 for inversions with constant wind speeds and 7 for varying wind speeds) despite the uniformity and large number of observations ($m = 1000$), consistent with real inversions (Varon et al., 2023).

Formatted: Justified, Line spacing: 1.5 lines

Deleted: 1

Table 1: Inversion parameters for simulation experiments

Inversion parameter	One-dimensional model	Two-dimensional model
Domain state vector dimension	20	285
Prior emission estimate	Randomly sampled from normal distribution with mean 25 ppb d ⁻¹ and standard deviation 5 ppb d ⁻¹	Modified Express Extension of the Gridded EPA inventory for 2020 ^(f)
Prior error standard deviation ^(a)	50%	50%
Prior boundary condition estimate	0 ppb ^(b)	Smoothed TROPOMI fields ^(g)
Boundary condition error standard deviation	15 ppb	10 ppb
Number of buffer elements	1 ^(c)	Clusters: 10 No clusters: 440 ^(h)
Buffer scale factor values ^(d)	Constant winds: 33 Varying winds: 47 ^(e)	1000
Observing system error standard deviation ^(a)	10 ppb	15 ppb ⁽ⁱ⁾

^(a) The error covariance matrices are assumed diagonal.

^(b) We optimize an incremental correction to the BC.

^(c) We optimize one buffer element that is a cluster of three native-resolution grid cells.

^(d) The buffer scale factor is the factor p applied to the prior error standard deviation in the buffer grid cell.

^(e) We calculate the buffer scale factor with Eq. (9).

^(f) We increase the Express Extension of the Gridded EPA inventory for 2020 (Maasakkers et al., 2023) by a factor of three so that the prior error statistics are consistent with the true emissions.

^(g) We use the IMI default boundary conditions, which are given by monthly mean TROPOMI observations smoothed over ≈ 1000 km.

^(h) We take as the inversion buffer zone the five concentric rings of $0.25^\circ \times 0.3125^\circ$ grid cells around the outer edge of the domain. We solve inversions using 10 buffer cluster elements and 440 native resolution buffer grid cells.

⁽ⁱ⁾ The IMI assumes a constant error standard deviation of 15 ppb for all observations aggregated into errors for the averaged observations accounting for error correlations following Chen et al. (2023). We use the errors generated by the IMI directly.




Figure 3: Exact and predicted influence of a constant BC bias on the posterior fluxes generated by a one-dimensional inverse model using different BC correction methods. The lines represent the posterior error induced by a 10 ppb BC perturbation as a function of the number of grid cells from the upwind boundary using constant (a) and varying (b) wind speeds. The error is given as the difference between inversions solved with the true and biased BCs, normalized by the prior fluxes ($\Delta x/x_A$). The error is shown for inversions solved with no BC correction method, the boundary method, and the buffer method. The buffer grid cell is not shown. The panels also show the preview. Only the first 13 of 20 grid cells are shown; all errors approach zero after this point. The varying wind speed is a see-sawing wind with mean equal to the constant wind speed.

Deleted: ^d

Deleted: Boundary condition mean

Deleted: ^e

Formatted: Superscript

Deleted: 0

Deleted: grid cells

Deleted: 1

Formatted: Superscript

Deleted: ^f

Deleted: ^b

Deleted: 4.8

Deleted: 6.8

Deleted: ^c

Deleted: ^g

Formatted: Not Superscript/ Subscript

Deleted: ^b

Deleted: ^c


Deleted: 10

Deleted: ^d

Deleted: ^e

Deleted: ^f

Deleted: ^g



Deleted:

Deleted: the BC perturbation and

Deleted: (

Deleted: Δc)

Deleted: mechanism

430 Figure 3 (panel a) shows the relative difference in the posterior fluxes, normalized by the prior fluxes, induced by a constant 10 ppb BC perturbation for an inversion with constant wind speeds. As expected from our theoretical analysis, the resulting error on average decreases as the distance from the upwind boundary increases. Exceptions to the decreasing trend result from normalizing by the prior fluxes. The preview (Eq. (12)) accurately predicts the error as expected from the specification of the wind speed. The diagnostic (Eq. (5); not shown) perfectly quantifies the no correction method error as expected given the specification of the BC bias.

440 Figure 3 (panel a) also shows the effect of the buffer and boundary methods, which are described in Table 1. The buffer method optimizes a buffer cluster of three native resolution grid cells. Increasing the buffer grid cell size decreases the magnitude of BC-induced errors but increases the computational cost. The prior uncertainties of the buffer element are increased by a factor of $10p$ where p is the scale factor for which the boundary and buffer methods are theoretically equivalent (Eq. (9)). Increasing p improves performance, but the error reduction quickly asymptotes so that a factor of 10 is sufficient to maximize error reduction in all tested cases. The boundary method optimizes an average correction to the BC.

445 Both the boundary and buffer methods reduce the error induced by the constant BC perturbation. The buffer method achieves larger uncertainty reductions than the boundary method as expected due to the choice of scale factor on the prior uncertainty. The uncorrected errors for the boundary method result from the decreased information available to correct the upwind fluxes, while the residual errors for the buffer method result from uncorrected BC biases. In all cases, the residual error magnitude, as measured by the root mean square error (RMSE) of the posterior fluxes compared to the inversion solved with the true BCs, is constant as a function of BC perturbation. This implies that the correction methods may increase posterior errors for grid cells close to the boundary in inversions with small BC biases. Excluding these grid cells from final analysis will minimize the effect of BC biases on the inversion.

455 The consistent performance of the metrics and of the correction methods results in part from the use of constant wind speeds, which is aligned with the assumptions used to derive the preview and the equivalence between the correction approaches. Figure 3 (panel b) shows the effect of varying wind speeds. The BC-induced bias consists of decaying upwind biases and structured downwind biases. The preview (Eq. (12)) accurately captures the average error decay over the domain, though it is unable to predict the structure of the downwind biases. The diagnostic (Eq. (5); not shown) perfectly predicts both the upwind and downwind biases due to its representation of transport and perfect knowledge of the BC bias. Both correction methods decrease the upwind biases, but only the boundary method decreases the downwind biases. Moreover, the RMSE of the posterior fluxes compared to the inversion solved with the true BCs is constant as a function of perturbation magnitude for the boundary method while the RMSE increases with perturbation magnitude for the buffer method. The buffer method is less able to decrease the variability of the BC-induced biases due to its dependence on wind speed (Eq. (9)).

Deleted: Because the posterior fluxes and metrics respond approximately linearly to constant BC biases (Eqs. (5), (10), and (14)), we show results normalized for the perturbation. We also normalize by the prior fluxes, which are unaffected by BC biases. ...

Deleted: ,

Deleted: which is calculated using the mean wind speed,

Deleted: captures

Deleted: decay

Deleted: from Eq. (5)

Deleted: †

Deleted: .

Deleted:

Deleted: Indeed, the methods perform almost identically as predicted by the steady state approximation (Eq. (10)). ...

Deleted: (first two terms of Eq. (6))

Deleted: (third term of Eq. (6))

Deleted:

Deleted: The boundary method corrects BC concentrations to within 1 ppb of the true BC, while the buffer method changes the buffer grid cell emissions unphysically by a factor of -6.6 ...

Deleted: Importantly, BC-induced errors decay over a similar number of grid cells regardless of buffer size so that multiple rows of buffer grid cells are preferable to large clusters. Combining the buffer and boundary methods produces results identical to the buffer approach....

Deleted: cy

Deleted: upwind decay but not

Deleted: while t

Deleted: and downwind

Deleted: The boundary method strongly reduces both the magnitude and the variability of the BC-induced errors while t...

Deleted: s

Deleted: the magnitude but not

Deleted: 10




Figure 4: Exact and predicted influence of a periodic BC bias on the posterior fluxes generated by a one-dimensional inverse model using different BC correction methods. Panel a shows the true and periodic BC. Panels b and c show the posterior error induced by the periodic BC as a function of the number of grid cells from the upwind boundary error is given as the difference between inversions solved with the true and biased BCs, normalized by the prior fluxes ($\Delta x/x_A$) for constant and varying wind speeds, respectively. The error is shown for inversions solved with no BC correction method, the buffer method, the boundary method, and the combination method. The buffer and boundary methods each optimize one additional element. The error is also shown for a version of the combination method that optimizes three instead of one boundary elements. The buffer grid cell is not shown. The panels also show the preview and diagnostic, which are calculated using the mean BC bias over the inversion period. The varying wind speed is a see-sawing wind with mean equal to the constant wind speed.

3.2 Varying boundary condition perturbations

Varying BC biases are unresolvable within inversions without prior knowledge of their structure. We consider variable BC biases to identify the types of biases that are most important to avoid, to understand the impact of correction methods, and to demonstrate the performance of the preview and diagnostic when the structure of the BC bias is not known. Because varying biases can be represented as the sum of periodic functions, we represent them with periodic BCs with varying y-intercept, amplitude, and period number (Supplement S3). The correction methods are implemented as in the constant perturbation case (Table 1 and Sect. 3.1) but with three BC elements optimized instead of one in the boundary method. This marginally increases the computational cost of the inversion but improves performance.

As in the inversion with a constant BC error and varying wind speeds, the periodic BC-induced bias consists of decaying upwind biases and structured downwind biases (Figure S1). The upwind biases are driven by the mean bias of the periodic perturbation over the inversion period. The spatial structure of the downwind biases is driven by the frequency of the BC error and the magnitude by the amplitude. This suggests that the correction methods will improve the upwind errors but not the downwind errors and that decreasing the amplitude of BC errors may reduce biases that are otherwise unresolvable.




Figure 5: Sensitivity of the BC-induced posterior error to changes in parameters in a one-dimensional inverse model. The vertical axis shows the influence length scale defined as the number of grid cells before which the prior-normalized posterior errors resulting from a 10 ppb BC bias decrease below 0.25 ($\Delta x/x_A < 0.25$). The lines show the influence length scale for the preview and for inversions solved with no BC correction method, the boundary method, and the buffer method. Shading gives the one standard deviation range for 100 inversions solved with different random prior fluxes. Panel (a) shows the influence length scale as a function of the information ratio R (Eq. (7)) that describes the ratio of observing system error variances to the prior error variances in concentration units. The information ratio $R = 1$ is marked by a vertical dashed line. Panel (b) shows the influence length scale as a function of grid cell residence time (τ).

Figure 4 shows the relative difference in the posterior fluxes, normalized by the prior fluxes, induced by a demonstration periodic perturbation with mean bias of 4 ppb over the inversion period (panel a and Supplement S3) for inversions with constant and varying wind speeds (panels b and c, respectively). The upwind biases are controlled by the mean bias with the error on average decreasing with the distance from the upwind boundary. The preview and diagnostic (Eqs. (13) and (5), respectively), which are calculated using the mean bias, capture these biases. The diagnostic also captures the downwind biases attributable to transport. Similarly, both correction methods decrease the upwind bias. The boundary method achieves modest error reductions because the BC-induced error is of similar magnitude to the error generated by the loss of information content due to the BC optimization. Downwind biases are only decreased by the boundary method when three or more BC elements are optimized.

3.3 Varying inversion parameters

BC-induced posterior biases are a function of the relative constraint from the observations compared to the prior fluxes as measured by the information ratio (R , Eq. (7)) and the grid cell residence time (τ , Eq. (6)). To determine how changes in these quantities influence BC-induced biases, we define the influence length scale for our one-dimensional model as the number of grid cells required for the BC-induced error normalized by the prior fluxes to decrease below 0.25 ($\Delta x/x_A < 0.25$). The influence length scale captures the strong dependence of the BC-induced errors on the number of grid cells from the domain boundary and reflects the additional computational cost associated with optimizing these grid cells. We vary the mean prior flux (from 20 to 50 ppb h^{-1}) and associated prior error standard deviations, observation density (from 50 to 200 observations per grid cell), and mean wind speed (from 3 to 10 $m s^{-1}$) and calculate the domain average information ratio and

Deleted: 7

Deleted: 1

Varying BC biases are unresolvable within inversions without prior knowledge of their structure. As a result, we consider the types of BC biases that are most important to avoid in an inversion. Because varying biases can be represented as the sum of periodic functions, we represent them with a series of periodic BCs with varying y-intercept, amplitude, and period number (Supplement S3). ...

Deleted: 1

Deleted: 2

Deleted: 8

Deleted: 7

Deleted: .

residence time for each inversion. Uncertainty is quantified as the standard deviation of the influence length scale across 100
565 inversions solved with random prior fluxes. All inversions use varying wind speeds and a 10 ppb BC perturbation, though
the results are robust for constant wind speeds.

Figure 5 (panel a) shows the influence length scale as a function of the information ratio for an inversion with no correction
570 method. As predicted from the one-dimensional model, solved with constant parameters (Eq. (8) and Fig. 2), the influence
length scale decreases with the information ratio. The largest rate of change occurs in the observation-limited regime as the
observational constraint shifts the BC-induced errors into the upwind grid cells. In the observation-rich case, the influence
length scale is relatively constant as a function of the information ratio. Figure 5 (panel b) also shows the inverse dependence
of the influence length scale on grid cell residence time as predicted by Eq. (8). As residence time increases, the influence
575 length scale decreases because the relative contribution of the fluxes compared to the BC increases in the observed
concentrations, improving the observational constraint.

Finally, we consider the performance of the metrics and correction methods. The preview (Eq. (12)) on average captures the
dependence of the BC-induced error (no correction method) on the information ratio and grid cell residence time. In almost
580 all cases, it marginally overestimates the influence length scale so that it represents an upper bound on error. As expected
from the prescription of the BC perturbation, the diagnostic (not shown) perfectly predicts the influence length scale
associated with no correction method. Both correction methods display functionally no dependence on the information ratio
and grid cell residence time. The boundary and buffer methods comparably reduce the influence length scale in all cases.

4 Two-dimensional numerical solution

We extend the framework and understanding derived from the analytical solution and one-dimensional model to a two-
585 dimensional demonstration inversion of TROPOMI-like methane column pseudo-observations over the Permian basin using
the GEOS-Chem transport model as implemented by the Integrated Methane Inversion (IMI; Varon et al., 2022; Estrada et
al., 2025; Nesser, 2025). The Permian Basin is the largest oil producing region in North America and the subject of many
inverse analyses (e.g., Zhang et al., 2020; Barkley et al., 2023; Varon et al., 2023; Vanselow et al., 2024). Table S1
summarizes the model parameters, Table 1 gives the inversion parameters, and Fig. 6 shows the inversion domain, true
590 emissions (panel a), prior emissions (panel b), and observation density as defined by real TROPOMI observations for May
2020 (panel c) for the $n = 285$ grid cells within the inversion domain. The true BC is given by the IMI's TROPOMI-based
smoothed BCs (Estrada et al., 2025). Assuming a constant wind speed of 5 m s^{-1} and using average values of the prior and
observing system error standard deviations, the domain average information ratio is $R = 0.03$, reflecting the heterogeneous
observational constraint.

- Deleted: 4
- Deleted: and two-box
- Deleted: s
- Deleted: 9
- Deleted: , Eq. (15),
- Deleted: 4
- Deleted: s
- Deleted: 9
- Deleted: and (15)
- Deleted: and metrics
- Deleted: generates a stable estimate of the influence length scale that underestimates the true influence length scale by about 50%...
- Deleted: , which is still adequate
- Deleted: to inform the choice of inversion domain
- Deleted: The preview influence length scale is also relatively insensitive to variations in the information ratio and residence time because it uses variable grid cell information ratios and residence times instead of domain averages. ...
- Deleted: given by th
- Deleted: e
- Deleted: error
- Deleted: outperforms the buffer cases
- Deleted: , reducing the influence length scale to one or fewer grid cells...
- Deleted: The buffer method reduces the influence length scale compared to the error associated with no correction mechanism in the observation-limited case but not the observation-rich case. ... As in the inversion with a constant BC perturbation and varying wind speeds, the BC-induced bias consists of decaying upwind biases and structured downwind biases. The magnitude of both the upwind and downwind errors depends on the mean BC bias and is largely independent of the bias frequency (Fig. S1). (... [1])
- Deleted: As in the inversion with a constant BC perturbatic (... [2])
- Deleted: CTM
- Deleted: (
- Deleted:)
- Deleted: (
- Deleted: 5
- Deleted: ,
- Deleted: which represents a best case scenario for BC variability

We apply a spatially variable BC bias of 7.5 ppb, 10 ppb, 10 ppb, and 12.5 ppb to the northern, southern, eastern, and western boundaries using the Jacobian matrix for each edge of the domain as generated by the IMI. Figure 6 (panel a) shows the resulting error in the posterior emissions normalized by the prior emissions. Unlike the BC-induced errors shown in the one-dimensional inversions, which used random prior fluxes, the distribution of the relative errors is strongly a function of the prior emissions. This likely results from the skewed distribution of the emissions across the domain, which limits the inversion's ability to correct fluxes near the domain edge.




Figure 6: Exact and predicted influence of a constant BC bias on a two-dimensional simulation inversion over the Permian basin in Texas using different BC correction methods. True emissions from the Environmental Defense Fund high-resolution inventory (a; Zhang et al., 2020) and true BCs from smoothed TROPOMI observations are used to generate pseudo-observations over the Permian basin (dashed line) with observational density (c) given by the TROPOMI methane observations for May 2020. The inversion is regularized by prior emissions given by the Express Extension of the Gridded EPA Inventory for 2020 multiplied by a factor of three so that the prior error statistics include the true emissions (b; Maasackers et al., 2023). The error induced by a spatially variable BC bias with mean 10 ppb is calculated as the difference between inversions solved with the true and biased BCs normalized by the prior fluxes and the perturbation to yield the approximate percent error induced per ppb of BC bias. The error is shown for inversions solved with no correction method (e), the boundary method (f), and the buffer method using ten buffer clusters (g) and five rows of native resolution grid cells (h). The dark black outlines show the buffer clusters or grid cells where the prior uncertainties are artificially inflated and for which posterior fluxes are calculated but not shown. The preview (d) is also shown.

Deleted: 5


Deleted: d

Deleted: percent

Deleted: per ppb of mean BC bias

Deleted: error standard deviations

Deleted: This dependence is consistent with the linear dependence of BC-induced errors on the information ratio, which is in turn proportional to the prior error variance, as predicted by the one-dimensional constant parameter inversion for $R < 1$ (Eq. (9)).



Deleted:

Deleted: 5

The preview (Eq. (12)) calculated with a BC uncertainty of 10 ppb is shown in Fig. 6 (panel d). We estimate the Jacobian matrix elements following Eq. (6) as implemented by Nesser et al. (2024) with a constant wind speed of 5 m s^{-1} , constant surface pressure of 1000 hPa, and a grid cell length scale given by the square root of each grid cell's area. We define the upwind length scale as the minimum distance from each grid cell's center to the border, dividing the domain into four regions based on proximity to each boundary. Upwind emissions and observation counts are calculated as the cumulative sum of the median value of each row or column between each grid cell and its closest boundary. Observing system errors are calculated by decreasing the 15 ppb single-observation uncertainty by the square root of the observation count. We apply a 5 ppb observing system error minimum to reflect transport and BC uncertainty.

Deleted: 5

Deleted: e

Deleted: For the preview, w

Deleted: 7

The preview models the decay in BC-induced error as distance from the domain boundary increases accounting for the prior emission distribution and observation density (Eq. (12), Sect. 2.3). As a result, the preview most accurately captures the effect of the BC bias nearest to the domain edge with a Pearson correlation coefficient of $r = 0.76$ between the BC-induced biases and the preview for grid cells within three rows of the boundary. The performance of the preview degrades near the center of the domain, though there is still good agreement between the BC-induced biases and the preview for the full domain ($r = 0.67$). The preview is larger than the BC-induced bias in most (63%) of the grid cells so that the preview represents an error upper-bound. The diagnostic (not shown) perfectly captures the errors across the full domain as expected from the specification of the BC uncertainty.

Deleted: of 15 ppb

Deleted: .

Deleted: combines the

Deleted: with information about

Figure 6 (panels f, g, and h) also shows the correction methods. For the buffer method, we test two sets of buffer grid cells within the five concentric rings of grid cells around the outer edge of the domain. The first (clusters) aggregates the individual grid cells into 10 large buffer clusters using K-means clustering. The second (no clusters) uses 440 individual grid cells as buffers. In all cases, we scale the prior error standard deviation for the buffer elements by a factor of $p = 1000$, which we find is sufficient to minimize BC-induced errors in all tested inversions. This very large, unphysical factor is chosen to compensate for the very small emissions around the edge of the Permian basin. For the boundary method, we optimize a mean bias correction along the northern, southern, eastern, and western boundaries.

Deleted: but is unable to predict the effect on emissions in the domain interior, consistent with the one-dimensional inversion with varying wind speeds. ...

Deleted: 5

Deleted: the

The correction results mirror those found in the one-dimensional varying wind speed example. All correction methods decrease the BC-induced errors. Despite relying on fewer than half of the observations used by the buffer method, the boundary method virtually eliminates BC-induced errors while avoiding unphysical flux corrections and significantly decreasing computational cost by reducing the domain size. The buffer method with no clusters outperforms the buffer method with clusters, suggesting that multiple rows of buffer grid cells are preferable to large clusters. Larger numbers of buffer grid cells may also better absorb biases with higher resolution spatial variability.

5 Conclusions

We developed and demonstrated a framework to predict and correct the effect of BC biases on the optimal (posterior) gridded surface fluxes generated by regional inversions of atmospheric trace gas observations using a ~~transport model~~. We proposed two metrics to predict BC-induced errors both before (preview) and after (diagnostic) the inversion is solved. The preview can inform the choice of inversion domain while the diagnostic improves posterior error quantification. We also considered two methods to correct BC biases as part of an inversion by optimizing the BC directly (boundary method) or by unphysically correcting grid cell fluxes outside the domain of interest (buffer method). Both methods can obtain identical error reductions in inversions with constant wind speeds, but the boundary method is more effective for inversions with variable wind speeds, provides a physical constraint, and reduces computational cost. Beyond the application to regional inversions of long-lived gases presented here, the framework is more generally applicable to the analysis of bias in the observations or ~~transport model~~ and the treatment of initial conditions in global inversions.

We demonstrated our theoretical framework using a simple one-dimensional model for the horizontal transport of an inert trace gas, which represents the worst-case scenario for the propagation of BC biases to the posterior flux estimate. BC-induced errors on average decay with increasing distance from the domain edge while smaller downwind systematic biases result from variability in the ~~transport model~~ wind speeds or BC biases. The length scale over which BC-induced errors have a significant effect on the posterior fluxes is minimized when the observations provide a strong constraint across the domain, which limits the bias to the most ~~upwind~~ grid cells. The preview ~~identifies the grid cells with the largest biases, supporting domain specification before the inversion is conducted~~, while the diagnostic perfectly predicts the BC-induced error when the BC perturbation is specified. The boundary ~~and buffer method both significantly reduce~~ the influence length scale, ~~but only the boundary method can decrease the magnitude of downwind errors~~. The results are robust for constant and variable BC biases.

We extended the framework to a two-dimensional demonstration inversion over the Permian Basin in Texas using model transport from the GEOS-Chem ~~transport model~~. ~~The BC-induced biases do not decay with distance from the boundary but instead correlate with the prior flux estimate due to the concentration of emissions in the center of the domain. Despite the difference in the distribution of BC-induced biases compared to the one-dimensional model, we find similar performance of the metrics and correction methods.~~ The preview accurately predicts the BC-induced errors, ~~with the best agreement occurring in the grid cells~~ closest to the domain boundary. The diagnostic accurately describes the errors across the full domain. ~~The boundary method functionally eliminates the BC-induced errors at much lower computational cost despite relying on fewer than half of the observations used in the buffer method. The buffer method decreases but does not eliminate BC-induced errors, with performance improving as the number of buffer clusters increases.~~

Deleted: chemical

Deleted: (CTM)

Deleted: high-resolution inversions of short-lived gases,

Deleted: CTM

Deleted: CTM

Deleted: upstream

Deleted: generates a stable estimate of the influence length scale

Deleted: significantly

Deleted: s

Deleted: and the magnitude of downwind errors in all cases

Deleted: while

Deleted: the buffer method

Deleted: s

Deleted:

Deleted: influence length scale only for observation-limited inversions...

Deleted: .

Deleted: CTM

Deleted: combines the expected decay in BC-induced errors as distance from the domain boundary increases with information about the prior flux distribution and observation density. As a result, the preview ...

Deleted: T

Code availability

The code and data for both the one-dimensional and two-dimensional simulation experiments are available at <https://doi.org/10.5281/zenodo.17417750> (Nesser, 2025). The IMI v2.0 used to generate the inputs for the two-dimensional example is available at https://github.com/geoschem/integrated_methane_inversion/releases/tag/imi-2.0.1.

Deleted: 15742048

Field Code Changed

Author contributions

HN and DJJ designed the study. HN, DJJ, and KWB contributed to the theoretical development. HN conducted the theoretical analysis and demonstration applications. MDT contributed to the proof of Eq. (8). KWB, DJV, CR, AT, FJCS, ER, and JDM discussed the results. HN, DJJ, and KWB wrote the paper with input from all authors.

Deleted: 9

835 Acknowledgments

HN was funded in part by an appointment to the NASA Postdoctoral Program at the Jet Propulsion Laboratory, California Institute of Technology, administered by Oak Ridge Associated Universities under contract with NASA. The research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D0004). This work was funded in part by ExxonMobil Technology and Engineering Company. We thank Kimberly Mueller for her feedback.

References

Balashov, N. V., Davis, K. J., Miles, N. L., Lauvaux, T., Richardson, S. J., Barkley, Z. R., and Bonin, T. A.: Background heterogeneity and other uncertainties in estimating urban methane flux: results from the Indianapolis Flux Experiment (INFLUX), *Atmospheric Chemistry and Physics*, 20, 4545–4559, <https://doi.org/10.5194/acp-20-4545-2020>, 2020.

845 Barkley, Z., Davis, K., Miles, N., Richardson, S., Deng, A., Hmiel, B., Lyon, D., and Lauvaux, T.: Quantification of oil and gas methane emissions in the Delaware and Marcellus basins using a network of continuous tower-based measurements, *Atmospheric Chemistry and Physics*, 23, 6127–6144, <https://doi.org/10.5194/acp-23-6127-2023>, 2023.

Brasseur, G. and Jacob, D. J.: *Modeling of atmospheric chemistry*, Cambridge university press, Cambridge New York, 2017.

850 Byrne, B., Liu, J., Bowman, K. W., Yin, Y., Yun, J., Ferreira, G. D., Ogle, S. M., Baskaran, L., He, L., Li, X., Xiao, J., and Davis, K. J.: Regional Inversion Shows Promise in Capturing Extreme-Event-Driven CO₂ Flux Anomalies but Is Limited by Atmospheric CO₂ Observational Coverage, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD040006, <https://doi.org/10.1029/2023JD040006>, 2024.

855 Estrada, L. A., Varon, D. J., Sulprizio, M., Nesser, H., Chen, Z., Balasus, N., Hancock, S. E., He, M., East, J. D., Mooring, T. A., Oort Alonso, A., Maasakkers, J. D., Aben, I., Baray, S., Bowman, K. W., Worden, J. R., Cardoso-Saldaña, F. J., Reidy, E., and Jacob, D. J.: Integrated Methane Inversion (IMI) 2.0: an improved research and stakeholder tool for

- monitoring total methane emissions with high resolution worldwide using TROPOMI satellite observations, *Geoscientific Model Development*, 18, 3311–3330, <https://doi.org/10.5194/gmd-18-3311-2025>, 2025.
- 860 Göckede, M., Turner, D. P., Michalak, A. M., Vickers, D., and Law, B. E.: Sensitivity of a subregional scale atmospheric inverse CO₂ modeling framework to boundary conditions, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2010JD014443>, 2010.
- Hancock, S. E., Jacob, D. J., Chen, Z., Nesser, H., Davitt, A., Varon, D. J., Sulprizio, M. P., Balasus, N., Estrada, L. A., Cazoria, M., Dawidowski, L., Diez, S., East, J. D., Penn, E., Randles, C. A., Worden, J., Aben, I., Parker, R. J., and
865 Maasakkers, J. D.: Satellite quantification of methane emissions from South American countries: a high-resolution inversion of TROPOMI and GOSAT observations, *Atmospheric Chemistry and Physics*, 25, 797–817, <https://doi.org/10.5194/acp-25-797-2025>, 2025.
- Karion, A., Lopez-Coto, I., Gourdji, S. M., Mueller, K., Ghosh, S., Callahan, W., Stock, M., DiGangi, E., Prinzevalli, S., and Whetstone, J.: Background conditions for an urban greenhouse gas network in the Washington, DC, and Baltimore
870 metropolitan region, *Atmospheric Chemistry and Physics*, 21, 6257–6273, <https://doi.org/10.5194/acp-21-6257-2021>, 2021.
- Lauvaux, T., Schuh, A. E., Uliasz, M., Richardson, S., Miles, N., Andrews, A. E., Sweeney, C., Diaz, L. I., Martins, D., Shepson, P. B., and Davis, K. J.: Constraining the CO₂ budget of the corn belt: exploring uncertainties from the assumptions in a mesoscale inverse system, *Atmospheric Chemistry and Physics*, 12, 337–354, <https://doi.org/10.5194/acp-12-337-2012>, 2012.
- 875 Lauvaux, T., Miles, N. L., Deng, A., Richardson, S. J., Cambaliza, M. O., Davis, K. J., Gaudet, B., Gurney, K. R., Huang, J., O’Keefe, D., Song, Y., Karion, A., Oda, T., Patarasuk, R., Razlivanov, I., Sarmiento, D., Shepson, P., Sweeney, C., Turnbull, J., and Wu, K.: High-resolution atmospheric inversion of urban CO₂ emissions during the dormant season of the Indianapolis Flux Experiment (INFLUX), *Journal of Geophysical Research: Atmospheres*, 121, 5213–5236, <https://doi.org/10.1002/2015JD024473>, 2016.
- 880 Maasakkers, J. D., McDuffie, E. E., Sulprizio, M. P., Chen, C., Schultz, M., Brunelle, L., Thrush, R., Steller, J., Sherry, C., Jacob, D. J., Jeong, S., Irving, B., and Weitz, M.: A Gridded Inventory of Annual 2012–2018 U.S. Anthropogenic Methane Emissions, *Environ. Sci. Technol.*, 57, 16276–16288, <https://doi.org/10.1021/acs.est.3c05138>, 2023.
- Nesser, H.: Boundary condition sensitivity simulation experiments (v2.0), <https://doi.org/10.5281/zenodo.17417750>, 2025.
- 885 Nesser, H., Jacob, D. J., Maasakkers, J. D., Lorente, A., Chen, Z., Lu, X., Shen, L., Qu, Z., Sulprizio, M. P., Winter, M., Ma, S., Bloom, A. A., Worden, J. R., Stavins, R. N., and Randles, C. A.: High-resolution US methane emissions inferred from an inversion of 2019 TROPOMI satellite data: contributions from individual states, urban areas, and landfills, *Atmospheric Chemistry and Physics*, 24, 5069–5091, <https://doi.org/10.5194/acp-24-5069-2024>, 2024.
- Rodgers, C. D.: *Inverse methods for atmospheric sounding: theory and practice*, World Scientific, Singapore, 2000.
- 890 Sargent, M., Barrera, Y., Nehr Korn, T., Hutyra, L. R., Gately, C. K., Jones, T., McKain, K., Sweeney, C., Hegarty, J., Hardiman, B., Wang, J. A., and Wofsy, S. C.: Anthropogenic and biogenic CO₂ fluxes in the Boston urban region, *Proceedings of the National Academy of Sciences*, 115, 7491–7496, <https://doi.org/10.1073/pnas.1803715115>, 2018.
- Sargent, M. R., Floerchinger, C., McKain, K., Budney, J., Gottlieb, E. W., Hutyra, L. R., Rudek, J., and Wofsy, S. C.: Majority of US urban natural gas emissions unaccounted for in inventories, *Proceedings of the National Academy of Sciences*, 118, e2105804118, <https://doi.org/10.1073/pnas.2105804118>, 2021.

- 895 Shen, L., Zavala-Araiza, D., Gautam, R., Omara, M., Scarpelli, T., Sheng, J., Sulprizio, M. P., Zhuang, J., Zhang, Y., Qu, Z., Lu, X., Hamburg, S. P., and Jacob, D. J.: Unravelling a large methane emission discrepancy in Mexico using satellite observations, *Remote Sensing of Environment*, 260, 112461, <https://doi.org/10.1016/j.rse.2021.112461>, 2021.
- Vanselow, S., Schneising, O., Buchwitz, M., Reuter, M., Bovensmann, H., Boesch, H., and Burrows, J. P.: Automated detection of regions with persistently enhanced methane concentrations using Sentinel-5 Precursor satellite data, *Atmospheric Chemistry and Physics*, 24, 10441–10473, <https://doi.org/10.5194/acp-24-10441-2024>, 2024.
- 900 Varon, D. J., Jacob, D. J., Sulprizio, M., Estrada, L. A., Downs, W. B., Shen, L., Hancock, S. E., Nesser, H., Qu, Z., Penn, E., Chen, Z., Lu, X., Lorente, A., Tewari, A., and Randles, C. A.: Integrated Methane Inversion (IMI 1.0): a user-friendly, cloud-based facility for inferring high-resolution methane emissions from TROPOMI satellite observations, *Geoscientific Model Development*, 15, 5787–5805, <https://doi.org/10.5194/gmd-15-5787-2022>, 2022.
- 905 Varon, D. J., Jacob, D. J., Hmiel, B., Gautam, R., Lyon, D. R., Omara, M., Sulprizio, M., Shen, L., Pendergrass, D., Nesser, H., Qu, Z., Barkley, Z. R., Miles, N. L., Richardson, S. J., Davis, K. J., Pandey, S., Lu, X., Lorente, A., Borsdorff, T., Maasackers, J. D., and Aben, I.: Continuous weekly monitoring of methane emissions from the Permian Basin by inversion of TROPOMI satellite observations, *Atmospheric Chemistry and Physics*, 23, 7503–7520, <https://doi.org/10.5194/acp-23-7503-2023>, 2023.
- 910 Wecht, K. J., Jacob, D. J., Frankenberg, C., Jiang, Z., and Blake, D. R.: Mapping of North American methane emissions with high spatial resolution by inversion of SCIAMACHY satellite data, *Journal of Geophysical Research: Atmospheres*, 119, 7741–7756, <https://doi.org/10.1002/2014JD021551>, 2014.
- Zhang, Y., Gautam, R., Pandey, S., Omara, M., Maasackers, J. D., Sadavarte, P., Lyon, D., Nesser, H., Sulprizio, M. P., Varon, D. J., Zhang, R., Houweling, S., Zavala-Araiza, D., Alvarez, R. A., Lorente, A., Hamburg, S. P., Aben, I., and Jacob, D. J.: Quantifying methane emissions from the largest oil-producing basin in the United States from space, *Sci. Adv.*, 6, eaaz5120, <https://doi.org/10.1126/sciadv.aaz5120>, 2020.
- 915

Page 14: [1] Deleted

Nesser, Hannah O (US 329G)

10/12/25 8:13:00 AM



Page 14: [2] Deleted

Nesser, Hannah O (US 329G)

10/11/25 11:55:00 AM

