

Climate field reconstructions for the North Atlantic region of annual, seasonal and monthly resolution spanning CE 1241-1970

Jesper Sjolte ¹ and Qin Tao ¹

¹Department of Geology – Quaternary Science, Lund University, Sölvegatan 12, 223 62, Lund, Sweden.

Correspondence: Jesper Sjolte (jesper.sjolte@geol.lu.se)

Abstract. The North Atlantic region is a key component of the climate system via ~~large-scale~~ large-scale atmosphere and ocean circulation. Climate field reconstructions can provide a long-term context for ongoing climate change and contribute to our understanding of climate dynamics, impact of external forcings, and act as references for model evaluation and baseline for natural variability. There are distinct differences in North Atlantic climate variability between the seasons in terms of climate modes and amplitude of the variance. Constraining long-term climate variability in sub-annual resolution is therefore needed for a more complete understanding of the governing processes. In this study, we present reconstructed climate in annual, seasonal ~~and seasonal~~, and monthly resolution based on a small high-quality network of proxy data combined with output from an isotope enabled climate model. Compared to earlier work, we have improved the methodology to obtain better skill across a larger area and more realistic variance of the reconstructed variables which include 2m temperature (T2m), sea surface temperature (SST), sea level pressure (SLP) and precipitation amount. Here we validate the reconstructions against reanalysis data, observed SST and eight long-term records of observed temperature. The reconstructed temperature correlates with up to 0.71 for seasonal data and 0.68 for annual data compared to reanalysis data, while the correlation is about 0.3 for monthly resolution. The skill for SLP shows the imprint of large-scale circulation for winter with more local pattern dominating for summer. This is also mirrored in the skill for precipitation. In addition, the reconstructed annual mean SST shows basin-wide skill for the North Atlantic, indicating relevance of the reconstruction to studies of atmosphere-ocean interaction. In summary, the results show the potential of assimilating a small high-quality network of proxy records.

1 Introduction

The climate of the past millennium is an important reference period for current and future climate change due to relatively abundant climate proxy data ~~and~~, while also spanning the onset of the industrialisation of society (Jungclauss et al., 2017). Despite the good coverage of proxy data, many challenges remain with respect to temporal resolution and spatial representation of climate reconstructions. Climate reconstructions ~~are often representing~~ often represent annual mean data (e.g., Tardif et al., 2019), or only the summer season (Büntgen et al., 2020), and can be limited to only one climate variable ~~only, e.g., e.g.~~ temperature. This is to a large extent due to the readily available tree-ring data for this period and/or uncertainties in dating and seasonality of other types of proxy data. However, key information ~~of the spatiotemporal about spatio-temporal~~ variability and dynamics is lost when choosing to target the annual mean for reconstructions, for example, due to seasonal differences in

the main climate modes (Sjolte et al., 2020). Several different approaches have been taken to reconstruct past climate modes of the North Atlantic region, most notably for reconstructions of the main mode of ~~winter~~ winter variability, namely the North Atlantic Oscillation (NAO). One approach is to specifically target NAO variability and reconstruct a time series of NAO (Trouet et al., 2009; Ortega et al., 2015; Michel et al., 2020), while others aim to reconstruct the pressure field and then extract the main modes (Luterbacher et al., 2004; Sjolte et al., 2018, 2020). The latter approach has the advantage of potentially separating different climate modes such as NAO and the Eastern Atlantic pattern, and thereby avoiding to assign variability in the proxy data to the reconstructed NAO which might be due to imprints of other climate modes.

Recently, climate field reconstructions of monthly resolution have also been produced (Valler et al., 2021, 2024). For this work instrumental data and historical documents have been included, which can constrain the variability on shorter time-scales than the seasonal resolution possible from climate proxy records. However, due to the lack of ~~the~~ instrumental data and historical documentation in the earlier part of the reconstructions the skill cannot be expected to be consistent from the early to the later part of the reconstructed time frame. This can ~~for example~~, for example, cause data-dependent shifts in the spatial patterns of atmospheric modes (Tao et al., 2023).

The skill of climate reconstructions is invariably linked to the climate signal preserved in proxy data. Quantifiable climate information can be found in archives preserving the variability of the isotopic composition of precipitation. In extratropical regions, the mean isotopic composition of precipitation is strongly correlated with the local temperature (Dansgaard, 1964). The isotopic composition is usually formulated as the relative deviation from the Vienna mean standard ocean water composition using delta notation, e.g. $\delta^{18}\text{O}$ for the relative abundance of ^{18}O in a water sample (Craig, 1961). Despite the linear spatial relation between $\delta^{18}\text{O}$ and temperature, there are large regional differences in the relationship between local climate and temporal variations in $\delta^{18}\text{O}$, which hampers a simple translation from $\delta^{18}\text{O}$ to temperature (e.g., Sjolte et al., 2011).

However, $\delta^{18}\text{O}$ remains one of the most important sources of paleoclimate variability from polar ice cores (Jouzel, 2013). The availability of seasonally resolved $\delta^{18}\text{O}$ ice core data is determined by the accumulation rate and annual layer thickness. The annual cycle of $\delta^{18}\text{O}$ is attenuated due to diffusion in the firn and the annual cycle is lost at low accumulation sites (Johnsen, 1977). While the annual cycle can be partly restored using mathematical back-diffusion (Johnsen et al., 2000), these limitations mean that only relatively few ice core sites can be used for millennium scale seasonal reconstructions.

~~While~~ Although the aforementioned advantages of seasonal reconstructions are clear, the main limitations lie in the availability of well-dated, ~~long-term seasonally resolved~~ seasonally resolved, long-term proxy data. As already mentioned above, tree-ring data ~~ticks~~ tick many of the boxes for ~~high quality~~ high-quality seasonal data. However, several caveats are still present when selecting tree-ring data for reconstructions. Firstly, in ~~term~~ terms of tree-ring width or maximum late wood density (MXD), not all trees are strongly sensitive to temperature (Büntgen et al., 2008). Secondly, ~~sampling~~ sample replication of tree-ring chronologies ~~vary~~ varies over time, typically with decreasing replication of older trees, which increases the uncertainty back in time (Ljungqvist et al., 2020). The recent decades have seen an increase in studies of oxygen isotope data measured on tree ring cellulose (Balting et al., 2021). However, the number of millennium-length isotope chronologies ~~are~~ is still quite limited, and the interpretation of the data is not straightforward due to ~~both~~ hydrological and biophysical influences on the isotope signal (e.g. Seftigen et al., 2011; Balting et al., 2021).

Gridded climate field reconstructions can be analysed in similar ways as climate model data or meteorological reanalysis data, which is very useful for comparing these different datasets. A range of methods can be used to produce climate field reconstructions, with the common feature being that climate model output is resampled for the best fit to climate proxy data, or using spatial statistics to infer climate patterns (see also Smerdon et al. (2023)). Challenges for [doing](#) climate field reconstructions include uneven distribution of proxy data sites. This tends to concentrate both skill and variance in the areas near the proxy data sites, which skews ~~the spatiotemporal~~ [spatio-temporal](#) variability and creates biases [in](#) the representation of the main modes of variability (Sjolte et al., 2020; Tao et al., 2023).

In this study (hereafter, SAT25) we use seasonally resolved climate proxy data and an isotope enabled climate model to produce climate reconstructions of annual, seasonal and monthly resolution for sea level pressure, temperature, and precipitation for summer and winter over the North Atlantic region covering the time period 1241-1970. The climate proxy data ~~includes~~ [include](#) Greenland ice core $\delta^{18}\text{O}$ data, tree ring MXD, blue intensity (BI), as well as tree ring cellulose $\delta^{18}\text{O}_{cell}$. Our study can be seen as an update to the previous reconstructions by Sjolte et al. (2018, 2020) (hereafter ~~;~~ SEA18 and SEA20) including expanded proxy data network and improved methodology. We evaluate the reconstructions against reanalysis data, long-term temperature observations, and compare with other reconstructions.

75

2 Data

2.1 Proxy and climate model data

We select the proxy data with the following criteria. Firstly, the proxy data should have a strong imprint of climate. In case of the MXD, tree-ring width (TRW) and BI data from tree-ring chronologies, this means high correlation to local temperature, while we look for a clear imprint of atmospheric circulation in the $\delta^{18}\text{O}$ -based records. Secondly, the proxy data needs to have 0–1-year dating uncertainty and a well-defined seasonality. To achieve consistent performance of the reconstruction, we only select proxy records that span the whole time frame of the reconstruction (CE 1241-1970) (Figure ~~1 and Table 1~~ [1 and Table 1](#) and [Table 1](#)). [The dating uncertainty of anchored tree-ring chronologies covering the past millennium is generally assumed to be zero. The ice core data are dated using layer counting and synchronised using volcanic reference horizons, and for the past millennium the dating uncertainty is estimated to be maximum 1 year \(Vinther et al., 2006b\). The volcanic dating horizons ensure that an entire ice core record is never shifted by one year. However, one cannot exclude that some records are off by up to one year in some intervals between dating horizons.](#)

We use the isotope enabled version of the climate model ECHAM5/MPIOM (Werner et al., 2016) run for the past 1200 years forced ~~with by~~ natural and anthropogenic forcings. This is the same model simulation used in SEA18 and SEA20.

The atmosphere component is run in a T31L19 configuration corresponding to $3.75^\circ \times 3.75^\circ$ horizontal resolution and 19 vertical layers, while the ocean component, MPIOM GR30L40, is set to 3° horizontal resolution with 40 layers. The model features isotope fractionation during all phase changes, including kinetic fractionation during evaporation and in

mixed-phase clouds.

95 2.2 Interpretation of $\delta^{18}\text{O}_{cell}$

The variability of $\delta^{18}\text{O}_{cell}$ is governed by several different processes, some of which have a high degree of co-variability. From the following analysis, we assume that the main control of $\delta^{18}\text{O}_{cell}$ is the isotopic composition of precipitation forming the groundwater, which comprises the source water for formation of xylem sap used to form cellulose during the growing season. Depending on the hydrological setting at the site of the tree-ring chronology, there can be a lagged isotope signal from precipitation formed prior to the growing season, or from climate driven intra-annual variability. The lag can be due to groundwater recharge during spring ~~melt-of-snow~~ snow melt or sap formed during late winter and spring (Wang et al., 2005). Our tests indicate that both $\delta^{18}\text{O}_{cell}$ records used in this study are sensitive to weather variability (SLP, T2m, precip. amount) during the growing season (Figure S1), as well as the weather variability of the winter preceding the growing season (Figure S2). The records ENGL and CZEC are mainly correlated to variability connected to the winter NAO, which can be explained by the JJA Palmer drought severity index (pdsi) at these sites being sensitive to the DJF NAO. This is due to moisture availability of the growing season being affected by the preceding winter precipitation, as well as land-atmosphere feedback mechanisms where weather patterns promoting summer drought are enhanced by low soil moisture of the previous winter season (Wang et al., 2011). Based on these findings we use the $\delta^{18}\text{O}_{cell}$ records as representing a mixture of summer and winter variability. The impact of large-scale circulation on weather variability and $\delta^{18}\text{O}$ in precipitation weakens in spring, while the maximum correlation between $\delta^{18}\text{O}_{cell}$ (ENG and CZEC, Table 1) and climate variables are seen in the growing season (Loader et al., 2020; Büntgen et al., 2021). We use JJA for representing the growing season and the extended winter season (Nov-Apr) to represent the winter preceding the growing season. Using the extended winter accounts both for impact of the winter circulation patterns, precipitation and composition groundwater for early spring sap formation. In addition, this choice of seasonal dependency the $\delta^{18}\text{O}_{cell}$ is linked to the Greenland ice core data through the extended winter $\delta^{18}\text{O}$, and to the conventional tree-ring data through the co-variability of JJA temperature and $\delta^{18}\text{O}$. While a mechanistic approach to connect the $\delta^{18}\text{O}_{cell}$ to $\delta^{18}\text{O}$ in precipitation is possible through forward modelling (e.g. Roden et al., 2000), this would also introduce new parameters with uncertainties specific to each site. Since the untreated $\delta^{18}\text{O}_{cell}$ in the first place has a clear imprint of atmospheric circulation for summer and winter, we choose to assume that the variability of $\delta^{18}\text{O}_{cell}$ primarily is governed by a mix of $\delta^{18}\text{O}$ in precipitation for both seasons.

120

2.3 Observations, reanalysis and ~~reconstructions~~ reconstructions for validation and comparison

We use a variety of data ranging from instrumental data to climate field reconstructions to evaluate our climate reconstruction. For assessing spatial coverage of skill, we use the 20th Century Reanalysis version 3 (20CRv3) (Slivinski et al., 2021). We compared the 20CRv3 precipitation against the CPC gridded precipitation dataset (Chen et al., 2008) and found them to perform comparably on seasonal timescales. Due to 20CRv3 also covering ocean grid points, we prefer the 20CRv3 for evaluation

125

of the reconstructed precipitation field, despite not being constrained directly by observed precipitation. As a supplement to evaluating the reconstruction with the 20CRv3 temperature we use long-term observed temperature from Greenland (Vinther et al., 2006a), Iceland (Jónsson, 1989), England (Legg et al., 2025), Denmark (Cappelen et al., 2021) and Sweden (Moberg et al., 2002; Bergström and Moberg, 2002), to achieve a longer overlapping time interval for the evaluation. To evaluate reconstructed sea surface temperature (SST) we use three different datasets ERSSTv5 (Huang et al., 2017), COBE2 (Hirahara et al., 2014), HadISST (Rayner et al., 2003) that have different properties due to different treatment of observational data and missing values. In ~~the discussion we go beyond addition to purely~~ observation-based datasets ~~and compare~~, we compare our to the climate field reconstruction ModeE-RA (Valler et al., 2024) of monthly resolution covering 1421-2008.

135 2.4 Methods

We use the analogue method to assimilate proxy data using an isotope enabled climate model. This setup can be described as assimilation with a static ensemble or static model prior. Our reconstruction is not calibrated against observations, and only relies on matching the model output with proxy data. For each site we evaluate the model output against the proxy data, linking simulated JJA temperature to conventional tree-data (MXD/BI/TRW), and simulated seasonal $\delta^{18}\text{O}$ with ice core $\delta^{18}\text{O}$ and $\delta^{18}\text{O}_{\text{cell}}$ (Table 2). We use different evaluation criteria of the model data for the summer and winter season. For the summer season we use a χ^2 -measure for the goodness of fit:

$$\chi_{sum}^2(t) = \sum_{i=1}^n \frac{(m(n, t') - p(n, t))^2}{\sigma_{mp}} \quad (1)$$

Where, m is the modelled anomaly and p is the proxy anomaly at a given site, n is the number of proxies, t is the year of the reconstruction, t' is the year of the model run, and σ_{mp} is the ~~error estimated from the~~ combined standard deviation of the model and proxy data. Dividing by σ_{mp} normalizes the data, which is effectively the same as normalizing all time series (proxy data and model output for proxy sites) prior to the matching procedure.

The imprint of large-scale circulation on the isotopic composition is very strong for winter and the amplitude of the $\delta^{18}\text{O}$ anomalies bears a signal in it-self. Part of this signal is the gradient between proxy sites (Sjolte et al., 2018). Since we are interested in preserving as much of the signal in the proxy data as possible, we do not normalize the data when fitting the model data to the proxy data. The measure for goodness of fit for winter is then:

$$\chi_{win}^2(t) = \sum_{i=1}^n (m(n, t') - p(n, t))^2 \quad (2)$$

Where, m is the modelled anomaly and p is the proxy anomaly at a given site, n is the number of proxies, t is the year of the reconstruction and t' is the year of the model run.

For the annual mean and monthly reconstruction, the proxies for summer and winter are matched simultaneously using Eq. 1. The monthly data from the best matching model years are then extracted, taking into account that the ice core data for winter

are representing months Nov-Apr when the annual mean for the calendar mean is calculated (i.e., Nov-Dec are assigned to previous calendar year compared to the matched winter proxy data centred in January).

In this study we calculated the ensemble mean using a logarithmic weighting function. Due to the bias-variance trade off, a high number of ensemble member reduces noise and increases skill in term of correlation but causes decrease in variability, as well as skewing the variability towards the main mode and causing mixed secondary modes in the SLP field. Before arriving at our current approach, we tested different ways calculate the ensemble mean: constant number of ensemble members with equal weighting (as SEA18/20), variable number of ensemble members based on match with proxy data, also in combination with a minimum number of ensemble members. Here we calculate the weighted mean based on the χ^2 fit (Eq. 1 & 2), which yields a smoother behaviour of reconstruction with less year-to-year difference in the statistical properties of the reconstruction compared to a variable number of ensemble members, while still allowing the best matching models years to have the most weight. The weighting is calculated as the logarithm of the normalized χ^2 -distance, where the least important ensemble member receives zero weighting (Figure S3). The χ^2 -distance as a function of number of ensemble members is similar to an exponential decay function and we apply the logarithm to achieve a more equal weighting of the ensemble members. Otherwise, the result would be skewed too much towards a few of the best fitting ensemble members, which results in more noise in the reconstruction. With the applied weighting 50% of the weight goes to the best matching 30% of the ensemble members.

~~Time-series are shown-~~

In our results we show time series as anomalies from the mean of the full-length data series, and monthly data are anomalies with respect to the mean annual cycle. We use Pearson correlation (r) and significance levels are determined using Student's t-test. For correlation maps we list the maximum correlation, mean significant correlation and number of grid cells with significant correlation below the figure title. The number of grid cells with significant is a rough indication of the spatial coverage of the skill, which will depend on the spatial auto-correlation, which in turn depends on season, latitude (area of grid cell) and type of climate variable.

3 Results

3.1 Ensemble reconstruction and initial evaluation

Before comparing to observed climate we evaluate the coherence between the reconstruction and the assimilated proxy data, and assess how many model analogues per year to use in the ensemble reconstruction. When determining number of ensemble members to include for each year there are several factors to consider. A high number of ensemble members reduces noise by smoothing the variability spatially and temporally, but ~~this also reduces and skews variability to~~ as mentioned above, the smoothing also makes it more difficult to separate the main modes of variability (e.g., NAO) climate variability. For example, with a high number of ensemble members, the East Atlantic and Scandinavian blocking pattern cannot be separated and more of the variability is assigned to the NAO. Conversely, a low number of ensemble members fail to span both the imperfect fit of the model to the proxy data, as well as the confounding noise of the proxy data.

We first consider the match between the reconstructed signal at the sites of the proxy data. There is high correlation between

the reconstruction and all proxy records, with a match to ice core $\delta^{18}\text{O}$ and $\delta^{18}\text{O}_{cell}$ in the range $r \sim 0.65-0.9$, and on average slightly better correlation between reconstructed temperature and conventional tree-ring records, though in similar range ($r \sim 0.6-0.9$). We tested the match with proxy data with increasing number of ensemble members. The maximum correlation to proxy data is achieved with 10-100 ensemble members per year, however the match to the proxy data only degrades slowly with up to 300 ensemble members per year due to the weighting with respect to χ^2 -distance to the proxy data (Figure S4). Using 300 ensemble members introduces spatial smoothing which filters out noise in particularly for areas far from the proxy sites (see Section 3.5). With this high number of ensemble members, we on one hand lose information on modes of atmospheric circulation, as mentioned in the introduction, but on the other hand gain skill for basin-wide variability for the North Atlantic region. We therefore provide two versions of the reconstruction, one using 150 ensemble members and one using 300 ensemble members. For simplicity we focus on the version with 300 ensemble-ensemble members as the differences are subtle and will be explored in future work.

Previous seasonal reconstructions (SEA18/20) only used ice core data, or introduced tree-ring data after a pre-selection of model analogues had been made using ice core data. Here we fully incorporate the tree-ring data. The match with ice core records can be achieved regardless if the tree-ring data is included or not, indicating that the analogue sampling pool is sufficiently large to provide relevant analogues, and that further constraining the climate variability with records in Europe does not hinder a good match over Greenland.

To further test the role of the tree-ring and ice core data, we made two new reconstructions based on our annual reconstruction. One based only on ice core data (RECON_{IC}) and once based only on tree-ring data (RECON_{TR}). We then extracted the data for tree-ring sites from RECON_{IC} and the ice core sites from RECON_{TR}. The results show that the prediction of tree-ring variability from RECON_{IC} has weak skill, although some correlations are significant (see Figure S5-8). The best correspondence is seen for Scandinavian tree-ring records (TAA, EFmean) and winter $\delta^{18}\text{O}$ correlated to CZEC $\delta^{18}\text{O}_{cell}$. Conversely, the prediction of ice core variability from RECON_{TR} also has weak skill (Figure S9 & S10), with the best correlation to GRIP $\delta^{18}\text{O}$ ($r = 0.33$) (Figure S9). This underlines the different properties of the proxy data due to differences in seasonality, region and what is recorded by the proxies. In broad terms, the main strength of the ice core data is to record large-scale atmospheric circulation for winter, while the tree-ring data capture temperature changes on a more local scale, but the wider availability of tree-ring data ensures a reasonable spatial coverage to capture a large-scale summer signal. Our main conclusion from this test is that the properties of the different sets of proxy data are complementary and constrain different seasons and regions of the reconstruction.

The uneven distribution of proxy data concentrates skill and variance near the sites of the proxy data, and calculating the ensemble mean dampens the overall variance. We bias correct the variance of the reconstruction by rescaling the standard deviation of each grid point with the corresponding standard deviation of the ECHAM5-wiso/MPIOM run. Our motivation is that the climate model acting as the model prior is the best estimate of the variance in locations not constrained by proxy data. This very simple correction improves the match between reconstructed variance compared to the variance for SLP, T2m and precipitation from 20CRv3 (Figures S5 & S6S11 & S12).

To test the long-term stability of the reconstruction we investigated temporal variations of the match with proxy data. The

average fit of model analogues is fairly constant over time, indicating that the quality of the climate reconstruction is consistent throughout 1241-1970. One reason for this consistency is the strict selection of proxy records that span the whole reconstruction, which means that ~~that~~ there are no methodological differences throughout the reconstruction. We found no clear preference with respect to selection of model analogues throughout the reconstruction. This is ~~eval~~-consistent with results for earlier versions of the reconstruction (SEA18/20).

230 3.2 Comparison to meteorological reanalysis

SEA25 is not calibrated to observations (see Section 2.4). The comparison to observed climate in this section is therefore an independent measure of skill. In addition, since we use a constant number of proxy data and no change in methodology throughout the reconstruction, we should, in theory, reach the same level of skill for any time period of the reconstruction.

Compared to the JJA and DJF mean for 20CRv3, the reconstruction shows coherent patterns of skill for SLP and T2m, while the skill for precipitation is patchier (Figure 22). Standout features are widespread significant correlation for the winter SLP and strong correlation over the eastern North Atlantic and Northern Europe for summer T2m (max. $r = 0.71$). In winter the level of correlation is tightly connected with the skill for reconstructed circulation patterns. This is especially clear for the winter SLP and T2m which resemble the spatial pattern of NAO-type variability, while summer SLP and T2m which resemble the spatial pattern of Scandinavian blocking-type variability. The weaker skill for summer circulation is due to a combination of less well constrained large-scale circulation during summer and biases in the JJA SLP patterns of the ECHAM5/MPIOM model (Sjolte et al., 2020).

The correlation patterns for the annual mean are, not surprisingly, a combination of features seen for summer and winter (Figure 3-3 a-c). While it is less pronounced than for DJF, the signature of NAO-type variability is still evident in the spatial patterns of the correlation. The maximum correlation ($r = 0.68$) is achieved for T2m with lower correlations for SLP and precipitation than for winter, indicating that auto-correlation in temperature aids higher skill in some areas, while the seasonal information is lost in the annual mean SLP and precipitation due to shifting circulation patterns and different impact of circulation on T2m and precipitation on a seasonal scale.

On monthly time scales the skill of the ~~reconstructing~~-reconstruction is lower with at maximum correlation of 0.34 for T2m (average for all months) (Figure 3-3 d-f). Despite lower skill, the correlation is still significant over large areas, due to the larger sampling pool (12 times as more sample points compared to annual data). The spatial patterns of the correlation are similar to that of the annual mean reconstruction. Reviewing the skill for individual months, correlations peak in February and August for T2m with maximum correlation of 0.55 and 0.70, respectively (Figure S7S13 & S14). For SLP the imprint of large-scale circulation is again evident during winter months (max. corr. = 0.38 for February) with lower and more localized skill during summer ~~:(not shown)~~. The skill for the monthly reconstruction varies regionally due to the seasonality of the proxy data and seasonal shifts in climate patterns, leading to summer months and winter months not having high skill at the same location. For example, this is part of the reason for the annual average monthly temperature correlations to remain low, although they peak

above 0.6 for temperature during July and August (S13).

3.3 Comparison to long-term temperature records

260 In addition to the comparison with 20CRv3 we compare to observed temperature from Greenland, Iceland, Central England, Denmark, and Sweden which are among the longest observed high-quality temperature records and span our study area well. For summer the high skill of the reconstruction seen in Fig. 2b is confirmed for sites in Sweden and England (Figure 44). It appears that there are periods of over and underestimated reconstructed temperature which could be a sign of underestimated multi-decadal variability, or it could be due to a different long-term trend in the reconstruction. However, for Uppsala, which is
265 close by Stockholm and has a longer observational record, the reconstruction shows the highest correlation to observed summer temperature ($r = 0.56$). It is also reaffirming that the reconstruction shows good correlation to the longest existing instrumental record, the Central England temperature ($r = 0.5$). The lower skill for Greenland sites during summer is similar to earlier studies, because Greenland ice core summer $\delta^{18}\text{O}$ is more sensitive to climate variability east of Greenland, also illustrated by good correlation to Iceland summer temperature ($r = 0.54$) (Vinther et al., 2010; Sjolte et al., 2020).
270 The highest skill during winter is seen for Greenland winter temperatures ($r \sim 0.6$), although we also see good performance for Stockholm ($r = 0.46$), despite having no proxy site nearby (Figure 55). Compared to summer there are only the two $\delta^{18}\text{O}_{\text{cell}}$ proxy sites available for the winter season besides the Greenland ice cores. However, this is partly counteracted by the stronger impact of large-scale circulation on regional climate variability enabling good skill also in the Eastern part of the study area. We observe that the variance for DJF is approximately double of the JJA variance. This is captured well for most sites by the
275 reconstruction, also showing that the variance correction is effective also compared to instrumental data. For annual mean data the correlation is more consistently in the range of 0.5-0.6 (Table 23), again, combining the characteristics of the summer and winter season, as well as capturing amplitude of the temperature variance (Figure S8S15). The monthly reconstruction has correlations of ~ 0.3 at all sites but for Central England which has lower correlation ($r = 0.21$). The lower skill for Central England is likely due to the winter variability being less coupled to the NAO. As the reconstruction is not constrained at monthly resolution, we cannot expect high skill on a month-to-month basis. What we do see is the
280 reconstruction capturing part of the modulation of the annual cycle seen in the observations. For example, for the cold winters during the 16th and 17th century in Scandinavia (Figure S9S16).

3.4 Comparison to the ModE-RA reanalysis

285 There are only few reconstructions of monthly resolution spanning more than a few centuries. In extension of the validation of the reconstruction against observation-based datasets, we have also compared our reconstruction to the ModE-RA reanalysis reaching back to 1421. In addition to proxy data, ModE-RA is assimilated based on instrumental data and historical documents. Comparing our reconstruction and ModE-RA temperature for the full overlapping time period 1421-1970, we see a fair correlation for the summer months, showing large areas of coherent correlation over the North Atlantic (Figure 7). In contrast,

290 the winter months show little in common between the two reconstructions. For SLP the coherence between ModE-RA and our reconstruction is generally low (not shown). Looking at different time periods it is clear that we find a better correspondence with the ModE-RA reanalysis our reconstruction towards the later part of the reconstruction (e.g., 20th century), where ModE-RA is incorporating instrumental data (not shown). This is especially true for the winter season, for both temperature and SLP.

295 3.5 Annual mean SST evaluation

The annual reconstruction, which targets the calendar year, enables direct comparison to more common annual reconstructions and is useful for analysing indices of sea surface temperature which are often defined on annual mean data. Our reconstruction uses a high number of ensemble members (300, $\sim 25\%$ of the available model analogues) introducing spatial smoothing to filter out noise in particular for areas far from the proxy sites. Using this high number of ensemble members, we on one hand
300 lose information on modes of atmospheric circulation, as mentioned in the introduction, but on the other hand gain skill for basin-wide variability for the North Atlantic region (Figure 66).

4 Discussion and conclusions

Compared to the SEA18/20 reconstructions, our new reconstruction shows several clear improvements in the performance for summer and winter. This includes more wide-spread skill for all reconstructed variables and more realistic variance due
305 to a simple, but effective, variance correction. With the introduction of the variance correction, the performance of the reconstruction is significantly improved in comparison with the 20CRv3 and observed temperature records. This indicates that the reconstruction will lend itself well to studies of forcing attribution, climate extremes and model evaluation given the more realistic amplitude of changes featured in the reconstruction. Furthermore, we included reconstructions in annual and monthly resolution, ~~which~~. The calendar year is a widely used format in other reconstructions, and we include this to make
310 comparisons to existing datasets more straight forward. Similarly, our monthly reconstruction can be useful in-comparison to other data-sets when compared or used with other data if a specific seasonality is targeted, which then can be a combination of any months. Finally, we also included reconstructed precipitation as an output variable.

~~There are only few reconstructions of monthly resolution spanning more than a few centuries.~~ In the validation of our reconstruction, we see a consistent match to the proxy data, as well as the long-term observed temperature. ~~In-extension of this we have also~~ There are only few reconstructions of monthly resolution spanning more than a few centuries and we have
315 compared our reconstruction to the recent ModE-RA reanalysis reaching back to 1421. In-addition to proxy data The comparison shows weak correlation between the two reconstructions, with lower correlation in the earlier part of the overlapping time frame, in particularly for the winter season. This might be due to fewer data constraining the ModE-RA is assimilated based on instrumental data and historical documents. Comparing our reconstruction and ModE-RA temperature for the full overlapping
320 time period 1421-1970 we see a fair correlation for the summer months, showing large areas of coherent correlation over the North Atlantic (Figure 7). In-contrast, the winter months show little in common between the two reconstructions. For SLP

the coherence between ModE-RA and our reconstruction is generally low (not shown). Looking at different time periods it is clear that we find a better correspondence with the ModE-RA reanalysis our reconstruction towards the later reanalysis in the earlier part of the reconstruction (e.g., 20th century), where ModE-RA is incorporating instrumental data (not shown). This is especially true for the winter season, for both temperature and SLP.

325 Reconstructing sub-annual variability remains challenging. This is to a large extent due to the lack of available proxy data for constraining the winter season. However, as shown in SEA18/20₂, even a low number of high-quality proxy records is sufficient to capture large-scale winter circulation if the method is optimized for this purpose. The winter variability of SEA25 is well constrained by the ice core $\delta^{18}\text{O}$ and $\delta^{18}\text{O}_{cell}$ throughout the reconstructed time frame.

330 The variability of the monthly data can also depend reconstruction also depends on how the model analogues are sampled. As explained in the Methods section, the model data for the monthly reconstruction is sampled based on the annual reconstruction. This means that the model output for summer and winter proxies are evaluated simultaneously is evaluated against the proxy data simultaneously, and we select the best matching model analogues for the whole year. We tested extracting monthly data based on the selection of model years from the seasonal reconstruction. ~~This made~~, where model output for summer and
335 winter is evaluated against the proxy data separately, and a different model analogue can be selected for representing summer and winter months, respectively. Comparing the two ways of selecting model analogues we see some slight differences where the SLP for winter months was slightly better based on the seasonal selection of model analogues, while the temperature generally was slightly better based on the annual selection (Figure [S7S14](#)). However, if indeed targeting seasonal means we do recommend using the JJA and DJF reconstructions.

340 Our method is based on the selection of model analogues that best match the proxy data. As we use a static ensemble, any year of the simulation is allowed to be matched to any year of the proxy data, which disrupts the continuity of the model simulation. The proxy data we have selected for SEA25 are sensitive to atmospheric variability, and the sampled model output and the reconstruction will attain the variance of atmospheric variability. For the SST reconstruction the analogue sampling means that the variance can be overestimated on at year-to-year basis, and the any long-term information on ocean currents from the model
345 run are not maintained in the resampled model output of the reconstruction. The overestimated SST variance is to a certain extend counteracted by the variance correction based on the model variance.

One of the novel aspects of this work is the inclusion of $\delta^{18}\text{O}_{cell}$ data from tree-ring chronologies. As summarized in the Introduction and Data sections, the signal recorded in $\delta^{18}\text{O}_{cell}$ is the result of complex biophysical processes and environmental variability. We choose not to attempt to include such complexity, and instead rely on empirical statistical relationships
350 between the $\delta^{18}\text{O}_{cell}$ signal and climate variables derived from observations. In our tests of the reconstruction the inclusion of the $\delta^{18}\text{O}_{cell}$ data is important for the performance of the reconstruction, for both summer and winter. It provides, so to speak, the European end of the NAO see-saw which then also constrains the variability of other weather patterns. While this shows that the $\delta^{18}\text{O}_{cell}$ data does contain important climate information even if it is based on simple assumptions, it does not exclude that more information could be obtained using forward modelling of the biophysical and hydrological processes.

355 The relatively strict selection of proxy data for this work means that we have a low number of proxy sites. New proxy data can be included in future versions of the reconstruction as it becomes available. We see a need for new millennium-length

$\delta^{18}\text{O}_{cell}$ records from Europe to constrain the winter variability further. In addition, using a simulation with an updated isotope enabled model with better spatial resolution for the data assimilation could lead to generally improved performance of the reconstruction.

360 *Data availability.* The SEA25 reconstructions are archived at Zenodo (10.5281/zenodo.15746008).

Author contributions. JS initiated and designed the research project. Method development, investigation and evaluation of reconstructions were done in collaboration by JS and QT. JS wrote the first draft of the manuscript which was revised after comments by QT.

Competing interests. The authors declare no competing interests.

Acknowledgements. This work was supported by the Swedish Energy Agency (Project No. 51375-1) and the strategic research program of
365 Modelling the Regional and Global Earth system (MERGE) hosted by the Faculty of Science at Lund University.

References

- Balting, D. F., Ionita, M., Wegmann, M., Helle, G., Schleser, G. H., Rimbu, N., Freund, M. B., Heinrich, I., Caldarescu, D., and Lohmann, G.: Large-scale climate signals of a European oxygen isotope network from tree rings, *Climate of the Past*, 17, 1005–1023, <https://doi.org/10.5194/cp-17-1005-2021>, publisher: Copernicus GmbH, 2021.
- 370 Bergström, H. and Moberg, A.: Daily Air Temperature and Pressure Series for Uppsala (1722–1998), *Climatic Change*, 53, 213–252, <https://doi.org/10.1023/A:1014983229213>, 2002.
- Büntgen, U., Frank, D., Wilson, R., Carrer, M., Urbinati, C., and Esper, J.: Testing for tree-ring divergence in the European Alps, *Global Change Biology*, 14, 2443–2453, <https://doi.org/10.1111/j.1365-2486.2008.01640.x>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2486.2008.01640.x>, 2008.
- 375 Büntgen, U., Arseneault, D., Boucher, E., Churakova (Sidorova), O. V., Gennaretti, F., Crivellaro, A., Hughes, M. K., Kirilyanov, A. V., Klippel, L., Krusic, P. J., Linderholm, H. W., Ljungqvist, F. C., Ludescher, J., McCormick, M., Myglan, V. S., Nicolussi, K., Pierrat, A., Oppenheimer, C., Reinig, F., Sigl, M., Vaganov, E. A., and Esper, J.: Prominent role of volcanism in Common Era climate variability and human history, *Dendrochronologia*, 64, 125–157, <https://doi.org/10.1016/j.dendro.2020.125757>, 2020.
- Büntgen, U., Urban, O., Krusic, P. J., Rybníček, M., Kolář, T., Kyncl, T., Ač, A., Koňasová, E., Čáslavský, J., Esper, J., Wagner, S.,
380 Saurer, M., Tegel, W., Dobrovolný, P., Cherubini, P., Reinig, F., and Trnka, M.: Recent European drought extremes beyond Common Era background variability, *Nat. Geosci.*, 14, 190–196, <https://doi.org/10.1038/s41561-021-00698-0>, bandiera_abtest: a Cg_type: Nature Research Journals Number: 4 Primary_atype: Research Publisher: Nature Publishing Group Subject_term: Atmospheric science;Climate change;Hydrology;Palaeoclimate Subject_term_id: atmospheric-science;climate-change;hydrology;palaeoclimate, 2021.
- Cappelen, J., Kern-Hansen, C., Laursen, E., Joergensen, P., and Joergensen, B.: Denmark – DMI Historical Climate Data Collection 1768–
385 2020, Tech. Rep. 21-02, DMI, 2021.
- Chen, M., Shi, W., Xie, P., Silva, V. B. S., Kousky, V. E., Wayne Higgins, R., and Janowiak, J. E.: Assessing objective techniques for gauge-based analyses of global daily precipitation, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009132>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2007JD009132>, 2008.
- Craig, H.: Isotopic Variations in Meteoric Waters, *Science*, 133, 1702–1703, <https://doi.org/10.1126/science.133.3465.1702>, publisher:
390 American Association for the Advancement of Science, 1961.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16, 436–468, <https://doi.org/10.1111/j.2153-3490.1964.tb00181.x>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.2153-3490.1964.tb00181.x>, 1964.
- Hirahara, S., Ishii, M., and Fukuda, Y.: Centennial-Scale Sea Surface Temperature Analysis and Its Uncertainty, *Journal of Climate*, 27, 57–75, <https://doi.org/10.1175/JCLI-D-12-00837.1>, place: Boston MA, USA Publisher: American Meteorological Society, 2014.
- 395 Huang, B., Thorne, P. W., Banzon, V. F., Boyer, T., Chepurin, G., Lawrimore, J. H., Menne, M. J., Smith, T. M., Vose, R. S., and Zhang, H.-M.: Extended Reconstructed Sea Surface Temperature, Version 5 (ERSSTv5): Upgrades, Validations, and Intercomparisons, *Journal of Climate*, 30, 8179–8205, <https://doi.org/10.1175/JCLI-D-16-0836.1>, place: Boston MA, USA Publisher: American Meteorological Society, 2017.
- Johnsen, S. J.: Stable isotope homogenization of polar firn and ice, *I.A.H.S.-A.I.S.H. PUBL.; U.S.A.; DA. 1977; NO 118; PP. 210-219; BIBL. 12 REF.; (ISOT. IMPURETES NEIGES GLACES. ACTES COLLOQ.; GRENOBLE; 1975), 118, 210–219, 1977.*
400

- Johnsen, S. J., Clausen, H. B., Cuffey, K. M., Hoffmann, G., Schwander, J., and Creyts, T.: Diffusion of stable isotopes in polar firn and ice : the isotope effect in firn diffusion, in: Physics of Ice Core Records, pp. 121–140, Hokkaido University Press, <http://hdl.handle.net/2115/32465>, 2000.
- Jónsson, T.: The observations of Jon Thorsteinsson in Nes and Reykjavik 1820–1854, Tech. rep., Icel. Met. Office Report, 1989.
- 405 Jouzel, J.: A brief history of ice core science over the last 50 yr, *Climate of the Past*, 9, 2525–2547, <https://doi.org/10.5194/cp-9-2525-2013>, publisher: Copernicus GmbH, 2013.
- Jungclauss, J. H., Bard, E., Baroni, M., Braconnot, P., Cao, J., Chini, L. P., Egorova, T., Evans, M., González-Rouco, J. F., Goosse, H., Hurrett, G. C., Joos, F., Kaplan, J. O., Khodri, M., Klein Goldewijk, K., Krivova, N., LeGrande, A. N., Lorenz, S. J., Luterbacher, J., Man, W., Maycock, A. C., Meinshausen, M., Moberg, A., Muscheler, R., Nehrbass-Ahles, C., Otto-Bliesner, B. I., Phipps, S. J., Pongratz, J., 410 Rozanov, E., Schmidt, G. A., Schmidt, H., Schmutz, W., Schurer, A., Shapiro, A. I., Sigl, M., Smerdon, J. E., Solanki, S. K., Timmreck, C., Toohey, M., Usoskin, I. G., Wagner, S., Wu, C.-J., Yeo, K. L., Zanchettin, D., Zhang, Q., and Zorita, E.: The PMIP4 contribution to CMIP6 – Part 3: The last millennium, scientific objective, and experimental design for the PMIP4 *past1000* simulations, *Geoscientific Model Development*, 10, 4005–4033, <https://doi.org/10.5194/gmd-10-4005-2017>, publisher: Copernicus GmbH, 2017.
- Legg, T., Packman, S., Caton Harrison, T., and McCarthy, M.: An Update to the Central England Temperature Series—HadCET v2.1, 415 *Geoscience Data Journal*, 12, e284, <https://doi.org/https://doi.org/10.1002/gdj3.284>, e284 GDJ-2024-07-0048.R1, 2025.
- Ljungqvist, F. C., Piermattei, A., Seim, A., Krusic, P. J., Büntgen, U., He, M., Kirilyanov, A. V., Luterbacher, J., Schneider, L., Seftigen, K., Stahle, D. W., Villalba, R., Yang, B., and Esper, J.: Ranking of tree-ring based hydroclimate reconstructions of the past millennium, *Quaternary Science Reviews*, 230, 106074, <https://doi.org/10.1016/j.quascirev.2019.106074>, 2020.
- Loader, N. J., Young, G. H. F., McCarroll, D., Davies, D., Miles, D., and Ramsey, C. B.: Summer precipitation for the England 420 and Wales region, 1201–2000 ce, from stable oxygen isotopes in oak tree rings, *Journal of Quaternary Science*, 35, 731–736, <https://doi.org/10.1002/jqs.3226>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/jqs.3226>, 2020.
- Luterbacher, J., Dietrich, D., Xoplaki, E., Grosjean, M., and Wanner, H.: European Seasonal and Annual Temperature Variability, Trends, and Extremes Since 1500, *Science*, 303, 1499–1503, <https://doi.org/10.1126/science.1093877>, publisher: American Association for the Advancement of Science, 2004.
- 425 Michel, S., Swingedouw, D., Chavent, M., Ortega, P., Mignot, J., and Khodri, M.: Reconstructing climatic modes of variability from proxy records using ClimIndRec version 1.0, *Geoscientific Model Development*, 13, 841–858, <https://doi.org/10.5194/gmd-13-841-2020>, publisher: Copernicus GmbH, 2020.
- Moberg, A., Bergström, H., Ruiz Krigsman, J., and Svanered, O.: Daily Air Temperature and Pressure Series for Stockholm (1756–1998), *Climatic Change*, 53, 171–212, <https://doi.org/10.1023/A:1014966724670>, 2002.
- 430 Ortega, P., Lehner, F., Swingedouw, D., Masson-Delmotte, V., Raible, C. C., Casado, M., and Yiou, P.: A model-tested North Atlantic Oscillation reconstruction for the past millennium, *Nature*, 523, 71–74, <https://doi.org/10.1038/nature14518>, 2015.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/https://doi.org/10.1029/2002JD002670>, eprint: 435 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2002JD002670>, 2003.
- Roden, J. S., Lin, G., and Ehleringer, J. R.: A mechanistic model for interpretation of hydrogen and oxygen isotope ratios in tree-ring cellulose, *Geochimica et Cosmochimica Acta*, 64, 21–35, [https://doi.org/10.1016/S0016-7037\(99\)00195-7](https://doi.org/10.1016/S0016-7037(99)00195-7), 2000.

- Seftigen, K., Linderholm, H. W., Loader, N. J., Liu, Y., and Young, G. H. F.: The influence of climate on $^{13}\text{C}/^{12}\text{C}$ and $^{18}\text{O}/^{16}\text{O}$ ratios in tree ring cellulose of *Pinus sylvestris* L. growing in the central Scandinavian Mountains, *Chemical Geology*, 286, 84–93, 440 <https://doi.org/10.1016/j.chemgeo.2011.04.006>, 2011.
- Sjolte, J., Hoffmann, G., Johnsen, S. J., Vinther, B. M., Masson-Delmotte, V., and Sturm, C.: Modeling the water isotopes in Greenland precipitation 1959–2001 with the meso-scale model REMO-iso, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2010JD015287>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2010JD015287>, 2011.
- Sjolte, J., Sturm, C., Adolphi, F., Vinther, B. M., Werner, M., Lohmann, G., and Muscheler, R.: Solar and volcanic forcing of North Atlantic 445 climate inferred from a process-based reconstruction, *Climate of the Past*, 14, 1179–1194, <https://doi.org/10.5194/cp-14-1179-2018>, publisher: Copernicus GmbH, 2018.
- Sjolte, J., Adolphi, F., Vinther, B. M., Muscheler, R., Sturm, C., Werner, M., and Lohmann, G.: Seasonal reconstructions coupling ice core data and an isotope-enabled climate model – methodological implications of seasonality, climate modes and selection of proxy data, *Climate of the Past*, 16, 1737–1758, <https://doi.org/10.5194/cp-16-1737-2020>, publisher: Copernicus GmbH, 2020.
- 450 Slivinski, L. C., Compo, G. P., Sardeshmukh, P. D., Whitaker, J. S., McColl, C., Allan, R. J., Brohan, P., Yin, X., Smith, C. A., Spencer, L. J., Vose, R. S., Rohrer, M., Conroy, R. P., Schuster, D. C., Kennedy, J. J., Ashcroft, L., Brönnimann, S., Brunet, M., Camuffo, D., Cornes, R., Cram, T. A., Domínguez-Castro, F., Freeman, J. E., Gergis, J., Hawkins, E., Jones, P. D., Kubota, H., Lee, T. C., Lorrey, A. M., Luterbacher, J., Mock, C. J., Przybylak, R. K., Pudmenzky, C., Slonosky, V. C., Tinz, B., Trewin, B., Wang, X. L., Wilkinson, C., Wood, K., and Wyszyński, P.: An Evaluation of the Performance of the Twentieth Century Reanalysis Version 3, *Journal of Climate*, 34, 455 1417–1438, <https://doi.org/10.1175/JCLI-D-20-0505.1>, publisher: American Meteorological Society Section: Journal of Climate, 2021.
- Smerdon, J. E., Cook, E. R., and Steiger, N. J.: The Historical Development of Large-Scale Paleoclimate Field Reconstructions Over the Common Era, *Reviews of Geophysics*, 61, e2022RG000782, <https://doi.org/10.1029/2022RG000782>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022RG000782>, 2023.
- Tao, Q., Sjolte, J., and Muscheler, R.: Persistent Model Biases in the Spatial Variability of Winter North Atlantic Atmospheric Circulation, *Geophysical Research Letters*, 50, e2023GL105231, <https://doi.org/10.1029/2023GL105231>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2023GL105231>, 2023.
- Tardif, R., Hakim, G. J., Perkins, W. A., Horlick, K. A., Erb, M. P., Emile-Geay, J., Anderson, D. M., Steig, E. J., and Noone, D.: Last Millennium Reanalysis with an expanded proxy database and seasonal proxy modeling, *Climate of the Past*, 15, 1251–1273, <https://doi.org/10.5194/cp-15-1251-2019>, publisher: Copernicus GmbH, 2019.
- 465 Trouet, V., Esper, J., Graham, N. E., Baker, A., Scourse, J. D., and Frank, D. C.: Persistent Positive North Atlantic Oscillation Mode Dominated the Medieval Climate Anomaly, *Science*, 324, 78–80, <https://doi.org/10.1126/science.1166349>, _eprint: <https://www.science.org/doi/pdf/10.1126/science.1166349>, 2009.
- Valler, V., Franke, J., Brugnara, Y., and Brönnimann, S.: An updated global atmospheric paleo-reanalysis covering the last 400 years, *Geoscience Data Journal*, n/a, <https://doi.org/10.1002/gdj3.121>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/gdj3.121>, 2021.
- 470 Valler, V., Franke, J., Brugnara, Y., Samakinwa, E., Hand, R., Lundstad, E., Burgdorf, A.-M., Lipfert, L., Friedman, A. R., and Brönnimann, S.: ModE-RA: a global monthly paleo-reanalysis of the modern era 1421 to 2008, *Sci Data*, 11, 36, <https://doi.org/10.1038/s41597-023-02733-8>, number: 1 Publisher: Nature Publishing Group, 2024.
- Vinther, B., Jones, P., Briffa, K., Clausen, H., Andersen, K., Dahl-Jensen, D., and Johnsen, S.: Climatic signals in multiple highly resolved stable isotope records from Greenland, *Quaternary Science Reviews*, 29, 522–538, 475 <https://doi.org/https://doi.org/10.1016/j.quascirev.2009.11.002>, 2010.

- Vinther, B. M., Andersen, K. K., Jones, P. D., Briffa, K. R., and Cappelen, J.: Extending Greenland temperature records into the late eighteenth century, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006810>, 2006a.
- Vinther, B. M., Clausen, H. B., Johnsen, S. J., Rasmussen, S. O., Andersen, K. K., Buchardt, S. L., Dahl-Jensen, D., Seierstad, I. K., Siggaard-Andersen, M.-L., Steffensen, J. P., Svensson, A., Olsen, J., and Heinemeier, J.: A synchronized dating of three Greenland ice cores throughout the Holocene, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/https://doi.org/10.1029/2005JD006921>, 2006b.
- 480 Wang, F., Arseneault, D., Boucher, E., Gennaretti, F., Yu, S., and Zhang, T.: Tropical volcanoes synchronize eastern Canada with Northern Hemisphere millennial temperature variability, *Nature Communications*, 13, 5042, <https://doi.org/10.1038/s41467-022-32682-6>, publisher: Nature Publishing Group, 2022.
- 485 Wang, G., Dolman, A. J., and Alessandri, A.: A summer climate regime over Europe modulated by the North Atlantic Oscillation, *Hydrology and Earth System Sciences*, 15, 57–64, <https://doi.org/10.5194/hess-15-57-2011>, publisher: Copernicus GmbH, 2011.
- Wang, K.-Y., Kellomäki, S., Zha, T., and Peltola, H.: Annual and seasonal variation of sap flow and conductance of pine trees grown in elevated carbon dioxide and temperature, *Journal of Experimental Botany*, 56, 155–165, <https://www.jstor.org/stable/24030897>, publisher: Oxford University Press, 2005.
- 490 Werner, M., Haese, B., Xu, X., Zhang, X., Butzin, M., and Lohmann, G.: Glacial–interglacial changes in H₂¹⁸O, HDO and deuterium excess – results from the fully coupled ECHAM5/MPI-OM Earth system model, *Geoscientific Model Development*, 9, 647–670, <https://doi.org/10.5194/gmd-9-647-2016>, publisher: Copernicus GmbH, 2016.
- Wilson, R., Anchukaitis, K., Briffa, K. R., Büntgen, U., Cook, E., D’Arrigo, R., Davi, N., Esper, J., Frank, D., Gunnarson, B., Hegerl, G., Helama, S., Klesse, S., Krusic, P. J., Linderholm, H. W., Myglan, V., Osborn, T. J., Rydval, M., Schneider, L., Schurer, A., Wiles, G.,
495 Zhang, P., and Zorita, E.: Last millennium northern hemisphere summer temperatures from tree rings: Part I: The long term context, *Quaternary Science Reviews*, 134, 1–18, <https://doi.org/10.1016/j.quascirev.2015.12.005>, 2016.




Figure 1. Sites for proxy data assimilated in the reconstruction. Geographical coordinates of sites and sources of data are listed in Table 1.




Figure 2. Point-wise correlation between reconstructed and 20CRv3 SLP, T2m and precipitation for JJA mean (a-c) and DJF mean (d-f) data. The maximum correlation, mean significant correlation and number of grid points with significant correlation in indicated for each subplot. Contour indicates $p = 0.01$.




Figure 3. Point-wise correlation between reconstructed and 20CRv3 SLP, T2m and precipitation for annual mean (a-c) and monthly mean (d-f) data. The maximum correlation, mean significant correlation and number of grid points with significant correlation in indicated for each subplot. Contour indicates $p = 0.01$.




Figure 4. Time series of reconstructed (blue) and observed (yellow) JJA temperature for Nuuk, Ilulissat, Qaqortoq, Stykkisholmur, Central England, Copenhagen, Stockholm, and Uppsala. The blue shading indicated +/- 1 std dev of the ensemble reconstructed temperature.





Figure 5. Time series of reconstructed (blue) and observed (yellow) DJF temperature for Nuuk, Ilulissat, Qaqortoq, Stykkisholmur, Central England, Copenhagen, Stockholm, and Uppsala. The blue shading indicated +/- 1 std dev of the ensemble reconstructed temperature.


SST annual corr (recon, ERSSTv5)

Max. = 0.65, Mean = 0.33 (n sig. = 266)



SST annual corr (recon, COBE2)

Max. = 0.64, Mean = 0.33 (n sig. = 385)



SST annual corr (recon, HadISST)

Max. = 0.57, Mean = 0.34 (n sig. = 322)




Figure 6. Reconstruction of annual sea surface temperature (SST) correlated point-wise to three different SST datasets based on observations (ERSSTv5 1870-1970, COBE2 1850-1970, HadISST 1854-1970). Contour indicates $p = 0.01$.




Figure 7. Point-wise correlation between SAT25 and Mode-RA monthly T2m. The maximum correlation, mean significant correlation and number of grid points with significant correlation in indicated for each subplot. Contour indicates $p = 0.01$.

Table 1. Proxy data sites assimilated in the reconstruction with coordinates and data sources listed.

Location	Site name	Lat. (°N)	Long. (°E)	Source
Scotland	SCOT	57.08	-3.44	(Wilson et al., 2016)
E Alps - Tyrol	TYR	47.30	12.30	(Wilson et al., 2016)
Jaemtland	JAEM	63.30	13.25	(Wilson et al., 2016)
Tjeggelvas, Arjeplog, Ammarnäs composite	TAA	65.54-66.36	16.06-18.12	(Wilson et al., 2016)
North Fenno	EFmean	66-69	19-32	(Wilson et al., 2016)
Forfjorddalen	FORF	68.47	15.43	(Wilson et al., 2016)
Tatra	TAT	48-49	19-20	(Wilson et al., 2016)
South Finland	SFIN	62	29	(Wilson et al., 2016)
Quebec	L105	50.80	-68.8	(Wang et al., 2022)
Quebec	L20	54.60	-71.2	(Wang et al., 2022)
Quebec	L135	56.70	-74	(Wang et al., 2022)
Cent. England	ENG	51.5	-1.035	(Loader et al., 2020)
Central Europe	CZEC	49.0	13.0	(Büntgen et al., 2021)
Greenland	Crete	71.12	-37.32	(Vinther et al., 2010)
Greenland	DYE-3 71	65.18	-43.83	(Vinther et al., 2010)
Greenland	DYE-3 79	65.18	-43.83	(Vinther et al., 2010)
Greenland	GRIP 89-1	72.58	-37.64	(Vinther et al., 2010)
Greenland	GRIP 89-3	72.58	-37.6	(Vinther et al., 2010)
Greenland	GRIP 93	72.58	-37.64	(Vinther et al., 2010)
Greenland	Milcent	70.30	-44.50	(Vinther et al., 2010)
Greenland	Renland	71.27	-26.73	(Vinther et al., 2010)

Table 2. ~~Correlation Proxy parameters and corresponding model variables used in matching model output to Long-term temperature observations Annual, JJA, DJF, monthly proxy data.~~ ~~Boldface marks significant correlations with $p < 0.01$.~~

<u>Model variable</u>	<u>JJA T2m</u>	<u>JJA $\delta^{18}\text{O}$</u>	<u>May-Oct $\delta^{18}\text{O}$</u>	<u>Nov-Apr $\delta^{18}\text{O}$</u>
<u>MXD/TRW/BI</u>	<u>X</u>			
<u>$\delta^{18}\text{O}_{\text{cell}}$</u>		<u>X</u>		<u>X</u>
<u>Ice core $\delta^{18}\text{O}$ summer</u>			<u>X</u>	
<u>Ice core $\delta^{18}\text{O}$ winter</u>				<u>X</u>

Table 3. Correlation to Long-term temperature observations for the annual, JJA, DJF, and monthly mean. Boldface marks significant correlations with $p < 0.01$.

Site	Annual	JJA	DJF	monthly
Nuuk	0.63	0.21	0.62	0.30
Ilulissat	0.62	0.13	0.61	0.31
Qaqortoq	0.55	0.16	0.63	0.34
Stykkishólmur	0.58	0.54	0.33	0.28
Central England	0.49	0.50	0.24	0.21
Copenhagen	0.55	0.55	0.42	0.31
Stockholm	0.53	0.48	0.46	0.32
Uppsala	0.45	0.56	0.39	0.27