

Asymmetric shifts in seasonal transition timing in a sub-Arctic river system under climate warming
No longer on schedule, the pattern is breaking apart: The Loss of Seasonal Synchrony in a Sub-Arctic River System Under Warming Climate


Abolfazl Jalali Shahrood^{1*}, Amirhossein Ahrari¹, Ali Torabi Haghighi¹

¹Department of Water, Energy and Environmental Engineering Research Unit, University of Oulu, Pentti Kaiterankatu 1, Oulu, P.O.Box 8000, 90570, Finland

*Correspondence to: Abolfazl Jalali Shahrood (contact@abolfazljalali.com); Present address: Helsinki, Finland (independent researcher).

Formatted: Superscript

Abstract. Climate warming is altering the timing and synchrony of snow and ice processes across northern river systems, yet long-term shifts in their seasonal dynamics remain insufficiently resolved. Here, we analyze a 5759-year daily record (1 Oct 1966–202330 Sep 2025) from the River Oulankajoki in northeastern Finland to characterize proxy-based seasonal transitions in discharge, snow depth, and air temperature. freeze-up, break-up, snow accumulation and melt, and key atmospheric temperature transition points. Using a rule-based detection framework, we identify earlier shifts in springtime events across snow, discharge, and temperature timings (significant for discharge and temperature). Using a process-based detection tool, we identify significant advances in spring-related events, including snow melt, ice break-up, and the seasonal shift from cold to warm temperatures. In contrast, autumn/winter transition timings for those variables such as freeze-up and snow-onset exhibit higher year-to-year variability and no consistent trends. The Cooling Dominant Period (CDP) has shortened. Over the record the thermal cooling transition shifted later than snow onset, and the thermal warming transition earlier than snow end. This indicates that thermal proxies are shifting faster than their surface-response counterparts at this site. The durations of cold season, ice cover, and snow melt periods have shortened, while warm and open-water seasons have lengthened. Moreover, the temporal gap between atmospheric warming and surface responses has increased in spring but contracted in autumn. These findings suggest not only a shift in seasonal timing but also a growing desynchronization between atmospheric conditions and cryo-hydrological processes, with implications for Arctic river ecology, ice forecasting, and flood risk under continued climate change.



1 Introduction

The Arctic's cryo-hydrological systems are experiencing significant changes driven by climate warming. Arctic regions are warming at a rate approximately two to four times faster than the global average, with the most pronounced warming occurring during autumn and winter months (García Criado et al., 2025; Masson-Delmotte et al., 2021; Prowse et al., 2007). Annual average temperatures in the Arctic are expected to rise by approximately 3.7°C by 2050, relative to the 1981–2000 baseline (Prowse et al., 2007). Snow cover extent and duration have generally declined across the Northern Hemisphere in recent decades (Bring et al., 2016), with snow depth and duration projected to continue their downward trends (Burrell et al., 2023). Specifically, Arctic snow cover duration has shortened by approximately 2–4 days per decade over the past 30–40 years, and spring snow cover extent has declined by over 30% since 1971 (Box et al., 2019). Snowmelt is occurring earlier in many regions, often accompanied by more frequent rain-on-snow events (Park et al., 2016). These changes have also affected the timing of spring freshets, which have shifted earlier in Eurasian basins at a rate of about 1.1 days per decade, while North American basins show no significant trends (Feng et al., 2021).

Cryo-hydrological processes in Arctic river systems are experiencing significant transformations driven by climate change. These widespread changes in river flow patterns are primarily driven by rising air temperatures (Pavelsky and Zarnetske, 2017). The timing and magnitude of spring river flows are shifting, with many Arctic rivers transitioning from a nival to a pluvial regime (i.e., snowmelt-dominated to rainfall-dominated) (Prowse et al., 2006). River ice, a critical element of the cryosphere that significantly impacts the global hydrological system, is highly sensitive to weather and hydrological conditions (Shen, 2016). This sensitivity is particularly evident in the Northern Hemisphere, where by 2010, major ice cover existed on 29% of total river lengths, and seasonal ice affected approximately 58% of river lengths (Bennett and Prowse, 2010). Globally,

Field Code Changed

the duration of river ice cover has declined ~~Specifically~~specifically over the past three decades (Newton and Mullan, 2021; Yang et al., 2020), with studies attributing these changes to global warming (Fukš, 2023).

An example is the Danube River, where increasing winter temperatures have reduced its ice cover duration by about 28 days per century (Ionita et al., 2018). Broader analyses using more than 400,000 Landsat images reveal a global average reduction in river ice extent from 10% to 7.5% over the last 30 years (Yang et al., 2020). The most substantial reductions in ice cover duration have occurred in northeastern North America, Central Europe, and areas surrounding the Tibetan Plateau (Fukš, 2023). Ice thickness has also shown widespread decline. For instance, major Arctic rivers in Russia experienced reductions in maximum ice thickness ranging from 2.3 to 12.6 cm per decade between 1955 and 2012 (Fukš, 2023). Numerous studies report overall decreases in river ice thickness across various regions (Nalbant and Sharma, 2023; Vuglinsky, 2017; Vuglinsky and Valatin, 2018).

The timing of river ice freeze-up and break-up has been significantly shifting. Many rivers and lakes around the world exhibit later freeze-up dates and earlier break-up times, which leads to a shorter annual ice cover period (Janowicz, 2010; Newton and Mullan, 2021; Rokaya et al., 2019; Sharma et al., 2022; Shiklomanov and Lammers, 2014; Takács and Kern, 2015; Yang et al., 2020). Long-term records spanning the past 150–200 years provide clear evidence of these trends across Northern Hemisphere lakes and rivers, with freeze-up occurring approximately 5.8 days per century later and break-up about 6.5 days per century earlier (Burrell et al., 2023). On average, ice break-up has moved earlier by about 0.6 days per decade since 1850, correlating with a global temperature increase of about 0.12°C per decade (Fukš, 2023; Magnuson et al., 2000). Freeze-up timing shows greater regional variability, which shows both delayed and earlier occurrences depending on location; however, the dominant global trend is toward later freeze-up dates (Fukš, 2023). Spring and autumn temperatures play critical roles in determining break-up and freeze-up timing, respectively (Brown et al., 2018). Research conducted in the Mackenzie River Delta suggests increasingly earlier spring break-up, with Siberian rivers exhibiting even more pronounced changes than rivers in North America (Podkowa et al., 2023) ~~(Kugler et al., 2010)~~. These shifts are driven primarily by rising springtime air temperatures and indirectly by enhanced river discharge from increased snowmelt (Bieniek et al., 2011; Brown et al., 2018). Shifts in river ice regimes are disrupting both ecosystems and human activities in the Arctic. Earlier ice break-up and reduced ice cover alter seasonal flooding, nutrient transport, and thermal conditions in aquatic systems, that might affect fish habitats and species composition (Prowse et al., 2006; Yang et al., 2020). These changes also threaten traditional practices, winter transportation, and water resource availability in northern communities (Brown et al., 2018; Fukš, 2023). On land, permafrost thaw is reshaping vegetation, while aquatic ecosystems experience species shifts and declines in fish condition (Lehnherr et al., 2018; Liljedahl et al., 2016).

River ice and cold-climate hydrological processes are important indicators of climate change in Arctic regions, yet their potential is limited by persistent data gaps. River discharge data offer a useful proxy, with the timing and magnitude of spring flows reflecting changes in ice break-up. Increased spring discharge typically accelerates break-up, while autumn and early winter flows influence freeze-up timing (Feng et al., 2021; Park et al., 2016). Among climate variables, air temperature is the dominant driver of ice regimes, strongly correlating with both freeze-up and break-up events (Ionita et al., 2018; Shiklomanov

80 and Lammers, 2014). The freezing index, based on cumulative sub-zero temperatures, is commonly used to estimate ice thickness and seasonal transitions. Specifically, spring temperatures play a critical role in predicting break-up dates (Park et al., 2016). Snow cover is another key proxy, with declining winter snow depths and shorter durations contributing to earlier break-up through reduced surface albedo and enhanced melt (Bring et al., 2016; Lesack et al., 2014). Together, these variables indicate the value of integrating hydrological and climatic data to monitor cryospheric change.

85 There is a significant shortage of consistent hydrological observations related to river ice across the Arctic, particularly in remote regions (Podkova et al., 2023) (Kugler et al., 2010). Long-term records of freeze-up and break-up dates, as well as ice thickness, are scarce and often fragmented, that limits the ability to detect robust climate trends (Feng et al., 2021; Park et al., 2016). The decline in river gauging stations since the mid-1980s, especially in Russia and Canada, has further reduced the availability of continuous datasets (Prowse et al., 2007). Inconsistencies in data collection methods and definitions, such as

90 differing criteria for the start or end of ice events, complicate trend analysis and comparisons across regions (Sharma et al., 2022; Yang et al., 2020). For example, some observations record initial break-up, while others note complete ice clearance, a process that may span several weeks (Prowse et al., 2007). These methodological disparities, combined with reduced ground-based monitoring, hinder efforts to separate climate change signals from natural variability and challenge the integration of datasets at the pan-Arctic scale.

95 Given the persistent data gaps and inconsistencies in direct river ice observations, proxy ~~variables~~~~indicators~~ such as air temperature, river discharge, and snow cover have become essential for assessing changes in ice dynamics. These proxies are particularly important in regions where long-term observational records are sparse or incomplete. In response to this challenge, the present study builds on the River Ice Timing Characteristics and Extremes (RiTICE) ~~tool~~~~framework developed in~~ MATLAB (Jalali Shahrood et al., 2023) ~~to characterize long-term proxy-based transition timing in the River Oulankajoki to~~ evaluate long-term ice break-up trends in the River Oulankajoki in northern Finland. RiTICE is a rule-based framework that detects Phase Change Timings (PCTs) from daily discharge, snow depth, and air temperature time series. RiTiCE estimates break-up timing using daily discharge data and From these PCTs, Intra-Variable Intervals (IVIs) and Cross-Variable Intervals (CVIs) are derived to describe timing relationships between thermally-derived (i.e., air temperature) and surface response (i.e., discharge and snow depth) transitions. In this study, PCTs are grouped into an *onset-group* (autumn/winter transitions) and a *release-group* (spring transitions). RiTiCE has been validated across multiple Arctic and sub-Arctic river systems, including the Tornionjoki, Kiiminkijoki, Kemijoki, and Tana rivers (Jalali Shahrood, 2023; Jalali Shahrood et al., 2024). RiTICE provides a consistent and scalable method for monitoring river ice phenology in regions lacking direct measurements. This study aims to answer the following research questions:

100 (1) How have the PCTs shifted over the observational record?

105 (2) How have the derived IVIs and CVIs changed, and do these changes differ between the onset-group and the release-group?

110 (3) How has the temporal coupling between thermally-derived PCTs and their surface response evolved in terms of the direction and magnitude of cross-variable timing offsets?

Formatted: Font: Italic, Complex Script Font: Italic

Formatted: Font: Italic, Complex Script Font: Italic

2 Methodology

2.1 Phase Change Timings (PCTs) Detection

RiTICE is applied to annual daily time series of discharge, snow depth, and air temperature to detect transition timings in each variable. To identify the onset of seasonal transitions in river and climate-related variables, we used the RiTiCE (River Ice Timing Characteristics and Extremes) framework, developed and implemented in MATLAB. RiTiCE requires annual daily time series. Input data must be prepared under consistent preprocessing rules and analyzed using variable-specific transition detection methods.

2.2 Data Preparation

The datasets must be prepared as follows:

- Leap days (29 February) must be removed to maintain a uniform 365-day structure across all years.
- Data were must be reorganized by Water Year (WY), defined from October 1st to September 30th.
- Note. Each water year is named for its starting calendar year; for example, WY1966 spans 1 October 1966 to 30 September 1967, and the full record WY1966–WY2024 covers 1 October 1966 to 30 September 2025.

RiTICE detects PCTs, IVIs, and CVIs from annual daily time series of discharge, snow depth, and air temperature. The timings may not necessarily correspond directly to observed processes such as ice formation, break-up, or snowmelt but were earlier tested against recorded ice condition data in previous studies (Jalali Shahrood et al., 2023, 2024). The timings at which a variable undergoes detected transition are referred to as PCTs and defined in Table 1.

From PCTs, two types of derived metrics are constructed: (i) IVIs which describe durations between PCTs within a single variable, and (ii) CVIs, which quantify signed timing offsets between two PCT dates from different variables. IVIs and CVIs are summarized in Tables S1–S2. We report only the primary IVIs defined between each variable’s PCTs; their complementary intervals are deterministic 365-day complements and therefore redundant.

Primary aim of RiTiCE is to detect the timing of an event (or known as Feature in this study) on annual daily time series with a generic mathematical approach. The timing is when a parameter changes its phase known as Phase Change Timing (i.e., PCT). The terms “Features” and “Periods” are defined in Table 1 and Table 2.

Table 1 Summary of Phase Change Timing (PCT) events detected by RiTiCE.

Event	Abbreviation	Definition in RiTiCE	Parameter	Method to detect
break up day	BUD	The day when river flow begins rising after a long low flow period, which signals	Discharge	DVD (Daily Value Difference)

Formatted: Normal

Formatted: Font: (Default) +Headings (Times New Roman)
Complex Script Font: +Headings (Times New Roman)

		the start of ice break up.		
freeze-up day	FUD	The start of the longest period of stable discharge before the break up day;	Discharge	DVD (Daily Value Difference)
Snow build-up day	SBD	The start of the longest continuous period of non zero snow depth	Snow depth	CSP (Continuous Snow Presence)
Snow melting day	SMD	The end of the longest continuous period of non zero snow depth	Snow depth	CSP (Continuous Snow Presence)
Temperature Transition Point	TTP ⁺	The day on which transition from cold to warm season occurs	Temperature	ZCAT (Zero Referenced Cumulative Area Transition)
Temperature Transition Point	TTP ⁻	The day on which transition from warm to cold season occurs	Temperature	ZCAT (Zero Referenced Cumulative Area Transition)

Table 2 Definitions of seasonal periods used in the analysis.

Period	Definition	Parameter
Cold Season	Period between TTP⁻ and TTP⁺	Temperature
Warm Season	Complementary period to the cold season	Temperature
Ice Cover	Period between FUD and BUD	Discharge

Open Water	Period between BUD and FUD	Discharge
Snow Cover	Period between SBD and SMD	Snow depth
No Snow	Period between SMD and SBD	Snow depth

145

150 Table 1 Summary of PCTs detected by RiTiCE, their definitions, variables used and methods applied. Legacy labels (e.g., FUD, BUD, SBD, SMD, TTP-/TTP+) are omitted to avoid misinterpretation of PCTs as direct representations of freeze-up, break-up, or snowmelt events.

PCT	Legacy PCT	Definition	Variable	Method
DSO (Discharge Stabilization Onset)	FUD	Start of the longest period of reduced discharge variability	Discharge	DVD (Daily Value Difference)
DSE (Discharge Stabilization End)	BUD	End of the longest period of reduced discharge variability	Discharge	DVD (Daily Value Difference)
CSO (Continuous Snow Onset)	SBD	Start of the longest continuous period of non-zero snow depth	Snow depth	LSS (Longest Snow Sequence)
CSE (Continuous Snow End)	SMD	End of the longest continuous period of non-zero snow depth	Snow depth	LSS (Longest Snow Sequence)
CCT (Cumulative Cooling Transition)	TTP-	Transition to cumulative cooling dominance	Temperature	ZCAT (Zero-Referenced Cumulative Area Transition)
CWT (Cumulative Warming Transition)	TTP+	Transition to cumulative warming dominance	Temperature	ZCAT (Zero-Referenced Cumulative Area Transition)

Formatted Table

Formatted

Figure 1 summarizes how RiTiCE detects PCTs and IVIs from daily time series. Three variables of discharge, snow depth, and air temperature are processed using variable-specific methods. The DVD method detects Discharge Stabilization Onset and End (DSO, DSE), LSS identifies Continuous Snow Onset and End (CSO, CSE), and ZCAT defines Cooling and Warming Transitions (CCT, CWT). Each PCT pair defines an IVI, Discharge Stability Period (DSP: DSO–DSE), Continuous Snow Period (CSP: CSO–CSE), and Cooling Dominant Period (CDP: CCT–CWT).




Figure 1 Overview of the PCT (Phase Change Timing) feature extraction framework.

Formatted: Font: (Default) Times New Roman, Bold, Com
Script Font: Times New Roman, Bold
Formatted: Normal, Don't keep with next




Fig. 1 Overview of the PCT (Phase Change Timing) feature extraction framework.

2.3-2 Discharge Phase Change Timings (PCTs): Discharge Stabilization Onset and End (DSO & DSE) River-Ice Features: Freeze-Up and Break-Up Days (FUD & BUD)

In RiTiCE, DSO and DSE represent transition timings inferred from changes in discharge variability rather than direct observations of river ice conditions. DSO and DSE capture changes in discharge variability that may be associated with ice-related processes, but are not equivalent to visually observed freeze-up or break-up dates as defined by the International Association for Hydraulic Research (IAHR, 1980); although they can be compared with those reported in earlier studies (Jalali Shahrood et al., 2023, 2024).

In RiTiCE, FUD and BUD refer to the hydrological signature of river ice freeze-up and break-up. This is distinct from the visual or observational break-up date (e.g., the first day a river is entirely ice-free) and freeze-up date (e.g., The date on which the water body was first observed to be completely frozen over), as defined by the International Association for Hydraulic Research (IAHR, 1980). Instead, BUD refers explicitly to the timing of the river ice break-up day which is defined as the day when the river's flow begins to rise after a long period of low and stable discharge, corresponding to the start of ice break-up and the transition toward spring flow conditions. On the other hand, RiTiCE defines FUD as the day of year on which the flow enters the stable (i.e., least fluctuations) before BUD occurs.

Field Code Changed

Discharge stabilization is detected using the Daily Value Difference (DVD) method. Daily changes in discharge were computed between consecutive days as forward differences:

BUD is detected using the Daily Value Difference (DVD) method. The change in discharge is calculated between each pair of consecutive days:

$$\Delta Q_t = Q_{t+1} - Q_t \quad (1)$$

Note. An equivalent backward-difference definition $\Delta Q_t = Q_t - Q_{t-1}$ uses the same consecutive-day differences with a one-day index shift and therefore yields the same stability structure; detected DSO/DSE are not materially sensitive to this choice.

For each WY (365 days), this yields $N = 364$ day-to-day changes in discharge ΔQ_t . Variability of ΔQ_t across the entire WY is summarized by:

The stable segment, bounded by the FUD and BUD, reflects sustained ice presence. To isolate this period, we computed the daily variation in discharge (DVD) and applied a threshold defined by the standard deviation ($\pm\sigma$). A binary mask was then generated for $|\text{DVD}| \leq \sigma$, from which the longest continuous segment was extracted to represent the ice-covered season.

$$\sigma_{\Delta Q} = \sqrt{\frac{1}{N} \sum_{t=1}^N (\Delta Q_t - \overline{\Delta Q})^2}, \quad (2)$$

$$\text{BUD} = \max\{t_k\} + 1 \text{ where } |\Delta Q_{t_k}| < \sigma_{\Delta Q} \quad (3)$$

$$\text{FUD} = \min\{t_k\} + 1 \text{ where } |\Delta Q_{t_k}| < \sigma_{\Delta Q} \quad (4)$$

We classify day-to-day changes as stable when $|\Delta Q_t| \leq \sigma_{\Delta Q}$. DSE is taken as the day of WY corresponding to the end of the longest contiguous sequence of stable daily changes in the full WY, with day indices obtained by applying a consistent one-day offset to the ΔQ_t series.

In a second step, discharge and day-to-day changes are restricted to the period before DSE, a separate variability threshold is computed from those pre-DSE day-to-day changes only, and DSO is taken as the start of the longest contiguous stable segment under that pre-DSE threshold (with one-day offset). A pre-DSE threshold is used because the full-water-year variability of day-to-day discharge changes is dominated by the spring freshet, which is the largest source of day-to-day change in the annual hydrograph. Under a single full-water-year threshold, the low-variability winter recession can satisfy the stability criterion only a day or two into the water year in some years, placing DSO implausibly early. Restricting the threshold to the pre-DSE differences, from which the freshet is excluded, yields a stricter variability scale that locates DSO at the onset of the sustained winter recession. DSO is constrained to occur before DSE. DSP is then defined as the interval between DSO and DSE ($\text{DSP} = \text{DSE} - \text{DSO}$). DSE is the end day of the longest full-water-year stable run, and DSO is the start day of the longest pre-DSE stable run; when several runs share the maximum length, the earliest is selected. In this record

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

Formatted: Font: Not Bold, Complex Script Font: Not Bold

the two runs lie within a single contiguous stable segment in all 59 water years, so DSP = DSE – DSO represents a single continuous discharge-stability period (Figure 2).

Formatted: Font: Not Bold, Complex Script Font: Not Bold

$$DSE = \max\{t_k\} + 1 \quad \text{where} \quad |\Delta Q_{t_k}| \leq \sigma_{\Delta Q} \quad (3)$$

$$DSO = \min\{s_k\} + 1 \quad \text{where} \quad |\Delta Q_{s_k}| \leq \sigma_{\Delta Q}^{\text{pre}} \quad (4)$$

Where:

Q_t : discharge on day t (m^3/s)

ΔQ_t : day-to-day change $Q_{t+1} - Q_t$ (m^3/s)

$N = 364$: number of day-to-day changes in a 365-day WY


$\overline{\Delta Q}$: mean of ΔQ_t over the WY

t_k = indices of the longest contiguous run in the full WY with $|\Delta Q_t| \leq \sigma_{\Delta Q}$

s_k = indices of the longest contiguous run before DSE with $|\Delta Q_t| \leq \sigma_{\Delta Q}^{\text{pre}}$

$\sigma_{\Delta Q}$: standard deviation of ΔQ_t over the full WY (used for DSE)

$\sigma_{\Delta Q}^{\text{pre}}$: standard deviation of ΔQ_t computed on day-to-day changes before DSE (used for DSO)



Formatted: Left

Where:

Q_t = Discharge on day t (m^3/s)

ΔQ_t = Daily change in discharge, $Q_{t+1} - Q_t$ (m^3/s)

N = Number of daily ΔQ_t values in the water year (364)

$\overline{\Delta Q}$ = Mean of all ΔQ_i values over the water year

$\sigma_{\Delta Q}$ = Standard deviation of the ΔQ_i series

t_k = Time indices of the longest contiguous run satisfying $|\Delta Q_i| < \sigma_{\Delta Q}$

BUD = Break Up Day, defined as the day index $\max\{t_k\} + 1$

FUD = Freeze Up Day, defined as the day index $\min\{t_k\} + 1$

Figure 2 Detection of Discharge Stabilization Onset (DSO) and Discharge Stabilization End (DSE) by the Daily Value Difference (DVD) method, illustrated for WY1966. (Left) Daily discharge with DSO (blue dashed line) and DSE (red dashed line); the Discharge Stability Period (DSP) is shaded blue and the complementary Discharge Variability Period (DVP) pink. (Centre) Day-to-day discharge changes (DVD) with both variability thresholds: the full-water-year $\pm\sigma$ band (green dashed), used to locate DSE as the end of the longest full-water-year stable run, and the stricter pre-DSE $\pm\sigma$ band (orange dotted), used to locate DSO as the start of the longest stable run before DSE. (Right) The two corresponding binary stability masks: the full-water-year mask (green, for DSE) and the pre-DSE mask (orange, for DSO); a day is stable (mask = 1) when its DVD value lies within the relevant band, otherwise unstable (mask = 0). The pre-DSE mask in the third panel follows the same binary logic of Full-WY mask (0 and 1) but for better readability its height is manually reduced in this figure.




Fig. 2 Detection of BUD (Break Up Day) and FUD (Freeze Up Day) using discharge data. (Left) Daily discharge with BUD (blue dashed line) and FUD (red dashed line). (Center) Daily Value Difference (DVD) method showing changes in discharge. (Right) Binary ice cover mask derived from periods where $|\text{DVD}| \leq \sigma$, used to define stable low-flow (ice covered) periods between BUD and FUD.

2.4.3 Snow Features: Build-Up and Melt Days (SBD & SMD) Snow Depth Phase Change Timings (PCTs): Continuous Snow Onset and End (CSO & CSE)

To determine the timing of persistent snow cover, we apply the Longest Snow Sequence (LSS) method (Figure 3). A binary snow presence mask is generated from daily snow depth values, where each day is assigned a value of 1 if snow is present ($S \neq 0$) and 0 otherwise:

To determine the timing of persistent seasonal snow cover, we apply the Continuous Snow Presence (CSP) method. A binary mask is generated based on daily snow depth values, where each day is assigned a value of 1 if snow is present ($S \neq 0$) and 0 otherwise. The longest continuous segment of days with a value of 1 in this mask identifies the main snow cover period. The first day of this segment is defined as the Snow Build-Up Day (SBD). The last day of this segment is defined as the Snow Melt Day (SMD):

$$M_t = H(S_t), \quad \text{where } H(x) = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \end{cases} \quad (55)$$

LSS identifies the longest $M_t=1$ segment and hence CSO and CSE; CSP is then the period between CSO and CSE:

$$[CSO = t_{\text{start}}, \quad CSE = t_{\text{end}}] \quad M_t = 1 \quad \forall t \in [t_{\text{start}}, t_{\text{end}}] \quad (6)$$

Longest continuous subsequence of days where $M_t = 1$,

$$[SBD = t_{\text{start}}, \quad SMD = t_{\text{end}}] \quad M_t = 1 \quad \forall t \in [t_{\text{start}}, t_{\text{end}}] \quad (6)$$

Where:

S_t = Snow depth on day t (in cm), for $t \in \{1, 2, \dots, 365\}$

M_t = Binary snow presence mask

$[t_{\text{start}}, t_{\text{end}}]$ = Indices of the longest continuous subsequence such that $M_t = 1$ for all $t \in [t_{\text{start}}, t_{\text{end}}]$

Where:

S_t = Snow depth on day t (in cm), for $t \in \{1, 2, \dots, 365\}$

M_t = Binary snow presence mask

$[t_{\text{start}}, t_{\text{end}}]$ = Indices of the longest continuous subsequence such that $M_t = 1$ for all $t \in [t_{\text{start}}, t_{\text{end}}]$

SBD = t_{start} (Snow Build-Up Day)

SMD = t_{end} (Snow Melt Day)

Formatted

Formatted




Figure 3 Detection of Continuous Snow Onset (CSO) and Continuous Snow End (CSE) from snow depth data. (Top) Daily snow depth with CSO (blue dashed line) and CSE (red dashed line). (Bottom) Binary snow presence mask used to identify the longest Continuous Snow Period (CSP).

Formatted: Caption




Fig. 3 Detection of SBD (Snow Build-Up Day) and SMD (Snow Melt Day) from snow depth data. (Top) Daily snow depth with SBD (blue dot) corresponding to the start of continuous snow cover and SMD (red dot) corresponding to the end. (Bottom) Binary snow presence mask (1 if snow depth > 0, else 0) used to identify the longest continuous snow covered period.

2.5.4 Temperature Features: Transition Points (TTP) & TTP⁺ Air Temperature Phase Change Timings (PCTs): Cumulative Cooling and Warming Transitions (CCT & CWT)

Temperature transitions are characterized using the Zero-Referenced Cumulative Area Transition (ZCAT) method, which captures the integrated thermal state by accumulating daily areas between the temperature curve and the 0 °C baseline (Figure 4). ZCAT is used because it returns a single within-year transition date for each of the cooling and warming phases directly from the cumulative curve. Freezing and thawing degree-day sums (FDD and TDD) quantify accumulated thermal energy in °C days but do not by themselves define a transition date without an additional threshold or breakpoint rule. FDD and TDD are therefore reported separately as complementary thermal context rather than as timing definitions. For each day t , the area between day t and $t + 1$ is computed using the trapezoidal rule:

305

To characterize the seasonal temperature transition, we apply the Zero-Referenced Cumulative Area Transition (ZCAT) method. This approach captures the integrated thermal state by accumulating daily areas between the temperature curve and the 0°C baseline.

For each day t , the area between day t and $t + 1$ is computed using the trapezoidal rule with a unit time step:

$$A_t = \frac{T_t + T_{t+1}}{2} \tag{77}$$

310

The cumulative temperature area is then given by:

$$C_t = \sum_{i=1}^t A_i \tag{88}$$

where:

315

- T_t = daily mean temperature on day t
- A_t = area contribution between day t and $t + 1$
- C_t = Cumulative curve
- CCT = the last zero-crossing of C_t prior to CWT, marking the transition from cumulative warming to cumulative cooling dominance.
- CWT = the day at which C_t reaches its minimum, marking the transition from cumulative cooling to cumulative warming dominance.

320

Note. In one water year (WY2009), C_t did not cross zero within the water-year window, and CCT defaulted to the first day of the water year. Inspection of the preceding water year showed that the corresponding zero-crossing fell at the end of September, just before the water-year boundary. Because this affects a single year, the water-year definition and the ZCAT procedure are kept unchanged in this study.

325

Formatted




Figure 4 Detection of Cumulative Cooling Transition (CCT) and Cumulative Warming Transition (CWT) using Zero-Referenced Cumulative Area Transition (ZCAT). (Top) Daily air temperature. (Bottom) Cumulative temperature curve, used to detect CCT & CWT. As A_t is computed between t and $t + 1$, reported transition days follow the same $+1$ index mapping.

where:

- T_t = daily mean temperature on day t
- A_t = area contribution between day t and $t + 1$
- C_t = Cumulative curve
- TTP^+ = the day corresponding to the minimum of the C_t , the transition from cold season to warm season.
- TTP^- = timing of the last zero-crossing of the C_t before TTP^+ , the transition from warm season to cold season.




Fig. 4 Identification of temperature transition points (TTP⁻ and TTP⁺) using cumulative temperature analysis. (Top) Daily air temperature with cold season highlighted; TTP⁻ (blue dot) signals the start of cumulative cooling, and TTP⁺ (red dot) signals the transition to cumulative warming. (Bottom) Cumulative temperature curve used to detect these inflection points.

2.5 Data

Daily air temperature and snow depth records were obtained from the Finnish Meteorological Institute (FMI) Open Data Portal (Finnish Meteorological Institute, 2026), specifically from the Kuusamo Kiutaköngäs weather station (N66°22'4", E29°19'40"), about 500 meters away from the Oulanka Research Station. These records span from October 1966 to September 2025 (WY1966–WY2024). Snow depth is recorded in whole centimeters. The FMI code –1 denotes a trace (snow present but below the 1 cm measurement resolution) and was recoded to 0 for the continuous-snow-cover analysis. A total of 35 non-consecutive days across the WY1966–WY2024 window (12 days with missing values on recorded dates and 23 absent calendar dates) were linearly interpolated to maintain a complete 365-day-per-year series. The observational accuracy for snow depth is ± 2 cm, and the records resolutions is 1 cm (whole centimeters). Since CSO and CSE are defined by the longest continuous run of snow presence, they are governed by the sustained winter snowpack rather than by ± 2 cm noise near the 1 cm presence

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Heading 2

threshold. Daily river discharge measurements were acquired from the Oulanka Research Station (daily mean), hosted by the Finnish Environment Institute (SYKE, 2026). Winter discharge values in the Hertta database are flagged as 'Redukoitu' (quality code '='), meaning they are estimates derived from post-processing corrections applied when ice conditions affect the rating curve relationship. The DVD-based proxies (DSO, DSE) are therefore derived from a quality-controlled rather than a directly measured record, and DSO/DSE should be interpreted as discharge regime transition proxies rather than direct ice phenology dates.

In addition to discharge, snow depth, and air temperature used for RiTiCE, we analyzed ancillary datasets of water temperature and ice thickness (available for limited periods). These records are operationally constrained (e.g., seasonal deployment, access/safety, and quality control) and therefore are not treated as direct freeze-up/break-up observations. Instead, they are used as operational context to assess whether the PCTs occur in a plausible seasonal neighborhood relative to these operations, and not as validation of the proxy timings (Figure S2). The ancillary datasets were obtained from SYKE open data platform (SYKE, 2026).

2.6 Trends and correlations analysis

Long-term monotonic trends in annual PCT, IVI, and CVI series, and in the hydroclimatic extremes, were tested using the Mann–Kendall test (Kendall, 1975; Mann, 1945)(Mann, 1945). To reduce potential inflation of significance arising from serial autocorrelation, each series was pre-whitened by removing the AR(1) component prior to testing (Von Storch, 1999)(Von Storch, 1999). This pre-whitened Mann–Kendall procedure is referred to as PW-MK throughout results. Statistical significance was evaluated at $p < 0.05$. Trend magnitude was quantified using Sen's slope (Sen, 1968; Theil, 1950)(Sen, 1968), reported in the native units of each metric (days yr^{-1} for PCTs and IVIs; signed days yr^{-1} for CVIs). As many series are tested, we controlled the false discovery rate within predefined families of related tests (the six PCT trends, the three IVI trends, the six CVI trends, the three seasonal air-temperature trends, the degree-day trends, and the hydroclimatic extreme-value trends) using the Benjamini–Hochberg procedure (Benjamini and Hochberg, 1995). We report the Benjamini–Hochberg q-value alongside the uncorrected p-value. Our main conclusions rest on trends that are individually strong and remain significant after this correction; trends that are nominally significant (uncorrected $p < 0.05$) but fall just above the corrected threshold ($q > 0.05$) are reported with their q-values and interpreted as nominal rather than as established trends.

Associations among PCTs and IVIs over the full record were assessed using Spearman rank correlations (ρ) (Spearman, 1904), computed on paired water-year values. Correlation p-values are reported alongside ρ . Within each Spearman correlation matrix we also applied the Benjamini–Hochberg correction. Every correlation interpreted here remains significant after correction. The record was also divided into an early period (WY1966–WY1994, $n=29$) and a late period (WY1995–WY2024, $n=30$) to describe changes in timing structure between the two halves of the study period. Early–late correlation differences were

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Heading 2

385 assessed with a permutation test (the 59 paired water-year values were pooled and randomly reassigned to the early and late
periods 20,000 times to build the null distribution of the difference in Spearman ρ) and are treated as descriptive. (Fisher, 1921)
To detect long-term monotonic trends in the key variables, we applied the Mann-Kendall trend test, a non-parametric method
widely used in hydrology and climatology. The significance of observed trends was evaluated at conventional confidence
levels ($P\text{-Value} < 0.05$). To further explore the relationships between seasonal PCT features, we calculated the Spearman
correlation coefficients among all six PCT markers including BUD, FUD, SMD, SBD, TTP⁺, and TTP⁻, over the full 57-year
390 record.

2.76 Study Area

River Oulankajoki (Figure 5), is a boreal river system located in northeastern Finland near the Arctic Circle (Arvola and
Nurmesniemi, 2000; Saraniemi et al., 2008). Situated within the boreal zone, it displays cold-climate hydrological
395 characteristics, including seasonal snow accumulation, extended ice cover, and distinct freeze-thaw cycles. Hydrologically,
the river is strongly seasonal (Blåfield et al., 2024; Saraniemi et al., 2008). The mean annual discharge is approximately 25.5
m³/s, with low flows dropping to around 3 m³/s in late winter and peak flows reaching up to 249 m³/s during the spring
snowmelt in May–June. The river typically freezes from mid-November to early May. Recent climatic trends show significant
warming in the region, with average temperatures rising by 0.61°C per decade and summer temperatures by 0.41°C per decade.
400 These trends are shortening the winter season and altering the basin’s overall hydroclimatic regime. Over the past five decades,
spring floods have weakened by 7%, while high-flow events in other seasons have increased by 10%. Annual minimum flows
have risen by 28%, that reflects both climatic and hydrological shifts (Blåfield et al., 2024).




Figure 5 Location of the River Oulankajoki basin.

Fig. 5 Location and of the River Oulankajoki basin.

Daily air temperature and snow depth records were obtained from the Finnish Meteorological Institute (FMI) Open Data Portal, specifically from the Kuusamo-Kiutaköngäs weather station (N66°22'4", E29°19'40"), about 500 meters away from the Oulanka Research Station. These records span from October 1966 to September 2023 and serve as key indicators of thermal and snowmelt dynamics affecting river-ice phenology. Daily river discharge measurements were acquired from the Oulanka Research Station, hosted by SYKE (Finnish Environment Institute).

3 Results


3.1 Distribution and Early–Late Shifts of PCTs Seasonal Hydroclimatic Patterns in a Reference Year

Over the 59-year record (WY1966–WY2024, $n=59$), the six PCTs ranged across distinct portions of the WY (Figure 6; embedded table). Among the onset-group, DSO ranged from day 15 to day 95 (median 49, $SD \pm 17.5$ d). CSO ranged from day

5 to day 70 (median 34, SD ± 14.3 d). CCT ranged from day 1 to day 88 (median 38, SD ± 20.9 d; the day 1 value corresponds to WY2009, see Section 2.4). Among the release-group, DSE ranged from day 196 to day 231 (median 216, SD ± 8.6 d). CSE ranged from day 206 to day 238 (median 224, SD ± 7.0 d). CWT ranged from day 177 to day 222 (median 201, SD ± 10.9 d). CCT showed the largest spread and CSE showed the smallest of all six PCTs.

Formatted: English (United States)

Formatted: Keep with next



PCT	Mean $\pm$ SD	Median [p10-p90]	Min	Max
DSO	50.0 $\pm$ 17.5	49 [27-72]	15	95
DSE	216.3 $\pm$ 8.6	216 [205-226]	196	231
CSO	34.7 $\pm$ 14.3	34 [19-51]	5	70
CSE	223.5 $\pm$ 7.0	224 [214-233]	206	238
CCT	38.0 $\pm$ 20.9	38 [12-66]	1	88
CWT	200.3 $\pm$ 10.9	201 [184-214]	177	222

Figure 6 Summary of PCTs over 59 years.

The record was divided into two approximately equal sub-periods to allow balanced comparison of early and late period distributions. Comparing the early period (WY1966–WY1994, n=29) with the late period (WY1995–WY2024, n=30), all three onset-group PCTs shifted to later days of the WY (Figure 7). DSO shifted by +2.5 d (median: day 48.0 to day 50.5). CSO shifted by +4.0 d (median: day 34.0 to day 38.0). CCT shifted by +17.5 d (median: day 30.0 to day 47.5). All three release-group PCTs shifted to earlier days of the WY. DSE shifted by –3.0 d (median: day 218.0 to day 215.0). CSE shifted by –1.5 d (median: day 224.0 to day 222.5). CWT shifted by –12.5 d (median: day 208.0 to day 195.5). Decadal-binned distributions of all six PCTs across consecutive 10-year blocks (WY1966–WY1975 through WY2016–WY2024) are provided in Supplementary Figure S1; in the decadal-binned distributions, CCT and CWT showed the most consistent directional

Formatted: Not Highlight

progression across successive decades, while DSO, CSO, DSE, and CSE varied across decadal blocks without a monotonic step pattern.

This study presents one of the most comprehensive long-term assessments of cryo-hydrological dynamics in a sub-Arctic river system, analyzing 57 years of in-situ observations from the River Oulankajoki. By defining and analyzing six seasonal (PCT) features and their corresponding durations, we observed a clear and systematic shift toward earlier thaw-related events and a broader restructuring of cold-season hydroclimatic regimes. The 1966 reference cycle illustrates the structurally ordered nature of seasonal transitions under stable cryo-hydrological conditions. In more recent decades, however, this sequence has grown increasingly variable. Rather than a simple forward shift of the seasonal calendar, we observe elongated transition windows. These results are consistent with Arctic-wide trends which shows earlier melt onset and later freeze-up.

In the reference year 1966, the seasonal progression of cryo-hydrological conditions showed distinct and coherent transitions in discharge, snow cover, and air temperature (Fig. 6). Freeze-Up Day (FUD) was identified on day 58 of the hydrological year, initiating a 166-day ice cover period that lasted until Break-Up Day (BUD) on day 223 (Fig. 6, top). Snow Build-Up Day (SBD) occurred earlier, on day 48, with snow depth remaining continuously above zero until Snow Melt Day (SMD) on day 225, resulting in a snow cover duration of 178 days (Fig. 6, middle). Air temperature dropped below freezing early in the year, with the warm-to-cold transition point (TTP⁻) occurring on day 11. The cold-to-warm transition point (TTP⁺) was reached on day 218, after which temperatures stayed consistently above freezing. This marked a 208-day cold season (Fig. 6, bottom). Full-year characterizations for all 57 years are provided in Appendix A.

Formatted: English (United Kingdom)




Figure 7 Early vs late period distributions of PCTs (day of WY) for WY1966–WY1994 (n=29) and WY1995–WY2024 (n=30): panel titles report the change in median timing (late – early, days). Top row shows onset-group PCTs and bottom row release-group PCTs. Left column shows discharge PCTs, middle snow depth PCTs, and right air temperature PCTs.

Formatted: Font: (Default) Times New Roman, Complex
Script Font: Times New Roman, English (United Kingdom)




Fig. 6 Hydroclimatic conditions in 1966 showing seasonal transitions. (Top) Discharge with break-up (BUD) and flow rise (FUD) days. (Middle) Snow depth with build up (SBD) and melt (SMD) days. (Bottom) Air temperature with cold to warm (TTP⁻) and warm to cold (TTP⁺) transitions.

3.2 Ordering Patterns and Interannual Variability of PCT SequencingTemporal Sequencing and Interannual Variability of Seasonal Features

The onset-group varied in ordering across the full record (WY1966–WY2024, n=59; Figure 8A). Five of the six possible orderings occurred at least once. The most frequent ordering was CSO→CCT→DSO, occurring in 25 years (42.4%). CCT→CSO→DSO occurred in 13 years (22.0%), CSO→DSO→CCT in 11 years (18.6%), CCT→DSO→CSO in 7 years (11.9%), and DSO→CSO→CCT in 3 years (5.1%). The ordering DSO→CCT→CSO did not occur in any year. Across the full record, DSO preceded CCT in 14 of 59 years (23.7%), and DSO preceded CSO in 7 of 59 years (11.9%). Three water years illustrate how these reversals arise (Figure S3). In WY1975, a short early snow spell fell outside the longest continuous snow segment, so CSO was set later and the order was CCT, DSO, CSO. In WY2011, discharge stabilized while the cumulative cooling transition was delayed, giving the order CSO, DSO, CCT. In WY2020, discharge stabilized early and the initial snow record again fell outside the continuous segment, giving the order DSO, CSO, CCT. The release group occupied a fixed global ordering in all 59 years: CWT→DSE→CSE occurred in 100% of WYs (Figure 8B). DSO, CSO, and CCT each ranged across global ranks 1, 2, and 3, with no PCT holding a fixed rank position in any year, while CWT held global rank 4, DSE rank 5, and CSE rank 6 in every year (Figure 8G).

475 Comparing the early (WY1966–WY1994, n=29; Figures 8C–D) and late (WY1995–WY2024, n=30; Figures 8E–F) sub-
480 periods, the release-group invariance (CWT→DSE→CSE, 100%) held in both periods without exception. Within the onset
group, the ordering CSO→CCT→DSO was the most frequent in both periods (37.9% early, tied with CCT→CSO→DSO at
37.9%; 46.7% late).

A central finding of this study is the asymmetry between spring and autumn transitions. The sequence of seasonal cryo-
hydrological features revealed a clear separation between early- and late-year events (Fig. 7). Snow Build-Up Day (SBD),
warm-to-cold transition (TTP⁻), and Freeze-Up Day (FUD) showed considerable year-to-year fluctuations in both order and
timing. In contrast, the cold-to-warm transition (TTP⁺), Break Up Day (BUD), and Snow Melt Day (SMD) maintained a fixed
sequence across all 57 years.

SBD and TTP⁻ frequently alternated between the first and second positions, with FUD occasionally preceding both.
485 Meanwhile, TTP⁺, BUD, and SMD maintained a stable order, consistently appearing in the final positions of the seasonal
progression. This persistent ordering reflects a strong temporal stability in spring-related markers compared to the more
variable autumn features (Fig. 7, bottom). This difference illustrates that spring transitions are increasingly governed by
consistent atmospheric warming, whereas autumn processes remain subject to localized and stochastic controls, such as

precipitation phase, surface insulation, and early snowfall events.





Figure 8 Interannual variability and asymmetry in the ordering of PCTs (WY1966–WY2024, n=59). (A) Onset-group ordering frequencies, full record. (B) Release-group order-pattern frequencies, full record. (C) Onset-group order-pattern frequencies, early period (WY1966–WY1994, n=29). (D) Release-group ordering frequencies, early period. (E) Onset-group ordering frequencies, late period (WY1995–WY2024, n=30). (F) Release-group ordering frequencies, late period. (G) Year-to-year PCT rank order changes across the full record (rank 1 = earliest in that water year).

Formatted: Font: (Default) Times New Roman, Complex Script Font: Times New Roman, English (United Kingdom)
Formatted: Caption



500 **Fig. 7 Annual ranks and temporal shifts of six cryo hydrological Phase Change Timing Features (1966–2022). (Top) rank distributions; (middle) frequency of shifts; (bottom) year-to-year order changes.**

505 **3.3 Spearman Correlations Among PCTs and IVIs - Full Record and Early-Late Comparison**
Correlations Among PCT Features and Seasonal Periods

Within the release-group, DSE and CSE showed the strongest PCT correlation ($r=0.74$, $p<0.001$), followed by DSE and CWT ($r=0.60$, $p<0.001$) and CSE and CWT ($r=0.53$, $p<0.001$) over the 59-year record. Within the onset-group, DSO and CSO ($r=0.46$, $p<0.001$), DSO and CCT ($r=0.48$, $p<0.001$), and CSO and CCT ($r=0.49$, $p<0.001$) were all significant and positive (Figure 9A).

In the early period (WY1966–WY1994, $n = 29$), the strongest correlations were DSE and CSE ($r = 0.82$, $p < 0.001$) and DSE and CWT ($r = 0.67$, $p < 0.001$). DSO and CSO ($r = 0.64$, $p < 0.001$) and CSE and CWT ($r = 0.55$, $p = 0.002$) were also significant (Figure 9B).

In the late period (WY1995–WY2024, $n = 30$), DSO and CCT, and CSO and CCT, increased to $r = 0.52$ ($p = 0.003$) and $r = 0.68$ ($p < 0.001$), respectively, while DSE and CSE, DSE and CWT, and CSE and CWT, decreased to $r = 0.60$ ($p < 0.001$), $r = 0.44$ ($p = 0.015$), and $r = 0.44$ ($p = 0.016$), respectively. DSO and DSE showed a nominally significant negative correlation ($r = -0.45$, $p = 0.013$) in the late period (Figure 9C).

For IVIs over the full record, all three pairings were significant and positive: DSP and CSP ($r = 0.55$, $p < 0.001$), DSP and CDP ($r = 0.57$, $p < 0.001$), and CSP and CDP ($r = 0.58$, $p < 0.001$) (Figure 9D). In the early period, DSP and CSP ($r = 0.61$, $p < 0.001$) and DSP and CDP ($r = 0.43$, $p = 0.021$) were significant; CSP and CDP did not reach significance ($r = 0.36$, $p = 0.052$) (Figure 9E). In the late period, DSP and CSP ($r = 0.4$, $p = 0.03$), DSP and CDP ($r = 0.54$, $p = 0.002$), and CSP and CDP ($r = 0.69$, $p < 0.001$) were all significant (Figure 9F), with CSP and CDP showing the strongest IVI correlation of either period. The early–late correlation differences are not significant under a permutation test and are described without implying a change in coupling.

Relationships among the six Phase Change Timing (PCT) features and seasonal period durations reveal structured associations across the 57-year record (Fig. 8). Among the PCT features, the strongest correlation was observed between Snow Melt Day (SMD) and Break Up Day (BUD), with a coefficient of ($r = 0.76$). Both features were also positively correlated with the cold-to-warm temperature transition point (TTP⁺), with values of ($r = 0.60$) for BUD and ($r = 0.54$) for SMD. In contrast, the warm-to-cold transition marker (TTP⁻) showed weaker and negative correlations with spring events, including ($r = -0.34$) with SMD and ($r = -0.28$) with BUD. Correlations among cold-season features were more moderate. Snow Build-Up Day (SBD) was positively associated with both FUD ($r = 0.44$) and TTP⁻ ($r = 0.48$).

The seasonal period duration matrix showed a strong inverse relationship between Cold Season and both No Snow ($r = -0.57$) and Open Water periods ($r = -0.55$). Cold Season duration was positively associated with Snow Cover ($r = 0.57$) and Ice Cover ($r = 0.55$), while Snow Cover and Ice Cover were also correlated ($r = 0.52$). No Snow and Open Water durations were moderately correlated ($r = 0.52$), and both were negatively related to Ice Cover and Cold Season. In contrast, the duration of the Snow Melt period exhibited weak correlations with all other periods, ranging from -0.05 to 0.17 .

Formatted: English (United Kingdom)





Figure 9 Spearman correlation matrices for PCTs (top row, panels A–C) and IVIs (bottom row, panels D–F) for the full record (WY1966–WY2024, n=59), early period (WY1966–WY1994, n=29), and late period (WY1995–WY2024, n=30).



Formatted: Font: (Default) Times New Roman, Complex Script Font: Times New Roman

Formatted: Caption

Formatted: Caption, Don't keep with next

~~Fig. 8 Correlation matrix of seasonal transition timing parameters. The values represent Spearman correlation coefficients between key phenological markers: BUD (Break Up Day), FUD (Freeze Up Day), SMD (Snow Melt Day), SBD (Snow Build Up Day), TTP⁺ (Cold to Warm Temperature Transition), and TTP⁻ (Warm to Cold Transition).~~

~~3.4 Long-term Trends in PCT, IVI, CVI, Temperature, Freeze–Thaw Degree-Days, and Hydroclimatic Extremes~~ ~~Long-term Trends in Phase Change Timings, Seasonal Durations, Hydroclimatic Indicators, and Lag Metrics~~

~~Among the six PCTs, CCT shifted later at +0.538 d/yr (PW-MK $p < 0.001$) and CWT shifted earlier at -0.357 d/yr ($p = 0.002$). DSE shifted earlier at -0.182 d/yr ($p = 0.023$) and CSE at -0.118 d/yr ($p = 0.038$). DSO (Sen = $+0.154$ d/yr, $p = 0.281$) and CSO (Sen = $+0.128$ d/yr, $p = 0.247$) showed no significant trend. Among the IVIs, CDP decreased at -0.871 d/yr ($p < 0.001$). DSP decreased at -0.333 d/yr ($p = 0.041$) under the SD-based threshold; because this DSP trend does not persist under an alternative variability scale, it is reported as indicative rather than as a robust trend. CSP showed a negative Sen slope (-0.237 d/yr) but did not reach significance ($p = 0.089$) (Figure 10). After within-family Benjamini–Hochberg correction, CCT ($q = 0.00385$), CWT ($q = 0.00663$), and DSE ($q = 0.0457$) remain significant among the PCTs and CDP ($q < 0.001$) among the IVIs, whereas CSE ($q = 0.0572$) and DSP ($q = 0.062$) fall just above the corrected threshold and are treated as nominal. Figure 9 summarizes long-term trends across four thematic categories of phase change timings, seasonal durations, hydroclimatic indicators, and lag metrics. In the phase change timings, statistically significant trends ($p < 0.05$) were detected for Break-Up Day (BUD), Snow-Melt Day (SMD), and both positive and negative Temperature Transition Points (TTP⁺, TTP⁻). BUD and SMD showed a shift toward earlier dates, indicating earlier river ice break-up and complete snowmelt. TTP⁺ has shifted earlier, while TTP⁻ has delayed. No significant trends were observed for Freeze-Up Day (FUD) and Snow-Build-Up Day (SBD), although both showed slight upward tendencies. For seasonal durations, significant shortening was detected in the Cold Season ($p < 0.000$), Ice Cover Duration ($p = 0.026$), and Snow-Melt Duration ($p = 0.001$), while the Warm Season ($p < 0.000$) and Open Water Duration ($p = 0.042$) lengthened. Snow Cover and No Snow durations increased but did not reach statistical significance ($p = 0.128$). The hydroclimatic indicators section shows that Minimum Discharge increased significantly ($p = 0.036$), whereas Maximum Discharge remained stable ($p = 0.918$). Ice Cover Average Discharge showed a near-significant trend ($p = 0.062$). Neither Minimum ($p = 0.073$) nor Maximum Air Temperature ($p = 0.184$) had significant trends, although both increased slightly. Cold Season Average Temperature showed a non-significant rise ($p = 0.346$). Maximum Snow Depth remained stable ($p = 0.660$). In terms of lag metrics, the time lag between Cold Season Start and Snow Cover Start significantly decreased ($p = 0.004$), while the lags between Cold Season End and both Ice Cover End ($p = 0.003$) and Snow Cover End ($p = 0.001$) increased significantly. The lag between Cold Season Start and Ice Cover Start showed a non-significant decline ($p = 0.096$).~~

Formatted: Keep with next




Figure 10 Long-term trends in Phase Change Timings (PCTs) and Intra-Variable Intervals (IVIs) (WY1966–WY2024; n=59). Panel titles report the PW-MK p-value, the Benjamini–Hochberg q-value (corrected within family), and Sen's slope; the q-value is shown in red where the trend remains significant after correction ($q < 0.05$).

The offset between CSE and CWT decreased at -0.231 d/yr ($p=0.011$), and the offset between CSO and CCT increased at $+0.333$ d/yr ($p=0.040$); both offsets are nominal after the multiple-testing correction ($q=0.068$ and $q=0.120$, respectively). The remaining four CVIs showed no significant trend, the offset between DSO and CSO ($\text{Sen}=+0.074$ d/yr, $p=0.496$), between DSO and CCT ($\text{Sen}=+0.222$ d/yr, $p=0.103$), between DSE and CSE ($\text{Sen}=0.000$ d/yr, $p=0.958$), and between DSE and CWT ($\text{Sen}=-0.167$ d/yr, $p=0.102$) (Figure 11).

Formatted: Keep with next

Formatted: Keep with next




Figure 11 Long-term trends in cross-variable timing offsets (CVIs) derived from PCT pairs (WY1966–WY2024; n=59). Panel titles report the PW-MK p-value, the Benjamini–Hochberg q-value (corrected within family), and Sen's slope; the q-value is shown in red where the trend remains significant after correction ($q < 0.05$).

585 Mean air temperature increased significantly in all three seasonal windows (Figure 12), with Oct–Dec at $+0.066$ °C/yr ($p=0.001$), Mar–May at $+0.037$ °C/yr ($p=0.0076$), and the Nov–Apr window at $+0.067$ °C/yr ($p=0.0041$). All three seasonal air-temperature trends remain significant after within-family correction.




Figure 12 Long-term trends in seasonal air temperature (AT) (WY1966–WY2024; n=59). Panel titles report the PW-MK p-value, the Benjamini–Hochberg q-value (corrected within family), and Sen's slope; the q-value is shown in red where the trend remains significant after correction ($q < 0.05$).

595

FDD accumulated from WY start to CCT increased at $+0.956\text{ }^{\circ}\text{C}\cdot\text{days}/\text{yr}$ ($p<0.01$). FDD accumulated from CCT to CWT decreased at $-13.9\text{ }^{\circ}\text{C}\cdot\text{days}/\text{yr}$ ($p<0.001$), and full water-year FDD decreased at $-12.5\text{ }^{\circ}\text{C}\cdot\text{days}/\text{yr}$ ($p<0.002$). TDD accumulated from WY start to CWT increased at $+0.76\text{ }^{\circ}\text{C}\cdot\text{days}/\text{yr}$ ($p=0.044$) and full water-year TDD increased at $+6.78\text{ }^{\circ}\text{C}\cdot\text{days}/\text{yr}$ ($p<0.003$). TDD accumulated from CCT to CWT showed no significant trend ($\text{Sen}=-0.104\text{ }^{\circ}\text{C}\cdot\text{days}/\text{yr}$, $p=0.278$) (Figure 13). After within-family correction, the three FDD trends and the full-water-year TDD trend ($q=0.00573$) remain significant, whereas the trend in TDD accumulated from the water-year start to CWT ($q=0.053$) falls just above the threshold and is treated as nominal.

600





Figure 13 Long-term trends in freezing and thawing degree-day (FDD/TDD) linked to the ZCAT transition timings (WY1966–WY2024; $n=59$). Panel titles report the PW-MK p-value, the Benjamini–Hochberg q-value (corrected within family), and Sen’s slope; the q-value is shown in red where the trend remains significant after correction ($q < 0.05$).

605

Annual maximum discharge ($p=0.896$), minimum discharge ($p=0.172$), and average discharge during the DSP ($p=0.178$) showed no significant trends. Maximum snow depth ($p=0.539$) and average snow depth during the CSP ($p=0.295$) showed no significant trends. Annual maximum air temperature ($p=0.156$) and average air temperature during the CDP ($p=0.261$) showed no significant trends. Annual minimum air temperature showed the strongest extreme-value trend ($+0.0596\text{ }^{\circ}\text{C}/\text{yr}$, uncorrected $p=0.0498$), but no extreme-value metric remains significant after the multiple-testing correction (Figure 14).

Formatted: Keep with next



610 **Figure 14** Long-term trends in hydroclimatic extreme-value and their average within IVIs (WY1966–WY2024; n=59). Panel titles report the PW-MK p-value, the Benjamini–Hochberg q-value (corrected within family), and Sen's slope; the q-value is shown in red where the trend remains significant after correction ($q < 0.05$).




Fig. 9 Mann-Kendall trend analysis from October 1966 to September 2023.

4 Discussion

Our results form a coherent picture of asymmetric change in proxy-based transition timing at the River Oulankajoki over 59 WYs. Thermally-derived transitions (CCT, CWT) shifted faster and further than their surface response transitions (DSO, CSO, DSE, CSE), and this asymmetry was more coherent and systematic in the release-group, where all three PCTs shifted earlier (significantly for CWT and DSE, and nominally for CSE after multiple-testing correction) with CWT shifting furthest, than in the onset-group, where only the thermally-derived CCT shifted significantly while surface proxies remained trendless. Among the onset group, the thermally-derived CCT shifted significantly (+0.538 d/yr, $p=0.001$), while the surface response proxies DSO and CSO showed no significant trend. At the period level, this contrast is clear; CCT shifted +17.5 d between the early and late sub-periods, compared with only +2.5 d for DSO and +4.0 d for CSO. In the release group, CWT shifted -12.5 d versus -3.0 d for DSE and -1.5 d for CSE. Air temperature is measured at a single point approximately 500 m from the gauge, while discharge integrates the entire upstream basin; this spatial mismatch can contribute to occasional ordering reversals in onset-group PCTs (e.g., DSO preceding CSO or CCT), particularly in years with early baseflow stabilization or delayed cumulative cooling. Figure S3 shows three example water years (WY1975, WY2011, and WY2020) in which these reversals occur. Among the release group, CWT, DSE, and CSE all shifted to earlier days, with CWT carrying the largest shift (-0.357 d/yr, $p=0.002$). These PCT shifts are consistent with the observed warming across all three seasonal windows: Oct-Dec mean

AT increased at $+0.066^{\circ}\text{C}/\text{yr}$ ($p=0.001$), Nov–Apr at $+0.067^{\circ}\text{C}/\text{yr}$ ($p=0.004$), and Mar–May at $+0.037^{\circ}\text{C}/\text{yr}$ ($p=0.008$), indicating that both the onset and release seasons warmed significantly over the record.

This difference is consistent with release-group PCTs being more tightly governed by progressive atmospheric warming, whereas onset-group PCTs remained subject to localized and stochastic controls, such as precipitation phase, surface insulation, and early snowfall events. Prior studies report that river ice release timing across North America and Europe has advanced at approximately 0.6 days per decade, closely linked to warming temperatures (Chen and She, 2020; Newton and Mullan, 2021), while onset timing trends remain spatially heterogeneous and weakly correlated with climate signals (Liston and Hall, 1995; Prowse and Bonsal, 2004), which is consistent with the pattern observed at this case study site. The sequencing analysis reinforces this structure; within the onset-group, CCT, DSO, and CSO exhibited flexible interannual rankings, with five of six possible orderings occurring across the 59-year record. Moreover, the dominant ordering $\text{CSO} \rightarrow \text{CCT} \rightarrow \text{DSO}$ increased in frequency from 37.9% in the early period to 46.7% in the late period, which suggests a gradual consolidation of onset-group sequencing consistent with CCT shifting progressively later relative to the surface proxies; since this ordering involves DSO, whose absolute timing is method-dependent, these frequencies are interpreted qualitatively. In contrast, the release-group maintained the invariant ordering $\text{CWT} \rightarrow \text{DSE} \rightarrow \text{CSE}$ in all 59 years. The correlation structure further supports this asymmetry; strong pairwise correlations among release-group PCTs (CWT and DSE: $r=0.60$; CWT and CSE: $r=0.53$; DSE and CSE: $r=0.74$) contrasted with weaker and more variable correlations among onset-group PCTs, particularly in the early period. Onset-group correlations were numerically higher in the late period ($\text{CSO} - \text{CCT}$ $r=0.68$, $\text{DSO} - \text{CCT}$ $r=0.52$) than in the early period, and some release-group correlations were numerically lower ($\text{DSE} - \text{CSE}$ $r=0.82$ to $r=0.60$), but none of these early–late differences is significant under a permutation test.

The CDP shortened at $-0.871 \text{ d}/\text{yr}$ ($p<0.001$), the largest IVI trend in the dataset, contracting simultaneously from both ends as CCT shifted later and CWT shifted earlier. FDD accumulated between CCT and CWT declined at $-13.9^{\circ}\text{C}\cdot\text{days}/\text{yr}$ ($p<0.001$), which indicates that the thermal winter narrowed in both duration and cold intensity. Notably, FDD accumulated from the WY start to CCT increased at $+0.956^{\circ}\text{C}\cdot\text{days}/\text{yr}$ ($p=0.010$), indicating that early-season cold accumulation intensified even as the overall cold period contracted. The CDP therefore narrowed primarily through a later onset of the cooling-dominant phase and an earlier end, rather than through a uniform reduction in cold intensity across the season. Full water-year FDD also declined ($-12.5^{\circ}\text{C}\cdot\text{days}/\text{yr}$, $p=0.002$), while full water-year TDD increased ($+6.78^{\circ}\text{C}\cdot\text{days}/\text{yr}$, $p=0.003$), which further supports the rebalancing of the annual thermal budget away from cold accumulation and toward warm accumulation driven primarily by the inward movement of CDP boundaries. DSP also decreased under the primary threshold ($-0.333 \text{ d}/\text{yr}$, $p=0.041$), which is consistent with DSE shifting earlier; however, this DSP trend is not robust to the discharge variability scale and is treated as indicative. CSP showed a negative Sen’s slope but did not reach significance ($p=0.089$), consistent with the smaller period-level shift in CSE (-1.5 d) relative to CWT (-12.5 d) between sub-periods. These results at this site are directionally in line with reported trends of earlier seasonal transitions across Arctic and sub-Arctic systems (Dauginis and

Brown, 2021; Fukš, 2023; Markus et al., 2009; Newton and Mullan, 2021; Podkowa et al., 2023; Sharma et al., 2016; Wang and Feng, 2024), though generalization beyond this site is not made here.

We also identified two changes in CVIs, the temporal offsets between thermally-derived and surface response transitions, both nominal after the multiple-testing correction. The interval between CSO and CCT increased at +0.333 d/yr ($p=0.040$), meaning CCT shifted later relative to CSO over the record. The interval between CSE and CWT shifted at -0.231 d/yr ($p=0.011$), which means CWT advanced faster than CSE. The interval between DSE and CWT shifted in the same direction but did not reach significance ($p=0.102$). The interval between DSE and CSE remained stationary (Sen= 0.000 d/yr, $p=0.958$), both DSE and CSE shifted almost in parallel throughout the record. A number of studies report that despite earlier atmospheric warming, surface responses have lagged in sub-Arctic and boreal systems (Schwartz et al., 2006; Stone et al., 2002). This growing offset between thermally-derived and surface response transitions in systems with similar thermal-nival structure, poses challenges for hydrological modelling and ecosystem forecasting, as degree-day and temperature-threshold models may increasingly misestimate the timing of surface-driven responses (Bjorkman et al., 2015; Groisman and Easterling, 1994; Siegel et al., 2022). The smaller shift in CSE relative to CWT (-1.5 d vs -12.5 d) may also partly reflect that CSO/CSE are based on terrestrial snow depth at a single station rather than snow conditions on the river surface.

Despite significant shifts in PCT timing and IVI duration, all discharge and snow depth magnitude metrics remained statistically unchanged over the 59-year record. Neither maximum discharge ($p=0.896$) nor minimum discharge ($p=0.172$) showed a significant trend. Average discharge during the DSP showed no significant trend ($p=0.178$). Maximum snow depth ($p=0.539$) and average snow depth during the CSP ($p=0.295$) also remained stable. Annual minimum air temperature showed a nominal increase ($+0.060$ °C/yr, uncorrected $p=0.0498$) that does not survive the multiple-testing correction, and annual maximum air temperature showed no significant trend ($p=0.156$). These findings align with studies reporting resilience in streamflow and snow depth magnitudes despite substantial hydroclimatic timing shifts (Harder et al., 2015; Shiklomanov et al., 2007), with increases in winter baseflow in northern basins attributed to permafrost thaw and deeper infiltration (St. Jacques and Sauchyn, 2009). Although some studies report increasing peak flows in certain regions (Burn et al., 2010; Byun et al., 2019), maximum discharge at this site remained stable over the record ($p=0.896$).

Changes in the timing of onset-group and release-group PCTs alter the seasonal windows available for sediment and nutrient transport, aquatic habitat structure, and biological processes. Earlier shifts in release-group PCTs and a contracting CDP affect the timing of ecological cues that depend on seasonal temperature and flow structure (Janowicz, 2010; Prowse et al., 2011). The stability and timing of the winter low-flow period bear on the reliability of winter infrastructure in northern communities (Prowse et al., 2007). Since the DSP trend is not robust to the discharge variability scale, we do not draw a conclusion about a change in the length of this period here.

All findings in this study are specific to the River Oulankajoki at the Oulanka gauging station over WY1966–WY2024. Winter discharge values in the source record are operationally corrected estimates (quality code 'Redukoitu'), meaning DSO and DSE

are derived from a quality-controlled rather than a directly measured series and should be interpreted accordingly as discharge regime transition proxies. The two discharge timings differ markedly in their sensitivity to the DVD variability scale. Replacing the SD-based threshold with an IQR-based alternative (IQR/1.349) displaces DSO by a median of 91 days (median of signed differences, later in almost all years) but DSE by a median of only 7 days (earlier in every year), because DSO marks the onset of stability immediately after the abrupt spring freshet, whereas DSE marks the end of the winter stable period along the gradual autumn–winter recession. DSE and its earlier trend are robust to this choice, whereas DSO, DSP, and all DSO-derived quantities are method-dependent and are interpreted qualitatively rather than as quantitative trend estimates; in particular, the DSP trend does not persist under the alternative scale. The qualitative onset-group result (flexible interannual ordering of DSO, CSO, and CCT with no coherent onset trend) and the release-group invariance (CWT→DSE→CSE in all 59 years) do not depend on the variability scale. Independent observational context was provided by ice thickness measurement records (WY1980–WY1994, n=15) and water temperature winter gap data (WY1970–WY2024, n=55); however, both datasets are operationally constrained and neither constitutes a direct freeze-up or break-up observation, their temporal relationship with the proxy PCTs shows considerable interannual spread (Figure S2), and they should be interpreted as plausibility checks rather than validation. The proxy framework used here is most appropriate for unregulated boreal rivers without frequent mid-winter melt events; applicability to regulated or hydropeaking systems would require re-evaluation of the DVD stability definition. This study presents one of the most comprehensive long-term assessments of cryo-hydrological dynamics in a sub-Arctic river system, analysing 57 years of in-situ observations from the River Oulankajoki. By defining and analysing six seasonal (PCT) features and their corresponding durations, we observed a clear and systematic shift toward earlier thaw-related events and a broader restructuring of cold-season hydroclimatic regimes.

A central finding of this study is the asymmetry between spring and autumn transitions. Spring PCT features including, SMD, BUD, and TTP⁺, showed statistically significant trends toward earlier timing. In contrast, autumn PCT features including, SBD, FUD, and TTP⁻, showed no significant long-term trends and demonstrated greater interannual variability, with frequent delays and timing fluctuations. This difference illustrates that spring transitions are increasingly governed by consistent atmospheric warming, whereas autumn processes remain subject to localized and stochastic controls, such as precipitation phase, surface insulation, and early snowfall events. This seasonal asymmetry is strongly supported by prior studies. Break-up dates across North America and Europe have advanced by ~0.6 days per decade, closely linked to spring warming (Chen and She, 2020; Newton and Mullan, 2021), while freeze-up trends remain spatially heterogeneous and weakly correlated with climate signals (Liston and Hall, 1995; Prowse and Bonsal, 2004). These dynamics are further evidenced in our feature sequencing analysis. SBD, TTP⁻, and FUD exhibit flexible interannual rankings, frequently switching order across years. In contrast, the sequence of TTP⁺, BUD, and SMD remains fixed. This consistency supports the interpretation that spring processes are more temporally constrained and climatically synchronized, while autumn markers retain internal autonomy. The correlation matrix reinforces this conclusion, strong interconnections among spring features (i.e., TTP⁺, BUD, and SMD) contrast with weak or inconsistent correlations among autumn indicators (i.e., SBD, TTP⁻, and FUD).

In addition to timing shifts, we have found evidence of a broader redistribution of seasonal durations. The cold season, ice cover, and snow melt periods have significantly shortened, while the warm season and open water durations have lengthened. Trends in snow cover and no snow durations were weak but positive which suggests increasing variability. This uneven redistribution reflects a restructuring of seasonal phase coherence. (Dauginis and Brown, 2021; Markus et al., 2009) Rather than a simple forward shift of the seasonal calendar, we observe elongated transition windows. These results are consistent with Arctic-wide trends which shows earlier melt onset and later freeze-up. This contributes to lengthened melt seasons of up to 2–3 weeks per decade (Dauginis and Brown, 2021; Markus et al., 2009). Satellite-based observations also highlight spatial heterogeneity in snow cover dynamics (Dye, 2002; Kim et al., 2018). The emergence of complex, nonlinear transitional phases, such as multi-stage ice freeze-up and snow melt onset without sustained thaw, reveals the increasing fragmentation of seasonal coherence across Arctic river systems (Kim et al., 2018; Markus et al., 2009).

We have detected evolving lag relationships between atmospheric temperature transitions and corresponding surface responses. The lag between cold season onset (i.e., TTP⁻) and snow accumulation (i.e., SBD) has decreased, this means a faster response to initial freezing. However, lags between the end of the cold season (i.e., TTP⁺) and the snow melt (i.e., SMD) and ice (i.e., BUD) have increased significantly, which means snow and ice are taking longer to respond to spring warming. A number of studies support this trend that despite earlier spring warming, snow melt has lagged expectations (Schwartz et al., 2006; Stone et al., 2002). This disconnection between air temperature and surface melt poses substantial challenges for both hydrological modeling and ecosystem forecasting. Traditional degree-day models may fail to predict the timing of snowmelt-driven runoff or spring ecological cues, as lagged surface responses become increasingly decoupled from atmospheric indicators (Bjorkman et al., 2015; Groisman and Easterling, 1994; Siegel et al., 2022).

Despite the pronounced shifts in cryospheric timing and duration, our analysis reveals relatively stable trends in hydrological extremes. Minimum discharge has increased modestly, while maximum discharge and snow depth have remained statistically unchanged. We observed a nearly significant increase in river flow during the ice-covered period ($p = 0.062$), which may suggest more water is moving beneath the ice. These findings align with studies reporting regional resilience in streamflow metrics despite substantial hydroclimatic shifts (Harder et al., 2015; Shiklomanov et al., 2007). Increases in winter baseflow observed in northern basins have been attributed to permafrost thaw and deeper infiltration (St. Jacques and Sauchyn, 2009). Although some studies report increasing spring peaks in certain regions (Burn et al., 2010; Byun et al., 2019), our results suggest that in many sub-Arctic systems, hydrological extremes remain buffered, potentially due to offsetting influences like soil water retention, increased evapotranspiration, or groundwater recharge (Bring et al., 2016).

Ecologically, changes in freeze-up and break-up timing alter sediment and nutrient transport, affect aquatic habitat structure, and misalign biological cues. Earlier phytoplankton blooms and zooplankton peaks may benefit fast-growing species but disrupt food web synchrony for others with more complex life cycles (Janowicz, 2010; Prowse et al., 2011). From a societal perspective, shorter and less reliable ice seasons threaten winter road networks, vital for northern communities and industries (Prowse et al., 2007). Mid-winter break-up events pose risks to infrastructure, navigation, and hydropower operations (Beltaos et al., 2007). While extended open water periods may reduce icebreaking costs, they also disrupt traditional practices, fish

migrations, and ecological timing. Changes in freeze-thaw sequences increase the likelihood of compound flood risks, particularly from rain-on-snow events or early break-up under high-flow conditions (Beltaos and Prowse, 2009) as our results show significantly increasing trend of minimum discharge during ice cover period. In that way we may observe more mechanical ice break-up in the near future, which leads to more hazardous floods than thermal ones. They tend to produce thicker ice jams and lead to higher water levels, which increases the risk of severe flooding and damage (Prowse and Stephenson, 1986). The dynamic nature of mechanical break-up events, which involves rapid discharge, moving ice, and powerful waves, can trigger intense erosion and sudden spikes in sediment transport (Beltaos and Burrell, 2021). Mechanical break-up events also involve large runoff volumes acting on thick, intact ice, that leads to extreme surges in water level (Prowse et al., 2007). Ecologically, ice-induced floods from mechanical break-up events can significantly disrupt both in-channel and riparian processes (Prowse and Culp, 2003).

Taken together, our results show that the cryo-hydrological system is not just shifting earlier in the year, it is being fundamentally reorganized. Instead of a simple seasonal shift, we see complex changes in how different processes (including timing, duration, and response delays) interact. In some cases, earlier melt doesn't mean faster melting it can actually stretch out the melt period but with less intensity (Pomeroy et al., 2015). This evolving cryo-hydrological complexity challenges the utility of temperature-based models and reinforces the need for improved process-based representations. Effective forecasting and resource management will require models that integrate thermal-hydraulic coupling, threshold effects, and lagged surface responses to remain robust under continued Arctic warming. We acknowledge persistent limitations in cryo-hydrological datasets. Long-term, spatially-dense monitoring networks remain sparse across the Arctic, particularly in Eurasia (Feng et al., 2021; Ionita et al., 2018). This constrains the ability to generalize findings, attribute change drivers, or validate remote-sensing products. Advancing our understanding will require sustained in-situ measurements, integration of high-resolution satellite data, and deployment of distributed sensor networks capable of capturing both surface and subsurface dynamics in cold environments.

5 Conclusions

This study provides a 5759-year assessment of seasonal cryo-hydrological dynamics in a sub-Arctic river system using a structured detection approach across six key timing metrics. By applying the RiTiCE framework to the River Oulankajoki, we document a coherent shift in release-group timings toward earlier dates (statistically significant for DSE and CWT, and nominal for CSE after multiple-testing correction), while onset-group PCTs (DSO, CSO) remain temporally unstable and trendless, and CCT shifted significantly later. This growing asymmetry is evident in proxy-detected transition timing at this site. We document a coherent and statistically significant shift in spring-related features—namely snow melt, river ice break-up, and the atmospheric cold-to-warm transition—toward earlier dates, while autumn-related features remain temporally

unstable and trendless. This divergence highlights a growing seasonal asymmetry in how Arctic rivers respond to climate forcing.

The analysis reveals that changes in phase transition timing are not occurring in isolation. The CDP has shortened, and the offset between CSE and CWT narrowed over the record (CWT advancing faster than CSE; nominal after the multiple-testing correction). The DSP also shortened under the primary threshold, but this DSP trend is reported as indicative because it is sensitive to the discharge variability scale. These results document differential rates of shift between thermally-derived and surface response proxies at this site, rather than a uniform seasonal shift. The analysis reveals that changes in phase transition timing are not occurring in isolation. Cold season, ice cover, and snow melt durations have shortened, whereas the warm season and open water periods have expanded. Lags between atmospheric transitions and surface responses have also evolved: spring surface processes are increasingly delayed relative to the onset of warming, while autumn reactions occur more promptly following cooling. These trends suggest a breakdown in the internal synchrony of cold season processes rather than a uniform seasonal shift.

Despite these structural changes, hydrological extremes such as peak discharge and maximum snow depth remain relatively stable. This suggests a buffered system response, possibly due to subsurface storage, evapotranspiration shifts, or groundwater compensation. However, a modest but consistent increase in minimum winter discharge points to enhanced under-ice flow, potentially linked to permafrost degradation or increased baseflow, which may elevate the risk of mechanical break-up events. The results demonstrate that proxy-detected seasonal transitions at this site show increasing differential rates of shift under continued warming. Hydrological transitions in Arctic river systems are becoming less predictable and more fragmented. The differential shifts between thermally-derived and surface proxy transitions pose a challenge for proxy-based forecasting of seasonal conditions. The progressive loss of seasonal coherence—both in timing and interprocess alignment—poses a challenge for forecasting river ice conditions, assessing flood risk, and managing aquatic ecosystems. These findings emphasize the need for long-term observational records and improved process-based models that incorporate not only temperature thresholds but also lagged hydrological responses and transitional variability under continued warming at this site and in systems with similar thermal-nival structure. Arctic warming.

Appendices

Appendix A

<https://figshare.com/s/d6de8875fd8151a80085>

Code and Data availability

The daily air temperature and snow depth records used in this study are publicly available from the Finnish Meteorological Institute (FMI) Open Data Portal (<https://en.ilmatieteenlaitos.fi/open-data>), and the daily mean river discharge records from the Finnish Environment Institute (SYKE) Hertta database (<https://wwwp2.ymparisto.fi/scripts/kirjaudu.asp>). The RiTiCE analysis code and the processed water-year input series generated for this study are available from the corresponding author upon reasonable request. All data supporting the findings of this study are publicly available at: FMI Open Data Portal: <https://en.ilmatieteenlaitos.fi/open-data> Finnish Environmental Institute (SYKE): <https://wwwp2.ymparisto.fi/scripts/kirjaudu.asp>

Author contribution

A.J.S. conceived the study, developed the methodology, processed and analyzed the data, wrote the manuscript, and prepared all figures. A.A. contributed to the interpretation of results and manuscript preparation. A.T.H. supervised the study.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

The authors gratefully acknowledge the Maa- ja vesiteknikan tuki ry (MVTT) for funding this research through grant no. 4802. We thank the Oulanka Research Station for long-term hydrological monitoring and infrastructure support. We are also grateful to SYKE (Finnish Environment Institute) and the Finnish Meteorological Institute (FMI) for providing the ~~discharge and meteorological~~ datasets. We appreciate the efforts of all personnel involved in field measurements and data collection over this ~~57~~59-year record. Artificial intelligence tools were used in a limited capacity to assist with code refinement, code commenting, instruction generation, and language editing of the manuscript. No scientific concepts, methodology, or analytical logic were generated by AI.

References

Arvola, L. and Nurmesniemi, A.: Seasonal dynamics of plankton in two boreal rivers in northern Finland, SIL Proceedings, 1922-2010, 27, 1928–1932, <https://doi.org/10.1080/03680770.1998.11901578>, 2000.

Benjamini, Y. and Hochberg, Y.: Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing, Journal of the Royal Statistical Society Series B: Statistical Methodology, 57, 289–300, <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>, 1995.

Bennett, K. E. and Prowse, T. D.: Northern Hemisphere geography of ice-covered rivers, Hydrological Processes, 24, 235–240, <https://doi.org/10.1002/hyp.7561>, 2010.

Bieniek, P. A., Bhatt, U. S., Rundquist, L. A., Lindsey, S. D., Zhang, X., and Thoman, R. L.: Large-scale climate controls of interior Alaska river ice breakup, *Journal of Climate*, 24, 286–297, 2011.

Bjorkman, A. D., Elmendorf, S. C., Beamish, A. L., Vellend, M., and Henry, G. H. R.: Contrasting effects of warming and increased snowfall on Arctic tundra plant phenology over the past two decades, *Global Change Biology*, 21, 4651–4661, <https://doi.org/10.1111/gcb.13051>, 2015.

Blåfield, L., Marttila, H., Kasvi, E., and Alho, P.: Temporal shift of hydroclimatic regime and its influence on migration of a high latitude meandering river, *Journal of Hydrology*, 633, 130935, 2024.

Box, J. E., Colgan, W. T., Christensen, T. R., Schmidt, N. M., Lund, M., Parmentier, F.-J. W., Brown, R., Bhatt, U. S., Euskirchen, E. S., and Romanovsky, V. E.: Key indicators of Arctic climate change: 1971–2017, *Environmental Research Letters*, 14, 045010, 2019.

Bring, A., Fedorova, I., Dibike, Y., Hinzman, L., Mård, J., Mernild, S. H., Prowse, T., Semenova, O., Stuefer, S. L., and Woo, M.-K.: Arctic terrestrial hydrology: A synthesis of processes, regional effects, and research challenges, *Journal of Geophysical Research: Biogeosciences*, 121, 621–649, 2016.

Brown, D. R., Brinkman, T. J., Verbyla, D. L., Brown, C. L., Cold, H. S., and Hollingsworth, T. N.: Changing river ice seasonality and impacts on interior Alaskan communities, *Weather, Climate, and Society*, 10, 625–640, 2018.

Burn, D. H., Sharif, M., and Zhang, K.: Detection of trends in hydrological extremes for Canadian watersheds, *Hydrological Processes*, 24, 1781–1790, <https://doi.org/10.1002/hyp.7625>, 2010.

Burrell, B. C., Beltaos, S., and Turcotte, B.: Effects of climate change on river-ice processes and ice jams, *International Journal of River Basin Management*, 21, 421–441, <https://doi.org/10.1080/15715124.2021.2007936>, 2023.

Byun, K., Chiu, C.-M., and Hamlet, A. F.: Effects of 21st century climate change on seasonal flow regimes and hydrologic extremes over the Midwest and Great Lakes region of the US, *Science of the Total Environment*, 650, 1261–1277, 2019.

Chen, Y. and She, Y.: Long-term variations of river ice breakup timing across Canada and its response to climate change, *Cold Regions Science and Technology*, 176, 103091, 2020.

Dauginis, A. A. and Brown, L. C.: Recent changes in pan-Arctic sea ice, lake ice, and snow-on/off timing, *The Cryosphere*, 15, 4781–4805, 2021.

Feng, D., Gleason, C. J., Lin, P., Yang, X., Pan, M., and Ishitsuka, Y.: Recent changes to Arctic river discharge, *Nature communications*, 12, 6917, 2021.

Finnish Meteorological Institute: Daily Air Temperature and Snow Depth Observations, Kuusamo Kiutaköngäs Weather Station, 2026.

Fukš, M.: Changes in river ice cover in the context of climate change and dam impacts: a review, *Aquat Sci*, 85, 113, <https://doi.org/10.1007/s00027-023-01011-4>, 2023.

García Criado, M., Myers-Smith, I. H., Bjorkman, A. D., Elmendorf, S. C., Normand, S., Aastrup, P., Aerts, R., Alatalo, J. M., Baeten, L., and Björk, R. G.: Plant diversity dynamics over space and time in a warming Arctic, *Nature*, 1–9, 2025.

Groisman, P. Y. and Easterling, D. R.: Variability and trends of total precipitation and snowfall over the United States and
890 Canada, *Journal of Climate*, 7, 184–205, 1994.

Harder, P., Pomeroy, J. W., and Westbrook, C. J.: Hydrological resilience of a Canadian Rockies headwaters basin subject to
changing climate, extreme weather, and forest management, *Hydrological Processes*, 29, 3905–3924,
<https://doi.org/10.1002/hyp.10596>, 2015.

IAHR: International Association for Hydraulic Research Association, *MULTILINGUAL ICE TERMINOLOGY*, 1980.

895 Ionita, M., Badaluta, C.-A., Scholz, P., and Chelcea, S.: Vanishing river ice cover in the lower part of the Danube basin—signs
of a changing climate, *Scientific Reports*, 8, 7948, 2018.

Jalali Shahrood, A.: Past, present, and future of river flow regime in Nordic region focusing on river ice break-up events, 2023.

Jalali Shahrood, A., Ahrari, A., Rossi, P. M., Klöve, B., and Torabi Haghighi, A.: RiTiCE: River Flow Timing Characteristics
and Extremes in the Arctic Region, *Water*, 15, 861, 2023.

900 Jalali Shahrood, A., Ahrari, A., Karjalainen, N., Klöve, B., and Haghighi, A. T.: Application of RiTiCE in understanding
hydro-meteorological controls on ice break-up patterns in River Tornionjoki, *Environ Monit Assess*, 196, 764,
<https://doi.org/10.1007/s10661-024-12910-w>, 2024.

Janowicz, J. R.: Observed trends in the river ice regimes of northwest Canada, *Hydrology Research*, 41, 462–470, 2010.

Kendall, M. G.: Rank correlation methods, 4th ed., Charles Griffin, London, 202 pp., 1975.

905 Lehnerr, I., St. Louis, V. L., Sharp, M., Gardner, A. S., Smol, J. P., Schiff, S. L., Muir, D. C., Mortimer, C. A., Michelutti,
N., and Tarnocai, C.: The world’s largest High Arctic lake responds rapidly to climate warming, *Nature communications*, 9,
1290, 2018.

Lesack, L. F., Marsh, P., Hicks, F. E., and Forbes, D. L.: Local spring warming drives earlier river-ice breakup in a large Arctic
delta, *Geophysical Research Letters*, 41, 1560–1567, 2014.

910 Liljedahl, A. K., Boike, J., Daanen, R. P., Fedorov, A. N., Frost, G. V., Grosse, G., Hinzman, L. D., Iijima, Y., Jorgenson, J.
C., and Matveyeva, N.: Pan-Arctic ice-wedge degradation in warming permafrost and its influence on tundra hydrology, *Nature
geoscience*, 9, 312–318, 2016.

Liston, G. E. and Hall, D. K.: Sensitivity of lake freeze-up and break-up to climate change: a physically based modeling study,
Annals of Glaciology, 21, 387–393, 1995.

915 Magnuson, J. J., Robertson, D. M., Benson, B. J., Wynne, R. H., Livingstone, D. M., Arai, T., Assel, R. A., Barry, R. G., Card,
V., and Kuusisto, E.: Historical trends in lake and river ice cover in the Northern Hemisphere, *Science*, 289, 1743–1746, 2000.

Mann, H. B.: Nonparametric tests against trend, *Econometrica: Journal of the econometric society*, 245–259, 1945.

Markus, T., Stroeve, J. C., and Miller, J.: Recent changes in Arctic sea ice melt onset, freezeup, and melt season length, *J.
Geophys. Res.*, 114, 2009JC005436, <https://doi.org/10.1029/2009JC005436>, 2009.

920 Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., and Gomis,
M. I.: Climate change 2021: the physical science basis, Contribution of working group I to the sixth assessment report of the
intergovernmental panel on climate change, 2, 2391, 2021.

Nalbant, M. A. and Sharma, S.: Investigating the impact of climate change on river ice thickness across the Northern Belt of United States, *ISH Journal of Hydraulic Engineering*, 29, 217–231, <https://doi.org/10.1080/09715010.2022.2050309>, 2023.

925 Newton, A. M. and Mullan, D. J.: Climate change and Northern Hemisphere lake and river ice phenology from 1931–2005, *The Cryosphere*, 15, 2211–2234, 2021.

Park, H., Yoshikawa, Y., Oshima, K., Kim, Y., Ngo-Duc, T., Kimball, J. S., and Yang, D.: Quantification of warming climate-induced changes in terrestrial Arctic river ice thickness and phenology, *Journal of Climate*, 29, 1733–1754, 2016.

Pavelsky, T. M. and Zametske, J. P.: Rapid decline in river icings detected in Arctic Alaska: Implications for a changing hydrologic cycle and river ecosystems, *Geophysical Research Letters*, 44, 3228–3235, <https://doi.org/10.1002/2016GL072397>, 930 2017.

Podkowa, A., Kugler, Z., Nghiem, S. V., and Brakenridge, G. R.: Ice Freeze-Up and Break-Up in Arctic Rivers Observed With Satellite L-Band Passive Microwave Data From 2010 to 2020, *Water Resources Research*, 59, e2022WR031939, <https://doi.org/10.1029/2022WR031939>, 2023.

935 Prowse, T., Alfredsen, K., Beltaos, S., Bonsal, B. R., Bowden, W. B., Duguay, C. R., Korhola, A., McNamara, J., Vincent, W. F., Vuglinsky, V., Walter Anthony, K. M., and Weyhenmeyer, G. A.: Effects of Changes in Arctic Lake and River Ice, *AMBIO*, 40, 63–74, <https://doi.org/10.1007/s13280-011-0217-6>, 2011.

Prowse, T. D. and Bonsal, B. R.: Historical trends in river-ice break-up: a review, *Hydrology Research*, 35, 281–293, 2004.

Prowse, T. D., Wrona, F. J., Reist, J. D., Gibson, J. J., Hobbie, J. E., Lévesque, L. M., and Vincent, W. F.: Climate change effects on hydroecology of Arctic freshwater ecosystems, *AMBIO: A Journal of the Human Environment*, 35, 347–358, 2006.

940 Prowse, T. D., Bonsal, B. R., Duguay, C. R., and Lacroix, M. P.: River-ice break-up/freeze-up: a review of climatic drivers, historical trends and future predictions, *Annals of Glaciology*, 46, 443–451, 2007.

Rokaya, P., Morales-Marín, L., Bonsal, B., Wheeler, H., and Lindenschmidt, K.-E.: Climatic effects on ice phenology and ice-jam flooding of the Athabasca River in western Canada, *Hydrological Sciences Journal*, 64, 1265–1278, <https://doi.org/10.1080/02626667.2019.1638927>, 945 2019.

Saraniemi, M., Huusko, A., and Tahkola, H.: Spawning migration and habitat use of adfluvial brown trout, *Salmo trutta*, in a strongly seasonal boreal river., *Boreal environment research*, 13, 2008.

Schwartz, M. D., Ahas, R., and Aasa, A.: Onset of spring starting earlier across the Northern Hemisphere, *Global Change Biology*, 12, 343–351, <https://doi.org/10.1111/j.1365-2486.2005.01097.x>, 2006.

950 Sen, P. K.: Estimates of the regression coefficient based on Kendall's tau, *Journal of the American statistical association*, 63, 1379–1389, 1968.

Sharma, S., Magnuson, J. J., Batt, R. D., Winslow, L. A., Korhonen, J., and Aono, Y.: Direct observations of ice seasonality reveal changes in climate over the past 320–570 years, *Sci Rep*, 6, 25061, <https://doi.org/10.1038/srep25061>, 2016.

Sharma, S., Filazzola, A., Nguyen, T., Imrit, M. A., Blagrove, K., Bouffard, D., Daly, J., Feldman, H., Feldsine, N., and Hendricks-Franssen, H.-J.: Long-term ice phenology records spanning up to 578 years for 78 lakes around the Northern Hemisphere, *Scientific Data*, 9, 318, 2022.

955

Shen, H. T.: River Ice Processes, in: *Advances in Water Resources Management*, edited by: Wang, L. K., Yang, C. T., and Wang, M.-H. S., Springer International Publishing, Cham, 483–530, https://doi.org/10.1007/978-3-319-22924-9_9, 2016.

Shiklomanov, A. I. and Lammers, R. B.: River ice responses to a warming Arctic—recent evidence from Russian rivers, *Environmental Research Letters*, 9, 035008, 2014.

Shiklomanov, A. I., Lammers, R. B., Rawlins, M. A., Smith, L. C., and Pavelsky, T. M.: Temporal and spatial variations in maximum river discharge from a new Russian data set, *Journal of geophysical research: Biogeosciences*, 112, 2007.

Siegel, J. E., Fullerton, A. H., and Jordan, C. E.: Accounting for snowpack and time-varying lags in statistical models of stream temperature, *Journal of Hydrology X*, 17, 100136, 2022.

965 Spearman, C.: The Proof and Measurement of Association between Two Things, *The American Journal of Psychology*, 15, 72, <https://doi.org/10.2307/1412159>, 1904.

St. Jacques, J.-M. and Sauchyn, D. J.: Increasing winter baseflow and mean annual streamflow from possible permafrost thawing in the Northwest Territories, Canada, *Geophysical Research Letters*, 36, 2009.

Stone, R. S., Dutton, E. G., Harris, J. M., and Longenecker, D.: Earlier spring snowmelt in northern Alaska as an indicator of climate change, *J. Geophys. Res.*, 107, <https://doi.org/10.1029/2000JD000286>, 2002.

970 SYKE: Daily River Discharge Measurements, Oulanka Research Station, 2026.

Takács, K. and Kern, Z.: Multidecadal changes in the river ice regime of the lower course of the River Drava since AD 1875, *Journal of Hydrology*, 529, 1890–1900, 2015.

Theil, H.: A rank-invariant method of linear and polynomial regression analysis, *Indagationes mathematicae*, 12, 173, 1950.

975 Von Storch, H.: Misuses of Statistical Analysis in Climate Research, in: *Analysis of Climate Variability*, edited by: Von Storch, H. and Navarra, A., Springer Berlin Heidelberg, Berlin, Heidelberg, 11–26, https://doi.org/10.1007/978-3-662-03744-7_2, 1999.

Vuglinsky, V.: Assessment of changes in ice regime characteristics of Russian lakes and rivers under current climate conditions, *Natural Resources*, 8, 416, 2017.

980 Vuglinsky, V. and Valatin, D.: Changes in ice cover duration and maximum ice thickness for rivers and lakes in the Asian part of Russia, *Natural Resources*, 9, 73, 2018.

Wang, X. and Feng, L.: Patterns and Trends in Northern Hemisphere River Ice Phenology from 2000 to 2021, *Remote Sensing of Environment*, 313, 114346, <https://doi.org/10.1016/j.rse.2024.114346>, 2024.

Yang, X., Pavelsky, T. M., and Allen, G. H.: The past and future of global river ice, *Nature*, 577, 69–73, 2020.

985