

# Enhancing particle number concentration modelling accuracy in China by incorporating various nucleation parameterization schemes into the CMAQ version 5.3.2 model

5 Jianjiong Mao<sup>1,2</sup>, Lei Jiang<sup>1</sup>, Zhicheng Feng<sup>1</sup>, Jingyi Li<sup>1</sup>, Yanhong Zhu<sup>1</sup>, Momei Qin<sup>1</sup>,  
Song Guo<sup>2,3</sup>, Min Hu<sup>2,3</sup>, Jianlin Hu<sup>1,3</sup>

<sup>1</sup> Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution Control, School of Environmental Science and Engineering, Nanjing University of Information Science & Technology, Nanjing 210044, China

10 <sup>2</sup> State Key Laboratory of Regional Environment and Sustainability, International Joint Laboratory for Regional Pollution Control, Ministry of Education (IJRC), College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

<sup>3</sup> Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, Nanjing University of Information Science & Technology, Nanjing 210044, China

15

20

*Correspondence to:* Jianlin Hu (jianlinhu@nuist.edu.cn)

25

**Abstract.** Particle number concentration (PNC) is a key parameter for assessing particles' effects on public health and climate. Although the Community Multiscale Air Quality (CMAQ) model has been widely used to investigate the particle mass concentrations in China, its capability of reproducing the PNC is unclear. After applying the default CMAQ model (version 5.3.2) in Beijing and Nanjing, the model underestimates PNC by 70% to 80% with the default Binary Homogeneous Nucleation (BHN) parameterization scheme. We then implement Ternary Homogeneous Nucleation (THN), Ion Mediated Nucleation (IMN), and ~~sulfuric acid~~-dimethylamine (DMA) nucleation parameterization schemes into the CMAQ model. Seven modelling scenarios are conducted to explore the model's performance on PNC. The results indicate that the scenario with combined IMN and DMA scheme (SumID scheme) yields the best agreement with the observations, enhancing PNC during new particle formation (NPF) events by 84% in Beijing and 36% in Nanjing. In SumID scheme, IMN contributes to 56.30% of the number concentrations in Beijing and 27.86% in Nanjing, while DMA accounts for 28.15% in Beijing and 29.27% in Nanjing, respectively. IMN nucleation contributed PNC exhibits a pronounced diurnal variation with higher concentrations during the day and lower levels at night. DMA pathway predominantly influences NPF events during the morning and evening peak. This study enhances the model's capability to accurately simulate NPF events and underscores the significant influence of IMN and DMA nucleation on PNC in ~~eastern cities~~.

Deleted: H<sub>2</sub>SO<sub>4</sub>

Deleted: China

**Keywords:** New particle formation; nucleation parameterization; number concentration; ion mediated nucleation; sulfuric acid-dimethylamine nucleation.

**Synopsis:** CMAQ model incorporated with IMN and DMA nucleation schemes significantly improves its modeling accuracy for particle number concentrations in ~~eastern cities~~.

Deleted: China

## 1. Introduction

Particulate matter (PM) significantly impacts regional air quality (Fuzzi et al., 2015), global climate change (Ning et al., 2024; Zhang et al., 2024), and public health (Hu et al., 2017a). Particle number concentration (PNC) is a critical parameter for assessing its environmental health and climate impacts.

New particle formation (NPF), a process ~~in which gaseous vapors in the atmosphere form critical molecular clusters that~~ grow into particles (Zhang et al., 2012; Wang et al., 2017), is a significant source of PNC, contributing ~30% of PNC at the surface and over 90% in the upper troposphere (Yu et al., 2020b; Zhao et al., 2024).

Deleted: that

Deleted: and

Laboratory and ambient observational studies have been conducted in various locations and environments to investigate the NPF processes. Sulfuric acid ( $\text{H}_2\text{SO}_4$ ), ammonia ( $\text{NH}_3$ ), ion, organic vapors, dimethylamine, nitric acid ( $\text{HNO}_3$ ), and iodic acid ( $\text{HIO}_3$ )/iodous acid ( $\text{HIO}_2$ ) are found to participate in the particle nucleation. A few nucleation theories and parameterizations have been proposed, such as  $\text{H}_2\text{SO}_4$ -water ( $\text{H}_2\text{O}$ ) Binary Homogeneous Nucleation (BHN) (Kulmala et al., 1998; Vehkamäki, 2002; Yu, 2006a, 2007),  $\text{H}_2\text{SO}_4$ - $\text{NH}_3$ - $\text{H}_2\text{O}$  Ternary Homogeneous Nucleation (THN) (Napari, 2002; Napari et al., 2002; Merikanto et al., 2007; Almeida et al., 2013), Ion Induced Nucleation (IIN) (Laakso, 2002; Liu et al., 2022), Ion Mediated Nucleation (IMN) (Yu and Turco, 1997, 2000, 2001; Yu, 2002, 2006b, 2010; Yu et al., 2008; Yu et al., 2010; Yu et al., 2018; Yu et al., 2020a),  $\text{H}_2\text{SO}_4$ -dimethylamine nucleation (DMA) (Loukonen et al., 2010; Kurten et al., 2014; Ruusuvoori et al., 2015; Olenius et al., 2017; Julin et al., 2018; Lu et al., 2020; Liu et al., 2021a; Liu et al., 2021b; Wang et al., 2021b), organic nucleation (Zhang et al., 2004; Fan et al., 2006; Wang et al., 2015; Yan et al., 2021),  $\text{HNO}_3$ - $\text{H}_2\text{SO}_4$ - $\text{NH}_3$  nucleation (Wang et al., 2022b) and  $\text{HIO}_3$ / $\text{HIO}_2$  nucleation (Zhang et al., 2022; Zhao et al., 2024).

Different nucleation theories are applicable to various environments, and multiple nucleation theories are interrelated and nucleate synergistically in some cases. For example, the IMN theory takes into account BHN, as well as the synergistic effects between THN and ions (Yu et al., 2018; Yu et al., 2020a).

Based on these studies, various nucleation parameterization schemes, which calculate nucleation rates based on precursor concentrations and environmental factors, have been developed and implemented into three-dimensional (3-D) chemical transport models (CTMs) to study NPF events or PNC in the atmosphere. The WACCM model shows that BHN underestimates number concentration in the upper

troposphere and lower stratosphere (English et al., 2011). The GEOS-Chem-APM model, utilizing the IMN parameters, suggests that IMN predominantly influences PNC in the troposphere (Yu et al., 2010). The NAQPMS-APM-VBS model demonstrates that organic nucleation can reproduce the summer NPF events in Beijing (Chen et al., 2019). By coupling all the different forms of nucleation parameterization schemes discussed above, the E3SM model indicates that H<sub>2</sub>SO<sub>4</sub>-dimethylamine nucleation dominates number concentration in 1 km height (Zhao et al., 2024). Based on this result, the Weather Research and Forecasting model coupled with chemistry (WRF-chem), established the same nucleation parameterization schemes, proves that HIO<sub>3</sub> nucleation is the main nucleated way in the future (Ning et al., 2024).

**Deleted:** The E3SM model shows that H<sub>2</sub>SO<sub>4</sub>-dimethylamine nucleation dominates number concentration in 1 km height.

**Deleted:** The WRF-chem model proves that HIO<sub>3</sub> nucleation is the main nucleated way in the future

The Community Multiscale Air Quality (CMAQ) model, known for its relatively comprehensive chemistry module, has been widely used for studying mass concentrations and sources of pollutants in China (Hu et al., 2016; Hu et al., 2017a; Hu et al., 2017b; Huang et al., 2018; Yang et al., 2019; Li et al., 2020; Ma et al., 2021; Mao, 2023; Li et al., 2024; Qin et al., 2025). However, its application in simulating number concentrations, particularly in modeling NPF, remains limited. In the CMAQ model version 4.0, the BHN parameterization by Kulmala et al. (1998) is utilized, while the CMAQ v5.0 incorporates the Vehkamäki (2002) BHN parameterization, which employs the Kelvin formula along with particle physics, hydrogen bond breakage, and surface tension to develop a theoretical equation. Studies using the default CMAQ model suggest BHN parameterization alone may not adequately resolve intense NPF events (Zhang et al., 2010b; Zhang et al., 2010c). Efforts have been made to incorporate other parameterization schemes into the CMAQ model. For examples, CMAQ simulations in the Northwest Pacific indicates that binary or ternary nucleation theories cannot fully reproduce all PNC and commonly underpredict number concentrations by approximately one order of magnitude on average (Elleman and Covert, 2009). Previous studies using the CMAQ also have compared four different nucleation theories, revealing that discrepancies in simulating PNC could exceed three orders of magnitude, with notable regional variations (Zhang et al., 2010a; Zhang et al., 2010c). When binary nucleation included secondary condensable organics, CMAQ predicts PNC consistent with daytime measurements in Northeast Houston in 2004 (Fan et al., 2006). Currently, there are no studies evaluating the effects of number concentration simulations of IMN and DMA in China.

**Deleted:** numerical

In this study, we first evaluate the performance of the default CMAQ v5.3.2 model on predicting PNC in Beijing and Nanjing during two NPF periods, and then implement three additional nucleation parameterization schemes into the model, aiming to improve CMAQ's accuracy and to better understand the key nucleation pathways in different urban regions of China.

Deleted: episodes,

## 120 2. Materials and Methods

### 2.1 Model description

In the default aerosol mode of CMAQ v5.3.2, the BHN nucleation parameterization of Vehkamäki (2002) is used (Zhang et al., 2010c). This parameterization is based on rigorous nucleation kinetics and the thermodynamically consistent version of the classical binary homogeneous nucleation theory. For simplified calculation, Vehkamäki (2002) fitted a polynomial form, which can be put in the aerosol mode in CTMs models. In this way, relative humidity ( $RH$ ), air temperature ( $T$ ), and  $H_2SO_4$  ( $N_a$ ) affect the NPF rates in the formula. The nucleation rate  $J$  is expressed as the exponent of the third-order polynomial of  $\ln(RH/100)$  and  $\ln(N_a)$ :

$$\begin{aligned}
 J [1 / (\text{cm}^3 \text{ s})] = & \exp \left\{ a(T, x^*) + b(T, x^*) \ln \frac{RH}{100} \right. \\
 & + c(T, x^*) \ln^2 \frac{RH}{100} + d(T, x^*) \ln^3 \frac{RH}{100} \\
 & + e(T, x^*) \ln(N_a) + f(T, x^*) \ln \frac{RH}{100} \ln(N_a) \\
 & + g(T, x^*) \ln^2 \frac{RH}{100} \ln(N_a) + h(T, x^*) [\ln(N_a)]^2 \\
 & \left. + i(T, x^*) \ln \frac{RH}{100} [\ln(N_a)]^2 + j(T, x^*) [\ln(N_a)]^3 \right\}, \quad (1)
 \end{aligned}$$

where the coefficients  $a(T, x^*) \dots j(T, x^*)$  is a function of temperature and the critical cluster mole fraction  $x^*$ .  $x^*$  is given as:

$$\begin{aligned}
 x^* = & 0.740997 - 0.00266379T \\
 & - 0.00349998 \ln(N_a) + 0.0000504022T \ln(N_a) \\
 & + 0.00201048 \ln \frac{RH}{100} - 0.000183289T \ln \frac{RH}{100} \\
 & + 0.00157407 \ln^2 \frac{RH}{100} - 0.0000179059T \ln^2 \frac{RH}{100} \\
 & + 0.000184403 \ln^3 \frac{RH}{100} - 1.50345 \cdot 10^{-6} T \ln^3 \frac{RH}{100}, \quad (2)
 \end{aligned}$$

Previous studies (Yu et al., 2017; Yan et al., 2018; Yan et al., 2021) have found that NPF events exhibit different nucleation mechanisms under varying conditions. Our initial application of the default CMAQ v5.3.2 model reveals that the BHN parameterization significantly underpredicts number concentration in both Beijing and Nanjing (Mao, 2023). Therefore, we implement three additional nucleation parameterizations, i.e., THN (Zhao et al., 2021a), IMN (Yu et al., 2018), DMA (Zhao et al., 2021a), into the CMAQ model nucleation parameterizations to enhance its ability to simulate PNC.

The IMN nucleation parameterization is affected by  $\text{H}_2\text{SO}_4$ ,  $\text{NH}_3$ ,  $RH$ ,  $T$ , ionization rate in the atmosphere, and aerosol surface area. The ionization rate is calculated based on the look-up table provided in Usoskin and Kovaltsov (2006), which accounts for factors include latitude, longitude, surface type, surface pressure, and air pressure. This comprehensive approach allows for a detailed accounting of ion concentrations involved in nucleation within the atmospheric environment (Yu and Turco, 1997, 2000, 2001). For investigating the nucleation processes involving ion clusters, a kinetic model that integrates new thermodynamic data, physical algorithms, and observational insights has been employed to calculate the IMN nucleation speeds (Yu, 2010). To incorporate this method into 3D-CTMs model, Yu et al. (2020a) developed a 6-D look-up table, which is used in this study to facilitate fast calculations. This look-up table encompasses the considerations of BHN, Binary IMN (BIMN), THN, and Ternary IMN (TIMN). In this study, we specifically employ the TIMN module (Table S1). The details about IMN nucleation parameterization are shown in Yu et al. (2018).

THN and DMA nucleation parameterizations used in this study are derived from Zhao et al. (2021a), which were developed and updated in several previous studies (Riccobono et al., 2014; Kirkby et al., 2016; Dunne et al., 2016; Gordon et al., 2017). In THN parameterizations,  $\text{H}_2\text{SO}_4$ ,  $\text{NH}_3$ , and  $T$  are considered, while  $RH$  is not included. In earlier classical binary and ternary nucleation parameterizations,  $RH$  is indispensable, which are based on classical kinetics and thermodynamic theory. The THN nucleation parameterization is given as:

$$J_{\text{THN}} = k_{\text{in}}(T) \cdot f_{\text{n}}([\text{NH}_3], [\text{H}_2\text{SO}_4]) \cdot [\text{H}_2\text{SO}_4]^{2.891024}, \quad (3)$$

$$\ln k_{\text{in}} = 182.4495 - \exp(1.203451 \times (\frac{T}{1000} + 4.188065)), \quad (4)$$

$$f_{\text{n}}([\text{NH}_3], [\text{H}_2\text{SO}_4]) = [\text{NH}_3] / (1.5703478 \times 10^{-6} + [\text{H}_2\text{SO}_4]^{2.891024} / [\text{NH}_3]^{8.003471}), \quad (5)$$

**Deleted:** includes BHN, Binary IMN (BIMN), THN, and Ternary IMN (TIMN) are all considered

**Deleted:** thermodynamically

The units for NH<sub>3</sub> and H<sub>2</sub>SO<sub>4</sub> concentrations are 10<sup>6</sup> cm<sup>-3</sup>, while T unit is Kelvin (K).

The DMA nucleation parameterization is affected by H<sub>2</sub>SO<sub>4</sub> and dimethylamine, without considering T and RH. The nucleation parameterization is given as:

$$J_{\text{DMA}} = 1.93 \cdot 10^{-28} ([\text{dimethylamine}] / (2.5 \cdot 10^7))^{4.36} [\text{H}_2\text{SO}_4]^{3.7}, \quad (6)$$

while dimethylamine and H<sub>2</sub>SO<sub>4</sub> concentrations units are cm<sup>-3</sup>. These four nucleation parameterizations have been implemented into CMAQ model.

## 2.2 Model applications

180 The revised CMAQ v5.3.2, configured with the gas-phase mechanism of SAPRC07tic and the aerosol module of AERO6i, is employed in this study to simulate NPF events. [These settings are currently the best choice for CMAQ to simulate gas precursors and aerosol chemistry \(Li et al., 2022a\)](#). The modeling domain covers China and the surrounding countries and regions with 197 × 127 grids shown in Figure S1a, with 36 km × 36 km horizontal grid resolution. The vertical resolution is 18 layers. The initial and boundary conditions are provided by the default profiles of CMAQ model. The simulated periods are 185 from March 9 to April 5 in 2016 and from April 13 to 20 in 2018 with the first two days as spin-up, reducing the effects of the initial conditions on the simulated results.

The meteorological fields are simulated by WRF v4.2.1 with the FNL reanalysis dataset. The FNL data are obtained from the U.S. National Centre for Atmospheric Research (NCAR), with a spatial resolution of 1.0° × 1.0° (<http://rda.ucar.edu/datasets/ds083.2/>). The physical parameterizations used in WRF 190 include the Thompson microphysical scheme; RRTMG longwave/shortwave radiation scheme; Noah land-surface scheme; MYJ boundary layer scheme; and modified Tiedtke cumulus parameterization scheme. The detailed configuration settings could be found in previous studies (Hu et al., 2016; Wang et al., 2021a; Mao et al., 2022).

195 The Multi-resolution Emission Inventory for China version 1.3 (MEIC v1.3) (Zheng et al., 2018), and Regional Emission inventory in ASia version 3.2 (REAS v3.2) (Kurokawa and Ohara, 2020) are used to provide the anthropogenic emissions. MEIC served as the anthropogenic emissions from China, and REAS served as the anthropogenic emissions from neighboring countries and regions. Biogenic emissions are generated using the Model for Emissions of Gases and Aerosols from Nature version 2.1

(MEGAN v2.1) (Guenther et al., 2012) for the simulated period. The open biomass burning emissions  
200 are processed using the Fire INventory for NCAR (FINN) during the study period (Wiedinmyer et al.,  
2011). In the CMAQ model, the PNC is modeled based on the assumption that it follows a three-  
lognormal distribution: Aitken-mode, Accumulation-mode, and Coarse-mode. The current emission  
inventory only has mass emissions of PM<sub>2.5</sub> (PM < 2.5 μm) and PM<sub>2.5-10</sub> (2.5 μm < PM < 10 μm) and  
does not contain emission information on the particle number emissions. The CMAQ model assumes  
205 that 99.9% of PM<sub>2.5</sub> mass in the accumulation mode and 0.1% in the Aitken mode. All mass emissions  
of PM<sub>2.5-10</sub> are in the coarse model. Then the mass emissions are converted to number emissions based  
on the size distribution parameters.

To estimate the dimethylamine concentrations, we adopt the method used in the study of Yu and Luo  
(2014) to include the emissions, chemical reactions and deposition of dimethylamine into the CMAQ  
210 model. We calculate the dimethylamine emissions by using the ratio of dimethylamine to NH<sub>3</sub> fluxes,  $6.6 \times 10^{-4}$   
(Gunnar and Paul, 1995) and approximate the spatial distribution and seasonal variations of  
dimethylamine emissions based on those of NH<sub>3</sub>. In the CMAQ model, we include the oxidation of  
dimethylamine by hydroxyl radicals (OH) and the reaction rate coefficient is  $6.52 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$   
(Carl, 1998). The dry and wet deposition, as well as horizontal and vertical transport, is also  
215 considered, following the approaches used for NH<sub>3</sub>.

We conduct seven simulations configured with different nucleation schemes (Table 1). In addition to  
individual simulations of the BHN, THN, IMN, and DMA parameterizations, we conduct one simulation  
with no nucleation (None), and one simulation considering the BHN, THN, and DMA parameterizations  
together (SumBTD), and another considering the IMN and DMA parameterizations together (SumID).

220 The rationale behind these combinations lies in the different nucleation theories. Since BHN represents  
the interaction between H<sub>2</sub>SO<sub>4</sub> and H<sub>2</sub>O, it is a fundamental theory in nucleation mechanisms that exists  
in most environments (Sipilä et al., 2010). THN represents the involvement of H<sub>2</sub>SO<sub>4</sub>, H<sub>2</sub>O, and NH<sub>3</sub> in  
nucleation, used to explain the higher nucleation rates in the atmosphere (Merikanto et al., 2007).  
Therefore, the binary and ternary nucleation scenarios are interconnected. But IMN nucleation accounts  
225 for the synergistic interactions among BHN, THN, and ions (Yu et al., 2018; Yu et al., 2020a).  
Consequently, this nucleation mechanism cannot be integrated with BHN and THN scenarios. Given that



230 DMA nucleation, containing H<sub>2</sub>SO<sub>4</sub> and dimethylamine, has been identified as the predominant nucleation mechanism in urban environments (Yao et al., 2018; Liu et al., 2021b; Wang et al., 2021b), this study specifically addresses urban nucleation mechanism. Therefore, the DMA nucleation scheme must be incorporated into the combined scenarios. Based on the discussed above, we have retained only two combined scenarios for further analysis. The “None” nucleation scheme means that the new particle formation is turned off and the PM number concentration is simulated without NPF events. All other simulations setting is performed with the same model configurations and inputs described above.

**Table 1. Experiment configuration.**

Experiment	Description
None	Without NPF mode
BHN	With BHN nucleation parameterization
THN	With THN nucleation parameterization
IMN	With IMN nucleation parameterization
DMA	With DMA nucleation parameterization
SumBTD	With BHN, THN and DMA nucleation parameterizations
SumID	With IMN and DMA nucleation parameterizations

235

### 2.3 Observation data

Observed number concentration were collected in Beijing (location in the Northern China) and in Nanjing (located in Southern China). Observation in Beijing is derived from previous studies (Cai et al., 2017; Wang et al., 2023) during March 11 to April 5, 2016. The measurement site is located on the campus of Tsinghua University (116°28'E, 39°48'N) (Figure S1b). The diethylene glycol scanning mobility particle spectrometers (DEG-SMPS) was used to measure sub-5 nm particle size distributions and a particle size distribution system (including a TSI aerodynamic particle sizer and two parallel SMPSs) was used to measure size distributions of particles from 3 nm (in electrical mobility diameter) to 10 μm (Liu et al., 2014; Cai et al., 2017). The observation result is shown in Figure S2, indicate 11 NPF events during the observation period. PNC observations in Nanjing were conducted from April 15 to 20, 2018, as reported by previous study (Lai et al., 2022), at the SORPES site (118°57'10"E,

245

32°07'14"N) in the Xianlin Campus of Nanjing University (Figure S1c). This period has 4 NPF events (15th to 17th, 19th). The two sets of instruments were employed to measure number concentration: a Differential Mobility Particle Sizer (DMPS) covering the size range of 6–800 nm, and a Particle Size Magnifier (PSM) for the 1–6 nm size distribution (Qi et al., 2015).

### 3. Results and Discussion

#### 3.1 Evaluation of PNC


Figure 1 presents the comparison of hourly predicted and observed PNC in Beijing and Nanjing. The default BHN nucleation parameterization significantly underestimates PNC in both Beijing 2016 and Nanjing 2018 periods. In the Beijing 2016 episode, the BHN parameterization yields normalized mean bias (NMB) of -0.70, normalized mean error (NME) of 0.70, and a low correlation coefficient (R) of 0.02. In the Nanjing 2018 episode, the CMAQ model with the default BHN has a NMB of -0.78, NME of 0.83, and R of -0.26. These indices are very similar to the 'None' case in which no NPF scheme is considered, indicating that BHN contributes marginally to PNC in these periods.

The THN, IMN, and DMA parameterization scheme individually improves the CMAQ performance on PNC. In Beijing, the THN parameterization yields NMB of -0.20, NME of 0.42, and a higher R of 0.60. The IMN parameterization has a lower NMB of -0.04, NME of 0.40 and R of 0.62. Although the DMA parameterization is suggested to be considered in urban (Cai et al., 2021; Liu et al., 2021b), it yields a low NMB value of -0.20, but the R value is also low (0.04). In Nanjing, the THN parameterization yields NMB of -0.51, NME of 0.71, R of 0.25. The IMN parameterization has a lower NMB of -0.47, NME of 0.70 and R of 0.38. The DMA parameterization exhibits NMB of -0.56 and NME of 0.83 along with a low R of -0.13. These schemes all demonstrate improvements compared to the None and BHN schemes, and the performance in Beijing with these schemes is generally better than in Nanjing.

In combined cases, the SumBTD has NMB of 0.07, NME of 0.56, R of 0.30 and the SumID has NMB of 0.16, NME of 0.55, R of 0.37 in Beijing. The SumBTD has NMB of -0.44, NME of 0.74, R of 0.13 and the SumID has NMB of -0.42, NME of 0.73, R of 0.27 in Nanjing. The SumID produce better results than SumBTD in general. These findings indicate that accurately capturing and resolving each NPF event under the current parameterization remains challenging. Nonetheless, compared to the default scheme, it

reduces discrepancies in magnitude, and is better to previous studies (Elleman and Covert, 2009; Zhang  
275 et al., 2010c).

Figure 2 shows the predicted and observed number concentrations in  $PM_1$  ( $PM < 1.0 \mu m$ ), Aitken-mode  
and Accumulation-mode during the whole sampling periods. The number concentration in  $PM_1$   
encompasses both the Aitken mode and the Accumulation mode. The Aitken mode pertains to particles  
280 smaller than  $0.1 \mu m$  and is more influenced by the nucleation parameterization schemes. Based on the  
definition of NPF events in previous studies (Wu et al., 2007; Guo et al., 2014; Wang et al., 2016; Peng  
et al., 2021), which observe a notable increase in PNC, we infer that the high-value data points depicted  
in the Figure 2 primarily correspond to the NPF events.




285

**Figure 1. Model performance of PM<sub>1</sub> number concentration in (1) Beijing and (2) Nanjing f. Each point corresponds to a 1-h average value. (a) None scenario. (b) BHN scenario. (c) THN scenario. (d) IMN scenario. (e) DMA scenario. (f) SumBTD scenario. (g) SumID scenario. The 1:1, 1:2, and 2:1 line are shown. NMB (red font): normalized mean bias; NME (blue font): normalized mean error; R (black font): correlation coefficient.**

290


In Beijing 2016 (Figure 2a), BHN underpredicts total number concentrations over an order of magnitude ( $1 \times 10^3 \text{ cm}^{-3}$ ), failing to capture intense NPF events ( $7 \times 10^4 \sim 2 \times 10^5 \text{ cm}^{-3}$ ). THN has a significant



Deleted:

improvement in PNC simulation although it underpredicts in low concentration ( $< 1 \times 10^4 \text{ cm}^{-3}$ ); DMA  
295 exhibit superior performance in capturing extreme values ( $> 1 \times 10^5 \text{ cm}^{-3}$ ), which are indicative of intense  
NPF events; THN and IMN has similar performance ( $3 \times 10^3 \sim 1 \times 10^5 \text{ cm}^{-3}$ ). Due to SumBTD and SumID  
considering multiple nucleation parameterizations, they provide improved simulations compared to other  
cases ( $8 \times 10^3 \text{ cm}^{-3} \sim 2 \times 10^5 \text{ cm}^{-3}$ ). However, SumBTD may be overestimated at high values or  
underestimated at low values relative to SumID and observational data, which makes SumID the best  
300 choice in these scenarios. Similarly, in Nanjing 2018 (Figure 2b), BHN significantly underpredicts PNC  
over an order of magnitude ( $1 \times 10^3 \text{ cm}^{-3}$ ). In contrast, THN, IMN and DMA display similar and great  
simulation performance ( $4 \times 10^3 \text{ cm}^{-3} \sim 3 \times 10^4 \text{ cm}^{-3}$ ). SumBTD and SumID nucleation parameterizations  
perform similarly and better than others ( $7 \times 10^3 \text{ cm}^{-3} \sim 9 \times 10^4 \text{ cm}^{-3}$ , Observation:  $7 \times 10^3 \text{ cm}^{-3} \sim 2 \times 10^5 \text{ cm}^{-3}$ ).  
Overall, SumID is capable of effectively capturing both common and intense NPF events in  $\text{PM}_{10}$   
305 number concentration. We subsequently use the SumID results to analyze the relative contributions of  
IMN and DMA parameterizations to PNC in the section 3.2.

In the Aitken-mode simulations for Beijing in 2016 (Figure 2c), BHN significantly underpredicts Aitken-  
mode concentration ( $2 \times 10^3 \text{ cm}^{-3} \sim 1 \times 10^4 \text{ cm}^{-3}$ ). The IMN, SumBTD and SumID demonstrate superior  
performance ( $7 \times 10^3 \text{ cm}^{-3} \sim 2 \times 10^5 \text{ cm}^{-3}$ , Observation:  $5 \times 10^3 \text{ cm}^{-3} \sim 2 \times 10^5 \text{ cm}^{-3}$ ). In Nanjing (Figure 2d),  
310 SumID performs similarly to SumBTD ( $4 \times 10^3 \text{ cm}^{-3} \sim 6 \times 10^4 \text{ cm}^{-3}$ , Observation:  $4 \times 10^3 \text{ cm}^{-3} \sim 2 \times 10^5 \text{ cm}^{-3}$ ),  
and better than others. Current research indicates that the involvement of amine gases (e.g.  $\text{NH}_3$ ,  
dimethylamine) (Kirkby et al., 2023; Ning et al., 2024; Feng et al., 2025) and atmospheric ions (Yu et al.,  
2018; Yu et al., 2020a) can promote the NPF. The main reason for the underestimation of BHN is that  
this nucleation scheme only considers  $\text{H}_2\text{SO}_4$  and  $\text{H}_2\text{O}$ , while SumID takes  $\text{H}_2\text{SO}_4$ , amine gases and ions  
315 into account simultaneously. For Accumulation-mode number concentrations, the seven nucleation  
parameterizations perform much better and align closely with observations in both Beijing (Figure 2e)  
( $2 \times 10^2 \text{ cm}^{-3} \sim 5 \times 10^3 \text{ cm}^{-3}$ , Observation:  $1 \times 10^2 \text{ cm}^{-3} \sim 1 \times 10^4 \text{ cm}^{-3}$ ) and Nanjing (Figure 2f) ( $2 \times 10^2 \text{ cm}^{-3}$   
 $\sim 5 \times 10^3 \text{ cm}^{-3}$ , Observation:  $5 \times 10^2 \text{ cm}^{-3} \sim 8 \times 10^3 \text{ cm}^{-3}$ ). The reason is that the Accumulation mode is  
primarily influenced by emissions, with only a small fraction of particulate matter being affected by the  
320 formation and growth of new particles.



**Figure 2. Comparison of (a), (b)  $PM_{10}$ , (c), (d) Aitken-mode, and (e), (f) Accumulation-mode number concentration in seven scenarios with observations during NPF events in (a, c, e) Beijing and (b, d, f) Nanjing. The dashed blue line is the median value, and red prism is the mean value.**

325

### 3.2 Contribution of different parameterizations to PNC


Figure 3 illustrates the observed hourly PNC alongside four different schemes: None, IMN, DMA, and SumID. As shown in Figure 3a, the None scheme significantly underestimates the PNC in Beijing.

330

Incorporating the DMA or IMN nucleation parameterizations markedly improves the modeled PNC. The IMN scheme, in particular, demonstrates greater improvement, with differences between the simulated and observed values remaining within  $5 \times 10^4 \text{ cm}^{-3}$  in Beijing. However, these schemes still fail to effectively capture the majority of NPF events. When both the IMN and DMA pathways are considered,


335

the SumID scheme successfully reproduces 11 NPF events (Figure S2), maintaining discrepancies within  $3 \times 10^4 \text{ cm}^{-3}$  during the observation periods. In Nanjing, the None scheme fails to adequately simulate number concentration, underestimating it by an order of magnitude, as shown in Figure 2b. The incorporation of the DMA parameterization scheme improves the simulation of number concentration during morning and evening peaks (especially on the April 15 and 19). By including the IMN scheme,



340 the model captures elevated daytime values on NPF days, aligning with observed peak times. Nonetheless, both schemes continue to underestimate concentrations ( $> 1 \times 10^5 \text{ cm}^{-3}$ ). When these two ways integrated (SumID), the model successfully replicates the trends of NPF events on the April 15 and 19, although it still underestimates concentrations by more than  $1 \times 10^5 \text{ cm}^{-3}$ . Among them, the events on the 16th and 17th are not captured by the model (Figure 3b).


345 The simulation results from Beijing and Nanjing underscore the necessity of considering both IMN and DMA nucleation processes. Furthermore, they suggest that nucleation mechanisms may vary between regions, indicating the importance of employing multiple nucleation parameterization schemes simultaneously. These findings demonstrate that, under the current set of seven schemes, accurately identifying and resolving each NPF event remains challenging, as shown in Figures 1 to Figure 3. Despite 350 these challenges, the adoption of the integrated approach (SumID scheme) significantly reduces discrepancies in magnitude compared to the default scheme. Consequently, due to its effectiveness in addressing these challenges, the SumID scheme is selected for further analysis.



355 **Figure 3. Observation and four schemes of None, IMN, DMA and SumID in hourly particle number concentrations ( $\text{cm}^{-3}$ ) in (a) Beijing and (b) Nanjing. Background shading represents nighttime periods.**

Figure 4 presents the distinct pathways of emissions, IMN, and DMA in contributing to the number concentration (%) within the SumID optimal scenario framework across the entire episode, as well as during NPF days and non-NPF days. Due to the inability to capture two NPF events on the [April 16th](#)


Deleted: of



Deleted:

and 17th in Nanjing, these events are excluded from the analysis. During the entire episode in Beijing (Figure 4a), emissions, IMN, and DMA contribute 15.55%, 56.30%, and 28.15% to the PNC, respectively. Therefore, IMN nucleation is the main nucleation way in Beijing. On NPF days (Figure 4b), the IMN nucleation pathway accounts for 61.31% of the PNC, while the DMA nucleation pathway contributes 31.23%. Conversely, on non-NPF days (Figure 4c), emissions represent 22.94% of the PNC. This indicates that although the conditions are considered non-NPF, nucleation still takes place. However, the NPF-related nanoparticles undergo rapid transformation through condensation and coagulation when interacting with high concentrations of particles from emissions with elevated ratios. This leaves no opportunity for the nanoparticles to grow over 10 nm (Wu et al., 2007; Guo et al., 2014; Wang et al., 2016; Peng et al., 2021). In Nanjing, across the entire episode, the contributions of emissions, IMN, and DMA pathways to the PNC are 42.87%, 27.86%, and 29.27%, respectively (Figure 4d). Since the majority of simulated days are NPF days, the IMN ratio during NPF days increases by only 4.33%, while emissions decrease by 2.47% (Figure 4e). Notably, the contribution of the DMA pathway decreases by 1.86% on NPF days but rises to 32.58% on non-NPF days (Figure 4f). The high ratios of primary emissions have a significant impact on nucleation pathways, accounting for 40.40% on NPF days and 47.23% on non-NPF days. Consequently, the proportions of the IMN and DMA nucleation pathways are constrained to 32.19% and 29.27% on NPF days, respectively, and 20.20% and 32.58% on non-NPF days. This phenomenon may also be attributed to a potential underestimation of NPF day simulations in Nanjing (Figure 3).

**Deleted:** However, the fine particles are rapidly removed due to condensation and coagulation with high concentrations of particles originating from emissions with elevated ratios




**Figure 4.** Pies show emission, IMN and DMA pathways contribution percentage (%) of SumID optimal scenario in whole episode, NPF days and non-NPF days.

Figure S3 illustrates the average diel variation of hourly contributions across entire periods, NPF days, and non-NPF days. The partitioning method is based on the days selected from the observed data. In Figure S3, IMN contributes a high number concentration in the daytime with a notable diurnal variation



390 both in Beijing and Nanjing across entire periods, NPF days, and non-NPF days, DMA dominates the  
 NPF events on the morning and evening peak, especially in Nanjing. At the night, emissions ratio  
 increases in Beijing and dominate over 90% in Nanjing. This is attributed to the low concentration of  
 $H_2SO_4$  in the absence of radiation (Lu et al., 2019) and DMA can promote nucleating in low  $H_2SO_4$   
 concentration (Guo et al., 2020; Liu et al., 2021b; Zhao et al., 2021b; Wang et al., 2022a). Figure 5 shows  
 395 the hourly averaged NPF rates of the two pathways during the NPF events in Beijing and Nanjing. During  
 Beijing NPF events (Figure 5a), IMN rates are generally higher than DMA, and dominates the SumID  
 nucleation rates in the daytime (Figure S3). In Nanjing, IMN and DMA pathways nucleation rates are  
 over  $10^{-2}$  on average (Figure 5b), while IMN has diurnal variation (Figure S3). The corresponding  
 contribution in number concentrations is shown in Figure 3.

400




**Figure 5.** NPF hourly particle nucleation rate ( $cm^{-3} s^{-1}$ ) of two nucleation parameterizations within SumID nucleation parameterization in (a) Beijing and (b) Nanjing. Background shading represents nighttime periods.

405

### 3.3 Discussion

As previously mentioned, IMN significantly impacts NPF events, contributing to over 50% of PNC in Beijing and 30% in Nanjing. During daytime periods, IMN dominates, accounting for over 90% of the PNC. In contrast, during morning and evening peak periods, DMA is more influential, comprising over  
 410 70% of the concentrations. These results are consistent with those of previous studies. The NAQPMS-

Deleted: during three types of days



Deleted:

APM model effectively captured the NPF events in Beijing during January 2006 (Chen et al., 2017) and September 2015 (Du et al., 2021), which can be attributed to IMN nucleation processes. The Geos-chem  
415 model with IMN nucleation processes also reproduced the NPF events in Beijing during March 2016 (Wang et al., 2023). Otherwise, variations in DMA nucleation parameters could influence PNC WRF-Chem modeling results in Beijing (Shen et al., 2024; Feng et al., 2025). The CMAQ model, which accounted for both IMN and DMA, reproduced NPF in Shanghai. The results indicated that DMA contributed to over 60% of the number concentration (Chang et al., 2023). The E3SM model,  
420 incorporating 11 nucleation pathways, indicated that eastern China was predominantly influenced by DMA nucleation at an altitude of 1 km, while H<sub>2</sub>SO<sub>4</sub>-NH<sub>3</sub>-H<sub>2</sub>O neutral and ion nucleation prevailed at altitudes of 2-6 km (Zhao et al., 2024). Hence, we conclude that the updated model results are reliable.

However, some issues persist with the updated model. First, the updated model can't capture all NPF events in Nanjing, especially on April 14. Previous studies have shown that H<sub>2</sub>SO<sub>4</sub>-organic nucleation  
425 (Chen et al., 2019; Zhao et al., 2024) and growth (Cai et al., 2016) are important in NPF modeling. We have not incorporated the parameterization for H<sub>2</sub>SO<sub>4</sub>-organic nucleation, which might result in underestimations in Nanjing. Second, observations (Guo et al., 2014; Wang et al., 2014; Lee et al., 2019; Cai et al., 2022; Kulmala et al., 2022; Kontkanen et al., 2022; Li et al., 2022b) found that the preexisted particulate, condensation sink, and growth rate will affect the new particle number concentration,  
430 especially under 10 nm. Some models have demonstrated that considering the microphysical processes of aerosols can significantly improve the accuracy of PNC simulations (Fountoukis et al., 2012; Zhao et al., 2024). Under high condensation sink, larger particles preferentially condense available vapors, thereby inhibiting the growth of smaller particles. Consequently, the small particles are eventually coagulated by the larger particles, thereby modifying the particle size distribution of the aerosols. But  
435 most of the CTMs don't have a relationship between condensation sink, growth rate, and new particle production module. Therefore, this can lead to an overprediction of PNCs under high condensation sink scenarios. Third, the emission of DMA gas was modeled using its ratio with NH<sub>3</sub>. Dry and wet deposition, along with horizontal and vertical transport, were handled using the same methodologies applied to NH<sub>3</sub>. This approach may have led to an overprediction of DMA gas concentrations. Last, the number emissions  
440 are converted from mass emissions based on the size distribution parameters. Currently, there are pan-European particle number emission inventories (Kulmala et al., 2011; Julin et al., 2018) and a U.S.

**Deleted:** Most of the CTMs don't have a relationship between condensation sink, growth rate, and new particle production module. In the CMAQ model, this can lead to an overprediction of PNCs under high condensation sink scenarios. In such cases, larger particles preferentially condense available vapors, thereby inhibiting the growth of smaller particles. Consequently, the small particles are eventually coagulated by the larger particles and subsequently lost from the system.

number emission inventory (Posner and Pandis, 2015). Recent research (Olin et al., 2022) indicates that accurately representing number emissions for particles is essential for simulating number concentrations effectively. Therefore, the number emission should be established in China in the future.

#### 4. Conclusion

455 After integrating multiple **nucleation** schemes into the aerosol module of CMAQ v5.3.2, including BHN, THN, IMN, DMA, SumBTD (BHN+THN+DMA), and SumID (IMN+DMA), the SumID nucleation scheme was identified as the optimal approach. This scheme enhances the simulation of number concentrations during periods in Beijing and Nanjing, particularly improving the representation of NPF events. Furthermore, the updated model can quantify the contributions of different nucleation pathways  
460 to total number concentrations and nucleation rates. ▼

**Deleted:** The results confirm the enhanced model's ability to accurately simulate NPF events and highlight the substantial impact of IMN and DMA nucleation on particle number concentrations in China.

The updated model can improve 36 % to 84 % number concentration simulation in Beijing and Nanjing during whole periods, whereas the default model, which employs the BHN parameterization, underestimates 70 % and 78 % compared to observations. The IMN contributes to 56.30 % of the number concentrations in Beijing and 27.86 % in Nanjing in SumID scheme during whole periods. In these  
465 periods, DMA accounts for 28.15 % and 29.27 % of the number concentration, respectively. In NPF days, IMN contributes 61.31 % and 32.19 % number concentrations in Beijing and Nanjing, which are increased 10 % compared with non-NPF days. Additionally, IMN nucleation pathway significantly enhance PNCs during the daytime, exhibiting a pronounced diurnal variation with higher concentrations during the day and lower levels at night. In contrast, DMA predominantly influences certain NPF events  
470 during the morning and evening peak. Overall, this study validates the enhanced model's capability to accurately simulate NPF events and underscores the significant influence of IMN and DMA nucleation pathways on particle number concentrations in **eastern cities**. ▼

**Deleted:** China

#### 475 Code and data availability.

The model code base used to generate the results for IMN and DMA nucleation (CMAQ version 5.3.2)

can be found on Zenodo at <https://doi.org/10.5281/zenodo.15739718> (Mao and Hu, 2025). Data will be made available on request to the corresponding author.

**Author contributions.**

485 J.H. conceived the research, J.M. and Z.F. conducted the model development and simulations, L.J., J.L., Y.Z., M.Q. contributed to data analyses. S.G., M.H. provided the observation data and result discussions. J.M. and J.H. wrote the manuscript, and all authors contributed to manuscript editing.

**Competing interests.**

The contact author has declared that none of the authors has any competing interests.

490 **Financial support.**

This study was supported by the Key Research and Development Program project (2022YFC3701000, task 5) and the National Natural Science Foundation of China (42021004).

## References

- 495 Almeida, J., Schobesberger, S., Kurten, A., Ortega, I. K., Kupiainen-Maatta, O., Praplan, A. P., Adamov, A., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurten, T., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Leiminger, M.,  
500 Leppa, J., Loukonen, V., Makhmutov, V., Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petaja, T., Riccobono, F., Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld, J. H., Simon, M., Sipila, M., Stozhkov, Y., Stratmann, F., Tome, A., Trostl, J., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P., Yli-Juuti, T., Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R., Vehkamäki, H., and Kirkby, J.: Molecular understanding of sulphuric acid-amine particle nucleation in the atmosphere, *Nature*, 502, 359-363, 10.1038/nature12663, 2013.
- Cai, C., Zhang, X., Wang, K., Zhang, Y., Wang, L., Zhang, Q., Duan, F., He, K., and Yu, S.-C.: Incorporation of new particle formation and early growth treatments into WRF/Chem: Model  
510 improvement, evaluation, and impacts of anthropogenic aerosols over East Asia, *Atmos. Environ.*, 124, 262-284, 10.1016/j.atmosenv.2015.05.046, 2016.
- Cai, R., Häkkinen, E., Yan, C., Jiang, J., Kulmala, M., and Kangasluoma, J.: The effectiveness of the coagulation sink of 3–10 nm atmospheric particles, *Atmos. Chem. Phys.*, 22, 11529-11541, 10.5194/acp-22-11529-2022, 2022.
- 515 Cai, R., Yang, D., Fu, Y., Wang, X., Li, X., Ma, Y., Hao, J., Zheng, J., and Jiang, J.: Aerosol surface area concentration: a governing factor in new particle formation in Beijing, *Atmos. Chem. Phys.*, 17, 12327-12340, 10.5194/acp-17-12327-2017, 2017.
- Cai, R., Yan, C., Yang, D., Yin, R., Lu, Y., Deng, C., Fu, Y., Ruan, J., Li, X., Kontkanen, J., Zhang, Q., Kangasluoma, J., Ma, Y., Hao, J., Worsnop, D. R., Bianchi, F., Paasonen, P., Kerminen, V.-M., Liu, Y., Wang, L., Zheng, J., Kulmala, M., and Jiang, J.: Sulfuric acid-amine nucleation in urban  
520 Beijing, *Atmos. Chem. Phys.*, 21, 2457-2468, 10.5194/acp-21-2457-2021, 2021.
- Carl, S. A. C., J. N.: Sequential Two (Blue) Photon Absorption by NO<sub>2</sub> in the Presence of H<sub>2</sub> as a Source of OH in Pulsed Photolysis Kinetic Studies: Rate Constants for Reaction of OH with CH<sub>3</sub>NH<sub>2</sub>, (CH<sub>3</sub>)<sub>2</sub>NH, (CH<sub>3</sub>)<sub>3</sub>N, and C<sub>2</sub>H<sub>5</sub>NH<sub>2</sub> at 295 K, *J. Phys. Chem. A*, 102, 8131-8141, 10.1021/jp9821937, 1998.
- 525 Chang, Y., Ling, Q., Ge, X., Yuan, X., Zhou, S., Cheng, K., Mao, J., Huang, D., Hu, Q., Lu, J., Cui, S., Gao, Y., Lu, Y., Zhu, L., Tan, W., Guo, S., Hu, M., Wang, H., Huang, C., Huang, R. J., Zhang, Y., and Hu, J.: Nonagricultural emissions enhance dimethylamine and modulate urban atmospheric nucleation, *Sci. Bull.*, 68, 1447-1455, 10.1016/j.scib.2023.05.033, 2023.
- 530 Chen, X., Wang, Z., Li, J., Chen, H., Hu, M., Yang, W., Wang, Z., Ge, B., and Wang, D.: Explaining the spatiotemporal variation of fine particle number concentrations over Beijing and surrounding areas in an air quality model with aerosol microphysics, *Environ. Pollut.*, 231, 1302-1313, 10.1016/j.envpol.2017.08.103, 2017.
- Chen, X., Yang, W., Wang, Z., Li, J., Hu, M., An, J., Wu, Q., Wang, Z., Chen, H., Wei, Y., Du, H., and  
535 Wang, D.: Improving new particle formation simulation by coupling a volatility-basis set (VBS) organic aerosol module in NAQPMS+APM, *Atmos. Environ.*, 204, 1-11, 10.1016/j.atmosenv.2019.01.053, 2019.

- Du, W., Dada, L., Zhao, J., Chen, X., Daellenbach, K. R., Xie, C., Wang, W., He, Y., Cai, J., Yao, L., Zhang, Y., Wang, Q., Xu, W., Wang, Y., Tang, G., Cheng, X., Kokkonen, T. V., Zhou, W., Yan, C.,  
540 Chu, B., Zha, Q., Hakala, S., Kurppa, M., Järvi, L., Liu, Y., Li, Z., Ge, M., Fu, P., Nie, W., Bianchi, F., Petäjä, T., Paasonen, P., Wang, Z., Worsnop, D. R., Kerminen, V.-M., Kulmala, M., and Sun, Y.: A 3D study on the amplification of regional haze and particle growth by local emissions, *npj Clim. Atmos. Sci.*, 4, 10.1038/s41612-020-00156-5, 2021.
- Dunne, E. M., Gordon, H., Kürten, A., Almeida, J., Duplissy, J., Williamson, C., Ortega, I. K., Pringle,  
545 K. J., Adamov, A., Baltensperger, U., Barmet, P., Benduhn, F., Bianchi, F., Breitenlechner, M., Clarke, A., Curtius, J., Dommen, J., Donahue, N. M., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Jokinen, T., Kangasluoma, J., Kirkby, J., Kulmala, M., Kupc, A., Lawler, M. J., Lehtipalo, K., Makhmutov, V., Mann, G., Mathot, S., Merikanto, J., Miettinen, P., Nenes, A., Onnela, A., Rap, A., Reddington, C. L. S., Riccobono, F., Richards, N. A.  
550 D., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Simon, M., Sipilä, M., Smith, J. N., Stozkhov, Y., Tomé, A., Tröstl, J., Wagner, P. E., Wimmer, D., Winkler, P. M., Worsnop, D. R., and Carslaw, K. S.: Global atmospheric particle formation from CERN CLOUD measurements, *Science*, 354, 1119 - 1124, 10.1126/science.aaf2649, 2016.
- Elleman, R. A. and Covert, D. S.: Aerosol size distribution modeling with the Community Multiscale  
555 Air Quality modeling system in the Pacific Northwest: 2. Parameterizations for ternary nucleation and nucleation mode processes, *J. Geophys. Res.*, 114, D11207, 10.1029/2009jd012187, 2009.
- English, J. M., Toon, O. B., Mills, M. J., and Yu, F.: Microphysical simulations of new particle  
formation in the upper troposphere and lower stratosphere, *Atmos. Chem. Phys.*, 11, 9303-9322,  
10.5194/acp-11-9303-2011, 2011.
- 560 Fan, J., Zhang, R., Collins, D., and Li, G.: Contribution of secondary condensable organics to new particle formation: A case study in Houston, Texas, *Geophys. Res. Lett.*, 33, L15802, 10.1029/2006gl026295, 2006.
- Feng, Z., Mao, J., Jiang, L., Qian, Y., Shang, D., Guo, S., Hu, M., and Hu, J.: Impacts of dimethylamine  
emissions on particle number concentration and cloud condensation nuclei in Beijing, *npj Clean*  
565 *Air*, 1, 11, 10.1038/s44407-025-00011-y, 2025.
- Fountoukis, C., Riipinen, I., Denier van der Gon, H. A. C., Charalampidis, P. E., Pilinis, C.,  
Wiedensohler, A., O'Dowd, C., Putaud, J. P., Moerman, M., and Pandis, S. N.: Simulating ultrafine  
particle formation in Europe using a regional CTM: contribution of primary emissions versus  
secondary formation to aerosol number concentrations, *Atmos. Chem. Phys.*, 12, 8663-8677,  
570 10.5194/acp-12-8663-2012, 2012.
- Fuzzi, S., Baltensperger, U., Carslaw, K., Decesari, S., Denier van der Gon, H., Facchini, M. C.,  
Fowler, D., Koren, I., Langford, B., Lohmann, U., Nemitz, E., Pandis, S., Riipinen, I., Rudich, Y.,  
Schaap, M., Slowik, J. G., Spracklen, D. V., Vignati, E., Wild, M., Williams, M., and Gilardoni, S.:  
Particulate matter, air quality and climate: lessons learned and future needs, *Atmos. Chem. Phys.*,  
575 15, 8217-8299, 10.5194/acp-15-8217-2015, 2015.
- Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A.,  
Dommen, J., Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C.,  
Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., Kürten, A., Lehtipalo, K., Makhmutov, V.,  
Molteni, U., Rissanen, M. P., Stozkhov, Y., Tröstl, J., Tsagkogeorgas, G., Wagner, R., Williamson,  
580 C., Wimmer, D., Winkler, P. M., Yan, C., and Carslaw, K. S.: Causes and importance of new particle  
formation in the present-day and preindustrial atmospheres, *J. Geophys. Res. Atmos.*, 122, 8739-

8760, 10.1002/2017jd026844, 2017.

Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an  
585 extended and updated framework for modeling biogenic emissions, *Geosci. Model. Dev.*, 5, 1471-1492, 10.5194/gmd-5-1471-2012, 2012.

Gunnar, W. S. and Paul, J. C.: Emission of Aliphatic Amines from Animal Husbandry and their Reactions: Potential Source of N<sub>2</sub>O and HCN, *J. Atmos. Chem.*, 22, 319-346, 1995.

Guo, S., Hu, M., Zamora, M. L., Peng, J., Shang, D., Zheng, J., Du, Z., Wu, Z., Shao, M., Zeng, L.,  
590 Molina, M. J., and Zhang, R.: Elucidating severe urban haze formation in China, *Proc. Natl. Acad. Sci. U. S. A.*, 111, 17373-17378, 10.1073/pnas.1419604111, 2014.

Guo, S., Hu, M., Peng, J., Wu, Z., Zamora, M. L., Shang, D., Du, Z., Zheng, J., Fang, X., Tang, R., Wu, Y., Zeng, L., Shuai, S., Zhang, W., Wang, Y., Ji, Y., Li, Y., Zhang, A. L., Wang, W., Zhang, F., Zhao, J., Gong, X., Wang, C., Molina, M. J., and Zhang, R.: Remarkable nucleation and growth of  
595 ultrafine particles from vehicular exhaust, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 3427-3432, 10.1073/pnas.1916366117, 2020.

Hu, J., Chen, J., Ying, Q., and Zhang, H.: One-year simulation of ozone and particulate matter in China using WRF/CMAQ modeling system, *Atmos. Chem. Phys.*, 16, 10333-10350, 10.5194/acp-16-10333-2016, 2016.

600 Hu, J., Huang, L., Chen, M., Liao, H., Zhang, H., Wang, S., Zhang, Q., and Ying, Q.: Premature Mortality Attributable to Particulate Matter in China: Source Contributions and Responses to Reductions, *Environ. Sci. Technol.*, 51, 9950-9959, 10.1021/acs.est.7b03193, 2017a.

Hu, J., Li, X., Huang, L., Ying, Q., Zhang, Q., Zhao, B., Wang, S., and Zhang, H.: Ensemble prediction of air quality using the WRF/CMAQ model system for health effect studies in China, *Atmos. Chem. Phys.*, 17, 13103-13118, 10.5194/acp-17-13103-2017, 2017b.

605 Huang, Y., Deng, T., Li, Z., Wang, N., Yin, C., Wang, S., and Fan, S.: Numerical simulations for the sources apportionment and control strategies of PM<sub>2.5</sub> over Pearl River Delta, China, part I: Inventory and PM<sub>2.5</sub> sources apportionment, *Sci. Total Environ.*, 634, 1631-1644, 10.1016/j.scitotenv.2018.04.208, 2018.

610 Julin, J., Murphy, B. N., Patoulias, D., Fountoukis, C., Olenius, T., Pandis, S. N., and Riipinen, I.: Impacts of Future European Emission Reductions on Aerosol Particle Number Concentrations Accounting for Effects of Ammonia, Amines, and Organic Species, *Environ. Sci. Technol.*, 52, 692-700, 10.1021/acs.est.7b05122, 2018.

Kirkby, J., Amorim, A., Baltensperger, U., Carslaw, K. S., Christoudias, T., Curtius, J., Donahue, N. M.,  
615 Haddad, I. E., Flagan, R. C., Gordon, H., Hansel, A., Harder, H., Junninen, H., Kulmala, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Lelieveld, J., Möhler, O., Riipinen, I., Stratmann, F., Tomé, A., Virtanen, A., Volkamer, R., Winkler, P. M., and Worsnop, D. R.: Atmospheric new particle formation from the CERN CLOUD experiment, *Nat. Geosci.*, 16, 948-957, 10.1038/s41561-023-01305-0, 2023.

620 Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J., Trostl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A. K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf, M., Kurten, A., Laaksonen, A., Lehtipalo, K.,  
625 Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Perakyla, O., Piel, F., Petaja, T., Praplan, A. P.,

Pringle, K., Rap, A., Richards, N. A., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipila, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tome, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533, 521-526, 10.1038/nature17953, 2016.

630 Kontkanen, J., Stolzenburg, D., Olenius, T., Yan, C., Dada, L., Ahonen, L., Simon, M., Lehtipalo, K., and Riipinen, I.: What controls the observed size-dependency of the growth rates of sub-10 nm atmospheric particles?, *Environ. Sci. Atmos.*, 2, 449-468, 10.1039/d1ea00103e, 2022.

635 Kulmala, M., Laaksonen, A., and Pirjola, L.: Parameterizations for sulfuric acid/water nucleation rates, *J. Geophys. Res. Atmos.*, 103, 8301-8307, 10.1029/97jd03718, 1998.

Kulmala, M., Cai, R., Stolzenburg, D., Zhou, Y., Dada, L., Guo, Y., Yan, C., Petaja, T., Jiang, J., and Kerminen, V. M.: The contribution of new particle formation and subsequent growth to haze formation, *Environ. Sci. Atmos.*, 2, 352-361, 10.1039/d1ea00096a, 2022.

640 Kulmala, M., Asmi, A., Lappalainen, H. K., Baltensperger, U., Brenguier, J. L., Facchini, M. C., Hansson, H. C., Hov, Ø., O'Dowd, C. D., Pöschl, U., Wiedensohler, A., Boers, R., Boucher, O., de Leeuw, G., Denier van der Gon, H. A. C., Feichter, J., Krejci, R., Laj, P., Lihavainen, H., Lohmann, U., McFiggans, G., Mentel, T., Pilinis, C., Riipinen, I., Schulz, M., Stohl, A., Swietlicki, E., Vignati, E., Alves, C., Amann, M., Ammann, M., Arabas, S., Artaxo, P., Baars, H., Beddows, D. C. S., Bergström, R., Beukes, J. P., Bilde, M., Burkhardt, J. F., Canonaco, F., Clegg, S. L., Coe, H., Crumeyrolle, S., D'Anna, B., Decesari, S., Gilardoni, S., Fischer, M., Fjaeraa, A. M., Fountoukis, C., George, C., Gomes, L., Halloran, P., Hamburger, T., Harrison, R. M., Herrmann, H., Hoffmann, T., Hoose, C., Hu, M., Hyvärinen, A., Hörrak, U., Iinuma, Y., Iversen, T., Josipovic, M., Kanakidou, M., Kiendler-Scharr, A., Kirkevåg, A., Kiss, G., Klimont, Z., Kolmonen, P., Komppula, M., Kristjánsson, J. E., Laakso, L., Laaksonen, A., Labonnote, L., Lanz, V. A., Lehtinen, K. E. J., Rizzo, L. V., Makkonen, R., Manninen, H. E., McMeeking, G., Merikanto, J., Minikin, A., Mirme, S., Morgan, W. T., Nemitz, E., O'Donnell, D., Panwar, T. S., Pawlowska, H., Petzold, A., Pienaar, J. J., Pio, C., Plass-Duelmer, C., Prévôt, A. S. H., Pryor, S., Reddington, C. L., Roberts, G., Rosenfeld, D., Schwarz, J., Seland, Ø., Sellegri, K., Shen, X. J., Shiraiwa, M., Siebert, H., Sierau, B., Simpson, D., Sun, J. Y., Topping, D., Tunved, P., Vaattovaara, P., Vakkari, V., Veefkind, J. P., Visschedijk, A., Vuollekoski, H., Vuolo, R., Wehner, B., Wildt, J., Woodward, S., Worsnop, D. R., van Zadelhoff, G. J., Zardini, A. A., Zhang, K., van Zyl, P. G., Kerminen, V. M., S Carslaw, K., and Pandis, S. N.: General overview: European Integrated project on Aerosol Cloud Climate and Air Quality interactions (EUCAARI) – integrating aerosol research from nano to global scales, *Atmos. Chem. Phys.*, 11, 13061-13143, 10.5194/acp-11-13061-2011, 2011.

650 Kurokawa, J. and Ohara, T.: Long-term historical trends in air pollutant emissions in Asia: Regional Emission inventory in ASia (REAS) version 3, *Atmos. Chem. Phys.*, 20, 12761-12793, 10.5194/acp-20-12761-2020, 2020.

655 Kurten, A., Jokinen, T., Simon, M., Sipila, M., Sarnela, N., Junninen, H., Adamov, A., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., Dommen, J., Donahue, N. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hakala, J., Hansel, A., Heinritzi, M., Hutterli, M., Kangasluoma, J., Kirkby, J., Laaksonen, A., Lehtipalo, K., Leiminger, M., Makhmutov, V., Mathot, S., Onnela, A., Petaja, T., Praplan, A. P., Riccobono, F., Rissanen, M. P., Rondo, L., Schobesberger, S., Seinfeld, J. H., Steiner, G., Tome, A., Trostl, J., Winkler, P. M., Williamson, C., Wimmer, D., Ye, P.,



- 670 Baltensperger, U., Carslaw, K. S., Kulmala, M., Worsnop, D. R., and Curtius, J.: Neutral molecular cluster formation of sulfuric acid-dimethylamine observed in real time under atmospheric conditions, *Proc. Natl. Acad. Sci. U. S. A.*, 111, 15019-15024, 10.1073/pnas.1404853111, 2014.
- Laakso, L.: Model studies on ion-induced nucleation in the atmosphere, *J. Geophys. Res.*, 107, 4427, 10.1029/2002jd002140, 2002.
- 675 Lai, S., Hai, S., Gao, Y., Wang, Y., Sheng, L., Lupascu, A., Ding, A., Nie, W., Qi, X., Huang, X., Chi, X., Zhao, C., Zhao, B., Shrivastava, M., Fast, J. D., Yao, X., and Gao, H.: The striking effect of vertical mixing in the planetary boundary layer on new particle formation in the Yangtze River Delta, *Sci. Total Environ.*, 829, 154607, 10.1016/j.scitotenv.2022.154607, 2022.
- Lee, S. H., Gordon, H., Yu, H., Lehtipalo, K., Haley, R., Li, Y., and Zhang, R.: New Particle Formation in the Atmosphere: From Molecular Clusters to Global Climate, *J. Geophys. Res. Atmos.*, 124, 7098-7146, 10.1029/2018jd029356, 2019.
- 680 Li, J., Xie, X., Li, L., Wang, X., Wang, H., Jing, S. a., Ying, Q., Qin, M., and Hu, J.: Fate of Oxygenated Volatile Organic Compounds in the Yangtze River Delta Region: Source Contributions and Impacts on the Atmospheric Oxidation Capacity, *Environmental Science & Technology*, 56, 11212-11224, 10.1021/acs.est.2c00038, 2022a.
- 685 Li, X., Zhao, B., Zhou, W., Shi, H., Yin, R., Cai, R., Yang, D., Dällenbach, K., Deng, C., Fu, Y., Qiao, X., Wang, L., Liu, Y., Yan, C., Kulmala, M., Zheng, J., Hao, J., Wang, S., and Jiang, J.: Responses of gaseous sulfuric acid and particulate sulfate to reduced SO<sub>2</sub> concentration: A perspective from long-term measurements in Beijing, *Science of The Total Environment*, 721, 137700, 10.1016/j.scitotenv.2020.137700, 2020.
- 690 Li, X., Li, Y., Cai, R., Yan, C., Qiao, X., Guo, Y., Deng, C., Yin, R., Chen, Y., Li, Y., Yao, L., Sarnela, N., Zhang, Y., Petaja, T., Bianchi, F., Liu, Y., Kulmala, M., Hao, J., Smith, J. N., and Jiang, J.: Insufficient Condensable Organic Vapors Lead to Slow Growth of New Particles in an Urban Environment, *Environ. Sci. Technol.*, 56, 9936-9946, 10.1021/acs.est.2c01566, 2022b.
- 695 Li, Z., Zhao, B., Yin, D., Wang, S., Qiao, X., Jiang, J., Li, Y., Shen, J., He, Y., Chang, X., Li, X., Liu, Y., Li, Y., Liu, C., Qi, X., Chen, L., Chi, X., Jiang, Y., Li, Y., Wu, J., Nie, W., and Ding, A.: Modeling the Formation of Organic Compounds across Full Volatility Ranges and Their Contribution to Nanoparticle Growth in a Polluted Atmosphere, *Environ. Sci. Technol.*, 58, 1223-1235, 10.1021/acs.est.3c06708, 2024.
- 700 Liu, J., Jiang, J., Zhang, Q., Deng, J., and Hao, J.: A spectrometer for measuring particle size distributions in the range of 3 nm to 10 μm, *Front. Environ. Sci. Eng.*, 10, 63-72, 10.1007/s11783-014-0754-x, 2014.
- Liu, L., Guo, S., Zhao, Z., and Li, H.: Free Energy Prediction of Ion-Induced Nucleation of Aqueous Aerosols, *J. Phys. Chem. A*, 126, 2407-2416, 10.1021/acs.jpca.1c09787, 2022.
- 705 Liu, L., Yu, F., Tu, K., Yang, Z., and Zhang, X.: Influence of atmospheric conditions on the role of trifluoroacetic acid in atmospheric sulfuric acid-dimethylamine nucleation, *Atmos. Chem. Phys.*, 21, 6221-6230, 10.5194/acp-21-6221-2021, 2021a.
- Liu, L., Yu, F., Du, L., Yang, Z., Francisco, J. S., and Zhang, X.: Rapid sulfuric acid-dimethylamine nucleation enhanced by nitric acid in polluted regions, *Proc. Natl. Acad. Sci. U. S. A.*, 118, e2108384118, 10.1073/pnas.2108384118, 2021b.
- 710 Loukonen, V., Kurtén, T., Ortega, I. K., Vehkamäki, H., Pádua, A. A. H., Sellegri, K., and Kulmala, M.: Enhancing effect of dimethylamine in sulfuric acid nucleation in the presence of water – a computational study, *Atmos. Chem. Phys.*, 10, 4961-4974, 10.5194/acp-10-4961-2010, 2010.

- Lu, Y., Liu, L., Ning, A., Yang, G., Liu, Y., Kurtén, T., Vehkamäki, H., Zhang, X., and Wang, L.:  
 715 Atmospheric Sulfuric Acid-Dimethylamine Nucleation Enhanced by Trifluoroacetic Acid, *Geophys. Res. Lett.*, 47, e2019GL085627, 10.1029/2019gl085627, 2020.
- Lu, Y., Yan, C., Fu, Y., Chen, Y., Liu, Y., Yang, G., Wang, Y., Bianchi, F., Chu, B., Zhou, Y., Yin, R.,  
 Baalbaki, R., Garmash, O., Deng, C., Wang, W., Liu, Y., Petäjä, T., Kerminen, V. M., Jiang, J.,  
 720 Kulmala, M., and Wang, L.: A proxy for atmospheric daytime gaseous sulfuric acid concentration in  
 urban Beijing, *Atmos. Chem. Phys.*, 19, 1971-1983, 10.5194/acp-19-1971-2019, 2019.
- Ma, L., Zhu, Y., Zheng, M., Sun, Y., Huang, L., Liu, X., Gao, Y., Shen, Y., Gao, H., and Yao, X.:  
 Investigating three patterns of new particles growing to the size of cloud condensation nuclei in  
 Beijing's urban atmosphere, *Atmos. Chem. Phys.*, 21, 183-200, 10.5194/acp-21-183-2021, 2021.
- Mao, J.: New particle formation and long-term trend modeling number concentration in Chinese typical  
 725 cities, Master, 10.27248/d.cnki.gnjqc.2023.000218, 2023.
- Mao, J. and Hu, J.: Code: Community Multiscale Air Quality (CMAQ) version 5.3.2 with Ion Mediated  
 Nucleation and sulfuric acid-dimethylamine Nucleation, Zenodo [code], 10.5281/zenodo.15739718,  
 2025.
- Mao, J., Li, L., Li, J., Sulaymon, I. D., Xiong, K., Wang, K., Zhu, J., Chen, G., Ye, F., Zhang, N., Qin,  
 730 Y., Qin, M., and Hu, J.: Evaluation of Long-Term Modeling Fine Particulate Matter and Ozone in  
 China During 2013–2019, *Front. Environ. Sci.*, 10, 872249, 10.3389/fenvs.2022.872249, 2022.
- Merikanto, J., Napari, I., Vehkamäki, H., Anttila, T., and Kulmala, M.: New parameterization of  
 sulfuric acid-ammonia-water ternary nucleation rates at tropospheric conditions, *J. Geophys. Res.*,  
 112, D15207, 10.1029/2006jd007977, 2007.
- 735 Napari, I.: Parametrization of ternary nucleation rates for H<sub>2</sub>SO<sub>4</sub>-NH<sub>3</sub>-H<sub>2</sub>O vapors, *J. Geophys. Res.*,  
 107, 4381, 10.1029/2002jd002132, 2002.
- Napari, I., Noppel, M., Vehkamäki, H., and Kulmala, M.: An improved model for ternary nucleation of  
 sulfuric acid–ammonia–water, *J. Chem. Phys.*, 116, 4221-4227, 10.1063/1.1450557, 2002.
- 740 Ning, A., Shen, J., Zhao, B., Wang, S., Cai, R., Jiang, J., Yan, C., Fu, X., Zhang, Y., Li, J., Ouyang, D.,  
 Sun, Y., Saiz-Lopez, A., Francisco, J. S., and Zhang, X.: Overlooked significance of iodic acid in  
 new particle formation in the continental atmosphere, *Proc. Natl. Acad. Sci. U. S. A.*, 121,  
 e2404595121, 10.1073/pnas.2404595121, 2024.
- Olenius, T., Halonen, R., Kurtén, T., Henschel, H., Kupiainen-Määttä, O., Ortega, I. K., Jen, C. N.,  
 Vehkamäki, H., and Riipinen, I.: New particle formation from sulfuric acid and amines:  
 745 Comparison of monomethylamine, dimethylamine, and trimethylamine, *J. Geophys. Res. Atmos.*,  
 122, 7103-7118, 10.1002/2017jd026501, 2017.
- Olin, M., Patoulias, D., Kuuluvainen, H., Niemi, J. V., Rönkkö, T., Pandis, S. N., Riipinen, I., and Dal  
 Maso, M.: Contribution of traffic-originated nanoparticle emissions to regional and local aerosol  
 levels, *Atmos. Chem. Phys.*, 22, 1131-1148, 10.5194/acp-22-1131-2022, 2022.
- 750 Peng, J., Hu, M., Shang, D., Wu, Z., Du, Z., Tan, T., Wang, Y., Zhang, F., and Zhang, R.: Explosive  
 Secondary Aerosol Formation during Severe Haze in the North China Plain, *Environ. Sci. Technol.*,  
 55, 2189-2207, 10.1021/acs.est.0c07204, 2021.
- Posner, L. N. and Pandis, S. N.: Sources of ultrafine particles in the Eastern United States, *Atmos.*  
*Environ.*, 111, 103-112, 10.1016/j.atmosenv.2015.03.033, 2015.
- 755 Qi, X., Ding, A., Nie, W., Petäjä, T., Kerminen, V., Herrmann, E., Xie, Y., Zheng, L., Manninen, H.,  
 Aalto, P., Sun, J., Xu, Z., Chi, X., Huang, X., Boy, M., Virkkula, A., Yang, X., Fu, C., and Kulmala,  
 M.: Aerosol size distribution and new particle formation in the western Yangtze River Delta of

- China: 2 years of measurements at the SORPES station, *Atmos. Chem. Phys.*, 15, 12445-12464, 10.5194/acp-15-12445-2015, 2015.
- 760 Qin, M., She, Y., Wang, M., Wang, H., Chang, Y., Tan, Z., An, J., Huang, J., Yuan, Z., Lu, J., Wang, Q., Liu, C., Liu, Z., Xie, X., Li, J., Liao, H., Pye, H. O. T., Huang, C., Guo, S., Hu, M., Zhang, Y., Jacob, D. J., and Hu, J.: Increased urban ozone in heatwaves due to temperature-induced emissions of anthropogenic volatile organic compounds, *Nature Geoscience*, 18, 50-56, 10.1038/s41561-024-01608-w, 2025.
- 765 Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Downard, A., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hansel, A., Junninen, H., Kajos, M., Keskinen, H., Kupc, A., Kurten, A., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petaja, T., Praplan, A. P., Santos, F. D., Schallhart, S., Seinfeld, J. H.,  
770 Sipilä, M., Spracklen, D. V., Stozhkov, Y., Stratmann, F., Tome, A., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Wimmer, D., Carslaw, K. S., Curtius, J., Donahue, N. M., Kirkby, J., Kulmala, M., Worsnop, D. R., and Baltensperger, U.: Oxidation products of biogenic emissions contribute to nucleation of atmospheric particles, *Science*, 344, 717-721, 10.1126/science.1243527, 2014.
- 775 Ruusuvoori, K., Hietala, P., Kupiainen-Määttä, O., Jokinen, T., Junninen, H., Sipilä, M., Kurtén, T., and Vehkamäki, H.: The charging of neutral dimethylamine and dimethylamine-sulfuric acid clusters using protonated acetone, *Atmos. Meas. Tech.*, 8, 2577-2588, 10.5194/amt-8-2577-2015, 2015.
- Shen, J., Zhao, B., Wang, S., Ning, A., Li, Y., Cai, R., Gao, D., Chu, B., Gao, Y., Shrivastava, M., Jiang, J., Zhang, X., and He, H.: Cluster-dynamics-based parameterization for sulfuric acid-dimethylamine nucleation: comparison and selection through box and three-dimensional modeling, *Atmos. Chem. Phys.*, 24, 10261-10278, 10.5194/acp-24-10261-2024, 2024.
- 780 Sipilä, M., Berndt, T., Petäjä, T., Brus, D., Vanhanen, J., Stratmann, F., Patokoski, J., Mauldin, R. L., Hyvärinen, A.-P., Lihavainen, H., and Kulmala, M.: The Role of Sulfuric Acid in Atmospheric Nucleation, *Science*, 327, 1243-1246, doi:10.1126/science.1180315, 2010.
- 785 Usoskin, I. G. and Kovaltsov, G. A.: Cosmic ray induced ionization in the atmosphere: Full modeling and practical applications, *J. Geophys. Res.*, 111, D21206, 10.1029/2006jd007150, 2006.
- Vehkamäki, H.: An improved parameterization for sulfuric acid-water nucleation rates for tropospheric and stratospheric conditions, *J. Geophys. Res.*, 107, 4622, 10.1029/2002jd002184, 2002.
- Wang, D., Guo, H., Cheung, K., and Gan, F.: Observation of nucleation mode particle burst and new  
790 particle formation events at an urban site in Hong Kong, *Atmos. Environ.*, 99, 196-205, 10.1016/j.atmosenv.2014.09.074, 2014.
- Wang, G., Zhang, R., Gomez, M. E., Yang, L., Levy Zamora, M., Hu, M., Lin, Y., Peng, J., Guo, S., Meng, J., Li, J., Cheng, C., Hu, T., Ren, Y., Wang, Y., Gao, J., Cao, J., An, Z., Zhou, W., Li, G., Wang, J., Tian, P., Marrero-Ortiz, W., Secrest, J., Du, Z., Zheng, J., Shang, D., Zeng, L., Shao, M.,  
795 Wang, W., Huang, Y., Wang, Y., Zhu, Y., Li, Y., Hu, J., Pan, B., Cai, L., Cheng, Y., Ji, Y., Zhang, F., Rosenfeld, D., Liss, P. S., Duce, R. A., Kolb, C. E., and Molina, M. J.: Persistent sulfate formation from London Fog to Chinese haze, *Proc. Natl. Acad. Sci. U. S. A.*, 113, 13630-13635, doi:10.1073/pnas.1616540113, 2016.
- Wang, J., Li, M., Li, L., Zheng, R., Fan, X., Hong, Y., Xu, L., Chen, J., and Hu, B.: Particle number size distribution and new particle formation in Xiamen, the coastal city of Southeast China in wintertime, *Sci. Total Environ.*, 826, 154208, 10.1016/j.scitotenv.2022.154208, 2022a.
- 800

- 805 Wang, K., Ma, X., Tian, R., and Yu, F.: Analysis of new particle formation events and comparisons to simulations of particle number concentrations based on GEOS-Chem-advanced particle microphysics in Beijing, China, *Atmos. Chem. Phys.*, 23, 4091-4104, 10.5194/acp-23-4091-2023, 2023.
- 810 Wang, M., Xiao, M., Bertozzi, B., Marie, G., Rorup, B., Schulze, B., Bardakov, R., He, X. C., Shen, J., Scholz, W., Marten, R., Dada, L., Baalbaki, R., Lopez, B., Lamkaddam, H., Manninen, H. E., Amorim, A., Ataei, F., Bogert, P., Brasseur, Z., Caudillo, L., De Menezes, L. P., Duplissy, J., Ekman, A. M. L., Finkenzeller, H., Carracedo, L. G., Granzin, M., Guida, R., Heinritzi, M., Hofbauer, V., Hohler, K., Korhonen, K., Krechmer, J. E., Kurten, A., Lehtipalo, K., Mahfouz, N. G. A., Makhmutov, V., Massabo, D., Mathot, S., Mauldin, R. L., Mentler, B., Muller, T., Onnela, A., Petaja, T., Philippov, M., Piedehierro, A. A., Pozzer, A., Ranjithkumar, A., Schervish, M., Schobesberger, S., Simon, M., Stozhkov, Y., Tome, A., Umo, N. S., Vogel, F., Wagner, R., Wang, D. S., Weber, S. K., Welti, A., Wu, Y., Zauner-Wieczorek, M., Sipila, M., Winkler, P. M., Hansel, A., 815 Baltensperger, U., Kulmala, M., Flagan, R. C., Curtius, J., Riipinen, I., Gordon, H., Lelieveld, J., El-Haddad, I., Volkamer, R., Worsnop, D. R., Christoudias, T., Kirkby, J., Mohler, O., and Donahue, N. M.: Synergistic HNO<sub>3</sub>-H<sub>2</sub>SO<sub>4</sub>-NH<sub>3</sub> upper tropospheric particle formation, *Nature*, 605, 483-489, 10.1038/s41586-022-04605-4, 2022b.
- 820 Wang, X., Li, L., Gong, K., Mao, J., Hu, J., Li, J., Liu, Z., Liao, H., Qiu, W., Yu, Y., Dong, H., Guo, S., Hu, M., Zeng, L., and Zhang, Y.: Modelling air quality during the EXPLORE-YRD campaign – Part I. Model performance evaluation and impacts of meteorological inputs and grid resolutions, *Atmos. Environ.*, 246, 118131, 10.1016/j.atmosenv.2020.118131, 2021a.
- 825 Wang, Z., Liu, Y., Wang, C., Jiang, S., Feng, Y., Huang, T., and Huang, W.: Multicomponent nucleation of malonic acid involved in the sulfuric acid - dimethylamine system and its atmospheric implications, *Atmos. Environ.*, 267, 118558, 10.1016/j.atmosenv.2021.118558, 2021b.
- 830 Wang, Z., Hu, M., Pei, X., Zhang, R., Paasonen, P., Zheng, J., Yue, D., Wu, Z., Boy, M., and Wiedensohler, A.: Connection of organics to atmospheric new particle formation and growth at an urban site of Beijing, *Atmos. Environ.*, 103, 7-17, 10.1016/j.atmosenv.2014.11.069, 2015.
- 835 Wang, Z., Wu, Z., Yue, D., Shang, D., Guo, S., Sun, J., Ding, A., Wang, L., Jiang, J., Guo, H., Gao, J., Cheung, H. C., Morawska, L., Keywood, M., and Hu, M.: New particle formation in China: Current knowledge and further directions, *Sci. Total Environ.*, 577, 258-266, 10.1016/j.scitotenv.2016.10.177, 2017.
- 840 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model. Dev.*, 4, 625-641, 10.5194/gmd-4-625-2011, 2011.
- 845 Wu, Z., Hu, M., Liu, S., Wehner, B., Bauer, S., Maßling, A., Wiedensohler, A., Petäjä, T., Dal Maso, M., and Kulmala, M.: New particle formation in Beijing, China: Statistical analysis of a 1-year data set, *J. Geophys. Res. Atmos.*, 112, 10.1029/2006JD007406, 2007.
- 850 Yan, C., Dada, L., Rose, C., Jokinen, T., Nie, W., Schobesberger, S., Junninen, H., Lehtipalo, K., Sarnela, N., Makkonen, U., Garmash, O., Wang, Y., Zha, Q., Paasonen, P., Bianchi, F., Sipilä, M., Ehn, M., Petäjä, T., Kerminen, V.-M., Worsnop, D. R., and Kulmala, M.: The role of H<sub>2</sub>SO<sub>4</sub>-NH<sub>3</sub> anion clusters in ion-induced aerosol nucleation mechanisms in the boreal forest, *Atmos. Chem. Phys.*, 18, 13231-13243, 10.5194/acp-18-13231-2018, 2018.
- 855 Yan, C., Yin, R., Lu, Y., Dada, L., Yang, D., Fu, Y., Kontkanen, J., Deng, C., Garmash, O., Ruan, J., Baalbaki, R., Schervish, M., Cai, R., Bloss, M., Chan, T., Chen, T., Chen, Q., Chen, X., Chen, Y.,

- Chu, B., Dällenbach, K., Foreback, B., He, X., Heikkinen, L., Jokinen, T., Junninen, H., Kangasluoma, J., Kokkonen, T., Kurppa, M., Lehtipalo, K., Li, H., Li, H., Li, X., Liu, Y., Ma, Q., Paasonen, P., Rantala, P., Pileci, R. E., Rusanen, A., Sarnela, N., Simonen, P., Wang, S., Wang, W., Wang, Y., Xue, M., Yang, G., Yao, L., Zhou, Y., Kujansuu, J., Petäjä, T., Nie, W., Ma, Y., Ge, M., He, H., Donahue, N. M., Worsnop, D. R., Veli-Matti, K., Wang, L., Liu, Y., Zheng, J., Kulmala, M., Jiang, J., and Bianchi, F.: The Synergistic Role of Sulfuric Acid, Bases, and Oxidized Organics Governing New-Particle Formation in Beijing, *Geophys. Res. Lett.*, 48, e2020GL091944, 10.1029/2020gl091944, 2021.
- 850 Yang, X., Wu, Q., Zhao, R., Cheng, H., He, H., Ma, Q., Wang, L., and Luo, H.: New method for evaluating winter air quality: PM<sub>2.5</sub> assessment using Community Multi-Scale Air Quality Modeling (CMAQ) in Xi'an, *Atmos. Environ.*, 211, 18–28, 10.1016/j.atmosenv.2019.04.019, 2019.
- 855 Yao, L., Garmash, O., Bianchi, F., and al., e.: Atmospheric new particle formation from sulfuric acid and amines in a Chinese megacity, *Science*, 361, 278–281, 10.1126/science.aao4839, 2018.
- Yu, F.: Altitude variations of cosmic ray induced production of aerosols: Implications for global cloudiness and climate, *J. Geophys. Res.*, 107, 1118, 10.1029/2001ja000248, 2002.
- 860 Yu, F.: Binary H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O homogeneous nucleation based on kinetic quasi-unary nucleation model: Look-up tables, *J. Geophys. Res.*, 111, D04201, 10.1029/2005jd006358, 2006a.
- Yu, F.: From molecular clusters to nanoparticles: second-generation ion-mediated nucleation model, *Atmos. Chem. Phys.*, 6, 5193–5211, 2006b.
- 865 Yu, F.: Improved quasi-unary nucleation model for binary H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O homogeneous nucleation, *J. Chem. Phys.*, 127, 054301, 10.1063/1.2752171, 2007.
- Yu, F.: Ion-mediated nucleation in the atmosphere: Key controlling parameters, implications, and look-up table, *J. Geophys. Res.*, 115, D03206, 10.1029/2009jd012630, 2010.
- 870 Yu, F. and Luo, G.: Modeling of gaseous methylamines in the global atmosphere: impacts of oxidation and aerosol uptake, *Atmos. Chem. Phys.*, 14, 12455–12464, 10.5194/acp-14-12455-2014, 2014.
- Yu, F. and Turco, R. P.: The role of ions in the formation and evolution of particles in aircraft plumes, *Geophys. Res. Lett.*, 24, 1927–1930, 10.1029/97gl01822, 1997.
- Yu, F. and Turco, R. P.: Ultrafine aerosol formation via ion-mediated nucleation, *Geophys. Res. Lett.*, 27, 883–886, 10.1029/1999gl011151, 2000.
- 875 Yu, F. and Turco, R. P.: From molecular clusters to nanoparticles: Role of ambient ionization in tropospheric aerosol formation, *J. Geophys. Res. Atmos.*, 106, 4797–4814, 10.1029/2000jd900539, 2001.
- Yu, F., Nadykto, A. B., Luo, G., and Herb, J.: H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O binary and H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O-NH<sub>3</sub> ternary homogeneous and ion-mediated nucleation: lookup tables version 1.0 for 3-D modeling application, *Geosci. Model. Dev.*, 13, 2663–2670, 10.5194/gmd-13-2663-2020, 2020a.
- 880 Yu, F., Wang, Z., Luo, G., and Turco, R.: Ion-mediated nucleation as an important global source of tropospheric aerosols, *Atmos. Chem. Phys.*, 8, 2537–2554, 2008.
- Yu, F., Luo, G., Nair, A. A., Schwab, J. J., Sherman, J. P., and Zhang, Y.: Wintertime new particle formation and its contribution to cloud condensation nuclei in the Northeastern United States, *Atmos. Chem. Phys.*, 20, 2591–2601, 10.5194/acp-20-2591-2020, 2020b.
- 885 Yu, F., Nadykto, A. B., Herb, J., Luo, G., Nazarenko, K. M., and Uvarova, L. A.: H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O-NH<sub>3</sub> ternary ion-mediated nucleation (TIMN): kinetic-based model and comparison with CLOUD measurements, *Atmos. Chem. Phys.*, 18, 17451–17474, 10.5194/acp-18-17451-2018, 2018.
- Yu, F., Luo, G., Bates, T. S., Anderson, B., Clarke, A., Kapustin, V., Yantosca, R. M., Wang, Y., and

- 890 Wu, S.: Spatial distributions of particle number concentrations in the global troposphere:  
Simulations, observations, and implications for nucleation mechanisms, *J. Geophys. Res.*, 115,  
D17205, 10.1029/2009jd013473, 2010.
- Yu, H., Ren, L., and Kanawade, V. P.: New Particle Formation and Growth Mechanisms in Highly  
Polluted Environments, *Curr. Pollut. Rep.*, 3, 245-253, 10.1007/s40726-017-0067-3, 2017.
- 895 Zhang, J., Gong, X., Crosbie, E., Diskin, G., Froyd, K., Hall, S., Kupc, A., Moore, R., Peischl, J.,  
Rollins, A., Schwarz, J., Shook, M., Thompson, C., Ullmann, K., Williamson, C., Wisthaler, A., Xu,  
L., Ziemba, L., Brock, C. A., and Wang, J.: Stratospheric air intrusions promote global-scale new  
particle formation, *Science*, 385, 210-216, doi:10.1126/science.adn2961, 2024.
- Zhang, R., Khalizov, A., Wang, L., Hu, M., and Xu, W.: Nucleation and growth of nanoparticles in the  
900 atmosphere, *Chem. Rev.*, 112, 1957-2011, 10.1021/cr2001756, 2012.
- Zhang, R., Suh, I., Zhao, J., Zhang, D., Fortner, E. C., Tie, X., Molina, L. T., and Molina, M. J.:  
Atmospheric New Particle Formation Enhanced by Organic Acids, *Sci. Rep.*, 304, 1487, 2004.
- Zhang, R., Xie, H. B., Ma, F., Chen, J., Iyer, S., Simon, M., Heinritzi, M., Shen, J., Tham, Y. J., Kurten,  
T., Worsnop, D. R., Kirkby, J., Curtius, J., Sipila, M., Kulmala, M., and He, X. C.: Critical Role of  
905 Iodous Acid in Neutral Iodine Oxoacid Nucleation, *Environ. Sci. Technol.*, 56, 14166-14177,  
10.1021/acs.est.2c04328, 2022.
- Zhang, Y., McMurry, P. H., Yu, F., and Jacobson, M. Z.: A comparative study of nucleation  
parameterizations: 1. Examination and evaluation of the formulations, *J. Geophys. Res.*, 115,  
D20212, 10.1029/2010jd014150, 2010a.
- 910 Zhang, Y., Liu, P., Liu, X., Pun, B., Seigneur, C., Jacobson, M. Z., and Wang, W.: Fine scale modeling  
of wintertime aerosol mass, number, and size distributions in central California, *J. Geophys. Res.*,  
115, 10.1029/2009jd012950, 2010b.
- Zhang, Y., Liu, P., Liu, X., Jacobson, M. Z., McMurry, P. H., Yu, F., Yu, S., and Schere, K. L.: A  
comparative study of nucleation parameterizations: 2. Three-dimensional model application and  
915 evaluation, *J. Geophys. Res.*, 115, D20213, 10.1029/2010jd014151, 2010c.
- Zhao, B., Fast, J. D., Donahue, N. M., Shrivastava, M., Schervish, M., Shilling, J. E., Gordon, H.,  
Wang, J., Gao, Y., Zaveri, R. A., Liu, Y., and Gaudet, B.: Impact of Urban Pollution on Organic-  
Mediated New-Particle Formation and Particle Number Concentration in the Amazon Rainforest,  
*Environ. Sci. Technol.*, 55, 4357-4367, 10.1021/acs.est.0c07465, 2021a.
- 920 Zhao, B., Donahue, N. M., Zhang, K., Mao, L., Shrivastava, M., Ma, P., Shen, J., Wang, S., Sun, J.,  
Gordon, H., Tang, S., Fast, J., Wang, M., Gao, Y., Yan, C., Singh, B., Li, Z., Huang, L., Lou, S., Lin,  
G., Wang, H., Jiang, J., Ding, A., Nie, W. i., Qi, X., Chi, X., and Wang, L.: Global variability in  
atmospheric new particle formation mechanisms, *Nature*, 10.1038/s41586-024-07547-1, 2024.
- Zhao, G., Zhu, Y., Wu, Z., Zong, T., Chen, J., Tan, T., Wang, H., Fang, X., Lu, K., Zhao, C., and Hu,  
925 M.: Impact of aerosol–radiation interaction on new particle formation, *Atmos. Chem. Phys.*, 21,  
9995-10004, 10.5194/acp-21-9995-2021, 2021b.
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang,  
Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since  
2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095-14111, 10.5194/acp-  
930 18-14095-2018, 2018.