

Transient Flow Patterns of an Annular-like Stratospheric Polar Vortex.

Huw C. Davies¹, Michael A. Sprenger¹

¹Institute for Atmospheric and Climate Science, ETHZ, Zurich, 8092, Switzerland

5 *Correspondence to:* Michael A. Sprenger (michael.sprenger@env.ethz.ch)

Abstract. A two-component study is undertaken of sub-planetary scale flow features associated with the wintertime polar-night vortex at upper-stratospheric levels. First, case studies based upon reanalysis data are presented to illustrate the scale, structure, evolution and dynamics of the features potential vorticity in three disparate flow settings along with signatures of some accompanying flow, thermal, and constituent variables. The features are shown to be present near the periphery of the vortex and are associated with the vortex's predilection to develop a structure akin to an annular-like band of enhanced potential vorticity. Second, theoretical considerations and numerical model simulations of perturbations of an annular band of enhanced absolute vorticity using a non-divergent barotropic model on a polar β -plane yield results consistent both with the occurrence and character of the observed sub-planetary scale features, and with the rapid reconfiguration of the band when subject to a planetary scale wave perturbation. The latter result suggests that that the break-up of the polar vortex at stratopause-levels can be facilitated by the synergetic combination of strong planetary-scale Rossby-wave forcing from below acting upon the vortex's annular potential vorticity band.

1. INTRODUCTION.

20 The general form of wintertime stratospheric polar vortex (SPV) is well established (see e.g. Waugh and Polvani, 2010; Schoeberl and Newman, 2015; Butchart, 2022). It extends from the lower stratosphere at ~ 15 km to well beyond the stratopause at ~ 50 km, attains a maximum wind strength often greater than 120 ms^{-1} at stratopause levels, and is characterized in the stratosphere by an inner cold core driven primarily by the lack of ozone-related radiative heating during the polar night.

25 On quasi-horizontal or isentropic surfaces, the SPV's lateral 'edge' has been defined in various ways (Waugh et al 2016; Manney et al 2021). Kinematically it has been associated with the iso-line demarking the strongest wind strength (Waugh et al 2016) or the maximum streamline density (Harvey et al 2002; Harvey et al 2009), cartographically as a feature-based configuration (Serra et al 2017; Lawrence & Manney, 2018), and dynamically as the zone of sharp lateral PV gradient (Butchart & Remsberg, 1986; Manney et al 1994; Nash et al 30 1996). The latter definition arises from regarding the vortex's lateral configuration as an inner core of high potential vorticity (PV) encased by a narrow zone of strong PV gradient that is itself encircled by broad and well-mixed surf zone of lower PV (McIntyre & Palmer, 1984). The adoption of one or another of the above criteria to define the vortex's periphery will depend upon its practical utility or its interpretative value for the specific purpose in hand.

35 The present study is geared to examining the sub-planetary and finer-scaled flow features that are contiguous to, or related to, the structure of the SPV's periphery, and to this end we utilize the PV perspective (Hoskins et al, 1985). The study is framed around the postulate that a, possibly fragmented, annulus of enhanced PV (- i.e. a maximum in the across-jet distribution of the PV) can exist near the periphery. This assertion is rendered plausible by two factors. First, a strong laterally confined jet with a relative vorticity maximum (say, ζ_{\max}) located 40 at a co-latitude $\chi (< 30^\circ)$ on the vortex's poleward side would connote a maximum of absolute vorticity provided

$(\zeta_{\max}/2\Omega) > 1/2\chi^2$, and this inequality is readily satisfied (- often by an order of magnitude) in the upper stratosphere. This in turn would connote a PV maximum (*sic* annulus of enhanced PV) if the variation of the stratification on an isentropic surface within the vortex is not comparably large. Second, the presence of such an annular structure on an isentropic surface is a necessary criterion for instability of axisymmetric quasi-geostrophic stratospheric flow (Charney and Stern, 1962), and the realization of the instability could result in a fragmented annulus.

The probable existence of such an annular structure in the polar stratosphere has long been recognized (Hartmann, 1983), and indeed a balanced flow response to a radiative equilibrium (Leovy, 1964) or radiatively determined (Shine, 1987) stratospheric state would possess such a maximum. Such a feature has also been detected in other planetary atmospheres (e.g. Sharkey et al 2021; Shultis et al, 2025) and is often a feature in the vicinity of the hurricane's eyewall (Schubert et al, 2018). Moreover, Davies and Spengler (2024) - hereafter denoted DS24 - drew attention to flow features at stratopause elevations that are akin to that of a fragmented annulus.

An annular structure of enhanced PV would be embedded within an SPV that itself exhibits a distinct life cycle (Lu et al 2021), and marked inter-annual variations (Butchart, 2022; Rao et al 2024) particularly those related to the occurrence and aftermath of a Sudden Stratospheric Warming (Baldwin et al 2021). This poses questions as to whether the annular structure or its fragmented counterpart is present at all phases of the life cycle, and what role (if any) it plays in the development of Sudden Stratospheric Warming (SSW) events that have themselves been pointedly linked to a variety of mechanisms and preconditioned states deemed to be more amenable to engendering such events (Plumb, 1981; Fredricksen, 1982; Smith, 1992; Matthewman and Esler, 2011; de la Camara, et al 2017; Lawrence and Manney, 2020; Yessimbet et al 2022).

Again, the annulus occupies the same spatial domain as two classes of shorter-term phenomena, and questions arise regarding the extent of their inter-dependence. One class can be viewed as intrinsic natural modes confined to, and governed by the structure of, the SPV itself such as planetary-scale zonally propagating waves (Venne & Stanford, 1979; Randel & Lait, 1991), smaller scale zonally propagating waves and synoptic-scale features (Harvey et al 2002; Lu et al 2021). The second class comprise planetary-scale Rossby waves (PRWs) (see e.g. Hitchcock and Haynes, 2016) and short-term smaller-scale gravity waves (GW) (see e.g. Holton & Alexander, 2000; Eichinger et al 2020) both of which propagate upward from lower elevations and can break on attaining large amplitude. Both are influenced by the SPVs structure, and the PRWs tend to break in the mid-to-upper stratosphere and the GWs at even higher elevations. On breaking they can heavily modulate the SPV's structure. Indeed, breaking PRWs (Hitchman and Huesmann, 2007; Greer et al, 2013) strip PV filaments off the vortex edge, create the surf zone (McIntyre & Palmer, 1984), and are endemic to the occurrence of SSW events (Martius et al, 2009). Likewise breaking gravity waves can serve as a drag (GWD) reducing the strength of the SPV's jet.

In this study attention is focused on the presence, role and dynamics of a PV annulus and/or its fragmented counterpart in the wintertime upper stratosphere. It is set against the backcloth of the series of questions posed above, and it has observational and theoretical components. For the observational component, illustrative case studies pertaining to the SPV's annular structure are presented for three different winter settings. For each case attention is drawn to the structure, temporal evolution, and dynamical character of the realized transient sub-planetary and planetary scale flow features. The three cases comprise: - (i) a mildly perturbed mid-winter vortex in the Northern Hemisphere, (ii) a comparatively undisturbed early winter vortex in the SH, and (iii) a highly disturbed NH vortex in mid-February associated with a developing SSW.

The theoretical component is prompted by noting that the observationally detected features (DS24) bear comparison with the break-up of a barotropic jet-like flow. The ready inference that a jet-like structure might satisfy a necessary condition for barotropic instability has spawned a fleet of studies for basic states akin to that

of:- the SPV's jet and the circumpolar jets of other planetary atmospheres (see for example, Pfister, 1979; Davies, 1981; Hartmann, 1983; Michaelangeli et al., 1987; Manney et al., 1988; Seviour et al 2017; Mitchell et al., 2021; 85
Waugh et al., 2023); the hurricane's eyewall (Schubert et al, 2018); and the inner region of extratropical cyclones (Leutwyler and Schaer, 2019). Key dimensionless parameters that govern the dynamics of these other systems can differ substantially from those that pertain to the stratosphere's polar night jet, but herein we consider only flow settings representative of the SPV. The strategy adopted here is to select an idealized theoretical model capable of 90
capturing the essence of rapidly growing wave patterns of a barotropic model for flow settings akin to that of a deep undisturbed SPV. The specific objective is not to replicate (or contrast) the results with that of the earlier studies, but rather to seek further basic understanding of, and insight upon, the dynamics of the observed features.

Both components are foreshadowed by the recognition that the presence of an annular PV feature in the upper stratosphere would influence the character of in-situ generated phenomena and of waves propagating into 95
the domain. For example, a change in the SPV's structure modifies its transmissivity to upward propagating waves (Hitchcock and Haynes, 2016). The strong positive PV gradient on the outer side of the annulus constitutes a columnary shaft that favours vertical PRW propagation (Abatzoglou and Magnusdottir, 2007). Contrariwise, the negative PV gradient on its inner side would inhibit the propagation into the vortex core, and thereby in principle help isolate the core from external effects. It is pertinent to note that this 'shielding of the vortex core' associated 100
with the PV gradients contrasts markedly with the dynamics at the counterpart tropopause-level sharp PV gradient (and accompanying jet stream) located at the break in the mid-latitude extratropical tropopause (Appenzeller and Davies, 1992).

The paper is arranged as follows. Section 2 sets out the observational case studies. Section 3 sets out the theoretical results derived using an idealized barotropic model to study the dynamics of both small and finite 105
amplitude perturbations of an SPV-like basic state. In a final section further remarks are made regarding the study's results and their possible implications.

2.0 Observed Flow Features.

Three case-studies are presented with the objective of capturing the essence of transient flow features associated 110
with an annular-like PV structure in the wintertime upper stratosphere.

2.1 Data.

Consideration of the flow at stratopause and higher elevations using reanalysis data sets are subject to constraints arising from the decrease in the number and resolution of sampled meteorological variables and the shortcomings 115
of the parent assimilation model at these elevations (- see the overview in S-RIP, 2022). Nevertheless, Reanalysis fields have been utilized, albeit with caveats, to examine a range of upper stratospheric diagnostic studies. Moreover, DS24 showed examples of three Reanalysis data sets, viz. ERA-Interim (Dee, et al 2011a, b), MERRA (Gelaro, 2017), and ERA-5 (Hersbach, 2020), capturing the same sub-planetary and synoptic-scale flow features.

Here, we proceed pragmatically, examine only levels at and below the prevailing stratopause and adopt 120
the ERA-Interim data set. In Appendix A illustrations are provided of the overall similarity of this study's key PV patterns as captured by the ERA-Interim and MERRA2 Reanalysis data sets.

2.2 Diagnostic Analysis.

The objective is to establish the existence, scale, structure, evolution and dynamics of the sub-planetary transient 125
flow features. To this end the instantaneous scale and structure of the PV features is first illustrated at a vertical series of pressure-level charts. In addition, for two of the three case studies, we exploit the PV perspective (Hoskins

et al 1983) by displaying a time-sequence of displays on pertinent isentropic surfaces to capture :- the time-development of the PV pattern; the possible Lagrangian-conservation of the PV of the individual features; and to assess the dynamics and possible instability of the flow. Note that the rapid vertical
130 increase of the background PV can serve to diffuse/sharpen the equatorward edge of an annular feature on a pressure surface located above/below the stratopause in comparison to its structure on a contiguous isentropic surface.


Further insight is provided by examining aspects of the contemporaneous distribution, on pertinent pressure-level surfaces, of other flow variables including the relative vorticity, divergence, potential temperature,
135 specific humidity and ozone mass mixing ratio. For each case study we present a summary of key features at the 10mb level followed by examples of the structure and evolution of the flow at upper stratospheric levels for specific time windows.

2.3 Features of a Mid-winter SPV

140 The Boreal Winter of 2004-2005 was comparatively non-descript and there wasn't an SSW event. At the 10mb level the time-latitude pattern of zonal mean velocity (Fig. 1a) increased almost monotonically to a maximum of $\sim 70 \text{ ms}^{-1}$ by the 19th of January and decreased almost uniformly thereafter. The corresponding time-longitude pattern of the meridional velocity at 60N (Fig. 1b) shows evidence of a strong $m = 2$ pattern by the middle of December. Its amplitude and phase remained comparatively constant for the remainder of the Winter although it
145 transiently weakened and retrogressed in early January. The onset of the wave and the forementioned transient retrogression was accompanied by a relatively strong upward EP fluxes (not shown).

At stratopause elevations, the winter's flow pattern in general showed notable, transient synoptic-scale variations. Here we select the 11th January as a not atypical day, and Figure 2 shows the instantaneous patterns of the PV and the relative vorticity on a sequence of pressure levels at 00Z 11th January. At all levels there are
150 planetary scale features (- zonal wavenumber $m = 1-3$) with evidence of finer-scale features embedded within an annular-like structure. The upper two rows (panels a-d) indicate the presence within or on the periphery of the SPV of localized regions of PV maxima with co-located but more diffuse regions of positive relative vorticity. These features resemble those identified in DS24 and are sub-planetary / synoptic in scale, and their relative vorticity values ($\sim 1.0 \cdot 10^{-4} \text{ s}^{-1}$) are comparable to that of the earth's rotation. Their spatial scale signifies: - (a) via
155 Stokes's circulation theorem, a circulation of $\sim 25 \text{ ms}^{-1}$, and (b) a supra-geostrophic flow. Thus, the contemporaneous existence of these synoptic-scale features (- hereafter referred to as SSS features) indicates that they are an integral dynamical aspect of the SPV's overall structure, and their amplitude and lateral width are indicators of the in-situ jet's sharpness. In principle the flow field attributable to these features can influence the spatial distribution of chemical constituents within the SPV's core and at its periphery.


160



165

Fig. 1 Panel (a) shows the zonal mean velocity as a function of latitude (30-90N) at 10hPa for the Boreal Winter of 2004-2005, and panel (b) depicts the meridional velocity field at 60N at the same level for the same winter season. (Based upon the NCEP/NCAR Reanalysis.)

170



175

Fig. 2 Depicted are polar stereographic projections poleward from 30N of the PV in units of $\text{K m}^2 \text{kg}^{-1} \text{s}^{-1}$ (left column) and the relative vorticity in units of s^{-1} (right column) for 00Z 11th January 2005 on the 1, 2, 5 and 10 mb surfaces. (Based upon the ERA-Interim Reanalysis.)

180


185 Panels (e-j) demonstrate that at this time most of the forementioned SSS features are vertically coherent
extending down to 10 hPa (~ 30 km) with the relative vorticity exhibiting comparable amplitude at all levels. This
contrasts with the expected increase with height of an upward propagating wave's velocity field as the density
decreases. Although the features are deep, their meridional width diminishes downwards so that they exhibit a
stalactite-like structure. Again, the character of the features, with their predominantly positive maxima of PV,
190 differ intrinsically from the patterns of the corresponding fields astride the tropopause level jet (Appenzeller and
Davies, 1992), and this again points to the different transient dynamics of the two jets.

Figure 3 displays, for the same time instant as Fig. 2, the distributions of the temperature on the 1, 2 and
3mb pressure surfaces (panels a-c), and the wind vector field (panel d), the specific humidity (panel e) and the
divergence patterns (panel f) on the 2mb surface. The temperature fields are comparatively smooth and indicate
that the SPV's cold pool weakened above 2 mb. In general, they do not exhibit the finer-scale features of the
195 corresponding PV and relative vorticity patterns, whereas the stratification / static stability (not shown) at the PV
maxima as determined in terms of the potential temperature difference between the different pressure levels, is
typically twice that of the adjacent areas. These two contrasting facets of the thermal structure are in harmony with
the forementioned vertical coherence of the PV features that attain a maximum amplitude at stratopause levels.

The wind vector field (Fig. 3d) pinpoints the location of the jet, and it is evident that the jet's overall
200 strength suffices to substantially conceal the wind signature of the individual SSS features. In passing we note that
a convenient *ad hoc* measure of the SPV's core is the domain located poleward of zero relative vorticity (- see Fig.
2 and Fig. 3d). In the ERA-Interim Reanalysis the specific humidity is inert and merely advected with the in-situ
winds and hence is akin to a flow tracer. Its distribution (Fig. 3e) is suggestive of a well-mixed SPV core that is
substantially encircled by an elongated streamer of low humidity values advected on a different potential
205 temperature surface.

The amplitude of the divergence field (Fig. 3f) is typically less than a quarter of the corresponding relative
vorticity (and the corresponding absolute vorticity) values of the SSS features, and its spatial pattern is more
reminiscent of buoyancy/gravity waves. This suggests that the dynamics of the sub-planetary scale features at
stratopause elevations has a strong quasi-barotropic component. This lends a measure of credence to our later
210 adoption of a barotropic theoretical model.


The prior development of the PV pattern depicted in Fig. (2c) is shown in Fig. (4). It displays a 6 hourly
time-sequence of the corresponding patterns to 00Z 11th January. The fragmented annulus is seen to evolve with
time and the individual embedded SSS features are readily identified exhibiting a measure of temporal coherence
as they are both advected by, and serve to modify, the in-situ flow. Note that the panels in Fig. 4 display the PV
215 pattern on the 2hPa surface. This renders ready comparison with the patterns of other variables shown on the 2hPa
surface in Figures 2 and 3. In the subsequent case studies the evolution of the PV pattern is shown on the
dynamically more instructive isentropic surfaces.



220

Fig. 3 Distribution of the temperature on the 1, 2 and 3mb surfaces (upper row) and the wind vector (ms^{-1}), specific humidity ($10^{-6} \text{kg kg}^{-1}$) and divergence (10^{-5}s^{-1}) on the 2mb surface (lower row) for the same time instant as Fig. 2. (Based upon the ERA-Interim Reanalysis.)

225



230

Fig. 4 Evolution of the PV ($\text{K m}^2 \text{kg}^{-1} \text{s}^{-1}$) on the 2hPa pressure surface at 6 hourly intervals from 00Z on the 8th January to 18Z on the 10th January.

235

2.4 Features of a nascent SPV

The Austral winter stratosphere is less subject to PRW forcing from below, and here we examine the SPV during late Spring / early Southern Hemisphere winter of 2018. On the 10mb pressure surface, the nascent SPV in April has a weak pole-centred PV anomaly, and by mid-May the jet's mean zonal flow had increased to $\sim 60 \text{ ms}^{-1}$ (Fig. 5a). In late April through early May there is evidence of a weak eastward propagating $m = 1$ wave (Fig. 5b) but it disappeared by mid-month. The accompanying polar temperature had decreased by mid-May to $\sim 200\text{K}$ with a pole-to-30S temperature difference of some 25K (not shown).

Above, at stratopause levels, the localized region of high PV at the SPV's core began to break apart in early May and thereafter had a fragmented structure. Here the focus is on the synoptic-scale features in mid-month. Fig. (6) shows the PV, relative vorticity, divergence, potential temperature, ozone mass mixing ratio, and wind vector patterns on the 1mb pressure surface at 00Z on 19th May. A train of isolated PV centres circumscribes the Hemisphere near 60S with another PV feature in the neighbourhood of the pole (Fig. 6a). The train is also readily identifiable in the relative vorticity pattern (Fig. 6b) with, as in the previous case, peak amplitudes greater than $1.2 \times 10^{-4} \text{ s}^{-1}$ - a value an order of magnitude stronger than that of the less organized divergence / gravity-wave field (Fig. 6c). The depth of the train itself extends downward at least over a scale depth with similar co-located patterns present on the 2mb and 3mb surfaces.

The potential temperature (θ) field (Fig. 6d) shows that at the 1mb level the train is accompanied by a co-located zonal band of low potential temperature ($\sim 1650\text{K}$). One possible inference is that the forementioned polar PV feature is located on a higher isentropic surface (around $\theta \sim 1800\text{K}$). Saliiently, the contemporaneous distribution of the ozone mass mixing ratio (Fig. 6e), shows not only a rich pattern of lower O3 values within the SPV's core, but also a string of high values coincident with the vortex train itself. The latter is consistent with the presence of raised isentropic surfaces beneath the features of enhanced PV. Finally, the wind vector pattern (- Fig. 6f) is consistent with the existence of strong shear on the poleward side of the jet and the presence of troughs linked to the stronger vortices of the train.

Figure (7) gives an indication of the prior development and movement of the isolated SSS features in the period from the 15th to the 20th May. It depicts at daily interval the PV pattern on the 1700 isentropic surface and shows that the features advect eastward with, and are deformed by, the in-situ flow. To a measure they retain their amplitude pointing to their predominantly adiabatic character. Their temporal development is reminiscent of the downstream growth of a vortex train forming on a barotropically unstable shear layer rather than a synchronous development of vortices on such a layer. A noteworthy feature is that Fig. (6a) and Fig. (7e) display a close correspondence of both the individual features and the rim-like structure on the pressure and isentropic surfaces.

270

275




Fig. 5 Counterpart of Fig. (1) but for April-June 2018 in the Southern Hemisphere at 10mb. The zonal velocity now pertains to the latitude range 30-90S, and the meridional velocity field is for 60S.

280

285

290





Fig. 6 Depiction of the PV($\text{Km}^2\text{kg}^{-1}\text{s}^{-1}$), relative vorticity (10^{-4} s^{-1}), divergence (10^{-5} s^{-1}), potential temperature (K), ozone mass mixing ratio ($10^{-6}\text{ kg kg}^{-1}$), and wind vector (ms^{-1}) patterns on the 1mb pressure surface at 00Z on 19th May. The panels are polar stereographic projections poleward from 30S, except for panel (f) which is only poleward from 60S.

295

300



305

Fig. 7 Evolution of the PV pattern at stratopause levels on the 1700K isentropic surface at daily intervals from 00Z on the 15th May to the 20th May 2018. The labels I, II etc. attached to the PV maxima indicate the successive locations of the individual features. (Units: $\text{Km}^2\text{kg}^{-1}\text{s}^{-1} * 10^2$)

310

2.5 Features related to an SSW.

A notable Sudden Stratospheric Warming occurred in February 2018, and it has subsequently been the subject of considerable scrutiny. This includes consideration of the event's development (Butler et al 2020), the contribution of GWD (Watanabe et al, 2022), its dynamics and predictability (Rao et al 2018; Karpechko et al. 2018; Knight et al 2020; Erner et al, 2020), the accompanying middle atmosphere chemical signature (Pérot and Orsolini, 2021), its difference to the 2019 SSW (Butler et al 2020), its impact upon the mesosphere (Wang et al, 2019), and role of surface conditions in its occurrence (Lü et al, 2020). Here we briefly consider the planetary-scale and annular features of this SSW.

320

Figure (8) is the counterpart of Fig. 1 but for the 2017-2018 Boreal Winter. It shows that at 10mb the zonal mean velocity (Fig 8a) strengthened during the last week of December, attained a maximum of $\sim 60 \text{ ms}^{-1}$ in early January, decelerated to $\sim 30 \text{ ms}^{-1}$ in the period from the 14th to the 19th of the month, and subsequently oscillated about this value before finally reversing around the 11th February. The time-longitude section for the meridional velocity at 60N (Fig. 8b) indicates the presence of significant planetary scale wave patterns. The $m=1$ wave present in December was supplemented by an $m = 2$ signal in January and subsequently supplanted by the

325

latter ahead of the SSW event. Polar temperature tended to decrease to the 10th January, increased by ~15K by the 19th, and oscillated thereafter until the onset of the SSW (not shown).


330 Above, at stratopause levels, the presence of the forementioned planetary scale waves were evidenced first by an off-pole PV pattern in December, and a dumb-bell off-pole pattern in January that distorted in the run up to the SSW. Figure (9) shows the patterns of the PV (left column), relative vorticity (centre column) and temperature (right column) on the 1mb (upper row), 2mb (middle row) and 10mb (lower row) surfaces at a time (00Z 12th Feb.) shortly after the SPV had satisfied the standard empirical criteria for an SSW. A distinctive and strong split of the SPV's PV distribution is evident at all three levels. The PV patterns on the upper two levels are similar (- as are the relative vorticity patterns), and the range of the temperature fields are also similar. The near-
335 pole temperatures are relatively low (high) at the upper levels (lower level). Together these flow and thermal patterns attest to the vertical coherency of an anomalous PV structure that is 'effectively' centred near or below the 2mb surface.

Figure 10 portrays the development of the PV pattern at upper and mid-stratospheric levels at 48-hour intervals immediately ahead of the satisfaction of the SSW criterion. The upper row depicts the patterns on the
340 1500K isentropic surface at 00Z on the 6, 8 and 10th February, and the lower row shows the corresponding patterns on the 850K isentropic surface.


On the higher surface an annular-like structure is evident on the 6th in the form of an off-pole elliptical structure with embedded sub-planetary scale features. It begins to fragment between the 6th and 8th, and by the 10th notable PV maxima have formed over Canada and western Europe. On the lower theta surface, the initially
345 compact elliptic PV structure (Fig. 10d), first distends (Fig. 10e) and then splits (Fig. 10f). An inference is that the vortex is disrupted earlier and more severely on a higher θ surface.

The corresponding thermal patterns (not shown) on the upper (2mb) and lower (10mb) levels possess a dipole structure of opposite polarity that reverses with time. From the 10th to the 12th February the near polar-temperature changes at the 1, 2, and 10mb levels are respectively -18K, -32K, and +16K.

350




355 Fig. 8 As for Fig. (1) but for the Winter of 2017-2018.



360

Fig. 9 Distribution of the PV($\text{Km}^2\text{kg}^{-1}\text{s}^{-1}$) in the left column, relative vorticity (10^{-4}s^{-1}) in the centre column, and potential temperature in the right column on the 1, 2, and 10mb pressure surfaces at 00Z 12th February (- shortly after the SSW).

365



370

Fig. 10 The upper row display the PV patterns in the upper stratosphere on the 1500K isentropic surface at 00Z for the 6, 8 and 10th February, and the lower row shows the corresponding patterns but on the 850K surface. The labels A,B,C etc. attached to individual PV maxima identify the successive location of the individual features. (Units: $\text{Km}^2\text{kg}^{-1}\text{s}^{-1} * 10^6$)

375

380 2.6 Assessment of the Results.

The foregoing case studies are for three fundamentally different flow settings and are focused on the finer-scaled flow features contiguous to, or related to, the structure of the Stratospheric Polar Vortex's periphery at stratopause levels. In particular, they sought to capture the essence of their scale, structure, temporal evolution, and dynamics.

The results show the features to be present in all three flow settings as synoptic-scale PV extrema located principally on the vortex's periphery in both hemispheres with associated relative vorticity values of $\sim 10^{-4}\text{s}^{-1}$ and extending typically at least a scale-depth below the stratopause. The temperature and horizontal divergence patterns do not have a comparable localized signal on pressure surfaces located at stratopause levels. Together these features are consistent with deep isolated PV features centred near the stratopause with associated strong static stability at their core and weak quasi-horizontal temperature gradients at their centre. Also taken together the amplitude and synoptic-scale of the vorticity centres and the significantly weaker amplitude of their divergence

390

suggests that the contribution of the horizontal divergence term to the development of the vertical component of vorticity is weak. This is not inconsistent with the adoption of an idealized barotropic model in the study's theoretical component (Section 3).

395 Again, the distinctive pattern of the ozone mixing ratio is consistent with the existence and dynamical signature of the isolated PV features whilst the space-time development of the PV pattern itself on key isentropic surfaces sheds light on their translation, development and conservation. In effect these transient sub-planetary and synoptic scale, predominantly monopole, positive PV (and relative vorticity) anomalies translate with and distort the ambient SPV's flow. These observationally based characterization of the PV features underlines the desirability of establishing the climatology of the features. In this context we note for example that at the 2mb level on every
400 January 15th of the 41 year ERA-Interim data set, SSS features and/or an annular-like structure were present on 24 days, the PV vortex was severely disrupted on another 16 days (- of which 11 were related to major or minor SSW events at 10 hPa), and on one day there was a single pole-centred PV vortex.

The presence of the features can in principle contribute to the distribution of chemical constituents within the core. Their existence near the core's periphery suggests that they can modify the perceived width of the SPV, potentially help mix air both within the core and across its periphery, and serve to diffuse the jet (Bowman and
405 Chen, 1994; Ishioka K., and Yoden, 1995; Mitsu and Yoden, 2001). The latter surmise points to their possible role, in addition to GWD, in reducing the jet strength at and above the stratopause. Again, the predominantly monopole structure of the features is consistent with their predilection to evolve from a banded structure of enhanced PV near the SPV's periphery and is strongly reminiscent of the growth to finite amplitude of perturbations of a barotropically unstable flow. Also, the break-up of the stratopause-level annular band of
410 enhanced PV during a Sudden Stratospheric Warming event opens the possibility that the band might participate in, and contribute to, the initial development of such an event.

415 3.0 Theoretical Model Studies.

Our objective here is to explore the character of perturbations that can evolve on an annular-like flow structure akin to that of a deep, axisymmetric and undisturbed SPV. To this end we parsimoniously adopt an idealized theoretical model capable of representing fundamental features of such a setting including the occurrence of barotropic instability. The model constitutes non-divergent barotropic flow on a polar β -plane. It can sustain a
420 basic state of a cylindrical (possibly unstable) jet, and it incorporates a suitable representation of the latitudinal variation of the Coriolis effect but excludes consideration of three-dimensional effects, and therefore best represents flows structures that are comparatively deep. These restrictions, together with the model's simple configuration, imply that our results will constitute a potentially insightful qualitative guide rather than providing quantitative definitive statements.

425

3.1 Model Configuration and Methods of Analysis

On a polar β -plane the earth's vorticity is given by $f = 2\Omega [1 - \frac{1}{2}(r^*/R)^2]$, where r^* denotes radial distance on the plane and R refers to the Earth's radius. Then the model's governing equation takes the form

$$D(Z^*)/Dt^* = 0, \tag{1}$$

430 where D/Dt^* denotes the material derivative for two-dimensional flow, and Z^* is the 'system-relative' absolute vorticity

$$Z^* = \zeta^* - 2\Omega [\frac{1}{2}(r^*/R)^2].$$

Here ζ^* denotes the relative vorticity and is such that $\zeta^* = \nabla^2 \psi^*$ with ψ^* denoting the stream function. In the following the dimensional starred variables are replaced by non-dimensional unstarred variables :- $r = r^*/R$, $Z = Z^*/(2\Omega)$, $\zeta = \zeta^*/(2\Omega)$, and an azimuthal velocity $V = V^*/(2\Omega R)$.

To facilitate the derivation of analytical results we let the ‘system relative’ absolute vorticity of the undisturbed circumpolar vortex comprise three concentric zones (see Fig.11) of respective uniform absolute vorticity (Z_I, Z_{II}, Z_{III}), given by

$$Z_I = 0, \quad \text{for } r < a, \quad (2a)$$

$$Z_{II} = C, \quad \text{for } a \leq r \leq b, \quad (2b)$$

$$Z_{III} = nC, \quad \text{for } r > b. \quad (2c)$$

This represents a relatively quiescent innermost core ($r < a$), an annular region ($a \leq r \leq b$) yielding an absolute vorticity maximum, and an outer domain ($r > b$) of weaker vorticity (i.e. $n < 0$).

The accompanying axisymmetric basic flow (V) takes the form

$$V_I = (1/8)r^3, \quad \text{for } r < a, \quad (3a)$$




Fig. 11 Panel (a) is a schematic of the polar β -plane and the geometry for the initial distribution of the vorticity.

Panel (b) shows the radial distribution of three concentric zones of uniform Z (-displayed as heavy dashed lines) such that there is annular band of enhanced Z located between ($a < r < b$).

$$V_{II} = \frac{1}{2}Cr [1 - (a/r)^2] + (1/8)r^3 \quad \text{for } a \leq r \leq b, \quad (3b)$$

$$V_{III} = \frac{1}{2}Cr [n + (b/r)^2 (\epsilon - n)] + (1/8)r^3 \quad \text{for } r > b, \quad (3c)$$

It is also pertinent to note that,

$$460 \quad (V_a/a) = (1/8)a^2, \quad (4)$$

$$(V_b/b) = 1/2[C\varepsilon + (1/4)b^2], \quad (5)$$

where (V_a/a) and (V_b/b) denote respectively the angular velocity at $r = a$ and b .

Here $\varepsilon = [1 - (a/b)^2]$ is a measure of the width of the annular region relative to its radial location, and the $(1/8)r^3$ term in Eqs (3a-c) is attributable to the β -effect. In effect the parameters (a, b, C, n) determine the basic state of the vortex core, with typically $C \leq 2$, $b \sim 1/2$, and a and b such that ε is capable of spanning across its
465 permissible $[0, 1]$ range.

The parameter (nC) determines the relative vorticity (ζ) distribution in the exterior (i.e. $\zeta_{III} = nC + 1/2 r^2$). Its value is bounded by the requirements that the absolute vorticity in the far-field be less than in the polar region ($nC < 0$), and the basic state flow should be inertially stable ($nC > -1$). A more stringent condition is the
470 requirements that at a radial distance, say $r = r_c$, the velocity is significantly reduced and ζ approaches zero (i.e. $nC \sim -1/2 r_c^2$). For such a reduction to occur by say $\sim 30N$ then typically $0.7 > r_c > 1.2$.

Figure 12 shows illustrative examples of the basic state flow profiles (V) representable by Eqs. (3). For the $r \leq b$ domain, the jet profiles correspond to maxima of 90 ms^{-1} and 135 ms^{-1} located at $b = 1/2$ ($\approx 60N$), for wide ($\varepsilon = 0.91$), intermediate ($\varepsilon = 0.75$) and narrow ($\varepsilon = 0.36$) annuli. Note that, for C in the range $(1/2 - 2)$ and ε in the
475 range $(3/4 - 1)$, then $v_b \approx 1/4(C\varepsilon)$, and the velocity profiles shown in the outer domain ($r > b$) span those for realistic values of r_c . In essence Fig. 12 shows that the parameter settings permit a range of appropriate velocity profiles.

Results are derived for the linear and non-linear response of perturbations of stipulated basic states. The linear component includes a normal mode analysis that establishes the growth rate and azimuthal wavenumber (m) of unstable modes, and an analysis of the response to a specific form of non-normal mode initial perturbation. The
480 non-linear response is simulated using a contour-dynamics technique (Dritschel, 1989) with adjustments of the standard contour advection technique to (a) account for the specification of the absolute as opposed to the relative vorticity and (b) circumvent the standard contour advection stipulation of non-zero absolute vorticity in the outermost domain. The forward integration is conducted using a 4th order Runge-Kutta scheme and, depending upon the purpose of the simulation, the initial state is perturbed by adding either weak amplitude random noise to
485 the location of the interfaces or a finite amplitude displacement to the location of the interface at $r = b$.

3.2 Normal Mode Analysis

3.2.1 Derivation of the Instability Criterion

Small amplitude perturbations of the governing equation (Eq.1) satisfy the linear equation,

$$490 \quad \{\partial/\partial t + (V/r)\partial/\partial\theta\}q' + u' \partial/\partial r (Z) = 0 \quad (6)$$

Here (q', u') refer to the perturbation vorticity and radial velocity with $q' = \nabla^2 \psi'$. Likewise, (Z, V) are, as indicated earlier, the basic state absolute vorticity and azimuthal velocity (Eqs. 2 and 3).

Adopting a 'PV Perspective' for the perturbations (Davies and Bishop, 1994) solutions of Eq. (6) are sought for stream-function perturbations (ψ') of the form

$$495 \quad \psi' = \psi'_a + \psi'_b,$$

where the two components represent wave perturbations propagating respectively on the vorticity discontinuities at the domain interfaces located at $r = a$ and b . The associated stream functions are

$$\psi'_{(a,b)} = F_{(a,b)}(r) \sin (m\theta - \omega t),$$

and for basic states of the form given by Eqs. (3), F has a r^m and r^{-m} dependency respectively within and exterior
 500 to the pertinent interface. On applying appropriate conditions at $r = 0$, as $r \rightarrow \infty$, and at the interfaces, the wave
 amplitudes at the two interfaces, $F_{(a,b)} = (A, B)$, are linked by the relationships,

$$(\omega/m)A = (V_a/a)A + \frac{1}{2}(C/m) [A + (1-\varepsilon)^{m/2}B], \quad (7a)$$

and $(\omega/m)B = (V_b/b)B - \frac{1}{2}(C/m)(1-n) [B + (1-\varepsilon)^{m/2}A]$ (7b).

The three-terms on the right-hand side in each of Eqs. (7a, b) correspond respectively to :- translation with the
 505 ambient angular velocity (Eqs. 3d, e); a ‘Rossby-wave type’ contribution of the in-situ wave; and a far-field
 ‘Rossby-wave type’ contribution due to the influence of a wave at one interface acting upon that at the other
 interface. In the absence of the latter far-field contributions, equivalence of the phase velocities would require -
 see Eqs. (4, 5)

$$\mathcal{D} = 0, \quad (8a)$$

510 where $\mathcal{D} = [m \varepsilon (1+\mu) - (2-n)]$ (8b)

$$\text{with } \mu = b^2/(4C).$$

We return to consider the import of Eq. (8a) later but proceed here to note that together Eqs (7a, b) yield the linear
 system’s wave-dispersion relationship. After some manipulation the instability criterion can be written as

$$\Delta > 0, \quad (9a)$$


515 with $\Delta = [-\mathcal{D}^2 + 4(1-n)(1-\varepsilon)^m]$, (9b)

and for unstable waves, the growth rate (σ) and angular phase speed (v) are given by,

$$\sigma = (1/4) C \sqrt{\Delta}, \quad (9c)$$

$$v = \frac{1}{2} [(V_a/a + V_b/b) + (nC)/m], \quad (9d)$$


In line with the necessary criteria for barotropic instability the annular region needs to be a maximum of absolute
 520 vorticity (see Eq. 9b), i.e. $(1-n) > 0$.



525 Fig. 12 A sample of the model's possible basic state velocity (V) profiles corresponding to a dimensional jet strength of 90 ms^{-1} (left) and 135 ms^{-1} (right) located at $b = \frac{1}{2}$ ($\approx 60\text{N}$). In each panel the three curves for $r < b$, depicted with long, intermediate and short dashes, refer respectively to wide ($\varepsilon = 0.91$), intermediate ($\varepsilon = 0.75$), and narrow ($\varepsilon = 0.36$) annuli, and the dotted line is the velocity profile, where applicable, in the $r < a$ sector. The two full curves beyond $r = b$ encompass the physically constrained band (i.e. $0.7 > r_c > 1.2$) of velocity profiles. (For the specific values of C see the legend to Fig. 13).

530

535



540 Fig.13 Growth rate (σ) displayed as a function of the wavenumber (m) for a jet of strength 90 ms^{-1} (left) and 135 ms^{-1} (right) located at $b = \frac{1}{2}$ ($\approx 60\text{N}$). Note the differing scale of the ordinates. The dashed red, dashed blue and thick black curves are for respectively: - narrow ($\varepsilon = 0.36$), intermediate ($\varepsilon = 0.7$); and wide ($\varepsilon = 0.91$) annular bands. For the 90 ms^{-1} jet, the two curves for each of the three categories the $[C, r_c]$ values are :- $[0.91, (0.85-0.95)$; $0.44, (0.89-0.93)$; $0.36, (0.85-0.95)$]. The corresponding values for the

545 135 ms^{-1} jet are:- $[1.44, (0.93-1.0)$; $0.69, (0.93-1.05)$; $0.57, (0.96-1.07)$].

3.2.2 Sample Growth Rate Curves.

A sample of the dependency of the instability upon the configuration of the basic state is illustrated in Fig. 13. It shows the growth rate (σ) as a function of the wavenumber (m) for settings corresponding to those referred to

550 earlier (- see Figs.12 and 13). In terms of conventional flow parameters, the growth rate is seen to increase markedly with jet strength, decrease strongly as the sharpness of the jet decreases on its poleward side (i.e. width of the annulus increases), and is less sensitive to the velocity profile in the surf zone (as determined by the value of r_c). In effect for these settings the sensitivity to the jet skewness is primarily governed by the structure within the core. These results are in qualitative agreement with, and complement those of, earlier barotropic instability

555 studies (see in particular Hartman, 1983) but a significant difference (- discussed further below) is the stability of the $m = 1$ or 2 modes. The large growth rate of the sub-planetary modes is notable since a pragmatic lower bound for an unstable mode's growth rate would be that σ should exceed τ , where $\tau \sim (10 \text{ days})^{-1}$ is the radiative decay timescale (Shine, 1987) of deep perturbations within the polar-night vortex in the upper stratosphere.

560

3.2.3 Dynamics of the Instability.

Figure 13 provides a useful, but non-exhaustive, display of the unstable modes dependence upon the basic state configuration. Further numerous computations using Eq. (9c) would enable the sensitivity to the externally specified parameters (such as b) or combination of those parameters (such as μ) to be empirically inferred. Here
 565 in contrast we seek to deductively determine those sensitivities by pinpointing the dynamics underlying of the instability and thereby concomitantly establishing the wavenumber of the most unstable mode.

To this end we adopt the PV-based phenomenological interpretation (Lighthill, 1967; Hoskins et al 1985) and mathematical formulation (Davies & Bishop, 1994) of barotropic and baroclinic instability. In this framework the present form of instability is viewed in terms of two counter propagating waves on the opposite-signed PV
 570 jumps at $r=a$ and $r=b$ that are held stationary relative to one another by the differing in-situ flow at their respective locations. In such a configuration the waves inter-dependency can promote mutual amplification and/or induce a modification of their azimuthal phase velocities.

The key to the instability dynamics comes from noting the implications of $\mathcal{D} = 0$. First, phase locking of the two waves is achieved without an inter-wave contribution (- see discussion leading to Eq. 8a) so that the
 575 two perturbations are in quadrature and their inter-wave influence is constrained to yield maximal growth. Second there is (see Eq. 9b) an azimuthal wave number (m_0) for which a wave perturbation is unstable with

$$m_0 = (2 - n) / [\varepsilon (1 + \mu)] \quad (10a)$$

In effect m_0 is the ratio of the sum of the two vorticity discontinuities to the difference in the angular velocities across the annular region. It can be reformulated as

$$580 \quad m_0 \varepsilon = 2 [1 + rc^2/(4C)] / [1 + b^2/(4C)] \quad . \quad (10b)$$

585




Fig. 14 Left panel displays λ in the space of $[(rc)^2, C]$ for $b = \frac{1}{2}$ with the realistic range for rc lightly shaded.

590 Right panel shows φ as a function of ε for selected values of the wave number m^* , and the permissible domain is confined to the lightly shaded wedge-shaped region with $(m^* \varepsilon) > 2.0$.

595

It follows that, for specified values of the external parameters, satisfaction of Eqs (10a, b) signifies the instability of the wavenumber m_0 . The accompanying growth rate is given by Eq. (9c), so that

$$\sigma = \frac{1}{2} C [(1 - n) (1 - \varepsilon)^m]^{1/2} \quad (11a)$$

600 In conjunction with Eq. (10b) this can be recast in the form

$$\sigma = \frac{1}{2} \{C [C + \frac{1}{2}(rc)^2] (1 - (m_0 \varepsilon)/m_0)^{m_0}\}^{1/2} \quad (11b)$$

The third consequence of setting $\mathcal{D} = 0$ is that m_0 provides (for $m_0 > 2$) an excellent estimate of the wavenumber ($m = m^*$) of the most unstable mode. This follows from (a) deducing that

$$605 \quad (m^*/m_0) = 1 + \lambda \varphi \quad (12)$$

$$\text{where } \lambda = [1 + \frac{1}{2} rc^2] / \{ [1 + rc^2/(4C)] [1 + b^2/(4C)] \}$$

$$\text{and } \varphi = (1 - \varepsilon)^{m^*} \ln(1 - \varepsilon) / (\varepsilon) .$$

and (b) noting from Fig. (14) that λ is less than or order 1 and $\varphi \sim -0.1$ for the range of pertinent basic state settings.

610 In summary the dynamics of the instability is encapsulated by the characteristics of the normal mode whose two interface waves are in quadrature (i.e. $\mathcal{D} = 0$). It is unstable, its wavenumber (m_0) is an accurate estimate of that of the most unstable mode (m^*) for $(m^*) > 2$, and it provides (via Eqs. 10) a compact description of the sensitivity of m^* to the configuration of the basic state. Again, it demonstrates that the occurrence of unstable high wavenumbers (say, $m^* > 8$) requires narrower annular regions (- small ε); unstable sub-planetary scale waves
615 (say, $3 < m^* < 8$) exist for a wide range of external settings; and the graver modes ($m = 1, 2$) are stable.

3.2.4 An Instability Ready Reckoner,

The two panels of Fig. (15) are based upon Eq. (10) and together constitute an instability ‘Ready Reckoner’. For a stipulated specification of the external parameters, the procedure is as follows :-

- 620 (i) Fig. (15a) determines the value of $(m_0 \varepsilon)$ as a function of rc^2 and C (- see Eq. 10b),
(ii) an estimate m_0 of the most unstable wavenumber is then known given the specification of ε ,
Furthermore, the factors influencing the growth rate and an estimate of that growth can be gleaned from inspection of Fig. (15b) and in conjunction with Eq. (11b). It is noteworthy that the growth rate is highly dependent upon the value of C , favoured by higher values of rc^2 , and is sensitive to the value of m_0 (- see Fig. 15b).

625 In effect the Ready Reckoner is an indicator of the scale and growth rate of the most unstable modes over the realistic range of external parameters. Moreover, the results, in conformity with the observational cases studies of Section 2, indicate that sub-planetary and synoptic-scale features can evolve preferentially for a wide-range of realistic annular flow settings.

630





Fig. 15 A sensitivity guide to the most unstable normal mode: Displayed in panel (a) is a plot of $(m_0 \varepsilon)$ as a function of $(r_c)^2$ for a range of C with $b = 1/2$. Again the realistic range for r_c is the lightly shaded region. Panel (b) depicts the growth factor $(1 - (m_0 \varepsilon)/m_0)^{m_0}$ as a function of the wavenumber m_0 .

3.2.5 Stability of the $m = 1, 2$ Modes

640 The character of the gravest modes ($m = 1, 2$) are not adequately represented by Eqs, (10) and (11). However, they are of particular interest in connection with the occurrence of an SSW, and therefore merit **special** attention. For these two modes the instability criterion, *sic.* the parameter Δ (Eqs. 9), equates to,

$$\Delta = - \{ \varepsilon^2 (1 + \mu)^2 - 2\varepsilon [n + \mu(2-n)] + n^2 \} \quad \text{for } m=1,$$

and $\Delta = - \{ \varepsilon^2 [\mu(2+\mu) + n] - \varepsilon [n + \mu(2-n)] + (n^2/4) \}$ for $m=2$,

645 where, as before, $\mu = b^2/(4C)$ and $n = -1/2(r_c^2/C)$.

For realistic values of μ and n , generally $\Delta < 0$, so that both modes are indeed stable. Notwithstanding Δ can be small ($0 > \Delta > -0.2$) for $m = 2$ across the entire range of ε values, and small for $m = 1$ as $\varepsilon \rightarrow 0$. (This will become a significant factor in the next sub-section when we consider a mechanism that runs counter to the non-growth of these normal modes.)

650

3.3 Initial-value Dependency.

Consider a configuration of our model system that at the initial instant comprises a pre-existing small amplitude wave on the outer ($r = b$) interface but with no perturbation on the inner ($r = a$) interface. The analogue stratospheric setting would be the combination of:- (a) the distortion of the outer interface by, say, a Rossby wave emanating from tropopause elevations and propagating upward preferentially along the waveguide formed by the sharp PV gradient on the polar vortex's outer edge; and (b) the inner interface being initially substantially undisturbed being protected by the reversed latitudinal PV gradient on an annular band's inner side.

To replicate the model's evolution of such a configuration we now allow the stream functions of the two interface waves to assume the form,

660
$$\psi'_1 = \mathcal{A}(t) \sin [m\theta - \delta_1(t)], \quad \text{on } r=a \quad (13a)$$

and
$$\psi'_2 = \mathcal{B}(t) \sin [m\theta - \delta_2(t)]. \quad \text{on } r=b \quad (13b)$$

In effect the waves are allowed to have a time-varying amplitudes (\mathcal{A}, \mathcal{B}) and phases (δ_1, δ_2) at respectively $r=(a, b)$, with $\mathcal{B}(0) = \mathcal{B}_0$ and $\mathcal{A}(0) = 0$. Thus, in contrast to the normal mode analysis which allows the time development of perturbations of a fixed spatial form, the amplitude/growth rate and shape of these perturbations evolve with time. Substitution into Eq. (6) yields the following triad of predictive equations,

665

$$\partial/\partial t (\mathcal{A}) = -1/2 (C\gamma) \mathcal{B} \sin \delta, \quad (14a)$$

$$\partial/\partial t (\mathcal{B}) = -1/2 (C\gamma) (1-n) \mathcal{A} \sin \delta, \quad (14b)$$

$$\mathcal{A}\mathcal{B} \partial/\partial t (\delta) = +1/2 C \{ (\mathcal{A}\mathcal{B}) \mathcal{D} - \gamma [(1-n) \mathcal{A}^2 + \mathcal{B}^2] \cos \delta \}, \quad (14c)$$

where $\gamma = (1 - \varepsilon)^{m/2} = (a/b)^m$, and δ is the relative phase-difference $\delta = (\delta_2 - \delta_1)$.

670 A trenchant inference is that Eqs (14a, b) indicate that the inner and outer waves can grow synchronously provided only that the basic state configuration satisfies the necessary (but not necessarily sufficient) normal mode criterion for instability, $(1-n) > 0$, and that δ remains in the range $[-\pi, 0]$ with maximum growth prevailing when $\delta = -\pi/2$. Significantly this transient growth prevails irrespective of the value of Δ , and pertains even for grave ($m = 1$ and 2) perturbations that are stable for normal mode perturbations.

675

The triad possess two temporal invariants,

$$\partial/\partial t [(1-n)\mathcal{A}^2 - \mathcal{B}^2] = 0, \quad (15a)$$

$$\text{and} \quad \partial/\partial t \left\{ \frac{1}{2}(\mathcal{D}/\gamma) [(\mathcal{A}^2 + \mathcal{B}^2)/(2-n)] - \mathcal{A}\mathcal{B}\cos\delta \right\} = 0. \quad (15b)$$

These invariants can be exploited to show that the growth of the wave on the inner interface ($r = a$) is given by,

$$680 \quad \partial/\partial t (\mathcal{A}) = (1/4)C [\Delta\mathcal{A}^2 + 4\gamma^2\mathcal{B}_0^2]^{1/2} \quad (16)$$

$$\text{so that} \quad \mathcal{A} = (2\gamma/\Delta^{1/2})\mathcal{B}_0 \sinh [\Delta^{1/2}T] \quad \text{for } \Delta > 0 \quad (17a)$$

$$\text{and} \quad \mathcal{A} = (2\gamma/\Delta^{*1/2})\mathcal{B}_0 \sin [\Delta^{*1/2}T] \quad \text{for } \Delta^* = -\Delta > 0 \quad (17b)$$

$$\text{where } T = (1/4)Ct$$

Note that for an unstable configuration ($\Delta > 0$), the factor $\Delta^{1/2}T$ is the normal mode growth rate (Eq. 9c), and the
685 coupled waves evolve toward a state of exponential growth. For a stable configuration ($\Delta^* > 0$) the amplitude attains a maximum of

$$\mathcal{A}_{\max} = 2(a/b)^m \mathcal{B}_0 / \Delta^{*1/2} \quad \text{at the time } t = 2\pi / (C\Delta^{*1/2}). \quad (17c)$$

For both settings, the initial short time evolution is given by

$$\mathcal{A} \approx 2(a/b)^m \mathcal{B}_0 T \quad (17d)$$

690

The dependency upon $(a/b)^m$ indicates that it is larger for a narrow annular region and planetary-scale modes. Pointedly this growth can be significant. For example, if $(a/b) = 1/2$ and $C = 1/2$, then in dimensional terms \mathcal{A} would attain the value of \mathcal{B}_0 in $\sim 2/3$ day for $m = 1$ and in $\sim 4/3$ days for $m = 2$, so that the wave-mean flow interaction would rapidly become non-linear. Together Eqs. (17c, d) establish the form of the basic state that would yield
695 significant transient growth of the \mathcal{A} wave.

The accompanying amplitude of the \mathcal{B} -wave and the phase-difference are,

$$\mathcal{B}^2 = (1-n)\mathcal{A}^2 + \mathcal{B}_0^2 \quad (17e)$$

$$\text{and} \quad \cos^2\delta = (1/4) (\mathcal{D}/\gamma)^2 (\mathcal{A}/\mathcal{B}_0)^2 / [1 + (1-n)(\mathcal{A}/\mathcal{B}_0)^2] \quad (17f)$$

Eq. (17e) implies that after significant growth the amplitude ratio of the two waves would be

$$700 \quad (\mathcal{A}/\mathcal{B}) = 1/(1-n)^{1/2} = 1/[1+r_c^2/(2C)]^{1/2}$$

so that for a weak vortex the \mathcal{B} wave would dominate.

A feature of Eq. (17f) is that, for a flow setting with $\mathcal{D} = 0$, the \mathcal{A} and \mathcal{B} waves are *ab initio* phase-locked with $\delta = -\pi/2$ and hence the optimum growth criterion is sustained (Eqs. 14a, b).

The seminal deduction of the above analysis is that, for an annular basic state, the initial growth rate
705 and wavenumber of the fastest growing non-normal mode can be, and generally is, significantly different to that of the fastest growing normal mode. Indeed, for a basic state that would yield a most unstable synoptic-scale normal mode, there can exist $m = 1$ and 2 non-normal mode perturbations whose initial growth rate surpasses that of the fastest normal mode.


A further implication arises because the stratosphere's transmissivity to upward propagating PRWs serves
710 as a 'scale-selection' mechanism only allowing these gravest waves to reach well into the stratosphere to disturb the vortex's rim at higher elevations. Thus, it is the gravest waves that could initiate the fore-mentioned significant 'scale-sensitive' growth.

715 **3.4 Non-linear Development.**

For the same model configuration, we examine the non-linear evolution of perturbations to the model's stipulated basic state. In Figure (16) the three rows are for different initial configurations each with a jet of 120 ms^{-1} but located at $b = \frac{1}{2}$ (- for the first two rows but with different annuli widths) and for the atypical value of $b = 0.67$ (third row). The three columns depict respectively the initial small amplitude randomly perturbed initial states (left column), an early phase of development (central column) and the mature non-linear (right column) patterns. In accord with the linear normal-mode instability theory, the upper two rows capture the development of the synoptic-scale most unstable wavenumbers ($m = 6$ and 9) that then leads to an aggregation of the bands into quasi-isolated mono-signed vortices each attached to its neighbours by thin filaments. The third row, for which linear theory indicates that only the $m = 3$ and 4 waves are unstable, shows that after an extended time (- the calculated growth rates are very weak) there is a complex break-up of the annular structure.

Figure (17) shows an example of the evolution of an initial finite amplitude $m = 2$ perturbations of the $r = b$ interface but with no perturbation of the inner ($r = a$) interface. This setting is that envisaged in the previous sub-section, and pertains for $(C, a, b, n) = (1.0, 0.447, 0.5, -0.5)$ that again corresponds to a jet of 120 ms^{-1} and perturbed by a wave with an amplitude of 0.03 . Again, in harmony with the results derived earlier, the evolution exhibits a notable growth over a period of 3.2 days.


735



740

Fig. 16 Evolution of three different annular bands subjected to initial small amplitude random perturbations. The initial states are shown in panels (a, d and g) for which the external parameters (a , b , C , n) take the respective values of $(0.434, 0.5, 1.71, -0.57)$, $(0.4, 0.5, 1.12, -0.63)$, and $(0.53, 0.67, 0.20, -0.65)$. In each case the jet strength is 120ms^{-1} . Panels (b, e and h) display an early quasi-linear development phase, and panels (c, f, i) capture a mature non-linear phase.

745



750

Fig. 17 Development of an annular band's absolute vorticity when the outer interface ($r = b$) is subject to an initial finite-amplitude $m = 2$ wave perturbation. The simulation is for (a, b, C, n) taking the values $(0.447, 0.5, 1.0, -0.5)$ with an initial wave amplitude of (0.03) . Panel (a) displays an initial state, and the patterns displayed in panel (b) and panel (c) are in dimensional terms after 1.3 and 3.2 days.

755

760

It results in the formation of two vortex structures each possessing a trailing filamentary structure, and the pattern
 765 and its subsequent development (not shown) bears a modicum of resemblance to the split classification pattern
 that can emerge during an SSW event. However, the simplicity of our model cautions against overstating the
 relationship to realized events.

It is also pertinent to note the results of two null simulations of this case (not shown). First for an initial
 small amplitude random noise perturbation of both interfaces, there evolves the (expected according to our normal
 770 mode theory) growth of a wave number $m = 12$. The second null simulation corresponds to an initial configuration
 comprising a jet of the same strength as before at $r = b$ but now with a uniform absolute vorticity distribution
 across the entire $r < b$ domain. When subjected to the same 0.03 amplitude wave perturbation at $r = b$, the
 configuration evolves to a steadily rotating ellipse with an aspect ratio close to unity. In effect, these ‘null’
 simulations indicate that the vortex breakdown is attributable explicitly to the existence of the annular band. This
 775 underlines and illustrates the significance of the results presented in sub-section 3.2.4, and points to the efficacy
 of the gravest-mode wave forcing to the break-up of an annular vortex.

3.5 Model Critique and Assessment of the Results.

This study’s examination of the possible barotropic instability of the Polar Night Jet was geared not to replicating
 780 (or refuting) the results of the previous numerous related studies but rather was aimed at seeking further
 understanding of the associated processes. Our results are constrained by the simplicity of the adopted polar β -
 plane configuration and non-divergent barotropic model, and here we first comment in turn upon these restrictions.

The polar β -plane is a reasonable assumption for flow features confined substantially to the polar and
 extra-tropical latitudes because the Coriolis effect is well represented at these latitudes. By way of comparison, for
 785 barotropic flow on the sphere:- (a) the Circulation Theorem imposes an integral constraint on the net globally area
 averaged relative vorticity, and (b) the requirement of inertial/symmetric stability of an axisymmetric basic state
 the relative vorticity must also be zero at the equator. On a polar β -plane there is no *à priori* equivalent to the
 integral constraint whilst a condition of zero relative vorticity imposed at the β -plane’s pseudo-equivalent of the
 equator (that is, $r_c = 1.4$) would imply for our model that a state of (unlikely) neutrality to symmetric perturbations
 790 pertains in the entire region between b and r_c (in effect the surf zone). Our approach has been to select a basic
 state absolute vorticity profile that ensures a relatively quiescent inner region (i.e. $Z_l = 0$, for $r < a$), a jet capable
 of spanning the range of observed values, and a value for the absolute vorticity in the outer region that is both
 consistent with inertial (as opposed to barotropic) stability and yields realistic values for the zonal velocity and /or
 the relative vorticity at our model’s equivalent of the sub-tropics.

795 The impact of setting, $Z_l = IC$, is that the key Eqs. (10, 11) are modified to the form,

$$m_0 = [2 - (l + n)] / [\varepsilon (1 + \mu - l)] \quad (18)$$

$$\sigma = \frac{1}{2} / C / [(1 - n) (1 - l) (1 - \varepsilon)^m]^{1/2} \quad (19)$$

In effect for $l \ll 1$ the changes are minor with a reduced growth rate. The reverse shear across the annulus attributed
 to the inner region inhibits the instability, and a similar effect was noted in a simpler setting by Dritschel (1989).

800 The second constraint referred to above is the limitation of non-divergence and barotropy. This alleviated
 somewhat at stratopause elevations by the deductions presented in Section 2 that were based upon the character of
 the observed features. Paldor (2021) provide a formal comparison of geostrophic, shallow water divergent and
 non-divergent representations of barotropic instability. In passing we again note that our earlier observational
 deductions render more challenging the specification of the barotropic ‘equivalent depth’ as adopted in some
 805 shallow water instability studies (Matthewman and Esler, 2011; Sharkey et al, 2021). More fundamentally the
 assumption of barotropy excludes consideration of baroclinic (c.f. Dickinson, 1983; Simmons 1974) and mixed

barotropic-baroclinic (c.f. Manney & Randel, 1993) instability at these elevations. In mitigation we note that the monopole character of the PV features points to a barotropic-like instability mechanism.

Positively, in comparison with some other studies, our model does not invoke quasi-geostrophy (c.f. Davies, 1981; Hartmann, 1983), incorporates an acceptable representation of the β -effect (c.f. Dritschel, 1989; Dritschel & Polvani, 1992). The adoption of the polar β -plane configuration, as opposed to spherical geometry (Hartman, 1983; Michaelangeli et al., 1987; Manney et al., 1988; Mitchell et al., 2021; Waugh et al., 2023) enables a simpler derivation and interpretation of analytical solutions for a wide range of relatively realistic basic states velocity fields. Also, our examination of the annular feature and its response to a large wave perturbation differs from that pursued by Matthewman and Esler (2011) who considered the development of a non-annular vortex to such forcing.

The dependency of the most unstable normal mode to the jet's strength and skewness and to the width of the annular region was noted earlier. More generally the linear normal mode analysis and the non-linear simulations indicate that, for a wide range of external parameters characterizing the model's jet structure, the modes of maximum growth possess space-time scales and structure consistent with that of the observed features. In effect barotropic instability provides an attractive mechanism to account for the observed sub-planetary scale features.

An apparent discrepancy is that our normal-mode graver modes ($m = 1, 2$) are stable in contrast with their instability as detected by Hartmann (1983). This can be linked to the form of the basic state adopted by Hartmann that can comprise a second weaker annulus of enhanced PV in mid-latitudes. This latter structure introduces two possible additional modes of instability associated either with the second annulus (with a most unstable mode of high wavenumber) or with the broad region of diminished PV located between the two annuli (with a most unstable mode of low wavenumber). This association was noted by Hartmann. The equivalent for our model configuration is an annulus of reduced absolute vorticity with a jet located at $r = a$ associated with ($C \rightarrow -1$; $l \rightarrow -1$; $n \ll 1$). Inspection of Eqs. (18, 19) suggests, and direct evaluation of Δ (- see sub-section 3.2.4) confirms, that the wavenumber $m = 2$ can be unstable and possess a large growth rate. The interpretation is that this mode's instability is aided by the basic state flow component within the annulus that is attributable to the "shear-effect" of the highly non-quiescent inner region.

Clearly the existence of this unstable mode in an upper-stratospheric setting depends crucially on the presence of an annulus of reduced PV. The evidence for such a feature is meagre although an instantaneous N-S section at a given longitude might (incorrectly) attribute such a structure to the existence of a trailing streamer of high PV emanating from the core and present within the surf zone at that longitude.

840 **4.0 Further Remarks.**

This study can be viewed as lending further credence to two assertions regarding the Stratospheric Polar Vortex. The first is that *the vortex's periphery can be populated with distinctive and dynamically significant sub-planetary scale flow features*. Their presence (DS24) and this study's more detailed description of their structure, temporal evolution and dynamics serves to augment and refine the well-established documentation of SPV's overall structure (see Waugh and Polvani, 2010; Schoeberl and Newman, 2015; Butchart, 2022).

The second is that *there is merit in considering the dynamics of a jet-like barotropically unstable flow configuration comprising an annular band of enhanced absolute vorticity to help account for the character of the forementioned observed features*. Our linear instability analysis of such a band indicates that, for a wide range of external parameters characterizing the model's jet structure, the readily calculated normal modes that yield

850 maximum growth rates possess space-time scales and structure consistent with that of the observed features. In effect barotropic instability provides an attractive mechanism to account for the observed sub-planetary scale features. In contrast the model's gravest planetary-scale normal modes ($m = 1, 2$) are shown to be stable and hence do not of themselves support a small amplitude instability hypothesis associated with SSW events.

855 In addition, the study relates to two other key themes of contemporary SPV studies. First, for mass and chemical composition budget studies and for dynamically studies, it has been customary to identify (- albeit as noted earlier in diverse ways) the SPV's 'edge'. The occurrence of sub-planetary scale features near the SPV's periphery, each associated with a distinctive flow feature, adds further complication to defining the 'edge'. Therefore, it would appear appropriate to characterize the vortex's periphery as possibly comprising "a rim of
860 finite-width, that is highly variable in both in space and time, and that corresponds to an annular or fragmented region of enhanced PV".

As noted earlier, this study bears upon the dynamics of SSW events that result from strong planetary-scale Rossby-wave ($m = 1$ or 2) forcing from below acting upon the SPV. Our model simulations depict the rapid break-up of an annular structure when subject to strong $m = 1$ or 2 wave-perturbation of its outer rim. It is therefore
865 attractive to conjecture that a PV annulus of the stratospheric polar vortex in conjunction with the forementioned planetary-scale wave forcing of its outer rim could facilitate the break of the SPV itself. In such a development, the vortex 'break-down' would result, not from a displacement or splitting of the SPV's core at stratopause levels, but rather from the aggregation into one or two distinct isolated entities of the annular band's PV on the SPV's periphery. Our linear theory indicates that the rapidity of the break-up is in part favoured by a narrow annular
870 band. However, direct observational corroboration of the conjecture is hampered in part by the almost ubiquitous presence of large deviations away from symmetric circumpolar flow of the SPV at stratopause levels.

The present study would benefit from more detailed reanalysis-based case studies that include an evaluation of the influence of the parent model's GWD parameterization at and above the stratopause. Likewise, it would be illuminating to perform numerical simulations with a primitive equation model possessing high
875 horizontal and vertical spatial resolution of potentially unstable basic flow states (- or realized atmospheric states) and undertaken with, and without, strong Rossby wave forcing from below.

880


Appendix A: Comparison of upper stratospheric PV features in two Reanalysis data sets.

The veracity of Reanalysis data sets at upper stratospheric levels is constrained by the accuracy of the dynamics and non-parameterized physics of the parent assimilation models, the adequacy of the raw data used for the analysis, the fidelity of any upper level phenomenologically based sub-grid scale Gravity Wave Drag (GWD)
885 parameterization scheme, and any ad hoc divergence damping scheme of the models. In so-far as the parameterized schemes are tuned to match the expected state of the upper atmosphere, they have an artefact element that might/could compensate for, or disguise, other physical effects.

The Figures recorded in the main text were derived from ERA-Interim Reanalysis (Dee et al, 2011a, b). Here we compare this data set's representation of the present study's key flow features with the corresponding
890 fields derived from the MERRA2 set (Gelaro et al, 2017). The two assimilation models have horizontal resolutions at polar-extratropical latitudes, number of vertical levels, and top-most level of respectively for ERA-Interim of (~ 79 km, 60, 0.1hPa), and for MERRA2 of (~ 50 km, 72, 0.01hPa). In addition, MERRA2 incorporates improved

stratospheric temperature data, and there are significant differences (see S-RIP 2022, in particular Tables 2.3 and 2.7) in the treatment of the uppermost levels.

895



900 Fig. A1. A comparison of the PV patterns derived with ECMWF-Interim data at 1, 2, 3, 5 and 10mb. The left panels Fig. 2 (a, b, c, e, g, i) are reproduced from the main text's Fig. 2, and in the right panels are the corresponding patterns derived from MERRA2 data. (Note the different colour scaling schemes for the two sets).

905 In Figures (A1, A2 and A3) the PV patterns displayed respectively in Fig 2 (a, c, e, f, g), Fig. 6a and Fig 9 (left column) are set alongside the corresponding MERRA2 patterns. The choice of PV as the elected variable is appropriate since it involves both the rotational component of the velocity and the thermal field whilst avoiding the divergent wind component that can be / is influenced by the differing treatment of inertia-gravity waves at stratopause levels in the various Reanalysis fields.

910

915




Fig. A2. As for Fig.(A1) but now comparing the PV pattern shown in Fig. (6a).

920




Fig. A3

As for Fig. (A1) but now comparing the PV pattern shown in Fig. (9).

The similarity of these seminal sub-planetary scale PV features in the ERA-Interim and MERRA2 is striking particularly considering the forementioned differences in the procedure for deriving the two Reanalyses. It testifies to the consistency of their representation of these key features and lends credence to their accuracy and to our adoption of the ERA-Interim data set.

Code and data availability. The NCAR-NCEP Reanalysis data was accessed via the NOAA Physical Sciences Laboratory (<http://psl.noaa.gov/>). The ERA-Interim data was accessed from the same source, and also directly from the ECMWF data repositories (Dee et al 2011b). The code for the Contour Dynamics simulation is set out in Dritschel (1989).

Author contributions. HCD designed and undertook the observational and theoretical components of the study in consultation with MAS, and MAS undertook the contour dynamics simulations. HCD wrote the paper, the figures were prepared jointly, and MAS edited the final version of the text.

Competing interests. The contact author has declared that neither of the authors has any competing interests.

Acknowledgements. The authors wish to thank Prof. Peter Hitchcock and Prof. Thomas Birner for their insightful and constructive comments. Access to the NCAR-NCEP and ERA-Interim Reanalysis data sets as set out above, is gratefully acknowledged as is the use of their software. Again, the availability of the original contour dynamics code is appreciated.

Financial support. This research has been supported by Prof. Heini Wernli's research group at the Institute of Atmospheric and Climate Science of the ETH Zurich.

References

- Abatzoglou, J. T., and Magnusdottir, G.: Wave breaking along the stratospheric polar vortex as seen in ERA-40 data. *Geophysical Research Letters*, 34(8), -. doi: L08812 Doi 10.1029/2007gl029509., 2007.
- Appenzeller, Ch. and Davies, H.C.: Structure of stratospheric intrusions into the troposphere. *Nature*, 358, 570-572, 1992.
- Baldwin, M. P., Ayarzagüena, B., Birner, T., Butchart, N., Butler, A. H., Charlton-Perez, A. J., et al.: Sudden Stratospheric warmings. *Reviews of Geophysics*, 59, e2020RG000708. <https://doi.org/10.1029/2020RG000708>, 2021.
- Bowman, K.P. and Chen, P.: Mixing by barotropic instability in a non-linear model. *J. Atmos. Sci.*, 51, 3692-3705, 1994.
- Butchart, N.: The stratosphere: a review of the dynamics and variability, *Weather Clim. Dynam.*, 3, 1237–1272, 2022.
- Butchart, N. and Remsberg, E. E.: The area of the stratospheric polar vortex as a diagnostic for tracer transport on an isentropic surface, *J. Atmos. Sci.*, 43, 1319–1339, 1986.
- Butler A.H., Lawrence Z.D., Lee S.H., Lillo S.P., Long C.S. : Differences between the 2018 and 2019 stratospheric polar vortex split events. *Q.J.R. Meteorol. Soc.* 146: 3503–3521. <https://doi.org/10.1002/qj.3858>, 2020.

- Charney, J.G. and Stern, M. E.: On the stability of internal jets in a rotating atmosphere. *J. Atmos. Sci.*, 19, 159-172, 1962.
- 980
- Davies H. C.: An interpretation of sudden warmings in terms of potential vorticity. *J. Atmos. Sci.*, 38, 428-445, 1981.
- Davies, H. C. and Bishop, C.: Eady edge waves and rapid development. *J. Atmos. Sci.*, 51, 1930-1946, 1994.
- Davies H.C. and Sprenger, M. A.: Cyclone-like features within the stratospheric polar-night vortex. *Geophys. Res. Letters*, 51, e2024GL109529, 2024.
- 985
- Davies, H.C., Sprenger, M.A. and Wernli, H.: Quasi-geostrophic Theory. In: Gerald R. North, John Pyle and Fuqing Zhang (Editors). *Encyclopedia of Atmospheric Sciences*. 3rd Edition., Vol. 2, 393-403. DOI:10.1016/B978-0-12-382225-3.00326-1, 2024.
- Dee, D. P., et al.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Quart. J. Roy. Met. Soc.*, 137, 553-597, 2011a.
- 990
- Dee, D.P., Uppala, S.M., Simmons, A.J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M.A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A.C.M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A.P., Monge-Sanz, B.M., Morcrette, J.J., Park, B.K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.N. and Vitart, F. : ERA-Interim global atmospheric reanalysis. [Dataset] Copernicus Climate Change Service (C3S) Climate Data Store (CDS), <https://doi.org/10.24381/cds.f2f5241d>. (2011b)
- 995
- de la Camara, A., Albers J.R., Birner, T., Garcia, R.R., Pitchcock, P., Kinnison, D.E. and Smith A.K.: Sensitivity of sudden stratospheric warmings to previous stratospheric conditions. *J. Atmos. Sci.*, 74, 2857–2877, 2017.
- 1000
- Dritschel, D.G.: On the stabilization of a two-dimensional vortex strip by adverse shear. *J. Fluid Mech.* 206, 193-221, 1989.
- Dritschel, D.G.: Contour dynamics and contour surgery: Numerical algorithms for extended, high-resolution modelling of vortex dynamics in two-dimensional, inviscid, incompressible flows. *Comput. Phys. Rep.*, 10, 77-146, 1989.
- 1005
- Dritschel, D.G. and Polvani, L.M.: The roll-up of vorticity strips on the surface of a sphere. *J. Fluid Mech.*, 234, 47-69, 1992.
- Dickinson, R.E.: Baroclinic instability of an unbounded zonal shear flow in a compressible atmosphere. *J. Atmos. Sci.*, 30, 1520-1527, 1973.
- 1010
- Eichinger, R., Garny, H., Sacha, P., Danker, J., Dietmuller, S. and Oberlander-Hayn, S: Effects of missing gravity waves on stratospheric dynamics; part 1: Climatology. *Climate Dynamics* 54, 3165-3183, 2020.
- Erner, I., Karpechko, A.Y., and Järvinen., H.J.: Mechanisms and predictability of sudden stratospheric warming in winter 2018. *Weather Clim. Dynam.*, 1, 657–674, 2020 <https://doi.org/10.5194/wcd-1-657-2020>
- Fredriksen, J.S.: Instability of the three-dimensional distorted stratospheric polar vortex at the onset of the sudden warming. *J. Atmos. Sci.*, 39, 2313-2329, 1982.
- 1015
- Gelaro, R., et al. : The Modern-Era Retrospective analysis for research and applications, version 2 (MERRA-2). *Journal of Climate*, 30, 5419–5454, 2017.
- Greer, K., Thayer, J.P. and Harvey, V.L.: A climatology of polar winter stratopause warmings and associated planetary wave breaking. *J. Geophys. Res.*, 108, 4168-4180, 2013.
- 1020
- Hartmann, D.L.: Barotropic instability of the polar night jet stream, *J. Atmos. Sci.*, 40, 817-835, 1983.
- Harvey, V.L., Pierce, R.B., Fairlie, T.D., and Hitchman, M.H.: A climatology of stratospheric polar vortices and

- anticyclones. *J. Geophys. Res.*, 107(D20), 4442, 2002.
- Harvey, V.L., Randall, C.E. and Hitchman, M.H., (2009), Breakdown of potential vorticity - based equivalent latitude as A vortex-centered coordinate in the polar winter mesosphere, *J. Geophys. Res.*, 114, D22105, doi:10.1029/2009JD012681.
- 1025
- Hersbach, H. et al., The ERA5 global reanalysis. *Quart. J. R. Meteorol. Soc.*, 146, 1999–2049, 2020.
- Hitchcock, P., and Haynes, P.: Stratospheric control of planetary waves, *Geophys. Res. Lett.*, 43, 11,884–11, 892, doi:10.1002/2016GL071372
- Hitchman, M.H. and Huesmann, A.S.: A seasonal climatology of Rossby wave breaking in the 320-2000K layer. *J. Atmos. Sci.*, 64, 1922-1932, 2007.
- 1030
- Holton J.R. and Alexander, M.J.: The role of waves in the transport circulation of the middle atmosphere. *Geophys Monograph Ser.*, 123, 21–35. 2000.
- Hoskins, B. J., McIntyre, M.E., and Robertson, A.W.: On the use and significance of isentropic potential vorticity maps. *Quart. J. Roy. Meteorol. Soc.*, 111, 877-946, 1985.
- 1035
- Ishioka K., and Yoden, S. : Non-linear aspects of a barotropically unstable polar vortex in a forced-dissipative system: Flow regimes and Tracer transport. *J. Met. Soc. Japan*, 201-212, 1995.
- Karpechko, A. Y., Charlton-Perez, A., Balmaseda, M., Tyrrell, N., and Vitart, F.: Predicting sudden stratospheric Warming 2018 and its climate impacts with a multimodel ensemble. *Geophysical Research Letters*, 45, 13,538–13,546. <https://doi.org/10.1029/2018GL081091>, 2018.
- 1040
- Knight, J., Scaife, A., Bett, P.E., Collier, T., Dunstone, N., Gordon, M., Hardiman, S., Hermanson, L., Kay, G., McLean, P., Pilling, C., Smith, D., Stringer, N., Thornton, H. and Walker, B. : Predictability of European Winters 2017/2018 and 2018/2019: Contrasting influences from the tropics and stratosphere. *Amos. Sci. Lett.* 22: e1009. <https://doi.org/10.1002/asl.100>, 2021.
- Kuchar, A., Öhlert M., Eichinger, R., and Jacobi, C.: Large-ensemble assessment of the Arctic stratospheric polar Vortex morphology and disruptions. *Weather Clim. Dynam.*, 5, 895–912, 2024. <https://doi.org/10.5194/wcd-5-895-2024>.
- 1045
- Lawrence, Z. D., & Manney, G. L.: Characterizing stratospheric polar vortex variability with computer vision techniques. *Journal of Geophysical Research: Atmospheres*, 123, 1510–1535., 2018. <https://doi.org/10.1002/2017JD027556>
- 1050
- Lawrence, Z.D. and Manney, G.L.: Does the Arctic stratospheric polar vortex exhibit signs of preconditioning prior to sudden stratospheric warmings? *J. Atmos. Sci.*, 77, 611-632, 2020.
- Leovy, C.: Radiative equilibrium of the mesosphere. *J. Atmos. Sci.*, 21, 238-248, 1964.
- Leutwyler, D. and Schär, C.: Barotropic Instability of a Cyclone Core at Kilometer-Scale Resolution. *J. Adv. Modeling Earth Syst.*, 11, 3390-3402, 2019.
- 1055
- Lighthill, M.J.: Waves in Fluids. *Comm. Pure & App. Maths*, 20, 267-293, 1967.
- Lu, Z., Li, F., Orsolini, Y.J., Gao, Y., and He, S. : Understanding of European Cold Extremes, Sudden Stratospheric Warming, and Siberian Snow Accumulation in the Winter of 2017/18. *J. Climate*, 33, 527-545, 2020.
- Lu, H., Gray, L.J., Martineau, P., King, J.V.C. and Bracegirdle, T.J.: Regime behavior in the upper stratosphere as a precursor of stratosphere–troposphere coupling in the Northern Winter. *J. Climate*, 34, 7677-7696, 1060 2021.
- Manney, G.L., Nathan, T.R., and Standford, J.L.: Barotropic stability of realistic stratospheric jets. *J. Atmos. Sci.*, 45, 2545-2555, 1988.
- Manney, G.L. and Nathan, N.R.: Stability of westward-moving waves in realistic stratospheric zonal flows. *J. Atmos. Sci.*, 47, 775-794, 1990.

- 1065 Manney, G.L., Butler A.H., Lawrence, Z.D., Wargan, K., and Santee, M.L.: What's in a name? On the use and significance of the term "Polar Vortex". *Geophys. Res. Letters*, 49, e2021GL097617, 2021.
- Martius, O., Polvani, L.M. and Davies, H.C.: Blocking precursors to stratospheric sudden warming events. *Geophys. Res. Letters*, 36, L14806, doi:10.1029/2009GL038776,2009.
- Matthewman, N. J., and Esler, J.G.: Sudden stratospheric warmings as self-tuning Resonances. Part I: Vortex
1070 splitting events. *J. Atmos. Sci.*, 68, 2481-2504, 2011.
- McIntyre, M.E., and Palmer T.P.: The 'surf-zone' in the stratosphere. *J. Atmos & Terres. Phys.* 46, 825-849, 1984.
- Michaelangeli, D.V. and Zurek, R.W. : Barotropic Instability of Midlatitude Zonal Jets on Mars, Earth and Venus. *J. Atmos. Sci.*, 44, 2031-2041, 1987.
- 1075 Mitchell, D. M., Scott, R.K., Seviour, W.J.M., Thomson, S.I., Waugh, D.W., Teanby, N.A., and Ball, E.R.: Polar vortices in planetary atmospheres. *Rev. Geophys*, 59, e2020RG000723, 2021.
- Mitzu, R. and Yoden, S.: Chaotic Mixing and transport barriers in an idealized stratospheric polar vortex. *J. Atmos. Sci.*, 58, 2616-2629, 2001.
- Nash, E. R., Newman, O.P.A., Rosenfield, J. E. and Schoeberl, M.R.: An objective determination of the polar
1080 vortex using Ertel's potential vorticity. *J. Geophys. Res.*, 101, 9471–9478, 1996.
- Paldor, N., Shamir, O. and Garfinkel, C.M.: Barotropic instability of a zonal jet on the sphere: from non-divergence through quasi-geostrophy to shallow water, *Geophysical Astrophysical Fluid Dynamics*, 115:1, 15-34, DOI:10.1080/03091929.2020.1724996, 2021.
- Perot, K., and Orsolini, Y.J.: Impact of the major SSWs of February 2018 and January 2019 on the middle
1085 atmospheric nitric oxide abundance. *J. Atmos & Solar-Terr Phys.*, 218, 2021.
<https://doi.org/10.1016/j.jastp.2021.105586>
- Pfister, L.: A theoretical study of three-dimensional barotropic instability with applications to the upper stratosphere. *J. Atmos. Sci.*, 36, 908-920, 1979.
- Plumb, R.A.: Instability of the distorted polar night vortex: A theory of stratospheric warmings. *J. Atmos. Sci.*, 38,
1090 2514-2531, 1981.
- Randel, W.J. & Lait, L.R.: Dynamics of the 4-day wave in the Southern Hemisphere polar stratosphere. *J. Atmos. Sci.*, 48, 2496-2508, 1991.
- Rao, J., Ren, R., Chen, H., Yu, Y., & Zhou, Y.: The stratospheric sudden warming event in February 2018 and its prediction by a climate system model. *Journal of Geophysical Research: Atmospheres*, 123, 13,332–
1095 13,345. <https://doi.org/10.1029/2018JD028908>.
- Rao, J., Garfinkel, C.I., and Butler A.H.: Stratospheric polar vortex variability. In 'Atmospheric Oscillations: Sources of Subseasonal-to-seasonal variability and predictability.' Chapter 14, 277-299, 2024. Elsevier.
- Schubert, W.H., Montgomery, M.T., Taft, R.K., Guinn, T.A., Fulton, S.R., Kossin, J.P., & Edwards, J.P.:
1100 Polygonal eyewalls, asymmetric eye contraction, and potential vorticity mixing in hurricanes. *J. Atmos. Sci.*, 56, 1197-1223, 2018.
- Serra, M., Sathe, P., Beron-Vera, F., and Haller, G.: Uncovering the edge of the polar vortex. *J. Atmos. Sci.*, 74, 3871-3885, 2017.
- Seviour, W.J.M., Waugh, D.W., and Scott, R.K.: The Stability of Mars's annular polar vortex. *J. Atmos. Sci.* 74, 1533-1547, 2017.
- 1105 Sharkey, J., Teanby, N.A., Sylvestre, M., Mitchell, D.M., Seviour, W.J.M., Nixon, C.A., and Irwin P.G.J.: Potential vorticity structure of Titan's polar vortices from Cassini CIRS observations, 2021, *Icar*, 354, 114030, 2021.

- Shine, K.P.: The middle atmosphere in the absence of dynamical heat fluxes. *Quart. J. Roy. Meteorol. Soc.*, 113, 603-633, 1987.
- 1110 Schoeberl, M. R. and Newman, P.A.: Polar Vortex. In: Gerald R. North, John Pyle and Fuqing Zhang (Editors). *Encyclopedia of Atmospheric Sciences*. 2nd Edition., Vol. 4, 12-17, 2015.
- Shultis, J., Seviour, W., Waugh D. and Toigo, A.: Transport and mixing in planetary polar vortices with annular and monopolar potential vorticity structures . *The Planetary Science Journal*, 6, 63, 2025.
- Simmons. A.J.: Baroclinic instability at the winter stratopause. *Quart. J. R. Met. Soc.*, 100, 531-540, 1974.
- 1115 Smith, A.K.: Preconditioning for Stratospheric Sudden Warmings: Sensitivity Studies with a Numerical Model. *J. Atmos. Sci.*, 49, 1003-1019, 1992.
- S-RIP (2022), SPARC Reanalysis Intercomparison Project (S-RIP) Final Report. M. Fujiwara, M., Manney, G.L., Gray, L.J. and Wright, J.S. (Eds.), SPARC Report No. 10, WCRP-17/2020,
- Venne, D. E. and Stanford., J.L.: Observations of a 4-day temperature wave in the polar winter stratosphere.
- 1120 *J. Atmos. Sci.*, 36, 2016-2019, 1979.
- Wang; Y., Shulga, V., Milinevsky, G., Patoka, A., Evtushevsky, O., Klekociuk, A., Han, W., Grytsai, A., Shulga, A., Myshenko, V., and Antyufeyev, O.: Winter 2018 major sudden stratospheric warming impact on midlatitude mesosphere from microwave radiometer measurements. *Atmos. Chem. Phys.*, 19, 10303–10317, 2019. <https://doi.org/10.5194/acp-19-10303-2019>
- 1125 Watanabe, S., Koshin, D., Noguchi, S., and Sato, K.: Gravity wave morphology during the 2018 sudden stratospheric warming simulated by a whole neutral atmosphere general circulation model. *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036718. <https://doi.org/10.1029/2022JD036718>, 2022.
- Waugh, D.M.: Fluid dynamics of polar vortices on Earth, Mars, and Titan. *Annu. Rev. Fluid Mech*, 55, 265-269,
- 1130 2023.
- Waugh., D.W. and Polvani, L.M.: Stratospheric Polar Vortices. In ‘The Stratosphere: Dynamics, Transport, and Chemistry. *Geophys. Monograph Series*, 190, 43-57, 2010.
- Waugh, D.W., Sobel.A.H., and Polvani, L.M.: What is the polar vortex and how does it influence weather? *Bull. Amer. Meteor. Soc.*, 98, 37-44, 2016.
- 1135 Yessimbet, K. , Shepherd, T.G., Osso, A.C. and Steiner, A.K.: Pathways of influence between Northern Hemisphere blocking and stratospheric polar vortex variability. *Geophys. Res. Letters*, 49, e2022GL100895. <https://doi.org/10.1029/2022GL100895> 2022.