

Continuous *in situ* monitoring of ~~the vertical and horizontal passage of a labeled~~labeled-water pulse through a boreal Scots pine forest: vertical and horizontal fluxes

John D. Marshall^{1,2,3}, Maren Dubbert⁴, Teresa E. Gimeno⁵, Ruth-Kristina Magh^{6,2}, Kathrin Kühnhammer⁷, David Dubbert⁴, Paul Koeniger⁸, Matthias Cuntz⁹, Matthias Beyer¹⁰

¹Department of Geosciences, University of Gothenburg, Gothenburg, Sweden

²Forest Ecology and Management, Swedish University of Agricultural Sciences (SLU), SE-901 83 Umeå, Sweden

³Department of Matters and Energy Fluxes, Global Change Research Institute, 60300 Brno, Czechia

⁴Isotope Biogeochemistry and Gas Fluxes, Leibniz Institute for Agricultural and Landscape Research (ZALF), 15374 Müncheberg, Germany

⁵CREAF, 08193 Bellaterra (Cerdanyola del Vallès), Catalonia, Spain,

⁶Terrestrial Ecohydrology, Friedrich Schiller University, 07749 Jena, Germany

⁷Ecosystem Physiology, Albert-Ludwigs-Universität Freiburg, Germany

⁸Federal Institute for Geosciences and Natural Resources (BGR), D-30655 Hannover, Germany

⁹Université de Lorraine, AgroParisTech, INRAE, UMR Silva, 54000 Nancy, France

¹⁰TU Braunschweig, IGÖ - Abt. Umweltgeochemie, 38106 Braunschweig, Germany

Corresponding Authors:

John D. Marshall john.marshall@gu.se

Matthias Beyer, matthias.beyer@tu-bs.de

Abstract

Water ~~labe~~labeled with stable isotopes provides a conservative tracer, being neither produced nor consumed, for water flowpaths within soils and root systems. We added a strong, evenly distributed ²H—~~HO~~²HHO label to one m² of soil surface and continuously monitored its passage downward into the soil and upward into the stems of surrounding trees, with the objective of illuminating spatiotemporal lateral root water uptake and overlap. The study was conducted during the historic drought of 2018 in a mature Scots pine (*Pinus sylvestris*) forest growing on sandy soil in northern Sweden. Continuous *in situ* isotopic measurements of tree xylem water evidenced root system overlap of six trees within the ~~labe~~labeled square meter.

This result is —consistent with previous estimates from labelled nutrient uptake —measurements at this site. However, —label uptake differed sharply among trees, even —within the same radius; 90% of the label was taken up by one of the two —trees closest to the labelled plot.— Horizontal transport rates in tree roots averaged $0.17 \pm 0.05 \text{ m d}^{-1}$, meaning that the arrival of label pulse in tree stems was delayed by —6-33 days from first tree to last. Root water uptake by trees appeared restricted to the upper 60 cm of mineral soil, even at the peak of the drought-; the volume-weighted average depth of uptake was $30.6 \pm 1.2 \text{ cm}$ throughout the study period. Label intensity of the mineral soil weakened throughout the drought—, consistent with the notion that the label was being dispersed or diluted.—~~Label —ed into pre-event water recovery was low (3-4%), —and we hypothesize that this was due to a significant upward flux of water into the organic surface horizons.—~~ Our —data provide a daily —and three-dimensional description of the passage of a ~~label —ed~~labeled water pulse, highlighting the heterogeneity in horizontal water ~~transport flowpaths~~distribution by roots and the uneven partitioning of label among individual trees in a boreal forest.

1. Introduction

—Trees take up enormous quantities of water from the soil to grow and survive. ~~Th~~—is This uptake is driven by atmospheric demand for water vapor (Campbell and Norman, 2012), and is constrained by soil moisture availability and root distributions (Bachofen et al., 2024; McCulloh et al., 2003). In forests, roots of different individuals, —and their mycorrhizae—, extend and overlap both vertically and horizontally, —thus sampling across heterogenous water availabilities —(Bachofen et al., 2024; Goldsmith et al., 2019), while also competing for water and nutrients —(Henriksson et al., 2021; Lutter et al., 2021). So far,— vertical patterns of root water uptake have been studied more often than horizontal patterns (Beyer et al., 2016; Jackson et al., 1996; Stocker et al., 2023).

In one of the rare studies investigating horizontal uptake patterns in forests, isotopical label uptake was detected as far as 3-6 m from the stem ~~was found (—(~~ (Sternberg et al., 2002; Sternberg et al., 2005). Whether vertical or horizontal, long-distance transport comes at a considerable carbon expense to the tree (Guswa, 2008)—, —), which must ultimately limit the benefits of long horizontal roots. The costs and benefits are complicated by the near universality of common mycorrhizal networks (Martin and van der Heijden, 2024; Simard and Durall, 2004), which could extend uptake zones beyond the range of individual root systems. At present, the physiological significance of these networks is insufficiently quantified (Henriksson et al., 2023; Martin and van der Heijden, 2024). Here, we use “root” water uptake as shorthand to refer to tree roots plus their mycorrhizal partners.

These observations raise questions about the degree of —overlap in uptake zones between individual trees —(Henriksson et al., 2021; Kulmatiski et al., 2020). —One might argue that, at the extreme, ~~there would be no~~ root systems might be so territorial that we observe no overlap in water uptake and the label would be detected in only one tree around the plot. Each tree's roots systems would system might grow as if ~~each~~it owned property, perhaps avoiding other roots when they ~~were~~are approached (Mahall and Callaway, 1992). ~~On~~At the other ~~hand, roots~~extreme, root systems might

overlap extensively, making use of the limited ~~space~~soil volume in their immediate vicinity, whether it is occupied by other roots or not. Most data suggest the latter (Kulmatiski et al., 2010; Lutter et al., 2021; Lwila et al., 2024), but there are also instances of ~~the~~ “some degree of territoriality” ~~implied by the former~~ (e.g., Kuiper and Coutts, 1992). ~~Quantification~~ If this is a continuous variable, then quantification of this overlap might be as simple as counting the labelled trees around a 1-m² labelling plot. Such quantification is needed for scaling water and nutrient uptake from individual trees to forest stands (Manoli et al., 2014), as well as for estimating belowground competition ~~for studies of tree drought response~~, in silviculture and production ecology (Lutter et al., 2021). ~~Finally, the~~ The overlap might reveal novel underground ecological processes, for example water sharing ~~among individuals~~ (Kakouridis et al., 2022) ~~or stealing among individuals, which would influence the benefits of water-use efficiency (Cohen, 1970).~~

Tracing water flowpaths within soils is challenging and often involves destructive sampling, which implies large disturbances to the soil column and potentially irreversible harm to the root system. *In situ* methods, based on online monitoring of soil and tree xylem water isotopic composition, allow for tracking water through the soil-plant-atmospheric continuum in real time (Beyer et al. 2020—). But so far, there have been few isotopic assessments of water uptake dynamics (Gessler et al., 2022; Kühnhammer et al., 2022, 2023; Landgraf et al., 2022; Seeger and Weiler, 2021; Volkmann et al., 2016).

We studied these questions using *in situ* methods in a boreal forest in northern Sweden during the “drought of the century” —in— the summer of 2018 (Gutierrez Lopez et al., 2021). This episode provided a unique condition wherein long horizontal or deep vertical roots might have been important, especially if they reached into distant water sources —(Bachofen et al., 2024). Such a shift in water sources might (Lindroth et al., 2020) or might not (Gessler et al., 2022) compensate for soil drying, holding canopy evapotranspiration constant (Bachofen *et al.*, 2024). Responses to this drought event have already been described elsewhere in terms of sap flow (Gutierrez Lopez et al., 2021), latent heat flux (Lindroth et al., 2020; Mensah et al., 2021), and vertical water uptake patterns (Gessler et al., 2022). Horizontal water uptake over the whole growing season was described in an earlier study at the same site (Henriksson et al., 2021), but the short-term dynamics during the peak of the drought and the recovery period have not been investigated previously.

Here, we monitored continuously and *in situ* soil and tree xylem water isotopic composition during the peak and drought recovery periods. This experiment followed the dynamics and spatial distributions of a point-labe—ling labeling with ²HHO in a mature pine forest, ~~where considerable root—interaction had previously been reported (Henriksson et al., 2021).~~ We aimed to quantify the interaction using—degree of territoriality in water uptake ~~patterns among co-occurring by counting~~

~~labelled~~ tree individuals—, to quantify rates of horizontal water transport through roots, and to describe the fate of a label ~~pulse~~ applied to the mineral soil surface of a deep sandy soil.

2. Materials and Methods

2.1 Study site

This study was conducted in a ~100-year-old naturally regenerated, homogeneous— forest dominated by Scots pine (*Pinus sylvestris* L.), in northern Sweden (Rosinedalsheden, 64°10'N, 19° 45'E, 145 m a.s.l.). Understory vegetation was composed of ericaceous shrubs (*Vaccinium myrtillus* L. and *Vaccinium vitis-idaea* L.) and a layer of mosses and lichens. In 2013, the leaf area index was 2.7 m²m⁻². The stand averaged 18.6 (sd = 0.1) cm in diameter and 17.5 (sd < 0.1) m in total height in 2013 (Lim et al., 2015). The climate is typical of the northern boreal zone: average annual temperature and precipitation are 2.4 (sd = 0.8) °C and —638 (sd = 107) mm, respectively (Klosterhalfen et al., 2023)—. Winters are cold and long; summers are usually cool and wet. The photosynthetically active period extends from mid-April to mid-November (Tarvainen et al., 2018; Vernay et al., 2020). A soil description was performed at a nearby forest—, approximately 6 km away, under similar stand and landscape conditions. The soil was labeled a Regosol (FAO 1988) that had formed on fluvial sediment. The texture was considered loamy sand to sandy loam with lenses of silt loam at 25-45 cm depth. The clay content was less than 11% —(Plamboeck et al., 1999)—. The sample plot was located approximately 30 m from an eddy covariance system installed on a tower above the uniform canopy (21.5 m above the ground)) (Lindroth et al., 2020; Zhao et al., 2022). The eddy covariance system measured latent energy fluxes from the canopy over the whole study period using H₂O mixing ratios measured with a gas analyzer (LI-7200, LI-COR Environmental, Lincoln, USA) at 20 Hz frequency (Jocher et al., 2017).

The study was conducted from June to September 2018, when northern Scandinavia, like much of central Europe, underwent an exceptional drought; in Scandinavia, it was the driest period in fifty years (Gutierrez Lopez et al., 2021; Lindroth et al., 2020). From snowmelt (18 May) until 23 July, a period of more than two months, the study area received only 56 mm precipitation. The drought was then broken by a series of strong rainfall events that delivered 85 mm in eight days. We applied the soil label in the middle of the drought period and tracked its passage through the last two weeks of the drought and into the wetter period afterward.

2.2. Continuous measurements of water isotopic composition

2.2.1. Experimental design

We selected 15 Scots pines within a circular plot of 12 m radius around the ~~labe~~labeled area (Fig. 1). To study horizontal transport of water, we applied isotopic tracer on 10th of July 2018 over a 1-m²

area in the center of the experimental plot. We present the isotopic data in δ -notation (Gonfiantini, 1978), with units of per mil (‰) relative to the VSMOW-SLAP scale. The tracer consisted of —a mixture of 25 L of deionized tap water ($\delta^2\text{H}$: -90‰) with 200 mL of ^2H -enriched water ($^2\text{H}_2\text{H}_2\text{O}$, 99.5—%), yielding a ^2H atom fraction of 0.0163 ($\delta^2\text{H} \cong 106,000\text{‰}$). The ^2H -enriched water was injected at the surface of the mineral soil (i.e., underneath the moss and humus layers) by forcing a rigid plastic tube through the organic layer and injecting water into the created hole using a syringe. To ensure homogeneous distribution of the tracer, we applied 70 mL at each of five points within each 12.5×12.5 cm grid across the 1-m² label area, totaling 320 injections (22.4 L). After injections on the grid were complete, we injected the small volume of remaining label randomly within the ~~labeled~~ area.

For the sake of completeness, it should be noted that two additional ~~label~~—~~inglabeling~~ experiments were attempted. The first, on the 6th of June, 2018 (soon after snowmelt), consisted of applying a weaker ($\delta^2\text{H} = 3500\text{‰}$) ~~label~~ to the same 1-m² plot. The label was briefly detected in the surface soils but did not show up in the trees at all. Therefore, we repeated the ~~label~~—~~inglabeling~~ with the stronger ($\delta^2\text{H} \cong 106,000\text{‰}$) label solution described above, on July 10th. The second experiment, on the 20th of July 2018, was designed to detect deep water uptake (Burgess et al., 1998; Caldwell and Richards, 1989; Dawson and Pate, 1996; Moreira et al., 2000). We applied 1 L of deionized water containing 2.94 mL of H_2^{18}O ($\delta^{18}\text{O} \cong 470\text{‰}$), at a depth of 50 cm, directly under the stems of five trees (see Beyer *et al.*, (2016, 2018) for further application details). We did not detect the ^{18}O label in any of these trees (data not shown). However, to avoid potential interferences between ~~label~~ ~~inglabeling~~ experiments, these trees were dropped from consideration. They are represented by purple symbols in Figure 1 and not addressed further.

2.3 Volumetric water content

—Volumetric soil —water content (VWC) was measured along a vertical profile inside the ~~labeled~~ area using TDR probes (5TM, Metergroup, Pullman, USA). The TDR probes were placed at 10, 30, 50, 70, and 100 cm soil depth. For our sandy soils, we used the factory-default calibration. We took particular care when installing —both the TDR and the isotope probes (see 2.3—.2) to avoid damaging roots during probe installation, hoping that the roots from all surrounding trees would continue to take up water normally from the ~~labeled~~ block of soil. We estimated total water uptake by measuring the daily decline in —VWC at each depth, multiplying VWC by the soil depth represented by that sample, and— accumulating the decline over all depths (the single-step, multi-layer water balance of Guderle and Hildebrandt, 2015):

$$\Sigma U = \Sigma (\Delta \text{VWC}_{10} * D_{10} + \Delta \text{VWC}_{30} * D_{30} + \Delta \text{VWC}_{50} * D_{50}) \quad \text{Eq. 1}$$


Where $\overline{UD}$ is ~~mean~~the sum of uptake ~~depth~~ per day, ΔVWC_n is the daily change in VWC at depth n , D_n is the thickness of the layer represented by each VWC (always 20 cm). This approach yields a water consumption rate (ΣU) in cm day^{-1} , integrated over the upper 60 cm of mineral soil. Because the uppermost TDR probe was located 10 cm deep in the mineral soil and was covered by the forest floor, we assumed there would be little contribution of soil surface evaporation to this measured change in VWC and that it would therefore be dominated by root uptake. We return to the broader topic of surface evaporation in the discussion.

We then measured the mean depth of uptake was by weighting $\Delta VWC_n \cdot D_n$ of each layer by its mean depth, summing, and dividing by the total uptake, thus:

$$UD = \frac{\Delta VWC_{10} \cdot D_{10} \cdot 10 + \Delta VWC_{30} \cdot D_{30} \cdot 30 + \Delta VWC_{50} \cdot D_{50} \cdot 50}{\Sigma U} \quad \text{Eq. 2}$$

Where UD is mean uptake depth and 10, 30, 50 cm are the depths of each VWC sensor. ΣU

~~is the sum of uptake (Eq. 1).~~



~~This approach yields a water consumption rate (ΣU) in cm day^{-1} , integrated over the upper 60 cm of mineral soil. Because the uppermost TDR probe was 10 cm deep in the mineral soil and covered by the forest floor, we assumed there would be little contribution of soil surface evaporation to this flux.~~




Fig. 1. Map of studied trees (numbered and yellow-filled circles). The 1-m² plot ~~labe~~ ~~edlabeled~~ with ²H₂O is shown in the center as a red square. The trees ~~label~~ ~~edlabeled~~ with H₂¹⁸O and subsequently dropped from the study are ~~shown in purple-filled with white~~. Two Picarro L2130i analyzers were placed in a ~~covered~~ box ~~at adjacent to the triangle plot center~~. Horizontal distance from the center of the ~~label~~ ~~edlabeled~~ plot is indicated by the blue concentric rings and blue numbers. Distances of individual trees from the plot center are presented in Table S1.

2.3. 2. In situ measurements of the soil and plant water isotopic composition

From June to September 2018, we measured the stable isotopic composition ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) and soil VWC across depth inside the 1m² plot. The isotopic composition was measured with custom-made equilibration probes inserted at depths of 5, 10, 15, 20, 30, 50, 70, and 100 cm depth in the center of the ~~label~~ ~~edlabeled~~ soil volume. The probes consisted of two PFA Teflon tubes (1/8" AD, 1/16" ID, Teddington AB, Skogås, Sweden) that were glued (Pattex Kraft-Mix, Henkel AG & Co. KGaA, Düsseldorf, Germany) into a gas-permeable membrane (Accurel GP V8/2HF, 3 M, Germany; 0.155 cm wall thickness, 0.55 cm i.d., 0.86 cm o.d.) of 10 cm length. One of the two Teflon tubes was inserted a short distance into the probe. It provided air drawn through a desiccant tower (Drierite, Hammond,

Xenia, OH, USA) reducing potential contamination with ambient water vapor. The second tube was placed at the opposite end of the probe to ensure that the air stream would equilibrate isotopically with soil —water before leaving the probe. This tube was connected to a Picarro 16-port distribution manifold (A0311, Picarro Inc., Santa Clara, USA). The manifold was connected to a Picarro L2130i CRDS (Cavity Ring-Down Spectroscopy) water isotope analyzer. —We verified that materials used did not affect measured isotopic composition by placing them in ambient air and comparing to measurements without soil probes. To prevent condensation, the PFA tubes were heated by electrical heating lines (TE's Raychem, TE Connectivity, Schaffhausen, Switzerland) wrapped in foam insulation tubes (Armacell, Münster, Germany).

We also measured the isotopic composition of tree xylem using the borehole equilibration method (Marshall et al., 2020). To measure xylem $\delta^2\text{H}$ and $\delta^{18}\text{O}$, a borehole was drilled all the way through the center of each tree. All trees were cored along the N-S axis at a height of 1.3 m using an electric drill with a 5-mm bit. We rinsed the boreholes with acetone to avoid the obstruction of cells by pitch (Marshall et al. 2020). ~~One end~~The inlet of the borehole was open to the atmosphere to allow equilibration of ambient water vapor with liquid xylem water evaporated inside the borehole. We have previously shown that equilibration occurs even if ambient water vapor is added at the borehole inlet (Marshall et al., 2020). ~~The other end~~Although the continuity of the xylem tracheids is broken by the drilling, the borehole walls remain in communication with the flowing water in the adjacent xylem (Marshall et al., 2020) such that arrival time of a water label matches predictions based on sap flow (pers. comm., Kühnhammer). The outlet of the borehole was coupled to 6-mm PFA tube (Teddington AB, Skogås, Sweden) using stainless steel connectors (Swagelok) plumbed to a rotary 16-port valve (Vici, Valco Europe, Schenkon, Switzerland). The valve was connected to a second Picarro L2130i CRDS water isotope analyzer. Again, electrical heating lines wrapped in foam insulation pipes were installed to prevent condensation in the PFA tubing and the valve. This setup allowed continuous measurements of water concentration and isotopic composition of the water vapor equilibrated inside the borehole and allowed us to calculate the isotopic composition of the xylem water (Marshall et al., 2020).

All trees and the water isotope values of the atmosphere were monitored sequentially. Automatic calibration of the analyzer was performed every five hours using the Picarro Standard Delivery Module (SDM). The SDM injects a water standard of known isotopic composition into the analyzer's vaporizer. Each of the standards was measured in vapor phase for 20 minutes at two flow rates: $0.08 \mu\text{L s}^{-1}$ and $0.05 \mu\text{L s}^{-1}$. The two injection rates produced different water ~~vapo~~vapor concentrations, which we used to correct for the concentration-dependence of the isotope measurements. —At each injection rate, we used two standards covering the range of expected variability in $\delta^2\text{H}$ —and $\delta^{18}\text{O}$: depleted deionized water (~~$\delta^2\text{H}$~~ $\delta^2\text{H}$ = -94.28 ‰ and $\delta^{18}\text{O}$ = -12.87 ‰) and a homemade $\delta^2\text{H}$ -~~enriched~~ standard ($\delta^2\text{H}$ —= 225.8 ‰, $\delta^{18}\text{O}$ = -12.70 ‰). Any drift in instrument readings would have been

detected and corrected for the multiple calibrations within each day. The isotopic composition of the two standards was determined on an isotope ratio mass spectrometer (DeltaV, Thermo Fisher) at the Stable Isotope Laboratory of the Swedish University of Agricultural Science (Umeå, Sweden). Each tree was measured for at least 30 minutes and the duration of the measurement interval for each tree was adjusted depending on the tube length. To avoid memory effects when the valve had just turned, we used only the data from the last five minutes in each valve position. To further avoid non-steady state conditions, as when condensation may have been present or stem temperatures were rapidly changing, we deleted values of water concentration with a standard deviation higher than 1,000 $\mu\text{mol mol}^{-1}$ or values of water concentration that were higher than 50,000 $\mu\text{mol mol}^{-1}$.

Borehole temperature is a critical parameter for calculating xylem isotopic composition because it is needed for the conversion of water isotope values in measured vapor phase to liquid xylem water (Eq. 1-3). Copper-constantan thermocouples (Omega Engineering, Norwalk, CT, USA) were installed at approximately 2 cm depth within the borehole and in contact with the surface of the xylem tissue to measure the temperature of the xylem surface. We further filtered all data for which difference in temperature between the borehole and the atmosphere was higher than 5 degrees, which occurred most often when the temperature was rapidly changing in the morning and evening. As above, this was done to reduce the effects of condensation within the borehole or the tubing and to avoid the risk of non-steady state temperatures conditions within the borehole.

2.4 —Data processing and analyses

2.4.1 —Calculation of $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of xylem water

After calibration and instrument drift-correction, the xylem liquid water $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values were estimated using the relations proposed by Majoube (1971) for describing the equilibrium fractionation between the liquid and vapor phases of $\delta^2\text{H}$ and $\delta^{18}\text{O}$ based on the temperature (T in K) measured by the thermocouples in the borehole:

$$-1000 \ln \alpha_{2H} = \frac{24.844 \times 10^6}{T^2(K)} - \frac{76.248 \times 10^3}{T(K)} + 52.612 \quad \text{Eq. 3}$$

$$-1000 \ln \alpha_{18O} = \frac{1.137 \times 10^6}{T^2(K)} \quad \text{Eq. 4}$$

The resulting α -values were multiplied by the raw isotope ratios (R, e.g., $^2\text{H}/^1\text{H}$) of equilibrated water vapor (R_{2Hvap}) to estimate liquid water isotopic ratio (R_{2Hliq}):

$$R_{2Hliq} = R_{2Hvap} * \alpha^2H \quad \text{Eq. 5}$$

They were then converted back to delta notation.

The soil liquid water values were estimated using a “regression” method that uses a polynomial calibration curve to simultaneously account for isotopic equilibration (Majoube, 1971), device-specific effects of water vapor concentration, and the interaction of the sampling probe with the soil (Beyer et al., 2018, 2020; Oerter and Bowen, 2017). Regressions were fitted to measurements of dried bulk soil from the field site, placed in impermeable bags and filled with liquid water of known isotopic composition (“standards”). As with the xylem, these calibration measurements were made each time the manifold cycled through the samples throughout the field campaign. The resulting regression equations were:

$$^2H_{liquid} = (2.631e-04 * ^2H_{vapor})^2 + 1.216 * ^2H_{vapor} - 0.001379 * H_2O + 119.9 \quad Eq. 6$$

with adjusted $R^2 = 0.9987$, and:

$$^{18}O_{liquid} = (-6.318e-09 * H_2O)^2 + 0.7895 * ^{18}O_{vapor} + 0.00005189 * H_2O + 5.754 \quad Eq. 7$$

with adjusted $R^2 = 0.85$.

2.4.2 Calculation of line-conditioned lc-excess

To detect subtle increases in the δ^2H -enriched water in the trees, we used the local meteoric water line to calculate line-conditioned deuterium-excess (lc-excess) (Landwehr and Coplen, 2006). The local meteoric water line was estimated from precipitation collected at the Svartberget field station (<https://www.icos-sweden.se/Svartberget>), 8 km away. There were 225 samples collected at least biweekly during the whole of 2017 and 2018. The resulting LMWL was:

$$\delta^2H = 1.32 + 7.44 * \delta^{18}O \quad Eq. 8$$

with $R^2 = 0.97$.

The lc-excess was then calculated as:

$$lc-excess = \delta^2H_{measured} - (1.32 + 7.44 * \delta^{18}O_{measured}) \quad Eq. 9$$

Values of $\delta^{18}\text{O}$ -excess above zero were interpreted as evidence of the uptake of $\delta^2\text{H}$ -enriched water. The advantage of this approach is that it removes background variation in $\delta^2\text{H}$ that is correlated with $\delta^{18}\text{O}$, revealing low levels of $\delta^2\text{H}$ -enrichment more clearly than the raw isotopic compositions. All analyses were performed using either excel or the base package of R (R Core Team (2023)).

3. Results

The summer of 2018 began with only one significant rainfall after the snowpack melted. During the drought that followed, ecosystem evapotranspiration fell by about 50% —but recovered immediately after the first rains (Fig. 2).





Fig. 2. Daily evapotranspiration at the Rosinedalheden boreal pine forest in calendar year 2018. The yellow box highlights the period of the drought. The drought began on day 174, more than a month after the snowpack melted (blue line), and was interrupted by only one small precipitation event (red bars). The drought ended with a series of large events beginning on day 204.

During the drought, the VWC fell at all depths within the upper 50 cm of soil (Fig. 3a). In contrast, the 70-cm depth showed almost no variation in VWC, consistent with our visual observations that there were few roots there. The isotopic $\delta^2\text{H}$ -enrichment, which amounted to 25

mm of water on this ~~plot~~, occurred on DOY 191, and was accompanied by small, short-lived increases in VWC at 10 and 30 cm. The depth-weighted VWC in the upper 60 cm of soil (Fig. 3b) declined by 0.57 mm day⁻¹ in the early part of the drought and 0.83 ~~m~~mm day⁻¹ during the latter part (both $R^2 > 0.99$). Mean water uptake depth (Fig. 3c) reflected the steep decline at 30 cm depth, averaging approximately 30 cm throughout the drought. Brief excursions were detected after the water additions due to ~~label~~inglabeling (DOY 191) and the first rains (DOY 203). In general, mean water uptake depth moved towards the surface during the drying event. After the rains began on DOY 203, VWC sharply increased at all depths down to 50 cm, with the sharpest increase detected at 30 cm.



330

331

332




Fig. 3. a) Soil VWC (VWC) by depth under the ~~label~~~~ed~~~~labeled~~ 1-m^2 area; b) Total water storage ~~—~~summed over ~~—~~60 cm of soil depth. Slopes of regression lines ~~—~~were fitted to the drought-induced decline, neglecting the day after the ~~labe~~~~ling~~~~—labeling~~ (m); c) ~~—~~ Mean daily water uptake depth. The strong positive excursions are associated with ~~label~~~~ing~~~~labeling~~ and rainfall events. In all panels, the drought period is shown by the yellow box and the ~~labe~~~~ling~~~~labeling~~ date is shown as a vertical, dashed red line at DOY 191.





Fig. 4. —Isotopic composition (δ^2H) of —soil water by depth showing a—) δ^2H of depths up to 15 cm, b—) δ^2H of depths 20 cm and below. Note the differences in y-axis scaling. . In both panels, the drought period is shown by the yellow box and the labelling date is shown as a vertical, dashed red line at DOY 191.

After the 3H —HO labe—ling 2H HO labeling, the isotopic signature observed in the 5-cm probe briefly rose above 35,000 ‰, in the 10-cm probe above 20,000 ‰, and in the 15-cm probe above 2,000‰ (Fig. 4a). The probes at 20, 30, 50, and 70 cm responded more slowly and less strongly (Fig. 4b—). We detected no increase in δ^2H at 100 cm depth by the end of August (DOY 240). Thus the

label was concentrated in the upper 10-15 cm, especially in the period immediately after the ~~label~~
~~ing-labeling~~. In these upper depths, the label intensity fell by around 1,000 % day⁻¹ over the
drought period and then dropped sharply when the rains arrived on day 204. Below, label intensity
increased at 20 and 30 cm even before the rains arrived. The probe at 50 cm detected the label weakly,
but the 70 cm probe detected it clearly; both the 50- and 70-cm probes responded about four weeks
after the ~~label~~ ~~ing-labeling~~. The $\delta^{18}\text{O}$ also increased at 5 and 10 cm after the ^2HHO labeling, but
the increase was at most ≈ 20 ‰ (Fig. S1), three orders of magnitude smaller than for $\delta^2\text{H}$ (Fig. 4a).






Fig. 5. Line-conditioned deuterium-excess (lc-excess) of xylem water in tree stems. Values above zero, denoted by the horizontal black line, show likely label uptake. Panel a—) Tree 1, near the soil-labelling plot; b—) Labeled trees with lc-excess at some point greater than zero; c—) Unlabeled trees, all more than 4 m from the center of the labelling plot. Note that the y-axes are on different scales and the zero-intercepts are at different locations on panel a vs panels b and c. In all panels, the drought period is shown by the yellow box and the labelling date is shown as a vertical, dashed red line at DOY 191.

Among the trees surrounding the 1-m² label plot, we detected the label in six (i.e. lc-excess > 0). Tree 1 showed by far the highest $\delta^2\text{H}$ values (Fig. 5a) and was responsible for 90% of the cumulative label water uptake in the study (Fig. S2). The $\delta^2\text{H}$ of the xylem sap of this tree (at 1.3 m height) began to change about four days after labelling and it reached its maximum (nearly 500 ‰) eight days after the labelling (Fig. 5a). The label intensity then decreased, maintaining

relatively high enrichment until the end of the experiment. In contrast, $\delta^{18}\text{O}$ of this tree displayed only a slight increase ($\approx 2\text{‰}$) during this period (Fig. S3). Four other trees on the plot increased $\delta^2\text{H}$ steadily over about two weeks after the labelling (Fig. 5b) reaching lc-excess values greater than zero. All these trees were within 4 m of the plot center. Three of the four trees within the 4-m radius began to respond before the rains started. Although it was more than 4 m from the plot center, Tree 12 also displayed lc-excess values greater than zero, but only briefly (Fig. 5b). The other trees that were more than 4 m from the plot center showed no response to the labelling (Fig. 5c).




Figure 6. — Distance from the plot center vs. number of days after labelling. — Only trees that responded to the labelling are shown.

Among the six trees that showed label uptake, the breakthrough time varied with distance (—Fig. 6). The first tree (Tree 1, 1.2 m from plot center, Table S1) required four days for label arrival and the

last (Tree 12, 6.7 m from plot center) required 33 days. The x-intercept was approximately equal to the label date (0 on this x-axis) and the slope of the regression line (0.172 (0.05) m day⁻¹) described the maximum horizontal transport rate.

4. Discussion

We added a strong $\delta^2\text{H}$ -labeled water pulse to 1 m² of the mineral soil surface in a mature boreal Scots pine forest and monitored water isotopic composition in soils and tree xylem continuously afterward. The label initially remained near the soil surface, but it gradually moved deeper into the soil and was eventually detected as deep as 70 cm. The label was also taken up by roots, as evidenced by the label presence in xylem water of six surrounding trees, including all trees within 4 m of the labeled plot center. The labeling occurred during a drought of rare intensity in the boreal north. We were able to observe labeling patterns both during the drought and after the rainfall events that relieved it.

Horizontal transport and xylem water isotope dynamics

The experiment was designed primarily to detect and quantify patterns of root (and mycorrhizal) overlap and dynamics of horizontal water uptake by trees. We found the label in stems of six surrounding trees, meaning that the root systems of individual trees do not express complete territoriality at this scale. We therefore conclude that all of these trees had roots in the labeled 1-m² plot. This is far more overlap than we observed in the crowns, which scarcely overlapped at all in the horizontal direction.

There are relatively few studies describing horizontal water uptake patterns in forest trees (Göttlicher et al., 2008; Henriksson et al., 2021; Lutter et al., 2021). Root distributions have been mapped in some temperate forests (Gao et al., 2021; Kuiper and Coutts, 1992; Lwila et al., 2024), but the presence of roots is not sufficient to predict water uptake (Göttlicher et al., 2008). Long-distance horizontal transport through overlapping root systems might increase uniformity in water availability among trees. It might also favour trees near openings by providing access to the unused soil water in the openings (Lutter et al., 2021; Moreira et al., 2000; Sternberg et al., 2002; Sternberg et al., 2005). This favouring might, as with deep vertical roots, provide a “lifeline” to trees growing under extreme drought conditions (Bachofen et al., 2024). However, it might also represent a significant carbon cost (Guswa, 2008), perhaps reducing partitioning to aboveground production (Fernandez-Tschieder et al., 2024; Marshall et al., 2023).

The rate of uptake varied among tree individuals. For example, the stem of Tree 2 was at the edge of the labeled plot (0.7 m from the center of the plot) and yet it showed far less label than Tree 1, which was slightly farther away (Figs. 5a and 5b). The label intensity detected in tree xylem became rather weak with greater distance from the plot, which

we addressed by calculating the lc -excess. This parameter corrected for event-based variation using the strong correlation between natural abundance of δ^2H and $\delta^{18}O$, allowing us to detect low levels of labeling. When we focused simply on the first appearance of the label, as defined by lc -excess greater than zero, we found that there was a correlation with distance from the plot (Fig. 6), suggesting that the label took some time to traverse the horizontal distance to the stems. The slope of this relationship, $0.17 (0.05) \text{ m day}^{-1}$, is an estimate of the maximum horizontal transport rate. For comparison, measurements of water flow through pine stems are typically around 1 m day^{-1} (Plamboeck et al., 1999; Tarvainen et al., 2018). Note also that the label appeared in Tree 1 in only four days. This suggests that the surface roots were alive and ready to begin water uptake, despite the severe drought (Marshall, 1986). The peak label intensity in Tree 1 appeared eight days after the labeling, perhaps due to the arrival of label via slower flow paths after its initial breakthrough (Gao et al., 2021; Seeger and Weiler, 2021; Werner et al., 2021). Alternatively, as the drought advanced, the tree could have increased the amount of water taken up from deeper soil layers (e.g., Gessler et al. 2022); however, our VWC data suggest that deep water uptake did not increase as a proportion of the total (Fig. 3c) late in the drought.

We also observed a brief, weak pulse in $\delta^{18}O$ of the surface soil horizons after the labeling. This might have occurred due to $\delta^{18}O$ -enrichment of the 2H_2O used for labeling (not measured), but we believe it is more likely interference in the spectrometer by the strong $^2H^1H^{16}O$ absorption peak with the adjacent $^1H^1H^{18}O$ peak, which has been observed in other studies using high concentrations of δ^2H (Beyer et al., 2016, 2018). In any case, the apparent $\delta^{18}O$ pulse was orders of magnitude weaker than the δ^2H pulse and can be considered negligible.

Ownership hypothesis

— Territoriality

The water uptake benefits of long horizontal roots could be assessed not only by the detection of label in surrounding trees, but also by the label intensity. One can imagine a continuum between complete domination of uptake by a single tree, analogous to owned and defended property, vs. complete sharing of the benefit, perhaps modified by radial distance from the tree stem. Although six trees were labeled by all trees within some radial distance from the tree stem. The distinction is important, in part, because conservation of soil water for later use would be favored by territorial rooting and disfavored by extensive root overlap (Cohen, 1970). Although six trees were labeled in our plot, we detected a strong preference for uptake by one of the two trees nearest the plot center, while the other nearest tree at a similar distance took up very little of the label. This preference remained remarkably constant even as the label pulse dissipated (Fig. S2). These results are similar to the cumulative 2H - HO label ing observed in tree rings of trees in a replicated experiment at the same site, also conducted in 2018 (S2).

In order to investigate this “~~ownership hypothesis~~” territoriality further, ~~w~~e have used the data from this earlier experiment to rank the trees from each of their plots according to label intensity (Henriksson et al., 2021) (Fig. 7). ~~Such a~~The ranking shows a strong preference for the ~~label~~ inglabeling of one or two trees, with much lighter ~~labe~~linglabeling in several more (Fig. 7b). Henriksson et al., (2021) found that ~~—~~the most heavily ~~label~~edlabeled trees were not necessarily those nearest the ~~labe~~ledlabeled plot. We interpret these and our results as evidence of ~~dominance~~territorial water uptake, but not to the point of exclusion, ~~leaning toward~~.

Our results may have been influenced by choosing a microsite with particularly high tree density (1600 trees per hectare relative to 1010 (SD= 125) trees per hectare in the defended property end of the water-sharing continuum introduced above. The stand (Lim et al., 2015)). This was done to facilitate access to the central isotopic analyzer. However, in an earlier study in the same stand with 1-m² water-labelled plots, Henriksson et al. also reported 8 (SE=2) labelled trees per plot. We therefore conclude that our results were not atypical.

The territoriality interpretation draws attention to the orientation of the trunk boreholes in our study. Ours were oriented north-south while the boreholes in Henriksson et al. (2021) were oriented toward the plot center. ~~—~~We initially believed that borehole orientation would not matter because ~~—~~an earlier report~~—~~ had shown that label was evenly distributed throughout the tree circumference rather quickly in white pine (White et al., 1985). More recent evidence from our site (Tarvainen et al., 2018) suggests that ~~this~~the stem label distribution is not ~~true~~—. ~~—~~uniform near the ground. Our ~~fixed~~ borehole orientation (N-S) ~~—~~may have contributed to the weak ~~labe~~linglabeling observed for ~~—T~~ ~~ree~~Tree 2 (on the western edge of the ~~label~~edlabeled plot) compared to ~~—~~Tree~~—~~ 1 (to the northeast). ~~We~~ despite the proximity of Tree 2 to the labelled plot. Therefore, we recommend orienting boreholes toward the ~~labelling plot~~center of small labelled plots in future work ~~studying horizontal water uptake patterns. However, even if the label intensity was weakened by the borehole orientation, the timing and magnitude of the label pulse among trees is interpretable based on the uniform timing (Figs. 5a, 5b) and distance effects (Fig. 6) in the trees where label was detected.~~




Fig. 7. Water isotopic composition ($\delta^2\text{H}$) of xylem wood in trees surrounding three ~~labelled~~ ~~edlabelled~~ 1-m² plots (data reprinted with permission from Henriksson et al., 2021). The trees surrounding each plot are ranked according to ~~label~~ ~~inglabeling~~ intensity. Filled symbols represent trees that were identified as “~~label~~ ~~edlabeled~~.” Panel a) shows the full data range and panel b) highlights the transition from lightly labelled to ~~unlabelled~~ ~~edunlabelled~~ trees.

Soil water content and vertical root water uptake

We interpret ~~daily declines in soil~~ VWC during the drought period as evidence of tree root water uptake throughout the upper 50 cm of mineral soil (Fig. 3a). The slopes of the declines at 10 and ~~50 cm~~ 50cm are similar, but that at 30 cm is steeper. There was no evidence of change in water content at ~~70 cm~~ 70cm, consistent with our informal observation that roots were rare below 50 cm

depth. The integrated water loss rate averaged $0.057 \text{ cm day}^{-1}$ in the early part of the drought and increased to $0.083 \text{ cm day}^{-1}$ in the last part, meaning that water uptake from these depths increased slightly as the drought proceeded. However, this conclusion is based on only three measurements across depth, the uppermost at 10 cm. Note that if there were evaporation in addition to root water uptake from the 10 cm depth, it would be included in water uptake depth estimate. Although there was almost certainly some evaporation from the injection sites at the surface of the mineral soil (0 cm depth), we speculate that evaporation was small at the measured 10 cm depth.

The maximum amount of label found in the soil was 4-5%, which is small in comparison to the amount added (see “Partial Label Recovery” in Suppl. Materials). However, we did not measure evaporation from the soil surface during the labelling and thereafter, which probably consumed a majority of the remainder. We tried to minimize surface evaporation by injecting at the mineral soil surface, but perhaps the injected water was drawn back up into the litter and moss and humus layers and above the topmost soil probes by matric forces during the injection.

After the injection, the labelled water was expected to flow downward under unsaturated conditions for as long as the water potential exceeded field capacity (Bonan, 2015). Remembering that the drought had already begun when the label was applied, the downward flow was expected to cease when field capacity was reached. One might expect a bit of short-range mixing by diffusion to continue after this point. This is consistent with the timing and intensity of the label peaks in Figs. 4a and 4b, where the label intensity decreased sharply between 15 cm (2027‰ maximum) and 20 cm depth (400‰ maximum, 8 days later). The added label could also have escaped detection by flowing down preferential flow paths (Clothier et al., 2008; Weiler, 2017), e.g. due to differences in soil density or along macropores or tree roots, leading to uneven infiltration and potentially bypassing probes and sensors. This might explain, for example, the higher maximum labelling at 70cm than at 50cm and at 30cm than at 20cm. However, the labelling patterns are largely consistent with the field-capacity scenario.

The current study ~~is similar to~~ follows on the classic study of Plamboeck et al. (1999), which was conducted at a nearby site with similar soils. Plamboeck et al. (1999)— estimated that, without irrigation, the 2-cm humus layer provided 10% of total uptake in July. When we estimated the mean depth of water uptake (Fig. 3c), it was remarkably stable throughout this severe drought, with a slight upward shift toward the end. —Our estimated mean uptake depth (30.6 (SE=1.2) cm) was similar to a depth-weighted estimate based on Plamboeck et al.’s (1999) isotopic data (30.3 (0.07) cm). Both our results and those of Plamboeck et al. —(1999) suggest a rather constant water uptake depth —as a drought develops- (Fig. 3c). This differs from a global meta-analysis, which found a downward shift in response to drought (Bachofen et al., 2024). If our trees cannot shift deeper under drought, then it

may make the broad horizontal root distribution even more important as an emergency source of soil water.

Label Recovery

We used both VWC and le excess to estimate recovery of the labeled water pulse. The integrated VWC increased by 0.72 mm after the addition of 25 mm labeled water, representing 3% of the water volume added. The excess 2H represented 16.2 μm (16.2 mL) immediately after the label addition. This value is about 4% of the 400 mL 2HH_2O that was added to the plot. The close agreement between these percentages is encouraging, but their small magnitude highlights our inability to close the water budgets.

We speculate that the low recoveries are due to water absorption and root water uptake by the desiccated soil, tree roots, and understorey vegetation roots in the layers above our uppermost probe (5 and 10 cm below the surface of the mineral soil for δ^2H and VWC , respectively). For illustration, using the maximum VWC measured by Plamboeck et al. (1999) on a nearby site with similar soils and vegetation, we estimate that the humus layer could retain 6 mm of precipitation, the 0–10 cm mineral horizon could retain 20 mm more. Thus, added together, these two layers could have effectively retained all the water we added. The large rainfall event that broke the drought (37.2 mm) resulted in an increase of only 2.8 mm in total water storage (as in Fig. 3b), approximately 8% of the expected value. If 30% were lost to interception, then 26 mm fell to the soil. This would mean that ground vegetation, roots, litter, and upper 5 cm of soil retained $26 - 2.8 = 23.2$ mm of precipitation, which is slightly less than the 25 mm we added as label and would have only allowed a small proportion of the label to reach our probes. Our choice to inject the label at the mineral soil surface ensured that we detected some of the label in the mineral soil, but it appears it was impacted by significant losses to the upper mineral and the organic layers as well.

Simplified borehole method

The method used to measure xylem water here included several simplifications of the borehole method used in previous studies. In particular, we did not use dry air (e.g., Kühnhammer et al., 2022) relying instead on the two-way exchange and equilibration of water vapor during the borehole passage (Marshall et al., 2020). In addition, we relied on the pump within the analyzer to draw air through the boreholes under negative pressure, apparently without significant leakage into the downstream airflows. This eliminates the need for the upstream pump used in some other applications (e.g., Kühnhammer et al., 2022). Finally, we did not use any screening material to inhibit microbial growth (Landgraf et al., 2022), relying instead on the natural defensive compartmentation of the xylem to serve that purpose (Shigo, 1984). Although a visible band of brown tissue formed around the borehole, our data showed that the borehole vapor continued to respond to xylem water changes

until the end of the experiment, more than three months after the boreholes were drilled. One complication that remained was the placement of an analyzer in the field. Several recent papers describe methods for field sampling and storage of water vapor for stable isotope analysis (Havranek et al., 2020; Herbstritt et al., 2023; Magh et al., 2022), which would mean sacrificing the continuous data streams collected here, but would circumvent the need for an analyzer in the field and provide opportunities for more replication.

Simplified borehole method

The method used to measure xylem water here included several simplifications of the borehole method used in previous studies. In particular, we did not use dry air (e.g., Kühnhammer et al., 2022) relying instead on the two-way exchange and equilibration of water vapor during the borehole passage (Marshall et al., 2020). In addition, we relied on the pump within the analyzer to draw air through the boreholes under negative pressure, apparently without significant air leakage into the downstream flows. This eliminates the need for the upstream pump used in some other applications (e.g., Kühnhammer et al., 2022). Finally, we did not use any screening material to inhibit microbial growth (Landgraf et al., 2022), relying instead on the natural defensive compartmentation of the xylem to serve that purpose (Shigo, 1984). Although a visible band of brown tissue formed around the borehole, our data showed that the borehole vapor continued to respond to xylem-water changes until the end of the experiment, more than three months after the boreholes were drilled. This is somewhat surprising given that the drilling must cavitate the xylem conduits that are cut, but our data, like previously published results with this method (Kühnhammer et al., 2021, 2022; Marshall et al., 2020), suggest that hydraulic contact with the xylem flow is maintained. As a reminder, we note that modeled results suggest that the equilibration is so fast that it would best reflect the isotopic composition of the water in the last few mm of the borehole (Marshall et al., 2020). Fortunately, this is where the flow is fastest and the circumference is greatest, both contributing to high proportions of the total stem flux.

One complication that remained was the placement of an analyzer in the field. Several recent papers describe methods for field sampling and storage of water vapor for stable isotope analysis (Havranek et al., 2020; Herbstritt et al., 2023; Magh et al., 2022), which would mean sacrificing the continuous data streams collected here, but would circumvent the need for an analyzer in the field and provide opportunities for more replication.

These results provide a dynamic description of soil water storage, transport, and root water uptake, including passage of the label through surrounding tree stems as they take up the label for transpiration. The timing and detection of the label are consistent with expectations insofar as it supports the notion that the root systems of several trees overlap on each square meter of ground

surface. However, the concentration of the label in a single tree was inconsistent with the notion of extensive horizontal sharing. This question deserves more attention as it influences the scaling of transpiration from trees to stands.

Author contributions: JM, MD, and MB conceptualized the work. MB and JM managed data curation and analysis. Funding support was organized by MB and MD. Methodology was organized by MB, DD, PK, KK, MC, and JM. JM wrote the initial draft, but all authors contributed to writing and visualization.

~~Acknowledgments:~~

Acknowledgments:

We thank Noelia Saavedra, Nils Henriksson, and Hyungwoo Lim (SLU-Umeå) for their help with the ~~labe—ling~~ labeling and measurement of these plots, and Nils Henriksson for permission to reanalyze his published data. Tomas Lundmark and the Svartberget Field Station of SITES provided housing for the group during the most intense part of this fieldwork. Tomas also provided the Picarro ~~analyz~~ ~~er~~ analyzer that we used for the tree measurements. We are also grateful to the Svartberget field staff for provision of the eddy covariance and meteorological data. Hector Correia (TU-Braunschweig) helped with the field work. RKM acknowledges the funding bodies Knut and Alice Wallenberg foundation within the project Future Forests KAW 2018.0259. She was also financially supported by the AquaDiva CRC 1076 within the project B02. MD was supported by DFG MD1688/1-1 and 6-1. TEG received funding from the Spanish Ministry of Science and Universities (MCIN/AEI/10.13039/501100011033/, projects: CNS2024-154609 and RYC2021-031759-I) and from the Catalan Government (SGR-Cat 2021 00849). KK was financially supported by the Volkswagen Foundation (contract no. [A122505](#); reference no. 92889 to MB) ~~and the German Research Foundation (DFG, SFB 1537 ECOSENSE).~~ MC acknowledges support from a grant by the French National Research Agency (ANR, ANR-21-CE02-0033-01) and from a grant overseen by ANR as part of the “Investissements d’Avenir” program (ANR-11-LABX-0002-01, Lab of Excellence ARBRE). ~~the French National Research Agency (ANR, ANR-21-CE02-0033-01) and from a grant overseen by ANR as part of the “Investissements d’Avenir” program (ANR-11-LABX-0002-01, Lab of Excellence ARBRE).~~

5. References:

644 Bachofen, C., Tumber-Dávila, S. J., Mackay, D. S., McDowell, N. G., Carminati, A., Klein, T.,
645 Stocker, B. D., Mencuccini, M., and Grossiord, C.: Tree water uptake patterns across the globe, *New*
646 *Phytol.*, 242, 1891–1910, <https://doi.org/10.1111/nph.19762>, 2024.

647 Beyer, M., Koeniger, P., Gaj, M., Hamutoko, J. T., Wanke, H., and Himmelsbach, T.: A deuterium-
648 based labeling technique for the investigation of rooting depths, water uptake dynamics and
649 unsaturated zone water transport in semiarid environments, *J. Hydrol.*, 533, 627–643,
650 <https://doi.org/10.1016/j.jhydrol.2015.12.037>, 2016.

651 Beyer, M., Hamutoko, J. T., Wanke, H., Gaj, M., and Koeniger, P.: Examination of deep root water
652 uptake using anomalies of soil water stable isotopes, depth-controlled isotopic labeling and mixing
653 models, *J. Hydrol.*, 566, 122–136, <https://doi.org/10.1016/j.jhydrol.2018.08.060>, 2018.

654 Beyer, M., Kühnhammer, K., and Dubbert, M.: In situ measurements of soil and plant water isotopes:
655 a review of approaches, practical considerations and a vision for the future, *Hydrol. Earth Syst. Sci.*,
656 24, 4413–4440, <https://doi.org/10.5194/hess-24-4413-2020>, 2020.

657 Burgess, S. S. O., Adams, M. A., Turner, N. C., and Ong, C. K.: The redistribution of soil water by
658 tree root systems, *Oecologia*, 115, 306–311, <https://doi.org/10.1007/s004420050521>, 1998.

659 Caldwell, M. M. and Richards, J. H.: Hydraulic lift: water efflux from upper roots improves
660 effectiveness of water uptake by deep roots, *Oecologia*, 79, 1–5, <https://doi.org/10.1007/BF00378231>,
661 1989.

662 Campbell, G. S. and Norman, J.: *An Introduction to Environmental Biophysics*, Springer Science &
663 Business Media, 296 pp., 2012.

664 Dawson, T. E. and Pate, J. S.: Seasonal water uptake and movement in root systems of Australian
665 phraeatophytic plants of dimorphic root morphology: a stable isotope investigation, *Oecologia*, 107,
666 13–20, <https://doi.org/10.1007/BF00582230>, 1996.

667 Fernandez-Tschieder, E., Marshall, J. D., and Binkley, D.: Carbon budget at the individual-tree scale:
668 dominant Eucalyptus trees partition less carbon belowground, *New Phytol.*, 242, 1932–1943,
669 <https://doi.org/10.1111/nph.19764>, 2024.

670 Gao, D., Joseph, J., Werner, R. A., Brunner, I., Zürcher, A., Hug, C., Wang, A., Zhao, C., Bai, E.,
671 Meusburger, K., Gessler, A., and Hagedorn, F.: Drought alters the carbon footprint of trees in soils—
672 tracking the spatio-temporal fate of ¹³C-labelled assimilates in the soil of an old-growth pine forest,
673 *Glob. Change Biol.*, 27, 2491–2506, <https://doi.org/10.1111/gcb.15557>, 2021.

674 Drought reduces water uptake in beech from the drying topsoil, but no compensatory uptake occurs
675 from deeper soil layers: <https://pub.epsilon.slu.se/26344/>, last access: 4 March 2022.

676 Gessler, A., Bächli, L., Rouholahnejad Freund, E., Treydte, K., Schaub, M., Haeni, M., Weiler, M.,
677 Seeger, S., Marshall, J., Hug, C., Zweifel, R., Hagedorn, F., Rigling, A., Saurer, M., and Meusburger,
678 K.: Drought reduces water uptake in beech from the drying topsoil, but no compensatory uptake
679 occurs from deeper soil layers, *New Phytol.*, 233, 194–206, <https://doi.org/10.1111/nph.17767>, 2022.

680 Goldsmith, G. R., Allen, S. T., Braun, S., Engbersen, N., González-Quijano, C. R., Kirchner, J. W.,
681 and Siegwolf, R. T. W.: Spatial variation in throughfall, soil, and plant water isotopes in a temperate
682 forest, *Ecohydrology*, 12, e2059, <https://doi.org/10.1002/eco.2059>, 2019.

683 Gonfiantini, R.: Standards for stable isotope measurements in natural compounds, *Nature*, 271, 534–
684 536, <https://doi.org/10.1038/271534a0>, 1978.

685 Götlicher, S. G., Taylor, A. F. S., Grip, H., Betson, N. R., Valinger, E., Högberg, M. N., and
686 Högberg, P.: The lateral spread of tree root systems in boreal forests: Estimates based on ^{15}N uptake
687 and distribution of sporocarps of ectomycorrhizal fungi, *For. Ecol. Manag.*, 255, 75–81,
688 <https://doi.org/10.1016/j.foreco.2007.08.032>, 2008.

689 Guderle, M. and Hildebrandt, A.: Using measured soil water contents to estimate evapotranspiration
690 and root water uptake profiles – a comparative study, *Hydrol. Earth Syst. Sci.*, 19, 409–425,
691 <https://doi.org/10.5194/hess-19-409-2015>, 2015.

692 Guswa, A. J.: The influence of climate on root depth: A carbon cost-benefit analysis, *Water Resour.*
693 *Res.*, 44, <https://doi.org/10.1029/2007WR006384>, 2008.

694 Gutierrez Lopez, J., Tor-ngern, P., Oren, R., Kozii, N., Laudon, H., and Hasselquist, N. J.: How tree
695 species, tree size, and topographical location influenced tree transpiration in northern boreal forests
696 during the historic 2018 drought, *Glob. Change Biol.*, 27, 3066–3078,
697 <https://doi.org/10.1111/gcb.15601>, 2021.

698 Havranek, R. E., Snell, K. E., Davidheiser-Kroll, B., Bowen, G. J., and Vaughn, B.: The Soil Water
699 Isotope Storage System (SWISS): An integrated soil water vapor sampling and multiport storage
700 system for stable isotope geochemistry, *Rapid Commun. Mass Spectrom.*, 34, e8783,
701 <https://doi.org/10.1002/rcm.8783>, 2020.

702 Henriksson, N., Lim, H., Marshall, J., Franklin, O., McMurtrie, R. E., Lutter, R., Magh, R.,
703 Lundmark, T., and Näsholm, T.: Tree water uptake enhances nitrogen acquisition in a fertilized boreal
704 forest – but not under nitrogen-poor conditions, *New Phytol.*, 232, 113–122,
705 <https://doi.org/10.1111/nph.17578>, 2021.

706 Henriksson, N., Marshall, J., Högberg, M. N., Högberg, P., Polle, A., Franklin, O., and Näsholm, T.:
707 Re-examining the evidence for the mother tree hypothesis – resource sharing among trees via
708 ectomycorrhizal networks, *New Phytol.*, 239, 19–28, <https://doi.org/10.1111/nph.18935>, 2023.

709 Herbstritt, B., Gralher, B., Seeger, S., Rinderer, M., and Weiler, M.: Technical note: Discrete in situ
710 vapor sampling for subsequent lab-based water stable isotope analysis, *Hydrol. Earth Syst. Sci.*, 27,
711 3701–3718, <https://doi.org/10.5194/hess-27-3701-2023>, 2023.

712 Jackson, R. B., Canadell, J., Ehleringer, J. R., Mooney, H. A., Sala, O. E., and Schulze, E. D.: A
713 global analysis of root distributions for terrestrial biomes, *Oecologia*, 108, 389–411,
714 <https://doi.org/10.1007/BF00333714>, 1996.

715 Jocher, G., Marshall, J., Nilsson, M. B., Linder, S., De Simon, G., Hörnlund, T., Lundmark, T.,
716 Näsholm, T., Ottosson Löfvenius, M., Tarvainen, L., Wallin, G., and Peichl, M.: Impact of Canopy
717 Decoupling and Subcanopy Advection on the Annual Carbon Balance of a Boreal Scots Pine Forest as
718 Derived From Eddy Covariance, *J. Geophys. Res. Biogeosciences*, 123, 303–325,
719 <https://doi.org/10.1002/2017JG003988>, 2017.

720 Kakouridis, A., Hagen, J. A., Kan, M. P., Mambelli, S., Feldman, L. J., Herman, D. J., Weber, P. K.,
721 Pett-Ridge, J., and Firestone, M. K.: Routes to roots: direct evidence of water transport by arbuscular
722 mycorrhizal fungi to host plants, *New Phytol.*, 236, 210–221, <https://doi.org/10.1111/nph.18281>,
723 2022.

724 Klosterhalfen, A., Chi, J., Kljun, N., Lindroth, A., Laudon, H., Nilsson, M. B., and Peichl, M.: Two-
725 level eddy covariance measurements reduce bias in land-atmosphere exchange estimates over a
726 heterogeneous boreal forest landscape, *Agric. For. Meteorol.*, 339, 109523,
727 <https://doi.org/10.1016/j.agrformet.2023.109523>, 2023.

728 Kühnhammer, K., Dahlmann, A., Iraheta, A., Gerchow, M., Birkel, C., Marshall, J. D., and Beyer, M.:
729 Continuous in situ measurements of water stable isotopes in soils, tree trunk and root xylem: Field
730 approval, *Rapid Commun. Mass Spectrom.*, 36, e9232, <https://doi.org/10.1002/rcm.9232>, 2022.

731 Kühnhammer, K., van Haren, J., Kübert, A., Bailey, K., Dubbert, M., Hu, J., Ladd, S. N., Meredith, L.
732 K., Werner, C., and Beyer, M.: Deep roots mitigate drought impacts on tropical trees despite limited
733 quantitative contribution to transpiration, *Sci. Total Environ.*, 893, 164763,
734 <https://doi.org/10.1016/j.scitotenv.2023.164763>, 2023.

735 Kuiper, L. C. and Coutts, M. P.: Spatial disposition and extension of the structural root system of
736 Douglas-fir, *For. Ecol. Manag.*, 47, 111–125, [https://doi.org/10.1016/0378-1127\(92\)90269-F](https://doi.org/10.1016/0378-1127(92)90269-F), 1992.

737 Kulmatiski, A., Beard, K. H., Verweij, R. J. T., and February, E. C.: A depth-controlled tracer
738 technique measures vertical, horizontal and temporal patterns of water use by trees and grasses in a
739 subtropical savanna, *New Phytol.*, 188, 199–209, <https://doi.org/10.1111/j.1469-8137.2010.03338.x>,
740 2010.

741 Kulmatiski, A., Adler, P. B., and Foley, K. M.: Hydrologic niches explain species coexistence and
742 abundance in a shrub–steppe system, *J. Ecol.*, 108, 998–1008, [https://doi.org/10.1111/1365-](https://doi.org/10.1111/1365-2745.13324)
743 2745.13324, 2020.

744 Landgraf, J., Tetzlaff, D., Dubbert, M., Dubbert, D., Smith, A., and Soulsby, C.: Xylem water in
745 riparian willow trees (*Salix alba*) reveals shallow sources of root water uptake by in situ monitoring of
746 stable water isotopes, *Hydrol. Earth Syst. Sci.*, 26, 2073–2092, [https://doi.org/10.5194/hess-26-2073-](https://doi.org/10.5194/hess-26-2073-2022)
747 2022, 2022.

748 Landwehr, J. M. and Coplen, T. B.: Line-conditioned excess: a new method for characterizing stable
749 hydrogen and oxygen isotope ratios in hydrologic systems, *International Conference on Isotopes in*
750 *Environmental Studies*, Monte-Carlo, Monaco, 132–135, 2006.

751 Lechthaler, S., Kiorapostolou, N., Pitacco, A., Anfodillo, T., and Petit, G.: The total path length’s
752 hydraulic resistance according to known anatomical patterns: what is the shape of the root-to-leaf
753 tension gradient along the plant’s longitudinal axis?, *J. Theor. Biol.*,
754 <https://doi.org/10.1016/j.jtbi.2020.110369>, 2020.

755 Lim, H., Oren, R., Palmroth, S., Tor-ngern, P., Mörling, T., Näsholm, T., Lundmark, T., Helmisaari,
756 H.-S., Leppälammi-Kujansuu, J., and Linder, S.: Inter-annual variability of precipitation constrains
757 the production response of boreal *Pinus sylvestris* to nitrogen fertilization, *For. Ecol. Manag.*, 348,
758 31–45, <https://doi.org/10.1016/j.foreco.2015.03.029>, 2015.

759 Lindroth, A., Holst, J., Linderson, M.-L., Aurela, M., Biermann, T., Heliasz, M., Chi, J., Ibrom, A.,
760 Kolari, P., Klemetsson, L., Krasnova, A., Laurila, T., Lehner, I., Lohila, A., Mammarella, I., Mölder,
761 M., Löfvenius, M. O., Peichl, M., Pilegaard, K., Soosaar, K., Vesala, T., Vestin, P., Weslien, P., and
762 Nilsson, M.: Effects of drought and meteorological forcing on carbon and water fluxes in Nordic
763 forests during the dry summer of 2018, *Philos. Trans. R. Soc. B Biol. Sci.*, 375, 20190516,
764 <https://doi.org/10.1098/rstb.2019.0516>, 2020.

765 Lutter, R., Henriksson, N., Lim, H., Blaško, R., Magh, R.-K., Näsholm, T., Nordin, A., Lundmark, T.,
766 and Marshall, J. D.: Belowground resource utilization in monocultures and mixtures of Scots pine and
767 Norway spruce, *For. Ecol. Manag.*, 500, 119647, <https://doi.org/10.1016/j.foreco.2021.119647>, 2021.

768 Lwila, A., Ammer, C., Gailing, O., Leinemann, L., and Mund, M.: Root overlap and allocation of
769 above- and belowground growth of European beech in pure and mixed stands of Douglas fir and
770 Norway spruce, *For. Ecosyst.*, 11, 100217, <https://doi.org/10.1016/j.fecs.2024.100217>, 2024.

771 Magh, R.-K., Gralher, B., Herbstritt, B., Kübert, A., Lim, H., Lundmark, T., and Marshall, J.:
772 Technical note: Conservative storage of water vapour – practical in situ sampling of stable isotopes in
773 tree stems, *Hydrol. Earth Syst. Sci.*, 26, 3573–3587, <https://doi.org/10.5194/hess-26-3573-2022>, 2022.

774 Mahall, B. E. and Callaway, R. M.: Root Communication Mechanisms and Intracommunity
775 Distributions of Two Mojave Desert Shrubs, *Ecology*, 73, 2145–2151,
776 <https://doi.org/10.2307/1941462>, 1992.

777 Majoube, M.: Fractionnement en oxygène 18 et en deutérium entre l’eau et sa vapeur, *J Chim Phys*,
778 68, 1423–1436, 1971.

779 Manoli, G., Bonetti, S., Domec, J.-C., Putti, M., Katul, G., and Marani, M.: Tree root systems
780 competing for soil moisture in a 3D soil–plant model, *Adv. Water Resour.*, 66, 32–42,
781 <https://doi.org/10.1016/j.advwatres.2014.01.006>, 2014.

782 Marshall, J. D.: Drought and shade interact to cause fine-root mortality in Douglas-fir seedlings, *Plant*
783 *Soil*, 91, 51–60, 1986.

784 Marshall, J. D., Brooks, J. R., and Lajtha, K.: Sources of variation in the stable isotopic composition
785 of plants, *Stable Isot. Ecol. Environ. Sci.*, 22–60, 2007.

786 Marshall, J. D., Cuntz, M., Beyer, M., Dubbert, M., and Kuehnhammer, K.: Borehole Equilibration:
787 Testing a New Method to Monitor the Isotopic Composition of Tree Xylem Water in situ, *Front. Plant*
788 *Sci.*, 11, <https://doi.org/10.3389/fpls.2020.00358>, 2020.

789 Marshall, J. D., Tarvainen, L., Zhao, P., Lim, H., Wallin, G., Näsholm, T., Lundmark, T., Linder, S.,
790 and Peichl, M.: Components explain, but do eddy fluxes constrain? Carbon budget of a nitrogen-
791 fertilized boreal Scots pine forest, *New Phytol.*, 239, 2166–2179, <https://doi.org/10.1111/nph.18939>,
792 2023.

793 Martin, F. M. and van der Heijden, M. G. A.: The mycorrhizal symbiosis: research frontiers in
794 genomics, ecology, and agricultural application, *New Phytol.*, 242, 1486–1506,
795 <https://doi.org/10.1111/nph.19541>, 2024.

796 McCulloh, K. A., Sperry, J. S., and Adler, F. R.: Water transport in plants obeys Murray’s law,
797 *Nature*, 421, 939–942, <https://doi.org/10.1038/nature01444>, 2003.

798 Mensah, C., Šigut, L., Fischer, M., Foltýnová, L., Jocher, G., Acosta, M., Kowalska, N., Kokrda, L.,
799 Pavelka, M., Marshall, J. D., Nyantakyi, E. K., and Marek, M. V.: Assessing the Contrasting Effects
800 of the Exceptional 2015 Drought on the Carbon Dynamics in Two Norway Spruce Forest Ecosystems,
801 *Atmosphere*, 12, 988, <https://doi.org/10.3390/atmos12080988>, 2021.

802 Moreira, M. Z., Sternberg, L. da S. L., and Nepstad, D. C.: Vertical patterns of soil water uptake by
803 plants in a primary forest and an abandoned pasture in the eastern Amazon: an isotopic approach,
804 *Plant Soil*, 222, 95–107, <https://doi.org/10.1023/A:1004773217189>, 2000.

805 Oerter, E. J. and Bowen, G.: In situ monitoring of H and O stable isotopes in soil water reveals
806 ecohydrologic dynamics in managed soil systems, *Ecohydrology*, 10, e1841,
807 <https://doi.org/10.1002/eco.1841>, 2017.

808 Plamboeck, A. H., Grip, H., and Nygren, U.: A hydrological tracer study of water uptake depth in a
809 Scots pine forest under two different water regimes, *Oecologia*, 119, 452–460,
810 <https://doi.org/10.1007/s004420050807>, 1999.

811 da S. L. Sternberg, L., Moreira, M. Z., and Nepstad, D. C.: Uptake of water by lateral roots of small
812 trees in an Amazonian Tropical Forest, *Plant Soil*, 238, 151–158, 2002.

813 Seeger, S. and Weiler, M.: Temporal dynamics of tree xylem water isotopes: in situ monitoring and
814 modeling, *Biogeosciences*, 18, 4603–4627, <https://doi.org/10.5194/bg-18-4603-2021>, 2021.

815 Shigo, A. L.: Compartmentalization: A Conceptual Framework for Understanding How Trees Grow
816 and Defend Themselves, *Annu. Rev. Phytopathol.*, 22, 189–214,
817 <https://doi.org/10.1146/annurev.py.22.090184.001201>, 1984.

818 Simard, S. W. and Durall, D. M.: Mycorrhizal networks: a review of their extent, function, and
819 importance, *Can. J. Bot.*, 82, 1140–1165, <https://doi.org/10.1139/b04-116>, 2004.

820 Sternberg, L. da S. L., Bucci, S., Franco, A., Goldstein, G., Hoffman, W. A., Meinzer, F. C., Moreira,
821 M. Z., and Scholz, F.: Long range lateral root activity by neo-tropical savanna trees, *Plant Soil*, 270,
822 169–178, <https://doi.org/10.1007/s11104-004-1334-9>, 2005.

823 Stocker, B. D., Tumber-Dávila, S. J., Konings, A. G., Anderson, M. C., Hain, C., and Jackson, R. B.:
824 Global patterns of water storage in the rooting zones of vegetation, *Nat. Geosci.*, 16, 250–256,
825 <https://doi.org/10.1038/s41561-023-01125-2>, 2023.

826 Tarvainen, L., Wallin, G., Lim, H., Linder, S., Oren, R., Ottosson Löfvenius, M., Råntfors, M., Tor-
827 ngern, P., and Marshall, J.: Photosynthetic refixation varies along the stem and reduces CO₂ efflux in
828 mature boreal *Pinus sylvestris* trees, *Tree Physiol.*, 38, 558–569,
829 <https://doi.org/10.1093/treephys/tpx130>, 2018.

830 Vernay, A., Tian, X., Chi, J., Linder, S., Mäkelä, A., Oren, R., Peichl, M., Stangl, Z. R., Tor-Ngern,
831 P., and Marshall, J. D.: Estimating canopy gross primary production by combining phloem stable
832 isotopes with canopy and mesophyll conductances, *Plant Cell Environ.*, 43, 2124–2142,
833 <https://doi.org/10.1111/pce.13835>, 2020.

834 Volkmann, T. H. M., Kühnhammer, K., Herbstritt, B., Gessler, A., and Weiler, M.: A method for in
835 situ monitoring of the isotope composition of tree xylem water using laser spectroscopy, *Plant Cell*
836 *Environ.*, 39, 2055–2063, <https://doi.org/10.1111/pce.12725>, 2016.

837 Werner, C., Meredith, L. K., Ladd, S. N., Ingrisch, J., Kübert, A., van Haren, J., Bahn, M., Bailey, K.,
838 Bamberger, I., Beyer, M., Blomdahl, D., Byron, J., Daber, E., Deleeuw, J., Dippold, M. A., Fudyma,
839 J., Gil-Loaiza, J., Honeker, L. K., Hu, J., Huang, J., Klüpfel, T., Krechmer, J., Kreuzwieser, J.,
840 Kühnhammer, K., Lehmann, M. M., Meeran, K., Misztal, P. K., Ng, W.-R., Pfannerstill, E., Pugliese,
841 G., Purser, G., Roscioli, J., Shi, L., Tfaily, M., and Williams, J.: Ecosystem fluxes during drought and
842 recovery in an experimental forest, *Science*, 374, 1514–1518, <https://doi.org/10.1126/science.abj6789>,
843 2021.

844 White, J. W. C., Cook, E. R., Lawrence, J. R., and Wallace S., B.: The DH ratios of sap in trees:
845 Implications for water sources and tree ring DH ratios, *Geochim. Cosmochim. Acta*, 49, 237–246,
846 [https://doi.org/10.1016/0016-7037\(85\)90207-8](https://doi.org/10.1016/0016-7037(85)90207-8), 1985.

847 Zhao, P., Chi, J., Nilsson, M. B., Löfvenius, M. O., Högberg, P., Jocher, G., Lim, H., Mäkelä, A.,
848 Marshall, J., Ratcliffe, J., Tian, X., Näsholm, T., Lundmark, T., Linder, S., and Peichl, M.: Long-term
849 nitrogen addition raises the annual carbon sink of a boreal forest to a new steady-state, *Agric. For.*
850 *Meteorol.*, 324, 109112, <https://doi.org/10.1016/j.agrformet.2022.109112>, 2022.

851