

Cosmogenic ${}^3\text{He}$ exposure dating in mafic rocks by ‘Virtual mineral separation’ of pyroxene

Marie Bergelin^{1,2}, Greg Balco^{3,1}, and Richard A. Ketcham⁴

¹Berkeley Geochronology Center, 2455 Ridge Road, Berkeley, CA 94709, USA

²Department of Earth and Planetary Science, University of California, Berkeley, 307 McCone Hall, Berkeley, CA 94720, USA.

³Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, CA 94550, USA.

⁴Jackson School of Geosciences, The University of Texas at Austin, 23 San Jacinto Blvd, Austin, Texas, 78712, USA

Correspondence to: Marie Bergelin (mbergelin@bgc.org)

Abstract. We describe a ‘virtual mineral separation’ method for measuring the cosmogenic ${}^3\text{He}$ concentration in pyroxene in mafic rocks that consist mainly of plagioclase and pyroxene, without physically separating the minerals. This approach is significantly faster and more cost-effective than the conventional method, which requires physical separation and purification of pyroxene grains by time-consuming and labor-intensive crushing, acid cleaning, magnetic separation, HF etching, and handpicking under a microscope. The premise of the method is that helium diffusivity is much higher in plagioclase than in pyroxene, so controlled preheating of a mixed whole-rock sample can degas ${}^3\text{He}$ from plagioclase while retaining all ${}^3\text{He}$ in pyroxene. A second heating step releases all ${}^3\text{He}$ from pyroxene for measurement. To then obtain a ${}^3\text{He}$ concentration in pyroxene rather than the whole rock, we determine the pyroxene weight fraction in the sample using X-ray computed tomography (CT). A comparison of ${}^3\text{He}$ concentrations in pyroxene measured using virtual mineral separation with those measured in the same samples by physical mineral separation in previous work shows no evidence of systematic bias between the methods. Virtual mineral separation greatly simplifies the workflow for ${}^3\text{He}$ exposure-dating of mafic rocks, reduces time, effort, and cost, and permits measurements on very small samples. This enables new emerging applications of exposure dating, such as quantifying stochastic surface processes, ecosystem studies, and potential subglacial bedrock exposure dating.

Deleted: effectively

1 Introduction

In this paper we develop a method for measuring the cosmogenic ${}^3\text{He}$ concentration in pyroxene in mafic rocks without physically separating the pyroxene. The stable cosmogenic nuclide ${}^3\text{He}$ is commonly used in mafic rocks for exposure dating, as it is retained in pyroxene and easily measured using a noble gas mass spectrometer (e.g. Kurz, 1986; Cerling, 1990; Balter-Kennedy et al., 2020). This makes cosmogenic ${}^3\text{He}$ exposure dating more economical compared to other costlier nuclides relying on accelerator mass spectrometry (such as cosmogenic ${}^{10}\text{Be}$, ${}^{26}\text{Al}$, and ${}^{36}\text{Cl}$). However, the sample preparation to obtain pure pyroxene separates is time-consuming and tedious, and often results in under-constrained inclusion of impurities (Bromley et al., 2014).

Deleted: entive

At present, the procedure for measuring cosmogenic ${}^3\text{He}$ in pyroxene has two steps: separation and purification of pyroxene
35 separates, followed by heating of the pyroxene separates under vacuum and analysis of the released helium in a noble gas mass spectrometer. The second step of this process (vacuum degassing and mass spectrometry) is rapid, efficient, and fully automated at many laboratories. For example, the system in use at Berkeley Geochronology Center (BGC) has a typical continuous throughput of 10-12 samples/day, with approximately 1 hour/day of operator time needed for sample changes, for a two-heating-step ${}^3\text{He}$ analysis with associated standards and blanks.

40 In contrast, the first step of pyroxene separation is time-consuming and labor-intensive, and requires crushing and sieving of a rock sample, acid cleaning of the crushed rock, heavy liquid and/or magnetic separation to concentrate pyroxene, and HF etching to remove adhering plagioclase. Pyroxene purification is typically finalized by additional magnetic separation and hand-picking (e.g. Kurz, 1986; Brook et al., 1995; Bruno et al., 1997; Licciardi et al., 1999; Schäfer et al., 1999; Blard et al.,
45 2009; Bromley et al., 2014), making the resulting purity subjective and difficult to quantify. This is critical as the presence of non-helium-retentive impurities such as plagioclase (Cerling, 1990) can lead to an overestimation of sample mass and, consequently, an underestimation of measured ${}^3\text{He}$ concentration. The process becomes inherently more tedious in fine-grained rocks where monocrystalline grains are difficult to obtain. None of these pyroxene separation steps are automated, several require hazardous chemicals, and some require a trained technician. Therefore, the primary limitation in collecting large data
50 sets of ${}^3\text{He}$ concentrations in pyroxene is the process of mineral separation.

Plagioclase, commonly found with pyroxene as the major constituents of many mafic rocks, was first determined by Cerling (1990) to be non-retentive of ${}^3\text{He}$, noting that this potentially causes problems for ${}^3\text{He}$ exposure dating in whole rock. However, in a mafic rock that predominantly consists of pyroxene and plagioclase, the contrast in ${}^3\text{He}$ retentivity could be advantageous.

55 In recent work, Bergelin et al. (2025) quantified the observation that ${}^3\text{He}$ is poorly retained in plagioclase by measuring diffusion kinetics of helium in both plagioclase and pyroxene. This result can be used to develop a method for degassing ${}^3\text{He}$ from plagioclase in a single heating step, without degassing the ${}^3\text{He}$ retained in the pyroxene, prior to cosmogenic ${}^3\text{He}$ analysis, leading to what we refer to as a ‘virtual mineral separation’. The required temperature and duration for such a separation can be determined using the measured diffusion kinetics (Bergelin et al., 2025) of both minerals and forward calculation of the
60 fraction release (Fechtig and Kalbitzer, 1966).

A virtual mineral separation method would eliminate the initial labor-intensive step of physically isolating pyroxene from whole-rock samples. Instead, ${}^3\text{He}$ is measured directly in pyroxene within a whole rock sample following a preheating which degasses ${}^3\text{He}$ from the less retentive plagioclase. However, doing so results in a ${}^3\text{He}$ concentration which has units of [atoms
65 ${}^3\text{He}$ in pyroxene] / [g rock]. Therefore, to use the virtual mineral separation method for an exposure dating application, it is necessary to convert this measurement to a ${}^3\text{He}$ concentration in pyroxene ([atoms ${}^3\text{He}$ in pyroxene] / [g pyroxene]). This

Deleted: ,

Deleted: is

Deleted: a

Deleted: study

Deleted: , we found the diffusivity between ${}^3\text{He}$ in plagioclase and pyroxene to be significantly different. This difference opens the possibility to degas ${}^3\text{He}$ from plagioclase in a single heating

requires independent determination of the pyroxene weight fraction ($[\text{g pyroxene}] / [\text{g rock}]$). To preserve the efficiency of the
75 virtual mineral separation, this procedure must also be rapid and efficient. X-ray computed tomography (CT) is advantageous
for this purpose as it can identify mineral grains and quantify shape and volume separately or within aggregates (Cooperdock
et al., 2022; e.g. Hofmann et al., 2021). A key advantage is that CT scanning can be applied directly to the same 100-mg-scale
aliquot degassed for ${}^3\text{He}$ measurements, and many aliquots can be scanned simultaneously, making it a cost-effective and high-
throughput step. Although CT scanning has not been used for this specific purpose before, the difference in density between
80 pyroxene ($3.2 - 3.5 \text{ g cm}^{-3}$) and plagioclase ($2.6-2.8 \text{ g cm}^{-3}$) indicates that they should be easily differentiable.

Deleted: , and we establish this below.

This project has multiple components to it, which we have divided into three separate sections that describe methods and
process validation results for each component. First, we use diffusion kinetics of He in pyroxene and plagioclase recently
85 measured by Bergelin et al. (2025) to design a heating schedule that separates ${}^3\text{He}$ in plagioclase from that in pyroxene and
perform experiments to show that it is effective. Second, we determine the weight fraction of pyroxene in the whole-rock
sample through CT scanning. And third, we apply the virtual mineral separation method to a large set of exposure-dating
samples and compare the results to existing ${}^3\text{He}$ measurements in pyroxene separates (Balter-Kennedy et al., 2020).

Deleted: previously measured

Deleted: pyroxene and

Deleted: its success against

The main objective of this study is to make measurements of ${}^3\text{He}$ in pyroxene from mafic rock faster and more efficient by a
90 virtual mineral separation method. For this purpose, we focus on the Ferrar Dolerite, an intrusive gabbroic rock found
throughout the Transantarctic Mountains (TAM) in Antarctica that predominantly consists of subequal parts of plagioclase
and pyroxene, with minor contributions from other mineral phases (Elliot and Fleming, 2021). Its commonly fine-grained
texture makes physical pyroxene separation challenging, making it an ideal candidate for the virtual mineral separation
approach. Due to its resistance to weathering, the Ferrar dolerite is one of the most common surface lithologies in Antarctica,
95 and therefore commonly used for exposure-dating studies. The success of this study could enable applications in quantifying
stochastic surface processes, ecosystem studies, and potential subglacial bedrock exposure dating. Further, it could be relevant
to various other lithologies in Antarctica and elsewhere and potentially transformative for science applications that require
large data sets of exposure ages. It would not be as transformative as in-situ exposure dating in the field (Farley et al., 2014),
but it would be a step in the same direction.

100 2 Method 1: Experimental design

2.1 Diffusion kinetics of pyroxene and plagioclase

The diffusion kinetics of noble gases in a mineral can be determined through stepwise degassing of a single grain irradiated
with neutrons or protons to produce an initial homogeneous distribution of gas. In a recent study, Bergelin et al. (2025) found
the diffusion kinetics of ${}^3\text{He}$ in irradiated plagioclase and pyroxene grains from the Ferrar Dolerite to be vastly different, as
105 expected from previous observations of missing ${}^3\text{He}$ in plagioclase (Eberhardt et al., 1966; Cerling, 1990). Here, we perform

Deleted:


Deleted: t

Deleted: , we

Deleted: W

an additional experiment to show how this difference in diffusion kinetics between plagioclase and pyroxene causes the
 115 minerals to degas ${}^3\text{He}$ separately. We conducted prograde step-heating experiments on (i) individual proton-irradiated
 plagioclase (LABCO-Plag) and pyroxene (LABCO-Px) grains obtained from the same sample set analysed in Bergelin et al.
 (2025) and (ii) a crushed whole-rock sample of Ferrar Dolerite (15-OTW-049-CHA, described by Bromley et al., 2024) that
contains a mixture of pyroxene and plagioclase grains, and has an apparent ${}^3\text{He}$ exposure age of 8.4 Ma, making the naturally
produced ${}^3\text{He}$ concentration adequate for a step-degassing experiment. The whole-rock sample was crushed and sieved to a
 120 grain size of 100-160 μm , then washed in water and leached in 10% HCl overnight at room temperature. Once dried, 7.06 mg
 of bulk sample grains were packed into a small (\sim 5mm) Pt-Ir alloy package. For all three experiments, we used a consistent
heating schedule and the exact experimental setup and measurement procedure described in Bergelin et al. (2025; see Table
S1).

125 In Fig. 1, we show the fractional release of ${}^3\text{He}$ during the three step-heating experiments, emphasizing the distinctive
 difference in gas release between plagioclase and pyroxene. At 260°C (step 29), the plagioclase sample had released 97% of
 its ${}^3\text{He}$, compared to only 2% from the pyroxene sample and 3% from the bulk whole-rock sample. Interestingly, the naturally
 130 irradiated bulk whole-rock grains (15-OTW-049-CHA) display a ${}^3\text{He}$ release pattern similar to that of pyroxene, despite
 consisting of subequal parts of plagioclase and pyroxene. Given that ${}^3\text{He}$ is incompletely retained in plagioclase at Earth surface
 temperatures, and the extended exposure time of this particular sample, most of the ${}^3\text{He}$ produced in plagioclase appears to
 have been lost during surface exposure (Bergelin et al., 2025).




135 **Figure 1** Step heating analysis of proton-irradiated, single grains of Ferrar dolerite plagioclase (blue), pyroxene (red), and naturally
 cosmic-ray irradiated, crushed Ferrar dolerite containing a mixture of plagioclase and pyroxene grains (black). Each of the three
 degassing experiments experienced the same heating schedule. (a) shows the fractional release of ${}^3\text{He}$ atoms, and (b) shows the
 temperature reached during each heating step. Note that the duration of each step varies, which is not shown in this figure (Table
 S1).

Deleted: identical

Deleted: following the same measurement procedure (see Table S1 for more details). Additionally, we applied the same heating schedule and measurement procedure to a crushed sample of Ferrar Dolerite (15-OTW-049-CHA, described by Bromley et al., 2014) that has not been separated, and therefore contains a mixture of pyroxene and plagioclase grains. This sample has an apparent ${}^3\text{He}$ exposure age of 8.39 ± 0.24 Ma, making the naturally produced ${}^3\text{He}$ concentration adequate for an extended step-degassing experiment.

Deleted: and analysed using the same procedure as the single mineral grains for step-heating degassing.




Deleted:

Deleted: red

Deleted: blue

We use the diffusion kinetics for plagioclase and pyroxene, as determined by Bergelin et al. (2025), to model the predicted fractional release of ^3He from these minerals during a single heating step (Fig. 2). The fractional gas release is calculated using a Multiple Diffusion Domain (MDD) model (Lovera et al., 1989; Lovera et al., 1997), following the fractional loss equations

155 of Fechtig and Kalbitzer (1966, equation 4a-c). While there is a large difference between ^3He diffusion kinetics for plagioclase and pyroxene, there are also smaller variations in diffusivity within each mineral group, due to compositional differences (Tremblay et al., 2017; Bergelin et al., 2025). Given that the single-step heating schedule is intended for bulk samples, where individual mineral grains may exhibit compositional heterogeneity, we evaluate the effective temperature and time ranges by considering both the highest and lowest diffusivities determined from each mineral group. The fractional release of ^3He from 160 plagioclase and pyroxene during a single-step heating schedule is then estimated based on the most diffusive pyroxene and most retentive plagioclase kinetics, which represents a least-effective-case scenario for designing the separation procedure. For plagioclase, this estimation uses the diffusion kinetics determined from grains of Ferrar Dolerite (ROB-plag; see discussion in Bergelin et al., 2025). However, for pyroxene, this includes diffusion kinetics obtained from grains of Ferrar dolerite pyroxene (LABCO-Px and ROB-Px) and gem-quality pyroxene grains (Bergelin et al., 2025).



165

Figure 2 Area of temperature and time combination between the most and least retentive ^3He mineral phases within pyroxene (red) and plagioclase (blue) for a fractional gas loss of 0.01 and 0.98, respectively. For plagioclase, the temperature-time range is based on the diffusion kinetics determined from grains of the Ferrar Dolerite, ROB-plag (Bergelin et al., 2025). For pyroxene, the range includes diffusion kinetics obtained for grains of Ferrar dolerite pyroxene (LABCO-Px and ROB-Px) and gem-quality pyroxene (GEM-Px) grains (Bergelin et al., 2025). The grain size radius used to determine the diffusion kinetics range from 108-168 μm . Any temperature-time combinations above the blue shade are expected to degas >0.98 , and any combinations below the red shade are expected to degas < 0.01 of the total ^3He in each mineral phase. The area below the red shade and above the blue shade represents the temperature and time field to separate ^3He released from plagioclase, without releasing ^3He from pyroxene. The black dot shows the temperature and time combination that we use in this experiment, and is suitable for a standard vacuum oven.

170

175 Figure 2 shows that a single heating step at 220 °C for a duration of 8 hr is theoretically sufficient to degas more than 98% of ^3He from plagioclase while retaining over 99% of ^3He in pyroxene, even for the least-effective case represented by the least retentive pyroxene and most retentive plagioclase. This results in an effective separation of ^3He between the two minerals. Although comparable separation efficiency can be achieved by, e.g., increasing the temperature and reducing the duration, the maximum temperature is constrained by the operational limit of standard vacuum ovens (< 250 °C).

Deleted: ^3He diffusion kinetics differ significantly between

Formatted: Superscript

Deleted: for

Deleted: exist

Deleted: providing the

Deleted:

Deleted:

Deleted: between ^3He released from plagioclase and that retained in pyroxene. ...

Deleted: ,

Deleted: h

Deleted: of

Deleted: . The grain size radius used to determine the diffusion kinetics range from 108-168 μm

Deleted: expected

Deleted: to separate ^3He released from plagioclase and pyroxene...

Deleted: .

2.2 Validation of established single-step heating schedule

To assess whether a single heating step effectively separates ${}^3\text{He}$ released from plagioclase and pyroxene, we compared ${}^3\text{He}$ concentration from multiple preheated and un-preheated individual grains of each mineral from the Ferrar Dolerite (ROB, Bergelin et al., 2025). These grains were proton-irradiated, resulting in an elevated ${}^3\text{He}$ concentration detectable within single grains, and immediately frozen post-irradiation until analysis to prevent gas loss ([the purpose of this procedure is described in Bergelin et al., 2025](#)). The ${}^3\text{He}$ concentrations from equivalent un-preheated grains of proton-irradiated plagioclase and pyroxene were obtained from degassing experiments already described in Bergelin et al. (2025). Here, we subjected similar grains from the same sample set to a single-preheating step prior to analysis for comparison, as described below.

205

The set of four preheated samples consists of two ROB-plagioclase grains and two ROB-pyroxene grains. Each grain was weighed before analysis, packed into a small Ta package, and placed in an AccuTemp 0.9 cft (AT09) vacuum oven. The oven was then put under vacuum at 2×10^{-1} torr to prevent oxidation of the sample and package or diffusion of atmospheric gases into the sample. In addition to the oven's built-in temperature monitoring, two type-K thermocouples were installed inside: 210 One was affixed to the oven shelf surface, while the other was placed inside a blank Ta package positioned adjacent to the samples. Both thermocouples were wired through the oven door and sealed with self-fusing silicone tape. The four samples were heated at 220°C for 8 hrs as established above.

215 The total ${}^3\text{He}$ concentration in all four preheated sample grains was then measured using the BGC 'Ohio' noble gas mass spectrometer following the procedure described in Balter-Kennedy et al. (2020). The total ${}^3\text{He}$ concentration in the un-preheated plagioclase and pyroxene grains was calculated from the total amount of ${}^3\text{He}$ measured during the step-heating diffusion experiment conducted by Bergelin et al. (2025; supplementary data). As the irradiation-induced ${}^3\text{He}$ production is likely to vary among the grains because of spatial variability in the proton flux, we do not expect identical ${}^3\text{He}$ concentrations in all sample grains. However, this assessment is meant to provide preliminary evidence of the effectiveness of the single-step 220 heating schedule.

Table 1 Measured ${}^3\text{He}$ in proton-irradiated plagioclase and pyroxene grains from Ferrar Dolerite sample ROB (Bergelin et al., 2025)

Sample name	Oven	Mass of grain (mg)	Total Measured ${}^3\text{He}$ (Atoms)	Calculated ${}^3\text{He}$ conc. (Atoms/g)	Source
<i>Plagioclase</i>					
ROB-Plag-Da	-	0.112	2.76×10^8	2.46×10^{12}	Bergelin et al. (2025)
ROB-Plag-Db	-	0.101	1.64×10^8	1.62×10^{12}	Bergelin et al. (2025)
ROB-Plag-A	Yes	0.057	7.97×10^4	1.40×10^9	This paper

ROB-Plag-B	Yes	0.135	5.43×10^6	4.02×10^{10}	This paper
Pyroxene					
ROB-Px-Da	-	0.119	5.56×10^8	4.68×10^{12}	Bergelin et al. (2025)
ROB-Px-A	Yes	0.153	8.88×10^8	5.80×10^{12}	This paper
ROB-Px-B	Yes	0.167	1.39×10^9	8.30×10^{12}	This paper

Table 1 shows the amount of ^3He measured in all seven sample grains. Preheating plagioclase grains prior to analysis decreases the ^3He concentration by at least two orders of magnitude, indicating a gas loss of more than 98%. In contrast, we find that pyroxene grains that were subjected to the same preheating had equivalent to slightly higher concentrations than the grains that were not preheated. These results are consistent with predictions in Fig. 2, confirming that a single 8-hour heating at 220°C effectively degasses ^3He from plagioclase while retaining it in pyroxene. When applied to bulk whole-rock material, this established single-step heating schedule provides the basis for the virtual mineral separation of ^3He in pyroxene from mafic rocks without physically having to separate and isolate pyroxene grains.

Deleted:

3 Method 2: Pyroxene fraction estimation

3.1 Sample description and preparation

We apply the virtual mineral separation method to a total of 36 samples that have known concentrations of ^3He in pyroxene from previous measurements in pyroxene separates (Balster-Kennedy et al., 2020; Bromley et al., 2024). All samples belong to the Ferrar dolerite, a mafic intrusive rock consisting primarily of calcic plagioclase and several orthopyroxenes and clinopyroxenes (Elliot and Fleming, 2021). These samples are from high-elevation surface boulders collected from various moraines near Roberts (ROB) and Otway (OTW) Massif in the central Transantarctic Mountains (TAM), and have apparent ^3He -in-pyroxene exposure ages ranging from 1 to 11 Ma (see Balster-Kennedy et al., 2020, and Bromley et al., 2024, for details).

Deleted: ^3He

240 Sample preparation for previously published ^3He measurements on pyroxene separates from these rocks (Balster-Kennedy et al., 2020; Bromley et al., 2024) involved the following steps: Crushed rock samples were sieved to a 125-250 μm grain size fraction, and boiled for 2 hr in 10% HNO_3 . Pyroxene grains were then separated from plagioclase using a water-based heavy liquid, followed by etching in 5% HF to dissolve any adhering plagioclase and remove the outer surface of the pyroxene grain. 245 The etched pyroxene grains were passed through a magnetic separator and handpicked under a binocular microscope to remove any remaining contaminants. Finally, pyroxene grains were packed into Ta packages for cosmogenic ^3He analysis.

In contrast, the virtual mineral separation method does not require physical separation and purification of the pyroxene grains. Instead, we hand-crushed and sieved a small rock sample to a 90-150 μm grain size fraction. [This smaller grain size fraction](#)


maximizes monocrystalline grains and is advantageous for CT scanning, as each grain is more likely to consist of only plagioclase or pyroxene. The grains were then rinsed and sonicated in water for 1 hr to remove any dust particles from crushing. After drying, 40-80 mg of the bulk whole-rock grains were directly packed into 1 mL vials for X-ray CT scanning. The sample 255 mass was selected based on prior measurements of ${}^4\text{He}$ in pyroxene separates to yield a total ${}^4\text{He}$ abundance within the pressure range of the gas standards used to calibrate the mass spectrometer (Balter-Kennedy et al., 2020; Bromley et al., 2024).

In addition to these 36 samples, we included several supplementary samples to evaluate the efficacy and limitations of the virtual mineral separation method. (i) To assess reproducibility, we prepared multiple aliquots from two whole-rock samples: 260 15-OTW-21-DUT and 15-OTW-47-CHA. (ii) To test whether crushing is necessary, we packaged aliquots of a whole rock chip (<100 mg) directly into a 1 mL vial from sample 16-ROB-052-BAS. This represents minimal sample processing and eliminates crushing. (iii) To validate the CT-based image processing workflow and the weight percent calculation, we included 265 a set of control samples derived from 15-ROB-56-NAC and 16-ROB-72-MNM, having known proportions of plagioclase and pyroxene grains. These control samples were prepared by crushing whole-rock material, followed by sieving, sonication in water, and leaching in 10% HCl. Mineral separation was then achieved by repeated passes through a magnetic separator. Pyroxene separates were subsequently etched in 5% HF and 1% HNO_3 to remove any adhering plagioclase. Final purification involved handpicking under a binocular microscope to remove any remaining impurities. Each control sample consisted of 100 mg of precisely weighed mineral separates, containing a known amount between 30-45% plagioclase.

Deleted: and

3.2 CT scan

270 The sample vials were CT scanned in batches of 7-8 samples at the University of Texas High-Resolution X-ray Computed Tomography Facility (UTCT) in Austin, TX, using the Zeiss Xradia 620 Versa system. Scans were performed with the X-ray source set to 80 kV and 10W and utilizing a low-energy beam filter, collecting 3201 projections through a 360° rotation with an acquisition time of 0.3s per projection and stage motion between projections (dithering) to reduce ring artifacts, with a total 275 scan time of 56 minutes per batch. A beam-hardening correction was applied during reconstruction with Zeiss software, which provided stacks of up to 1200 16-bit TIFF images with a voxel resolution of 7.99 μm . Each voxel value reflects the X-ray attenuation, which is a function of density, atomic number, and the X-ray energy spectrum, and is represented in a greyscale ranging from 0 to 65,535. From each image stack, each sample was individually cropped and processed (Fig. 3).



280 Figure 3 Computed tomography (CT) image and voxel classification of whole rock grain sample of Ferrar dolerite, 15-ROB-028-
 COL (Z). CT images (a) and (b) show the vertical and horizontal scans of the sample packed in a 1 mL vial. These grey-scale CT
images show the three mineral phases: oxides (white), pyroxene (light grey), and plagioclase (dark grey), surrounded by air (black).
 The histogram from a representative subvolume (c) shows the distribution of CT number and the classification of each mineral
 phase determined and highlighted in red for (d) oxides, (e) pyroxene, and (f) plagioclase. The workflow outlined represents the
 workflow applied to all samples.

285

Deleted: at

For each sample, we observe a frequency distribution of CT numbers with multiple characteristic peaks of distinct X-ray attenuation (Fig. 3c), that we interpret as air, plagioclase (plag), and pyroxene (px), with a tail of high-density minerals (ox; interpreted as oxides). Although distinct CT number variations were observed within individual mineral groups (suggesting potential sub-classification), we did not attempt to differentiate between mineral phases beyond the three primary groups, as this was mostly manifested by a wider peak in the histogram (e.g., pyroxene phase in Fig. 3c), rather than two separate peaks.

Deleted: fig

We created a workflow using the [freely distributed](#) software Blob3D, which was written for CT data processing (Ketcham, 2005; Ketcham and Mote, 2019). The segmentation of the CT data to differentiate pyroxene, plagioclase, and other phases is not straightforward, and cannot be done accurately utilizing simple histogram analysis. Although the voxel size is $\sim 8 \mu\text{m}$, the boundary between any two phases, or a phase and air, spans multiple voxels due to the finite resolution of the data; for the data in this study, the point-spread function (PSF) width was measured to be ~ 3.5 voxels, meaning that it takes 3.5 voxels to span 95% of the transition between adjacent phases (Ketcham et al., 2010). Thus, for example, a 120- μm grain of pyroxene has a 3.5-voxel (28 μm) transition to adjacent air, spanning the outer 23% of the grain radially and 40% of its volume, as well as a region beyond the grain that corresponds to 50% of its volume. Averaging between a high-attenuation phase and air in this region produces CT numbers overlapping those of lower-attenuation phases; for example, assigning one range of CT numbers to pyroxene and a lower range to plagioclase would produce volumetrically significant shells or atolls of voxels misclassified as plagioclase surrounding pyroxene grains.

We thus utilize a segmentation strategy based on selecting grains based on the CT numbers of their centers, and then expanding the selections to capture the inner part of their blurred region, or a threshold and expand operation (Ketcham, 2005). Working from highest-attenuation to lowest-attenuation phase, a CT number range is defined that encompasses the end-member CT numbers found in the grain cores, which is broadened to include lower CT numbers that remain unique to that phase to also capture the initial part of the transition region. To capture the transition region to its midpoint, the selections are then expanded by 1 voxel to a limit defined by the average core value and air. After each phase is captured, its voxels are removed from further consideration, and an additional region beyond the grain boundary is also removed to eliminate atolls. The removed voxels are given a temporary CT number that will not be captured during the initial threshold step during segmentation of the next-lower-attenuation phase, but can be captured in the expansion step. A final remove islands operation is used to eliminate remaining isolated voxels and stringers. Each segmentation was verified visually, and the parameters of each step were adjusted as necessary to optimize grain differentiation.

For the two whole rock chips (16-ROB-052-BASb,c), boundaries were assumed to be rock rather than air, therefore, CT number thresholds were defined midway between the mineral of interest and that of the lower-attenuation mineral class value. When identifying lower-attenuation phases, voxels assigned to higher-attenuation mineral classes were excluded, and their

two-voxel radius values were replaced with transitional voxel values, ensuring these were treated as boundary voxels rather than misclassified mineral phases (see details [of segmentation](#) in Fig. 3d-f).

325 The range in CT numbers, which is dependent on the scan parameters and energies, was consistent within each batch. To evaluate whether threshold values from one batch could be applied across others, sample 15-ROB-056-[NACb \(VV\)](#) was
330 scanned in three different batches for intercomparison of CT numbers. By applying the same established image processing workflow, we achieve a standard deviation of 0.27 wt.% for both pyroxene and plagioclase and < 0.01 wt.% for oxide. This reproducibility supports the use of a single threshold for each mineral classification across all samples, with minor modifications as needed (See details for each sample in Table S2).


Deleted: BAS

We also did a higher-resolution scan (4 μm voxel size, only one vial in field of view, duration 36 minutes) of a single test specimen to estimate resolution effects. While the resulting data were clearer, similar processing achieved results within 0.7 wt.% of the 8 μm scan data.

335 3.3 Pyroxene weight percent estimation

The CT image processing yields voxel percentages for each mineral phase, which are converted to weight percent using mineral-specific densities. For the plagioclase group, we apply a density of 2.73 g cm^{-3} , which is that of anorthite (An85) and characteristic of the Ferrar dolerite plagioclase. For the oxide phases, we use the density of ilmenite (4.72 g cm^{-3}), a common accessory mineral in Ferrar Dolerite (Elliot and Fleming, 2021). Although pyroxene composition varies between clinopyroxene
340 and orthopyroxene within individual samples, previous work (Bergelin et al., 2024) shows minimal variability in bulk pyroxene chemistry among Ferrar Dolerite samples collected from similar localities. [We show this with the mineral composition of pyroxene separates determined from XRF analysis](#) from samples collected from Roberts and Otway Massif (Bergelin et al., 2024, Table 4) in Fig. 4. For the pyroxene phases, we therefore apply the average density for these samples of 3.45 g cm^{-3} . Even if endmember densities were used, the resulting variation would be less than 5%, which falls within the typical uncertainty
345 range for cosmogenic nuclide dating.

Deleted: To
Deleted: , we
Deleted: present
Deleted:
Deleted: data
Deleted: from



355 **Figure 4** Ternary diagram of wollastonite-ferrosilite-enstatite showing the pyroxene composition from 10 Ferrar dolerite samples obtained from Roberts and Otway Massif in Antarctica by Balter-Kennedy et al. (2020) and analyzed for XRF by Bergelin et al. (2024). The XRF analysis was done on bulk samples of pyroxene separates and, therefore, most likely includes multiple pyroxene phases.

Deleted: data
Deleted: are from
Deleted: analysis

3.4 Validation of weight percent estimation

360 The initial workflow method for the image processing was developed using the control samples 15-ROB-056-NACa and b, yielding <1.1 wt.% deviations between known and estimated values (Table 2). Thresholds were then further adjusted to fit the samples having unknown mass percent. To validate the adjusted workflow, we applied it to two known ratio samples from 15-ROB-072-MNM (a and b) and repeated scans of 15-ROB-056-NACb, achieving < 1.0 wt.% variation in mass estimation.

Table 2 Estimated mineral phase fraction in control samples of mixed pyroxene and plagioclase of known mass ratio for establishing and updating the CT scan voxel classification workflow.

ID	sample name	sample mass (mg)	px mass (mg)	plag mass (mg)	known wt.% px	known wt.% plag	voxel % (oxid)	voxel % (px)	voxel % (plag)	combined sample density (g/cm3)	calc. wt.% oxid	calc. wt.% px	calc. wt.% plag	relative diff (px)	relative diff (plag)
<i>Initial workflow</i>															
UU	15-ROB-056-NACa-mix30	101.0	30.1	70.9	29.8	70.2	0.010	3.6	10.7	2.91	0.12	29.6	70.3	0.01	0.00
VV (1)	15-ROB-056-NACb-mix40	101.1	40.1	61.0	39.6	60.4	0.010	3.9	8.0	2.97	0.14	38.0	61.9	0.04	0.03
<i>Updated workflow</i>															
UU*	15-ROB-056-NACa-mix30	101.0	30.1	70.9	29.8	70.2	0.002	3.4	9.6	2.92	0.02	31.0	69.0	0.04	0.02
VV															
(1)*	15-ROB-056-NACb-mix40	101.1	40.1	61.0	39.6	60.4	0.002	3.7	7.1	2.98	0.03	39.9	60.0	0.01	0.01
VV (2)	15-ROB-056-NACb-mix40	101.1	40.1	61.0	39.6	60.4	0.002	3.9	7.6	2.98	0.03	39.4	60.6	0.01	0.00
VV (3)	15-ROB-056-NACb-mix40	101.1	40.1	61.0	39.6	60.4	0.002	5.0	9.6	2.98	0.03	39.7	60.2	0.00	0.00

XX	16-ROB-072- <u>MNM</u> a-mix35	100.4	35.1	65.4	34.9	65.1	0.003	4.5	11.1	2.94	0.03	34.1	65.9	0.02	0.01
YY	16-ROB-072- <u>MNM</u> b-mix45	106.7	45.3	61.5	42.4	57.6	0.009	6.5	11.7	2.99	0.08	41.1	58.8	0.03	0.02

* Sample re-evaluated using updated workflow

4 Method 3: Cosmogenic ${}^3\text{He}$ measurements

4.1 Cosmogenic ${}^3\text{He}$ analysis

370 We measured cosmogenic ${}^3\text{He}$ abundances in whole-rock samples of Ferrar dolerite. Following CT scanning, samples were carefully transferred from small 1 mL vials to Ta packets and weighed. On average, > 99.5 wt.% of the sample grains were successfully recovered. The samples were then placed centrally on a shelf within the vacuum oven, pumped down to 2×10^{-1} Torr, and heated at 220°C for 8 hr to degas ${}^3\text{He}$ from plagioclase (see sect. 2.2 for details). After cooling, the samples were loaded in batches of 12-14 into the sample chamber attached to the ‘Ohio’ noble gas mass spectrometer at BGC, and pumped 375 down to high vacuum ($<10^{-7}$ Torr) for analysis.

Deleted: s

Formatted: Superscript

Formatted: Heading 2

Helium extraction and analysis follow the procedure used for separated pyroxene grains, as described by Balter-Kennedy et al. (2020). Because the whole-rock samples were not HF-etched, ${}^4\text{He}$ concentrations are expected to be approximately 60% higher (Bromley et al., 2024). To keep total ${}^4\text{He}$ pressure within the range of available calibration standards, we used three 380 heating steps to completely degas helium, rather than the standard two-step procedure. We included an initial 3 min heating at 600°C, releasing roughly 50% of the retained helium, followed by a 12 min heating at 1200°C and a final heating of 15 min at 1300°C. The last heating step typically contains 1-5% of the total helium released and ensures complete extraction.

Deleted: at of

Deleted:

Deleted: ages

Deleted: a

Process blanks measured on empty Ta packets within each batch were $<10^5$ atoms of ${}^3\text{He}$ and $<10^{10}$ atoms of ${}^4\text{He}$, 385 corresponding to less than 0.8 % and 0.1 % of that measured in the samples, respectively. Replicate measurements of the CRONUS-P intercomparison standard (Blard et al., 2015) during analysis yielded an average ${}^3\text{He}$ concentration of $4.827 \pm 0.032 \times 10^9$ atoms g^{-1} . This is slightly lower than that measured during the previous study of ${}^3\text{He}$ in pyroxene separates ($5.03 \pm 0.15 \times 10^9$ atoms g^{-1} ; Balter-Kennedy et al., 2020). For comparison, all data sets are normalized to the accepted CRONUS-P value of 5.02×10^9 atoms g^{-1} (Blard et al., 2015).

390 4.2 Results

Deleted: 5

Deleted: 2

Deleted: 3

commonly differ by more than the nominal measured uncertainty (see discussion below), there is no systematic bias between the two sets of measurements.

405

Table 3 Measured ${}^3\text{He}$ concentration in whole-rock Ferrar dolerite samples. Columns show the sample ID used in CT scanning, sample name, sample mass, CT scanned voxel percentage determined from the workflow detailed in fig. 3, calculated weight percentage for each mineral phase based on mineral density, and the measured cosmogenic ${}^3\text{He}$ concentration in virtually-separated (this study) and pyroxene-separated (Balter-Kennedy et al., 2020; Bromley et al., 410 2024) samples. All ${}^3\text{He}$ measurements are corrected to CRONUS-P = 5.02×10^9 atoms g⁻¹.

ID	sample name	sample mass (g)	voxel % (oxide)	voxel % (px)	voxel % (plag)	combined sample density	wt.% oxide	wt.% px	wt.% plag	px mass (g)	virtual separated $[{}^3\text{He}]$ +/- (10^9 atoms g ⁻¹) ^a	physical separated $[{}^3\text{He}]$ +/- (10^9 atoms g ⁻¹) ^a
A	15-OTW-001-MOG	79.64	0.19	5.6	10.8	3.00	1.8	39.0	59.2	31.03	4.26 ± 0.17	4.13 ± 0.12
B	15-OTW-002-MOG	50.06	0.28	5.0	9.9	3.01	2.9	38.0	59.1	19.03	1.187 ± 0.046	1.118 ± 0.032
C	15-OTW-003-MOG	82.20	0.15	5.0	9.6	3.00	1.6	39.0	59.3	32.08	2.274 ± 0.090	2.405 ± 0.070
D	15-OTW-004-MOG	45.99	0.17	5.7	10.1	3.01	1.6	41.0	57.3	18.87	3.46 ± 0.13	3.31 ± 0.10
E	15-OTW-017-DUT	82.43	0.24	5.2	10.1	3.00	2.4	38.5	59.1	31.77	2.731 ± 0.040	3.135 ± 0.093
F	15-OTW-020-DUT	83.73	0.27	4.6	10.4	2.98	2.8	34.7	62.5	29.04	3.073 ± 0.047	3.48 ± 0.10
G	15-OTW-021-DUTa	82.63	0.36	5.6	10.3	3.02	3.4	39.4	57.2	32.53	3.725 ± 0.043	4.1 ± 0.12
H	15-OTW-021-DUTb	81.77	0.25	4.7	11.2	2.97	2.5	34.1	63.5	27.87	3.742 ± 0.043	4.1 ± 0.12
I	15-OTW-021-DUTc	80.74	0.36	5.4	11.7	2.99	3.3	35.4	61.3	28.59	3.825 ± 0.045	4.1 ± 0.12
J	15-OTW-022-DUT	81.58	0.33	5.3	11.9	2.99	3.0	35.0	62.0	28.53	1.746 ± 0.023	1.955 ± 0.063
K	15-OTW-025-JOS	82.73	0.24	6.2	11.0	3.01	2.1	40.7	57.2	33.68	1.996 ± 0.026	2.310 ± 0.067
L	15-OTW-026-JOS	84.52	0.27	5.2	11.1	2.99	2.6	36.4	61.1	30.74	3.577 ± 0.047	3.82 ± 0.12
M	15-OTW-027-JOS	83.77	0.18	6.1	9.1	3.04	1.8	44.9	53.3	37.62	4.022 ± 0.048	4.56 ± 0.15
N	15-OTW-032-JOS	82.96	0.18	7.0	10.1	3.04	1.6	45.9	52.4	38.11	2.125 ± 0.028	2.212 ± 0.070
O	15-OTW-039-MON	55.78	0.12	5.3	11.8	2.97	1.1	36.0	62.9	20.08	6.043 ± 0.087	6.9 ± 0.2
P	15-OTW-044-CHA	70.24	0.28	4.9	9.9	3.00	2.9	37.5	59.6	26.32	10.52 ± 0.14	10.23 ± 0.30
Q	15-OTW-046-CHA	82.13	0.23	6.4	11.6	3.01	2.0	40.4	57.6	33.22	8.34 ± 0.12	9.03 ± 0.26
R	15-OTW-047-CHAa	70.20	0.23	4.8	11.7	2.97	2.2	33.6	64.2	23.57	7.63 ± 0.10	7.77 ± 0.23
S	15-OTW-047-CHAb	71.08	0.28	4.8	11.6	2.97	2.7	33.6	63.7	23.88	7.51 ± 0.10	7.77 ± 0.23
T	15-OTW-047-CHAc	71.66	0.25	4.7	11.8	2.96	2.4	32.7	64.9	23.46	7.67 ± 0.10	7.77 ± 0.23
U	15-OTW-049-CHA	59.90	0.36	6.9	10.8	3.04	3.1	43.1	53.8	25.80	9.38 ± 0.12	9.52 ± 0.28
V	15-OTW-051-CHA	83.76	0.39	5.5	11.7	3.00	3.5	35.9	60.6	30.10	9.70 ± 0.12	9.25 ± 0.27
X	15-OTW-053-OLD	81.21	0.29	6.9	10.1	3.05	2.6	45.2	52.2	36.74	11.15 ± 0.13	11.20 ± 0.32
Y	15-ROB-009-MZB	43.64	0.59	4.7	13.5	2.98	5.0	29.2	65.8	12.74	9.38 ± 0.12	9.09 ± 0.26
Z	15-ROB-028-COL	49.80	0.05	7.4	10.1	3.04	0.4	47.7	51.8	23.77	12.56 ± 0.14	13.71 ± 0.40
AE	15-ROB-058-NAC	41.29	0.25	6.1	8.4	3.06	2.7	46.5	50.8	19.20	1.413 ± 0.019	1.357 ± 0.039
OE	15-ROB-064-MUS	40.02	0.17	7.9	10.0	3.06	1.4	49.1	49.5	19.64	0.766 ± 0.011	0.667 ± 0.019
AO	16-ROB-002-NLI	80.44	0.21	6.0	7.6	3.08	2.4	48.9	48.7	39.37	0.4654 ± 0.0066	0.438 ± 0.013
AA	16-ROB-032-WIN	41.71	0.18	5.8	9.2	3.03	1.8	43.4	54.7	18.12	2.357 ± 0.031	2.40 ± 0.07

Formatted: Superscript

BB	16-ROB-042-SSU	42.48	0.44	6.8	11.1	3.05	3.7	42.2	54.1	17.91	2.388 \pm 0.034	2.195 \pm 0.064
CC	16-ROB-052-BASa	82.32	0.36	6.3	10.6	3.04	3.2	41.7	55.0	34.35	2.785 \pm 0.034	2.588 \pm 0.075
FF	16-ROB-054-BAS	41.59	0.15	5.9	8.6	3.04	1.6	45.7	52.6	19.02	2.384 \pm 0.029	2.527 \pm 0.073
GG	16-ROB-069-MNM	42.34	0.74	5.8	9.3	3.09	7.2	40.8	52.1	17.26	0.3850 \pm 0.0095	0.402 \pm 0.015
HH	16-ROB-071-MNM	40.92	0.49	6.3	9.3	3.08	4.7	44.1	51.2	18.06	0.4866 \pm 0.0081	0.537 \pm 0.020
II	16-ROB-072-MNM	41.50	0.28	6.2	6.9	3.11	3.2	51.4	45.4	21.33	0.3531 \pm 0.0068	0.3101 \pm 0.0090
JJ	16-ROB-082-WIN	83.39	0.46	4.0	10.2	2.99	4.9	31.3	63.8	26.12	0.4409 \pm 0.0068	0.410 \pm 0.012
KK	16-ROB-084-WIN	70.52	0.66	4.7	10.4	3.03	6.5	33.7	59.8	23.77	0.727 \pm 0.010	0.633 \pm 0.018
LL	16-ROB-085-WIN	42.70	0.24	5.1	6.7	3.08	3.1	47.5	49.4	20.27	0.758 \pm 0.011	0.791 \pm 0.025
MM	16-ROB-106-WAL	42.47	0.39	6.9	7.3	3.12	4.0	52.1	43.9	22.13	2.217 \pm 0.030	2.357 \pm 0.068
NN	16-ROB-118-BBY	41.85	0.28	5.5	7.9	3.06	3.1	45.1	51.7	18.89	2.183 \pm 0.030	2.286 \pm 0.071

Whole-rock Chips

DD	16-ROB-052-BASb	81.62	0.10	4.7	9.0	2.99	1.1	39.5	59.3	32.26	2.814 \pm 0.041	2.59 \pm 0.08
EE	16-ROB-052-BASc	81.66	0.11	5.3	15.5	2.93	0.9	30.1	69.1	24.57	3.027 \pm 0.047	2.59 \pm 0.08

* For samples with more than one measurement, the concentration represents the error-weighted mean and the standard error.






Figure 5 Comparison of cosmogenic ${}^3\text{He}$ concentration measured in virtual-separated pyroxene samples (Bergelin: this study) and pyroxene-separated samples (Balter: Balter-Kennedy et al., 2020). (a) compares the measured cosmogenic ${}^3\text{He}$ concentration in pyroxene and (b) shows the relative difference between the two methods. Error bars show 1-sigma uncertainties. Note that both data sets have been normalized to CRONUS-P (Blair et al., 2015) as described in section 4.1.

5 Discussion

5.1 Reproducibility

We analyzed multiple aliquots from two samples to evaluate both the homogeneity of pyroxene fractionation within a sample and the reproducibility of measured ${}^3\text{He}$ concentration. Aliquots of 15-OTW-047-CHA show consistent pyroxene content,



varying less than 1% by weight, indicating homogeneity between aliquots. In contrast, aliquots from sample 15-OTW-021-DUT exhibit greater variability and heterogeneity, with pyroxene content differing up to 5% by weight.

430 Additionally, aliquots b and c of sample 16-ROB-052-BAS are small rock chips, where aliquot a is a crushed, mixed grain sample. While aliquot a contains 42% pyroxene by weight, the rock chips showed both lower and more variable pyroxene content. Notably, aliquot 16-ROB-052-BASc displayed only 30% pyroxene, and is among the lowest percentages measured (Table 3).

435 The combined sample density computed from the CT-estimated mineral composition for each sample ranges between 2.93 - 3.13 g cm⁻³ with an average and standard deviation of 3.02 +/- 0.05, which is in agreement with previously reported density for Ferrar dolerite (2.96-3.06 g cm⁻³: Margerison et al., 2005). However, the observed variability in pyroxene percentage between aliquots and range in combined density among samples does not exclude a potential systematic density segregation of plagioclase and pyroxene grains during crushing, sieving, washing, and handling of the samples, leading to a likely 440 enrichment of heavier pyroxene grains and, consequently, an increased density. This highlights the importance of performing CT scans for each sample aliquot prior to cosmogenic ³He analysis, as we cannot assume uniformity across aliquots of either rock chips or crushed grain samples. That is, the sample being CT scanned must be the exact same sample that is analyzed in the mass spectrometer.

445 In Fig. 6, we show variability in replicate ³He measurements obtained from (i) the two pyroxene-separated and virtually-separated methods, and (ii) multiple aliquots of the same sample measured using the same method. The scatter between the pyroxene-separated and virtually-separated ³He measurements exceeds the internal scatter observed for the pyroxene-separated method only (Fig. 2b in Balter-Kennedy et al., 2020). However, despite the limited number of virtually-separated replicates available, this method tends to display low internal scatter and reproducibility compared to that of the pyroxene-separated 450 method. This suggests that the scatter observed in Fig. 5 is related to sample-specific differences between the material analyzed using the two methods, rather than internal methodological inconsistency.





Figure 6 Relative standard deviation of replicated cosmogenic ${}^3\text{He}$ analysis for the pyroxene-separated and virtual-separated method. White dots show replicated measurements on pyroxene-separated samples obtained from Balter-Kennedy et al. (2020, Fig 2b), where the grey dot shows the replicated CRONUS-P measurement analyzed with the pyroxene-separated samples. Blue dots show the replicated measurement on aliquots of virtual-separated samples from this study, where the light blue dot shows the replicated CRONUS-P measurements analyzed with the virtual-separated samples. The red dots show the relative standard deviation of the paired measurements from the two methods on the same sample. The size of the circle indicates the number of samples used in the calculated relative standard deviation. To summarize, both methods have better internal replicability than the replicability between the two methods.

5.2 Inclusion of ilmenite as a trace mineral

Ferrar dolerite is known to contain trace amounts of oxide minerals, most commonly ilmenite (TiFeO_3) (Elliot and Fleming, 2021). During pyroxene separation, most iron oxides readily dissolve in HCl. However, ilmenite, like pyroxene, is less soluble in HCl and more resistant to HF etching, and may remain present in the pyroxene-separated samples despite additional purification steps such as handpicking under a binocular microscope or further magnetic separation following HF etching. If not properly accounted for, its persistence can influence the measured cosmogenic ${}^3\text{He}$ concentration.

Ilmenite ~~exhibits~~ similar or potentially greater helium retentivity compared to pyroxene, but is mostly, if not completely, degassed at temperatures used for the extraction of helium in pyroxene. We show this through a simple experiment involving sample 16-ROB-069-MNM. We prepared two aliquots from purified pyroxene separates, and two aliquots from the magnetic fraction (estimated *by eye* to contain 25–35% ilmenite) and analyzed the ${}^3\text{He}$ and ${}^4\text{He}$ concentration across the three heating steps used for complete degassing analysis (Table 4). We find that measured ${}^3\text{He}$ concentrations in samples with high ilmenite concentrations were ~~lower~~ than those in pure pyroxene separates, which is in agreement *with* that observed by Margerison et al. (2005).

Deleted: 6

Deleted: displays

Deleted: distinguishably

Table 4 Measured ${}^3\text{He}$ and ${}^4\text{He}$ concentration in aliquots from the pyroxene separated and magnetic separated (containing pyroxene and oxides) fraction of sample 16-ROB-69, and the fractional release from each analytical heating step.

Sample name	Aliquot	Sample mass (mg)	Heating time (min)	Heating Temp (C)	Measured ${}^3\text{He}$ (Matoms/g)	${}^3\text{He}$ % release	Total measured ${}^3\text{He}$ (Matom/g)	Measured ${}^4\text{He}$ (Tatom/g)	${}^4\text{He}$ % release	Total measured ${}^4\text{He}$ (Tatom/g)
16-ROB-069-MNM-px (Pyroxene)	a	25.36	3	600	244.8 ± 6.2	65.2	375.7 ± 7.2	359.1 ± 8.2	52.6	682 ± 11
			12	1200	124.2 ± 3.6	33.1		312.6 ± 7.2	45.8	
			15	1300	6.7 ± 0.8	1.8		10.7 ± 0.25	1.6	
16-ROB-069-MNM-px (Pyroxene)	b	24.72	3	600	242.8 ± 5.9	63.8	380.8 ± 7.1	301.92 ± 6.9	50.3	601 ± 10
			12	1200	134.9 ± 3.8	35.4		287.5 ± 6.6	47.9	
			15	1300	3.1 ± 0.7	0.8		11.37 ± 0.26	1.9	
16-ROB-069-MNM-px-ox (Pyroxene and oxide)	a	24.66	3	600	160 ± 4.3	43.7	366 ± 7.2	530 ± 12	39.3	1349 ± 21
			12	1200	188.8 ± 5.7	51.6		750 ± 17	55.6	
			15	1300	17.2 ± 1	4.7		68.1 ± 1.6	5.1	
16-ROB-069-MNM-px-ox (Pyroxene and oxide)	b	24.56	3	600	126.9 ± 3.7	35.4	358.5 ± 6.7	383.4 ± 8.8	31.4	1219 ± 20
			12	1200	205.2 ± 5.4	57.2		767 ± 18	62.9	
			15	1300	26.5 ± 1.3	7.4		68.5 ± 1.6	5.6	

Deleted: ¶

... [1]


The apparent cosmogenic ${}^3\text{He}$ production rate in ilmenite is $93.6 \text{ atoms g}^{-1} \text{ yr}^{-1}$, corresponding to a production ratio of 0.78 relative to pyroxene (Larsen et al., 2019). That means, for samples with, e.g., 30% ilmenite content, we would expect to measure a concentration that is about 7% less than that of pure pyroxene-separates. This is generally in agreement with that observed (Table 4), suggesting that ${}^3\text{He}$ in ilmenite does completely degas during analysis. However, the increased release of helium at higher temperature heating steps indicates that ilmenite retains helium more effectively than pyroxene, which is in agreement with limited diffusion data for ${}^4\text{He}$ in pyroxene and ilmenite (Futagami et al., 1993). As a result, failing to recognize ilmenite contamination of a supposed pyroxene sample may lead to an underestimation of the cosmogenic ${}^3\text{He}$ concentration in pyroxene, due to the lower production rate in ilmenite, or, potentially, a heating schedule sufficient for degassing pyroxene but not ilmenite.

In the virtual mineral separation method, we can correct for the presence of ilmenite using the high-attenuation phase (oxide) in the CT scans. That is, of course, based on the assumption that the high-density oxide content is all presumed to be ilmenite. In Fig. 7a, we compare the measured ${}^3\text{He}$ concentrations between the pyroxene-separated and virtual-separated methods, where we correct the mass to include the oxide weight fraction and account for the difference in cosmogenic ${}^3\text{He}$ production rate (Larsen et al., 2019). When considering the pyroxene weight percent only, we get a mean ratio (Bergelin $[{}^3\text{He}]$ / Balter $[{}^3\text{He}]$) and standard deviation of 0.99 ± 0.08 , compared to 0.94 ± 0.08 if we assume that the oxides are part of the ${}^3\text{He}$ contributing

Deleted: the

mass. This apparent systematic underestimation is not completely consistent with the results above, but it is evident that marginally better agreement is obtained by assuming that the oxide fraction does not contribute ${}^3\text{He}$. Regardless, we find no correlation between the oxide phase weight percentage from the CT scan and the measured ${}^3\text{He}$ concentration ratio (Fig. 7b).

505 This indicates that accounting for the oxide phase contamination, such as ilmenite, does not improve the agreement between the two methods, and ilmenite is therefore not the primary source of the scatter observed in Figs. 6 and 7b.



510 Figure 7 the relative difference between measured cosmogenic ${}^3\text{He}$ concentration in virtual-separated pyroxene (Bergelin: this study) and pyroxene separates (Balter: Balter-Kennedy et al., 2020), compared to (a) measured cosmogenic ${}^3\text{He}$ concentration in pyroxene, and (b) oxide weight percentage. Grey data show the ratio where virtual-separated ${}^3\text{He}$ concentrations are determined from the pyroxene weight as indicated in fig. 5, where the concentration is simply N/w_{px} , and where N is the ${}^3\text{He}$ amount (atoms), and w_{px} is the pyroxene weight. The red data show the virtual-separated ${}^3\text{He}$ concentration determined from the weight of pyroxene and ilmenite and corrected for the difference in production rate, such that the concentration is $N/(w_{\text{px}} + w_{\text{ox}} \times 0.78)$, and where w_{ox} is the oxide weight.


5.3 Sources of scatter in the data set

We see more scatter between the pyroxene-separated and virtual-separated methods than expected from measurement uncertainty. This is evident in Fig. 5b (the relative difference is indistinguishable from 1) and Fig. 6, which show no systematic bias, but do show scatter exceeding measurement uncertainty. Assuming that the mass-spectrometric ${}^3\text{He}$ measurements 520 themselves are unbiased, the difference in apparent ${}^3\text{He}$ concentration in pyroxene in the same sample obtained by physical and virtual mineral separation must stem from either incomplete purification of pyroxene in the physical case, or incorrect quantifications of pyroxene weight in the virtual case. As both are possible and difficulties in pyroxene purification are routinely observed (Bromley et al., 2014), we do not know a priori which value is more accurate.

525 In addition to ilmenite, other mineral inclusions can contribute to variability in the measured ${}^3\text{He}$ concentration. Due to the fine-grained, microcrystalline texture of Ferrar dolerite, physical separation of pure pyroxene grains is challenging, often resulting in pyroxene separates including accessory minerals such as plagioclase, ilmenite, and other minor phases. These inclusions affect cosmogenic ${}^3\text{He}$ exposure dating: plagioclase contributes to mass but without hosting ${}^3\text{He}$ atoms, while oxides

Deleted: and instead implies that ${}^3\text{He}$ produced in the oxide fraction was not detected in the measurements, so marginally more accurate results are obtained by assuming that the oxide fraction does not contribute ${}^3\text{He}$

Deleted:



Deleted:

Deleted:

Deleted: 6

Deleted: 6

Deleted: 5b (the relative difference is indistinguishable from 1)

Deleted: where we observe no systematic bias

Deleted: derived from physical and virtual mineral separation

such as ilmenite contribute helium but at a lower production rate, both leading to scatter and a systematic underestimation of ^3He concentration in pyroxene separates. Such bias would result in a ratio systematically >1 in Fig. 5b, which is not the case.

The CT-based virtual mineral separation method can resolve these issues by identifying and excluding non-pyroxene minerals, 545 which would be expected to improve analytical accuracy. However, some minor uncertainties and potential sources of scatter remain due to misclassification of CT voxels and/or varying diffusivity behavior in bulk sample grains. For example, when determining the workflow for identifying the mineral phases in the CT image analyses, the estimated mineral abundance deviated by up to 4% from the known value (Table 2), indicating occasional voxel misclassification, but also perhaps reflecting that the separates used to construct the control samples were themselves not perfect. We note that a perfect segmentation is 550 not possible, due to both the resolution limitations of the data and the simplified 3-phase model we assume. Further, the degassing process is sensitive to mineral composition and grain size. We account for this by applying the highest determined retentivity for ^3He in plagioclase and the lowest in pyroxene when determining the oven heating schedule. However, inter-grain variability may still result in percent-level ^3He loss from pyroxene and retention in plagioclase during oven degassing and before ^3He analysis, resulting in internal scatter in the data. Further, both the CT segmentation method and conversion of 555 voxel volume to mass are affected by the compositional variability for each mineral phase, adding to the internal scatter.

Deleted: both

Other potential sources of discrepancy between the pyroxene-separation and virtually-separation methods, related to the sample-specific scatter observed in Fig. 6, include heterogeneous distribution of ^3He within a sample and the presence of 560 noncosmogenic ^3He from nucleogenic production. While heterogeneity of ^3He in a sample arises from the decrease in production rate with sample thickness, crushing and sample preparation should assure thorough mixing. However, the virtual mineral separation process involves limited sample processing of very small aliquots (<100 mg), so this could result in some variability. The noncosmogenic component of ^3He in pyroxene has been estimated in various studies to be less than 10^7 atoms 565 g^{-1} (Margerison et al., 2005; Kaplan et al., 2017; Eaves et al., 2018). However, for samples exposed for more than a million years, this contribution is less than 3% of the total ^3He concentrations in samples, so we see no means by which variability in noncosmogenic ^3He could significantly affect our results.

Formatted: Superscript

Most errors we envision for the two methods would lead to a systematic bias one way or the other, including contamination in pyroxene-separates, and incorrect estimation of pyroxene weight percent in virtual-separates. As we do not see a systematic bias, no single one of these possibilities can explain the scatter, but we cannot exclude that one or more of them is present. To 570 summarize, we are not able to attribute percent-level variability between the two separation methods to a specific cause, and furthermore are not able to assess whether the physically- or virtually-separated results are more accurate, but this does not affect the conclusion that no systematic bias between the two methods is evident.

575 **5.4 Advantages and disadvantages of ‘virtual mineral separation’****Deleted: 6**


The virtual-mineral-separation method significantly reduces sample processing time by eliminating labor-intensive steps such as extensive crushing, sieving, magnetic separation, acid leaching, and etching. However, the exact processing time and efficiency highly depend on the number of samples processed in parallel and the specific arrangement of a particular laboratory. In general, hand-crushing, sieving, and rinsing of multiple small (< 1 g) samples, followed by drying and packaging, takes roughly several hours of effort over two to three days. The needed time for the CT scan of a batch of 7-8 samples and subsequent image processing takes roughly 3-4 hours, given that the workflow for uniquely classifying each mineral phase using Blob 3D software has already been established. Degassing helium from plagioclase in the vacuum oven for 8 hr at 220°C prior to analysis in the mass spectrometer takes a day. However, besides loading and unloading the oven, this process is hands-free.

585

To further enhance the efficiency of this method, we directly packaged and analyzed aliquots of whole rock chips collected directly in the field, omitting any physical processing. However, some analytical limitations arise from minimizing the sample processing to this extent.

590 The concentration of ${}^4\text{He}$ in a sample is largely influenced by its nucleogenic production through the decay of uranium (U) and thorium (Th). For the Ferrar Dolerite, dated to 183 Ma (Burgess et al., 2015), this results in a ${}^4\text{He}/{}^3\text{He}$ ratio ranging from 2.6×10^4 to 2.2×10^6 . During standard pyroxene separation, HF leaching removes the surface-implanted ${}^4\text{He}$ from the grain boundary, significantly reducing the total ${}^4\text{He}$ concentration (Bromley et al., 2014). In the virtual-separation method, where HF leaching is omitted, we therefore observe an increase in ${}^4\text{He}$ concentration (Fig. 8 and Table S3). An increased ${}^4\text{He}$ concentration, in turn, limits the ${}^3\text{He}$ signal available for analysis because of the need to adjust sample sizes such that the total helium pressures remain within a calibrated range. Although there exist strategies to improve ${}^3\text{He}$ measurements at very low ${}^3\text{He}/{}^4\text{He}$ ratios (Amidon and Farley, 2010), they are not routinely used. However, this issue is also somewhat specific to the


595

Deleted: and**Deleted: es****Deleted: 1**

605 **Figure 8 Comparison of cosmogenic ^{40}He concentration measured in virtual-separated pyroxene samples (Bergelin: this study) and
pyroxene-separated samples (Balter: Balter-Kennedy et al., 2020).**


The analysis of whole rock chip aliquots (16-ROB-052-ASb,c) presents additional challenges when both crushing and HF leaching are omitted from the sample preparation. We found that gas cleanup and helium purification were more challenging for measurement on rock chips, most likely because unprocessed rock chips contain higher concentrations of hydrated and/or volatile-containing minerals or weathering products, that were apparently removed during crushing and rinsing. Thus, 610 developing a routine analytical workflow for whole rock chips, limiting the sample processing to only involve collecting and packaging a single < 100 mg rock chip into 1 ml vials, would most likely require additional cleaning measures for released gas, such as attaching a liquid nitrogen cold trap or incorporating additional getters to absorb CO_2 and H_2O .

615 However, the CT data of the whole-rock chips does provide additional context that is lost when the sample is crushed, which may help with both understanding the material and improving the segmentation (Fig. 9). The pyroxene is clearly zoned, predominantly consisting of a brightening toward grain rims or relict grain boundaries most likely reflecting exchange of Mg for Fe or Ca, and explaining the breadth of the pyroxene histogram peak (Fig. 3c). Multiple high-attenuation phases are evident, and grain attenuation and/or morphology could help differentiate them. Multiple low-attenuation phases are also evident, with the darker regions possibly reflecting alkali feldspar amidst the slightly brighter plagioclase. Rock chips may also be easier to 620 collect and transport, and can record a better-constrained depth interval than crushed larger samples. Finally, segmentation of rock-chip data is more straightforward than for crushed grains, as the extensive blurring between small fragments and air is avoided.



Deleted: as

Deleted: .



Deleted:

Figure 9 CT image of whole-rock chip 16-ROB-52-BASb (DD; image slice 358). The rock chip measures ~4 mm in diameter. Pyroxene (px) shows zoning consisting of brightening towards the grain boundary. Multiple high-attenuation phases are observed for oxides (ox), as well as multiple low-attenuation phases are observed for plagioclase (plag).

Deleted: 5

630 5.5 Implications of virtual mineral separation for cosmogenic nuclide applications

Deleted: 6

Virtual mineral separation greatly simplifies the workflow for ${}^3\text{He}$ measurements in mafic lithologies, and, therefore, has the potential to enable several new Earth science applications as follows.

635 Exposure-dating of surface features typically involves single boulders or clasts, where many of the processes that affect the exposure ages are inherently stochastic and quantized. For example, a boulder has either tipped over or cracked after it was emplaced, or it has not. Therefore, accurately dating a landform subjected to stochastic disturbance requires comparing an observed distribution of exposure ages to that predicted by a process model (e.g. Balco, 2011, 2020; Applegate et al., 2010; Applegate et al., 2012). The greater the number of exposure ages that can be obtained from a single landform and used in such a comparison, the more robust the age estimate. In Antarctica, Ferrar Dolerite is one of the most dominant surface lithologies 640 in ice-free areas due to its resistance to weathering, so is commonly extremely abundant on glacial moraines, drift surfaces, colluvial aprons, and other surface features whose age is of interest and that are subject to stochastic processes of both cosmogenic-nuclide inheritance and postdepositional disturbance. The virtual mineral separation method of exposure dating applied to these features would enable faster, more efficient, and more complete characterization of frequency distributions of apparent exposure ages, which in turn would improve understanding of inheritance distributions, quantification of 645 postdepositional disturbance frequency, and better estimates of the true ages of these features.

650 The virtual mineral separation method allows collection of extremely small (< 1g) sample mass, rather than the conventional size of hundreds of grams. That is, a single rock chip (100 mg) can be collected from boulders across a field and directly packaged into 1mL vials for analysis, [given that the specific requirements for the lithology are met as described below](#). This improves speed in the field, reduces visual impacts, and reduces logistical requirements, especially for Antarctic applications or for remote sites where samples are often carried out by foot. In addition, minimizing sample size is important for sample-limited applications such as subglacial bedrock exposure dating. For example, small clasts of a lithology presumed to be Ferrar 655 dolerite were collected from the base of the Taylor Dome ice core in 1994 (Grootes et al., 1994; Steig et al., 2000). If this material had not been subsequently lost, the virtual mineral separation method could have allowed detection and quantification of surface exposure in these clasts with minimal consumption of the samples.

660 Measuring multiple cosmogenic nuclides in a single sample is valuable for various applications of exposure dating. Multiple-nuclide systematics (e.g., ${}^{36}\text{Cl}/{}^{10}\text{Be}/{}^3\text{He}$ ratios in pyroxene) allow corrections for surface weathering and erosion, as well as quantifying when and how often a surface has experienced burial. However, measurement of ${}^{10}\text{Be}$ and ${}^{36}\text{Cl}$ in Ferrar Dolerite pyroxenes (Bergelin et al., 2024; Swanger et al., 2025) is costly and labor-intensive, requiring extensive mineral separation

and chemical extraction. As a screening tool, virtual mineral separation enables large quantities of cosmogenic ${}^3\text{He}$ measurements across a landscape and identifies an optimal subset of sample candidates for multiple-nuclide studies.

665 Finally, cost, speed, and efficiency are critical limits for emerging applications of exposure dating in Antarctic surface process studies. Recent research on dispersal and succession in Antarctic terrestrial ecosystems has focused on the importance of surface disturbance by past glaciation of currently ice-free areas (Lyons et al., 2016; Diaz et al., 2020, and references therein), and on the use of exposure-dating to quantify the timing and frequency of glacial (and other) disturbance as a means of
670 understanding ecosystem dynamics. Since these landscape disturbance properties are, again, inherently stochastic in nature, large quantities of data are needed to characterize the large-scale relationship of surface disturbance to species dispersal and diversity. Virtual mineral separation would permit collection of such large data sets of exposure ages of surface clasts, or, potentially, could be applied to soils or sediments derived from weathering of Ferrar dolerite to collect depth-profile data needed to quantify subsurface mixing of soils (Diaz et al., 2020; Putkonen et al., 2008; Schaller et al., 2009).

675 In this study, we specifically choose to focus on Ferrar dolerite because it presents a simple case for which (i) the mineralogy predominantly comprises of a non-helium-retentive plagioclase and helium-retentive pyroxene with minor other mineral constituents, and (ii) a large data set of ${}^3\text{He}$ measurements on pyroxene separates already exists. In addition, the ubiquity of Ferrar dolerite surface clasts throughout the Transantarctic Mountains provides numerous potential field applications.
680 However, other mafic lithologies, such as gabbroic rocks or potentially fine-grained basalts where mineral separation is near impossible, may also be suitable, providing they share similar mineralogical characteristics. The key requirement is that the rock be composed of minerals with strongly contrasting helium retentivity, such that any minerals that do not completely retain helium over geologic time scales of interest can be degassed without degassing the retentive minerals. It must also be possible to identify the retentive minerals accurately enough to determine their proportion in the rock and calculate an appropriate ${}^3\text{He}$ production rate. Thus, rocks that include relatively retentive, high-density minerals such as pyroxene, olivine (Trull et al., 1991; Shuster et al., 2004), or oxides (Futagami et al., 1993) with relatively diffusive, low-density minerals such as quartz (Shuster and Farley, 2005; Tremblay et al., 2014) or most feldspars (Bergelin et al., 2025) are likely suitable for virtual mineral separation. On the other hand, lithologies containing significant amounts of magmatic helium (which is not significant in the Ferrar dolerite) may present complications requiring additional processing steps (e.g., crushing operations to degas gas or fluid inclusions) and limiting the advantages of virtual separation.

690 Conclusion

We have shown that the cosmogenic ${}^3\text{He}$ concentration in pyroxene in mafic rocks can be measured without physically separating the pyroxene. First, we explored the diffusion kinetics of ${}^3\text{He}$ in plagioclase and pyroxene to establish that a single heating step of 220°C for 8 hr is sufficient to reach complete separation between ${}^3\text{He}$ released from plagioclase and that retained

Deleted: The key requirement is that the rock mainly be composed of minerals with strongly contrasting helium retentivity. If additional helium-retentive minerals, like ilmenite or olivine (Trull et al., 1991; Futagami et al., 1993; Shuster et al., 2004) are present, or non-retentive minerals with different diffusion kinetics (e.g., other feldspars; Bergelin et al., 2025), then the lithology is likely suitable for virtual mineral separation. On the other hand

Deleted: 1

Deleted: 7

in pyroxene. Second, we determined the weight percentage fraction of pyroxene in crushed whole-rock samples through CT scanning. This step is important for all samples, as we cannot assume homogeneity between aliquots, and to apply the virtual mineral separation to exposure dating applications. Third, we compared the virtual mineral separation approach with existing ^3He measurements in pyroxene separates (Balter-Kennedy et al., 2020).

We find that virtual mineral separation yields ^3He concentrations in pyroxene that are indistinguishable from those derived from physical mineral separation, with no evidence of systematic bias. While observed differences between the results of the two methods exceed estimated analytical uncertainty, these discrepancies appear to be stochastic rather than systematic. This makes it difficult to assess which method is preferable, as both include unquantified uncertainties that can either under- or overestimate the measured nuclide concentration. However, a lack of systematic bias between the methods combined with percent-level differences, even if the latter are poorly explained, lead us to conclude that the virtual mineral separation method is accurate for cosmogenic ^3He exposure dating.

The virtual mineral separation method offers a more efficient approach to measure the cosmogenic ^3He concentration in pyroxene in mafic rocks, requiring limited physical processing time. This makes collection of large quantities of cosmogenic ^3He data for exposure dating relatively easy, fast, and cost-effective, which in turn enables new emerging applications of exposure dating, such as quantifying stochastic surface processes, ecosystem studies, and potential subglacial bedrock exposure dating.

Code and data availability

All data described in the paper are included in the supplement.

Author contribution

MB carried out the experiments and measurements, with assistance from GB and RAK in experimental design and data reduction. MB led the data interpretation and prepared the manuscript with contributions from all authors.

Competing interests

Greg Balco is a member of the editorial board of Geochronology.

Acknowledgments

The LLNL portion of this work was carried out under Contract DE-AC52-07NA27344; this is LLNL-JRNL-2007614.

Deleted: -DRAFT

Financial support

This research has been supported by the National Science Foundation (grant no. 2139497) and by the Ann and Gordon Getty Foundation

735 References

Amidon, W. H. and Farley, K. A.: Mass spectrometric ${}^3\text{He}$ measurement in ${}^4\text{He}$ -rich phases: Techniques and limitations for cosmogenic ${}^3\text{He}$ dating of zircon, apatite, and titanite, *Geochemistry, Geophysics, Geosystems*, 11, n/a-n/a, 10.1029/2010gc003178, 2010.

Applegate, P. J., Urban, N. M., Laabs, B. J. C., Keller, K., and Alley, R. B.: Modeling the statistical distributions of cosmogenic exposure dates from moraines, *Geosci Model Dev*, 3, 293-307, 10.5194/gmd-3-293-2010, 2010.

740 Applegate, P. J., Urban, N. M., Keller, K., Lowell, T. V., Laabs, B. J. C., Kelly, M. A., and Alley, R. B.: Improved moraine age interpretations through explicit matching of geomorphic process models to cosmogenic nuclide measurements from single landforms, *Quaternary Research*, 77, 293-304, 10.1016/j.yqres.2011.12.002, 2012.

Balco, G.: Contributions and unrealized potential contributions of cosmogenic-nuclide exposure dating to glacier chronology, 1990–2010, *Quaternary Science Reviews*, 30, 3-27, 10.1016/j.quascirev.2010.11.003, 2011.

745 Balco, G.: Glacier Change and Paleoclimate Applications of Cosmogenic-Nuclide Exposure Dating, *Annual Review of Earth and Planetary Sciences*, 48, 21-48, 10.1146/annurev-earth-081619-052609, 2020.

Balmer-Kennedy, A., Bromley, G., Balco, G., Thomas, H., and Jackson, M. S.: A 14.5-million-year record of East Antarctic Ice Sheet fluctuations from the central Transantarctic Mountains, constrained with cosmogenic ${}^3\text{He}$, ${}^{10}\text{Be}$, ${}^{21}\text{Ne}$, and ${}^{26}\text{Al}$, *The Cryosphere*, 14, 2647-2672, 10.5194/tc-14-2647-2020, 2020.

750 Bergelin, M., Balco, G., Corbett, L. B., and Bierman, P. R.: Production rate calibration for cosmogenic ${}^{10}\text{Be}$ in pyroxene by applying a rapid fusion method to ${}^{10}\text{Be}$ -saturated samples from the Transantarctic Mountains, Antarctica, *Geochronology*, 6, 491-502, 10.5194/gchron-6-491-2024, 2024.

Bergelin, M., Gorin, A. L., Balco, G., and Cassata, W. S.: Diffusion kinetics of ${}^3\text{He}$ in pyroxene and plagioclase and applications to cosmogenic exposure dating and paleothermometry in mafic rocks, *Geochronology*, 7, 493-511, 10.5194/gchron-7-493-2025, 2025.

755 Blard, P. H., Lavé, J., Farley, K. A., Fornari, M., Jiménez, N., and Ramirez, V.: Late local glacial maximum in the Central Altiplano triggered by cold and locally-wet conditions during the paleolake Tauca episode (17–15ka, Heinrich 1), *Quaternary Science Reviews*, 28, 3414-3427, 10.1016/j.quascirev.2009.09.025, 2009.

Blard, P. H., Balco, G., Burnard, P. G., Farley, K. A., Fenton, C. R., Friedrich, R., Jull, A. J. T., Niedermann, S., Pik, R., 760 Schaefer, J. M., Scott, E. M., Shuster, D. L., Stuart, F. M., Stute, M., Tibari, B., Winckler, G., and Zimmermann, L.: An inter-laboratory comparison of cosmogenic ${}^3\text{He}$ and radiogenic ${}^4\text{He}$ in the CRONUS-P pyroxene standard, *Quat Geochronol*, 26, 11-19, 10.1016/j.quageo.2014.08.004, 2015.

Bromley, G., Balco, G., Jackson, M., Balmer-Kennedy, A., and Thomas, H.: East Antarctic Ice Sheet Variability In The Central 765 Transantarctic Mountains Since The Mid Miocene, *Clim. Past Discuss*, [Preprint], in review, 10.5194/cp-2024-21, 2024.

Bromley, G. R. M., Winckler, G., Schaefer, J. M., Kaplan, M. R., Licht, K. J., and Hall, B. L.: Pyroxene separation by HF leaching and its impact on helium surface-exposure dating, *Quat Geochronol*, 23, 1-8, 10.1016/j.quageo.2014.04.003, 2014.

Brook, E. J., Brown, E. T., Kurz, M. D., Ackert, R. P., Raisbeck, G. M., and Yiou, F.: Constraints on age, erosion, and uplift of Neogene glacial deposits in the Transantarctic Mountains determined from in situ cosmogenic ${}^{10}\text{Be}$ and ${}^{26}\text{Al}$, *Geology*, 23, 770 1063-1066, 10.1130/0091-7613(1995)023<1063:Coaeau>2.3.Co;2, 1995.

Bruno, L. A., Baur, H., Graf, T., Schlu, C., Signer, P., and Wieler, R.: Dating of Sirius Group tillites in the Antarctic Dry Valleys with cosmogenic ${}^3\text{He}$ and ${}^{21}\text{Ne}$, *Earth and Planetary Science Letters*, 147, 37-54, 1997.

Burgess, S. D., Bowring, S. A., Fleming, T. H., and Elliot, D. H.: High-precision geochronology links the Ferrar large igneous province with early-Jurassic ocean anoxia and biotic crisis, *Earth and Planetary Science Letters*, 415, 90-99, 775 10.1016/j.epsl.2015.01.037, 2015.

Cerling, T. E.: Dating Geomorphologic Surfaces Using Cosmogenic ${}^3\text{He}$, *Quaternary Research*, 33, 148-156, 10.1016/0033-5894(90)90015-d, 1990.

Cooperdock, E. H. G., Hofmann, F., Tibbets, R. M. C., Carrera, A., Takase, A., and Celestian, A. J.: Technical note: Rapid phase identification of apatite and zircon grains for geochronology using X-ray micro-computed tomography, *Geochronology*, 4, 501-515, 10.5194/gchron-4-501-2022, 2022.

Diaz, M. A., Li, J., Michalski, G., Darrah, T. H., Adams, B. J., Wall, D. H., Hogg, I. D., Fierer, N., Welch, S. A., Gardner, C. B., and Lyons, W. B.: Stable Isotopes of Nitrate, Sulfate, and Carbonate in Soils From the Transantarctic Mountains, Antarctica: A Record of Atmospheric Deposition and Chemical Weathering, *Frontiers in Earth Science*, 8, 10.3389/feart.2020.00341, 2020.

Eaves, S. R., Collins, J. A., Jones, R. S., Norton, K. P., Tims, S. G., and Mackintosh, A. N.: Further constraint of the in situ cosmogenic ${}^{10}\text{Be}$ production rate in pyroxene and a viability test for late Quaternary exposure dating, *Quat Geochronol*, 48, 121-132, 10.1016/j.quageo.2018.09.006, 2018.

Eberhardt, P., Geiss, J., and Grögl, N.: Distribution of rare gases in the pyroxene and feldspar of the Khor Temiki meteorite, *Earth and Planetary Science Letters*, 1, 7-12, 10.1016/0012-821X(66)90096-3, 1966.

Elliott, D. H. and Fleming, T. H.: Chapter 2.1b Ferrar Large Igneous Province: petrology, Geological Society, London, *Memoirs*, 55, 93-119, 10.1144/m55-2018-39, 2021.

Farley, K. A., Malespin, C., Mahaffy, P., Grotzinger, J. P., Vasconcelos, P. M., Milliken, R. E., Malin, M., Edgett, K. S., Pavlov, A. A., Hurowitz, J. A., Grant, J. A., Miller, H. B., Arvidson, R., Beegle, L., Calef, F., Conrad, P. G., Dietrich, W. E., Eigenbrode, J., Gellert, R., Gupta, S., Hamilton, V., Hassler, D. M., Lewis, K. W., McLennan, S. M., Ming, D., Navarro-Gonzalez, R., Schwenger, S. P., Steele, A., Stolper, E. M., Sumner, D. Y., Vaniman, D., Vasavada, A., Williford, K., Wimmer-Schweingruber, R. F., and Team, M. S. L. S.: In situ radiometric and exposure age dating of the martian surface, *Science*, 343, 1247166, 10.1126/science.1247166, 2014.

Fechtig, H. and Kalbitzer, S.: The diffusion of argon in potassium-bearing solids, in: *Potassium–Argon Dating*, edited by: Schaeffer, O. A., and Zahringer, J., Heidelberg, Springer, 68-106, 10.1007/978-3-642-87895-4_4, 1966.

Futagami, T., Ozima, M., Nagai, S., and Aoki, Y.: Experiments on thermal release of implanted noble gases from minerals and their implications for noble gases in lunar soil grains, *Geochimica et Cosmochimica Acta*, 57, 3177-3194, 10.1016/0016-7037(93)90302-d, 1993.

Grootes, P. M., Steig, E. J., and Stuwer, M.: Taylor Ice Dome study 1993-1994: An ice core to bedrock, *Antarctic Journal of the United States*, 29, 79-81, 1994.

Hofmann, F., Cooperdock, E. H. G., West, A. J., Hildebrandt, D., Strößner, K., and Farley, K. A.: Exposure dating of detrital magnetite using ${}^3\text{He}$ enabled by microCT and calibration of the cosmogenic ${}^3\text{He}$ production rate in magnetite, *Geochronology*, 3, 395-414, 10.5194/gchron-3-395-2021, 2021.

Kaplan, M. R., Lichti, K. J., Winckler, G., Schaefer, J. M., Bader, N., Mathieson, C., Roberts, M., Kassab, C. M., Schwartz, R., and Graly, J. A.: Middle to Late Pleistocene stability of the central East Antarctic Ice Sheet at the head of Law Glacier, *Geology*, 45, 963-966, 10.1130/g39189.1, 2017.

Ketcham, R. A.: Computational methods for quantitative analysis of three-dimensional features in geological specimens, *Geosphere*, 1, 10.1130/ges00001.1, 2005.

Ketcham, R. A. and Mote, A. S.: Accurate Measurement of Small Features in X-Ray CT Data Volumes, Demonstrated Using Gold Grains, *Journal of Geophysical Research: Solid Earth*, 124, 3508-3529, 10.1029/2018jb017083, 2019.

Ketcham, R. A., Slottke, D. T., and Sharp, J. M.: Three-dimensional measurement of fractures in heterogeneous materials using high-resolution X-ray computed tomography, *Geosphere*, 6, 499-514, 10.1130/ges00552.1, 2010.

Kurz, M. D.: In situ production of terrestrial cosmogenic helium and some applications to geochronology, *Geochimica et Cosmochimica Acta*, 50, 2855-2862, 10.1016/0016-7037(86)90232-2, 1986.

Larsen, I. J., Farley, K. A., and Lamb, M. P.: Cosmogenic ${}^3\text{He}$ production rate in ilmenite and the redistribution of spallation ${}^3\text{He}$ in fine-grained minerals, *Geochimica et Cosmochimica Acta*, 265, 19-31, 10.1016/j.gca.2019.08.025, 2019.

Licciardi, J. M., Kurz, M. D., Clark, P. U., and Brook, E. J.: Calibration of cosmogenic ${}^3\text{He}$ production rates from Holocene lava flows in Oregon, USA, and effects of the Earth's magnetic field, *Earth and Planetary Science Letters*, 172, 261-271, 10.1016/s0012-821X(99)00204-6, 1999.

825 Lovera, O. M., Richter, F. M., and Harrison, T. M.: The $^{40}\text{Ar}/^{39}\text{Ar}$ thermochronometry for slowly cooled samples having a distribution of diffusion domain sizes, *Journal of Geophysical Research: Solid Earth*, 94, 17917-17935, 10.1029/JB094iB12p17917, 1989.

Lovera, O. M., Grove, M., Mark Harrison, T., and Mahon, K. I.: Systematic analysis of K-feldspar step heating results: I. Significance of activation energy determinations, *Geochimica et Cosmochimica Acta*, 61, 3171-3192, 10.1016/s0016-7037(97)00147-6, 1997.

830 Lyons, W. B., Deuerling, K., Welch, K. A., Welch, S. A., Michalski, G., Walters, W. W., Nielsen, U., Wall, D. H., Hogg, I., and Adams, B. J.: The Soil Geochemistry in the Beardmore Glacier Region, Antarctica: Implications for Terrestrial Ecosystem History, *Sci Rep*, 6, 26189, 10.1038/srep26189, 2016.

Margerison, H. R., Phillips, W. M., Stuart, F. M., and Sugden, D. E.: Cosmogenic ^3He concentrations in ancient flood deposits from the Coombs Hills, northern Dry Valleys, East Antarctica: interpreting exposure ages and erosion rates, *Earth and Planetary Science Letters*, 230, 163-175, 10.1016/j.epsl.2004.11.007, 2005.

835 Putkonen, J., Balco, G., and Morgan, D.: Slow regolith degradation without creep determined by cosmogenic nuclide measurements in Arena Valley, Antarctica, *Quaternary Research*, 69, 242-249, 10.1016/j.yqres.2007.12.004, 2008.

Schäfer, J. M., Ivy-Ochs, S., Wieler, R., Leya, I., Baur, H., Denton, G. H., and Schlüchter, C.: Cosmogenic noble gas studies in the oldest landscape on earth: surface exposure ages of the Dry Valleys, Antarctica, *Earth and Planetary Science Letters*, 167, 215-226, 10.1016/s0012-821x(99)00029-1, 1999.

840 Schaller, M., Ehlers, T. A., Blum, J. D., and Kallenber, M. A.: Quantifying glacial moraine age, denudation, and soil mixing with cosmogenic nuclide depth profiles, *Journal of Geophysical Research*, 114, 10.1029/2007jf000921, 2009.

Shuster, D. L. and Farley, K. A.: Diffusion kinetics of proton-induced ^{21}Ne , ^3He , and ^4He in quartz, *Geochimica et Cosmochimica Acta*, 69, 2349-2359, 10.1016/j.gca.2004.11.002, 2005.

845 Shuster, D. L., Farley, K. A., Sisterson, J. M., and Burnett, D. S.: Quantifying the diffusion kinetics and spatial distributions of radiogenic ^4He in minerals containing proton-induced ^3He , *Earth and Planetary Science Letters*, 217, 19-32, 10.1016/s0012-821x(03)00594-6, 2004.

Steig, E. J., Morse, D. L., Waddington, E. D., Stuiver, M., Grootes, P. M., Mayewski, P. A., Twickler, M. S., and Whitlow, S. I.: Wisconsinan and holocene climate history from an ice core at taylor dome, western ross embayment, Antarctica, *Geografiska Annaler: Series A, Physical Geography*, 82, 213-235, 10.1111/j.0435-3676.2000.00122.x, 2000.

850 Swanger, K. M., Babcock, E., Winsor, K., Lamp, J. L., and Winckler, G.: Moraines and dead ice in Taylor Valley, Antarctica, record retreat of alpine and outlet glaciers from Marine Isotope Stage 5 to 4, *Arctic, Antarctic, and Alpine Research*, 57, 10.1080/15230430.2025.2478696, 2025.

Tremblay, M. M., Shuster, D. L., and Balco, G.: Diffusion kinetics of ^3He and ^{21}Ne in quartz and implications for cosmogenic noble gas paleothermometry, *Geochimica et Cosmochimica Acta*, 142, 186-204, 10.1016/j.gca.2014.08.010, 2014.

Tremblay, M. M., Shuster, D. L., Balco, G., and Cassata, W. S.: Neon diffusion kinetics and implications for cosmogenic neon paleothermometry in feldspars, *Geochimica et Cosmochimica Acta*, 205, 14-30, 10.1016/j.gca.2017.02.013, 2017.

855 Trull, T. W., Kurz, M. D., and Jenkins, W. J.: Diffusion of cosmogenic ^3He in olivine and quartz: implications for surface exposure dating, *Earth and Planetary Science Letters*, 103, 241-256, 10.1016/0012-821x(91)90164-d, 1991.

Page 18: [1] Deleted

Marie Bergelin

10/27/25 4:45:00 PM