

Low and Consistent Asymmetry Parameters in Arctic and Mid-latitude Cirrus

Emma Järvinen¹ and Franz Martin Schnaiter^{1,2}

¹Institute for Atmospheric and Environmental Research, University of Wuppertal, Wuppertal, Germany

²schnaiTEC GmbH, Wuppertal, Germany

Correspondence: Emma Järvinen (jaervinen@uni-wuppertal.de)

Abstract. Cirrus clouds play a critical role in the Earth’s radiation budget, yet their shortwave optical properties remain poorly constrained. The asymmetry parameter (g), which governs the angular distribution of scattered light, is particularly sensitive to ice crystal morphology, a property that varies widely in cirrus. To provide observational constraints on the magnitude of g and to investigate its relationship with ice microphysical properties, we analysed simultaneous in situ measurements of particle morphology and angular light scattering using the Particle Habit Imaging and Polar Scattering (PHIPS) probe. These measurements were conducted during the Cirrus in High Latitudes (CIRRUS-HL) campaign in June and July 2021, which sampled both mid-latitude and Arctic cirrus across a range of cloud types and temperatures down to -63°C . We found that cirrus in both regions exhibited a consistently low median asymmetry parameter of 0.737. The observed g values were largely insensitive to variations in temperature, humidity, and crystal aspect ratio, and showed only minor variation across ice habits. A systematic decrease in g with increasing particle size was identified, median g ranging from 0.776 for sub- $30\text{ }\mu\text{m}$ particles in mid-latitude cirrus to minimum values of 0.718 and 0.713 for $175\text{ }\mu\text{m}$ particles in mid-latitude and Arctic cirrus, respectively. The measured values are significantly lower than those commonly used in current radiative transfer schemes, suggesting that cirrus clouds may contribute less to net atmospheric warming than often assumed. These results provide improved observational constraints for the representation of ice cloud optical properties in climate models and support efforts to reduce uncertainties in cirrus cloud radiative forcing.

1 Introduction

To improve the representation of bulk radiative properties of cirrus clouds in climate models and remote sensing algorithms, it is important to have a good understanding of the ice crystal single-scattering properties (Mishchenko et al., 1996; Zhang et al., 1999; Yang et al., 2001). The key single-scattering properties used in a two-stream radiative transfer formalism are the extinction coefficient, single-scattering albedo and the asymmetry parameter (g), from which the latter has a complicated relationship with ice crystal morphology in the solar spectrum. Ice crystal morphological properties, like crystal size, habit as well as degree and type of crystal complexity, have been observed to vary greatly between different cloud types, in-cloud temperature, and geographical location (e.g. Heymsfield et al., 2017; Wolf et al., 2018; Woods et al., 2018; Lawson et al., 2019).

25 Due to this high morphological variability, constraining the single-scattering properties of ice crystals in numerical simulations remains a major challenge. Numerical ray-tracing studies for the solar spectral range predict a wide variety of g values, typically between 0.74 and 0.94, with higher values associated with pristine hexagonal habits and lower values with complex particle shapes or surface roughness (e.g., Jaquinta et al., 1995; Macke et al., 1996b, 1998; Um and McFarquhar, 2007; Yang et al., 2008). However, it is important to correctly quantify g as biases can lead to a significant uncertainty in estimations of the
30 ice cloud radiative effect in climate models (Järvinen et al., 2018; Yi et al., 2013) or in retrieved cirrus microphysical properties from remote sensing observations (Yang et al., 2008; Yi et al., 2017a). Vogelmann and Ackerman (1995) estimated that g must be known within 2-5% in order to constrain computed shortwave fluxes to within approximately 5%.

Observational studies are needed to constrain the magnitude of g in radiative transfer simulations. Unlike numerical estimates of the ice crystal asymmetry parameter, measurements indicate a more moderate variability in the cirrus optical prop-
35 erties. Studies estimating the asymmetry parameter from radiometric observations have reported a low asymmetry parameter around 0.7 in a case study over the midwest USA (Stephens et al., 1990) and previous satellite-based multi-angle observations have reported values for the visible asymmetry parameter below 0.8 (van Diedenhoven et al., 2013; Wang et al., 2014; van Diedenhoven et al., 2014; Cole et al., 2014; van Diedenhoven, 2021). Similarly, lidar observations provide another means of retrieving the asymmetry parameter, with recent studies also reporting values around 0.76 (Gil-Díaz et al., 2025).

40 The limitation of estimating g from remote sensing observations is their dependence on an optical model. The retrieved values are inherently constrained by the model, meaning they can never be lower than the smallest g allowed by the model. As a result, retrievals frequently converge on this lower limit (Järvinen et al., 2023). For example, studies that identify the severely roughened hexagonal aggregate model as the best fit for observations consistently retrieve g values around 0.75 in the visible (Cole et al., 2014; Järvinen et al., 2018; Wang et al., 2018; Forster and Mayer, 2022)

45 A more direct approach to measure g is by using a polar nephelometer. So far, measurements of three airborne polar nephelometers have been reported in the literature: the Polar Nephelometer (PN; Gayet et al., 1998), the Cloud Integrating Nephelometer (CIN; Gerber et al., 2000), and the Particle Habit Imaging and Polar Scattering (PHIPS) probe (Abdelmonem et al., 2016; Schnaiter et al., 2018). These measurements have shown median or average g values between 0.73 and 0.79 (Gerber et al., 2000; Auriol et al., 2001; Garrett et al., 2001, 2003; Gayet et al., 2004, 2006; Febvre et al., 2009; Gayet et al., 2012;
50 van Diedenhoven et al., 2013). However, direct measurements of g have been limited to only a few airborne campaigns and to mostly tropical deep convective or midlatitude systems. So far, the only in situ measurement reported in Arctic cirrus has been one case study of a cirrus spissatus band with a cloud top temperature of -46°C (Garrett et al., 2001). Moreover, prior studies have typically focused on bulk optical properties, with limited information on how g relates to simultaneously measured microphysical properties of the same ice crystal population.

55 The objective of this study is to expand the direct in situ observations of g in cirrus by analyzing a comprehensive, multi-flight dataset from both mid-latitude and Arctic cirrus, collected during the CIRRUS in High Latitudes (CIRRUS-HL) campaign using the PHIPS instrument. This dataset enables a unique dual-regional comparison within the same season and includes single-particle measurements that allow for a direct linkage between ice crystal microphysical properties and their angular light-scattering characteristics. Beyond quantifying the magnitude of g , this study investigates its dependence on key microphysical

parameters, including effective radius, ice water content, habit, and aspect ratio. By combining bulk and particle-resolved analyses, the results provide new observational constraints for developing physically consistent optical parameterisations for radiative transfer applications.

2 Methodology

2.1 The CIRRUS-HL airborne campaign

The CIRRUS-HL airborne campaign took place from 6 June to 28 July 2021 using the High Altitude and Long Range (HALO) research aircraft (Jurkat-Witschas et al., 2025). The original aim of the campaign was to investigate the microphysical and optical properties of Arctic cirrus using in situ and remote sensing instrumentation, with operations initially planned from Kiruna, Sweden. However, due to pandemic-related restrictions, the campaign base was relocated to Oberpfaffenhofen, Germany. This logistical change broadened the scientific scope to include both mid-latitude and Arctic cirrus (see Fig. 1), as Arctic flight operations now required long transit legs and refueling stops in Scandinavia or Iceland. The revised campaign strategy focused on contrasting cirrus clouds in different regions and meteorological conditions. Accordingly, flights were designed to sample natural mid-latitude cirrus in diverse weather regimes, including warm conveyor belt systems, high-pressure in situ cirrus, and convective outflow, as well as cirrus potentially influenced by aviation emissions. The time frame of the campaign coincided with pandemic air traffic restrictions, resulting in a 65 to 80% reduction in daily commercial flight activity compared to the same time frame in 2019 (De La Torre Castro et al., 2023). The campaign period also overlapped with extreme rainfall events in central Europe, resulting in fatal flooding in Germany and Belgium (Lehmkuhl et al., 2022).




Figure 1. Trajectories of CIRRUS-HL measurement flights (F02 - F23) included to the analysis. Research flights targeting Arctic cirrus were performed as double flights with a refuelling stop. These were F05/F06, F08/F09 and F10/F11.

2.2 Microphysical and optical measurements

During CIRRUS-HL in situ measurements of the ice crystal microphysical and optical properties were conducted with the PHIPS airborne cloud instrument (Abdelmonem et al., 2016; Schnaiter et al., 2018). PHIPS is a combination of a single-

80 particle stereo-microscopic imager and a polar nephelometer, which allow for a combined analysis of optical and microphysical properties at the level of individual ice crystals.

2.2.1 Derivation of microphysical properties from stereo-microscopic images

The PHIPS stereo-microscopic imager consists of two CCD camera and microscope assemblies with an angular viewing distance of 120° each acquiring 1360 by 1024 pixel bright field microscopic images of individual cloud particles at a maximum
85 acquisition rate of 10 Hz. During CIRRUS-HL a magnification setting of 6x was used for both camera assemblies corresponding to an optical resolution of about $6\text{ }\mu\text{m}$ and a field of view of approximately $1.1\times 1.5\text{ mm}$, limiting the maximum observable particle size to about 1.5 mm. For each image pair a set of microphysical parameters were retrieved using an algorithm discussed in Schön et al. (2011). These microphysical parameters include the maximum dimension (D_{max}), area-equivalent diameter (i.e., equivalent diameter of a sphere having the same cross-sectional area, D_{eq}), projected area (A_{proj}),
90 and the aspect ratio (AR).

The stereo-microscopic images were also manually inspected and assigned to a habit class. A detailed description of the habit classification scheme can be found in the supplement of Järvinen et al. (2023). While the original habit classification scheme includes 18 categories, for the purposes of this study we reduced the habit classes to 9 cirrus-relevant classes. These include: *plate* (encompassing also sector and skeleton plates), *column*, *frozen droplet*, *side plane*, *bullet rosette*, *mixed rosette*,
95 *capped column*, and *other*, which contains crystals that could not be confidently assigned to a specific habit. Additionally, we introduced an *aggregate* class to account for particles composed of multiple crystals, even though it does not represent a single-crystal habit in the traditional sense.

To account for measurement artefacts, a separate *shattered* class was introduced for particles identified as fragments from shattering events. Shattering was identified through manual visual inspection of stereo-microscopic images from PHIPS. Particles were classified as shattered when multiple small fragments (typically $<100\text{ }\mu\text{m}$) were observed within a single image
100 frame, which has a size of approximately $1.1\text{ mm} \times 1.5\text{ mm}$. The likelihood of multiple unconnected ice particles naturally occurring within such a small field of view is considered low for cirrus conditions, where particle concentrations and clustering tendencies are limited. Additionally, larger crystals were classified as shattered if they exhibited broken edges, truncated branches, or jagged outlines inconsistent with physically intact ice crystal habits. These particles were excluded from further
105 analysis (see Sections 2.2.2 and 2.2.4). Representative stereo-images for each habit class and the shattered class are provided in the supplementary material.

2.2.2 Calculation of particle size distribution and bulk microphysical properties

Although the stereo-microscopic imager is limited to a maximum acquisition rate of 10 Hz, the trigger system of the PHIPS instrument can detect particles at a maximum acquisition rate of 13 kHz. The sensitive area of the instrument is determined by
110 the overlap of the trigger system field of view and the scattering laser intensity distribution and varies between approximately 0.3 and 0.9 mm^2 , depending on the size of the particle (Waitz et al., 2021). This size-dependent sensitive area together with the aircraft's true air speed is used to calculate the sampling volume of PHIPS. Due to the high velocity of a jet aircraft, compression

of air occurs near probe and wing edges. To obtain the true ambient sampling volume, a thermodynamic correction described by Weigel et al. (2016) is applied.

115 The sizing of individual ice crystals is primarily based on the stereo-microscopic images. The images of the same particle from two viewing angles are used to derive the mean D_{eq} and A_{proj} per particle. In case a particle trigger event did not generate a stereo-image due to the limitations of the imaging rate, D_{eq} and A_{proj} were derived based on the relationship between the particle geometrical size and partial scattering cross section measured by the nephelometer as discussed in Waitz et al. (2021). Figures S2 and S3 show the D_{eq} and A_{proj} calibration for the CIRRUS-HL campaign. During CIRRUS-HL flights, typically
120 70-80% of the particle triggers also produced stereo-images, which means that on average the particle sizing was mostly based on the stereo-microscopic images. However, during four flights only below 50% of the triggers produced a stereo-image, in which cases the sizing was mostly based on the nephelometer.

For the calculation of particle size distributions, particles were sorted into logarithmically spaced bins between 15 and 700 μm in D_{eq} . Number concentrations were computed by dividing the number of particles in each bin by the corresponding
125 sampling volume. The bulk microphysical properties ice water content (IWC), extinction coefficient (β_{ext}), effective radius (r_{eff}), and mass-equivalent spherical radius (r_m) were derived using the particle-resolved size and area information.

The extinction coefficient (β_{ext}) was calculated based on the A_{proj} of individual particles, measured in discrete size bins. For each size bin i , the projected area concentration $dA_{proj,i}$ was determined as

$$dA_{proj,i} = \frac{\sum_k A_{proj,i,k}}{V_i}, \quad (1)$$

130 where $A_{proj,i,k}$ is the projected area of the k -th particle in bin i , and V_i is the corresponding sampling volume for that bin. The summation is performed over all detected particles k within the bin.

The total extinction coefficient was then obtained by summing over all size bins:

$$\beta_{ext} = 2 \cdot \sum_i dA_{proj,i}, \quad (2)$$

where the factor of 2 assumes randomly oriented particles and geometric optics (Bohren and Huffman, 1998).

135 The IWC was calculated in a similar way by summing the masses of individual particles detected within each size bin. The mass $m_{i,k}$ of each particle k in bin i was estimated from its projected area $A_{proj,i,k}$ using the empirical power-law relationship derived for mid-latitude ice crystals from Baker and Lawson (2006):

$$m_{i,k} = 0.115 \cdot A_{proj,i,k}^{1.218}, \quad (3)$$

where $A_{proj,i,k}$ is the projected area of each particle k in bin i .

140 The mass concentration per bin dm_i was then calculated as:

$$dm_i = \frac{\sum_k m_{i,k}}{V_i}, \quad (4)$$

where V_i is the bin-specific sampling volume.

The total IWC was obtained by summing over all size bins:

$$\text{IWC} = \sum_i dm_i. \quad (5)$$

145 The effective radius r_{eff} was derived using:

$$r_{\text{eff}} = \frac{3}{2} \cdot \frac{\text{IWC}}{\rho_{\text{ice}} \cdot \beta_{\text{ext}}}, \quad (6)$$

where ρ_{ice} is the bulk density of ice.

The mass-equivalent spherical radius $r_{\bar{m}}$ was calculated as:

$$r_{\bar{m}} = \left(\frac{3\text{IWC}}{4\pi N \rho_{\text{ice}}} \right)^{1/3}, \quad (7)$$

150 where N is the total number concentration. The mass-equivalent sphere radius $r_{\bar{m}}$ represents the radius of a sphere with the average mass per particle of the ensemble, and is useful for linking two-moment microphysical schemes to bulk optical properties (Baran et al., 2025).

2.2.3 Removal of shattering events

Two types of particle size distributions (PSDs) were generated: one standard PSD including all detected particles (i.e., all trigger events), and another including only the subset of imaged particles that were classified as intact based on objective visual criteria from stereo-microscopic images (i.e., not affected by shattering). For the standard PSD, potential shattering events were filtered by applying an interarrival time threshold of 0.1 ms, based on the appearance of a distinct secondary mode in the interarrival time distribution (see Fig. S1).

We emphasize that a visual classification step is essential, as the inter-arrival time method alone cannot detect all shattering events. This is because shattering fragments arriving within very short temporal separations (on the order of a few microseconds) fall below the electronic dead time of the PHIPS detection system. At typical aircraft speeds, this corresponds to a spatial separation of roughly 2 mm. Consequently, shattering fragments that arrive nearly simultaneously within this small area cannot be distinguished by timing information alone and may appear as independent particles in the nephelometer signal. However, these shattering events can be captured by the imager, which has a frame size of approximately 1.5 mm × 1.1 mm.

165 Shattering events not captured by the interarrival threshold were identified by manually inspecting the stereo-scopic image frames and using reproducible morphological criteria. A particle was classified as shattered if: (1) multiple unconnected fragments smaller than approximately 100 μm appeared within the same image frame (on one or both stereo views), or (2) a single particle displayed clear morphological evidence of breakage—such as missing hexagonal facets, truncated arms, or irregular, jagged boundaries.

170 These features are physically implausible for naturally occurring cirrus ice crystals within the small PHIPS sampling volume, given typical cirrus number concentrations and spatial distributions. Visual inspection therefore provides an independent, physically justified means to identify residual shattering events that cannot be filtered algorithmically. Representative examples of such shattering cases are shown in the Supplementary Information.




Figure 2. The mean PHIPS particle size distribution (PSD) and projected area size distribution for mid-latitude and Arctic cirrus. The solid lines show the mean based on 1-Hz PHIPS data in cirrus conditions ($T < -38^\circ\text{C}$ and $IWC > 10^{-5} \text{ g m}^{-3}$) and the dashed lines the mean for only crystals with stereo-images that were manually classified as intact particles.

The effect of manual classification is shown in Figure 2, which compares the mean particle size and projected area distributions before and after removal of shattering artefacts for mid-latitude and Arctic cirrus. The largest differences occur for $D_{eq} < 100 \mu\text{m}$, where the interarrival time method alone overestimates the number concentration due to undetected fragments. After removal based on image inspection, the number concentration decreases toward smaller sizes, producing a more physically consistent distribution.

This refinement primarily affects parameters sensitive to small particles, such as the mean mass radius r_m , while quantities weighted by projected area or extinction (e.g., r_{eff}) remain largely unchanged (see Fig. S5). The manually screened dataset thus provides a more physically realistic representation of the shape of the size distribution of the true ice particle population but is underestimating the number concentration and IWC as only a subset of the triggering events with a corresponding stereo-microscopic image are included in the analysis.

2.2.4 Retrieval of ice asymmetry parameter

The g was retrieved from PHIPS measurements following the framework of Xu et al. (2022). In this method, the measured angular scattering function between 18° – 170° is represented by a truncated series of Legendre polynomials. The underlying assumption of the method is that at scattering angles greater than 18° , the measured signal is dominated by geometrical-optics (GO) contributions—namely refraction and reflection—with only minor contributions from diffraction (Takano and Liou, 1989; Macke et al., 1996b). The unmeasured forward range below 18° is assumed to contain the bulk of the diffraction peak, which is estimated from the measured particle size using a parameterised function derived from scalar diffraction theory.

The GO component of the phase function is reconstructed over the full angular range by fitting the measured data with a truncated Legendre polynomial expansion. The Legendre coefficients are obtained using Gauss–Legendre quadrature, which ensures optimal numerical accuracy. To keep all quadrature nodes within the PHIPS angular range, the maximum degree of the Legendre expansion is restricted to seven. This truncation implies that the method is applicable only to smooth phase functions that can be adequately represented by a low-order polynomial expansion.

We evaluated this requirement for our dataset (see Appendix A) and found that 98% of the measured angular scattering functions show no distinct features such as 22° halo peaks. An uncertainty analysis was performed using 1612 ray-tracing simulations of hexagonal ice crystals with varying aspect ratios, sizes, and surface roughness, generated with the Macke et al. (1996b) ray-tracing code. This closure study demonstrates that, under realistic mixtures of ice crystal morphologies, the retrieval uncertainty of g ranges from approximately -0.001 to +0.02 when halo features are absent from the phase functions. A detailed description of the underlying assumptions, validation tests, and uncertainty quantification is provided in Appendix A.

3 Results

To analyse cirrus microphysical and optical properties by geographical region, the CIRRUS-HL observations were classified into mid-latitude and Arctic cirrus using a latitude threshold of 65°N. Cirrus clouds were defined based on a combination of in-cloud temperature below -38°C and IWC greater than 10^{-5} g m^{-3} , derived from the 1 Hz PHIPS data. The campaign covered a wide geographical range, from 38.2°N to 75.6°N, with the highest cirrus cloud sampled at 14.5 km altitude and the coldest cirrus temperature recorded at -62.8°C. In total, approximately 5625 km of mid-latitude cirrus and 1060 km of Arctic cirrus were sampled, reflecting the greater spatial extent and coverage of mid-latitude observations during the campaign.

3.1 Ice crystal habits in mid-latitudes and Arctic cirrus

Figure 3 shows the fraction of habits as a function of D_{eq} for the mid-latitude and Arctic cirrus cases. The most frequently occurring ice crystal habit class in both regions was the “other” category, accounting for 45% and 66% of all classified particles in mid-latitudes and Arctic, respectively. The high occurrence is especially pronounced for crystals smaller than 100 μm , for which over half of the measured particles fell into this group. In such cases, the classification to the “other” category can be due to the limited optical resolution available for morphological identification. Beyond this technical limitation, the “other” category also reflects the inherent complexity and variability of cirrus ice crystals, which often do not conform to traditional habit classifications (see Fig. 4 for example PHIPS images).

Irregular and compact ice crystals, broadly corresponding to our “other” category, have frequently been reported as the dominant habit type in both mid-latitude and Arctic cirrus (Garrett et al., 2001; Lawson et al., 2019; Wolf et al., 2018). For example, Wolf et al. (2018) observed that compact or irregular habits accounted for 50% and 90% of the crystal populations in Arctic liquid-origin and in-situ cirrus, respectively. In addition to observations from high-resolution imaging instruments, scanning electron microscope analyses have also confirmed the dominance of irregular shapes in cirrus clouds, even among small ice crystals (Magee et al., 2021).




Figure 3. Ice crystal habit distribution as a function of the area equivalent diameter (D_{eq}) for mid-latitude and Arctic cirrus. The numbers indicate the number of ice crystal stereo-image pairs included in the habit analysis per size bin.

The most notable difference between mid-latitude and Arctic cirrus was a more frequent occurrence of aggregation in mid-latitudes compared to the Arctic. For mid-latitude cirrus aggregates was the second most frequent habit class, accounting for 32% of all classified crystals, whereas in Arctic cirrus they accounted only for 7% of all classified crystals. In mid-latitudes the fraction of aggregates was observed to clearly increase towards larger sizes with the highest fraction of 81% found for the largest size group included in the analysis, between 400 and 500 μm . The more frequent occurrence of aggregates in mid-latitude cirrus can be explained through the more dynamic atmosphere at the lower latitudes (Barahona et al., 2017) and higher initial ice number concentration (Sourdeval et al., 2018), potentially promoting aggregation.

In mid-latitude cirrus most of the aggregates were consisting of plates (34%), suggesting a convective origin, or of crystals classified as "other" (27%). A large fraction of aggregates consisted of frozen droplets (25%) but these, together with the simultaneously observed single frozen droplets, originated from research flight 12 targeting a strong overshooting convective system (see Schäfler and Rautenhaus, 2023). In contrast, frozen droplets and their aggregates were rarely detected in other mid-latitude cases, indicating that their occurrence was closely linked to specific convective conditions.

The second notable difference between the two regions was a higher occurrence of bullet and mixed rosettes in the Arctic. Bullet and mixed rosettes comprised 19% of all classified crystals and dominated the population larger than 200 μm , comprising 56–89% in that size range. This makes them the second most frequent habit type in Arctic cirrus. A similar dominance of bullet rosettes as the second most common habit was also reported in Arctic conditions by Wolf et al. (2018). For mid-latitude cirrus bullet and mixed rosettes were common but less frequent comprising 11% of all crystals and reaching maximum occurrence frequencies of 25–28% in the size range of 250–400 μm .

Other identified habits in both regions, each contributing between 2 and 9% to the total population, included side planes, as well as single columns. In contrast, single plates were rare in Arctic cirrus (0.5%) compared to 3.5% in mid-latitude cirrus.




Figure 4. Example ice crystal microscopic images recorded by PHIPS from mid-latitude and Arctic cirrus as a function of their area equivalent diameter (D_{eq}). The distribution of the different habits corresponds to Figure 3.

3.2 Cirrus microphysical and optical properties in mid-latitudes and Arctic

Figure 5 presents the frequency distribution of key bulk microphysical properties of cirrus (N , IWC , β_{ext} and r_{eff}) along with g as a function of latitude, derived from PHIPS observations for the size range of 15 to 700 μm . The selected microphysical parameters are commonly used as input for radiative transfer schemes and were derived from 10-second averaged PHIPS particle size distributions. The asymmetry parameter was calculated from populations of 50 consecutive ice crystals and includes only manually inspected and classified intact (non-shattered) particles. Similarly, r_{eff} was derived using only non-shattered, imaged particles to ensure accurate representation of individual crystal morphology. In contrast, N , β_{ext} and IWC were calculated using the full dataset, including particles without images, since applying the same shattering removal would lead to a systematic underestimation of these quantities. The minimum detection limits for the 10-second averages were approximately 0.5 L^{-1} for N , 0.005 km^{-1} for β_{ext} and 0.05 mg m^{-3} for IWC , based on the instrument's sampling volume and estimated particle mass.

The median total concentration of ice crystals in the 15 to 700 μm size range varied between 1 and 52 L^{-1} and showed no clear latitudinal trend. However, slightly higher overall median values were observed in mid-latitude cirrus (11 L^{-1}) compared to Arctic cirrus (8 L^{-1}) (Table 1). Mean values differed more noticeably, with 120 L^{-1} in the mid-latitudes and 8 L^{-1} in the Arctic. This difference was primarily driven by occasional high concentrations exceeding 100 L^{-1} in the mid-latitude dataset, which were not observed in the Arctic. However, these elevated concentrations are likely influenced by particle shattering effects.

Similar to the total ice number concentration, both the IWC and β_{ext} values did not exhibit a significant latitudinal trend between 42°N and 75°N, where most observations were conducted. Median IWC values ranged from 0.2 to 10 mg m^{-3} , with an overall median of 2.9 mg m^{-3} and 3.0 mg m^{-3} for mid-latitudes and Arctic cirrus, respectively. Also the mean values were similar for mid-latitude and Arctic cirrus, 2.6 and 2.4 mg m^{-3} , respectively, which confirms that the higher number concentrations observed in the mid-latitudes are from small ice crystals that do not contribute significantly to the IWC . Median β_{ext} values ranged from 0.04 to 0.4 km^{-1} , with the same overall median value for both mid-latitude and Arctic of 0.13 km^{-1} .

The only microphysical parameter that shows a clear latitudinal dependence was r_{eff} , with median values increasing from approximately 24 μm within latitudes from 39°N to 42°N to 45 μm within latitudes above 70°N. An exception to this trend was observed near 62°N, where median effective radii around 50 μm were recorded in a liquid-origin system sampled off the coast of Norway during the double flight F18/F19. A similar trend towards higher r_{eff} values in more pristine environments was observed by Gayet et al. (2004) during the Interhemispheric Differences in Cirrus Properties From Anthropogenic Emissions (INCA) field experiments, where the average northern hemispheric effective radius (36 μm) was smaller than that observed in the southern hemisphere (42 μm).

In contrast to the trend in r_{eff} , g showed no systematic latitudinal variation, with median values ranging from 0.710 to 0.745. Notably, the median asymmetry parameter was almost identical in both regions (0.74) indicating a remarkable similarity in the bulk optical scattering characteristics.




Figure 5. Frequency distribution of total concentration (N), ice water content (IWC), extinction coefficient (β_{ext}), effective radius (r_{eff}) and asymmetry parameter (g) observations from the PHIPS as a function of latitude. Microphysical observations are 10-second average values and asymmetry parameter is retrieved for only particles with stereo-images where shattering artefacts are excluded. The black solid lines mark the median line. Latitude bin widths are 1° for microphysical observations and 2° for the asymmetry parameter observations. The histograms show the distribution of the values for all mid-latitude (ML) and Arctic cases using a latitude threshold of 65°N . The dashed line over the histograms indicate the median value.

Table 1. Summary statistics (median, mean, and percentiles) of microphysical and optical properties derived from PHIPS measurements in mid-latitude (ML) and Arctic cirrus clouds. Shown are total number concentration (N), ice water content (IWC), effective radius (r_{eff}), and asymmetry parameter (g). The analysis includes particles in the PHIPS measurement range of 15 to 700 μm .

Variable	Statistic	ML	Arctic
$N \text{ [L}^{-1}\text{]}^1$	Median	10.9	8.2
	Mean	120.1	8.1
	25th percentile	3.0	2.5
	75th percentile	45.4	25.6
	5th percentile	0.9	0.8
	95th percentile	187.3	83.1
$IWC \text{ [mg m}^{-3}\text{]}^1$	Median	2.9	3.0
	Mean	2.6	2.4
	25th percentile	0.7	1.1
	75th percentile	8.8	6.1
	5th percentile	0.13	0.19
	95th percentile	35.1	13.9
$\beta_{\text{ext}} \text{ [km}^{-1}\text{]}^1$	Median	0.13	0.13
	Mean	0.13	0.11
	25th percentile	0.040	0.051
	75th percentile	0.39	0.26
	5th percentile	0.0087	0.011
	95th percentile	1.5	0.62
$r_{\text{eff}} \text{ [\mu m]}$	Median	35.9	37.4
	Mean	36.1	38.2
	25th percentile	30.9	33.1
	75th percentile	41.1	42.5
	5th percentile	23.8	27.8
	95th percentile	49.2	51.2
g	Median	0.737	0.738
	Mean	0.736	0.735
	25th percentile	0.723	0.724
	75th percentile	0.749	0.746
	5th percentile	0.702	0.705
	95th percentile	0.772	0.759

¹Lower detection limits for 10-second averages are approximately 0.5 L^{-1} for N , 0.05 mg m^{-3} for IWC and 0.005 km^{-1} for β_{ext} .

The mid-latitude cases covered a broader range of cirrus conditions, while Arctic sampling was limited to five predominantly liquid-origin cirrus systems north of 65°N. Due to pandemic-related base relocation to Germany, Arctic flights required long transits and thus targeted synoptic cirrus with predictable development: a warm frontal cirrus (F05), two liquid-origin systems (F08/F09), a jet-stream-associated liquid-origin cirrus (F10/F11), and a mix of liquid-origin and in-situ cirrus over Iceland (F13/F14, Fig. 1). This focus introduces a bias toward liquid-origin cirrus, which have been shown to exhibit higher *IWC* compared to in situ-formed cirrus (Luebke et al., 2016). This sampling bias should be taken into account when interpreting observed latitudinal trends in cirrus microphysical and optical properties.

To better account for thermodynamic influences on cirrus properties, it is useful to examine the observations as a function of in-cloud temperature rather than geographic location. Figure 6 shows the statistical distribution of N , *IWC*, β_{ext} , r_{eff} and g , comparing mid-latitude (red) and Arctic (blue) observations. Median N was observed to be higher in the mid-latitudes for temperatures down to approximately −55°C. At colder temperatures, Arctic values become comparable to, or even exceed, those in the mid-latitudes. However, the observed differences are not statistically significant at the coldest temperatures below −55°C and between −44°C and −48°C (two-sample t-test, $p > 0.05$).

Both *IWC* and β_{ext} values were generally similar in both regions at temperatures above −55°C and showed a decreasing trend with decreasing temperature. At colder temperatures values observed in Arctic cirrus exceeded those observed in mid-latitude, however, due to the limited number of observations this difference is not statistically significant ($p > 0.05$). The decreasing trend in *IWC* with decreasing temperature is well documented also in previous studies (Krämer et al., 2016; Jeggle et al., 2025).

The r_{eff} was mostly larger in the Arctic compared to the mid-latitudes with median values exceeding those in the mid-latitudes by 5 μm and showing a decreasing trend with temperature for temperatures down to −55°C. At temperatures colder than −55°C, the r_{eff} of Arctic cirrus increases again, showing up to twice as large r_{eff} . The observed differences were found to be statistically significant in the temperature ranges between −38°C and −40°C, and between −44°C and −52°C.

The asymmetry parameter showed a slight increasing trend towards colder temperatures with values around 0.72 measured at −40°C and values between 0.75 and 0.76 at −60°C. Somewhat lower values were observed in Arctic cirrus at temperatures below −55°C, but as with the other parameters, the observed differences at colder temperatures are not statistically significant.

Inspection of Table 1 shows that the microphysical properties derived from the PHIPS instrument are in general agreement with previously reported values in the mid-latitudes and Arctic (Luebke et al., 2016; Heymsfield et al., 2017; Wolf et al., 2018). The median N values are systematically higher than those reported in the literature based on optical array probes with larger sampling volumes (Luebke et al., 2016; Heymsfield et al., 2017; De La Torre Castro et al., 2023). This discrepancy arises because low concentrations, below approximately 0.5 L^{-1} , are not detected with PHIPS within the 10-second averaging period. Values of N and *IWC* obtained from optical array probe measurements during the campaign are discussed in detail by De La Torre Castro et al. (2023) and Jurkat-Witschas et al. (2025). Furthermore, the observed r_{eff} in both regions are significantly lower than the values reported by De La Torre Castro et al. (2023) for the same study. This discrepancy is most likely due to their assumption of a simplified spherical ice particle shape to derive this quantity from the measured particle size distribution.




Figure 6. Statistical analysis of the observed total number concentration (N), ice water content (IWC), extinction coefficient (β_{ext}), effective radius (r_{eff}), and asymmetry parameter (g) as a function of in-cloud temperature in degree Celsius. The solid line shows the median values and the shaded area the interquartile range for mid-latitude (ML; red) and Arctic (blue) observations using a latitude threshold of 65° .

3.3 Sensitivity of g to ice microphysical and environmental factors

To examine how g relates to environmental and microphysical properties, we analysed simultaneous measurements of in-cloud temperature, relative humidity over ice (RH_i), and key microphysical parameters. All variables were averaged over the same time intervals used for the g retrievals, each corresponding to 50 consecutive intact ice crystals (averaging times of 44 s–4 min, 25th–75th percentile, or 3–20 L sampled volume). The analysed microphysical quantities include D_{eq} , aspect ratio, IWC , r_{eff} , and N , all derived from the manually cleaned PHIPS data. In addition, we included the mass-equivalent spherical radius ($r_{\bar{m}}$) defined by Baran et al. (2025). This parameter represents the radius of a sphere with the mean mass of the ice crystal population and has been proposed as a more physically consistent quantity for linking two-moment microphysical schemes with optical properties. In contrast to the effective radius r_{eff} , $r_{\bar{m}}$ does not require assumptions about the ice particle surface area.

Table 2 summarises the Pearson correlation coefficients between g and the investigated parameters. Among the tested parameters, the strongest correlation was observed with D_{eq} ($r = -0.42$), followed by $r_{\bar{m}}$ ($r = -0.34$) and r_{eff} ($r = -0.30$), all indicating moderate negative correlations. These results suggest that g decreases with increasing particle size. Only weak correlations were found with temperature, RH_i , AR, and N .

Figure 7 shows the frequency distribution of the measured g as a function of r_{eff} for ML and Arctic cirrus cases. The same figure for $r_{\bar{m}}$ is shown in Supplementary Information (Fig. S15). It can be seen that g values vary between 0.65 and 0.79. In the ML dataset, the median g decreases from 0.74 to 0.72 with increasing r_{eff} up to approximately $r_{eff}=40 \mu\text{m}$, beyond which the trend reverses and the median g increases with particle size. However, this upward trend at larger sizes is not statistically robust due to the limited number of observations in that range. In Arctic cirrus, the data are more sparse overall, but a consistent decrease in the median g from 0.74 to 0.70 with increasing r_{eff} is observed across the measured range.

In radiative transfer simulations, ice optical properties are frequently parameterised as a function of r_{eff} . Figure 7 shows four commonly used parameterisations of g at a wavelength band centered at 532 nm, corresponding to the PHIPS light scattering measurements. The first parameterisation, developed by Fu (1996), is based on columnar crystals with single scattering properties calculated using an improved geometric optics method. The asymmetry parameter is expressed as a function of a

Table 2. Pearson correlation coefficients (r) and p-values for the relationship between g and various atmospheric and microphysical parameters. Moderate correlations are highlighted.

Variable	r	p-value
Temperature	-0.21	0.000
RH_i	-0.07	0.146
Aspect ratio	-0.11	0.019
D_{eq}	-0.42	0.000
$\log(IWC)$	-0.08	0.081
r_{eff}	-0.30	0.000
$r_{\bar{m}}$	-0.34	0.000
N	-0.10	0.022




Figure 7. Frequency distribution of asymmetry parameter (g) observations as a function of effective radius (r_{eff}). Both the asymmetry parameter and effective radius are retrieved only for particles with stereo-microscopic image that were not classified as shattering. The black line indicated the median value for the effective radius bins with a bin width of $3 \mu\text{m}$. Existing short wave optical parameterisations are overlaid to the measurement data. These are the parameterisation by Fu (1996) (Fu; blue line), general habit mixture (GHM, dashed black line; Cole et al., 2013), the severely roughened column aggregate model used in MODIS Collection 6 retrievals (MC6; Platnick et al., 2016) and the optical parameterisation used in ECHAM6 general circulation model (ECHAM; Pincus and Stevens, 2013).

generalised effective size, which can be related to r_{eff} via Eq. 3.12 in Fu (1996). The second parameterisation represents the
335 General Habit Mixture (GHM), which assumes a combination of nine crystal habits with severe surface roughness ($\sigma = 0.5$)
(Cole et al., 2013). The third is the MODIS Collection 6 (MC6) parameterisation, which assumes aggregates composed of

severely roughened columns (Platnick et al., 2016). The fourth is the parameterisation implemented in the ECHAM6 general circulation model (Pincus and Stevens, 2013).

Among the four, only the MC6 parameterisation—assuming complex, severely roughened aggregates with a fixed asymmetry parameter of 0.75—captures the upper range of the observed values, but does not capture the median values or the lower range. The remaining parameterisations overestimate g across the entire size range and fail to reproduce majority of the measured values. Moreover, all four parameterisations predict an increasing trend in the g with increasing r_{eff} , which contrasts with the observational data showing a clear decreasing trend.

3.3.1 Analysis based on individual particle classes

A key advantage of PHIPS as a single particle polar nephelometer is its ability to directly link the microphysical and optical properties of individual ice crystals. This allows for the selection of particles with specific microphysical characteristics, such as size or aspect ratio, and enables the construction of “artificial” particle populations consisting of crystals with similar properties. This approach results in a very detailed investigation of how individual microphysical parameters influence g , isolated from the natural microphysical variability present in real cloud segments. This approach was recently used to retrieve the asymmetry parameter of bullet rosettes from PHIPS data captured during CIRrus-HL (Wagner et al., 2024).

Figure 8 presents a statistical analysis of g for all ice crystals measured during CIRrus-HL that were manually classified as intact. The ice crystals were grouped based on their D_{eq} (for bin definitions, see Waitz et al., 2021), AR (defined as the smaller of the two aspect ratios from the stereo images) using a bin width of 0.1, and simultaneously measured ice supersaturation (S_{ice}), also using a bin width of 0.1. Each bin contains a minimum of 100 ice crystals. To retrieve g , the crystals within each bin were further divided into groups of 50 consecutive particles, and one g was calculated per group. The number of these groups is indicated in the figure. Additionally, a single g was retrieved for all crystals within each bin, with the corresponding value and uncertainty shown as markers.

Of all the variables, only D_{eq} shows a clear decreasing trend with g . This is consistent with the results from the previous analysis, where cloud volumes were analysed instead of particle classes. The highest median values ($g = 0.776$), were measured for mid-latitude ice crystals with D_{eq} between 15 and 30 μm , while the lowest median values ($g = 0.718$) were observed for D_{eq} between 150 and 200 μm . For larger particles, the g increases again, although this trend is not statistically robust due to fewer observations.

Figure 9 further examines this size dependence of g by separating the data according to ice crystal habit. The analysis was conducted for the seven most frequently occurring habit categories: aggregate, column, side plane, frozen droplet, bullet rosette, plate, and the “other” category. For each habit–size combination, a median g was retrieved for each size bin, provided that at least 50 intact crystals were available in the bin.

The general size dependence of g observed in Figure 8 is seen across most habits, except for frozen droplets that only cover the two smallest size bins. A maximum difference of 0.05 in g is found between columns, which show the lowest values, and aggregates, which exhibit the highest values. This difference exceeds the measurement uncertainty and is therefore considered to be significant. Side planes, bullet rosettes, plates, and crystals in the “other” category all show similar g values within the




Figure 8. Statistical analysis of the asymmetry parameter (g) for mid-latitude (ML) and Arctic cirrus as a function of (a) particle area equivalent diameter (D_{eq}), (b) aspect ratio, and (c) ice supersaturation (S_{ice}). The box indicates the 25th and 75th percentiles (quartiles), the line the 50th percentile (median) and the whiskers the 5th and 95th percentiles of the g values retrieved for samples of 50 crystals. The markers indicate the asymmetry parameter retrieval for all crystals within the bin with error bars denoting the retrieval uncertainty. Sample sizes corresponding the number of asymmetry parameter retrievals for groups of 50 particles for each category are annotated in the plots.

uncertainty range for sizes below 200 μm . For larger bullet rosettes, g increases and becomes comparable to that of aggregates of similar size.




Figure 9. Mean asymmetry parameter, g , as a function of particle diameter for various ice crystal habits, including aggregates, plates, columns, frozen droplets, side planes, and bullet rosettes. The shaded areas represent the retrieval uncertainty in asymmetry parameter for each habit.

No significant difference was found in the median g between mid-latitude and Arctic cirrus with median g between 0.724 and 0.758 observed for each AR and S_{ice} category. In general, the variability in g was greater in the mid-latitudes across all categories (g varied between 0.686 and 0.815 for groups of 50 crystals) compared to the Arctic (g varied between 0.690 and 0.787). However, this may be partly attributed to the smaller sample size in the Arctic.

4 Discussion

Despite differences in geographic location and the ice crystal habit distributions, the g values showed no significant variation between mid-latitude and Arctic cirrus but consistently low median values were observed. Low g values below 0.8 have been systematically measured also by other airborne polar nephelometers for the visible spectral range. Measurements with the CIN instrument have shown g values around 0.74 ± 0.03 for Arctic ice clouds (Gerber et al., 2000; Garrett et al., 2001) and values around 0.75 ± 0.01 for tropical anvil cirrus (Garrett et al., 2003). In mid-latitudes, median g values between 0.769 and 0.79 have been measured using the PN instrument (Gayet et al., 2004; Febvre et al., 2009; Auriol et al., 2001; Gayet et al., 2012).

Our results show that g is largely insensitive to variations in atmospheric conditions (temperature and relative humidity), as well as to most microphysical properties of ice crystals. This finding is consistent with several previous observational studies. For example, Febvre et al. (2009) reported similar optical properties for natural cirrus and aged contrails, despite differences

in their size distributions. Likewise, Garrett et al. (2001) found that g values were insensitive to habit variations in Arctic cirrus. During the INCA experiment, similar g values were derived for the northern and southern hemispheres (Gayet et al., 2004, 2006), with only a weak temperature dependence noted by Gayet et al. (2006).

390 Here, the only parameter for which a moderate correlation with g was observed is particle size: we find that g values tend to decrease with increasing D_{eq} , r_{eff} or $r_{\bar{m}}$. This behaviour differs from the results of Gayet et al. (2004), who reported an increase in g with effective diameter, whereas Garrett et al. (2003) observed no clear dependence between g and r_{eff} for small anvil ice crystals with $r_{eff} < 20 \mu\text{m}$. Interestingly, the aspect ratio does not appear to influence g in our data, which stands in contradiction to theoretical studies predicting a strong dependence between g and AR , even for roughened crystals (Iaquinta
395 et al., 1995; van Diedenhoven et al., 2014).

The low median g value of 0.737 cannot be explained by pristine hexagonal shapes (Iaquinta et al., 1995; Schmitt et al., 2006). Rather, a significant degree of surface roughness or other deformations are needed in order to numerically reproduce our observations (Macke et al., 1996a; Yang et al., 2013; Wagner et al., 2024). Given that the 95th percentiles of the g distributions are 0.772 for mid-latitude and 0.759 for Arctic cirrus, we conclude that the majority of the measured ice crystals exhibit a
400 significant degree of morphological complexity. These findings suggest that crystal complexity plays a more dominant role in determining g than any other microphysical property. The observed size dependence of g may also reflect an increasing degree of crystal complexity in larger ice particles.

In-situ measurements have indicated g values above 0.8 only for a few cases. Garrett et al. (2001) reported g values of 0.81 for sublimating ice crystals at RH_i around 87%, and Gayet et al. (2012) found similar g values for RH_i below 80%. Febvre
405 et al. (2009) observed g values of 0.827 for young contrails, with a mean RH_i of 118%. In our dataset, we did not observe an increase in g at lower RH_i values. However, our observations included only a limited number of ice crystals at RH_i below 80%. Therefore, the effect of sublimation on g remains inconclusive and needs further investigation.

A limitation of this study is that g retrievals were restricted to ice crystals with D_{eq} below $300 \mu\text{m}$. This constraint arises because larger particles frequently saturated the first few measurement channels of our polar nephelometer, preventing a reliable
410 retrieval of g . However, as shown in the particle size and area distributions in Figure 2 and the cumulative distributions in Figure S1, ice crystals smaller than $300 \mu\text{m}$ account for 97% (99%) of the total number and 83% (94%) of the total cross-sectional area in mid-latitude (Arctic) cirrus. Based on this, we consider our retrieved g values to be representative of the cirrus cloud populations sampled during the CIRRUS-HL campaign.

Our results show a substantial discrepancy between the parameterisations of g used in climate models and the values mea-
415 sured in this study (see Fig. 7). This difference has significant implications for inferred cloud reflectivity or albedo, which can be demonstrated using a simple approximation of the plane albedo for thin cirrus clouds (Meador and Weaver, 1980)

$$R = \frac{\omega_0}{2\mu_0} (1 - g) \tau, \quad (8)$$

where ω_0 is the single-scattering albedo, μ_0 is the cosine of the solar zenith angle, and τ is the optical thickness. For constant optical thickness and solar angle, the plane albedo is directly proportional to $1 - g$. Therefore, a decrease of g from 0.8 to
420 the measured 0.737 corresponds to an increase in albedo of approximately 32%, making cirrus clouds more reflective at solar

wavelengths. This indicates that an overestimation of g can lead to a significant bias in the estimation of the radiative effect of cirrus clouds. Additionally, such an overestimation can introduce errors in the retrieval of cirrus optical depth from space-borne radiance measurements (Yang et al., 2008; Yi et al., 2017a).

Studies using general circulation models (GCMs) or Moderate Resolution Imaging Spectroradiometer (MODIS) retrievals to simulate top-of-the-atmosphere (TOA) fluxes have shown that incorporating optical parameterisations with lower g values in the visible spectrum leads to a closer agreement between these TOA flux simulations and TOA flux estimates from direct measurements of radiances using space-borne remote sensing (Kristjánsson et al., 2000; Yi et al., 2017b). However, Ren et al. (2023) found that optical parameterisations with low g values around and below 0.75 overestimated both shortwave (SW) and longwave (LW) fluxes at TOA compared to observations from the Clouds and the Earth's Radiant Energy System (CERES). Their study emphasised the need for further optimisation of the scattering phase function to achieve radiative closure. Yi et al. (2017b) also showed that lowering g (by changing the optical parameterisation from MC5 to MC6) significantly reduced the estimated TOA SW flux from MODIS observations. This reduction resulted from lower cloud optical thickness retrieved by the MC6 model, which was subsequently used as input to the radiative transfer simulations (Yi et al., 2017a).

Currently, cirrus and ice containing ice clouds have been estimated to exert a globally and annually averaged net warming effect of $2\text{--}5 \text{ Wm}^{-2}$ (Hong et al., 2016; Yi et al., 2017b; L'Ecuyer et al., 2019). Modelling studies using GCMs have shown that lowering the g values in visible from values around 0.8 to 0.75 will decrease the net ice CRE by $1\text{--}2 \text{ Wm}^{-2}$ (Järvinen et al., 2018; Yi et al., 2013; Yi, 2022). Our results suggest that an additional decrease of the g to values below 0.75 might be needed, which would lead to an even higher bias in CRE.

A potential bias in CRE estimates is highly significant in the context of the Earth's energy balance. The current estimate of the global radiative imbalance is on the order of 0.5 to 1 Wm^{-2} (Loeb et al., 2021), meaning that even small biases in cirrus CRE can mask or amplify the signals of anthropogenic forcing. Cirrus clouds cover up to 30% of the globe at any given time (Stubenrauch et al., 2013), and errors in their radiative properties can accumulate spatially and temporally, introducing systematic uncertainty into global climate models. For instance, an overestimation of the asymmetry parameter g leads to an underestimation of cirrus reflectivity, allowing more solar radiation to reach the surface. In global climate models, this can trigger dynamic feedbacks: increased surface heating may destabilize the atmosphere, enhance convection, and promote the formation of low-level clouds. In parallel, GCMs are often tuned to maintain top-of-atmosphere energy closure (Mauritsen et al., 2012). If cirrus reflectivity is systematically too low, the tuning can be made at the expense of physical realism. Improving the representation of cirrus optical properties, particularly g , is therefore essential for accurately quantifying cloud–radiation interactions and reducing uncertainty in climate projections.

5 Summary

We analysed ice crystal microphysical and correlated angular light scattering properties of mid-latitude and Arctic cirrus clouds using in situ measurements performed with the PHIPS airborne cloud probe during the CIRRUS-HL campaign. Ice crystal number concentration and ice water content showed no clear latitudinal trends, with similar median values of 11 L^{-1}

and 2.8 mg m^{-3} for mid-latitude cirrus and 8 L^{-1} and 3.0 mg m^{-3} for Arctic cirrus. In contrast, the effective radius increased
455 toward higher latitudes. Furthermore, clear regional differences in ice crystal habit distributions were observed: aggregates
were the dominant habit in mid-latitude cirrus for crystals larger than $200 \mu\text{m}$, while bullet and mixed rosettes dominated Arctic
cirrus. Smaller crystals in both regions were primarily complex in shape and unclassifiable using standard habit categories.

Despite the differences in habits, both mid-latitude and Arctic cirrus exhibited nearly identical asymmetry parameter values,
with the same median asymmetry parameter of 0.737 for both regions and observed values as low as 0.686 (1th percentile). The
460 asymmetry parameter was largely insensitive to environmental conditions such as temperature and relative humidity, as well as
to ice crystal aspect ratio and bulk microphysical parameters including IWC and number concentration. Moderate correlations
were found only with particle size.

The relatively low asymmetry parameter values cannot be represented using optical models for idealised pristine crystals.
Comparison with four commonly used radiative transfer parameterisations revealed that most overestimated asymmetry pa-
465 rameter by up to 0.1. Only the MODIS Collection 6 parameterisation, which assumes severely roughened columnar crystals,
captured the upper range of the observed values. Given the inverse relationship between the asymmetry parameter and cloud
reflectivity, overestimating asymmetry parameter from 0.737 to 0.800 can lead to an underestimation of cirrus albedo by up to
32%, with implications for both radiative forcing estimates and satellite retrievals of optical depth. Such biases are particularly
important because cirrus clouds have estimated to exert a net warming effect of $2\text{--}5 \text{ W m}^{-2}$ globally, and reducing asymmetry
470 parameter in climate models by just 0.05 has been shown to lower this warming by $1\text{--}2 \text{ W m}^{-2}$. These discrepancies highlight
the need for improved observational constraints and parameterisations of asymmetry parameter to better quantify the radiative
role of cirrus clouds and reduce uncertainties in climate projections.

Data availability. Processed PHIPS microphysical data and single-particle light scattering data will be available in the HALO database
(<https://halo-db.pa.op.dlr.de/mission/125>) upon publication. The stereo-microscopic images are available from the authors upon request.

475 **Appendix A: Accuracy and Assumptions of the Asymmetry Parameter Retrieval**

A1 Overview of the retrieval framework

Representing the scattering phase function as a series of Legendre polynomials is a common approach in atmospheric sciences
and is widely used in radiative-transfer and remote-sensing applications (Wiscombe, 1977; Hu et al., 2000). This formulation
is convenient because the Legendre polynomials are harmonic functions of $\cos\theta$ and, thus, provide an efficient mathematical
480 basis for describing the angular dependence of scattered light. As a general rule, the number of Legendre terms, N , required
to accurately represent a phase function increases with the particle size parameter, x . For example, in case of large ice crystals
($x > 100$) the narrow forward peak requires a high number of expansion terms, up to several thousands, to resolve the original
phase function.

To reduce some of this complexity, the first step in our retrieval framework is to separate the diffraction contribution from the geometrical-optics (GO) component of scattering. Although this decomposition is not fundamental from the electromagnetic perspective, it is a well-established and practical approximation for large atmospheric ice particles and widely used in conventional GO methods (e.g., Takano and Liou, 1989; Macke et al., 1996b). Studies using improved GO methods that avoid this artificial separation have shown convergence toward conventional ray-tracing results for the large size parameters relevant to our study (Yang and Liou, 1996).

This separation is also convenient because the diffraction contribution within the PHIPS measurement range (18° – 170°) can be considered negligible. This assumption is supported by theoretical work showing that for large ice particles, diffraction intensity decreases by more than three orders of magnitude beyond 18° (e.g., Hesse, 2008; Chunlei and Keya, 1996; Travis et al., 2002). In our retrieval framework, the diffraction component is calculated using scalar diffraction theory for circular apertures. For particles within the PHIPS size range, the resulting diffraction asymmetry parameter g_d varies between 0.995 and 1.0 (Xu et al., 2022). Because g_d is so close to unity, the influence of particle shape on g_d after orientation averaging can be considered negligible. This assumption was verified by comparing our circular-aperture parameterisation with that of van Diedenhoven et al. (2014), which assumes hexagonal particles with unity aspect ratio. The resulting difference corresponds to an uncertainty of approximately 0.001 in the total asymmetry parameter.

Consequently, the measured PHIPS angular range can be treated as purely GO scattering. To obtain the GO contribution to g the measurement needs to be extrapolated to the missing forward and backscattering ranges. This is done by fitting a truncated Legendre polynomial expansion to the measured angular intensities. The GO scattering phase function $P_{GO}(\theta)$ is expressed as a finite series of Legendre polynomials $P_l(\cos\theta)$:

$$P_{GO}(\theta) \approx \sum_{l=0}^N (2l+1) \hat{c}_l P_l(\cos\theta), \quad (\text{A1})$$

where $\hat{c}_l$ are the Legendre expansion coefficients of degree l , and N is the maximum degree of the expansion. The factor $2l+1$ follows the standard normalization of Legendre polynomials over the interval $[-1, 1]$.

The coefficients $\hat{c}_l$ are obtained through orthogonal projection of the measured phase function onto the Legendre basis. In practice, this integration is carried out numerically using Gauss–Legendre quadrature, which provides optimal accuracy by evaluating the integrand at specific quadrature nodes x_i corresponding to the roots of $P_{N+1}(x_i)$:

$$\hat{c}_l = \frac{1}{2} \int_{-1}^1 P_{GO}(\arccos(x)) P_l(x) dx \approx \frac{1}{2} \sum_{i=1}^N w_i I(\arccos(x_i)) P_l(x_i), \quad (\text{A2})$$

where w_i are the Gauss–Legendre weights associated with each node and $I(\arccos(x_i))$ the measured intensity interpolated to the Gauss–Legendre quadrature nodes x_i .

In our retrieval, the number of quadrature nodes is set to eight, corresponding to a Legendre expansion of eight terms ($N = 7$). This truncation ensures that all quadrature nodes remain within the PHIPS angular coverage (18° – 170°) and avoids extrapolation into unmeasured regions.

515 Using eight terms implies that the fitted Legendre expansion can accurately represent only smooth angular scattering functions. Therefore, the retrieval method is not applicable to pristine ice crystals that exhibit sharp features in their phase functions. A preliminary uncertainty analysis of this approach was presented by Xu et al. (2022), where the authors compared the retrievals results to true g values using simulations of hexagonal ice crystals with varying degree of surface roughness and aspect ratio. However, the uncertainty analysis was based on a limited number of crystal geometries. In the present work, we extend this
520 analysis using a substantially larger set of simulated ice crystal morphologies to refine the uncertainty estimate and confirm the validity of the truncation choice.

A2 Updated uncertainty analysis

To extend the analysis of Xu et al. (2022), we performed additional closure tests using the ray-tracing model of Macke et al. (1996b). The new simulations include 1612 orientation-averaged configurations of hexagonal ice crystals with sizes between
525 20 and 500 μm and aspect ratios ranging from 0.68 to 0.85, motivated by the PHIPS microphysical observations. Surface roughness was implemented using Gaussian tilted-facet perturbations with roughness parameters (σ) between 0 and 0.9 (interval of 0.1). In addition, internal air inclusions were simulated using the mean free path (MFP) approach (Macke et al., 1996a), with MFP values between 10 and 10^8 (no inclusions).

The purpose of this investigation was to evaluate how the retrieval error in g depends on the smoothness of the scattering
530 phase function. To achieve this, the models were grouped according to their roughness parameter σ , which is directly related to the smoothness of the calculated scattering function (higher σ corresponds to smoother phase functions). Since natural cirrus clouds exhibit a diversity of shapes and aspect ratios, the simulations were averaged over ten different crystal configurations before calculating g . Although this ensemble size represents only a moderate level of morphological variability, sensitivity tests (not shown) confirmed that increasing the number of configurations had no effect on the results.

535 For each case, two values of g were computed: (i) the “true” value derived directly from the full simulated phase functions and (ii) the “retrieved” value obtained from the “true” phase function truncated to the PHIPS angular range (18° – 170°), corresponding to the angular coverage of the parabolic mirror detectors used in the PHIPS nephelometer. The truncation was carried out by integrating the simulated scattering intensity over the solid angle of each detector. Each detector covers an angular range around its central viewing angle with a half-angle acceptance of 3.5° , corresponding to a solid angle of 0.0117 sr. The simulated
540 phase functions, computed with a 1° angular resolution, were integrated over each detector’s acceptance using trapezoidal numerical integration.

Figure A1 illustrates the validation results for simulations with a low roughness parameter ($\sigma = 0.1$). The simulated phase functions still contain a modest halo feature that is clearly visible even in the truncated phase function at the PHIPS angular resolution. It should be noted that the PHIPS measurements in Panel (c) do not show a halo peak, even in the 1-99 percentile
545 range, suggesting that higher roughness parameters are needed to model the measured phase functions. Panel (b) shows that




Figure A1. Validation of the asymmetry parameter retrieval for simulations with a roughness parameter $\sigma = 0.1$. The dataset includes 180 orientation-averaged crystal configurations, with g values retrieved for sets of 10 orientations each. Panel (a) compares the true complexity parameter $C_{p,true}$ with the retrieved C_p , with the dashed line indicating the 1:1 relation. Panel (b) shows a heatmap of the retrieval error $g_{true} - g_{retrieved}$ as a function of C_p , with red dashed lines marking ± 0.001 error bounds. Panel (c) displays measured scattering phase functions for mid-latitude (blue) and Arctic (red) cirrus, including median, 25–75%, and 1–99% percentiles, with gray curves representing ray-tracing simulations of individual crystal configurations truncated to PHIPS measurement range.

the difference between the “true” and “retrieved” g can reach up to 0.08 but decreases rapidly with increasing complexity parameter C_p . The C_p parameter quantifies the decay rate of the Legendre expansion coefficients (Xu et al., 2022), with larger values corresponding to more featureless phase functions. While Xu et al. (2022) previously suggested that the retrieval is accurate within an absolute error of 0.001 for $C_p \geq 0.4$, our extended simulations reveal that for relatively pristine particles (550 $\sigma = 0.1$), the retrieved C_p values derived from the truncated phase functions can exceed 0.4 (as shown in Panel (a)), making the C_p parameter an unreliable quality indicator for assessing retrieval accuracy for lightly roughened crystals.

Figure A2 presents the corresponding results for $\sigma = 0.2$. Increasing the surface roughness from 0.1 to 0.2 smooths out the halo feature in the simulations, yielding featureless phase functions that more closely resemble the measured data (Fig. A2c). The retrieval error in g is now below 0.02, although the retrieved values tend to be slightly biased toward lower g . Further




Figure A2. Same as Figure A1, but for simulations with a roughness parameter $\sigma = 0.2$.

555 increasing the roughness to $\sigma = 0.3$ or higher values (not shown) reduces the retrieval error below 0.005 and also changes the bias sign, resulting in a slight overestimation of the true g .

These extended simulations confirm that the retrieval algorithm applied to the PHIPS angular range (18° – 170°) is reliable only for sufficiently smooth phase functions or sufficiently roughened ice crystals (in our simulations, $\sigma \geq 0.2$). We found that relatively pristine particles ($\sigma = 0.1$) yield C_p values above the previously proposed C_p threshold of 0.4 (Xu et al., 2022),
 560 indicating that C_p cannot be used as a criterion for assessing retrieval accuracy. However, in all cases where the retrieval error exceeded 0.02, the simulated phase functions exhibited distinct halo features that would be clearly detectable at the angular resolution of PHIPS. Since such features are entirely absent from the measured phase functions (see panel c in Figs. A1 and A2), we conclude that the natural ice crystals in our dataset exhibit sufficient surface roughness for the retrieval method to remain valid. Based on these results, we adopt a more conservative uncertainty estimate of -0.001 to $+0.02$ for asymmetry
 565 parameter retrievals derived from PHIPS measurements compared to the value given in Xu et al. (2022).

Author contributions. EJ conceptualised the study. MS designed the measurement concept of PHIPS and led the instrument development. EJ and MS collected the PHIPS data during the CIRRUS-HL campaign. EJ performed the data analysis and wrote the manuscript. MS contributed to the interpretation of the results and provided comments on the manuscript.

Competing interests. Martin Schnaiter and Emma Järvinen are affiliated with schnaiTEC GmbH, the manufacturer of the PHIPS instrument.
570 Martin Schnaiter is additionally employed part-time by schnaiTEC GmbH.

Acknowledgements. The authors would like to thank the HALO crew for supporting the operations during the CIRRUS-HL campaign and the science and weather forecast teams for flight planning. This work was funded by the Helmholtz Association's Initiative and Networking Fund (grant agreement no. VH-NG-1531).

References

- 575 Abdelmonem, A., Järvinen, E., Duft, D., Hirst, E., Vogt, S., Leisner, T., and Schnaiter, M.: PHIPS–HALO: the airborne Particle Habit Imaging and Polar Scattering probe – Part 1: Design and operation, *Atmos. Meas. Tech.*, 9, 3131–3144, 2016.
- Auriol, F., Gayet, J.-F., Febvre, G., Jourdan, O., Labonnote, L., and Brogniez, G.: In Situ Observation of Cirrus Scattering Phase Functions with 22° and 46° Halos: Cloud Field Study on 19 February 1998, *Journal of the Atmospheric Sciences*, 58, 3376 – 3390, [https://doi.org/10.1175/1520-0469\(2001\)058<3376:ISOOCS>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<3376:ISOOCS>2.0.CO;2), 2001.
- 580 Baker, B. and Lawson, R. P.: Improvement in Determination of Ice Water Content from Two-Dimensional Particle Imagery. Part I: Image-to-Mass Relationships, *Journal of Applied Meteorology and Climatology*, 45, 1282 – 1290, <https://doi.org/10.1175/JAM2398.1>, 2006.
- Barahona, D., Molod, A., and Kalesse, H.: Direct estimation of the global distribution of vertical velocity within cirrus clouds, *Scientific Reports*, 7, 6840, 2017.
- Baran, A. J., Manners, J., Field, P. R., Furtado, K., and Hill, A.: A consistent coupling of two-moment microphysics and bulk ice optical properties, and its impact on radiation in a regional weather model, *Quarterly Journal of the Royal Meteorological Society*, n/a, e5025, <https://doi.org/https://doi.org/10.1002/qj.5025>, 2025.
- 585 Bohren, C. F. and Huffman, D. R.: *Absorption and Scattering of Light by Small Particles*, Wiley-VCH, Weinheim, Germany, ISBN 978-0-471-29340-8, 1998.
- Chunlei, L. and Keya, Y.: Calculation of ice crystal diffraction, *Advances in Atmospheric Sciences*, 13, 340–348, <https://doi.org/10.1007/BF02656851>, 1996.
- 590 Cole, B. H., Yang, P., Baum, B. A., Riedi, J., C.-Labonnote, L., Thieuleux, F., and Platnick, S.: Comparison of PARASOL Observations with Polarized Reflectances Simulated Using Different Ice Habit Mixtures, *Journal of Applied Meteorology and Climatology*, 52, 186 – 196, <https://doi.org/10.1175/JAMC-D-12-097.1>, 2013.
- Cole, B. H., Yang, P., Baum, B. A., Riedi, J., and C.-Labonnote, L.: Ice particle habit and surface roughness derived from PARASOL polarization measurements, *Atmospheric Chemistry and Physics*, 14, 3739–3750, <https://doi.org/10.5194/acp-14-3739-2014>, 2014.
- 595 De La Torre Castro, E., Jurkat-Witschas, T., Afchine, A., Grewe, V., Hahn, V., Kirschler, S., Krämer, M., Lucke, J., Spelten, N., Wernli, H., Zöger, M., and Voigt, C.: Differences in microphysical properties of cirrus at high and mid-latitudes, *Atmospheric Chemistry and Physics*, 23, 13 167–13 189, <https://doi.org/10.5194/acp-23-13167-2023>, 2023.
- Febvre, G., Gayet, J.-F., Minikin, A., Schlager, H., Shcherbakov, V., Jourdan, O., Busen, R., Fiebig, M., Kärcher, B., and Schumann, U.: On optical and microphysical characteristics of contrails and cirrus, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/https://doi.org/10.1029/2008JD010184>, 2009.
- 600 Forster, L. and Mayer, B.: Ice crystal characterization in cirrus clouds III: retrieval of ice crystal shape and roughness from observations of halo displays, *Atmospheric Chemistry and Physics*, 22, 15 179–15 205, <https://doi.org/10.5194/acp-22-15179-2022>, 2022.
- Fu, Q.: An Accurate Parameterization of the Solar Radiative Properties of Cirrus Clouds for Climate Models, *Journal of Climate*, 9, 2058 – 2082, [https://doi.org/10.1175/1520-0442\(1996\)009<2058:AAPOTS>2.0.CO;2](https://doi.org/10.1175/1520-0442(1996)009<2058:AAPOTS>2.0.CO;2), 1996.
- 605 Garrett, T. J., Hobbs, P. V., and Gerber, H.: Shortwave, single-scattering properties of arctic ice clouds, *Journal of Geophysical Research: Atmospheres*, 106, 15 155–15 172, <https://doi.org/https://doi.org/10.1029/2000JD900195>, 2001.
- Garrett, T. J., Gerber, H., Baumgardner, D. G., Twohy, C. H., and Weinstock, E. M.: Small, highly reflective ice crystals in low-latitude cirrus, *Geophysical Research Letters*, 30, <https://doi.org/https://doi.org/10.1029/2003GL018153>, 2003.

- 610 Gayet, J.-F., Auriol, F., Oshchepkov, S., Schröder, F., Duroure, C., Febvre, G., Fournol, J.-F., Crépel, O., Personne, P., and Daugereon, D.: In situ measurements of the scattering phase function of stratocumulus, contrails and cirrus, *Geophysical Research Letters*, 25, 971–974, <https://doi.org/https://doi.org/10.1029/98GL00541>, 1998.
- Gayet, J.-F., Ovarlez, J., Shcherbakov, V., Ström, J., Schumann, U., Minikin, A., Auriol, F., Petzold, A., and Monier, M.: Cirrus cloud microphysical and optical properties at southern and northern midlatitudes during the INCA experiment, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/https://doi.org/10.1029/2004JD004803>, 2004.
- 615 Gayet, J.-F., Shcherbakov, V., Mannstein, H., Minikin, A., Schumann, U., Ström, J., Petzold, A., Ovarlez, J., and Immler, F.: Microphysical and optical properties of midlatitude cirrus clouds observed in the southern hemisphere during INCA, *Quarterly Journal of the Royal Meteorological Society*, 132, 2719–2748, <https://doi.org/https://doi.org/10.1256/qj.05.162>, 2006.
- Gayet, J.-F., Mioche, G., Bugliaro, L., Protat, A., Minikin, A., Wirth, M., Dörnbrack, A., Shcherbakov, V., Mayer, B., Garnier, A., and
- 620 Gourbeyre, C.: On the observation of unusual high concentration of small chain-like aggregate ice crystals and large ice water contents near the top of a deep convective cloud during the CIRCLE-2 experiment, *Atmospheric Chemistry and Physics*, 12, 727–744, <https://doi.org/10.5194/acp-12-727-2012>, 2012.
- Gerber, H., Takano, Y., Garrett, T. J., and Hobbs, P. V.: Nephelometer Measurements of the Asymmetry Parameter, Volume Extinction Coefficient, and Backscatter Ratio in Arctic Clouds, *Journal of the Atmospheric Sciences*, 57, 3021 – 3034, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0469(2000)057<3021:NMOTAP>2.0.CO;2)
- 625 [0469\(2000\)057<3021:NMOTAP>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<3021:NMOTAP>2.0.CO;2), 2000.
- Gil-Díaz, C., Sicard, M., Sourdeval, O., Saiprakash, A., Muñoz Porcar, C., Comerón, A., Rodríguez-Gómez, A., and Oliveira, D. C. F. D. S.: Study of optical scattering properties and direct radiative effects of high-altitude cirrus clouds in Barcelona, Spain, with 4 years of lidar measurements, *Atmospheric Chemistry and Physics*, 25, 3445–3464, <https://doi.org/10.5194/acp-25-3445-2025>, 2025.
- Hesse, E.: Modelling diffraction during ray tracing using the concept of energy flow lines, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 109, 1374–1383, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2007.11.002>, x Conference on Electromagnetic and Light Scattering by Non-Spherical Particles, 2008.
- 630 Heymsfield, A. J., Krämer, M., Luebke, A., Brown, P., Cziczó, D. J., Franklin, C., Lawson, P., Lohmann, U., McFarquhar, G., Ulanowski, Z., and Tricht, K. V.: Cirrus Clouds, *Meteorological Monographs*, 58, 2.1 – 2.26, <https://doi.org/10.1175/AMSMONOGRAPHIS-D-16-0010.1>, 2017.
- 635 Hong, Y., Liu, G., and Li, J.-L. F.: Assessing the Radiative Effects of Global Ice Clouds Based on CloudSat and CALIPSO Measurements, *Journal of Climate*, 29, 7651 – 7674, <https://doi.org/10.1175/JCLI-D-15-0799.1>, 2016.
- Hu, Y.-X., Wielicki, B., Lin, B., Gibson, G., Tsay, S.-C., Stamnes, K., and Wong, T.: δ -Fit: A fast and accurate treatment of particle scattering phase functions with weighted singular-value decomposition least-squares fitting, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 65, 681–690, [https://doi.org/https://doi.org/10.1016/S0022-4073\(99\)00147-8](https://doi.org/https://doi.org/10.1016/S0022-4073(99)00147-8), 2000.
- 640 Jaquinta, J., Isaka, H., and Personne, P.: Scattering Phase Function of Bullet Rosette Ice Crystals, *Journal of Atmospheric Sciences*, 52, 1401 – 1413, [https://doi.org/10.1175/1520-0469\(1995\)052<1401:SPFOBR>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<1401:SPFOBR>2.0.CO;2), 1995.
- Järvinen, E., Jourdan, O., Neubauer, D., Yao, B., Liu, C., Andreae, M. O., Lohmann, U., Wendisch, M., McFarquhar, G. M., Leisner, T., and Schnaiter, M.: Additional global climate cooling by clouds due to ice crystal complexity, *Atmospheric Chemistry and Physics*, 18, 15 767–15 781, <https://doi.org/10.5194/acp-18-15767-2018>, 2018.
- 645 Järvinen, E., Nehlert, F., Xu, G., Waitz, F., Mioche, G., Dupuy, R., Jourdan, O., and Schnaiter, M.: Investigating the vertical extent and short-wave radiative effects of the ice phase in Arctic summertime low-level clouds, *Atmospheric Chemistry and Physics*, 23, 7611–7633, <https://doi.org/10.5194/acp-23-7611-2023>, 2023.

- Jeggle, K., Neubauer, D., Binder, H., and Lohmann, U.: Cirrus formation regimes – data-driven identification and quantification of mineral dust effect, *Atmospheric Chemistry and Physics*, 25, 7227–7243, <https://doi.org/10.5194/acp-25-7227-2025>, 2025.
- 650 Jurkat-Witschas, T., Voigt, C., Groß, S., Kaufmann, S., Sauer, D., De la Torre Castro, E., Krämer, M., Schäfler, A., Afchine, A., Attinger, R., Bartolome Garcia, I., Beer, C. G., Bugliaro, L., Clemen, H.-C., Dekoutsidis, G., Ehrlich, A., Grawe, S., Hahn, V., Hendricks, J., Järvinen, E., Klimach, T., Krüger, K., Krüger, O., Lucke, J., Luebke, A. E., Marsing, A., Mayer, B., Mayer, J., Mertes, S., Moser, M., Müller, H., Nenakhov, V., Pöhlker, M., Pöschl, U., Pörtge, V., Rautenhaus, M., Righi, M., Röttenbacher, J., Rubin-Zuzic, M., Schaefer, J., Schnaiter, M., Schneider, J., Schumann, U., Spelten, N., Stratmann, F., Tomsche, L., Wagner, S., Wang, Z., Weber, A., Wendisch, M., Wernli, H.,
- 655 Wetzel, B., Wirth, M., Zahn, A., Ziereis, H., and Zöger, M.: CIRRUS-HL: Picturing high- and mid-latitude summer cirrus and contrail cirrus above Europe with airborne measurements aboard the research aircraft HALO, *Bulletin of the American Meteorological Society*, accepted, in press, 2025.
- Järvinen, E., van Diedenhoven, B., Magee, N., Neshyba, S., Schnaiter, M., Xu, G., Jourdan, O., Delene, D., Waitz, F., Lolli, S., and Kato, S.: Ice Crystal Complexity and Link to the Cirrus Cloud Radiative Effect, chap. 3, pp. 47–85, American Geophysical Union (AGU), ISBN 9781119700357, <https://doi.org/https://doi.org/10.1002/9781119700357.ch3>, 2023.
- 660 Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Meyer, J., Zoeger, M., Smith, J., Herman, R. L., et al.: A microphysics guide to cirrus clouds–Part 1: Cirrus types, *Atmospheric Chemistry and Physics*, 16, 3463–3483, 2016.
- Kristjánsson, J. E., Edwards, J. M., and Mitchell, D. L.: Impact of a new scheme for optical properties of ice crystals on climates of two GCMs, *Journal of Geophysical Research: Atmospheres*, 105, 10 063–10 079, <https://doi.org/https://doi.org/10.1029/2000JD900015>, 2000.
- 665 Lawson, R. P., Woods, S., Jensen, E., Erfani, E., Gurganus, C., Gallagher, M., Connolly, P., Whiteway, J., Baran, A. J., May, P., Heymsfield, A., Schmitt, C. G., McFarquhar, G., Um, J., Protat, A., Bailey, M., Lance, S., Muehlbauer, A., Stith, J., Korolev, A., Toon, O. B., and Krämer, M.: A Review of Ice Particle Shapes in Cirrus formed In Situ and in Anvils, *Journal of Geophysical Research: Atmospheres*, 124, 10 049–10 090, <https://doi.org/https://doi.org/10.1029/2018JD030122>, 2019.
- Lehmkuhl, F., Schüttrumpf, H., Schwarzbauer, J., Brüll, C., Dietze, M., Letmathe, P., Völker, C., and Hollert, H.: Assessment of the 2021
- 670 summer flood in Central Europe, *Environmental Sciences Europe*, 34, 107, <https://doi.org/10.1186/s12302-022-00685-1>, 2022.
- Loeb, N. G., Johnson, G. C., Thorsen, T. J., Lyman, J. M., Rose, F. G., and Kato, S.: Satellite and Ocean Data Reveal Marked Increase in Earth’s Heating Rate, *Geophysical Research Letters*, 48, e2021GL093 047, <https://doi.org/https://doi.org/10.1029/2021GL093047>, 2021.
- Luebke, A. E., Afchine, A., Costa, A., Groöb, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Krämer, M.: The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, *Atmospheric Chemistry and Physics*,
- 675 16, 5793–5809, <https://doi.org/10.5194/acp-16-5793-2016>, 2016.
- L’Ecuyer, T. S., Hang, Y., Matus, A. V., and Wang, Z.: Reassessing the Effect of Cloud Type on Earth’s Energy Balance in the Age of Active Spaceborne Observations. Part I: Top of Atmosphere and Surface, *Journal of Climate*, 32, 6197 – 6217, <https://doi.org/10.1175/JCLI-D-18-0753.1>, 2019.
- Macke, A., Mishchenko, M. I., and Cairns, B.: The influence of inclusions on light scattering by large ice particles, *Journal of Geophysical*
- 680 *Research: Atmospheres*, 101, 23 311–23 316, <https://doi.org/https://doi.org/10.1029/96JD02364>, 1996a.
- Macke, A., Mueller, J., and Raschke, E.: Single scattering properties of atmospheric ice crystals, *Journal of the Atmospheric Sciences*, 53, 2813–2825, 1996b.
- Macke, A., Francis, P. N., McFarquhar, G. M., and Kinne, S.: The Role of Ice Particle Shapes and Size Distributions in the Single Scattering Properties of Cirrus Clouds, *Journal of the Atmospheric Sciences*, 55, 2874 – 2883, [https://doi.org/10.1175/1520-0469\(1998\)055<2874:TROIPS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<2874:TROIPS>2.0.CO;2), 1998.
- 685

- Magee, N., Boaggio, K., Staskiewicz, S., Lynn, A., Zhao, X., Tusay, N., Schuh, T., Bandamede, M., Bancroft, L., Connelly, D., Hurler, K., Miner, B., and Khoudary, E.: Captured cirrus ice particles in high definition, *Atmospheric Chemistry and Physics*, 21, 7171–7185, <https://doi.org/10.5194/acp-21-7171-2021>, 2021.
- Mauritsen, T., Stevens, B., Roeckner, E., Crueger, T., Esch, M., Giorgetta, M., Haak, H., Jungclaus, J., Klocke, D., Matei, D., Mikolajewicz, U., Notz, D., Pincus, R., Schmidt, H., and Tomassini, L.: Tuning the climate of a global model, *Journal of Advances in Modeling Earth Systems*, 4, <https://doi.org/https://doi.org/10.1029/2012MS000154>, 2012.
- Meador, W. E. and Weaver, W. R.: Two-Stream Approximations to Radiative Transfer in Planetary Atmospheres: A Unified Description of Existing Methods and a New Improvement, *Journal of Atmospheric Sciences*, 37, 630 – 643, [https://doi.org/10.1175/1520-0469\(1980\)037<0630:TSATRT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<0630:TSATRT>2.0.CO;2), 1980.
- 695 Mishchenko, M. I., Rossow, W. B., Macke, A., and Lacis, A. A.: Sensitivity of cirrus cloud albedo, bidirectional reflectance and optical thickness retrieval accuracy to ice particle shape, *Journal of Geophysical Research: Atmospheres*, 101, 16 973–16 985, <https://doi.org/https://doi.org/10.1029/96JD01155>, 1996.
- Pincus, R. and Stevens, B.: Paths to accuracy for radiation parameterizations in atmospheric models, *Journal of Advances in Modeling Earth Systems*, 5, 225–233, <https://doi.org/https://doi.org/10.1002/jame.20027>, 2013.
- 700 Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks, P. A., Holz, R. E., et al.: The MODIS cloud optical and microphysical products: Collection 6 updates and examples from Terra and Aqua, *IEEE Transactions on Geoscience and Remote Sensing*, 55, 502–525, 2016.
- Ren, T., Yang, P., Loeb, N. G., Smith Jr., W. L., and Minnis, P.: On the Consistency of Ice Cloud Optical Models for Spaceborne Remote Sensing Applications and Broadband Radiative Transfer Simulations, *Journal of Geophysical Research: Atmospheres*, 128, e2023JD038 747, <https://doi.org/https://doi.org/10.1029/2023JD038747>, 2023.
- 705 Schmitt, C. G., Iaquinta, J., and Heymsfield, A. J.: The Asymmetry Parameter of Cirrus Clouds Composed of Hollow Bullet Rosette-Shaped Ice Crystals from Ray-Tracing Calculations, *Journal of Applied Meteorology and Climatology*, 45, 973 – 981, <https://doi.org/10.1175/JAM2384.1>, 2006.
- Schnaiter, M., Järvinen, E., Abdelmonem, A., and Leisner, T.: PHIPS-HALO: the airborne particle habit imaging and polar scattering probe – Part 2: Characterization and first results, *Atmos. Meas. Tech.*, 11, 341–357, 2018.
- 710 Schäfler, A. and Rautenhaus, M.: Interactive 3D Visual Analysis of Weather Prediction Data Reveals Midlatitude Overshooting Convection during the CIRRUS-HL Field Experiment, *Bulletin of the American Meteorological Society*, 104, E1426 – E1434, <https://doi.org/10.1175/BAMS-D-22-0103.1>, 2023.
- Schön, R., Schnaiter, M., Ulanowski, Z., Schmitt, C., Benz, S., Möhler, O., Vogt, S., Wagner, R., and Schurath, U.: Particle Habit Imaging Using Incoherent Light: A First Step toward a Novel Instrument for Cloud Microphysics, *Journal of Atmospheric and Oceanic Technology*, 28, 493 – 512, <https://doi.org/10.1175/2011JTECHA1445.1>, 2011.
- 715 Sourdeval, O., Gryspeerdt, E., Krämer, M., Goren, T., Delanoë, J., Afchine, A., Hemmer, F., and Quaas, J.: Ice crystal number concentration estimates from lidar–radar satellite remote sensing – Part 1: Method and evaluation, *Atmospheric Chemistry and Physics*, 18, 14 327–14 350, <https://doi.org/10.5194/acp-18-14327-2018>, 2018.
- 720 Stephens, G. L., Tsay, S.-C., Stackhouse, P. W., and Flatau, P. J.: The Relevance of the Microphysical and Radiative Properties of Cirrus Clouds to Climate and Climatic Feedback, *Journal of Atmospheric Sciences*, 47, 1742 – 1754, [https://doi.org/10.1175/1520-0469\(1990\)047<1742:TROTMA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1990)047<1742:TROTMA>2.0.CO;2), 1990.

- Stubenrauch, C. J., Rossow, W. B., Kinne, S., Ackerman, S., Cesana, G., Chepfer, H., Girolamo, L. D., Getzewich, B., Guignard, A., Heidinger, A., Maddux, B. C., Menzel, W. P., Minnis, P., Pearl, C., Platnick, S., Poulsen, C., Riedi, J., Sun-Mack, S., Walther, A., Winker, D., Zeng, S., and Zhao, G.: Assessment of Global Cloud Datasets from Satellites: Project and Database Initiated by the GEWEX Radiation Panel, *Bulletin of the American Meteorological Society*, 94, 1031 – 1049, <https://doi.org/10.1175/BAMS-D-12-00117.1>, 2013.
- Takano, Y. and Liou, K.-N.: Solar Radiative Transfer in Cirrus Clouds. Part I: Single-Scattering and Optical Properties of Hexagonal Ice Crystals, *Journal of Atmospheric Sciences*, 46, 3 – 19, [https://doi.org/10.1175/1520-0469\(1989\)046<0003:SRTICC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1989)046<0003:SRTICC>2.0.CO;2), 1989.
- Travis, L., Mishchenko, M. I., and Lacis, A. A.: Scattering, Absorption, and Emission of Light by Small Particles, <https://api.semanticscholar.org/CorpusID:49399284>, 2002.
- Um, J. and McFarquhar, G. M.: Single-Scattering Properties of Aggregates of Bullet Rosettes in Cirrus, *Journal of Applied Meteorology and Climatology*, 46, 757 – 775, <https://doi.org/10.1175/JAM2501.1>, 2007.
- van Diedenhoven, B.: Variation of Ice Microphysical Properties With Temperature and Humidity at Tops of Convective Clouds, *Geophysical Research Letters*, 48, e2021GL093 673, <https://doi.org/https://doi.org/10.1029/2021GL093673>, e2021GL093673 2021GL093673, 2021.
- van Diedenhoven, B., Cairns, B., Fridlind, A. M., Ackerman, A. S., and Garrett, T. J.: Remote sensing of ice crystal asymmetry parameter using multi-directional polarization measurements Part 2: Application to the Research Scanning Polarimeter, *Atmospheric Chemistry and Physics*, 13, 3185–3203, <https://doi.org/10.5194/acp-13-3185-2013>, 2013.
- van Diedenhoven, B., Ackerman, A. S., Cairns, B., and Fridlind, A. M.: A Flexible Parameterization for Shortwave Optical Properties of Ice Crystals, *Journal of the Atmospheric Sciences*, 71, 1763 – 1782, <https://doi.org/10.1175/JAS-D-13-0205.1>, 2014.
- van Diedenhoven, B., Fridlind, A. M., Cairns, B., and Ackerman, A. S.: Variation of ice crystal size, shape, and asymmetry parameter in tops of tropical deep convective clouds, *Journal of Geophysical Research: Atmospheres*, 119, 11,809–11,825, <https://doi.org/https://doi.org/10.1002/2014JD022385>, 2014.
- Vogelmann, A. M. and Ackerman, T. P.: Relating Cirrus Cloud Properties to Observed Fluxes: A Critical Assessment, *Journal of Atmospheric Sciences*, 52, 4285 – 4301, [https://doi.org/10.1175/1520-0469\(1995\)052<4285:RCCPTO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<4285:RCCPTO>2.0.CO;2), 1995.
- Wagner, S. W., Schnaiter, M., Xu, G., Nehlert, F., and Järvinen, E.: Light scattering and microphysical properties of atmospheric bullet rosette ice crystals, *EGU sphere*, 2024, 1–27, <https://doi.org/10.5194/egusphere-2024-3316>, 2024.
- Waitz, F., Schnaiter, M., Leisner, T., and Järvinen, E.: PHIPS–HALO: the airborne Particle Habit Imaging and Polar Scattering probe – Part 3: Single-particle phase discrimination and particle size distribution based on the angular-scattering function, *Atmospheric Measurement Techniques*, 14, 3049–3070, <https://doi.org/10.5194/amt-14-3049-2021>, 2021.
- Wang, C., Yang, P., Dessler, A., Baum, B. A., and Hu, Y.: Estimation of the cirrus cloud scattering phase function from satellite observations, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 138, 36–49, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2014.02.001>, 2014.
- Wang, Y., Hioki, S., Yang, P., King, M. D., Di Girolamo, L., Fu, D., and Baum, B. A.: Inference of an Optimal Ice Particle Model through Latitudinal Analysis of MISR and MODIS Data, *Remote Sensing*, 10, <https://doi.org/10.3390/rs10121981>, 2018.
- Weigel, R., Spichtinger, P., Mahnke, C., Klingebiel, M., Afchine, A., Petzold, A., Krämer, M., Costa, A., Molleker, S., Reutter, P., Szakáll, M., Port, M., Grulich, L., Jurkat, T., Minikin, A., and Borrmann, S.: Thermodynamic correction of particle concentrations measured by underwing probes on fast-flying aircraft, *Atmospheric Measurement Techniques*, 9, 5135–5162, <https://doi.org/10.5194/amt-9-5135-2016>, 2016.
- Wiscombe, W. J.: The Delta–M Method: Rapid Yet Accurate Radiative Flux Calculations for Strongly Asymmetric Phase Functions, *Journal of Atmospheric Sciences*, 34, 1408 – 1422, [https://doi.org/10.1175/1520-0469\(1977\)034<1408:TDMRYA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1408:TDMRYA>2.0.CO;2), 1977.

- 760 Wolf, V., Kuhn, T., Milz, M., Voelger, P., Krämer, M., and Rolf, C.: Arctic ice clouds over northern Sweden: microphysical properties studied with the Balloon-borne Ice Cloud particle Imager B-ICI, *Atmospheric Chemistry and Physics*, 18, 17371–17386, <https://doi.org/10.5194/acp-18-17371-2018>, 2018.
- Woods, S., Lawson, R. P., Jensen, E., Bui, T., Thornberry, T., Rollins, A., Pfister, L., and Avery, M.: Microphysical properties of tropical tropopause layer cirrus, *Journal of Geophysical Research: Atmospheres*, 123, 6053–6069, 2018.
- 765 Xu, G., Schnaiter, M., and Järvinen, E.: Accurate Retrieval of Asymmetry Parameter for Large and Complex Ice Crystals From In-Situ Polar Nephelometer Measurements, *Journal of Geophysical Research: Atmospheres*, 127, 1–19, <https://doi.org/10.1029/2021jd036071>, 2022.
- Yang, P. and Liou, K. N.: Geometric-optics–integral-equation method for light scattering by nonspherical ice crystals, *Appl. Opt.*, 35, 6568–6584, <https://doi.org/10.1364/AO.35.006568>, 1996.
- Yang, P., Gao, B.-C., Baum, B. A., Wiscombe, W. J., Hu, Y. X., Nasiri, S. L., Soulen, P. F., Heymsfield, A. J., McFarquhar, G. M., and
770 Miloshevich, L. M.: Sensitivity of cirrus bidirectional reflectance to vertical inhomogeneity of ice crystal habits and size distributions for two Moderate-Resolution Imaging Spectroradiometer (MODIS) bands, *Journal of Geophysical Research: Atmospheres*, 106, 17267–17291, <https://doi.org/https://doi.org/10.1029/2000JD900618>, 2001.
- Yang, P., Zhang, Z., Kattawar, G. W., Warren, S. G., Baum, B. A., Huang, H.-L., Hu, Y. X., Winker, D., and Iaquinta, J.: Effect of Cavities on the Optical Properties of Bullet Rosettes: Implications for Active and Passive Remote Sensing of Ice Cloud Properties, *Journal of Applied
775 Meteorology and Climatology*, 47, 2311 – 2330, <https://doi.org/10.1175/2008JAMC1905.1>, 2008.
- Yang, P., Bi, L., Baum, B. A., Liou, K.-N., Kattawar, G. W., Mishchenko, M. I., and Cole, B.: Spectrally Consistent Scattering, Absorption, and Polarization Properties of Atmospheric Ice Crystals at Wavelengths from 0.2 to 100 μ m, *Journal of the Atmospheric Sciences*, 70, 330–347, <https://doi.org/10.1175/JAS-D-12-039.1>, 2013.
- Yi, B.: Diverse cloud radiative effects and global surface temperature simulations induced by different ice cloud optical property parameter-
780 izations, *Scientific Reports*, 12, 10539, 2022.
- Yi, B., Yang, P., Baum, B. A., L’Ecuyer, T., Oreopoulos, L., Mlawer, E. J., Heymsfield, A. J., and Liou, K.-N.: Influence of Ice Particle Surface Roughening on the Global Cloud Radiative Effect, *Journal of the Atmospheric Sciences*, 70, 2794 – 2807, <https://doi.org/10.1175/JAS-D-13-020.1>, 2013.
- Yi, B., Rapp, A. D., Yang, P., Baum, B. A., and King, M. D.: A comparison of Aqua MODIS ice and liquid water cloud physical and
785 optical properties between collection 6 and collection 5.1: Pixel-to-pixel comparisons, *Journal of Geophysical Research: Atmospheres*, 122, 4528–4549, <https://doi.org/https://doi.org/10.1002/2016JD025586>, 2017a.
- Yi, B., Rapp, A. D., Yang, P., Baum, B. A., and King, M. D.: A comparison of Aqua MODIS ice and liquid water cloud physical and optical properties between collection 6 and collection 5.1: Cloud radiative effects, *Journal of Geophysical Research: Atmospheres*, 122, 4550–4564, <https://doi.org/https://doi.org/10.1002/2016JD025654>, 2017b.
- 790 Zhang, Y., Macke, A., and Albers, F.: Effect of crystal size spectrum and crystal shape on stratiform cirrus radiative forcing, *Atmospheric Research*, 52, 59–75, [https://doi.org/https://doi.org/10.1016/S0169-8095\(99\)00026-5](https://doi.org/https://doi.org/10.1016/S0169-8095(99)00026-5), 1999.