



Mesoscale eddies heterogeneously modulate CO₂ fluxes in eddy-rich regions of the Southern Ocean

Mariana Salinas-Matus^{1,2,3}, Nuno Serra³, Fatemeh Chegini³, and Tatiana Ilyina^{3,4,1}

¹Ocean Biogeochemistry, Climate Variability Department, Max Planck Institute for Meteorology, Hamburg, Germany

²International Max Planck Research School on Earth System Modelling, Hamburg, Germany

³Modeling the carbon cycle in the Earth system, Department of Earth System Sciences, Universität Hamburg, Hamburg, Germany

⁴Earth system modeling, Helmholtz-Zentrum Hereon, Geesthacht, Germany

Correspondence: Mariana Salinas-Matus (mariana.salinas@mpimet.mpg.de)

Abstract. Mesoscale eddies are known to influence the Southern Ocean biogeochemistry. However, the distinct contributions of cyclonic and anticyclonic eddies to air-sea CO₂ fluxes, as well as their longer-term effects remain poorly studied. We present results from a 27-year global eddy-resolving ocean-biogeochemical simulation. We used the Okubo-Weiss parameter to classify the modeled flow regimes into cyclonic and anticyclonic eddies, peripheries, and the surrounding background waters. Our results reveal a heterogeneous influence of eddies depending on the region, driven by regional differences in eddy intensity and the gradients in background properties. The factors controlling CO₂ fluxes within eddies follow the same degree of importance as in background waters, with $\Delta p\text{CO}_2$ being the dominant factor. This is driven primarily by changes in dissolved inorganic carbon. Our analysis shows that eddies act as a persistent carbon sink on decadal timescales, while their influence on shorter timescales is more variable and strongly shaped by eddy polarity. Anticyclonic and cyclonic eddies and periphery account for around 10 % of the Southern Ocean's carbon uptake, with anticyclonic eddies showing the highest carbon uptake per unit area. The ability of eddies to absorb carbon computed in our results is consistent with recent observational estimates, confirming that the model realistically represents the influence of mesoscale eddies on CO₂ fluxes. Above all, our results underscore the role of mesoscale eddies in enhancing carbon uptake across the Southern Ocean.

1 Introduction

The Southern Ocean exhibits high mesoscale eddy activity (Chelton et al., 2011b; Petersen et al., 2013). These mesoscale eddies influence the physical properties of seawater by transporting heat and salinity across regions and into the ocean's interior (Morrison et al., 2013; Griffies et al., 2015; Morrison et al., 2016; von Storch et al., 2016). Similarly, these eddies play a substantial role in the transport of biogeochemical properties (Chelton et al., 2011a; Dufour et al., 2015; Schütte et al., 2016; Moreau et al., 2017; Dawson et al., 2018; Patel et al., 2020; Su et al., 2021; Lévy et al., 2023). Notably, the mass transport induced by eddies is comparable to large-scale transport (Zhang et al., 2014). Moreover, eddies play a significant role in shaping Southern Ocean future climate projections (Farneti et al., 2010; Dufour et al., 2015; Hogg et al., 2015; Bilgen and Kirtman, 2020; van Westen and Dijkstra, 2021; Putrasahan et al., 2021). However, the study of mesoscale eddies has been



limited by the lack of biogeochemical datasets that can capture the high eddy activity in the region, both from observational and modeling perspectives. Observational studies are limited by the sparse spatial and temporal coverage of measurements (Gray, 2023; Dong et al., 2024), while eddy-resolving simulations are limited by their high computational cost (Hewitt et al., 2022; Guo and Timmermans, 2024).

Despite growing research interest, the role of cyclone and anticyclonic eddies in modulating air-sea CO₂ fluxes remains inconclusive. Regional and short-term oceanographic campaigns have identified that mesoscale eddies exhibit diverse influence on carbon uptake. It has been shown that both cyclonic and anticyclonic eddies can function as carbon sinks (Jones et al., 2017; Orselli et al., 2019a; Ford et al., 2023). Conversely, other studies have reported inverse patterns, where anticyclonic eddies release CO₂ to the atmosphere, while cyclonic eddies serve as carbon sinks (Pezzi et al., 2021; Kim et al., 2022). Additionally, it has been shown that there are seasonal differences in the carbon uptake depending on the eddy polarity (Song et al., 2016; Jones et al., 2017).

Only recently have studies begun to investigate the effects of mesoscale processes on air-sea CO₂ fluxes at larger spatial scales (Keppler et al., 2024; Guo and Timmermans, 2024; Li et al., 2025). Keppler et al. (2024) combined mesoscale eddies identified from satellite altimeter with data from biogeochemical Argo floats, showing on average that both cyclonic and anticyclonic eddies contribute to the carbon uptake in the Southern Ocean. However, cyclonic eddies were associated with a weaker carbon uptake or more outgassing, whereas anticyclonic eddies were associated with stronger carbon uptake or less outgassing. Li et al. (2025), using an observation-based machine learning approach, showed that anticyclonic eddies substantially enhance CO₂ uptake on average, whereas cyclonic eddies contribute less consistently. From a modeling perspective, Guo and Timmermans (2024) used an eddy-resolving global biogeochemical simulation to isolate the effect of mesoscale activity. Their findings indicate that mesoscale dynamics account for over 30 % of the total variance of the air-sea CO₂ fluxes in energetic regions and can act either as sources or sinks depending on the region, showing the importance of mesoscale dynamics from a global perspective. However, their study did not distinguish between the specific contributions of cyclonic and anticyclonic eddies. To date, no modeling study has explicitly examined the potentially contrasting impacts of cyclonic and anticyclonic eddies on CO₂ fluxes across the Southern Ocean, where this distinction may be particularly relevant.

This distinction may be important because cyclonic and anticyclonic eddies possess contrasting physical structures that are expected to influence air-sea CO₂ fluxes differently. Anticyclonic eddies are high-pressure systems characterized by positive vorticity in the southern hemisphere, which causes isopycnals to be pushed downward. This induces water downwelling, resulting in higher surface temperatures compared to surrounding waters, a mechanism known as eddy-pumping. Higher temperature reduces the solubility of CO₂ but the downwelling effect may transport carbon to deeper ocean layers. The opposite pattern occurs in cyclonic eddies, which are characterized by negative vorticity in the southern hemisphere, uplifted isopycnals, and water upwelling (McGillicuddy, 2016; Jones et al., 2017; Ford et al., 2023).

The regional differences within the Southern Ocean add another layer of complexity when attempting to generalize eddy behavior (McGillicuddy, 2016). For instance, the Agulhas Current system, south of Africa, is highly energetic and contributes significantly to the global thermohaline circulation via eddies and filaments involved in the Agulhas leakage (Richardson, 2007; Biastoch et al., 2009; Beal et al., 2011; Haarsma et al., 2011). The Brazil-Malvinas Confluence Zone, located where the Brazil



and Malvinas Currents meet, is characterized by strong temperature and salinity gradients, which create high-energy eddies and thermohaline fronts (Barré et al., 2006; Mason et al., 2017; Souza et al., 2021). In contrast, the region South of Tasmania is mostly influenced by the Antarctic Circumpolar Current (ACC), which produces a complex frontal system and robust eddy activity, influencing the water mass exchange between major ocean basins and the deep ocean (Rintoul and Sokolov, 2001). These distinct regional settings suggest that eddies impacts on CO₂ fluxes may not be uniform across the Southern Ocean, highlighting the need to investigate how mesoscale eddies operate in different oceanographic contexts.

Our objective in this work is to determine how anticyclonic and cyclonic eddies, periphery, and the surrounding background waters modulate air-sea CO₂ fluxes, to identify the physical and biogeochemical mechanisms driving these fluxes, and to examine how these mechanisms vary across different eddy-rich regions and the entire Southern Ocean. To achieve this, we use a global eddy-resolving ocean–biogeochemical model. Our model simulation is long enough to study the influence of eddies on CO₂ fluxes across timescales ranging from daily to decadal.

2 Methods

2.1 Model description

We use the ocean model ICON-O (Korn, 2017; Korn et al., 2022) incorporating the ocean biogeochemistry model HAMOCC (Six and Maier-Reimer, 1996; Ilyina et al., 2013). ICON-O solves the hydrostatic Boussinesq equations of the large-scale ocean dynamics with a free surface. The horizontal grid is composed of triangular cells with an Arakawa C-type staggering of variables. The grid is generated by subdividing the 20 triangular faces of an icosahedron inscribed into the sphere. The model computes tracer advection using a flux-corrected algorithm, which combines a low-order upwind scheme for stability with a higher-order scheme for accuracy. Vertical mixing is parametrized using a prognostic equation for turbulent kinetic energy, where the mixing efficiency is determined by a variable mixing length that adapts to local ocean conditions (Korn, 2017; Korn et al., 2022).

HAMOCC computes biological and chemical sources and sinks of biogeochemical tracers in the water column and sediment (Six and Maier-Reimer, 1996; Ilyina et al., 2013). Primary producers are represented by bulk phytoplankton and cyanobacteria, with growth limited by temperature, light, and nutrient availability. Grazing by zooplankton is restricted to bulk phytoplankton. Organic matter is divided into sinking particulate detritus and dissolved organic matter. All marine organic compounds are assumed to share a uniform elemental composition, based on a modified Redfield ratio. The biogeochemical tracers are advected and diffused by the ocean model. The ocean model also provides temperature, salinity and sea ice concentration to compute transformation rates, constants and fluxes. Atmospheric pressure, wind speed, and shortwave radiation are prescribed.

The air-sea exchange of CO₂ depends on the surface layer piston velocity, solubility factor, and pCO₂. The pCO₂ is determined from the total dissolved inorganic carbon (DIC), total alkalinity, temperature, and salinity. Total DIC is taken as the sum of all dissociated inorganic carbon species concentrations. Changes of DIC are driven by photosynthesis, grazing, zooplankton excretion, remineralization of dissolved organic matter and detritus, carbonate calcium production and dissolution, and air-sea flux.



2.2 Experiment design

We performed 27 years (1995-2022) of global simulations, saving daily average output. The nominal grid resolution varies between 8.4 and 10 km. The vertical grid is structured and non-uniform. We use 72 vertical layers, the thickness of the layers growing gradually with depth. The upper 300 m are represented by 29 layers. ERA5 is used as atmospheric forcing (Hersbach et al., 2020).

Due to the substantial computational time required for spin-up at high resolution and the complexity of HAMOCC, a direct spin-up at 10 km resolution was impractical. Therefore, a cascade strategy was implemented. First, the spin-up was conducted at a lower resolution of 40 km. A long spin-up of approximately 3,000 years was performed for the ocean's physical conditions with ICON-O only. The spin-up for the HAMOCC biogeochemistry model at pre-industrial conditions was carried out for around 900 years, until the upper ocean drift stabilized. The 40km resolution model was then run for the transient period until the year 1990. The physical and biogeochemical results from the coarser resolution were then used to initialize the 10 km resolution model. To allow the ocean conditions to adapt, only the ocean model was run for 5 years. After this adaptation period, the biogeochemical conditions were added. A tuning of the biogeochemical components in the 10 km resolution model was performed before the production phase of the simulation. This cascade strategy ensures an adequate representation of upper ocean mesoscale and regional processes, which are the main focus of this study. Furthermore, all model drifts in the control run remain sufficiently small in accordance with the CMIP6 protocol (Jones et al., 2016).

2.3 Eddy detection method

To detect mesoscale eddies and deformation-dominated areas (Fig. 1b and 2a), we implemented an algorithm based on the Okubo-Weiss parameter (OW) (Okubo, 1970; Weiss, 1991). OW measures the relative dominance of rotation (vorticity) against deformation (strain):

$$OW = \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right)^2 + \left(\frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right)^2 - \left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right)^2 \quad (1)$$

Normal Strain Shear Strain Vorticity

Four flow regimes were defined using OW: anticyclonic eddy cores, cyclonic eddy cores, periphery, and quiescent "background". The "background" regime is defined using a threshold of $\pm 0.05 \times 10^{-9} s^{-1}$, encompassing all OW values within this range. Vorticity-dominated regions (eddy flow regimes) are identified as closed contours with negative OW values smaller than $-0.05 \times 10^{-9} s^{-1}$. Relative vorticity is then used to distinguish between anticyclonic and cyclonic eddy cores. Deformation-dominated regions, referred to here as the periphery regime, are defined as areas with positive OW values exceeding $0.05 \times 10^{-9} s^{-1}$. These regions occur at the interacting regions between eddies and between eddies and large-scale currents, or along meanders and filaments, where the flow is stretched or compressed rather than rotating. Figure 2a provides a snapshot illustrating the defined flow regimes.



120 Eddies were not tracked, so the daily eddy-covered areas may represent either the persistence of the same eddies or the appearance of new ones from one day to the next. Additionally, the eddy lifetime was not considered in the analysis. Composites of each flow regime were generated for studying the overall effect of the flow regimes on the CO₂ flux. Here, a composite refers to the area-weighted average of relevant variables across all identified instances of a given flow regime, providing a representative picture of the typical conditions associated with each flow regime.

125 The net CO₂ flux is influenced by processes operating across different timescales (Gu et al., 2023). To determine the frequency range at which mesoscale eddies have the greatest impact on CO₂ flux, we applied a Fourier analysis to decompose the eddy composites into distinct frequency bands: (1) intra-annual variations above 16 months and up to 27 years (M16), (2) annual variations between 8 and 16 months (M8-M16), (3) intra-annual variations between 4 and 8 months (M4-M8), (4) intra-annual variations between 1 and 4 months (M1-M4), and (5) submonthly variations shorter than 1 month, down to daily

130 changes (M1). This study is the first to use a dataset that enables a time decomposition of eddy influence on CO₂ that ranges from decadal variability to daily changes.

3 Results and discussion

3.1 Simulated mesoscale eddy characteristics

Our 10 km resolution simulation captures mesoscale eddy activity (Fig. 1a and b) in the Southern Ocean, which is defined

135 here as the ice-free ocean south of 30°S. The model simulates high vorticity values (positive and negative) in the ACC region, the Agulhas Retroflexion region, the Brazil-Malvinas Confluence region, and the coast and south of Australia. Those regions also present the largest vorticity variability, highlighting intense mesoscale dynamics (Fig. 1a). The detected eddies cannot be directly compared with eddies tracked from observations. However, our model's eddy counts are consistent with the distribution of eddy kinetic energy attributed to eddies reported by Chelton et al. (2011b). Moreover, the distribution of detected eddies

140 aligns with the Southern Ocean's eddy hotspots reported by Frenger et al. (2015) (Fig. 1b).

The average daily number of eddies varies across regions, ranging from 11 to 38 eddies present each day. There is little to no difference between the number of anticyclonic and cyclonic eddies within each region. In terms of spatial coverage, mesoscale flow regimes occupy between 10 - 20% of the regions (Table 1). Satellite observations suggest this field typically covers 25-30% of the ocean surface, regardless of the size of the region (Chaigneau et al., 2009). It is important to consider

145 that the model has a limited ability to simulate the full eddy activity in the highest latitudes, where the deformation radii are well below 10 km.

Based on eddy spatial density in the model, we define three eddy-rich regions: the Brazil-Malvinas Confluence region, the Agulhas Retroflexion region and the South of Tasmania region (Fig. 1b, orange rectangles). The selection is made due to the heterogeneous nature of the Southern Ocean which makes it difficult to generalize the effect of eddies in the CO₂ flux.

150 For instance, the eddy intensity differs between regions. The Agulhas Retroflexion region presents the strongest and most persistent anticyclonic and cyclonic eddy intensity. Eddies in the Brazil-Malvinas Confluence region are less intense but with higher variability. Anticyclones in the Tasmania region are the least intense among the three regions but with the smallest



Region	total area (10 ⁶ km ²)	Area covered by mesoscale flow regimes %	Daily number of eddies	
			Anticyclones	Cyclones
Agulhas Retroflection	14.3	16.7	38	32
Brazil-Malvinas Confluence	10.2	9.9	14	11
South Tasmania	4.8	19.7	23	22

Table 1. Summary of eddy statistics for each region. Mesoscale flow regimes include anticyclonic eddies, cyclonic eddies, and periphery.

variability, while cyclonic intensity is similar to the Brazil-Malvinas Confluence region but with persistent intensity (Fig. 1c). Frenger et al. (2015) showed that the intensity is stronger in the Agulhas Retroflection, followed by the Brazil-Malvinas
155 Confluence, and weakest south of Tasmania.




Figure 1. Overview of modeled mesoscale eddies in the simulation period (1996–2022): (a) Mean and standard deviation of vorticity, with an approximate location of the ACC, plotted based on surface height, (b) Locations of detected anticyclonic and cyclonic eddy centroids, defined regions (orange rectangles), and eddy centroids census within 1° × 1° grid cells, and (c) Relative vorticity composites for anticyclonic and cyclonic eddies within selected regions.




Figure 2. Snapshot of the identified mesoscale dynamics and tracer or tracer flux anomalies in the Agulhas Retroflexion region (defined in Fig. 1b) on 03 June 2014. (a) Okubo-Weiss parameter, delineating three flow regimes: vorticity-dominated (green), deformation-dominated (violet), and "background" (grey). (b) Sea Surface Temperature (SST) anomaly, (c) air-sea CO₂ flux anomaly, and (d) Surface DIC anomaly. In all panels, contours outline anticyclonic and cyclonic eddies.

3.2 Heterogeneous dissolved inorganic carbon and temperature patterns across flow regimes in eddy-rich regions

The selected regions exhibit distinct characteristics, influenced by their latitudinal positions, ocean current, and frontal systems, and mesoscale activity. In the Agulhas Retroflexion and Brazil-Malvinas Confluence regions, both situated at similar latitudes, a pronounced spatial pattern emerges (Fig. 3a). The pronounced spatial pattern is marked by warmer temperatures, higher salinity, and lower pCO₂ and DIC concentrations to the north, contrasting sharply with colder, fresher, and carbon-rich waters to the south. The Tasmania region, located farther south, experiences water upwelling that brings carbon-rich waters from the deep to the surface, resulting in colder temperatures and high DIC concentrations throughout the year (Fig. 3a). The overall behavior of the simulated temperature and DIC is consistent with the oceanographic knowledge of the Southern Ocean (Talley et al., 2011). Mesoscale activity plays a substantial role in modulating carbon and temperature patterns (Fig. 2b-d).

In our simulations, flow regimes exhibit distinct characteristics that influence the CO₂ flux (Fig. 2b-d), with contrasting vertical structures of cyclonic and anticyclonic eddies emerging (Fig. 3b), consistent with previous findings (Keppler et al., 2024). Anticyclones, which tend to induce water downwelling, exhibit higher temperatures and lower DIC concentrations. In contrast, cyclones exhibit lower temperatures and higher DIC concentrations, mostly driven by water upwelling (Fig. 3b). The periphery regime shows intermediate properties, representing the transition zones surrounding both anticyclonic and cyclonic eddies. In the Agulhas Retroflexion region, the "background" shows slightly lower temperatures and higher DIC concentrations compared with cyclonic eddies. This contrast is influenced by the Indian Ocean origin of the eddies, which transport warmer waters into the region. However, in the Brazil Confluence and Tasmania regions, "background" conditions exhibit temperature and DIC levels between those observed for anticyclonic and cyclonic eddies (Fig. 3b).

The strong relationship between temperature, DIC, and gas solubility directly influences the exchange of CO₂ between the ocean and the atmosphere. The Southern Ocean's overall CO₂ uptake capacity is evident in our simulation. An outgassing band



between 40°S and 55°S, particularly in the Atlantic and Indian sectors, is associated with the high DIC concentration in the Southern Ocean upwelling region. North of 60°S, the CO₂ flux exhibits relatively low variability compared to the polar region but with notable localized signals influenced by mesoscale structures (Fig. 3c).

In the Agulhas Retroflection and Brazil-Malvinas Confluence regions, all four regimes show a mean distribution indicating carbon uptake, with a wider spread toward stronger uptake in both cyclonic and anticyclonic eddies. However, the distribution of cyclonic eddies and "background" shifts slightly toward carbon outgassing events (Fig. 3c).

In contrast, the Tasmania region presents a different pattern. The "background" shows a mean data distribution close to zero CO₂ flux, while cyclonic, anticyclonic, and periphery regimes indicate carbon release. Cyclonic eddies exhibit the highest outgassing. However, all regimes present a broad spread extending towards higher carbon uptake (Fig. 3c). The outgassing pattern might be associated with the predominance of eddies in zones influenced by upwelling carbon-rich waters (Fig. 3a) and the overall lower eddy intensity in the region (Fig. 1c).

3.3 Enhanced CO₂ uptake by anticyclonic eddies

The integrated CO₂ flux over the 27 analyzed years indicates that mesoscale flow regimes (anticyclones, cyclones, and periphery) have a greater capacity for carbon uptake compared to the "background" (Fig. 4a). When accounting for the area they occupy, these regimes are responsible for 15–22% of the carbon uptake within the defined regions, and 10% of the total carbon uptake in the entire Southern Ocean (Table 2).

Among these regimes, anticyclonic eddies show an enhanced ability to take up CO₂ (Fig. 4a). In particular, in the Agulhas Retroflection and Brazil-Malvinas Confluence regions, anticyclonic eddies exhibit the highest carbon uptake capacity, whereas the "background" has the lowest carbon uptake per unit area. In these two regions, cyclonic eddies also exhibit an enhanced ability to take up carbon compared with the "background", although not as intense as anticyclonic eddies (Fig. 4a). Anticyclonic and cyclonic eddies have the ability to take up more, or outgas less, carbon compared to the "background" (Fig. 4c). This is explained by the stronger eddy-intensity (Fig. 1c) and the stronger vertical gradient in those two regions (Fig. 3b).

Tasmania region exhibits a contrasting pattern, as the "background" has the highest CO₂ uptake (Fig. 4a). Here, both anticyclonic and cyclonic eddies take up less or outgas more carbon (Fig. 4c). This pattern is influenced by the weaker eddy intensity (Fig. 1c) and the characteristics of the region. Most eddies in the Tasmania region are located in the high-DIC band, which limits the ocean's capacity for CO₂ uptake (Fig. 3b). Nevertheless, anticyclonic and cyclonic eddies take up carbon in the entire analyzed period (Fig. 4a), a pattern influenced by sporadic but intense carbon uptake events (Fig. 3).

The periphery regime displays an intermediate carbon uptake between anticyclonic and cyclonic eddies. However, when considering the area it covers, the periphery regime is the mesoscale regime that contributes the most to net CO₂ uptake, accounting for around 60–70% of the carbon uptake generated by the mesoscale flow regimes. Given this substantial contribution, understanding the physical and biogeochemical drivers of CO₂ uptake in the periphery regime is an important direction for future research. Meanwhile, the combined contribution of anticyclonic and cyclonic eddies represents about 20–30% of the mesoscale regimes' carbon uptake (Table 2). This is particularly relevant given the observed trend of increasing eddy kinetic energy (EKE) in the Southern Ocean over recent decades (Hogg et al., 2015). If this trend continues, it could lead to a more




Figure 3. Overview of sea surface temperature (SST), surface DIC, and CO₂ flux in the Southern Ocean and composite quantities for flow regimes within the defined regions: (a) Mean and standard deviation of SST and DIC; (b) Vertical temperature and DIC profile composites for the flow regimes within the defined regions, (c) Mean and standard deviation of CO₂ flux, along with composite CO₂ fluxes for the flow regimes within the defined regions.

210 eddy-active Southern Ocean, with an expansion of the area influenced by mesoscale features. Such changes may potentially enhance the ocean's capacity for CO₂ uptake in the future, as larger regions become dominated by eddies.

The influence of mesoscale eddies on CO₂ flux has been identified through in situ data. Previous studies have shown that Agulhas anticyclonic eddies have a stronger CO₂ uptake ability than their surrounding water (Orselli et al., 2019b), which contributes to the faster acidification of the region (Orselli et al., 2019a). In the Tasmania region, Jones et al. (2017) identified
215 eddies as hotspots for carbon uptake. In the Brazil-Malvinas Confluence, a single anticyclonic eddy was identified as a CO₂ source to the atmosphere (Pezzi et al., 2021).

Our model results are generally consistent with these observational findings. We observe a notably enhanced carbon uptake by anticyclones in the Agulhas region compared to the other regions, and eddies in the Tasmania region also act as persistent CO₂ sinks. In the Brazil-Malvinas Confluence, a direct comparison is not possible since our analysis focuses on the overall



220 eddy effect over time rather than on individual eddy events. However, it is worth mentioning that both anticyclonic and cyclonic eddies do exhibit episodes of CO₂ outgassing (Fig. 3c).

Despite regional heterogeneity, we find that the overall enhanced ability of anticyclonic eddies is a persistent feature across the entire Southern Ocean. Anticyclonic eddies take up more or outgas less carbon than the background, while cyclonic eddies exhibit the opposite pattern (Fig. 4c). This is consistent with studies using observation-based datasets (Keppler et al., 2024; Li et al., 2025). This contrasting behavior may be attributed to eddy-pumping. In anticyclonic eddies, this mechanism facilitates the transfer of carbon to deeper ocean layers, enhancing the ocean's capacity for carbon uptake over time. In contrast, cyclonic eddies may act to reduce carbon uptake by limiting this downward transport.

The strong agreement between the model and observations underscores the critical role of mesoscale eddies in modulating CO₂ fluxes. This agreement indicates that the model is effectively capturing the key physical and biogeochemical processes involved. Moreover, the model provides a more detailed and continuous representation of eddy dynamics, offering a more robust signal than observations alone, which may be spatially or temporally limited.

Table 2. Area-integrated carbon uptake contributions by region and mesoscale flow regime. The value beneath each region name indicates the total carbon uptake (including background, eddies, and periphery). The values in parentheses represent the contribution of each flow regime as a percentage, first relative to the total uptake (in bold), and second relative to the combined uptake from all mesoscale flow regimes (eddies + periphery).

Carbon uptake [Pg C yr ⁻¹]				
Region	Mesoscale flow regimes	Disentangling Mesoscale flow regimes		
		Anticyclones	Cyclones	Periphery
Agulhas Retroflection 0.14±0.08	0.032 ± 0.016 (22±11%)	0.009 ± 0.003 (27 ± 11%)	0.003 ± 0.002 (10 ± 7%)	0.020 ± 0.010 (62 ± 31%)
Brazil-Malvinas Confluence 0.10±0.07	0.013 ± 0.009 (13±9%)	0.002 ± 0.001 (15 ± 7%)	0.001 ± 0.001 (7 ± 7%)	0.010 ± 0.007 (74 ± 51%)
South Tasmania 0.02±0.05	0.003 ± 0.010 (15±50%)	0.0006 ± 0.001 (20 ± 62%)	0.0002 ± 0.001 (8 ± 60%)	0.002 ± 0.006 (71 ± 206%)
Entire SO 1.17±0.60	0.118 ± 0.063 (10±5%)	0.020 ± 0.010 (18 ± 9%)	0.010 ± 0.008 (9 ± 7%)	0.086 ± 0.045 (72 ± 38%)

The analysis suggest that eddies act as a persistent carbon sink on decadal timescales (M16) (Fig. 4d, e, and f). In the Agulhas Retroflection and Brazil-Malvinas Confluence regions, both anticyclonic and cyclonic eddies exhibit enhanced CO₂ uptake relative to the "background" (Fig. 4d and e). In contrast, eddies in the Tasmania region do not absorb more carbon than the "background" (Fig. 4f). Nevertheless, since the integrated CO₂ flux in this region remains an overall carbon uptake, these eddies persistently take up carbon despite the high concentrations of DIC in that region. This is particularly important since the recent trend has proposed increased eddy activity in the Southern Ocean (Hogg et al., 2015; Martínez-Moreno et al., 2021) and the growing influence of eddies on heat distribution in a warming ocean (He et al., 2023).




Figure 4. (a) Time-integrated CO_2 flux composites for flow regimes across the defined regions, including relative contributions of CO_2 flux drivers (solubility, pCO_2 and, piston velocity) to the variability. (b) For the Agulhas Retroflection region, though the other regions follow the same pattern, relative contributions of these drivers across different frequency bands (M16: interannual variations above 16 months, M8-M16: annual variations between 8 and 16 months, M4-M8: intra-annual variations between 4 and 8 months, M1-M4: intra-annual variations between 1 and 4 months, and M1: submonthly variations). (c) Composite anomalies of total CO_2 flux relative to the "background" for anticyclonic and cyclonic eddies. (d–f) Composite anomalies of CO_2 flux across different frequency bands in three regions: (d) Agulhas Retroflection, (e) Brazil–Malvinas Confluence, and (f) south of Tasmania.

At intra-annual frequencies (M1 to M8-M16), the carbon flux anomalies are generally smaller but exhibit greater variability, indicating rapid events of carbon uptake and release. However, a pattern emerges: anticyclones tend to absorb more carbon or release less carbon than the "background", while cyclones typically absorb less or release more carbon. This pattern is the most pronounced in the Agulhas region, followed by the Brazil-Malvinas Confluence, and is much weaker in the Tasmania region. The contrast between cyclones and anticyclones is particularly distinguishable in the M1-M4 and M4-M8 bands. These frequency bands are linked to seasonal differences or rapid events such as phytoplankton blooms (Fig. 4d, e, and f). Keppler et al. (2024) identified a clear seasonal pattern, with cyclonic eddies in the ACC exhibiting outgassing in fall and carbon uptake in spring, while anticyclonic eddies north of the ACC showed enhanced uptake during spring.



3.4 DIC as the key regulator of oceanic $p\text{CO}_2$ and its main role on CO_2 flux

The CO_2 flux is influenced by three components: (1) solubility factor (S_c), driven by temperature and salinity, (2) piston velocity (k_w), driven by wind speed and temperature, and (3) the air-sea $p\text{CO}_2$ difference ($\Delta p\text{CO}_2 = p\text{CO}_2^{\text{ocean}} - p\text{CO}_2^{\text{atm}}$).

250 Appendix A details how each term (S_c , k_w , and $\Delta p\text{CO}_2$) is isolated for the analysis.

Among these factors, k_w and $\Delta p\text{CO}_2$ are the dominant contributors to the total CO_2 flux (Fig. 4a). In the Agulhas Retroflection and Brazil-Malvinas Confluence regions, the influence of $\Delta p\text{CO}_2$ is larger in cyclonic eddies, periphery, and "background". However, in anticyclonic eddies, k_w has a stronger impact. This may result from reduced variability in surface $\Delta p\text{CO}_2$, as anticyclones contain warmer, lower-nutrient water, which limits biological activity and reduces $p\text{CO}_2$ disequilibrium with the atmosphere. By contrast, in the Tasmania region $\Delta p\text{CO}_2$ consistently dominates across all flow regimes. The contribution of S_c remains minimal across all regions and flow regimes, ranging from 10-13% (Fig. 4a).

The time-frequency analysis of the effect of the CO_2 flux terms reveals a shift in the relative importance of these drivers. At higher frequencies, k_w becomes the dominant factor influencing CO_2 flux, while at lower frequencies, the influence of $\Delta p\text{CO}_2$ increases (Fig. 4b). Despite these variations, $\Delta p\text{CO}_2$ remains a key driver of the total flux. (Fig. 4a). Our results align with those of Gu et al. (2023), which reported that, at higher frequencies (subseasonal timescales), wind explains the majority of the global CO_2 flux anomaly, while at lower frequencies (seasonal, interannual and decadal timescales) $\Delta p\text{CO}_2$ is the dominant factor. Our study expands on the factors that control the CO_2 flux in eddies, showing that k_w tends to play a greater or comparable role in eddies than in the "background" regime.

In our simulation, the $p\text{CO}_2^{\text{atm}}$ is prescribed, thus variations in $\Delta p\text{CO}_2$ are primarily driven by changes in $p\text{CO}_2^{\text{ocean}}$. The $p\text{CO}_2^{\text{ocean}}$ is driven by changes in DIC, alkalinity, sea surface salinity, and SST. To decompose the contributions of these factors, we applied a decomposition method (Takahashi et al., 1993), detailed in Appendix B. In the three selected regions, DIC and SST were the primary drivers of $\Delta p\text{CO}_2^{\text{ocean}}$. The contributions of alkalinity and salinity were relatively small; therefore, salinity was grouped with SST to represent the physical drivers, and alkalinity was grouped with DIC to represent the biogeochemical drivers. Because the salinity contribution was minor, the combined SST and salinity component is referred to as the thermal component, while the DIC and alkalinity component is referred to as the non-thermal component.

The regression of the total flux on the non-thermal component, primarily driven by DIC, exhibits high R^2 values (Fig. 5), indicating that a larger proportion of $p\text{CO}_2^{\text{ocean}}$ variability is explained by this component. Its dominance is present in the total signal as well across almost all frequency bands (except M4-M8) and regions. In contrast, the thermal component shows consistently low R^2 values (Fig. 5a, b), suggesting a less relevant role in explaining the variability. It is worth mentioning that the thermal component counteracts the influence of the non-thermal component.

Other studies have found that DIC is the dominant driver of oceanic $\Delta p\text{CO}_2$ in the Southern Ocean (Landschützer et al., 2016; Lerner et al., 2021). Our results reinforce this finding and further show that the drivers of oceanic $p\text{CO}_2$ in eddies are largely consistent with those in the surrounding "background" waters. Our results show that DIC is the dominant contributor to the non-thermal component of the $p\text{CO}_2^{\text{ocean}}$, which in turn exerts the strongest influence on air-sea CO_2 fluxes. Therefore, DIC is a key driver of CO_2 fluxes in the Southern Ocean eddies.




Figure 5. Contribution of the thermal and non-thermal components to the total oceanic ΔpCO_2 across flow regimes in the defined regions. The coefficient of determination (R^2) was calculated from linear regressions between each component (thermal and non-thermal) and the total ΔpCO_2 , quantifying the proportion of variability explained by each. Regressions were performed separately for each component using time series of the composites within each region.

4 Summary and conclusion

The purpose of our analysis was twofold: to characterize the role of mesoscale oceanographic features in modulating CO_2 fluxes across three eddy-rich regions of the Southern Ocean, and to investigate the physical and biogeochemical drivers that control these fluxes. We present a 27-year long high-resolution simulation, which is notably long for an ocean-biogeochemical model of this kind. Our simulation enables, for the first time, to assess the influence of eddies on CO_2 fluxes across a broad range of timescales, from daily to decadal variability. The model successfully captures mesoscale activity, reproducing the distinct characteristics of eddies: anticyclones associated with higher temperatures and lower DIC concentration, while cyclones exhibit lower temperatures and higher DIC concentrations.

We find that eddies act as a persistent carbon sink at decadal scales. At shorter timescales, mean flux anomalies are smaller but exhibit higher variability, with anticyclonic and cyclonic eddies displaying distinct behaviors. Anticyclones typically enhance carbon uptake or suppress outgassing, whereas cyclones more often reduce uptake or promote outgassing. These highlight the importance of eddy type in shaping short-term air-sea CO_2 fluxes.

Beyond these temporal dynamics, our results show that the mesoscale flow regimes (anticyclones, cyclones, and periphery) are more effective in taking up carbon than the "background", accounting for about 10% of the total carbon uptake in the Southern Ocean, despite the smaller region they occupy. Anticyclonic eddies are especially effective, emerging as the flow



regime with the highest carbon uptake per unit area. These results are relevant given current trends suggesting that eddies may become more widespread in the future in the Southern Ocean.

Across all timescales and regions, $\Delta p\text{CO}_2$ plays a central role in modulating the CO_2 flux, outweighing the influence of piston velocity (wind) and solubility factor. We identified the non-thermal component (primarily driven by DIC) as the dominant control on oceanic $\Delta p\text{CO}_2$. This finding is consistent with previous studies on the Southern Ocean but extends our understanding by explicitly identifying the drivers of CO_2 fluxes within eddies. Our results indicate that the factors regulating oceanic $\Delta p\text{CO}_2$ in eddies are largely consistent with those in the surrounding "background" waters. However, given the differences we observe in CO_2 uptake between eddies and the "background", further analysis is needed to determine the processes driving DIC variability within eddies and their impact on carbon fluxes.

The pattern of anticyclonic and cyclonic eddies in modulating CO_2 fluxes in the Southern Ocean in our simulation is consistent with recent observational estimates. This confirms that the model realistically represents the influence of mesoscale eddies on CO_2 fluxes. This agreement supports the model's ability to capture key physical and biogeochemical processes driving carbon uptake in the Southern Ocean. Moreover, the overall ability of eddies to absorb carbon confirms the importance of mesoscale eddies in modulating the CO_2 fluxes in the Southern Ocean, a region recognized as one of the most critical for global carbon sequestration.

Code availability. Python scripts to reproduce the analysis are available at: <https://doi.org/10.17617/3.LQT15X>

Author contributions. TI conceived the original idea. MSM and NS designed the study and led the analysis. FC, TI and members of TI's group developed the strategy for running the model. FC and MSM carried out the model simulations. MSM wrote the manuscript, with all authors contributing to its review and editing.

Competing interests. The authors declare that they have no competing interests.

Acknowledgements. MSM was supported by the German Research Foundation (DFG) under Germany's Excellence Strategy—EXC 2037 "CLICCS—Climate, Climatic Change, and Society"—Project Number 390683824. TI and FC were supported by the EU Horizon 2020 research and innovation program under grant agreement 101003536 (project ESM2025—Earth System Models for the Future). Computational resources were provided by the German Climate Computing Center (DKRZ). We thank Lucas Casaroli for providing the initial conditions used to initialize the 10 km resolution simulation. We are grateful to David Nielsen for his internal review, which helped improve the manuscript. We also thank Jacqueline Behncke for her valuable comments and feedback on the manuscript.



Appendix A: ocean-atmosphere CO₂ flux terms effect

The air-sea CO₂ flux is determined by the solubility factor (S_c), influenced by temperature and salinity, the piston velocity (k_w), influenced by wind speed and temperature, and the air-sea pCO₂ difference ($\Delta pCO_2 = pCO_2^{ocean} - pCO_2^{atm}$).

$$CO_{2flux} = S_c k_w \Delta pCO_2 \quad (A1)$$

To isolate the effects of each term, we used the long-term mean state of the system as a baseline. By keeping the baseline constant for each driver, we could subtract its effect from the total CO₂ flux to determine the individual contributions (Chikamoto and DiNezio, 2021).

$$CO_{2flux} \approx F_{piston} + F_{solubility} + F_{\Delta pCO_2} \quad (A2)$$

$$F_{piston} = k_w(t) S_c(t) \Delta pCO_2(t) - \overline{k_w(t_b)} S_c(t) \Delta pCO_2(t) \quad (A3)$$

$$F_{solubility} = k_w(t) S_c(t) \Delta pCO_2(t) - k_w(t) \overline{S_c(t_b)} \Delta pCO_2(t) \quad (A4)$$

$$F_{\Delta pCO_2} = k_w(t) S_c(t) \Delta pCO_2(t) - k_w(t) S_c(t) \overline{\Delta pCO_2(t_b)} \quad (A5)$$

$$(A6)$$

where $\overline{k_w(t_b)}$, $\overline{S_c(t_b)}$, and $\overline{\Delta pCO_2(t_b)}$ are the respective baseline values, defined as their long-term means.

The combined effects of each driver should approximate the total CO₂ flux. The reconstruction is an approximation because it does not account for the effects of sea ice, although our analysis was restricted to sea-ice-free regions.

Appendix B: Decomposition of pCO_2^{ocean}

Since pCO_2^{air} is prescribed in the simulation, the primary variations in ΔpCO_2 are driven by changes in pCO_2^{sea} . The pCO_2^{sea} is influenced by variations in DIC, alkalinity, sea surface salinity, and temperature. To decompose the pCO_2^{sea} , we applied a

widely used decomposition method (Takahashi et al., 1993).

$$\Delta pCO_2^{ocean} = \Delta pCO_2^{DIC} + \Delta pCO_2^{alk} + \Delta pCO_2^{SST} + \Delta pCO_2^{SSS} \quad (B1)$$

$$\Delta pCO_2^{DIC} = \Delta DIC * \gamma_{DIC} * pCO_{2Ref} / \overline{DIC} \quad (B2)$$

$$\Delta pCO_2^{alk} = \Delta alk * \gamma_{alk} * pCO_{2Ref} / \overline{alk} \quad (B3)$$

$$\Delta pCO_2^{SSS} = \Delta SSS * \gamma_{SSS} * pCO_{2Ref} / \overline{SSS} \quad (B4)$$

$$\Delta pCO_2^{SST} = (\exp^{\Delta SST * \gamma_{SST}} * pCO_{2Ref}) - pCO_{2Ref} \quad (B5)$$

$$(B6)$$

The values of γ_X refer to the Revelle factors. γ_{SSS} and γ_{SST} are 1 and 0.0423; γ_{DIC} and γ_{alk} are computed daily as (Sarmiento and Gruber, 2006):



$$\gamma_{DIC} = \frac{3 \, alk \, DIC - 2 \, DIC^2}{(2 \, DIC - alk) (alk - DIC)} \quad (B7)$$

$$350 \quad \gamma_{alk} = -\frac{alk^2}{(2 \, DIC - alk) (alk - DIC)} \quad (B8)$$

$$(B9)$$



References

- Barré, N., Provost, C., and Saraceno, M.: Spatial and temporal scales of the Brazil-Malvinas Current confluence documented by simultaneous MODIS Aqua 1.1-km resolution SST and color images, *Advances in Space Research*, 37, 770–786, <https://doi.org/10.1016/j.asr.2005.09.026>, 2006.
- Beal, L. M., Ruijter, W. P. D., Biastoch, A., Zahn, R., Cronin, M., Hermes, J., Lutjeharms, J., Quartly, G., Tozuka, T., Baker-Yeboah, S., Bornman, T., Cipollini, P., Dijkstra, H., Hall, I., Park, W., Peeters, F., Penven, P., Ridderinkhof, H., and Zinke, J.: On the role of the Agulhas system in ocean circulation and climate, <https://doi.org/10.1038/nature09983>, 2011.
- Biastoch, A., Böning, C. W., Schwarzkopf, F. U., and Lutjeharms, J. R.: Increase in Agulhas leakage due to poleward shift of Southern Hemisphere westerlies, *Nature*, 462, 495–498, <https://doi.org/10.1038/nature08519>, 2009.
- Bilgen, S. I. and Kirtman, B. P.: Impact of ocean model resolution on understanding the delayed warming of the Southern Ocean, *Environmental Research Letters*, 15, <https://doi.org/10.1088/1748-9326/abbc3e>, 2020.
- Chaigneau, A., Eldin, G., and Dewitte, B.: Eddy activity in the four major upwelling systems from satellite altimetry (1992–2007), *Progress in Oceanography*, 83, 117–123, <https://doi.org/10.1016/j.pocean.2009.07.012>, 2009.
- Chelton, D. B., Gaube, P., Schlax, M. G., Early, J. J., and Samelson, R. M.: The Influence of Nonlinear Mesoscale Eddies on Near-Surface Oceanic Chlorophyll, <https://www.science.org>, 2011a.
- Chelton, D. B., Schlax, M. G., and Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Progress in Oceanography*, 91, 167–216, <https://doi.org/10.1016/j.pocean.2011.01.002>, 2011b.
- Chikamoto, M. O. and DiNezio, P.: Multi-Century Changes in the Ocean Carbon Cycle Controlled by the Tropical Oceans and the Southern Ocean, *Global Biogeochemical Cycles*, 35, <https://doi.org/10.1029/2021GB007090>, 2021.
- Dawson, H. R., Strutton, P. G., and Gaube, P.: The Unusual Surface Chlorophyll Signatures of Southern Ocean Eddies, *Journal of Geophysical Research: Oceans*, 123, 6053–6069, <https://doi.org/10.1029/2017JC013628>, 2018.
- Dong, Y., Bakker, D. C. E., Bell, T. G., Yang, M., Landschützer, P., Hauck, J., Rödenbeck, C., Kitidis, V., Bushinsky, S. M., and Liss, P. S.: Direct observational evidence of strong CO₂ uptake in the Southern Ocean, <https://www.science.org>, 2024.
- Dufour, C. O., Griffies, S. M., de Souza, G. F., Frenger, I., Morrison, A. K., Palter, J. B., Sarmiento, J. L., Galbraith, E. D., Dunne, J. P., Anderson, W. G., and Slater, R. D.: Role of mesoscale eddies in cross-frontal transport of heat and biogeochemical tracers in the Southern Ocean, *Journal of Physical Oceanography*, 45, 3057–3081, <https://doi.org/10.1175/JPO-D-14-0240.1>, 2015.
- Farneti, R., Delworth, T. L., Rosati, A. J., Griffies, S. M., and Zeng, F.: The role of mesoscale eddies in the rectification of the Southern ocean response to climate change, *Journal of Physical Oceanography*, 40, 1539–1557, <https://doi.org/10.1175/2010JPO4353.1>, 2010.
- Ford, D. J., Tilstone, G. H., Shutler, J. D., Kitidis, V., Sheen, K. L., Dall’Olmo, G., and Orselli, I. B.: Mesoscale Eddies Enhance the Air-Sea CO₂ Sink in the South Atlantic Ocean, *Geophysical Research Letters*, 50, <https://doi.org/10.1029/2022GL102137>, 2023.
- Frenger, I., Münnich, M., Gruber, N., and Knutti, R.: Southern Ocean eddy phenomenology, *Journal of Geophysical Research: Oceans*, 120, 7413–7449, <https://doi.org/10.1002/2015JC011047>, 2015.
- Gray, A. R.: The Four-Dimensional Carbon Cycle of the Southern Ocean, *Annual Review of Marine Science*, 41, 53, <https://doi.org/10.1146/annurev-marine-041923>, 2023.
- Griffies, S. M., Winton, M., Anderson, W. G., Benson, R., Delworth, T. L., Dufour, C. O., Dunne, J. P., Goddard, P., Morrison, A. K., Rosati, A., Wittenberg, A. T., Yin, J., and Zhang, R.: Impacts on ocean heat from transient mesoscale eddies in a hierarchy of climate models, *Journal of Climate*, 28, 952–977, <https://doi.org/10.1175/JCLI-D-14-00353.1>, 2015.



- Gu, Y., Katul, G. G., and Cassar, N.: Multiscale Temporal Variability of the Global Air-Sea CO₂ Flux Anomaly, *Journal of Geophysical Research: Biogeosciences*, 128, <https://doi.org/10.1029/2022JG006934>, 2023.
- Guo, Y. and Timmermans, M. L.: The Role of Ocean Mesoscale Variability in Air-Sea CO₂ Exchange: A Global Perspective, *Geophysical Research Letters*, 51, <https://doi.org/10.1029/2024GL108373>, 2024.
- Haarsma, R. J., Campos, E. J., Drijfhout, S., Hazeleger, W., and Severijns, C.: Impacts of interruption of the Agulhas leakage on the tropical Atlantic in coupled ocean-atmosphere simulations, *Climate Dynamics*, 36, 989–1003, <https://doi.org/10.1007/s00382-009-0692-7>, 2011.
- 395 He, Q., Zhan, W., Cai, S., Du, Y., Chen, Z., Tang, S., and Zhan, H.: Enhancing impacts of mesoscale eddies on Southern Ocean temperature variability and extremes, *Proceedings of the National Academy of Sciences of the United States of America*, 120, <https://doi.org/10.1073/pnas.2302292120>, 2023.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G. D., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 400 Hewitt, H., Fox-Kemper, B., Pearson, B., Roberts, M., and Klocke, D.: The small scales of the ocean may hold the key to surprises, <https://doi.org/10.1038/s41558-022-01383-9>, 2022.
- Hogg, A. M. C., Meredith, M. P., Chambers, D. P., Abrahamsen, E. P., Hughes, C. W., and Morrison, A. K.: Recent trends in the Southern Ocean eddy field, *Journal of Geophysical Research: Oceans*, 120, 257–267, <https://doi.org/10.1002/2014JC010470>, 2015.
- Ilyina, T., Six, K. D., Segschneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations, *Journal of Advances in Modeling Earth Systems*, 5, 287–315, <https://doi.org/10.1029/2012MS000178>, 2013.
- 410 Jones, C. D., Arora, V., Friedlingstein, P., Bopp, L., Brovkin, V., Dunne, J., Graven, H., Hoffman, F., Ilyina, T., John, J. G., Jung, M., Kawamiya, M., Koven, C., Pongratz, J., Raddatz, T., Randerson, J. T., and Zaehle, S.: C4MIP-The Coupled Climate-Carbon Cycle Model Intercomparison Project: Experimental protocol for CMIP6, *Geoscientific Model Development*, 9, 2853–2880, <https://doi.org/10.5194/gmd-9-2853-2016>, 2016.
- 415 Jones, E. M., Hoppema, M., Strass, V., Hauck, J., Salt, L., Ossebaar, S., Klaas, C., van Heuven, S. M., Wolf-Gladrow, D., Stöven, T., and de Baar, H. J.: Mesoscale features create hotspots of carbon uptake in the Antarctic Circumpolar Current, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 138, 39–51, <https://doi.org/10.1016/j.dsr2.2015.10.006>, 2017.
- Keppler, L., Eddebbar, Y. A., Gille, S. T., Guisewhite, N., Mazloff, M. R., Tamsitt, V., Verdy, A., and Talley, L. D.: Effects of Mesoscale Eddies on Southern Ocean Biogeochemistry, *AGU Advances*, 5, <https://doi.org/10.1029/2024AV001355>, 2024.
- 420 Kim, D., Lee, S. E., Cho, S., Kang, D. J., Park, G. H., and Kang, S. K.: Mesoscale eddy effects on sea-air CO₂ fluxes in the northern Philippine Sea, *Frontiers in Marine Science*, 9, <https://doi.org/10.3389/fmars.2022.970678>, 2022.
- Korn, P.: Formulation of an unstructured grid model for global ocean dynamics, *Journal of Computational Physics*, 339, 525–552, <https://doi.org/10.1016/j.jcp.2017.03.009>, 2017.
- Korn, P., Brüggemann, N., Jungclaus, J. H., Lorenz, S. J., Gutjahr, O., Haak, H., Linardakis, L., Mehlmann, C., Mikolajewicz, U., Notz, D., Putrasahan, D. A., Singh, V., von Storch, J. S., Zhu, X., and Marotzke, J.: ICON-O: The Ocean Component of the ICON Earth
- 425



- System Model—Global Simulation Characteristics and Local Telescoping Capability, *Journal of Advances in Modeling Earth Systems*, 14, <https://doi.org/10.1029/2021MS002952>, 2022.
- Landschützer, P., Gruber, N., and Bakker, D. C.: Decadal variations and trends of the global ocean carbon sink, *Global Biogeochemical Cycles*, 30, 1396–1417, <https://doi.org/10.1002/2015GB005359>, 2016.
- 430 Lerner, P., Romanou, A., Kelley, M., Romanski, J., Ruedy, R., and Russell, G.: Drivers of Air-Sea CO₂ Flux Seasonality and its Long-Term Changes in the NASA-GISS Model CMIP6 Submission, *Journal of Advances in Modeling Earth Systems*, 13, <https://doi.org/10.1029/2019MS002028>, 2021.
- Li, X., Gan, B., Zhang, Z., Cao, Z., Qiu, B., Chen, Z., and Wu, L.: Oceanic uptake of CO₂ enhanced by mesoscale eddies, Tech. rep., <https://www.science.org>, 2025.
- 435 Lévy, M., Couespel, D., Haëck, C., Keerthi, M. G., Mangolte, I., and Prend, C. J.: The Impact of Fine-Scale Currents on Biogeochemical Cycles in a Changing Ocean, <https://doi.org/10.1146/annurev-marine-020723>, 2023.
- Martínez-Moreno, J., Hogg, A. M. C., England, M. H., Constantinou, N. C., Kiss, A. E., and Morrison, A. K.: Global changes in oceanic mesoscale currents over the satellite altimetry record, *Nature Climate Change*, 11, 397–403, <https://doi.org/10.1038/s41558-021-01006-9>, 2021.
- 440 Mason, E., Pascual, A., Gaube, P., Ruiz, S., Pelegrí, J. L., and Delepoulle, A.: Subregional characterization of mesoscale eddies across the Brazil-Malvinas Confluence, *Journal of Geophysical Research: Oceans*, 122, 3329–3357, <https://doi.org/10.1002/2016JC012611>, 2017.
- McGillicuddy, D. J.: Mechanisms of Physical-Biological-Biogeochemical Interaction at the Oceanic Mesoscale, *Annual Review of Marine Science*, 8, 125–159, <https://doi.org/10.1146/annurev-marine-010814-015606>, 2016.
- Moreau, S., Penna, A. D., Llort, J., Patel, R., Langlais, C., Boyd, P. W., Matear, R. J., Phillips, H. E., Trull, T. W., Tilbrook, B., Lenton,
- 445 A., and Strutton, P. G.: Eddy-induced carbon transport across the Antarctic Circumpolar Current, *Global Biogeochemical Cycles*, 31, 1368–1386, <https://doi.org/10.1002/2017GB005669>, 2017.
- Morrison, A. K., Saenko, O. A., Hogg, A. M. C., and Spence, P.: The role of vertical eddy flux in Southern Ocean heat uptake, *Geophysical Research Letters*, 40, 5445–5450, <https://doi.org/10.1002/2013GL057706>, 2013.
- Morrison, A. K., Griffies, S. M., Winton, M., Anderson, W. G., and Sarmiento, J. L.: Mechanisms of Southern Ocean heat uptake and
- 450 transport in a global eddying climate model, *Journal of Climate*, 29, 2059–2075, <https://doi.org/10.1175/JCLI-D-15-0579.1>, 2016.
- Okubo, A.: Horizontal dispersion of floatable particles in the vicinity of velocity singularities such as convergences*, 1970.
- Orselli, I. B., Goyet, C., Kerr, R., de Azevedo, J. L., Araujo, M., Galdino, F., Touratier, F., and Garcia, C. A.: The effect of Agulhas eddies on absorption and transport of anthropogenic carbon in the South Atlantic Ocean, *Climate*, 7, <https://doi.org/10.3390/CLI7060084>, 2019a.
- Orselli, I. B., Kerr, R., Azevedo, J. L., Galdino, F., Araujo, M., and Garcia, C. A.: The sea-air CO₂ net fluxes in the South Atlantic Ocean
- 455 and the role played by Agulhas eddies, *Progress in Oceanography*, 170, 40–52, <https://doi.org/10.1016/j.pocean.2018.10.006>, 2019b.
- Patel, R. S., Llort, J., Strutton, P. G., Phillips, H. E., Moreau, S., Pardo, P. C., and Lenton, A.: The Biogeochemical Structure of Southern Ocean Mesoscale Eddies, *Journal of Geophysical Research: Oceans*, 125, <https://doi.org/10.1029/2020JC016115>, 2020.
- Petersen, M. R., Williams, S. J., Maltrud, M. E., Hecht, M. W., and Hamann, B.: A three-dimensional eddy census of a high-resolution global ocean simulation, *Journal of Geophysical Research: Oceans*, 118, 1759–1774, <https://doi.org/10.1002/jgrc.20155>, 2013.
- 460 Pezzi, L. P., de Souza, R. B., Santini, M. F., Miller, A. J., Carvalho, J. T., Parise, C. K., Quadro, M. F., Rosa, E. B., Justino, F., Sutil, U. A., Cabrera, M. J., Babanin, A. V., Voermans, J., Nascimento, E. L., Alves, R. C., Munchow, G. B., and Rubert, J.: Oceanic eddy-induced modifications to air–sea heat and CO₂ fluxes in the Brazil-Malvinas Confluence, *Scientific Reports*, 11, <https://doi.org/10.1038/s41598-021-89985-9>, 2021.



- Putrasahan, D. A., Gutjahr, O., Haak, H., Jungclaus, J. H., Lohmann, K., Roberts, M. J., and von Storch, J. S.: Effect of Resolving Ocean
465 Eddies on the Transient Response of Global Mean Surface Temperature to Abrupt 4xCO₂ Forcing, *Geophysical Research Letters*, 48,
<https://doi.org/10.1029/2020GL092049>, 2021.
- Richardson, P. L.: Agulhas leakage into the Atlantic estimated with subsurface floats and surface drifters, *Deep-Sea Research Part I: Oceanographic Research Papers*, 54, 1361–1389, <https://doi.org/10.1016/j.dsr.2007.04.010>, 2007.
- Rintoul, S. R. and Sokolov, S.: Baroclinic transport variability of the Antarctic Circumpolar Current south of Australia (WOCE repeat section
470 SR3), *Journal of Geophysical Research: Oceans*, 106, 2815–2832, <https://doi.org/10.1029/2000jc900107>, 2001.
- Sarmiento, J. L. and Gruber, N.: *Ocean biogeochemical dynamics*, in: *Ocean biogeochemical dynamics*, Princeton university press, 2006.
- Schütte, F., Karstensen, J., Krahmann, G., Hauss, H., Fiedler, B., Brandt, P., Visbeck, M., and Körtzinger, A.: Characterization of "dead-zone"
eddies in the eastern tropical North Atlantic, *Biogeosciences*, 13, 5865–5881, <https://doi.org/10.5194/bg-13-5865-2016>, 2016.
- Six, K. D. and Maier-Reimer, E.: Effects of plankton dynamics on seasonal carbon fluxes in an ocean general circulation model, *Global*
475 *Biogeochemical Cycles*, 10, 559–583, <https://doi.org/10.1029/96GB02561>, 1996.
- Song, H., Marshall, J., Munro, D. R., Dutkiewicz, S., Sweeney, C., McGillicuddy, D. J., and Hausmann, U.: Mesoscale modulation of air-sea
CO₂ flux in Drake Passage, *Journal of Geophysical Research: Oceans*, 121, 6635–6649, <https://doi.org/10.1002/2016JC011714>, 2016.
- Souza, R., Pezzi, L., Swart, S., Oliveira, F., and Santini, M.: Air-sea interactions over eddies in the Brazil-malvinas confluence, *Remote*
Sensing, 13, <https://doi.org/10.3390/rs13071335>, 2021.
- 480 Su, J., Strutton, P. G., and Schallenberg, C.: The subsurface biological structure of Southern Ocean eddies revealed by BGC-Argo floats,
Journal of Marine Systems, 220, <https://doi.org/10.1016/j.jmarsys.2021.103569>, 2021.
- Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., and Sutherland, S. C.: Seasonal variation of CO₂ and nutrients in the high-
latitude surface oceans: A comparative study, *Global Biogeochemical Cycles*, 7, 843–878, <https://doi.org/10.1029/93GB02263>, 1993.
- Talley, L. D., Pickard, G. L., Emery, W. J., and Swift, J. H.: Chapter 4 - Typical Distributions of Water Characteristics, in: *Descriptive Physical*
485 *Oceanography (Sixth Edition)*, edited by Talley, L. D., Pickard, G. L., Emery, W. J., and Swift, J. H., pp. 67–110, Academic Press, Boston,
sixth edition edn., ISBN 978-0-7506-4552-2, <https://doi.org/https://doi.org/10.1016/B978-0-7506-4552-2.10004-6>, 2011.
- van Westen, R. M. and Dijkstra, H. A.: Ocean eddies strongly affect global mean sea-level projections, *Science Advances*, 7, eabf1674, 2021.
- von Storch, J. S., Haak, H., Hertwig, E., and Fast, I.: Vertical heat and salt fluxes due to resolved and parameterized meso-scale Eddies,
Ocean Modelling, 108, 1–19, <https://doi.org/10.1016/j.ocemod.2016.10.001>, 2016.
- 490 Weiss, J.: *The dynamics of enstrophy transfer in two-dimensional hydrodynamics**, 1991.
- Zhang, Z., Wang, W., and Qiu, B.: Oceanic mass transport by mesoscale eddies, *Science*, 345, 322–324,
<https://doi.org/10.1126/science.1252418>, 2014.