

Balloon-borne Stratospheric Vertical Profiling of Carbonyl Sulfide and Evaluation of Ozone Scrubbing Materials

5 Alessandro Zanchetta¹, Steven van Heuven¹, Joram Hooghiem^{1,6}, Rigel Kivi³, Thomas Laemmel⁴, Michel Ramonet⁴, Markus Leuenberger⁵, Peter Nyfeler⁵, Sophie L. Baartman⁶, Maarten Krol^{6,7}, and Huilin Chen^{1,2}

¹ Center for Isotope Research (CIO), Energy and Sustainability Research Institute Groningen (ESRIG), University of Groningen, Groningen, the Netherlands

10 ² Joint International Research Laboratory of Atmospheric and Earth System Sciences, School of Atmospheric Sciences, Nanjing University, Nanjing, China

³ Space and Earth Observation Centre, Finnish Meteorological Institute (FMI), Sodankylä, Finland

⁴ Laboratoire des Sciences du Climat et l'Environnement (LSCE), CEA - CNRS - UVSQ - University Paris-Saclay, Gif sur Yvette, France

15 ⁵ Climate and environmental Physics, Physics Institute and Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

⁶ Meteorology and Air Quality, Wageningen University & Research, Wageningen, the Netherlands

⁷ Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, the Netherlands

20 Corresponding to Huilin.Chen@nju.edu.cn or Huilin.Chen@rug.nl

Abstract

Carbonyl sulfide (COS) is a low abundance atmospheric trace gas that has a tropospheric lifetime of 2-2.5 years, allowing it to reach the stratosphere, where it undergoes photolysis and reactions with •OH and O• radicals, generating precursors of stratospheric aerosols. Vertical profiling of COS has rarely been realised, especially for stratospheric observations. In this study, we introduce a new technique for continuous and discrete vertical profiling of COS based on the analysis of air samples collected by AirCore, the LIghtweight Stratospheric Air (LISA) sampler, and its scaled-up version BigLISA, in three campaigns in Trainou (2019), Kiruna (2021) and Sodankylä (2023) using a Quantum Cascade Laser Spectrometer (QCLS). To eliminate potential COS measurement biases, we have investigated the efficiency of different scrubbers based on cotton and squalene for removing ozone (O₃) and assessed their potential impacts on COS measurement. Furthermore, we examined the influence of different inlet configurations and O₃ scrubbers on the retrieved COS profiles, and found no significant impact within the uncertainties. We found that the differences with the averaged profiles obtained from the Atmospheric Chemistry Experiment – Fourier Transform Spectrometer (ACE-FTS) and the measured AirCore profiles and LISA samples were less than 10% (± 50 ppt) at both mid and polar latitudes. Differences between our observations and COS observations from the *S*Pectromètre *I*nfraRouge *d'*Absorption à *L*asers *E*mbarqués (SPIRALE) ranged from 10% to 15%, with both methods

35

showing similar COS trends over altitude. Moreover, we found squalene-based scrubbers to be suitable for quantitative O₃ removal. Both AirCore and the LISA samplers are lightweight and suitable for routine balloon-borne COS profiling, providing useful observations for stratospheric research and validation of COS retrievals from remote sensing techniques.

1. Introduction

40 Carbonyl sulfide (COS, also referred at as OCS) is an odorless and colorless gas species (Ferm, 1957). It is the most abundant sulfur-containing gas species in the atmosphere, with a tropospheric mole fraction of 350-500 parts per trillion (ppt, pmol/mol) in the unpolluted free troposphere (Berry et al., 2013; Remaud et al., 2023). It has been suggested as a proxy to partition photosynthetic uptake of CO₂ from respiration, to improve the quantification of carbon fluxes between atmosphere and vegetation (Campbell et al., 2008; Montzka et al., 2007; Sandoval-Soto et al., 2005; Stimler et al., 2009; Whelan et al., 2018).
45 Given its relatively long tropospheric lifetime of 2-2.5 years (Ma et al., 2021; Montzka et al., 2007; Remaud et al., 2023), COS can reach the stratosphere, where it is converted to CO₂ and sulfur dioxide (SO₂), a precursor of stratospheric aerosols, by photolysis and reactions with •OH and O• radicals (Brühl et al., 2012; Chin and Davis, 1995; Krysztofiak et al., 2015). COS is considered to likely be the largest contributor to stratospheric sulfur aerosols during volcanic quiescent periods (Brühl et al., 2012; Crutzen, 1976; Kremser et al., 2016).

50

Currently, observations of stratospheric COS vertical profiles and/or total columns are performed by ground- and satellite-based remote sensing (Barkley et al., 2008; Bernath, 2005; Toon et al., 2018) and by deploying balloon-borne spectrometers (Krysztofiak et al., 2015; Toon et al., 2018). The collection of air samples for COS analysis was only carried out sporadically or only in the upper troposphere/lowermost stratosphere (10-12 km) (Engel and Schmidt, 1994; Karu et al., 2023). In this
55 paper, we present new techniques to collect continuous and discrete stratospheric air samples, based on the balloon-borne AirCore (Karion et al., 2010) and the LIghtweight Stratospheric Air (LISA) (Hooghiem et al., 2018) and BigLISA samplers, respectively, paired with a Quantum-Cascade Laser Spectrometer (QCLS, Aerodyne Research Inc., MA, USA, model TILDAS-CS) for COS measurements (Kooijmans et al., 2016; Stimler et al., 2009). These methods allow analysis of collected air samples with minimal preparation and treatment, which reduces risks of contamination during sampling and storage.

60

Possible impacts of stratospheric ozone (O₃) on collected air samples for COS observations have been reported in previous studies (Engel and Schmidt, 1994), as have the impacts of pollution-induced tropospheric O₃ on other reduced sulfur compounds, such as dimethyl sulfide (DMS) or carbon disulfide (CS₂) (Andreae et al., 1993; Hofmann et al., 1992; Persson and Leck, 1994), known precursors of atmospheric COS (Whelan et al., 2018). O₃, a strong oxidant, may in fact react with
65 reduced sulfur compounds causing a variable, yet possibly significant reduction in their abundance (Andreae et al., 1993; Engel and Schmidt, 1994; Hofmann et al., 1992; Persson and Leck, 1994). Recent tests with a limit of detection around 50 ppt, however, reported no influence of O₃ on neither DMS, nor CS₂, but found positive interference on measurements of Volatile

Organic Compounds (VOCs) containing carbonyl groups (Ernle et al., 2023). Since stratospheric O₃ is more abundant than pollution-induced tropospheric O₃ (in particular between 15 and 35 km of altitude, where O₃ mixing ratios may reach roughly 10 ppm) (World Meteorological Organization (WMO), 1999), its impact on air samples for COS observations may be significant. Therefore, we have investigated different techniques to remove O₃ before sampling, and assessed their potential impacts on the mole fractions of COS and other trace gases. In particular, we have investigated different O₃ scrubbers and their scrubbing efficacy and effect on COS, also by deploying different inlets on the aforementioned samplers. Furthermore, we show a comparison of measured continuous and discrete COS samples with previous observations. A particular focus is set on a cross-validation comparison with SPIRALE's in situ spectrometry (Krysztofiak et al., 2015) and ACE-FTS remote sensing COS observations (Bernath, 2005; Glatthor et al., 2017; Schmidt et al., 2024; Velazco et al., 2011).

2. Materials and methods

2.1 Samplers

The three different samplers deployed to collect stratospheric air will be presented in this section. All instruments flew under weather balloons that typically reached altitudes of 30 to 35 km. The presented data was collected in different campaigns, namely the RINGO campaign in Trainou (TRN, France, 2019), the HEMERA campaign (Schuck et al., 2025) in Kiruna (KRN, Sweden, 2021) and the OSTRICH campaign at the Fourier Transform Spectrometer (FTS) site (Kivi and Heikkinen, 2016) in Sodankylä (SOD, Finland, 2023). An overview of campaigns and samplers is reported in Table 1. Air samples collected by all devices were analysed on a QCLS, which will be described in Sect. 2.2.

2.1.1 AirCore

The AirCore sampler was first introduced by Karion et al. (2010) to retrieve CO₂ and CH₄ vertical profiles. It consists of a long, stainless-steel tube usually shaped as a coil, internally coated with Sulfinert® to prevent reactions or adsorption of gas species with the tube walls (Karion et al., 2010; Membrive et al., 2017). When used to retrieve vertical profiles, AirCore sampling is realised passively and continuously along the coil at ambient temperature (Karion et al., 2010; Membrive et al., 2017; Wagenhäuser et al., 2021), differently from the discrete samples that can be collected by cryosamplers that, however, need to be cooled down to 27 K with liquid neon before flight (Laube et al., 2010; Schmidt et al., 1987). Before the flight, the coil is filled with a known gas mixture, which will later help identifying the starting point of the AirCore profiles during analysis. During ascent, the coil empties through one open end due to decreasing ambient pressure. After the balloon bursts or is cut away, the instrument collects air during descent while ambient pressure is increasing, without using a pump. Knowing sampling pressure and temperature, and assuming pressure equilibrium during the filling process, each aliquot of moles of air can be calculated for each sampling altitude interval. These can be associated to the aliquot of analysed moles of air in the coil, allowing the retrieval of vertical profiles (Karion et al., 2010; Membrive et al., 2017; Tans, 2022); however, the selected start and end point of each analysis, air mixing inside the coil, sample loss, and general fill dynamics may all be causes for deviations

from this approximation (Membrive et al., 2017; Tans, 2022; Wagenhäuser et al., 2021). Detailed discussions of fill dynamics and uncertainties treatment can be found in Tans (2022) and Membrive et al. (2017), respectively. In this study, we followed the approach described by Membrive et al. (2017) to retrieve the altitude mapping and the relative uncertainties along the vertical profiles, with one exception, as follows.

As reported in Table 1, flight SOD3 included a double-sided sampling AirCore, property of the University of Bern, which was flown and analysed by our group. One half of this 200 m long AirCore was equipped with an oxygen-spiking system, programmed to inject 5 minuscule shots of O₂ in the coil as altitude markers at 21045, 17005, 11837, 7870 and 4592 m. Although our QCLS is not capable of measuring O₂, these injections were visible as COS anomalies along the profile (see Sect. 4.1). Therefore, the altitude mapping for this AirCore was realised by matching the COS spikes with the reported spiking altitudes. Moreover, the cotton scrubber installed on one of the two inlets likely adsorbed water (H₂O) before the ascent phase, which was then taken in the AirCore at the beginning of the descent, mixing tropospheric H₂O with stratospheric air at the highest altitude. This required a dilution correction, followed by a matrix effect correction inferred from the correlation of different species with H₂O mole fraction.

At the altitude ceiling of the balloon flight, the AirCore's coil still contains part of the fill gas. This remaining fill gas can mix with the sampled air at the top of the profile. Similarly, air from the lowest part of the sampled profile mix with the gas used to push the air out of the coil during analysis. Therefore, the highest and the lowest parts of the profiles are flagged and are not used for further analysis.

For the double-sided AirCore deployed on flight SOD3, fill gas flagging starts at lower altitude compared to other AirCores. Given the design of this AirCore, air is collected from both ends of the coil, allowing the simultaneous sampling of two profiles. Therefore, the top of the profiles and the remaining fill gas are located at the centre of the coil and not at one of its ends. For all other AirCores, the top of the profile is the first part to be analysed, while for this sampler it will have to travel 100 m through the coil before reaching the analyser. Thus, during analysis the top of the profile and the remaining fill gas have a longer time to get mixed compared to other AirCores. Moreover, the resulting gas mixture may also experience a stronger smearing effect due to its path through the coil. Altogether, this results in a larger portion of the profile that cannot be considered for analysis compared to other AirCores.

Some AirCores experienced COS contamination due to specific design features (e.g., differential pressure sensors along the coil or glue connections). Consequently, the contaminated COS mole fractions have been removed and are shown as gaps. The causes of contamination are discussed in Sect. 4.1.

130 2.1.2 LISA sampler

The LISA sampler used in this study is a further miniaturised (55 L package size) and light-weight (2.9 kg) version of the original sampler developed by Hooghiem et al. (2018). The instrument is battery-powered and is controlled by a microcontroller, which also logs GPS, pressures, temperatures, and general instrument status. Differently from AirCore, LISA actively pumps ambient air into four different 2.5L multi-layer foil (MLF) bags (type 30228-U, Supelco Inc., USA) through a custom-made manifold. The sampling is performed during the ascent phase of the flight, since the vertical speed is slower than during descent and this allows for a higher vertical resolution of the vertical sampling. The valves of the MLF bags are opened and closed by servos. The sampling pressure intervals for each bag are programmed before the flight and limited to an absolute sample pressure of 280 hPa to prevent bag burst after sampling. This is necessary because the ambient pressure continues to decrease during the remainder of the ascent, reaching about 10 hPa at 30 km altitude. The sampling pressures and the derived sampling altitude intervals and sample volumes are reported in Table S1 in the Supplement. The mid-points of sample collections are calculated considering the dependency of pump performance on the ambient pressure and the bag filling status, as described by Hooghiem et al. (2018).

Unusually high COS mole fractions were measured in laboratory tests and in some collected samples, which we speculate are due to outgassing from polymers (Lee and Brimblecombe, 2016). Therefore, during the SOD campaign in 2023, we performed pre-conditioning of the MLF bags differently from what Hooghiem et al. (2018) described. Before each flight, MLF bags were not only filled and vacuumed with purified N₂, but filled and vacuumed repeatedly with air from a cylinder of synthetic air mixed with low mole fractions of CH₄, CO₂ and CO, which was meant to simulate stratospheric air conditions and contained 0 ppb N₂O and 0 ppt COS. This was done to prevent outgassing of different gas species, and in particular COS, from the polymers composing the MLF bags. The gas mixture used to flush the bags was measured on the QCLS before filling the bags and when it was pumped out from them, allowing a control of potential contaminations under stratospheric sampling conditions. After the LISA sampler was recovered and brought back to the laboratory in the field, LISA air samples were transferred to glass flasks and stored for later analyses of COS and other trace gas species. Here we present the analysis results of the air samples left in the sampling bags, when present, directly after the sample transfer from the MLF bags to glass flasks (these latter were not analysed on the QCLS). The leftover volume of one of these samples was insufficient for analysis (SOD3 – L4), while two others showed unusually high mole fractions for several of the analysed gas species (SOD2 – L4, SOD5 – L3), in spite of the pre-conditioning of these bags before flight, most likely due to contamination with tropospheric air during sampling and/or while transferring the samples. Based on the insufficient sample volume and/or the anomalies in the measured gases, these three samples were labelled as outliers and will not be presented in this work.

160 2.1.3 BigLISA

BigLISA is a larger-volume and functionally improved variant of the LISA sampler, with a mass of roughly 12 kg. On the HEMERA missions in KRN, two independently operating BigLISA samplers were flown in a single enclosure on a high-payload balloon (Schuck et al., 2025). Each BigLISA unit consists of a central box containing electronics and pneumatic components, and 6 externally mounted 10 L MLF bags (type 30229-U, Supelco Inc., USA). The BigLISA pneumatic hardware
165 consists of a two-stage pump, a flow-reversing valve system that allows purging, a manifold and 6 closable MLF bags, all powered, monitored and controlled from a single control board.

Two stage-pumping is attained by connecting the four heads of two double-headed pumps (model NMP830.1.2KPDC-B HP, KNF, Germany) in a three to one configuration. This attains a high flow rate and a high compression factor. Laboratory tests
170 confirmed the potential to reach a compression ratio of 25 at low ambient pressures. However, potentially due to unfavourably dimensioned connecting tubing, the compression ratio obtained during flight ranged between 2 and 5 times over ambient pressure.

Operation and logging are provided by a custom-made Printed Circuit Board (PCB) running an ESP32 microcontroller
175 (Adafruit Industries, model HUZZAH32). Temperature is monitored at the pump heads, battery packs, power converters and within the larger BigLISA outer housing. In each pack, before and between pumping operations, the two 3-way valves are actuated as necessary to maintain in-pack temperature above 5 °C, preventing loss of battery capacity and conceivable pump stalling. Power for electronics, pumps and valves is provided by two packs (for redundancy) of eight Saft LSH14 Li-SOCl₂ cells.

180 Housing for the two BigLISA units was provided by a customised high strength but lightweight aluminium frame, that was able to withstand the high accelerations potentially experienced during parachute deployment. In this structure, the two BigLISA packs are mounted centrally, surrounded by the 6 + 6 MLF bags, which are individually suspended using tie-wraps on concentric stainless-steel wires. Protection from wind and radiation is provided by 1 mm thick aluminium sheeting. GPS receivers and sampling inlet pumping lines are led out radially at the top of the package.

185 During ascent, starting at 120 hPa (~15 km), bags were (re-)evacuated by the two-stage pump. This procedure removes conceivable traces of tropospheric air from the bags, and tests for plumbing integrity. From 30 hPa (~25 km), the manifold was flushed with ambient air to clear it of residual tropospheric air and water vapour. Just prior to collecting a sample, the respective bag would be repeatedly filled with a tiny amount of air and evacuated, to dilute any residual air in it. Sampling
190 took place during the descent, for as long as it took to reach 200 hPa of sample pressure, but never more than 1800 seconds (less for lower samples), and would stop when the next sample was due to be collected. The sampling altitude intervals are reported in Table S1 in the Supplement and are shown in Figure 2. The mid-points of sampling altitudes were calculated similarly to Hooghiem et al. (2018), considering the decreased pump performance when compression becomes necessary to

195 fill the bag. However, given that BigLISA sampled during descent, this effect was counteracted by the increase in ambient pressure during sampling. Therefore, the mid-points of BigLISA fall more towards the average of the sampling interval than the ones of LISA. A number of BigLISA samples (KRN - BL7, BL10, BL11 and BL12) showed unusually high mole fractions in multiple measured species. Similarly to LISA, these samples were labelled as outliers due to potential contamination, and therefore removed from the analysis.

200 **Table 1: Overview of the instruments deployed for COS sampling, the launch location, samplers' sizes and additional features, and the inlet features in different campaigns. The reported diameters (\varnothing) refer to the outer diameter of the tubing. The flight code helps identifying which instruments have been deployed on the same balloons, and is used to refer to these flights in the text.**

Location	Flight code	Flight date and takeoff time (UTC)	Apogee	Instrument	Instrument features	Inlet features
Trainou (FRA) 47°58' N 2°06' E	TRN1	17/06/2019 12:30	34.6 km	AirCore	23 m x \varnothing 8 mm + 46 m x \varnothing 4 mm V ~ 1600 cm ³	Free inlet
	TRN2	18/06/2019 8:00	32.6 km	AirCore	36 m x \varnothing 3/16" + 38 m x \varnothing 1/8" V ~ 830 cm ³	Mg(ClO ₄) ₂ dryer
	TRN3	18/06/2019 10:39	34.9 km	AirCore	23 m x \varnothing 8 mm + 46 m x \varnothing 4 mm V ~ 1600 cm ³	Free inlet
	TRN4	20/06/2019 07:07	32.3 km	AirCore	36 m x \varnothing 3/16" + 38 m x \varnothing 1/8" V ~ 830 cm ³	Free inlet
Kiruna (SWE) 67°53' N 21°04' E	KRN	12/08/2021 21:18	33.1 km	AirCore	37 m x \varnothing 3/16" + 39 m x \varnothing 1/8" V ~ 860cm ³	Mg(ClO ₄) ₂ dryer
	KRN	12/08/2021 21:18	33.1 km	AirCore	36 m x \varnothing 3/16" + 38 m x \varnothing 1/8" V ~ 830 cm ³	Free inlet

	KRN	12/08/2021 21:18	33.1 km	BigLISA	12 x 10 L MLF-bags	Mg(ClO ₄) ₂ dryer
Sodankylä (FIN) 67°22' N 26°37' E	SOD1	02/08/2023 7:00	29.7 km	AirCore	40 m x Ø 1/4" + 58 m x Ø 1/8" V ~ 1400 cm ³ Differential pressure sensors	Mg(ClO ₄) ₂ dryer
	SOD2	02/08/2023 14:50	30.1 km	LISA	4 x 2.5 L MLF-bags, pre-conditioned	Mg(ClO ₄) ₂ dryer
	SOD3	05/08/2023 10:56	26.3 km	LISA	4 x 2.5 L MLF-bags, pre-conditioned	Free inlet
	SOD3	05/08/2023 10:56	26.3 km	AirCore	200 m x Ø 1.5 mm" V ~ 1400 cm ³ Double-sided sampling	One side free inlet, one side cotton-based O ₃ scrubber
	SOD4	06/08/2023 9:03	29.8 km	LISA	4 x 2.5 L MLF-bags, pre-conditioned	Cotton- based O ₃ scrubber
	SOD5	08/08/2023 6:17	29.3 km	AirCore	40 m x Ø 1/4" + 58 m x Ø 1/8" V ~ 1400 cm ³ Differential pressure sensors	Mg(ClO ₄) ₂ dryer and cotton-based O ₃ scrubber

2.2 Quantum Cascade Laser Spectrometer (QCLS)

The trace gas analyser used to perform COS measurements in all campaigns is a dual laser QCLS by Aerodyne Research Inc. (BillERICA, MA, USA), operating in the mid-infrared frequencies. This technique was firstly introduced for COS measurements by Stimler et al. (2009) and further developed by Kooijmans et al. (2016). The QCLS employed in this study has also been used in previous UAV- and aircraft-borne tropospheric active AirCore measurements for CH₄ and N₂O (Tong et al., 2023; Vinković et al., 2022), as well as in situ tropospheric COS observations (Zanchetta et al., 2023). The QCLS can measure CH₄, CO₂, N₂O, CO, COS, H₂O, and O₃ simultaneously. Its cavity is controlled at a temperature of 298 K and a pressure of ~66 hPa

(50 Torr). The QCLS measures at a constant mass flow of 50 mL min^{-1} and the measured data is output at 1 Hz. For COS, the precision (1σ) falls usually between 15 and 25 ppt at 1 Hz, depending on the laboratory conditions (e.g., ambient temperature stability). The cell of the QCLS has a volume of 150 mL, which at 50 Torr corresponds to an effective cavity volume of ~ 10 mL. The spatial resolution of AirCore measurements in this configuration is roughly 2000 m at 28 km, 300 m at 15 km and 200 m at 10 km altitude, which resembles the resolution ranges presented by Membrive et al. (2017). The QCLS is controlled with a custom-made frontend, operated via a dedicated software, which allows switching between different inlets without causing changes in pressure through the system. Overall, the instrument achieves a precision better than 0.6 ppb for CH_4 , 0.2 ppm for CO_2 , 0.12 ppb for N_2O , 1 ppb for CO, 20 ppt for COS, 20 ppm for H_2O , and 100 ppb for O_3 .


As reported in Zanchetta et al. (2023), field standard cylinders are calibrated against NOAA standards (NOAA-2004 COS scale) in the laboratory before and after each measurement period to test for drift in the molar fraction of gas species. The COS mole fraction measurements of nine cylinders are available, and five cylinders changed by less than 2.5 ppt yr^{-1} , two cylinders decreased by $\sim 10 \text{ ppt yr}^{-1}$ and two cylinders decreased by $\sim 30 \text{ ppt yr}^{-1}$. The four cylinders that drifted more than 10 ppt yr^{-1} were not used as reference cylinders in the data processing. All of the cylinders were uncoated aluminium cylinders, which, according to experience at NOAA, are more prone to COS mole fraction drift than Aculife-treated aluminium cylinders. More details on the instrumental calibration, precision and stability can be found in Kooijmans et al. (2016).

2.3 Ozone scrubbing materials

O_3 is a reactive gas species that can be found at mole fractions up to about 8 ppm in the stratosphere, where it is formed by the interaction of atmospheric O_2 with UV radiation (Bernhard et al., 2023). O_3 is a strong oxidant and can react with other trace gases, including reduced sulfur compounds such as dimethyl sulfide (DMS) and carbon disulfide (CS_2) (Andreae et al., 1985; Hofmann et al., 1992; Persson and Leck, 1994). Moreover, the oxidation of DMS and CS_2 , indirect precursors of COS, was reported as a potential bias of tropospheric COS measurements obtained after cryogenic sampling (Hofmann et al., 1992). On top of this, the amount of COS was found to be lower if sampled in presence of O_3 (Engel and Schmidt, 1994). Therefore, a number of oxidant removal substances, such as manganese dioxide (MnO_2), cotton wadding, sodium carbonate (Na_2CO_3), potassium hydroxide (KOH) or a KI/glycerol/Vitex solution have been deployed to remove oxidants during cryogenic sampling, in both tropospheric and stratospheric applications (Andreae et al., 1985; Engel and Schmidt, 1994; Hofmann et al., 1992; Persson and Leck, 1994 and references therein). Among these, scrubbers based on cotton wadding were tested and proved to be effective for tropospheric cryogenic samples (Hofmann et al., 1992; Persson and Leck, 1994); however, only MnO_2 has been deployed for stratospheric applications (Engel and Schmidt, 1994). Therefore, we tested multiple substances for their O_3 scrubbing efficiency and their influence on the mole fractions of the analysed gases. During the first laboratory tests, we found that MnO_2 interacted strongly with multiple tracers. In order to find a suitable O_3 scrubber that would perform well under stratospheric conditions, we conducted a series of experiments, as described below.

240

We designed an experimental setup (Figure 1) that obtains an air mixture with high O₃ at low pressure and low temperature, to assess the performance of O₃ scrubbers under stratospheric air conditions. Air from a laboratory-made synthetic air (~79% N₂, ~21% O₂) cylinder containing mole fractions of the measurable tracers, close to stratospheric conditions (1435.7 ppb CH₄, 0 ppb N₂O, 392.67 ppm CO₂, 0 ppt COS, and 0 ppb CO), was flowed through a custom-made Y-shaped quartz tube, which
245 passed through a UV lamp that served as an O₃ generator. With a QCLS-controlled mass flow of 50 mL min⁻¹ and a path length of about 6 inches (~ 15.2 cm) through UV radiation, it was possible to generate up to 3500 – 4000 ppb O₃. The air could be directed to each side alternately using a three-way valve, allowing measurements of the air after O₃ generation, with or without an oxidant scrubber. The tube passed through a polystyrene box, where 193 K freeze packs could be inserted to simulate low- to mid-stratospheric temperatures (air temperature could reach as low as roughly 213 K). The flow and pressure of the air in
250 the sampling lines were controlled by the QCLS frontend, with the mass flow typically maintained at 50 mL min⁻¹ and the pressure reaching as low as 250 hPa. A bypass channel directly connected to the QCLS was also available for reference measurements.



255 **Figure 1: The experimental setup for the O₃ scrubber testing.** The black circles represent three-way valves. Apart from the quartz tube, all connections were realised with stainless steel tubing.

The initial tests focused only on the performance of O₃ scrubbers based on cotton, supported by available research on the consumption of O₃ by contact with fabrics (Andreae et al., 1993; Coleman et al., 2008; Hofmann et al., 1992; Persson and Leck, 1994). Surgical cotton wool (approximately 3 - 10 g of cotton) was inserted on one side of the Y-shaped quartz tube.
260 While cotton scrubbers seemed to work initially (Table 3), in later tests it was noticed that cotton would quickly lose efficiency, in particular after some storage time. While “new” cotton pads still exhibited some O₃ scrubbing capacity, the same pads, after storage (“old” cotton), showed almost no scrubbing ability.

It was observed that the efficiency of O₃ scrubbing changed significantly depending on whether nitrile gloves were used or not
265 while handling the cotton pads: the scrubbing efficiency increased when the pads were handled with bare hands. This led to the hypothesis that O₃ was mostly removed by reaction with skin oils rather than with the cotton itself. This hypothesis was

further supported by existing literature (Coffaro and Weisel, 2022; Coleman et al., 2008; Zhou et al., 2016). Following Coffaro and Weisel (2022), knowing that squalene, a triterpene, accounts for 12% of skin oils composition (Picardo et al., 2009), squalene-based scrubbers were also tested. Consequently, the potential impacts of the reaction between O₃ and squalene on other trace gases were investigated. The squalene-based O₃ scrubbers consisted of one drop of laboratory-quality squalene (Sigma-Aldrich, ≥ 98%) deposited with a Pasteur pipette on glass wool, which had been previously proven inert to the analysed trace gases.

We designed the final experimental setup with five possible configurations:

- a. No O₃ generation, air flowed through the control channel
- b. O₃ generation, air flowed through the control channel
- c. No O₃ generation, air flowed through the cotton/squalene scrubber
- d. O₃ generation, air flowed through the cotton/squalene scrubber
- e. Total bypass channel

The data from the experimental time series was selected and designated to the respective configuration. These configurations were then used as categorizations to perform an ANOVA test, which would eventually corroborate significant differences between each species' mole fraction, depending on the experimental configuration. The results are presented and discussed in the following paragraphs. The ANOVA test results on the effect of the squalene scrubbers on other tracers are reported in Sect. S3 in the Supplement.

The possible coincidental removal of COS by squalene and cotton was assessed just briefly with the same experimental setup described above, placing squalene in one channel and cotton in the other. For this test, a tropospheric compressed air mixture (2201.3 ppb CH₄, 338.9 ppb N₂O, 430.10 ppm CO₂, 747 ppt COS, and 180.6 ppb CO) was employed. The findings for COS and O₃ are presented at the end of the Sect. 3.2.

2.4 Datasets for COS profiles comparison and validation

For validation purposes, our COS profiles were compared to the SPIRALE results reported by Krysztofiak et al. (2015), as well as a selection from the Atmospheric Chemistry Experiment – Fourier Transform Spectrometer (ACE-FTS, v5.3) dataset (Bernath, 2005; Boone et al., 2023; Schmidt et al., 2024; Velazco et al., 2011).

2.4.1 SPIRALE

The in situ balloon-borne SPIRALE spectrometer presented by Krysztofiak et al. (2015) was deployed in two flights at polar latitudes (Kiruna, Sweden, 67°53' N 21°04' E) in 2009 and 2011. The resulting COS profiles cover altitudes between 14.3 – 21.6 km and 18.5 – 22.0 km, respectively. Given the different altitudinal resolution, both the measured AirCore profiles and

the SPIRALE results were averaged in 0.5 km bins to calculate the difference between COS mole fractions. A comparison
300 between AirCore and SPIRALE COS profiles is shown in Figure 7.

2.4.2 ACE-FTS

ACE-FTS is a satellite-borne spectrometer measuring altitude profile information for temperature, pressure, and mole fractions of several gas species (Bernath, 2005; Glatthor et al., 2017; Velazco et al., 2011), including COS, by sun occultation. Each profile of ACE-FTS contains 1 km resolution data from 0-150 km altitude. To have a comparable dataset with the measured
305 AirCore profiles, COS ACE-FTS observations collected between June and September in the 2012 – 2024 period were selected, with latitudes ranging between 45-49° N for TRN and 65-69° N for KRN and SOD, resulting in 502 and 1681 COS profiles, respectively. These selected profiles were then averaged for both latitudinal ranges. The COS profiles measured with AirCore in this study are averaged over 1 km intervals to obtain a comparable dataset. The resulting averaged profiles and their comparison with the observed AirCore profiles are shown in Figure 10.

310 3. Results

3.1 COS observations


Figure 2 shows the COS profiles measured from AirCores and the LISA/BigLISA samples of the campaigns reported in Table 1. For all campaigns, different inlets were deployed. Different inlet configurations are reported in Table 1 and in the figure captions.

315

The tropospheric COS mole fractions vary from flight to flight, ranging between about 400 to 500 ppt. The thermal tropopause height (World Meteorological Organization, 1957) was between about 10.6 and 11.5 km at mid latitudes (TRN, June 2019, Fig. 1a), and between 9.9 and 10.9 km at polar latitudes (KRN, August 2021, Fig. 1b and SOD, 2023, Fig. 1c), respectively (see Fig. S7-S15 in Sect. S4 of the supplement). The COS stratospheric sink is clearly noticeable for all campaigns above the
320 tropopause and is further discussed in Sect. 4.1. At polar latitudes, the COS mole fraction decreases from an all-campaigns average of 421 ± 26 ppt below 17 km to 180 ± 15 ppt between 20 and 22 km. At mid-latitudes, the mole fractions decrease from a campaign average of 488 ± 12 ppt below 17 km to 220 ± 59 ppt between 20 and 22 km.

As reported in Table 2, the BigLISA samples measured in KRN in 2021 show consistently higher mole fractions compared to
325 the AirCore profiles. By contrast, the LISA samples measured in SOD in general show good agreement with the continuous profiles. The largest difference was found in SOD3, between a collection of free-inlet LISA samples and an AirCore equipped with a cotton scrubber obtained from the same flight. The possible explanations for these differences are discussed in Sect. 4.1.2.

Measured COS profiles and samples during different campaigns



330

Figure 2: Measured COS AirCore (AC) profiles and LISA/BigLISA samples in different sampling campaigns.

Table 2: COS mole fraction difference between AirCore and (Big)LISA samples that flew on the same days. The AirCore COS mole fraction is calculated as the COS average over (Big)LISA's sampling altitude range.

(Big)LISA sample name	Altitude range (km)	Δ COS (ppt) AirCore – (Big)LISA	Inlets features	Average Δ COS (ppt)
KRN – BL3	20.1 – 22.5	-48	AirCore: Mg(ClO ₄) ₂ dryer BigLISA: Mg(ClO ₄) ₂ dryer	-80 ± 50
KRN – BL9	18.0 – 19.9	-125		
KRN – BL4	17.7 – 19.0	-133		
KRN – BL5	15.9 – 16.4	-78		
KRN – BL6	14.5 – 14.7	-16		
KRN – BL3	20.1 – 22.5	-52	AirCore: free inlet BigLISA: Mg(ClO ₄) ₂ dryer	-53 ± 20
KRN – BL9	18.0 – 19.9	-68		
KRN – BL4	17.7 – 19.0	-70		
KRN – BL5	15.9 – 16.4	-53		
KRN – BL6	14.5 – 14.7	-21		

SOD1 – L1	14.1 – 14.6	-32	AirCore: Mg(ClO ₄) ₂ dryer LISA: Mg(ClO ₄) ₂ dryer	-16 ± 24
SOD1 – L2	16.8 – 18.0	-27		
SOD1 - L3	21.5 – 22.7	11		
SOD3 – L1	14.2 – 14.6	17	AirCore: free inlet LISA: free inlet	31 ± 20
SOD3 – L2	17.0 – 17.9	46		
SOD3 – L3	21.6 – 23.5	No AirCore data		
SOD3 – L1	14.2 – 14.6	25	AirCore: cotton, without dryer LISA: free inlet	/
SOD3 – L2	17.0 – 17.9	No AirCore data		
SOD3 – L3	21.6 – 23.5	No AirCore data		

3.2 Tests on O₃ scrubbing materials

335 As reported in Table 3, the initial tests on cotton O₃ scrubbers (performed in 2023) resulted in the quantitative removal of about 3.5 ppm O₃, lasting up to roughly 2.5 to 40 L of air with a flow rate of 50 mL min⁻¹, depending on the amount of cotton used. However, months later, cotton pads from the same bag lost their scrubbing capacity after only about 0.05 to 0.75 L of air, soon showing a clear drop in performance and allowing progressively more O₃ to pass through. As explained in Sect. 2.3, it was observed that the efficiency of O₃ removal was significantly higher when the cotton was handled with bare hands and eventually led to squalene-based scrubbers.

340

We found that one drop of squalene from a Pasteur pipette on glass wool was sufficient to quantitatively remove O₃ up to roughly 12 L of air containing 3.5 ppm O₃ at 50 mL min⁻¹, without showing any sign of O₃ breakthrough (and we speculate it could have possibly scrubbed for an even longer duration). When no O₃ was generated, no significant differences were observed for COS, indicating that neither cotton, nor squalene caused COS contaminations when interacting with the air samples.

345


Table 3: Summary of the first experiments regarding O₃ scrubbers, which eventually led to the choice of focusing on squalene-based scrubbers for the laboratory tests.

Scrubber type	Capacity of scrubbing ~3.5 ppm O ₃ (air volume)	Influence on O ₃	Influence on COS
“Old” cotton, handled with gloves (1 cotton pad, ~ 0.3 g of cotton)	~ 50 – 750 mL	Quantitative removal at the beginning, followed by progressive O ₃ breakthrough	Not significant, as long as the scrubber worked

“Old” cotton, handled with bare hands (1 cotton pad, ~ 0.3 g of cotton)	≥ 5000 mL	Quantitative removal	Not significant
“New” cotton, handled with gloves (1 cotton pad, ~ 0.3 g of cotton)	~ 800 - 2500 mL	Quantitative removal at the beginning, followed by progressive O ₃ increase	Not significant, as long as the scrubber worked
Squalene (1 Pasteur-pipette drop, ~ 0.05 mL, on rock wool)	≥ 12000 mL	Quantitative removal	Not significant
No scrubber	-	~3.5 ppm measured	0 – 40 ppt increase

350

(a) ANOVA test results for COS



(b) ANOVA test results for O₃




Figure 3: ANOVA test representation for COS (panel a, top) and O₃ (panel b, bottom). The red lines represent the median, while the edges of the blue boxes correspond to the 25th (bottom) and 75th (top) percentile. Black whiskers extend to most extreme data point.

355

The results of the ANOVA test on squalene-based scrubbers are represented in Figure 3 for both COS and O₃. The corresponding p-values are reported in Table S6 and S7 in the Supplement. It is clear that squalene removes O₃ quantitatively: when in place, O₃ mole fraction shows no significant difference between the scrubbing squalene and the configurations where O₃ is not generated (p-value > 0.05).

360

Regarding COS, significant differences (p-value < 0.05) were found between the configuration where O₃ flows through without any scrubber and all other configuration. However, it is noteworthy that significant differences in COS mole fraction were also observed between the configuration where O₃ is generated and then removed by squalene, and the bypass configuration. The air mixture used for this experiment was prepared in the laboratory and contained 0 ppt COS.

365

The effect of squalene on CO₂, CO, N₂O and CH₄ was tested just briefly and is shown in Fig. S1-S4 and Tables S2-S5 in Sect. S2 of the Supplement. Overall, CO trends resembled very closely the ones observed for COS, while no significant variance was observed for all other tracers.

370 The effects of different O₃ scrubbers on COS and O₃, tested on tropospheric compressed air, are shown in Figure 4 and Figure 5. The p-values of the ANOVA test for each species are reported in Table S8 and Table S9 in the Supplement.

When O₃ was not generated, none of the two scrubbers showed significant differences with the bypass channel for either COS, or O₃. Clearly, the O₃ scrubbing was performed quantitatively by the squalene-based scrubber, while the cotton-based scrubber was likely saturated and could not remove the vast majority of the produced ozone. When O₃ was produced in the UV lamp, 375 similarly to what happened during the tests on air simulating stratospheric conditions, we measured higher COS molar fractions both with cotton and with squalene.





Figure 4: Effects of squalene and cotton O₃ scrubbers on a compressed tropospheric air mixture containing COS.



380 **Figure 5: effects of squalene and cotton O₃ scrubbers on O₃ in a compressed tropospheric air mixture.**

4. Discussion

4.1 Collected COS profiles and (Big)LISA samples

4.1.1 COS profiles

The AirCore data presented in Figure 2 show continuously sampled stratospheric COS profiles collected with balloon-borne instruments. Previous stratospheric or Upper Troposphere/Lowermost Stratosphere (UT/LMS) observations were realised with discrete whole-air sampling (Engel and Schmidt, 1994; Karu et al., 2023), in situ spectrometers (Gurganus et al., 2024; Kloss et al., 2021; Kryzstofiak et al., 2015; Leung et al., 2002; Wofsy et al., 2017; Wofsy, 2011) or remote sensing (Glatthor et al., 2017; Toon et al., 2018; Velazco et al., 2011; Yousefi et al., 2019). LISA and BigLISA represent lightweight additions to these methods. For validation purposes, AirCore, LISA and BigLISA observations have been compared with SPIRALE in situ observation (Kryzstofiak et al., 2015) and the ACE-FTS remote sensing observations (Bernath, 2005; Velazco et al., 2011).

385
390

The occasional data gaps shown in the AirCore profiles have different reasons, as summarised in Table 4. TRN4 and the KRN AirCore equipped with Mg(ClO₄)₂ had tubing of two different diameters, which were connected to each other using a sleeve adapter into which they were glued using Loctite Super Attak glue. SOD1 and SOD5 AirCores, instead, were equipped with

395 differential pressure sensors (Amsys, model AMS5915_0050_D_B) and showed signs of COS contamination at the sensors' location. SOD3 AirCore showed COS spikes related to the O₂ altitude-mapping technique employed in one of its halves. COS outgassing from sulfur-containing polymers, in particular rubbers, has been reported in several studies (Cadle and Williams, 1978; Levine et al., 2023; Pos and Berresheim, 1993). We speculate that the polymers constituting the glue, or components of valves (e.g., O-rings) and differential pressure sensors, may have caused the COS outgassing. In the case of SOD3, another possibility could be direct oxidation reactions of reduced sulfur gas species with O₂.
400

Nonetheless, the AirCore profiles we presented are similar to observations reported by previous studies: in TRN, COS mole fraction first decreases from tropospheric values of about 510 ppt up to around 10.5 km, to about 420 ppt at 17 km. Then, it undergoes a faster decrease to 92 – 203 ppt at 22 km. This is consistent with observations reported in previous studies at comparable latitudes (Leung et al., 2002; Toon et al., 2018). Although altitude itself does not provide a precise proxy for quantitative comparisons of vertical profiles, this agreement is a strong suggestion of robust stratospheric COS measurements.
405

At polar latitudes, we observed larger tropospheric variability. The profiles showed tropospheric COS mole fractions ranging between 400 – 470 ppt. Glatthor et al. (2017) reported values at 5 km altitude as low as 330 ppt for COS, during northern summer months at polar latitudes between 2003 – 2012, while Toon et al. (2018) showed a value around 410 ppt under similar conditions. Regarding the stratospheric part of the profile, we observed slight increases or stable values up to 13 – 15 km range (350 – 450 ppt), followed by decreases down to 95 – 130 ppt around 22 km. Most flights from the SOD campaign, in particular SOD1 and SOD5, were characterised by variable lapse rates above the tropopause (see Sect. S4 in the Supplement) which we suspect may be an indicator for COS convective transport and mixing in the lowermost stratosphere. Nevertheless, the observed stratospheric COS range seems consistent with previous studies. Leung et al. (2002) reported 440 ppt below 14 km, decreasing to 120 ppt at 22 km. Glatthor et al. (2017) reports around 490 ppt at 13 km, followed by a decrease to 140 ppt at 22 km. Krysztofiak et al. (2015) reported 420 ± 100 ppt COS below 17 km, decreasing to 150 ppt at 22 km. A more quantitative comparison between our profiles and Krysztofiak et al. (2015) and ACE-FTS (Bernath, 2005; Velazco et al., 2011; Yousefi et al., 2019) observations is presented in Sect. 4.2 and 4.3, respectively.
410
415

420 4.1.2 (Big)LISA samples

The samples from KRN BigLISA show clear signs of contamination, in particular above 20 km (Figure 2). We believe this may be due to COS outgassing from some plastic components of the MLF bags used to collect the samples, such as the O-ring in the valves, or simply tropospheric air that remained trapped inside the bag. Although all the deployed bags are indicated as suitable for sulfur compounds, they are not recommended for low-ppm volatile organic compounds due to background levels (Sigma Aldrich, 2025). This might have also influenced our COS measurements, perhaps due to spectroscopic effects. The contamination appears to be inversely proportional to the sampling pressure and the collected sample volume. Unfortunately,
425

it has not been possible to assess the cause of this contamination precisely. Given these circumstances and the impossibility of applying any correction to these results, BigLISA will be left out of the discussion and comparisons with other datasets.

430 As described in Sect. 2.1.2, during the SOD campaign we introduced a pre-treatment technique that has mitigated this issue for COS, based on previous laboratory tests. Filling and vacuuming the bags with a stratospheric-mimicking gas seemed to have reduced the contamination significantly for most LISA samples, as previously reported in Figure 2 and Table 2. However, in some cases and in particular for smaller samples collected at higher altitudes (SOD2 – L4, SOD5 – L3), this pre-conditioning was not sufficient to prevent biases that are likely ascribable to the bags employed during the campaign or may perhaps be due
435 to contamination during sampling or during the transferring procedure of the samples in glass flasks. For the remaining samples, when LISA flew on the same balloon as one of our AirCores (e.g. SOD1, SOD3), the largest average difference of 31 ± 15 ppt was found when both instruments were flown with a free inlet. Although some variability can be observed between different LISA samples at similar altitudes, their COS mole fraction falls well within the range of the AirCore profiles (see Sect. 4.1.3, 4.2 and 4.3). Overall, LISA samples may need further improvement regarding the sampling and measurement
440 protocols. However, the results presented in this study suggest that this technique should be suitable to obtain reliable COS measurements and that the pre-treatment of MLF bags prevents, or at least reduces, COS contamination due to outgassing from the inner layers of these sample containers.

4.1.3 Datasets consistency

Figure 6 shows the measured COS AirCore profiles and LISA samples from all campaigns, plotted against altitude above
445 tropopause (see Sect. S4 in the Supplement). TRN1 is not presented in this figure, since the tropopause height could not be estimated due to missing temperature and relative humidity data. Differences of up to ~ 100 ppt can be clearly seen between the measured profiles. However, these differences are not constant with altitude and do not show any clear trend over the time span of the campaigns. Moreover, these differences do not show any clear relationship with the different inlets employed.

The most likely explanation for differences between the measured profiles may reside in stratospheric horizontal transport
450 from different latitudes (Toon et al., 2018). The tight correlation between CH₄ and N₂O (Sect. S5 in the Supplement) suggests that the day-to-day variability can likely be ascribed to atmospheric transport (Kondo et al., 1996; Plumb, 2007; Plumb and Ko, 1992). Moreover, long-term changes in COS seasonal cycle, sources and sinks (Belviso et al., 2022; Schmidt et al., 2024; Sturges et al., 2001) may affect its stratospheric abundance, although no significant stratospheric trends have been observed in the last decades (Barkley et al., 2008; Coffey and Hannigan, 2010; Rinsland et al., 2008; Toon et al., 2018).

455 Other possible causes of the differences between the profiles may reside in a combination of instrumental uncertainties, altitude mapping algorithms, sample loss after landing, contaminations and instrumental features (e.g., inlets, different air mixing in the AirCore coil during sampling). Overall, it is difficult to assess the contribution of each of these parameters quantitatively. Instrumental uncertainties and altitude mapping are consistent for our results. No quantification of sample loss is available, but given the AirCore design, this should mostly affect the tropospheric part of the profiles. Some clear contaminations

460 affecting AirCore profiles (Table 4) and (Big)LISA samples were marked as outliers, but it is still possible that more subtle effects affected the samples, in particular the (Big)LISA data. These may include mixing with dead volumes of tropospheric air or fill gas, impurities in the deployed scrubbers or effects due to the instrumental components (e.g., O-rings, tubing, bags). However, even if these uncertainties are hard to quantify, they may account for up to a few cm³, against collected sample volumes ranging roughly between 800 – 1600 cm³. Therefore, we assume that differences due to instrumental effects remain
 465 marginal, while we believe that the day-to-day variability and long-term trends in COS mole fractions are the most important cause for the observed differences.

Table 4: Data gaps caused by clear contamination effects and their respective causes in different AirCore profiles.

Flight identifier and AirCore inlet	Data gaps	Contamination points along the coil
TRN4 – Mg(ClO ₄) ₂	9.2 – 13.9 km	Glue connection between tubes of different diameter in the coil
KRN – Mg(ClO ₄) ₂	12.7 – 13.9 km	Glue connection between tubes of different diameter in the coil
SOD1 – Mg(ClO ₄) ₂	11.9 – 12.8 km	Differential pressure sensors
SOD3 – cotton	4.0 – 5.0 km 7.4 – 8.3 km 11.1 – 12.3 km 16.5 – 17.9 km 20.4 – 21.3 km	Valve employed in O ₂ spiking technique to retrieve altitude, or reactions of other species with O ₂ itself
SOD5 – Mg(ClO ₄) ₂ and cotton	12.4 – 13.7 km	Differential pressure sensors


470 The two SOD3 profiles sampled with University of Bern’s double-sided AirCore are of particular interest, since one side was equipped with a cotton-based O₃ scrubber and the other side was left with a free inlet. As reported in Table 4, the spiking technique deployed by the University of Bern caused COS anomalies in one of the profiles. Moreover, as explained in Sect. 2.1.1, the effect of fill gas was amplified for this sampler: the correlation between CH₄ and N₂O (Fig. S18 in the Supplement) shows a clear deviation for N₂O mole fractions lower than 280 ppb (corresponding to an altitude of roughly 18 km). This
 475 prevented a reasonable analysis of possible effects of O₃ and/or of the efficiency of the cotton scrubber. However, the resulting (shortened) COS profiles show very good accordance with the profiles obtained from other AirCores and small differences ranging between 17 – 46 ppt with the comparable LISA samples (Table 2).

Since it has been shown that O₃ daily variability can affect oxidizable species (Dirksen et al., 2011; Li et al., 2021), we also
 480 speculated that part of the differences between the measured profiles may be ascribed to a daily cycle, similarly to what happens

for O₃ (Frith et al., 2020; Li et al., 2021; Schranz et al., 2018; Studer et al., 2014). Suggestions of a correlation between COS and O₃ stratospheric chemistry and abundance have been presented in previous studies (Engel and Schmidt, 1994). •OH and O•, known to cause COS loss (Brühl et al., 2012; Chin and Davis, 1995; Kryzstofiak et al., 2015), are part of the Chapman cycle of O₃ (Frith et al., 2020; Studer et al., 2014). O₃ daily variability is reported to change depending on seasonality (daily patterns in solar radiation, photolysis), latitude, temperature and atmospheric pressure level (Frith et al., 2020; Studer et al., 2014). However, reaction kinetics and the current estimates of stratospheric COS sinks of 50 ± 15 GgS yr⁻¹ due to photolysis (Whelan et al., 2018) do not support the hypothesis of a detectable daily cycle with the methods presented in this study. On top of that, since we are uncertain about the effectiveness of the cotton-based scrubbers deployed during our campaigns and as we have no O₃ measurements, we cannot support this hypothesis.

490

Overall, we consider our results to be a trustworthy representation of the COS stratospheric conditions during sampling. In fact, in spite of the observed differences between the retrieved profiles, all profiles show similar trends when compared with one another and with the LISA samples. All measurements show the expected reduction due to a stratospheric sink clearly. Their agreement with SPIRALE and ACE-FTS observations will be discussed in Sect. 4.2 and 4.3, respectively.



495


Figure 6: COS profiles and samples from all campaigns, plotted against altitude above tropopause.

4.2 Comparison with SPIRALE observations


Figure 7 and Figure 8 show a comparison plot of AirCore and LISA observations with the in situ SPIRALE COS observations realised by Krysztofiak et al. (2015) in 2009 and 2011. It is evident that SPIRALE observations are only available between 14.5 and 22 km. These observations have a higher spatial resolution over the vertical column (3 to 5 m), but are associated to lower precision compared to AirCore COS measurements analysed by QCLS. Moreover, both techniques measure profiles that are specific for the location and the time where the measurement occurs. In this case, both studies have data collected at polar latitudes, although at different times of the year and, most importantly, between 10 and 14 years apart from each other. However, some studies reported no significant trends for stratospheric COS in recent years (Barkley et al., 2008; Coffey and Hannigan, 2010; Rinsland et al., 2008; Toon et al., 2018), while others report significant trends that vary over time and latitude (Bernath et al., 2020; Glatthor et al., 2017; Hannigan et al., 2022; Kremser et al., 2015; Lejeune et al., 2017). Given the discrepancies between existing literature, more regular observations would be highly desirable to better understand observed trends in stratospheric COS over time. Lightweight instruments such as AirCore and LISA could provide a valuable tool to retrieve balloon-borne stratospheric samples more frequently, at a relatively low cost. To realise a qualitative comparison, both AirCore profiles and SPIRALE observations have been averaged over 0.5 km bins. Using the averaged bins, we calculated the difference between AirCore and SPIRALE. As shown in Figure 7 and Figure 8, most AirCore profiles and all the LISA samples collected at altitudes comparable to the SPIRALE datasets fall within the uncertainty range of both SPIRALE profiles. The horizontal error bars reported in Figure 7 represent the uncertainty of QCLS measurements, while the ones in Figure 8 are the standard deviation obtained from the averaging of SPIRALE over LISA's sampling altitude intervals and AirCore profiles over 0.5 km altitude intervals. The shaded areas in Figure 8 correspond to SPIRALE's errors.

Between 15.5 and 17.5 km, differences between 40 – 110 ppt can be seen between both KRN and SOD AirCore profiles and the 2009 SPIRALE data. However, in this range (in particular between 15.8 and 16.2 km), SPIRALE measured a COS spike that reached up to 577 ppt, a rather unusual mole fraction for these altitudes, which is reflected also in a previous comparison between SPIRALE and ACE-FTS (see Fig. 5 in Krysztofiak et al., 2015). Moreover, as reported in Fig. 5 of Krysztofiak et al. (2015), SPIRALE results fall generally above the averaged ACE-FTS observations. Unfortunately, only COS measurements are available from SPIRALE and it is not possible to verify this idea with observations of other tracers. Nevertheless, considering day-to-day variability (e.g. discrepancies in tropopause height, or the above-cited air transport), the 12 to 14 years differences between the campaigns and the possible long-term trends occurred in this time span (Bernath et al., 2020; Glatthor et al., 2017; Hannigan et al., 2022; Kremser et al., 2015; Lejeune et al., 2017), the qualitative comparisons suggest a strong agreement between the datasets, with the AirCore and LISA results being generally lower than SPIRALE results in 2009.

With regard to SPIRALE observations in 2011, more marked differences are found in comparison with the averaged AirCore
 530 COS mole fractions from the KRN and SOD campaigns. This could be due to the limited altitudinal range of these observations
 and due to the different sampling season (April for this SPIRALE measurements, August for our campaigns), on top of the
 reasons mentioned earlier for the differences between our data and the SPIRALE 2009 observations. Nevertheless, at polar
 latitudes we observe COS mole fraction decreasing from an all-campaign average of 421 ± 26 ppt below 17 km to 170 ± 41
 ppt between 20 – 22 km (see Figure 7). This is comparable to the decrease from 460 to 150 ppt reported by Krysztofiak et al.
 535 (2015) and from 440 to 120 ppt reported by and Leung et al. (2002).



540 **Figure 7: Measured COS profiles and LISA samples at polar latitudes (Kiruna - KRN, Sodankylä - SOD), compared with the SPIRALE's in situ observations from flights in KRN (red and yellow shading representing COS \pm std. dev.), presented by Krysztofiak et al. (2015). The dotted horizontal lines signal the intervals within which the average COS mole fraction is calculated (see Figure 8). The horizontal error bars represent the uncertainty of QCLS measurements.**




545 **Figure 8: Measured COS profiles and LISA samples at polar latitudes (Kiruna - KRN, Sodankylä - SOD), compared with the SPIRALE's in situ observations from flights in KRN (red and yellow shading representing $\text{COS} \pm \text{std. dev.}$), presented by Krysztofiak et al. (2015). The dotted horizontal lines signal the intervals within which the average COS mole fraction is calculated. Panels (a) and (b) show the difference between the measured AirCore profiles and the SPIRALE results. The error bars correspond to the standard deviation obtained from the averaging of SPIRALE over LISA's sampling altitude intervals, and over AirCore profiles over 0.5 km altitude intervals.**

4.3 Comparison with ACE-FTS observations

550 Figure 9 and Figure 10 show the comparison of the measured TRN, KRN and SOD AirCore profiles with the ACE-FTS v5.3 (Boone et al., 2023; Schmidt et al., 2024) averaged observations. The agreement between the averaged ACE-FTS profiles and the AirCore profiles is rather coherent at both mid and polar latitudes. The TRN data shows good agreement between ACE-FTS and AirCore profiles, in particular in the stratospheric part of the profile, where most differences fall within the ± 50 ppt range. The biggest discrepancies are found in the tropospheric part, where seasonal variabilities and daily variations are more pronounced. At polar latitudes, the profiles measured in KRN in 2021 show better agreement with the ACE-FTS results, while 555 the SOD measurements generally resulted in higher COS mole fractions above 15 km altitude, with differences up to ~ 80 ppt. This may be due to specific conditions during the period when the flights were performed, such as different atmospheric transport patterns, or the uncertainties in the AirCore methodology described in Sect. 4.1.3. Similarly to what has been described in Sect. 4.2, differences between ACE-FTS and AirCore and LISA samples were calculated. In this case, AirCore profiles were averaged over 1 km intervals to make them comparable to the ACE-FTS resolution.

Most profiles show higher COS mole fractions in the 14 – 24 km range at polar latitudes and in the 16 – 24 km range at mid-latitudes when compared to ACE-FTS averages. Glatthor et al. (2017) reported a difference up to 100 ppt between COS mole fraction retrieved from MIPAS remote sensing and the ACE-FTS ones between 13 – 16 km. Velazco et al. (2011) found COS mole fractions obtained from the JPL MkIV interferometer 15% higher than ACE-FTS profiles, while Krysztofiak et al. (2015) reported consistency between SPIRALE and ACE-FTS within 11% at polar latitudes and a positive difference of 15 – 20% at mid-latitudes, taking into account both instrumental uncertainties. The discrepancy we observe between AirCore profiles and ACE-FTS is in the range of 80 ppt and between 17 – 22 km. Nonetheless, some profiles show lower COS mole fractions in the same ranges. Therefore, the differences are either due to day-to-day variability, or due to instrumental issues, such as different scrubbers that were employed in different flights.



570

Figure 9: Comparison of the AirCore profiles in Trainou (TRN) with ACE-FTS average over summer months between 45 – 49° N in the 2012 – 2024 period. In panel (a), the shaded yellow area represents the averaged ACE-FTS results $\pm 1\sigma$. The dotted horizontal lines signal the intervals within which the average is calculated. In panel (b), the shaded area corresponds to ACE-FTS 0 difference, $\pm 1\sigma$. The error bars in panel (b) correspond to the standard deviation obtained averaging AirCore profiles over 0.5 km altitude intervals.

575




Figure 10: Comparison of the AirCore profiles (Kiruna - KRN, Sodankylä - SOD) and LISA samples with ACE-FTS average over summer months at 65 – 69° N in the 2012 – 2024 period. In panel (a), the shaded yellow area represents the averaged ACE-FTS results $\pm 1\sigma$. The dotted horizontal lines signal the intervals within which the AirCores' average is calculated. In panel (b), the shaded area corresponds to ACE-FTS 0 difference, $\pm 1\sigma$. The error bars in panel (b) correspond to the standard deviation obtained averaging AirCore profiles over 0.5 km altitude intervals.

4.4 O₃ scrubbing experiments

Our study showed that cotton had only limited O₃ destruction efficiency over time (Table 3), in spite of its employment as an O₃ scrubber for reduced sulfur compounds in previous studies (Andreae et al., 1985; Hofmann et al., 1992; Persson and Leck, 1994). Existing literature reports reactions of cellulose (the primary constituent of cotton) with O₃ leading to the formation of carbonyl and carboxyl groups on cellulose itself (Valls et al., 2022; Zhang et al., 2024). We speculate that the high abundance of O₃ (up to about 3500 ppb) may have saturated the cotton rapidly. This abundance is comparable to that of the stratospheric O₃ layer (Ansmann et al., 2022). Moreover, when the cotton was used several months after the first opening of its package, it is possible that its exposure to atmospheric O₃ (or other oxidants) may have compromised its performance.

As reported in the previous paragraph, squalene was confirmed to be an effective and efficient O₃ scrubbing substance, as we expected after consulting existing literature (Coffaro and Weisel, 2022; Coleman et al., 2008; Zhou et al., 2016). Squalene

scrubbed O₃ in laboratory tests even down to 250 hPa and around 213 K. Unfortunately, since its testing began after the
595 presented campaigns, squalene-based scrubbers have never been deployed in actual fieldwork.

As shown in Figure 3, COS mole fraction was found significantly higher when O₃ was generated both with and without the
squalene scrubber. This is in contrast with the O₃-induced COS loss reported in previous studies (Engel and Schmidt, 1994)
and resembles the positive interference on carbonyl-bearing VOCs measurements reported by Ernle et al. (2023). Assuming
600 the glass to be inert, this observation implies that COS is produced either (i) at the UV lamp during O₃ generation, likely from
traces of VOC or other impurities in the supply gases or (ii) in the tubing downstream of the quartz glass by reactions between
O₃ and wall contaminations, before reaching the scrubber. When tested on compressed tropospheric air, we could observe the
same trends for COS and, although a small decrease was observed for squalene when compared to the bypass channel, it was
not statistically significant. We believe the COS enhancement is due to oxidation of other compounds (e.g., CS₂, DMS) in
605 presence of UV radiation, and not due to the scrubbers themselves. The larger variance for the tests on the scrubbers without
O₃ generation is due to some spectroscopic effects that affected the instrument precision during part of the tests (not shown).
Either way, no sign of negative biases on COS was measured for either of the two scrubbers.

Overall, it is unclear whether COS may have been simultaneously removed by O₃ and produced by the reaction of O₃ with
610 COS precursors that may have been present as impurities in the air mixture employed for the experiments. This limits the
direct comparison with stratospheric field studies (Engel and Schmidt, 1994). Nevertheless, COS mole fraction was
significantly lower when ozonated air was measured after passing through the squalene scrubber (or through the cotton
scrubber, before saturation) than when it was measured after the control channel. We speculate that new volatile compounds
containing carbonyl groups, possibly impurities in the compressed air mixture used for the experiment and/or products of
615 reactions between impurities and O₃, may have influenced the QCLS spectrum, biasing COS measurements. Different studies
report the creation of carbonyl and carboxyl groups on both saturated and unsaturated carbon polymers (Cataldo, 2001; Valls
et al., 2022; Zhang et al., 2024; Zhou et al., 2016). However, these reaction products should not contain sulfur compounds.
Nevertheless, squalene reduced this bias when compared to the configuration without O₃ scrubbing. Therefore, we believe that
most of this bias could have been produced either by products of photochemical reactions due to the UV-lamp employed for
620 O₃ generation, or by reactions of O₃ with impurities or with experimental components. In particular, a phenomenon described
as “ozone cracking” is known to affect polymers, such as (vulcanised) rubbers (Cataldo, 2001; Salomon and Van Bloois, 1963;
Tse, 2007) and polymers that may contain sulfur. Therefore, we also speculate that the reaction of O₃ with squalene and its
consequent removal may have mitigated these reactions, reducing the production of COS during the performed analyses.
Overall, there is no indication that squalene would negatively bias COS measurements, while the presence of O₃ could be
625 detrimental for the measurements, in particular when in presence of unsaturated polymers, which may be prone to degradation.
Unfortunately, the effect of O₃ on long-term stored samples was not investigated within this study.

5. Conclusion

This study presented in situ stratospheric COS observations based on collected air samples using two new techniques, AirCore
630 and (Big)LISA samplers. The collected continuous and discrete stratospheric samples were analysed with a QCLS in the CIO
laboratory at the University of Groningen. The results obtained with both techniques closely resemble the stratospheric trends
retrieved from previous discrete samples and in situ spectroscopic observations. Moreover, we found differences typically
ranging between ± 50 ppt between AirCore data and averaged ACE-FTS data obtained with remote sensing, although we
observe higher COS estimations when approaching low COS abundances when compared to ACE-FTS. We found that, when
635 deploying MLF bags to measure COS, it is necessary to pre-treat the bags before flight to prevent COS contamination due to
outgassing from the polymers constituting the bag. More frequent balloon-borne measurements with AirCore and (Big)LISA
could provide valuable insights regarding stratospheric COS trends, among other species, and support the existing remote
sensing observational network.

640 We also found that cotton-based O₃ scrubbers may have limited efficiency, especially when cotton has been in contact with
the air for several months. Squalene-based scrubbers showed excellent O₃ scrubbing performances and seemed to have no
effect on COS abundance and may become a valuable addition to stratospheric samplers that require O₃ removal.

We found no clear evidence that stratospheric O₃ causes positive or negative biases in COS measurements, since no repeatable
645 differences were found in our samples while deploying different sorts of inlets with different O₃ scrubber materials. However,
the observed COS mole fractions showed some day-to-day variability, a hypothesis which may be ascribed to stratospheric
transport or instrumental biases. Consistent differences in COS profiles point to observed transport variability, hypothesis that
may be corroborated by modelling efforts. The investigation on the effects of O₃ on air samples, in particular containing
reduced sulfur species, could be facilitated by the deployment of the squalene-based scrubbers.

650

Data availability

The data used in this work are available from <https://doi.org/10.5281/zenodo.15749915> (Zanchetta et al., 2025).

Author contribution

655 HC and MK conceived the concept, SvH, AZ and HC designed the experiment, AZ, SvH, JH, RK, TL, MR, ML, PN, SLB,
and HC collected the data. AZ, SvH, and HC wrote the manuscript with contribution from all authors.

Competing interests

Huilin Chen is a member of the editorial board of Atmospheric Measurement Techniques.

660

Acknowledgements

We are grateful for the support during the preparation of the campaigns by Bert Kers, Marcel de Vries and Marc Bleeker at the Center of Isotope Research. We would like to thank the colleagues who collaborated during the campaigns, especially Maria Elena Popa, Johannes Laube and Johannes Degen. Moreover, we would like to thank the European Union project HEMERA for funding the campaign in Kiruna in 2021, which was realised thanks to the collaborative effort of CNES, and ATMO ACCESS for funding the Sodankylä campaign in 2023. We thank Gisele Krysztofiak and Valery Catoire (LPC2E) for providing their in situ balloon-borne SPIRALE spectrometer observations made in 2009 and 2011. The dataset is maintained by the French national centre for Atmospheric data and services AERIS. The AirCore flights were partially supported by the ESA project FRM4GHG.

670

Financial support

This research was supported by the ERC advanced funding scheme (AdG 2016 project no. 742798, project abbreviation COS-OCS), and by the Ruisdael Observatory infrastructure cofinanced by the Dutch Research Council (NWO, Grant No. 184.034.015) and ICOS Netherlands. This work was also supported by the Natural Science Foundation of China (42475115) and by the Fundamental Research Funds for the Central Universities (14380235).

675

References

- Andreae, M. O., Ferek, R. J., Bermond, F., Byrd, K. P., Engstrom, R. T., Hardin, S., Houmère, P. D., LeMarrec, F., Raemdonck, H., and Chatfield, R. B.: Dimethyl sulfide in the marine atmosphere, *J. Geophys. Res. Atmospheres*, 90, 12891–12900, <https://doi.org/10.1029/JD090iD07p12891>, 1985.
- 680 Andreae, T. W., Andreae, M. O., Bingemer, H. G., and Leck, C.: Measurements of dimethyl sulfide and H_2S over the western North Atlantic and the tropical Atlantic, *J. Geophys. Res. Atmospheres*, 98, 23389–23396, <https://doi.org/10.1029/91JD03016>, 1993.
- 685 Ansmann, A., Ohneiser, K., Chudnovsky, A., Knopf, D. A., Eloranta, E. W., Villanueva, D., Seifert, P., Radenz, M., Barja, B., Zamorano, F., Jimenez, C., Engelmann, R., Baars, H., Griesche, H., Hofer, J., Althausen, D., and Wandinger, U.: Ozone depletion in the Arctic and Antarctic stratosphere induced by wildfire smoke, *Atmospheric Chem. Phys.*, 22, 11701–11726, <https://doi.org/10.5194/acp-22-11701-2022>, 2022.
- 690 Barkley, M. P., Palmer, P. I., Boone, C. D., Bernath, P. F., and Suntharalingam, P.: Global distributions of carbonyl sulfide in the upper troposphere and stratosphere, *Geophys. Res. Lett.*, 35, L14810, <https://doi.org/10.1029/2008GL034270>, 2008.

- Belviso, S., Remaud, M., Abadie, C., Maignan, F., Ramonet, M., and Peylin, P.: Ongoing Decline in the Atmospheric COS Seasonal Cycle Amplitude over Western Europe: Implications for Surface Fluxes, *Atmosphere*, 13, 812, <https://doi.org/10.3390/atmos13050812>, 2022.
- 695 Bernath, P. F.: Atmospheric Chemistry Experiment (ACE): Mission overview, *Geophys. Res. Lett.*, 32, L15S01, <https://doi.org/10.1029/2005GL022386>, 2005.
- Bernath, P. F., Steffen, J., Crouse, J., and Boone, C. D.: Sixteen-year trends in atmospheric trace gases from orbit, *J. Quant. Spectrosc. Radiat. Transf.*, 253, 107178, <https://doi.org/10.1016/j.jqsrt.2020.107178>, 2020.
- 700 Bernhard, G. H., Bais, A. F., Aucamp, P. J., Klekociuk, A. R., Liley, J. B., and McKenzie, R. L.: Stratospheric ozone, UV radiation, and climate interactions, *Photochem. Photobiol. Sci.*, 22, 937–989, <https://doi.org/10.1007/s43630-023-00371-y>, 2023.
- Berry, J., Wolf, A., Campbell, J. E., Baker, I., Blake, N., Blake, D., Denning, A. S., Kawa, S. R., Montzka, S. A., Seibt, U., Stimler, K., Yakir, D., and Zhu, Z.: A coupled model of the global cycles of carbonyl sulfide and CO₂: A possible new window on the carbon cycle, *J. Geophys. Res. Biogeosciences*, 118, 842–852, <https://doi.org/10.1002/jgrg.20068>, 2013.
- 705 Boone, C. D., Bernath, P. F., and Lecours, M.: Version 5 retrievals for ACE-FTS and ACE-imagers, *J. Quant. Spectrosc. Radiat. Transf.*, 310, 108749, <https://doi.org/10.1016/j.jqsrt.2023.108749>, 2023.
- Brühl, C., Lelieveld, J., Crutzen, P. J., and Tost, H.: The role of carbonyl sulphide as a source of stratospheric sulphate aerosol and its impact on climate, *Atmospheric Chem. Phys.*, 12, 1239–1253, <https://doi.org/10.5194/acp-12-1239-2012>, 2012.
- 710 Cadle, S. H. and Williams, R. L.: Gas and Particle Emissions from Automobile Tires in Laboratory and Field Studies, *J. Air Pollut. Control Assoc.*, 28, 502–507, <https://doi.org/10.1080/00022470.1978.10470623>, 1978.
- Campbell, J. E., Carmichael, G. R., Chai, T., Mena-Carrasco, M., Tang, Y., Blake, D. R., Blake, N. J., 715 Vay, S. A., Collatz, G. J., Baker, I., Berry, J. A., Montzka, S. A., Sweeney, C., Schnoor, J. L., and Stanier, C. O.: Photosynthetic Control of Atmospheric Carbonyl Sulfide During the Growing Season, *Science*, 322, 1085–1088, <https://doi.org/10.1126/science.1164015>, 2008.
- Cataldo, F.: On the ozone protection of polymers having non-conjugated unsaturation, *Polym. Degrad. Stab.*, 72, 287–296, [https://doi.org/10.1016/S0141-3910\(01\)00017-9](https://doi.org/10.1016/S0141-3910(01)00017-9), 2001.
- 720 Chin, M. and Davis, D. D.: A reanalysis of carbonyl sulfide as a source of stratospheric background sulfur aerosol, *J. Geophys. Res. Atmospheres*, 100, 8993–9005, <https://doi.org/10.1029/95JD00275>, 1995.

- Coffaro, B. and Weisel, C. P.: Reactions and Products of Squalene and Ozone: A Review, *Environ. Sci. Technol.*, 56, 7396–7411, <https://doi.org/10.1021/acs.est.1c07611>, 2022.
- 725 Coffey, M. T. and Hannigan, J. W.: The temporal trend of stratospheric carbonyl sulfide, *J. Atmospheric Chem.*, 67, 61–70, <https://doi.org/10.1007/s10874-011-9203-4>, 2010.
- Coleman, B. K., Destailats, H., Hodgson, A. T., and Nazaroff, W. W.: Ozone consumption and volatile byproduct formation from surface reactions with aircraft cabin materials and clothing fabrics, *Atmos. Environ.*, 42, 642–654, <https://doi.org/10.1016/j.atmosenv.2007.10.001>, 2008.
- 730 Crutzen, P. J.: The possible importance of CSO for the sulfate layer of the stratosphere, *Geophys. Res. Lett.*, 3, 73–76, <https://doi.org/10.1029/GL003i002p00073>, 1976.
- Dirksen, R. J., Boersma, K. F., Eskes, H. J., Ionov, D. V., Bucsela, E. J., Levelt, P. F., and Kelder, H. M.: Evaluation of stratospheric NO₂ retrieved from the Ozone Monitoring Instrument: Intercomparison, diurnal cycle, and trending, *J. Geophys. Res.*, 116, D08305, <https://doi.org/10.1029/2010JD014943>,
735 2011.
- Engel, A. and Schmidt, U.: Vertical profile measurements of carbonylsulfide in the stratosphere, *Geophys. Res. Lett.*, 21, 2219–2222, <https://doi.org/10.1029/94GL01461>, 1994.
- Ernle, L., Ringsdorf, M. A., and Williams, J.: Influence of ozone and humidity on PTR-MS and GC-MS VOC measurements with and without a Na₂ S₂ O₃ ozone scrubber, *Atmospheric Meas. Tech.*, 16, 1179–
740 1194, <https://doi.org/10.5194/amt-16-1179-2023>, 2023.
- Ferm, R. J.: The Chemistry Of Carbonyl Sulfide, *Chem. Rev.*, 57, 621–640, <https://doi.org/10.1021/cr50016a002>, 1957.
- Frith, S. M., Bhartia, P. K., Oman, L. D., Kramarova, N. A., McPeters, R. D., and Labow, G. J.: Model-based climatology of diurnal variability in stratospheric ozone as a data analysis tool, *Atmospheric Meas. Tech.*, 13, 2733–2749, <https://doi.org/10.5194/amt-13-2733-2020>, 2020.
745
- Glatthor, N., Höpfner, M., Leyser, A., Stiller, G. P., Von Clarmann, T., Grabowski, U., Kellmann, S., Linden, A., Sinnhuber, B.-M., Krysztofiak, G., and Walker, K. A.: Global carbonyl sulfide (OCS) measured by MIPAS/Envisat during 2002–2012, *Atmospheric Chem. Phys.*, 17, 2631–2652, <https://doi.org/10.5194/acp-17-2631-2017>, 2017.
- 750 Gurganus, C., Rollins, A. W., Waxman, E., Pan, L. L., Smith, W. P., Rei Ueyama, Feng, W., Chipperfield, M. P., Atlas, E. L., Schwarz, J. P., Lee, S., and Thornberry, T. D.: Highlighting the impact of anthropogenic OCS emissions on the stratospheric sulfur budget with in-situ observations, *Authorea Authorea*, <https://doi.org/10.22541/essoar.172801406.62154439/v1>, 2024.

- 755 Hannigan, J. W., Ortega, I., Shams, S. B., Blumenstock, T., Campbell, J. E., Conway, S., Flood, V.,
Garcia, O., Griffith, D., Grutter, M., Hase, F., Jeseck, P., Jones, N., Mahieu, E., Makarova, M., De
Mazière, M., Morino, I., Murata, I., Nagahama, T., Nakijima, H., Notholt, J., Palm, M., Poberovskii, A.,
Rettinger, M., Robinson, J., Röhling, A. N., Schneider, M., Servais, C., Smale, D., Stremme, W., Strong,
K., Sussmann, R., Te, Y., Vigouroux, C., and Wizenberg, T.: Global Atmospheric OCS Trend Analysis
760 From 22 NDACC Stations, *J. Geophys. Res. Atmospheres*, 127, e2021JD035764,
<https://doi.org/10.1029/2021JD035764>, 2022.
- Hofmann, U., Hofmann, R., and Kesselmeier, J.: Cryogenic trapping of reduced sulfur compounds using
a nafion drier and cotton wadding as an oxidant scavenger, *Atmospheric Environ. Part Gen. Top.*, 26,
2445–2449, [https://doi.org/10.1016/0960-1686\(92\)90374-T](https://doi.org/10.1016/0960-1686(92)90374-T), 1992.
- Hooghiem, J. J. D., de Vries, M., Been, H. A., Heikkinen, P., Kivi, R., and Chen, H.: LISA: a
765 lightweight stratospheric air sampler, *Atmospheric Meas. Tech.*, 11, 6785–6801,
<https://doi.org/10.5194/amt-11-6785-2018>, 2018.
- Karion, A., Sweeney, C., Tans, P., and Newberger, T.: AirCore: An Innovative Atmospheric Sampling
System, *J. Atmospheric Ocean. Technol.*, 27, 1839–1853, <https://doi.org/10.1175/2010JTECHA1448.1>,
2010.
- 770 Karu, E., Li, M., Ernle, L., Brenninkmeijer, C. A. M., Lelieveld, J., and Williams, J.: Carbonyl Sulfide
(OCS) in the Upper Troposphere/Lowermost Stratosphere (UT/LMS) Region: Estimates of Lifetimes
and Fluxes, *Geophys. Res. Lett.*, 50, e2023GL105826, <https://doi.org/10.1029/2023GL105826>, 2023.
- Kivi, R. and Heikkinen, P.: Fourier transform spectrometer measurements of column CO₂ at Sodankylä,
Finland, *Geosci. Instrum. Methods Data Syst.*, 5, 271–279, <https://doi.org/10.5194/gi-5-271-2016>, 2016.
- 775 Kloss, C., Tan, V., Leen, J. B., Madsen, G. L., Gardner, A., Du, X., Kulesa, T., Schillings, J., Schneider,
H., Schrade, S., Qiu, C., and Von Hobe, M.: Airborne Mid-Infrared Cavity enhanced Absorption
spectrometer (AMICA), *Atmospheric Meas. Tech.*, 14, 5271–5297, <https://doi.org/10.5194/amt-14-5271-2021>, 2021.
- Kondo, Y., Schmidt, U., Sugita, T., Engel, A., Koike, M., Amedieu, P., Gunson, M. R., and Rodriguez,
780 J.: NO_y correlation with N₂O and CH₄ in the midlatitude stratosphere, *Geophys. Res. Lett.*, 23, 2369–
2372, <https://doi.org/10.1029/96GL00870>, 1996.
- Kooijmans, L. M. J., Uitslag, N. A. M., Zahniser, M. S., Nelson, D. D., Montzka, S. A., and Chen, H.:
Continuous and high-precision atmospheric concentration measurements of COS, CO₂, CO and H₂O
using a quantum cascade laser spectrometer (QCLS), *Atmospheric Meas. Tech.*, 9, 5293–5314,
785 <https://doi.org/10.5194/amt-9-5293-2016>, 2016.

- Kremser, S., Jones, N. B., Palm, M., Lejeune, B., Wang, Y., Smale, D., and Deutscher, N. M.: Positive trends in Southern Hemisphere carbonyl sulfide, *Geophys. Res. Lett.*, 42, 9473–9480, <https://doi.org/10.1002/2015GL065879>, 2015.
- 790 Kremser, S., Thomason, L. W., von Hobe, M., Hermann, M., Deshler, T., Timmreck, C., Toohey, M.,
Stenke, A., Schwarz, J. P., Weigel, R., Fueglistaler, S., Prata, F. J., Vernier, J.-P., Schlager, H., Barnes, J.
E., Antuña-Marrero, J.-C., Fairlie, D., Palm, M., Mahieu, E., Notholt, J., Rex, M., Bingen, C.,
Vanhellemont, F., Bourassa, A., Plane, J. M. C., Klocke, D., Carn, S. A., Clarisse, L., Trickl, T., Neely,
R., James, A. D., Rieger, L., Wilson, J. C., and Meland, B.: Stratospheric aerosol-Observations,
795 processes, and impact on climate: Stratospheric Aerosol, *Rev. Geophys.*, 54, 278–335,
<https://doi.org/10.1002/2015RG000511>, 2016.
- Krysztofiak, G., Té, Y. V., Catoire, V., Berthet, G., Toon, G. C., Jégou, F., Jeseck, P., and Robert, C.:
Carbonyl Sulphide (OCS) Variability with Latitude in the Atmosphere, *Atmosphere-Ocean*, 53, 89–101,
<https://doi.org/10.1080/07055900.2013.876609>, 2015.
- Laube, J. C., Engel, A., Bönisch, H., Möbius, T., Sturges, W. T., Braß, M., and Röckmann, T.: Fractional
800 release factors of long-lived halogenated organic compounds in the tropical stratosphere, *Atmospheric
Chem. Phys.*, 10, 1093–1103, <https://doi.org/10.5194/acp-10-1093-2010>, 2010.
- Lee, C.-L. and Brimblecombe, P.: Anthropogenic contributions to global carbonyl sulfide, carbon
disulfide and organosulfides fluxes, *Earth-Sci. Rev.*, 160, 1–18,
<https://doi.org/10.1016/j.earscirev.2016.06.005>, 2016.
- 805 Lejeune, B., Mahieu, E., Vollmer, M. K., Reimann, S., Bernath, P. F., Boone, C. D., Walker, K. A., and
Servais, C.: Optimized approach to retrieve information on atmospheric carbonyl sulfide (OCS) above
the Jungfraujoch station and change in its abundance since 1995, *J. Quant. Spectrosc. Radiat. Transf.*,
186, 81–95, <https://doi.org/10.1016/j.jqsrt.2016.06.001>, 2017.
- Leung, F. T., Colussi, A. J., Hoffmann, M. R., and Toon, G. C.: Isotopic fractionation of carbonyl sulfide
810 in the atmosphere: Implications for the source of background stratospheric sulfate aerosol, *Geophys.
Res. Lett.*, 29, <https://doi.org/10.1029/2001GL013955>, 2002.
- Levine, Y., Chetrit, E., Fishman, Y., Siyum, Y., Rabaev, M., Fletcher, A., and Tartakovsky, K.: A novel
approach to plasticizer content calculation in an acrylonitrile-butadiene rubber real-time aging study
(NBR), *Polym. Test.*, 124, 108091, <https://doi.org/10.1016/j.polymertesting.2023.108091>, 2023.
- 815 Li, K.-F., Khoury, R., Pongetti, T. J., Sander, S. P., Mills, F. P., and Yung, Y. L.: Diurnal variability of
stratospheric column NO₂ measured using direct solar and lunar spectra over Table Mountain,
California (34.38° N), *Atmospheric Meas. Tech.*, 14, 7495–7510, [https://doi.org/10.5194/amt-14-7495-](https://doi.org/10.5194/amt-14-7495-2021)
2021, 2021.

- 820 Ma, J., Kooijmans, L. M. J., Cho, A., Montzka, S. A., Glatthor, N., Worden, J. R., Kuai, L., Atlas, E. L., and Krol, M. C.: Inverse modelling of carbonyl sulfide: implementation, evaluation and implications for the global budget, *Atmospheric Chem. Phys.*, 21, 3507–3529, <https://doi.org/10.5194/acp-21-3507-2021>, 2021.
- 825 Membrive, O., Crevoisier, C., Sweeney, C., Danis, F., Hertzog, A., Engel, A., Bönisch, H., and Picon, L.: AirCore-HR: a high-resolution column sampling to enhance the vertical description of CH₄ and CO₂, *Atmospheric Meas. Tech.*, 10, 2163–2181, <https://doi.org/10.5194/amt-10-2163-2017>, 2017.
- Montzka, S. A., Calvert, P., Hall, B. D., Elkins, J. W., Conway, T. J., Tans, P. P., and Sweeney, C.: On the global distribution, seasonality, and budget of atmospheric carbonyl sulfide (COS) and some similarities to CO₂, *J. Geophys. Res.*, 112, D09302, <https://doi.org/10.1029/2006JD007665>, 2007.
- 830 Persson, Cecilia. and Leck, Caroline.: Determination of Reduced Sulfur Compounds in the Atmosphere Using a Cotton Scrubber for Oxidant Removal and Gas Chromatography with Flame Photometric Detection, *Anal. Chem.*, 66, 983–987, <https://doi.org/10.1021/ac00079a009>, 1994.
- Picardo, M., Ottaviani, M., Camera, E., and Mastrofrancesco, A.: Sebaceous gland lipids, *Dermatoendocrinol.*, 1, 68–71, <https://doi.org/10.4161/derm.1.2.8472>, 2009.
- 835 Plumb, R. A.: Tracer interrelationships in the stratosphere, *Rev. Geophys.*, 45, <https://doi.org/10.1029/2005RG000179>, 2007.
- Plumb, R. A. and Ko, M. K. W.: Interrelationships between mixing ratios of long-lived stratospheric constituents, *J. Geophys. Res. Atmospheres*, 97, 10145–10156, <https://doi.org/10.1029/92JD00450>, 1992.
- 840 Pos, W. H. and Berresheim, H.: Automotive tire wear as a source for atmospheric OCS and CS₂, *Geophys. Res. Lett.*, 20, 815–817, <https://doi.org/10.1029/93GL00972>, 1993.
- 845 Remaud, M., Ma, J., Krol, M., Abadie, C., Cartwright, M. P., Patra, P., Niwa, Y., Rodenbeck, C., Belviso, S., Kooijmans, L., Lennartz, S., Maignan, F., Chevallier, F., Chipperfield, M. P., Pope, R. J., Harrison, J. J., Vimont, I., Wilson, C., and Peylin, P.: Intercomparison of Atmospheric Carbonyl Sulfide (TransCom-COS; Part One): Evaluating the Impact of Transport and Emissions on Tropospheric Variability Using Ground-Based and Aircraft Data, *J. Geophys. Res. Atmospheres*, 128, e2022JD037817, <https://doi.org/10.1029/2022JD037817>, 2023.
- 850 Rinsland, C. P., Chiou, L., Mahieu, E., Zander, R., Boone, C. D., and Bernath, P. F.: Measurements of long-term changes in atmospheric OCS (carbonyl sulfide) from infrared solar observations, *J. Quant. Spectrosc. Radiat. Transf.*, 109, 2679–2686, <https://doi.org/10.1016/j.jqsrt.2008.07.008>, 2008.

- Salomon, G. and Van Bloois, F.: The mechanism of ozone cracking. I. Ozone cracking of plastic films, *J. Appl. Polym. Sci.*, 7, 1117–1132, <https://doi.org/10.1002/app.1963.070070325>, 1963.
- 855 Sandoval-Soto, L., Stanimirov, M., Von Hobe, M., Schmitt, V., Valdes, J., Wild, A., and Kesselmeier, J.: Global uptake of carbonyl sulfide (COS) by terrestrial vegetation: Estimates corrected by deposition velocities normalized to the uptake of carbon dioxide (CO₂), *Biogeosciences*, 2, 125–132, <https://doi.org/10.5194/bg-2-125-2005>, 2005.
- Schmidt, M., Bernath, P., Boone, C., Lecours, M., and Steffen, J.: Trends in atmospheric composition between 2004–2023 using version 5 ACE-FTS data, *J. Quant. Spectrosc. Radiat. Transf.*, 325, 109088, <https://doi.org/10.1016/j.jqsrt.2024.109088>, 2024.
- 860 Schmidt, U., Kulesa, G., Klein, E., Röth, E.-P., Fabian, P., and Borchers, R.: Intercomparison of balloon-borne cryogenic whole air samplers during the MAP/GLOBUS 1983 campaign, *Planet. Space Sci.*, 35, 647–656, [https://doi.org/10.1016/0032-0633\(87\)90131-0](https://doi.org/10.1016/0032-0633(87)90131-0), 1987.
- Schranz, F., Fernandez, S., Kämpfer, N., and Palm, M.: Diurnal variation in middle-atmospheric ozone observed by ground-based microwave radiometry at Ny-Ålesund over 1 year, *Atmospheric Chem. Phys.*, 18, 4113–4130, <https://doi.org/10.5194/acp-18-4113-2018>, 2018.
- 865 Schuck, T. J., Degen, J., Keber, T., Meixner, K., Wagenhäuser, T., Ghysels, M., Durry, G., Amarouche, N., Zanchetta, A., Van Heuven, S., Chen, H., Laube, J. C., Baartman, S. L., Van Der Veen, C., Popa, M. E., and Engel, A.: Measurement report: Greenhouse gas profiles and age of air from the 2021 HEMERA-TWIN balloon launch, *Atmospheric Chem. Phys.*, 25, 4333–4348, <https://doi.org/10.5194/acp-25-4333-2025>, 2025.
- 870 SupelTM-Inert Multi-Layer Foil: <https://www.sigmaldrich.com/NL/en/product/supelco/30228u>, last access: 30 April 2025.
- Stimler, K., Nelson, D., and Yakir, D.: High precision measurements of atmospheric concentrations and plant exchange rates of carbonyl sulfide using mid-IR quantum cascade laser, *Glob. Change Biol.*, <https://doi.org/10.1111/j.1365-2486.2009.02088.x>, 2009.
- 875 Studer, S., Hocke, K., Schanz, A., Schmidt, H., and Kämpfer, N.: A climatology of the diurnal variations in stratospheric and mesospheric ozone over Bern, Switzerland, *Atmospheric Chem. Phys.*, 14, 5905–5919, <https://doi.org/10.5194/acp-14-5905-2014>, 2014.
- 880 Sturges, W. T., Penkett, S. A., Barnola, J., Chappellaz, J., Atlas, E., and Stroud, V.: A long-term record of carbonyl sulfide (COS) in two hemispheres from firn air measurements, *Geophys. Res. Lett.*, 28, 4095–4098, <https://doi.org/10.1029/2001GL013958>, 2001.
- Tans, P.: Fill dynamics and sample mixing in the AirCore, *Atmospheric Meas. Tech.*, 15, 1903–1916, <https://doi.org/10.5194/amt-15-1903-2022>, 2022.

- 885 Tong, X., Van Heuven, S., Scheeren, B., Kers, B., Hutjes, R., and Chen, H.: Aircraft-Based AirCore Sampling for Estimates of N₂O and CH₄ Emissions, *Environ. Sci. Technol.*, 57, 15571–15579, <https://doi.org/10.1021/acs.est.3c04932>, 2023.
- Toon, G. C., Blavier, J.-F. L., and Sung, K.: Atmospheric carbonyl sulfide (OCS) measured remotely by FTIR solar absorption spectrometry, *Atmospheric Chem. Phys.*, 18, 1923–1944, <https://doi.org/10.5194/acp-18-1923-2018>, 2018.
- 890 Tse, M. F.: Ozone cracking and flex cracking of crosslinked polymer blend compounds, *J. Appl. Polym. Sci.*, 103, 2183–2196, <https://doi.org/10.1002/app.25139>, 2007.
- Valls, C., Cusola, O., and Roncero, M. B.: Evaluating the potential of ozone in creating functional groups on cellulose, *Cellulose*, 29, 6595–6610, <https://doi.org/10.1007/s10570-022-04694-4>, 2022.
- 895 Velazco, V. A., Toon, G. C., Blavier, J.-F. L., Kleinböhl, A., Manney, G. L., Daffer, W. H., Bernath, P. F., Walker, K. A., and Boone, C.: Validation of the Atmospheric Chemistry Experiment by noncoincident MkIV balloon profiles, *J. Geophys. Res.*, 116, D06306, <https://doi.org/10.1029/2010JD014928>, 2011.
- Vinković, K., Andersen, T., De Vries, M., Kers, B., Van Heuven, S., Peters, W., Hensen, A., Van Den Bulk, P., and Chen, H.: Evaluating the use of an Unmanned Aerial Vehicle (UAV)-based active AirCore system to quantify methane emissions from dairy cows, *Sci. Total Environ.*, 831, 154898, <https://doi.org/10.1016/j.scitotenv.2022.154898>, 2022.
- 900 Wagenhäuser, T., Engel, A., and Sitals, R.: Testing the altitude attribution and vertical resolution of AirCore measurements with a new spiking method, *Atmospheric Meas. Tech.*, 14, 3923–3934, <https://doi.org/10.5194/amt-14-3923-2021>, 2021.
- 905 Whelan, M. E., Lennartz, S. T., Gimeno, T. E., Wehr, R., Wohlfahrt, G., Wang, Y., Kooijmans, L. M. J., Hilton, T. W., Belviso, S., Peylin, P., Commane, R., Sun, W., Chen, H., Kuai, L., Mammarella, I., Maseyk, K., Berkelhammer, M., Li, K.-F., Yakir, D., Zumkehr, A., Katayama, Y., Ogée, J., Spielmann, F. M., Kitz, F., Rastogi, B., Kesselmeier, J., Marshall, J., Erkkilä, K.-M., Wingate, L., Meredith, L. K., He, W., Bunk, R., Launois, T., Vesala, T., Schmidt, J. A., Fichot, C. G., Seibt, U., Saleska, S., Saltzman, E. S., Montzka, S. A., Berry, J. A., and Campbell, J. E.: Reviews and syntheses: Carbonyl sulfide as a multi-scale tracer for carbon and water cycles, *Biogeosciences*, 15, 3625–3657, <https://doi.org/10.5194/bg-15-3625-2018>, 2018.
- 915 Wofsy, S., Daube, B., Jimenez-Pizarro, R., Kort, E., Pittman, J., Park, S., Commane, R., Xiang, B., Santoni, G., Jacob, D., Fisher, J., Pickett-Heaps, C., Wang, H., Wecht, K., Wang, Q., Stephens, B., Shertz, S., Watt, A., Romashkin, P., Campos, T., Haggerty, J., Cooper, W., Rogers, D., Beaton, S., Hendershot, R., Elkins, J., Fahey, D., Gao, R., Schwarz, J., Moore, F., Montzka, S., Perring, A., Hurst, D., Miller, B., Sweeney, C., Oltmans, S., Hintsä, E., Nance, D., Dutton, G., Watts, L., Spackman, J., Rosenlof, K., Ray, E., Hall, B., Zondlo, M., Diao, M., Keeling, R., Bent, J., Atlas, E., Lueb, R., and M.

- J. Mahoney (Deceased): HIPPO Combined Discrete Flask and GC Sample GHG, Halocarbon, and Hydrocarbon Data. Version 1.0 (1.0), https://doi.org/10.3334/CDIAC/HIPPO_012, 2017.
- 920 Wofsy, S. C.: HIAPER Pole-to-Pole Observations (HIPPO): fine-grained, global-scale measurements of climatically important atmospheric gases and aerosols, *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.*, 369, 2073–2086, <https://doi.org/10.1098/rsta.2010.0313>, 2011.
- World Meteorological Organization (WMO): *Meteorology – A three-dimensional science: Second session of the Commission for Aerology*, 1957.
- 925 World Meteorological Organization (WMO): *Scientific assessment of ozone depletion, 1998*, edited by: National Oceanic and Atmospheric Administration, National Aeronautics and Space Administration, United Nations Environment Programme, World Meteorological Organization, and European Commission, National Oceanic and Atmospheric Administration : National Aeronautics and Space Administration ; United Nations Environment Programme ; World Meteorological Organization ;
- 930 European Commission, Washington, DC : Nairobi, Kenya : Geneva, Switzerland : Brussels, Belgium, 1 pp., 1999.
- Yousefi, M., Bernath, P. F., Boone, C. D., and Toon, G. C.: Global measurements of atmospheric carbonyl sulfide (OCS), OC34S and O13CS, *J. Quant. Spectrosc. Radiat. Transf.*, 238, 106554, <https://doi.org/10.1016/j.jqsrt.2019.06.033>, 2019.
- 935 Zanchetta, A., Kooijmans, L. M. J., Van Heuven, S., Scifo, A., Scheeren, H. A., Mammarella, I., Karstens, U., Ma, J., Krol, M., and Chen, H.: Sources and sinks of carbonyl sulfide inferred from tower and mobile atmospheric observations in the Netherlands, *Biogeosciences*, 20, 3539–3553, <https://doi.org/10.5194/bg-20-3539-2023>, 2023.
- Zhang, L., Fu, J., Gao, W., Li, Y., and Fan, X.: Revealing the structural variation and degradation mechanism of cellulose during ozone oxidation treatment, *Ind. Crops Prod.*, 219, 119101, <https://doi.org/10.1016/j.indcrop.2024.119101>, 2024.
- 940 Zhou, S., Forbes, M. W., Katrib, Y., and Abbatt, J. P. D.: Rapid Oxidation of Skin Oil by Ozone, *Environ. Sci. Technol. Lett.*, 3, 170–174, <https://doi.org/10.1021/acs.estlett.6b00086>, 2016.