

# 1 Extending the range and reach of physically-based Greenland ice 2 sheet sea-level projections

3 Heiko Goelzer<sup>1</sup>, Constantijn J. Berends<sup>2</sup>, Fredrik Boberg<sup>3</sup>, Gael Durand<sup>4</sup>, Tamsin Edwards<sup>5</sup>, Xavier  
4 Fettweis<sup>6</sup>, Fabien Gillet-Chaulet<sup>4</sup>, Quentin Glaude<sup>6,7</sup>, Philippe Huybrechts<sup>8</sup>, Sébastien Le clec'h<sup>8</sup>, Ruth  
5 Mottram<sup>3</sup>, Brice Noel<sup>6</sup>, Martin Olesen<sup>3</sup>, Charlotte Rahlves<sup>1,9</sup>, Jeremy Rohmer<sup>10</sup>, Michiel van den Broeke<sup>2</sup>,  
6 Roderik S.W. van de Wal<sup>2,11</sup>

7 <sup>1</sup> NORCE Research, Bjerknes Centre for Climate Research, Bergen, Norway

8 <sup>2</sup> Institute for Marine and Atmospheric research Utrecht, Utrecht University, Utrecht, the Netherlands

9 <sup>3</sup> Danish Meteorological Institute (DMI), Copenhagen, Denmark

10 <sup>4</sup> Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, 38000 Grenoble, France

11 <sup>5</sup> Department of Geography, King's College London, London, UK

12 <sup>6</sup> Laboratory of Climatology, Department of Geography, SPHERES research unit, University of Liège, Liège, Belgium

13 <sup>7</sup> Applied Computer Electronics Laboratory, University of Liège, Liège, Belgium

14 <sup>8</sup> Vrije Universiteit Brussel, Earth System Sciences and Departement Geografie, Pleinlaan 2, Brussel, Belgium

15 <sup>9</sup> Department of Earth Science, University of Bergen, Bjerknes Centre for Climate Research, Bergen, Norway

16 <sup>10</sup> BRGM, 3 av. C. Guillemin, 45060 Orléans, France

17 <sup>11</sup> Faculty of Geosciences, Department of Physical Geography, Utrecht University, Utrecht, the Netherlands

18 *Correspondence to:* Heiko Goelzer (heig@norceresearch.no)

19 **Abstract.** We present an ensemble of physically-based ice sheet model projections for the Greenland ice sheet (GrIS) that was  
20 produced as part of the European project PROTECT. Our ice sheet model (ISM) simulations are forced by high-resolution  
21 regional climate model (RCM) output and other climate model forcing, including a parameterisation for the retreat of marine-  
22 terminating outlet glaciers. The experimental design builds on the Ice Sheet Model Intercomparison Project for CMIP6  
23 (ISMIP6) protocol and extends it to more fully account for uncertainties in sea-level projections. We include a wider range of  
24 CMIP6 climate model output, more climate change scenarios, several climate downscaling approaches, a wider range of  
25 sensitivity to ocean forcing and we extend projections beyond the year 2100 up to year 2300, including idealised overshoot  
26 scenarios. GrIS sea-level rise contributions range from 16-76 mm (SSP1-2.6/RCP2.6), 22-163 mm (SSP2-4.5) and 27-354 mm  
27 (SSP5-8.5/RCP8.5) in the year 2100 (relative to 2014). The projections are strongly dependent on the climate scenario,  
28 moderately sensitive to the choice of RCM, and relatively insensitive to the ice sheet model choice. In year 2300, contributions  
29 reach 49 to 3127 mm, indicative of large uncertainties and a potentially very large long-term response. Idealised overshoot  
30 experiments to 2300 produce sea-level contributions in a range from 49 to 201 mm, with the ice sheet seemingly stabilised in  
31 a third of the experiments. Repeating end of the 21<sup>st</sup> century forcing until 2300 results in contributions of 58-163 mm (repeated  
32 SSP1-2.6), 98-218 mm (repeated SSP2-4.5) and 282-1230 mm (repeated SSP5-8.5). The largest contributions of more than  
33 3000 mm by year 2300 are found for extreme scenarios of extended SSP5-8.5 with unabated warming throughout the 22<sup>nd</sup> and  
34 23<sup>rd</sup> century. We also extend the ISMIP6 forcing approach backwards over the historical period and successfully produce  
35 consistent simulations in both past and future for three of the four ISMs. The ensemble design of ISM experiments is geared

36 towards the subsequent use of emulators to facilitate statistical interpretation of the results and produce probabilistic projections  
37 of the GrIS contribution to future sea-level rise.

## 38 1 Introduction

39 The Greenland ice sheet (GrIS) has transitioned from a near zero overall mass balance before the early 1990s to rapidly  
40 increasing mass loss that is ongoing today (van den Broeke et al., 2017). The driving mechanism of this change can be largely  
41 attributed to atmospheric and oceanic warming surrounding the ice sheet, which is amplified in the Arctic region compared to  
42 the global mean (Rantanen et al. 2022). This makes the ice sheet the currently largest single cryospheric contributor to global  
43 mean sea-level rise (e.g. Fox-Kemper et al. 2021). Projecting the future evolution of the GrIS is therefore an important element  
44 in providing sea-level practitioners with relevant information for adaptation planning and providing policy makers with  
45 guidance concerning the urgent need for mitigation, in line with the PROTECT project goals (Durand et al. 2022).

46 Projections of ice sheet contributions to future sea-level rise have recently been organised into a global community effort under  
47 the guidance of the Ice Sheet Model Intercomparison Project for CMIP6 (ISMIP6, <https://climate-cryosphere.org/about-ismip6/>). The initiative provided projections for both the Greenland and Antarctic ice sheets that served as the main source of  
48 the ice sheet sea-level projections (Fox-Kemper et al. 2021) in the latest report of the IPCC, AR6. The ISMIP6 GrIS projections  
49 (Goelzer et al., 2020a) used an experimental protocol (Nowicki et al., 2016, 2020) that included a regional climate model  
50 (RCM) to dynamically downscale global climate scenarios to the ice sheet scale. While only one RCM was used for the  
51 projections in ISMIP6 for feasibility reasons, recent work has revealed that different RCMs can show widely different  
52 behaviour under future climate change (Glaude et al., 2024). This strongly suggests that fully characterising uncertainties in  
53 ice sheet projections requires a broader sampling of climate forcing uncertainty, in particular pertaining to the downscaling  
54 process. In addition, the first wave of ISMIP6 projections was forced by output from CMIP5 models (Goelzer et al., 2020a)  
55 and a subsequent update used a small subset of the then available CMIP6 models forcing (Payne et al. 2021), both under only  
56 two scenarios (RCP8.5/SSP5-8.5 and RCP2.6/SSP1-2.6). Extending the forcing to a broader range of downscaled CMIP6  
57 climate forcing and scenarios was therefore another major concern.

58 The ISMIP6 projections started with year 2015 and the protocol did not provide any guidance on how ice sheet modellers  
59 should initialise to that starting point. This led to a wide range of ice sheet histories preceding the GrIS projections, with  
60 numerous models not matching observed mass changes (Goelzer et al., 2020a; The IMBIE Team, 2020; Aschwanden et al.,  
61 2021). This also created challenges in presenting and combining projected ice sheet changes with observed changes in the  
62 AR6. Since then, it has become a priority to consider the historical experiment as part of the simulation and it has been shown  
63 that extending the ISMIP6 forcing protocol over the historical period can produce GrIS projections consistent with observed  
64 mass changes (Rahlfes et al., 2025a).

65 While physically-based ice sheet model simulations like those produced by ISMIP6 now form an important basis of sea-level  
66 change projections, statistical tools are needed to generalise the results and make meaningful inferences. This need arises  
67


largely from the limited sampling of climate model forcing, model physics and parameter choices that remain relatively sparse due to practical limitations, computational cost and feasibility. A consequence is that some projections nowadays heavily rely on emulators (e.g. Edwards et al., 2021; Rohmer et al. 2022; Rohmer et al. 2025; Edwards et al., 2026) to help their interpretation. Designing ice sheet experiments that can serve both direct interpretation of the result and feeding into emulators has become an important consideration.

This paper presents a new set of physically-based GrIS sea-level projections designed to extend the ISMIP6 effort in several aspects and to inform a next generation of emulator-based projections. We describe the experimental protocol, forcing and models (Sect. 2), present results (Sect. 3) and close with a discussion (Sect. 4) and conclusions (Sect. 5).

## 2 Experimental setup

### 2.1 Experimental protocol

The ice sheet model experiments largely follow the ISMIP6 protocol for GrIS projections, which is documented in detail elsewhere (Nowicki et al., 2016, 2020; Goelzer et al., 2018, 2020a). Here, we provide a brief summary of the main principles and differences. Output from selected CMIP Earth system models (ESMs) serves as boundary conditions for i) RCMs (producing the surface mass and energy balance and near-surface climate fields) and ii) for a retreat parameterization for marine-terminating outlet glaciers (Slater et al., 2019; 2020). These, in turn, provide forcing for ice sheet models (Figure 1).



**Figure 1.** General forcing approach for Greenland ice sheet model projections

88 **2.1.1 Forcing approach**


89 The forcing from three RCMs, namely MAR (Delhasse et al., 2020), HIRHAM (Motttram et al., 2017) and RACMO (Noël et  
90 al., 2018) and from a statistical downscaling approach (Noël et al., 2022), is provided in the form of annual surface mass  
91 balance (SMB) and surface temperature (ST) anomalies relative to the period 1960-1989 (red box in Figure 2). In addition, we  
92 provide estimates of local annual vertical SMB and ST gradients, that are used to propagate dynamic ice sheet elevation  
93 changes when updating SMB and ST. The SMB forcing applied in the ice sheet model at a given time t is:

94

$$95 \text{SMB}(x,y,t) = \text{SMBref}(x,y) + a\text{SMB}(x,y,t) + d\text{SMBdz}(x,y,t) * dz(t), \tag{1}$$

96

97 Where SMBref [mm yr<sup>-1</sup>] is the surface mass balance used by each individual ice sheet model during initialisation, aSMB [mm  
98 yr<sup>-1</sup>] is the SMB anomaly, dSMBdz [mm yr<sup>-1</sup> m<sup>-1</sup>] is the vertical gradient of SMB and dz [m] is the elevation change since the  
99 start of the experiment. A similar approach applies to ST that can be used as boundary condition for the evolution of ice  
100 temperature in the model. Differences in SMB and ST stem from the various ESMs and scenarios that are downscaled.  
101 The retreat of marine-terminating outlet glaciers is parameterised as an empirically derived function of ocean thermal forcing  
102 (from the ESMs) and runoff (from the RCMs), both identified as the main drivers for melting of marine-based calving fronts  
103 (green box in Figure 2, Slater et al., 2019; 2020). This is a relatively crude approximation for the complex interaction between  
104 glaciers and the ocean that is poorly understood and difficult to resolve in large-scale ice sheet models. The uncertainty in this  
105 forcing is captured by a retreat parameter κ that can be expressed probabilistically and is sampled at its median (med), 25<sup>th</sup>  
106 (high) and 75<sup>th</sup> (low) percentile value (like in ISMIP6) and in extension at its 5<sup>th</sup> and 95<sup>th</sup> percentile value (cf. Figure 5a in  
107 Slater et al., 2019). In some cases, we have added control experiments without any prescribed retreat that are labelled ‘pno’.  
108



110 **Figure 2.** Surface mass balance and retreat forcing

## 111 2.2 Regional climate model forcing

112 The emphasis of this project is to extend the range of available forcings to a larger number of CMIP6 ESMs, scenarios (SSP1-  
 113 2.6, SSP2-4.5, SSP5-8.5) and to provide surface mass balance forcing from several RCMs: MAR (Delhasse et al., 2020),  
 114 RACMO (Noël et al., 2018) and HIRHAM (Mottram et al., 2017). In addition, we use forcing produced by a statistical  
 115 downscaling approach (SDBN1, Noël et al., 2016, 2020, 2022), which has been used here to translate ESM forcing from  
 116 CESM2-WACCM directly to the ice sheet scale. We consider this approach similar in capability to an RCM in terms of the  
 117 forcing it provides. An overview of available forcing data is given in Table 1. Corresponding retreat mask forcing can be  
 118 constructed given sufficient climate model output and additional ESM ocean data, typically retrieved from the CMIP archives  
 119 (<https://esgf-node.llnl.gov/projects/cmip6/>). Forcing with MAR version 3.9 was produced for ISMIP6 and remained available  
 120 for ice sheet simulations under PROTECT.

121

122 **Table 1.** SMB forcing data available for ice sheet modellers. MARv3.9 output was produced for ISMIP6.

CMIP	ESM	SSP1-2.6 RCP2.6	SSP2-4.5	SSP5-8.5 RCP8.5
CMIP6	CESM2	MARv3.12 RACMO2.3p2	MARv3.12 RACMO2.3p2	MARv3.12
	CESM2-Leo <sup>†</sup>			MARv3.9 MARv3.12 RACMO2.3p2 HIRHAM5
	CESM2-WACCM	SDBN1 <sup>‡</sup>		SDBN1 <sup>3</sup>
	CNRM-CM6-1			MARv3.9 MARv3.12
	CNRM-ESM2-1			MARv3.9 MARv3.12
	EC-Earth3	HIRHAM5		HIRHAM5
	IPSL-CM6A-LR			MARv3.12
	MPI-ESM1	MARv3.12	MARv3.12	MARv3.12
	NorESM2-MM		MARv3.12	MARv3.12
	UKESM1-0-LL		MARv3.12	MARv3.12
	UKESM1-0-LL-Robin <sup>†</sup>			MARv3.12
CMIP5	ACCESS1.3			MARv3.9 MARv3.12
	CSIRO-Mk3.6.0			MARv3.9
	HadGEM2-ES			MARv3.9
	IPSL-CM5-MR			MARv3.9

	MIROC5	MARv3.9		MARv3.9
	NorESM1-M			MARv3.9

<sup>†</sup> pre-CMIP6 ensemble member. <sup>‡</sup>Direct statistical downscaling of CESM2-WACCM (Noël et al., 2022).

### 2.2.1 Required climate model output data

The data required to produce ice sheet model forcing were developed during ISMIP6 in collaboration with the developers of MAR, the only RCM used to generate projections for the project at the time. This includes extension of the RCM forcing beyond the observed ice sheet mask and producing output needed for vertical adjustment of the forcing to a changing ice sheet topography. In MAR this is done with the same statistical downscaling method used to produce results at 1 km resolution (Franco et al., 2012) as done in the GrSMBMIP intercomparison (Fettweis et al., 2020). In RACMO and SDBN1, vertical gradients were estimated following Noël et al. (2016) combining statistically downscaled SMB components with surface elevation and ice mask from the Greenland Ice Mapping Project (GIMP) DEM (Howat et al., 2014), down-sampled to 1 km spatial resolution. Vertical gradients were first computed on ice-covered grid cells using SMB components and surface elevation of the current grid-cell and at least five (up to eight) neighbours and further extrapolated outside the ice sheet to cover the tundra region. In HIRHAM5, gradients are produced at a 5 km horizontal resolution using an updated subsurface scheme (Langen et al., 2017). These gradients are subsequently bilinearly interpolated to the 1 km MAR grid. Outside the observed ice mask, extrapolation to cover the tundra is performed via distance-weighted averaging, followed by smoothing using weighted averages of the grid points, including the eight surrounding points.

To facilitate use of RCMs and other downscaled climate forcing in PROTECT and other projects, we outline a detailed list of required data in Appendix A.

### 2.3 Forcing dataset preparation

Output from RCMs and ESMs is collected and processed using methods established during ISMIP6. The aim is to provide a consistent forcing dataset for ice sheet modellers in a familiar format. It requires interpolation of RCM output to a common grid at 1 km resolution, calculating anomalies and adjusting units and file formats. Retreat mask forcing is produced based on the initial ice sheet mask for each individual participating ice sheet model and version. All data are provided in NetCDF format following the ISMIP6 guidelines (<https://thegithub.org/groups/ismip6/wiki/ISMIP6-Projections-Greenland>).

### 2.4 Participating ice sheet models

The ensemble includes four numerical ice sheet models that are routinely run by the participating partners for GrIS simulations (IMAU, VUB, IGE, NORCE). A brief overview of the model characteristics is given in Table 2 and short model descriptions are given below.

151 **Table 2.** Ice sheet model names and characteristics. SIA - Shallow ice approximation to the force balance, SSA - Shallow shelf  
152 approximation, HO – higher order approximation (Fürst et al. 2013), DIVA - variationally derived, depth-integrated approximation  
153 (Goldberg 2011).

Group-Model	Type	Resolutions (km)	Variants
IMAU-IMAUICE	SIA-SSA, regular grid	10, 16, 20, 30, 40	Sliding law, spinup
VUB-GISM	Regular grid	5	HO, SIA
IGE-ELMER	SSA, finite element	1 – 6 (variable)	Sliding law
NORCE-CISM	DIVA, regular grid	2, 4, 8, 16	Initial and historical SMB

154

#### 155 2.4.1 IMAU-IMAUICE

156 The model (Berends et al., 2022) is initialised using a hybrid approach, combining a basal inversion method (Berends et al.,  
157 2023) with a paleoclimate spin-up. During the inversion phase of the initialisation, spatial patterns in basal slipperiness are  
158 iteratively adjusted until the modelled ice sheet reaches a stable state that closely matches the observed present-day ice sheet  
159 geometry (Morlighem et al., 2017) and surface velocity (doi: 10.24381/cds.0b96b838). The prescribed climate is fixed at  
160 present-day conditions: monthly mean values of 2-m air temperature and total precipitation, which are obtained from the 1950-  
161 1980 mean of the ERA5 reanalysis (Hersbach et al., 2020). The SMB is calculated from these quantities using the IMAU-ITM  
162 model, which is calibrated to RACMO2.3p2 over the 1979-2014 period (Fettweis et al., 2020). The steady-state geometry and  
163 basal slipperiness resulting from the inversion phase are then used to initialise the model during the last interglacial, 120,000  
164 years ago. The climate evolution of the last glacial cycle is then prescribed using a matrix method (Berends et al., 2018), based  
165 on different pre-calculated GCM output for the different IMAU-ICE versions: either HadCM3 (Singarayer and Valdes, 2010),  
166 CCSM (Brady et al., 2013), or the PMIP3 best-performing ensemble mean (Scherrenberg et al., 2023). Climate evolution  
167 during the historical period is approximated by forcing the climate matrix with the Law Dome ice-core CO<sub>2</sub> record (MacFarling  
168 Meure et al., 2006), subjected to a 60-year smoothing representing the delayed response of the climate to changes in CO<sub>2</sub>.

#### 169 2.4.2 VUB-GISM

170 VUB-GISM (Huybrechts, 2002; Fürst et al., 2013; 2015) is configured either with the higher order or a shallow ice  
171 approximation to the force balance. GISM was initialised to the present-day geometry by assimilation of the observed ice  
172 thickness (Le clec'h et al., 2019). A steady state was assumed for the starting date of December 1989 using the 1960–1989  
173 mean SMB from MAR forced by the ERA5 meteorological reanalysis climate. The iterative initialization method optimised  
174 both the basal sliding coefficient in unfrozen areas and the rate factor in Glen's flow law for frozen areas. The ice temperature  
175 and the initial velocity field needed in the initialization procedure were derived from a glacial spin-up with a freely evolving  
176 geometry over the last two glacial cycles with a synthesised temperature record based on ice-core data from Dome C, NGRIP,  
177 GRIP and GISP2 (Fürst et al., 2015). For this spin-up experiment, a PDD model was used with an observed precipitation field

178 derived from the Bales et al. (2009) surface accumulation for the period 1950–2000 and scaled by 5% per degree. The ice  
179 temperature and velocity fields from the “free geometry present-day” were rescaled to the observed ice thickness (Morlighem  
180 et al., 2017) and excluded peripheral ice (Citterio et al., 2013). The historical experiment is run from January 1990 to December  
181 2014 using the yearly SMB from MAR forced by ERA5 meteorological reanalysis. For the projections, the standard retreat  
182 forcing from the ISMIP6 protocol is applied.

### 183 **2.4.3 IGE-ELMER**

184 The model is initialised using an inverse control method as in Gillet-Chaulet et al. (2012) to calibrate the basal friction  
185 coefficient field. For the momentum equations, we solve the shelfy-stream approximation with a sub-grid parameterization of  
186 the friction for partially grounded elements. The vertically averaged viscosity is constant in all simulations and is initialised  
187 using the temperature field coming from a palaeo-spin-up (125 kyr) of the SICOPOLIS model. The basal friction coefficient  
188 is constant in all transient simulations and is initialised with the control method so that the mismatch between observed and  
189 modelled surface velocities is minimum. As observations, we use a composite from the NASA Making Earth System Data  
190 Records for Use in Research Environments (MEaSUREs) Greenland Ice Sheet Velocity Map (V1) (Joughin et al., 2010). The  
191 ice sheet topography is initialised using the IceBridge BedMachine Greenland V3 data set (Morlighem et al., 2017). The ice  
192 sheet model is then relaxed for 20 years using a constant surface mass balance given by the 1960–1989 mean SMB from the  
193 regional climate model MAR v3.12 forced with ERA5 (Fettweis et al., 2017). The calving front positions are fixed during the  
194 relaxation. We use an anisotropic mesh with a horizontal resolution ranging for 1 to 6 km. For the projections, the standard  
195 ISMIP6 protocol is applied and we test the sensitivity to different friction laws: a linear friction law, Weertman friction law  
196 with  $m=1/3$  and a parameterised Coulomb friction law.

### 197 **2.4.4 NORCE-CISM**

198 The Community Ice Sheet Model (CISM; Lipscomb et al., 2019) is run using a depth-integrated higher-order velocity solver  
199 based on Goldberg (2011) and a basal-sliding law based on Schoof (2005). The ice sheet is initialised with present-day  
200 thickness and bed topography (Morlighem et al., 2017) and an idealised temperature profile. CISM is then spun up for 5 000  
201 years with surface mass balance and surface temperature from a 1960–1989 climatology provided by the MAR regional climate  
202 model (Fettweis et al., 2017) and with basal heat fluxes from Shapiro and Ritzwoller (2004). During the spin-up, the model is  
203 nudged toward present-day thickness by adjusting friction coefficients in a basal-sliding power law. There is no dependence  
204 of basal sliding on basal temperature or water pressure. All floating ice is assumed to calve immediately. For partly grounded  
205 cells at the marine margin, basal shear stress is weighted using a grounding-line parameterization. By the end of the spin-up,  
206 the ice thickness, temperature and velocity fields are very close to steady-state and closely match the provided observed  
207 geometry and also the observed horizontal velocity, which is not used during initialisation. For the historical period (1960–  
208 2014), the model is run forward with SMB and surface temperature anomalies, including lapse-rate corrections, from the MAR  
209 simulation that provided the background climatology and with retreat forcing of various sensitivities. Basal friction coefficients




210 are held fixed at the values obtained during the spin-up. The different CISM model versions used here differ by the horizontal  
211 grid resolution (2 - 16 km), by the RCM version used for spinup and historical run (MARv3.9 vs MARv3.12) and by the  
212 sensitivity of the retreat parameterisation applied over the historical period.  
213

## 214 **2.5 Experiments**

### 215 **2.5.1 Ice sheet initialisation and historical run**

216 Under ISMIP6 protocol, ice sheet modellers were free to initialise their model as they wish, with the aim to produce a present-  
217 day state of the ice sheet that is close to observations. This procedure may involve a historical experiment that brings the ice  
218 sheet into a state that is assigned to the end of 2014. In contrast to this freedom in setting up the model, the projections 2015-  
219 2100 that then follow are very tightly constrained by the forcing. This is also the case for the retreat forcing, which takes the  
220 individual 2014 ice mask as a reference and provides masks that impose the position of the (retreated) calving fronts forward  
221 in time. For PROTECT we have extended this approach by providing retreat forcing before 2015 that is calculated from  
222 reconstructions of past runoff and ocean thermal forcing (see Rahlves et al., 2025a). This allows for a consistent forcing of the  
223 models in past and future and considers historical retreat of the outlet glaciers, which was an important source of mass loss  
224 after 1990 (The IMBIE Team, 2020). We can now interpret the experiment leading up to 2015 as a real historical simulation.  
225 The ISMIP6 practice of removing the results of an unforced control experiment from the projections is therefore not needed  
226 here. Figure 3 illustrates the 2014 state of selected model versions in comparison with observations (BedMachine v3,  
227 Morlighem et al., 2017).

228 The practice of including the historical experiment as part of the experimental design (which was not the case for ISMIP6)  
229 should ultimately imply that any variation in the ISM modelling choices should be represented in this experiment. As a  
230 consequence, each model variant would in principle require a separate historical experiment, so that modelling choices remain  
231 consistent at the beginning of the projections (here in year 2015). At the beginning of the project, we did not apply this  
232 constraint and most ice sheet model runs were conducted with a single historical experiment (like in ISMIP6) with medium  
233 retreat sensitivity, which then branches into projections with different sensitivity. We later conducted some experiments with  
234 consistent retreat forcing sensitivity with NORCE-CISM (cf. list of ISM experiments in Appendix B).



**Figure 3.** Ice thickness comparison. (a) Present day observations (Morlighem et al., 2017). (b-e) Difference of modelled 2014 ice thickness compared to observations for one model version per group. The root mean square (RMS) difference to the observations in m is given in the lower right of panels b-e.

## 2.5.2 Future projections to year 2100

The future projections from 2015 to 2100 follow the ISMIP6 forcing protocol with SMB anomalies and retreat forcing applied as described in Sec. 2.1.1. With the available forcing described in Sec. 2.3, we obtained output from 14 different global models, forced with three different scenarios (SSP1-2.6, SSP2-4.5, SSP5-8.5) and downscaled with three RCMs and one statistical downscaling method.

## 2.5.3 Extensions after year 2100

Few CMIP6 models have carried out the scenarioMIP extensions (O'Neill et al., 2016) to 2300, and even fewer have provided 6-hourly output typically required for RCMs to downscale the data. We have currently only three examples of ice sheet forcing with what we will refer to as ‘natural extensions’ beyond 2100 from the ESMs IPSL-CM6A-LR (for scenarioMIP SSP5-8.5-ext) and CESM2-WACCM (for scenarioMIP SSP5-8.5-ext and scenarioMIP SSP1-2.6-ext). While SSP1-2.6-ext stabilises to a CO<sub>2</sub> concentration well below 500 ppm, SSP5-8.5-ext stabilises towards a CO<sub>2</sub> concentration of about 2200 ppm, roughly double its value at year 2100. CESM2-WACCM has been statistically downscaled with SDBN1 (not requiring 6-hourly output) and IPSL-CM6A-LR has been dynamically downscaled with variants of MARv3.12. There is one extension to 2200 with MARv3.12 downscaling IPSL-CM6A-LR under scenario SSP5-8.5/SSP5-8.5-ext using the same approach as for the other experiments. The MAR modellers questioned the validity of continuing to downscale the relatively strong climate forcing from IPSL-CM6A-LR SSP5-8.5-ext at a fixed present-day topography beyond 2200, given that the ice sheet geometry should have considerably changed by then, hence impacting SMB (Delhasse et al., 2024). We have therefore performed two additional pilot experiments with different topography updates extending to 2300 (MARv3.13-e05 and MARv3.13-e55). The construction of these forcings is described in more detail in Appendix C. The retreat mask forcing can in principle be constructed in the same way as for the experiments extending to 2100. However, the underlying assumptions of the parameterisation may not hold for the very large retreat distances produced under sustained very strong warming to 2300. Because of that we have

260 already limited the retreat sensitivity to the 25-75 percentile range for the natural extensions, but caution that these simulations  
261 show higher uncertainty.


262 In addition to the natural extensions, we have designed schematic extensions of the forcing data to the year 2300 to evaluate  
263 the longer-term response of the ice sheet for a broader range of ESMs. The first set of extensions is carried out by repeating  
264 the forcing of the last ten available years (2091-2100) in randomised order and keeping the retreat mask of year 2100 constant.  
265 For a second type of schematic extension, we have designed overshoot scenarios mimicking SSP5-3.4-OS by reusing the  
266 regular SSP5-8.5 forcing before 2100 and simulating a climate cooling and corresponding increase of the SMB until 2300.  
267 These overshoot scenarios are constructed using global mean temperature as a proxy for the temperature and SMB evolution  
268 by sampling existing yearly forcing data until 2055 and reorganising them to new time series until 2300. The shape of the  
269 global temperature proxy evolution is parameterised and has been calibrated to a few existing ESM results (CESM2-WACCM,  
270 IPSL-CM6A-LR, MRI-ESM2-0) for overshoot scenario SSP5-3.4-OS. The resulting time series are illustrated in Appendix C.  
271 Aside from the obvious shortcoming that the latter two are schematic extensions, the formulation of the retreat forcing implies  
272 a constant mask for stabilising the forcing, which may underestimate the retreat. Furthermore, another problem on this  
273 timescale in general may be that the climate response to changing ice sheet geometry is not properly accounted for. Alternative  
274 prolongations could be envisioned and thus the current approaches should be considered pilot experiments and not a guide to  
275 produce realistic scenarios.

## 276 **2.6 Ensemble design**

277 The collection of forcing data covers a wide range of variations across different ESMs and greenhouse gas (GHG) scenarios,  
278 but ultimately represents an ‘ensemble of opportunity’. This is even more true for the selection of RCMs and ISMs, which is  
279 limited to available models in the consortium. PROTECT has therefore conceptualised and operated a modelling strategy from  
280 the beginning that embeds the physically-based modelling into a wider framework allowing for a statistically meaningful  
281 probabilistic interpretation of the results.

282 In order to facilitate the sampling strategy in that framework, experiments in the ensemble are labelled by 6 characteristics that  
283 are colour-coded in Figure 4 and given in square brackets in the following. Setting up a specific ice sheet simulation requires  
284 climate forcing (SMB and ST) for a given choice [Global model, GHG scenario, Regional model] and retreat mask forcing for  
285 a given choice [Global model, GHG scenario, Regional model, Retreat sensitivity]. We furthermore have different ice sheet  
286 models built on a certain [Code base] (here referred to by the ISM name) and they are operated using certain modelling  
287 [Choices] (initialization strategy, approximations, parameterizations, parameter choices). In our current approach, different  
288 sets of modelling choices are summarised and assigned to a specific model version number. However, the impact of specific  
289 modelling choices could be further analysed e.g., by using the technique described by Rohmer et al. (2022; 2025).

290



**Figure 4.** Forcing and model options relevant for the larger ensemble design.


The current set of results discussed below is a broad sampling of the available forcings and parameter choices, intended to cover a wide range of possible projections and their uncertainties. Based on feedback from the researchers running emulators using these results, we have iteratively updated the ensemble with additional simulations to refine the sampling for specific choices where needed. The repeated extensions and overshoot scenarios are examples of additional experiments that were deemed important to improve the emulator performance for predictions up to 2300.

### 3 Results

The following results are presented as an overview of available ISM simulations and provide insight into the typical ranges and main uncertainties. We have produced 1472 individual ice sheet model projections that form the ensemble of GrIS results. An overview of the used ISM model versions is given in Table A2. Sea-level contributions are calculated taking into account density (and bedrock adjustment for IMAUICE) following Goelzer et al. (2020b).


Figure 5 illustrates the typical time-dependence of the projections from the ensemble, with output from one model version per group under the range of ESM and RCM forcing with median outlet glacier retreat sensitivity. It also shows historical simulations of various lengths for the different ISMs. Under this forcing, which includes scenarios SSP1-2.6, SSP2-4.5 and SSP5-8.5 for various ESMs, all sea-level contributions are increasing and positive by the year 2100. Judging by average mass loss rates over the last 30 years, none of the simulations shows signs of ice sheet stabilisation (zero or positive mass change)

309 towards the end of the experiment, but rather continued mass loss, suggesting larger to much larger contributions for time  
 310 scales beyond 2100.



311  
 312 **Figure 5.** Projected sea-level contribution from the GrIS until 2100 from a subset of experiments with all four participating ice sheet models  
 313 (one model version per group), median retreat sensitivity and forcings produced specifically for PROTECT (MARv3.12, RACMO2.3p2,  
 314 HIRHAM5). The aim of this figure is to illustrate the range and distribution of the projections, not individual members.


315 Results for the year 2100 of the whole ensemble of projections with all available scenarios, ESMs, RCMs and ISMs under five  
 316 different retreat sensitivities (5th percentile, high, med, low, 95th percentile) are summarised in Figure 6. We have not included  
 317 results for the experiments that continue to 2300 here, which are instead shown in Figure 9 with results at the respective ends  
 318 of the simulations. The contributions in the year 2100 (relative to 2014) lie in a range between 16 and 354 mm, with the largest  
 319 numbers from experiments that combine high climate sensitivity (UKESM1-0-LL variants) and very high retreat sensitivity  
 320 (5th percentile). The corresponding global mean temperature anomalies as diagnosed from the ESMs are given in Figure S1  
 321 in the supplement.



**Figure 6.** Overview of produced GrIS sea-level projections for the year 2100 from 4 ice sheet models (23 different model versions) and 5 retreat parameter settings. Triangular light green markers for NORCE-CISM indicate experiments with extreme values of the retreat parameter  $\kappa$  in the 5<sup>th</sup> percentile (upward-pointing) and 95<sup>th</sup> percentile (downward-pointing).

Figure 7 illustrates ISM results of the runs to 2100 sorted by different categories. The comparison between ISMs, RCMs, scenarios and CMIP iterations shows primarily the sampling frequency across the ensemble. Unequal sampling limits the direct interpretation of the results, but some conclusions can be drawn, nevertheless. The range of results for the different ISMs is largely similar (Figure 7a) and only larger for CISM because a wider range of retreat parameters (5<sup>th</sup> - 95<sup>th</sup> percentile range) was sampled with this model. Simulations forced with regional models MAR and HIRHAM5 (Figure 7b) show higher contributions under high climate forcing compared to RACMO, which is in line with SMB results discussed recently (Glaude et al, 2024): the contrasted response to warming of the utilised RCMs primarily stems from differences in projected runoff, which is amplified by the positive melt-albedo feedback.

335 The full scenario ranges (Figure 7c) of projected sea-level contributions from the GrIS by the year 2100 (relative to year 2014)  
 336 are 16-76 mm (SSP1-2.6/RCP2.6), 22-163 mm (SSP2-4.5) and 27-354 mm (SSP5-8.5/RCP8.5). For the narrower range of the  
 337 retreat parameter (25<sup>th</sup> - 75<sup>th</sup> percentile range as in ISMIP6 and performed by all ISMs), the (upper) scenario ranges are reduced  
 338 to 22-127 mm (SSP2-4.5) and 27-265 mm (SSP5-8.5/RCP8.5). In summary, these results indicate a very large range of sea-  
 339 level contributions in particular under forcing scenario RCP8.5/SSP5-8.5. Figure 7d shows an increased sensitivity from  
 340 CMIP5 to CMIP6, confirming earlier results (e.g. Hofer et al., 2020; Payne et al., 2021), although unequal sampling is an  
 341 additional factor.  
 342




343 **Figure 7.** Sea-level contribution by year 2100 from the GrIS sorted by a) ice sheet model, b) regional climate model, c) scenario and d)  
 344 CMIP ensemble. The colour legend is the same for all panels. Scenarios labelled ‘2.6’ in c) include SSP1-2.6 and RCP2.6 and ‘8.5’ includes  
 345 SSP5-8.5 and RCP8.5. Triangular light green markers for NORCE-CISM indicate experiments with extreme values of the retreat parameter  
 346  $\kappa$  in the 5<sup>th</sup> percentile (upward-pointing) and 95<sup>th</sup> percentile (downward-pointing).  
 347

348

349 Uncertainty in the projections arises from the climate forcing (different ESMs, scenarios), the translation of the forcing to the  
 350 ice sheet scale (RCMs/downscaling, retreat parameterisation) and from the ISMs themselves. We have quantified these  
 351 uncertainty ranges by comparing experiments with one of the factors modified at a time and averaging over available subsets.  
 352 Under SSP5-8.5/RCP8.5 forcing, the ESM choice explains a range of 130 mm (cf. Figure 7c), compared to a range of 84 mm  
 353 for RCMs (cf. Figure 7b), 50 mm for ISMs (cf. Figure 7a) and 13 mm for retreat forcing (25<sup>th</sup> - 75<sup>th</sup> percentile range).  
 354 Figure 8a shows results for a schematic prolongation to 2300 for one of the ice sheet models with repeated SMB forcing and  
 355 constant retreat mask after year 2100. It illustrates that sea-level contributions from Greenland continue to increase well beyond  
 356 year 2100 even under stabilised forcing. Contributions can exceed 1.2 m (under very high retreat forcing) by 2300 for  
 357 prolonged SSP5-8.5/RCP8.5 but may stabilise for prolonged SSP1-2.6/RCP2.6 somewhere below 200 mm. The scenario  
 358 ranges with repeated forcing are 58-163 mm (repeated SSP1-2.6), 98-218 mm (repeated SSP2-4.5) and 282-1230 mm (repeated  
 359 SSP5-8.5).  
 360 Results from the schematic overshoot scenarios, mimicking SSP5-3.4-OS (Figure 8b) with sea-level contributions at year 2300  
 361 in a range between 49 and 201 mm, show stabilisation for three out of the nine experiments (CESM2-WACCM SDBN1,  
 362 CESM2-Leo RACMO2.3p2, NorESM2-MM MARv3.12), while the others have an ongoing near-linear mass loss trend at the  
 363 end of the experiments by 2300. The natural extensions to 2300 (Figure 8c) for CESM2-WACCM SDBN1 show a range  
 364 between 92 mm (SSP1-2.6) and 3127 mm (SSP5-8.5), indicating a strong dependence on the climate forcing, large  
 365 uncertainties and a potentially very large long-term response. Results for IPSL-CM6A-LR SSP5-8.5-ext show that including  
 366 a topography update (MARv3.13-e55) leads to a 6% larger contribution in 2300 compared to calculating the SMB for a fixed  
 367 surface elevation (MARv3.13-e50). This is in addition to the parameterised SMB-height feedback active in both experiments.  
 368 For the natural extensions (Figure 8c) we also show the corresponding global mean temperature anomalies as diagnosed from  
 369 the ESMs (Figure 8d) to put the results for the extreme warming scenarios into perspective.






370

371 **Figure 8.** Extensions to 2300 with NORCE-CISM for various ESMs/RCMs with median values of retreat parameter  $\kappa$  for a) repeated forcing  
 372 after 2100 and b) overshoot scenario mimicking SSP5-3.4-OS. c) Natural extensions with CMIP6 forcing until 2300 with median (solid),  
 373 high(dot-dashed) and low (dotted) values of retreat parameter  $\kappa$ . For scenario SSP1-2.6, experiments with various values for  $\kappa$  are largely  
 374 overlapping and difficult to distinguish. Extensions to 2200 are overlapping with the respective continuations to 2300 and are not shown. d)  
 375 ESM global mean temperature anomaly relative to 1960-1989 for the experiments in c.

376 Figure 9 summarizes results at the end of the experiments for all schematic prolongations to 2300 (overshoot: o2300 and  
 377 repeat: r2300) and also includes the natural extensions to 2300 for CESM2-WACCM and IPSL-CM6A-LR. Note that results  
 378 are displayed on a logarithmic vertical axis.  
 379



**Figure 9.** Results for extensions to the year 2300 with NORCE-CISM for various ESMs/RCMs. o2300 - overshoot scenario, r2300 - repeated forcing, x2300 – regular ScenarioMIP extension. The rightmost experiment forced with MARv3.13-e55 has a topography update during the MAR simulation. Note the logarithmic vertical axis.

## 4 Discussion

For the scenarios and forcings covered in both ensembles and for the same range of the retreat parameter (25<sup>th</sup> - 75<sup>th</sup> percentile range), our ranges of projected sea-level contributions at 2100 (16-67 mm for SSP1-2.6/RCP2.6, 27-265 mm for SSP5-8.5/RCP8.5) largely overlap with ISMIP6 results (11-58 mm for SSP1-2.6/RCP2.6, 35-250 mm for SSP5-8.5/RCP8.5; Payne et al., 2021). Slightly different ranges here are due to using a subset of ISMs, but also due to the incorporation of additional ESMs and RCMs. Including a wider range of the retreat parameter (5<sup>th</sup> - 95<sup>th</sup> percentile range) has led to a larger upper end of the full scenario ranges presented here as 16-76 mm (SSP1-2.6/RCP2.6), 22-163 mm (SSP2-4.5) and 27-354 mm (SSP5-8.5/RCP8.5). We have added 159 experiments for the intermediate scenario SSP2-4.5, that was not represented in ISMIP6. Inclusion of these results of an intermediate scenario does not increase the total range of projections but adds additional information for possible future emulation. Results for experiments with the same climate model (CESM2-Leo) but different RCMs (MAR, RACMO, HIRHAM) mirror the results from a comparison of the underlying SMB (Glaude et al., 2024), with considerable differences in the projected sea-level contribution due to the choice of RCM. In addition, a larger relative contribution from experiments forced with HIRHAM here compared to the SMB results (Glaude et al., 2024) is related to the

way SMB is extended beyond the ice sheet mask and how the vertical gradients are determined for parameterising the SMB-height feedback. In combination, this highlights the urgent need to include uncertainty due to climate downscaling from global to ice sheet scale in the projections, which was likely under-represented in ISMIP6 due to the use of only one RCM and only one method to take the SMB-height feedback into account.

Uncertainties in the projections in this ‘ensemble of opportunity’ arise from sampling of ESMs, RCMs, ISMs and retreat parameters, which implies that statistically meaningful interpretation of the raw model output is challenging. We have therefore mostly limited the interpretation of results to typical ranges and leave finer-grained analysis to downstream efforts (e.g. Rohmer et al., 2024, 2025; Edwards et al., 2021; 2026). Under the high forcing scenario SSP5-8.5/RCP8.5, global climate model uncertainty (here choice of ESM) dominates and explains a total range of 130 mm in the projections to 2100. This is compared to a range of 84 mm for choice of RCM (not sampled in ISMIP6), and 50 mm for the choice of ISM, which is similar to ISMIP6, despite the smaller number of ISMs in the present work. The range of 13 mm for retreat forcing (25<sup>th</sup> - 75<sup>th</sup> percentile range) is slightly smaller compared to ISMIP6 (19 mm), but increases considerably to 52 mm when extending to the 5<sup>th</sup> - 95<sup>th</sup> percentile range that we have additionally explored here.

Extending the forcing and the simulations backwards over the historical period is an important improvement compared to ISMIP6 and will eventually allow for a better comparison between observations and models. We have not attempted here to perform specific experiments that quantify the effect of including a real historical experiment on the projections, but results by Rahlves et al. (2025a) give indications that the impact on the projected sea-level contribution is minor.

Schematic extensions with repeated forcing to 2300 from a subset of ESMs (and only one ice sheet model) show an upper range of contributions exceeding 1.2 m for prolonged SSP5-8.5/RCP8.5 and potentially stabilising contributions for prolonged SSP1-2.6/RCP2.6 below 25 cm. These results are given with the caveat of the underlying schematic experimental setup and a limited ensemble size. In comparison, regular ScenarioMIP extensions under scenario SSP5-8.5-ext that we have for two global models (IPSL-CM6A-LR, CESM2-WACCM) produce contributions in the year 2300 exceeding 2.5 m and 3 m, respectively. This is in strong contrast to results under CESM2-WACCM SSP1-2.6-ext with only 92 mm, underlining that the climate scenario is the dominant source of uncertainty on this timescale. We also emphasise that the natural extensions to 2300 lead to considerably higher contributions compared to the extensions with repeated forcing, indicating that these scenarios are very different and shouldn’t be conflated. It also underlines an urgent need for more ESM output going to 2300 or even beyond. On these timescales and under such high forcing, feedbacks between ice sheet and climate and how they are taken into account become first-order effects and introduce large uncertainties. In our standalone ice sheet modelling approach, the lack of proper climate feedbacks is an important limitation that may be addressed with interactive coupling of ESMs and ISMs (e.g. Muntjewerf et al., 2020; Smith et al., 2021; Goelzer et al., 2025). In addition, the retreat forcing approach has to be considered with caution for extended time periods in particular under high forcing scenarios. Combined, these results indicate that modifications to the ISMIP6 forcing protocols and new methods to account for a changing ice sheet geometry (e.g. Goelzer et al., 2020c; Delhasse et al., 2024; Rahlves et al., 2025b) are needed for robust standalone ice sheet simulations well beyond year 2100. Nevertheless, the experiments with repeated forcing give an approximate idea of how stabilising forcing (and

climate) at different levels could play out. On the considered timescale, stabilising the forcing has the effect of stabilising the rate of change, not the ice sheet itself (unless the rate is close to zero). Results from the schematic overshoot scenarios, mimicking SSP5-3.4-OS, were added specifically to provide an emulator with additional, complementary information on ice sheet changes under forcing that does not follow a continuous increase in temperature. Results under this forcing show that three (CESM2-WACCM SDBN1, CESM2-Leo RACMO2.3p2, NorESM2-MM MARv3.12) out of the nine experiments with different climate model forcing produce what seems like a stabilising GrIS towards 2300. Creating this ensemble with a relatively small group of ice sheet modellers bears the risk of underestimating an important part of the ISM uncertainty. We anticipate this potential gap to be closed by ISMIP7 and other follow-up initiatives. The advantage of a smaller group of modelers that we have exploited in this work lies in a more flexible and adaptable experimental design.

## 5 Conclusions

We have produced a large ensemble of Greenland ice sheet projections with four different ice sheet models under various forcings drawn from a wide range of ESMs, scenarios, RCMs, and retreat parameters. Uncertainty in the ice sheet models is furthermore sampled with various model versions that differ by horizontal grid resolution, applied sliding law, and initial states. Under high forcing, the largest contributor to the uncertainty is the choice of ESM, followed by the RCM and ISM. RCM uncertainty, or more generally, uncertainties in the climate downscaling process need to be better quantified in the future. This contribution to the European project PROTECT extends the projections of ISMIP6 in several important regards, with an additional, intermediate scenario, several different RCMs, and more CMIP6 models. Results from different extensions up to 2300 give a perspective on challenges for standalone simulations on this time scale.

## Appendix A: Required climate model output for ice sheet forcing

This section describes the climate model output required to construct ISMIP6-type ice sheet forcing for Greenland ice sheet projections.

**Surface mass balance:** annual cumulative SMB [ $\text{mm yr}^{-1}$  w.e.]

Like most variables, the SMB needs to be extended outside of the observed ice sheet mask to accommodate ice sheet models with a slightly larger than observed footprint. See main text for details on how this was done in the different downscaling procedures.

**Vertical gradients of runoff:** annual mean slope of the local runoff-elevation gradients [ $\text{mm yr}^{-1}$  w.e.  $\text{m}^{-1}$ ].

This variable is needed to parameterise the SMB-height feedback in ice sheet models (via  $d\text{SMB}/dz$  in Eq. 1). The gradients are expected to be predominantly negative as runoff generally declines with elevation and should be masked to 0 where no runoff is present. This variable has to be relatively smooth. Using gradients in runoff rather than gradients in SMB to

parameterize the SMB-height feedback is chosen because precipitation does not have consistent gradients with elevation. The variable needs to be extended outside of the observed ice sheet margin.

**Surface temperature:** annual mean surface temperature [ $^{\circ}\text{C}$ ]

This variable should represent as best as possible the temperature evolution at the upper ice surface as it is used to force the thermodynamic ice sheet solution at the upper boundary. In climate models with detailed snow physics, this can be, for example, a deep firn temperature. In the absence of detailed climate model output at that level, the skin temperature or even the 2m air temperature may be acceptable workarounds. The variable needs to be extended outside of the observed ice sheet margin.

**Vertical gradients of surface temperature:** annual mean **slope** of the local temperature-elevation gradient [ $^{\circ}\text{C m}^{-1}$ ].

This variable is used to apply a lapse-rate correction of the surface temperature boundary condition with changing surface elevation. This variable should be relatively smooth and needs to be extended outside of the observed ice sheet margin.

**Runoff:** monthly cumulative runoff [ $\text{mm yr}^{-1}$  w.e.].

This variable is used in combination with ocean thermal forcing to derive the outlet glacier retreat parameterization. As it is based on the observed geometry, this is the only variable that does not need to be extended over the tundra.

**Ocean thermal forcing:** We need to know the exact model version of the forcing ESM so we can extract matching ocean data from the CMIP archive.

Because we calculate anomalies relative to the period 1960-1989, SMB and ST have to cover the historical period (1960-2014) in addition to the projection period (2015-2100). All other data should cover at least the projection period (2015-2100). In addition, for climate forcing data to be used for ice sheet model initialisation and historical experiments, it should be provided over the historical period from 1950 under ERA5 forcing or another reanalysis product.

## Appendix B: List of ISM projections

**Table B1.** Ice sheet model versions and number of experiments. Bold model versions are shown in Figure 3 and 5.

Linear - linear sliding law, Weertman - Weertman sliding law ( $m=1/3$ ), ZI - Zoet-Iverson sliding law, RC - regularised Coulomb sliding law, PMIP3 - PMIP3 ensemble mean forcing for spinup, HadCM3 - HadCM3 forcing for spinup, CCSM - CCSM forcing for spinup, MARv3.9 - initialised with MARv3.9, MARv3.12 - initialised with MARv3.12, Num - number of experiments for different forcings (ESM, scenario, RCM, retreat).

Group	Model	Resolution (km)	Variant	Num
IGE	ElmerIce2	1 - 6	Linear	14
	<b>ElmerIce3</b>	1 - 6	Weertman	15
IMAU	IMAUICE1	40	ZI, PMIP3	57
	IMAUICE2	30	ZI, PMIP3	57

	IMAUICE3	20	ZI, PMIP3	57
	<b>IMAUICE5</b>	10	ZI, PMIP3	57
	IMAUICE6	20	ZI, HadCM3	57
	IMAUICE7	20	ZI, CCSM	57
	IMAUICE8	20	RC, PMIP3	57
NORCE	CISM02-MAR39	2	MARv3.9	36
	CISM04-MAR312	4	MARv3.12	48
	<b>CISM04-MAR39</b>	4	MARv3.9	69
	CISM04c-MAR39	4	MARv3.9, consistent <sup>†</sup>	95
	CISM04e-MAR312	4	MARv3.12, extension to 2200	5
	CISM04-MAR312	4	MARv3.12	4
	CISM08-MAR312	8	MARv3.12	48
	CISM08-MAR39	8	MARv3.9	115
	CISM08c-MAR39	8	MARv3.9, consistent <sup>†</sup>	95
	CISM16-MAR312	16	MARv3.12	48
	CISM16-MAR39	16	MARv3.9	100
	CISM16c-MAR312	16	MARv3.12, consistent <sup>†</sup>	110
	CISM16oc-MAR39	16	MARv3.9, overshoot to 2300	45
	CISM16t-MAR39	16	MARv3.9, repeat to 2300	65
	CISM16tc-MAR39	16	MARv3.9, repeat to 2300, consistent <sup>†</sup>	59
	CISM16xc-MAR12	16	MARv3.12, extension to 2300, consistent <sup>†</sup>	24
VUB	<b>GISMHOMv1</b>	5	Higher-order model	57
	GISMSIAv1	5	Shallow ice approximation	21

488 <sup>†</sup> retreat sensitivity consistent between historical and projection.

## 489 **Appendix C: Construction of extensions until 2300.**

490 **Extensions under climate forcing IPSL-CM6A-LR SSP5-8.5** have been downscaled with MARv3.13, which is largely  
491 similar to v3.12. The only difference is a small correction of albedo as a function of the liquid water content of the surface  
492 snowpack. Experiment MARv3.13-e05 uses SMB forcing produced at a fixed topography, as for the other experiments. In  
493 addition, we have experiment MARv3.13-e55, which uses SMB forcing produced at a changing topography. The topography  
494 change was produced by running two iterations between MAR and CISM with consecutive update of SMB and topography.  
495 The processing steps were the following:

- 496 1. Run MARv3.13 forced with IPSL-CM6A-LR SSP5-8.5 to 2300, where a quarter of the cumulated SMB anomaly is  
497 used to update the topography. This underestimates the topography change compared to a theoretical fully-coupled  
498 experiment by around a factor 4, so it is close to no update of the topography.
- 499 2. Run CISM with the SMB in 1.

500 3. Run MARv3.13 forced with IPSL-CM6A-LR SSP5-8.5 to 2300 with topography changes taken every 10 years from  
501 2070 forward from 2.

502 4. Run CISM with the SMB in 3.

503

504 Schematic extension of forcing between 2100 and 2300 based on existing data until 2100.

505 **Repeat scenarios.** The forcing until 2100 is the same as the corresponding scenario. From 2101 – 2300 the forcing is randomly  
506 repeated by shuffling the last 10 years of existing data (2091-2100). The following indices are used.

507 year = 2101, 2102, 2103, [...], 2298, 2299, 2300 ;  
508 shuffled\_time\_repeat = 2093, 2099, 2095, 2100, 2092, 2097, 2098, 2094,  
509 2091, 2096, 2100, 2097, 2099, 2095, 2096, 2091, 2093, 2098, 2094, 2092,  
510 2099, 2095, 2094, 2096, 2091, 2097, 2100, 2093, 2092, 2098, 2100, 2095,  
511 2098, 2094, 2093, 2097, 2092, 2099, 2096, 2091, 2097, 2098, 2099, 2093,  
512 2095, 2100, 2092, 2096, 2091, 2094, 2098, 2091, 2094, 2100, 2099, 2092,  
513 2093, 2096, 2095, 2097, 2100, 2094, 2091, 2096, 2095, 2093, 2092, 2099,  
514 2097, 2098, 2100, 2098, 2091, 2096, 2093, 2092, 2099, 2094, 2097, 2095,  
515 2094, 2097, 2095, 2098, 2093, 2092, 2096, 2099, 2100, 2091, 2097, 2095,  
516 2092, 2094, 2100, 2098, 2099, 2091, 2096, 2093, 2093, 2091, 2096, 2095,  
517 2097, 2099, 2098, 2092, 2094, 2100, 2097, 2100, 2098, 2096, 2091, 2094,  
518 2099, 2092, 2093, 2095, 2091, 2099, 2100, 2093, 2095, 2094, 2092, 2098,  
519 2096, 2097, 2094, 2097, 2095, 2099, 2092, 2098, 2096, 2093, 2100, 2091,  
520 2094, 2098, 2093, 2097, 2092, 2100, 2096, 2095, 2091, 2099, 2095, 2091,  
521 2096, 2100, 2094, 2097, 2093, 2092, 2098, 2099, 2091, 2094, 2092, 2097,  
522 2096, 2100, 2098, 2093, 2099, 2095, 2096, 2091, 2094, 2093, 2098, 2097,  
523 2092, 2100, 2095, 2099, 2098, 2091, 2100, 2092, 2097, 2094, 2096, 2093,  
524 2099, 2095, 2095, 2096, 2091, 2100, 2099, 2093, 2094, 2092, 2097, 2098;  
525  
526


527 **Overshoot scenarios.**

528 The schematic overshoot scenario mimicking SSP5-3.4-OS are based on the global temperature evolution as illustrated in  
529 Figure C1 Until year 2055, the forcing is the same as SSP5-8.5. From year 2056 – 2165, the temperature decreases similarly  
530 to the increase between 2030 and 2055 but backwards at 0.25 the rate (drawing four years for one). From 2156 on we shuffle  
531 and repeat the forcing earlier in the experiment, drawn from the time window 2026 – 2038.

532 Forcing until 2055 is the same as the corresponding SSP5-8.5 scenario. From 2056 – 2300 the following indices are used.

533 year = 2056, 2057, 2058, [...], 2298, 2299, 2300 ;  
534 shuffled\_time\_overshoot = 2056, 2056, 2055, 2054, 2053, 2055, 2054, 2053,  
535 2052, 2054, 2053, 2052, 2051, 2053, 2052, 2051, 2050, 2052, 2051, 2050,  
536 2049, 2051, 2050, 2049, 2048, 2050, 2049, 2048, 2047, 2049, 2048, 2047,  
537 2046, 2048, 2047, 2046, 2045, 2047, 2046, 2045, 2044, 2046, 2045, 2044,  
538 2043, 2045, 2044, 2043, 2042, 2044, 2043, 2042, 2041, 2043, 2042, 2041,  
539 2040, 2042, 2041, 2040, 2039, 2041, 2040, 2039, 2038, 2040, 2039, 2038,  
540 2037, 2039, 2038, 2037, 2036, 2038, 2037, 2036, 2035, 2037, 2036, 2035,  
541 2034, 2036, 2035, 2034, 2033, 2035, 2034, 2033, 2032, 2034, 2033, 2032,  
542 2031, 2033, 2032, 2031, 2030, 2032, 2031, 2030, 2029, 2033, 2038, 2031,  
543 2037, 2029, 2035, 2028, 2034, 2036, 2030, 2027, 2032, 2035, 2031, 2037,

2029, 2030, 2033, 2032, 2034, 2028, 2038, 2027, 2036, 2032, 2031, 2027,  
 2037, 2034, 2033, 2036, 2035, 2029, 2030, 2038, 2028, 2027, 2034, 2029,  
 2030, 2033, 2036, 2031, 2032, 2035, 2038, 2028, 2037, 2034, 2038, 2027,  
 2036, 2037, 2029, 2035, 2031, 2030, 2032, 2033, 2028, 2036, 2031, 2027,  
 2032, 2030, 2038, 2028, 2034, 2035, 2033, 2029, 2037, 2036, 2028, 2030,  
 2027, 2038, 2032, 2037, 2033, 2034, 2029, 2031, 2035, 2028, 2030, 2035,  
 2033, 2029, 2037, 2034, 2031, 2038, 2032, 2027, 2036, 2032, 2031, 2027,  
 2030, 2028, 2034, 2037, 2035, 2033, 2036, 2038, 2029, 2029, 2036, 2035,  
 2033, 2037, 2028, 2027, 2031, 2038, 2034, 2032, 2030, 2038, 2027, 2033,  
 2037, 2030, 2034, 2036, 2028, 2031, 2029, 2032, 2035, 2038, 2027, 2030,  
 2031, 2034, 2035, 2037, 2036, 2032, 2029, 2033, 2028 ;



**Figure C1.** Illustration of the construction of schematic overshoot scenarios mimicking SSP5-3.4-OS. To construct the forcing, SMB and retreat forcing are drawn from existing annual forcing files (not shown). Instead, the figure shows the sequence of global mean temperature anomaly drawn from each individual original ESM temperature time series.

## Data Availability

The forcing data will be provided in ISMIP6 format on a public archive. It consists of SMB and ST anomalies and their respective vertical gradients that are generic for all ice sheet models. The retreat mask forcing is produced specifically for each individual ice sheet model version and is maintained by the modellers.

The provided ice sheet model data consists of the most important diagnostic output at annual time resolution, such as ice thickness, bedrock and surface topography, horizontal velocities and integral mass balance terms. We are following the ISMIP6 data request format (<https://thehub.org/groups/ismip6/wiki/ISMIP6-Projections-Greenland>). For common analysis, ice sheet model output was conservatively interpolated to a standard 5 km diagnostic grid. These model output data will be made available on a public archive, while the raw ice sheet model output is the responsibility of the individual modelling groups.

Projected sea-level contributions will be provided on a public archive.

## Author contributions



572 HG designed the experimental setup with input from TE, prepared and distributed the forcing data, collected and processed  
573 the output data, analysed the results and produced the figures. XF, QG, MvdB, BN, RM, MO, FB produced climate forcing  
574 data. HG, CR, CJB, FGC and SL conducted ice sheet model experiments. HG wrote the manuscript with input from all co-  
575 authors.

576  
577 **Competing interests**  
578 At least one of the (co-)authors is a member of the editorial board of The Cryosphere.

579  
580 **Disclaimer**  
581 Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps,  
582 institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every  
583 effort to include appropriate place names, the final responsibility lies with the authors.

584  
585 **Acknowledgements**  
586 We acknowledge the World Climate Research Programme and its Working Group on Coupled Modelling for coordinating and  
587 promoting CMIP5 and CMIP6. We thank the climate modelling groups for producing and making available their model output  
588 and the Earth System Grid Federation (ESGF) for archiving the CMIP data. We thank the ISMIP6/7 steering committee and  
589 community for defining and providing a framework for this work. Resources were provided by Sigma2 - the National  
590 Infrastructure for High Performance Computing and Data Storage in Norway through projects NN8085K, NN8006K,  
591 NS5011K, NS8006 K and NS8085K. B.N. is a Research Associate of the Fonds de la Recherche Scientifique de Belgique–  
592 F.R.S.-FNRS.

593  
594 **Financial support**  
595 This research has received funding from the European Union's Horizon 2020 research and innovation programme under grant  
596 agreement 869304 (PROTECT) and has been supported by the Research Council of Norway under project 324639 (GREASE).

597  
598 **References**  
599 Aschwanden, A., Bartholomaus, T. C., Brinkerhoff, D. J., and Truffer, M.: Brief communication: A roadmap towards credible  
600 projections of ice sheet contribution to sea level, The Cryosphere, 15, 5705–5715, <https://doi.org/10.5194/tc-15-5705-2021>,  
601 2021.

602 Bales, R. C., Guo, Q., Shen, D., McConnell, J. R., Du, G., Burkhart, J. F., Spikes, V. B., Hanna, E., and Cappelen, J.: Annual  
603 accumulation for Greenland updated using ice core data developed during 2000–2006 and analysis of daily coastal  
604 meteorological data, *Journal of Geophysical Research Atmospheres*, 114, <https://doi.org/10.1029/2008JD011208>, 2009.

605 Berends, C. J., de Boer, B., and van de Wal, R. S. W.: Application of HadCM3@Bristolv1.0 simulations of paleoclimate as  
606 forcing for an ice-sheet model, ANICE2.1: set-up and benchmark experiments, *Geosci. Model Dev.*, 11, 4657–4675,  
607 <https://doi.org/10.5194/gmd-11-4657-2018>, 2018.

608 Berends, C. J., Goelzer, H., Reerink, T. J., Stap, L. B., and van de Wal, R. S. W.: Benchmarking the vertically integrated ice-  
609 sheet model IMAU-ICE (version 2.0), *Geosci. Model Dev.*, 15, 5667–5688, <https://doi.org/10.5194/gmd-15-5667-2022>,  
610 2022.

611 Berends, C. J., Van De Wal, R. S. W., Van Den Akker, T., and Lipscomb, W. H.: Compensating errors in inversions for  
612 subglacial bed roughness: same steady state, different dynamic response, *Cryosphere*, 17, 1585–1600,  
613 <https://doi.org/10.5194/tc-17-1585-2023>, 2023.

614 Brady, E. C., Otto-Bliesner, B. L., Kay, J. E., and Rosenbloom, N.: Sensitivity to Glacial Forcing in the CCSM4, *Journal of*  
615 *Climate*, 26, 1901–1925, <https://doi.org/10.1175/JCLI-D-11-00416.1>, 2013.

616 Citterio, M. and Ahlstrøm, A. P.: Brief communication “The aerophotogrammetric map of Greenland ice masses,” *The*  
617 *Cryosphere*, 7, 445–449, <https://doi.org/10.5194/tc-7-445-2013>, 2013.

618 Delhasse, A., Kittel, C., Amory, C., Hofer, S., van As, D., S. Fausto, R., and Fettweis, X.: Brief communication: Evaluation  
619 of the near-surface climate in ERA5 over the Greenland Ice Sheet, *The Cryosphere*, 14, 957–965,  
620 <https://doi.org/10.5194/tc-14-957-2020>, 2020.

621 Delhasse, A., Beckmann, J., Kittel, C., and Fettweis, X.: Coupling MAR (Modèle Atmosphérique Régional) with PISM  
622 (Parallel Ice Sheet Model) mitigates the positive melt–elevation feedback, *The Cryosphere*, 18, 633–651,  
623 <https://doi.org/10.5194/tc-18-633-2024>, 2024.

624 Durand, G., van den Broeke, M. R., Le Cozannet, G., Edwards, T. L., Holland, P. R., Jourdain, N. C., Marzeion, B., Mottram,  
625 R., Nicholls, R. J., Pattyn, F., Paul, F., Slangen, A. B. A., Winkelmann, R., Burgard, C., van Calcar, C. J., Barré, J.-B.,  
626 Bataille, A., and Chapuis, A.: Sea-Level Rise: From Global Perspectives to Local Services, *Frontiers in Marine Science*,  
627 Volume 8-2021, 2022.

628 Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E.,  
629 Smith, C. J., McKenna, C. M., Simon, E., Abe-Ouchi, A., Gregory, J. M., Larour, E., Lipscomb, W. H., Payne, A. J.,  
630 Shepherd, A., Agosta, C., Alexander, P., Albrecht, T., Anderson, B., Asay-Davis, X., Aschwanden, A., Barthel, A., Bliss,  
631 A., Calov, R., Chambers, C., Champollion, N., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Felikson, D., Fettweis, X.,  
632 Fujita, K., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A.,  
633 Huss, M., Huybrechts, P., Immerzeel, W., Kleiner, T., Kraaijenbrink, P., Le clec’h, S., Lee, V., Leguy, G. R., Little, C. M.,  
634 Lowry, D. P., Malles, J. H., Martin, D. F., Maussion, F., Morlighem, M., O’Neill, J. F., Nias, I., Pattyn, F., Pelle, T., Price,  
635 S. F., Quiquet, A., Radić, V., Reese, R., Rounce, D. R., Rückamp, M., Sakai, A., Shafer, C., Schlegel, N. J., Shannon, S.,

Smith, R. S., Straneo, F., Sun, S., Tarasov, L., Trusel, L. D., Van Breedam, J., van de Wal, R., van den Broeke, M., Winkelman, R., Zekollari, H., Zhao, C., Zhang, T., and Zwinger, T.: Projected land ice contributions to twenty-first-century sea level rise, *Nature*, 593, 74–82, <https://doi.org/10.1038/s41586-021-03302-y>, 2021.

Edwards, T. L. et al.: PROTECT sea-level projections. in prep, 2026.

Fettweis, X., Box, J. E., Agosta, C., Amory, C., Kittel, C., Lang, C., van As, D., Machguth, H., and Gallée, H.: Reconstructions of the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model, *The Cryosphere*, 11, 1015–1015, <https://doi.org/10.5194/tc-11-1015-2017>, 2017.

Fettweis, X., Hofer, S., Krebs-Kanzow, U., Amory, C., Aoki, T., Berends, C. J., Born, A., Box, J. E., Delhasse, A., Fujita, K., Gierz, P., Goelzer, H., Hanna, E., Hashimoto, A., Huybrechts, P., Kapsch, M.-L., King, M. D., Kittel, C., Lang, C., Langen, P. L., Lenaerts, J. T. M., Liston, G. E., Lohmann, G., Mernild, S. H., Mikolajewicz, U., Modali, K., Mottram, R. H., Niwano, M., Noël, B., Ryan, J. C., Smith, A., Streffing, J., Tedesco, M., van de Berg, W. J., van den Broeke, M., van de Wal, R. S. W., van Kampenhout, L., Wilton, D., Wouters, B., Ziemen, F., and Zolles, T.: GrSMBMIP: intercomparison of the modelled 1980–2012 surface mass balance over the Greenland Ice Sheet, *The Cryosphere*, 14, 3935–3958, <https://doi.org/10.5194/tc-14-3935-2020>, 2020.

Fox-Kemper, B., Hewitt, H. T., Xiao, C., Adalgeirsdottir, G., Drijfhout, S. S., Edwards, T. L., Golledge, N. R., Hemer, M., Kopp, R. E., Krinner, G., Mix, A., Notz, D., Nowicki, S., Nurhati, I. S., Ruiz, L., Sallee, J.-B., Slangen, A. B. A., and Yu, Y.: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, United Kingdom and New York, NY, USA, 1211–1362, <https://doi.org/10.1017/9781009157896.011>, 2021.

Franco, B., Fettweis, X., Lang, C., and Erpicum, M.: Impact of spatial resolution on the modelling of the Greenland ice sheet surface mass balance between 1990–2010, using the regional climate model MAR, *Cryosphere*, 6, 695–711, <https://doi.org/10.5194/tc-6-695-2012>, 2012.

Fürst, J. J., Goelzer, H., and Huybrechts, P.: Effect of higher-order stress gradients on the centennial mass evolution of the Greenland ice sheet, *Cryosphere*, 7, 183–199, <https://doi.org/10.5194/tc-7-183-2013>, 2013.

Fürst, J. J., Goelzer, H., and Huybrechts, P.: Ice-dynamic projections of the Greenland ice sheet in response to atmospheric and oceanic warming, *Cryosphere*, 9, 1039–1062, <https://doi.org/10.5194/tc-9-1039-2015>, 2015.

Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice sheet contribution to sea-level rise from a new-generation ice-sheet model, *Cryosphere*, 6, 1561–1576, <https://doi.org/10.5194/tc-6-1561-2012>, 2012.

Glaude, Q., Noel, B., Olesen, M., Van den Broeke, M., van de Berg, W. J., Mottram, R., Hansen, N., Delhasse, A., Amory, C., Kittel, C., Goelzer, H., and Fettweis, X.: A Factor Two Difference in 21st-Century Greenland Ice Sheet Surface Mass

669 Balance Projections From Three Regional Climate Models Under a Strong Warming Scenario (SSP5-8.5), *Geophysical*  
670 *Research Letters*, 51, e2024GL111902, <https://doi.org/10.1029/2024GL111902>, 2024.

671 Goelzer, H., Nowicki, S., Edwards, T., Beckley, M., Abe-Ouchi, A., Aschwanden, A., Calov, R., Gagliardini, O., Gillet-  
672 Chaulet, F., Golledge, N. R., Gregory, J., Greve, R., Humbert, A., Huybrechts, P., Kennedy, J. H., Larour, E., Lipscomb,  
673 W. H., Le clec'h, S., Lee, V., Morlighem, M., Pattyn, F., Payne, A. J., Rodehacke, C., Rückamp, M., Saito, F., Schlegel,  
674 N., Seroussi, H., Shepherd, A., Sun, S., van de Wal, R., and Ziemen, F. A.: Design and results of the ice sheet model  
675 initialisation experiments initMIP-Greenland: an ISMIP6 intercomparison, *The Cryosphere*, 12, 1433–1460,  
676 <https://doi.org/10.5194/tc-12-1433-2018>, 2018.

677 Goelzer, H., Nowicki, S., Payne, A., Larour, E., Seroussi, H., Lipscomb, W. H., Gregory, J., Abe-Ouchi, A., Shepherd, A.,  
678 Simon, E., Agosta, C., Alexander, P., Aschwanden, A., Barthel, A., Calov, R., Chambers, C., Choi, Y., Cuzzone, J., Dumas,  
679 C., Edwards, T., Felikson, D., Fettweis, X., Golledge, N. R., Greve, R., Humbert, A., Huybrechts, P., Le clec'h, S., Lee,  
680 V., Leguy, G., Little, C., Lowry, D. P., Morlighem, M., Nias, I., Quiquet, A., Rückamp, M., Schlegel, N.-J., Slater, D. A.,  
681 Smith, R. S., Straneo, F., Tarasov, L., van de Wal, R., and van den Broeke, M.: The future sea-level contribution of the  
682 Greenland ice sheet: a multi-model ensemble study of ISMIP6, *The Cryosphere*, 14, 3071–3096, [https://doi.org/10.5194/tc-](https://doi.org/10.5194/tc-14-3071-2020)  
683 [14-3071-2020](https://doi.org/10.5194/tc-14-3071-2020), 2020a.

684 Goelzer, H., Coulon, V., Pattyn, F., de Boer, B., and van de Wal, R.: Brief communication: On calculating the sea-level  
685 contribution in marine ice-sheet models, *The Cryosphere*, 14, 833–840, <https://doi.org/10.5194/tc-14-833-2020>, 2020b.

686 Goelzer, H., Noël, B. P. Y., Edwards, T. L., Fettweis, X., Gregory, J. M., Lipscomb, W. H., van de Wal, R. S. W., and van den  
687 Broeke, M. R.: Remapping of Greenland ice sheet surface mass balance anomalies for large ensemble sea-level change  
688 projections, *The Cryosphere*, 14, 1747–1762, <https://doi.org/10.5194/tc-14-1747-2020>, 2020c.

689 Goelzer, H., Langebroek, P. M., Born, A., Hofer, S., Haubner, K., Petrini, M., Leguy, G., Lipscomb, W. H., and Thayer-  
690 Calder, K.: Interactive coupling of a Greenland ice sheet model in NorESM2, *Geosci. Model Dev.*, 18, 7853–7867,  
691 <https://doi.org/10.5194/gmd-18-7853-2025>, 2025.

692 Goldberg, D. N.: A variationally derived, depth-integrated approximation to a higher-order glaciological flow model, *Journal*  
693 *of Glaciology*, 57, 157–170, <https://doi.org/10.3189/002214311795306763>, 2011.

694 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers,  
695 D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De  
696 Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A.,  
697 Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti,  
698 G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal*  
699 *of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

700 Hofer, S., Lang, C., Amory, C., Tedstone, A., Fettweis, X., Kittel, C., and Delhasse, A.: Greater Greenland Ice Sheet  
701 contribution to global sea level rise in CMIP6, *Nature Communications*, 1–11, [https://doi.org/10.1038/s41467-020-20011-](https://doi.org/10.1038/s41467-020-20011-8)  
702 [8](https://doi.org/10.1038/s41467-020-20011-8), 2020.

703 Howat, I. M., Negrete, A., and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface  
 704 elevation data sets, *The Cryosphere*, 8, 1509–1518, <https://doi.org/10.5194/tc-8-1509-2014>, 2014.

705 Huybrechts, P.: Sea-level changes at the LGM from ice-dynamic reconstructions of the Greenland and Antarctic ice sheets  
 706 during the glacial cycles, *Quaternary Science Reviews*, 21, 203–231, [https://doi.org/10.1016/S0277-3791\(01\)00082-8](https://doi.org/10.1016/S0277-3791(01)00082-8),  
 707 2002.

708 Huybrechts, P., Gregory, J. M., Janssens, I., and Wild, M.: Modelling Antarctic and Greenland volume changes during the  
 709 20th and 21st centuries forced by GCM time slice integrations, *Global and Planetary Change*, 42, 83–105,  
 710 <https://doi.org/10.1016/j.gloplacha.2003.11.011>, 2004.

711 Joughin, I., Smith, B. E., Howat, I. M., Scambos, T., and Moon, T.: Greenland flow variability from ice-sheet-wide velocity  
 712 mapping, *Journal of Glaciology*, 56, 415–430, <https://doi.org/10.3189/002214310792447734>, 2010.

713 Langen, P. L., Fausto, R. S., Vandecrux, B., Mottram, R. H., and Box, J. E.: Liquid Water Flow and Retention on the Greenland  
 714 Ice Sheet in the Regional Climate Model HIRHAM5: Local and Large-Scale Impacts, *Frontiers in Earth Science*, Volume  
 715 4-2016, 2017.

716 Le clec’h, S., Quiquet, A., Charbit, S., Dumas, C., Kageyama, M., and Ritz, C.: A rapidly converging initialisation method to  
 717 simulate the present-day Greenland ice sheet using the GRISLI ice sheet model (version 1.3), *Geosci. Model Dev.*, 12,  
 718 2481–2499, <https://doi.org/10.5194/gmd-12-2481-2019>, 2019.

719 Lipscomb, W. H., Price, S. F., Hoffman, M. J., Leguy, G. R., Bennett, A. R., Bradley, S. L., Evans, K. J., Fyke, J. G., Kennedy,  
 720 J. H., Perego, M., Ranken, D. M., Sacks, W. J., Salinger, A. G., Vargo, L. J., and Worley, P. H.: Description and evaluation  
 721 of the Community Ice Sheet Model (CISM) v2.1, *Geoscientific Model Development*, 12, 387–424,  
 722 <https://doi.org/10.5194/gmd-12-387-2019>, 2019.

723 MacFarling Meure, C., Etheridge, D., Trudinger, C., Steele, P., Langenfelds, R., van Ommen, T., Smith, A., and Elkins, J.:  
 724 Law Dome CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O ice core records extended to 2000 years BP, *Geophysical Research Letters*, 33,  
 725 <https://doi.org/10.1029/2006GL026152>, 2006.

726 Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A.,  
 727 Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R.,  
 728 Mayer, L., Mouginot, J., Noël, B. P. Y. Y., O’Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon,  
 729 P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., Zinglensen, K. B., Cofaigh, C. Ó., Palmer, S., Rysgaard,  
 730 S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen,  
 731 K. B.: BedMachine v3: Complete bed topography and ocean bathymetry mapping of Greenland from multi-beam echo  
 732 sounding combined with mass conservation, *Geophysical Research Letters*, 44, 11,051–11,061,  
 733 <https://doi.org/10.1002/2017GL074954>, 2017.

734 Mottram, R., Boberg, F., Langen, P., Yang, S., Rodehacke, C., Christensen, J. H., and Madsen, M. S.: Surface mass balance  
 735 of the Greenland ice sheet in the regional climate model HIRHAM5: Present state and future prospects, *Low Temperature*  
 736 *Science*, 75, 105–115, <https://doi.org/10.14943/lowtemsci.75.105>, 2017.

737 Muntjewerf, L., Petrini, M., Vizcaino, M., Ernani da Silva, C., Sellevold, R., Scherrenberg, M. D. W., Thayer-Calder, K.,  
738 Bradley, S. L., Lenaerts, J. T. M., Lipscomb, W. H., and Lofverstrom, M.: Greenland Ice Sheet Contribution to 21st Century  
739 Sea Level Rise as Simulated by the Coupled CESM2.1-CISM2.1, *Geophysical Research Letters*, 47,  
740 <https://doi.org/10.1029/2019GL086836>, 2020.

741 Noël, B., van de Berg, W. J., Machguth, H., Lhermitte, S., Howat, I., Fettweis, X., and van den Broeke, M. R.: A daily, 1 km  
742 resolution data set of downscaled Greenland ice sheet surface mass balance (1958–2015), *The Cryosphere*, 10, 2361–2377,  
743 <https://doi.org/10.5194/tc-10-2361-2016>, 2016.

744 Noël, B., Van De Berg, W. J., Van Wessem, J. M., Van Meijgaard, E., Van As, Di., Lenaerts, J. T. M., Lhermitte, S., Munneke,  
745 P. K., Smeets, C. J. P. P., Van Uft, L. H., Van De Wal, R. S. W., and Van Den Broeke, M. R.: Modelling the climate and  
746 surface mass balance of polar ice sheets using RACMO2 - Part 1: Greenland (1958-2016), *Cryosphere*, 12, 811–831,  
747 <https://doi.org/10.5194/tc-12-811-2018>, 2018.

748 Noël, B., van Kampenhout, L., van de Berg, W. J., Lenaerts, J. T. M., Wouters, B., and van den Broeke, M. R.: Brief  
749 communication: CESM2 climate forcing (1950–2014) yields realistic Greenland ice sheet surface mass balance, *The*  
750 *Cryosphere*, 14, 1425–1435, <https://doi.org/10.5194/tc-14-1425-2020>, 2020.

751 Noël, B., Lenaerts, J. T. M., Lipscomb, W. H., Thayer-Calder, K., and van den Broeke, M. R.: Peak refreezing in the Greenland  
752 firn layer under future warming scenarios, *Nature Communications*, 13, 6870, [https://doi.org/10.1038/s41467-022-34524-](https://doi.org/10.1038/s41467-022-34524-x)  
753 [x](https://doi.org/10.1038/s41467-022-34524-x), 2022.

754 Nowicki, S. M. J., Payne, A., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., and Shepherd,  
755 A.: Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6, *Geoscientific Model Development*, 9,  
756 4521–4545, <https://doi.org/10.5194/gmd-9-4521-2016>, 2016.

757 Nowicki, S., Payne, A. J., Goelzer, H., Seroussi, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Alexander, P., Asay-Davis,  
758 X. S., Barthel, A., Bracegirdle, T. J., Cullather, R., Felikson, D., Fettweis, X., Gregory, J., Hatterman, T., Jourdain, N. C.,  
759 Kuipers Munneke, P., Larour, E., Little, C. M., Morlinghem, M., Nias, I., Shepherd, A., Simon, E., Slater, D., Smith, R.,  
760 Straneo, F., Trusel, L. D., van den Broeke, M. R., and van de Wal, R.: Experimental protocol for sealevel projections from  
761 ISMIP6 standalone ice sheet models, *The Cryosphere*, 14, 2331-2368-2331–2368, [https://doi.org/10.5194/tc-14-2331-](https://doi.org/10.5194/tc-14-2331-2020)  
762 [2020](https://doi.org/10.5194/tc-14-2331-2020), 2020.

763 O'Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.-  
764 F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario Model Intercomparison Project  
765 (ScenarioMIP) for CMIP6, *Geosci. Model Dev.*, 9, 3461–3482, <https://doi.org/10.5194/gmd-9-3461-2016>, 2016.

766 Payne, A. J., Nowicki, S., Abe-Ouchi, A., Agosta, C., Alexander, P., Albrecht, T., Asay-Davis, X., Aschwanden, A., Barthel,  
767 A., Bracegirdle, T. J., Calov, R., Chambers, C., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Edwards, T. L., Felikson,  
768 D., Fettweis, X., Galton-Fenzi, B. K., Goelzer, H., Gladstone, R., Golledge, N. R., Gregory, J. M., Greve, R., Hattermann,  
769 T., Hoffman, M. J., Humbert, A., Huybrechts, P., Jourdain, N. C., Kleiner, T., Munneke, P. K., Larour, E., Le clec'h, S.,  
770 Lee, V., Leguy, G., Lipscomb, W. H., Little, C. M., Lowry, D. P., Morlighem, M., Nias, I., Pattyn, F., Pelle, T., Price, S.

771 F., Quiquet, A., Reese, R., Rückamp, M., Schlegel, N. J., Seroussi, H., Shepherd, A., Simon, E., Slater, D., Smith, R. S.,  
 772 Straneo, F., Sun, S., Tarasov, L., Trusel, L. D., Van Breedam, J., van de Wal, R., van den Broeke, M., Winkelmann, R.,  
 773 Zhao, C., Zhang, T., and Zwinger, T.: Future Sea Level Change Under Coupled Model Intercomparison Project Phase 5  
 774 and Phase 6 Scenarios From the Greenland and Antarctic Ice Sheets, *Geophysical Research Letters*, 48, 1–8,  
 775 <https://doi.org/10.1029/2020GL091741>, 2021.

776 Rahlves, C., Goelzer, H., Born, A., and Langebroek, P. M.: Historically consistent mass loss projections of the Greenland ice  
 777 sheet, *The Cryosphere*, 19, 1205–1220, <https://doi.org/10.5194/tc-19-1205-2025>, 2025a.

778 Rahlves, C., Goelzer, H., Born, A., and Langebroek, P. M.: Investigating the multi-millennial evolution and stability of the  
 779 Greenland ice sheet using remapped surface mass balance forcing, *EGUsphere*, 2025, 1–23,  
 780 <https://doi.org/10.5194/egusphere-2025-2192>, 2025b.

781 Rantanen, M., Karpechko, A. Yu., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen,  
 782 A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Communications Earth & Environment*, 3,  
 783 168, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.

784 Rohmer, J., Thieblemont, R., Le Cozannet, G., Goelzer, H., and Durand, G.: Improving interpretation of sea-level projections  
 785 through a machine-learning-based local explanation approach, *The Cryosphere*, 16, 4637–4657, [https://doi.org/10.5194/tc-](https://doi.org/10.5194/tc-16-4637-2022)  
 786 [16-4637-2022](https://doi.org/10.5194/tc-16-4637-2022), 2022.

787 Rohmer, J., Goelzer, H., Edwards, T., Le Cozannet, G., and Durand, G.: Drawing lessons for multi-model ensemble design  
 788 from emulator experiments: application to future sea level contribution of the Greenland ice sheet, *EGUsphere* [preprint],  
 789 <https://doi.org/10.5194/egusphere-2025-52>, 2025.

790 Scherrenberg, M. D. W., Berends, C. J., Stap, L. B., and Van De Wal, R. S. W.: Modelling feedbacks between the Northern  
 791 Hemisphere ice sheets and climate during the last glacial cycle, *Climate of the Past*, 19, 399–418,  
 792 <https://doi.org/10.5194/cp-19-399-2023>, 2023.

793 Schoof, C.: The effect of cavitation on glacier sliding, *Proceedings of the Royal Society A: Mathematical, Physical and*  
 794 *Engineering Sciences*, 461, 609–627, <https://doi.org/10.1098/rspa.2004.1350>, 2005.

795 Shapiro, N. M. and Ritzwoller, M. H.: Inferring surface heat flux distributions guided by a global seismic model: particular  
 796 application to Antarctica, *Earth and Planetary Science Letters*, 223, 213–224, <https://doi.org/10.1016/j.epsl.2004.04.011>,  
 797 2004.

798 Singarayer, J. S. and Valdes, P. J.: High-latitude climate sensitivity to ice-sheet forcing over the last 120kyr, *Quaternary*  
 799 *Science Reviews*, 29, 43–55, <https://doi.org/10.1016/j.quascirev.2009.10.011>, 2010.

800 Slater, D. A., Straneo, F., Felikson, D., Little, C. M., Goelzer, H., Fettweis, X., and Holte, J.: Estimating Greenland tidewater  
 801 glacier retreat driven by submarine melting, *The Cryosphere*, 13, 2489–2509, <https://doi.org/10.5194/tc-13-2489-2019>,  
 802 2019.



803 Slater, D. A., Felikson, D., Straneo, F., Goelzer, H., Little, C. M., Morlighem, M., Fettweis, X., and Nowicki, S.: Twenty-first  
804 century ocean forcing of the Greenland ice sheet for modelling of sea level contribution, *The Cryosphere*, 14, 985–1008,  
805 <https://doi.org/10.5194/tc-14-985-2020>, 2020.

806 Smith, R. S., Mathiot, P., Siahayan, A., Lee, V., Cornford, S. L., Gregory, J. M., Payne, A. J., Jenkins, A., Holland, P. R.,  
807 Ridley, J. K., and Jones, C. G.: Coupling the U.K. Earth System Model to Dynamic Models of the Greenland and Antarctic  
808 Ice Sheets, *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002520,  
809 <https://doi.org/10.1029/2021MS002520>, 2021.

810 The IMBIE Team, Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs,  
811 K., Joughin, I., Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., A. G., Agosta, C., Ahlstrøm, A., Babonis,  
812 G., Barletta, V. R., Bjørk, A. A., Blazquez, A., Bonin, J., Colgan, W., Csatho, B., Cullather, R., Engdahl, M. E., Felikson,  
813 D., Fettweis, X., Forsberg, R., Hogg, A. E., Gallee, H., Gardner, A., Gilbert, L., Gourmelen, N., Groh, A., Gunter, B.,  
814 Hanna, E., Harig, C., Helm, V., Horvath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P. L., Lecavalier,  
815 B., Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mottram, R., Mouginit, J.,  
816 Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noël, B., Otsuka, I., Pattie, M. E., Peltier, W. R., Pie, N.,  
817 Rietbroek, R., Rott, H., Sandberg Sørensen, L., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schröder, L., Seo, K. W.,  
818 Simonsen, S. B., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., van de Berg, W. J., van der Wal, W., van  
819 Wessem, M., Vishwakarma, B. D., Wiese, D., Wilton, D., Wagner, T., Wouters, B., and Wuite, J.: Mass balance of the  
820 Greenland Ice Sheet from 1992 to 2018, *Nature*, 579, 233–239, <https://doi.org/10.1038/s41586-019-1855-2>, 2020.

821 van den Broeke, M., Box, J., Fettweis, X., Hanna, E., Noël, B., Tedesco, M., van As, D., van de Berg, W. J., and van  
822 Kampenhout, L.: Greenland Ice Sheet Surface Mass Loss: Recent Developments in Observation and Modeling, *Current*  
823 *Climate Change Reports*, 3, 345–356, <https://doi.org/10.1007/s40641-017-0084-8>, 2017.