

Reply to the comments on the manuscript

”Ground-based observations of periodic temperature fluctuations in the mesopause region with periods larger than 2 days”

by Christoph Kalicinsky, Robert Reisch, and Peter Knieling

We thank the reviewers for their helpful comments and recommendations. In the following, we discuss the issues addressed by the reviewers and explain our opinions and the modifications of our manuscript.

We enumerate the comments and repeat them in bold face. The modifications of the manuscript are displayed in the marked-up manuscript version as coloured text. Deleted parts are shown in red and new or modified text parts in blue.

1 Comments Reviewer 1

Thank you for the much-improved paper. There are still some concerns needs to be addressed.

1. **It is not clear, how exactly the fit was done for each year. The example in section 2.2 (Figure 1) showed yearly fit lasted for 365 days centered in the summer. While moving LSP window was introduced in section 2.3, two months at each end of the year were added to the fit. So, the fit was applied to Nov-Feb (total 16 months) centered in summer? However, for the argument of this new fit, it was compared with a fit centered in winter. This is also the case for Figure 7. It is very confusing. Please clearly state how the fit was applied to the data.**

The best possible detrending of the data is the prerequisite for the LSP analysis (see reply to reviewer 2). As the seasonal variation differs from year to year, one fit for all years at once is not advisable. Thus, the fit is calculated separately for each year. As we want to analyse all possible windows, it is necessary to add at least 30 days at both sides of the year. Then the first and last day of the year can be used as centre day for the LSP with a window length of 60 days. In order to reduce possible discontinuities for the different yearly fits at the transition points (begin and end of the year), we used two extra month at both sides of the time interval. Comparisons showed that this lead to a reasonable agreement to a fit centred in winter and thus the transition region is fitted well. We introduced a new figure that illustrates our approach and the comparison to the fits centred in winter.

The data in Fig. 7 are derived in the same way but centred in winter.

2. **I don't follow the discussion on the quasi-quadrennial oscillation. The higher LSP power for the 3rd proxy does not mean anything, as none of them are significant peaks. It is not enough to conclude that the strength of waves does not contribute to this oscillation. What is the physical meaning of a 4-year oscillation on the length of the event?**

We reduced the statements on the quasi-4-oscillation to not overinterpret the results.

3. **I don't understand why the authors tried so hard to pin a Rossby wave mode to each planetary wave observed. It is, as the authors stated, beyond of the**

scope of this paper.

We reduced the assignments to possible Rossby wave modes as it is beyond the scope of the paper. We just wanted to say that such Rossby wave modes are possible candidates for our observations as for example shown in the case of the Zhao et al. (2019) study. But additional data are necessary to prove this.

4. Please reduce the use of the conjunction words. Some of the long sentences are really hard to understand.

We carefully looked through our manuscript and reduced the number of long sentences.

Minor comments:

Line 7: is typically "larger": what do you mean by larger, scale, amplitude or periods? Or the authors mean occurred more frequently and larger amplitude? In the paper, "large/minor" activities also appeared which is very confusing.

Line 8-9 and some other places: for periods, longer/shorter instead of "larger/smaller"

Line 8-10: Not sure how this <10 day waves fit in the picture.

Line 9: "these waves": it is not clear if it refers to waves with periods shorter than 20 days or 10 days?

Line 115-124: This paragraph is talking about the fit, not LSP. Please move it to the fit session and clarify the method.

Line 176: remove "also due to"

Line 177: remove "also"

Line 301: amplitude, do you mean period?

Line 448: remove one of "the"

Please check the different uses of: analysis, analyze, analyses and correct them throughout the paper.

We corrected all minor comments.

Reply to the comments on the manuscript

”Ground-based observations of periodic temperature fluctuations in the mesopause region with periods larger than 2 days”

by Christoph Kalicinsky, Robert Reisch, and Peter Knieling

We thank the reviewers for their helpful comments and recommendations. In the following, we discuss the issues addressed by the reviewers and explain our opinions and the modifications of our manuscript.

We enumerate the comments and repeat them in bold face. The modifications of the manuscript are displayed in the marked-up manuscript version as coloured text. Deleted parts are shown in red and new or modified text parts in blue.

1 Comments Reviewer 2

The authors have made some of the recommended corrections and significant improvements, however I make some further comments with reference to my previous review:

2. **A declining trend of the amplitude of the annual cycle is determined from the yearly fit of a seasonal cycle. It would be of interest to the reader to know what rate of decline is obtained. Only one year is shown in Fig 1 (1997) with its fit. The fit would be discontinuous over a year boundary because previous and subsequent years are derived separately and this makes no physical sense. Line 93. A seasonal fit to the whole timeseries is what I am asking for. The residuals will show year to year variations and the trend about the long term continuous climatology. In what way does this have “implications on further analyses” ?**

The trend of the amplitude of the annual cycle is about -0.5 K/decade (Kalicinsky et al., 2024). We added this information to the manuscript.

We reduced the discontinuities at the year boundaries by adding the additional months to the data for each year. We added a new figure to the manuscript to illustrate the approach (see also reply to reviewer 1).

It is correct that one fit the the whole time series would lead to residuals that still include year to year variations and the trend about the long term climatology. But such underlying trends that remain in addition to the fluctuations which we want to analyse will lead to various misleading effects when using the LSP (see point 3. for more explanations). Thus, the best possible detrending for each year is the prerequisite for the LSP analysis and we focused on this and accepted small possible discontinuities at the year boundaries.

3. **I would argue that one fit for all years would precisely capture the changes in seasonal behaviour for each year, and make the years comparable. I don’t believe that this process would result in “wrong or biased LSP results” because the longest periodicity derived is 60 days and you fit only the first 3 annual terms (365,183,122 days) in the seasonal fit. Demonstrate that this**

biases the LSP results.

The LSP power is the reduction in sum of squares when fitting a sinusoid to the data. Such a fit of a sinusoid is influenced when there is an additional trend remaining after the subtraction of the seasonal fit. This would be the case when one fit for the whole time series would be used. As we are interested in the fluctuations around the seasonal variation in each year such a remaining trend would be something we clearly want to avoid.

We identified 3 possible problems related to such underlying trends. They are described and illustrated with artificial examples in the following.

1. A trend can lead to a reduction of the normalized power in comparison to the same time series without a trend. The trend is an additional part of the total variance of the time series and so the oscillation explains less of this total variance. As a consequence the significance of the result is reduced as well because of the reduced power. This can lead to a missed event.




Figure 1: Left: Time series of a sinusoid with a period of 10 days with (red) and without (blue) additional trend. Right: Resulting LSP for the two time series with same colours. The horizontal dashed line marks the maximum power achievable for a single sinusoid.

2. A trend can also lead to a shift of the period where the maximum power is observed. This would lead to a bias of the results.




Figure 2: Left: Time series of a sinusoid with a period of 50 days with (red) and without (blue) additional trend. Right: Resulting LSP for the two time series with same colours. The vertical dashed line marks the period of 50 days.

3. In some cases for periods with a period length close to the window length the amplitude of the oscillation is overestimated as the fitted sinusoid tries to compensate also for the trend. This would lead to a wrong estimation of the amplitude.




Figure 3: Left: Time series of a sinusoid with a period of 55 days with (red) and without (blue) additional trend. Right: Resulting amplitudes for the two time series with same colours. The vertical dashed line marks the period of 55 days and the horizontal dashed line marks the expected and true amplitude of 3K.

In total, a remaining trend can lead to miss detections, wrong results or biases in the results. We clearly want to avoid all of these known problems and thus decided to treat each year separately.

4. I note the rephrasing of the text but this section is still difficult to interpret.

We rephrased the part again.

- 7. I remain of the opinion (along with reviewer 1) that the first method of counting wave occurrence is invalid. Fig 3a) and associated commentary should be removed.**

We removed the figure and associated comments.

- 8. The seasonal dependence figure has been reduced to periods less than or greater than 20 days. How was this division chosen ? What is the significance of 20 days ?. Line 232 (tracked changes) still refers to periods below 10 days.**
The 20 days were chosen by hand as they lie in the gap between the two regimes of shorter periods and longer periods. Here, we count the 28 days events to the longer periods and the quasi-2-day, quasi-5-day, quasi-10-day, and quasi-16-day events to the shorter periods. But to make the division and the differences more clear, we made a new figure where we additionally displayed the differences for periods < 10 days and periods between 10 and 20 days. Now one can see that events for the periods < 10 days show the maxima at the equinoxes and observations in the summer, whereas the events for the periods between 10 and 20 days do neither show clear maxima at the equinoxes nor maxima in winter. The observations are more equally distributed over the whole year with no detections in summer.

Ground-based observations of periodic temperature fluctuations in the mesopause region with periods ~~larger~~ longer than 2 days

Christoph Kalicinsky^{1,*}, Robert Reisch^{1,2,*}, and Peter Knieling¹

¹Institute for Atmospheric and Environmental Research, University of Wuppertal, Germany

²Institute of Climate and Energy Systems: Stratosphere (ICE-4), Research Center Jülich, Germany

*These authors contributed equally to this work.

Correspondence: Christoph Kalicinsky (kalicins@uni-wuppertal.de)

Abstract. We analysed more than 30 years (1988–2021) of $\Theta_{HOH}^*(3,1)$ rotational temperatures observed from Wuppertal, Germany, with respect to periodic fluctuations (2 to 60 d) using the Lomb-Scargle periodogram. The main type of fluctuation observed in the last decades shows a period of about 28 d ~~and is probably a Rossby wave (1,4) mode~~. Other periods which are frequently found in the observations lie in the period ranges around 2 d, 5 to 6 d, 8 to 12 d, and around 15 d and can likely be
5 assigned to the quasi-2-day, the quasi-5-day, the quasi-10-day, and the quasi-16-day wave, respectively.

The ~~wave activity is typically larger~~ occurrence frequency of the waves is typically higher in winter time than in summer time because of the different wave filtering in summer and winter. This winter to summer difference holds for waves with ~~larger longer~~ periods, but it breaks off in the case of ~~smaller shorter~~ periods below about 20 d. The occurrence frequency of these waves with periods below 20 d exhibit two smaller maxima around the equinoxes. ~~Thereby the~~ A further division of these
10 observations shows that waves with periods below 10 d account for the majority of observations in the ~~months from month~~ April to September. Waves with periods between 10 d and 20 d are more equally observed throughout the year except for summer.

The long-term behaviour of the wave activity ~~shows a quasi-bidecadal and a quasi-quadrennial oscillation. A further analysis suggests that the~~ indicates a quasi-bidecadal oscillation is mainly, which is likely driven by the amplitude of the waves, ~~whereas~~
15 ~~the quasi-quadrennial oscillation is likely caused by changes in the length of the events.~~

1 Introduction

Planetary waves are large scale global phenomena that are known to have an important role for the global circulation due to transport and deposition of momentum for a long time (e.g. Salby , 1984; Andrews et al, 1987; Volland , 1988). They are typically generated at lower altitudes, propagate upwards, and are even able to reach the mesosphere and lower thermosphere
20 (MLT) region under certain conditions (e.g. Holton , 1984; Laštovička , 1997; Smith , 2003; Sassi et al., 2012). Ground-based observations of wind, temperatures, and airglow in the MLT region proved as a good way to observe planetary or planetary-like waves with different periods and monitor their temporal evolution (e.g. Espy et al., 1997; Yoshida et al., 1999; Bittner et al., 2000; Luo et al., 2000; Takahashi et al., 2002; Kishore et al., 2004; Espy et al., 2005; French et al., 2005; Takahashi et al., 2005; Höppner and Bittner , 2007; Stockwell et al., 2007; López-González et al., 2009; Day and Mitchell, 2010a, b; Hecht et

25 al., 2010; French and Kelkociuk , 2011; Takahashi et al., 2013; Egito et al., 2018; Zhao et al., 2019; Reisin , 2021). In the MLT region waves with largely different periods have been observed in the past. These periods range from only a few days in the case of very fast waves with period up to 4 d (e.g. Yoshida et al., 1999; Takahashi et al., 2005; López-González et al., 2009; Hecht et al., 2010; Egito et al., 2018; Reisin , 2021) to periods in the range of almost a week to some weeks (5 to 30 days) (e.g. Espy et al., 1997; Luo et al., 2000; French et al., 2005; Jarvis , 2006; Day and Mitchell, 2010a, b; French and Kelkociuk ,
30 2011; Takahashi et al., 2013; Zhao et al., 2019) to periods even longer than 30 days (Espy et al., 2005; Stockwell et al., 2007; French and Kelkociuk , 2011). Some of these fluctuations at specific periods can be assigned to different Rossby wave modes. These modes are influenced by the distribution of zonal background winds and appear in the presence of such winds in rather specific period ranges such as the Rossby wave (1,4) mode at about 28 d ~~and the (1,3) mode at about 16~~ (e.g. Kasahara , 1980; Salby , 1981a, b).

35 Compared to satellite observations which only observe a local point a few times a day, ground-based observations have the advantage of measuring nearly continuously. Furthermore, some ground-based instruments have been operating for several decades as the instrument can be maintained continuously which is not possible for satellite instruments. Thus, ground-based instruments and their corresponding long-time records are very suitable not only for the detection of waves but also for the analysis of the long-term evolution and the occurrence frequencies of waves with different periods. The observations by the
40 GRIPS (GRound-based Infrared P-branch Spectrometer) instruments at Wuppertal provide one of the longest temperature time series for the mesopause region around the whole globe which has been used in several different studies (e.g. Bittner et al., 2000; Oberheide et al., 2006; Höppner and Bittner , 2007; Offermann et al., 2010; Kalicinsky et al., 2016, 2018, 2024). The observations at Wuppertal have already been used to analyse planetary waves in the past by using wavelet transform and wave proxies (Bittner et al., 2000; Höppner and Bittner , 2007). Höppner and Bittner (2007) observed a long-term periodic behaviour
45 of the wave activity with a period of roughly 20 years similar to the Hale cycle. In their study they used the standard deviation of the temperature residuals (after subtracting the seasonal variations) as proxy for the planetary wave activity as at least a large part of these temperature fluctuations are thought to be caused by planetary or planetary-like wave activity (Bittner et al., 2000; Höppner and Bittner , 2007).

In our new study we now have the large advantage of an extended time series as Höppner and Bittner (2007) only used ob-
50 servations until 2005. Furthermore, we used a different technique based on the Lomb-Scargle periodogram to detect periodic fluctuations ~~and~~ which is well suited to handle time series with data gaps (Kalicinsky et al., 2020). Compared to the previous studies this technique no longer requires data assimilation before analysis to deal with the data gaps (Bittner et al., 2000; Höppner and Bittner , 2007). The paper is structured as follows. Sect. 2 describes the measurements technique and the data as well as the method to detect the periodic fluctuations with some improvements made during this study. The results of the analysis with
55 respect to the occurrence frequencies of waves with different periods and the long-term behaviour of planetary scale waves are presented in Sect. 3. These results are discussed and compared to previous results and different other observations in Sect. 4. Finally, Sect. 5 summarizes the results.

2 Measurements and data analysis

This section summarises all important information regarding the measurements and data analysis. First, the instruments and the measurement technique are described. Second, the detrending of the OH* rotational temperatures by using a seasonal fit is explained. In the last subsection the moving Lomb Scargle method is explained. This method is used to analyse the residual temperatures with respect to periodic fluctuations in the period range between 2 and 60 d. In contrast to Kalicinsky et al. (2020), where the method is described for the first time, we improved the method in this study to overcome some of the drawbacks of the former method. These improvements are detailed in Sect. 2.3.

2.1 GRIPS instruments

The temperature observations used in this study were derived from the measurements of two GRIPS (GROund-based Infrared P-branch Spectrometer) instruments, namely GRIPS-II and GRIPS-N. GRIPS-II started its measurements in the early 1980s with temporary measurements and the continuous operation started in mid-1987. It stopped working in 2011 because of a detector failure. The GRIPS-N instrument is the follow-up instrument of GRIPS-II and continues the measurements until present (Kalicinsky et al., 2018, 2024). Both instruments were operated at Wuppertal, Germany (51° N, 7° E). GRIPS-II ~~is~~ was a Czerny-Turner spectrometer with a Ge detector cooled by liquid nitrogen (see e.g. Bittner et al., 2002, for instrument details). The GRIPS-N instrument is also a Czerny-Turner spectrometer equipped with a thermoelectrically cooled InGaAs detector. The instrument has very similar optical properties as the GRIPS-II instrument, which makes it a suitable replacement instrument (Kalicinsky et al., 2018, 2024).

Both instruments measure three emission lines of the OH*(3,1) band in the near infrared region (1.524 μm –1.543 μm), namely the P₁(2), P₁(3), and P₁(4) lines. The layer of excited OH molecules is located at about 87 km height and has a layer thickness of approximately 9 km (Oberheide et al., 2006; Offermann et al., 2010). The measurements were carried out every night, except in nights with cloudy conditions. Favourable measurement conditions occur on about 220 nights per year (Oberheide et al., 2006; Offermann et al., 2010). The relative intensities of the P₁(2), P₁(3), and P₁(4) lines are used to derive rotational temperatures (see Bittner et al., 2000, and references therein). The OH* rotational temperature may deviate from the kinetic temperature, especially in cases of higher vibrational levels. However, rotational temperatures that were determined from emissions originating from the OH*(3,1) band are expected to be close to the kinetic temperatures (Noll et al., 2015). In addition, the analysis of periodic fluctuations does not require absolute values.

2.2 Seasonal variations of OH* rotational temperatures

In order to analyse periodic fluctuations with periods of a few days to a few weeks one has to remove the seasonal variations first as these variations have large amplitudes and otherwise would mask the periodic fluctuations of interest. The seasonal variations can be described by an annual, a semi-annual and a ter-annual cycle (Bittner et al., 2000; Kalicinsky et al., 2024).

The full description of the seasonal fit is

$$T = T_0 + \sum_{i=1}^3 A_i \cdot \sin\left(\frac{2 \cdot \pi \cdot i}{365.25}(t + \phi_i)\right), \quad (1)$$

90 where T_0 is the annual average temperature, t is the time in days of year, and A_i , ϕ_i are the amplitudes and phases of the sinusoids. These fit parameters vary from year to year, e.g. there is a declining trend of the amplitude of the annual cycle of about -0.5 K decade⁻¹ (see Kalicinsky et al., 2024, for more details). Because of these variations a fit according to Eq. 1 is used to derive residual temperatures for each year separately. One fit for the whole time series of temperatures for more than 35 years is not advisable, since the resulting residuals after subtraction of such a fit would still contain some variations (e.g. trends or year to year variations) with implications on ~~further analyses.~~ Fig. the further analysis. The residual temperatures are analysed with a moving Lomb Scargle method and this method requires the best possible detrending of the data with respect to the seasonal variations in each year to avoid misleading results (see Sect. 2.3). Additionally, the analysis is performed such that the centre days of the time windows used always cover a complete year. To ensure that each day of a year can act as a centre day the time series used for the calculation of the residuals has to be larger than the year itself and we added two extra months at either side of the year. Then a window centred around the first and the last day of the year can be used. Another benefit of this approach is a better constrained fit at the beginning and the end of the year, since the behaviour of the observations before and after the year of interest is considered. Therefore, the new fit for the larger time interval typically compares better to a fit that is calculated for the complete winter only (fit time interval: July 1st to June 30th (of the next year)) at the beginning and end of the year. This also leads to a reduction of discontinuities at the year borders compared to a fit using the data of one year only. Fig. 1 shows an example ~~for the seasonal variations. The black dots show the observed OH* rotational temperatures and the red curve shows the fit curve. In the lower panel of~~ of the seasonal fit for the year 1997. The data are shown in a 16 month interval centred in summer. In addition to the seasonal fit in this time interval (red curve in Fig. 1) the fits to the data from July 1st to June 30th of the previous and following year are shown as blue curves. In the transition region from one year to the next (solid blue line in Fig. 1) the agreement between the fit for the year itself (red curve in Fig. 1) and the two fits centred in winter (solid blue line in Fig. 1 ~~the~~) is reasonably good. The vertical dashed lines mark the time interval of 14 months which is the largest interval used in the moving Lomb Scargle method (see Sect. 2.3). The residual temperatures (data - ~~fit~~ curve) ~~are shown. These residual temperatures are used to derive periodic fluctuations. Seasonal variations of the OH* rotational temperatures in the year 1997. The black dots show the OH* rotational temperatures and the red curve shows the fit curve according to Eq. 1. In the lower panel the residual temperatures (data - fit curve) are displayed.~~) that are analysed are shown in the lower panel of Fig. 1.

105
110
115

2.3 Moving Lomb-Scargle periodogram

The periodic fluctuations are analysed with the moving Lomb-Scargle periodogram (Kalicinsky et al., 2020). ~~This~~ The method is based on the original Lomb-Scargle periodogram (LSP) ~~, a~~ (Lomb, 1976; Scargle, 1982). This method that can detect periodic fluctuations in all kind of time series even in time series with unequal spacing, which is the case when data gaps are

120




Figure 1. Seasonal variations of the OH* rotational temperatures in the year 1997 with two months added at both sides. The black dots show the OH* rotational temperatures and the red curve shows the fit curve according to Eq. 1 to the 16 months interval. The blue curves show the fits to the data from July 1st to June 30th of the previous and following year. The vertical dashed lines mark the 14 month interval centred in summer. In the lower panel the residual temperatures (data - fit curve) of this 14 month interval are displayed.

present (Lomb, 1976; Seargle, 1982). Thus, it is very suitable to analyse OH*(3,1) rotational temperature time series that exhibit irregular data gaps due to weather conditions. In the approach described by Kalicinsky et al. (2020) a time window of fixed length, for example 60 d, is used for the analysis. The starting point is the beginning of the time series and the window is shifted by the minimum sampling step (here one day) until the end of the time series is reached. For the data points within each of these individual time windows, that all include a different part of the complete time series, a LSP is calculated separately. In this way periodic fluctuations can be detected as well as the time in combination with the temporal evolution of the periods and amplitudes of these fluctuations can be observed (see Kalicinsky et al., 2020, for a complete description of the method). ~~Due to the variation of the seasonal fits from year to year, one fit for the whole time series of temperatures for more than 35 years is not advisable. Thus, the residuals are calculated for each year separately. The following LSP analysis is performed such that the centre days of the time windows always cover a complete year. To ensure that each day of a year can act as a centre day the time series used for the calculation of the residuals has to be larger than the year itself and we added two extra months at either side of the year. This enables a LSP for each day of the year as centre day even for window lengths of several weeks and also accounts for the variations of the seasonal fits. Another benefit of this approach is a better constrained fit at the beginning and the end of the year, since the behaviour of the observations before and after the year of interest is considered. Therefore, the new fit for the larger time interval typically compares better to a fit that is calculated for the complete winter only (fit time interval: July 1st to June 30th (of the next year)) at the beginning and end of the year.~~ The use of a fixed window length also has a drawback, which is mainly noticeable in the case of signals with periods much smaller shorter than the window length. It is not very likely that such a signal with a period of only a few days remains in atmospheric temperatures for several weeks.

When the time window is much longer than the duration of such an event the results using the LSP are damped, i.e. the period and amplitude is a mean value over the whole time interval ~~and~~. In such a case the amplitude is typically much smaller than it would be when ~~only the shorter duration length of the event~~ a shorter time window would be used ~~as time window~~. ~~In the worst case events with small~~. As a consequence events with short periods and a small duration may be missed in the analysis. In order to avoid this drawback we introduced a varying window length here. The window length changes ~~in the same way as~~ simultaneously to the analysed period, i.e. a smaller window is used for ~~smaller shorter~~ periods and vice versa. The window length can only be reduced to a certain threshold value, because of ~~the significance of the results and two important points~~. First, the data gaps ~~that~~ should not exceed the length of the window used. ~~Below this threshold value the window length stays constant~~. As Second, the significance of the results depends on the number of data points (e.g. Cumming et al. , 1999; Zechmeister and Kürster , 2009; Kalicinsky et al., 2020) ~~it is not possible to use which hinders the use of~~ too small windows. The significance of a result for the same signal decreases if less data points were incorporated in the analysis, i.e. the possibility to produce such a signal by chance e.g. due to noise is larger in the case of less data points. Therefore, a signal that accounts for the same variance of the total time series could be significant if more data points were used for the analysis ~~and not significant for less data points~~. In total, the minimum window length has to be a trade-off. On the one hand, it has to be large enough to avoid problems with too many data gaps and worse significance ~~compared to a larger window~~ and, on the other hand, it has to be small enough to detect events with ~~smaller periods that occur in the time range of days to one or two weeks~~ short periods and short duration. ~~Below the threshold value the window length stays constant~~.

We also used the number of cycles of an oscillation at a given period as a factor to calculate the window lengths and the stepping. A factor of one means the analysed period and the time window are always the same. In the case of the factor of 2 the window length is twice as large as the period analysed using this window, i.e. two cycles of an oscillation with a certain period would fit in the time window used for the analysis. The minimum window length (divided by the factor) still defines the minimum period below which the time window stays constant. For example the scaling factor of 2 sets the minimum period to 12 d up to which the periods below are analysed with the defined window length of 25 d. After the period of 12 d the rounded double window length corresponding to the analysed period is used, e.g. 26 d for 12.6 d, 27 d for 13.2 d, 30 d for 14.9 d, 40 d for 19.9 d and 120 d for 60 d. The scaling of the minimum period has been done to ensure a softer transition to larger window lengths.

The significance of the results is analysed using the false alarm probability (FAP). The FAP gives the probability that a certain peak with a certain normalized power can occur just by chance somewhere in the whole analysed period range. It is determined using an empirical expression where the coefficients were determined by using Monte Carlo simulations. By using this expression the significance for each peak can be evaluated depending on the window length, period range and number of data points (see Kalicinsky et al., 2020, for more details). Compared to the former approach described in Kalicinsky et al. (2020) the range of window lengths is now larger, since the window length changes in relation to the analysed period. In the former evaluation of the significance in dependence of the window length only lengths of 30 d and larger have been considered. In this study we also use smaller window lengths. Therefore we reevaluated the significance in dependence of the used window length and the analysed period range in a larger range of window lengths (10 to 90 d) and for more representations. Here we simulated 100

times 10000 representations instead of only one time 10000 representations as done in Kalicinsky et al. (2020). In this way
175 we obtained new results that give refined coefficients for Eq. 6 in Kalicinsky et al. (2020) with an improved estimation of the
uncertainties. The new values of coefficients are $2.98 \pm 0.02 \text{ dd}^{-1}$ for the slope and $-0.23 \pm 0.01 \text{ d}^{-1}$ for the intercept. The
given uncertainties are one times the standard deviation of their mean. Thus, the new results show that the former results by
Kalicinsky et al. (2020) were by chance at the end of the 3σ range for both values. Additionally, small differences may occur
from the wider range used for the window lengths.

180 A prerequisite for the whole method is a good detrending with respect to the seasonal variations in each year. Otherwise,
remaining underlying trends will influence the results. Three different problems are known to us. First, a trend can lead to a
reduction of the normalized power in comparison to the same time series without trend. As a consequence the significance of
the result is reduced as well because of the reduced power. This can lead to a missed event. Second, a trend can also lead to
a shift of the period where the maximum power is observed. This would lead to a bias of the results. Third, in some cases for
185 periods with a period length close to the window length the amplitude of the oscillation is overestimated as the fitted sinusoid
tries to compensate also for the trend. This would lead to a wrong estimation of the amplitude. Thus, we used the fitting
approach as described in Sect. 2.2 to avoid these problems.

Fig. 2 shows a comparison between the new results using the varying window and the old results with a fixed window length of
60 d for the example year 2005. The values used for the analysis with varying window length are a minimum window length
190 of 25 d and a factor of 1.0. The window length stays constant at a period of 25 d and below and monotonically increases with
increasing period above this value. Obviously, the new results allow for more variability of the amplitude as the smoothing
effect is smaller. Because of these lower fluctuations and ~~also due to~~ higher significance levels in the case of more data points
used for the analysis, it ~~also~~ may occur that the time interval showing significant results is slightly larger in the time domain
for the results with the fixed window. This effect mainly shows up at ~~larger-longer~~ periods. The benefit of the new method
195 arises at ~~smaller-shorter~~ periods. Thus, in the result for the improved method many more significant events are detected (see
Fig. 2a). These events are typically very short in duration. In the case of a fixed window with too large size these events will be
smoothed such as they are not detectable any more (see Fig. 2b). Thus, with the method described by Kalicinsky et al. (2020)
significant events could be missed and the improved method presented here gives a large benefit.

3 Results

200 This section is divided into two parts. In subsection 3.1 the periods of different fluctuations that are typically caused by planetary
waves are analysed. This includes the occurrence frequencies of the observed periods and also seasonal differences of the
occurrence frequencies for specific period ranges. Subsection 3.2 focuses on the long-term evolution of the wave activity, which
mainly deals with the question if there is a long-term trend or even a long-term periodic behaviour of the wave activity. All
presented figures were deduced from LSP results that have been calculated with the new approach with varying window length.
205 The minimum window length was 25 d and the factor 1. Only periodic fluctuations that have been classified as significant




Figure 2. Comparison of the LSP results for the amplitude in the year 2005. **a** Improved method with varying time window. The minimum window length used for the analysis was 25 d, i.e. below a period of 25 d the window length was constant at 25 d and monotonically increased above this threshold period. **b** Former method with fixed window length of 60 d. The x-axes show the centre days of the time windows and the y-axis the period and frequency, respectively. The amplitude displayed is colour coded. The black contour lines mark the region of significant results.


events using the FAP and our empirical equation to calculate it (see Sect. 2.3) were taken into account, i.e. only results inside the contour lines in Fig. 2a were used.

3.1 Occurrence frequencies of waves

~~We use two different ways to count the wave events. First, In the first step~~ we simply counted the days of each individual event, i.e. all centre days that lie inside a contour line and, thus, show a significant event. ~~This is a measure of the importance of waves at different periods with respect to the total time dominated by these waves.~~ As events with larger longer period tend to also last longer, ~~it is expected that there is an overall increase of the counted days with increasing period. Therefore, we used a second way and we~~ divided the counts in each bin by the mean period of the corresponding bin. This leads to an observation

frequency in terms of cycles and removes the dependency on the period. These results ~~can be used to compare~~ show the relative importance of the waves at different periods. Furthermore, it still accounts for the length of the individual events, i.e. longer events including more cycles still get a stronger weight compared to shorter events. This would not be the case if the events themselves would be counted only.

Figure 3 shows the occurrence frequency of events with certain periods ~~for the two different ways of counting~~. ~~Figure 3a shows the expected overall increase of the occurrence frequency with increasing period when simply days are counted. Nevertheless, at some specific periods events occur more often than events at other periods. Here, the counted days clearly separate from the expected overall increase of these days with increasing period. These~~ The preferred periods are about 28 d, 15 d, 12 d, 10 d, 8 d and in the period range of 5 to 6 d. ~~The same peaks are still present after dividing the peaks height by the mean periods of the corresponding bins, but with different relative heights compared to each other (see Fig. 3b). In addition to the previously mentioned peaks also a clear peak at about 2, and 2 d. Thereby, the most events were observed at periods of about 28 dis present which is very small when counting days only. Furthermore, the relative importance of the waves in the period range around 15. Additionally, there are numerous detections of waves with periods longer than 30 dis reduced, but here no clear maximum can be observed.~~



ab

Figure 3. Occurrence frequency of events in dependency of the period. ~~a The histogram shows the number of centre days of the LSP analysis with significant results plotted against the corresponding period. b~~ The histogram shows the number of centre days of the LSP analysis with significant results divided by the mean periods of the corresponding bins.

The seasonal dependency of the occurrence frequencies is largely dependent on the period itself. Figure 4 shows the number of centre days with significant events plotted against the month for different period ranges. In the upper panel of Fig. 4 the period range is larger longer or equal to 20 d (left) and shorter than 20 d (right). Obviously, events with these periods more often occur periods < 20 d occur more often in winter time than in summer time, ~~whereby~~. In this period range the largest numbers were observed in January and February and the lowest numbers in June. In the case of events with shorter periods

the situation is largely different. ~~The lower panel of Fig. 4 shows the same type of plots for~~ In the period range below $\lesssim 20$ d
235 ~~. In this period range~~ the largest number of events were detected around the equinoxes (late March and late September). The
maximum number of ~~significant~~ days with significant events were observed in April and September. Thereby, the number of
days with significant events around the equinox in fall is larger than in spring time. Note here, that this important result is only
visible by using the new approach with the varying window length. When using the former approach described in Kalicinsky
et al. (2020) a large number of events with ~~smaller-shorter~~ periods are missed and the clear seasonal structure with the maxima
240 around the equinoxes is obscured. The lower panel of Fig. 4 shows a further subdivision of the shorter periods. The results for
the period range below 10 d is show in the lower left panel and the results for the period range between 10 d and 20 d is shown
in the lower right panel. The contribution of events with periods below 10 d is partly very high. In the months from May to
June these waves account for nearly all observations. In the months around the equinoxes (April, August, and September) the
proportion of the waves with the ~~smallest-shortest~~ periods is still larger than 60%. ~~In the other months the proportions typically~~
245 ~~vary around 50% and the waves with periods < 10 d still account for the majority of observations in the whole period range~~
< 20 d. The results in the period range between 10 and 20 d do neither show clear maxima at the equinoxes nor maxima in
winter. The observations are more equally distributed over the whole year with no detections in summer.

The distribution of the observed amplitudes of the events is shown in Fig. 5. Note here that all single amplitudes that
correspond to centre days within an event are displayed, i.e. one event would have its own distribution of amplitudes. The
250 mean amplitude of all events is 7.9 K with a standard deviation of 2.8 K. The smallest amplitudes that belong to a significant
fluctuation are slightly above 2 K and the largest observed amplitudes are nearly 16 K. When the amplitudes are plotted for
different period ranges (not shown), no obvious difference can be seen. Thus, the amplitudes are not significantly larger for
events with ~~smaller-or-larger-shorter or longer~~ periods. Note here that there might be some reduction of the amplitudes due to
vertical averaging, especially for waves with ~~smaller-shorter~~ periods and smaller vertical wavelengths. In the majority of the
255 cases waves with periods even below about 10 d exhibit vertical wavelengths that are a multiple of the FWHM of the OH layer
(e.g. Buriti et al., 2005; Ern et al., 2013; Yamazaki and Matthias, 2019; Reisin, 2021). Very small vertical wavelengths below
e.g. 20 km are only reported for a very minor portion of all observations (Reisin, 2021). Thus, the averaging effect should be
small or maybe even negligible.

3.2 Long-term evolution of wave activity

260 The second part of this study deals with the long-term evolution of the wave activity. In former studies this long-term evolution
was analysed by using the standard deviation of the residual temperatures (data - seasonal fit) for each single year as a proxy
for the wave activity in this year (e.g. Höppner and Bittner, 2007). The authors observed a quasi-bidecadal oscillation of the
wave activity in this study. The standard deviation as proxy has the drawback that all kind of fluctuations are included in this
quantity and not only the significant ones. Because of this, we use the mean amplitude of the significant events as a new proxy.
265 This new proxy also has a small drawback, because it does not include the length of the events, only the strength. Thus, we
also use the amplitude weighted sum of significant days as another proxy, i.e. the mean amplitude times the number of days at
which a significant event was detected. As all of the three proxies have similarities and also differences the comparison of the




Figure 4. Seasonal dependency of the occurrence frequencies of events with different periods. In the upper panel of the figure the number of centre days with significant results are plotted against the month for the period range ≥ 20 d (left) and the period range < 20 d (right). In a further subdivision of the second period range is shown in the lower panel. The results for the period range < 10 d is shown in the lower left panel and the results for the period range between 10 d and 20 d is shown in the lower right panel.

individual LSPs can help to gain information on the importance of the length and the strength of the wave events on certain periodic behaviours. Figure 6a shows the time series of all three different quantities (with mean subtracted). The periodicity of the time series is analysed with the LSP and the results are shown in Fig. 6b with the same colours as used in Fig. 6a. The two time series in the two upper panels in Fig. 6a, the standard deviation and the mean amplitude, show a rather similar behaviour with corresponding times of maxima and minima. Only at some points, for example in the years 2007 and 2009, there are larger differences. The LSPs for these two time series show also peaks at very similar periods (see Fig. 6b). The main peak in the long-period range is located at about 20 y in both cases, whereby the maximum in the LSP for the mean amplitude lies slightly below 20 y (red curve in Fig. 6b) and in the case of the standard deviation slightly above this value (black curve in Fig. 6b). Due to the uncertainty (the width of the peaks) this difference is not significant. These two peaks at about 20 years are




Figure 5. Distribution of the observed amplitudes. All single amplitudes that correspond to centre days within an event are displayed.

not significant at a 95% confidence level with respect to the complete analysed period range (FAP). This is a common problem with this rather conservative approach in the case when several similar large peaks occur in a periodogram, but does not mean that the oscillations are not real. With respect to the single frequency the significant level for the peak at 20 years is almost 280 95% in the case of the standard deviation and better than 95% for the mean amplitude, i.e. it is rather uncertain that a peak with that height occurs at exactly this period just by chance. As we are searching for a peak at a period of 20 years because of the former study by Höppner and Bittner (2007), this second way is valid here. The time series of the weighted sum of significant days does not show the same long periodic fluctuation as the other two quantities (lower panel of Fig. 6a and blue curve in Fig. 6b). The difference between the time series of the mean amplitude and that of the weighted sum is simply the number of days with significant results. Consequently, the fact that only one time series shows the long-term oscillation implies that 285 only the amplitude of the events shows these long-periodic behaviour and not the number of days. As the standard deviation is a measure of the amplitudes of all fluctuations within the analysed time interval, the standard deviation shows almost the same long-term behaviour. Furthermore, this also indicates that in most years the significant fluctuations dominate the standard deviation of the residual temperatures.

290 The peak that has the largest power in all of the periodograms is the peak at about 4 years in the LSP of the weighted sum of days. As before the peak is not significant with respect to the complete period range but highly significant with respect to the single frequency. A fit to the time series gives a period of nearly exactly 4 years. As ~~peaks at a period of about 4 are only seen with largely reduced power in the LSP for the standard deviation and with even less power in the LSP for the mean amplitude, the conclusion here is opposite to the quasi-bidecadal oscillation and suggests that the length of the events shows a~~ quasi-quadrennial oscillation only one proxy shows a clear peak, we do not further interpret the results.




Figure 6. Long-term behaviour of the wave activity. **a** The upper panel shows the time series of the yearly standard deviation of the temperature residuals after subtracting the seasonal fit. In the middle panel the time series of the mean amplitude of all significant events in a single year is displayed. The lower panel shows the product of the mean amplitude and the sum of the centre days with a significant result in the LSP. **b** The LSP for the three time series are shown with the same colours as of the time series themselves (black: standard deviation, blue: weighted sum, red: mean amplitude).

4 Discussion

The discussion is divided into two parts. First, the ~~possible origin of the observed fluctuations is discussed and observed~~ fluctuations are compared to other results ~~. Additionally, and~~ the seasonality of the wave ~~activity occurrence~~ activity occurrence is examined. In the second part, the long-term behaviour of the wave activity and the fluctuation of this activity itself is discussed.

300 4.1 Periodic fluctuations

The highest peak in the histogram occurs at a period of around 28 d (about 25 to 30 days in Fig. 3), ~~which~~. This shows that period fluctuations in this period range are very common. Waves in this period range are observed at the largest number of days (not shown) as well as for the largest number of cycles. Thus, these waves are the most important ones in relative and absolute terms. And indeed, in a larger number of winters an event with an average period in this range takes place. Fig. 7 shows an example for such an event. The time series of the temperature residuals over the winter 1997/1998 is shown as black curve in Fig. 7a. Obviously, between the days 160 and 260 a large periodic fluctuation is present. The amplitude of this fluctuation increases at the beginning and reaches maximum values around day 215. Then it decreases again. The red curve shows a sinusoid with varying amplitude fitted to the data. The resulting mean amplitude is about 26 d and, thus, falls in the period range from about 25 to 30 d. The LSP results in Fig. 7b shows some deviation from this mean period at the beginning of the event, where the period of the maximum peak in the LSP lies slightly above 30 d, but only for a few days. Therefore, we would still categorize the whole event as a quasi-28 day wave. A very similar event was observed over Antarctica at Rothera station in the winter 2014 (Zhao et al., 2019). The authors used ground-based and satellite data to analyse the vertical and horizontal structure of this event and find that it was consistent with the Rossby wave (1,4) mode. This is also consistent with theory where the Rossby wave (1,4) mode has a period of about 28 d in the presence of zonal background winds (Kasahara, 1980). Zhao et al. (2019) state in their study that the propagation in the MLT region is severely limited in summer because of the low phase speed of this type of waves. They also refer to a study by Sassi et al. (2012) who found that the propagation to higher altitudes is expected to be transient because of the effect of the background winds and the wind filtering. This could also explain that we observe the 28 d wave typically in winter and not in summer. There is another possible effect on the temperature in the mesopause region that shows a periodic behaviour in the right same period range: the 27-day solar cycle. But the known amplitudes of this influence are considerably smaller than the ones observed here. Beig et al. (2008) reviewed studies concerning the influence of the 27-day solar cycle on temperatures in the mesosphere and lower thermosphere and reported values less than 4 K ~~and typically smaller~~. Typically the values in the mesopause region are smaller than in the mesosphere itself. Other studies also show periodic fluctuations in coincidence with the 27-day solar cycle (a time-lag is present) with amplitudes smaller than 1 K (von Savigny et al., 2012; Thomas et al., 2015; Rong et al., 2020). Thus, the amplitudes of the fluctuations observed here are much larger than what is expected in the case of the 27-day solar cycle. We also analysed the GRIPS OH(3,1) rotational temperatures with respect to the influence of the 27-day solar cycle and observed similar amplitudes (< 1 K) than in previous studies. Furthermore, we excluded time intervals in winter with large wave activity strong events in the 27 d period range from the analysis and observed no significant differences to the previous results (personal communication C. v. Savigny). This led us to the conclusion that the influence of the 27-day solar cycle on the temperatures and the fluctuation with periods of about 28 d that partly show amplitudes larger greater than 10 K are two completely independent phenomena. Hence, we believe that the Rossby (1,4) mode could be a possible explanation for our observations. Since we only have local observations from one single observation site and neither information on the horizontal nor information on the vertical structure, this identification is a guess and not a proven fact. Additional data would be necessary here and in the following cases to obtain a proven identification of




Figure 7. Example for an event with a large temperature fluctuation with a period of around 28 d. **a** The temperature time series after subtracting the seasonal variations is shown as black curve (the seasonal fit was centred in winter). The red curve is the fit to the data in the middle part of the time series. The x-axis show days since July 1st, 1997. **b** LSP result for the amplitudes of periodic fluctuations over the winter 1997/1998. The x-axis show the centre days of the windows since July 1st, 1997 and the y-axis show the period and frequency of the fluctuations, respectively. The amplitude is displayed colour coded. The LSP was calculated with a fixed window of 60 d to focus on the long periods and slightly smooth the results for better visibility. Furthermore, the influence of the partly larger data gaps is reduced.

Another period range that is frequently observed for planetary waves in the MLT region lies in the region around 16 d (e.g. Espy et al., 1997; Luo et al., 2000; French et al., 2005; Jarvis, 2006; Day and Mitchell, 2010b; French and Kelkociuk, 2011; Takahashi et al., 2013). We also observe a smaller peak in the histogram in the period range around 15 d (see Fig. 3). ~~Waves in this region may be an observation of a Rossby wave (1,3) mode as shown for winds (Kasahara, 1980; Salby, 1981b) and temperatures (Espy et al., 1997). The range of periods that were predicted in the presence of zonal background winds range from 12 to 20 (Salby, 1981b) and 16 to 19 (Kasahara, 1980), respectively. Thus, although our observations peak at about~~

~~15, the observed period range between about 14 to 20 is still in this range and observations of this Rossby wave mode are a possibility, but for a proved identification additional data are necessary~~ and we would assign this as a quasi-16-day wave, too. Note here ~~also~~ that the resolution of the LSP (FWHM) which is about $1/T$ (see Cumming et al. , 1999), where T is the length of the time window (in this period range 25 d). This means that single observations with periods of 15 d or 16 d are not significantly different. Another wave that has frequently been mentioned in literature is the quasi-5-day wave (e.g. Wu et al., 1994; Jarvis , 2006; Day and Mitchell, 2010a). In our observations we observe also a larger number of significant events in this period range. More precisely, we observe two peaks, one at about 5 d and one at about 6 d. However, due to the resolution of the LSP, these two peaks would not be significantly different for single detections. Thus, we would assign all of these ~~observation observations~~ to the quasi-5-day wave. The ~~Rossby wave (1,1) mode has been shown to appear in this period range for winds (Kasahara , 1980; Salby , 1981b) and the predicted range of periods for the observation of this mode is reported to be 4.4 to 5.7 (Salby , 1981b). This range is fitting quite well to our observations which makes this specific Rossby wave mode a candidate for our observations.~~ The quasi-10-day wave is also a known wave type in the atmosphere (e.g. Jarvis , 2006; Takahashi et al., 2013; Forbes and Zhang , 2015). In our observation observations we see a very prominent peak at about 10 d and two additional peaks nearby at slightly above 8 d and slightly below 12 d. According to the resolution of the LSP results we would assign all observation in this range to the quasi-10-day wave. It has been shown for winds that the Rossby wave (1,2) mode appears in this period range in the presence of a background wave (Kasahara , 1980; Salby , 1981b) with a predicted period range that goes from 8.3 to 10.6 (Salby , 1981b). In the case of temperatures between 20 and 100 observations in this range to the quasi-10-day wave ~~was also connected to a Rossby wave mode. Hence, such a Rossby wave mode is a possible explanation for at least a part of our observations in this period range.~~ The quasi-2-day wave is also a prominent feature in the MLT region (e.g. Wu et al., 1993; Takahashi et al., 2005; López-González et al., 2009; Hecht et al., 2010; Yue et al., 2012; Reisin , 2021). We also observed enhanced activity observed an enhanced occurrence frequency in this period range ; ~~which gets clearly visible when the counted days are normalized by the mean period of the bin (see Fig. 3b)), too.~~ Relative to its period length the quasi-2-day wave is a prominent feature of our observations. ~~This type of wave is can also be a Rossby wave mode with zonal wave number 3 (Wu et al., 1993; López-González et al., 2009) and the normal mode in this case is (3,0) (Kasahara , 1980; Yue et al., 2012). This type of Rossby wave mode is possibly the explanation of our observations, too.~~ The last major part of events is detected We also observed many events in the period range above 30 d. ~~As mentioned above, the large increase is at least partly caused by the fact that periodic fluctuations with longer periods typically also lasts longer which enhances the days at which these waves are observed. Nonetheless, it does not explain the observation itself.~~ Planetary waves with very similar periods have also been reported for other ~~observations~~ observation sites in Antarctica (Espy et al., 2005; Stockwell et al., 2007; French and Kelkociuk , 2011).

Observations of different Rossby wave modes could possibly explain a number of our observations, e.g. the Rossby wave mode (1,3) 16-day-wave (Kasahara , 1980; Salby , 1981b; Espy et al., 1997) or the Rossby wave mode (3,0) the quasi-2-day wave (Kasahara , 1980; Wu et al., 1993; López-González et al., 2009; Yue et al., 2012), and cause the observations to appear primarily in certain period ranges. However, a better constraint of the origin of the waves and their complete ~~structure~~ structure would need additional data and further investigations which is beyond the scope of this paper.

The seasonal variation of the ~~wave-activity-occurrence frequency of the waves~~ varies for different types of waves. In the period range below 20 d we observe two maxima ~~in the activity~~ around the spring equinox in April and in late summer (August/September), whereby the major part of the observations ~~aound-around~~ the equinoxes and in summer stem from waves with periods below 10 d (compare Fig. 4). In the case of waves with longer periods (> 20 d) a large maximum in winter and late fall is observed ~~whereas the activity~~. Observations in summer and other seasons ~~is-are~~ comparable small (compare Fig. 4).

A larger ~~wave-activity-occurrence frequency~~ for waves with ~~smaller-shorter~~ periods in other seasons than winter has also been observed by others. Reisin (2021) analysed the seasonality of the quasi-2-day wave and observed ~~higher-activity-a larger number of events~~ in Southern hemisphere summer than winter. The analysis by López-González et al. (2009) revealed higher ~~activity-occurrence probabilities~~ for the quasi-2-day wave observed in OH temperatures in the period from about January to August, ~~whereas~~. in the same study the probability for the quasi-5-day wave shows local maxima in March/April, August/September and November. Wu et al. (1994) analysed two years of HDRI (High Resolution Doppler Imager) data and observed wave events of the quasi-5-day wave mainly in April/May and in September to October. Radar observations over India showed larger activities of the 6.5-day wave around the equinoxes (April/May and September/October) (Kishore et al., 2004). Riggins et al. (2006) also observed larger events in April/May when analysing SABER data over a time period of three years. Observations in both hemispheres at polar sites showed strong wave activity for the 5-day wave in winter and late summer (August/September) but no significant events at equinoxes (Day and Mitchell, 2010a). In a former study of the GRIPS observations at Wuppertal the authors also reported a larger amount of events in summer time in the case of planetary waves with ~~smaller-shorter~~ periods (Bittner et al., 2000). In total, the seasonal variation of the wave activity in the case of ~~smaller~~ ~~shorter~~ periods observed at Wuppertal agrees quite well with other observations. The observation of large planetary wave events in summer cannot be explained with the excitation of these waves in the troposphere and their propagation up to the MLT region, because the mean stratospheric flow in summer prevents these upward propagation of most waves (e.g. Charney and Drazin, 1961). Thus, other mechanisms are necessary to explain the observations. Two main mechanisms are discussed in previous work. On the one hand, the waves can be excited at higher altitudes and propagate upwards afterwards and, on the other hand, ducting from the winter hemisphere to the summer hemisphere could take place (e.g. Bittner et al., 2000; Riggins et al., 2006; Day and Mitchell, 2010a, and references therein). In the period range between 10 and 20 d we observe ~~activity~~ ~~wave events~~ in most of the seasons except for summer (~~not shown~~). The quasi-16-day wave and a part of the observations of the quasi-10-day wave are mainly responsible for this ~~enhancements~~. Most of the other studies found in literature also report on similar seasonal distributions. For the quasi-10-day wave Forbes and Zhang (2015) presented maximum activity in winter and around the equinoxes in mid-latitudes. The analysis of OH temperatures by López-González et al. (2009) showed the by far largest ~~activity-occurrence probability of~~ the quasi-10-day in September and October and some smaller ~~activity-probability~~ in the time period from February to June ~~and no~~. No observations in July and August are reported in the study. In the case of the quasi-16-day wave the picture is quite similar with ~~larger-stronger~~ activity in winter and only minor activity in summer months (Luo et al., 2000; Day and Mitchell, 2010b). Nonetheless, the quasi-16-day wave has also been observed in summer (e.g. Espy et al., 1997). Discussed mechanisms are again local phenomena and ducting from the winter to the summer hemisphere (see Espy et al., 1997; Luo et al., 2000, and references therein).

In the case of wave events with periods ~~larger~~longer than 20 d we observe a large summer to winter differences~~with a~~. A huge number of events ~~that~~ took place in winter time or late fall and only a very small number of events in summer. As already mentioned this is expected because of the wave filtering in summer that prevents the wave propagation to the MLT region.

415 These findings are also confirmed by previous other studies. The observation of the ~~large~~-28 d wave event reported by Zhao et al. (2019) also took place in winter. Similarly, Stockwell et al. (2007) report that the largest activity of waves in the period range from 30 to 50 d is observed in late winter.

4.2 Long-term behaviour of wave activity

We see a main long-periodic fluctuation with a period of about 20 years in the long-term behaviour of the wave activity. This

420 can be seen in two of the three different proxies. Of the two new proxies only the yearly mean amplitude of the significant events shows a clear long-term behaviour. Therefore, the standard deviation, which is mainly determined by the amplitude of the significant fluctuations in most years, also shows a very similar long-term behaviour. The fact, that the time series of the weighted sum of days with significant results does not show the same long-period fluctuation suggests that in years with higher activity during the quasi-bidecadal oscillation (for example mid-1990s) not more events are expected but events with larger

425 amplitudes than in the years with lower activity (for example around 2005).

The quasi-bidecadal oscillation of the standard deviation has also been observed by Höppner and Bittner (2007) for a shorter time interval of observations (until 2005). Thus, their findings agree well with ours and the wave activity observed by the standard deviation proxy continues in this quasi-bidecadal oscillatory way. Jarvis (2006) also observed a quasi-bidecadal oscillation in the wave activity of the 5-day planetary wave. Like us Jarvis (2006) observed the quasi-bidecadal oscillation

430 in the residual planetary wave amplitudes. The observed change lies in the range of about 10% which is in line with our observations here, where we see a change of roughly ± 1 K and a mean amplitude of about 8 K. In contrast to our findings, where we see the quasi-bidecadal oscillation in the mean over all periods in the range from 2 to 60 d, Jarvis (2006) observed the long-term fluctuation for the 5-day planetary wave only and not for the 10-day and 16-day planetary wave.

In former studies of the OH*(3,1) rotational temperatures we already observed the quasi-bidecadal oscillation in yearly mean

435 temperature observations (Kalicinsky et al., 2016, 2018, 2024). The phase of the quasi-bidecadal oscillation of the temperature observations is nearly the same as that of the mean amplitude and the standard deviation of the residuals, respectively. All of the time series show maxima around the mid-1990s and around 2015/16 in addition to a minimum around 2005. This means the larger amplitudes of the significant wave events occur together with enhanced yearly mean temperatures and vice versa. ~~As~~
~~we~~We also observed the quasi-bidecadal oscillation in other altitudes such as the mesosphere and stratosphere with alternating

440 sign from one atmospheric layer to the other (stratosphere and lower thermosphere (GRIPS OH*(3,1) rotational temperatures) are in phase and the mesosphere is shifted by 180°) (Kalicinsky et al., 2018) and also on ground (Kalicinsky and Koppmann, 2022)~~, this~~. This temperature oscillation in the atmosphere might influence other atmospheric parameters such as the wind and, therefore, also influence the wave filtering. Likely, only in certain years the conditions are favourable for waves with smaller amplitudes to reach higher altitudes, whereas in other years only waves with amplitudes that are large enough reach

445 the observation altitude of the GRIPS instrument. But a complete analysis of this hypothesis will require additional data and is

beyond the scope of this paper.

Additionally, we found ~~large~~ some indications for a quasi-quadrennial oscillation of the weighted sum of days of significant events, which is a proxy for wave activity that also largely takes into account the ~~the length of the events. As this oscillation is observed in a reduced way in the time series of the standard deviation and even more reduced in that of the mean amplitude,~~
450 ~~the conclusion for the quasi-quadrennial oscillation is opposite to that for the quasi-bidecadal oscillation. The length of the events shows a long-term periodic behaviour with a period of about 4 and not the amplitude, i.e. the strength of the events.~~ French et al. (2020) observed a very similar quasi-quadrennial oscillation for the OH/OH* rotational temperatures above Davis, Antarctica. The phase of the two oscillations is also quite similar with the Wuppertal oscillation slightly preceding by up to one year. French et al. (2020) discussed a potential role of planetary or tidal waves in ~~the~~ their quasi-quadrennial oscillation.
455 This would agree well with our observations and could possibly explain the time lag. Nonetheless, a complete analysis would require additional data and is beyond the scope of this paper. Furthermore, a longer time series and a potentially increased significance of the results would be beneficial as well.

5 Summary and conclusions

We analysed more than 30 years of OH(3,1) rotational temperatures observed from Wuppertal with respect to periodic fluctua-
460 tions in the period range from 2 to 60 d. Fluctuations with a period around 28 d are the main fluctuation observed in this era. ~~These fluctuations are probably Rossby waves (1,4) mode that appear with a period of about 28 in ground-based measurements in presence of a zonal background wind.~~ Other period ranges which are often detected in the observations are around 2 d, 5 to 6 d, 8 to 12 d, and around 15 d.

Most of the ~~wave activity is~~ waves are observed in winter time because of the different wave filtering in summer and winter,
465 i.e. the conditions in winter are typically more favourable for waves to reach higher altitudes. This winter to summer difference is not universal for all waves. It holds for waves with larger longer periods, but it breaks off in the case of smaller shorter periods below about 20 d. These waves with smaller shorter periods occur more evenly distributed across the year with even a larger number of events around the equinoxes. Thereby waves with periods below 10 d are nearly completely responsible for the observations in summer and to a large degree for that around equinoxes.

470 The mean amplitude of all observed significant events is about 8 K, whereby the amplitudes range from about 2 to 16 K. A significant difference for the distribution of the amplitudes dependent on the period range was not observed.

The analysis of the long-term behaviour of the wave activity revealed a quasi-bidecadal ~~and a quasi-quadrennial oscillation.~~
The quasi-bidecadal oscillation. This oscillation is observed in two wave proxies, the standard deviation of the temperature residuals and the mean amplitude of the significant events within a year. Since the last wave proxy, the amplitude weighted
475 sum of days with significant results, does not show this fluctuation, the conclusion is that the amplitude is the quantity that shows the quasi-bidecadal oscillation. This means, that in certain years not more events but events with larger amplitudes are expected, whereas in other years the mean amplitude of the events is smaller. The quasi-bidecadal oscillation is in phase with the oscillation of the yearly mean temperatures themselves. A connection between changes in the background temperature field

and, thus, also other parameters, may influence the wave filtering and lead to the observation of the quasi-bidecadal fluctuation
480 of the wave activity. ~~The quasi-quadrennial oscillation is primarily observed in the weighted sum of the days of significant
485 events, which leads to the conclusion that the length of the events show this oscillation rather than the amplitude.~~

Data availability. The OH(3,1) rotational temperatures which were derived from the GRIPS observations at Wuppertal can be obtained by request to the corresponding author.

Author contributions. CK conceptualised the study. RR performed the analyses of OH(3,1) rotational temperatures under intensive discus-
485 sion with CK. PK provided the OH(3,1) rotational temperatures. The article was written by CK with contributions from all coauthors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This research was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – 519284835. We gratefully acknowledge the two anonymous reviewers that helped to certainly improve the manuscript.

References

- 490 Andrews, D., Holton, J., and Leovy, C.: Middle atmosphere dynamics, Academic Press, London, 1987.
- Beig, G., Scheer, J., Mlynczak, M. G., and Keckhut, P.: Overview of the temperature response in the mesosphere and lower thermosphere to solar activity, *Rev. Geophys.*, 46, RG3002, doi:10.1029/2007RG00236, 2008.
- Bittner, M., Offermann, D., and Graef, H.H.: Mesopause temperature variability above a midlatitude station in Europe, *J. Geophys. Res.*, 105, 2045–2058, doi:10.1029/1999JD900307, 2000.
- 495 Bittner, M., Offermann, D., Graef, H.H., Donner, M., and Hamilton, K.: An 18-year time series of OH* rotational temperatures and middle atmosphere decadal variations, *J. Atmos. Sol. Terr. Phys.*, 64, 1147–1166, doi:10.1016/S1364-6826(02)00065-2, 2002.
- Buriti, R. A., Takahashi, H., Lima, L. M., and Medeiros, A. F.: Equatorial planetary waves in the mesosphere observed by airglow periodic oscillations, *Adv. Space Res.*, 35, 2031–2036, doi:10.1016/j.asr.2005.07.012., 2005.
- Charney, J. G., and Drazin, P. G.: Propagation of planetary-scale disturbances from lower into the upper atmosphere, *J. Geophys. Res.*, 66, 500 83–109, 1961.
- Cumming, A., Marcy, G.W., and Butler, R.P.: The lick planet search: detectability and mass thresholds, *Astrophysical Journal*, 526, 890–915, doi:10.1086/308020, 1999.
- Day, K. A., and Mitchell, N. J.: The 5-day wave in the Arctic and Antarctic mesosphere and lower thermosphere, *J. Geophys. Res.*, 115, D01109, doi:10.1029/2009JD012545, 2010.
- 505 Day, K. A. and Mitchell, N. J.: The 16-day wave in the Arctic and Antarctic mesosphere and lower thermosphere, *Atmos. Chem. Phys.*, 10, 1461–1472, doi:10.5194/acp-10-1461-2010, 2010
- Egito, F., Buriti, R. A., Frago Medeiros, A., and Takahashi, H.: Ultrafast Kelvin waves in the MLT airglow and wind, and their interaction with the atmospheric tides, *Ann. Geophys.*, 36, 231–241, doi:10.5194/angeo-36-231-2018, 2018.
- Ern, M., Preusse, P., Kalisch, S., Kaufmann, M., and Riese, M.: Role of gravity waves in the forcing of quasi two-day waves in the meso- 510 sphere: An observational study, *J. Geophys. Res. Atmos.*, 118, 3467–3485, doi:10.1029/2012JD018208, 2013.
- Espy, P. J., Stegman, J., and Witt, G.: Interannual variations of the quasi-16-day oscillation in the polar summer mesospheric temperature, *J. Geophys. Res.*, 102(D2), 1983–1990, doi:10.1029/96JD02717, 1997.
- Espy, P. J., Hibbins, R. E., Riggin, D. M., and Fritts, D. C.: Mesospheric planetary waves over Antarctica during 2002, *Geophys. Res. Lett.*, 32, L21804, doi:10.1029/2005GL023886, 2005.
- 515 Forbes, J. M., and Zhang, X.: Quasi-10-day wave in the atmosphere, *J. Geophys. Res. Atmos.*, 120, 11079–11089, doi:10.1002/2015JD023327, 2015.
- French, W. J. R., Burns, G. B., and Espy, P. J.: Anomalous winter hydroxyl temperatures at 69°S during 2002 in a multiyear context, *Geophys. Res. Lett.*, 32, L12818, doi:10.1002/2015JD023327, 2005.
- French, W. J. R., and Klekociuk, A. R.: Long-term trends in Antarctic winter hydroxyl temperatures, *J. Geophys. Res.*, 116, D00P09, 520 doi:10.1029/2011JD015731, 2011.
- French, W. J. R., Klekociuk, A. R., and Mulligan, F. J.: Analysis of 24 years of mesopause region OH rotational temperature observations at Davis, Antarctica – Part 2: Evidence of a quasi-quadrennial oscillation (QO) in the polar mesosphere, *Atmos. Chem. Phys.*, 20, 8691–8708, doi:10.5194/acp-20-8691-2020, 2020.

- Hecht, J. H., Walterscheid, R. L., Gelin, L. J., Vincent, R. A., Reid, I. M., and Woithe, J. M.: Observations of the phase-locked 2 day wave
525 over the Australian sector using medium-frequency radar and airglow data, *J. Geophys. Res.*, 115, D16115, doi:10.1029/2009JD013772,
2010.
- Holton, J. R.: The Generation of Mesospheric Planetary Waves by Zonally Asymmetric Gravity Wave Breaking, *J. Atmos. Sci.*, 41, 3427 –
3430, doi:10.1175/1520-0469(1984)041<3427:TGOMPW>2.0.CO;2, 1984.
- Höppner, K., and Bittner, M.: Evidence for solar signals in the mesopause temperature variability?, *J. Atmos. Sol. Terr. Phys.*, 69, 431–448,
530 doi:10.1016/j.jastp.2006.10.007, 2007.
- Jarvis, M.J.: Planetary wave trends in the lower thermosphere – Evidence for 22-year solar modulation of the quasi 5-day wave, *J. Atmos.
Sol. Terr. Phys.*, 68, 1902–1912, doi:10.1016/j.jastp.2006.02.014, 2006.
- Kalicinsky, C., Knieling, P., Koppmann, R., Offermann, D., Steinbrecht, W., and Wintel, J.: Long-term dynamics of OH* temperatures over
central Europe: trends and solar correlations, *Atmos. Chem. Phys.*, 16, 15 033–15 047, doi:10.5194/acp-16-15033-2016, 2016.
- 535 Kalicinsky, C., Peters, D. H. W., Entzian, G., Knieling, P., and Matthias, V.: Observational evidence for a quasi-bidecadal oscillation in the
summer mesopause region over Western Europe, *J. Atmos. Sol. Terr. Phys.*, 178, 7–16, doi:10.1016/j.jastp.2018.05.008, 2018.
- Kalicinsky, C., Reisch, R., Knieling, P., and Koppmann, R.: Determination of time-varying periodicities in unequally spaced time series of
OH* temperatures using a moving Lomb-Scargle periodogram and a fast calculation of the false alarm probabilities, *Atmos. Meas. Tech.*,
13, 467–477, doi:10.5194/amt-13-467-2020, 2020.
- 540 Kalicinsky, C., and Koppmann, R.: Multi-decadal oscillations of surface temperatures and the impact on temperature increases, *Sci. Rep.*,
12, 19895, doi:10.1038/s41598-022-24448-3, 2022.
- Kalicinsky, C., Kirchhoff, S., Knieling, P., and Zlotos, L.O.: Long-term variations in the mesopause region derived from OH*(3,1) rotational
temperature observations at Wuppertal, Germany, from 1988 – 2022, *Adv. Space Res.*, 73, 3398–3407, doi:10.1016/j.asr.2023.08.045,
2024.
- 545 Kasahara, A.: Effect of zonal flows on the free oscillations of a barotropic atmosphere, *J. Atmos. Sci.*, 37(5), 917–929, doi:10.1175/1520-
0469(1980)037<0917:EOZFOT>2.0.CO;2, 1980.
- P. Kishore, Namboothiri, S. P., Igarashi, K., Gurubaran, S., Sridharan, S., Rajaram, R., and Venkat Ratnam, M.: MF radar observations of
6.5-day wave in the equatorial mesosphere and lower thermosphere, *J. Atmos. Sol.-Terr. Phys.*, 66, doi:10.1016/j.jastp.2004.01.026, 2004.
- Laštovička, J.: Observations of tides and planetary waves in the atmosphere-ionosphere system, *Adv. Space Res.*, 20, 1209–1222,
550 doi:10.1016/S0273-1177(97)00774-6, 1997.
- Lomb, N.R.: Least-squares frequency analysis of unequally spaced data, *Astrophysics and Space Science*, 39, 447–462, 1976.
- López-González, M. J., Rodríguez, E., García-Comas, M., Costa, V., Shepherd, M. G., Shepherd, G. G., Aushev, V. M., and Sargoytchev, S.:
Climatology of planetary wave type oscillations with periods of 2–20 days derived from O2 atmospheric and OH(6-2) airglow observations
at mid-latitude with SATI, *Ann. Geophys.*, 27, 3645–3662, doi:10.5194/angeo-27-3645-2009, 2009.
- 555 Luo, Y., Manson, A. H., Meek, C. E., Meyer, C. K., and Forbes, J. F.: The quasi 16-day oscillations in the mesosphere and lower thermosphere
at Saskatoon (52°N, 107°W), 1980–1996, *J. Geophys. Res.*, 105(D2), 2125–2138, doi:10.1029/1999JD900979, 2000.
- Noll, S., Kausch, W., Kimeswenger, S., Unterguggenberger, S., and Jones, A. M.: OH populations and temperatures from simultaneous
spectroscopic observations of 25 bands, *Atmos. Chem. Phys.*, 15, 3647–3669, doi:doi.org/10.5194/acp-15-3647-2015, 2015.
- Oberheide, J., Offermann, D., Russell III, J.M., and Mlynarczyk, M.G.: Intercomparison of kinetic temperature from 15 μm CO₂ limb
560 emissions and OH*(3,1) rotational temperature in nearly coincident air masses: SABER, GRIPS, *Geophys. Res. Lett.*, 33, L14811,
doi:10.1029/2006GL026439, 2006.

- Offermann, D., Hoffmann, P., Knieling, P., Koppmann, R., Oberheide, J., and Steinbrecht, W.: Long-term trend and solar cycle variations of mesospheric temperature and dynamics, *J. Geophys. Res.*, 115, D18127, doi:10.1029/2009JD013363, 2010.
- Reisin, E. R.: Quasi-two-day wave characteristics in the mesopause region from airglow data measured at El Leoncito (31.8°S, 69.3°W), *J. Atmos. Sol.-Terr. Phys.*, 218, 105613, doi:10.1016/j.jastp.2021.105613, 2021.
- 565 Rigglin, D. M., Liu, H.-L., Lieberman, R. S., Roble, R. G., Russell III, J. M., Mertens, C. J., Mlyneczek, M. G., Pancheva, D., Franke, S. J., Murayama, Y., Manson, A. H., Meek, C. E., and Vincent, R. A.: Observations of the 5-day wave in the mesosphere and lower thermosphere, *J. Atmos. Sol.-Terr. Phys.*, 68, 323-339, doi:10.1016/j.jastp.2005.05.010, 2006.
- Rong, P., von Savigny, C., Zhang, C., Hoffmann, C. G., and Schwartz, M. J.: Response of middle atmospheric temperature to the 27 d solar cycle: an analysis of 13 years of microwave limb sounder data, *Atmos. Chem. Phys.*, 20, 1737–1755, doi:10.5194/acp-20-1737-2020, 2020.
- 570 Salby, M. L.: Rossby Normal Modes in Nonuniform Background Configurations. Part I: Simple fields, *J. Atmos. Sci.*, 38, 1803–1826, 1981.
- Salby, M. L.: Rossby Normal Modes in Nonuniform Background Configurations. Part II: Equinox and Solstice Conditions, *J. Atmos. Sci.*, 38, 1827–1840, 1981.
- 575 Salby, M.: Survey of planetary-scale travelling waves: the state of theory and observation, *Rev. Geophys. Space Phys.*, 22, 209–236, 1984
- Sassi, F., Garcia, R. R., and Hoppel, K. W.: Large-scale Rossby normal modes during some recent Northern Hemisphere winters, *J. Atmos. Scien.*, 69(3), 820–839, doi:10.1175/JAS-D-1-0103.1, 2012.
- Scargle, J.D.: Studies in astronomical time series analysis. II. Statistical aspects of spectral analysis of unevenly spaced data, *Astrophysical Journal*, 263, 835–853, 1982.
- 580 Smith, A. K.: The Origin of Stationary Planetary Waves in the Upper Mesosphere, *J. Atmos. Sci.*, 60, 3033 – 3041, doi:10.1175/1520-0469(2003)060<3033:TOOSPW>2.0.CO;2, 2003.
- Stockwell, R. G., Rigglin, D. M., French, W. J. R., Burns, G. B., and Murphy, D. J.: Planetary waves and intraseasonal oscillations at Davis, Antarctica, from undersampled time series, *J. Geophys. Res.*, 112, D21107, doi:10.1029/2006JD008034, 2007.
- Takahashi, H., Buriti, R. A., Gobbi, D., Batista, P. P.: Equatorial planetary wave signatures observed in mesospheric airglow emissions, *J. Atmos. Sol.-Terr. Phys.*, 64, 1263–1272, doi:10.1016/S1364-6826(02)00040-8, 2002.
- 585 Takahashi, H., Lima, L. M., Wrasse, C. M., Abdu, M. A., Batista, I. S., Gobbi, D., Buriti, R. A., and Batista, P. P.: Evidence on 2–4 day oscillations of the equatorial ionosphere h'F and mesospheric airglow emissions, *Geophys. Res. Lett.*, 32, L12102, doi:10.1029/2004GL022318, 2005.
- Takahashi, H., Shiokawa, K., Egito, F., Murayama, Y., Kawamura, S., Wrasse, C. M.: Planetary wave induced wind and airglow oscillations in the middle latitude MLT region, *J. Atmos. Sol.-Terr. Phys.*, 98, 97–104, doi:10.1016/j.jastp.2013.03.014, 2013.
- 590 Thomas, G. E., Thurairajah, B., Hervig, M. E., and von Savigny: Solar-induced 27-day variations of mesospheric temperature and water vapor from the AIM SOFIE experiment: Drivers of polar mesospheric cloud variability, *J. Atmos. Sol.-Ter. Phys.*, 134, 56–68, doi:10.1016/j.jastp.2015.09.015, 2015.
- Volland, H.: *Atmospheric Tidal and Planetary Waves*. Kluwer Academic Publishers, Boston, 1988.
- 595 von Savigny, C., Eichmann, K.-U., Robert, C. E., Burrows, J. P., and Weber, M.: Sensitivity of equatorial mesopause temperatures to the 27-day solar cycle, *Geophys. Res. Lett.*, 39, L21804, doi:10.1029/2012GL053563, 2012.
- Wu, D. L., Hays, P. B., Skinner, W. R., Marshall, A. R., Burrage, M. D., Lieberman, R. S., and Orland, D. A.: Observations of the quasi 2-day wave from the High Resolution Doppler Imager on Uars, *Geophys. Res. Lett.*, 20(24), 2853–2856, doi:10.1029/93GL03008, 1993.

- 600 Wu, D. L., Hays, P. B., and Skinner, W. R.: Observations of the 5-day wave in the mesosphere and lower thermosphere, *Geophys Res. Lett.*, 21, 2733–2736, doi:10.1029/94GL02660, 1994.
- Yamazaki, Y., and Matthias, V.: Large-amplitude quasi-10-day waves in the middle atmosphere during final warmings. *Journal of Geophysical Research: Atmospheres*, 124, 9874–9892. doi:10.1029/2019JD030634, 2019.
- Yoshida, S., Tsuda, T., Shimizu, A., and Nakamura, T.: Seasonal variations of 3.0~3.8-day ultra-fast Kelvin waves observed with a meteor wind radar and radiosonde in Indonesia, *Earth Planet Space*, 51, 675–684, doi:10.1186/BF03353225, 1999.
- 605 Yue, J., Liu, H.-L., and Chang, L. C.: Numerical investigation of the quasi 2 day wave in the mesosphere and lower thermosphere, *J. Geophys. Res.*, 117, D05111, doi:10.1029/2011JD016574, 2012.
- Zechmeister, M., and Kürster, M.: The generalised Lomb-Scargle periodogram - A new formalism for the floating-mean and Keplerian periodograms, *Astronomy and Astrophysics*, 496, 577–584, doi:10.1051/0004-6361:200811296, 2009.
- 610 Zhao, Y., Taylor, M. J., Pautet, P.-D., Moffat-Griffin, T., Hervig, M. E., Murphy, D. J., French, W. J. R., Liu, H. L., Pendleton Jr., W. R., and Russell III, J. M.: Investigating an unusually large 28-day oscillation in mesospheric temperature over Antarctica using ground-based and satellite measurements. *J. Geophys. Res.: Atmos.*, 124, 8576–8593. doi:10.1029/2019JD030286, 2019.