

In situ production of hybrid N₂O in dust-rich Antarctic ice

Lison Soussaintjean¹, Jochen Schmitt¹, Joël Savarino², J. Andy Menking^{3,4}, Edward J. Brook⁵, Barbara Seth¹, Vladimir Lipenkov⁶, Thomas Röckmann⁷, Hubertus Fischer¹

5

¹Climate and Environmental Physics, Physics Institute and Oeschger Centre for Climate Change Research, University of Bern, Bern, 3012, Switzerland

²University Grenoble Alpes, CNRS, IRD, Grenoble INP, INRAE, IGE, Grenoble, 38000, France

10 ³Australian Antarctic Program Partnership, Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Australia

⁴Environmental Research Unit Commonwealth Scientific and Industrial Research Organisation Aspendale, Victoria, Australia

⁵College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR 97331, USA

⁶Climate and Environmental Research Laboratory, Arctic and Antarctic Research Institute, Saint Petersburg, 199397, Russia


15 ⁷Institute for Marine and Atmospheric research Utrecht (IMAU), Utrecht University, 3584CC Utrecht, the Netherlands

Correspondence to: Lison Soussaintjean (lison.soussaintjean@unibe.ch)

Abstract. Nitrous oxide (N₂O) is a potent greenhouse gas involved in the destruction of stratospheric ozone. Past atmospheric mixing ratios of N₂O are archived in ice cores; however, the presence of in situ N₂O production in dust-rich Antarctic ice complicates their accurate reconstruction, especially during glacial periods. This production occurs in extremely cold ice and without sunlight. This study aims to understand the reaction producing N₂O in Antarctic ice by identifying the precursors and the reaction pathway. We compared the oxygen and nitrogen bulk and position-specific isotope composition of in situ N₂O in ice cores to the isotopic composition of nitrate (NO₃⁻), a possible precursor of N₂O. The ¹⁵N signature of NO₃⁻ is fully transferred into the central N atom (N^α) of in situ N₂O, but it is not transferred into the terminal N atom (N^β), resulting in a 50 % transfer of the ¹⁵N signature of NO₃⁻ into the bulk ¹⁵N isotopic composition. These findings suggest that the in situ N₂O production involves two different nitrogen precursors present in ice: the central N atom (N^α) originates from NO₃⁻ and the terminal N atom (N^β) from a different precursor not yet identified. Oxygen isotope analysis shows that NO₃⁻ cannot be the only reservoir for the O atom of in situ N₂O. Temperature, pH, and absence of sunlight in Antarctic ice point to an abiotic N-nitrosation reaction. The limiting factor of the reaction is probably associated with mineral dust and might be Fe²⁺, reducing NO₃⁻ to NO₂⁻ or the precursor of the N^β atom. The site preference (SP) values of in situ N₂O are highly variable between different ice cores and depend on the bulk ¹⁵N isotopic composition of N₂O, itself depending on the ¹⁵N isotopic composition of the NO₃⁻ precursor. This finding is unexpected because SP is usually determined by the production pathway through symmetric reaction intermediates that mix the N atoms in α and β positions and average out their isotopic difference. In contrast, our results provide the first evidence of a hybrid N₂O production

35 pathway involving an asymmetric intermediate that preserves the distinct ^{15}N signatures of two different precursors – one contributing to the N^α atom and the other to the N^β atom.

1 Introduction



40 **Figure 1.** N_2O mixing ratio, N_2O bulk nitrogen isotopic composition, and Ca^{2+} concentrations that represent the mineral dust
content, measured in the TALDICE (Schilt et al., 2010a), EDC (Schilt et al., 2010b), EDML (Fischer et al., 2019), and Vostok
(Miteva et al., 2007) ice cores. Because of its low dust content, the TALDICE record in black is considered to represent the best
estimate of atmospheric N_2O levels (see Sect. 3.2.2). Stars correspond to samples affected by in situ production and represent the
total N_2O measured, i.e., atmospheric N_2O plus in situ N_2O . Grey boxes mark the dust-rich sections of the EDC ice core
45 corresponding to the Last Glacial Maximum (LGM), the Marine Isotope Stage (MIS) 4 and 6 cold periods.

Nitrous oxide (N₂O) is a potent greenhouse gas whose global warming potential is 273 times higher than that of carbon dioxide (CO₂) on a 100-year timescale (IPCC, 2021). As a result of the anthropogenic alteration of the nitrogen cycle (Gruber and Galloway, 2008), the N₂O atmospheric mixing ratio has risen by 23 % compared to pre-industrial levels (Rubino et al., 2019), reaching 338 ppb in December 2024 (Lan et al., 2025) and contributing around 6 % to the radiative forcing from long-lived greenhouse gases. N₂O is also the primary source of stratospheric nitrogen oxides (NO_x) that destroy ozone (Ravishankara et al., 2009).

Soils and aquatic environments are the main sources of atmospheric N₂O. Production of N₂O occurs both during denitrification, the reduction of nitrate (NO₃⁻) to nitrite (NO₂⁻) and ultimately molecular nitrogen (N₂), and as a by-product of nitrification, where ammonium (NH₄⁺) is oxidized to hydroxylamine (NH₂OH) and then converted to NO₂⁻ and NO₃⁻ (Baggs, 2011; Bange, 2008). N₂O is also produced during the reduction of NO₂⁻ to N₂O in nitrifier denitrification, where nitrifying microorganisms reduce NO₂⁻ to N₂O similarly to heterotrophic denitrifiers (Wrage-Mönnig et al., 2018). These reactions are performed by bacteria, archaea and fungi, but abiotic reactions have also been documented (Zhu-Barker et al., 2015). These include chemo-denitrification (Jones et al., 2015; Tischer et al., 2022), which is the reduction of NO₃⁻ or NO₂⁻ by Fe(II), and NH₂OH oxidation by Fe(III) or Mn(IV).

The composition of the atmosphere in the past can be reconstructed by analyzing air trapped in polar and high-altitude ice cores. Ice cores are the only direct atmospheric archive and have provided continuous CO₂ and CH₄ records over the last 800,000 years covering eight glacial-interglacial cycles (Loulergue et al., 2008; Lüthi et al., 2008) and a partial record of N₂O (Schilt et al., 2010b). In addition to N₂O mixing ratios, measurement of the stable isotopic composition of N₂O is used for source attribution of N₂O emissions. In the current state of the climate system, natural sources account for 57 % of total N₂O emissions, while anthropogenic sources contribute the remaining 43 % (Tian et al., 2020). Natural sources are approximately 58 % emissions from soils, 38 % from marine ecosystems, and the remaining 4 % from lightning and atmospheric production (Tian et al., 2020). N₂O emissions from both land and marine ecosystems may have varied in the past. Because sources differ in their isotopic signatures, past changes in terrestrial and marine emissions can be reconstructed using the isotopic composition of N₂O archived in ice cores (Fischer et al., 2019; Menking et al., 2020; Schilt et al., 2014).

Beyond bulk isotope analyses ($\delta^{15}\text{N}$ and $\delta^{18}\text{O}$), position-specific measurements provide even more detailed insights into N₂O production mechanisms. Because N₂O is an asymmetric molecule, the nitrogen isotopic composition of the two N atoms can deviate from each other, and this difference can be measured separately at the central N atom (N^α, bonded to oxygen) and the terminal N atom (N^β). From these values, the site preference (SP = $\delta^{15}\text{N}^{\alpha} - \delta^{15}\text{N}^{\beta}$) can be calculated. Because SP reflects intramolecular isotope partitioning during N-N bond formation, it is primarily controlled by the reaction mechanism and the structure of the last intermediate rather than by the isotopic composition of the precursor (Frame and Casciotti, 2010; Sutka et al., 2003, 2006; Toyoda et al., 2005). As a result, SP values are often characteristic of specific N₂O formation pathways and can remain constant even when precursor $\delta^{15}\text{N}$ values vary widely. In contrast to bulk $\delta^{15}\text{N}$, which integrates source and fractionation effects, SP provides mechanistic information on how N₂O is formed and is therefore widely used to discriminate between N₂O production pathways; SP values are typically negative for bacterial denitrification and positive for

nitrification (Toyoda et al., 2017). Using this tool, Prokopiou et al. (2018) showed that SP values increased since preindustrial times, pointing to a relative shift from denitrification to nitrification, consistent with agricultural emissions playing a major role in the N₂O increase. Similarly, Menking et al. (2025) demonstrated that the increase in N₂O concentrations during the transition from the Last Glacial Maximum to the Holocene reflected contributions from both nitrification and denitrification, whereas the N₂O decrease during the Younger Dryas was driven by reduced nitrification. However, the atmospheric signal of greenhouse gases archived in ice cores can be altered (Anklin et al., 1995; Delmas, 1993; Lee et al., 2020; Mühl et al., 2023). This is particularly true for N₂O: the N₂O mixing ratio in glacial period ice from Antarctica shows large scatter between nearby samples from the same core and different values between ice cores for the same age (Fig.1) (Flückiger et al., 2002; Sowers, 2001; Stauffer et al., 2003), which is inconsistent with the homogeneous global atmospheric mixing ratio of N₂O resulting from its 123-year atmospheric lifetime (Prather et al., 2015) and the low-pass filtering of the atmospheric signal through the slow bubble enclosure process. Because these anomalies were observed for ice core analyses using two distinct gas extraction methods – wet extraction by melting (Schilt et al., 2010b) and dry extraction by grating the ice (Sowers, 2001) – it was concluded that N₂O production occurred already in the ice sheet and not during the analysis. This “in situ production” or excess N₂O in dust-rich sections of polar ice cores has two major consequences for the climatic interpretation. First, as the production process has not yet been identified, and the excess N₂O has not yet been quantified, we do not have access to past atmospheric N₂O mixing ratios during most past glacial periods (Schilt et al., 2010b), where high dust contents are found in the ice. Secondly, samples affected by in situ production deviate from the isotopic composition of atmospheric N₂O (Fig.1), which compromises source attribution (Fischer et al., 2019; Schilt et al., 2014). It is therefore necessary to understand the N₂O production process(es) in the ice to systematically identify affected samples and thus avoid misinterpretation. Several detection methods were implemented to scrutinize the N₂O record (Flückiger et al., 2004; Spahni et al., 2005), but they were only used to discard the samples likely to be affected. Understanding the production processes could potentially provide a reliable way of quantifying the fraction of N₂O produced in situ and to correct the measured signal with the aim to obtain the past atmospheric signal.

This work also has relevance for nitrogen cycle processes in extreme environments. Very few studies have focused on N₂O production under extreme conditions as those encountered in the Antarctic ice sheet, where temperatures reach down to -60 °C, pressures are enormous, and reaction time scales can be of the order of several thousand years. Priscu et al. (2008) measured very high N₂O mixing ratios in a permanently ice-covered lake in the Dry Valleys, Antarctica, which they attributed to microbial nitrification. Investigating N₂O production in Antarctic ice is an opportunity to explore whether microorganisms can be metabolically active at -60 °C. Although previous studies suggested potential N₂O production by bacteria (Flückiger et al., 2002; Schilt et al., 2010b; Sowers, 2001), there is currently no evidence to support microbial activity in Antarctic ice as a source for in situ N₂O. High microbial counts have only been found in a few of the Vostok ice core samples where in situ production was reported (Sowers, 2001). Also, abiotic reactions may cause N₂O formation in polar ice. In the Dry Valleys, Antarctica, Samarkin et al. (2010) demonstrated abiotic N₂O production by chemo-denitrification in the Don Juan Pond soils.

115 This study uses isotope analysis to characterize in situ N₂O in various ice cores. The background of the study, the extreme environmental conditions in the polar environment, and the potential consequences for the reactions involved are presented in Sect. 2. Based on the strong enrichment in ¹⁵N observed in some samples affected by in situ N₂O production (Fig. 1), we hypothesize that NO₃⁻, which can also be highly enriched in ¹⁵N in ice, is one of the nitrogen precursors for in situ N₂O. To test this hypothesis, we measured the isotopic composition of N₂O and NO₃⁻ in the same ice core samples and calculated the isotopic signature of in situ N₂O (Sects. 3 and 4). Position-specific nitrogen isotope analysis of N₂O was carried out to further constrain the reaction pathway(s) involved. The potential mechanisms for in situ N₂O production are discussed in Sect. 5.

2 Background – Potential precursors and reaction rate of N₂O production in polar ice

For both Greenland and Antarctic ice cores, N₂O production is inferred in ice-core sections corresponding to glacial periods, while there is no in situ production apparent for the Holocene and other interglacial and interstadial periods (Flückiger et al., 2004; Schilt et al., 2010b). The chemical composition of the ice differs significantly between glacial and interglacial periods, particularly in terms of its mineral dust content. Dust concentrations are significantly higher during glacial periods than during interglacial periods (factor of 30-100) in Antarctic and Greenland ice cores (Fuhrer et al., 1999; Lambert et al., 2012). In fact, N₂O excess production is only observed in dust-rich ice and, in Antarctica specifically, it increases with higher dust concentrations (Fig. 1). This points to a reaction involving at least one compound contained in or associated with the dust.

The excess N₂O produced in Antarctic ice is on average 0.4 nmol N.kg⁻¹. The main nitrogen compounds that are generally considered as N₂O precursors are NO₃⁻ and NH₄⁺. The typical concentrations of NO₃⁻ and NH₄⁺ in ice are above 320 nmol N.kg⁻¹ and 55 nmol N.kg⁻¹, respectively (Kaufmann et al., 2010; Röthlisberger et al., 2000a), and therefore more than sufficient to produce the observed excess N₂O. However, neither NO₃⁻ nor NH₄⁺ spontaneously react to N₂O. These precursors need to be activated by a chemical or biochemical agent, such as iron II (Fe²⁺) (Zhu-Barker et al., 2015) or bacteria, respectively. Both Fe²⁺ (Spolaor et al., 2012, 2013; Traversi et al., 2004) and bacteria (Miteva et al., 2016; Rohde et al., 2008; Sowers, 2001) were detected in polar ice.


Previous research has shown that samples affected by in situ production exhibit isotopic deviations from the atmospheric signature that differ in both direction and magnitude across ice cores (Fischer et al., 2019; Menking et al., 2025; Schilt et al., 2014; Sowers, 2001). It has been observed that the ¹⁵N signatures of total N₂O – i.e. atmospheric plus in situ N₂O – depend on snow accumulation: sites with low accumulation show enrichment in ¹⁵N compared to the atmospheric signature (Sowers, 2001), while sites with higher accumulation show depletion in ¹⁵N (Fischer et al., 2019) (Fig. 1). A similar dependence of isotopic composition on accumulation rate is observed for NO₃⁻. The accepted reason is that after deposition some of the NO₃⁻ undergoes photolysis in the photic zone, corresponding approximately to the first meter of the surface snow. Kinetic isotope effects for photolysis favor the loss of the lighter isotopologues (i.e. containing ¹⁴N), resulting in ¹⁵N enrichment of the remaining NO₃⁻ (Blunier et al., 2005; Frey et al., 2009). The fraction of photolyzed NO₃⁻ and the degree of isotopic

enrichment depends on the time NO_3^- spends in the photic zone before it is archived in the ice by progressive burial. At low-accumulation sites, the duration within the photic zone is longer and a large fraction of NO_3^- is photolyzed leading to a highly ^{15}N -enriched NO_3^- archived in the ice, reaching values of $\delta^{15}\text{N}_{\text{bulk}} = 300 \text{ ‰}$ at Concordia (Erbland, 2011; Erbland et al., 2013). In ice core samples affected by in situ N_2O production, the presence of high $\delta^{15}\text{N}_{\text{bulk}}$ values of N_2O along with ^{15}N -enriched NO_3^- strongly suggests that in situ N_2O production uses N from NO_3^- . The isotopic composition of NH_4^+ has never been measured in Antarctic ice cores, because large amounts of ice are required at the low NH_4^+ concentrations, and the samples are highly sensitive to NH_3 contamination (Lamothe et al., 2023). Although a low contamination protocol has been developed by Lamothe et al. (2023), it was used for samples with higher NH_4^+ concentrations, where the blanks are small compared to the sample concentration. Nonetheless, the hypothesis of NH_4^+ as an N_2O precursor is discussed in Sect. 5.1.

The production of N_2O might already start near the surface of the ice sheet, in the snow and firn, but most of the produced N_2O would be lost to the atmosphere via air exchange. In the diffusive zone of the firn column and below the air lock-in depth (Battle et al., 1996), exchange of the firn air with the atmosphere is limited or excluded and the produced N_2O is partly or fully preserved in the ice. To be able to observe in situ N_2O in ice cores, the production must continue close to or below the lock-in depth (located between 50 to 100 m, depending on the location and conditions of the ice sheet). For example, during the LGM at Vostok, Antarctica, the lock-in depth was around 94 m, and the age of the ice at this depth was 5200 years (Table 1). This means that at Vostok, N_2O was still produced several thousand years after the precursors were deposited onto the ice sheet. This observation shows that either it takes time for the precursors to come into contact, for example by diffusion in the ice or ice recrystallization, or that the reaction is very slow due to low temperatures. In the latter case, it would not be possible to reproduce such slow reaction kinetics in the laboratory. This observation also shows that the reaction does not need light.

3 Samples and Methods

3.1 Ice core samples



170

Figure 2. Location of the ice core drilling sites presented in this study. Maps created using the Quantarctica (Matsuoka et al., 2018) and QGreenland (Moon et al., 2023) QGIS data packages.

To gain a comprehensive understanding of the N_2O production process(es) in diverse environmental settings, we analyzed
175 ice core samples from different sites with different characteristics: variable dust contents, variable dust chemical
compositions, variable snow accumulation rates and therefore isotopic signatures of NO_3^- , and variable drilling site
temperatures. These characteristics are reported in Table 1. Calcium (Ca^{2+}) concentrations are used as a proxy for dust
content because they have been found to be correlated with mineral dust concentrations (Ruth et al., 2002). Figure 2 shows
the location of the drilling sites.

180 We analyzed samples from five Antarctic ice cores: the Talos Dome Ice Core (TALDICE), the European Project for Ice
Coring in Antarctica (EPICA) Dome C core (EDC), the EPICA Dronning Maud Land core (EDML), the Vostok ice core,
and the main transect of the Taylor Glacier (TG) blue ice area. Additionally, we analyzed samples from one Greenland ice
core, the North Greenland Ice Core Project ice core (NGRIP). The TALDICE ice core was selected for its low dust content
(and expected low degree of in situ N_2O) compared to other Antarctic ice cores, the EDC and Vostok ice cores for their low
185 snow accumulation rate, the EDML ice core for its relatively higher snow accumulation rate. The NGRIP ice core was
analyzed to compare the isotopic signature of in situ N_2O in Greenland and Antarctic ice.

We measured the stable isotopic composition ($\delta^{15}\text{N}_{\text{bulk}}$ and $\delta^{18}\text{O}$) of N_2O in the TALDICE, EDC, EDML, Vostok, TG and NGRIP ice cores, and the stable isotopic composition ($\delta^{15}\text{N}$ and $\delta^{18}\text{O}$) of NO_3^- in all samples except NGRIP. We also carried out position-specific isotope analysis of the central ($\delta^{15}\text{N}^\alpha$) and terminal ($\delta^{15}\text{N}^\beta$) nitrogen atoms in N_2O in the Vostok and TG ice cores.

The samples date from either the Last Glacial Maximum (LGM) and glacial-interglacial transition (14 to 29 ka), Marine Isotope Stage 4 (MIS4, 57 to 71 ka), or Marine Isotope Stage 6 (MIS6, 130 to 191 ka), which are all dust-rich glacial periods and are therefore prone to in situ production of N_2O . We selected samples younger than 150 ka because the N_2O atmospheric baseline recorded in the dust-poor TALDICE ice core and used in our mass balance approach (see Sect. 3.2.2) does not extend to older periods.

Table 1. Characteristics of the ice core drilling sites presented in this study.

	TG[†]	Vostok	EDC	EDML	TALDICE	NGRIP
Latitude (°)	-77.7598 ^b	-78.4642	-75.1000	-75.0019	-72.8302	75.1000
Longitude (°) https://add.scar.org/	161.7212 ^b	106.8370	123.3326	0.0663	159.2004	-42.32
Elevation (MSL) (https://add.scar.org/)	580	3488	3233	2892	2320 ^c	2917
accumulation rate^{*.a} in ice equivalent per year (cm)	0.8 ^b	1.2	1.4	3.7	4.5	6.3
Age maximum^a (ka)	150	408	813	145	343 ^d	125
Annual mean snow surface temperature[*] (°C)	-46 [†]	-66 ^c	-60 ^c	-53 ^c	-44 ^c	-46 ^f
Maximum depth where the N_2O production occurred (m)	-	300	400	770	750	1600
Temperature range of the ice where the production occurred (bore hole temperature) (°C)	-	-56.6 - -55.2 ^g	-53.6 - -51.6 ^h	~ -40 ⁱ	-	-30 - -32 ^j
Ice age – Gas age difference (Δage)^{*.a} (yr)	3350 ^b	5170	3960	1590	1000	890
Air lock-in depth^a (m)	29 [†]	94	83	84	70	79
Ca²⁺ concentration[*] (ng.g⁻¹) measured by continuous	21±15 ^b	-	35±21 ^l	47±37 ^m	16±9 ⁿ	292±187 ^o

flow analysis (CFA)^k						
Fe concentration* (ng.g⁻¹) <i>[measurement technique]</i> Fe^{2+*} Fe^{3+*}	-	-	~0.5 ^p [CFA] ~13 ^p [ICP-SFMS]	-	~5 ^q [CFA] ~3 ^q [CFA] ~2 ^q [CFA]	-
Dominant source of dust	Southern South America (SSA) ^r	SSA ^s	SSA ^t	SSA ^t	SSA ^u	East Asian deserts ^v

[†]special case: horizontal core. LGM mean snow surface temperature estimated based on $\delta^{18}\text{O}(\text{H}_2\text{O})$ data. LGM lock-in depth calculated using the Herron-Langway firn model with delta age and temperature inputs for the TG accumulation zone near

200 Taylor Dome (Herron and Langway, 1980).

*averaged over the period 15 – 30 ka

^aBouchet et al. (2023)

^bBaggenstos et al. (2018)

^cBuiron et al. (2011)

205 ^dCrotti et al. (2021)

^eMarkle and Steig (2022)

^fKindler et al. (2014)

^gSalamatin et al. (1994)

^hRitz et al. (1982)

210 ⁱWilhelms et al. (2007)

^jTarasov and Peltier (2003)

^kRöthlisberger et al. (2000b)

^lLambert et al. (2012)

^mFischer et al. (2007)

215 ⁿSchüpbach et al. (2014)

^oBigler (2004)

^pTraversi et al. (2004)

^qSpolaor et al. (2013)

^rAarons et al. (2017)

220 ^sDelmonte et al. (2004)

^tMarino et al. (2009)

^uDelmonte et al. (2010)

^vRuth et al. (2003)

3.2 N₂O analysis

225 3.2.1 Measurement of N₂O mixing ratio and isotopic composition

Prior to analysis, the samples were decontaminated by removing ~5 mm of the outer surface potentially affected by diffusion of modern air and chemical contaminants. Scraping was performed with a precleaned knife and wearing polyethylene gloves to minimize NO₃⁻ contamination (see Sect. 3.3).

The mixing ratio and bulk isotopic composition of N₂O were measured at the University of Bern, Switzerland, using the
230 method described in detail in Schmitt et al. (2014). Briefly, the air was extracted by melting the ice core samples with infrared light in a glass vessel under high vacuum. Water vapor was removed from the air sample with a cold trap, CO₂ was removed using Ascarite™, and N₂O, CH₄, and other trace gases were separated from the bulk air components (N₂, O₂, and Ar) using a cold trap filled with activated carbon. N₂O and CH₄ were then separated on a gas chromatography column, and N₂O was analyzed with an IsoPrime isotope ratio mass spectrometer (IRMS). The results were converted to international
235 isotope scales. Isotopic compositions are expressed in δ values where $\delta = \frac{R_{sample}}{R_{reference}} - 1$, with R referring to ¹⁵N/¹⁴N and ¹⁸O/¹⁶O ratios. The international references are N₂ in air for nitrogen and Vienna Standard Mean Ocean Water (VSMOW) for oxygen. The analytical precision of N₂O mixing ratio, $\delta^{15}N_{bulk}$ and $\delta^{18}O$ values is 3 ppb, 0.3 ‰ and 0.4 ‰ respectively.

We analyzed the position-specific isotopic composition of N₂O from ice core samples at Oregon State University following gas extraction, N₂O purification, IRMS analysis, and data reduction methods described in detail in Menking et al. (2025).
240 Briefly, the air was extracted by grating the ice core samples at -60°C to open the enclosed air bubbles (Bauska et al., 2016), i.e., using a so-called dry extraction device, without melting the ice. N₂O was purified and pre-concentrated using a series of progressively smaller-volume cold traps and gas chromatography separation of the trapped N₂O from residual CO₂. Isotopic measurements were performed using a Thermo Delta V Plus IRMS, where m/z 44, 45, and 46 and m/z 30 and 31 were monitored simultaneously for N₂O isotopes and N₂O fragment (NO) isotopes, respectively. The position-specific isotopic
245 composition of N₂O, i.e. $\delta^{15}N^{\alpha}$ (central-position N atom), $\delta^{15}N^{\beta}$ (terminal-position N atom) and site preference (SP), is defined as:

$$\delta^{15}N_{bulk} = \frac{\delta^{15}N^{\alpha} + \delta^{15}N^{\beta}}{2} \quad (1)$$

$$SP = \delta^{15}N^{\alpha} - \delta^{15}N^{\beta} \quad (2)$$

The $\delta^{15}N_{bulk}$, $\delta^{15}N^{\alpha}$ and $\delta^{18}O$ values were measured and the $\delta^{15}N^{\beta}$ and SP values were calculated using Eq. (1) and (2),
250 respectively. The precision of the measurement is 3 ppb for N₂O mixing ratio, ± 0.4 ‰ for $\delta^{15}N_{bulk}$, ± 0.6 ‰ for $\delta^{15}N^{\alpha}$, ± 0.8 ‰ for $\delta^{15}N^{\beta}$, ± 1.3 ‰ for SP , and ± 0.6 ‰ for $\delta^{18}O$.

3.2.2 Calculation of the mixing ratio and bulk isotopic composition of the in situ N₂O fraction

The N₂O mixing ratio and isotopic composition that we measured represent the total N₂O extracted from the ice cores, i.e., a mixture of atmospheric N₂O plus any in situ N₂O. For our study, we need to calculate the mixing ratio and isotopic composition of in situ N₂O only. To calculate the in situ values from the measured values, we used the following mass balance approach for each sample:

$$[N_2O]_{in\ situ} = [N_2O]_{meas} - [N_2O]_{atm} \quad (3)$$

$$\delta_{in\ situ} = \frac{\delta_{meas} * [N_2O]_{meas} - \delta_{atm} * [N_2O]_{atm}}{[N_2O]_{in\ situ}} \quad (4)$$

where $[N_2O]_{meas}$ is the N₂O mixing ratio measured in the ice core sample, $[N_2O]_{in\ situ}$ is the mixing ratio of in situ N₂O in that sample, and $[N_2O]_{atm}$ is the true atmospheric mixing ratio of N₂O at the gas age of the sample; the same terminology applies for delta values.

One issue with this mass balance approach is the need to know $[N_2O]_{atm}$ and δ_{atm} , given that most N₂O records are affected by in situ production. We used the N₂O record from the dust-poor TALDICE ice core, which is the best estimate of an atmospheric baseline. TALDICE is not significantly impacted by in situ N₂O production but is limited in time reconstruction (~150 ka). Low in situ contribution is supported by the observation that the TALDICE N₂O mixing ratios during the dust-rich LGM and MIS4 periods generally align with the NGRIP N₂O mixing ratios, after in situ N₂O outliers are removed from the NGRIP record (Fig. A1 in Appendix A). Previous studies identified the high-resolution NGRIP record as representative of atmospheric N₂O variations (Flückiger et al., 2004; Schilt et al., 2010a, 2013). As atmospheric N₂O mixing ratios are globally homogeneous while in situ N₂O production is ice-core dependent, the agreement between the TALDICE and NGRIP records supports the atmospheric nature of the TALDICE record. This is particularly significant because the NGRIP and TALDICE ice cores, drilled in Greenland and Antarctica, respectively, have very different chemical compositions that would lead to different in situ production features. To associate each sample with the atmospheric values matching its gas age, we applied a cubic smoothing spline interpolation with a 2000-year smoothing window to the measured TALDICE data, which has a temporal resolution of approximately 300 years over the LGM and 600 years over MIS4.

We acknowledge that there could be a small in situ production of N₂O in the TALDICE ice core, isotopically undetectable if the isotopic signature of in situ N₂O is close to the atmospheric one. We conducted a sensitivity study to evaluate the impact of this possibility (Fig. C1 and C2 in Appendix C). Assuming that N₂O production is proportional to dust content, we estimated the in situ N₂O mixing ratios by multiplying the calcium concentrations – used as a dust proxy – by a constant N₂O production factor. These in situ N₂O mixing ratios were subtracted from the TALDICE N₂O mixing ratios to obtain a corrected record that was used in the mass balance approach as an atmospheric baseline. This approach assumes a constant total air content (TAC) across all samples. While variations in TAC affect the conversion from Ca²⁺-based N₂O estimates (in ng.g⁻¹) to mixing ratios (ppb), this effect is relatively minor (within ±10 %) and does not dominate the overall uncertainty in our mass balance calculation. Our sensitivity study varied the production factor from 0 to 1 ppb-N₂O/ng.g⁻¹-Ca²⁺ (Fig. C1 and C2 in Appendix C). A production factor above 0.5 ppb-N₂O/ng.g⁻¹-Ca²⁺ is unlikely, as it would result in significantly

285 lower N₂O mixing ratios in TALDICE compared to NGRIP. Increasing production factors in TALDICE result in isotopic signatures of in situ N₂O in the other ice cores that are closer to atmospheric values. Importantly, assuming small production factors of 0.1 to 0.2 ppb-N₂O/ng.g⁻¹-Ca²⁺, which represents the upper range of expected values for TALDICE, results in only minimal changes in regression slopes when comparing the isotopic compositions of in situ N₂O and NO₃⁻. Thus, we conclude that the interpretations that follow remain valid even if the assumption that TALDICE represents the true atmospheric N₂O
290 baseline is not entirely fulfilled.

3.2.3 Calculation of the position-specific isotopic composition of the in situ N₂O fraction

We used the same mass balance approach as in Sect. 3.2.3 to calculate the in situ $\delta^{15}\text{N}^{\alpha}$ values in the Vostok and TG samples, with Eq. (3) and (4) in which the δ values are substituted with $\delta^{15}\text{N}^{\alpha}$. The in situ $\delta^{15}\text{N}^{\beta}$ values were calculated with Eq. (1) ($\delta^{15}\text{N}^{\beta}_{\text{in situ}} = 2 \delta^{15}\text{N}_{\text{bulk in situ}} - \delta^{15}\text{N}^{\alpha}_{\text{in situ}}$) and in situ SP values with Eq. (2) ($\text{SP}_{\text{in situ}} = \delta^{15}\text{N}^{\alpha}_{\text{in situ}} - \delta^{15}\text{N}^{\beta}_{\text{in situ}}$). One
295 problem with this approach is that the TALDICE record does not include position-specific information. As the atmospheric reference $\delta^{15}\text{N}^{\alpha}_{\text{atm}}$, we therefore used the average of the values measured by Menking et al. (2025) in samples from the TG ice core during the period 16 to 21 ka, as these values vary little over time during the last glacial period and because these TG samples were shown to be unaffected by in situ production by comparison with TALDICE and NGRIP.

3.2.4 Uncertainties

300 There are several sources of error in our approach, including measurement uncertainties in N₂O mixing ratios and isotopic composition, uncertainties in the TALDICE N₂O spline interpolation, uncertainties in gas-age estimates – which in turn impact the chronological alignment between ice cores – and potential small in situ production in TALDICE. We used a Monte Carlo method for error propagation by running 1000 simulations to determine the mixing ratio and isotopic composition of in situ N₂O. We varied the variables of Eq. (3) and (4) within their uncertainty ranges (Table B1 in Appendix
305 B). The derived uncertainties were calculated as the standard deviation of the results of the 1000 simulations.

As expected, the uncertainty increases for low in situ N₂O mixing ratios (see Eq. 4). To avoid too large uncertainties, we excluded samples with calculated in situ N₂O mixing ratios below 20 ppb. This threshold was chosen because a large part of the dataset exhibits in situ mixing ratios below 50 ppb. Applying a higher cutoff would remove a substantial number of samples and significantly limit our ability to investigate in situ N₂O production processes. For samples with in situ N₂O
310 mixing ratios close to 20 ppb, the propagated uncertainties are large (up to several tens of per mil). Nevertheless, these samples were included in this study because in situ N₂O production in ice, although small in absolute mixing ratio, can significantly alter the measured N₂O isotopic composition (Fig. 3a) and these isotopic deviations should be documented.

3.3 Measurement of NO_3^- concentration and isotopic composition

After wet extraction and dry extraction of N_2O at the University of Bern and Oregon State University, respectively, the
315 meltwater and melted ice chips from ice-core samples were used for nitrate isotope analysis. To avoid any chemical reaction
and hence production or consumption of NO_3^- , the collected samples were refrozen and stored at $-25\text{ }^\circ\text{C}$.


To assess the potential contamination impact of the gas extraction techniques on NO_3^- , we measured the concentration and
isotopic composition of NO_3^- in ultrapure ice samples spiked with a controlled amount of NO_3^- isotope standard after gas
extraction as described above. Additionally, we compared the concentration and isotopic composition of NO_3^- in duplicate
320 ice core samples, one analyzed directly and the other analyzed after gas extraction (Table D1 in Appendix D). These tests
show that the gas extraction techniques do not lead to measurable contamination, loss, or $\delta^{15}\text{N}$ isotopic fractionation of NO_3^-
in the collected samples. Although $\delta^{18}\text{O}(\text{NO}_3^-)$ values are slightly affected, the impact is negligible relative to their typical
variability observed in ice cores.

The method used for nitrate isotope analysis is described in detail in previous publications (Erbland et al., 2013; Kaiser et al.,
325 2007; Morin et al., 2009). Briefly, the NO_3^- concentration was measured by ion chromatography using a Dionex Integrion
system; NO_3^- in the samples was preconcentrated using an AG 1-X8 anion exchange resin in the chloride form. After the
sample was drained and the NO_3^- ions were quantitatively trapped onto the resin, NO_3^- was eluted from the resin with 6 mL
of 1M NaCl solution in three portions of 2 mL. NO_3^- was then converted to N_2O through bacterial denitrification, using a
strain of *Pseudomonas aureofaciens*. The bacteria were injected in 2 mL aliquots into 20 mL headspace vials. To remove air
330 and dissolved N_2O , the vials were purged for 3 h with pure helium. The concentrated NO_3^- samples were added to the vials
in volumes adjusted to obtain 100 nmol of NO_3^- , and were allowed to denitrify overnight. For isotope analysis, a continuous
He flow was used to transfer the produced N_2O from the headspace vial and carry it through a purification line. The N_2O
sample was passed through columns of perchlorate, Ascarite, and Supelco Purge Trap type F to remove water, CO_2 , and
VOCs, respectively. Following this purification step, the purified N_2O was decomposed to N_2 and O_2 on a gold catalyst kept
335 at $850\text{ }^\circ\text{C}$, and the isotopic compositions of the obtained N_2 and O_2 were measured with a Thermo Fischer MAT 253 IRMS.
The results were corrected for blank contribution and calibrated on international scales following the procedure described in
Erbland et al. (2013). Measurement errors are $\pm 0.8\text{ }‰$ for $\delta^{15}\text{N}$ and $\pm 1.0\text{ }‰$ for $\delta^{18}\text{O}$.

4 Results

4.1 Bulk isotopic signatures of total N_2O and in situ N_2O

340



345 **Figure 3. Nitrogen and oxygen isotopic composition of total N_2O measured in samples from different ice cores (a) and nitrogen and oxygen isotopic composition calculated for in situ N_2O in the same samples (b). The mixing ratio of in situ N_2O is represented by the size of the markers. The grey cross represents the average isotopic signature of the TALDICE samples and reflects the isotopic composition of atmospheric N_2O . In panel (b), one TALDICE sample shows in situ N_2O production associated with an exceptionally high dust peak (see Text). Note that the scale of the axes is different for panel (a) and (b).**

Figure 3a illustrates the measured isotopic signatures of total N_2O from various ice cores. The TALDICE data, representing atmospheric N_2O , clusters around a mean $\delta^{15}N_{\text{bulk}}$ value of $+11.3 \pm 0.7$ ‰ (1σ standard deviation) and a mean $\delta^{18}O$ value of $+45.7 \pm 0.9$ ‰. These data cover the last 140-kyr interval and their standard deviation is only about two times the measurement error, hence the detectable temporal variability of the isotopic signature of N_2O is small. In contrast, other ice cores show wider ranges, both in $\delta^{15}N_{\text{bulk}}$ and $\delta^{18}O$: the deviations from the atmospheric range and relationship with elevated N_2O mixing ratios indicate that these samples are substantially affected by in situ production.

355 Importantly, the magnitude and direction of deviations from the atmospheric signature are ice core specific. Because the in situ N_2O fraction shapes the observed isotopic deviations, we closely examine the isotopic signature of in situ N_2O in Fig. 3b, which presents the results of our mass balance calculation. Individual samples in the NGRIP ice core in Greenland show the highest in situ N_2O production, up to 380 ppb, while Antarctic ice cores like Vostok, EDML, EDC, and TG show in situ production in the order of 40 ppb, which corresponds to 20 % of the atmospheric N_2O mixing ratios during glacial periods.


360 The isotopic signature of in situ N_2O is highly variable, ranging from -37 ‰ to $+189$ ‰ for $\delta^{15}N_{\text{bulk}}$ and from -41 ‰ to $+100$ ‰ for $\delta^{18}O$. Not only does in situ N_2O have a distinct isotopic signature in each ice core, it also exhibits a variable isotopic signature within a given ice core. This suggests that the isotopic signature of the precursor(s) varies. Interestingly, all the ice cores except EDML show a negative correlation between $\delta^{15}N_{\text{bulk}}(N_2O_{\text{in situ}})$ and $\delta^{18}O(N_2O_{\text{in situ}})$ (Fig. 3b) while EDML shows a positive correlation.

365 While we used the TALDICE N₂O record to represent atmospheric N₂O, it is important to note that one sample within this ice core record is affected by in situ production (Fig. 3b). By comparison with the TALDICE spline, this particular sample contains 31 ppb of in situ N₂O with a $\delta^{15}\text{N}_{\text{bulk}}(\text{N}_2\text{O}_{\text{in situ}})$ value of -5 ‰ and a $\delta^{18}\text{O}(\text{N}_2\text{O}_{\text{in situ}})$ value of +67 ‰. Despite this anomaly, there are compelling reasons to use TALDICE as a reliable record for atmospheric values. Firstly, only one sample out of the 192 measured was found to be affected by in situ production, indicating that such occurrences are rare. Indeed, this sample is characterized by an exceptionally high dust content of 188 ng.g⁻¹ of Ca²⁺, which is 280 % higher than typical dust peaks observed in the TALDICE ice core during the LGM. Secondly, the amount of in situ N₂O in the affected sample is relatively small compared to this high dust content: it corresponds to a production factor of 0.16 ppb of N₂O per ng.g⁻¹ of Ca²⁺, which has little influence on our results, as shown by our sensitivity study (Fig. C2 in Appendix C).

370 We stress that the mass balance approach used to calculate the isotopic signature of in situ N₂O leads to substantial uncertainties due to error propagation from atmospheric N₂O estimates and analytical precision. These uncertainties are not shown in Fig. 3 for readability but are reported in Figs. 4 – 7 and are explicitly taken into account in the following comparisons between $\delta^{15}\text{N}(\text{N}_2\text{O}_{\text{in situ}})$ and $\delta^{15}\text{N}(\text{NO}_3^-)$, including the regression analyses.

4.2 Comparison of isotopic compositions of NO₃⁻ and in situ N₂O

In this section, we compare the measured isotopic composition of NO₃⁻ and the calculated isotopic composition of in situ N₂O to test our hypothesis that NO₃⁻ is a precursor for in situ produced N₂O. The analyses come from the same ice samples: N₂O was measured in the extracted air and subsequently used to calculate the in situ N₂O isotopic signature, while NO₃⁻ was measured in the sample meltwater collected after air extraction. To compare the data, we used the regression method by York et al. (2004) that accounts for errors in both the x and y variables and their potential correlation.



385 **Figure 4. Relation between $\delta^{15}\text{N}_{\text{bulk}}(\text{N}_2\text{O}_{\text{in situ}})$ and $\delta^{15}\text{N}(\text{NO}_3^-)$. The dashed line represents the linear regression.**

Figure 4 shows the bulk nitrogen isotopic composition of in situ N_2O and the nitrogen isotopic composition of NO_3^- measured in the EDML, EDC, TG and Vostok ice cores. There is a clear positive correlation between $\delta^{15}N_{\text{bulk}}(N_2O_{\text{in situ}})$ and $\delta^{15}N(NO_3^-)$ with a R^2 value of 0.86 which points to nitrate being a key precursor for the in situ produced N_2O . However, the slope is significantly different from 1, and in fact close to one half (0.54 ± 0.02), i.e., roughly one of the two N atoms of the in situ produced N_2O originates from NO_3^- .




Figure 5. Relation between $\delta^{18}O(N_2O_{\text{in situ}})$ and $\delta^{18}O(NO_3^-)$.

395

Figure 5 shows the comparison between the $\delta^{18}O$ values of NO_3^- and in situ N_2O . There is no correlation between these two variables across or within sites, and no general trend towards ^{18}O enrichment or depletion of in situ N_2O relative to NO_3^- . At Vostok and EDC, $\delta^{18}O(NO_3^-)$ values are around $+45 \pm 6$ ‰ (1σ), while $\delta^{18}O(N_2O_{\text{in situ}})$ values are approximately $+31 \pm 17$ ‰. At TG, $\delta^{18}O(NO_3^-)$ values are higher at $+68 \pm 4$ ‰, and $\delta^{18}O(N_2O_{\text{in situ}})$ values reach $+93 \pm 11$ ‰. At EDML, $\delta^{18}O(NO_3^-)$ values are the highest at $+87 \pm 6$ ‰, whereas $\delta^{18}O(N_2O_{\text{in situ}})$ values are the lowest, at $+17 \pm 11$ ‰.

The very high $\delta^{18}O(N_2O_{\text{in situ}})$ values observed at TG may appear anomalous compared to other sites, but we are confident that they do not result from a bias in the measurements or calculation. Even without applying the mass balance calculation, the measured N_2O at TG already shows $\delta^{18}O$ values higher than the atmospheric signature in samples affected by in situ production (Fig. 3a). This indicates that the in situ N_2O at TG is indeed enriched in $\delta^{18}O$. A similar enrichment is observed in the NGRIP ice core. Importantly, these elevated $\delta^{18}O$ values were measured using two different extraction methods and two different IRMS instruments at the University of Bern and Oregon State University, which strengthens our confidence that the signal is robust and not an artefact of a specific analytical setup.


405

In summary, despite the clear correlation between $\delta^{15}\text{N}_{\text{bulk}}(\text{N}_2\text{O}_{\text{in situ}})$ and $\delta^{15}\text{N}(\text{NO}_3^-)$ values suggesting NO_3^- as a precursor, the incomplete transfer of both its nitrogen and oxygen isotopic compositions to in situ N_2O indicates that NO_3^- alone does not fully account for the nitrogen and oxygen sources of in situ N_2O . To gain deeper insights into the reaction mechanisms, we turn our focus to the position-specific isotopic composition of in situ N_2O .

4.3 Site preference of ^{15}N in in situ N_2O

Figure 6 shows clear correlations of $\delta^{15}\text{N}^\alpha$, $\delta^{15}\text{N}^\beta$, and SP values versus $\delta^{15}\text{N}_{\text{bulk}}$ of total N_2O (a, c) and in-situ N_2O (b, d) measured in the TG and Vostok.

415



420

Figure 6. Position-specific isotopic composition measured for total (atmospheric + in situ) N_2O and calculated for in situ N_2O in the Vostok and TG ice cores. The panels show the relation between $\delta^{15}\text{N}^\alpha$, $\delta^{15}\text{N}^\beta$ and $\delta^{15}\text{N}_{\text{bulk}}$ of total N_2O (atmospheric + in situ) (a) and in situ N_2O (b), and between site preference (SP) and $\delta^{15}\text{N}_{\text{bulk}}$ of total N_2O (atmospheric + in situ) (c) and in situ N_2O (d). The size of the markers on panel (b) and (d) are scaled to the amount of in situ N_2O in the sample. In panel (b), TG samples are in the

bottom-left corner. The crosses represent the unaffected atmospheric values measured by Menking et al. (2025) in dust-poor Taylor Glacier samples from the period 16 to 21 ka.

In situ N_2O at Vostok exhibits high $\delta^{15}N^{\alpha}$ values from +95 ‰ to +152 ‰ and much lower $\delta^{15}N^{\beta}$ values from +13 ‰ to +43 ‰. In contrast, in situ N_2O at TG has low $\delta^{15}N^{\alpha}$ values from -50 ‰ to -39 ‰ and low $\delta^{15}N^{\beta}$ values from -34 ‰ to -18 ‰, with $\delta^{15}N^{\alpha}$ being lower than the $\delta^{15}N^{\beta}$ values. The Vostok and TG samples exhibit very different $SP(N_2O_{in\ situ})$ values; +59 ‰ to +122 ‰ and -22 ‰ to -8 ‰, respectively. These values differ significantly from those of atmospheric N_2O which range in SP from +17 ‰ to +24 ‰ across last glacial termination, as reconstructed from dust-poor sections of TG ice (Menking et al., 2025). The $SP(N_2O_{in\ situ})$ values at Vostok are very high and variable, and are outside the range of typical reported values from -11 ‰ to +37 ‰ (Toyoda et al., 2017; Zhu-Barker et al., 2015). Note that the $SP(N_2O_{in\ situ})$ values at Vostok seem to be correlated to the $\delta^{15}N_{bulk}(N_2O_{in\ situ})$ values with a slope of 1.8 ± 0.7 , whereas SP values in field and culture studies have been shown to be generally independent of $\delta^{15}N_{bulk}$ (Frame and Casciotti, 2010; Sutka et al., 2003, 2006; Toyoda et al., 2005). For TG, there is too little data to identify a potential correlation.

Interestingly, the $SP(N_2O_{in\ situ})$ values at TG are negative. Although the low amounts of in situ N_2O at TG result in large uncertainties in the calculated in situ signatures, the low SP signal is already visible in the measured (total) N_2O data (Fig. 6c), independent of any mass-balance calculation. While Vostok samples affected by in situ production show total SP values higher than the atmospheric signature, TG samples show the opposite pattern, with total SP values lower than the atmospheric signature. This opposite deviation indicates that the low SP values observed at TG reflect a real feature of in situ N_2O production rather than an artefact of the calculation.

440




Figure 7. Relation between $\delta^{15}N^{\alpha}(N_2O_{in\ situ})$ and $\delta^{15}N(NO_3^-)$ (a) and between $\delta^{15}N^{\beta}(N_2O_{in\ situ})$ and $\delta^{15}N(NO_3^-)$ (b).

445 In Fig. 7, the $\delta^{15}\text{N}^{\alpha}$ and $\delta^{15}\text{N}^{\beta}$ values of in situ N_2O are compared with $\delta^{15}\text{N}$ values of NO_3^- measured in the TG and Vostok ice cores on the same samples. When considering both TG and Vostok samples, $\delta^{15}\text{N}^{\alpha}(\text{N}_2\text{O}_{\text{in situ}})$ shows a strong positive correlation with $\delta^{15}\text{N}(\text{NO}_3^-)$ (slope = 1.0 ± 0.1 , $R^2 = 0.97$). This relationship also holds when considering the Vostok data alone, indicating that the correlation is robust even within a single ice core. In contrast, $\delta^{15}\text{N}^{\beta}(\text{N}_2\text{O}_{\text{in situ}})$ does not show a statistically significant correlation with $\delta^{15}\text{N}(\text{NO}_3^-)$ for either site.

450 5 Discussion

5.1 Production of hybrid N_2O

The strong correlation between $\delta^{15}\text{N}_{\text{bulk}}(\text{N}_2\text{O}_{\text{in situ}})$ and $\delta^{15}\text{N}(\text{NO}_3^-)$ indicates that NO_3^- is involved in the in situ N_2O formation. However, the slope is (0.54 ± 0.02) (Fig. 4), whereas a slope of 1 would be expected if both nitrogen atoms in N_2O originate from NO_3^- in the ice. Therefore, our results suggest that in situ N_2O is not derived from NO_3^- alone, and does
455 not originate from a single precursor (we use the term “single-precursor N_2O ” hereafter, see Fig. 9). The strong correlation of $\delta^{15}\text{N}^{\alpha}(\text{N}_2\text{O}_{\text{in situ}})$ and $\delta^{15}\text{N}(\text{NO}_3^-)$ with a slope of 1.0 ± 0.1 (Fig. 7a), implies that the central N atom in N_2O (N^{α}), originates exclusively from NO_3^- archived in the ice. At the same time, the $\delta^{15}\text{N}^{\beta}(\text{N}_2\text{O}_{\text{in situ}})$ values are not correlated to $\delta^{15}\text{N}(\text{NO}_3^-)$, indicating that the terminal N atom (N^{β}) originates from a different nitrogen source. The slope of the linear regression between $\text{SP}(\text{N}_2\text{O}_{\text{in situ}})$ and $\delta^{15}\text{N}_{\text{bulk}}(\text{N}_2\text{O}_{\text{in situ}})$ (1.8 ± 0.7) is consistent, within uncertainty, with transfer of a variable N
460 isotopic composition from NO_3^- to N^{α} and transfer of a constant N isotopic composition from another precursor to N^{β} . Indeed, by substitution of equations Eq. (1) and (2), one would expect $\text{SP} = 2 * \delta^{15}\text{N}_{\text{bulk}} - 2 * \delta^{15}\text{N}^{\beta}$, where $\delta^{15}\text{N}^{\beta}$ is relatively constant within each ice core, although its specific value differs between cores (Fig. 7b). Our results suggest that the N_2O produced in situ is thus “hybrid N_2O ”, defined by its two nitrogen atoms originating from two distinct nitrogen precursors, one of them being NO_3^- in ice (Fig. 9).

465 To assess whether microbial N_2O reduction to N_2 could provide an alternative explanation for the observed isotopic patterns of in situ N_2O – such as high SP values correlated with $\delta^{15}\text{N}_{\text{bulk}}$ or high $\delta^{18}\text{O}$ values – we examined the relationship between SP and $\delta^{18}\text{O}$ in our samples (Fig. 8). Previous studies showed that N_2O reduction produces a characteristic increase in both SP and $\delta^{18}\text{O}$ in the residual N_2O , with data falling along a “reduction line” defined by the ratio of the fractionation factors (median slope ≈ 0.36 ; Lewicka-Szczebak et al., 2015; Yu et al., 2020). In contrast, in both the Vostok and Taylor Glacier ice
470 cores, SP increases while $\delta^{18}\text{O}$ decreases, and the data clearly do not fall along the expected reduction line (Fig. 8). This dual isotope plot for in situ N_2O is therefore incompatible with N_2O reduction. We note that published reduction lines were derived from studies conducted at warmer temperatures than those in ice cores (Yu et al., 2020). Very low temperatures could modify the fractionation factors; however, colder conditions would be expected to increase the fractionation for both SP and $\delta^{18}\text{O}$, still resulting in a positive slope. The mismatch between our data and the reduction line therefore remains
475 regardless of potential temperature effects. A second line of evidence comes from the correlation between $\delta^{15}\text{N}(\text{NO}_3^-)$ and $\delta^{15}\text{N}^{\alpha}(\text{N}_2\text{O}_{\text{in situ}})$, which shows a slope of ~ 1 . Because N_2O reduction preferentially breaks N-O bonds of light isotopes, the

residual N₂O becomes enriched in ¹⁵N at the α position, resulting in a steeper slope than observed here. Taken together, these observations show that microbial N₂O reduction cannot explain the isotopic signatures in the ice cores, and that N₂O consumption does not occur in the ice. The isotopic signature of in situ N₂O is therefore explained by hybrid N₂O production.

480




Figure 8. Dual isotope plot of $\delta^{18}\text{O}$ and site preference (SP) values of in situ N₂O. Theoretical N₂O reduction lines (thin black lines) are shown for comparison, with intercepts calculated assuming that samples with the lowest SP values are unaffected by N₂O reduction. The slope of the reduction line corresponds to the median of published fractionation ratios for SP versus $\delta^{18}\text{O}$ during N₂O reduction, as reported by Yu et al. (2020).

485

5.1.1 Potential reaction pathways

A large number of studies have reported the production of hybrid N₂O through various reaction pathways, including N-nitrosation reactions (Spott et al., 2011) and the decomposition of ammonium nitrate (NH₄NO₃) (Rubasinghege et al., 2011). N-nitrosation reactions generally involve the replacement of a hydrogen atom on a nucleophilic precursor by a nitroso (-N=O) group (Spott et al., 2011). Production of N₂O occurs during N-nitrosation under acidic conditions, when the nitrosating agent nitrite (NO₂⁻) reacts with the nucleophile hydroxylamine (NH₂OH) to form nitroxyl (HNO). Two HNO molecules then dimerize to produce nitrous oxide (N₂O) and water (Spott et al., 2011). This reaction can be abiotic or mediated by bacteria, archaea, or fungi. Other nucleophilic species can react with NO₂⁻ to produce N₂O, such as azide (N₃⁻), ammonium (NH₄⁺), hydrazine (N₂H₄), or salicylhydroxamic acid (C₇H₇NO₃) (Spott et al., 2011).

490

495

Another pathway for hybrid N₂O production is the decomposition of ammonium nitrate (NH₄NO₃), either by thermal decomposition or by a light-initiated reaction involving the photoreduction of NO₃⁻ to NO₂ coupled with the oxidation of NH₄⁺ to NH₂, which react with each other to produce N₂O (Rubasinghege et al., 2011). However, several arguments challenge the feasibility of the NH₄NO₃ decomposition in ice. Firstly, the ice environment lacks sufficient heat for thermal decomposition, and light penetration is minimal within the deep firn and ice, raising questions about the availability of

500 sufficient energy for these reactions. The high activation energy of the decomposition would make the reaction very slow at the ice temperature. Additionally, if N_2O were produced from NH_4NO_3 , the oxygen atom in N_2O would originate from NO_3^- and inherit its isotopic composition, yet the $\delta^{18}\text{O}$ values of in situ N_2O do not correlate with those of NO_3^- (Fig. 5). Finally, NO_3^- and NH_4^+ are present in dust-poor polar ice as well, where no in situ N_2O production is observed, and similarly, there is not necessarily in situ N_2O production everywhere in dust-rich ice from Greenland despite sufficient NO_3^- and NH_4^+ . In our
505 view, an N-nitrosation reaction is more likely than a NH_4NO_3 decomposition reaction for N_2O production in ice.

5.1.2 Source of central nitrogen atom N^{α}

In the case of in situ N_2O produced by N-nitrosation, the central nitrogen atom (N^{α}) may originate from NO_2^- after reduction of NO_3^- . However, it is unlikely that this NO_2^- derives directly from NO_3^- photolysis in the near-surface snowpack. NO_3^- photolysis produces both gaseous NO_2 and NO_2^- ion, with NO_2^- accounting for $\sim 10\%$ of the photolysis products (Meusinger
510 et al., 2014; Warneck and Wurzinger, 1988). Photolysis occurs at the snow surface only, where sunlight penetrates and where air exchange with the atmosphere is still active. Any N_2O produced through photolysis-driven pathways in this zone would therefore be largely released to the atmosphere and not preserved in the ice core record. In addition, photolysis enriches the remaining NO_3^- in ^{15}N while producing NO_2^- that is depleted in ^{15}N , with its isotopic composition depending on the extent of photolysis. If photolysis-derived NO_2^- were a precursor of in situ N_2O , the $\delta^{15}\text{N}^{\alpha}$ signature of in situ N_2O would
515 reflect both the initial $\delta^{15}\text{N}(\text{NO}_3^-)$ and the extent of photolysis, and would not be directly proportional to the $\delta^{15}\text{N}$ values of NO_3^- archived in the ice. Instead, our data show a strong proportionality between $\delta^{15}\text{N}(\text{NO}_3^-)$ and $\delta^{15}\text{N}(\text{N}_2\text{O}_{\text{in situ}})$, indicating that the NO_2^- precursor must form deeper in the ice from archived NO_3^- that already carries its final, post-photolysis isotopic signature, rather than from photolysis-derived NO_2^- produced in the surface snowpack. Therefore, the reduction of NO_3^- to NO_2^- occurring within the ice requires a reducing agent. Such a reducing agent is likely associated with mineral dust, which
520 is consistent with the observation that in situ N_2O production occurs only in dust-rich ice. Iron II (Fe^{2+}) is likely involved in this reduction step (see Sect. 5.2).

5.1.3. Source of terminal nitrogen atom N^{β}

Our work does not yet allow us to identify the nucleophilic precursor for N^{β} , especially because the presence and concentrations of most N-bearing compounds in ice have not yet been investigated. Nevertheless, NH_4^+ represents a potential
525 candidate for supplying N^{β} in the case of N-nitrosation reactions. Atmospheric NH_4^+ exhibits $\delta^{15}\text{N}$ values ranging from -20‰ to $+25\text{‰}$ (Chen et al., 2022), rather similar to the $\delta^{15}\text{N}^{\beta}$ values of in situ N_2O in Vostok and TG with average $\delta^{15}\text{N}^{\beta}$ values of -27‰ and $+24\text{‰}$, respectively. These site differences could be due to post-depositional processes altering the isotopic composition of NH_4^+ differently at different drilling sites. The nitrogen isotopic composition of NH_4^+ has been measured only in one alpine glacier ice core, with $\delta^{15}\text{N}$ values ranging from -15‰ to $+5\text{‰}$ for the years 2013-2017
530 (Lamothe et al., 2023). To further investigate this hypothesis, we suggest measuring the nitrogen isotopic composition of NH_4^+ in Antarctic ice cores, specifically from Vostok and Taylor Glacier, to compare with the $\delta^{15}\text{N}^{\beta}$ values of in situ N_2O .

5.2 Limiting factor of the reaction

One key observation is that the elevated N₂O mixing ratios during glacial periods are of similar magnitude over the last 800 kyr in the EDC ice core with no increase with depth/age (Schilt et al., 2010b). This observation suggests that N₂O in situ production does not increase further over time and is essentially finished at the LGM (20 to 26.5 ka). Therefore, we hypothesize a mechanism that is limited by the concentration or transport/diffusion of a reactant. The limiting factor cannot be NO₃⁻ as the concentrations remain high over time. Thus, it could be the precursor of the terminal N atom (N^β) or the agent reducing NO₃⁻ to NO₂⁻.

Indeed, because N-nitrosation reactions typically involve NO₂⁻ rather than NO₃⁻ (Spott et al., 2011), a prior conversion of NO₃⁻ to NO₂⁻ is required. While this step can be carried out by denitrifying organisms, the extremely low temperatures and acidic conditions in Antarctic ice make an abiotic process more likely (see Sect. 5.5). NO₃⁻ can be abiotically reduced to NO₂⁻ by redox-active metal ions such as Fe²⁺ or Mn²⁺ (Zhu-Barker et al., 2015). This hypothesis is supported by the presence of Fe²⁺ in ice cores (Spolaor et al., 2012, 2013). Baccolo et al. (2021) found decreasing Fe²⁺ concentrations with depth in the TALDICE core, and below 1500 m only Fe³⁺ is present in the form of jarosite (KFe³⁺₃(SO₄)₂(OH)₆), a mineral that forms through chemical weathering of aeolian dust under acidic conditions in deep ice. These findings indicate that Fe²⁺ undergoes slow post-depositional oxidation in the ice. One possible explanation is that NO₃⁻ oxidizes dust-derived Fe²⁺, which could explain the observed link between dust content and in situ N₂O production. In this scenario, Fe²⁺ is the limiting factor for N₂O production, as it is progressively consumed during the conversion from NO₃⁻ to NO₂⁻.

5.3 Site preference constraints on the N₂O production mechanism

Table 2. Comparison of the site preference (SP) range measured for hybrid N₂O in a previous study with SP ranges calculated for in situ N₂O in the Vostok and TG ice cores.

Reaction pathway	SP range (‰)	Suspected intermediate species	Reference
Abiotic oxidation of NH ₂ OH by NO ₂ ⁻ (production of hybrid N ₂ O)	[+34 ; +35]	Symmetric intermediate: cis-hyponitrite (⁻ ONNO ⁻)	(Heil et al., 2014)
N ₂ O in situ production at Vostok	[+57.2 ; +187.4]	Asymmetric intermediate (unknown)	This study
N ₂ O in situ production at TG	[-17.3 ; -6.9]	Asymmetric intermediate (unknown)	This study

Since many reaction pathways have specific SP values, the SP signature is commonly used to attribute N₂O production pathway to a specific pathway. High SP values ranging from +22 ‰ to +37 ‰ are reported for nitrification (Toyoda et al., 2002, 2017), fungal denitrification (Toyoda et al., 2017), and abiotic reactions such as NO₂⁻ reduction and NH₂OH oxidation (Heil et al., 2015; Toyoda et al., 2005). Several studies suggested that the similarity of the SP signatures for these distinct

formation pathways is attributable to a common intermediate species – hyponitrous acid HONNOH_{cis} (Heil et al., 2015; Toyoda et al., 2002, 2005, 2017). During decomposition, this intermediate preferentially breaks at the ^{14}N -O bond over the ^{15}N -O bond, enriching the central N atom (N^α) in ^{15}N and explaining high SP values (Toyoda et al., 2002) (Fig. 9a). In contrast, bacterial and nitrifier denitrification produce N_2O with low SP values from -11 ‰ to 0 ‰ (Toyoda et al., 2017), which are thought to derive from a different intermediate. Modeling work suggested a trans-hyponitrous structure that decomposes preferentially into $^{15}\text{N}^{14}\text{NO}$ due to kinetic isotope effects, thus explaining the low SP values of these formation pathways (Fehling, 2012) (Fig. 9a). Overall, in these cases, SP values are therefore determined by the decomposition step of the last intermediate into N_2O and are independent of the $\delta^{15}\text{N}$ values of the precursor (Fig. 9).

For in situ N_2O , however, the SP values for the Vostok and TG ice cores do not fall into either typical low or high SP range and are highly variable (Table 2). The constant-SP pathways discussed above produce single-precursor N_2O , but in the case of hybrid N_2O production, one would expect a priori that the SP would no longer be a unique mechanism-dependent signature. Instead, for hybrid reactions the SP should be influenced by the difference between the $\delta^{15}\text{N}$ signatures of the two precursors, if the central N atom (N^α) is derived from one precursor and the terminal N atom (N^β) from the other. However, Heil et al. (2014) demonstrated that hybrid N_2O from the nitrosation of NH_2OH by NO_2^- is also characterized by a constant SP value of +34 ‰ (Table 2) regardless of the precursor signatures. The authors concluded that the NH_2OH -nitrosation mechanism also involves the symmetric hyponitrous intermediate (Fig. 9b). In fact, since this intermediate is symmetrical, the N atoms from each precursor can either take the α -position or the β -position in the final N_2O molecule; the difference in their $\delta^{15}\text{N}$ values then has no influence on the SP value (Fig. 9b).

Because of the variability of the $\text{SP}(\text{N}_2\text{O}_{in situ})$ values and their dependence on the $\delta^{15}\text{N}(\text{NO}_3^-)$ signature, we therefore deduce that the production mechanism of in situ N_2O does not involve the hyponitrous symmetrical intermediate. In Fig. 9c, we propose a new reaction mechanism that involves an asymmetric intermediate, resulting in a hybrid N_2O molecule in which the N atoms at the α and β positions are derived from a specific precursor. N^α and N^β retain the two individual signatures of the precursors rather than losing them as discussed above. The high variability of SP values is then attributable to the variability of $\delta^{15}\text{N}^\alpha$ values (Fig. 6), which originates from the wide range of $\delta^{15}\text{N}(\text{NO}_3^-)$ values (Fig. 7).

We note that all published SP datasets used for comparison were obtained at ambient temperatures, whereas in situ N_2O production in Antarctic ice occurs at very low temperatures. Such low temperatures imply extremely slow reaction rates, which may alter the magnitude of isotope fractionation and make direct comparison of absolute SP values with ambient-temperature experiments difficult. To our knowledge, no SP measurements exist under such cold conditions. However, our interpretation does not rely on comparing absolute SP values. Instead, we focus on whether SP is constant or variable with respect to the $\delta^{15}\text{N}$ signature of the precursor; this property should be independent of the absolute magnitude of isotope fractionation. Although lower temperatures may increase kinetic isotope effects and shift absolute SP values, they do not change whether SP remains constant (as in reactions involving symmetrical intermediates) or varies with the precursor isotopic composition (as expected when an asymmetrical intermediate forms). Thus, the comparison of the mechanisms remains valid even without low-temperature experimental data from previous studies.

Several mechanisms could potentially explain why the intermediate of in situ N_2O production is asymmetric, even though its exact chemical structure remains unknown. Firstly, the precursor of the β -position N atom could be different from NH_2OH . Although most studies on hybrid N_2O production report a reaction between NH_2OH and NO_2^- (Frame et al., 2017; Spott et al., 2011; Stieglmeier et al., 2014; Terada et al., 2017), other nucleophilic precursors have been reported as precursors of hybrid N_2O . Hydrazine (N_2H_4), for example, forms the asymmetrical intermediate $HO-N=N-NH_2$ (Perron et al., 1976). A second possibility is that very low temperatures modify the structure or stability of the intermediate normally formed from NH_2OH and NO_2^- at ambient temperature conditions, favoring an asymmetrical species and thereby generating the observed dependence of SP on the precursor $\delta^{15}N$ values.

The same hybrid production mechanism proposed for in situ N_2O production in Antarctic ice may occur at Don Juan Pond (DJP), Antarctica. Like in ice cores, N_2O formation at DJP reported by Samarkin et al. (2010) occurs under very low temperatures and involves NO_3^- and especially NO_2^- , with higher production rates for NO_2^- as precursor. The authors demonstrated that N_2O extracted from DJP soil is abiotically produced. Moreover, they measured very low and highly variable SP values at DJP, down to -45 ‰, which also fall outside the typical SP range. This variability in SP values and similar environmental conditions support the possibility of a shared, likely abiotic, mechanism of hybrid N_2O production in Antarctic ice and at DJP.




Figure 9. Production mechanisms for single-precursor N_2O (a) and hybrid N_2O (b), (c). Two mechanisms for hybrid N_2O production are proposed: panel (b) explains constant SP values, while panel (c), which explains the observed SP dependence on the $\delta^{15}N$ values of the precursor, is more likely to represent the in situ production pathway. (b) Mechanism with a symmetric intermediate, here the *cis*-hyponitrous acid $HONNOH$. Depending on which N-O bond of the symmetric intermediate breaks, the NO_2 -derived and NH_2OH -derived N atoms can each end up either in the central position (α) or in the terminal position (β) in N_2O . The percentage of molecules 1 and 2, and consequently the site preference, only reflects the preferential cleavage of the ^{14}N -O bond over the ^{15}N -O bond during the last step of N_2O formation. Therefore, hybrid N_2O exhibits a consistent site preference that is independent of the $\delta^{15}N$ values of the precursors (Heil et al., 2014). In this case the O atom is derived from one precursor or the

620 other. (c) Mechanism with an asymmetric intermediate. In this case, the O atom and the central N atom (N^{α}) are always derived from NO_2^- and the terminal N atom (N^{β}) from NH_2OH . The two N atoms retain the $\delta^{15}\text{N}$ signatures of their individual precursors. The site preference is therefore variable, as it depends on the difference between the $\delta^{15}\text{N}$ values of the two precursors. This mechanism matches our observations but to our knowledge has not yet been reported in the literature. Note that the precursors and intermediates shown in this figure are only examples used to illustrate our point about SP values. It is one hypothesis among others for the mechanism of in situ N_2O production as the nitrogen source of the terminal nitrogen N^{β} in N_2O (in yellow) remains unknown. During in situ production, Fe^{2+} carried by dust may reduce NO_3^- to NO_2^- and act as the limiting factor once fully oxidized to Fe^{3+} .

5.4 Source of oxygen in in situ N_2O

625 The $\delta^{18}\text{O}$ signature of in situ N_2O does not reflect the $\delta^{18}\text{O}$ of NO_3^- , indicating that additional processes modify the oxygen isotopic composition during or prior to N_2O formation; below, we therefore discuss possible mechanisms that could explain this decoupling, while noting that the dominant process remains uncertain.

Considering an N-nitrosation pathway, the observed $\delta^{18}\text{O}(\text{N}_2\text{O}_{\text{in situ}})$ values might be explained by the combined processes of NO_3^- reduction and NO_2^- isotopic equilibration with H_2O (Casciotti et al., 2007). Since N-nitrosation typically involves NO_2^- rather than NO_3^- , it is likely that NO_3^- is first reduced to NO_2^- . Several studies have shown that NO_2^- is prone to exchange its O atoms with the O atoms of water molecules in aqueous solution (Bunton et al., 1959). The water from Antarctic ice is strongly depleted in ^{18}O , with $\delta^{18}\text{O}(\text{H}_2\text{O})$ values of approximately -62 ‰ at Vostok (Lorius et al., 1985), -56 ‰ at EDC (Landais and Stenni, 2021), -42 ‰ at EDML (EPICA Community Members, 2010) and -42 ‰ as well at TG (Baggenstos et al., 2018). Thus, incorporation of O atoms from H_2O into N_2O through exchange with NO_2^- could explain the ^{18}O depletion
630 observed in in situ N_2O compared to NO_3^- in the EDML, EDC, and Vostok ice cores (Fig. 5). However, the $\delta^{18}\text{O}(\text{N}_2\text{O}_{\text{in situ}})$ values are not correlated to $\delta^{18}\text{O}(\text{H}_2\text{O})$ values and the difference between $\delta^{18}\text{O}(\text{N}_2\text{O}_{\text{in situ}})$ and $\delta^{18}\text{O}(\text{NO}_3^-)$ is significantly larger at EDML than at EDC and Vostok. This may indicate an incomplete exchange of O atoms between NO_2^- and H_2O , with a larger fraction of exchanged O atoms at EDML than EDC and Vostok.

In contrast, $\delta^{18}\text{O}(\text{N}_2\text{O}_{\text{in situ}})$ values at TG are unexpectedly higher than $\delta^{18}\text{O}(\text{NO}_3^-)$, suggesting that oxygen exchange with
640 water alone cannot explain the $\delta^{18}\text{O}(\text{N}_2\text{O}_{\text{in situ}})$ values. In this case, the observed enrichment in ^{18}O compared to NO_3^- might be due to the “branching effect” associated with NO_3^- reduction. This effect results from the incomplete incorporation of the oxygen pool from NO_3^- to N_2O (Casciotti et al., 2007). Indeed, only two out of three O atoms from NO_3^- are transferred into NO_2^- , then one out of two from NO_2^- to N_2O . Since ^{16}O atoms are preferentially lost, NO_2^- and N_2O become enriched in ^{18}O . For the abiotic reduction of NO_2^- to N_2O by Fe^{2+} , the branching effect results in positive isotope effect of about +30 ‰ at 25
645 °C (Buchwald et al., 2016) which is likely even larger at very low temperatures. Although the same branching effect probably occurs at EDC, Vostok, and EDML, the fraction of exchanged O atoms might be smaller at TG, making the branching effect relatively more pronounced in the final $\delta^{18}\text{O}(\text{N}_2\text{O}_{\text{in situ}})$ values.

The reason why the fraction of exchanged O atoms would vary among ice cores is not fully understood. It likely depends on the relative rates of oxygen isotope equilibration and the N-nitrosation reaction, both of which are temperature- and pH-
650 dependent. Casciotti et al. (2007) showed that oxygen exchange is faster under acidic conditions, and Su et al. (2019) reported that low pH also accelerates abiotic nitrosation reactions. However, measurement constraints of pH remain limited.

Although glacial Antarctic ice is generally acidic and Greenland ice is typically more alkaline, pH measurements are not available for the specific samples investigated here, making it difficult to interpret the differences in $\delta^{18}\text{O}(\text{N}_2\text{O}_{\text{in situ}})$ among Antarctic sites in terms of pH effects. Moreover, even if pH were measured in the liquid phase (meltwater), it would be difficult to directly infer the pH of the solid ice matrix, as pH is formally defined only for liquid solutions. In addition, the relevant reactions likely occur at the surfaces of dust particles (where the reduction of NO_3^- to NO_2^- may take place) rather than in the bulk ice, and the chemical conditions in these micro-environments may differ from those of the surrounding ice. To conclude, the combination of different ice chemical compositions and environmental conditions during the reaction could influence the fraction of O atoms exchanged between NO_2^- and H_2O . While the relative contributions of oxygen exchange and branching effect to the $\delta^{18}\text{O}(\text{N}_2\text{O}_{\text{in situ}})$ values remain unclear, overall, these processes may mask the transfer of the original $\delta^{18}\text{O}(\text{NO}_3^-)$ signature into in situ N_2O .

5.5 Likelihood of an abiotic production of N_2O

In situ N_2O production in polar ice has often been attributed to microbial activity. Sowers (2001) reported two N_2O maxima in the Vostok ice core at around 150 ka that coincided with elevated bacterial counts. Similarly, Rohde et al. (2008) found that in the GISP2 Greenland ice core, most of the samples affected by in situ production were associated with large cell counts.

However, several observations challenge the hypothesis of a microbial in situ production. First, the correlation between microbial counts and N_2O maxima may not be causal, as the dust content could be a confounding variable: mineral particles are the main carriers of microbial cells to the ice sheet (Miteva, 2008; Miteva et al., 2016), so both microbial concentrations and in situ N_2O are positively correlated with dust content in Antarctic ice cores. Second, because the low temperatures affect microbial metabolism and thus limit the reaction, one would expect the increase in ice temperature with depth to result in an increase in excess N_2O as well. However, such an increase is not observed (Schilt et al., 2010b). Third, microbial activity generally requires the presence of liquid water. Although thin water channels called “liquid veins” can form during ice crystal growth due to concentration of acidic impurities at crystal interfaces (Barletta et al., 2012; Dani et al., 2012), the upper hundreds of meters where N_2O is produced are too cold for such features to exist at sites like Vostok (Dani et al., 2012). Fourth, our SP and $\delta^{18}\text{O}$ data indicate the absence of microbial N_2O reduction in the ice (Sect. 5.1). While absence of N_2O reduction alone does not rule out microbial N_2O production, these two processes are generally linked in terrestrial and aquatic environments, where N_2O produced by bacterial denitrification is typically at least partly reduced to N_2 . Fifth, no functional genes involved in bacterial and archaeal nitrification and denitrification were detected by PCR amplification in the NEEM Greenland ice core during the last glacial period (Miteva et al., 2016). Finally, Antarctic ice is an acidic environment (EPICA community members, 2004; Wolff et al., 1997), but the review by Spott et al. (2011) indicates that hybrid N_2O production by biotic nitrosation generally occurs under neutral pH conditions, within a range of 6 to 7.5. Only one study reported production of hybrid N_2O by ammonia-oxidizing archaea at a moderately low pH of 5.5 (Jung et al., 2019). In

contrast, Su et al. (2019) showed that abiotic production of hybrid N₂O is enhanced at pH ≤ 5. Given the above constraints, an abiotic mechanism is the most plausible explanation for in situ N₂O production.

An abiotic reaction triggered by low pH could also explain the differences in N₂O production observed between Antarctic and Greenland ice. In Antarctic ice cores, N₂O production occurs consistently throughout dust-rich depths and the amount of in situ N₂O is roughly correlated with the amount of dust (Schilt et al., 2010b, a). In contrast, Greenland ice shows erratic N₂O production with a finite number of outliers of very high N₂O mixing ratio (Flückiger et al., 2004). In Greenland ice cores it occurs in specific depth intervals with moderate dust concentrations, while sections with very high dust concentrations rarely show in situ production. This very dusty Greenland ice is alkaline due to the deposition of alkaline dust (calcium carbonates), which can neutralize acids in the ice (Biscaye et al., 1997; Mayewski et al., 1994; Wolff et al., 1997), a phenomenon not observed in Antarctic ice which is acidic (EPICA community members, 2004; Wolff et al., 1997). The generally higher pH of Greenland ice (Rasmussen et al., 2023) may inhibit in situ N₂O production, potentially explaining the absence of widespread production in these cores. Isolated outliers with elevated N₂O mixing ratios occur at depths where the ice is more acidic, for instance near volcanic horizons, as they are often found close to sulfate peaks.


6 Conclusion

Using bulk and position-specific isotope analyses, we have improved the understanding of N₂O production in Antarctic ice while also raising new questions about its nature. Our results show that N₂O produced in ice is hybrid N₂O, with its central N atom (N^α) derived from NO₃⁻ and its terminal N atom (N^β) from a different precursor. This hybrid N₂O is likely produced abiotically through an N-nitrosation reaction, in which NO₃⁻ is first reduced to NO₂⁻ that then reacts with a nucleophilic compound that supplies N^β in N₂O. Fe²⁺ contained in dust particles may be the agent that reduces NO₃⁻ to NO₂⁻. In addition to the precursor of the terminal N atom, this reducing agent could be a limiting factor in the reaction. Low pH may also be a necessary condition for this process. To better understand the pH control on N₂O production, future work should investigate the isolated in situ N₂O outliers in Greenland ice. The sources of the terminal N atom (N^β) and the O atom in in situ N₂O remain uncertain. One possibility is that N^β originates from NH₄⁺, potentially converted to NH₂OH; measuring the isotopic composition of NH₄⁺ in Antarctic ice cores could help testing this hypothesis. For the O atom, it may be derived from two different oxygen pools – either from NO₃⁻ or from H₂O through oxygen exchange with NO₂⁻.

Our work revealed that the isotopic composition of in situ N₂O is highly variable even within a single ice core. As a result, N₂O records from ice cores cannot be accurately corrected for in situ production using a fixed isotopic signature in a mass balance approach. A more robust correction method would be to measure NO₃⁻ isotopic composition alongside total N₂O in the same samples, as we did in this study. Since the in situ N₂O signature reflects the variability of NO₃⁻ isotopes, this method would help estimate the isotopic signature of in situ N₂O in affected samples. We could not apply this method to the existing N₂O records because the meltwater at that time was not collected after N₂O analysis, thus the NO₃⁻ isotopic composition cannot be measured in these specific samples.

This study also highlights the importance of accounting for in situ N₂O production when using N₂O isotopes for source attribution. In some ice cores, the isotopic signature of in situ N₂O deviates drastically from the atmospheric signal. Even small amounts of in situ N₂O can significantly impact the measured isotopic composition and thus alter paleoclimatic interpretations. For example, for atmospheric N₂O at a mixing ratio of 200 ppb and in situ N₂O at 5 ppb, with $\delta^{15}\text{N}_{\text{bulk}}(\text{N}_2\text{O}_{\text{in situ}})$ as high as +70 ‰ as we observed in the Vostok ice core, this small in situ contribution would shift the $\delta^{15}\text{N}_{\text{bulk}}(\text{N}_2\text{O}_{\text{total}})$ by +1.5 ‰. Such a shift would result in a 16 % increase in estimated marine N₂O emissions and a change in marine-to-terrestrial emission ratio from 0.39 to 0.45, which is close to the entire change observed over the last 21 kyr (Fischer et al., 2019). Therefore, identifying and excluding all samples affected by in situ production is essential to avoid misinterpretation. Our results reveal a previously unidentified N₂O production pathway likely involving an asymmetric reaction intermediate. This process may not be limited to ice but could also occur in other environments such as soils or aquatic systems, suggesting it may contribute to atmospheric N₂O more broadly. This has important implications for the interpretation of SP values. The SP(N₂O_{in situ}) values observed in this study show unusual variability and strong dependence on the isotopic composition of the precursors, compared to SP values reported in previous studies (Heil et al., 2014; Toyoda et al., 2017). These results may reflect the production of hybrid N₂O via an asymmetric intermediate, where N^α and N^β originate from distinct precursors and retain the isotopic signature of their respective sources. In this case, the SP values depend on the difference between the isotopic signatures of the two sources. This mechanism complicates the use of SP to trace N₂O pathways, as they are not always characterized by a constant SP value. We therefore recommend further investigation of SP values in hybrid N₂O under different environmental conditions and reaction mechanisms.

Appendix A – Can the TALDICE data be used as an atmospheric reference?



735

Figure A1. N₂O concentrations measured in the TALDICE (Schilt et al., 2010a), NGRIP (Flückiger et al., 2004; Schilt et al., 2010a, 2013), and EDC (Schilt et al., 2010b) ice cores over the last 120 kyr (a) and over the last 40 kyr (b). In situ N₂O outliers have been removed from the NGRIP record. The EDC ice core is affected by in situ production of N₂O during the LGM and MIS4.

Appendix B – Uncertainties in the mass balance calculation

740 **Table B1.** Uncertainty ranges used in the Monte-Carlo calculation of the uncertainty of in situ N_2O isotopic signature. Here we report the uncertainties due to differences in gas age scales of TALDICE and other ice cores, as the atmospheric values in the mass balance calculation are defined as the spline values matching the age of the measured sample.

Variable	Uncertainty range
Gas age	0.7 kyr
N_2O concentration of the atmospheric spline	4 ppb
$\delta^{15}N$ values of the N_2O atmospheric spline	0.3 ‰
$\delta^{15}N^a$ value of atmospheric N_2O (Menking et al., 2025)	1.7 ‰
$\delta^{18}O$ values of the N_2O atmospheric spline	0.6 ‰
Measured N_2O concentration	4 ppb
Measured $\delta^{15}N(N_2O)$ values	0.3 ‰
Measured $\delta^{15}N^a(N_2O)$ values	0.6 ‰
Measured $\delta^{18}O(N_2O)$ values	0.6 ‰

Appendix C – Sensitivity study: potential in situ contribution in TALDICE N_2O

745




Figure C1. Spline interpolation of TALDICE N₂O concentrations, corrected for potential in situ production proportional to dust content. The different curves correspond to varying production factors, which link dust concentration to the amount of in situ N₂O produced. (a): 0 to 140 kyr BP, (b): 14 to 38 kyr BP.

750




Figure C2. Sensitivity study testing the impact of in situ N₂O production in TALDICE proportional to Ca²⁺ concentrations, with production factors (PF) ranging from 0 to 1. PF = 0 means no N₂O production. (a) Relation between δ¹⁵N_{bulk}(N₂O_{in situ}) and δ¹⁵N(NO₃⁻). (b) Relation between δ¹⁵N^α(N₂O_{in situ}) and δ¹⁵N(NO₃⁻).

755 **Appendix D – Assessment of potential NO₃⁻ contamination, loss, or isotopic fractionation during N₂O extraction from ice core samples**

Table D1. NO₃⁻ concentration and isotopic composition of the same samples with and without N₂O extraction.

		NO ₃ ⁻ concentration (ng.g ⁻¹)		NO ₃ ⁻ isotopic composition			
				δ ¹⁵ N (‰)		δ ¹⁸ O (‰)	
		No N ₂ O extraction	After N ₂ O extraction	No N ₂ O extraction	After N ₂ O extraction	No N ₂ O extraction	After N ₂ O extraction
Ultrapure water with NO ₃ ⁻ isotopic standard USGS32 (Bohlke et al., 1993)		80	75.2	+180 (reference value)	+175.7 ± 0.5	+25.4 ± 0.2 (reference value)	+34.7 ± 1.7
Duplicate ice core samples	B37 108	55.7	53.0	51.9 ± 0.4	54.8 ± 0.4	48.4 ± 2.3	59.2 ± 2.3
	EDC 1841	35.6	35.1	175.8 ± 0.5	176.0 ± 0.4	38.2 ± 1.7	40.5 ± 2.3

	EDML 1768	93.0	90.3	46.9 ± 0.4	50.9 ± 0.4	76.1 ± 2.3	84.1 ± 2.3
	EDML 976	64.8	64.6	63.3 ± 0.5	65.0 ± 0.5	78.5 ± 1.7	93.9 ± 1.7

760 **Data Availability.**

The data presented in this study are available on PANGAEA at: <https://doi.pangaea.de/10.1594/PANGAEA.986638>

Author Contribution.

The concept of the study was developed by JSc, HF, and LS. The bulk isotope analyses of N₂O were carried out by LS and BS. LS performed the position-specific isotope analyses of N₂O under the guidance of JM and EB and the NO₃⁻ isotope
765 analyses under the guidance of JSa. JSa, VL, and EB provided ice core samples. LS and JSc analyzed the data. TR, JM, and EB provided input on SP measurements. HF managed and supervised the project. LS wrote the manuscript with contributions from all authors.

Competing interests.

The contact author has declared that none of the authors has any competing interests.

770 **Acknowledgements.**

This publication was generated in the frame of the DEEPICE project. The project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 955750

Financial support.

This research has been supported by the Horizon 2020 research and innovation programme (grant no. 955750) and by the
775 Swiss National Science Foundation (grant no. 200020_200328).

References

- Aarons, S. M., Aciego, S. M., Arendt, C. A., Blakowski, M. A., Steigmeyer, A., Gabrielli, P., Sierra-Hernández, M. R., Beaudon, E., Delmonte, B., Baccolo, G., May, N. W., and Pratt, K. A.: Dust composition changes from Taylor Glacier (East Antarctica) during the last glacial-interglacial transition: A multi-proxy approach, *Quaternary Science Reviews*, 162, 60–71, 780 <https://doi.org/10.1016/j.quascirev.2017.03.011>, 2017.
- Anklin, M., Barnola, J.-M., Schwander, J., Stauffer, B., and Raynaud, D.: Processes affecting the CO₂ concentrations measured in Greenland ice, *Tellus B*, 47, 461–470, <https://doi.org/10.1034/j.1600-0889.47.issue4.6.x>, 1995.
- Baccolo, G., Delmonte, B., Niles, P. B., Cibin, G., Di Stefano, E., Hampai, D., Keller, L., Maggi, V., Marcelli, A., Michalski, J., Snead, C., and Frezzotti, M.: Jarosite formation in deep Antarctic ice provides a window into acidic, water- 785 limited weathering on Mars, *Nat Commun*, 12, 436, <https://doi.org/10.1038/s41467-020-20705-z>, 2021.
- Baggenstos, D., Severinghaus, J. P., Mulvaney, R., McConnell, J. R., Sigl, M., Maselli, O., Petit, J., Grente, B., and Steig, E. J.: A Horizontal Ice Core From Taylor Glacier, Its Implications for Antarctic Climate History, and an Improved Taylor Dome Ice Core Time Scale, *Paleoceanog and Paleoclimatol*, 33, 778–794, <https://doi.org/10.1029/2017PA003297>, 2018.
- Baggs, E. M.: Soil microbial sources of nitrous oxide: recent advances in knowledge, emerging challenges and future 790 direction, *Current Opinion in Environmental Sustainability*, 3, 321–327, <https://doi.org/10.1016/j.cosust.2011.08.011>, 2011.
- Bange, H. W.: Gaseous Nitrogen Compounds (NO, N₂O, N₂, NH₃) in the Ocean, in: *Nitrogen in the Marine Environment*, Elsevier, 51–94, <https://doi.org/10.1016/B978-0-12-372522-6.00002-5>, 2008.
- Barletta, R. E., Priscu, J. C., Mader, H. M., Jones, W. L., and Roe, C. H.: Chemical analysis of ice vein microenvironments: II. Analysis of glacial samples from Greenland and Antarctica, *J. Glaciol.*, 58, 1109–1118, 795 <https://doi.org/10.3189/2012JoG12J112>, 2012.
- Battle, M., Bender, M., Sowers, T., Tans, P. P., Butler, J. H., Elkins, J. W., Ellis, J. T., Conway, T., Zhang, N., Lang, P., and Clarket, A. D.: Atmospheric gas concentrations over the past century measured in air from firn at the South Pole, *Nature*, 383, 231–235, <https://doi.org/10.1038/383231a0>, 1996.
- Bauska, T. K., Baggenstos, D., Brook, E. J., Mix, A. C., Marcott, S. A., Petrenko, V. V., Schaefer, H., Severinghaus, J. P., 800 and Lee, J. E.: Carbon isotopes characterize rapid changes in atmospheric carbon dioxide during the last deglaciation, *Proc. Natl. Acad. Sci. U.S.A.*, 113, 3465–3470, <https://doi.org/10.1073/pnas.1513868113>, 2016.
- Bigler, M.: Hochauflösende Spurenstoffmessungen an polaren Eisbohrkernen: Glazio-chemische und klimatische Prozessstudien, <https://doi.org/10.57694/7211>, 2004.
- Biscaye, P. E., Grousset, F. E., Revel, M., Van Der Gaast, S., Zielinski, G. A., Vaars, A., and Kukla, G.: Asian provenance 805 of glacial dust (stage 2) in the Greenland Ice Sheet Project 2 Ice Core, Summit, Greenland, *J. Geophys. Res.*, 102, 26765–26781, <https://doi.org/10.1029/97JC01249>, 1997.
- Blunier, T., Floch, G. L., Jacobi, H., and Quansah, E.: Isotopic view on nitrate loss in Antarctic surface snow, *Geophysical Research Letters*, 32, 2005GL023011, <https://doi.org/10.1029/2005GL023011>, 2005.

- Bohlke, J. K., Gwinn, C. J., and Coplen, T. B.: NEW REFERENCE MATERIALS FOR NITROGEN-ISOTOPE-RATIO MEASUREMENTS, *Geostandards and Geoanalytical Research*, 17, 159–164, <https://doi.org/10.1111/j.1751-908X.1993.tb00131.x>, 1993.
- Bouchet, M., Landais, A., Grisart, A., Parrenin, F., Prié, F., Jacob, R., Fourré, E., Capron, E., Raynaud, D., Lipenkov, V. Y., Loutre, M.-F., Extier, T., Svensson, A. M., Martinerie, P., Leuenberger, M. C., Jiang, W., Ritterbusch, F., Lu, Z.-T., and Yang, G.-M.: The Antarctic ice core chronology (AICC2023), <https://doi.org/10.1594/PANGAEA.961017>, 2023.
- Buchwald, C., Grabb, K., Hansel, C. M., and Wankel, S. D.: Constraining the role of iron in environmental nitrogen transformations: Dual stable isotope systematics of abiotic NO₂⁻ reduction by Fe(II) and its production of N₂O, *Geochimica et Cosmochimica Acta*, 186, 1–12, <https://doi.org/10.1016/j.gca.2016.04.041>, 2016.
- Buiron, D., Chappellaz, J., Stenni, B., Frezzotti, M., Baumgartner, M., Capron, E., Landais, A., Lemieux-Dudon, B., Masson-Delmotte, V., Montagnat, M., Parrenin, F., and Schilt, A.: TALDICE-1 age scale of the Talos Dome deep ice core, East Antarctica, *Clim. Past*, 7, 1–16, <https://doi.org/10.5194/cp-7-1-2011>, 2011.
- Bunton, C. A., Llewellyn, D. R., and Stedman, G.: Oxygen exchange between nitrous acid and water, *J. Chem. Soc.*, 568, <https://doi.org/10.1039/jr9590000568>, 1959.
- Casciotti, K. L., Böhlke, J. K., McIlvin, M. R., Mroczkowski, S. J., and Hannon, J. E.: Oxygen Isotopes in Nitrite: Analysis, Calibration, and Equilibration, *Anal. Chem.*, 79, 2427–2436, <https://doi.org/10.1021/ac061598h>, 2007.
- Chen, Z.-L., Song, W., Hu, C.-C., Liu, X.-J., Chen, G.-Y., Walters, W. W., Michalski, G., Liu, C.-Q., Fowler, D., and Liu, X.-Y.: Significant contributions of combustion-related sources to ammonia emissions, *Nat Commun*, 13, 7710, <https://doi.org/10.1038/s41467-022-35381-4>, 2022.
- Crotti, I., Landais, A., Stenni, B., Bazin, L., Parrenin, F., Frezzotti, M., Ritterbusch, F., Lu, Z.-T., Jiang, W., Yang, G.-M., Fourré, E., Orsi, A., Jacob, R., Minster, B., Prié, F., Dreossi, G., and Barbante, C.: An extension of the TALDICE ice core age scale reaching back to MIS 10.1, *Quaternary Science Reviews*, 266, 107078, <https://doi.org/10.1016/j.quascirev.2021.107078>, 2021.
- Dani, K. G. S., Mader, H. M., Wolff, E. W., and Wadham, J. L.: Modelling the liquid-water vein system within polar ice sheets as a potential microbial habitat, *Earth and Planetary Science Letters*, 333–334, 238–249, <https://doi.org/10.1016/j.epsl.2012.04.009>, 2012.
- Delmas, R. J.: A natural artefact in Greenland ice-core CO₂ measurements, *Tellus B*, 45, 391–396, <https://doi.org/10.1034/j.1600-0889.1993.t01-3-00006.x>, 1993.
- Delmonte, B., Basile-Doelsch, I., Petit, J.-R., Maggi, V., Revel-Rolland, M., Michard, A., Jagoutz, E., and Grousset, F.: Comparing the Epica and Vostok dust records during the last 220,000 years: stratigraphical correlation and provenance in glacial periods, *Earth-Science Reviews*, 66, 63–87, <https://doi.org/10.1016/j.earscirev.2003.10.004>, 2004.
- Delmonte, B., Baroni, C., Andersson, P. S., Schoberg, H., Hansson, M., Aciego, S., Petit, J., Albani, S., Mazzola, C., Maggi, V., and Frezzotti, M.: Aeolian dust in the Talos Dome ice core (East Antarctica, Pacific/Ross Sea sector): Victoria Land

- versus* remote sources over the last two climate cycles, *J Quaternary Science*, 25, 1327–1337, <https://doi.org/10.1002/jqs.1418>, 2010.
- EPICA community members: Eight glacial cycles from an Antarctic ice core, *Nature*, 429, 623–628, <https://doi.org/10.1038/nature02599>, 2004.
- 845 EPICA Community Members: Stable oxygen isotopes of ice core EDML, <https://doi.org/10.1594/PANGAEA.754444>, 2010.
- Erbland, J.: Contraintes isotopiques sur l'interprétation de l'enregistrement en nitrate dans la carotte de glace de Vostok, Université de Grenoble, 2011.
- Erbland, J., Vicars, W. C., Savarino, J., Morin, S., Frey, M. M., Frosini, D., Vince, E., and Martins, J. M. F.: Air–snow
850 transfer of nitrate on the East Antarctic Plateau – Part 1: Isotopic evidence for a photolytically driven dynamic equilibrium in summer, *Atmos. Chem. Phys.*, 13, 6403–6419, <https://doi.org/10.5194/acp-13-6403-2013>, 2013.
- Fehling, C.: Mechanistic Insights from the ¹⁵N-Site Preference of Nitrous Oxide utilizing High Resolution Near-Infrared cw Cavity Ringdown Spectroscopy and Density Functional Theory Calculations, PhD Thesis, Kiel University, 130 pp., 2012.
- Fischer, H., Fundel, F., Ruth, U., Twarloh, B., Wegner, A., Udisti, R., Becagli, S., Castellano, E., Morganti, A., Severi, M.,
855 Wolff, E., Littot, G., Röthlisberger, R., Mulvaney, R., Hutterli, M. A., Kaufmann, P., Federer, U., Lambert, F., Bigler, M., Hansson, M., Jonsell, U., De Angelis, M., Boutron, C., Siggaard-Andersen, M.-L., Steffensen, J. P., Barbante, C., Gaspari, V., Gabrielli, P., and Wagenbach, D.: Reconstruction of millennial changes in dust emission, transport and regional sea ice coverage using the deep EPICA ice cores from the Atlantic and Indian Ocean sector of Antarctica, *Earth and Planetary Science Letters*, 260, 340–354, <https://doi.org/10.1016/j.epsl.2007.06.014>, 2007.
- 860 Fischer, H., Schmitt, J., Bock, M., Seth, B., Joos, F., Spahni, R., Lienert, S., Battaglia, G., Stocker, B. D., Schilt, A., and Brook, E. J.: N₂O changes from the Last Glacial Maximum to the preindustrial – Part 1: Quantitative reconstruction of terrestrial and marine emissions using N₂O stable isotopes in ice cores, *Biogeosciences*, 16, 3997–4021, <https://doi.org/10.5194/bg-16-3997-2019>, 2019.
- Flückiger, J., Monnin, E., Stauffer, B., Schwander, J., Stocker, T. F., Chappellaz, J., Raynaud, D., and Barnola, J.-M.: High-
865 resolution Holocene N₂O ice core record and its relationship with CH₄ and CO₂: High-resolution Holocene N₂O ice core record, *Global Biogeochem. Cycles*, 16, 10-1-10–8, <https://doi.org/10.1029/2001GB001417>, 2002.
- Flückiger, J., Blunier, T., Stauffer, B., Chappellaz, J., Spahni, R., Kawamura, K., Schwander, J., Stocker, T. F., and Dahl-Jensen, D.: N₂O and CH₄ variations during the last glacial epoch: Insight into global processes., *Global Biogeochem. Cycles*, 18, n/a-n/a, <https://doi.org/10.1029/2003GB002122>, 2004.
- 870 Frame, C. H. and Casciotti, K. L.: Biogeochemical controls and isotopic signatures of nitrous oxide production by a marine ammonia-oxidizing bacterium, *Biogeosciences*, 7, 2695–2709, <https://doi.org/10.5194/bg-7-2695-2010>, 2010.
- Frame, C. H., Lau, E., Nolan, E. J., Goepfert, T. J., and Lehmann, M. F.: Acidification Enhances Hybrid N₂O Production Associated with Aquatic Ammonia-Oxidizing Microorganisms, *Front. Microbiol.*, 7, <https://doi.org/10.3389/fmicb.2016.02104>, 2017.

- 875 Frey, M. M., Savarino, J., Morin, S., Erbland, J., and Martins, J. M. F.: Photolysis imprint in the nitrate stable isotope signal in snow and atmosphere of East Antarctica and implications for reactive nitrogen cycling, *Atmos. Chem. Phys.*, 9, 8681–8696, <https://doi.org/10.5194/acp-9-8681-2009>, 2009.
- Fuhrer, K., Wolff, E., and Johnsen, S. J.: Timescales for dust variability in the Greenland Ice Core Project (GRIP) ice core in the last 100,000 years, *J. Geophys. Res.*, 104, 31043–31052, <https://doi.org/10.1029/1999JD900929>, 1999.
- 880 Gruber, N. and Galloway, J. N.: An Earth-system perspective of the global nitrogen cycle, *Nature*, 451, 293–296, <https://doi.org/10.1038/nature06592>, 2008.
- Heil, J., Wolf, B., Brüggemann, N., Emmenegger, L., Tuzson, B., Vereecken, H., and Mohn, J.: Site-specific ^{15}N isotopic signatures of abiotically produced N_2O , *Geochimica et Cosmochimica Acta*, 139, 72–82, <https://doi.org/10.1016/j.gca.2014.04.037>, 2014.
- 885 Heil, J., Liu, S., Vereecken, H., and Brüggemann, N.: Abiotic nitrous oxide production from hydroxylamine in soils and their dependence on soil properties, *Soil Biology and Biochemistry*, 84, 107–115, <https://doi.org/10.1016/j.soilbio.2015.02.022>, 2015.
- Herron, M. M. and Langway, C. C.: Firm Densification: An Empirical Model, *J. Glaciol.*, 25, 373–385, <https://doi.org/10.3189/S0022143000015239>, 1980.
- 890 IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, , In Press, <https://doi.org/10.1017/9781009157896>, 2021.
- Jones, L. C., Peters, B., Lezama Pacheco, J. S., Casciotti, K. L., and Fendorf, S.: Stable Isotopes and Iron Oxide Mineral Products as Markers of Chemodenitrification., *Environ. Sci. Technol.*, 49, 3444–3452, <https://doi.org/10.1021/es504862x>, 2015.
- 895 Jung, M.-Y., Gwak, J.-H., Rohe, L., Gieseemann, A., Kim, J.-G., Well, R., Madsen, E. L., Herbold, C. W., Wagner, M., and Rhee, S.-K.: Indications for enzymatic denitrification to N_2O at low pH in an ammonia-oxidizing archaeon, *The ISME Journal*, 13, 2633–2638, <https://doi.org/10.1038/s41396-019-0460-6>, 2019.
- Kaiser, J., Hastings, M. G., Houlton, B. Z., Röckmann, T., and Sigman, D. M.: Triple Oxygen Isotope Analysis of Nitrate Using the Denitrifier Method and Thermal Decomposition of N_2O , *Anal. Chem.*, 79, 599–607, <https://doi.org/10.1021/ac061022s>, 2007.
- 900 Kaufmann, P., Fundel, F., Fischer, H., Bigler, M., Ruth, U., Udisti, R., Hansson, M., de Angelis, M., Barbante, C., Wolff, E. W., Hutterli, M., and Wagenbach, D.: Ammonium and non-sea salt sulfate in the EPICA ice cores as indicator of biological activity in the Southern Ocean, *Quaternary Science Reviews*, 29, 313–323, <https://doi.org/10.1016/j.quascirev.2009.11.009>, 2010.
- 905 Kindler, P., Guillevic, M., Baumgartner, M., Schwander, J., Landais, A., and Leuenberger, M.: Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core, *Clim. Past*, 10, 887–902, <https://doi.org/10.5194/cp-10-887-2014>, 2014.
- Lambert, F., Bigler, M., Steffensen, J. P., Hutterli, M., and Fischer, H.: Centennial mineral dust variability in high-resolution ice core data from Dome C, Antarctica, *Clim. Past*, 8, 609–623, <https://doi.org/10.5194/cp-8-609-2012>, 2012.

- Lamothe, A., Savarino, J., Ginot, P., Soussaintjean, L., Gautier, E., Akers, P. D., Caillon, N., and Erbland, J.: An extraction method for nitrogen isotope measurement of ammonium in a low-concentration environment, *Atmos. Meas. Tech.*, 16, 4015–4030, <https://doi.org/10.5194/amt-16-4015-2023>, 2023.
- Lan, X., Thoning, K., Dlugokencky, E., and NOAA Global Monitoring Laboratory: Trends in globally-averaged CH₄, N₂O, and SF₆ determined from NOAA Global Monitoring Laboratory measurements (Version 2025-04), <https://doi.org/10.15138/P8XG-AA10>, 2025.
- Landais, A. and Stenni, B.: Water stable isotopes (dD, d18O) from EPICA Dome C ice core (Antarctica) (0-800 ka), <https://doi.org/10.1594/PANGAEA.934094>, 2021.
- Lee, J. E., Edwards, J. S., Schmitt, J., Fischer, H., Bock, M., and Brook, E. J.: Excess methane in Greenland ice cores associated with high dust concentrations, *Geochimica et Cosmochimica Acta*, 270, 409–430, <https://doi.org/10.1016/j.gca.2019.11.020>, 2020.
- Lewicka-Szczebak, D., Well, R., Bol, R., Gregory, A. S., Matthews, G. P., Misselbrook, T., Whalley, W. R., and Cardenas, L. M.: Isotope fractionation factors controlling isotopocule signatures of soil-emitted N₂ O produced by denitrification processes of various rates, *Rapid Comm Mass Spectrometry*, 29, 269–282, <https://doi.org/10.1002/rcm.7102>, 2015.
- Lorius, C., Jouzel, J., Ritz, C., Merlivat, L., Barkov, N. I., Korotkevitch, Y. S., and Kotlyakov, V.: Delta 18O record versus age for 150 ka in ice core Vostok, <https://doi.org/10.1594/PANGAEA.860950>, 1985.
- Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola, J.-M., Raynaud, D., Stocker, T. F., and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH₄ over the past 800,000 years, *Nature*, 453, 383–386, <https://doi.org/10.1038/nature06950>, 2008.
- Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura, K., and Stocker, T. F.: High-resolution carbon dioxide concentration record 650,000–800,000 years before present, *Nature*, 453, 379–382, <https://doi.org/10.1038/nature06949>, 2008.
- Marino, F., Castellano, E., Nava, S., Chiari, M., Ruth, U., Wegner, A., Lucarelli, F., Udisti, R., Delmonte, B., and Maggi, V.: Coherent composition of glacial dust on opposite sides of the East Antarctic Plateau inferred from the deep EPICA ice cores, *Geophysical Research Letters*, 36, 2009GL040732, <https://doi.org/10.1029/2009GL040732>, 2009.
- Markle, B. R. and Steig, E. J.: Improving temperature reconstructions from ice-core water-isotope records, *Clim. Past*, 18, 1321–1368, <https://doi.org/10.5194/cp-18-1321-2022>, 2022.
- Mayewski, P. A., Meeker, L. D., Whitlow, S., Twickler, M. S., Morrison, M. C., Bloomfield, P., Bond, G. C., Alley, R. B., Gow, A. J., Meese, D. A., Grootes, P. M., Ram, M., Taylor, K. C., and Wumkes, W.: Changes in Atmospheric Circulation and Ocean Ice Cover over the North Atlantic During the Last 41,000 Years, *Science*, 263, 1747–1751, <https://doi.org/10.1126/science.263.5154.1747>, 1994.
- Menking, J. A., Brook, E. J., Schilt, A., Shackleton, S., Dyonisius, M., Severinghaus, J. P., and Petrenko, V. V.: Millennial-Scale Changes in Terrestrial and Marine Nitrous Oxide Emissions at the Onset and Termination of Marine Isotope Stage 4, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2020GL089110>, 2020.

- Menking, J. A., Lee, J. E., Brook, E. J., Schmitt, J., Soussaintjean, L., Fischer, H., Kaiser, J., and Rice, A.: Glacial-Interglacial and Millennial-Scale Changes in Nitrous Oxide Emissions Pathways and Source Regions, *Global Biogeochemical Cycles*, 39, e2024GB008287, <https://doi.org/10.1029/2024GB008287>, 2025.
- 945 Meusinger, C., Berhanu, T. A., Erbland, J., Savarino, J., and Johnson, M. S.: Laboratory study of nitrate photolysis in Antarctic snow. I. Observed quantum yield, domain of photolysis, and secondary chemistry, *The Journal of Chemical Physics*, 140, 244305, <https://doi.org/10.1063/1.4882898>, 2014.
- Miteva, V.: Bacteria in Snow and Glacier Ice, in: *Psychrophiles: from Biodiversity to Biotechnology*, Springer Berlin Heidelberg, Berlin, Heidelberg, 31–50, https://doi.org/10.1007/978-3-540-74335-4_3, 2008.
- 950 Miteva, V., Sowers, T., and Brenchley, J.: Production of N₂O by Ammonia Oxidizing Bacteria at Subfreezing Temperatures as a Model for Assessing the N₂O Anomalies in the Vostok Ice Core, *Geomicrobiology Journal*, 24, 451–459, <https://doi.org/10.1080/01490450701437693>, 2007.
- Miteva, V., Sowers, T., Schüpbach, S., Fischer, H., and Brenchley, J.: Geochemical and Microbiological Studies of Nitrous Oxide Variations within the New NEEM Greenland Ice Core during the Last Glacial Period, *Geomicrobiology Journal*, 33, 647–660, <https://doi.org/10.1080/01490451.2015.1074321>, 2016.
- 955 Morin, S., Savarino, J., Frey, M. M., Domine, F., Jacobi, H.-W., Kaleschke, L., and Martins, J. M. F.: Comprehensive isotopic composition of atmospheric nitrate in the Atlantic Ocean boundary layer from 65°S to 79°N, *J. Geophys. Res.*, 114, D05303, <https://doi.org/10.1029/2008JD010696>, 2009.
- 960 Mühl, M., Schmitt, J., Seth, B., Lee, J. E., Edwards, J. S., Brook, E. J., Blunier, T., and Fischer, H.: Methane, ethane, and propane production in Greenland ice core samples and a first isotopic characterization of excess methane, *Clim. Past*, 19, 999–1025, <https://doi.org/10.5194/cp-19-999-2023>, 2023.
- Perron, J. R., Stedman, G., and Uysal, N.: Kinetic and product study of the reaction between nitrous acid and hydrazine, *J. Chem. Soc., Dalton Trans.*, 2058, <https://doi.org/10.1039/dt9760002058>, 1976.
- 965 Prather, M. J., Hsu, J., DeLuca, N. M., Jackman, C. H., Oman, L. D., Douglass, A. R., Fleming, E. L., Strahan, S. E., Steenrod, S. D., Søvde, O. A., Isaksen, I. S. A., Froidevaux, L., and Funke, B.: Measuring and modeling the lifetime of nitrous oxide including its variability, *J. Geophys. Res. Atmos.*, 120, 5693–5705, <https://doi.org/10.1002/2015JD023267>, 2015.
- 970 Priscu, J. C., Christner, B. C., Dore, J. E., Popp, B. N., Casciotti, K. L., and Lyons, W. B.: Supersaturated N₂O in a perennially ice-covered Antarctic lake: Molecular and stable isotopic evidence for a biogeochemical relict, *Limnology & Oceanography*, 53, 2439–2450, <https://doi.org/10.4319/lo.2008.53.6.2439>, 2008.
- Prokopiou, M., Sapart, C. J., Rosen, J., Sperlich, P., Blunier, T., Brook, E., Van De Wal, R. S. W., and Röckmann, T.: Changes in the Isotopic Signature of Atmospheric Nitrous Oxide and Its Global Average Source During the Last Three Millennia, *JGR Atmospheres*, 123, <https://doi.org/10.1029/2018JD029008>, 2018.
- 975 Rasmussen, S. O., Dahl-Jensen, D., Fischer, H., Fuhrer, K., Hansen, S. B., Hansson, M., Hvidberg, C. S., Jonsell, U., Kipfstuhl, S., Ruth, U., Schwander, J., Siggaard-Andersen, M.-L., Sinnl, G., Steffensen, J. P., Svensson, A. M., and Vinther,

- B. M.: Ice-core data used for the construction of the Greenland Ice-Core Chronology 2005 and 2021 (GICC05 and GICC21), *Earth Syst. Sci. Data*, 15, 3351–3364, <https://doi.org/10.5194/essd-15-3351-2023>, 2023.
- Ravishankara, A. R., Daniel, J. S., and Portmann, R. W.: Nitrous Oxide (N₂O): The Dominant Ozone-Depleting Substance
980 Emitted in the 21st Century, *Science*, 326, 123–125, <https://doi.org/10.1126/science.1176985>, 2009.
- Ritz, C., Lliboutry, L., and Rado, C.: Analysis of a 870 m deep temperature profile at Dome C, *Annals of Glaciology*, 3, 284–289, 1982.
- Rohde, R. A., Price, P. B., Bay, R. C., and Bramall, N. E.: In situ microbial metabolism as a cause of gas anomalies in ice, *Proceedings of the National Academy of Sciences*, 105, 8667–8672, <https://doi.org/10.1073/pnas.0803763105>, 2008.
- 985 Röthlisberger, R., Hutterli, M. A., Sommer, S., Wolff, E. W., and Mulvaney, R.: Factors controlling nitrate in ice cores: Evidence from the Dome C deep ice core, *J. Geophys. Res.*, 105, 20565–20572, <https://doi.org/10.1029/2000JD900264>, 2000a.
- Röthlisberger, R., Bigler, M., Hutterli, M., Sommer, S., Stauffer, B., Junghans, H. G., and Wagenbach, D.: Technique for
990 Continuous High-Resolution Analysis of Trace Substances in Firn and Ice Cores, *Environ. Sci. Technol.*, 34, 338–342, <https://doi.org/10.1021/es9907055>, 2000b.
- Rubasinghege, G., Spak, S. N., Stanier, C. O., Carmichael, G. R., and Grassian, V. H.: Abiotic Mechanism for the Formation of Atmospheric Nitrous Oxide from Ammonium Nitrate, *Environ. Sci. Technol.*, 45, 2691–2697, <https://doi.org/10.1021/es103295v>, 2011.
- Rubino, M., Etheridge, D. M., Thornton, D. P., Howden, R., Allison, C. E., Francey, R. J., Langenfelds, R. L., Steele, L. P.,
995 Trudinger, C. M., Spencer, D. A., Curran, M. A. J., Van Ommen, T. D., and Smith, A. M.: Revised records of atmospheric trace gases CO₂, CH₄, N₂O, and d¹³C-CO₂ over the last 2000 years from Law Dome, Antarctica, *Earth Syst. Sci. Data*, 11, 473–492, <https://doi.org/10.5194/essd-11-473-2019>, 2019.
- Ruth, U., Wagenbach, D., Bigler, M., Steffensen, J. P., Röthlisberger, R., and Miller, H.: High-resolution microparticle profiles at NorthGRIP, Greenland: case studies of the calcium–dust relationship, *Ann. Glaciol.*, 35, 237–242,
1000 <https://doi.org/10.3189/172756402781817347>, 2002.
- Ruth, U., Wagenbach, D., Steffensen, J. P., and Bigler, M.: Continuous record of microparticle concentration and size distribution in the central Greenland NGRIP ice core during the last glacial period, *J. Geophys. Res.*, 108, 2002JD002376, <https://doi.org/10.1029/2002JD002376>, 2003.
- Salamatin, A. N., Lipenkov, V. Y., and Blinov, K. V.: Vostok (Antarctica) climate record time-scale deduced from the
1005 analysis of a borehole-temperature profile, *Annals of Glaciology*, 20, 207–214, 1994.
- Samarkin, V. A., Madigan, M. T., Bowles, M. W., Casciotti, K. L., Priscu, J. C., McKay, C. P., and Joye, S. B.: Abiotic nitrous oxide emission from the hypersaline Don Juan Pond in Antarctica, *Nature Geosci*, 3, 341–344, <https://doi.org/10.1038/ngeo847>, 2010.

- Schilt, A., Baumgartner, M., Schwander, J., Buiron, D., Capron, E., Chappellaz, J., Loulergue, L., Schüpbach, S., Spahni, R.,
1010 Fischer, H., and Stocker, T. F.: Atmospheric nitrous oxide during the last 140,000 years, *Earth and Planetary Science Letters*,
300, 33–43, <https://doi.org/10.1016/j.epsl.2010.09.027>, 2010a.
- Schilt, A., Baumgartner, M., Blunier, T., Schwander, J., Spahni, R., Fischer, H., and Stocker, T. F.: Glacial–interglacial and
millennial-scale variations in the atmospheric nitrous oxide concentration during the last 800,000 years, *Quaternary Science
Reviews*, 29, 182–192, <https://doi.org/10.1016/j.quascirev.2009.03.011>, 2010b.
- 1015 Schilt, A., Baumgartner, M., Eicher, O., Chappellaz, J., Schwander, J., Fischer, H., and Stocker, T. F.: The response of
atmospheric nitrous oxide to climate variations during the last glacial period, *Geophysical Research Letters*, 40, 1888–1893,
<https://doi.org/10.1002/grl.50380>, 2013.
- Schilt, A., Brook, E. J., Bauska, T. K., Baggenstos, D., Fischer, H., Joos, F., Petrenko, V. V., Schaefer, H., Schmitt, J.,
Severinghaus, J. P., Spahni, R., and Stocker, T. F.: Isotopic constraints on marine and terrestrial N₂O emissions during the
1020 last deglaciation, *Nature*, 516, 234–237, <https://doi.org/10.1038/nature13971>, 2014.
- Schmitt, J., Seth, B., Bock, M., and Fischer, H.: Online technique for isotope and mixing ratios of CH₄, N₂O, Xe and mixing
ratios of organic trace gases on a single ice core sample, *Atmos. Meas. Tech.*, 7, 2645–2665, <https://doi.org/10.5194/amt-7-2645-2014>, 2014.
- Schüpbach, S., Federer, U., Kaufmann, P. R., Albani, S., Barbante, C., Stocker, T. F., and Fischer, H.: Calcium and sodium
1025 concentrations of the Talos Dome ice core, *Antarctica*, <https://doi.org/10.1594/PANGAEA.833044>, 2014.
- Sowers, T.: N₂O record spanning the penultimate deglaciation from the Vostok ice core, *J. Geophys. Res.*, 106, 31903–
31914, <https://doi.org/10.1029/2000JD900707>, 2001.
- Spahni, R., Chappellaz, J., Stocker, T. F., Loulergue, L., Hausammann, G., Kawamura, K., Flückiger, J., Schwander, J.,
Raynaud, D., Masson-Delmotte, V., and Jouzel, J.: Atmospheric Methane and Nitrous Oxide of the Late Pleistocene from
1030 Antarctic Ice Cores, *Science*, 310, 1317–1321, <https://doi.org/10.1126/science.1120132>, 2005.
- Spolaor, A., Vallelonga, P., Gabrieli, J., Cozzi, G., Boutron, C., and Barbante, C.: Determination of Fe²⁺ and Fe³⁺ species
by FIA-CRC-ICP-MS in Antarctic ice samples, *J. Anal. At. Spectrom.*, 27, 310–317, <https://doi.org/10.1039/C1JA10276A>,
2012.
- Spolaor, A., Vallelonga, P., Cozzi, G., Gabrieli, J., Varin, C., Kehrwald, N., Zennaro, P., Boutron, C., and Barbante, C.: Iron
1035 speciation in aerosol dust influences iron bioavailability over glacial-interglacial timescales, *Geophysical Research Letters*,
40, 1618–1623, <https://doi.org/10.1002/grl.50296>, 2013.
- Spott, O., Russow, R., and Stange, C. F.: Formation of hybrid N₂O and hybrid N₂ due to codenitrification: First review of a
barely considered process of microbially mediated N-nitrosation, *Soil Biology and Biochemistry*, 43, 1995–2011,
<https://doi.org/10.1016/j.soilbio.2011.06.014>, 2011.
- 1040 Stauffer, B., Flückiger, J., Monnin, E., Nakazawa, T., and Aoki, S.: Discussion of the reliability of CO₂, CH₄ and N₂O
records from polar ice cores, *Mem. Natl Inst. Polar Res*, 139–152, 2003.

- Stieglmeier, M., Mooshammer, M., Kitzler, B., Wanek, W., Zechmeister-Boltenstern, S., Richter, A., and Schleper, C.: Aerobic nitrous oxide production through N-nitrosating hybrid formation in ammonia-oxidizing archaea, *The ISME Journal*, 8, 1135–1146, <https://doi.org/10.1038/ismej.2013.220>, 2014.
- 1045 Su, Q., Domingo-Félez, C., Jensen, M. M., and Smets, B. F.: Abiotic Nitrous Oxide (N₂O) Production Is Strongly pH Dependent, but Contributes Little to Overall N₂O Emissions in Biological Nitrogen Removal Systems, *Environ. Sci. Technol.*, 53, 3508–3516, <https://doi.org/10.1021/acs.est.8b06193>, 2019.
- Sutka, R. L., Ostrom, N. E., Ostrom, P. H., Gandhi, H., and Breznak, J. A.: Nitrogen isotopomer site preference of N₂O produced by *Nitrosomonas europaea* and *Methylococcus capsulatus* Bath, *Rapid Comm Mass Spectrometry*, 17, 738–745, 1050 <https://doi.org/10.1002/rcm.968>, 2003.
- Sutka, R. L., Ostrom, N. E., Ostrom, P. H., Breznak, J. A., Gandhi, H., Pitt, A. J., and Li, F.: Distinguishing Nitrous Oxide Production from Nitrification and Denitrification on the Basis of Isotopomer Abundances, *Appl Environ Microbiol*, 72, 638–644, <https://doi.org/10.1128/AEM.72.1.638-644.2006>, 2006.
- Tarasov, L. and Peltier, W. R.: Greenland glacial history, borehole constraints, and Eemian extent, *J. Geophys. Res.*, 108, 2001JB001731, <https://doi.org/10.1029/2001JB001731>, 2003. 1055
- Terada, A., Sugawara, S., Hojo, K., Takeuchi, Y., Riya, S., Harper, W. F., Yamamoto, T., Kuroiwa, M., Isobe, K., Katsuyama, C., Suwa, Y., Koba, K., and Hosomi, M.: Hybrid Nitrous Oxide Production from a Partial Nitrifying Bioreactor: Hydroxylamine Interactions with Nitrite, *Environ. Sci. Technol.*, 51, 2748–2756, <https://doi.org/10.1021/acs.est.6b05521>, 2017.
- 1060 Tian, H., Xu, R., Canadell, J. G., Thompson, R. L., Winiwarter, W., Suntharalingam, P., Davidson, E. A., Ciais, P., Jackson, R. B., Janssens-Maenhout, G., Prather, M. J., Regnier, P., Pan, N., Pan, S., Peters, G. P., Shi, H., Tubiello, F. N., Zaehle, S., Zhou, F., Arneeth, A., Battaglia, G., Berthet, S., Bopp, L., Bouwman, A. F., Buitenhuis, E. T., Chang, J., Chipperfield, M. P., Dangal, S. R. S., Dlugokencky, E., Elkins, J. W., Eyre, B. D., Fu, B., Hall, B., Ito, A., Joos, F., Krummel, P. B., Landolfi, A., Laruelle, G. G., Lauerwald, R., Li, W., Lienert, S., Maavara, T., MacLeod, M., Millet, D. B., Olin, S., Patra, P. K., Prinn, R. G., Raymond, P. A., Ruiz, D. J., van der Werf, G. R., Vuichard, N., Wang, J., Weiss, R. F., Wells, K. C., Wilson, C., Yang, 1065 J., and Yao, Y.: A comprehensive quantification of global nitrous oxide sources and sinks, *Nature*, 586, 248–256, <https://doi.org/10.1038/s41586-020-2780-0>, 2020.
- Tischer, J., Zopfi, J., Frey, C., Magyar, P. M., Brand, A., Oswald, K., Jegge, C., Frame, C. H., Miracle, M. R., Sòria-Perpinyà, X., Vicente, E., and Lehmann, M. F.: Isotopic signatures of biotic and abiotic N₂O production and consumption in the water column of meromictic, ferruginous Lake La Cruz (Spain), *Limnology & Oceanography*, 67, 1760–1775, 1070 <https://doi.org/10.1002/lno.12165>, 2022.
- Toyoda, S., Yoshida, N., Miwa, T., Matsui, Y., Yamagishi, H., Tsunogai, U., Nojiri, Y., and Tsurushima, N.: Production mechanism and global budget of N₂O inferred from its isotopomers in the western North Pacific, *Geophysical Research Letters*, 29, <https://doi.org/10.1029/2001GL014311>, 2002.

- 1075 Toyoda, S., Mutobe, H., Yamagishi, H., Yoshida, N., and Tanji, Y.: Fractionation of N₂O isotopomers during production by denitrifier, *Soil Biology and Biochemistry*, 37, 1535–1545, <https://doi.org/10.1016/j.soilbio.2005.01.009>, 2005.
- Toyoda, S., Yoshida, N., and Koba, K.: Isotopocule analysis of biologically produced nitrous oxide in various environments, *Mass Spectrometry Reviews*, 36, 135–160, <https://doi.org/10.1002/mas.21459>, 2017.
- Traversi, R., Barbante, C., Gaspari, V., Fattori, I., Largiuni, O., Magaldi, L., and Udisti, R.: Aluminium and iron record for the last 28 kyr derived from the Antarctic EDC96 ice core using new CFA methods, *Ann. Glaciol.*, 39, 300–306, <https://doi.org/10.3189/172756404781814438>, 2004.
- 1080 Warneck, P. and Wurzinger, C.: Product quantum yields for the 305-nm photodecomposition of nitrate in aqueous solution, *J. Phys. Chem.*, 92, 6278–6283, <https://doi.org/10.1021/j100333a022>, 1988.
- Wilhelms, F., Sheldon, S. G., Hamann, I., and Kipfstuhl, S.: Implications for and findings from deep ice core drillings an example: The ultimate tensile strength of ice at high strain rates, *Physics and Chemistry of Ice (The proceedings of the International Conference on the Physics and Chemistry of Ice held at Bremerhaven, Germany on 23-28 July 2006)* The Royal Society of Chemistry Special Publication No. 311, 635–639, 2007.
- 1085 Wolff, E. W., Moore, J. C., Clausen, H. B., and Hammer, C. U.: Climatic implications of background acidity and other chemistry derived from electrical studies of the Greenland Ice Core Project ice core, *J. Geophys. Res.*, 102, 26325–26332, <https://doi.org/10.1029/96JC02223>, 1997.
- 1090 Wrage-Mönnig, N., Horn, M. A., Well, R., Müller, C., Velthof, G., and Oenema, O.: The role of nitrifier denitrification in the production of nitrous oxide revisited, *Soil Biology and Biochemistry*, 123, A3–A16, <https://doi.org/10.1016/j.soilbio.2018.03.020>, 2018.
- York, D., Evensen, N. M., Martínez, M. L., and De Basabe Delgado, J.: Unified equations for the slope, intercept, and standard errors of the best straight line, *American Journal of Physics*, 72, 367–375, <https://doi.org/10.1119/1.1632486>, 2004.
- 1095 Yu, L., Harris, E., Lewicka-Szczebak, D., Barthel, M., Blomberg, M. R. A., Harris, S. J., Johnson, M. S., Lehmann, M. F., Liisberg, J., Müller, C., Ostrom, N. E., Six, J., Toyoda, S., Yoshida, N., and Mohn, J.: What can we learn from N₂O isotope data? – Analytics, processes and modelling, *Rapid Comm Mass Spectrometry*, 34, e8858, <https://doi.org/10.1002/rcm.8858>, 2020.
- 1100 Zhu-Barker, X., Cavazos, A. R., Ostrom, N. E., Horwath, W. R., and Glass, J. B.: The importance of abiotic reactions for nitrous oxide production, *Biogeochemistry*, 126, 251–267, <https://doi.org/10.1007/s10533-015-0166-4>, 2015.