

Holocene temperatures in southwestern Greenland controlled by topography, ice sheet proximity and oceanic conditions

Sudip Acharya^{1,*}, Allison A. Cluett^{1,2}, Amy L. Grogan¹, Jason P. Briner¹, Isla S. Castañeda³, Elizabeth K. Thomas¹

¹ Department of Earth Sciences, State University of New York at Buffalo, Buffalo, NY, USA

² University of California, Santa Cruz and NOAA Southwest Fisheries Science Center, CA, USA

³ University of Massachusetts Amherst, Amherst, MA, USA

*Correspondence to: Sudip Acharya (sudipach@buffalo.edu)

Abstract. The Holocene Thermal Maximum (HTM), a period during the Early and Middle Holocene when Greenland experienced a warmer climate than pre-industrial times, provides an ideal opportunity to test the sensitivity of the Greenland Ice Sheet to prolonged warmth. However, available climate reconstructions from the region provide a contrasting picture of the HTM—several reconstructions show an earlier HTM between the Early- to Middle-Holocene, while others show a delayed HTM between the Middle- to Late-Holocene. This discrepancy may be due to either the seasonal sensitivity of the proxies or to spatio-temporal climate variations. Here we generate five new Holocene timeseries of branched glycerol dialkyl glycerol tetraether (brGDGT)-inferred mean air temperature for months above freezing from lakes along a latitudinal transect in southwestern Greenland, yielding a total of seven Holocene brGDGT-inferred timeseries in this region. Decoupled Holocene trends between the ratio caldarchaeol to crenarchaeol and methylation of branched tetraethers index (MBT'_{SMe}) and the isomer ratio of 5- to 6-methyl brGDGTs (IR_{6Me}) in individual lakes suggest brGDGTs in these lakes are primarily influenced by temperature. Lake model sensitivity tests suggest minimal intra-lake variation in both the seasonal production window of brGDGT and the sensitivity of studied lakes to air temperature changes. Moreover, Holocene trends in the caldarchaeol-to-crenarchaeol ratio and reconstructed temperature are decoupled, suggesting that regional climate is the primary mechanism influencing these timeseries. Five timeseries. Six of the brGDGT timeseries suggest a temperature maximum between approx. 7.5 and 5 ka, following peak summer insolation and in agreement with many regional reconstructions. A coastal site experienced a temperature maximum between approx. 5 to 3 ka, coinciding with nearby sea surface temperature reconstructions, suggesting this site is influenced strongly by ocean-atmosphere heat exchange. A site far from both the coast and the Greenland Ice Sheet suggests peak warmth in the Early Holocene. These results, based on a single proxy, suggest that southwestern Greenland largely experienced similar Holocene summer temperature trends, but with local temperature variations occurred due to caused by proximity to the ice sheet and the ocean. Further investigations quantifying seasonal sensitivity between proxies and local effects (e.g., site-specific systematics, and proximity to the ice sheet and the ocean) may reveal more similarities among proxy timeseries than previously reconstructed—help reconcile the differences and refine Holocene climate reconstructions.

1 Introduction

35 Arctic warming has profound and far-reaching impacts on the region's environment (~~Rantanen et al., 2022; Thoman et al., 2023~~). For example, the Greenland Ice Sheet has been melting at an increasing rate, causing sea-level rise (~~Dutton et al., 2015; Frederikse et al., 2020~~) (Dutton et al., 2015; Frederikse et al., 2020). Ocean circulation and biogeochemical cycles have also undergone significant changes (~~Vihma et al., 2016; Wrona et al., 2016~~) (Vihma et al., 2016; Wrona et al., 40 2016). Nutrient input into lakes ~~have~~has increased due to permafrost thawing (Reyes and Loughheed, 2015), leading to summer stratification and enhanced microbial respiration (~~Antoniades et al., 2024; Jane et al., 2023; Klanten et al., 2023; Prowse et al., 2011~~) (Antoniades et al., 2024; Jane et al., 2023; Klanten et al., 2023; Prowse et al., 2011). These changes raise serious concerns about the future of the Arctic environment as global warming continues. However, available climate model projections show inter-model differences (~~IPCC, 2021; Seneviratne and Hauser, 2020~~) (IPCC, 2021; Seneviratne and Hauser, 2020), making it 45 difficult to accurately predict the future. To improve our understanding of the sensitivity of the Arctic system, glaciers, and the water cycle to future climate changes and to improve climate model capability, a long-term perspective on past climate changes and causal mechanisms is required (Axford et al., 2021; Briner et al., 2016).

The Holocene epoch (11.7 thousand years before present [ka] to 0 ka) on Greenland is characterized by millennia of relatively warm climate during the Early to Middle Holocene, likely warmer than preindustrial, and subsequent gradual cooling until 50 1850 CE, generally tracking boreal summer insolation (Axford et al., 2021; Briner et al., 2016; Kaufman et al., 2020). During the Early and Middle Holocene, the region around Greenland experienced major environmental changes: mountain glaciers were either smaller or entirely absent (~~Larocca et al., 2020; Larocca and Axford, 2022; Larsen et al., 2017; Schweinsberg et al., 2017, 2019~~) (Larocca et al., 2020; Larocca and Axford, 2022; Larsen et al., 2017; Schweinsberg et al., 2017, 2019), and the Greenland Ice Sheet was smaller compared to its current extent (Briner et al., 2020; Larsen et al., 2015; Leger et al., 2024), 55 potentially causing changes in atmospheric circulation (Masson-Delmotte et al., 2005; Thomas et al., 2016, 2023). However, the patterns of Holocene climate change in Greenland are still poorly resolved. This is in part due to site-specific climate, proxy-to-proxy, and proxy-model disagreement (~~Axford et al., 2021; Briner et al., 2016; Liu et al., 2014; Martin et al., 2024~~) (Axford et al., 2021; Briner et al., 2016; Liu et al., 2014; Martin et al., 2024). Therefore, quantifying temperature variations throughout Greenland during this epoch is of particular interest.

60 The Southwestern Greenland Ice Sheet (defined herein as the region from Nuussuaq to Cape Farewell, Figure 1) is highly sensitive to Holocene climate and environmental perturbations, due to its proximity to Atlantic Ocean deepwater formation, with attendant variability in heat and moisture transport (~~Downs et al., 2020; Lazier et al., 2002; Lesnek et al., 2020; Yang et al., 2016; Young et al., 2020~~) (Downs et al., 2020; Lazier et al., 2002; Lesnek et al., 2020; Yang et al., 2016; Young et al., 2020). Holocene temperature ~~timeseries~~time series from the ice-free areas in this region are mainly derived from lacustrine 65 proxies, including pollen, chironomids, and biomarkers (Axford et al., 2021; Briner et al., 2016; Gajewski, 2015; Kaufman,

2004). Although most of the existing time series show gradual cooling throughout the Late Holocene, they disagree on the timing and magnitude of the temperature maximum, with evidence for an earlier and larger temperature maximum farther north, but with differences between archives and proxies (~~Axford et al., 2021; Briner et al., 2016; Gajewski, 2015; Kaufman, 2004; Larocca and Axford, 2022~~)(Axford et al., 2021; Briner et al., 2016; Gajewski, 2015; Kaufman, 2004; Larocca and Axford, 2022).

Reconstructions of Greenland Ice Sheet margin positions between Ilulissat and Kangerlussuaq suggest rapid retreat and warmest summers between ~10.4 and 9.1 ka (Lesnek et al., 2020), a time period not covered by many other terrestrial archives. An annual temperature reconstruction based on argon and nitrogen isotopes from the Greenland Ice Sheet Project (GISP2) ice core reveals a Holocene thermal maximum (HTM) at ~7.9 to 6 ka, or 10 to 7.5 ka when considering possible elevation changes with temperature ~3 to 5 °C warmer than the recent decades (~~Axford et al., 2021; Kobashi et al., 2017~~)(Axford et al., 2021; Kobashi et al., 2017) (Fig. 1; point 6). Chironomid-inferred July air temperatures from North Lake near Ilulissat (Fig. 1, point 10) ~~show~~ the warmest period of the past 7 ka between ~6 and 4 ka, with temperatures ~2 to 3 °C higher than preindustrial (Axford et al., 2013). A branched glycerol dialkyl glycerol tetraether (brGDGT)-inferred summer lake water temperature time series from Lake Gus between Kangerlussuaq and Sisimiut reveals maximum temperature of the past 9 ka from 9 to 6 ka (Cluett et al., 2023) (Fig. 1, point 11), whereas brGDGTs from Lake 578 in southern Greenland reveals the warmest period in the past 11 ka from ~7.5 to 4.5 ka, with ice-free season temperature ~4 to 6 °C higher than preindustrial in both timeseries (Schneider et al., 2024) (Fig. 1, point 14). Alpine glaciers ~~from~~ in southern Greenland ~~were at~~ reached their smallest extent between ~7.3- and 7.1 ka, implying the warmest temperature in the past ~10.5 ka, with air temperature only ~1 to 2 °C above present (Larocca et al., 2020) (Fig. 1; ~~point~~points 15, 18). Pollen-based reconstructions from Lake Qipisarqo suggest that the thermal maximum of the past 8.5 ka occurred between ~7 and 5 ka, with summer air temperatures up to ~5 °C higher than pre-industrial (Frechette and de Vernal, 2009) (Fig. 1; point 12). In contrast, nearby pollen-based reconstructions from Spongilla Sø and Comarum Sø imply the warmest summers of the past 10.5 ka occurred after 5.2 ka, with temperatures ~1 to 2 °C higher than present (Fredskild, 1973; Gajewski, 2015) (Fig. 1; ~~point~~points 14 and 16). Additionally, marine reconstructions indicate the HTM between ~6 and 2 ka, with summer sea surface temperatures (SST) up to ~5 °C higher than present (~~Axford et al., 2021; Hansen et al., 2020; Ouellet-Bernier et al., 2014~~)(Axford et al., 2021; Hansen et al., 2020; Ouellet-Bernier et al., 2014).

Several mechanisms have been called upon to explain ~~the different~~differences in timing and magnitude of peak warmth: differing seasonal ~~sensitivity~~sensitivities of proxies, ice-sheet meltwater ~~influence causing~~influencing delayed warmth in marine archives, and vegetation lags causing delays in pollen records (Axford et al., 2021; Briner et al., 2016). Some of these mechanisms can be tested by generating time series from the same proxy in similar archives along a spatial gradient, thus avoiding possible differences caused by the archive/proxy.

Here we developed five new ~~timeseries~~time series of brGDGT-inferred mean air temperatures for months above freezing (MAF) from ~~five~~ lakes along a latitudinal transect from 64 to 69° N (~~Fig. 1~~) in southwestern Greenland, spanning the past 9

ka- (Fig. 1). BrGDGTs are membrane-spanning lipids of bacteria (Sinninghe Damsté et al., 2000) and they are well preserved in lake sediments (Castañeda and Schouten, 2011; Schouten et al., 2013). ~~BrGDGTs~~BrGDGT distributions are primarily influenced by temperature (Raberg et al., 2021), and therefore, this proxy has shown promise as a quantitative high-latitude terrestrial paleotemperature proxy (~~e.g., Cluett et al., 2023; Lindberg et al., 2022; Schneider et al., 2024; Thomas et al., 2018~~)(e.g., Cluett et al., 2023; Lindberg et al., 2022; Schneider et al., 2024; Thomas et al., 2018). However, the influence of site-specific factors such as sub-oxic conditions (~~van Bree et al., 2020; Raberg et al., 2025; Weber et al., 2018; Wu et al., 2021~~)(van Bree et al., 2020; Raberg et al., 2025; Weber et al., 2018; Wu et al., 2021), lake ice cover duration (Shanahan et al., 2013), and microbial ecology (~~Liang et al., 2024~~)(Liang et al., 2024) should also be taken into consideration before reconstructing temperature. To investigate the influence of site-specific variations such as mixing regimes that may cause sub-oxic conditions, duration of ice cover, and differing sensitivity of each lake to air temperature change, we ran the lake energy, hydrologic, and isotopic mass balance model developed by (Dee et al., 2018; Hostetler and Bartlein, 1990) and updated by (Morrill et al., 2019) for each lake under modern and perturbed climate scenarios. Overall, we aim to investigate (1) Holocene temperature evolution in southwestern Greenland, and (2) Possible causes of temporal differences in temperature maxima.

2 Materials and methods

2.1 Study area and coring

We studied five lakes in southwestern Greenland with informal names: Pluto, N3, Rosaea, Marshall, and Bullet (Fig. 1). None of the lakes currently ~~receives~~receives ice-sheet meltwater. Table 1 has detailed geographical and morphological features of each lake. Lakes Pluto (69.109° N, 51.032° W, 190 m asl; Fig. 1; point 1) and N3 (68.636° N, 50.981° W, 59 m asl; Fig. 1; point 2) are ~~situated~~ south of Ilulissat, western Greenland. Watershed vegetation consists of dwarf shrub heath, dominated by *Salix* sp. and *Betula* sp. (Bennike, 2000). On July 27, 2010, during core collection, depth profiles of temperature, conductivity, salinity, pH and dissolved oxygen were measured in Lake N3 (Corcoran et al., 2021; Thomas et al., 2016). During the field expedition in July 2009 (Pluto Lake) and late July 2010 (N3), both lakes had small inflowing streams and active channelized outflows. For more details, see Thomas et al., (2016; 2020).

Table 1: Geographical and morphological characteristics including site ID in Fig. 1, latitude (lat), longitude (lon), elevation (m asl), catchment area (km²), maximum depth (m), surface area (km²), and closest estimated distances from Ice Sheet and Ocean (km²) of study lakes (Pluto, N3, Rosaea, Marshall, and Bullet), Lake Gus (Cluett et al., 2023) and Lake 578 (Schneider et al., 2024).

Site ID. in Fig 1.	Lake name	Lat (°)	Lon (°)	Elevation (m asl)	Catchment area (km²)	Maximum depth (m)	Surface area (km²)	Distance from modern ice sheet (km)	Distance from ocean (km)
1	Pluto	69.109	-51.032	190	1.21	4	0.07	36	2

2	N3	68.636	-50.981	59	0.72	16	0.09	14	2
11	Gus	67.032	-52.427	300	4.61	6	0.09	100	65
3	Rosaea	66.982	-53.718	228	1.03	9	0.01	160	3
4	Marshall	64.464	-49.431	862	0.56	8	0.02	<0.5	150
5	Bullet	63.982	-49.537	944	0.30	8	0.02	1.5	100
13	578	61.080	-45.609	155	0.82	16	0.06	35	76

125


Lake Rosaea (66.982° N, 53.718° W; Fig. 1; point 3) is ~~situated~~ approx. 5 km north of Sisimiut ~~on~~in western Greenland and 3 km from the coastline ~~with~~of the Davis Strait. Watershed vegetation consists of shrubs, including *S. glauca* and *S. herbacea* on hill slopes, while *Empetrum nigrum*, *Betula nana*, and *Rhododendron tomentosum* are present on low-lying ground. Patches of snowpack were observed around the lake and at the higher elevations during the field expedition on July 18–21, 2018. Also, during that summer, the lake was fed by multiple active channelized inflow streams, and it had an outflow stream.

Marshall (64.464° N, 49.431° W, 862 m asl; Fig. 1; point 4) and Bullet (63.982° N, 49.537° W, 944 m asl; Fig. 1; point 5) lakes are ~~situated~~ inland of Nuuk in southwestern Greenland. Marshall Lake is <2 km outboard of the modern ice sheet margin and received ice-sheet meltwater briefly during the historical maximum extent. Bullet Lake is located approx. 60 km south ~~from~~of Marshall Lake. Lake catchment consists of vegetation including *Vaccinium uliginosum*, *E. nigrum*, *S. herbacea*, and *Sphagnum* sp. Both Marshall and Bullet ~~lakes~~Lake catchments include shrubs and grasses on hillslopes and *Eriophorum scheuchzeri*, and *Sphagnum* sp. around the shoreline. ~~Both~~Neither Marshall ~~and~~nor Bullet lakes ~~do not have channelized~~had channelised inflow ~~and~~or outflow streams during the summer 2017 field season, although water flowed into and out of the lakes in rocky low-lying parts of the catchment.

The climate in coastal southwestern Greenland is influenced by atmospheric and oceanic conditions ~~(Box et al., 2023; Westergaard-Nielsen et al., 2020)~~(Box et al., 2023; Westergaard-Nielsen et al., 2020). In the summer, moisture from North America contributes to the region, while in the winter, evaporation from proximal seas becomes the dominant moisture source (Cluett et al., 2021; Nusbaumer et al., 2019). Precipitation ~~amount~~ is relatively high ~~in~~along the southernmost ~~part of the~~ coast and decreases with increasing latitude (Cluett et al., 2021). From June to September, ~~the~~ temperature is above freezing, while ~~during their~~ winter ~~months~~, it drops well below freezing. Temperature generally ~~declined~~declines northward along the coast and with increasing elevation, where snow cover and radiative cooling are more pronounced (Karami et al., 2017). The West Greenland coastal current carries warm water from the North Atlantic Ocean to Baffin Bay, and influences ~~the~~ climate in coastal southwestern Greenland ~~(Westergaard-Nielsen et al., 2020)~~(Westergaard-Nielsen et al., 2020), whereas the southernmost land on western Greenland is influenced by the ~~east~~East Greenland current, which carries cold Arctic water around the southern tip of the island- (Fig. 1).

150 **2.2 Sediment core chronology**

^{14}C -ages and age-depth models for the sediment cores from lakes N3 and Pluto are published by Thomas et al. (2016) and Thomas et al. (2020). Here, we updated the age-depth models for the lakes employing the IntCal20 calibration curve (Fig. S1–2). Additionally, we developed age-depth models for Marshall, Bullet and Rosaea lakes (Fig. S3–5). The age-depth model for Marshall Lake is derived from ^{14}C ages ~~from~~of seven aquatic moss macrofossils, ~~and~~ for Bullet and Rosaea Lakes ~~derived~~ from nine aquatic macrofossils (Tables S1–S3). All ~~the~~ age-depth models were calculated ~~using~~with rbacon ~~within~~in the GeoChronR package, using IntCal20 (Blaauw and Christen, 2011; McKay et al., 2021).






Figure 1: Map of Greenland and surrounding area showing study lakes in red circles (1–5), sites mentioned in the text in blue diamonds (6–20) and ocean currents; WGC: West Greenland, EGC: East Greenland, IG: Irminger, BC: Baffin, and LC: Labrador currents. Site numbered as follows: (1) Pluto, (2) N3, (3) Rosaea, (4) Marshall (5) Bullet (6) GISP2 (Kobashi et al., 2017)(Kobashi et al., 2017), (7) Sermikassak, (8) Qangattaq ice cap (9) Saqqap Tasersua (Schweinsberg et al., 2017, 2019) (10) North Lake (Axford et al., 2013) (11) Lake Gus (Cluett et al., 2023) (12) Qipisarqo (Frechette and de Vernal, 2009; Kaplan et al., 2002) (13) Lake 578 (Schneider et al., 2024) (14) Spongilla Sø (Fredskild, 1973; Gajewski, 2015) (15) Alakariqssoq glaciers (Larocca et al., 2020) (16) Comarum Sø (Fredskild, 1973; Gajewski, 2015) (17) N14 (Andresen et al., 2004), (18) Quvnerit glaciers (Larocca et al., 2020), and marine sediment cores (19) MSM343300 (Ouellet-Bernier et al., 2014) and (20) CC70 (Gibb et al., 2015)(Gibb et al., 2015). The background colour represents modern summer sea surface temperatures (SSTs) (Fisher et al., 2023; Moon et al., 2023)(Fisher et al., 2023; Moon et al., 2023).

2.3 Biomarker analysis

We used the method described in Cluett et al. (2023) for the extraction and purification of GDGTs. Briefly, total lipids from homogenized 0.5 to 2 g of bulk sediment samples were extracted using a Dionex Accelerated Solvent Extractor 200 with 30 ml of dichloromethane (DCM): methanol 9:1 (volume:volume, v:v). Afterwards, 0.1 µg of a C₄₆ glycerol dialkyl glycerol tetraether (GDGT) was added to the total lipid extracts (TLE). TLEs were separated into neutral and acid fractions using aminopropyl silica gel columns eluted with DCM:isopropanol (2:1 v:v) and acids with 4% acetic acid in DCM, respectively. Neutral fractions were further split into apolar, ketone, and polar fractions using alumina columns eluted with hexane:DCM (9:1, v:v), hexane:DCM (1:1, v:v), and DCM:MeOH (1:1, v:v), respectively. GDGTs contained in the polar fraction were filtered using 0.45 µm PTFE filters. GDGTs were analyzed on an Agilent 1260 Ultra High-Performance Liquid Chromatograph (UHPLC, Agilent 1260) coupled to an Agilent 6120 Mass Selective Detector (MSD) in selected ion monitoring (SIM) mode in the Biogeochemistry Laboratory at the University of Massachusetts Amherst. For the separation of 5-methyl and 6-methyl isomers, we used two coupled waters UHPLC columns (150 mm * 2.1mm*1.7 µm) and followed the procedure described by Hopmans et al. (2016). Chromatography was performed applying a three-phase isocratic solvent gradient using 100% hexane (solvent A) and hexane:isopropanol (9:1, v:v, solvent B). Peak integration was done using the “chromatoPy” package in Python (Otiniano et al., 2025)(Otiniano et al., 2025). We determined the GDGT concentrations in relation to the C₄₆ internal standard.

We analyzed 23 samples from Pluto Lake at ~4 cm resolution, 69 samples from Lake N3 at ~3 cm resolution, 44 samples from Rosaea Lake at ~6 cm resolution, 23 samples from Marshall Lake at ~4 cm resolution, and 22 samples from Bullet Lake at ~7 cm resolution. All samples were from 0.5-cm-thick subsamples of sediment cores. The GDGT time series from Lake N3 is at 100-year-resolution for the past 8 ka. Those from lakes Pluto, Marshall, and Bullet provide 400-year-resolution, covering from 9 to 2 ka, 9 to 1 ka, and 9 to 1 ka, respectively. The GDGT time series from Lake Rosaea offers a 200-year-resolution, spanning from 9 to 1 ka.

2.4 Lake modelling

We ran sensitivity tests of the lake energy and water balance model developed by (Dee et al., 2018; Hostetler and Bartlein, 1990) and modified by (Morrill et al., 2019).Morrill et al. (2019). We used ERA5 climate data (air temperature, relative humidity, wind speed, incoming surface shortwave radiation, downward longwave radiation, surface pressure, and precipitation amount) averaged over the four grid boxes nearest each lake for January 1, 1994, to December 31, 2024, as meteorological input data (Muñoz Sabater, 2019). All the meteorological inputs were at a 6-hourly timestep except total precipitation amount, which is hourly timesteps summed to a total 6-hourly precipitation amount. Lake surface areas were estimated using Google Earth, and catchment areas were estimated using the Arctic DEM in QGIS (Porter et al., 2022). Depth-area slices at a resolution of 1 meter were estimated using bathymetry of each lake (Zhao et al., 2021), which was determined either using sonar or by depth sounding with a weighted measuring tape. We ran the lake model for 30 years, with a 10-year spin-up period at the beginning using 1994 meteorological input data.

We calibrated and validated the lake model using observation data from Lake 578 in southwestern Greenland (Zhao et al., 2021). The lake has six-hourly temperature data for three years, from July 2016 to August 2019, collected from multiple depths, and is similar in geographic location and size to the lakes in our study, for which we have little to no observational data. To calibrate the model and identify the best-performing parameter for the lake's thermodynamic balance (snow albedo, slush albedo, shortwave extinction coefficient, neutral drag coefficient, sediment albedo, sediment conductivity, and sediment specific heat), we follow established methods (Cluett et al., 2023; Longo et al., 2020; Morrill et al., 2019). An ensemble of 1,000 parameter sets was generated by simultaneously sampling each parameter from a uniform distribution using Latin Hypercube Sampling. Initial parameter distributions were wide but physically plausible ranges for natural lakes. The model was then run for each parameter set with ERA5 forcing data, and the resulting simulations were compared with observed lake temperature profiles from 2016 and 2017 (Fig. S6). The 30 parameter sets that yielded the lowest root mean squared error (RMSE) were selected as the best-performing configurations.

We then conducted a validation step: we ran the lake model using the identified best-performing parameters and the ERA5 forcing data. The simulated lake temperatures were compared with Lake 578 temperature observations from 2018 and 2019 (Fig. S7). While the model estimates a slight delay in spring warming relative to observations, it accurately reproduces observed ice-free season water temperatures at different depths (2.5, 5, 7.5, 10, and 14.5 m) (Fig. S7). Furthermore, the lake model successfully simulates the timing of ice-free season stratification in the lake from 2018 to 2019, albeit with an overestimated stratification duration (Fig. S8).

~~For lakes Bullet, Marshall, Pluto, Rosaea, and 578, we used the model parameters as for Lake Gus (Cluett et al., 2023), as these lakes have similar depths and comparable surface areas. For Lake N3, we used the lake model parameters defined by Corcoran et al., (2021) for the lake. We acknowledge that using the same lake model parameters might lead to data-model mismatches with lake conditions, however for the purpose of sensitivity tests on orbital scales, such changes can only have a minor impact (Fig. S17–S19). To evaluate the sensitivity of lakes to changes in air temperature we perturbed the ERA5 input annual, and summer (defined as June to September) temperature by ± 10 , ± 6 , ± 4 , and ± 2 °C. To evaluate the impact of shading on the lake water temperature, we perturbed the incoming short wave radiation by ± 10 , ± 25 , and $\pm 50\%$. We calculated shading by surrounding topography on for each lake using GDAL in python using Arctic DEM.~~ The model successfully simulates the lake being isothermal at the beginning and end of the ice-free season (Fig. S8). These results demonstrate that the model effectively simulates the thermodynamic processes, including temperature and stratification in Lake 578, despite a minor cold bias in spring. These results provide confidence that the subsequent sensitivity experiments examining lake surface temperature and ice-free-season stratification duration responses to prescribed changes in air temperature are reliable, although it should be noted that the ice-free-season isothermal duration might be underestimated in the model. In other words, the model is

biased towards conditions favoring ice-free season stratification and thus bottom water anoxic conditions, a key consideration for interpreting model results.

Next, we validated these parameters for Lake N3, the lake in our study with an ice-free-season temperature profile. We used the 30 best parameter sets identified in the calibration for Lake 578, and ERA5 data to simulate Lake N3 temperature for 30 years, including 2010, when we have temperature profile observations. There are minimal differences among parameter sets, and all agree well with observed temperatures at the surface and at depths in the lake (Fig. S9).

Given the similar size, morphometry, and climate conditions of Lake 578, Lake N3, and our other study lakes, we used the average of the 30 best-performing parameter values identified for Lake 578 to simulate all our study lakes. We used the average parameter values, rather than 30 individual parameter sets, as conducting sensitivity tests on all lakes using all 30 sets would be prohibitively computationally intensive. Furthermore, the changes forced by large climatological shifts dwarf the inter-parameter differences. Thus, we have used the best available data, to our knowledge, to both calibrate and validate the model for lakes that are similar in size and geographic location to the study sites. We believe that the parameter values we use for sensitivity tests in our study lakes provide accurate estimates of their thermodynamic responses, including lake temperature and stratification. We acknowledge that using the same lake model parameters may lead to slight mismatches between absolute and modelled temperature values; however, for sensitivity tests at orbital scales, such changes can have only a minor impact (Fig. S21–S23). To evaluate the sensitivity of lakes to changes in air temperature, we perturbed the ERA5 input annual and summer (June to September) temperatures by ± 10 , ± 6 , ± 4 , and ± 2 °C.

To evaluate the impact of shading on lake water temperature, we perturbed incoming short-wave radiation by $\pm 10\%$, $\pm 25\%$, and $+50\%$. We calculated shading from surrounding topography for each lake in Python using GDAL, with the Arctic DEM, for daytime (sunset to sunrise) at 15-minute time steps every 10 days from April to September under modern insolation parameters. Wind direction at each lake was calculated using 6-hourly ERA5 climate data from January 1, 1994, to December 31, 2024.

3. Results and discussion

3.1 ~~Anoxia, sources~~ Sources of brGDGTs and temperature implications

~~Total brGDGT concentration in these lake sediments from southwestern Greenland was about 6 to 10 times higher than total isoprenoid GDGT (isoGDGT), with both revealing a similar downcore pattern in all lakes, peaking in the early Holocene and declining towards the Late Holocene (Fig. S6–S10). Caldarchaeol (GDGT-0) is the most abundant isoGDGT (Fig. S11a), followed by crenarchaeol (GDGT-4) and GDGT-1, whereas GDGT-2, GDGT-3, and the crenarchaeol isomer (GDGT-4') were below detection limit in many samples, restricting TEX₈₆-based temperature reconstruction (Schouten et al., 2002).~~

Caldarchaeol is produced in anaerobic conditions in lake water or sediments by methanogenic *Euryarcheota* or heterotrophic *Bathyarchaea* (Buckles et al., 2013; Sinninghe Damsté et al., 2009), whereas crenarchaeol is produced in aerobic lake water and sediments by *Thaumarchaeota* (Baxter et al., 2021; Besseling et al., 2018). The activity of anaerobic archaeal communities is higher during winter in ice-covered lakes (Baxter et al., 2021; Blaga et al., 2011; Butler et al., 2019; Massé et al., 2019; Sinninghe Damsté et al., 2009). Therefore the ratio of caldarchaeol to crenarchaeol (Cald/Cren) may be a sensitive indicator of winter suboxic conditions in the water column and/or lake sediments, where the caldarchaeol is produced (Blaga et al., 2009). Cald/Cren ranged from 1 to 857 in all study lakes, with generally low values in lakes Marshall and Rosaea and high values in lakes Bullet, N3 and Pluto (Fig. 2a). Cald/Cren was higher from 10 to 6 ka in lakes Pluto, Rosaea, Bullet, and Marshall, which may indicate lower oxygen availability (Acharya et al., 2023; Baxter et al., 2021; Blaga et al., 2009) (Fig. 2a). After approx. 6 ka, the Cald/Cren remained relatively low and stable, indicating oxic lake sediments and/or water column. Cald/Cren in Lake N3 had higher values indicating suboxic conditions from 7.5 to 5 ka, and lags the decrease recorded in other lakes by about a millennium (Fig. 2a). We hypothesize that this could be related to a greater maximum depth (~17 m) of N3 compared to other lakes (<10 m water depth). Deeper conditions favor greater stratification, which could prolong suboxic conditions (Kerimoglu and Rinke, 2013). This Early Holocene timing of high Cald/Cren is similar to results from other Arctic lakes (Cluett et al., 2023; McFarlin et al., 2023; Schneider et al., 2024) and is probably caused by prolonged thermal stratification due to longer ice cover duration and/or summer warming due to high summer insolation (Klanten et al., 2023; MacIntyre et al., 2018).

BrGDGTs in lake sediments can be allochthonous—produced in the catchment soil and transported to the lake via runoff erosion, and autochthonous—produced in the water column and surface sediment (Baxter et al., 2021; Tierney et al., 2012). BrGDGTs produced in these diverse environments can have different responses to environmental conditions, directly impacting the temperature reconstruction (Acharya et al., 2023; Martin et al., 2020). Therefore, understanding the sources of brGDGTs is crucial before temperature reconstruction. We evaluated the sources of brGDGTs in southwestern Greenland lakes by comparing their distributions with those from high latitude (>50 °N) soil, peat, and lake sediment samples (Raberg et al., 2022). BrGDGTs from southwestern Greenland lakes have different fractional abundance of hexa-, penta- and tetra-methylated brGDGTs than those in peat and soil samples but are similar to high latitude lake surface sediment samples (Fig. S11), suggesting lacustrine origin. Furthermore, brGDGT IIIa, which is produced in higher relative abundance in lake water than in soils (Weber et al., 2018), is abundant in the southwestern Greenland lake sediments, similar to high latitude lake surface sediments, and greater than in high latitude soils (Fig. S11). This corroborates the autochthonous production of brGDGTs in southwestern Greenland lakes, although there remains a possibility of minor brGDGT contributions from the surrounding catchment.

Air temperature in southwestern Greenland is above freezing only from June to September, which causes lakes to become ice free and causes increases in lake water column nutrient content and light and oxygen availability (Shanahan et al., 2013; Zhao

et al., 2021), in turn causing high aquatic production. Although the exact source organism(s) of lacustrine brGDGTs is still unclear, brGDGT production in Arctic lakes is higher during the ice-free season than the ice covered season (Coleord et al., 2015; Shanahan et al., 2013; Zhao et al., 2021). The brGDGT indices that can indicate production under suboxic conditions, HP5 and fractional abundance of brGDGT IIIa (%IIIa) (Weber et al., 2018; Yao et al., 2020), were high in all the study lakes from 10 to 8 ka, low between 8 and 5 ka, and gradually increasing thereafter (Fig. 2b and Fig. S6–10). In lakes Bullet, Marshall, and Pluto, correlations between Cald/Cren and both HP5 and %IIIa were not significant (Table S4), suggesting a decoupling between sub-oxic conditions inferred from isoGDGT derived Cald/Cren and brGDGT derived HP5 and %IIIa. In Lake N3, we observed a significant negative correlation between Cald/Cren and %IIIa, indicating antiphase sub-oxic conditions inferred from isoGDGT and brGDGT proxies (Fig. S6–S10 and table S4). In contrast, Lake Rosaea had significant positive correlations between Cald/Cren and both HP5 and %IIIa. However, the Cald/Cren ratio in Lake Rosaea remained below 45 in all samples except the last sample where ratio is <110, suggesting relatively oxie conditions, although it is possible that suboxia influenced the brGDGTs distributions in Lake Rosaea (Fig. S8). Based on the evidence for seasonal production and the contrasting trends in these lake time series, we interpret the br and iso GDGTs to be predominantly produced during different seasons: brGDGTs in ice-free season and isoGDGTs in ice-covered season (Baxter et al., 2021; Blaga et al., 2011; van Bree et al., 2020; Loomis et al., 2014b; Shanahan et al., 2013). This contrast in seasonal production may be enhanced when caldarchaeol production is high due to winter lake water sub-oxia. In lakes N3, Pluto, and Marshall brGDGT IIIa'' (Weber et al., 2015) is below detection limit in all samples. Lake Rosaea had brGDGTs IIIa'' in, one Early Holocene and four Middle Holocene samples. In Lake Bullet, brGDGT IIIa'' concentration was below detection limit prior to 7 ka. After 7 ka, concentration increased, reaching a maximum between 6 and 3 ka (Fig. S10). Afterwards, concentration remained low. The Holocene trends between Cald/Cren and IIIa'' are decoupled with each other, while trend between IIIa'' and HP5 and %IIIa are similar, supporting our inferences of different production seasonality of GDGTs.




Figure 2: GDGT-based indices in study lakes. (A) Ratio of caldarchaeol to crenarchaeol (Cald/Cren) (Blaga et al., 2009), (B) Ratio of 5-methyl-hexamethylated and pentamethylated brGDGTs with no cyclopentane rings (HP5) (Yao et al., 2020), and (C) Methylation of Branched Tetraethers index (MBT'_{SME}) (De Jonge et al., 2014). Red triangles in (C) indicate samples with $IR_{6Me} > 0.4$, which are excluded from temperature reconstructions, due to likely non-thermal influence (Bauersachs et al., 2023; Novak et al., 2025).

Lake model simulations under 30 years of modern climate suggest the occurrence of stratification (i.e., mixing depth is less than the 75% of maximum lake depth) in all the study lakes during winter and early in the ice-free season (Fig. S14), as observed today in Arctic lakes (Antoniades et al., 2024; Klanten et al., 2024; Saros et al., 2016). Sensitivity test suggests that the duration of ice-free season stratification remain constant or increases as annual or JJAS temperature increases (Fig. 3g and Fig. S14). However, the simulations show that, no matter the climate forcing, the lakes do not remain stratified throughout the entire ice-free season (Fig. 3h and Fig. S14). This means that, even if the study lakes are stratified during early spring, the lake water fully mixes during the late summer and autumn, a phenomenon observed today in Arctic lakes with similar size and chemical composition (Antoniades et al., 2024; Klanten et al., 2023, 2024; Lindborg et al., 2016). The duration of the isothermal mixing period changes similarly in all the study lakes in response to annual or JJAS air temperature model sensitivity tests, except for Lake N3 and Lake 578 (Fig. 3h and Fig. S14). In these lakes, the duration of the isothermal mixing period shows minor variations with perturbation in annual or JJAS air temperature, probably because of their larger surface area and depth compared to other study lakes. Overall, these model simulations suggest that while summer stratification does occur in these lakes, the duration of stratification is relatively short and therefore likely unable to support summer suboxic conditions, similar to Arctic lakes that experience summer stratification today (Antoniades et al., 2024; Klanten et al., 2024). The lakes in this study are relatively shallow and have low dissolved solids compared to some Arctic lakes that today experience extended


summer stratification and hypolimnetic sub-oxia (Saros et al., 2016), making the lakes in this study more susceptible to mixing by solar heating and wind. Winter sub-oxia would have a minimal impact on brGDGTs, which are primarily produced during the ice-free period (Loomis et al., 2014b; Shanahan et al., 2013; Zhao et al., 2021). Therefore, we assume that brGDGTs in the study lakes, mainly produced during summer, are minimally influenced by changes in sub-oxic conditions during the Holocene.

While there are several lacustrine brGDGT temperature calibrations available globally (Martínez Sosa et al., 2021; Raberg et al., 2021; Zhao et al., 2023), regionally (Bauersachs et al., 2023; Dang et al., 2018; Otiniano et al., 2023, 2024; Russell et al., 2018), and site specifically (Zhao et al., 2021; Bittner et al., 2022; Feng et al., 2019), only Raberg et al. (2021), Zhao et al. (2021), and Otiniano et al. (2023; 2024) are based on the Hopmans et al., (2016) HPLC method separating GDGTs isomers and explicitly consider Arctic seasonal climate. Therefore, we assess these calibrations to infer temperature in the southwestern Greenland lakes. The Raberg et al. (2021) and Otiniano et al. (2023; 2024) calibrations estimate mean air temperature for the months above freezing (MAF) while Zhao et al. (2021) estimates epilimnion lake water temperature for the ice-free season (Ice free season lake water temperature, IFS LWT). For all analyzed samples, the reconstructed MAF based on equation 11 in Raberg et al. (2021) yields a temperature range from -3.9 to 11.6 °C; Otiniano et al. (2024) yields a temperature range from 3.8 to 8.6 °C; and Otiniano et al. (2023) yields a temperature range from 5.8 to 9.9 °C, while IFS LWT reconstrued using Zhao et al. (2021) yields a temperature range from -5.9 to 21.6 °C (Fig. S13). For lakes Marshall, N3, Pluto, and Rosaea, all four calibrations generated similar temperature trends throughout the Holocene, while the absolute values are different (Fig. S13). For Bullet Lake, the MBT'_{SME}-based calibrations (Otiniano et al. (2024), Otiniano et al. (2023), and Zhao et al. (2021)) yielded similar trends, whereas Raberg et al. (2021) yielded a divergent trend, in addition to the calibrations yielding different absolute values (Fig. S13, black time series). The discrepancy between these MBT'_{SME}-based calibrations and Raberg et al. (2021) calibration in Bullet Lake could be due brGDGTs Ib, which weighs more on Raberg et al., 2021 than MBT'_{SME}-based calibrations, is more abundance in Bullet Lake (Fig. S11).

Observations of Arctic lakes show that water temperature changes are smaller and delayed compared to air temperature (Kettle et al., 2004; Piccolroaz et al., 2020). The lake model sensitivity tests that we conduct here, align with observations, they suggest a lake temperature response of approx. 0.4 to 0.6 °C for every 1 °C annual or JJAS air temperature change (Fig. 4 and Fig. S16). Considering this relationship between air and lake temperature responses, the Zhao et al. (2021) calibration provides an excessively high estimation of the air temperature change during the Holocene, ranging from +10 to +15 °C. The estimation of temperature of the months above freezing (MAF-T) based on Otiniano et al. (2024; 2023) and Raberg et al. (2021) are more reasonable for southwestern Greenland. Therefore, since the three calibrations are equally valid for this region, we used the average of these calibrations to estimate the MAF-T in our study lakes (Otiniano et al., 2023, 2024; Raberg et al., 2021). We estimate the error as the standard deviation between these three calibrations. Root mean square error for each calibration is 1.97 °C (Raberg et al., 2021), 1.16 °C (Otiniano et al., 2023), and 1.2 °C (Otiniano et al., 2024).

370

The isomer ratio of 5- to 6-methyl brGDGTs (IR_{6Me}) ranges from 0.13 to 0.51 across all study lakes (Fig. S6–10). Previously, a non-thermal effect on lacustrine brGDGTs has been identified when IR_{6Me} is greater than 0.4 (Bauersachs et al., 2023; Novak et al., 2025). In Lake Pluto two samples during the Early Holocene and in Lake Bullet the bottom sample have IR_{6Me} greater than 0.4 (Fig. 2). Consequently, all samples with IR_{6Me} greater than 0.4 were excluded from temperature reconstructions. For Lake Gus, we employed the same screening for soil-derived brGDGTs (excluding all samples with <30% hexa-methylated GDGTs) as used in Cluett et al., (2023).




375

Figure 3: Simulations of fraction of ice cover, duration of ice free season (IFS), surface water temperature, IFS stratification, and IFS mixing period. (a) Simulations of the fraction of the lake covered by ice in study lakes. Line is the median value of the 30-year simulation under modern climate. (b) to (h) Sensitivity tests where annual temperature was changed, shown as anomalies relative to the simulation under modern climate. Each point denotes the median value of the 30-year simulation. Gray shading represents the

380

standard deviation for the 30 years simulated for Lake N3. Standard deviation is on the same order for other lakes. (b) to (c) Response of lake ice cover and mixing dynamics to changes in annual air temperature. (b) Onset date of ice free conditions. (c) IFS duration. (d) IFS stratification duration (stratification defined as mixing depth less than 75% of maximum lake depth). (e) IFS isothermal mixing duration (defined as mixing depth greater than 75% of maximum lake depth). (f) to (h) Relationship between IFS duration and lake surface water temperature averaged across (f) the entire year (g) the IFS and (h) JJAS.



385

Figure 4: Sensitivity of surface water (a–c) and average depth profile temperatures (d–f) to changes in annual air temperature. (a) and (d) annual, (b) and (e) IFS, (c) and (f) JJAS water temperature [°C]. Each point denotes the median value of the 30 year simulation.


390

Total brGDGT concentration in these lake sediments from southwestern Greenland is about 6-10 times higher than total isoprenoid GDGT (isoGDGT), with both showing a similar downcore pattern across all lakes, peaking in the Early Holocene and declining towards the Late Holocene (Fig. S10–S14). Caldarchaeol (GDGT-0) is the most abundant isoGDGT (Fig. S15a), followed by crenarchaeol (GDGT-4) and GDGT-1, whereas GDGT-2, GDGT-3, and the crenarchaeol isomer (GDGT-4') were below detection limit in many samples, restricting TEX₈₆-based temperature reconstruction (Schouten et al., 2002).

BrGDGTs in lake sediments can be allochthonous – produced in the catchment soil and transported to the lake via runoff erosion, and autochthonous – produced in the water column and/or surface sediment (Baxter et al., 2021; Tierney et al., 2012). BrGDGTs produced in these diverse environments can exhibit different responses to environmental conditions, thereby directly impacting temperature reconstruction (Acharya et al., 2023; Martin et al., 2020). Therefore, understanding the sources of brGDGTs is crucial before temperature reconstruction. We evaluated the sources of brGDGTs in southwestern Greenland lakes by comparing their distributions with those of high-latitude (>50 °N) soil, peat, and lake sediment samples (Raberg et al., 2022). BrGDGTs from southwestern Greenland lakes have different fractional abundance of hexa-, penta- and tetra-methylated brGDGTs than those in peat and soil samples but are similar to high-latitude lake surface sediment samples (Fig. S16), suggesting lacustrine origin. Furthermore, brGDGT IIIa, which is produced in higher relative abundance in lake water than in soils (Weber et al., 2018), is abundant in the southwestern Greenland lake sediments, similar to high-latitude lake surface sediments, and greater than in high-latitude soils (Fig. S15b). This corroborates the autochthonous production of brGDGTs in southwestern Greenland lakes, although a minor contribution from the surrounding catchment cannot be ruled out.

While several lacustrine brGDGT–temperature calibrations are available globally (Martínez-Sosa et al., 2021; Raberg et al., 2021; Zhao et al., 2023), regionally (Bauersachs et al., 2023; Dang et al., 2018; Otiniano et al., 2023, 2024; Russell et al., 2018), and site-specifically (Bittner et al., 2022; Feng et al., 2019; Zhao et al., 2021), only Raberg et al. (2021), Zhao et al. (2021), and Otiniano et al. (2023; 2024) are based on the Hopmans et al., (2016) HPLC method separating GDGTs isomers and explicitly consider Arctic seasonal climate. The Raberg et al. (2021) and Otiniano et al. (2023; 2024) calibrations estimate mean air temperature for the months above freezing (MAF-T) while Zhao et al. (2021) estimate epilimnion lake water temperature for the ice-free season (Ice-free season lake water temperature, IFS LWT) (Fig. S17). Given that Otiniano et al. (2024) provide the most recent calibration for Arctic lakes, using the extensive datasets synthesised from Otiniano et al. (2023), Raberg et al. (2021) and Zhao et al. (2021) and have the lowest prediction error (RMSE = 1.0 °C), we applied this calibration to estimate MAF-T in this study. This calibration yields a Holocene temperature range of 3.8–8.6 °C for the study lakes (Fig. 2). Although brGDGT distributions in Arctic lakes are primarily controlled by temperature (Otiniano et al., 2024; Raberg et al., 2021; Zhao et al., 2021), recent studies suggest they can also be influenced by site-specific factors, e.g., production period of brGDGTs and oxygenation (Harning et al., 2025; Raberg et al., 2025). We discuss the potential impact of these site-specific factors on reconstructed temperatures in Section 3.3.

The isomer ratio of 5- to 6-methyl brGDGTs (IR_{6Me}) ranges from 0.13 to 0.51 across all study lakes (Fig. S10–14). Previously, a non-thermal effect on lacustrine brGDGTs has been identified when IR_{6Me} exceeds 0.4 (Bauersachs et al., 2023; Novak et al., 2025). In Lake Pluto, two Early Holocene samples, and in Lake Bullet, the bottom sample has IR_{6Me}> 0.4 (Fig. S10–14). Consequently, all samples with IR_{6Me}> 0.4 were excluded from temperature reconstructions.



425 **Figure 2: Reconstructed mean air temperatures of the months above freezing (MAF-T) in (a) Lakes N3 and Pluto at 69 °N, (b) Lakes**
Gus (Cluett et al., 2023) and Rosaea at 67 °N, (c) Lakes Marshall and Bullet at 64 °N, and (d) Lake 578 at 61 °N (Schneider et al.,
2024). **For timeseries in a–c, the bold line indicates the median modelled age, and the light and dark shading represent the 95th**
percentile and interquartile age uncertainty, respectively. Numbers in parentheses next to site names refer to site IDs shown in Fig.
Gray line is the 1:1 line. Gray shading represents the standard deviation for the 30 years simulated for Lake N3. Standard deviation
430 **is on the same order for other lakes.**

1.

3.2 Holocene temperature variability

Reconstructed MAF-T shows both similarities and differences between time series, with differences not following a latitudinal
435 gradient. Reconstructed MAF-T varies from 5.95 to 9.68.3 °C for Pluto, 6.25.0 to 9.27.8 °C for N3, 4.85 to 8.47.7 °C for

Rosaea, 6.15.8 to 9.78.6 °C for Marshall, and 4.53.8 to 8.16.4 °C for Bullet, respectively (Fig. 5). ~~In lakes Pluto and Bullet, the temperature timeseries start at approx. 9 ka, and reveal a trend towards warm conditions (Fig. 5a).2).~~ In lakes Marshall and Rosaea, the temperature ~~timeseries~~ time series begin from approx. 9.5 ka, with relatively warm conditions indicated in Marshall Lake, while cold conditions prevail in Lake Rosaea (Fig. 5b2b and c). In Marshall Lake, the temperature decreased around 8.5 ka and then increased, peaking at approx. 7.1 ka.7 ka. In lakes Pluto and Bullet, the temperature time series begins at approx. 9 ka and show a trend towards warmer conditions, peaking in Pluto at 6 ka and in Bullet at 7.8 ka (Fig. 2). In Lake N3, the temperature ~~timeseries~~ time series begins at 8 ka ~~and reveals~~ with relatively warm conditions. In lakes N3, Marshall, and Pluto, the maximum temperature occurred between 7.5 ~~to~~ and 5 ka (Fig. 52). Immediately following the peak warmth, MAF-T in these three lakes reveals a gradual cooling trend until approx. 4.5 ka, followed by warm conditions until 3 ka, then gradual cooling. ~~In Bullet Lake, the temperature trends are muted compared to the other lakes, perhaps in part because the three calibrations yield contrasting trend (Fig. S13, black lines). The temperature maximum began at the same time as these other sites, around 8 ka, but lasted longer, until 2 ka, followed by cooling. Overall, the pattern of reconstructed MAF-T from Pluto, N3, Bullet, and Marshall lakes, spanning 5° latitude in southwestern Greenland, is similar to the annual temperature reconstruction from Greenland Ice Sheet Project 2 ice core (Fig. 6b), albeit with differences likely caused by spatio-temporal variation and resolution of the timeseries (Kobashi et al., 2017). Furthermore, the long-term temperature trends observed in N3, Pluto, and Marshall lakes is comparable to brGDGT~~ In Rosaea, the temperature maximum began at the same time as at these other sites, around 7.5 ka, but lasted longer until 4 ka, followed by cooling until 1 ka. In Bullet Lake, the maximum temperature occurred between 7.8 ka and 4 ka, followed by cooling until 3 ka, a warming at 2 ka, and then cooling towards the present (Fig. 2c).

Overall, the pattern of reconstructed MAF-T in our study lakes, spanning 5° latitude in southwestern Greenland, is similar to the brGDGTs-inferred MAF-T timeseries from Lake Gus (Cluett et al., 2023) and Lake 578 (Schneider et al., 2024) ~~(Freehette and de Vernal, 2009) (Figs. 5 and 6).~~, although the timeseries from Lake Gus reveals a peak of warmth two millennia earlier (from 9 to 7 ka) than other timeseries (7.5 to 5 ka) (Figs. 2 and 6). Next, we discuss the potential influence of site-specific factors on the seven brGDGT-inferred temperature time series in the region.

~~In contrast to the other timeseries, the reconstructed temperature in Lake Rosaea reveals consistent cool conditions from 9 to 7 ka, warming from 7 to 5 ka, and peak warmth between approx. 5 and 3 ka, lagging by two millennia peak warmth inferred from brGDGTs at six other southwestern Greenland lakes (Fig. 5b). This delayed peak warmth is synchronous with a biogenic silica record from Qipisarqo lake (Fig. 6i) (Kaplan et al., 2002) and pollen-based records from southern Greenland (not shown) (Freehette and de Vernal, 2009; Fredskild, 1973; Gajewski, 2015). Next, we discuss possible mechanisms explaining this temporal discrepancy in thermal maxima.~~




Figure 53.3 Possible influence of site-specific factors on reconstructed temperatures

Reconstructed mean air temperatures of the months above freezing (MAF-T) in (a) Lakes N3 and Pluto at 69°N, (b) Lakes Gus (Cluett et al., 2023) and Rosaea at 67°N, (c) Lakes Marshall and Bullet at 64°N, and (d) Lake 578 at 61°N (Schneider et al., 2024). Bold lines indicate average MAF-T inferred from three high-latitude lacustrine brGDGTs calibrations (Otiniano et al., 2023, 2024; Raberg et al., 2021), and shading represents standard deviation of these calibrations. Numbers in parentheses next to site names refer to site IDs shown in Fig. 1.


3.3 Possible mechanisms causing temporal differences in peak warmth among time series on western Greenland

BrGDGT-inferred MAF-T timeseries from southwestern Greenland reveal show a lower absolute value and at Lake Bullet. Moreover, the magnitude of the Holocene temperature change is lower at Lakes N3, Gus, and Bullet, and a delayed thermal maximum higher at Lake Rosaea, compared to than at Pluto, Lake 578, and Marshall. Additionally, the MAF-T time series

from Lake Gus shows peak warmth two millennia earlier than those from other study lakes. To interpret these variations, we explored and explore the potential mechanisms influencing impact of site-specific factors on reconstructed temperatures, we examined possible mechanisms that could influence the brGDGT-timeseries-inferred temperature time series from the seven lakes in southwestern Greenland.

3.3.1 Production period of brGDGTs

Lake ice regulates the availability of nutrients, light, and oxygen, thereby influencing the production period of bacteria producing brGDGTs. Therefore, the duration and timing of the ice-free season can influence the production period of brGDGTs (Loomis et al., 2014a; Miller et al., 2018). (Loomis et al., 2014; Miller et al., 2018) and, consequently, reconstructed temperatures. Simulations of all the study seven southwestern Greenland lakes under 30 years of modern climate suggest that all lakes exhibit < 2-week inter-lake variations in the timing of the onset of the ice-free season (Fig. 3a). Duration The duration of the ice-free season is approx. <20 days shorter in lakes Bullet and Marshall, probably likely due to their higher elevation, and colder temperatures and approx. 20 days longer in Lake 578 than, due to the lower elevation and higher temperature compared to lakes Pluto, N3, Gus, and Rosaea (Fig. 3a). Additionally, both the onset date and duration of the ice-free season respond similarly in all seven lakes to perturbations in annual air temperature (Figs. 3b–c). Annual, ice-free season and JJAS surface water temperature of all the lakes vary similarly with changes in duration of the ice-free season (Figs. 3d–e, 3f–h). These results suggest that the temperature response of the studied lakes to the same forcing is similar, and therefore, if brGDGTs are produced during the ice-free season, their temperature response should be similar among lakes if atmospheric temperature change is similar.






Figure-6

Figure 3: Simulations of fraction of ice cover, duration of ice-free season (IFS), surface water temperature, IFS stratification, and IFS mixing period. (a) Simulations of the fraction of the lake covered by ice in study lakes. Line is the median value from the 30-year simulation under the modern climate. (b) to (h) Sensitivity tests where annual temperature is changed, shown as anomalies relative to the simulation under modern climate. Each point denotes the median value of the 30-year simulation. Grey shading represents the standard deviation for the 30 years simulated for Lake N3. Standard deviation is on the same order as for other lakes. (b) to (e) Response of lake ice-cover and mixing dynamics to changes in annual air temperature. (b) Onset date of ice-free conditions. (c) IFS duration. (d) IFS stratification duration (stratification defined as mixing depth less than 75% of maximum lake depth). (e) IFS isothermal mixing duration (defined as mixing depth greater than 75% of maximum lake depth). (f) to (h) Relationship between IFS duration and lake surface water temperature averaged across (f) the entire year (g) the IFS and (h) JJAS. + Holocene temperature history from southwestern Greenland. (a) Smaller than present mountain glaciers (Schweinsberg et al., 2017, 2019), (b) Annual air temperature [°C] inferred from gas fractionation from Greenland Ice Sheet Project 2 ice core, with the black line

~~indicating elevation corrected values and the grey line indicating elevation uncorrected values. (Axford et al., 2021; Kobashi et al., 2017), (e) chironomid-inferred July air temperature [°C] from North Lake (Axford et al., 2013), brGDGT-inferred mean air temperatures for months above freezing (MAF-T) [°C] from (d) lakes N3 and Pluto (This study), (e) Lake Gus (Cluett et al., 2023), (f) lakes Marshall and Bullet (This study), brGDGT-inferred MAF-T [°C] from (g) Lake 578 (Schneider et al., 2024), (h) Lake Rosaea (This study), (i) Biogenic silica as an indicator of temperature from Lake Qipisarqo (Kaplan et al., 2002), Dinoecyst-inferred sea surface temperatures (SST) [°C] from (j) Disko Bugt core MSM343300 (Ouellet-Bernier et al., 2014), and (k) Baffin Bay core CC70 (Gibb et al., 2015) (l) Greenland Ice Sheet surface area ($\times 10^{12}$) [m^2] (Larsen et al., 2015) and (m) 21st June insolation at 65°N [W m^{-2}] (Laskar et al., 2004). Numbers in parentheses next to site name refer to site IDs shown in Fig. 1.~~

For some lakes, simulations under 10 °C lower magnitude of temperature change than modern resulted in year-round ice cover; therefore, these simulations were not plotted.

3.3.2 Sensitivity of the lake system to air temperature change

The sensitivity of the study lakes to air temperature may vary due to differences in morphological characteristics and thermodynamics, potentially leading to variations in the reconstructed Holocene MAF-T (Dokulil et al., 2021; Kettle et al., 2004; Piccolroaz et al., 2020). However, lake model simulations indicate that the annual, IFS, and JJAS surface water temperatures of the seven lakes show similar responses to changes in annual air temperature (Fig. 4). The lake model also demonstrates a comparable, but slightly muted, response to JJAS air temperature perturbations (Fig. S20). These findings suggest that, despite morphological differences, the surface water temperatures of the studied lakes are equally sensitive to variations in air temperature. The lake model sensitivity tests conducted here indicate a lake temperature response of approximately 0.4 to 0.6 °C for every 1 °C change in annual or JJAS air temperature, aligning with observations in Arctic lakes (Dokulil et al., 2021; Kettle et al., 2004; Piccolroaz et al., 2020). Since brGDGTs in lakes can be produced in surface waters, water columns, or sediment (Buckles et al., 2014; Yao et al., 2020; Zhao et al., 2021), we also examined the response of the average depth profile temperature of the lakes to perturbations in annual and JJAS air temperature (Fig. 4d–f and Fig. S20d–f). Similar to surface temperature, all lakes display comparable responses, except lakes N3 and 578, which show a slightly muted response, especially in the JJAS temperature profile, likely due to their greater maximum depths. Therefore, the increased depth of lakes N3 and 578 may have resulted in a slightly lower amplitude (roughly < 0.5 °C) of Holocene temperature variations compared to the other lakes (Fig. 2). Nonetheless, about 1.5 °C lower absolute temperatures in Bullet Lake, and an earlier thermal maximum in Gus compared to the other lakes cannot be attributed to sensitivity to air temperature forcing.




Figure 4: Sensitivity of surface water (a–c) and average depth profile temperatures (d–f) to changes in annual air temperature. (a) and (d) annual, (b) and (e) IFS, (c) and (f) JJAS water temperature [°C]. Each point denotes the median value of the 30-year simulation. Similar to the onset and duration of the ice-free season, the response of lake water temperature during the ice-free season may vary among study lakes due to their different morphological characteristics and thermodynamics. Yet, lake model simulation reveals that the annual, IFS, and JJAS surface water temperatures of the seven lakes exhibit similar responses to changes in annual air temperature (Fig. 4). The lake model exhibited a similar, but slightly muted, response to JJAS air temperature perturbations (Fig. S16). This result suggests that despite morphological differences, the surface water temperature of the studied lakes, is equally sensitive to changes in air temperature. Since brGDGTs in lakes can be produced in surface waters, water columns, and/or sediment (Acharya et al., 2023; Buckles et al., 2014; Yao et al., 2020; Zhao et al., 2021), we also examined the response of the average depth profile temperature of the lakes to perturbations in annual and JJAS air temperature (Fig. 4 d–f and Fig. S16 d–f). As with the surface temperature, all the lakes exhibit similar responses, except lakes N3 and 578, which has a slightly muted response compared other lakes, especially in the JJA temperature profile, probably caused by greater maximum depth compared to other lakes. Therefore, the greater depth of lakes N3 and 578 may have led to slightly lower amplitude (approx. $<1^{\circ}\text{C}$) Holocene temperature variations compared to the other lakes (Fig. 5). However,

several millennia of delayed thermal maximum in Lake Rosaea and approx. 2 °C lower absolute temperatures in Bullet Lake compared to other lakes cannot be explained by sensitivity to air temperature forcing.

3.3.3 North-South pattern in timing of temperature maxima

To explore the north-south pattern in the timing of temperature maxima, we compared all the available brGDGT-based temperature reconstructions from southwestern Greenland including from Lake Gus (Fig. 1; point 11) (Cluett et al., 2023) and Lake 578 (Fig. 1; point 13) (Schneider et al., 2024). The grey line is the 1:1 line. Grey shading represents the standard deviation for the 30 years simulated for Lake N3. Standard deviation is on the same order as for other lakes. For some lakes, simulations under 10 °C lower magnitude of temperature change than modern resulted in year-round ice cover; therefore, these simulations were not plotted.

3.3.4 Anoxia

Aquatically produced brGDGTs in the study lakes may be influenced by anoxic or suboxic conditions, potentially biasing temperature reconstructions (Acharya et al., 2023; Harning et al., 2025; Raberg et al., 2025). To assess this potential impact, we used the caldarchaeol-to-crenarchaeal ratio (Cald/Cren), a proxy for suboxic conditions in the water column and/or lake sediments where caldarchaeol is produced (Baxter et al., 2021; Blaga et al., 2009; Sinninghe Damsté et al., 2022). The Cald/Cren ratio ranged from 1 to 857 in all study lakes, with generally low values in lakes Marshall and Rosaea and high values in lakes Bullet, N3 and Pluto (Fig. 5). The ratio was higher from 10 to 7 ka in lakes Rosaea, Bullet, Marshall, and Pluto, suggesting reduced oxygen availability, while afterwards, it was lower, indicating higher oxygen availability. Similarly, the Cald/Cren ratio was higher in Lake N3 from 8 to 5 ka, suggesting lower oxygen availability before decreasing to low levels.


Correlations between Cald/Cren and reconstructed temperatures were weak or non-significant in Lakes N3 and Marshall throughout the Holocene (Table S4), indicating that suboxic conditions had little influence on reconstructed temperatures. In contrast, significant negative correlations were present in Lakes Rosaea, Bullet, and Pluto (Table S4), suggesting potential redox-related effects on reconstructed temperatures during periods of inferred suboxia (10–7 ka). Similar to our study sites, Lake 578 contains high Cald/Cren values during the Early Holocene (Schneider et al., 2024). This was interpreted to indicate anoxia during the ice-covered period and/or early in the ice-free season, and likely did not influence the brGDGT-inferred temperatures due to mixing during the ice-free season (Schneider et al., 2024). Together these seven lakes cover agreement between the latitudinal gradient Lake 578 brGDGT-inferred temperature with temperature from ~61°N. Other proxy time series suggests that anoxia is unlikely to 69°N (Table 1). If we assume a north-south pattern in the timing main factor controlling brGDGT distributions in Lake 578.

Furthermore, the lake model sensitivity tests under Holocene climate conditions (± 5 °C) indicate that stratification occurs early in the ice-free season in all study lakes but does not persist throughout the entire ice-free period (Fig. 3 and magnitude Fig.

S18). As ice-free season duration increases, the duration of the isothermal mixing increases and the duration of stratification remains constant. This is due to the relatively shallow depths and low dissolved-solids concentrations in these lakes, which make them prone to mixing by solar heating and wind. Such mixing during summer is observed today in Arctic lakes with similar size and chemical composition (Antoniades et al., 2024; Klanten et al., 2023, 2024; Lindborg et al., 2016). Due to the possible influence of oxygenation on brGDGT-distributions, we cannot completely rule out the effects of anoxia/suboxia and possible cold-bias on MAF-T in lakes Pluto, Bullet, Rosaea, Marshall, and N3 during the periods high Cald/Cren from 10 to 7 ka. Yet, based on the lake model sensitivity tests and evidence from Lake 578, we suggest that the impact of anoxia/suboxia on brGDGTs-inferred MAF-T is minimal.

We next examine brGDGT compounds that have been interpreted in some settings to indicate suboxic conditions. Although the ratio of 5-methyl hexa-methylated and penta-methylated brGDGTs with no cyclopentane rings (HP5) index and the fractional abundance of brGDGT IIIa (%IIIa) are suggested as indicators of anoxia in large and permanently stratified lakes (Weber et al., 2018; Yao et al., 2020), both 5-methyl hexa-methylated and penta-methylated brGDGTs exhibit a strong temperature dependence in Arctic lakes (Martínez-Sosa et al., 2021; Otiniano et al., 2023, 2024; Raberg et al., 2021) and can also be influenced by soil-derived input (Acharya et al., 2023; Martin et al., 2019). In Bullet Lake, Cald/Cren showed a non-significant correlation with both HP5 and %IIIa (Table S4). In Lake Rosaea, significant positive correlations were observed between Cald/Cren and both HP5 and %IIIa (Table S4). In Marshall and Pluto lakes, Cald/Cren was significantly positively correlated with HP5 but not with %IIIa. In Lake N3, no significant correlation was found with HP5, whereas a significant negative correlation was observed with %IIIa. As isoGDGTs and brGDGTs are produced by different microbial communities: brGDGTs by bacteria and isoGDGTs by archaea, these variable relationships may reflect seasonal and/or spatial differences in the niche of these microbial communities, which are subject to future study. Notably, in all our study lakes, both HP5 and %IIIa show a significant negative relationship with reconstructed temperature, aligning with previous studies that these proxies have a strong temperature dependence in Arctic settings.

Similarly, brGDGT IIIa'' has been suggested to be influenced by anoxia in a permanently stratified lake (Weber et al., 2015, 2018), but it shows no significant correlation with oxygen in Arctic lakes (Raberg et al., 2025). Moreover, brGDGT IIIa'' is present only in lakes Bullet and Rosaea, and the Holocene trends between brGDGT IIIa'' and cald/cren in these lakes are decoupled (Fig. S12 and S14). This evidence suggests that brGDGT IIIa'' is also not a reliable indicator of anoxia in these lakes.



610 **Figure 5: Variations of the ratio of caldarchaeol to crenarchaeol (Cald/Cren) in the study lakes.**

3.3.4 North-South pattern in timing of temperature maxima

A north-south gradient in the magnitude and timing of the Holocene thermal maximum has been suggested in Greenland, with a greater magnitude and earlier thermal maximum in the north and a lower magnitude and later thermal maximum in the south, caused by Arctic Amplification (Axford et al., 2021; Briner et al., 2016; Kaufman, 2004). If this pattern holds true in southwestern Greenland, the lakes at the highest latitudes in the our study transect (Pluto and N3) should exhibit higher magnitude magnitudes and earlier temperature maxima compared to than lakes at lower latitudes (Rosaea and Gus, 67°N; Marshall and Bullet, 64°N; and 578, 61°N). However, the timing of the temperature maximum at the southernmost site, Lake 578 (8 to 5 ka)), is similar to that slightly earlier or within age uncertainty of lakes that of N3 and Pluto at 69 °N and Lake

Marshall at 64 °N, occurring between *ca.* 7.5 to 5 ka (Fig. 52). Moreover, Lake Bullet, situated at a similar latitude to that of Lake Marshall, exhibits a prolonged temperature maximum between *ca.* 7.8 to 24 ka. The ~~time-series~~ timeseries that shows a ~~delayed~~ ~~an earlier~~ temperature maximum, between ~~ca. 59~~ and ~~36~~ ka, occurs in the middle of the transect: at Lake ~~Rosaea~~ Gus (67°N). This ~~delayed~~ ~~earlier~~ timing in temperature maximum at Lake ~~Rosaea~~ Gus is not observed in the timeseries from Lake ~~Gus~~ Rosaea, at the same latitude, where peak temperatures occurred between *ca.* ~~9 to 7.5~~ and 3 ka (Cluett et al., 2023). A similar ~~timing of the~~ thermal maximum ~~timing~~ among six of the seven brGDGT ~~timeseries~~ time series from southwestern Greenland suggests that Arctic Amplification was *likely* not the primary mechanism setting the timing of peak warmth in this region. ~~In addition, Lake Rosaea must be primarily responding to different primary forcing mechanisms than the other lakes.~~

3.3.45 Ice- sheet and ocean proximity

Lakes Bullet and Marshall are <2 km ~~distance to from~~ the margin of the modern Greenland Ice Sheet, whereas lakes N3, 578, and Pluto are > 10 km, and lakes Gus and Rosaea are > 100 km away (Table 1). Deglaciation from lakes Gus, Rosaea, Marshall, Bullet, and 578 occurred before 10 ka (Leger et al., 2024). The margin of the Greenland Ice Sheet was relatively stable from 9 to 6 ka near lakes Marshall, Bullet, and 578, while it was receding inland from lakes Rosaea and Gus. Since 9 ka, the margin of the ice sheet was always at least 50 km inland from lakes Rosaea and Gus. By contrast, lakes N3 and Pluto deglaciated at ~8 ka (Leger et al., 2024; Young et al., 2013), and the margin of the Greenland Ice Sheet was <3 km aerial distance from these lakes until 7.5 ka. The margin of the southwestern Greenland Ice Sheet retreated inland of its present-day margin around 6 ka and returned to its present margin around 1850 AD (Leger et al., 2024), although the little available evidence suggests the retreat was <10 km (Briner et al., 2014; Cronauer et al., 2016; Lesnek et al., 2020). This suggests that lakes Bullet, Marshall, Pluto, and N3 could have been significantly influenced by the proximity of the ice sheet during the Holocene, while Lakes Rosaea and Gus are minimally influenced, and Lake 578 might be moderately influenced. Thus, the fact that peak warmth occurred after 8 ka, following ice sheet retreat from Lakes Marshall, Bullet, N3, and Pluto, suggests that the ice sheet may have kept local conditions cold. ~~However, Marshall~~ In contrast, peak warmth at Lake ~~experienced moderately warm conditions at 9 ka, a period that most of these other timeseries do not capture, suggesting that ice sheet proximity may not have been~~ Gus occurred earlier, consistent with its more distal position relative to the ~~primary factor influencing MAF T at these lakes.~~ ice margin. Lake Rosaea ~~is an obvious outlier in this ice sheet proximity pattern: it was far from the ice sheet for most of the Holocene, so should be minimally influenced by katabatic winds, but has a low magnitude of temperature change and the~~ latest, however, shows a later thermal maximum ~~of all the records. Thus, temperature~~ than at Lake ~~Rosaea is likely not influenced by its distance from the Greenland Ice Sheet~~ Gus, despite its distal position relative to the ice margin, indicating that additional regional factors contributed to relatively cold conditions between ~9 and 7.5 ka.

Ocean conditions influence coastal climate by affecting ocean-atmosphere heat transfer (Bashmachnikov et al., 2023). A warmer ocean releases heat into the atmosphere, resulting in warmer coastal climates, ~~and vice versa with cold; conversely, a~~ colder ocean ~~conditions~~ leads to cooler coastal climates. Lakes Bullet and Marshall are ≥100 km, and lakes 578 and Gus are

>60 km from the open ocean (Table 1). Thus, the ocean influence on these lakes is likely minimal. Lakes N3, Pluto, and Rosaea are situated near the ocean (<3 km). Modern wind directions at Lake Rosaea are predominantly southerly, meaning that most ~~airmasses~~air masses at Lake Rosaea come directly from the Labrador Sea and Davis Strait (Fig. 7a6a). About 20% of winds at Lake Rosaea are northerly, meaning they pass over Baffin Bay prior to arriving at Lake Rosaea. In contrast, at lakes Pluto and

655 N3, wind direction is more variable, meaning that half of the ~~airmasses originate from land and half from Disko Bugt (Fig. 7a).~~The pattern of reconstructed lake water temperature from Lake Rosaea is similar to summer sea surface temperature (SST) timeseries from Disko Bugt and Baffin Bay (Fig. 6j, k), as well as other North Atlantic SST time series, which reach maxima in the Middle and Late Holocene (Gibb et al., 2015; Hansen et al., 2020; Ouellet-Bernier et al., 2014; Saini et al., 2022). This similarity implies that temperature at Lake Rosaea is more strongly influenced by oceanic conditions via ocean-atmosphere




660 heat exchange than the other lakes in this transect. Summer SSTs in eastern Baffin Bay were cold during the Early Holocene, probably caused by meltwater from the rapidly retreating Greenland Ice Sheet and a weaker western Greenland currentair masses originate from land and half from Disko Bugt (Fig. 6a). This suggests that Lake Rosaea is strongly influenced by the influenced by oceanic conditions via ocean-atmosphere heat exchange, while lakes N3 and Pluto are moderately influenced. During the Early Holocene, summer sea surface temperatures (SST) in Disko Bugt and Baffin Bay (Fig. 7j, k), as well as other

665 North Atlantic SST time series, were cold, likely due to meltwater from the rapidly retreating Greenland Ice Sheet and a weaker western Greenland Current (~~Gibb et al., 2015; Hansen et al., 2020~~).(Gibb et al., 2015; Hansen et al., 2020). SSTs rose during the ~~middle-Middle~~ Holocene ~~related~~due to decreased meltwater from the Greenland Ice Sheet and increased influence of warm Atlantic water masses (Ouellet-Bernier et al., 2014; Perner et al., 2013). The ~~pattern of reconstructed temperature pattern~~MAF-T in Lake Rosaea is similar to the SST time series from ~~Lakes Pluto~~Disko Bugt and ~~N3~~exhibits a closer resemblance to the

670 ~~lakes~~Baffin Bay. This suggests that ~~are far from~~cold SSTs in the ~~ocean~~. These lakes may have been less affected by Holocene ~~Ocean~~region likely led to cold conditions ~~than~~in Lake Rosaea ~~due to local topography, including proximity from 10 to the ice sheet and their position in interior Disko Bugt.~~

675 3.3.5 Topographic conditions

Steep topography surrounding a lake can shade the lake, influencing the direct short-wave radiation reaching lake surface, thereby regulating the lake surface water temperature. To investigate the mechanisms for the differences in absolute temperature and magnitude of Holocene temperature variations in the five study lakes, we examined the topographic conditions around each lake. Shading calculations show that of the five lakes, Pluto Lake is the least shaded by surrounding topography and Bullet Lake is the most shaded (15% less direct shortwave radiation than Pluto Lake) (Fig. 7b6b). Lakes N3 and Marshall are approx. 5% more shaded than Pluto Lake, whereas Lake Rosaea is 10% more shaded than Pluto Lake (Fig. 7b6b). This means that the topography surrounding lakes N3, Marshall, Rosaea, and Bullet ~~cause~~~~causes~~ them to receive correspondingly less solar radiation compared to Pluto, with Bullet receiving the least. We assess the impact of shading on lake water thermodynamics by perturbing the magnitude of incoming shortwave radiation in the lake model. These simulations suggest that a 10% decrease in incoming ~~short-wave~~~~shortwave~~ radiation can suppress the mean ~~ice-free season~~ lake surface temperature during the ice-free season by approx. 1.5 °C and a 25% decrease by 4 °C (Fig. 7c). Therefore, we suggest that the slight (~~-1-2 °C~~) difference in absolute temperature values between the lakes Pluto, N3, Marshall and Rosaea could be related to local topography and variations in incoming shortwave radiation to the lakes. Approx. approx. 2 °C lower absolute temperature and approx. 0.5 °C lower magnitude of temperature change in Bullet Lake may be due to the lowest incoming shortwave radiation, compared to the other study lakes. ~~There may also be other factors causing differences between these time series, but we believe we have assessed the primary mechanisms here.~~






Figure 76: (a) Wind strength and direction in the study lakes from April to September, based on six-hourly ERA5 data from 1994 to 2024. (b) Percentage change in average summer (April to September) direct radiation on study lakes relative to Pluto Lake, which is the least shaded of the five. (c) Sensitivity of average surface water temperature to changes in incoming shortwave radiation. Each point denotes the median value of the 30-year simulation. Gray shading represents the standard deviation for the 30 years simulated for Lake N3. Standard deviation is on the same order as for other lakes.

4 Holocene temperature history of southwestern Greenland

The majority of brGDGT-inferred MAF-T timeseries from southwestern Greenland (N3, Pluto, Marshall, Bullet and 578) document peak warmth in the Middle Holocene, between ca. 7 and 5 ka, following maximum summer insolation. Timeseries from Lake Gus reveal peak warmth between ca. 9 to 6 ka, earlier than many other time series. An earlier deglaciation at Lake Gus (before ca. 10 ka), and minimal influence from the Greenland Ice Sheet and coast during the Holocene, likely have caused

earlier warmth, Rosaea, and 578) document peak warmth in the Middle Holocene, between *ca.* 7.5 and 5 ka, following maximum summer insolation (Laskar et al., 2004). The timeseries from Lake Marshall shows warmth at *ca.* 9 ka, followed by cooling at *ca.* 8.5 ka. However, 25 samples from the Early Holocene in Lake Gus were removed due to evidence of high input of soil-derived brGDGTs (Cluett et al., 2023), so it is possible that other samples in that section have some minor influence of soil-derived brGDGTs, causing elevated inferred MAF-T. The timeseries from Lake Marshall shows warmth at *ca.* 9 ka, followed by cooling *ca.* 8 ka and subsequent warming, similar to the regional ice-sheet retreat rate reconstructions (Lesnek et al., 2020), suggesting this timeseries captured Early Holocene warm conditions not captured in most lake records, since they start later. Immediately after peak warmth, temperature reconstructions in all study lakes show a gradual cooling trend towards the Late Holocene. This pattern of temperature change can also be inferred from alpine glacier reconstructions from Immediately after peak warmth, reconstructed temperature in all these lakes shows a gradual cooling trend towards the Late Holocene. Overall, the pattern of reconstructed MAF-T from these lakes in southwestern Greenland is similar to the annual temperature reconstruction from the Greenland Ice Sheet Project 2 ice core (Fig. 7b), albeit with differences likely due to spatio-temporal variation and the resolution of the timeseries (Kobashi et al., 2017). Furthermore, the pattern of temperature change documented in southwestern Greenland can also be inferred from alpine glacier reconstructions throughout southwestern Greenland, which experienced high retreat rates or completely disappeared during the Early Holocene and reappeared during the Late Holocene (Fig. 6a7a) (Larocca et al., 2020; Larsen et al., 2017; Lasher et al., 2020; Schweinsberg et al., 2017, 2019). Furthermore, Holocene vegetation in southwestern Greenland indicates the warmest summer conditions in the Holocene between 7.2 and 6.3 ka, followed by cooling (Fredskild, 1983), a pattern similar to timeseries from lakes N3, Pluto, Rosaea, Marshall, and Bullet, and 578 (Schneider et al., 2024). Similarly, the occurrence of warmth-demanding ostracodes in lake sediments at the head of Kangerlussuaq (inland from Lake Gus) indicates peak Holocene summer temperatures from *c.* 7.0 to 6.5 ka (Bennike et al., 2010). The Early Holocene peak warmth and the subsequent cooler Late Holocene documented by the most of these new brGDGTs-derived time series and other summer-sensitive proxy time series in southwestern Greenland align with the temperature pattern observed in many proxy time series from Greenland and Arctic Canada, albeit with spatial and temporal variability (Axford et al., 2021; Briner et al., 2016; Kaufman, 2004; Larocca and Axford, 2022)(Axford et al., 2021; Briner et al., 2016; Kaufman, 2004; Larocca and Axford, 2022). Similarly, the biogenic silica (BSi) record from Lake Qipisarqo reveals a thermal maximum between 7 and 3 ka, akin to the brGDGTs-inferred MAF-T timeseries from southwestern Greenland (Andresen et al., 2004).

In contrast, time series from Lake Gus reveal a peak in warmth between *ca.* 9 and 6 ka, earlier than in many other time series. An earlier deglaciation at Lake Gus (before *ca.* 10 ka) and minimal influence from the Greenland Ice Sheet and coast during the Holocene likely caused earlier warmth. However, ~25 samples from the Early Holocene in Lake Gus were removed due to evidence of high input of soil-derived brGDGTs (Cluett et al., 2023), so it is possible that other samples in that section Lake Rosaea experienced a thermal maximum between approx. 5 to 3 ka, coinciding with nearby SST reconstructions (Fig. 4j, k) (Gibb et al., 2015; Ouellet-Bernier et al., 2014). Similarly, the biogenic silica (BSi) record from Lake Qipisarqo (Fig. 3i),

pollen-based reconstructions from Lakes Comarum Cø, and Kløft Sø, all coastal sites that are influenced by the East Greenland Current, reveal a delayed thermal maximum, after 6 ka, in accord with marine temperature timeseries (Fig. 3j, k) (Andresen et al., 2004; Fredskild, 1973; Gajewski, 2015; Kaplan et al., 2002). This suggests the influence of ocean conditions on coastal temperatures via ocean-atmosphere heat exchange.

exerted a minor influence on the inferred MAF-T. Although deglaciation at Lake Rosaea also occurred earlier (before 10 ka), the MAF-T reconstruction from Lake Rosaea does not show the Early Holocene warm signal observed at Lake Gus. This is likely due to a strong marine influence at Lake Rosaea, where air masses arrive directly from the Labrador Sea and Davis Strait and cold SST during 10 to 7 ka. In contrast to brGDGT-inferred MAF-T timeseries from southwestern Greenland, pollen-based reconstructions from lakes Comarum Cø and Kløft Sø in southern Greenland show a delayed thermal maximum after 6 ka (Fredskild, 1973; Gajewski, 2015). This could be due to marine influence and/or vegetation lags (Axford et al., 2021).

The magnitude of peak warmth in brGDGT-inferred MAF-T ~~showshows~~ both similarities and differences between time series. BrGDGT-inferred MAF-T timeseries from lakes Pluto, N3, ~~Gus, Rosaea,~~ and 578 ~~each~~ show on average $1.4 \pm 0 \pm 0.5.2$ °C higher temperatures during the peak warmth (from 7.5 to 5 ka) relative to ~~average between 2 ka to the present. Marshall Lake shows approx. 0.7 °C higher temperatures compared to the other lakes. The relatively warmer temperature in 1994–2024.~~ Although the timing of peak warmth in Lake Gus occurred earlier (from 9 to 6 ka), the magnitude of warming was similar to that observed at lakes Pluto, N3, and 578 (1.4 °C). Peak warmth at Rosaea, Bullet, and Marshall was approx. 0.5, 2, and 4 °C, respectively, higher than from lakes Pluto, N3, and 578. Warmer temperatures in Lake Rosaea are likely due to ocean influence, while in Lakes Marshall ~~Lake is~~ and Bullet, they are likely due to ~~its~~their higher ~~elevation~~elevations, which would have ~~caused~~led to a higher rate of warming (Pepin et al., 2015). ~~Lake Bullet shows relatively stable temperatures over the Holocene, perhaps due to lowest incoming shortwave radiation, compared to the other study lakes, although uniquely contrasting trends among the three brGDGT temperature calibrations for the Lake Bullet time series may also explain the relative lack of trend. Overall, the magnitude of peak warmth in lakes Pluto, N3, Gus, Marshall, and Bullet are lower than the chironomid-inferred July air temperature from North Lake showing 2–3 °C warmer conditions between 7 and 5 ka relative to average between 2 ka to the present (Axford et al., 2013). This difference in magnitude is likely related to (Pepin et al., 2015). However, it is also possible that modern temperatures at these lakes are not well represented, as no direct observations are available. We therefore relied on ERA5 reanalysis data, which have a relatively coarse grid resolution ($\sim 25 \times 25$ km) and may not fully capture local conditions due to Greenland's complex topography. Overall, the magnitude of peak warmth in lakes Pluto, N3, and 578 is similar to the ice sheet elevation change corrected annual air temperature [°C] inferred from gas fractionation from GISP2 ice core demonstrating ~ 1 °C warmer conditions from 7.5 to 5 ka relative to the averages of 1988–2015 (Axford et al., 2021; Kobashi et al., 2017). Slightly cooler temperatures during the Holocene peak warmth in mean annual air temperature at GISP2 compared to MAF-T in lakes Pluto, N3, Gus, Rosaea, and 578 could be due to the seasonality of the reconstructed temperature. Moreover, the magnitude of peak warmth in lakes Pluto, N3, and 578 is lower than the chironomid-inferred July air temperature~~

770 from North Lake, indicating 2–3°C warmer conditions between 7 and 5 ka relative to the present (Axford et al., 2013). This
difference in magnitude is likely related to the seasonality of reconstructed temperature (July for Chironomids, months above
freezing for brGDGTs) (Eggermont and Heiri, 2012). ~~The strongest solar radiation in July caused the higher magnitude of
warming compared to the MAF-T. Further, the magnitude of peak warmth in lakes Pluto, N3, Gus, Rosaea, and 578 is lower
than the ice sheet elevation change corrected annual air temperature [°C] inferred from gas fractionation from GISP2 ice core
demonstrating —1.3°C warmer conditions from 7 to 5 ka relative to the 2 to 0 ka average (Axford et al., 2021; Kobashi et al.,
2017). This is surprising as the Holocene annual air temperature show a lower warming than the MAF-T (Axford et al., 2021;
Briner et al., 2016). The higher magnitude of Holocene peak warmth in mean annual air temperate in GISP2 and MAF-T in
lakes Pluto, N3, Gus, Rosaea, and 578 could be due higher elevation of GISP2 causing higher summer warming during peak
warmth, than the coastal Greenland leading to overall higher magnitude of mean annual air temperature change in GISP2.~~
780 Further investigation of proxy time series considering seasonality of temperature change and covering a wider spatial range
~~across over the Greenland may help to~~The strongest solar radiation in July led to a greater magnitude of warming than in the
MAF-T. Further investigation of proxy time series, accounting for temperature seasonality and covering a wider spatial range
across Greenland, may help reconcile these differences.

Although there are differences in timing and magnitude of peak warmth in brGDGT-inferred MAF-T timeseries for
785 southwestern Greenland, these timeseries do not follow a latitudinal pattern. This is surprising, considering the pronounced
warming in higher latitudes compared to low latitudes due to local feedbacks and remote teleconnections observed during the
past century (England et al., 2021; Sweeney et al., 2023) and during the Holocene (Axford et al., 2021; Briner et al., 2016;
Kaufman, 2004). The absence of Arctic Amplification in southwestern Greenland brGDGT-inferred from MAF-T timeseries
in the past 9 ka could be because of several factors: (1) the latitudinal gradient across the study sites may be too small to
790 observe temperature differences using this proxy; (2) the temporal resolution of the time series may be too coarse to detect
subtle shifts in the timing of the thermal maximum; or (3) other local or regional influences, including strong ocean and
atmosphere northward heat transport (Dufour et al., 2016) and the absence of Holocene summer sea ice in this region (~~Gibb et
al., 2015~~)(Gibb et al., 2015), which may have been overprinted the expected signal from Arctic amplification. Future high-
resolution proxy timeseries covering a larger latitudinal range and using the same proxy, reflecting similar seasonality, could
795 help test this hypothesis.




Figure 7: Holocene temperature history from southwestern Greenland. (a) Smaller-than-present mountain glaciers (Schweinsberg et al., 2017, 2019), (b) Annual air temperature [°C] inferred from gas fractionation from Greenland Ice Sheet Project 2 ice core, with the black line indicating elevation-corrected values and the grey line indicating elevation-uncorrected values. (Axford et al., 2021; Kobashi et al., 2017), (c) chironomid-inferred July air temperature [°C] from North Lake (Axford et al., 2013), brGDGT-inferred mean air temperatures for months above freezing (MAF-T) [°C] from (d) lakes N3 and Pluto (This study), (e) Lake Gus (Cluett et

al., 2023), (f) lakes Marshall and Bullet (This study), brGDGT-inferred MAF-T [°C] from (g) Lake 578 (Schneider et al., 2024), (h) Lake Rosaea (This study), (i) Biogenic silica as an indicator of temperature from Lake Oipisargo (Kaplan et al., 2002), Dinocyst-inferred sea surface temperatures (SST) [°C] from (j) Disko Bugt core MSM343300 (Ouellet-Bernier et al., 2014), and (k) Baffin Bay core CC70 (Gibb et al., 2015), (l) Greenland Ice Sheet surface area ($\times 10^{12}$) [m²] (Larsen et al., 2015) and (m) 21st June insolation at 65 °N [W m⁻²] (Laskar et al., 2004). **5 Summary and conclusions**

We compareFor timeseries in d, e, f, and h the bold line indicates the median modelled age, and the light and dark shading represent the 95th percentile and interquartile age uncertainty, respectively. Numbers in parentheses next to the site name refer to the site IDs shown in Fig. 1.

5 Conclusions

We generated five new (from Pluto, N3, Rosaea, Marshall, and Bullet Lake) and compiled two published (from Gus Lake and Lake 578) brGDGT-inferred MAF-T time series spanning the past 9 ka from southwestern Greenland. The timeseries from lakes Pluto, N3, Marshall, Bullet, and 578 have similar trends, withLake model sensitivity tests suggest minimal intra-lake variation in both the seasonal production window of brGDGTs and the sensitivity of the studied lakes to air temperature changes. Further, anoxic/suboxic conditions had a minimal impact on the brGDGT distributions in these lakes. These results suggest that regional climate is the primary mechanism influencing these time series. The time series from all the lakes except Gus showed a temperature maxima between ca. 7 and 5 ka, 1.0 ± 0.5 °C higher temperatures relative to the 2 to 0 ka average, and then 7.5 and 5 ka, followed by gradual cooling towards the Late Holocene, similar to the many other regional timeseriestime series, following maximum annual insolation at 65 °N. The timeseries from Lake Gus reveals peak warmth between ca. 9 to 6 ka, earlier than many other time series, likely due to minimal influence from the coast and the Greenland Ice Sheet. TheIn addition, the time series from Lake Marshall suggests a warm Early Holocene, cooling around 8.5 ka, and subsequent warming, which is similar to the few available proxy records of Early Holocene temperature inat the ice-free margins of Greenland (Lesnek et al., 2020).

In contrast to the other timeseries, Lake Rosaea reveals a temperature maximum between ca. 5 and 3 ka, lagging peak warmth in other timeseries by two millennia. Lake model simulations suggest that proxy systematics, i.e., the production season of brGDGTs and the sensitivity of each lake system to climate changes, likely did not cause these temporal differences. Moreover, we do not observe a north-south trend in the timing of the thermal maximum in southwestern Greenland. Instead, the timing of peak warmth seems to be related to regional scale factors: proximity to the ice sheet and ocean. Lake Rosaea is situated at the coast, with prevailing southerly winds from the ocean. Holocene temperature change at Lake Rosaea is synchronous with regional marine timeseries that indicate weaker West Greenland Current and cooler summer ocean conditions during the Early and middle Holocene caused by melt water from Greenland Ice Sheet (Axford et al., 2021; Gibb et al., 2015; Hansen et al., 2020; Ouellet-Bernier et al., 2014). Temperature at Lake Rosaea rose during the middle Holocene when SSTs increased, due to decreased melt water flow from the Greenland Ice Sheet, which reached its minimum extent around 5 to 3 ka (Briner et al., 2014) and increased influence of warm Atlantic water masses (Gibb et al., 2015; Hansen et al., 2020; Ouellet-Bernier et al.,

835 2014). Similar to Lake Rosaea, proxy timeseries from other coastal sites, including Lake Qipisarqo, Camarum Sø, and Kloft Sø, show a delayed thermal maximum, in accord with SST timeseries (Freehette and de Vernal, 2009; Fredskild, 1973; Gajewski, 2015; Kaplan et al., 2002). We therefore suggest that temperature reconstructions from coastal sites where prevailing winds are onshore may be strongly influenced by ocean conditions via ocean-atmosphere heat exchange.

Overall, bGDGT-inferred MAF-T timeseries suggest that local variations in temperature, influenced by the. The timeseries from Lake Gus revealed peak warmth between *ca.* 9 and 6 ka, earlier than many other time series, likely due to minimal influence of the ocean and the Greenland Ice Sheet, as it is far from both. Although Lake Rosaea was minimally affected by the Greenland Ice Sheet margin during the Holocene, the MAF-T reconstruction from the lake does not show the thermal maximum between 9 and 6 ka, as observed at Lake Gus, but rather indicates that peak warmth occurred between 7.5 and 5 ka. This is likely due to a strong marine influence at Lake Rosaea, where air masses arrive directly from the Labrador Sea and Baffin Bay, which experienced low SSTs during 10 to 7 ka.

The peak warmth in lakes Pluto, N3, Gus, and 578 was approx. 1.4 ± 0.2 °C warmer than in 1994–2024. While in lakes Rosaea, Bullet, and Marshall, it was approx. 2, 3, and 5 °C warmer than in 1994–2024, respectively. The higher magnitude of peak warmth in lakes Bullet and Marshall is likely related to their higher elevation, whereas the magnitude of temperature change at Lake Rosaea may be influenced by its proximity to the ocean.

850 Overall, our results highlight that the majority of temperature time series from southwestern Greenland exhibit a similar pattern over the Holocene, albeit with local variations in magnitude and timing caused by topography and proximity to the ice sheet and ocean, caused discrepancies in timing and magnitude of Holocene temperature in southwestern Greenland. The ocean. Further, the seasonality (e.g., MAF, summer, annual) of reconstructed temperatures seems important when comparing the magnitude of Holocene temperature change between the timeseries using across time series derived from different proxies. Therefore, further investigation of proxy time series to quantify seasonal variability and the impact of local factors, (e.g., site-specific, topographic, ice-sheet proximity, and oceanic conditions,) will likely reveal broad similarities among proxy timeseries, helping to reconcile differences and refine Holocene climate reconstructions.

Data availability

860 All the data will be available at in the Palaeoclimatology data portal at the National Centers for Environmental Information for Palaeoclimatology data portal.

Authors contribution

We thank the people of Greenland for providing access to their **Landland**. The study was conceptualized by SA and EKT. **JPB and EKT obtained funding and led field seasons**. AC, ALG, and ISC conducted the laboratory analysis. SA and EKT prepared the manuscript with contributions from all co-authors.

865 Competing interests

The authors declare that they have no conflict of interest

Acknowledgements

We thank John Ryan-Henry, Kayla Hollister, Owen Cowling, Nancy Leon, and Jeff Salacup for their help in lipid biomarker analysis. We gratefully acknowledge Dr. Gerard A. Otiniano and Dr. Samuel Mark for discussions.

870 Financial support

This research was funded by National Science Foundation grant OPP-1504267 and OPP-2106971 to EKT and JPB, EAR-IF 1652274 to EKT and NSF Graduate Research Fellowship 1645677 to AAC.

References

- 875 Acharya, S., Zech, R., Strobel, P., Bliedtner, M., Prochnow, M., and De Jonge, C.: Environmental controls on the distribution of GDGT molecules in Lake Höglwörth, Southern Germany, Organic Geochemistry, 186, 104689, **2023**, <https://doi.org/10.1016/j.orggeochem.2023.104689>, 2023.
- Andresen, C. S., Björck, S., Bennike, O., and Bond, G.: Holocene climate changes in southern Greenland: evidence from lake sediments, Journal of Quaternary Science, 19, 783–795, <https://doi.org/10.1002/jqs.886>, 2004.
- 880 Antoniades, D., Klanten, Y., Cameron, E., Garcia-Oteyza, J., Oliva, M., and Vincent, W. F.: Evidence of summer thermal stratification in extreme northern lakes, Arctic Science, 10, 225–232, <https://doi.org/10.1139/as-2023-0037>, 2024.
- Axford, Y., Losee, S., Briner, J. P., Francis, D. R., Langdon, P. G., and Walker, I. R.: Holocene temperature history at the western Greenland Ice Sheet margin reconstructed from lake sediments, Quaternary Science Reviews, 59, 87–100, <https://doi.org/10.1016/j.quascirev.2012.10.024>, 2013.
- 885 Axford, Y., De Vernal, A., and Osterberg, E. C.: Past Warmth and Its Impacts During the Holocene Thermal Maximum in Greenland, Annu. Rev. Earth Planet. Sci., 49, 279–307, <https://doi.org/10.1146/annurev-earth-081420-063858>, 2021.
- Bashmachnikov, I. L., Raj, R. P., Golubkin, P., and Kozlov, I. E.: Heat Transport by Mesoscale Eddies in the Norwegian and Greenland Seas, Journal of Geophysical Research: Oceans, 128, e2022JC018987, <https://doi.org/10.1029/2022JC018987>, 2023.

- 890 Bauersachs, T., Schubert, C. J., Mayr, C., Gilli, A., and Schwark, L.: Branched GDGT-based temperature calibrations from Central European lakes, *Science of The Total Environment*, 167724, <https://doi.org/10.1016/j.scitotenv.2023.167724>, 2023.
- Baxter, A. J., van Bree, L. G. J., Peterse, F., Hopmans, E. C., Villanueva, L., Verschuren, D., and Sinninghe Damsté, J. S.: Seasonal and multi-annual variation in the abundance of isoprenoid GDGT membrane lipids and their producers in the water column of a meromictic equatorial crater lake (Lake Chala, East Africa), *Quaternary Science Reviews*, 273, <https://doi.org/10.1016/j.quascirev.2021.107263>, 2021.
- 895 Bennike, O.: Palaeoecological studies of Holocene lake sediments from west Greenland, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 155, 285–304, [https://doi.org/10.1016/S0031-0182\(99\)00121-2](https://doi.org/10.1016/S0031-0182(99)00121-2), 2000.
- Bennike, O., Anderson, N. J., and McGowan, S.: Holocene palaeoecology of southwest Greenland inferred from macrofossils in sediments of an oligosaline lake, *J Paleolimnol*, 43, 787–798, <https://doi.org/10.1007/s10933-009-9368-x>, 2010.
- 900 ~~Besseling, M. A., Hopmans, E. C., Boschman, R. C., Sinninghe Damsté, J. S., and Villanueva, L.: Benthic archaea as potential sources of tetraether membrane lipids in sediments across an oxygen minimum zone, *Biogeosciences*, 15, 4047–4064, <https://doi.org/10.5194/bg-15-4047-2018>, 2018.~~
- Bittner, L., De Jonge, C., Gil-Romera, G., Lamb, H. F., Russell, J. M., and Zech, M.: A Holocene temperature (brGDGT) record from Garba Guracha, a high-altitude lake in Ethiopia, *Biogeosciences*, 19, 5357–5374, <https://doi.org/10.5194/bg-19-5357-2022>, 2022.
- 905 Blaauw, M. and Christen, J. A.: Flexible Paleoclimate Age-Depth Models Using an Autoregressive Gamma Process, *Bayesian Anal*, 6, 457–474, <https://doi.org/10.1214/ba/1339616472>, 2011.
- Blaga, C. I., Reichart, G.-J., Heiri, O., and Sinninghe Damsté, J. S.: Tetraether membrane lipid distributions in water-column particulate matter and sediments: a study of 47 European lakes along a north–south transect, *Journal of Paleolimnology*, 41, 523–540, <https://doi.org/10.1007/s10933-008-9242-2>, 2009.
- 910 ~~Blaga, C. I., Reichart, G. J., Vissers, E. W., Lotter, A. F., Anselmetti, F. S., and Sinninghe Damsté, J. S.: Seasonal changes in glycerol dialkyl glycerol tetraether concentrations and fluxes in a perialpine lake: Implications for the use of the TEX86 and BIT proxies, *Geochimica et Cosmochimica Acta*, 75, 6416–6428, <https://doi.org/10.1016/j.gca.2011.08.016>, 2011.~~
- Box, J. E., Nielsen, K. P., Yang, X., Niwano, M., Wehrlé, A., van As, D., Fettweis, X., Køltzow, M. A. Ø., Palmason, B., Fausto, R. S., van den Broeke, M. R., Huai, B., Ahlstrøm, A. P., Langley, K., Dachauer, A., and Noël, B.: Greenland ice sheet rainfall climatology, extremes and atmospheric river rapids, *Meteorological Applications*, 30, e2134, <https://doi.org/10.1002/met.2134>, 2023.
- 915 van Bree, L. G. J., Peterse, F., Baxter, A. J., De Crop, W., van Grinsven, S., Villanueva, L., Verschuren, D., and Sinninghe Damsté, J. S.: Seasonal variability and sources of in situ brGDGT production in a permanently stratified African crater lake, *Biogeosciences*, 17, 5443–5463, <https://doi.org/10.5194/bg-17-5443-2020>, 2020.
- 920 Briner, J. P., Kaufman, D. S., Bennike, O., and Kosnik, M. A.: Amino acid ratios in reworked marine bivalve shells constrain Greenland Ice Sheet history during the Holocene, *Geology*, 42, 75–78, <https://doi.org/10.1130/G34843.1>, 2014.
- Briner, J. P., McKay, N. P., Axford, Y., Bennike, O., Bradley, R. S., De Vernal, A., Fisher, D., Francus, P., Fréchette, B., Gajewski, K., Jennings, A., Kaufman, D. S., Miller, G., Rouston, C., and Wagner, B.: Holocene climate change in Arctic Canada and Greenland, *Quaternary Science Reviews*, 147, 340–364, <https://doi.org/10.1016/j.quascirev.2016.02.010>, 2016.

- 925 Briner, J. P., Cuzzzone, J. K., Badgley, J. A., Young, N. E., Steig, E. J., Morlighem, M., Schlegel, N.-J., Hakim, G. J., Schaefer, J. M., Johnson, J. V., Lesnek, A. J., Thomas, E. K., Allan, E., Bennike, O., Cluett, A. A., Csatho, B., De Vernal, A., Downs, J., Larour, E., and Nowicki, S.: Rate of mass loss from the Greenland Ice Sheet will exceed Holocene values this century, *Nature*, 586, 70–74, <https://doi.org/10.1038/s41586-020-2742-6>, 2020.
- 930 ~~Buckles, L. K., Villanueva, L., Weijers, J. W., Verschuren, D., and Damsté, J. S.: Linking isoprenoidal GDGT membrane lipid distributions with gene abundances of ammonia-oxidizing Thaumarchaeota and uncultured crenarchaeotal groups in the water column of a tropical lake (Lake Challa, East Africa), *Environmental Microbiology*, 15, 2445–62, <https://doi.org/10.1111/1462-2920.12118>, 2013.~~
- 935 ~~Buckles, L. K.,~~ Weijers, J. W. H., Verschuren, D., and Sinninghe Damsté, J. S.: Sources of core and intact branched tetraether membrane lipids in the lacustrine environment: Anatomy of Lake Challa and its catchment, equatorial East Africa, *Geochimica et Cosmochimica Acta*, 140, 106–126, <https://doi.org/10.1016/j.gca.2014.04.042>, 2014.
- ~~Butler, T. M., Wilhelm, A. C., Dwyer, A. C., Webb, P. N., Baldwin, A. L., and Techtmann, S. M.: Microbial Community Dynamics During Lake Ice Freezing, *Sci Rep*, 9, 6231, <https://doi.org/10.1038/s41598-019-42609-9>, 2019.~~
- Castañeda, I. S. and Schouten, S.: A review of molecular organic proxies for examining modern and ancient lacustrine environments, *Quaternary Science Reviews*, 30, 2851–2891, <https://doi.org/10.1016/j.quascirev.2011.07.009>, 2011.
- 940 Cluett, A. A., Thomas, E. K., Evans, S. M., and Keys, P. W.: Seasonal Variations in Moisture Origin Explain Spatial Contrast in Precipitation Isotope Seasonality on Coastal Western Greenland, *JGR Atmospheres*, 126, e2020JD033543, <https://doi.org/10.1029/2020JD033543>, 2021.
- 945 Cluett, A. A., Thomas, E. K., McKay, N. P., Cowling, O. C., Castañeda, I. S., and Morrill, C.: Lake Dynamics Modulate the Air Temperature Variability Recorded by Sedimentary Aquatic Biomarkers: A Holocene Case Study From Western Greenland, *JGR Biogeosciences*, 128, e2022JG007106, <https://doi.org/10.1029/2022JG007106>, 2023.
- ~~Coleord, D. E., Cadieux, S. B., Brassell, S. C., Castañeda, I. S., Pratt, L. M., and White, J. R.: Assessment of branched GDGTs as temperature proxies in sedimentary records from several small lakes in southwestern Greenland, *Organic Geochemistry*, 82, 33–41, <https://doi.org/10.1016/j.orggeochem.2015.02.005>, 2015.~~
- 950 Corcoran, M. C., Thomas, E. K., and Morrill, C.: Using a Paired Chironomid $\delta^{18}\text{O}$ and Aquatic Leaf Wax $\delta^2\text{H}$ Approach to Reconstruct Seasonality on Western Greenland During the Holocene, *Paleoceanog and Paleoclimatol*, 36, e2020PA004169, <https://doi.org/10.1029/2020PA004169>, 2021.
- Cronauer, S. L., Briner, J. P., Kelley, S. E., Zimmerman, S. R. H., and Morlighem, M.: ^{10}Be dating reveals early-middle Holocene age of the Drygalski Moraines in central West Greenland, *Quaternary Science Reviews*, 147, 59–68, <https://doi.org/10.1016/j.quascirev.2015.08.034>, 2016.
- 955 Dang, X., Ding, W., Yang, H., Pancost, R. D., Naafs, B. D. A., Xue, J., Lin, X., Lu, J., and Xie, S.: Different temperature dependence of the bacterial brGDGT isomers in 35 Chinese lake sediments compared to that in soils, *Organic Geochemistry*, 119, 72–79, <https://doi.org/10.1016/j.orggeochem.2018.02.008>, 2018.
- 960 ~~De Jonge, C., Stadnitskaia, A., Hopmans, E. C., Cherkashov, G., Fedotov, A., and Sinninghe Damsté, J. S.: In situ produced branched glycerol dialkyl glycerol tetraethers in suspended particulate matter from the Yenisei River, Eastern Siberia, *Geochimica et Cosmochimica Acta*, 125, 476–491, <https://doi.org/10.1016/j.gca.2013.10.031>, 2014.~~
- Dee, S. G., Russell, J. M., Morrill, C., Chen, Z., and Neary, A.: PRYSM v2.0: A Proxy System Model for Lacustrine Archives, *Paleoceanog and Paleoclimatol*, 33, 1250–1269, <https://doi.org/10.1029/2018PA003413>, 2018.

- Dokulil, M. T., de Eyto, E., Maberly, S. C., May, L., Weyhenmeyer, G. A., and Woolway, R. I.: Increasing maximum lake surface temperature under climate change, *Climatic Change*, 165, 56, <https://doi.org/10.1007/s10584-021-03085-1>, 2021.
- 965 Downs, J., Johnson, J., Briner, J., Young, N., Lesnek, A., and Cuzzone, J.: Western Greenland ice sheet retreat history reveals elevated precipitation during the Holocene thermal maximum, *The Cryosphere*, 14, 1121–1137, <https://doi.org/10.5194/tc-14-1121-2020>, 2020.
- Druckenmiller, M. L., Thoman, R. L., and Moon, T. A.: NOAA Arctic Report Card 2025, <https://doi.org/10.25923/nrzf-j897>, 2025.
- 970 Dufour, A., Zolina, O., and Gulev, S. K.: Atmospheric Moisture Transport to the Arctic: Assessment of Reanalyses and Analysis of Transport Components, *Journal of Climate*, 29, 5061–5081, <https://doi.org/10.1175/JCLI-D-15-0559.1>, 2016.
- Dutton, A., Carlson, A. E., Long, A. J., Milne, G. A., Clark, P. U., DeConto, R., Horton, B. P., Rahmstorf, S., and Raymo, M. E.: Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, 349, aaa4019, <https://doi.org/10.1126/science.aaa4019>, 2015.
- 975 Eggermont, H. and Heiri, O.: The chironomid-temperature relationship: expression in nature and palaeoenvironmental implications, *Biological Reviews*, 87, 430–456, <https://doi.org/10.1111/j.1469-185X.2011.00206.x>, 2012.
- England, M. R., Eisenman, I., Lutsko, N. J., and Wagner, T. J. W.: The Recent Emergence of Arctic Amplification, *Geophysical Research Letters*, 48, e2021GL094086, <https://doi.org/10.1029/2021GL094086>, 2021.
- 980 Feng, X., Zhao, C., D’Andrea, W. J., Liang, J., Zhou, A., and Shen, J.: Temperature fluctuations during the Common Era in subtropical southwestern China inferred from brGDGTs in a remote alpine lake, *Earth and Planetary Science Letters*, 510, 26–36, <https://doi.org/10.1016/j.epsl.2018.12.028>, 2019.
- *Fisher, M., *Stafford, T., Moon, T., and Thurber, A.: QGreenland, , <https://doi.org/10.5281/zenodo.8326507>, 2023.
- Frechette, B. and de Vernal, A.: Relationship between Holocene climate variations over southern Greenland and eastern Baffin Island and synoptic circulation pattern, *Clim. Past*, 2009.
- 985 Frederikse, T., Landerer, F., Caron, L., Adhikari, S., Parkes, D., Humphrey, V. W., Dangendorf, S., Hogarth, P., Zanna, L., Cheng, L., and Wu, Y.-H.: The causes of sea-level rise since 1900, *Nature*, 584, 393–397, <https://doi.org/10.1038/s41586-020-2591-3>, 2020.
- Fredskild, B.: Studies in the Vegetational History of Greenland, *Palaeobotanical Investigations of some Holocene Lake and Bog deposits.*, Meddr Grønland, 198, <https://doi.org/10.7146/mog.v198.147748>, 1973.
- 990 Fredskild, B.: The Holocene vegetational development of the Godthåbsfjord area, West Greenland, *Meddelelser om Grønland. Geoscience*, 10, 28 pp.-28 pp., <https://doi.org/10.7146/moggeosci.v10i.140322>, 1983.
- Gajewski, K.: Quantitative reconstruction of Holocene temperatures across the Canadian Arctic and Greenland, *Global and Planetary Change*, 128, 14–23, <https://doi.org/10.1016/j.gloplacha.2015.02.003>, 2015.
- 995 Gibb, O. T., Steinhauer, S., Fréchet, B., de Vernal, A., and Hillaire-Marcel, C.: Diachronous evolution of sea surface conditions in the Labrador Sea and Baffin Bay since the last deglaciation, *The Holocene*, 25, 1882–1897, <https://doi.org/10.1177/0959683615591352>, 2015.

- Hansen, K. E., Giraudeau, J., Wacker, L., Pearce, C., and Seidenkrantz, M.-S.: Reconstruction of Holocene oceanographic conditions in eastern Baffin Bay, *Climate of the Past*, 16, 1075–1095, <https://doi.org/10.5194/cp-16-1075-2020>, 2020.
- 000 Harning, D. J., Sacco, S., Raberg, J. H., Ardenghi, N., Sepúlveda, J., Shapiro, B., Miller, G. H., and Geirsdóttir, Á.: Both redox potential and climate control molecular proxies in Icelandic Holocene lake sediments, *Commun Earth Environ*, 6, 763, <https://doi.org/10.1038/s43247-025-02701-7>, 2025.
- Hopmans, E. C., Schouten, S., and Sinninghe Damsté, J. S.: The effect of improved chromatography on GDGT-based palaeoproxies, *Organic Geochemistry*, 93, 1–6, <https://doi.org/10.1016/j.orggeochem.2015.12.006>, 2016.
- 005 Hostetler, S. W. and Bartlein, P. J.: Simulation of lake evaporation with application to modeling lake level variations of Harney-Malheur Lake, Oregon, *Water Resources Research*, 26, 2603–2612, <https://doi.org/10.1029/WR026i010p02603>, 1990.
- IPCC: AR6 Climate Change 2021: The Physical Science Basis, Intergovernmental pannel on climate change, Geneva, 2021.
- Jane, S. F., Mincer, J. L., Lau, M. P., Lewis, A. S. L., Stetler, J. T., and Rose, K. C.: Longer duration of seasonal stratification contributes to widespread increases in lake hypoxia and anoxia, *Global Change Biology*, 29, 1009–1023, <https://doi.org/10.1111/gcb.16525>, 2023.
- 010 Kaplan, M. R., Wolfe, A. P., and Miller, G. H.: Holocene Environmental Variability in Southern Greenland Inferred from Lake Sediments, *Quat. res.*, 58, 149–159, <https://doi.org/10.1006/qres.2002.2352>, 2002.
- Karami, M., Hansen, B. U., Westergaard-Nielsen, A., Abermann, J., Lund, M., Schmidt, N. M., and Elberling, B.: Vegetation phenology gradients along the west and east coasts of Greenland from 2001 to 2015, *Ambio*, 46, 94–105, <https://doi.org/10.1007/s13280-016-0866-6>, 2017.
- 015 Kaufman, D.: Holocene thermal maximum in the western Arctic (0–180°W), *Quaternary Science Reviews*, 23, 529–560, <https://doi.org/10.1016/j.quascirev.2003.09.007>, 2004.
- Kaufman, D., McKay, N., Routson, C., Erb, M., Dätwyler, C., Sommer, P. S., Heiri, O., and Davis, B.: Holocene global mean surface temperature, a multi-method reconstruction approach, *Sci Data*, 7, 201, <https://doi.org/10.1038/s41597-020-0530-7>, 2020.
- 020 ~~Kerimoglu, O. and Rinke, K.: Stratification dynamics in a shallow reservoir under different hydro-meteorological scenarios and operational strategies, *Water Resources Research*, 49, 7518–7527, <https://doi.org/10.1002/2013WR013520>, 2013.~~
- Kettle, H., Thompson, R., Anderson, N. J., and Livingstone, D. M.: Empirical modeling of summer lake surface temperatures in southwest Greenland, *Limnology and Oceanography*, 49, 271–282, <https://doi.org/10.4319/lo.2004.49.1.0271>, 2004.
- 025 Klanten, Y., Couture, R. -M., Christoffersen, K. S., Vincent, W. F., and Antoniades, D.: Oxygen Depletion in Arctic Lakes: Circumpolar Trends, Biogeochemical Processes, and Implications of Climate Change, *Global Biogeochemical Cycles*, 37, e2022GB007616, <https://doi.org/10.1029/2022GB007616>, 2023.
- Klanten, Y., MacIntyre, S., Fitzpatrick, C., Vincent, W. F., and Antoniades, D.: Regime Shifts in Lake Oxygen and Temperature in the Rapidly Warming High Arctic, *Geophysical Research Letters*, 51, e2023GL106985, <https://doi.org/10.1029/2023GL106985>, 2024.
- 030 Kobashi, T., Menviel, L., Jeltsch-Thömmes, A., Vinther, B. M., Box, J. E., Muscheler, R., Nakaegawa, T., Pfister, P. L., Döring, M., Leuenberger, M., Wanner, H., and Ohmura, A.: Volcanic influence on centennial to millennial Holocene Greenland temperature change, *Sci Rep*, 7, 1441, <https://doi.org/10.1038/s41598-017-01451-7>, 2017.

- Larocca, L. J. and Axford, Y.: Arctic glaciers and ice caps through the Holocene: a circumpolar synthesis of lake-based reconstructions, *Climate of the Past*, 18, 579–606, <https://doi.org/10.5194/cp-18-579-2022>, 2022.
- 1035 Larocca, L. J., Axford, Y., Bjørk, A. A., Lasher, G. E., and Brooks, J. P.: Local glaciers record delayed peak Holocene warmth in south Greenland, *Quaternary Science Reviews*, 241, 106421, <https://doi.org/10.1016/j.quascirev.2020.106421>, 2020.
- Larsen, N. K., Kjær, K. H., Lecavalier, B., Bjørk, A. A., Colding, S., Huybrechts, P., Jakobsen, K. E., Kjeldsen, K. K., Knudsen, K.-L., Odgaard, B. V., and Olsen, J.: The response of the southern Greenland ice sheet to the Holocene thermal maximum, *Geology*, 43, 291–294, <https://doi.org/10.1130/G36476.1>, 2015.
- 1040 Larsen, N. K., Strunk, A., Levy, L. B., Olsen, J., Bjørk, A., Lauridsen, T. L., Jeppesen, E., and Davidson, T. A.: Strong altitudinal control on the response of local glaciers to Holocene climate change in southwest Greenland, *Quaternary Science Reviews*, 168, 69–78, <https://doi.org/10.1016/j.quascirev.2017.05.008>, 2017.
- Lasher, G. E., Axford, Y., Masterson, A. L., Berman, K., and Larocca, L. J.: Holocene temperature and landscape history of southwest Greenland inferred from isotope and geochemical lake sediment proxies, *Quaternary Science Reviews*, 239, 106358, 1045 <https://doi.org/10.1016/j.quascirev.2020.106358>, 2020.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M., and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *A&A*, 428, 261–285, <https://doi.org/10.1051/0004-6361:20041335>, 2004.
- Lazier, J., Hendry, R., Clarke, A., Yashayaev, I., and Rhines, P.: Convection and restratification in the Labrador Sea, 1990–2000, *Deep Sea Research Part I: Oceanographic Research Papers*, 49, 1819–1835, [https://doi.org/10.1016/S0967-0637\(02\)00064-X](https://doi.org/10.1016/S0967-0637(02)00064-X), 2002. 1050
- Leger, T. P. M., Clark, C. D., Huynh, C., Jones, S., Ely, J. C., Bradley, S. L., Diemont, C., and Hughes, A. L. C.: A Greenland-wide empirical reconstruction of paleo ice sheet retreat informed by ice extent markers: PaleoGrIS version 1.0, *Clim. Past*, 20, 701–755, <https://doi.org/10.5194/cp-20-701-2024>, 2024.
- Lesnek, A. J., Briner, J. P., Young, N. E., and Cuzzone, J. K.: Maximum Southwest Greenland Ice Sheet Recession in the 1055 Early Holocene, *Geophysical Research Letters*, 47, e2019GL083164, <https://doi.org/10.1029/2019GL083164>, 2020.
- Liang, J., Chevalier, M., Liu, K., Perfumo, A., Wang, M., Xie, H., Hou, J., Herzsuh, U., and Chen, F.: Discrepancies in lacustrine bacterial lipid temperature reconstructions explained by microbial ecology, *Commun Earth Environ*, 5, 1–14, <https://doi.org/10.1038/s43247-024-01925-3>, 2024.
- Lindberg, K. R., Daniels, W. C., Castañeda, I. S., and Brigham-Grette, J.: Biomarker proxy records of Arctic climate change during the Mid-Pleistocene transition from Lake El’gygytyn (Far East Russia), *Clim. Past*, 18, 559–577, 1060 <https://doi.org/10.5194/cp-18-559-2022>, 2022.
- Lindborg, T., Rydberg, J., Tröjbom, M., Berglund, S., Johansson, E., Löfgren, A., Saetre, P., Nordén, S., Sohlenius, G., Andersson, E., Petrone, J., Borgiel, M., Kautsky, U., and Laudon, H.: Biogeochemical data from terrestrial and aquatic ecosystems in a periglacial catchment, West Greenland, *Earth System Science Data*, 8, 439–459, <https://doi.org/10.5194/essd-8-439-2016>, 2016. 1065
- Liu, Z., Zhu, J., Rosenthal, Y., Zhang, X., Otto-Bliesner, B. L., Timmermann, A., Smith, R. S., Lohmann, G., Zheng, W., and Elison Timm, O.: The Holocene temperature conundrum, *Proceedings of the National Academy of Sciences*, 111, E3501–5, <https://doi.org/10.1073/pnas.1407229111>, 2014.

- 070 Longo, W. M., Huang, Y., Russell, J. M., Morrill, C., Daniels, W. C., Giblin, A. E., and Crowther, J.: Insolation and greenhouse gases drove Holocene winter and spring warming in Arctic Alaska, *Quaternary Science Reviews*, 242, 106438, 2020.
- Loomis, S. E., Russell, J. M., Eggermont, H., Verschuren, D., and Sinninghe Damsté, J. S.: Effects of temperature, pH and nutrient concentration on branched GDGT distributions in East African lakes: Implications for paleoenvironmental reconstruction, *Organic Geochemistry*, 66, 25–37, <https://doi.org/10.1016/j.orggeochem.2013.10.012>, 2014a2014.
- 075 ~~Loomis, S. E., Russell, J. Martin, C., Ménot, G., Thouveny, N., Davtian, N., Andrieu-Ponel, V., Reille, M., Heurreux, A. M., D'Andrea, W. J., and Sinninghe Damsté, J. S.: Seasonal variability~~ Bard, E.: Impact of branched glycerol dialkyl glycerol tetraethers (brGDGTs) human activities and vegetation changes on the tetraether sources in a temperate lake system, *Geochimica et Cosmochimica Acta*, 144, 173–187 Lake St Front (Massif Central, France), *Organic Geochemistry*, 135, 38–52, <https://doi.org/10.1016/j.gea.2014.08.027>, 2014b.
- 080 ~~MacIntyre, S., Cortés, A., and Sadro, S.: Sediment respiration drives circulation and production of CO₂ in ice covered Alaskan arctic lakes, *Limnol Oceanogr Letters*, 3, 302–310, <https://doi.org/10.1002/lol2.10083>, 2018~~ *orggeochem.2019.06.005*, 2019.
- Martin, C., Ménot, G., Thouveny, N., Peyron, O., Andrieu-Ponel, V., Montade, V., Davtian, N., Reille, M., and Bard, E.: Early Holocene Thermal Maximum recorded by branched tetraethers and pollen in Western Europe (Massif Central, France), *Quaternary Science Reviews*, 228, <https://doi.org/10.1016/j.quascirev.2019.106109>, 2020.
- 085 Martin, K. C., Buizert, C., Brook, E., Williams, O. L., Edwards, J. S., Riddell-Young, B., Fudge, T. J., Mederbel, F., Beaudette, R., Severinghaus, J., Oyabu, I., Kawamura, K., Kirk, M., Koldtoft, I., Steffensen, J. P., and Blunier, T.: Greenland Ice Cores Reveal a South-To-North Difference in Holocene Thermal Maximum Timings, *Geophysical Research Letters*, 51, e2024GL111405, <https://doi.org/10.1029/2024GL111405>, 2024.
- Martínez-Sosa, P., Tierney, J. E., Stefanescu, I. C., Dearing Crampton-Flood, E., Shuman, B. N., and Routson, C.: A global Bayesian temperature calibration for lacustrine brGDGTs, *Geochimica et Cosmochimica Acta*, 305, 87–105, <https://doi.org/10.1016/j.gca.2021.04.038>, 2021.
- 090 ~~Massé, S., Botrel, M., Walsh, D. A., and Maranger, R.: Annual nitrification dynamics in a seasonally ice covered lake, *PLOS ONE*, 14, e0213748, <https://doi.org/10.1371/journal.pone.0213748>, 2019.~~
- 095 Masson-Delmotte, V., Landais, A., Stievenard, M., Cattani, O., Falourd, S., Jouzel, J., Johnsen, S. J., Dahl-Jensen, D., Sveinbjornsdottir, A., White, J. W. C., Popp, T., and Fischer, H.: Holocene climatic changes in Greenland: Different deuterium excess signals at Greenland Ice Core Project (GRIP) and NorthGRIP, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/10.1029/2004JD005575>, 2005.
- ~~McFarlin, J. M., Axford, Y., Kuseh, S., Masterson, A. L., Lasher, G. E., and Osburn, M. R.: Aquatic plant wax hydrogen and carbon isotopes in Greenland lakes record shifts in methane cycling during past Holocene warming, *Sci. Adv.*, 9, eadh9704, <https://doi.org/10.1126/sciadv.adh9704>, 2023.~~
- 100 McKay, N. P., Emile-Geay, J., and Khider, D.: geoChronR – an R package to model, analyze, and visualize age-uncertain data, *Geochronology*, 3, 149–169, <https://doi.org/10.5194/gchron-3-149-2021>, 2021.
- Miller, D. R., Habicht, M. H., Keisling, B. A., Castañeda, I. S., and Bradley, R. S.: A 900-year New England temperature reconstruction from in situ seasonally produced branched glycerol dialkyl glycerol tetraethers (brGDGTs), *Climate of Past*, 2018.
- 105 Moon, T., Fisher, M., Stafford, T., and Harden, L.: QGreenland (3.0.0), <https://doi.org/10.5281/zenodo.12823307>, 2023.

- Morrill, C., Meador, E., Livneh, B., Liefert, D. T., and Shuman, B. N.: Quantitative model-data comparison of mid-Holocene lake-level change in the central Rocky Mountains, *Climate Dynamics*, 53, 1077–1094, <https://doi.org/https://doi.org/10.1007/s00382-019-04633-3>, 2019.
- 1110 Muñoz Sabater, J.: ERA5-Land hourly data from 1950 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS), <https://doi.org/10.24381/cds.e2161bac>, 2019.
- Novak, J. B., Russell, J. M., Lindemuth, E. R., Prokopenko, A. A., Pérez-Angel, L., Zhao, B., Swann, G. E. A., and Polissar, P. J.: The Branched GDGT Isomer Ratio Refines Lacustrine Paleotemperature Estimates, *Geochem Geophys Geosyst*, 26, e2024GC012069, <https://doi.org/10.1029/2024GC012069>, 2025.
- 1115 Nusbaumer, J., Alexander, P. M., LeGrande, A. N., and Tedesco, M.: Spatial Shift of Greenland Moisture Sources Related to Enhanced Arctic Warming, *Geophysical Research Letters*, 46, 14723–14731, <https://doi.org/10.1029/2019GL084633>, 2019.
- Otiniano, G., ~~Thomas, Elizabeth. K., Castañeda, I. S., Acharya, S., and Mark, S.: chromatPy, 2025.~~
- ~~Otiniano, G.~~ A., Porter, T. J., Buceta, R. E., Bergman, M. E., and Phillips, M. A.: Climatic and environmentally driven variability in lacustrine brGDGT distributions at local to regional scales in Alaska and northwestern Canada, *Organic Geochemistry*, 181, <https://doi.org/10.1016/j.orggeochem.2023.104604>, 2023.
- 1120 Otiniano, G. A., Porter, T. J., Phillips, M. A., Juutinen, S., Weckström, J. B., and Heikkilä, M. P.: Reconstructing warm-season temperatures using brGDGTs and assessing biases in Holocene temperature records in northern Fennoscandia, *Quaternary Science Reviews*, 329, 108555, <https://doi.org/10.1016/j.quascirev.2024.108555>, 2024.
- 125 Otiniano, G. A., Thomas, E. K., Castañeda, I. S., Acharya, S., Sharma, Y., and Mark, S. Z.: chromatPy: An Open-Source Tool for Chromatographic Peak Deconvolution and Integration Applied to Glycerol Dialkyl Glycerol Tetraethers, *Paleoceanography and Paleoclimatology*, 40, e2025PA005316, <https://doi.org/10.1029/2025PA005316>, 2025.
- Ouellet-Bernier, M.-M., De Vernal, A., Hillaire-Marcel, C., and Moros, M.: Paleooceanographic changes in the Disko Bugt area, West Greenland, during the Holocene, *The Holocene*, 24, 1573–1583, <https://doi.org/10.1177/0959683614544060>, 2014.
- 130 Pepin, N., Bradley, R. S., Diaz, H. F., Baraer, M., Caceres, E. B., Forsythe, N., Fowler, H., Greenwood, G., Hashmi, M. Z., Liu, X. D., Miller, J. R., Ning, L., Ohmura, A., Palazzi, E., Rangwala, I., ~~Schöner~~Schoner, W., Severskiy, I., Shahgedanova, M., Wang, M. B., Williamson, S. N., Yang, D. Q., and ~~Mountain Research~~Mt Res Initiative EDW Working ~~Group~~Grp: Elevation-dependent warming in mountain regions of the world, *Nature ~~Clim~~Climate Change*, 5, 424–430, <https://doi.org/10.1038/nclimate2563>, 2015.
- Perner, K., Moros, M., Jennings, A., Lloyd, J., and Knudsen, K.: Holocene palaeoceanographic evolution off West Greenland, *The Holocene*, 23, 374–387, <https://doi.org/10.1177/0959683612460785>, 2013.
- 1135 Piccolroaz, S., Woolway, R. I., and Merchant, C. J.: Global reconstruction of twentieth century lake surface water temperature reveals different warming trends depending on the climatic zone, *Climatic Change*, 160, 427–442, <https://doi.org/10.1007/s10584-020-02663-z>, 2020.
- 1140 Porter, C., Howat, I., Noh, M.-J., Husby, E., Khuvis, S., Danish, E., Tomko, K., Gardiner, J., Negrete, A., Yadav, B., Klassen, J., Kelleher, C., Cloutier, M., Bakker, J., Enos, J., Arnold, G., Bauer, G., and Morin, P.: ArcticDEM - Strips, Version 4.1 (V1), <https://doi.org/10.7910/DVN/C98DVS>, 2022.

- Prowse, T., Alfredsen, K., Beltaos, S., Bonsal, B., Duguay, C., Korhola, A., McNamara, J., Pienitz, R., Vincent, W. F., Vuglinsky, V., and Weyhenmeyer, G. A.: Past and Future Changes in Arctic Lake and River Ice, *AMBIO*, 40, 53–62, <https://doi.org/10.1007/s13280-011-0216-7>, 2011.
- 145 Raberg, J. H., Harning, D. J., Crump, S. E., de Wet, G., Blumm, A., Kopf, S., Geirsdóttir, Á., Miller, G. H., and Sepúlveda, J.: Revised fractional abundances and warm-season temperatures substantially improve brGDGT calibrations in lake sediments, *Biogeosciences*, 18, 3579–3603, <https://doi.org/10.5194/bg-18-3579-2021>, 2021.
- Raberg, J. H., Miller, G. H., Geirsdóttir, Á., and Sepúlveda, J.: Near-universal trends in brGDGT lipid distributions in nature, *Science advances*, 8, eabm7625, <https://doi.org/10.1126/sciadv.abm7625>, 2022.
- 150 Raberg, J. H., de Wet, G. A., Geirsdóttir, Á., Sepúlveda, J., and Miller, G. H.: Oxygen Depletion in Lake Waters May Skew brGDGT-Inferred Temperatures by More Than 10°C, *Geophysical Research Letters*, 52, e2024GL113562, <https://doi.org/10.1029/2024GL113562>, 2025.
- Rantanen, M., Karpechko, A. Yu., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun Earth Environ*, 3, 168, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- 155 Reyes, F. R. and Loughheed, V. L.: Rapid Nutrient Release from Permafrost Thaw in Arctic Aquatic Ecosystems, *Arctic, Antarctic, and Alpine Research*, 47, 35–48, <https://doi.org/10.1657/AAAR0013-099>, 2015.
- Russell, J. M., Hopmans, E. C., Loomis, S. E., Liang, J., and Sinninghe Damsté, J. S.: Distributions of 5- and 6-methyl branched glycerol dialkyl glycerol tetraethers (brGDGTs) in East African lake sediment: Effects of temperature, pH, and new lacustrine paleotemperature calibrations, *Organic Geochemistry*, 117, 56–69, <https://doi.org/10.1016/j.orggeochem.2017.12.003>, 2018.
- 160 ~~Saini, J., Stein, R., Fahl, K., Weiser, J., Hebbeln, D., and Madaj, L.: Holocene variability in sea ice conditions in the eastern Baffin Bay Labrador Sea—A north–south biomarker transect study, *Boreas*, 51, 553–572, <https://doi.org/10.1111/bor.12583>, 2022.~~
- ~~Saros, J. E., Northington, R. M., Osburn, C. L., Burpee, B. T., and John Anderson, N.: Thermal stratification in small arctic lakes of southwest Greenland affected by water transparency and epilimnetic temperatures, *Limnology and Oceanography*, 61, 1530–1542, <https://doi.org/10.1002/lno.10314>, 2016.~~
- 165 Schneider, T., Castañeda, I. S., Zhao, B., Krüger, S., Salacup, J. M., and Bradley, R. S.: Tracing Holocene temperatures and human impact in a Greenlandic Lake: Novel insights from hyperspectral imaging and lipid biomarkers, *Quaternary Science Reviews*, 339, 108851, <https://doi.org/10.1016/j.quascirev.2024.108851>, 2024.
- Schouten, S., Hopmans, E. C., Schefuß, E., and Damste, J. S. S.: Distributional variations in marine crenarchaeotal membrane lipids: a new tool for reconstructing ancient sea water temperatures?, *Earth and Planetary Science Letters*, 204, 265–274, 2002.
- 170 Schouten, S., Hopmans, E. C., and Damste, J. S. S.: The organic geochemistry of glycerol dialkyl glycerol tetraether lipids: A review, *Organic Geochemistry*, 54, 19–61, <https://doi.org/10.1016/j.orggeochem.2012.09.006>, 2013.
- Schweinsberg, A. D., Briner, J. P., Miller, G. H., Bennike, O., and Thomas, E. K.: Local glaciation in West Greenland linked to North Atlantic Ocean circulation during the Holocene, *Geology*, 45, 195–198, <https://doi.org/10.1130/G38114.1>, 2017.
- 175 Schweinsberg, A. D., Briner, J. P., Licciardi, J. M., Bennike, O., Lifton, N. A., Graham, B. L., Young, N. E., Schaefer, J. M., and Zimmerman, S. H.: Multiple independent records of local glacier variability on Nuussuaq, West Greenland, during the Holocene, *Quaternary Science Reviews*, 215, 253–271, <https://doi.org/10.1016/j.quascirev.2019.05.007>, 2019.

- Seneviratne, S. I. and Hauser, M.: Regional Climate Sensitivity of Climate Extremes in CMIP6 Versus CMIP5 Multimodel Ensembles, *Earth's Future*, 8, e2019EF001474, <https://doi.org/10.1029/2019EF001474>, 2020.
- 180 Shanahan, T. M., Hughen, K. A., and Van Mooy, B. A. S.: Temperature sensitivity of branched and isoprenoid GDGTs in Arctic lakes, *Organic Geochemistry*, 64, 119–128, <https://doi.org/10.1016/j.orggeochem.2013.09.010>, 2013.
- Sinninghe Damsté, J. S., Hopmans, E. C., Pancost, R. D., Schouten, S., and Geenevasen, J. A.: Newly discovered non-isoprenoid glycerol dialkyl glycerol tetraether lipids in sediments, *Chemical Communications*, 1683–1684, 2000.
- 185 Sinninghe Damsté, J. S., ~~Ossebaar~~Weber, Y., Zopfi, J., ~~Abbas, B., Schouten, S., Lehmann, M. F., and Verschuren, D.: Fluxes~~Niemann, H.: Distributions and ~~distributionsources~~ of ~~tetraether lipids~~isoprenoid GDGTs in ~~an equatorial African lake: Constraints on the application of the TEX86 palaeothermometer~~Lake Lugano and ~~BIT index in lacustrine settings, Geochimica et Cosmochimica Acta, 73, 4232–4249~~other central European (peri-)alpine lakes: Lessons for their use as paleotemperature proxies, *Quaternary Science Reviews*, 277, <https://doi.org/10.1016/j.gea.2009.04.022>, 2009~~quascirev.2021.107352~~, 2022.
- 190 Sweeney, A. J., Fu, Q., Po-Chedley, S., Wang, H., and Wang, M.: Internal Variability Increased Arctic Amplification During 1980–2022, *Geophysical Research Letters*, 50, e2023GL106060, <https://doi.org/10.1029/2023GL106060>, 2023.
- ~~Thoman, R. L., Moon, T. A., and Druckenmiller, M. L.: NOAA Arctic Report Card 2023: Executive Summary, https://doi.org/10.25923/SVFA-K694, 2023.~~
- 195 Thomas, E. K., Briner, J. P., Ryan-Henry, J. J., and Huang, Y.: A major increase in winter snowfall during the middle Holocene on western Greenland caused by reduced sea ice in Baffin Bay and the Labrador Sea, *Geophysical Research Letters*, 43, 5302–5308, <https://doi.org/10.1002/2016GL068513>, 2016.
- Thomas, E. K., Castañeda, I. S., McKay, N. P., Briner, J. P., Salacup, J. M., Nguyen, K. Q., and Schweinsberg, A. D.: A Wetter Arctic Coincident With Hemispheric Warming 8,000 Years Ago, *Geophysical Research Letters*, 45, <https://doi.org/10.1029/2018GL079517>, 2018.
- 200 Thomas, E. K., Cluett, A. A., Erb, M. P., McKay, N. P., Briner, J. P., Castañeda, I. S., Corcoran, M. C., Cowling, O. C., Gorbey, D. B., Lindberg, K. R., Prince, K. K., and Salacup, J.: Early Holocene Laurentide Ice Sheet Retreat Influenced Summer Atmospheric Circulation in Baffin Bay, *Geophysical Research Letters*, 50, <https://doi.org/10.1029/2023gl103428>, 2023.
- 205 Tierney, J. E., Schouten, S., Pitcher, A., Hopmans, E. C., and Sinninghe Damsté, J. S.: Core and intact polar glycerol dialkyl glycerol tetraethers (GDGTs) in Sand Pond, Warwick, Rhode Island (USA): Insights into the origin of lacustrine GDGTs, *Geochimica et Cosmochimica Acta*, 77, 561–581, <https://doi.org/10.1016/j.gca.2011.10.018>, 2012.
- Vihma, T., Screen, J., Tjernström, M., Newton, B., Zhang, X., Popova, V., Deser, C., Holland, M., and Prowse, T.: The atmospheric role in the Arctic water cycle: A review on processes, past and future changes, and their impacts, *Journal of Geophysical Research: Biogeosciences*, 121, 586–620, <https://doi.org/10.1002/2015JG003132>, 2016.
- 210 Weber, Y., De Jonge, C., Rijpstra, W. I. C., Hopmans, E. C., Stadnitskaia, A., Schubert, C. J., Lehmann, M. F., Sinninghe Damsté, J. S., and Niemann, H.: Identification and carbon isotope composition of a novel branched GDGT isomer in lake sediments: Evidence for lacustrine branched GDGT production, *Geochimica et Cosmochimica Acta*, 154, 118–129, <https://doi.org/10.1016/j.gca.2015.01.032>, 2015.
- 215 Weber, Y., Sinninghe Damsté, J. S., Zopfi, J., De Jonge, C., Gilli, A., Schubert, C. J., Lepori, F., Lehmann, M. F., and Niemann, H.: Redox-dependent niche differentiation provides evidence for multiple bacterial sources of glycerol tetraether lipids in lakes, *Proceedings of the National Academy of Sciences*, 115, 10926–10931, <https://doi.org/10.1073/pnas.1805186115>, 2018.

- Westergaard-Nielsen, A., Hansen, B. U., Elberling, B., and Abermann, J.: Greenland Climates, in: *Encyclopedia of the World's Biomes*, edited by: Goldstein, M. I. and DellaSala, D. A., Elsevier, Oxford, 539–550, <https://doi.org/10.1016/B978-0-12-409548-9.11750-6>, 2020.
- 1220 Wrona, F. J., Johansson, M., Culp, J. M., Jenkins, A., Mård, J., Myers-Smith, I. H., Prowse, T. D., Vincent, W. F., and Wookey, P. A.: Transitions in Arctic ecosystems: Ecological implications of a changing hydrological regime, *Journal of Geophysical Research: Biogeosciences*, 121, 650–674, <https://doi.org/10.1002/2015JG003133>, 2016.
- Wu, J., Yang, H., Pancost, R. D., Naafs, B. D. A., Qian, S., Dang, X., Sun, H., Pei, H., Wang, R., Zhao, S., and Xie, S.: Variations in dissolved O₂ in a Chinese lake drive changes in microbial communities and impact sedimentary GDGT distributions, *Chemical Geology*, 579, <https://doi.org/10.1016/j.chemgeo.2021.120348>, 2021.
- 1225 Yang, Q., Dixon, T. H., Myers, P. G., Bonin, J., Chambers, D., van den Broeke, M. R., Ribergaard, M. H., and Mortensen, J.: Recent increases in Arctic freshwater flux affects Labrador Sea convection and Atlantic overturning circulation, *Nat Commun*, 7, 10525, <https://doi.org/10.1038/ncomms10525>, 2016.
- 1230 Yao, Y., Zhao, J., Vachula, R. S., Werne, J. P., Wu, J., Song, X., and Huang, Y.: Correlation between the ratio of 5-methyl hexamethylated to pentamethylated branched GDGTs (HP5) and water depth reflects redox variations in stratified lakes, *Organic Geochemistry*, 147, <https://doi.org/10.1016/j.orggeochem.2020.104076>, 2020.
- Young, N. E., Briner, J. P., Rood, D. H., Finkel, R. C., Corbett, L. B., and Bierman, P. R.: Age of the Fjord Stade moraines in the Disko Bugt region, western Greenland, and the 9.3 and 8.2 ka cooling events, *Quaternary Science Reviews*, 60, 76–90, <https://doi.org/10.1016/j.quascirev.2012.09.028>, 2013.
- 1235 Young, N. E., Briner, J. P., Miller, G. H., Lesnek, A. J., Crump, S. E., Thomas, E. K., Pendleton, S. L., Cuzzone, J., Lamp, J., Zimmerman, S., Caffee, M., and Schaefer, J. M.: Deglaciation of the Greenland and Laurentide ice sheets interrupted by glacier advance during abrupt coolings, *Quaternary Science Reviews*, 229, 106091, <https://doi.org/10.1016/j.quascirev.2019.106091>, 2020.
- 1240 Zhao, B., Castañeda, I. S., Bradley, R. S., Salacup, J. M., De Wet, G. A., Daniels, W. C., and Schneider, T.: Development of an in situ branched GDGT calibration in Lake 578, southern Greenland, *Organic Geochemistry*, 152, 104168, <https://doi.org/10.1016/j.orggeochem.2020.104168>, 2021.
- Zhao, B., Russell, J. M., Tsai, V. C., Blaus, A., Parish, M. C., Liang, J., Wilk, A., Du, X., and Bush, M. B.: Evaluating global temperature calibrations for lacustrine branched GDGTs: Seasonal variability, paleoclimate implications, and future directions, *Quaternary Science Reviews*, 310, 108124, <https://doi.org/10.1016/j.quascirev.2023.108124>, 2023.