



Atmospheric Chemistry Experiment (ACE) Winds

Matthew Wyatt¹, Peter F. Bernath^{1,2,3}, Chris Boone³, Léo Lavy², and Ryan Johnson¹

¹Department of Physics, Old Dominion University, Norfolk, VA, United States, 23529

²Department of Chemistry and Biochemistry, Old Dominion University, Norfolk, VA, United States, 23529

³Department of Chemistry, University of Waterloo, Waterloo, ON, Canada, N2L 3G1

Correspondence: Matthew Wyatt (mwyat001@odu.edu)

Abstract. The Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS) uses limb geometry to measure transmittance spectra of Earth's atmosphere by solar occultation. Line-of-sight wind speeds can be derived via Doppler shifts of molecular lines in infrared spectra. The wind look direction angles relative to geodetic North are derived from geometry. We validate the new ACE version 5.3 (v.5.3) line-of-sight winds with MIGHTI and meteor radar vector wind observations and find a ± 15 m/s sunrise/sunset shift above 80 km. We also compare line-of-sight winds from ACE-FTS v.5.2 and v.5.3 with vector winds from the MERRA-2, HWM14, and WACCM-X models. A ± 15 m/s sunrise-sunset bias persists in v.5.3 winds above 80 km but decreases to less than ± 5 m/s below 50 km. The v.5.3 wind speed profiles have improved relative to v.5.2 at all altitudes. Over 20 years of ACE wind speeds can be used to test atmospheric models.

1 Introduction

The Atmospheric Chemistry Experiment (ACE) mission on board the Canadian satellite SCISAT is used for remote sensing Earth's atmosphere (Bernath et al., 2005; Bernath, 2017). The mission uses limb geometry during solar occultations to measure the transmission spectra of the atmosphere. Using the Sun as a light source, the high-resolution Fourier transform spectrometer (ACE-FTS) on board collects spectra as the Sun rises and sets from the perspective of the satellite. Line-of-sight wind speeds can be derived from the Doppler shift of spectral lines in the limb measurements. This is possible due to the frequency stability of the Michelson interferometer. The major advantages of ACE winds are the large altitude range, 18 to 140 km, and the longevity of the mission, starting in 2004 and still ongoing.

ACE wind speed retrievals have been described in detail for v.5.2 of ACE-FTS processing (Boone et al., 2021). A validation of v.5.2 line-of-sight winds has been carried out by comparing against meteor radar and remote sensing data (Johnson et al., 2024). Version 5.3 processing has been released for ACE-FTS data products, which includes an update to the retrieval of line-of-sight wind speeds. A detailed explanation of the new wind retrieval process is included in this work.

Horizontal wind speeds are available in the troposphere and lower stratosphere through measurements from airplanes (Khelif et al., 1999) and balloons (Duruiseau et al., 2017; Kumer et al., 2014), ground-based lidar (Martner et al., 1993), and satellite



lidar from the ADM-Aeolus (Atmospheric Dynamics Mission Aeolus) (Stoffelen et al., 2005; Lux et al., 2020). Vector wind measurements in the upper mesosphere lower thermosphere (UMLT) can be recorded from space using Doppler shifts in airglow lines such as from atom oxygen. Examples of this are the TIMED Doppler Interferometer (TIDI) instrument on the Thermosphere Ionosphere Mesosphere Energetics and Dynamics (TIMED) satellite (Killeen et al., 2006) and previously from the Michelson Interferometer for Global-High-resolution Thermospheric Imaging (MIGHTI) instrument on the Ionospheric Connection Explorer (ICON) satellite (Englert et al., 2017) and the Wind Imaging Interferometer (WINDII) on the Upper Atmosphere Research Satellite (UARS) (Shepherd et al., 1993). Also on UARS, the High Resolution Doppler Imager (HRDI) used O₂ emission to measure winds in UMLT (Hays et al., 1994; Grassl et al., 1995). Line-of-sight winds near the mesopause have also been derived from the Doppler shift in O₂ emission lines by the Microwave Limb Sounder (MLS) instrument on the Aura satellite (Wu et al., 2008). Note that, since many of these missions are inactive, ACE winds are especially valuable in this region.

There have previously been technical issues preventing wind measurements in the middle atmosphere (around 30 km to 70 km) (Baumgarten, 2010; Rüfenacht et al., 2018). There has been some success with ground-based lidar (Baumgarten, 2010; Liu et al., 2002) and microwave Doppler wind radiometers (Kishore Kumar et al., 2015), but only up to the lower stratosphere. Sounding rockets (Schmidlin et al., 1985) can be used but are expensive and do not provide continuous measurements over long periods of time. Ground-based meteor radar (Liu et al., 2002; Tang et al., 2021) can also provide winds in the mesosphere. There have been successful space-based missions that have provided line-of-sight winds in the middle atmosphere, such as the Atmospheric Trace Molecule Spectroscopy (ATMOS) experiment flown on the Spacelab 3 shuttle (van Cleef et al., 1987) and the Superconducting Submillimeter-Wave Limb Emission Sounder (SMILES) mission (Baron et al., 2013) on the International Space Station.

The Modern-Era Retrospective Analysis for Research and Applications, version 2 (MERRA-2) produced by NASA's Global Modeling and Assimilation Office (GMAO) is an atmospheric reanalysis model based on modern satellite observations. MERRA-2 provides various data collections that contain information about many climate indicators, including atmospheric wind speeds. An in depth explanation of the model is available from Gelaro et al. (2017). MERRA-2 uses the Goddard Earth Observing System (GEOS) atmospheric model (Rienecker et al., 2008; Molod et al., 2015) and the Gridpoint Statistical Interpolation (GSI) analysis scheme (Kleist et al., 2009). The model uses a cubed-sphere horizontal discretization giving an approximate resolution of $0.5^\circ \times 0.625^\circ$ and 72 hybrid-eta (altitude) levels (Putman and Lin, 2007). These levels reach from the surface up to 0.01 hPa (around 75 km) (Gelaro et al., 2017), which overlaps the lower half of ACE data.

Horizontal Wind Model Version: 2014 (HWM14) is an empirical climatology model of horizontal winds ranging from the troposphere up through the thermosphere. A detailed description of the climatology is available from Drob et al. (Drob et al., 2015). The model began as HWM87 (Hedin et al., 1988) and has had multiple iterations since, each time adding more data from new instruments (Drob et al., 2015; Hedin et al., 1996; Drob et al., 2008). The newest version, HWM14, uses $\sim 73 \times 10^6$ observation measurements from 44 different instruments (Drob et al., 2015). The climatology uses a set of spherical harmonics to provide a statistical view of vector winds ranging from near the surface up to 1000 km (Drob et al., 2015). Because HWM14 is an empirical climatology, vector winds can be found for any given latitude and longitude.



The Whole Atmosphere Community Climate Model with thermosphere and ionosphere eXtension (WACCM-X) is produced by the National Center for Atmospheric Research (NCAR). WACCM-X is a comprehensive numerical model of the whole atmosphere, ranging from Earth's surface up to around 700 km (Liu et al., 2018). The model is an altitude extension of WACCM6 (Gettelman et al., 2019), which reaches up to ~ 140 km. WACCM outputs many climate and weather data products and is unique in that the model can be coupled with others to include ocean, sea ice, and land components (Liu et al., 2018; Gettelman et al., 2019). Using CESM2 as a framework, WACCM is a mesh of NCAR projects: High Altitude Observatory (HAO) in the upper atmosphere, Atmospheric Chemistry Observations & Modeling (ACOM) in the middle atmosphere, and Climate & Global Dynamics (CGD) in the lower atmosphere. WACCM-X provides global vector winds on a $1.9^\circ \times 2.5^\circ$ grid.

In Section 2 of this paper we provide a brief explanation of ACE v.5.2 line-of-sight wind speed processing and a detailed explanation of v.5.3. Following, in Section 3 we validate ACE v.5.3 line-of-sight wind speeds with MIGHTI and meteor radar data. We then compare both v.5.2 and v.5.3 to the MERRA-2, HWM14, and WACCM-X models in Section 4. In the final section we provide concluding remarks.

2 ACE-FTS Line-of-Sight Wind Retrievals

2.1 ACE v.5.2 Winds

The retrieval process for ACE-FTS v.5.2 wind profiles has previously been explained in depth by Boone et al. (2021), so only a short explanation is provided here. For each limb measurement, the atmosphere is partitioned into 4 km regions above 45 km. Below that, increasingly smaller regions are used that decrease to 2 km in size. This is done to account for refraction distorting the disk of the Sun during a measurement. A forward model calculation is then used to derive a spectrum at a tangent height near the center of each partition. This forward model is based on pressure, temperature, and volume mixing ratio (VMR) profiles taken from sr10063 (sr for sunrise and 10063 for the orbit number since launch). The observed spectra and sr10063 are cross correlated to derive wind speeds from the Doppler shift in the spectra. The wind speeds are then interpolated onto a 1 km grid using a cubic spline. Finally, the wind profiles are calibrated using line-of-sight winds from the Canadian Meteorological Center (CMC) between 19 and 24 km altitude. The final product is line-of-sight wind speeds between 18 and 140 km at a given time, location, and heading. The heading is the angle between the 'look-direction' of the instrument and geodetic North, which is calculated by the Systems Tool Kit software package (STK).

The models and instruments we compare with provide vector wind speeds, not line-of-sight wind speeds. We are able to convert the vector winds into line-of-sight winds using

$$\text{Line-of-Sight Wind Speed} = U \cdot \cos(\theta - 90^\circ) + V \cdot \cos(\theta), \quad (1)$$

where U and V are the zonal (West-East) and meridional (South-North) components, respectively, from the given model or instrument, and θ is the heading of the ACE-FTS instrument.



2.2 ACE v.5.3 Winds

90 Version 5.3 explicitly retrieves altitude profiles for line-of-sight wind speed simultaneously with retrieving the VMR profiles of molecules contributing to the spectrum. The retrievals follow the usual procedure described previously for VMRs (Boone et al., 2023), but with a piecewise linear interpolation for wind speed onto the 1-km altitude grid employed for forward model calculations. Deriving wind speed using a retrieval, rather than using the wavenumber offset of a given measurement as an estimate of the Doppler shift at that measurement's tangent height (as was done in previous processing versions), should better
95 capture the altitude variation of wind speed when there is significant structure in the profile. With the retrieval approach, the calculated spectrum employed in the analysis is also more representative of the measured spectrum compared to the "generic" calculation from a single occultation (sr10063) in a given altitude "bin" that was employed in the v.5.2 analysis.

The altitude range was divided into 5 overlapping segments. The microwindows for a given altitude segment targeted single bands or sets of closely spaced bands (including overlapping hot bands) for a particular isotopologue to maximize internal
100 consistency. It was found that observed shifts for lines from different isotopologues or lines from molecules without distinct bands (like H₂O) were not always consistent. The microwindows employed in the different altitude segments are provided in Table A1 of the Appendix. All measurements that fall within the wavenumber, altitude, and minimum transmittance ranges are employed in the retrieval. The filter for minimum transmittance is used to avoid lines that are saturated or where the signal is too close to the noise, either of which could obscure Doppler shift information in the measurement. The upper altitude limit
105 of a particular segment is determined by the maximum tangent height within the altitude range that has at least three lines satisfying the filter conditions, to avoid having an abnormally large noise feature being included in the analysis when the signal falls below the nominal noise level.

Wind retrieval results for the lowest altitude segment (termed "segment 1" in Table A1) exhibit excessively large variability, possibly a consequence of pressure shifts and deviations of line shapes from the assumed Voigt profile becoming more promi-
110 nent at higher pressures. Therefore, in this segment line-of-sight wind speed is determined using the average wavenumber shifts of the lines, similar to the approach employed in previous processing versions, rather than using the retrieved results. Attempts to push the analysis lower in altitude often yielded relatively large gradients below 18 km, which are not expected in that altitude region, suggesting that systematic errors from pressure-related issues (such as pressure shifts and line shapes) become excessive when pressure gets too high. Therefore, the lower altitude limit of the analysis remained at 18 km.

115 Wind profiles in the lowest two altitude segments have persistent biases compared to the results in the three highest segments, as shown in Fig. 1. Thus, fixed shifts are applied to the wind profiles in the lowest two segments: -35 m/s in segment 1 and +35 m/s in segment 2. In overlap regions between segments, results are calculated from an unweighted average of the two profiles. When no measurement occurs in the overlap region between segments, there is a small altitude gap in the reported wind profile. As with previous processing versions, the final wind profile is shifted such that the results between 18 and 24 km
120 match the expectations from an analysis run of the Canadian weather model at Environment and Climate Change Canada (Buehner et al., 2015).




Figure 1. Wind retrieval for sr10063 (measured June 25th, 2005, at latitude 37.6° S and longitude 161.4° E). **a)** The “raw” retrieval, with large steps for the profiles in the bottom two altitude segments. **b)** Internal consistency is improved after shifting the profiles in the bottom two altitude segments

2.2.1 Calculating Look Direction

Comparing ACE-FTS winds to vector (zonal and meridional) wind information from other sources requires knowledge of the look direction of the instrument relative to either a northward-pointing vector (the positive direction for meridional winds) or an eastward-pointing vector (the positive direction for zonal winds). Previous processing versions employed the Systems Tool Kit (STK) software package to generate “heading” information (the angle relative to a north-pointing vector) associated with the vector extending from the measurement tangent point to the satellite (Boone et al., 2021). However, lacking an STK built-in function for this purpose, the calculation script represented a kludge of the software that appears to have had accuracy issues, particularly when the measurement tangent point approached the poles.

An alternative approach for the heading calculation is available, based on the fact that ACE ephemerides (satellite position and velocity information) in the Earth-centered Earth-fixed J2000 coordinate system are available on time scales of one minute or less for the duration of the ACE mission. Geodetic information (latitude, longitude, and altitude) for ACE-FTS tangent points, expressed relative to the WGS84 reference frame, can be readily converted into J2000 coordinates (Zhu, 1994). With satellite and tangent point locations available in the same cartesian coordinate system, heading information can be directly calculated from first principles.

Given the geodetic coordinate of a tangent point in the WGS84 reference frame (ϕ, λ, h) , where ϕ is latitude, λ is longitude, and h is altitude, the corresponding coordinate of the point in the J2000 cartesian frame, (x_t, y_t, z_t) , can be calculated using the following algorithm (Zhu, 1994):

$$x_t = (N + h) \cos \phi \cos \lambda, \quad (2)$$

$$y_t = (N + h) \cos \phi \sin \lambda, \quad (3)$$

$$z_t = \left((1 - e^2)N + h \right) \sin \phi, \quad (4)$$



where e is the eccentricity of the ellipsoid describing Earth's sea level (e^2 is 0.00669437999014 for WGS84), and the radius of curvature (N) at latitude ϕ is calculated as follows:

$$N = \frac{a}{\sqrt{1 - e^2 \sin^2 \phi}}, \quad (5)$$

145 where a is the radius at the equator, 6378.137 km in WGS84.

The “look vector” is then simply the difference between the two points

$$(x_t - x_s)\hat{i} + (y_t - y_s)\hat{j} + (z_t - z_s)\hat{k}, \quad (6)$$

where (x_s, y_s, z_s) is the coordinate of the satellite in J2000, and \hat{i} , \hat{j} , and \hat{k} are unit vectors in the x , y , and z directions, respectively. However, determining the angle relative to vector winds requires calculating the projection of this vector onto a
150 plane normal to the tangent point that contains either an eastward-pointing or northward-pointing vector. Due to the symmetry of the assumed ellipsoid shape, it is more convenient to find a plane normal to the tangent point that contains an eastward-pointing vector. The equation for the normal plane can be expressed as follows:

$$normal(x, y, z) = \frac{\partial f}{\partial x}(x_t, y_t, z_t) \cdot [x - x_t] + \frac{\partial f}{\partial y}(x_t, y_t, z_t) \cdot [y - y_t] + \frac{\partial f}{\partial z}(x_t, y_t, z_t) \cdot [z - z_t], \quad (7)$$

where, because at a given z the “surface” of interest consists of a circle in the x/y plane:

$$155 \quad f(x, y, z) = x^2 + y^2 + g(z). \quad (8)$$

The derivative of the function $g(z)$ effectively represents the tilt of the plane as a function of z . However, the angle between the projection of the look vector on the plane and a vector in the plane at constant z (which will be parallel to the East/West direction vector at the tangent point) is not sensitive to the tilt of the plane in the z direction. Therefore, for simplicity, $g(z)$ can be set to a constant, chosen to be zero. This makes $f(x, y, z)$ a cylinder with its axis along the z -axis in J2000.

160 A vector orthogonal to the plane is therefore:

$$\frac{\partial f}{\partial x}(x_t, y_t, z_t)\hat{i} + \frac{\partial f}{\partial y}(x_t, y_t, z_t)\hat{j} + \frac{\partial f}{\partial z}(x_t, y_t, z_t)\hat{k} = 2x_t\hat{i} + 2y_t\hat{j} + 0\hat{k}. \quad (9)$$

The projection of the satellite's position on the plane (x_p, y_p, z_p) can be found using the dot product of the look vector with the (normalized) vector orthogonal to the plane:

$$\beta = \frac{\left((x_t - x_s)\hat{i} + (y_t - y_s)\hat{j} + (z_t - z_s)\hat{k} \right) \cdot (2x_t\hat{i} + 2y_t\hat{j})}{(2x_t\hat{i} + 2y_t\hat{j}) \cdot (2x_t\hat{i} + 2y_t\hat{j})} = \frac{2x_t(x_t - x_s) + 2y_t(y_t - y_s)}{4x_t^2 + 4y_t^2}, \quad (10)$$

$$165 \quad x_p = x_s + \frac{\partial f}{\partial x}(x_t, y_t, z_t)\beta = x_s + 2\beta x_t, \quad (11)$$

$$y_p = x_s + \frac{\partial f}{\partial y}(x_t, y_t, z_t)\beta = y_s + 2\beta y_t, \quad (12)$$

$$z_p = x_s + \frac{\partial f}{\partial z}(x_t, y_t, z_t)\beta = z_s. \quad (13)$$



The angle (θ) between the projected vector and an eastward-pointing vector in the plane then becomes

$$\theta = \tan^{-1} \left(\frac{z_p - z_t}{\sqrt{(x_p - x_t)^2 + (y_p - y_t)^2 + (z_p - z_t)^2}} \right). \quad (14)$$

170 Unfortunately, there is an ambiguity regarding whether the vector is directed to the East or to the West. To determine this, the longitude of the satellite (ϕ_s) is calculated:

$$\phi_s = 2 \tan^{-1} \left(\frac{\sqrt{x_s^2 + y_s^2} - x_s}{y_s} \right). \quad (15)$$

By simple geometry, if this longitude is West of the longitude of the measurement tangent point, the projected vector is directed westward from the tangent point, and if it is East of the tangent point, the projected vector is eastward. Knowing the angle of
175 the projected vector relative to an eastward-pointing vector, the heading angle reported in the ACE-FTS wind data (relative to a northward-pointing vector) can be readily determined.

Figure 2 compares the heading angle (relative to a northward pointing vector) determined using the new approach to values determined previously using STK for sunset occultations in 2019. Both approaches exhibit occasional apparent jumps in heading when the East/West pointing direction can be difficult to decipher, such as when the satellite is on the opposite side
180 of the pole from the tangent point. Overall, there is a general agreement in how the heading angle varies over the course of the year, but differences between the two angles often exceed 20° . Agreement is best when the heading is close to either north/south or East/West and worst when the tangent point is near one of the poles. The STK calculations exhibit significantly greater variability.




Figure 2. Calculated heading angle of the ACE-FTS line-of-sight vector for sunset occultations in 2019. Calculations for v.5.3 winds using J2000 coordinates for both the satellite and the tangent point are shown in blue, while calculations from previous ACE-FTS wind versions using STK are shown in orange.



3 ACE v.5.3 Validation

185 3.1 MIGHTI vs ACE v.5.3

The MIGHTI instrument uses a Michelson interferometer and utilizes the Doppler Asymmetric Spatial Heterodyne (DASH) technique to determine shifts in emission lines (Englert et al., 2017). The emission lines observed are the red ($O(^1D-^3P)$ 630.0 nm) and green ($O(^1S-^1D)$ 557.7 nm) lines of the oxygen atom. We only compare with winds derived from the green line data products, as the red line is less frequent below 150 km. MIGHTI uses two sensors that record measurements 5 to 190 8 minutes apart. These measurements can be used to determine vector winds due to being orthogonal to each other, with MIGHTI-A observing at 45° and MIGHTI-B at 135° relative to the satellite's motion.

A detailed comparison with ACE v.5.2 line-of-sight winds with MIGHTI vector winds is available in Johnson et al. (2024). We provide a similar comparison of ACE v.5.3 against v05 MIGHTI using 207 of the same sunrise and 215 sunset coincidences between 2019 and 2022 used in the previous paper. Those coincidences are measurements within 2.5° latitude and 5° longitude 195 in location and 2 hours in time. In the case that there were multiple MIGHTI measurements that fit this criteria for one ACE occultation, the spatially closest measurement was used.

In Fig. 3, we show a comparison between an individual ACE sunrise and sunset occultation and the coincident MIGHTI measurement. It is important to note that the heading angle differs in v.5.2 and v.5.3, meaning that the converted MIGHTI winds would be different for each. However, the resulting speeds are very similar, so we display the line-of-sight winds converted 200 using v.5.3 heading angles. There is similar agreement for both v.5.2 and v.5.3 with MIGHTI when looking at the sunrise coincidence. The sunset coincidence shows slightly better agreement using v.5.3.




Figure 3. Comparison between individual ACE v.5.2 and v.5.3 (a) sunrise and (b) sunset coincidences with MIGHTI measurements. MIGHTI wind speeds were converted using the ACE v.5.3 heading angle.

In an effort to remove outliers and find trends in the data, we use the same restrictions as Data Set 2 from Johnson et al. (2024). Data Set 2 used only the coincidences where the average difference between ACE and MIGHTI wind speeds was



less than 60 m/s, leaving us with 184 sunrises and 196 sunsets. The average of these MIGHTI and ACE measurements are presented in Fig. 4. Looking at the sunrises, there is similar profile agreement at all altitudes for both v.5.2 and v.5.3. The sunset comparison shows good agreement for both versions from 90 to 110 km and then deviates. There is a sunrise-sunset bias of around ± 15 m/s, shown in pink in Fig. 4.




Figure 4. Average wind speeds of (a) 184 sunrise and (b) 196 sunset ACE v.5.2 and v.5.3 coincidences with MIGHTI measurements. MIGHTI wind speeds were converted using the ACE v.5.3 heading angles.

3.2 Meteor Radar vs ACE v.5.3

The Institute of Geology and Geophysics of the Chinese Academy of Science has five meteor radar stations in China providing horizontal wind profiles: Mohe at 52.5° N, 122.3° E; Beijing at 40.3° N, 116.2° E; Wuhan at 30.5° N, 114.6° E; Ledong at 18.7° N, 109.2° E; and Sanya at 18.3° N, 109.6° E. The stations use VHF radar and five receivers to track the motion of meteoric trails. The measurements are reported hourly and provide an all-sky coverage (Tang et al., 2021). A detailed comparison of meteor radar data with ACE v.5.2 is provided in Johnson et al. (2024). A comparison with ACE v.5.3 is provided here, using a similar process. The spatial range for a coincidence is the same as used with MIGHTI: 2.5° latitude and 5° longitude in location and 2 hours in time.

An individual comparison between an ACE sunrise and sunset occultation with the corresponding meteor radar measurements is shown in Fig. 5. These occultations were chosen to be shown as the coincident meteor radar measurement covers a large altitude range, whereas some only provide data in a small altitude window. We again only show the meteor radar winds converted using the v.5.3 heading angles. There is excellent profile agreement for both the sunrise and sunset comparisons.




Figure 5. Comparison between individual ACE v.5.2 and v.5.3 (a) sunrise and (b) sunset coincidences with meteor radar measurements. Meteor radar wind speeds were converted using the ACE v.5.3 heading angle.

220 Unlike the comparison with MIGHTI, the number of coincidences here is small, so the entirety of the coincident data set is considered. There were 26 sunrise and 12 sunset coincidences. Of those, 18 were from the Mohe station, 12 from Beijing, and 4 from Wuhan and Ledong each. The averages of these coincidences are shown in Fig. 6. For the sunrises, we again see well matching profiles. The sunsets we see general agreement but with ACE showing more prominent features. The sunrise-sunset bias of $\sim \pm 15$ m/s is evident in this comparison.




Figure 6. Average wind speeds of (a) 26 sunrise and (b) 12 sunset ACE v.5.2 and v.5.3 coincidences with meteor radar measurements. Meteor radar wind speeds were converted using the ACE v.5.3 heading angles.



225 4 Comparisons with Models

4.1 MERRA-2 vs ACE

MERRA-2 provides instantaneous 3-dimensional 3-hourly horizontal vector winds. The model is best constrained up to ~ 40 km, but extends up to ~ 70 km. Because the model is a reanalysis, we are able compare individual ACE occultations as well as an average of measurements with the corresponding MERRA-2 data. To do this, we find the nearest MERRA-2 time to
230 the ACE occultation. Then, we bilinearly interpolate the nearest coordinates of MERRA-2 vector winds to the given latitude and longitude of the ACE occultation. The MERRA-2 vector winds are then converted to line-of-sight winds. We again only display the MERRA-2 profiles that are calculated using the v.5.3 heading angles.

A sample of individual sunrise and sunset occultation comparisons at northern, tropical, and southern latitudes can be seen in Fig. 7. These occultations were chosen as they show particularly strong altitude profile agreement, however, other occultations
235 show similar results. We see a decrease in bias for both sunrise and sunset occultations in v.5.3 compared with v.5.2 as well as better matching profile. These improvements are best seen below ~ 40 km, where MERRA-2 is better constrained. Near 52 km in Fig. 7(a), there is a sharp eastward increase in ACE v.5.3 wind speed. This shift occurs at the boundary of segments 2 and 3. There was probably an issue at the top of the segment 2 retrieval or at the bottom of the segment 3 retrieval for this occultation.




Figure 7. Individual ACE line-of-sight wind speeds compared with the corresponding MERRA-2 wind speeds. Plots (a-c) display sunrise comparisons taken during 2019 in the northern, tropical, and southern latitude regions, respectively. Plots (d-f) display sunset comparisons in the same regions above. (a) sr86572 taken at 46° N on September 7th, 2019. (b) sr84637 taken at 1° N on April 29th, 2019. (c) sr85581 taken at 45° S on July 2nd, 2019. (d) ss86041 taken at 45° N on August 2nd, 2019. (e) ss86206 taken at 1° S on August 13th, 2019. (f) ss86293 taken at 45° S on August 19th, 2019.

240 All ACE line-of-sight wind speed measurements in 2019 and their corresponding MERRA-2 wind speeds are averaged and compared in Fig. 8. There is a large profile disagreement between 40 and 60 km for ACE v.5.2 sunrises that now shows



good agreement in v.5.3. Comparing the sunsets, we see similar profile agreement but a decrease in bias, particularly at lower altitudes.




Figure 8. (a) Average wind speed profiles from all ACE sunrise measurements in 2019 for ACE v.5.2 (3535 occultations), ACE v.5.3 (3535 occultations), and MERRA-2. (b) Average wind speed profiles from all ACE sunset measurements in 2019 for ACE v.5.2 (3419 occultations), ACE v.5.3 (3422 occultations), and MERRA-2. MERRA-2 wind speeds were converted using the ACE v.5.3 heading angles.

We are also able to derive a sunrise-sunset bias by comparing with MERRA-2. To do this, we shift the MERRA-2 altitude to the nearest ACE altitude (maximum of ± 0.5 km) and find the difference between ACE and MERRA-2 wind speeds. The differences are then averaged over all measurements by ACE in 2019 and the results are shown in Fig. 9(a) for v.5.2 and (c) for v.5.3. We then subtract the sunrise and sunset averages from the total average to find the bias. The average bias at each altitude is displayed in Fig. 9(b) for v.5.2 and (d) for v.5.3. We find that MERRA-2 and ACE differ by less than 10 m/s up to 50 km with v.5.3, which is about half of the difference with v.5.2. We also find that the sunrise-sunset bias has strongly improved. It is now within ± 5 m/s up through 50 km, but largely within ± 3 m/s. Above 50 km the bias does grow to $\sim \pm 10$ m/s, as seen in v.5.2. However, this higher altitude region is not well constrained within the MERRA-2 reanalysis.




Figure 9. (a) Average difference between ACE v.5.2 (sunrise (blue), sunset (red), and both (black)) and MERRA-2 winds for each ACE wind speed measurement in 2019. (b) ACE sunrise-sunset bias, shown as the difference between sunrise and sunset averages and the total average in (a). Plots (c) and (d) are the same as (a) and (b), respectively, but for ACE v.5.3 processing.

4.1.1 Zonal Winds

Twice a year, around the equinoxes, the line-of-sight is such that the ACE satellite measures the East-West (or West-East) wind component, called the zonal wind. A substantial number of ACE occultations with a line-of-sight within plus or minus 10° of East-West (or West-East) can serve to map the zonal wind. For the year 2019, 427 sunrise occultations from March 1st (17° S, heading 278°) to April 5th (82° S, heading 260°) and 483 sunset occultations from September 6th (81° S, heading 100°) to October 19th (41° N, heading 80°) fall within the $\pm 10^\circ$ zonal wind category in the southern hemisphere. These occultations are more densely distributed near the pole. For 2° latitude bins, all wind profiles were averaged and the resulting maps of the $\pm 10^\circ$ zonal wind intensity as a function of altitude and latitude are shown in Fig. 10(a) and (c). The March-April wind signs were inverted so that positive zonal winds (in red) indicate winds moving from West to East. Note that the heading angle is exactly 270° (respectively 90°) for latitude around 73° S (71° S), which correspond to March 20th (September 23rd).

ACE measures the polar vortex near 30 km altitude in September-October. As expected, the vortex is weak in March-April. In September-October, the mesosphere is characterized by strong positive zonal winds in the whole hemisphere. It is well known that winds of the thermosphere are variable with local time. Because of ACE observation geometry, the sampled local times for sunrises (March-April) are typically between 6:00 am and 9:00 am and the sampled local times for sunsets (September-October) are typically between 15:00 pm and 18:00 pm. For sunrises in March-April, the thermosphere zonal



winds are positive around 100 km altitude and negative above. At sunsets in September-October, strong negative zonal winds are observed at 60° S and altitude 130 km.

270 MERRA-2 reanalysis wind vectors were sampled for each of the occultations and projected into the corresponding ACE line-of-sight direction. The latitude-bin-averaged MERRA-2 winds are shown in Fig. 10(b) and (d). For the September-October period, the MERRA-2 winds below 40 km are in good agreement with ACE, with errors lower than 15 m/s. Above 40 km, the MERRA-2 winds are weaker by about 20 m/s compared to ACE. For the March-October period, the MERRA-2 winds mostly agree with the ACE winds with errors less than 10 m/s. The strength of the negative zonal winds around 20° S and altitude 40 km is overestimated by MERRA-2 by around 15 m/s.

275 Overall, the ACE data show sharper wind features than MERRA-2 in the stratosphere and an intriguing wave-like pattern spanning 80° S to 20° S in both time periods. This pattern could be an artifact due to the smooth varying line-of-sight angle with latitude combined with winds in the non-tangent layers. At altitudes above 80 km, ACE provides a unique measurement of the complex dynamics of the thermosphere.




Figure 10. ACE measurements of the $\pm 10^\circ$ zonal winds as a function of latitude and altitude (a, c) and corresponding MERRA-2 winds (b, d) for year 2019. The September-October observations (a, b) correspond to sunset solar occultations with local time at the tangent point between 6:00 am and 9:00 am and the March-April observations (c, d) correspond to sunrise solar occultations with local time at the tangent point between 15:00 pm and 18:00 pm.

4.2 HWM14 vs ACE

HWM14 outputs for all of 2019 on a $5^\circ \times 5^\circ$ grid with 15 minute time increments using observational geomagnetic ap indexes were retrieved from the Community Coordinated Modeling Center (CCMC) based out of NASA Goddard Space Flight Center (GSFC). The data ranges from the surface up to 1000 km in 10 km intervals.

Since HWM14 is a climatology, we only compare the average of many ACE and corresponding HWM14 wind speeds. We again compare all ACE wind speed measurements available from 2019. We first separate the ACE occultations into northern (latitude $> 45^\circ$ N), tropical (latitude between 30° S and 30° N), and southern (latitude $< 45^\circ$ S) regions based on latitude. It is not reasonable for a model to predict wind patterns within the polar vortex during winter, so measurements during those periods are removed. Similar to MERRA-2, we then find the nearest HWM14 time to a given ACE occultation and bilinearly interpolate the vector winds of the nearby HWM14 coordinates to the given ACE coordinates. The HWM14 vector winds are then converted to line-of-sight winds.



290 This average of ACE v.5.2 and v.5.3 wind speeds from 2019 and their corresponding HWM14 wind speeds are shown in
Fig. 11. The displayed HWM14 wind speeds are calculated using the v.5.3 heading angles. We see general profile agreement
for both processing versions, with v.5.3 being slightly improved. ACE v.5.2 shows more prominent features than HWM14,
best seen at the ~ 90 and ~ 60 km features in Fig. 11(a) and (e), respectively. ACE v.5.3 still displays a prominent feature, but
better matches the model in both cases. The sunrise-sunset bias is not clear at lower altitudes, but does become more notable
295 above ~ 80 km.




Figure 11. Average line-of-sight wind speeds from ACE measurements in 2019 for ACE v.5.2, ACE v.5.3, and HWM14. HWM14 wind speeds were converted using the ACE v.5.3 heading angles. **(a-c)** are from sunrise measurements in the northern (latitude $> 45^\circ$ N), tropical (latitude between 30° S and 30° N), and southern (latitude $< 45^\circ$ S) regions, respectively. **(d-f)** match the same regions above, but for sunset measurements.

4.3 WACCM-X vs ACE

Specified Dynamics WACCM-X Version 2.2 provides global vector winds in 3 hour intervals. WACCM-X vector winds were compared with ACE-FTS line-of-sight winds for all of 2019 in a similar fashion as HWM14. The resulting line-of-sight winds are shown in Fig. 12. There is notably less agreement between both ACE versions and the WACCM-X profiles compared with
300 HWM14. Both ACE sunrise and sunset averages in the northern and southern latitude region show additional features not found in the WACCM-X model, although v.5.3 agrees more than v.5.2. The model and ACE data show the best profile agreement in the tropical region. The eastward bias in v.5.2 has also slightly decreased with v.5.3, particularly at lower altitudes.




Figure 12. Average line-of-sight wind speeds from ACE measurements in 2019 for ACE v.5.2 (black), ACE v.5.3 (red), and WACCM-X (blue). WACCM-X wind speeds were converted using the ACE v.5.3 heading angles. (a-c) are from sunrise measurements in the northern (latitude $> 45^\circ$ N), tropical (latitude between 30° S and 30° N), and southern (latitude $< 45^\circ$ S) regions, respectively. (d-f) match the same regions above, but for sunset measurements.

5 Conclusions

ACE v.5.3 wind speeds have been validated by instrument observations from MIGHTI and meteor radar. Wind speed profiles from ACE v.5.3 for sunrise and sunset occultations show improved profile agreement with MIGHTI and meteor radar measurements. The ± 15 m/s sunrise-sunset bias previously found is still seen at these higher altitudes (above 80 km).

ACE v.5.2 and v.5.3 line-of-sight wind speeds have been compared with vector winds from the MERRA-2, HWM14, and WACCM-X models. The new wind speeds have better profile agreement than v.5.2 does with each model. The v.5.3 processing has particularly strong profile agreement with MERRA-2 below 40 km, where the model is well constrained. The MERRA-2 comparison shows a decreased sunrise-sunset bias of less than ± 5 m/s below 50 km. The previously found ± 15 m/s bias is still seen with v.5.3 at higher altitudes.

The ACE-FTS instrument on the Canadian SCISAT satellite now provides line-of-sight wind speeds between 18 and 140 km dating back to 2004 with v.5.2 and the improved v.5.3 processing. There is a lack of wind speed measurements within this altitude region from other sources, making ACE wind speeds a valuable addition for model evaluation.



Altitude Segment	Center (cm ⁻¹)	Width (cm ⁻¹)	Altitude Range (km)	Transmittance Range	Isotopologue
1	2602.30	0.4	18-35	0.4-0.97	¹⁸ OCO
1	2604.50	0.8	18-35	0.4-0.97	¹⁸ OCO
1	2606.00	0.8	18-35	0.4-0.97	¹⁸ OCO
1	2607.00	0.8	18-35	0.4-0.97	¹⁸ OCO
1	2608.00	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2609.80	0.4	18-35	0.4-0.97	¹⁸ OCO
1	2610.50	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2610.90	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2611.60	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2612.40	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2616.10	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2616.84	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2617.50	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2619.01	0.9	18-35	0.4-0.97	¹⁸ OCO
1	2620.50	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2621.25	1.1	18-35	0.4-0.97	¹⁸ OCO
1	2623.40	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2624.80	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2627.00	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2629.80	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2632.05	1.0	18-35	0.4-0.97	¹⁸ OCO
1	2636.30	1.0	18-35	0.4-0.97	¹⁸ OCO
2	1889.02	0.4	30-45	0.65-0.96	CO ₂
2	1891.97	0.3	30-50	0.65-0.96	CO ₂
2	1896.17	0.3	30-55	0.65-0.96	CO ₂
2	1897.69	0.3	30-55	0.65-0.96	CO ₂
2	1898.13	0.3	30-45	0.65-0.96	CO ₂
2	1899.17	0.3	35-60	0.65-0.96	CO ₂
2	1899.29	0.2	30-45	0.65-0.96	CO ₂
2	1902.15	0.3	35-60	0.65-0.96	CO ₂
2	1905.05	0.3	35-60	0.65-0.96	CO ₂
2	1905.37	0.3	30-45	0.65-0.96	CO ₂
2	1906.43	0.3	35-60	0.65-0.96	CO ₂
Continued on next page					



Altitude Segment	Center (cm ⁻¹)	Width (cm ⁻¹)	Altitude Range (km)	Transmittance Range	Isotopologue
2	1908.37	0.3	30-45	0.65-0.96	CO ₂
2	1909.45	0.3	35-60	0.65-0.96	CO ₂
2	1911.01	0.3	35-60	0.65-0.96	CO ₂
2	1912.53	0.3	45-60	0.65-0.96	CO ₂
2	1914.11	0.3	35-60	0.65-0.96	CO ₂
2	1915.47	0.3	35-60	0.65-0.96	CO ₂
2	1917.05	0.3	35-60	0.65-0.96	CO ₂
2	1920.11	0.3	35-60	0.65-0.96	CO ₂
2	1924.77	0.3	35-60	0.65-0.96	CO ₂
2	1927.81	0.3	30-60	0.65-0.96	CO ₂
2	1929.35	0.3	30-60	0.65-0.96	CO ₂
2	1930.85	0.3	30-50	0.65-0.96	CO ₂
2	1933.91	0.3	30-60	0.65-0.96	CO ₂
2	1935.16	0.4	30-60	0.65-0.96	CO ₂
2	1936.47	0.3	30-55	0.65-0.96	CO ₂
2	1941.05	0.3	30-50	0.65-0.96	CO ₂
2	1950.67	0.3	30-55	0.65-0.96	CO ₂
2	1966.85	0.3	30-55	0.65-0.96	CO ₂
2	1970.31	0.3	30-55	0.65-0.96	CO ₂
2	1971.77	0.3	30-50	0.65-0.96	CO ₂
2	1975.05	0.3	35-45	0.65-0.96	CO ₂
3	2039.93	0.3	50-77	0.65-0.97	CO ₂
3	2042.95	0.3	50-77	0.65-0.97	CO ₂
3	2044.45	0.3	50-77	0.65-0.97	CO ₂
3	2045.95	0.3	50-77	0.65-0.97	CO ₂
3	2047.51	0.3	50-77	0.65-0.97	CO ₂
3	2049.03	0.3	50-77	0.65-0.97	CO ₂
3	2050.55	0.3	50-77	0.65-0.97	CO ₂
3	2052.09	0.3	50-77	0.65-0.97	CO ₂
3	2053.57	0.3	50-77	0.65-0.97	CO ₂
3	2055.15	0.3	50-77	0.65-0.97	CO ₂
3	2056.69	0.3	50-77	0.65-0.97	CO ₂
3	2058.29	0.3	50-77	0.65-0.97	CO ₂
Continued on next page					



Altitude Segment	Center (cm ⁻¹)	Width (cm ⁻¹)	Altitude Range (km)	Transmittance Range	Isotopologue
3	2059.73	0.3	50-77	0.65-0.97	CO ₂
3	2061.25	0.3	50-77	0.65-0.97	CO ₂
3	2062.87	0.3	50-77	0.65-0.97	CO ₂
3	2066.01	0.3	60-77	0.65-0.97	CO ₂
3	2067.51	0.3	60-77	0.65-0.97	CO ₂
3	2070.61	0.3	60-77	0.65-0.97	CO ₂
3	2072.17	0.3	60-77	0.65-0.97	CO ₂
4	2232.95	0.3	70-80	0.67-0.975	¹³ CO ₂
4	2237.27	0.3	70-90	0.67-0.975	¹³ CO ₂
4	2239.39	0.3	70-90	0.67-0.975	¹³ CO ₂
4	2243.59	0.3	70-95	0.67-0.975	¹³ CO ₂
4	2245.65	0.3	70-95	0.67-0.975	¹³ CO ₂
4	2248.71	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2249.66	0.3	70-95	0.67-0.975	¹³ CO ₂
4	2250.49	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2251.35	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2251.65	0.3	70-95	0.67-0.975	¹³ CO ₂
4	2252.31	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2253.19	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2253.75	0.3	70-100	0.67-0.975	¹³ CO ₂
4	2254.09	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2254.91	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2255.59	0.3	70-105	0.67-0.975	¹³ CO ₂
4	2257.60	0.4	85-105	0.67-0.975	¹³ CO ₂
4	2260.18	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2261.20	0.4	85-110	0.67-0.975	¹³ CO ₂
4	2261.93	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2262.75	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2263.21	0.3	70-110	0.67-0.975	¹³ CO ₂
4	2264.95	0.3	70-110	0.67-0.975	¹³ CO ₂
4	2266.07	0.3	70-85	0.67-0.975	¹³ CO ₂
4	2268.53	0.3	85-110	0.67-0.975	¹³ CO ₂
4	2270.29	0.3	90-110	0.67-0.975	¹³ CO ₂
Continued on next page					



Altitude Segment	Center (cm ⁻¹)	Width (cm ⁻¹)	Altitude Range (km)	Transmittance Range	Isotopologue
4	2272.03	0.3	70-110	0.67-0.975	¹³ CO ₂
4	2273.73	0.3	90-110	0.67-0.975	¹³ CO ₂
4	2275.41	0.3	70-110	0.67-0.975	¹³ CO ₂
4	2277.07	0.3	90-110	0.67-0.975	¹³ CO ₂
4	2278.60	0.4	80-110	0.67-0.975	¹³ CO ₂
4	2280.33	0.3	80-110	0.67-0.975	¹³ CO ₂
4	2281.85	0.3	85-110	0.67-0.975	¹³ CO ₂
4	2284.33	0.3	80-110	0.67-0.975	¹³ CO ₂
4	2285.86	0.3	80-110	0.67-0.975	¹³ CO ₂
4	2287.33	0.3	90-110	0.67-0.975	¹³ CO ₂
4	2288.79	0.3	85-110	0.67-0.975	¹³ CO ₂
4	2290.25	0.3	90-110	0.67-0.975	¹³ CO ₂
4	2291.79	0.3	80-110	0.67-0.975	¹³ CO ₂
4	2293.09	0.3	90-110	0.67-0.975	¹³ CO ₂
4	2294.53	0.3	90-110	0.67-0.975	¹³ CO ₂
4	2299.87	0.3	90-110	0.67-0.975	¹³ CO ₂
4	2302.29	0.3	90-110	0.67-0.975	¹³ CO ₂
4	2303.47	0.3	90-110	0.67-0.975	¹³ CO ₂
5	2300.50	0.4	95-120	0.67-0.985	CO ₂
5	2302.65	0.3	95-120	0.67-0.985	CO ₂
5	2306.95	0.3	95-140	0.67-0.985	CO ₂
5	2309.03	0.3	100-140	0.67-0.985	CO ₂
5	2309.63	0.3	95-105	0.67-0.985	CO ₂
5	2311.11	0.3	105-140	0.67-0.985	CO ₂
5	2311.59	0.3	95-105	0.67-0.985	CO ₂
5	2313.15	0.3	95-140	0.67-0.985	CO ₂
5	2314.29	0.3	95-110	0.67-0.985	CO ₂
5	2315.21	0.3	95-140	0.67-0.985	CO ₂
5	2316.07	0.3	95-115	0.67-0.985	CO ₂
5	2317.19	0.3	110-140	0.67-0.985	CO ₂
5	2319.10	0.4	95-140	0.67-0.985	CO ₂
5	2319.89	0.3	95-115	0.67-0.985	CO ₂
5	2320.75	0.3	95-115	0.67-0.985	CO ₂

Continued on next page



Altitude Segment	Center (cm ⁻¹)	Width (cm ⁻¹)	Altitude Range (km)	Transmittance Range	Isotopologue
5	2321.11	0.3	105-140	0.67-0.985	CO ₂
5	2321.61	0.3	95-115	0.67-0.985	CO ₂
5	2322.51	0.3	95-115	0.67-0.985	CO ₂
5	2323.11	0.3	95-140	0.67-0.985	CO ₂
5	2323.37	0.3	95-115	0.67-0.985	CO ₂
5	2324.24	0.3	95-115	0.67-0.985	CO ₂
5	2325.00	0.4	95-140	0.67-0.985	CO ₂
5	2328.63	0.3	95-140	0.67-0.985	CO ₂
5	2329.39	0.3	95-115	0.67-0.985	CO ₂
5	2330.17	0.3	95-115	0.67-0.985	CO ₂
5	2330.55	0.3	105-140	0.67-0.985	CO ₂
5	2331.03	0.3	95-115	0.67-0.985	CO ₂
5	2331.85	0.3	95-115	0.67-0.985	CO ₂
5	2332.35	0.3	95-140	0.67-0.985	CO ₂
5	2332.65	0.3	95-115	0.67-0.985	CO ₂
5	2333.37	0.3	95-115	0.67-0.985	CO ₂
5	2334.15	0.3	105-140	0.67-0.985	CO ₂
5	2337.59	0.3	100-140	0.67-0.985	CO ₂
5	2338.93	0.3	95-110	0.67-0.985	CO ₂
5	2339.37	0.3	95-140	0.67-0.985	CO ₂
5	2339.65	0.3	95-115	0.67-0.985	CO ₂
5	2341.07	0.3	110-140	0.67-0.985	CO ₂
5	2343.39	0.3	95-115	0.67-0.985	CO ₂
5	2344.09	0.3	95-115	0.67-0.985	CO ₂
5	2344.37	0.3	110-140	0.67-0.985	CO ₂
5	2344.77	0.3	95-115	0.67-0.985	CO ₂
5	2346.05	0.3	95-140	0.67-0.985	CO ₂
5	2347.59	0.3	110-140	0.67-0.985	CO ₂
5	2348.31	0.3	95-115	0.67-0.985	CO ₂
5	2348.97	0.3	95-115	0.67-0.985	CO ₂
5	2349.55	0.3	95-115	0.67-0.985	CO ₂
5	2350.01	0.3	100-140	0.67-0.985	CO ₂
5	2351.45	0.3	110-140	0.67-0.985	CO ₂
Continued on next page					



Altitude Segment	Center (cm^{-1})	Width (cm^{-1})	Altitude Range (km)	Transmittance Range	Isotopologue
5	2352.31	0.3	95-115	0.67-0.985	CO ₂
5	2354.43	0.3	105-140	0.67-0.985	CO ₂
5	2355.43	0.3	95-115	0.67-0.985	CO ₂
5	2355.89	0.3	105-140	0.67-0.985	CO ₂
5	2357.31	0.3	105-140	0.67-0.985	CO ₂
5	2358.73	0.3	105-140	0.67-0.985	CO ₂
5	2361.47	0.3	100-140	0.67-0.985	CO ₂
5	2362.79	0.3	105-140	0.67-0.985	CO ₂
5	2364.11	0.3	95-140	0.67-0.985	CO ₂
5	2365.39	0.3	100-140	0.67-0.985	CO ₂
5	2366.65	0.3	95-140	0.67-0.985	CO ₂
5	2367.87	0.3	95-140	0.67-0.985	CO ₂
5	2369.09	0.3	95-140	0.67-0.985	CO ₂
5	2370.27	0.3	95-140	0.67-0.985	CO ₂
5	2371.43	0.3	95-140	0.67-0.985	CO ₂
5	2372.55	0.3	95-140	0.67-0.985	CO ₂
5	2373.67	0.3	95-140	0.67-0.985	CO ₂
5	2374.75	0.3	95-140	0.67-0.985	CO ₂
5	2375.79	0.3	95-140	0.67-0.985	CO ₂
5	2376.83	0.3	95-140	0.67-0.985	CO ₂
5	2377.85	0.3	95-140	0.67-0.985	CO ₂
5	2378.83	0.3	95-140	0.67-0.985	CO ₂
5	2379.79	0.3	95-140	0.67-0.985	CO ₂
5	2380.71	0.3	95-140	0.67-0.985	CO ₂
5	2381.61	0.3	95-140	0.67-0.985	CO ₂
5	2382.49	0.3	95-140	0.67-0.985	CO ₂
5	2383.35	0.3	95-140	0.67-0.985	CO ₂

Table A1. Microwindow sets employed in the 5 altitude segments for the wind retrieval from ACE-FTS measurements, along with the allowed range for minimum transmittance in the window used to filter saturated or excessively weak lines from the analysis.

Author contributions. **Matthew Wyatt:** Formal analysis, Writing - original draft. **Peter F. Bernath:** Supervision, Writing - review and editing. **Chris Boone:** Investigation, Writing - original draft. **Léo Lavvy:** Formal analysis, Writing - original draft. **Ryan Johnson:** Investigation.



Competing interests. The authors declare that they have no conflict of interest.

325 *Acknowledgements.* The ACE mission is funded by the Canadian Space Agency (9F045-230412/001). Some support was provided by NASA
ACMAP, Atmospheric Composition Modeling and Analysis Program (80NSSC23K0999). PB acknowledges RB for productive discussion.



References

- Systems Tool Kit version 10; software used for satellite orbit propagation, <https://www.ansys.com/products/missions/ansys-stk>.
- Baron, P., Murtagh, D. P., Urban, J., Sagawa, H., Ochiai, S., Kasai, Y., Kikuchi, K., Khosrawi, F., Körnich, H., Mizobuchi, S., Sagi, K., and Yasui, M.: Observation of horizontal winds in the middle-atmosphere between 30° S and 55° N during the northern winter 2009–2010, *Atmospheric Chemistry and Physics*, 13, 6049–6064, <https://doi.org/10.5194/acp-13-6049-2013>, 2013.
- Baumgarten, G.: Doppler Rayleigh/Mie/Raman lidar for wind and temperature measurements in the middle atmosphere up to 80 km, *Atmospheric Measurement Techniques*, 3, 1509–1518, <https://doi.org/10.5194/amt-3-1509-2010>, 2010.
- Bernath, P.: The Atmospheric Chemistry Experiment (ACE), *Journal of Quantitative Spectroscopy and Radiative Transfer*, 186, 3–16, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2016.04.006>, satellite Remote Sensing and Spectroscopy: Joint ACE-Odin Meeting, October 2015, 2017.
- Bernath, P. F., McElroy, C. T., Abrams, M. C., Boone, C. D., Butler, M., Camy-Peyret, C., Carleer, M., Clerbaux, C., Coheur, P.-F., Colin, R., DeCola, P., DeMazière, M., Drummond, J. R., Dufour, D., Evans, W. F. J., Fast, H., Fussen, D., Gilbert, K., Jennings, D. E., Llewellyn, E. J., Lowe, R. P., Mahieu, E., McConnell, J. C., McHugh, M., McLeod, S. D., Michaud, R., Midwinter, C., Nassar, R., Nichitiu, F., Nowlan, C., Rinsland, C. P., Rochon, Y. J., Rowlands, N., Semeniuk, K., Simon, P., Skelton, R., Sloan, J. J., Soucy, M.-A., Strong, K., Tremblay, P., Turnbull, D., Walker, K. A., Walkty, I., Wardle, D. A., Wehrle, V., Zander, R., and Zou, J.: Atmospheric Chemistry Experiment (ACE): Mission overview, *Geophysical Research Letters*, 32, <https://doi.org/https://doi.org/10.1029/2005GL022386>, 2005.
- Boone, C., Bernath, P., and Lecours, M.: Version 5 retrievals for ACE-FTS and ACE-imagers, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 310, 108 749, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2023.108749>, 2023.
- Boone, C. D., Steffen, J., Crouse, J., and Bernath, P. F.: Line-of-Sight Winds and Doppler Effect Smearing in ACE-FTS Solar Occultation Measurements, *Atmosphere*, 12, <https://doi.org/10.3390/atmos12060680>, 2021.
- Buehner, M., McTaggart-Cowan, R., Beaulne, A., Charette, C., Garand, L., Heillette, S., Lapalme, E., Laroche, S., Macpherson, S. R., Morneau, J., and Zadra, A.: Implementation of Deterministic Weather Forecasting Systems Based on Ensemble–Variational Data Assimilation at Environment Canada. Part I: The Global System, *Monthly Weather Review*, 143, 2532 – 2559, <https://doi.org/10.1175/MWR-D-14-00354.1>, 2015.
- Drob, D. P., Emmert, J. T., Crowley, G., Picone, J. M., Shepherd, G. G., Skinner, W., Hays, P., Niciejewski, R. J., Larsen, M., She, C. Y., Meriwether, J. W., Hernandez, G., Jarvis, M. J., Sipler, D. P., Tepley, C. A., O'Brien, M. S., Bowman, J. R., Wu, Q., Murayama, Y., Kawamura, S., Reid, I. M., and Vincent, R. A.: An empirical model of the Earth's horizontal wind fields: HWM07, *Journal of Geophysical Research: Space Physics*, 113, <https://doi.org/https://doi.org/10.1029/2008JA013668>, 2008.
- Drob, D. P., Emmert, J. T., Meriwether, J. W., Makela, J. J., Doornbos, E., Conde, M., Hernandez, G., Noto, J., Zawdie, K. A., McDonald, S. E., Huba, J. D., and Klenzing, J. H.: An update to the Horizontal Wind Model (HWM): The quiet time thermosphere, *Earth and Space Science*, 2, 301–319, <https://doi.org/https://doi.org/10.1002/2014EA000089>, 2015.
- Duruiseau, F., Huret, N., Andral, A., and Camy-Peyret, C.: Assessment of the ERA-Interim Winds Using High-Altitude Stratospheric Balloons, *Journal of the Atmospheric Sciences*, 74, 2065 – 2080, <https://doi.org/10.1175/JAS-D-16-0137.1>, 2017.
- Englert, C. R., Harlander, J. M., Brown, C. M., Marr, K. D., Miller, I. J., Stump, J. E., Hancock, J., Peterson, J. Q., Kumler, J., Morrow, W. H., Mooney, T. A., Ellis, S., Mende, S. B., Harris, S. E., Stevens, M. H., Makela, J. J., Harding, B. J., and Immel, T. J.: Michelson Interferometer for Global High-Resolution Thermospheric Imaging (MIGHTI): Instrument Design and Calibration, , 212, 553–584, <https://doi.org/10.1007/s11214-017-0358-4>, 2017.



- 365 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R.,
Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R.,
Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and
Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419 –
5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 370 Gettelman, A., Mills, M. J., Kinnison, D. E., Garcia, R. R., Smith, A. K., Marsh, D. R., Tilmes, S., Vitt, F., Bardeen, C. G., McInerney, J., Liu,
H.-L., Solomon, S. C., Polvani, L. M., Emmons, L. K., Lamarque, J.-F., Richter, J. H., Glanville, A. S., Bacmeister, J. T., Phillips, A. S.,
Neale, R. B., Simpson, I. R., DuVivier, A. K., Hodzic, A., and Randel, W. J.: The Whole Atmosphere Community Climate Model Version 6
(WACCM6), *Journal of Geophysical Research: Atmospheres*, 124, 12 380–12 403, <https://doi.org/https://doi.org/10.1029/2019JD030943>,
2019.
- 375 Grassl, H. J., Skinner, W. R., Hays, P. B., Gell, D. A., Burrage, M. D., Ortland, D. A., Marshall, A. R., and Abreu, V. J.: Atmospheric wind
measurements with the high-resolution Doppler imager, *Journal of Spacecraft and Rockets*, 32, 169–176, <https://doi.org/10.2514/3.26590>,
1995.
- Hays, P. B., Wu, D. L., and Team, T. H. S.: Observations of the Diurnal Tide from Space, *Journal of Atmospheric Sciences*, 51, 3077 – 3093,
[https://doi.org/10.1175/1520-0469\(1994\)051<3077:OOTDTF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<3077:OOTDTF>2.0.CO;2), 1994.
- 380 Hedin, A., Fleming, E., Manson, A., Schmidlin, F., Avery, S., Clark, R., Franke, S., Fraser, G., Tsuda, T., Vial, F., and Vincent, R.:
Empirical wind model for the upper, middle and lower atmosphere, *Journal of Atmospheric and Terrestrial Physics*, 58, 1421–1447,
[https://doi.org/https://doi.org/10.1016/0021-9169\(95\)00122-0](https://doi.org/https://doi.org/10.1016/0021-9169(95)00122-0), 1996.
- Hedin, A. E., Spencer, N. W., and Killeen, T. L.: Empirical global model of upper thermosphere winds based on atmosphere and dynamics
Explorer satellite data, *Journal of Geophysical Research; (United States)*, 93:A9, <https://doi.org/10.1029/JA093iA09p09959>, 1988.
- 385 Johnson, R., Bernath, P., and Boone, C.: Validation of line-of-sight winds from ACE-FTS solar occultation measurements, *Journal of Quan-
titative Spectroscopy and Radiative Transfer*, 314, 108 870, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2023.108870>, 2024.
- Khelif, D., Burns, S. P., and Friehe, C. A.: Improved Wind Measurements on Research Aircraft, *Journal of Atmospheric and Oceanic
Technology*, 16, 860 – 875, [https://doi.org/10.1175/1520-0426\(1999\)016<0860:IWMORA>2.0.CO;2](https://doi.org/10.1175/1520-0426(1999)016<0860:IWMORA>2.0.CO;2), 1999.
- Killeen, T. L., Wu, Q., Solomon, S. C., Ortland, D. A., Skinner, W. R., Niciejewski, R. J., and Gell, D. A.: TIMED Doppler Interferometer:
Overview and recent results, *Journal of Geophysical Research: Space Physics*, 111, <https://doi.org/https://doi.org/10.1029/2005JA011484>,
390 2006.
- Kishore Kumar, G., Kishore Kumar, K., Baumgarten, G., and Ramkumar, G.: Validation of MERRA reanalysis upper-level winds
over low latitudes with independent rocket sounding data, *Journal of Atmospheric and Solar-Terrestrial Physics*, 123, 48–54,
<https://doi.org/https://doi.org/10.1016/j.jastp.2014.12.001>, 2015.
- Kleist, D. T., Parrish, D. F., Derber, J. C., Treadon, R., Wu, W.-S., and Lord, S.: Introduction of the GSI into the NCEP Global Data
395 Assimilation System, *Weather and Forecasting*, 24, 1691 – 1705, <https://doi.org/10.1175/2009WAF2222201.1>, 2009.
- Kumer, V.-M., Reuder, J., and Furevik, B. R.: A Comparison of LiDAR and Radiosonde Wind Measurements, *Energy Procedia*, 53, 214–
220, <https://doi.org/https://doi.org/10.1016/j.egypro.2014.07.230>, eERA DeepWind’ 2014, 11th Deep Sea Offshore Wind RD Conference,
2014.
- 400 Liu, A. Z., Hocking, W. K., Franke, S. J., and Thayaparan, T.: Comparison of Na lidar and meteor radar wind measurements at Starfire
Optical Range, NM, USA, *Journal of Atmospheric and Solar-Terrestrial Physics*, 64, 31–40, [https://doi.org/https://doi.org/10.1016/S1364-6826\(01\)00095-5](https://doi.org/https://doi.org/10.1016/S1364-6826(01)00095-5), 2002.



- 405 Liu, H.-L., Bardeen, C. G., Foster, B. T., Lauritzen, P., Liu, J., Lu, G., Marsh, D. R., Maute, A., McInerney, J. M., Pedatella, N. M., Qian, L., Richmond, A. D., Roble, R. G., Solomon, S. C., Vitt, F. M., and Wang, W.: Development and Validation of the Whole Atmosphere Community Climate Model With Thermosphere and Ionosphere Extension (WACCM-X 2.0), *Journal of Advances in Modeling Earth Systems*, 10, 381–402, <https://doi.org/https://doi.org/10.1002/2017MS001232>, 2018.
- Lux, O., Lemmerz, C., Weiler, F., Marksteiner, U., Witschas, B., Rahm, S., Geiß, A., and Reitebuch, O.: Intercomparison of wind observations from the European Space Agency’s Aeolus satellite mission and the ALADIN Airborne Demonstrator, *Atmospheric Measurement Techniques*, 13, 2075–2097, <https://doi.org/10.5194/amt-13-2075-2020>, 2020.
- 410 Martner, B. E., Wuertz, D. B., Stankov, B. B., Strauch, R. G., Westwater, E. R., Gage, K. S., Ecklund, W. L., Martin, C. L., and Dabberdt, W. F.: An Evaluation of Wind Profiler, RASS, and Microwave Radiometer Performance, *Bulletin of the American Meteorological Society*, 74, 599 – 614, [https://doi.org/10.1175/1520-0477\(1993\)074<0599:AEOWPR>2.0.CO;2](https://doi.org/10.1175/1520-0477(1993)074<0599:AEOWPR>2.0.CO;2), 1993.
- Molod, A., Takacs, L., Suarez, M., and Bacmeister, J.: Development of the GEOS-5 atmospheric general circulation model: evolution from MERRA to MERRA2, *Geoscientific Model Development*, 8, 1339–1356, <https://doi.org/10.5194/gmd-8-1339-2015>, 2015.
- 415 Putman, W. M. and Lin, S.-J.: Finite-volume transport on various cubed-sphere grids, *Journal of Computational Physics*, 227, 55–78, <https://doi.org/https://doi.org/10.1016/j.jcp.2007.07.022>, 2007.
- Rienecker, M. M., Suarez, M., Todling, R., Bacmeister, J., Takacs, L., Liu, H.-C., Gu, W., Sienkiewicz, M., Koster, R., Gelaro, R., Stajner, I., and Nielsen, J.: The GEOS-5 Data Assimilation System - Documentation of Versions 5.0.1, 5.1.0, and 5.2.0, Tech. rep., NASA Tech. Rep., <https://gmao.gsfc.nasa.gov/pubs/docs/Rienecker369.pdf>, NASA/TM-2008-104606, 2008.
- 420 Rüfenacht, R., Baumgarten, G., Hildebrand, J., Schranz, F., Matthias, V., Stober, G., Lübken, F.-J., and Kämpfer, N.: Intercomparison of middle-atmospheric wind in observations and models, *Atmospheric Measurement Techniques*, 11, 1971–1987, <https://doi.org/10.5194/amt-11-1971-2018>, 2018.
- Schmidlin, F., Carlson, M., Rees, D., Offermann, D., Philbrick, C., and Widdel, H. U.: Wind structure and variability in the middle atmosphere during the November 1980 Energy Budget Campaign, *Journal of Atmospheric and Terrestrial Physics*, 47, 183–193, [https://doi.org/https://doi.org/10.1016/0021-9169\(85\)90133-3](https://doi.org/https://doi.org/10.1016/0021-9169(85)90133-3), 1985.
- 425 Shepherd, G. G., Thuillier, G., Gault, W. A., Solheim, B. H., Hersom, C., Alunni, J. M., Brun, J.-F., Brune, S., Charlot, P., Cogger, L. L., Desaulniers, D.-L., Evans, W. F. J., Gattinger, R. L., Girod, F., Harvie, D., Hum, R. H., Kendall, D. J. W., Llewellyn, E. J., Lowe, R. P., Ohrt, J., Pasternak, F., Peillet, O., Powell, I., Rochon, Y., Ward, W. E., Wiens, R. H., and Wimperis, J.: WINDII, the wind imaging interferometer on the Upper Atmosphere Research Satellite, *Journal of Geophysical Research: Atmospheres*, 98, 10 725–10 750, <https://doi.org/https://doi.org/10.1029/93JD00227>, 1993.
- 430 Stoffelen, A., Pailleux, J., Källén, E., Vaughan, J. M., Isaksen, I., Flamant, P., Wergen, W., Andersson, E., Schyberg, H., Culoma, A., Meynart, R., Endemann, M., and Ingmann, P.: THE ATMOSPHERIC DYNAMICS MISSION FOR GLOBAL WIND FIELD MEASUREMENT, *Bulletin of the American Meteorological Society*, 86, 73 – 88, <https://doi.org/10.1175/BAMS-86-1-73>, 2005.
- Tang, Q., Zhou, Y., Du, Z., Zhou, C., Qiao, J., Liu, Y., and Chen, G.: A Comparison of Meteor Radar Observation over China Region with Horizontal Wind Model (HWM14), *Atmosphere*, 12, <https://doi.org/10.3390/atmos12010098>, 2021.
- 435 van Cleef, G. W., Shaw, J. H., and Farmer, C. B.: Zonal winds between 25 and 120 km obtained from solar occultation spectra, *Geophysical Research Letters*, 14, 1266–1268, <https://doi.org/https://doi.org/10.1029/GL014i012p01266>, 1987.
- Wu, D. L., Schwartz, M. J., Waters, J. W., Limpasuvan, V., Wu, Q., and Killeen, T. L.: Mesospheric doppler wind measurements from Aura Microwave Limb Sounder (MLS), *Advances in Space Research*, 42, 1246–1252, <https://doi.org/https://doi.org/10.1016/j.asr.2007.06.014>, 2008.

<https://doi.org/10.5194/egusphere-2025-3116>

Preprint. Discussion started: 10 July 2025

© Author(s) 2025. CC BY 4.0 License.



- 440 Zhu, J.: Conversion of Earth-centered Earth-fixed coordinates to geodetic coordinates, IEEE Transactions on Aerospace and Electronic Systems, 30, 957–961, <https://doi.org/10.1109/7.303772>, 1994.