



Challenges in Simulating Ozone Depletion Events in the Arctic Boundary Layer: A Case Study Using ECHAM/MESSy for Spring 2019/20

Stefanie Falk¹, Luca Reißig¹, Bianca Zilker², Andreas Richter², and Björn-Martin Sinnhuber¹

¹Institute of Meteorology and Climate Research, Atmospheric Trace Gases and Remote Sensing, Karlsruhe Institute of Technology, Karlsruhe, Germany

²Institute of Environmental Physics, University of Bremen, Bremen, Germany

Correspondence: Stefanie Falk (stefanie.falk@kit.edu)

Abstract. Ozone depletion events (ODEs) and bromine explosions (BEs) occur regularly in the springtime polar boundary layer. ODEs alter the oxidation capacity of the polar boundary layer and promote formation of toxic mercury. We investigated Arctic ODEs and BEs in 2019/20 using the chemistry-climate model ECHAM/MESSy v2.55.2, nudged with ERA5 reanalysis data. Model results were evaluated against surface ozone measurements, satellite-derived tropospheric BrO vertical column densities (VCDs), and in situ data from the MOSAiC expedition. The model underestimated boundary layer (BL) height during shallow BL conditions, coinciding with a warm surface temperature bias (2–10 K), particularly below -10°C , likely inherited from ERA5. An updated model configuration, incorporating more realistic multi-year sea ice and relaxed bromine release thresholds, improved agreement with coastal ozone observations (Eureka, Utqiagvik) but still failed to reproduce strong ODEs observed during MOSAiC. Modeled surface BrO mixing ratios were overestimated, while BrO VCDs were underestimated, suggesting that simply increasing Br₂ emissions does not resolve discrepancies. A weaker collocation between modeled BrO VCDs and ODEs aligns with prior airborne studies and may reflect tropospheric chemical and transport processes rather than stratospheric contamination. Despite decreasing Arctic sea ice extent and increasing BrO VCDs, long-term records from Alert, Utqiagvik, and Zeppelin show a decline in strong ODE frequency since 2000. This suggests that bromine emissions from first-year sea ice (FYSIC) alone may not fully account for observed ODE variability, and that additional climate-sensitive mechanisms may modulate Arctic ozone chemistry. Long-term model integrations are recommended to better understand these trends.

1 Introduction

The Arctic is subject to rapid changes due to the warming climate (AMAP, 2012; Rantanen et al., 2022). The impact of this change is reflected by the decrease of sea ice extent and thickness over the past decades (Stroeve and Notz, 2018; Lindsay and Schweiger, 2015) which caused a decline in multi-year sea ice and a stronger seasonality of the ice cover (Haine and Martin, 2017). Sea ice extent and thickness are projected to further decrease in all future projections (Notz and SIMIP Community, 2020) which may open new routes for cargo ships towards the end of the 21st century with an associated increase in Arctic



air pollution (Yumashev et al., 2017). The composition of the Arctic troposphere in winter and spring is strongly affected by long-range transport from mid-latitudes, while the polar dome prevents this transport in summer (Bozem et al., 2019). Primary pollutants include NO_x and CO, which are precursor substances for the formation of tropospheric ozone (O₃) after polar sunrise. Annual mean ozone volume mixing ratios (VMRs) at the surface are relatively low in the Arctic ($\chi_{\text{O}_3} \leq 30$ ppbV) compared to the mid-latitudes of the Northern Hemisphere and show a distinct seasonal cycle with a polar winter maximum (October–February) and a polar summer minimum (March/April–August/September) (Helmig et al., 2007; Whaley et al., 2023).

Tropospheric ozone VMR has been observed below the detection limit on sites near the Arctic Ocean in springtime (March–May) for decades. These periods of $\chi_{\text{O}_3} \leq 5$ ppb, sometimes extending for several days, were first recognized by Bottenheim et al. (1986) in Alert (Canada), who coined the term Ozone Depletion Events (ODEs). Later, satellite observations revealed coincident plumes of enhanced bromine monoxide (BrO) vertical column densities (VCDs) extending over synoptic scales in both Arctic and Antarctic regions (Richter et al., 1998; Wagner and Platt, 1998; Richter et al., 2002). This was evidence that bromine (Br) chemistry leads to the destruction of ozone during ODEs (Barrie et al., 1988; McConnell et al., 1992; Platt and Hönninger, 2003). The associated chemical mechanism also alters the oxidation capacity of the polar boundary layer and affects formation of toxic, oxidized mercury (Coburn et al., 2016; Wang et al., 2019c).

With polar sunrise, molecular bromine (Br₂) is photo-dissociated (Eq. (R1)) and subsequently destroys O₃ forming BrO (Eq. (R2)):



The resulting BrO can then self-react to form both Br and Br₂ (Eq. (R3)) or react with hydroperoxyl (HO₂) to form hypobromous acid (HOBr) (Eq. (R4)), which photo-dissociates to Br (Eq. (R5)) starting the cycle again:



These reactions terminate with the formation of reservoir species such as hydrogen bromide (HBr) that are not efficient at forming reactive bromine. The main source of inorganic bromine in the polar regions is sodium bromide (NaBr) from sea salt.



It is constantly emitted from the open ocean or open leads in the sea ice. Sea salt aerosols release Br_2 into the gas-phase (Yang et al., 2005). A minor contribution comes from organic bromine (for example CHBr_3 , CH_2Br_2 , CH_3Br) emitted from ocean and sea ice (Stemmler et al., 2015; Abrahamsson et al., 2018). Recycling of inorganic bromine is necessary to sustain ODEs (Abbatt et al., 2012). This can involve heterogeneous and multi-phase reactions of, for example, HOBr or bromine nitrate (BrNO_3) on salty ice surfaces or in salty solutions (Sander et al., 2006):



Interhalogen reactions are able to turn BrCl into Br_2 :



For each brominated trace gas in Eqs. (R6)–(R7) getting in contact with an icy surface of high salinity, twice the number of bromine atoms is released. This exponential increase is called bromine explosion (BE).

While this mechanism is well understood through lab experiments (Oldridge and Abbatt, 2011) and box modeling that also includes deeper snow layers (Toyota et al., 2014), the picture is far from complete. Accurate predictions on a larger scale still pose a challenge for Chemistry Transport (CTMs) and Chemistry-Climate Models (CCMs) in their default setup (Whaley et al., 2023).

Observational evidence indicates that BEs occur in two distinct synoptic situations: stable boundary layer (BL) with little mixing, and turbulent mixing during blizzards (Jones et al., 2009). This is reflected by two distinct mechanisms for bromine release implemented into models: (1) surface bulk snow and ice (Toyota et al., 2011, e.g.) and (2) blowing snow emissions (Yang et al., 2008, 2010). Marelle et al. (2021) showed that bromine emissions from bulk snow and ice dominate over those from blowing snow in triggering ODEs while blowing snow is an important additional source of sea salt aerosols.

Latest satellite observations of BrO VCDs with a high spatial resolution (Seo et al., 2019, 2020) indicate that snow-covered first-year sea ice in combination with open leads and sea ice cracks may act as the primary source of bromide during BEs which is in contrast to earlier findings (Ridley et al., 2003).

This work aims at a better understanding and characterization of the bromine release from sea ice and surface snow and the subsequent BEs and ODEs in the global CCM ECHAM/MESSy Atmospheric Chemistry (EMAC) model (Jöckel et al., 2005, 2016). To assess the model's performance in the central Arctic, we utilize observational data from the 2019/20 season, corresponding to the Multidisciplinary drifting Observatory for the Study of Arctic Climate (MOSAIC) expedition. In Section 2, we describe the EMAC model and present the observational data sets used for model evaluation. The observational data



consist of O_3 VMRs at Arctic ozone monitoring sites and satellite BrO VCD at high resolution from the TROPOMI instrument onboard the Copernicus Sentinel-5P satellite. Subsequently, we identify the best model setup based on spring 2019 (Section 3).

80 Results using this best setup are compared to observational data for spring 2020 especially accounting for measurements taken during the MOSAiC expedition. Implications on pan-Arctic and site-specific climatological ozone depletion as well as BrO VCDs are presented in Section 4. Finally, we summarize and discuss our results and give a brief outlook in Section 5.

2 Methods and data

2.1 Model description and setup

85 We use the Modular Earth Submodel System (MESSy) v2.55.2 in combination with basemodel ECHAM5.2 (ECHAM/MESSy) referred to as EMAC. MESSy is a software framework that combines Earth system components, such as atmosphere, land, ocean, and subsystems of these, in a modular way (Jöckel et al., 2005, 2016). This modularity enables the deployment of different basemodels with an identical numerical implementation of geophysical and -chemical processes and parameterization.

We base the EMAC model setup on REF-C1SD of the Chemistry-Climate Model Initiative (CCMI) 2 (Orbe et al., 2020).

90 We use specified dynamics (SD) for the historical period 2019/20, with nudged surface pressure (logarithmic) and nudged temperature, divergence, and vorticity fields from above the boundary layer (model level 8) up to a pressure level of 1 hPa. Sea Surface Temperatures (SSTs) and Sea Ice Cover (SIC) are also prescribed. Nudging data were generated from the European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysis v5 (ERA5) (Hersbach et al., 2020).

All simulations were conducted in a T42L90MA resolution with the spectral truncation T42 that translates to a $(2.7851 \times$
95 $2.8125)^\circ$ regular Gaussian grid in the polar regions (latitude $> \pm 68^\circ$). Our model setup comprises 90 hybrid pressure levels up to 0.01 hPa. These hybrid pressure levels are terrain following, resulting in about 1–3 levels within the Arctic spring-time BL.

We use the Module Efficiently Calculating the Chemistry of the Atmosphere (MECCA) (Sander et al., 2019) with the CCMI-base-02 chemistry mechanism in which mercury and iodine chemistry are switched off. This chemical mechanism comprises 261 gas-phase, 12 heterogeneous, and 80 photolysis reactions (Supplement A). Heterogeneous reactions are mainly restricted
100 to the stratosphere and use a prescribed aerosol surface concentration climatology (1996–2005) computed with the MESSy submodel MADE (Ackermann et al., 1998). We depict the most relevant cycles for ODEs grouped by families in a reaction network (Fig. 1). The NO_x cycle can be net-zero regarding ozone destruction and production.

In the following, we briefly describe the most relevant MESSy submodels that handle trace gas emissions and sinks. For a complete list of submodels used in this study, please consult Supplement B.

105 Trace gas emissions that depend on the state of the Earth system components (atmosphere, land, and ocean) are computed during the model integration in the MESSy submodel ONEMIS Kerkweg et al. (2006b). Therein, the subroutine AirSnow (Falk and Sinnhuber, 2018) handles the emission of bromine from sea ice and snow in the polar regions following the scheme suggested by Toyota et al. (2011). The dry deposition flux of O_3 on first-year sea ice (FY) triggers Br_2 emissions. Assuming that a considerable amount of bromide (Br^-) has already been pushed out of multi-year sea ice (MY), recycling is limited
110 by the amount of inorganic bromine (HBr, $BrNO_3$, and HOBr) deposition from gas-phase. A similar assumption is used for




Figure 1. Schematic of bromine explosions (BEs) and catalytic ozone depletion reaction network separated by families. The emission of Br_2 and BrCl triggers the ozone destruction leading to ODEs given the availability of light for photolysis.

snow on land (LS). The distribution of MY sea ice is prescribed. This scheme introduces a critical temperature T_{crit} effectively acting as a seasonal limit and a critical solar zenith angle (θ_{crit}) as a trigger that separates dark from sun-lit conditions with two molar yields (ϕ_1 , ϕ_2) parameterizing in-snow photochemistry.

The submodel also computes a sea salt aerosol mass flux from a prescribed aerosol surface concentration climatology. The corresponding bromine flux is derived by scaling this flux with a Cl to Br ratio of 667 and assuming half of this flux is released into the gas-phase (Yang et al., 2008).

Ocean-to-atmosphere fluxes of organic substances dissolved in seawater (e.g. dimethylsulfate (DMS), bromoform CHBr_3 and dibromomethane CH_2Br_2) (Pozzer et al., 2006) are computed by the AIRSEA submodel. The flux for each species is calculated from concentration gradients in the uppermost ocean layer and the lowermost atmosphere layer. The species-specific Henry constant defines the respective solubility. The corresponding concentrations of brominated very short-lived substances (VSLS) in ocean waters are taken from Wang et al. (2019b).

The submodels OFFEMIS and TNUDGE handled emissions from preprocessed 4D forcing fields (Kerkweg et al., 2006b). Emission inventories of greenhouse gases (GHGs), ODS, and ozone precursors are based on the German Aerospace Centre (Deutsches Zentrum für Luft- und Raumfahrt, DLR) CCMi2 inventory (available until 2019). GHGs included are N_2O , CH_4 , and CO_2 . ODS include a wide range of halons, chlorofluorocarbons (CFCs), and halogenated CFCs (HCFCs). Also included are potential ozone precursors like Volatile Organic Compounds (VOCs) and Non-Methane-Hydro-Carbons (NMHCs) from biomass and agricultural waste burning as well as in fossil fuel emissions from traffic and non-traffic consisting of VOCs,




Figure 2. Locations of the Arctic ozone monitoring sites used in this work. The Research vessel Polarstern’s drift track during the MOSAiC expedition (March–April–May 2020) is indicated in blue–red–orange, respectively. A spring 2020 mean sea ice age distribution from the Integrated Climate Data Center (ICDC) age of sea ice product (Tschudi et al., 2024) is shown.

NMHCs, NO, CO, SO₂, NH₃ plus aerosol SO₄ derived from SO₂, organic carbon (OC) and black carbon (BC) emissions. These CCMI2 inventories are not available beyond 2019 for our model, so we substituted the missing data with 2019 emissions.

130 Emissions of the minor VSLS (CHCl₂Br, CHClBr₂, CH₂ClBr) are based on Warwick et al. (2006). The submodel TNUDGE nudges the surface concentrations concentrations of the long-lived halons CF₃Br and CF₂ClBr, and CH₃Br towards observed mixing ratios.

Dry deposition is the most important way of removing non-soluble chemical substances from the atmosphere. The corresponding dry deposition resistances (v^{dd}) are computed by the submodel DDEP (Kerkweg et al., 2006a) following the Wesely-
135 scheme (Wesely, 1989). For most species, there are no observations of surface resistance (r_c). Estimates are calculated based on the Henry coefficient, a reactivity coefficient, and r_c of O₃ and sulfur dioxide (SO₂).

Wet deposition of soluble species is calculated via the SCAV submodel (Tost et al., 2006). It considers convective and large-scale precipitation separately as well as three modes of scavenging from ice-, liquid- and gas-phase.

2.2 Observational data

140 For evaluating our model results, we use surface ozone VMR from a network of Arctic monitoring sites, satellite observations of BrO column VCD from the TROPospheric Monitoring Instrument (TROPOMI) onboard the European Space Agency (ESA) Copernicus Sentinel-5 Precursor satellite, and atmosphere and trace gas observational records from the Multidisciplinary drifting Observatory for the Study of Arctic Climate (MOSAiC) expedition (2019/20). The ozone monitoring sites and MOSAiC drift track locations’ are shown in Fig. 2.

**Table 1.** Referred ozone monitoring sites in the Arctic and data periods used in this work.

Station	Period	Coordinates		
		latitude (° N)	longitude (° E)	altitude (m)
Alert [†]	1988–2013	82.499	-62.341	204.0
Eureka [†]	2018–2023	79.983	-85.95	610.0
Nord [†]	2001–2022	81.600	-16.67	20.0
Utqiagvik [‡]	1973–2023	71.323	-156.611	11.0
Summit [†]	2000–2024	72.578	-38.459	3238.0
Mt. Zeppelin [†]	1989–2024	78.910	11.888	474.0

Data source: [†] NILU - EBAS, [‡] NOAA - Global Monitoring Laboratory

145 2.2.1 Ozone monitoring sites

There are several monitoring sites providing long- and short-term χ_{O_3} records in the Arctic as listed in Table 1. For studying the similarity between modeled and observed χ_{O_3} , and possible changes due to a changing climate, we use long-term data from the National Oceanic and Atmospheric Administration (NOAA) atmospheric monitoring station Barrow at Utqiagvik, Alaska (USA) (McClure-Begley et al., 2024) spanning 5 decades, the over 3 decade long ozone record from Mt. Zeppelin, 150 Ny-Ålesund (Spitsbergen), and almost 30 years of observations from Alert, Nunavut (Canada). We supplement these with data records from sites located at Eureka, Nunavut (Canada), and Nord and Summit both located in Greenland (Denmark). All data, except for Barrow station, have been downloaded from the EBAS website (Norsk institutt for luftforskning (NILU)) in 1-hourly resolution. All model output was interpolated to the sites' locations using the MESSy submodel SCOUT.

2.2.2 TROPOMI

155 The ESA Copernicus Sentinel 5-Precursor satellite was launched in October 2017 with a designed lifetime of 7 years. It is on a sun-synchronous orbit with local time of ascending node is at 13:30 h. The TROPOMI instrument has a near-nadir resolution of $3.5 \times 7 \text{ km}^2$ ($3.5 \times 5.5 \text{ km}^2$ since July 2022) and a swath width of 2,600 km. Its level 2 data products include O_3 (total and tropospheric column, profile), NO_2 (total and tropospheric column), SO_2 (total column), and carbon monoxide CO (total column). For this study, BrO total and tropospheric columns have been retrieved with an optimized and adapted Differential 160 Optical Absorption Spectroscopy (DOAS) retrieval algorithm that was developed for earlier satellite missions (Seo et al., 2019, 2020). To isolate the tropospheric from the total column, a stratospheric BrO climatology (Theys et al., 2009) has been used. As no total column air-mass factor (AMF) for TROPOMI is currently available, a stratospheric AMF has been applied for the total columns. For the tropospheric columns, a simplified approach was used assuming a bright surface (albedo of 0.9) and a BrO surface layer of 400 m thickness. This affects the retrieval over dark surfaces, e.g. ocean, and boreal forest, by reducing 165 the amount of reconstructed BrO VCD (Choi et al., 2012).




Figure 3. TROPOMI BrO VCD for April 2019. Stratospheric BrO from a stratospheric climatology by Theys et al. (2009) was subtracted from TROPOMI total column to derive the tropospheric contribution. (a) Total column; (b) Tropospheric column.

In Fig. 3 TROPOMI monthly mean BrO total and tropospheric column VCD are shown for April 2019. The tropospheric column VCD (Fig. 3(b)) indicates BrO enhancement over the whole Arctic Ocean but most prominently over the Canadian archipelago and the Kara/Laptev Sea ($\text{VCDBrO} = (3 - 4) \cdot 10^{13} \text{ molec cm}^{-2}$).

2.2.3 MOSAiC expedition

170 The MOSAiC expedition was the largest scientific expedition in the Arctic to date. From September 2019 to October 2020, the German research vessel Polarstern drifted with the sea ice in the Central Arctic. In the course of this mission, a multitude of interdisciplinary experiments was conducted, including the measurement of meteorological conditions (Jozef et al., 2023) as well as O_3 (Angot et al., 2022) (Mahajan, 2022), and BrO (Benavent et al., 2022). The position of Polarstern during Spring 2020 is shown in Fig. 2.

175 The modeled surface temperature and BL height interpolated to the drift track are in general in good agreement with the observations (Fig. 4). This demonstrates that nudging with ERA5 reproduces the observed weather and meteorological conditions. Below -10°C , however, a difference in modeled and observed surface temperatures becomes apparent ($-2^\circ\text{C} < \Delta T_{\text{surf}} < -10^\circ\text{C}$). This bias increases with decreasing temperatures and is most prominent when BL heights are below 100 m. This could be caused by the ERA5 nudging data in combination with the relatively low vertical model resolution in the BL.

180 However, nudging only starts above model level 8. Wang et al. (2019a) have identified a regionally varying bias in both ERA-interim and ERA5 that increases at low temperatures (most notably below -20°C) compared with buoy observations which is similar to the bias observed in our model experiments.




Figure 4. Comparison of meteorological observations during the MOSAiC expedition with model results mapped to the MOSAiC drift track. (a) Temperature; (b) BL. The dashed red line represents $T_{crit} = -15^{\circ}\text{C}$

3 Model parameter sensitivity and setup

In this section, we describe the sensitivity of the AirSnow algorithm to critical parameters and boundary conditions. We introduce a more realistic multi-year sea ice concentration derived from the Integrated Climate Data Center (ICDC) age of sea ice product (Section 3.1) and implement a sigmoidal relaxation of the temperature threshold (Section 3.2). We test the performance of the AirSnow algorithm and find the best model setup (Section 3.3). To make sure that the model is not over-tuned, we exclusively use the period January–July 2019 for improving the model skill in terms of O_3 at observation sites.

All model experiments are listed in Table 2. In contrast to the release version in MESSy v2.55.2, we include two critical bug fixes concerning the temperature (conversion from Celsius to Kelvin) and solar zenith angle (sign flip) thresholds. We have already reported and fixed these in the release candidate of MESSy v2.56.

3.1 Multi-year sea ice cover fraction

By construction, the amount of Br_2 emitted using the Toyota scheme is sensitive to the assumed age of sea ice. Therefore, we expect more Br_2 emissions from regions with a large first-year sea ice concentration (FYSIC). Falk and Sinnhuber (2018) derived a multi-year sea ice concentration (MYSIC) from ERA-interim with the assumption of a static multi-year sea ice distribution computed from the SIC at the seasonal minimum of the previous year. We used the same assumption to derive MYSIC from ERA5 SIC and compared this with an age of sea ice (AoSI) product provided by the Integrated Climate Data Center (ICDC) (Tschudi et al., 2024).



Table 2. List of model experiments. Our MESSy v2.55.2 includes two critical bug fixes in the AirSnow mechanism that have been integrated into the MESSy release candidate. MYSIC based on ICDC Age of Arctic Sea Ice has been made available to the MESSy community. Br₂ release from prescribed sea salt aerosols was diagnostic only by default.

Exp.	Model version	MECCA chemistry	Period	AirSnow	MYSIC	T_{crit}	Sea salt Br ₂	notes
ref	2.55.2	CCMI2-base-01	2019-01 –2020-07	off	-	-	diagnostic	
sfa002	2.55.2	CCMI2-base-01	2019-01 –2020-07	on	ERA5	–15 °C	diagnostic	model intercomp.
sfa008	2.55.2	CCMI2-base-01	2019-01 –2019-07	on	ERA5	–15 °C	diagnostic	+ frozen lakes masked + sigmoidal T_{crit}
sfa010	2.55.2	CCMI2-base-01	2019-01 –2019-07	on	ICDC	–15 °C	diagnostic	+ frozen lakes masked + sigmoidal T_{crit}
sfa011	2.55.2	CCMI2-base-01	2019-01 –2019-07	on	ICDC	–10 °C	diagnostic	+ frozen lakes masked + sigmoidal T_{crit}
sfa012	2.55.2	CCMI2-base-01	2019-01 –2019-07	on	ICDC	–2.5 °C	diagnostic	+ frozen lakes masked + sigmoidal T_{crit}
sfa013	2.55.2	CCMI2-base-01	2019-01 –2019-07	on	ICDC	–10 °C	diagnostic	+ frozen lakes like LS + sigmoidal T_{crit}
sfa017	2.55.2	CCMI2-base-01	2019-01 –2020-07	on	ICDC	–10 °C	interactive	+ frozen lakes like LS + sigmoidal T_{crit}

ICDC AoSI uses sea ice drift data from satellite observations to assign an age to the individual ice floats for which drift trajectories are computed. Each grid cell with at least 15 % SIC is treated as a Lagrangian particle and traced every week. The co-existence of ice of varying ages in one grid cell prefers the survival of the older ice because younger and thinner sea ice deforms and melts more easily. This causes an overestimation of the multi-year sea ice cover.

To derive a MYSIC, we summed over all ICDC age classes ≥ 1 year and average the bi-weekly data over one month. The original resolution of 12.5×12.5 km was then remapped onto the target resolution (e.g. T42, T106, Appendix Fig A1).

Compared to ERA5-derived MYSIC, ICDC AoSI indicates distinctively less multi-year sea ice in the Central Arctic Ocean and the Canadian archipelago but more east of Greenland, north of Spitsbergen and Alaska (Fig. 5(a)). To identify which of the two MYSIC estimate is more realistic, we computed the total area covered by multi-year sea ice. The area derived from ERA5 is consistently $\sim 30\%$ larger than from ICDC AoSI over the period 1980 to 2020 (Appendix Fig. A2(a)). The total area of multi-year sea ice based on ICDC AoSI amounts to $2.5 \cdot 10^6$ km² in the 2020s, which is consistent with the range given by Regan et al. (2023).

We conducted two model experiments that differ only in the applied MYSIC (ERA5, ICDC AoSI) and found that the lower the MYSIC in a grid cell the more Br₂ and the less BrCl emissions from ice and snow are simulated (Fig. 5(b)). The opposite




Figure 5. Influence of MYSIC on AirSnow emission fluxes. (a) MYSIC difference ICDC AoSI-ERA5. (b) Br₂ (left) and BrCl (right) flux difference between a model experiment with ICDC AoSI and ERA5.

is true for grid cells with larger MYSIC. This means that strong local sources of Br₂ due to small-scale sea ice inhomogeneities may appear smeared out in our applied model resolution.

215 For our model experiments, we computed BrO VCD from χ_{BrO} and used the model potential vorticity (PV) tropopause index to separate tropospheric and stratospheric columns. We sampled the modeled total and tropospheric VCD at 13–14 h local time and calculated monthly averages to compare with the satellite retrieval. The mean modeled stratospheric contribution to the total BrO VCD in April was rather uniform in all experiments and amounts to $(2.5 - 3) \cdot 10^{13} \text{ molec cm}^{-2}$ (Appendix Fig. B2).

The TROPOMI BrO VCD for April 2019 (Fig. 3(a, b)) indicates hotspot regions over the Canadian archipelago and the Kara/Laptev Sea. This pattern is well reproduced in experiments with ICDC AoSI-derived MYSIC (Appendix Fig. B3(c, d)). 220 The ERA5-derived MYSIC shows an additional hotspot at the Alaskan coast (Appendix Fig. B3(a, b)) which is absent in the satellite VCD. Therefore, we conclude that MYSIC derived from ICDC AoSI is the better choice.

3.2 Temperature threshold

The threshold temperature T_{crit} constrains the occurrences of BEs temporally in springtime. Toyota et al. (2011) noted that 225 there is no T_{crit} that optimizes modeled ozone at all monitoring sites simultaneously. We propose that relaxing this temperature threshold might improve the agreement without implementing a detailed snow microphysics and snow chemistry scheme (Toyota et al., 2014) in a global CCM. For this purpose, we implemented a sigmoidal temperature dependency of the form

$$f_{\text{crit}}(T) = \frac{1}{1 + e^{T - T_{\text{crit}}}}, \quad (1)$$

with T the surface temperature in a given grid cell. The Br₂ and BrCl emission fluxes are then scaled with $f_{\text{crit}}(T)$ and reach 230 10% and 90% of the maximum flux at $T_{\text{crit}} \pm 2.2^\circ\text{C}$, respectively. Br₂ emissions thus already increase where temperatures are slightly above T_{crit} , most notably in coastal regions.

We then looked at model results at $T_{\text{crit}} \in \{-15, -10, -2.5\}^\circ\text{C}$. $T_{\text{crit}} = -2.5^\circ\text{C}$ means allowing bromine emission also at temperatures around the freezing point of freshwater. With a lower threshold, the modeled Br₂ fluxes consistently show an increase over first-year sea ice (Fig. 6). Emissions from multi-year sea ice regions, indicated by the BrCl fluxes, also display an



235 increase but remain two orders of magnitude lower than emissions from first-year sea ice regardless of the choice of T_{crit} . At $T_{\text{crit}} = -2.5^\circ\text{C}$, the model predicts Br_2 emissions in the Gulf of Bothnia, the White Sea, and the Sea of Okhotsk (Fig. 6(c)), though this does not result in enhanced BrO VCD (Fig. 6(f)). For a site at Utö, a Finnish island in the Baltic Sea, Jalkanen and Manninen (1996) reported a springtime bromine enhancement compared to chlorine which they attributed this to sea salt aerosol. No coincident ozone depletion is evident from their data. At $T_{\text{crit}} \geq -2.5^\circ\text{C}$ the BrO VCD in the hot spot regions is
240 closest to the TROPOMI VCD, though more enhanced BrO VCDs are found at the coast east of Greenland and East Siberia than observed. As indicated above, this could be a relict of the relatively low model resolution and associated land-sea mask that does not resolve all topographic features and congruent sub-grid temperature variance. Possibly, the stratospheric BrO climatology and simple AMF used in the TROPOMI retrieval also plays a role for the absolute values. Overall, our modeled BrO VCD in 2019 agrees best with satellite observations in both amount and spatial pattern of tropospheric BrO VCD at
245 $T_{\text{crit}} \geq -2.5^\circ\text{C}$.

3.3 Model skill and best setup

To decide on the best model setup, we evaluated the model skill in terms of the coefficient of determination of the linear regression (Pearson correlation coefficient-squared, R^2) and root-mean-square-error (RMSE) at the Arctic ozone monitoring sites (Section 2.2.1) for spring 2019.

250 The resulting R^2 and RMSE for 2019 at Utqiagvik, Eureka, Summit, and Zeppelin are listed in Table A1. The closer R^2 is to 1, the better the correlation. The closer the RMSE is to 0, the smaller the difference between the observed and modeled time series. For the corresponding histograms with fitted linear regression curves, consult Appendix Fig. A4.

We confirmed that there is not a single combination of parameters that optimizes both R^2 and RMSE for all sites. The model experiment with MYSIC from ICDC AoSI and $T_{\text{crit}} = -15^\circ\text{C}$ (sfa010) performs best with respect to R^2 ($\langle R^2 \rangle = 0.198$). The
255 experiments with $T_{\text{crit}} = -10^\circ\text{C}$ (sfa011, sfa017) are best in terms of RSME ($\langle \text{RMSE} \rangle = 0.365$). The treatment of freshwater lakes like land snow decreases the model skill slightly compared to the experiments where these are excluded (compare [sfa013, sfa017] with [sfa010, sfa011]). We will need to investigate this further. An experiment with $T_{\text{crit}} = -2.5^\circ\text{C}$ performs well at Zeppelin, which may suggest that temperatures may be too high in that particular region probably due to the low horizontal and vertical resolution (smoothed topography).

260 Based on these quantitative results, we decided on the setup with MYSIC derived from ICDC AoSI, $T_{\text{crit}} = -10^\circ\text{C}$, and Br_2 emissions from sea salt enabled (sfa017). Although qualitatively, tropospheric BrO VCD showed a better with $T_{\text{crit}} = -2.5^\circ\text{C}$.

4 Results

Our special focus lies on the spring 2020 for which ozone monitoring station records, TROPOMI satellite observations, and data from the MOSAiC expedition are available. In this section, we show the surface ozone VMR time series at different sites
265 for our best model setup and compute probability density functions (PDFs) of χ_{O_3} from observation and model results to look for indications of a possible climate change impact on ODEs (Section 4.1). We compare the modeled pan-Arctic pattern of




Figure 6. Br₂ and BrCl flux integral April 2019 and monthly mean tropospheric BrO VCD at 13 – 14 h local time in April 2019. The EMAC total column BrO has been split into tropospheric and stratospheric contributions using the modeled PV-tropopause height on hourly basis. Model experiments comprise $T_{\text{crit}} = (-15, -10, -2.5)^\circ\text{C}$ in ascending order. (a, b) $T_{\text{crit}} = -15^\circ\text{C}$; (c, d) $T_{\text{crit}} = -10^\circ\text{C}$; (e, f) $T_{\text{crit}} = -2.5^\circ\text{C}$.

ozone depletion and BrO VCD with TROPOMI retrieved BrO VCD (Section 4.3) and use the MOSAiC data to judge the overall model skill in the Central Arctic qualitatively (Section 4.2). Our best model setup comprises MYSIC from ICDC AoSI and a sigmoidal temperature threshold function centered at $T_{\text{crit}} = -10^\circ\text{C}$.



270 4.1 Arctic ozone monitoring sites

4.1.1 Ozone time series 2019/20

For the Arctic ozone monitoring sites Eureka, Nord, Summit, Utqiagvik, and Zeppelin, we show the 2019/20 composite time series of modeled (ref, sfa017) and observed χ_{O_3} in Fig. 7. We find that the effect of AirSnow on ozone VMR is contained temporally. As soon as the trigger conditions for ODEs are not satisfied anymore, O_3 VMR returns quickly to the values of the reference simulation. It is evident that AirSnow qualitatively improves the EMAC model capabilities of capturing tropospheric ozone in the Arctic spring at all sites. Key features like the late April/early May ODE in 2019 at Eureka and Utqiagvik and the late March/late April ODEs at Utqiagvik in 2019 match reasonably well in their timing but not in the observed strength of the ozone depletion. An ODE in March at Eureka station coinciding with a pronounced dip in ozone at Summit is not reproduced. The EMAC model generally underestimates χ_{O_3} in Arctic winter. Falk and Sinnhuber (2018) have shown that a higher surface resistance of ozone over ice and snow ($r_c = 10000 \text{ sm}^{-1}$) could improve this. In 2000, especially in March and May, ODEs at Eureka and Utqiagvik were captured less well, also reflected by a reduced model skill compared to 2019 (see Tab. B1). This was potentially caused by the recycling of 2019 emission inventories. As a secondary pollutant, tropospheric ozone depends on precursor substances like NO_x , CO, and VOCs. COVID-19 policies reduced emissions of these which had a measurable effect on tropospheric ozone (Venter et al., 2020).

Beyond Arctic springtime, we found peak ozone VMR at Utqiagvik in June 2019 and 2020 that are much larger than observed. These peak values are likely caused by large wildfires raging within the Arctic cycle in 2019 (Descals et al., 2022). These usually contribute to ozone precursors and cause episodes of enhanced tropospheric ozone (Cofer et al., 1990; Lindskog et al., 2007; Karlsson et al., 2013). However, Baker et al. (2016) have reported that attenuation of solar radiation and aging of aerosols are potentially underestimated in chemistry transport models causing photolysis rates and therefore in situ ozone production to be overestimated compared to observations.

For reference, surface ozone time series including only the bug fixes are shown in Appendix Fig. B1.

4.1.2 Ozone climatology

We assume that the amount of Br_2 emitted and, subsequently, the strength of ODEs should be highest for a high FYSIC (Falk and Sinnhuber, 2018). Around 2007, the amount of multi-year sea ice in the Arctic prominently dropped (Regan et al., 2023) (Appendix Fig. A2). Hence, we would expect an increased occurrence of ODEs after 2007. To test this, we divided the observational datasets with long-term records into two periods: (1) start of record until 2000 and (2) 2000 until the end of each record. For these two periods, we computed PDFs of observed χ_{O_3} as normalized histograms with 1 ppbV binning (Fig. 8). The Arctic ozone monitoring sites with long-term records are Alert, Utqiagvik, and Zeppelin. Error bars denote the year-by-year variance of the observational data using standard deviation.

In March, observations at Alert and Zeppelin display close-to-normal distributions peaking between 40 – 45 ppbV with a small tail towards low χ_{O_3} . These tails become larger throughout April and March and transform the distribution into a close-to-equal distribution which can be interpreted as an ODE fingerprint. The distribution at Utqiagvik shifts to a close-to-equal




Figure 7. Time series comparing observational records of χ_{O_3} at Arctic ozone monitoring sites with EMAC model output (ref, sfa017) interpolated to the station location. The periods March–May are highlighted in linen. The dashed red line indicates the 5 ppb threshold for ODEs. (a) Eureka; (b) Nord; (c) Summit; (d) Utqiaġvik; (e) Zeppelin.

distribution already in March following its more southern location. This is in line with previously identified temporally varying frequency distributions of χ_{O_3} in ship (Jacobi et al., 2010) and airplane (Ridley et al., 2003) expeditions in the Central Arctic.



305 In the observational period (2), all sites' distributions display a significant increase of higher χ_{O_3} compared to period (1) in spring. In period (2) data at Utqiagvik show the onset of a return to a close-to-normal distribution already in May compared to period (1). This means that conditions became less favorable for ODEs in recent decades. In April and May, the ODE bins ($\chi_{\text{O}_3} \leq 10$ ppb) at all stations are significantly (more than 1σ) less populated in time period (2). Especially, at Zeppelin almost no ODEs occur in period (2) while atmospheric background O_3 has increased. Although the instrumental uncertainty
310 in earlier years is usually larger, these data imply a significantly less frequent occurrence of strong or continuous ODEs in recent decades. This means that either the role of bromine emissions from FYSIC in promoting ODEs is overstated, or that other climate-sensitive factors are counteracting the observed increase in BrO VCD (Bougoudis et al., 2020) in terms of ozone depletion.

The modeled distributions differ from observations beyond the expected year-to-year variance. The reference simulation
315 follows a normal distribution for almost all sites and months. Only at Utqiagvik in March a slight tendency towards an equal distribution can be found. While generally too high, χ_{O_3} at Zeppelin is too low in March. Using AirSnow, the distributions get closer to observations for some months and sites. The best agreement is achieved at Alert and Zeppelin in April and Utqiagvik in March. However, a general shift towards lower ozone at all locations leads to an underestimation of χ_{O_3} above 37 ppbV. In March, there is only little change at Alert, Utqiagvik, and Zeppelin. In May, the PDF for Alert shifts towards lower values and
320 is more equally distributed. We conclude that χ_{O_3} distributions are captured more correctly, when accounting for the bromine emissions from ice and snow.

4.2 Ozone and bromine monoxide in the Central Arctic

In Section 2.2.3, we found that modeled surface temperature and BL agree reasonably well with observations during the MOSAiC expedition. In the following, we will compare modeled χ_{O_3} and χ_{BrO} with in situ observations during the MOSAiC
325 expedition for the leg March–May 2020.

We interpolated χ_{O_3} and χ_{BrO} for the model experiments ref, sfa002, and sfa017 onto the MOSAiC drift track and compared these with observations (Fig. 9). We found an ozone depletion of up to 20 ppbV with our best setup (sfa017), though the observed ODEs ($\chi_{\text{O}_3} \leq 5$ ppb) in the Central Arctic are not reproduced in any of our model experiments. During the modeled partial ODEs in April, much more surface BrO is produced than observed, while tropospheric BrO VCD remains lower than
330 observed. In May, χ_{BrO} is better reproduced. This suggests that more bromine per depleted ozone is needed in our model than observations suggest.

We further need to investigate the contributions of neglected processes like blowing snow (Yang et al., 2008; Marelle et al., 2021) and missing chemical pathways including iodine chemistry (Chameides and Davis, 1980; Benavent et al., 2022) to ODEs. In addition, photochemistry at large solar zenith angles ($100^\circ \leq \theta_\odot \leq 80^\circ$) and high surface albedo could be underestimated.

335 4.3 Pan-Arctic implications

The pattern of observed and modeled BEs show a considerable year-by-year variability. In 2020, TROPOMI tropospheric BrO VCD displays an extended area of increased BrO from the northern tip of the Canadian archipelago to the Laptev Sea




Figure 8. Histogram of χ_{O_3} for EMAC experiments sfa017 and ref (2019/20) compared to long-term observations separated into two periods: start of record to 2000 and 2000 to end of record. (a) Alert; (b) Utqiagvik; (c) Zeppelin.




Figure 9. Evaluation of EMAC experiments interpolated onto MOSAiC drift track. Compared to MOSAiC observations. (a) O_3 ; (b) BrO . Note that χ_{BrO} were not present in the reference experiment.




Figure 10. Monthly mean BrO VCD at 13 – 14 h local time in April 2020. The EMAC total column BrO has been split into tropospheric and stratospheric contributions using the modeled tropopause height on hourly basis. A stratospheric climatology (Theys et al., 2009) was subtracted from TROPOMI total column. (a) TROPOMI (total column); (b) TROPOMI (tropospheric column); (c) sfa017 (total column); (d) sfa017 (tropospheric column).

(BrO VCD $\geq 4 \cdot 10^{13}$ molec cm $^{-2}$). In the model experiment, this region decomposes into three hotspots: the northern tip of Greenland and the Canadian archipelago (BrO VCD = $(2.5 - 4) \cdot 10^{13}$ molec cm $^{-2}$), north of the Beaufort Sea, and the Laptev/Kara Sea, north of Novaya Zemliya, (BrO VCD = $(0.5 - 1.5) \cdot 10^{13}$ molec cm $^{-2}$) respectively. A modeled corridor of lower BrO enhancement coincides with the maximum of MYSIC. This implies a need to reconsider the assumptions regarding bromine recycling on multi-year sea ice. The modeled BrO VCDs are up to a factor of four lower than observed. As shown in Section 3, these scale with the amount of Br $_2$ emitted.

The monthly maximum of the difference in surface ozone (Fig. 11(a)) displays major hotspots ($\Delta O_{3\max} = (30 - 40)$ ppbV) in the Canadian archipelago/Baffin Bay/north of Greenland and the Laptev Sea/Eastern Arctic Ocean. These hotspots of ozone




Figure 11. Monthly (a) maximum and (b) mean ozone depletion in April 2020 computed from difference between experiment sfa017 and the reference simulation.

depletion colocate partly with regions of high BrO VCD (Fig. 10(d)). Ozone depletion on a pan-Arctic scale appears relatively weak on a monthly average (Fig. 11(b)) and never exceeds 17 ppbV in April compared to an ozone background of (29 ± 8) ppbV. Hence, modeled surface ozone is only depleted by $60 \pm 3\%$ on average. This confirms the result that ozone depletion is too weak not only along the MOSAiC drift track (Section 4.2) but also on larger scales. The hotspots of maximum depletion are located in the Canadian archipelago and the Central Arctic (North of the Aleutian Islands). In March and May, the major hotspot region is as well located in the Canadian archipelago, while the secondary hotspots vary both in strength and location (Appendix Fig. B4).

In the mid-latitudes, an average ozone reduction of 0–5 ppbV compared to the reference simulation is predicted. This average ozone reduction could be due to an advection of ozone depleted as well as bromine enriched air masses from the Arctic. In this regard, we found a pronounced reduction maximum of ozone of up to 25 ppbV in the North Atlantic between Iceland and the Faroe Islands. This signal is colocated with prescribed shipping emissions of CO, NO_x, NH₃, and SO₂ in that area and indicates that chemical ozone depletion could be locally enhanced by advected bromine from the Arctic.

5 Summary and conclusions

We used the chemistry-climate model ECHAM/MESSy v2.55.2 nudged with ERA5 data to study ozone depletion and bromine explosions in the Arctic in spring 2019/20. We compared model predictions with observations from ozone monitoring sites, TROPOMI satellite BrO VCD, and meteorological and atmospheric tracer data taken during the MOSAiC expedition.

The MOSAiC data revealed that the modeled boundary layer height was underestimated in periods of observed shallow BL (≤ 100 m). During these, the difference in observed and modeled surface temperatures was most prominent. Observed surface temperatures below -10°C appeared 2–10 K warmer in our model. This bias increased towards lower temperatures. This is in accordance with Wang et al. (2019a) who showed a regionally varying, warm bias at low temperatures in ERA5 compared to buoy data, most notably below -20°C . Although, for stability reasons, the lowermost 8 model levels are not nudged, this bias



could be inherited from ERA5. The relatively low vertical resolution amounting to 1–3 levels in the Arctic BL in our model setup didn't allow for resolving very shallow BL conditions.

With an improved model setup including a more realistic multi-year sea ice cover and a relaxation of the temperature threshold for bromine release, ozone depletion was better captured at ozone monitoring sites at the Arctic coast (Eureka, Utqiagvik). However, in contrast to observations, only partial ODEs ($(5 \leq \chi_{\text{O}_3} \leq 20)$ ppbv) were simulated for the MOSAiC expedition. The modeled BrO surface VMR was much higher than observed in April, while modeled BrO VCD was lower than observed. The discrepancies with observed BrO indicate that adding more Br₂ emission to the model might not be the solution to accurately model ODEs.

In general, we found a weaker correlation between modeled and observed χ_{O_3} at the Arctic ozone monitoring sites in 2020 compared to 2019. This was potentially caused by the recycling of 2019 emission inventories. As a secondary pollutant, tropospheric ozone depends on precursor substances like NO_x, CO, and VOCs. COVID-19 policies reduced emissions of these which had a measurable effect on tropospheric ozone (Venter et al., 2020).

Qualitatively, the spatial colocation between modeled monthly mean tropospheric BrO VCDs and the maximum ozone depletion appeared to be weaker in the Eastern Arctic than in the Canadian archipelago. A weak colocation agrees well with airborne O₃ and BrO observations by Ridley et al. (2003); Salawitch et al. (2010) who reported no or little ozone depletion in regions with enhanced BrO VCD, but rather attributed these findings to variations and uncertainties in the stratospheric contribution in the total column satellite retrieval. By separating tropospheric and stratospheric BrO VCD at the modeled tropopause instead of subtracting a stratospheric BrO climatology, we found the strongest increase in BrO VCD due to BEs in the troposphere. The weak colocation could, therefore, be an inherent feature of chemistry and advection in the troposphere.

On a long-term perspective, Arctic sea ice age, extent, and thickness have been decreasing strongly in the recent decades. This decrease should cause an increase in the occurrence of ODEs and BEs. However, the Arctic ozone monitoring stations with the longest records (Alert, Utqiagvik, and Zeppelin) showed a significant decrease in the probability of $\chi_{\text{O}_3} \leq 5$ ppb post year 2000. At the same time, the probability to observe higher χ_{O_3} increased. These data imply a significantly less frequent occurrence of strong or continuous ODEs in recent decades. This suggests that the role of bromine emissions from FYSIC in promoting ODEs may be overstated, or that other climate-sensitive factors are counteracting the observed increase in BrO VCD (Bougoudis et al., 2020) in terms of ozone depletion. Given the deficiency in reproducing very low χ_{O_3} , a model integration over several decades could be useful to identify these factors.

Toyota et al. (2014) have demonstrated in a modeling study that snow photochemistry in the photic zone, which is leading to the formation of HOBr, contributes more to Br₂ emissions than reactions in the snow skin layer. Following these results, Zhai et al. (2023) suggested parameterizing emissions from deeper snow layers in dependency of the solar zenith angle and showed that these emissions are important to interpret bromide records in Greenlandic ice cores. Such emissions are currently neglected in our model. Beside the parameterization of snow photochemistry, Zhai et al. (2023) also introduced a more physically motivated criterion relating snow albedo and snowmelt. Snowmelt probably is the final terminator of BEs, but experiments have shown that snow-metamorphism under non-melting temperature gradients already reduces the emission potential of bromine from snow (Edebeli et al., 2020). In combination with the shown discrepancies between observed and modeled χ_{O_3} and χ_{BrO}



during the MOSAiC expedition, this suggests that the assumptions of an infinite bromide source and instantaneous recycling may need to be revised. Including a multi-layer snow model with snow metamorphism and explicit snow chemistry could potentially improve the prediction of the bromine emission.

405 Dry deposition resistances are essential for the removal of atmospheric trace gases. Conceptually, these resistances parameterize the uptake of trace gases on different surfaces. Therefore, dry deposition has to be reevaluated when surface reactions are explicitly included or parameterized. For ozone, Barten et al. (2023) reported a high median dry deposition resistance ($r_c \propto 20000 \text{ sm}^{-1}$) over the Central Arctic sea ice which is one order of magnitude larger than what we applied in our simulations. For MESSy v2.52, Falk and Sinnhuber (2018) had shown an improved agreement of ozone VMR between model and
410 observation for $r_c = 10000 \text{ sm}^{-1}$.

Benavent et al. (2022) have shown the importance of iodine chemistry on the loss of tropospheric ozone during the MOSAiC campaign compared to bromine. Iodine chemistry is currently neglected in our chemistry mechanism and an inclusion would be subject to future work.

In summary, several key areas should be explored to advance our understanding of ozone and bromine chemistry in the
415 Arctic. First, contributions of bromide distribution in deeper snow layers to bromine emissions and snow metamorphism should be considered, as they play a crucial role in the cycling of bromine and ozone depletion in the region. Comparison with measured bromide in deeper snow layers will help to validate source strengths and sinks in atmospheric models. Second, the surface resistance of ozone and bromine species on different surfaces, such as ice, snow, and the ocean, needs further revision. More consistent parameterization of these resistances is necessary to better represent these reactive species. Third,
420 comprehensive heterogeneous and multiphase chemistry in the marine boundary layer is too computationally expensive in our current global CCM setup. Potentially, machine learning approaches could be used to bridge the gap from detailed box modeling studies to global scales. Finally, a higher model resolution could provide a more detailed representation of processes on inhomogeneous surfaces in the atmosphere-ocean-ice nexus, enhancing predictions of ozone and bromine concentrations, and contributing to a deeper understanding of their roles in the Arctic atmosphere.

425 *Code availability.* The Modular Earth Submodel System (MESSy) is being continuously further developed and applied by a consortium of institutions. The usage of MESSy and access to the source code is licenced to all affiliates of institutions who are members of the MESSy Consortium. Institutions can become a member of the MESSy Consortium by signing the MESSy Memorandum of Understanding. More information can be found on the MESSy Consortium website (<http://www.messy-interface.org>). The study presented here was performed based on MESSy version 2.55.2. Compared to MESSy v2.55.2, the updated AirSnow algorithm also includes two critical bug fixes concerning
430 the temperature and solar zenith angle thresholds. It has been integrated into the MESSy release candidate and will be available in the next official release (version 2.56). MYSIC based on ICDC Age of Arctic Sea Ice has been contributed to the MESSy community.

Python scripts used for data processing can be made available through github.



Data availability. Data from model experiment sfa002 will be available through the Arctic Bromine Model Intercomparison project <https://github.com/Arctic-Bromine-Model-Intercomparison/arctic-bromine-model-intercomp> led by Jennie Thomas (IGE, Grenoble).

435 Model data from the sensitivity studies presented here can be made available upon request.

TROPOMI BrO VCD data can be made available upon request from A. Richter and B. Zilker (University of Bremen).

Surface ozone data from Alert, Eureka, Nord, Summit, and Zeppelin used in this study were accessed from EBAS (<https://ebas.nilu.no>) hosted by NILU. Specifically, this includes data affiliated with the frameworks: AMAP, EMEP, GAW-WDCRG. Original data providers include: Audra McClure-Begley and Irina Petropavlovskikh (NOAA-ESRL), Wenche Aas (NILU), Mike Shaw (EC/AES), Rune Keller
440 (NERI). Surface ozone data from Utqiagvik (Barrow) were provided by NOAA Global Monitoring Laboratory, Boulder, Colorado, USA (<https://gml.noaa.gov>).

ICDC Age of Sea Ice product is distributed by CEN, University of Hamburg.

Data used in this manuscript was produced as part of the international Multidisciplinary drifting Observatory for the Study of the Arctic Climate (MOSAiC) with the tag MOSAiC20192020.

445 – Ozone (Angot et al., 2022)

– Bromine monoxide (Mahajan, 2022)

– Temperature, boundary layer height (Jozef et al., 2023)

The results contain modified Copernicus Climate Change Service information 2020. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains.



450 Appendix A: Methods




Figure A1. Maps of multi-year sea ice cover (MYSIC) in April from ICDC Age of Sea Ice (AoSI) product (middle, T42; right, T106) compared to a constant, annual mean MYSIC estimated from ERA5 sea ice cover (SIC; left) at seasonal low. T106 illustrates the gain in sea ice structure by a slightly higher model resolution. (a) 2019; (b) 2020.




Figure A2. Time series of multi-year Sea Ice covered area. All grid cells not equal zero have been selected for each year. MYSIC differ by 30%.




Figure A3. Br₂ and BrCl flux integral April 2019 with MYSIC from (a) ERA5 SIC; (b) ICDC AoSI.



Table A1. Model skill (R^2 and RMSE) at different Arctic ozone monitoring sites for different model experiments for spring 2019. For the respective 2d histograms and regressions see Appendix Fig. A4.

Experiment	Test	Utqiagvik	Eureka	Summit	Zeppelin
ref	R^2	0.19	0.01	0.22	0.01
	RMSE	0.45	0.96	0.18	0.34
sfa002	R^2	0.28	0.16	0.30	0.00
	RMSE	0.36	0.67	0.17	0.33
sfa008	R^2	0.29	0.17	0.29	0.00
	RMSE	0.36	0.66	0.17	0.33
sfa010	R^2	0.32	0.19	0.28	0.00
	RMSE	0.36	0.65	0.17	0.34
sfa011	R^2	0.28	0.17	0.30	0.01
	RMSE	0.34	0.63	0.17	0.32
sfa012	R^2	0.23	0.03	0.26	0.18
	RMSE	0.32	0.74	0.18	0.31
sfa013	R^2	0.28	0.17	0.28	0.01
	RMSE	0.34	0.64	0.17	0.32
sfa017	R^2	0.28	0.17	0.28	0.01
	RMSE	0.33	0.63	0.18	0.32




Figure A4. Sensitivity test. 2D histogram and linear fit of $\chi\text{O}_3^{\text{model}}$ vs $\chi\text{O}_3^{\text{obs}}$ for 2019, March–May. Shown are root-mean-square error (RMSE) and coefficient of determination R^2 . From top to bottom: model experiments as given in Tab. 2. Ozone monitoring sites from left to right: Utqiagvik, Eureka, Summit, and Zeppelin.



Appendix B: Results




Figure B1. Time series comparing observational records of χ_{O_3} at Arctic ozone monitoring sites with EMAC model output (ref, sfa002) interpolated to the station location. The periods March–May are highlighted in linen. The dashed red line indicates the 5 ppb threshold for ODEs. (a) Eureka; (b) Nord; (c) Summit; (d) Utqiaġvik; (e) Zeppelin.



Table B1. Model skill (R^2 and RMSE) at different Arctic ozone monitoring sites for different model experiments in 2020.

Experiment	Test	Utqiaġvik	Eureka	Nord	Summit	Zeppelin
ref	R^2	0.06	0.03	0.09	0.17	0.00
	RMSE	0.87	1.17	0.64	0.17	0.39
sfa002	R^2	0.10	0.05	0.05	0.20	0.07
	RMSE	0.79	0.87	0.61	0.17	0.34
sfa017	R^2	0.11	0.10	0.01	0.23	0.15
	RMSE	0.71	0.74	0.62	0.16	0.30




Figure B2. Monthly mean stratospheric BrO column at 13 – 14 h local time in April 2019. Shown are experiments (a) sfa008, (b) sfa010, (c) sfa013, and (d) sfa017.




Figure B3. BrO column at 13 – 14h local time in April 2019. (a) sfa008 (total column); (b) sfa008 (tropospheric column); (c) sfa010 (total column); (d) sfa010 (tropospheric column); (e) sfa013 (total column); (f) sfa013 (tropospheric column); (g) sfa017 (total column); (dh) sfa017 (tropospheric column). The modeled BrO total columns have been split into tropospheric and stratospheric columns using the modeled tropopause height on hourly basis.




Figure B4. Monthly (a) maximum and (b) mean ozone depletion in March and (c) maximum and (d) mean ozone depletion May 2020 computed from difference between experiment sfa017 and the reference simulation.



Author contributions. SF wrote and edited the manuscript, performed the model simulations, analyzed the model results for the ozone monitoring sites, bromine fluxes, and vertical column densities, and implemented and tested the model improvements. LR contributed to the analysis of MOSAiC data including the remapping of the drift track as well as pan-Arctic ozone, improvements to the setup, and identification of model bugs. BZ performed the TROPOMI analysis. AR provided the TROPOMI data. BMS proposed the project ‘BromoPole’ and contributed with insights and ideas. All authors contributed to the discussion and writing of the manuscript.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

Acknowledgements. This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) ‘517648310’. This work was performed on the HoreKa supercomputer funded by the Ministry of Science, Research and the Arts Baden-Württemberg. The authors would like to thank the German Federal Ministry of Education and Research and the German federal states (<http://www.nhr-verein.de/en/our-partners>) for supporting this work as part of the National High-Performance Computing (NHR) joint funding program. We gratefully acknowledge the funding by the DFG – project no. 268020496 - TRR 172, within the Transregional Collaborative Research Center “ArctiC Amplification: Climate relevant Atmospheric and SurfaCe Processes, and Feedback Mechanisms (AC)3”.

The authors like to thank Patrick Jöckel (DLR) for his helpful comments and insights regarding the MESSy framework and submodel setup. Special thanks go to Ole Kirner (SimLab, KIT) and Sören Johansson (IMKASF) for valuable help for the model and machine setup. Thanks to ICDC, CEN, University of Hamburg for data support, especially Stefan Kern. We thank all those who contributed to MOSAiC and made this endeavour possible (Nixdorf et al., 2021).

We thank Jennie Thomas (CNRS, Institut des Géosciences de l’Environnement, Grenoble, France) for organizing the Arctic Bromine Model Intercomparison project, which is a part of CATCH (the Cryosphere and Atmospheric Chemistry) and to which we contributed with parts of this work.

ChatGPT has been utilized to generate the manuscript abstract and title based on the Section “discussions and conclusions” as well as word smithing the outlook summary from a list of key points.



References

- Abbatt, T. P. D., Thomas, J. L., Abrahamsson, K., Boxxe, C., Granfors, A., Jones, A. E., King, M. D., Saiz-Lopez, A., Shepson, P. B., Sodeau, J., Toohey, D. W. and Toubin, C., von Glasow, R., Wren, S. N., and Yang, X.: Halogene activation via interactions with environmental ice and snow in the polar lower troposphere and other regions, *Atmos. Chem. Phys.*, 12, 6237–6271, <https://doi.org/10.5194/acp-12-6237-2012>, 2012.
- Abrahamsson, K., Granfors, A., Ahnoff, M., Cuevas, C. A., and Saiz-Lopez, A.: Organic bromine compounds produced in sea ice in Antarctic winter, *Nature Communications*, 9, <https://doi.org/10.1038/s41467-018-07062-8>, 2018.
- Ackermann, I. J., Hass, H., Memmesheimer, M., Ebel, A., Binkowski, F. S., and Shankar, U.: Modal aerosol dynamics model for Europe: development and first applications, *Atmospheric Environment*, 32, 2981–2999, [https://doi.org/https://doi.org/10.1016/S1352-2310\(98\)00006-5](https://doi.org/https://doi.org/10.1016/S1352-2310(98)00006-5), 1998.
- AMAP: Arctic Climate Issues 2011: Changes in Arctic Snow, Water, Ice and Permafrost, Overview Report, SWIPA 2011, Arctic Monitoring and Assessment Programme (AMAP), Oslo, 978-82-7971-073-8, 2012.
- Angot, H., Blomquist, B., Howard, D., Archer, S., Bariteau, L., Beck, I., Helmig, D., Hueber, J., Jacobi, H.-W., Jokinen, T., Laurila, T., Posman, K., Quéléver, L., Shupe, M. D., Schmale, J., and Boyer, M.: Ozone dry air mole fractions measured during MOSAiC 2019/2020 (merged dataset), <https://doi.org/10.1594/PANGAEA.944393>, 2022.
- Baker, K., Woody, M., Tonnesen, G., Hutzell, W., Pye, H., Beaver, M., Pouliot, G., and Pierce, T.: Contribution of regional-scale fire events to ozone and PM_{2.5} air quality estimated by photochemical modeling approaches, *Atmos. Environ.*, 140, 539–554, <https://doi.org/10.1016/j.atmosenv.2016.06.032>, 2016.
- Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J., and Rasmussen, R. A.: Ozone destruction and photochemical reactions at polar sunrise in the lower Arctic atmosphere, *Nature*, 334, 138–141, <https://doi.org/10.1038/334138a0>, 1988.
- Barten, J. G., Ganzeveld, L. N., Steeneveld, G.-J., Blomquist, B. W., Angot, H., Archer, S. D., Bariteau, L., Beck, I., Boyer, M., von der Gathen, P., Helmig, D., Howard, D., Hueber, J., Jacobi, H.-W., Jokinen, T., Laurila, T., Posman, K. M., Quéléver, L., Schmale, J., Shupe, M. D., and Krol, M. C.: Low ozone dry deposition rates to sea ice during the MOSAiC field campaign: Implications for the Arctic boundary layer ozone budget, *Elementa-Sci. Anthropol.*, 11, 00 086, <https://doi.org/10.1525/elementa.2022.00086>, 2023.
- Benavent, N., Mahajan, A. S., Li, Q., Cuevas, C. A., Schmale, J., Angot, H., Jokinen, T., Quéléver, L. L. J., Blechschmidt, A.-M., Zilker, B., Richter, A., Serna, J. A., Garcia-Nieto, D., Fernandez, R. P., Skov, H., Dumitrascu, A., Simões Pereira, P., Abrahamsson, K., Bucci, S., Duetsch, M., Stohl, A., Beck, I., Laurila, T., Blomquist, B., Howard, D., Archer, S. D., Bariteau, L., Helmig, D., Hueber, J., Jacobi, H.-W., Posman, K., Dada, L., Daellenbach, K. R., and Saiz-Lopez, A.: Substantial contribution of iodine to Arctic ozone destruction, *Nat. Geosci.*, 15, 770–773, <https://doi.org/10.1038/s41561-022-01018-w>, 2022.
- Bottenheim, J. W., Gallant, A. G., and Brice, K. A.: Measurements of NO_y Species and O₃ at 82-Degrees-N Latitude, *Geophys. Res. Lett.*, 13, 113–116, <https://doi.org/10.1029/GL013i002p00113>, 1986.
- Bougoudis, I., Blechschmidt, A.-M., Richter, A., Seo, S., Burrows, J. P., Theys, N., and Rinke, A.: Long-term time series of Arctic tropospheric BrO derived from UV–VIS satellite remote sensing and its relation to first-year sea ice, *Atmos. Chem. Phys.*, 20, 11 869–11 892, <https://doi.org/10.5194/acp-20-11869-2020>, 2020.
- Bozem, H., Hoor, P., Kunkel, D., Köllner, F., Schneider, J., Herber, A., Schulz, H., Leaitch, W. R., Aliabadi, A. A., Willis, M. D., Burkart, J., and Abbatt, J. P. D.: Characterization of transport regimes and the polar dome during Arctic spring and summer using in situ aircraft measurements, *Atmos. Chem. Phys.*, 19, 15 049–15 071, <https://doi.org/10.5194/acp-19-15049-2019>, 2019.



- 510 Chameides, W. L. and Davis, D. D.: Iodine - Its Possible Role In Tropospheric Photochemistry, *J. Geophys. Res.-Oceans Atmos.*, 85, 7383–7398, <https://doi.org/10.1029/JC085iC12p07383>, 1980.
- Choi, S., Wang, Y., Salawitch, R. J., Canty, T., Joiner, J., Zeng, T., Kurosu, T. P., Chance, K., Richter, A., Huey, L. G., Liao, J., Neuman, J. A., Nowak, J. B., Dibb, J. E., Weinheimer, A. J., Diskin, G., Ryerson, T. B., da Silva, A., Curry, J., Kinnison, D., Tilmes, S., and Levelt, P. F.: Analysis of satellite-derived Arctic tropospheric BrO columns in conjunction with aircraft measurements during ARCTAS and ARCPAC, *Atmos. Chem. Phys.*, 12, 1255–1285, <https://doi.org/10.5194/acp-12-1255-2012>, 2012.
- 515 Coburn, S., Dix, B., Edgerton, E., Holmes, C. D., Kinnison, D., Liang, Q., ter Schure, A., Wang, S., and Volkamer, R.: Mercury oxidation from bromine chemistry in the free troposphere over the southeastern US, *Atmos. Chem. Phys.*, 16, 3743–3760, <https://doi.org/10.5194/acp-16-3743-2016>, 2016.
- Cofer, W. R., Levine, J. S., Winstead, E. L., and Stocks, B. J.: Gaseous emissions from Canadian boreal forest fires, *Atmos. Environ.*, 24, 1653–1659, [https://doi.org/10.1016/0960-1686\(90\)90499-D](https://doi.org/10.1016/0960-1686(90)90499-D), 1990.
- 520 Descals, A., Gaveau, D. L. A., Verger, A., Sheil, D., Naito, D., and Peñuelas, J.: Unprecedented fire activity above the Arctic Circle linked to rising temperatures, *Sci.*, 378, 532–537, <https://doi.org/10.1126/science.abn9768>, 2022.
- Edebeli, J., Trachsel, J. C., Avak, S. E., Ammann, M., Schneebeli, M., Eichler, A., and Bartels-Rausch, T.: Snow heterogeneous reactivity of bromide with ozone lost during snow metamorphism, *Atmos. Chem. Phys.*, 20, 13 443–13 454, [https://doi.org/10.5194/acp-20-13443-](https://doi.org/10.5194/acp-20-13443-2020)
- 525 2020, 2020.
- Falk, S. and Sinnhuber, B.-M.: Polar boundary layer bromine explosion and ozone depletion events in the chemistry-climate model EMAC v2.52: implementation and evaluation of AirSnow algorithm, *Geosci. Model Dev.*, 11, 1115–1131, [https://doi.org/10.5194/gmd-11-1115-](https://doi.org/10.5194/gmd-11-1115-2018)
- 2018, 2018.
- Haine, T. W. N. and Martin, T.: The Arctic-Subarctic sea ice system is entering a seasonal regime: Implications for future Arctic amplification, *Sci. Rep.*, 7, 2045–2322, <https://doi.org/10.1038/s41598-017-04573-0>, 2017.
- 530 Helmig, D., Oltmans, S. J., Carlson, D., Lamarque, J.-F., Jones, A., Labuschagne, C., Anlauf, K., and Hayden, K.: A review of surface ozone in the polar regions, *Atmos. Environ.*, 41, 5138–5161, <https://doi.org/10.1016/j.atmosenv.2006.09.053>, recent Investigations of Snow Photochemistry and Air-Snow Exchange at Summit, Greenland, 2007.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 535 Jacobi, H. W., Morin, S., and Bottenheim, J. W.: Observation of widespread depletion of ozone in the springtime boundary layer of the central Arctic linked to mesoscale synoptic conditions, *J. Geophys. Res. Atmos.*, 115, <https://doi.org/10.1029/2010JD013940>, 2010.
- 540 Jalkanen, L. and Manninen, P.: Multivariate data analysis of aerosols collected on the Gulf of Finland, *Environmetrics*, 7, 27–38, [https://doi.org/10.1002/\(SICI\)1099-095X\(199601\)7:1<27::AID-ENV159>3.0.CO;2-3](https://doi.org/10.1002/(SICI)1099-095X(199601)7:1<27::AID-ENV159>3.0.CO;2-3), 1996.
- Jones, A. E., Anderson, P. S., Begoin, M., Brough, N., Hutterli, M. A., Marshall, G. J., Richter, A., Roscoe, H. K., and Wolff, E. W.: BrO, blizzards, and drivers of polar tropospheric ozone depletion events, *Atmos. Chem. Phys.*, 9, 4639–4652, [https://doi.org/10.5194/acp-9-](https://doi.org/10.5194/acp-9-4639-2009)
- 545 4639-2009, 2009.



- Jozef, G., Klingel, R., Cassano, J. J., Maronga, B., de Boer, G., Dahlke, S., and Cox, C. J.: Lower atmospheric properties relating to temperature, wind, stability, moisture, and surface radiation budget over the central Arctic sea ice during MOSAiC, <https://doi.org/10.1594/PANGAEA.957760>, 2023.
- Jöckel, P., Sander, R., Kerkweg, A., Tost, H., and Lelieveld, J.: Technical Note: The Modular Earth Submodel System (MESSy) - a new
550 approach towards Earth System Modeling, *Atmos. Chem. Phys.*, 5, 433–444, <https://doi.org/10.5194/acp-5-433-2005>, 2005.
- Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C. A. M., Brinkop, S., Cai, D. S., Dyroff, C., Eckstein, J., Frank, F., Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, M., Meul, S., Neumaier, M., Nuetzel, M., Oberlaender-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth System Chemistry integrated Modelling (ES-CiMo) with the Modular Earth Submodel System (MESSy) version 2.51, *Geosci. Model Dev.*, 9, 1153–1200, <https://doi.org/10.5194/gmd-9-1153-2016>, 2016.
555
- Karlsson, P., Ferm, M., Tømmervik, H., Hole, L., Karlsson, G., Ruoho-Airola, T., Aas, W., Hellsten, S., Akselsson, C., Mikkelsen, T., and Nihlgård, B.: Biomass burning in eastern Europe during spring 2006 caused high deposition of ammonium in northern Fennoscandia, *Environ. Pollut.*, 176C, 71–79, <https://doi.org/10.1016/j.envpol.2012.12.006>, 2013.
- Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöckel, P.: Technical Note: An implementation of the dry removal
560 processes DRY DEPosition and SEDimentation in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 4617–4632, <https://doi.org/10.5194/acp-6-4617-2006>, 2006a.
- Kerkweg, A., Sander, R., Tost, H., and Jöckel, P.: Technical note: Implementation of prescribed (OFFLEM), calculated (ONLEM), and pseudo-emissions (TNUDGE) of chemical species in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 3603–3609, <https://doi.org/10.5194/acp-6-3603-2006>, 2006b.
- 565 Lindsay, R. and Schweiger, A.: Arctic sea ice thickness loss determined using subsurface, aircraft, and satellite observations, *The Cryosphere*, 9, 269–283, <https://doi.org/10.5194/tc-9-269-2015>, 2015.
- Lindskog, A., Karlsson, P., Grennfelt, P., Solberg, S., and Forster, C.: An exceptional ozone episode in northern Fennoscandia, *Atmos. Environ.*, 41, 950–958, <https://doi.org/10.1016/j.atmosenv.2006.09.027>, 2007.
- Mahajan, A.: Substantial contribution of iodine to Arctic ozone destruction - data, <https://doi.org/10.17632/BN7Y7TZ4MFZ.1>, 2022.
- 570 Marelle, L., Thomas, J. L., Ahmed, S., Tuite, K., Stutz, J., Dommergue, A., Simpson, W. R., Frey, M. M., and Baladima, F.: Implementation and Impacts of Surface and Blowing Snow Sources of Arctic Bromine Activation Within WRF-Chem 4.1.1, *J. Adv. Model. Earth Syst.*, 13, <https://doi.org/10.1029/2020MS002391>, 2021.
- McClure-Begley, A., Petropavlovskikh, I., and Oltmans, S.: Barrow Atmospheric Baseline Observatory, 1973–2023-10, <https://doi.org/http://dx.doi.org/10.7289/V57P8WBF>, last accessed July 2024, 2024.
- 575 McConnell, J. C., Henderson, G. S., Barrie, L., Bottenheim, J., Niki, H., Langford, C. H., and Tempelton, E. M. J.: Photochemical Bromine Production Implicated in Arctic Boundary-Layer Ozone Depletion, *Nature*, 355, 150–152, <https://doi.org/10.1038/355150a0>, 1992.
- Nixdorf, U., Dethloff, K., Rex, M., Shupe, M., Sommerfeld, A., Perovich, D. K., Nicolaus, M., Heuzé, C., Rabe, B., Loose, B., Damm, E., Gradinger, R., Fong, A., Maslowski, W., Rinke, A., Kwok, R., Spreen, G., Wendisch, M., Herber, A., Hirsekorn, M., Mohaupt, V., Frickenhans, S., Immerz, A., Weiss-Tuider, K., König, B., Mengedocht, D., Regnery, J., Gerchow, P., Ransby, D., Krumpfen, T., Morgenstern, A., Haas, C., Kanzow, T., Rack, F. R., Saitzev, V., Sokolov, V., Makarov, A., Schwarze, S., Wunderlich, T., Wurr, K., and Boetius, A.: MOSAiC Extended Acknowledgement, <https://doi.org/10.5281/ZENODO.5541624>, 2021.
- 580 Norsk institutt for luftforskning (NILU): Database for observation data of atmospheric chemical composition and physical properties, online, <http://ebas.nilu.no/>, last accessed July 2024.



- Notz, D. and SIMIP Community: Arctic Sea Ice in CMIP6, *Geophys. Res. Lett.*, 47, e2019GL086749, <https://doi.org/10.1029/2019GL086749>, 2020.
- Oldridge, N. W. and Abbatt, J. P. D.: Formation of Gas-Phase Bromine from Interaction of Ozone with Frozen and Liquid NaCl/NaBr Solutions: Quantitative Separation of Surficial Chemistry from Bulk-Phase Reaction, *J. Phys. Chem. A*, 115, 2590–2598, <https://doi.org/10.1021/jp200074u>, 2011.
- Orbe, C., Plummer, D. A., Waugh, D. W., Yang, H., Jöckel, P., Kinnison, D. E., Josse, B., Marecal, V., Deushi, M., Abraham, N. L., Archibald, A. T., Chipperfield, M. P., Dhomse, S., Feng, W., and Bekki, S.: Description and Evaluation of the specified-dynamics experiment in the Chemistry-Climate Model Initiative, *Atmos. Chem. Phys.*, 20, 3809–3840, <https://doi.org/10.5194/acp-20-3809-2020>, 2020.
- Platt, U. and Hönninger, G.: The role of halogen species in the troposphere, *Chemosphere*, 52, 325–338, [https://doi.org/10.1016/S0045-6535\(03\)00216-9](https://doi.org/10.1016/S0045-6535(03)00216-9), naturally Produced Organohalogenes, 2003.
- Pozzer, A., Jöckel, P., Sander, R., Williams, J., Ganzeveld, L., and Lelieveld, J.: Technical Note: The MESSy-submodel AIRSEA calculating the air-sea exchange of chemical species, *Atmospheric Chemistry and Physics*, 6, 5435–5444, <https://doi.org/10.5194/acp-6-5435-2006>, 2006.
- Rantanen, M., Karpechko, A., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun. Earth Environ.*, pp. 1–10, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- Regan, H., Rampal, P., Ólason, E., Boutin, G., and Korosov, A.: Modelling the evolution of Arctic multiyear sea ice over 2000–2018, *The Cryosphere*, 17, 1873–1893, <https://doi.org/10.5194/tc-17-1873-2023>, 2023.
- Richter, A., Wittrock, F., Ladstätter-Weissenmayer, A., and Burrows, J.: Gome measurements of stratospheric and tropospheric BrO, *Adv. Space. Res.*, 29, 1667–1672, [https://doi.org/10.1016/S0273-1177\(02\)00123-0](https://doi.org/10.1016/S0273-1177(02)00123-0), 2002.
- Richter, A., Wittrock, F., Eisinger, M., and Burrows, J. P.: GOME observations of tropospheric BrO in northern hemispheric spring and summer 1997, *Geophys. Res. Lett.*, 25, 2683–2686, <https://doi.org/10.1029/98GL52016>, 1998.
- Ridley, B. A., Atlas, E. L., Montzka, D. D., Browell, E. V., Cantrell, C. A., Blake, D. R., Blake, N. J., Cinquini, L., Coffey, M. T., Emmons, L. K., Cohen, R. C., DeYoung, R. J., Dibb, J. E., Eisele, F. L., Flocke, F. M., Fried, A., Grahek, F. E., Grant, W. B., Hair, J. W., Hannigan, J. W., Heikes, B. J., Lefer, B. L., Mauldin, R. L., Moody, J. L., Shetter, R. E., Snow, J. A., Talbot, R. W., Thornton, J. A., Walega, J. G., Weinheimer, A. J., Wert, B. P., and Wimmers, A. J.: Ozone depletion events observed in the high latitude surface layer during the TOPSE aircraft program, *J. Geophys. Res. Atmos.*, 108, 1–22, <https://doi.org/10.1029/2001JD001507>, 2003.
- Salawitch, R. J., Canty, T., Kurosu, T., Chance, K., Liang, Q., da Silva, A., Pawson, S., Nielsen, J. E., Rodriguez, J. M., Bhartia, P. K., Liu, X., Huey, L. G., Liao, J., Stickel, R. E., Tanner, D. J., Dibb, J. E., Simpson, W. R., Donohoue, D., Weinheimer, A., Flocke, F., Knapp, D., Montzka, D., Neuman, J. A., Nowak, J. B., Ryerson, T. B., Oltmans, S., Blake, D. R., Atlas, E. L., Kinnison, D. E., Tilmes, S., Pan, L. L., Hendrick, F., Van Roozendaal, M., Kreher, K., Johnston, P. V., Gao, R. S., Johnson, B., Bui, T. P., Chen, G., Pierce, R. B., Crawford, J. H., and Jacob, D. J.: A new interpretation of total column BrO during Arctic spring, *Geophys. Res. Lett.*, 37, 9, <https://doi.org/10.1029/2010GL043798>, 2010.
- Sander, R., Burrows, J., and Kaleschke, L.: Carbonate precipitation in brine - a potential trigger for tropospheric ozone depletion events, *Atmos. Chem. Phys.*, 6, 4653–4658, <https://doi.org/10.5194/acp-6-4653-2006>, 2006.
- Sander, R., Baumgaertner, A., Cabrera-Perez, D., Frank, F., Gromov, S., Groöb, J. U., Harder, H., Huijnen, V., Jöckel, P., Karydis, V. A., Niemeyer, K. E., Pozzer, A., Riede, H., Schultz, M. G., Taraborrelli, D., and Tauer, S.: The community atmospheric chemistry box model CAABA/MECCA-4.0, *Geosci. Model Dev.*, 12, 1365–1385, <https://doi.org/10.5194/gmd-12-1365-2019>, 2019.



- Seo, S., Richter, A., Blechschmidt, A. M., Bougoudis, I., and Burrows, J. P.: First high-resolution BrO column retrievals from TROPOMI, *Atmos. Meas. Tech.*, 12, 2913–2932, <https://doi.org/10.5194/amt-12-2913-2019>, 2019.
- Seo, S., Richter, A., Blechschmidt, A. M., Bougoudis, I., and Burrows, J. P.: Spatial distribution of enhanced BrO and its relation to meteorological parameters in Arctic and Antarctic sea ice regions, *Atmos. Chem. Phys.*, 20, 12 285–12 312, <https://doi.org/10.5194/acp-20-12285-2020>, 2020.
- Stemmler, I., Hense, I., and Quack, B.: Marine sources of bromoform in the global open ocean - global patterns and emissions, *Biogeosciences*, 12, 1967–1981, <https://doi.org/10.5194/bg-12-1967-2015>, 2015.
- Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons, *Environ. Res. Lett.*, 13, 1–23, <https://doi.org/10.1088/1748-9326/aade56>, 2018.
- Theys, N., Van Roozendaal, M., Errera, Q., Hendrick, F., Daerden, F., Chabrillat, S., Dorf, M., Pfeilsticker, K., Rozanov, A., Lotz, W., Burrows, J., Lambert, J., Goutail, F., Roscoe, H., and Demazière, M.: Atmos. Chem. Phys. A global stratospheric bromine monoxide climatology based on the BASCOE chemical transport model, *Atmos. Chem. Phys.*, 9, 831–848, <https://doi.org/10.5194/acp-9-831-2009>, 2009.
- Tost, H., Jöckel, P. J., Kerkweg, A., Sander, R., and Lelieveld, J.: Technical note: A new comprehensive SCAVenging submodel for global atmospheric chemistry modelling, *Atmos. Chem. Phys.*, 6, 565–574, 2006.
- Toyota, K., McConnell, J. C., Lupu, A., Neary, L., McLinden, C. A., Richter, A., Kwok, R., Semeniuk, K., Kaminski, J. W., Gong, S. L., Jarosz, J., Chipperfield, M. P., and Sioris, C. E.: Analysis of reactive bromine production and ozone depletion in the Arctic boundary layer using 3-D simulations with GEM-AQ: inference from synoptic-scale patterns, *Atmos. Chem. Phys.*, 11, 3949–3979, <https://doi.org/10.5194/acp-11-3949-2011>, 2011.
- Toyota, K., McConnell, J. C., Staebler, R. M., and Dastoor, A. P.: Air-snowpack exchange of bromine, ozone and mercury in the springtime Arctic simulated by the 1-D model PHANTAS - Part 1: In-snow bromine activation and its impact on ozone, *Atmos. Chem. Phys.*, 14, 4101–4133, <https://doi.org/10.5194/acp-14-4101-2014>, 2014.
- Tschudi, M., Meier, W. N., Stewart, J. S., Fowler, C., and Maslanik, J.: EASE-Grid Sea Ice Age, Version 4.1, [northern hemisphere, 1983–2023], Tech. rep., NASA National Snow and Ice Data Center Distributed Active Archive Center, Boulder, Colorado USA, <https://doi.org/https://doi.org/10.5067/UTAV7490FEPB>, last accessed August 2023, 2024.
- Venter, Z. S., Aunan, K., Chowdhury, S., and Lelieveld, J.: COVID-19 lockdowns cause global air pollution declines, *Proc. Natl. Acad. Sci.*, 117, 18 984–18 990, <https://doi.org/10.1073/pnas.2006853117>, 2020.
- Wagner, T. and Platt, U.: Satellite mapping of enhanced BrO concentrations in the troposphere, *Nature*, 395, 486–490, <https://doi.org/10.1038/26723>, 1998.
- Wang, C., Graham, R. M., Wang, K., Gerland, S., and Granskog, M. A.: Comparison of ERA5 and ERA-Interim near-surface air temperature, snowfall and precipitation over Arctic sea ice: effects on sea ice thermodynamics and evolution, *Cryosphere*, 13, 1661–1679, <https://doi.org/10.5194/tc-13-1661-2019>, 2019a.
- Wang, S., Kinnison, D., Montzka, S. A., Apel, E. C., Hornbrook, R. S., Hills, A. J., Blake, D. R., Barletta, B., Meinardi, S., Sweeney, C., Moore, F., Long, M., Saiz-Lopez, A., Fernandez, R. P., Tilmes, S., Emmons, L. K., and Lamarque, J.-F.: Ocean Biogeochemistry Control on the Marine Emissions of Brominated Very Short-Lived Ozone-Depleting Substances: A Machine-Learning Approach, *J. Geophys. Res. Atmos.*, 124, 12 319–12 339, <https://doi.org/10.1029/2019JD031288>, 2019b.



- Wang, S., Mcnamara, S. M., Moore, C. W., Obrist, D., Steffen, A., Shepson, P. B., Staebler, R. M., Raso, A. R. W., Pratt, K. A., and
Performed, K. A. P.: Direct detection of atmospheric atomic bromine leading to mercury and ozone depletion, *ATMOSPHERIC, AND*
660 *PLANETARY SCIENCES*, 116, <https://doi.org/10.18739/A2D79598P>, 2019c.
- Warwick, N. J., Pyle, J. A., Carver, G. D., Yang, X., Savage, N. H., O'Connor, F. M., and Cox, R. A.: Global modeling of biogenic bromo-
carbons, *J. Geophys. Res.-Atmos.*, 111, <https://doi.org/10.1029/2006JD007264>, 2006.
- Wesely, M. L.: Parameterization Of Surface Resistances To Gaseous Dry Deposition In Regional-Scale Numerical-Models, *Atmos. Environ.*,
23, 1293–1304, [https://doi.org/10.1016/0004-6981\(89\)90153-4](https://doi.org/10.1016/0004-6981(89)90153-4), 1989.
- 665 Whaley, C. H., Law, K. S., Hjorth, J. L., Skov, H., Arnold, S. R., Langner, J., Pernov, J. B., Bergeron, G., Bourgeois, I., Christensen, J. H.,
Chien, R.-Y., Deushi, M., Dong, X., Effertz, P., Faluvegi, G., Flanner, M., Fu, J. S., Gauss, M., Huey, G., Im, U., Kivi, R., Marelle, L.,
Onishi, T., Oshima, N., Petropavlovskikh, I., Peischl, J., Plummer, D. A., Pozzoli, L., Raut, J.-C., Ryerson, T., Skeie, R., Solberg, S.,
Thomas, M. A., Thompson, C., Tsigaridis, K., Tsyro, S., Turnock, S. T., von Salzen, K., and Tarasick, D. W.: Arctic tropospheric ozone:
assessment of current knowledge and model performance, *Atmos. Chem. Phys.*, 23, 637–661, <https://doi.org/10.5194/acp-23-637-2023>,
670 2023.
- Yang, X., Cox, R. A., Warwick, N. J., Pyle, J. A., Carver, G. D., O'Connor, F. M., and Savage, N. H.: Tropospheric bromine chemistry and
its impacts on ozone: A model study, *J. Geophys. Res. Atmos.*, 110, 1–18, <https://doi.org/https://doi.org/10.1029/2005JD006244>, 2005.
- Yang, X., Pyle, J. A., and Cox, R. A.: Sea salt aerosol production and bromine release: Role of snow on sea ice, *Geophys. Res. Lett.*, 35, 1–5,
<https://doi.org/https://doi.org/10.1029/2008GL034536>, 2008.
- 675 Yang, X., Pyle, J. A., Cox, R. A., Theys, N., and Van Roozendaal, M.: Snow-sourced bromine and its implications for polar tropospheric
ozone, *Atmos. Chem. Phys.*, 10, 7763–7773, <https://doi.org/10.5194/acp-10-7763-2010>, 2010.
- Yumashev, D., van Hussen, K., Gille, J., and Whiteman, G.: Towards a balanced view of Arctic shipping: estimating economic impacts of
emissions from increased traffic on the Northern Sea Route, *Climatic Change*, 143, 143–155, <https://doi.org/10.1007/s10584-017-1980-6>,
2017.
- 680 Zhai, S., Swanson, W., McConnell, J. R., Chellman, N., Opel, T., Sigl, M., Meyer, H., Wang, X., Jaeglé, L., Stutz, J., Dibb, J. E., Fujita, K.,
and Alexander, B.: Implications of Snowpack Reactive Bromine Production for Arctic Ice Core Bromine Preservation, *J. Geophys. Res.*
Atmos., 128, e2023JD039 257, <https://doi.org/10.1029/2023JD039257>, 2023.