



CMIP7 Data Request: Land and Land Ice Priorities and Opportunities

5 Yue Li¹, Gang Tang^{2,3}, Eleanor O'Rourke⁴, Samar Minallah⁵, Martim Mas e Braga⁶, Sophie Nowicki⁷,
Robin S. Smith⁸, David M. Lawrence⁵, George C. Hurtt⁹, Daniele Peano¹⁰, Gesa Meyer¹¹, Birgit Hassler¹²,
Jiafu Mao¹³, Yongkang Xue¹, Martin Jukes¹⁴

¹Department of Geography, University of California, Los Angeles, 90095, USA

²School of Geography, Earth and Atmospheric Sciences, The University of Melbourne, Melbourne, 3010, Australia

³Department of Biogeochemical Signals, Max Planck Institute for Biogeochemistry, Jena, Germany

⁴CMIP International Project Office, ECSAT, Harwell Science & Innovation Campus, OX11 0FD, UK

10 ⁵Climate and Global Dynamics Laboratory, NSF National Center for Atmospheric Research, Boulder, CO 80301, USA

⁶British Antarctic Survey, Cambridge, CB3 0ET, UK

⁷Department of Earth Sciences, University at Buffalo, Buffalo, 14260, USA

⁸National Centre for Atmospheric Science, Department of Meteorology, University of Reading, Reading, RG6 6ES, UK

⁹Department of Geographical Sciences, University of Maryland, College Park, 21403, USA

15 ¹⁰CMCC Foundation - Euro-Mediterranean Center on Climate Change, Bologna, 40127, Italy.

¹¹Environment and Climate Change Canada, Climate Research Division, Victoria, V8N 1V8, Canada

¹²Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Physik der Atmosphäre, Weßling, 82234, Germany

¹³Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37830, USA

¹⁴University of Oxford, and UKRI STFC, UK

20 *Correspondence to:* Yue Li (yli@geog.ucla.edu)

Abstract. The Land and Land Ice Theme in the Coupled Model Intercomparison Project Phase 7 (CMIP7) represents the current understanding of physical processes in land surface ecosystems, hydrology, cryosphere, and their physical interactions with other Earth system components. Simulations from Earth system models (ESMs) could provide crucial information for assessing planetary safety, such as critical tipping elements, and be used to inform climate risks for improving climate impact assessments and policy decisions. This paper presents a collaborative effort to identify scientific opportunities in the Land and Land Ice Theme of the CMIP7 Data Request. The proposed opportunities build upon advances in ESMs, including new freshwater system and land ice processes being included in CMIP7, as well as the scientific community's demand for high-frequency and sub-grid-scale land surface outputs. In total, 25 variable groups that contain 716 variables have been identified to be potentially available to the broad scientific audience for performing analysis in land-atmosphere coupling, hydrological processes and freshwater systems, glacier and ice sheet mass balance and their influence on the sea levels, land use, and plant phenology. Key reflections from this data request effort include advocacy for closer engagement between the user community and modeling groups, reduction in the technical barriers to tracking existing parameters and defining new variables, and more streamlined variable management. These will be essential to enhance the usability and reliability of CMIP7 outputs for climate and Earth system research and applications to a broad audience that relies on the CMIP7 endeavor.

25
30



35 1 Introduction

Land and Land Ice (including glaciers, ice sheets, and ice shelves) processes are essential components in Earth system models (ESMs) to simulate and project future climate change and assess climate change impacts (Eyring et al., 2016, Lee et al., 2021). Key physical processes within the Land and Land Ice Theme include energy and hydrological interactions associated with land-atmosphere feedbacks, land use and land cover changes, vegetation, and cryosphere dynamics. Since the Coupled Model
40 Intercomparison Project Phase 6 (CMIP6), grand challenges have emerged in Earth system modeling to integrate these processes and capture interactions within planetary boundaries (Richardson et al., 2023) and the potential tipping elements that could result in dangerously rapid or irreversible changes to the Earth system (Armstrong et al., 2022). These have posed a higher requirement for improving ESMs with available observational data and improved understanding of Land and Land Ice processes. In this context, new “Opportunities” need to be identified for CMIP7 models to map variables that meet the needs
45 of scientific analyses and societal use and impact assessments. Building upon the provisional planning of the Baseline Climate Variables (BCVs) of CMIP7 (Jukes et al., 2025), this paper develops a collection of harmonized variables within the Land and Land Ice Theme that are available for the Land and Land Ice communities to work with and tackle these challenges.

Land-atmosphere (L-A) interactions represent a critical pathway through which land and land-ice processes regulate the
50 climate system through surface-air energy and water exchanges. Enhanced understanding of L-A interactions at various spatiotemporal scales has the potential to improve Earth system simulations of climate extremes, such as heat extremes (Dirmeyer et al., 2021; Biess et al., 2024) and tropical extreme rainfall (Negron-Juarez et al., 2024). These extreme events, including the intensification of subdaily wildfires due to meteorological dryness in the recent two decades (Balch et al., 2022; Luo et al., 2024a), have been reported to lead to significant societal consequences such as increased mortality from wildfire
55 air pollution (Park et al., 2024) and threatened energy systems (Zhao et al., 2023). Since CMIP6, significant progress has been made in evaluating daily and subdaily precipitation against observational data (Norris et al., 2021; Wehner et al., 2021; Lei et al., 2022; Dong et al., 2023). However, the reliance on observation-based reanalysis datasets, which provide high-frequency data, has revealed that key L-A interactions—such as their role in regional monsoon systems (Chauhan et al., 2023) and climate extremes like drought progression (Herrera-Estrada et al., 2019; Wu and Dirmeyer, 2020; Schumacher et al., 2022; Yoon et
60 al., 2023) and heatwaves (Chen et al., 2025)—are still not well represented in models, partially due to insufficient temporal resolution. To bridge this gap, ESMs must incorporate high-frequency (e.g., 3-hourly) output for L-A metrics, ensuring robust comparisons with observation and reanalysis data and improving the representation of rapid land-surface feedbacks and extreme event development, such as the morning flux and afternoon rainfall (Findell et al., 2024).

65 Hydrological processes are central to land and land-ice dynamics, with water connecting all components of the Earth system. Accurate and comprehensive representation of hydrological processes in ESMs is essential for assessing the water cycle, freshwater systems, and the associated feedback with the other components. Typical CMIP6 hydrological variables include



70 surface water balance, such as precipitation, evapotranspiration, runoff, and soil moisture (Cook et al., 2020). With the past
CMIP cycles, progress has been made to improve model representation of hydrological processes and highlight the need to
advance the assessment of freshwater systems, including lakes (Minallah and Steiner 2021; Briley et al. 2021; Notaro et al.
2022), artificial reservoirs (e.g., big dams, Vanderkelen et al., 2022), rivers (Lehner et al. 2019), estuaries (Sun et al., 2017;
Vettoretti et al., 2024), groundwater (Wu et al. 2021), permafrost (Li et al., 2010; Koven et al. 2013; Wang et al., 2017; Burke
et al. 2020), snowpack (Sun et al., 1999; Kouki et al. 2022; Zhang et al. 2022), and glaciers (Shrestha et al., 2015). It is thus
75 essential to propose a comprehensive CMIP7 hydrological variable set, that allows for accurately assessing and predicting
future changes in freshwater resources, water availability, and hydrometeorological extremes, and support hydrological
modeling applications in ESM frameworks.

Glacier and ice sheet (including ice cap and floating shelf) processes, are essential for understanding land ice interactions with
the atmosphere and ocean, making them a priority for CMIP7 Earth system modeling. Glaciers and ice sheets play a crucial
80 role in sea-level rise, freshwater availability, and climate feedbacks (e.g., ice-albedo feedback), yet their future evolution with
climate warming remains highly uncertain (Fox-Kemper et al. 2021). Mountain and polar glaciers, including those in
Greenland and (sub-)Antarctica (RGI Consortium 2023), were the primary contributors to 20th-century sea-level rise (Gregory
et al. 2013; Oppenheimer et al. 2019) and are expected to remain dominant drivers in the 21st century (Fox-Kemper et al.
2021). They also sustain ecosystems (Drenkhan et al., 2023), regulate hydroclimates (Milner et al. 2017), and provide vital
85 freshwater resources (Hock et al. 2019b; Huss and Hock 2018). Their sensitivity to climate change makes them regional tipping
elements, increasing risks of geohazards such as glacial lake outburst floods and avalanches (Wolken et al. 2021; Hock and
Truffer 2024). Ice sheet mass loss further accelerates sea-level rise, alters ocean circulation, and could trigger cascading Earth
system tipping points (Golledge et al 2019, Armstrong et al., 2022). To enhance projections of cryosphere dynamics in ESMs,
it is necessary to provide comprehensive variable groups in CMIP7 to assess the mass balance, contributions to sea-level rise,
90 and ice-climate feedbacks of glaciers, ice sheets, and ice shelves. Leveraging CMIP7 data will enhance understanding of glacier
and ice sheet responses to climate change, reducing uncertainties and strengthening global climate impact assessments.

Land-use change is a major driver of land-atmosphere and land-ice interactions, requiring improved representation in ESMs.
Human land use and management have profoundly altered the Earth's surface, driving changes in climate (Li et al., 2022b),
95 carbon dynamics (Hogan et al., 2025), water cycle (Sterling et al., 2013; Mao et al., 2015; te Wierik et al., 2021), and surface
energy fluxes (Lawrence et al., 2022). As global demands continue to increase for resources such as food and energy, the
expected intensification of land use will exert climatic and biogeochemical impacts. Within the context of CMIP6 and the
Land Use Model Intercomparison Project (LUMIP, Lawrence et al. 2016), land use impacts on climate and biogeochemical
cycling have been extensively assessed. Example studies that relied on the CMIP6-LUMIP simulations include quantification
100 of the biophysical climate effects of land use and land cover change (LULCC) generally (Boysen et al., 2020; Tang et al.,
2023; Luo et al., 2024b; Li et al., 2025a) and irrigation specifically (Al-Yaari et al., 2022), as well as an assessment of the role



of agricultural processes, especially fertilization, in amplifying the annual cycle of land-atmosphere CO₂ exchange (Lombardozzi et al., 2025). A major limitation in assessing the impacts of LULCC is the lack of sub-grid-scale modeled data that accurately reflect land surface heterogeneity, which is largely driven by human land transformations. Provision of this sub-grid information will allow for detailed ongoing assessment into the LULCC impacts on climate and the carbon cycle, even when only all-forcing historical runs are available.

Last but not least, vegetation on land is a key land process that influence the Earth system simulations through the cycles of energy (e.g., surface albedo), water, carbon, and nutrient cycles (e.g., nitrogen) and are thus crucial for CMIP7 ESM simulations. The variability in timing and duration of a plant's active season (i.e. phenology) influences the exchanges of energy, carbon, and water between land and atmosphere as observed in the last decades (e.g. Richardson et al., 2013; Keenan et al., 2014; Park et al., 2016). Similarly, vegetation greening unveils the relationship between land and atmosphere (e.g. Zhu et al., 2016; Munier et al., 2018; Winkler et al., 2021; Yu et al., 2021). Consequently, it is paramount to assess and understand the ability of ESMs to accurately reproduce the vegetation phenology and quantity. This evaluation can be performed using the Leaf Area Index (LAI) variable as done in previous studies (e.g. Murray-Tortarolo et al., 2013; Peano et al., 2021,2025; Li et al., 2022a). However, ESMs estimate LAI through the Plant Functional Types (PFT) discretization, which differs in each model. For this reason, it is relevant to assess plant phenology at a standard vegetation type level to increase the understanding of model limitations (e.g. Li et al., 2024b).

This theme paper will focus on model output variables required for the physical Land and Land Ice analysis. Biogeochemical and nutrient interactions are generally out-of-scope and are included in the Earth system theme (see McPartland et al., 2025) - except for the land use Opportunity due to the tight coupling between physical variables and biogeochemical fluxes. Full pressure-level atmospheric dynamics variables are included in the BCVs and Atmosphere theme. The CMIP experiment communities that are related to the data request from this theme include the High Resolution Model Intercomparison Project (HighResMIP, <https://highresmip.org/>, Roberts et al., 2025), the Ice Sheet Model Intercomparison Project for CMIP7 (ISMIP7, <https://climate-cryosphere.org/about-ismip6/>, Nowicki et al., 2024), the Land Use Model Intercomparison Project (LUMIP, <https://www.cesm.ucar.edu/projects/cmip6/lumip>, Lawrence et al., 2016), the Snow Model Intercomparison Project (SnowMIP, <https://www.wcrp-climate.org/modelling-wgcm-mip-catalogue/modelling-wgcm-mips-2/276-modelling-wgcm-catalogue-snowmip>, Krinner et al., 2018), the Irrigation Impacts Model Intercomparison Project (IRRMIP, <https://hydr.vub.be/projects/irrmip>, Yao et al., 2025), the Tipping Point Model Intercomparison Project (TIPMIP, <https://tipmip.pik-potsdam.de/>, Walton et al., 2025), the Marine Ice Sheet Ocean Model Intercomparison Project (MISOMIP, <https://misomip.github.io/>, De Rydt et al., 2024), the Glacier Model Intercomparison Project (GlacierMIP, <https://climate-cryosphere.org/glaciermip/>, Hock et al., 2019a), and Paleoclimate Modeling Intercomparison Project (PMIP, <https://pmip.lsce.ipsl.fr/>, Kageyama et al., 2018).



135 **2 Approach and methodology**

The Land and Land Ice Theme author team was recruited via an open call between 13 February and 8 March 2024 (<https://wcrp-cmip.org/cmip7-land-ice-call/>). Members were sought from across the Land and Land Ice communities to gather variable requirements for the CMIP7 Data Request (overview available at Mackallah et al, 2025). Applications were reviewed by LUMIP and ISMIP representatives alongside two members of the Data Request Task Team. A diverse final group of 21
140 committee members was chosen, including representatives from Climate and Cryosphere (CliC) project, Global Energy and Water Exchanges (GEWEX) program, PMIP, GlacierMIP, TIPMIP, Reduced Complexity Mode Intercomparison Project (RCMIP), the Fire Model Intercomparison Project (FireMIP), and the Global Climate Observing System-Terrestrial Observation Panel for Climate (GCOS-TOPC) with theme members spanning a range of geographical regions, career stages, and CMIP experiences, keeping in mind necessary expertise as well as geographic and gender diversity.

145

Many members of this team were not involved in previous data requests, but all were creators and/or users of past CMIP or land/ice sheet model data. Consequently, new ideas and opinions (see section 4) are combined with user experiences from CMIP6 and earlier CMIP rounds of data requests to identify:

- What was effective, e.g., the new Opportunities related to land ice as in the results.
- 150 • What was not, e.g., variables proposed by Opportunities that are already included in the baseline climate variable (see Juckes et al., 2025).
- Information about which variables were downloaded, e.g., download frequency available from Airtable, and which were used for major publications and assessment reports such as the United Nations Intergovernmental Panel on Climate Change (IPCC) Sixth Assessment Report (AR6).

155 Every member of the Land and Land Ice Theme was provided with a variable list that organized variables (as well as their downloading frequency in CMIP6) using several categories such as energy balance, water cycle, and land ice associated group. This helped each Opportunity proposer to avoid redundancy with existing variables or variable groups.

The team first convened on 19 July 2024, with community engagement activities beginning in parallel with the first public
160 consultation. Author team members were requested to utilize their networks as community representatives to gather scientific requirements for the Land and Land Ice Theme of the CMIP7 Data Request. Author team members used the Airtable cloud platform to propose the scientific Opportunities associated with the Land and Land Ice Theme. For each Opportunity, proposers had to define variable groups with different priority which included variables with specific requirements for the variable dimensions (e.g., latitude, longitude, and pressure levels), spatial and temporal average approach, and other property
165 information such as frequency (e.g., 3 h or daily). In the CMIP Data Request, variables are constructed by combining a “physical parameter” (with an attached CF standard name as in Hassell et al., 2017) with additional metadata to describe its spatial and temporal sampling (Mackallah et al., 2025). Opportunity proposers also had to propose a new physical parameter




if a new variable did not have a CF standard name (i.e., a string that identifies a geophysical quantity in a data variable) via Github webpage (<https://github.com/cf-convention/vocabularies/issues>).

170

In the first phase of consultation six Opportunities were submitted with the initial selection of variables and their technical definitions (Fig. 1). The author team met every two to three weeks to discuss the submitted Opportunities, identify any remaining gaps, integrate input from the wider community, highlight areas requiring cross thematic input and focus on variable group development and refinement.

175



180

Figure 1: Schematic diagram illustrating the Land and Land Ice Theme Opportunities. Land-atmosphere coupling Opportunity covers high-frequency (3hr) surface radiation, water, and energy fluxes. Hydrology Opportunity contains variables covering hydrological processes, water resources, and freshwater systems. Glacier and Ice sheet Opportunities cover the mass balance, energy, and water input to climate system. Land use and plant phenology Opportunities cover variables associated with sub-grid land use tiles and vegetation types, respectively. Figure originally published at Li et al., 2025b (see <https://doi.org/10.5281/zenodo.15681950>)

3 Information management and decision making



185 A harmonization sprint, involving all thematic teams, was held in September 2024, which resulted in the merging of several Opportunities within and across themes, and designated a lead theme for each Opportunity. The Land and Land Ice Theme moved forward, leading six Opportunities (see Table 1) plus the Land and Land Ice relevant variables of the Rapid Evaluation Framework opportunity (full details on this Opportunity can be found in Dingley et al, 2025). Following the v1.0 release in November 2024 the team focused on finalizing variable groups, supporting processing of new variables, and contributing to cross-thematic meetings on topics including time subsets. Regular team meetings continued, focused on Opportunity or variable specific requirements. Team members engaged proactively across themes, particularly with the Earth System thematic author team through a dedicated liaison member.

195 Collaborative spreadsheets were utilized to gather input between meetings with some members of the team interacting directly with the Airtable online database from a very early stage with IPO support and Data Request Task Team liaison members updating the Airtable records as needed. A systematic variable review was conducted during the Data Request’s Phase 2 and Phase 3 public consultation periods to address comments, rectify errors and highlight remaining outstanding issues to the team. Author team members also contributed to cross-thematic meetings focused on issues such as consistent use of time subsets, ensemble sizes, or prioritization of variable groups within Opportunities. By the end of the data request efforts, a total of 25 variable groups that contain 716 variables had been identified (Table 1).

Table 1. The final list of Land and Land Ice Theme Opportunities. The Opportunity ID indicates the identifier of each Land and Land Ice Theme Opportunity in the full CMIP7 Data Request content release of v1.2.

ID	Opportunity Title	Variable Groups	Number of experiments	Number of Variables
1	Accurate assessment of land-atmosphere coupling	2	14	24
38	Assessments for Hydrological Processes, Water Resources, and Freshwater Systems	8	21	199
35	Glacier changes, drivers, and impacts	5	16	176
39	Ice sheet mass loss, contributions to sea level rise and freshwater flux input to the climate system	7	26	284
43	Land use change	1	15	27
53	Plant Phenology	2	15	6
55	Rapid Evaluation Framework	1	52	20 (Land and Land Ice relevant)



4 Results: Content included in the CMIP7 Data Request

205 Given the important role of these land and land ice processes in ESM simulations in CMIP7, the Land and Land Ice Theme proposed the following scientific questions that could be addressed by the CMIP7 Data Request. These variables proposed by the Opportunities in Table 1 could be used by the Land and Land Ice Theme communities to analyze and comprehensively advance the understanding of the Earth planetary boundary. Specifically, the proposed Opportunities aim to address:

- How do high-frequency land-atmosphere coupling processes influence surface energy fluxes and water movement through the soil-plant-atmosphere continuum and how does this imply for the dynamics of climate extremes, and how can improved model simulations reduce diurnal and long-term uncertainties in these interactions?
- How do hydrological processes influence water availability, distribution, and feedback mechanisms across different Earth system components?
- How do past, present, and future changes in mountain and polar glaciers influence sea-level rise, regional hydroclimates, freshwater resources, and geohazards under different climate scenarios?
- How will the Greenland and Antarctic ice sheets evolve under future climate scenarios, will they act as tipping points in the Earth system, and how does their mass loss contribute to sea level rise, and impact on ocean circulation?
- How do LULCC, including land management practices, impact climate, biogeochemical cycles, and surface fluxes and stocks of carbon and water, as well as energy fluxes? Furthermore, in what ways can these changes contribute to climate change mitigation?
- How do vegetation type and daily frequency data reveal the seasonality and long-term trends of plant phenology that influence terrestrial carbon, energy, and water fluxes, and how can improved modeling of phenological processes enhance ESM simulations of land-atmosphere interactions?

4.1 Accurate assessment of land-atmosphere coupling (ID 1)

225 Land and atmosphere are closely coupled due to the surface soil and vegetation status providing critical boundary conditions for the surface energy flux, such as the latent and sensible heat fluxes. Changes of these surface energy and water fluxes will interact with and feedback to the atmospheric thermodynamics and circulation, which influence surface air temperature, humidity, and precipitation. Previous studies have shown that critical land-atmosphere coupling process, such as dynamics in evaporative fraction (i.e., the ratio of latent heat flux to the total turbulent heat fluxes) will influence the rainfall at sub-daily (e.g., Findell et al., 2011), seasonal (e.g., Koster et al., 2004; Seneviratne et al., 2006), and annual timescales (Li et al., 2022b). Despite the current CMIP6 monthly output of surface energy and water fluxes, soil, and vegetation variables, 3-hourly output in land-atmosphere coupling is urgently needed for the next-generation of global ESMs, which will provide the opportunity of being compared to the observation-based reanalysis data and flux towers (Findell et al., 2024). High-frequency land-atmosphere data are valuable to understand, and accurately model, key processes related to short-period (e.g., diurnal scale) but high intensity climate extremes such as heatwaves, droughts, and the onset of rainfall.



This Opportunity will provide 3-hourly data of critical variables in land-atmosphere coupling from the surface to specified lower-tropospheric pressure levels (Table 2). Critical local land-atmosphere coupling metrics can be computed based on variables from the group landatm_coupling_3hA, such as the use of mixing diagrams (Santanello et al., 2009, 2011), two-
240 legged metrics (Seo and Dirmeyer, 2022), and the triggering feedback strength (Findell et al., 2015). In another group with medium priority, high-frequency (i.e., 3 hourly) variables include solar radiation, partitioning of vegetation water fluxes (e.g., transpiration), soil moisture at different depths, and the pressure-level atmospheric variables (e.g., eastward wind). These variables will significantly enhance the exploration of mechanisms through which soil and vegetation influence the surface energy balance and affect the local atmospheric state. The variables in this Opportunity can also be used to evaluate ESM
245 simulations using the high-frequency observational data (Findell et al., 2024).

Two time subsets of 20 year simulations of historical and future scenarios have been requested. This is because previous studies suggest that a minimum of 10 years of data are required for robust statistical analysis (Findell et al., 2015). These two sets of high-frequency model output in land-atmosphere coupling will provide opportunities for model evaluations using direct
250 observations from field campaigns and measurements from reanalysis data. The multi-model intercomparisons enabled by these variables will demonstrate the modeling uncertainties in land-atmosphere coupling processes and be informative for the other MIPs that include land-atmosphere processes (e.g., HighResMIP). Lastly, this Opportunity will help to identify sources of process uncertainty, leading toward solutions for improvement of sub-daily soil and vegetation processes in the models.

255 **Table 2. Variable groups needed for ID 1 Accurate assessment of land-atmosphere coupling.**

Variable group name	Reason for inclusion
landatm_coupling_3hA	This variable group allows for adequate characterization of sub-daily coupling process between the land and the atmosphere.
landatm_coupling_3hr_medium	This variable group will add variables that estimate deeper soil moisture information and component terms of net radiation and evapotranspiration.

4.2 Assessments for hydrological processes, water resources, and freshwater systems (ID 38)

Water is the essential element linking all Earth system components – the atmosphere, hydrosphere, biosphere, geosphere, and cryosphere. Therefore, accurate representation of hydrological processes in ESMs is important to assess the distribution and
260 availability of freshwater resources and the feedback mechanisms between water, energy, and biogeochemical cycles (Clark et al., 2015). Climate change has significant and far-reaching impacts on water resources and freshwater systems, affecting their availability, quality, and distribution across the globe (Huntington, 2005). Changing temperatures and precipitation patterns, hydroclimatic shifts, and increasing risks of extreme events (floods and droughts) pose major challenges for ecosystems, agriculture, human health, and food and water security (Konapala et al., 2020).

265



This Opportunity is aimed at leveraging CMIP data for assessments, projections, and predictions of hydrological processes, water resources, and freshwater systems, including rivers and estuaries, lakes and inland water bodies, groundwater, snowpack, permafrost, and glaciers. The variables identified in this Opportunity are of interest to a large number of hydrological applications, including (1) comprehensive assessments of regional hydroclimates and their past, present, and future states, (2) assessments on the usability of Earth system models for water resources applications, (3) evaluating whether these processes/systems are adequately represented in the modelling frameworks, (4) offering opportunities for model intercomparisons across these topics (e.g., SnowMIP, IRRMIP, etc.), (5) providing input for hydrology, lake, snow, and other models, and (6) climate impact assessments of freshwater resources and water availability (Table 3).

The main variable group for this Opportunity is *hydrology* which includes variables related to (1) lakes and inland water bodies (sea/lake surface water temperatures, lake fraction, lake depth, etc.) that are used in assessments of lake-atmosphere interactions and usability of CMIP models for large lakes (Minallah and Steiner 2021, Briley et al. 2021, Notaro et al. 2022), (2) runoff to assess model hydrological sensitivity and improve streamflow projections (Lehner et al. 2019), (3) snow water equivalent and soil moisture to assess terrestrial water storage (Wu et al. 2021), and (4) permafrost and snow to study their sensitivity to climate change (Koven et al. 2013, Burke et al. 2020, Kouki et al. 2022, Zhang et al. 2022). Additionally, water budget variables (evaporative fluxes, soil moisture, runoff, potential evapotranspiration) are important for assessments of water availability and security, hydrometeorological extremes, and for hydrological simulations using watershed models.

Table 3. Variable groups needed for ID 38 Assessments for Hydrological Processes, Water Resources, and Freshwater Systems.

Variable group name	Reason for inclusion
hydrology	Aggregates variables related to terrestrial water storage, lakes, rivers and estuaries, runoff, groundwater, permafrost, snow, and glaciers.
WaterResources_subdaily WaterResources_daily WaterResources_monthly	Include potential evapotranspiration, runoff components, and soil moisture at different temporal resolutions (from sub-daily to monthly) for assessments of hydrological extremes, specifically floods and droughts.
snow_allfreq	Includes snow-related variables at fixed, daily, and monthly steps, important for cryosphere-hydrology assessments.
baseline_daily baseline_monthly	Include water cycle variables (e.g., precipitation, evaporation fluxes, etc.), and other important quantities (winds, temperatures) required for hydrology, snow, lake, and other models.



4.3 Glacier changes, drivers, and impacts (ID 35)

This Opportunity aims at glaciological assessments across timescales in the past, present, and future. Glaciers here are defined as mountain and polar glaciers, including peripheral glaciers in Greenland and (sub-)Antarctica that are decoupled from the ice sheets, as identified in the Randolph Glacier Inventory (RGI Consortium 2023). These glaciers were considered the primary contributors to 20th-century sea level rise from the cryosphere (Gregory et al. 2013; Oppenheimer et al. 2019). Based on the future scenarios in the IPCC AR6, glaciers are expected to remain the primary drivers of sea-level rise during the 21st century, in most scenarios exceeding the contributions from the two ice sheets (Fox-Kemper et al. 2021). Additionally, they have significance in sustaining ecosystems and modulating regional hydroclimates (Milner et al. 2017). They are a substantial freshwater resource, especially in mountainous regions where glaciers are important for water and food security for a large population (Hock et al. 2019b; Huss and Hock 2018). Through feedback in the mass balance response to climate change, glaciers are regional-scale tipping elements in the Earth system (Abrams et al. 2023; Davies et al. 2024) and are a driving factor in related geohazards, including glacial lake outburst floods, slope failures, avalanches, and ice falls (Wolken et al. 2021; Hock and Truffer 2024).

The variables requested for this Opportunity are critical as inputs for the glacier modelling community and to study the impacts of climate change on glaciers and glacier changes on other Earth system components (Table 4). The Glacier Model Intercomparison Project (GlacierMIP, Hock et al. 2019a) aims to advance our understanding of these impacts and reduce uncertainties in glacier projections through a coordinated intercomparison effort of global-scale glacier models. The results from the first two phases of GlacierMIP have informed the IPCC Special Report on the Cryosphere and the Oceans in a Warming Climate (IPCC 2019; Hock et al. 2019b) and IPCC AR6 report (Marzeion et al. 2020). The core variable group for this Opportunity is `landice_glaciers_allfreq`, which requests variables at daily and monthly temporal resolutions necessary for glacial mass balance computations, including temperature, precipitation fluxes, and energy budget quantities. Additional applications of this Opportunity are related to mountain cryosphere, glacio-hydrology, and local to regional-scale geohazards associated with glacier changes under different scenarios. For accurate partitioning between glacier-melt, snowmelt, and baseflow contributions to runoff in land and hydrological models at sub-seasonal to seasonal timescales, variables from terrestrial hydrology and snow processes at daily resolutions are also requested (`snow_allfreq`). Variable groups `baseline_fixed`, `baseline_daily`, and `baseline_monthly` host other key variables relevant for these assessments, including land cover types, surface elevation, winds, and heat fluxes.

Table 4. Variable groups needed for ID 35 Glacier changes, drivers, and impacts.

Variable group name	Reason for inclusion
---------------------	----------------------



landice_glaciers_allfreq	Core variable group for the Opportunity with variables required for GlacierMIP and other modeling activities
baseline_daily baseline_monthly	Include variables for glacier-climate impact assessments (e.g. precipitation fluxes, temperature, winds, etc.) at daily and monthly timesteps
snow_allfreq	Important variables for snow-glacier interface

4.4 Ice sheet mass loss, contributions to sea level rise and freshwater flux input to the climate system (ID 39)

320 The mass loss from the ice sheets of Greenland and Antarctica is a key contributor to future sea level rise, and their freshwater
input to the climate system provides an important forcing for ocean circulation and carbon and heat uptake. Sea level rise
affects every landmass on Earth and is one of the most challenging aspects of climate change to mitigate or reverse. Ice sheets
have also been theorized to be a key tipping element in the Earth system, with the potential for irreversible changes in this
century which may trigger a cascade of other tipping elements around the globe. Characterizing and understanding the future
325 evolution of the ice sheets and their interactions with the global climate system thus has the potential for huge societal impact.
There are many aspects of modelling and projecting large scale ice sheet change that are not yet understood sufficiently, so it
is important to leverage the broad spectrum of information in CMIP7 to add to our ability to understand and project how ice
sheets may change in the future.

330 For CMIP6, ISMIP6 (Nowicki et al. 2016, 2020) requested land ice data for the first time within CMIP for key variables
needed to capture the glaciated/ice sheet realms of the Greenland and Antarctic ice sheets. Variables requested in this
Opportunity build upon the ISMIP6 data request and will primarily be used to create boundary conditions for a large ensemble
of historical and scenario simulations conducted by standalone Ice Sheet Models (ISMs) coordinated by ISMIP7 (Table 5).
Not only will these variables be used to generate traditional surface mass balance and ice shelf basal melt boundary conditions
335 at the high spatial resolutions required for ice sheet modeling, but they will also be used to allow the influence of changing
climate on calving, ice shelf damage and collapse to be taken into account. Some ESMs in CMIP7 will conduct novel
simulations with interactive ice sheets (i.e., in a coupled setup), and the variables in this Opportunity are also specified to allow
the influence of such coupling on climate evolution and sea-level projections to be assessed. The combined output from
standalone ISM and coupled ESM simulations will allow the ice-sheet contributions to sea level rise implied by the
340 ScenarioMIP protocols to be projected, along with their model and scenario uncertainties. In addition to ISMIP7, other MIPs
concerned with ice sheet behavior, primarily SOFIA (Southern Ocean Freshwater Input from Antarctica; Swart et al., 2023),
TIPMIP (Walton et al., 2025) and MISOMIP (De Rydt et al., 2024), may use these variables.

345 **Table 5. Variable groups needed for ID 39 Ice sheet mass loss, contributions to sea level rise and freshwater flux input
to the climate system.**



Variable group name	Reason for inclusion
landice_global_allfreq	Surface variables from ESMs for creating forcings for standalone ISM simulations using a range of approaches, and for comparing polar climate evolution with coupled ESM-ISM simulations.
landice_greenland_allfreq landice_antarctica_allfreq	Variables from ISMs to diagnose ice sheet evolution and sea level contributions from Greenland and Antarctica, respectively, both for standalone ISM and coupled ESM-ISM simulations.
ocean_grid	Accurate ocean model grid dimensions are required to create some boundary forcings and diagnose sea level contributions.
baseline_monthly baseline_fixed	baseline variables are required to supplement specific requests in landice_global_allfreq.

4.5 Land use change (ID 43)

Human land-use activities have resulted in large changes to the Earth's surface, with resulting implications for climate. In the future, land-use activities are likely to expand and intensify further to meet growing demands for food, fiber, and energy. LUMIP aims to further advance understanding of the impacts of LULCC on climate, specifically addressing the following questions. (1) What are the effects of LULCC on climate and biogeochemical cycling (past–future)? (2) What are the impacts of land management on surface fluxes of carbon, water, and energy, and are there regional land-management strategies that can help mitigate climate change? These questions are particularly relevant in investigations of low emission scenarios, which typically rely on significant land-based mitigation to achieve climate targets.

The unique variable group requested for this Opportunity is the ‘landuse’ variable group (Table 6), which requests a subset of key biogeophysical and biogeochemical variables for five land use tiles (defined below). This variable group addresses the challenges of analyzing the effects of LULCC on physical and biogeochemical states of land and their interactions with the atmosphere. The underlying rationale for this request is that informative sub-grid-scale data that reflects the strong heterogeneity of the land surface, heterogeneity that is significantly driven by human transformation of the land, and which is calculated by land models, is not routinely archived. For CMIP6, LUMIP (Lawrence et al. 2016) requested sub-grid data for the first time within CMIP for key biogeophysical and biogeochemical variables. This subgrid output was utilized in several LUMIP studies (e.g., Tang et al., 2023). Reporting data by land use tile enables assessment of land-use carbon fluxes, including not only analysis of differences in fluxes between land-use and no land-use experiments, but also within a single land-use experiment, utilizing bookkeeping approaches (Houghton et al., 2012) that allow a more direct comparison of observed and modeled carbon inventories.



370 For the Land Use Opportunity, the sub-grid data request is for five land-use categories: (1) primary land (including bare ground and vegetated wetlands), (2) secondary land, (3) cropland, (4) pastureland, and (5) urban. Other sub-grid land surface categories such as lakes, rivers, and glaciers are excluded from this request. This set of land-use sub-grid reporting units closely corresponds to land-use categories that will be in the historical land-use reconstructions and future scenarios provided by the Land Use Harmonization forcing product (LUH3, updated from LUH2 – Hurtt et al., 2020). Primary (i.e., natural vegetation never affected by LULCC activity) and secondary (i.e., natural vegetation that has previously been harvested or abandoned
375 agricultural land with potential to regrow) are separated (they were combined in CMIP6 Data Request) to capture the unique behavior of the next generation of land models that distinguish between these two land types. This variable group is requested for historical and scenario simulations and for any specific Land Use Model Intercomparison Project (LUMIP) simulations.

Table 6. Variable groups needed for ID 43 Land use change.

Variable group name	Reason for inclusion
landuse	Subgrid land-use tile data that is critical for robust assessment of land use change in historical and scenario simulations.

380

4.6 Plant Phenology (ID 53)

Accurately modelling plant phenology seasonality and long-term trends in Earth system models is essential for effectively simulating and understanding terrestrial ecosystem fluxes of carbon, energy, and water with the atmosphere. In this framework, the “plant phenology” Opportunity aims to expand the data available for assessing these relations (Table 7). For this reason, a
385 set of variables is requested at vegetation type level and daily frequency, high enough to capture the observed and expected changes in phenophases timings (e.g. Park et al., 2016). The vegetation type (vegtype) coordinate was already available in CMIP6 and associated with the landCoverFrac variable. In the CMIP6 definition, the vegtype coordinate is a model-dependent dimension. In CMIP7, the vegtype coordinate, instead, requests values for seven specific vegetation categories (namely broadleaf_deciduous_trees, broadleaf_evergreen_trees, needleleaf_deciduous_trees, needleleaf_evergreen_tree,
390 natural_grasses, crops, and shrubs) to enhance the comparability among models and processes understanding (e.g. Li et al., 2024b).

The Plant Phenology Opportunity proposes two variable groups. The high-priority variable group (plant_phenology_high) contains the three principal variables used in evaluating the plant phenology: laiVgt, gppVgt, and LandCoverFrac. LAI is one
395 of the main variables used to estimate the timing of phenophases (e.g. Jolly et al., 2005; Richardson et al., 2012; Anav et al., 2013; Murray-Tortarolo et al., 2013; Peano et al., 2019, 2021, 2025; Song et al., 2021; Li et al., 2022a; Li et al., 2023), while Gross Primary Production (GPP) estimates the exchange of carbon between land and atmosphere and provide another metric to evaluate the relation between vegetation and climate (e.g. Marsh et al., 2025). The landCoverFrac variable, instead, provides the relative influence of each vegetation type on the total grid cell. These primary variables allow us to link the biases in



400 productivity with changes in phenology (e.g. Li et al., 2024a). The secondary variables (plant_phenology_medium variable
group) provide single components of land-atmosphere carbon fluxes at the vegtype level, namely Net Primary Productivity
(nppVgt), Heterotrophic Respiration (rhVgt), and Autotrophic Respiration (raVgt). This allows a deeper investigation of
sources of biases in the land-atmosphere carbon fluxes. This Opportunity applies to historical, land-hist, and scenario
simulations. The historical and land-hist simulations provide a benchmark to evaluate the model plant phenology biases (e.g.
405 Peano et al., 2021, 2025) while scenarios supply future changes.

Table 7. Variable groups needed for ID 53 Plant Phenology.

Variable group name	Reason for inclusion
plant_phenology_high	Subgrid vegetation type-level data, which is essential for assessing plant phenology simulated by models.
plant_phenology_medium	Subgrid vegetation type-level data providing components for assessing carbon fluxes associated with the simulated plant phenology.

4.7 Rapid Evaluation Framework (REF) (ID 55)

410 The CMIP Rapid Evaluation Framework (Hoffman et al., 2025) was created to evaluate and benchmark the newly available
CMIP7 Assessment Fast Track simulations as soon as they are uploaded to ESGF with metrics and diagnostics that are
available through different open-source evaluation and benchmarking tools. This Opportunity contains the set of variables that
are needed for the planned diagnostics and metrics for the REF (CMIP Model Benchmarking Task Team, 2024) (Table 8). The
suggested metrics/diagnostics for the REF to be available for all CMIP7 Assessment Fast Track experiments are in the first
415 instance very basic evaluations and are not expected to require very specific variables. The exact selection of variables was
also made consistent with the model evaluation diagnostics in Chapter 3 of the latest IPCC report (Eyring et al., 2021). Due to
the fixed timeline for the CMIP7 Assessment Fast Track simulations, there is only a short time period for the technical
implementation of the REF, and therefore the available metrics and diagnostics in this first version of the REF will be limited
to a temporal resolution of monthly mean data and about five metrics/diagnostics per realm based on a community selection.
420 The realms were chosen specifically to be consistent with the realms used for the data request. Find more information about
the REF Opportunity in Dingley et al. (2025).

Table 8. Variable groups needed for ID 55 Rapid Evaluation Framework.

Variable group name	Reason for inclusion
ref_land_and_landice	This is the set of variables that would be needed for the planned land and land ice diagnostics and metrics for the Rapid Evaluation Framework. The variable group will be



	linked with the "Rapid Evaluation Framework" Opportunity and is essential for the evaluation of the new CMIP7 Assessment Fast Track simulations on a routine basis.
--	---

425 5 Discussion

5.1 Prioritisation process and updates on CMIP7

The Land and Land Ice Theme author team is comprised of members directly from modelling centers and leads several model intercomparison projects. Therefore, the prioritization process is straightforward as most of the Opportunity proposers had experience of generating a comprehensive variable list within the Land and Land Ice Theme. Specific prioritization processes such as Accurate assessment of land-atmosphere coupling (ID 1) follows a recent peer-reviewed article that highlights the high-frequency land-atmosphere interaction variables (Findell et al., 2024). This Opportunity also contains the low-troposphere atmospheric variables that were reviewed by the Atmosphere Author Team (see more details in Appendix A). Another example that shows the cross-theme collaboration is the Assessments for Hydrological Processes, Water Resources, and Freshwater Systems (ID 38). The Land and Land Ice Theme author team worked with the Impacts & Adaptation team and Earth System team to decide to merge Water Security and Freshwater Ecosystem Services (ID 67) into ID 38 to focus on the water resources. ID 38 is fundamental since this Opportunity contains comprehensive hydrological variables on land. Other Opportunities such as, Glacier changes, drivers, and impacts (ID 35), Ice sheet mass loss, contributions to sea level rise and freshwater flux input to the climate system (ID 39), Land use change (ID 43), and Plant phenology (ID 53), were proposed from either modeling centres or the leadership team from corresponding MIPs and thus were quickly accepted by the cross-theme review.

For CMIP7, these new Opportunities will support a more comprehensive integration of Land and Land Ice Theme components, enable the identification of new feedback loops associated with improved Land and Land Ice processes, and provide critical surface boundary conditions for other Themes included in the CMIP7 Data Request. During the time of CMIP6, processes such as the land ice was firstly introduced to CMIP ESMs (Nowicki et al., 2016), which has been widely applied to studies that focus on important aspects in the Earth climate system such as the change in the Antarctic Ice Sheet (Stokes et al., 2022) and global sea-level rise (Edwards et al., 2021) under future climate scenarios. Similarly, studies that apply land use experiments from the CMIP6-LUMIP experiments (e.g., Li et al., 2022b) also serve as a benchmark to keep improving processes on the land surface, which include energy, water, carbon, and other biogeochemical cycles. In CMIP7 these separate processes will be integrated together and new opportunities such as the TIPMIP will offer chances to identify new feedback loops. For example, the glacier melting-induced changes in the ocean circulation over the Atlantic may generate climate teleconnections on the climate system in the Amazonian rainforest (Van Westen et al., 2024), which could be compared among CMIP7 ESMs by including the full feedback loop of considering the climate warming on ice sheet changes. Studies on these



climate change feedbacks and their interactions with land use changes may become feasible with improved Land and Land Ice processes in CMIP7.

455 **5.2 Key reflections and challenges from the CMIP7 Data Request process**

The CMIP7 Data Request has seen significant improvements over the CMIP6 era, particularly in process organization and community engagement. However, challenges remain regarding the detailed variable requests. For instance, two key Opportunities within the Land and Land Ice Theme—the accurate assessment of land-atmosphere coupling (ID 1) and the evaluation of hydrological processes, water resources, and freshwater systems (ID 38)—require variables with sub-daily
460 resolution. This inherently demands increased data storage capacity, an updated framework for reliable data sharing, and robust data validation before publication. Additionally, the land-use change Opportunity (ID 43) requires sub-grid information, necessitating more sophisticated model outputs. These requirements may discourage modeling groups from contributing to certain Opportunities. However, at this stage in the data request process - before modeling centers determine which variables they will provide for CMIP7 - it remains uncertain how widely these Opportunities will be adopted. A balance must be struck
465 between scientific needs, which should have already been addressed in the public consultation process, and the level of detail that can realistically be achieved. Moving forward, prior to finalizing the list of Opportunities, more intensive direct consultations between Opportunity leads and modeling groups may be necessary to ensure the effective implementation of CMIP7.

470 Another critical consideration is the quality control of requested data, which may need to be specified before finalizing the data request. For example, a recent study on land carbon and nitrogen mass conservation based on CMIP6 data (Tang et al., 2025a) reports significant mass imbalances arising solely from the C4MIP-requested variables. The primary causes include misinterpretation of the data request by ESM teams (e.g., incomplete data submissions) and the lack of a harmonized variable request across different MIP activities, leading to overlapping variables.

475 In the CMIP7 Data Request, issues with variables having similar definitions persist. For example, overlaps exist between variables requested under the land-use change Opportunity (ID 43) and the carbon cycle Opportunity (ID 15), potentially perpetuating the mass conservation issues observed in CMIP6. This problem is now recognized by data users, particularly those in the simple climate model (emulator) community, as emulator development relies heavily on the quality of ESM outputs
480 (Tang et al., 2025b). While some community-led initiatives aim to streamline data reporting from ESM teams to broaden data applications (Tang et al., 2025a), such efforts are not yet integrated into the CMIP7 Data Request process.

The challenge lies in the substantial effort required to harmonize the outputs that ESMs can produce, the variables expected by different MIP Opportunities, and the overlapping variables of interest across these Opportunities. While this may be
485 considered less critical than addressing the scientific questions that are the primary focus of the current data request, data



quality is fundamental to robust analysis. A unified data request would greatly enhance both data reporting and usage in the future, making it an issue worth greater attention moving forward. We understand that a “unified” or “harmonized” data request requires input not only from the CMIP7 Data Request Task Team (for reviewing the opportunities), but also from the MIPs (for updating the variable groups), the modeling centers (regarding which variables can realistically be produced in their models), and the data users (regarding the challenges they face when using the data). Achieving this will require a detailed review of the requested variables and feedback from all these parties. This represents a considerable amount of work and has, so far, been lacking in the CMIP7 Data Request process. However, given its importance to the entire climate modeling and data production workflow, we argue that establishing a robust feedback and refinement loop for variables should be a high priority in the near future, probably from the CMIP level rather than simply from the Data Request Task Team.

495 **5.3 Outstanding gaps in Earth system processes**

Despite advances in ESM simulations and the expanding suite of variables available in CMIP7, several critical gaps remain in the representation of land and land ice processes. One major area requiring further development is plant hydraulics, which governs water movement through the soil-plant-atmosphere continuum. While many land models include simplified representations of soil moisture and transpiration, they often lack explicit simulations of plant stem and leaf water potentials—
500 key physiological metrics that influence drought responses and plant mortality (Xu et al., 2016; Kennedy et al., 2019; Li et al., 2021; Yao et al., 2024). Observational datasets such as the sap flow from the SAPFLUXNET exist to evaluate these processes (Poyatos et al., 2020), but their incorporation into ESM diagnostics remains limited partially due to the scientific challenge to perform comprehensive and rigorous data-model integration. It also depends on whether the significance of these water-related processes is identified for various scientific modeling purpose (e.g., moisture conditions for fuels in fire modeling in ESMs).
505 Improved integration of plant hydraulics into ESMs would enable better assessments of hydrological shifts, aiding in the prediction of drought-induced forest dieback and its cascading effects on carbon, water and energy cycles.

As land models increasingly integrate demographic vegetation capabilities (Fisher et al., 2017; Koven et al., 2020), a more refined set of diagnostics is needed to capture complex vegetation dynamics, including competition, mortality, and recovery processes. However, many land models within ESMs still lack explicit representations of key biophysical mechanisms regulating vegetation structure and function, including hydraulic limits on water transport, physical constraints on tree stability, and temperature-dependent physiological thresholds. Moreover, standardized benchmarks for key biophysical outputs—such as canopy structure, vegetation height, leaf area dynamics, and surface roughness—remain underdeveloped, hindering model validation against remote sensing and field observations. Without these advancements, ESMs may struggle to accurately
515 simulate vegetation-climate feedbacks, particularly under future climate stressors.

Similarly, land ice processes remain underrepresented, particularly in the coupling between dynamic ice sheets and ocean interactions. Current models often exclude or simplify key mechanisms such as ocean cavities below ice shelves, subglacial



freshwater fluxes, and the role of aerosol deposition in modulating ice sheet albedo and melt. The omission of these processes
520 limits the accuracy of sea-level rise projections and ice-ocean feedbacks, which are critical for understanding long-term climate
stability.

Additionally, groundwater systems remain a weakly integrated component in ESMs, despite their significant role in modulating
soil moisture, runoff, and terrestrial water storage. Improved representation of groundwater dynamics is necessary to better
525 capture long-term drought resilience, subsurface hydrological feedbacks, and interactions between deep water storage and
surface climate variability. Similarly, accounting for the terrain characteristics is important to accurately represent terrestrial
processes and quantify energy and water budgets (Clark et al. 2015, Fan et al. 2019). However, ESMs typically run at
resolutions of around 100 kilometers, which prevents the capture of detailed topographic and landscape effects. This results in
large uncertainties in simulating the hydrological and radiative processes and assessing their spatiotemporal variability in
530 mountainous regions and complex terrains that are characterized by prominent topographic variations in surface elevation,
steepness (slope), and the orientation of the land surface (aspect).

Beyond the traditional land, hydrology, and cryosphere components, solid Earth processes are another frontier in Earth system
modeling that require greater attention. Processes such as isostatic adjustments (Whitehouse 2018), geothermal heat fluxes
535 (Lösing et al., 2020), and tectonic activity influence surface deformation, hydrological pathways (Michalek et al., 2023), and
even cryosphere stability (Albrecht et al., 2024), yet they remain largely excluded from ESM frameworks. The potential climate
implications of these interactions—including their role in modulating geohazards, altering water systems, and influencing
permafrost and ice sheet dynamics—underscore the need for a more integrated Earth system perspective. As CMIP7 advances,
addressing these gaps will be essential to improving climate projections, refining impact assessments, and reducing
540 uncertainties in future Earth system responses.

6 Conclusions

In this paper, we present the progress and evolution of the CMIP7 Data Request concerning land and land ice processes. The
updated request includes six refined Opportunities addressing distinct scientific questions. These Opportunities—focusing on
land-atmosphere coupling, glacier changes, freshwater processes, ice sheet loss and sea level rise, land use change, and plant
545 phenology—reflect the urgent needs of various research communities. This work serves as a valuable resource for modeling
groups, guiding potential modifications or enhancements to their output to better align with evolving scientific priorities.

While the scientific significance of these data request Opportunities is fundamental, effective variable management (e.g., e.g.,
removal of duplicates, harmonization of similar variables through improved definitions), robust data validation (e.g., mass
550 conservation check), and efficient data distribution are equally critical. The primary goal of the data request, and of modeling



groups producing data, is to facilitate downstream analysis, which necessitates consistency between requested variables and model outputs. However, CMIP6 data validation at the user end has demonstrated the challenges in ensuring this consistency.

555 Looking ahead, greater involvement of modeling groups—who best understand the capabilities of their models—should be prioritized during the data request design phase. Additionally, a comprehensive review of both existing and newly requested variables should be conducted before finalizing the next-generation data request. This review should extend beyond individual themes or Opportunities to encompass the entire variable space, considering the complex interactions within the climate system. Furthermore, community tools such as ESMValTool should be developed or updated to support the harmonization of variables within the whole Earth system and improve data consistency.

560



Appendix A: Opportunity processing

Following the first consultation phase, the submitted Opportunities related to the Land and Land Ice Theme were reviewed during author team meetings and assessed for their scientific value, clarity and selection of variables. If deemed appropriate, Opportunities were accepted, otherwise they were rejected or possibly merged with other Opportunities. Potential gaps and the creation of variable groups were identified in author team meetings, discussed with the proposer and relevant communities and brought to the attention of cross thematic theme coordinators if Opportunities were relevant to other themes and required their input. Comments from other themes or the public consultation were addressed and following the cross-thematic harmonization sprint some Opportunities with overlapping interests were merged.

The Opportunities led by the Land and Land Ice Theme as well as key decisions for each Opportunity are listed in the following table.

Action taken	Opportunity title	Meeting decision made	Notes from consultation and cross thematic	Notes from Author team
Accepted				
ID 1	Accurate assessment of land-atmosphere coupling	Author team meeting 25/10/2024	Cross-thematic group review requested additional description and impacts information. The atmosphere review confirmed that the Opportunity is appropriate and that high-frequency data is required.	Cross thematic and atmosphere team review points were addressed - more details were added to the description, expected impacts were explained and the variable groups were reviewed and soil moisture at different levels was added. Review of atmosphere variables was also conducted.
ID 35	Glacier changes, drivers, and impacts	Author team meeting 25/10/2024	The cross-thematic review suggested investigating the need for a more specific variable group. It was also suggested to include	The landice_glaciers_allfreq variable group was added in response to review. Discussion of potential merger and decision to



			mountain glaciers in the title and the possibility of merging with Hydrology and Fresh Water Systems (ID 38), Water Resources (ID 67) and Water Cycle (ID 66), which could be consolidated into two Opportunities.	retain a distinct glacier Opportunity and to re-name (from original Glacier geometry changes and glacio-hydrology).
ID 39	Ice sheet mass loss, contributions to sea level rise and freshwater flux input to the climate system	Author team meeting 25/10/2024	No comments from cross thematic or consultation review.	The Opportunity was clearly justified. Only a piControl timeslice was to be added.
ID 43	Land use change	Author team meeting 03/01/2025	No comments from cross thematic or consultation review.	The variable definitions for variables on land use tiles were modified.
ID 53	Plant phenology	Author team meeting 25/10/2024	The cross-thematic review accepted the scientific proposition, but noted that variables were not provided and that daily outputs are a high demand, although need was recognised.	An issue with the missing Vegetation_type variable group was resolved.
ID 38	Assessments for Hydrological Processes, Water Resources, and Freshwater Systems	Author team meeting 25/10/2024	The cross-thematic review suggested consolidating this Opportunity (previously Hydrology and Freshwater Systems), the	After consideration of potential for proposed merges, author team decided, together with input from Impacts & Adaptation team that ID 67



			<p>Water Resources (ID 67) and Water Cycle (ID 66) into two Opportunities, where one is focused on the water cycle and the other on land/water resources. The proposers of all three Opportunities were requested to make the necessary edits and a merge with ID 67 and Climate impact assessments on freshwater ecosystems (ID 15) was requested.</p>	<p>(see below) be merged while ID 66, in discussion with and led by the Earth System team, should remain standalone as it is a fundamental component together with energy and carbon cycles. A soil temperature profile was also identified to be included for permafrost.</p>
MERGED				
67	Water Security and Freshwater Ecosystem Services	Author team meeting 25/10/2024	<p>Cross thematic review suggested this Opportunity, Hydrology and Fresh Water Systems (ID 38) and Water cycle (ID 66) should be consolidated into two Opportunities with suggestion is one focused on water cycle, the other focused on land/water resources.</p>	<p>After considerable discussion with the Impacts and Adaptation team this Opportunity was merged into ID 38 (see above).</p>



575 Appendix B: Variable description

Variables that are newly introduced in CMIP7 related to Land and Land Ice Theme Opportunities are listed below. Details about the coordinate specifications regarding the time and spatial requirements for the variable’s outputs can be seen in the release v1.2 of the CMIP7 Data Request (<https://wcrp-cmip.org/cmip7-data-request-v1-2/>)

New physical parameter	CF standard name	Description	Further detail to aid compute	Coordinate specifications
3hr.mrso10	mass_content_of_water_in_soil_layer	Soil moisture at 3hr but for 0-1m	The variable is a high-frequency output at 3 hour interval and should be identical to 0-1m soil moisture.	sdepth10, 3hr
3hr.srfrad	surface_net_downward_radiative_flux	Net radiative flux at surface	The variable is a high-frequency output at 3 hour interval and should be identical to the net balance of surface downward and upward radiation for shortwave and longwave radiation.	3hr
E3hrPt.hus6	specific_humidity	Specific humidity on 6 pressure levels in the lower troposphere	This should be identical to variable (hus) but with specified vertical pressure levels from 950 hPa to 700 hPa with an interval of 50h Pa for vertical layers (6 vertical layers)	Global field (6 pressure levels), plev6, 3hr
E3hrPt.ta6	air_temperature	Air temperature on 6 pressure levels in the lower troposphere	This should be identical to variable (ta) but with specified vertical pressure levels from 950 hPa to 700 hPa with an interval of 50h Pa for vertical layers (6 vertical layers)	Global field (6 pressure levels), plev6, 3hr



E3hrPt.ua6	eastward_wind	Zonal wind (positive in a eastward direction) on 6 pressure levels in the lower troposphere	This should be identical to variable (ua) but with specified vertical pressure levels from 950 hPa to 700 hPa with an interval of 50h Pa for vertical layers (6 vertical layers)	Global field (6 pressure levels), plev6, 3hr
E3hrPt.va6	northward_wind	Meridional wind (positive in a northward direction) on 6 pressure levels in the lower troposphere	This should be identical to variable (va) but with specified vertical pressure levels from 950 hPa to 700 hPa with an interval of 50h Pa for vertical layers (6 vertical layers)	Global field (6 pressure levels), plev6, 3hr
E3hrPt.wap6	lagrangian_tendency_of_air_pressure	Omega ($=dp/dt$) on 6 pressure levels in the lower troposphere	This should be identical to variable (wap) but with specified vertical pressure levels from 950 hPa to 700 hPa with an interval of 50h Pa for vertical layers (6 vertical layers)	Global field (6 pressure levels), plev6, 3hr
Eday.gppVgt	gross_primary_productivity_of_biomass_expressed_as_carbon	The rate of synthesis of biomass from inorganic precursors by autotrophs ("producers") expressed as the mass of carbon which it contains. For example, photosynthesis in plants or phytoplankton. The producers also respire some of this biomass and the difference is referred	The variable is aggregated from the subgrid land structure into vegetation type level based on the categories: broadleaf_deciduous_trees, broadleaf_evergreen_trees, needleleaf_deciduous_trees, needleleaf_evergreen_tree, natural_grasses, crops, shrubs.	Global (single level) with extra dimension vegtype



		to as the net primary production. Reported on land-use tiles.		
Eday.laiVgt	leaf_area_index	A ratio obtained by dividing the total upper leaf surface area of vegetation by the (horizontal) surface area of the land on which it grows.	The variable is aggregated from the subgrid land structure into vegetation type level based on the categories: broadleaf_deciduous_trees, broadleaf_evergreen_trees, needleleaf_deciduous_trees, needleleaf_evergreen_tree, natural_grasses, crops, shrubs.	Global (single level) with extra dimension vegtype
Eday.nppVgt	net_primary_productivity_of_biomass_expressed_as_carbon	"Production of carbon" means the production of biomass expressed as the mass of carbon which it contains. Net primary production is the excess of gross primary production (rate of synthesis of biomass from inorganic precursors) by autotrophs ("producers"), for example, photosynthesis in plants or phytoplankton, over the rate at which the autotrophs themselves respire some of this	The variable is aggregated from the subgrid land structure into vegetation type level based on the categories: broadleaf_deciduous_trees, broadleaf_evergreen_trees, needleleaf_deciduous_trees, needleleaf_evergreen_tree, natural_grasses, crops, shrubs.	Global (single level) with extra dimension vegtype



		<p>biomass. "Productivity" means production per unit area. The phrase "expressed_as" is used in the construction $A_{\text{expressed_as}} B$, where B is a chemical constituent of A. It means that the quantity indicated by the standard name is calculated solely with respect to the B contained in A, neglecting all other chemical constituents of A.</p>		
Eday.raVgt	surface_upward_mass_flux_of_carbon_dioxide_expressed_as_carbon_due_to_plant_respiration	Carbon mass flux per unit area into atmosphere due to autotrophic respiration on land (respiration by producers) [see rh for heterotrophic production]. Calculated on vegetation type.	The variable is aggregated from the subgrid land structure into vegetation type level based on the categories: broadleaf_deciduous_trees, broadleaf_evergreen_trees, needleleaf_deciduous_trees, needleleaf_evergreen_tree, natural_grasses, crops, shrubs.	Global (single level) with extra dimension vegtype
Eday.rhVgt	surface_upward_mass_flux_of_carbon_dioxide_expressed_as_carbon_due_to_heterotrophic_respiration	Carbon mass flux per unit area into atmosphere due to heterotrophic respiration on land (respiration by consumers), calculated on vegetation type.	The variable is aggregated from the subgrid land structure into vegetation type level based on the categories: broadleaf_deciduous_trees, broadleaf_evergreen_trees, needleleaf_deciduous_trees,	Global (single level) with extra dimension vegtype



			needleleaf_evergreen_tree, natural_grasses, crops, shrubs.	
Emon.albsrfc	surface_albedo	Albedo of the surface	This is a global single level field representing the grid cell averaged albedo (averaged over all subgrid-scale structures and all wavebands)	longitude latitude time
fx.depthl	depth	Depth of lakes.	If this quantity is present in the model. If computed via volume and area, then this is lake volume divided by lake area	fx
fx.sfilkf	area_fraction	Fraction of horizontal land grid cell area occupied by lake.		typelkins, fx
Ofx.dxt0		The linear extent of the cell in the x direction of the horizontal grid centered at t-points (points for tracers such as temperature, salinity, etc.).	Not applicable to unstructured grids.	fx
Ofx.dxu0		The linear extent of the cell in the x direction of the horizontal grid centered at u-points (points for velocity in the x-direction).	Not applicable to unstructured grids.	fx
Ofx.dxv0		The linear extent of the cell in the x direction of the horizontal grid centered at v-points (points for velocity in the y-direction).	Not applicable to unstructured grids.	fx
Ofx.dy0		The linear extent of the cell in the y direction of the horizontal grid centered at t-points (points for tracers such as temperature, salinity, etc.).	Not applicable to unstructured grids.	fx
Ofx.dyu0		The linear extent of the cell in the y direction of the horizontal grid centered at u-points (points for velocity in the x-direction).	Not applicable to unstructured grids.	fx



Ofx.dyvo		The linear extent of the cell in the y direction of the horizontal grid centered at v-points (points for velocity in the y-direction).	Not applicable to unstructured grids.	fx
Omon.thkcelluo	cell_thickness	The time varying thickness of ocean cells centered at u-points (points for velocity in the x-direction).	"Thickness" means the vertical extent of a layer. "Cell" refers to a model grid-cell.	mon
Omon.thkcellvo	cell_thickness	The time varying thickness of ocean cells centered at v-points (points for velocity in the y-direction).	"Thickness" means the vertical extent of a layer. "Cell" refers to a model grid-cell.	mon

580 Code and data availability

The variables and their metadata included latest CMIP7 Assessment Fast Track Data Request can be accessed at <https://doi.org/10.5281/zenodo.14774070>. At the time of this publication, the latest major release is v1.2 (Data Request Task Team, 2025b; accessed at <https://doi.org/10.5281/zenodo.15116894>), and the latest minor release is v1.2.1 (Data Request Task Team, 2025c; accessed at <https://doi.org/10.5281/zenodo.15288187>). Previous versions include v1.0beta (Data Request Task Team, 2024a; accessed at <https://doi.org/10.5281/zenodo.14832540>), v1.0 (Data Request Task Team, 2024b; accessed at <https://doi.org/10.5281/zenodo.14832541>), and v1.1 (Data Request Task Team, 2025a; accessed at <https://doi.org/10.5281/zenodo.14774071>).

Author contribution

YL and GT led the conceptualization, investigation, methodology, supervision and visualization with data curation, methodology and visualization contributions from SM, SN, RS, DL, DP and EOR. YL and GT led the writing of this manuscript with contributions from all authors. EOR provided resources and project administration support.

Competing interests

DML is on the editorial board of *Geoscientific Model Development*. The authors declare that they have no other conflict of interest.

595 Acknowledgements

The Land and Land Ice Theme author team acknowledge the valuable contributions from a widespread scientific community and public who participated in the CMIP7 Data Request effort and public consulting processes. We thank the Land and Land



Ice Theme Committee members, including Fang Li, Ben Marzeion, and Manmeet Singh. We also thank James Anstey, Beth Dingley, Bo Huang, Sifan Koriche, and Alex Ruane for their helpful comments on the draft.

600 **Financial support**

YL acknowledges support from the UCLA Society of Hellman Fellows award. GT acknowledges support from the Guest Program at the Max Planck Institute for Biogeochemistry (MPI-BGC), as well as the Science Abroad Travelling Scholarships and the School of Geography, Earth and Atmospheric Sciences Graduate Researcher Travelling Funds from The University of Melbourne. The CMIP IPO (EOR) is hosted by the European Space Agency, with staff provided on contract by HE Space
605 Operations Ltd. SM and DML are supported by the NSF National Center for Atmospheric Research (NCAR), which is a major facility sponsored by the U.S. National Science Foundation under Cooperative Agreement No. 1852977. MMB has been supported by the British Antarctic Survey, Natural Environment Research Council (NERC), Polar Science for a Sustainable Planet programme. SN was supported by the NASA Sea Level Change Team and MAP programs. RS was supported by the UK Natural Environment Research Council through the TerraFIRMA: Future Impacts, Risks and Mitigation Actions in a
610 changing Earth System project, Grant reference NE/W004895/1. The work of BH was supported by the European Union's Horizon 2020 research and innovation program (grant no. 101003536; ESM2025–Earth System Models for the Future). JM was supported by the Reducing Uncertainties in Biogeochemical Interactions through Synthesis and Computing Scientific Focus Area (RUBISCO SFA) project funded through the Earth and Environmental Systems Sciences Division of the Biological and Environmental Research Office in the US Department of Energy (DOE) Office of Science. Oak Ridge National Laboratory
615 (ORNL) is supported by the Office of Science of the DOE under Contract No. DE-AC05-00OR22725.



References

- Abrams, J.F., S.J. Lade, J.F. Donges, J.E. Buxton (eds): ‘Section 2: Tipping Points Impacts’ in [T.M. Lenton, D.I. Armstrong McKay, S. Loriani, J.F. Abrams, S.J. Lade, J.F. Donges, M. Milkoreit, T. Powell, S.R. Smith, C. Zimm, J.E. Buxton, L. Laybourn, A. Ghadiali, J. G. Dyke (eds), *The Global Tipping Points Report 2023*. University of Exeter, Exeter, UK.], 2023.
- Albrecht, T., Bagge, M., and Klemann, V.: Feedback mechanisms controlling Antarctic glacial-cycle dynamics simulated with a coupled ice sheet–solid Earth model, *The Cryosphere*, 18, 4233–4255, <https://doi.org/10.5194/tc-18-4233-2024>, 2024.
- Al-Yaari, A., Ducharne, A., Thiery, W., Cheruy, F., and Lawrence, D.: The role of irrigation expansion on historical climate change: Insights from CMIP6. *Earth's Future*, 10, e2022EF002859. <https://doi.org/10.1029/2022EF002859>, 2022
- Anav, A., Friedlingstein, P., Kidston, M., Bopp, L., Ciais, P., Cox, P., Jones, C., Jung, M., Myneni, R., and Zhu, Z.: Evaluating the land and ocean components of the global carbon cycle in the CMIP5 Earth System models, *Journal of Climate*, 26, 6801–6843, <https://doi.org/10.1175/jcli-d-12-00417.1>, 2013.
- Armstrong McKay, D.I., Staal, A., Abrams, J.F., Winkelmann, R., Sakschewski, B., Loriani, S., Fetzer, I., Cornell, S.E., Rockström, J. and Lenton, T.M.: Exceeding 1.5 C global warming could trigger multiple climate tipping points. *Science*, 377(6611), p.eabn7950, <https://doi.org/10.1126/science.abn7950>, 2022.
- Balch, J.K., Abatzoglou, J.T., Joseph, M.B. et al: Warming weakens the night-time barrier to global fire, *Nature*, 602, 442–448 (2022). <https://doi.org/10.1038/s41586-021-04325-1>, 2022.
- Biess, B., Gudmundsson, L. and Seneviratne, S.I.: Assessing global and regional trends in spatially co-occurring hot or wet annual maxima under climate change. *Earth's Future*, 12(4), p.e2023EF004114. <https://doi.org/10.1029/2023EF004114>, 2024.
- Boysen, L. R., Brovkin, V., Pongratz, J., Lawrence, D. M., Lawrence, P., Vuichard, N., Peylin, P., Liddicoat, S., Hajima, T., Zhang, Y., Rocher, M., Delire, C., Séférian, R., Arora, V. K., Nieradzik, L., Anthoni, P., Thiery, W., Laguë, M. M., Lawrence, D., and Lo, M.-H.: Global climate response to idealized deforestation in CMIP6 models, *Biogeosciences*, 17, 5615–5638, <https://doi.org/10.5194/bg-17-5615-2020>, 2020.
- Briley, L., Rood, R., Notaro, M.: Large lakes in climate models: A Great Lakes case study on the usability of CMIP5. *Journal of Great Lakes Research*. 47. <https://doi.org/10.1016/j.jglr.2021.01.010>, 2021.



650 Burke, E. J., Zhang, Y., and Krinner, G.: Evaluating permafrost physics in the Coupled Model Intercomparison Project 6
(CMIP6) models and their sensitivity to climate change, *The Cryosphere*, 14, 3155–3174, [https://doi.org/10.5194/tc-14-3155-](https://doi.org/10.5194/tc-14-3155-2020)
2020, 2020.

Chauhan, T., Devanand, A., Roxy, M.K. et al. River interlinking alters land-atmosphere feedback and changes the Indian
655 summer monsoon. *Nat Commun* 14, 5928, <https://doi.org/10.1038/s41467-023-41668-x> , 2023

Chen, X., Wang, J., Pan, F., Song, Y., Liang, J., Huang, N., Jiang, K., Gao, R., Men, J., Bi, P., Zhang, F., Huang, Z., Huang,
B., and Pan, Z.: Land-atmosphere feedback exacerbated the mega heatwave and drought over the Yangtze River Basin of
China during summer 2022. *Agricultural and Forest Meteorology*, <https://doi.org/10.1016/j.agrformet.2024.110321> , 2025

660

Clark, M. P., Fan, Y., Lawrence, D. M., Adam, J. C., Bolster, D., Gochis, D. J., Hooper, R. P., Kumar, M., Leung, L. R.,
Mackay, D. S., Maxwell, R. M., Shen, C., Swenson, S. C., and Zeng, X.: Improving the representation of hydrologic processes
in Earth System Models, *Water Resources Research*, 51, 5929–5956, <https://doi.org/10.1002/2015wr017096>, 2015.

665 CMIP Model Benchmarking Task Team.: CMIP AR7 Fast Track Diagnostics list for the Rapid Evaluation Framework [Data
set]. Zenodo. <https://doi.org/10.5281/zenodo.14284375>, 2024

Cook, B.I., Mankin, J.S., Marvel, K., Williams, A.P., Smerdon, J.E., and Anchukaiti, K.J.: Twenty-first Century drought
projections in the CMIP6 forcing scenarios. *Earth's Future*, 8, no. 6, e2019EF001461, doi:10.1029/2019EF001461, 2020

670

Davies, B., McNabb, R., Bendle, J. et al. Accelerating glacier volume loss on Juneau Icefield driven by hypsometry and melt-
accelerating feedbacks. *Nat Commun* 15, 5099, <https://doi.org/10.1038/s41467-024-49269-y>, 2024.

Data Request Task Team. (2024a). CMIP-Data-Request/CMIP7_DReq_Content: Data request content for v1.0beta (released
675 21 Oct 2024) (v1.0beta). Zenodo. <https://doi.org/10.5281/zenodo.14832540>

Data Request Task Team. (2024b). CMIP-Data-Request/CMIP7_DReq_Content: Data request content for v1.0 (released 22
Nov 2024) (v1.0). Zenodo. <https://doi.org/10.5281/zenodo.14832541>

680 Data Request Task Team. (2025a). CMIP-Data-Request/CMIP7_DReq_Content: Data request content for v1.1 (released 30
Jan 2025) (v1.1). Zenodo. <https://doi.org/10.5281/zenodo.14774071>



Data Request Task Team (2025b). CMIP-Data-Request/CMIP7_DReq_Content: Data request content for v1.2 (released 1 Apr 2025) (v1.2). Zenodo. <https://doi.org/10.5281/zenodo.15116894>

685

Data Request Task Team (2025c). CMIP-Data-Request/CMIP7_DReq_Content: Data request content for v1.2.1 (released 26 Apr 2025) (v1.2.1). Zenodo. <https://doi.org/10.5281/zenodo.15288187>

De Rydt, J., Jourdain, N. C., Nakayama, Y., van Caspel, M., Timmermann, R., Mathiot, P., Asay-Davis, X. S., Seroussi, H.,
690 Dutrieux, P., Galton-Fenzi, B., Holland, D., and Reese, R.: Experimental design for the Marine Ice Sheet–Ocean Model
Intercomparison Project – phase 2 (MISOMIP2), *Geosci. Model Dev.*, 17, 7105–7139, [https://doi.org/10.5194/gmd-17-7105-](https://doi.org/10.5194/gmd-17-7105-2024)
2024, 2024.

Dingley, B., Anstey, J. A., Abalos, M., Abraham, C., Bergman, T., Bock, L., Hassler, L., Kramer, R. J., Luo, F., O’Connor, F.,
695 M., Šácha, P., Simpson, I. R., Wilcox, L.J., Zelinka, M. D.: Atmosphere Theme Data Request for CMIP7, 2025, in preparation

Dirmeyer, P., Balsamo, G., Blyth, E., Morrison, R., and Cooper, H.: Land-Atmosphere Interactions Exacerbated the Drought
and Heatwave Over Northern Europe During Summer 2018. *AGU Advances*. 2. 10.1029/2020AV000283, 2021.

700 Drenkhan, F., Buytaert, W., Mackay, J.D. et al.: Looking beyond glaciers to understand mountain water security. *Nat Sustain* 6,
130–138, <https://doi.org/10.1038/s41893-022-00996-4>, 2023

Dong W, Krasting JP, Guo H.: Analysis of precipitation diurnal cycle and variance in multiple observations, CMIP6 models,
and a series of GFDL-AM4. 0 simulations, *Journal of Climate*, Dec 15;36(24):8637-55, [https://doi.org/10.1175/JCLI-D-23-](https://doi.org/10.1175/JCLI-D-23-0268.1)
705 0268.1, 2023.

Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E.,
Smith, C. J., McKenna, C. M., Simon, E., Abe-Ouchi, A., Gregory, J. M., Larour, E., Lipscomb, W. H., Payne, A. J., Shepherd,
A., Agosta, C., Alexander, P., Albrecht, T., Anderson, B., Asay-Davis, X., Aschwanden, A., Barthel, A., Bliss, A., Calov, R.,
710 Chambers, C., Champollion, N., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Felikson, D., Fettweis, X., Fujita, K., Galton-
Fenzi, B. K., Gladstone, R., Golledge, N. R., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huss, M., Huybrechts,
P., Immerzeel, W., Kleiner, T., Kraaijenbrink, P., Clec’h, S. L., Lee, V., Leguy, G. R., Little, C. M., Lowry, D. P., Malles, J.-
H., Martin, D. F., Maussion, F., Morlighem, M., O’Neill, J. F., Nias, I., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Radić,
V., Reese, R., Rounce, D. R., Rückamp, M., Sakai, A., Shafer, C., Schlegel, N.-J., Shannon, S., Smith, R. S., Straneo, F., Sun,
715 S., Tarasov, L., Trusel, L. D., Van Breedam, J., Van De Wal, R., Van Den Broeke, M., Winkelmann, R., Zekollari, H., Zhao,



- C., Zhang, T., and Zwinger, T.: Projected land ice contributions to twenty-first-century sea level rise, *Nature*, 593, 74–82, <https://doi.org/10.1038/s41586-021-03302-y>, 2021.
- Eyring, V., Bony, S., Meehl, G.A., Senior, C.A., Stevens, B., Stouffer, R.J. and Taylor, K.E., Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*, 9(5), 1937-1958., 2016
- Eyring, V., N.P. Gillett, K.M. Achuta Rao, R. Barimalala, M. Barreiro Parrillo, N. Bellouin, C. Cassou, P.J. Durack, Y. Kosaka, S. McGregor, S. Min, O. Morgenstern, and Y. Sun.: Human Influence on the Climate System. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 423–552, doi: 10.1017/9781009157896.005, 2021.
- Fan, Y., Clark, M., Lawrence, D. M., Swenson, S., Band, L. E., Brantley, S. L., et al.: Hillslope hydrology in global change research and Earth system modeling. *Water Resources Research*, 55, 1737–1772. <https://doi.org/10.1029/2018WR023903>, 2019.
- Findell, K., Gentine, P., Lintner, B. and Guillod, B.: Data Length Requirements for Observational Estimates of Land-Atmosphere Coupling Strength. *Journal of Hydrometeorology*. 16. 150406064716003. 10.1175/JHM-D-14-0131.1., 2015
- Findell, K. L., Gentine, P., Lintner, B. R., and Kerr, C.: Probability of afternoon precipitation in eastern United States and Mexico enhanced by high evaporation, *Nat. Geosci.*, 4, 434–439, 2011.
- Findell, K. L., Yin, Z., Seo, E., Dirmeyer, P. A., Arnold, N. P., Chaney, N., Fowler, M. D., Huang, M., Lawrence, D. M., Ma, P.-L., and Santanello Jr., J. A.: Accurate assessment of land–atmosphere coupling in climate models requires high-frequency data output, *Geosci. Model Dev.*, 17, 1869–1883, <https://doi.org/10.5194/gmd-17-1869-2024>, 2024.
- Fisher RA, Koven CD, Anderegg WRL, et al.: Vegetation demographics in Earth System Models: A review of progress and priorities. *Glob Change Biol.* ,24, 35–54. <https://doi.org/10.1111/gcb.13910>, 2018.
- Fox-Kemper, B., H.T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S.S. Drijfhout, T.L. Edwards, N.R. Golledge, M. Hemer, R.E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I.S. Nurhati, L. Ruiz, J.-B. Sallée, A.B.A. Slangen, and Y. Yu.: Ocean,



- 750 Cryosphere and Sea Level Change. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1211–1362, doi: 10.1017/9781009157896.011, 2021.
- 755 Golledge, N.R., Keller, E.D., Gomez, N. et al.: Global environmental consequences of twenty-first-century ice-sheet melt. *Nature* 566, 65–72, <https://doi.org/10.1038/s41586-019-0889-9>, 2019.
- Gregory, J. M., White, N. J., Church, J. A., Bierkens, M. F. P., Box, J. E., van den Broeke, M. R., Cogley, J. G., Fettweis, X.,
760 Hanna, E., Huybrechts, P., Konikow, L. F., Leclercq, P. W., Marzeion, B., Oerlemans, J., Tamisiea, M. E., Wada, Y., Wake, L. M., and van de Wal, R. S. W.: Twentieth-Century Global-Mean Sea Level Rise: Is the Whole Greater than the Sum of the Parts?. *Journal of Climate*, 26(13), 4476–4499. <https://doi.org/10.1175/JCLI-D-12-00319.1>, 2013.
- Hassell, D., Gregory, J., Blower, J., Lawrence, B.N. and Taylor, K.E.: A data model of the Climate and Forecast metadata
765 conventions (CF-1.6) with a software implementation (cf-python v2. 1). *Geosci. Model Dev.*, 10, 4619–4646, <https://doi.org/10.5194/gmd-10-4619-2017>, 2017.
- Herrera-Estrada, J. E., Martinez, J. A., Dominguez, F., Findell, K. L., Wood, E. F., and Sheffield, J.: Reduced Moisture
770 Transport Linked to Drought Propagation Across North America, *Geophysical Research Letters*, vol. 46, no. 10, pp. 5243–5253, doi:10.1029/2019GL082475, 2019.
- Hock, R., Bliss, A., Marzeion, B., Giesen, R.H., Hirabayashi, Y., Huss, M., Radić, V. and Slangen, A.B. :GlacierMIP–A model
intercomparison of global-scale glacier mass-balance models and projections. *Journal of Glaciology*, 65(251), pp.453–467.
453–467. doi:10.1017/jog.2019.22, 2019a.
- 775 Hock, R., G. Rasul, C. Adler, B. Cáceres, S. Gruber, Y. Hirabayashi, M. Jackson, A. Kääh, S. Kang, S. Kutuzov, Al. Milner, U. Molau, S. Morin, B. Orlove, and H. Steltzer: High Mountain Areas. In: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate* [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegria, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]. Cambridge University Press, Cambridge, UK and
780 New York, NY, USA, pp. 131–202. <https://doi.org/10.1017/9781009157964.004>, 2019b.
- Hock, R., and Truffer, M.: Glacier hazards: Will they change in the future? *Arctic, Antarctic, and Alpine Research*, 56(1). <https://doi.org/10.1080/15230430.2024.233599>, 2024.



785 Hoffman, F., Hassler, B., Swaminathan, R., Lewis, J., Andela, B., Collier, N., Hegedús, D., Lee, J., Pascoe, C., Pflüger, M.,
Stockhouse, M., Ullrich, P., Xu, M., Bock, L., Chun, L., Gier, B. K., Kelley, D. I., Lauer, A., Lenhardt, J., Schlund, M.,
Sreeush, M. G., Weigel, K., Blockley, E., Beadling, R., Beucher, R., Dugassa, D. D., Lembo, V., Lu, J., Brands, S., Tjiputra,
J., Malinina, E., Mederios, B., Soccimarro, E., Walton, J., Kershaw, P., Marquez, A. L., Roberts, M. J., O'Rourke, E., Dingley,
B., Turner, B., Hewitt, H., and Dunne, J. P.: Rapid Evaluation Framework for the CMIP7 Assessment Fast Track [submitted
790 to GMD]

Hogan, J.; Lichstein, J.; Helmer, E.; Craig, M.; Fricke, E.; Henrich, V., et al.: Anthromes and forest carbon responses to global
change, *Plants People Planet*, 7, 1027–1042, <https://doi.org/10.1002/ppp3.10609>, 2025.

795 Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., Le Quéré, C., and Ramankutty,
N.: Carbon emissions from land use and land-cover change, *Biogeosciences*, 9, 5125–5142, [https://doi.org/10.5194/bg-9-5125-](https://doi.org/10.5194/bg-9-5125-2012)
2012, 2012.

Huntington, T. G.: Evidence for intensification of the global water cycle: Review and synthesis, *Journal of Hydrology*, 319,
800 83–95, <https://doi.org/10.1016/j.jhydrol.2005.07.003>, 2005.

Hurtt, G. C., Chini, L., Sahajpal, R., Frohking, S., Bodirsky, B. L., Calvin, K., Doelman, J. C., Fisk, J., Fujimori, S., Klein
Goldewijk, K., Hasegawa, T., Havlik, P., Heinemann, A., Humpenöder, F., Jungclaus, J., Kaplan, J. O., Kennedy, J., Krisztin,
T., Lawrence, D., Lawrence, P., Ma, L., Mertz, O., Pongratz, J., Popp, A., Poulter, B., Riahi, K., Shevliakova, E., Stehfest, E.,
805 Thornton, P., Tubiello, F. N., van Vuuren, D. P., and Zhang, X.: Harmonization of global land use change and management
for the period 850–2100 (LUH2) for CMIP6, *Geosci. Model Dev.*, 13, 5425–5464, <https://doi.org/10.5194/gmd-13-5425-2020>,
2020.

Huss, M., Hock, R. Global-scale hydrological response to future glacier mass loss. *Nature Clim Change* 8, 135–140,
810 <https://doi.org/10.1038/s41558-017-0049-x>, 2018.

IPCC, 2019: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.-O. Pörtner, D.C. Roberts, V.
Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama,
N.M. Weyer (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, 755 pp.
815 <https://doi.org/10.1017/9781009157964>, 2019.



- Jolly, W. M., Nemani, R., and Running, S. W.: A generalized, bioclimatic index to predict foliar phenology in response to climate, *Global Change Biology*, 11, 619–632, <https://doi.org/10.1111/j.1365-2486.2005.00930.x>, 2005.
- 820 Juckes, M., Taylor, K. E., Antonio, F., Brayshaw, D., Buontempo, C., Cao, J., Durack, P. J., Kawamiya, M., Kim, H., Lovato, T., Mackallah, C., Mizielinski, M., Nuzzo, A., Stockhause, M., Visioni, D., Walton, J., Turner, B., O'Rourke, E., and Dingley, B.: Baseline Climate Variables for Earth System Modelling, *Geosci. Model Dev.*, 18, 2639–2663, <https://doi.org/10.5194/gmd-18-2639-2025>, 2025.
- 825 Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclaus, J. H., Otto-Bliesner, B. L., Peterschmitt, J.-Y., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Brierley, C., Crucifix, M., Dolan, A., Fernandez-Donado, L., Fischer, H., Hoperft, P. O., Ivanovic, R. F., Lambert, F., Lunt, D. J., Mahowald, N. M., Peltier, W. R., Phipps, S. J., Roche, D. M., Schmidt, G. A., Tarasov, L., Valdes, P. J., Zhang, Q., and Zhou, T.: The PMIP4 contribution to CMIP6 – Part 1: Overview and over-arching analysis plan, *Geosci. Model Dev.*, 11, 1033–1057, <https://doi.org/10.5194/gmd-11-1033-2018>, 2018.
- 830 Keenan, T. F., Gray, J., Friedl, M. A., Toomey, M., Bohrer, G., Hollinger, D. Y., Munger, J. W., O'Keefe, J., Schmid, H. P., Wing, I. S., Yang, B., and Richardson, A. D.: Net carbon uptake has increased through warming-induced changes in temperate forest phenology, *Nat. Clim. Change*, 4, 598–604, <https://doi.org/10.1038/NCLIMATE2253>, 2014.
- 835 Kennedy, D., Swenson, S., Oleson, K. W., Lawrence, D. M., Fisher, R., Lola da Costa, A. C., and Gentine, P.: Implementing plant hydraulics in the Community Land Model, version 5. *Journal of Advances in Modeling Earth Systems*, 11, 485–513. <https://doi.org/10.1029/2018MS001500>, 2019.
- Konapala, G., Mishra, A. K., Wada, Y., and Mann, M. E.: Climate change will affect global water availability through
840 compounding changes in seasonal precipitation and evaporation, *Nature Communications*, 11, <https://doi.org/10.1038/s41467-020-16757-w>, 2020.
- Koster, R. D., Dirmeyer, P. A., Guo, Z., Bonan, G., Chan, E., Cox, P., Gordon, C. T., Kanae, S., Kowalczyk, E., Lawrence, D., Liu, P., Lu, C. H., Malyshev, S., McAvaney, B., Mitchell, K., Mocko, D., Oki, T., Oleson, K., Pitman, A., Sud, Y. C., ...
845 GLACE Team: Regions of strong coupling between soil moisture and precipitation. *Science (New York, N.Y.)*, 305(5687), 1138–1140. <https://doi.org/10.1126/science.1100217>, 2004.



Kouki, K., Räisänen, P., Luojus, K., Luomaranta, A., and Riihelä, A.: Evaluation of Northern Hemisphere snow water equivalent in CMIP6 models during 1982–2014, *The Cryosphere*, 16, 1007–1030, <https://doi.org/10.5194/tc-16-1007-2022>,
850 2022.

Koven, C. D., Riley, W.J., and Stern, A.: Analysis of Permafrost Thermal Dynamics and Response to Climate Change in the CMIP5 Earth System Models. *J. Climate*, 26, 1877–1900, <https://doi.org/10.1175/JCLI-D-12-00228.1>, 2013.

855 Koven, C. D., Knox, R. G., Fisher, R. A., Chambers, J. Q., Christoffersen, B. O., Davies, S. J., Detto, M., Dietze, M. C., Faybishenko, B., Holm, J., Huang, M., Kovenock, M., Kueppers, L. M., Lemieux, G., Massoud, E., McDowell, N. G., Muller-Landau, H. C., Needham, J. F., Norby, R. J., Powell, T., Rogers, A., Serbin, S. P., Shuman, J. K., Swann, A. L. S., Varadharajan, C., Walker, A. P., Wright, S. J., and Xu, C.: Benchmarking and parameter sensitivity of physiological and vegetation dynamics using the Functionally Assembled Terrestrial Ecosystem Simulator (FATES) at Barro Colorado Island,
860 Panama, *Biogeosciences*, 17, 3017–3044, <https://doi.org/10.5194/bg-17-3017-2020>, 2020.

Krinner, G., Derksen, C., Essery, R., Flanner, M., Hagemann, S., Clark, M., Hall, A., Rott, H., Brutel-Vuilmet, C., Kim, H., Ménard, C. B., Mudryk, L., Thackeray, C., Wang, L., Arduini, G., Balsamo, G., Bartlett, P., Boike, J., Boone, A., Chérury, F., Colin, J., Cuntz, M., Dai, Y., Decharme, B., Derry, J., Ducharme, A., Dutra, E., Fang, X., Fierz, C., Ghattas, J., Gusev, Y.,
865 Haverd, V., Kontu, A., Lafaysse, M., Law, R., Lawrence, D., Li, W., Marke, T., Marks, D., Ménégoz, M., Nasonova, O., Nitta, T., Niwano, M., Pomeroy, J., Raleigh, M. S., Schaedler, G., Semenov, V., Smirnova, T. G., Stacke, T., Strasser, U., Svenson, S., Turkov, D., Wang, T., Wever, N., Yuan, H., Zhou, W., and Zhu, D.: ESM-SnowMIP: assessing snow models and quantifying snow-related climate feedbacks, *Geosci. Model Dev.*, 11, 5027–5049, <https://doi.org/10.5194/gmd-11-5027-2018>,
2018.

870

Lawrence, D., Coe, M., Walker, W., Verhot, L., Vandecar, K.: The Unseen Effects of Deforestation: Biophysical Effects on Climate., *Frontiers in Forests and Global Change*, 5,. doi:10.3389/ffgc.2022.756115, 2022.

Lawrence, D. M., Hurtt, G. C., Arneth, A., Brovkin, V., Calvin, K. V., Jones, A. D., Jones, C. D., Lawrence, P. J., de Noblet-Ducoudré, N., Pongratz, J., Seneviratne, S. I., and Shevliakova, E.: The Land Use Model Intercomparison Project (LUMIP) contribution to CMIP6: rationale and experimental design, *Geosci. Model Dev.*, 9, 2973–2998, <https://doi.org/10.5194/gmd-9-2973-2016>, 2016.
875

Lee, J.-Y., J. Marotzke, G. Bala, L. Cao, S. Corti, J.P. Dunne, F. Engelbrecht, E. Fischer, J.C. Fyfe, C. Jones, A. Maycock, J.
880 Mutemi, O. Ndiaye, S. Panickal, and T. Zhou: Future Global Climate: Scenario-Based Projections and Near-Term Information. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the*

Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 553–672, doi:10.1017/9781009157896.006, 2021.

885

Lehner, F., Wood, A.W., Vano, J.A. et al. The potential to reduce uncertainty in regional runoff projections from climate models. *Nat. Clim. Chang.* 9, 926–933, <https://doi-org.cuucar.idm.oclc.org/10.1038/s41558-019-0639-x>, 2019.

890 Lei, M., Li, J., Zhang, L., Deng, C., Li, Y., and He, J.: Inconsistent frequency trends between hourly and daily precipitation during warm season in mainland of China. *Geophysical Research Letters*, 49(19), <https://doi.org/10.1029/2022GL100277>, 2022

Lösing, M., Ebbing, J., and Szwillus, W.: Geothermal Heat Flux in Antarctica: Assessing Models and Observations by Bayesian Inversion”, *Frontiers in Earth Science*, 8, Art. no. 105, doi:10.3389/feart.2020.00105, 2020.

895

Li, L., Yang, Z.-L., Matheny, A. M., Zheng, H., Swenson, S. C., Lawrence, D. M., et al.: Representation of plant hydraulics in the Noah-MP land surface model: Model development and multiscale evaluation. *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002214. <https://doi.org/10.1029/2020MS002214>, 2021.

900

Li, X., Ault, T., Evans, C. P., Lehner, F., Carrillo, C. M., Donnelly, A., Crimmins, T., Gallinat, A. S., and Schwartz, M. D.: Diverging Northern Hemisphere Trends in meteorological versus ecological Indicators of spring onset in CMIP6, *Geophysical Research Letters*, 50, <https://doi.org/10.1029/2023gl102833>, 2023.

905 Li, X., Ault, T., Richardson, A. D., Frohking, S., Herrera, D. A., Friedl, M. A., Carrillo, C. M., and Evans, C. P.: Northern hemisphere land-atmosphere feedback from prescribed plant phenology in CESM, *Journal of Climate*, <https://doi.org/10.1175/jcli-d-23-0179.1>, 2024a.

910 Li, X., Carrillo, C. M., Ault, T., Richardson, A. D., Friedl, M. A., and Frohking, S.: Evaluation of leaf phenology of different vegetation types from local to hemispheric scale in CLM, *Journal of Geophysical Research Biogeosciences*, 129, <https://doi.org/10.1029/2024jg008261>, 2024b.

Li, X., Melaas, E., Carrillo, C. M., Ault, T., Richardson, A. D., Lawrence, P., Friedlingstein, M. A., Seyednasrollah, B., Lawrence, D. M., and Young, A. M.: A comparison of land surface phenology in the Northern Hemisphere derived from



915 satellite remote sensing and the Community Land Model, *Journal of Hydrometeorology*, 23(6), 859–873,
<https://doi.org/10.1175/JHM-D-21-0169.1>, 2022a.

Li, Q., S. Sun, and Y. Xue, : Analyses and development of a hierarchy of frozen soil models for cold region study, *Journal of Geophysical Research-Atmosphere*. 115, D03107, doi:10.1029/2009JD012530, 2010.

920

Li, Y., Brando, P.M., Morton, D.C. et al.: Deforestation-induced climate change reduces carbon storage in remaining tropical forests, *Nat Commun* 13, 1964, <https://doi.org/10.1038/s41467-022-29601-0>, 2022b.

925 Li, Y., Huang, B., Tan, C., Zhang, X., Cherubini, F., and Rust, H. W.: Investigating the global and regional response of drought to idealized deforestation using multiple global climate models, *Hydrol. Earth Syst. Sci.*, 29, 1637–1658,
<https://doi.org/10.5194/hess-29-1637-2025>, 2025a.

Li, Y., et al. Schematic Diagram Illustrating the Land and Land Ice Theme Opportunities in CMIP7 Assessment Fast Track Data Request. Zenodo, 17 June 2025, doi:10.5281/zenodo.15681951, 2025b.

930

Lombardozzi, D.L., Wieder, W.R., Keppel-Aleks, G. et al.: Agricultural fertilization significantly enhances amplitude of land-atmosphere CO₂ exchange. *Nat Commun* 16, 1742. <https://doi.org/10.1038/s41467-025-56730-z>, 2025.

935 Luo, K., Wang, X., de Jong, M., and Flannigan, M.: Drought triggers and sustains overnight fires in North America. *Nature* 627, 321–327, <https://doi.org/10.1038/s41586-024-07028-5>, 2024a.

Luo, X., Ge, J., Cao, Y., Liu, Y., Yang, L., Wang, S., and Guo, W.: Local and nonlocal biophysical effects of historical land use and land cover changes in CMIP6 models and the intermodel uncertainty. *Earth's Future*, 12, e2023EF004220. <https://doi.org/10.1029/2023EF004220>, 2024b.

940

Mackallah, C., Jukes, M., Anstey, J., Pascoe, C., Rigoudy, G., Moine, M.-P., Lovato, T., Pamment, A., Kawamiya, M., Bergman, T., Schupfner, M., Koven, C., Lam, T., Dingley, B., O'Rourke, E., Turner, B., Ellis, D., and Mizielinski, M.: CMIP7 Data Request: a transparent community-led approach leveraging interactive web tools and enhanced CMIP governance [in preparation]

945

Mao, J., Fu, W., Shi, X., Ricciuto, D.M., Fisher, J.B., Dickinson, R.E., Wei, Y., Shem, W., Piao, S., Wang, K. and Schwalm, C.R.: Disentangling climatic and anthropogenic controls on global terrestrial evapotranspiration trends. *Environmental Research Letters*, 10(9), p.094008., 10.1088/1748-9326/10/9/094008 , 2015.



950 Marsh, H., Jin, H., Duan, Z., Holst, J., Eklundh, L., and Zhang, W.: Plant Phenology Index leveraging over conventional vegetation indices to establish a new remote sensing benchmark of GPP for northern ecosystems, *International Journal of Applied Earth Observation and Geoinformation*, 136, 104289, <https://doi.org/10.1016/j.jag.2024.104289>, 2025.

Marzeion, B., R. Hock, B. Anderson, A. Bliss, N. Champollion, K. Fujita, M. Huss, W. Immerzeel, P. Kraaijenbrink, J-H. Malles, F. Maussion, Valentina Radic, D. R. Rounce, A. Sakai, S. Shannon, R. van de Wal, H. Zekollari: Partitioning the Uncertainty of Ensemble Projections of Global Glacier Mass Change. *Earth's Future* 12, e2019EF001470, doi: 10.1029/2019EF001470, 2020.

McPartland, M.Y., Lovato, T.L., Koven, C., Wilson, J.D., Turner, B., Petrik, C.M., Licón-Saláiz, J., Li, F., Lhardy, F., Clement Kinney, J., Kawamiya, M., Gillett, N.P., Fall, C.M.N., Danek, C., Brierley, C.M., Bastos, A., Andrews, O.: CMIP7 Data Request: Earth System Priorities and Opportunities [in preparation]

Michalek, A.T., Villarini, G., and Husic, A.: Climate change projected to impact structural hillslope connectivity at the global scale, *Nat Commun* 14, 6788, <https://doi.org/10.1038/s41467-023-42384-2>, 2023.

965 Milner, A.M., K. Khamis, T.J. Battin, J.E. Brittain, N.E. Barrand, L. Füreder, S. Cauvy-Fraunié, G.M. Gíslason, D. Jacobsen, D.M. Hannah, A.J. Hodson, E. Hood, V. Lencioni, J.S. Ólafsson, C.T. Robinson, M. Tranter, L.E. Brown, Glacier shrinkage driving global changes in downstream systems, *Proc. Natl. Acad. Sci. U.S.A.* 114 (37) 9770-9778, <https://doi.org/10.1073/pnas.1619807114> (2017).

970 Minallah, S., and A. L. Steiner: Analysis of the Atmospheric Water Cycle for the Laurentian Great Lakes Region Using CMIP6 Models. *J. Climate*, 34, 4693–4710, <https://doi.org/10.1175/JCLI-D-20-0751.1>, 2021.

Munier, S., Carrer, D., Planque, C., Camacho, F., Albergel, C., Calvet, J.-C., Satellite Leaf Area Index: Global Scale Analysis of the Tendencies Per Vegetation Type Over the Last 17 Years, *Remote Sensing*, 10(3), 424, <https://doi.org/10.3390/rs10030424>, 2018.

Murray-Tortarolo, G., Anav, A., Friedlingstein, P., Sitch, S., Piao, S., Zhu, Z., Poulter, B., Zaehle, S., Ahlström, A., Lomas, M., Levis, S., Viovy, N., and Zeng, N.: Evaluation of land surface models in reproducing satellite-derived LAI over the high-latitude northern hemisphere. Part I: uncoupled DGVMs, *Remote Sens.*, 5, 4819–4838, <https://doi.org/10.3390/rs5104819>, 2013.



- 985 Negrón-Juárez, R., Wehner, M., Dias, M., Ullrich, P., Chambers, J., and Riley, W.: Coupled model intercomparison project phase 6 (CMIP6) high resolution model intercomparison project (HighResMIP) bias in extreme rainfall drives underestimation of amazonian precipitation. *Environmental Research Communications*. 6. 10.1088/2515-7620/ad6ff9, 2024.
- Norris, J., Hall, A., Neelin, J., Thackeray, C., and Chen, D.: Evaluation of the Tail of the Probability Distribution of Daily and Subdaily Precipitation in CMIP6 Models. *Journal of Climate*. 34. 1-61. 10.1175/JCLI-D-20-0182.1., 2021
- 990 Notaro, M., J. Jorns, and L. Briley,: Representation of Lake–Atmosphere Interactions and Lake-Effect Snowfall in the Laurentian Great Lakes Basin among HighResMIP Global Climate Models. *J. Atmos. Sci.*, 79, 1325–1347, <https://doi.org/10.1175/JAS-D-21-0249.1>, 2022.
- 995 Nowicki, S. M. J., Payne, A., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., and Shepherd, A.: Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6, *Geosci. Model Dev.*, 9, 4521–4545, <https://doi.org/10.5194/gmd-9-4521-2016>, 2016.
- 1000 Nowicki, S., Goelzer, H., Seroussi, H., Payne, A. J., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Alexander, P., Asay-Davis, X. S., Barthel, A., Bracegirdle, T. J., Cullather, R., Felikson, D., Fettweis, X., Gregory, J. M., Hattermann, T., Jourdain, N. C., Kuipers Munneke, P., Larour, E., Little, C. M., Morlighem, M., Nias, I., Shepherd, A., Simon, E., Slater, D., Smith, R. S., Straneo, F., Trusel, L. D., van den Broeke, M. R., and van de Wal, R.: Experimental protocol for sea level projections from ISMIP6 stand-alone ice sheet models, *The Cryosphere*, 14, 2331–2368, <https://doi.org/10.5194/tc-14-2331-2020>, 2020.
- 1005 Nowicki, S., Payne, A.J., Seroussi, H.L., et al., AGU Fall Meeting Abstracts, Preparing for the Ice Sheet Model Intercomparison Project for CMIP7 (ISMIP7), 38, C31D-038, 2024
- 1010 Oppenheimer, M., B.C. Glavovic, J. Hinkel, R. van de Wal, A.K. Magnan, A. Abd-Elgawad, R. Cai, M. Cifuentes-Jara, R.M. DeConto, T. Ghosh, J. Hay, F. Isla, B. Marzeion, B. Meyssignac, and Z. Sebesvari,: Sea Level Rise and Implications for Low-Lying Islands, Coasts and Communities. In: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 321-445. <https://doi.org/10.1017/9781009157964.006>, 2019.
- 1015 Park, T., Ganguly, S., Tømmervik, H., Euskirchen, E. S., Høgda, K.-A., Karlsen, S. R., Brovkin, V., Nemani, R. R., and Myneni, R. B.: Changes in growing season duration and productivity of northern vegetation inferred from long-term remote sensing data, *Environ. Res. Lett.*, 11, 084001, <https://doi.org/10.1088/1748-9326/11/8/084001>, 2016.



- 1020 Park, C.Y., Takahashi, K., Fujimori, S. et al. Attributing human mortality from fire PM_{2.5} to climate change. *Nat. Clim. Chang.* 14, 1193–1200, <https://doi.org/10.1038/s41558-024-02149-1>, 2024
- 1025 Peano, D., Materia, S., Collalti, A., Alessandri, A., Anav, A., Bombelli, A., and Gualdi, S: Global variability of simulated and observed vegetation growing season. *Journal of Geophysical Research: Biogeosciences*, 124, 3569–3587. <https://doi.org/10.1029/2018JG004881>, 2019.
- 1030 Peano, D., Hemming, D., Materia, S., Delire, C., Fan, Y., Joetzjer, E., Lee, H., Nabel, J. E. M. S., Park, T., Peylin, P., Wårlind, D., Wiltshire, A., and Zaehle, S.: Plant phenology evaluation of CRESCENDO land surface models – Part 1: Start and end of the growing season, *Biogeosciences*, 18, 2405–2428, <https://doi.org/10.5194/bg-18-2405-2021>, 2021.
- 1035 Peano, D., Hemming, D., Delire, C., Fan, Y., Lee, H., Materia, S., Nabel, J. E. M. S., Park, T., Wårlind, D., Wiltshire, A., and Zaehle, S.: Plant phenology evaluation of CRESCENDO land surface models. Part II: Trough, peak, and amplitude of growing season, *EGUsphere [preprint]*, <https://doi.org/10.5194/egusphere-2024-4114>, 2025.
- Poyatos, R., Granda, V., Flo, V., Adams, M. A., et al.: Global transpiration data from sap flow measurements: the SAPFLUXNET database, *Earth Syst. Sci. Data*, 13, 2607–2649, <https://doi.org/10.5194/essd-13-2607-2021>, 2021.
- 1040 RGI Consortium: Randolph Glacier Inventory - A Dataset of Global Glacier Outlines, Version 7.0. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. doi:10.5067/f6jmovy5navz. Online access: <https://doi.org/10.5067/f6jmovy5navz>, 2023.
- 1045 Richardson, A. D., Keenan, T. F., Migliavacca, M., Ryu, Y., Sonnentag, O., and Toomey, M.: Climate change, phenology, and phenological control of vegetation feedbacks to the climate system, *Agr. Forest Meteorol.*, 169, 156–173, <https://doi.org/10.1016/j.agrformet.2012.09.012>, 2013.
- 1050 Richardson, A. D., Anderson, R. S., Arain, M. A., Barr, A. G., Bohrer, G., Chen, G., Chen, J. M., Ciais, P., Davis, K. J., Desai, A. R., Dietze, M. C., Dragoni, D., Garrity, S. R., Gough, C. M., Grant, R., Hollinger, D. Y., Margolis, H. A., McCaughey, H., Migliavacca, M., Monson, R. K., Munger, J. W., Poulter, B., Raczka, B. M., Ricciuto, D. M., Sahoo, A. K., Schaefer, K., Tian, H., Vargas, R., Verbeeck, H., Xiao, J., and Xue, Y.: Terrestrial biosphere models need better representation of vegetation phenology: results from the North American Carbon Program Site Synthesis, *Global Change Biology*, 18, 566–584, <https://doi.org/10.1111/j.1365-2486.2011.02562.x>, 2012.



Richardson, K., Steffen, W., Lucht, W., Bendtsen, J., Cornell, S.E., Donges, J.F., Drüke, M., Fetzer, I., Bala, G., Von Bloh, W. and Feulner, G.: Earth beyond six of nine planetary boundaries. *Science advances*, 9(37), p.eadh2458, <https://doi.org/10.1126/sciadv.adh2458>, 2023.

1055 Roberts, M. J., Reed, K. A., Bao, Q., Barsugli, J. J., Camargo, S. J., Caron, L.-P., Chang, P., Chen, C.-T., Christensen, H. M., Danabasoglu, G., Frenger, I., Fučkar, N. S., ul Hasson, S., Hewitt, H. T., Huang, H., Kim, D., Kodama, C., Lai, M., Leung, L.-Y. R., Mizuta, R., Nobre, P., Ortega, P., Paquin, D., Roberts, C. D., Scoccimarro, E., Seddon, J., Treguier, A. M., Tu, C.-Y., Ullrich, P. A., Vidale, P. L., Wehner, M. F., Zarzycki, C. M., Zhang, B., Zhang, W., and Zhao, M.: High-Resolution Model Intercomparison Project phase 2 (HighResMIP2) towards CMIP7, *Geosci. Model Dev.*, 18, 1307–1332, 1060 <https://doi.org/10.5194/gmd-18-1307-2025>, 2025.

Santanello, J. A., Peters-Lidard, C. D., and Kumar, S. V.: Diagnosing the sensitivity of local Land–Atmosphere coupling via the soil Moisture–Boundary layer interaction, *Journal of Hydrometeorology*, 12, 766–786, <https://doi.org/10.1175/jhm-d-10-05014.1>, 2011.

1065 Santanello, J. A., Peters-Lidard, C. D., Kumar, S. V., Alonge, C., and Tao, W.-K.: A modeling and observational framework for diagnosing local Land–Atmosphere coupling on diurnal time scales, *Journal of Hydrometeorology*, 10, 577–599, <https://doi.org/10.1175/2009jhm1066.1>, 2009.

1070 Seo, E. and Dirmeyer, P. A.: Understanding the diurnal cycle of land–atmosphere interactions from flux site observations, *Hydrol. Earth Syst. Sci.*, 26, 5411–5429, <https://doi.org/10.5194/hess-26-5411-2022>, 2022.

Schumacher, D.L., Keune, J., Dirmeyer, P. et al.: Drought self-propagation in drylands due to land–atmosphere feedbacks. *Nat. Geosci.* 15, 262–268, <https://doi.org/10.1038/s41561-022-00912-7>, 2022.

1075 Seneviratne, S. I., Lüthi, D., Litschi, M., and Schär, C.: Land-atmosphere coupling and climate change in Europe. *Nature*, 443(7108), 205–209. <https://doi.org/10.1038/nature05095>, 2006.

1080 Shrestha, M., Koike, T., Hirabayashi, Y., Xue, Y., Wang, L., Rasul, G., and Ahmad, B.: Integrated simulation of snow and glacier melt in water and energy balance-based, distributed hydrological modeling framework at Hunza River Basin of Pakistan Karakoram region, *Journal of Geophysical Research Atmospheres*, 120, 4889–4919, <https://doi.org/10.1002/2014jd022666>, 2015.



1085 Song, X., Wang, D.-Y., Li, F., and Zeng, X.-D.: Evaluating the performance of CMIP6 Earth system models in simulating
global vegetation structure and distribution, *Advances in Climate Change Research*, 12, 584–
595, <https://doi.org/10.1016/j.accre.2021.06.008>, 2021.

1090 Sterling, S., Ducharne, A., and Polcher, J.: The impact of global land-cover change on the terrestrial water cycle. *Nature Clim
Change* 3, 385–390, <https://doi.org/10.1038/nclimate1690>, 2013

1095 Stokes, C. R., Abram, N. J., Bentley, M. J., Edwards, T. L., England, M. H., Foppert, A., Jamieson, S. S. R., Jones, R. S., King,
M. A., Lenaerts, J. T. M., Medley, B., Miles, B. W. J., Paxman, G. J. G., Ritz, C., Van De Fliedrt, T., and Whitehouse, P. L.:
Response of the East Antarctic Ice Sheet to past and future climate change, *Nature*, 608, 275–
286, <https://doi.org/10.1038/s41586-022-04946-0>, 2022.

1095 Sun, Q., Whitney, M. M., Bryan, F. O., and Tseng, Y.-H.: A box model for representing estuarine physical processes in Earth
system models, *Ocean Modelling*, 112, 139–153, <https://doi.org/10.1016/j.ocemod.2017.03.004>, 2017.

1100 Sun, S., Jin, J., and Xue, Y.: A simplified layer snow model for global and regional studies, *Journal of Geophysical Research
Atmospheres*, 104, 19587–19597, 1999.

1105 Swart, N., Martin, T., Beadling, R., Chen, J.-J., England, M. H., Farneti, R., Griffies, S. M., Hatterman, T., Haumann, F. A.,
Li, Q., Marshall, J., Muilwijk, M., Pauling, A. G., Purich, A., Smith, I. J., and Thomas, M.: The Southern Ocean Freshwater
release model experiments Initiative (SOFIA): Scientific objectives and experimental design, *EGUsphere [preprint]*,
<https://doi.org/10.5194/egusphere-2023-198>, 2023.

1110 Tang, T., Lee, X., Zhang, K., Cai, L., Lawrence, D. M., and Shevliakova, E.: Biophysical Impact of Land-Use and Land-Cover
Change on Subgrid Temperature in CMIP6 Models. *Journal of Hydrometeorology*, 24(3), 373–
388. <https://doi.org/10.1175/JHM-D-22-0073.1>, 2023

Tang, G., Nicholls, Z., Jones, C., Gasser, T., Norton, A., Ziehn, T., Romero-Prieto, A., and Meinshausen, M.: Investigating
carbon and nitrogen conservation in reported CMIP6 Earth system model data, *Geosci. Model Dev.*, 18, 2111–2136,
<https://doi.org/10.5194/gmd-18-2111-2025>, 2025a.

1115 Tang, G., Nicholls, Z., Norton, A., Zaehle, S., and Meinshausen, M.: Synthesizing global carbon–nitrogen coupling effects –
the MAGICC coupled carbon–nitrogen cycle model v1.0, *Geosci. Model Dev.*, 18, 2193–2230, <https://doi.org/10.5194/gmd-18-2193-2025>, 2025b.



Vanderkelen, I., Gharari, S., Mizukami, N., Clark, M. P., Lawrence, D. M., Swenson, S., Pokhrel, Y., Hanasaki, N., van
1120 Griensven, A., and Thiery, W.: Evaluating a reservoir parametrization in the vector-based global routing model mizuRoute
(v2.0.1) for Earth system model coupling, *Geosci. Model Dev.*, 15, 4163–4192, <https://doi.org/10.5194/gmd-15-4163-2022>,
2022.

Van Westen, R. M., Kliphuis, M., and Dijkstra, H. A.: Physics-based early warning signal shows that AMOC is on tipping
1125 course, *Science Advances*, 10, <https://doi.org/10.1126/sciadv.adk1189>, 2024.

Vettoretti, G., Nuterman, R., and Jochum, M.: Impacts of Parameterizing Estuary Mixing on the Large-Scale Circulations in
the Community Earth System Model. *J. Climate*, 37, 4461–4479, <https://doi.org/10.1175/JCLI-D-23-0365.1>, 2024.

Walton, J., Swaminathan, R., Jones, C., Dittus, A., Liddicoat, S., Rumbold, S., and Smith, R.: Assessing the Realism of the
1130 TIPMIP ESM idealized Experiment Protocol, EGU General Assembly 2025, Vienna, Austria, 27 Apr–2 May 2025, EGU25-
4403, <https://doi.org/10.5194/egusphere-egu25-4403>, 2025.

Wang, L., Zhou, J., Qi, J., Sun, L., Yang, K., Tian, L., Lin, Y., Liu, W., Shrestha, M., Xue, Y., Koike, T., Ma, Y., Li, X., Chen,
1135 Y., Chen, D., Piao, S., and Lu, H.: Development of a land surface model with coupled snow and frozen soil physics, *Water
Resources Research*, 53, 5085–5103, <https://doi.org/10.1002/2017wr020451>, 2017.

Wehner, M., Lee, J., Risser, M., Ullrich, P., Gleckler, P., and Collins, W.D.: Evaluation of extreme sub-daily precipitation in
high-resolution global climate model simulations, *Philos Trans A Math Phys Eng Sci.* 2021 Apr 19;379(2195):20190545. doi:
1140 10.1098/rsta.2019.0545

Whitehouse, P. L.: Glacial isostatic adjustment modelling: historical perspectives, recent advances, and future directions, *Earth
Surf. Dynam.*, 6, 401–429, <https://doi.org/10.5194/esurf-6-401-2018>, 2018.

te Wierik, S.A., Cammeraat, E.L., Gupta, J. and Artzy-Randrup, Y.A., Reviewing the impact of land use and land-use change
on moisture recycling and precipitation patterns. *Water Resources Research*, 57(7),
1145 p.e2020WR029234. <https://doi.org/10.1029/2020WR029234>, 2021.

Winkler, A. J., Myneni, R. B., Hannart, A., Sitch, S., Haverd, V., Lombardozzi, D., Arora, V. K., Pongratz, J., Nabel, J. E. M.
1150 S., Goll, D. S., Kato, E., Tian, H., Arneth, A., Friedlingstein, P., Jain, A. K., Zaehle, S., and Brovkin, V., Slowdown of the



greening trend in natural vegetation with further rise in atmospheric CO₂, *Biogeosciences*, 18, 4985–5010, <https://doi.org/10.5194/bg-18-4985-2021>, 2021.

1155 Wolken, G. J. et al.: Glacier and Permafrost Hazards. <https://doi.org/10.25923/v40r-0956>, 2021.

Wu, R., M. Lo, and B. R. Scanlon: The Annual Cycle of Terrestrial Water Storage Anomalies in CMIP6 Models Evaluated against GRACE Data. *J. Climate*, 34, 8205–8217, <https://doi.org/10.1175/JCLI-D-21-0021.1>, 2021.

1160 Wu, J. and Dirmeyer, P. A.: Drought Demise Attribution Over CONUS, vol. 125, no. 4, <https://doi.org/10.1029/2019jd031255>, 2020.

Xu, X., Medvigy, D., Powers, J.S., Becknell, J.M. and Guan, K.: Diversity in plant hydraulic traits explains seasonal and inter-annual variations of vegetation dynamics in seasonally dry tropical forests. *New Phytol*, 212: 80-95. <https://doi.org/10.1111/nph.14009>, 2016.

1165 Yao, Y., Ciais, P., Joetzjer, E., Hong, S., Li, W., Zhu, L., and Viovy, N.: Future drought-induced tree mortality risk in Amazon rainforest. *Earth's Future*, 12, e2023EF003740. <https://doi.org/10.1029/2023EF003740> , 2024.

1170 Yao, Y., Ducharne, A., Cook, B. I., De Hertog, S. J., Aas, K. S., Arboleda-Obando, P. F., Buzan, J., Colin, J., Costantini, M., Decharme, B., Lawrence, D. M., Lawrence, P., Leung, L. R., Lo, M.-H., Devaraju, N., Wieder, W. R., Wu, R.-J., Zhou, T., Jägermeyr, J., McDermid, S., Pokhrel, Y., Elling, M., Hanasaki, N., Muñoz, P., Nazarenko, L. S., Otta, K., Satoh, Y., Yokohata, T., Jin, L., Wang, X., Mishra, V., Ghosh, S., and Thiery, W.: Impacts of irrigation expansion on moist-heat stress based on IRRMIP results, *Nature Communications*, 16, <https://doi.org/10.1038/s41467-025-56356-1>, 2025.

1175 Yoon, D., Kang, T., Cha, D.-H., Song, C.K., Lee, M.I., Min, K.H., et al.: Role of land–atmosphere interaction in the 2016 Northeast Asia heat wave: Impact of soil moisture initialization. *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037718. <https://doi.org/10.1029/2022JD037718>, 2023

1180 Yu, L., Y. Xue, I. Diallo, 2021: Vegetation greening in China and its effect on summer regional climate, *Science Bulletin*, 66, 13-17. <https://doi.org/10.1016/j.scib.2020.09.003>, 2021.

Zhao, W., Zhu, B., Davis, S.J., Ciais, P., Hong, C., Liu, Z. and Gentine, P., Reliance on fossil fuels increases during extreme temperature events in the continental United States. *Communications Earth & Environment*, 4, 473, <https://doi.org/10.1038/s43247-023-01147-z>, 2023.



1185

Zhang, H., Zhang, F., Zhang, G., and Yan, W.: Why do CMIP6 models fail to simulate snow depth in terms of temporal change and high mountain snow of China skillfully? *Geophysical Research Letters*, 49, e2022GL098888. <https://doi.org/10.1029/2022GL098888>, 2022.

1190 Zhu, Z., Piao, S., Myneni, R.B., Huang, M., Zeng, Z., Canadell, J.G., Ciais, P., Sitch, S., Friedlingstein, P., Arneeth, A. and Cao, C., Greening of the Earth and its drivers. *Nature climate change*, 6(8), pp.791-795., <https://doi.org/10.1038/nclimate3004>, 2026.