

Investigating the impact of sub-ice shelf melt on ~~the~~ Antarctic ice sheet spin-up and projections

Fan Gao^{1,2}, Qiang Shen^{1,2}, Hansheng Wang^{1,2}, Tong Zhang³, Liming Jiang^{1,2}, Yan Liu⁴, C K Shum⁵, Yan An^{1,2}, and Xu Zhang^{1,2}

5 ¹State Key Laboratory of Precision Geodesy, Innovation Academy for Precision Measurement Science and Technology, Chinese Academy of Sciences, Wuhan 430077, China

²College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

³State Key Laboratory of Earth Surface Processes and Disaster Risk Reduction, Faculty of Geographical Science, Beijing Normal University, Beijing 100875, China

10 ⁴State Key Laboratory of Remote Sensing Science, College of Global Change and Earth System Science, Beijing Normal University, Beijing 100875, China

⁵Division of Geodetic Science, School of Earth Sciences, The Ohio State University, Columbus, Ohio, 43210, USA

Correspondence to: Qiang Shen (cl980606@whigg.ac.cn) and Tong Zhang (tongzhangice@gmail.com)

Abstract. Sub-ice shelf melting is critical for the stability of the Antarctic ~~Ice Sheet~~ ice sheet, as it influences ice-
15 shelf buttressing that reduces grounded ice flow. Previous studies have emphasized that uncertainties in the state of
sub-ice shelf melting contribute to uncertainties in future sea-level projections. To better understand how sub-ice shelf
melt rates affect model initialization and predictions, we adopt a single ice-sheet model (PISM) and investigate two
different sub-ice shelf melt rate schemes during model spin-ups. We then drive the Antarctic ~~Ice Sheet~~ ice sheet into
the future using identical environmental forcings. We find that, despite closely matched steady-state geometries
20 achieved through the spin-up process with different sub-ice shelf melt rates, the prognostic simulations reveal
significantly divergent ice mass changes, particularly in marine ice sheet regions. By 2100, the difference in global
sea-level contributions from the Antarctic ~~Ice Sheet~~ ice sheet can be as large as ~57%, primarily from West Antarctica.
This discrepancy arises because the spin-up initialization method alters the ice sheet's dynamic state, such as basal
friction and thermal regimes, leading to ~~varied differences in~~ ice-sheet mass changes ~~over time~~. Therefore, this study
25 underscores the importance of sub-ice shelf melting and ice-sheet model initialization methods in reducing
uncertainties in predicting the Antarctic ~~Ice Sheet's~~ ice sheet's future.

1 Introduction

A substantial majority of Antarctica's grounded ice discharges through its fringing ice shelves, which provide critical
buttressing to upstream ice ~~mainly~~ through two primary mechanisms: lateral shear stresses along sidewalls and basal
30 resistance forces at pinning points on topographic highs (Schoof, 2007; Goldberg et al., 2009; Feldmann ~~&~~and
Levermann, 2023; Feldmann et al., 2024; Miles ~~&~~and Bingham, 2024). Ice shelves are highly vulnerable to oceanic

forcing due to the near-flotation elevation exposes them to warm seawater, which causes enhanced basal melting (Bindschadler et al., 2013; Depoorter et al., 2013; Li et al., 2023). Observations reveal accelerating ocean-driven thinning of Antarctic ice shelves over recent decades (Paolo et al., 2015; Rignot et al., 2019), where enhanced basal melting reduces buttressing effects and promotes grounding-line retreat, which collectively represent the primary driver of increased ice discharge (Jacobs et al., 2011; Pritchard et al., 2012; Seroussi et al., 2014; Jourdain et al., 2020; Reese et al., 2020). Particularly on retrograde bed slopes, such retreat may trigger Marine Ice Sheet Instability (MISI), a critical feedback mechanism that is often identified as a decisive factor in the collapse of the West Antarctica (Schoof, 2007; Hill et al., 2024). This process may amplify [Antarctica's](#) contribution to global sea-level rise by 0.5 to 0.8 meters of sea-level equivalent (m SLE) this century (Ritz et al., 2015).

Methods for ice-sheet models to represent sub-ice shelf melting include linear/non-linear and local/non-local dependency thermal forcing parameterizations (Martin et al., 2011; Favier et al., 2019; Lowry et al., 2021), ice-shelf cavity models developed from box or plume models (Lazeroms et al., 2018; PICO, Reese et al., 2018; Favier et al., 2019; PICOP, Pelle et al., 2019), empirical approximations (Cornford et al., 2015; Cornford et al., 2020), basin-averaged melt estimates (Seroussi et al., 2019), and spatially partitioned quadratic parameterizations (ISMIP6 protocol, Jourdain et al., 2020). The initMIP-Antarctica experiments revealed that ice-sheet model responses exhibit significant divergence due to variations in initial basal melting conditions, with the resulting range accounting for 5 % to 125 % of the total mass change (Seroussi et al., 2019, 2020). This pronounced model spread underscores persistent challenges in accurately representing sub-ice shelf oceanic processes during ice-sheet model initialization (Pritchard et al., 2012; Alevropoulos-Borrill et al., 2020), and may propagate into projection uncertainties, particularly for ice dynamics influenced by oceanic forcing. [For instance, simulations with the Yelmo ice-sheet model indicate that the Antarctic ice sheet's sea-level contribution is highly sensitive to the ice-ocean interaction under varying basal melt coefficients, with several projections reaching up to 3 m SLE by 2500 \(Juarez-Martinez et al., 2024\).](#)

Previous model intercomparison projects (e.g., initMIP-Antarctica, [Seroussi et al., 2019](#)) combined ice-sheet models with varying numerical complexities and initialization methods, making it difficult to attribute uncertainties to specific sources; our study isolates the impact of oceanic conditions by using a single ice-sheet model with identical initialization except for the basal melting scheme. Zhang et al. (2024) addressed this limitation by adopting a single ice-sheet model (Community Ice Sheet Model, CISM; Lipscomb et al., 2019; Berdahl et al., 2023) to investigate the impacts of geothermal heat flux and basal sliding conditions on Greenland Ice Sheet initialization. Extending this approach and considering the crucial role of ice shelves in Antarctica, we propose conducting similar experiments for the Antarctic [Ice Sheet](#) (AIS) using a single ice-sheet model and initialization method to assess the impacts of sub-ice shelf melt rates. This focused investigation will address two key questions: (1) How do varying sub-ice shelf melt rates impact the model initialization state? (2) How does this initial state affect long-term AIS projections?

Therefore, in this paper, we consider two different sub-ice shelf melt rates schemes (Section 2) in the Parallel Ice Sheet Model by first spinning-up and then projecting the AIS. The structure of this paper is organized as follows: Section 2 details the methodological approach and experimental design for projections. Section 3 and Section 4

provide comprehensive results and discuss the implications for ice dynamics and sea-level rise projections, while Section 5 ~~analyzes~~analyses uncertainties in the model initialization and projection.

2 Model and Methods

70 We conduct ice-sheet simulations using the Parallel Ice Sheet Model (PISM v.1.0) (Bueler et al., 2007; Martin et al., 2011; Winkelmann et al., 2011; Albrecht et al., 2020), an open-source, three-dimensional thermomechanical coupled model that integrates ice dynamics and thermodynamics. PISM employs a hybrid stress balance strategy (Martin et al., 2011; Winkelmann et al., 2011) by combining the Shallow Ice Approximation (SIA) for grounded ice (Gudmundsson, 2003; Bueler et al., 2007; Pollard ~~&and~~ DeConto, 2012) and the Shallow Shelf Approximation (SSA) 75 for floating ice (Hindmarsh, 2006; Bueler ~~&and~~ Brown, 2009; Pollard ~~&and~~ DeConto, 2012). In PISM, basal resistance—which directly governs sliding velocities—is calculated using a generalized power law that ranges from plastic Coulomb sliding to a linear sliding law (Bueler and Brown, 2009; Winkelmann et al., 2011; Garbe et al., 2020). The grounding-line migration in PISM is ~~optimized~~determined through a sub-grid scheme, which interpolates key physical variables such as basal shear stress, basal melt rate, and basal friction based on spatial gradients across the 80 interface between grounded and floating cells (Feldmann et al., ~~2017~~2014; Nowicki et al., 2020). This approach reduces physical gradients across the grounding line and simulates a more realistic and dynamic representation of the ice margin (Leguy et al., 2014; Golledge et al., 2015).

We utilize the BedMachine v.3 dataset (Morlighem et al., 2019) for initial topography, encompassing ice thickness and bedrock topography. Air temperature and precipitation inputs are derived from RACMO 2.3p2, averaged over 85 1979–2014 (van Wessem et al., 2018). Surface mass balance is calculated using a degree-day model (Ohmura, 2001; Calov ~~&and~~ Greve, ~~2017~~2005), with near-surface temperature locally adjusted based on elevation changes using a correction factor of $0.008^{\circ}\text{C}/\text{m}^{-1}$ (Pittard et al., 2022). The PISM ocean module provides the sub-ice shelf temperature and mass flux to the ice dynamics core via two different schemes. Sub-ice shelf temperature is applied as a Dirichlet boundary condition in the energy conservation code, while sub-ice shelf mass flux enters as a source in the mass 90 conservation equation. ~~In our experiments (Martin et al., 2011).~~

For model spin-up, the mass flux is derived either directly from observed basal melt rates or indirectly through a parameterization using ocean temperature and salinity, depending on the available oceanic data. Where directly using observed basal melt rates (Rignot et al., 2013; Fig. 1) and ice-shelf basal temperature (Chambers et al., 2021; Fig. 1), the sub-ice shelf mass flux is ~~computed~~given by ~~Eq. 1~~:

$$95 \quad S_1 = \rho_i B, \quad (1)$$

where ρ_i indicates the ice density, and B represents the sub-ice shelf melt rates. For simulations driven by Southern Ocean temperature and salinity (Schmidtke et al., 2014; Fig. 2) as implemented in LOW21 (Lowry et al., 2021), the mass flux at ice-ocean interface—representing the coupled melting effect of ocean temperature and salinity—is obtained indirectly through a simplified linear thermal forcing (TF-linear) parameterization (Beckmann and Goosse, 100 2003; Martin et al., 2011) ~~following Eq. 2~~:

$$S_2 = \rho_{sw} c_m \gamma_T F_{melt} (T_s - T_f) / (L_i \rho_i), \quad (2)$$

where ρ_{sw} denotes the seawater density, c_m represents the specific heat capacity of the ocean mixed layer, L_i refers to the latent heat of phase change for ice, γ_T represents the thermal exchange velocity between seawater and ice (assigned $\gamma_T = 10^{-4}$; [Hellmer and Olbers, 1989](#); Holland and Jenkins, 1999; [Hellmer and Olbers, 1989](#)), F_{melt} is a model parameter (assigned $F_{melt} = 5 \times 10^{-3}$; [Beckmann and Goosse, 2003](#)), T_s is the vertically averaged ocean temperature between 200 m and 1000 m depth along the continental slope (assigned $T_s = 271.45$ K, representing relatively warm water masses of the coastal current ([Holland and Jenkins, 1999](#); [Beckmann and Goosse, 2003](#); [Martin et al., 2011](#)), and), T_f denotes the freezing temperature of seawater at depth z_b beneath the ice shelf:

$$T_f = 273.15 + 0.0939 - 0.057S_0 + 7.64 \times 10^{-4}z_b, \quad (3)$$

where S_0 denotes the specified ocean salinity (35 psu), z_b as the elevation (generally negative) of the base of the ice shelf. During the initialization procedure, to evaluate the specific role of oceanic conditions, we conducted two experiments using PISM: Experiment “S2” replicates the single simulation from LOW21 that used the best fit parameter set (the one minimizing mismatch with observations), employing a thermodynamic parameterization (Eq. 2) to estimate sub-ice shelf melt rates. Experiment (Fig. 3a): experiment “S1” uses the same model configuration—including all parameters, stress balance approximation, resolution, topography, and atmospheric conditions—but replaces the basal melting scheme with observed basal melt rates derived from satellite altimetry (ICESat-1), radar (OIB and ALOS PALSAR), and model outputs (RACMO2), based on Eq. 1. While experiment “S2” replicates the single simulation from LOW21 that used the best-fit parameter set (the one minimizing mismatch with observations), employing a thermodynamic parameterization (Eq. 2) to estimate sub-ice shelf melt rates.




Figure 1: Ocean conditions used in S1. (a) observation of sub-ice shelf basal melt rates ([Rignot et al., 2013](#)); (b) Temperature field beneath ice shelves ([Chambers et al., 2021](#)).




Figure 2: Comparison of sub-ice shelf melt rates between S1 and S2. (a) Sub-ice shelf melt rates derived from the TF-linear parameterization (S2). (b) Difference in basal melt rates used in S1 and S2, with three black boxes highlighting regions of interest: (c) Thwaites Basin, (d) Wilkes Land, and (e) George V Land Terre Adelie.

We applied a “multi-stage” spin-up procedure (Golledge et al., 2015; Lowry et al., 2021) to achieve a pseudo-equilibrium ice-sheet state under constant climate conditions, with a 16 km spatial resolution: (1) a brief 10-year smoothing utilizing the shallow ice approximation, (2) a 250,000-year simulation to allow thermal evolution run with the “no mass” option, which fixes the ice geometry and allows the enthalpy field to reach thermal evolve to equilibrium, (3) a 1,500-year model run incorporating full model physics, including the application of sub-ice shelf melt rates to constrain ice dynamics, and (4) a 65-year historical run to connect initialization and prediction, during which the current ice thickness is reconstructed. Further, based

Based on the initialized model state and the optimal parameter set from S1, we conduct projection experiments from 2015 by turning on or off the sub-grid grounding-line scheme in PISM. The “sub-grid scheme on (SGO) scenario” incorporated sub-grid melt interpolation near grounding lines, accelerating grounding-line retreat in our coarse-resolution model, while the “sub-grid scheme off (SGF) scenario” ignored melt in partially floating cells, yielding more conservative mass loss estimates (Albrecht et al., 2011; Golledge et al., 2015; Nowicki et al., 2020). ~~To~~

In projection, to ensure that differences in projections results originated solely from the model spin-up, we employed the same daily-resolution climate forcing forcings as LOW21 (Lowry et al., 2021), derived from the CMIP5 IPSL-CM5A-MR RCP2.6/8.5 (2015-2100; Barthel et al., 2020; Payne et al., 2021; Nowicki et al., 2021) ~~and~~. Furthermore, we also conducted the experiments to include SSP scenarios for Antarctic ice-sheet evolution, using daily climate forcings from the CMIP6 CNRM-CM6-1 SSP1-2.6/5-8.5 product (Nowicki et al., 2016; Kamworapan et al., 2021; Nowicki et al., 2021) spanning 2015–2100. These daily data were pre-processed by averaging to produce annual

mean forcings, which were then used to drive the ice-sheet model. This setup allowed us to assess and compare Antarctica's contribution to global mean sea-level rise by 2100. The basal melting scheme was parameterized using

the same linear thermodynamic framework for the ice-shelf-ocean boundary layer as that employed in LOW21. ~~This~~
~~specifically, the approach explicitly resolves heat and freshwater exchange processes at the ice-ocean~~
~~interface, defined in Eq. 2, which is driven by oceanic forcing under different~~ the annual-mean spatial distribution and
time series of absolute salinity and temperature fields derived from the RCP/SSP scenarios ~~from~~ (Jourdain et al., 2020;
Nowicki et al., 2020) for 2015 to 2100;

$$S_{proj} = \rho_{sw} c_m \gamma_T F_{melt} (T_{s_proj} - T_{f_proj}) / (L_i \rho_i) \quad (4)$$

here, S_{proj} , T_{s_proj} , and T_{f_proj} refer to values calculated by projected ocean forcings. Therefore, following the two
basal melt schemes that yielded distinct initial ice-sheet states and produced two projections (Fig. 3b): the S1-based
projection simulates future Antarctic evolution under RCP/SSP scenarios, starting from the S1-initialized state using
observed basal melt rates. While the S2-based projection utilizes the RCP-based sea-level contributions provided by
LOW21, which were generated using the same parameter configuration as S2 reproduction initialization.




Figure 3: Overview of model initialization and projection processes. The schematic summarizes the model setup during (a)
spin-up and (b) projection, using observed basal melt rates together with ice-shelf basal temperature (S1, orange box in a) and
Southern Ocean temperature and salinity (S2, yellow box in a; LOW21). The projections initialized from these two states are
denoted as S1-based and S2-based (LOW21), respectively.

3 Model Initialization Results

3.1 Comparison with the case of different sub-ice shelf melt rates

165 We validated the simulated ice ~~thickness and ice surface velocity results-sheet geometry~~ from both S1 and S2 using
observational datasets (BedMachine v.3; MEaSURES Phase-Based Antarctica Ice Velocity Map v.1). The ~~difference~~
~~in results show that both experiments achieve comparable accuracy against observations: the~~ root mean square
~~error/errors~~ (RMSE) for ~~the two experiments, derived from comparison against observation, is 2 m for~~ ice thickness
170 ~~are approximately 89 m (S1) and 3 m y⁻¹ 87 m (S2), differing by 2 m; the RMSE for ice surface velocity- are roughly~~
~~270 m y⁻¹ (S1) and 267 m y⁻¹ (S2), differing by 3 m y⁻¹.~~ Figure ~~34~~ shows the differences between model results and
observations: the left column displays discrepancies for S1 simulations, while the right column shows those for S2.
The comparison shows that the mass distribution and ice flow of S1 closely match those of S2. To better highlight the
differences between the two experiments in ice thickness and velocity over Thwaites Glacier, we selected a
representative transect where discrepancies were most pronounced. Along this transect, we compare the grounding-
175 line positions, ice thickness, and surface velocity profiles between S1 and S2.

We also compared the results of two experiments against those from the Antarctic ice-sheet model initialization
(initMIP-Antarctica) that employed the PISM (Seroussi et al., 2019). Both studies align with ensemble trends in ice
mass, ice sheet area, ice shelf area, and potential sea-level contributions. S1 simulations exhibit minor differences in
total ice sheet mass (-6% to +11%) and ice area (-7% to -1%) compared to initMIP-Antarctica. Notably, deviations in
180 potential sea-level contributions are more pronounced (-17% to -2%), while ice shelf area discrepancies reach 44%
relative to the DMI_PISM simulation (Fig. ~~45~~). Overall, despite minor differences in other metrics between our two
experiments and the initMIP-Antarctica ensemble simulations, the spin-up ice volumes (potential sea-level
contributions) in both S1 ($25.81 \times 10^6 \text{ km}^3$) and S2 ($25.77 \times 10^6 \text{ km}^3$) exhibit close agreement with the observed total
ice volume (BedMachine v.3; $26(\pm 0.4) \times 10^6 \text{ km}^3$). This validates the robustness of our initialization configuration and
185 ~~lends us gives~~ confidence for future projection experiments.





Figure 34: Comparing simulated initial state with observations. Modeled ice thickness (m) and surface ice velocity (m y^{-1}) at the end of spin-up. The left column shows ice thickness and ice surface velocity results from S1, alongside their difference from observation (Morlighem et al., 2019; Mougnot et al., 2019). The right column shows the corresponding results from S2, sharing common color bars with S1.

190



195 **Figure 45: Percentage deviations in steady-state metrics (ice mass, ice sheet area, ice shelf area, potential sea-level contribution) relative to the initMIP-Antarctica PISM-based ensemble.** The vertical axis represents the percentage deviation between the results of S1, S2, and the simulations from participating institutions (*_PISM, where * denotes the institution abbreviation).

3.2 Differences in Marine Ice-sheet regions


200 There are significant velocity differences in three marine-based regions characterized by retrograde bed slopes (Fig 56): Thwaites Basin (TB) in the West Antarctica (WAIS), Wilkes Land (WL), and George V Land–Terre Adelie (GVL) in the East Antarctica (EAIS). These regions are particularly susceptible to MISI due to their subglacial topography (Joughin et al., 2014; Mengel & Levermann, 2014; Greenbaum et al., 2015).

In the Thwaites Basin, the observed sub-ice shelf melt rates used in S1 (reaching 17.7 m y^{-1} beneath Thwaites ice shelf, Fig. 1) exceed S2’s parameterized values by approximately 5 m y^{-1} (Fig. 2). This higher basal melting weakens the

205 ice-shelf buttressing effect and accelerates the grounded ice flow, with a corresponding 74 m y^{-1} RMSE difference
from S2 (Fig. 5f6f). Compared to S2, this results in around 40 m more ice thinning near the grounding line and an
approximately 30 km more grounding-line retreat (Fig. 78), demonstrating an intrinsic connection within the ice-sheet
system. This systemic response is further evidenced by widespread thickening upstream, with a mean anomaly of 49.5
m. The ice volume above flotation in S1 shows a 5.5-fold bias reduction (-0.59%, 1.19 m SLE) compared to S2 (-
210 3.28%, 1.16 m SLE), aligning closely with observations ($1.20 \pm 0.02 \text{ m SLE}$, Table 1).

In the Wilkes Land, the Totten Glacier exhibits increased ice flow under observed melt rates, yielding a 44 m y^{-1} lower
RMSE in S1 relative to S2 (Fig. 5g6g), leading to regional mean ice thinning of 38.5 m. The faster flow of Totten
Glacier strengthens lateral resistance along its boundaries with adjacent glaciers, subsequently reducing ice flux into
the Voyeykov and Moscow Ice Shelves (Gagliardini et al., 2010; Van Der Veen et al., 2014). This dynamic response
215 is consistent with the simulated mean thickness anomaly of +39.2 m across these regions. The ice volume above
flotation bias in WL decreases to -4.61% (6.63 m SLE) in S1 results, compared to -5.14% (6.59 m SLE) in S2,
achieving a 10% improvement relative to the observed $6.95 \pm 0.09 \text{ m SLE}$ (Table 1).

In the George V Land–Terre Adelie, the enhanced flow of the Ninnis Ice Shelf in S1 results in increased ice discharge
and regional mean thinning (33.7 m). Conversely, the Cook and Mertz Ice Shelves and their upstream glaciers
220 experience reduced ice flux, causing regional ice thickness to increase by 25.5 m on average relative to S2. S1
simulations demonstrate a reduced ice volume above flotation bias of -5.23% (3.35 m SLE) in WL, outperforming -
5.42% (3.34 m SLE) of S2 and reflecting closer agreement with the observed $3.53 \pm 0.04 \text{ m SLE}$ (Table 1).



225 **Figure 56: Comparison of spin-up ice thickness and velocity misfits between S1 and S2.** (a–c) Ice thickness differences (S1 relative to S2) in the TB, WL, and GVL, respectively. (f–h) Ice surface velocity misfits in the TB, WL, and GVL, respectively. The difference in root mean square error (ΔRMSE) between S1 and S2, compared to observations ($\Delta\text{RMSE} = \text{RMSE}_{S1} - \text{RMSE}_{S2}$). (d) and (e) present the deviations in ice thickness and surface velocity between two experiments, with three black boxes highlighting regions showing the most significant discrepancies. Grounding lines: S2 (blue), S1 (red), and observed data (black) from BedMachine v.3 (Morlighem et al., 2019). The profile line locations corresponding to Fig. 78 are in Thwaites Glacier: (a), (d) brown; (f) white.

230

A comparison of simulation results between S1 and S2 for three marine ice sheet basins (TB, WL, and GVL; Table 1) reveals that the ice-sheet model driven by observed sub-ice shelf melt rates achieves slightly better alignment with observations. Although the RMSE of ice surface velocity in the TB shows an increase of 74 m y^{-1} in S1 compared to S2 (Fig. 56f), the bias in ice volume above flotation decreases by approximately 2.8%, while the biases for WL and GVL are reduced by 0.5% and 0.2% (Table 1), respectively.

235


Table 1: Ice volume above flotation (m SLE) in three marine ice sheet basins after spin-up, simulated at 16 km resolution. “S1” denotes the experiment using observed basal melt rates based on Eq. 1, while “S2” refers to the simulation that replicates LOW21 using the thermodynamic parameterization in Eq. 2.

Basins	Observation	S1	Misfit	S2	Misfit
Thwaites Basin (TB)	1.20 (± 0.02)	1.19	-0.59%	1.16	-3.28%
Wilkes Land (WL)	6.95 (± 0.09)	6.63	-4.61%	6.59	-5.14%
George V Land Terre Adelie (GVL)	3.53 (± 0.04)	3.35	-5.23%	3.34	-5.42%

3.3 Marine Ice Sheet dynamics during model spin-up

240 In the WAIS, particularly for glaciers adjacent to the Amundsen Sea Embayment, the subglacial bedrock topography lying below sea level amplifies the sensitivity to ocean-driven forcings (Pritchard et al., 2012). Previous studies stated that the Aurora Subglacial Basin in WL and the Wilkes Subglacial Basin in GVL, in the EAIS, are characterized by extensive sedimentary basins that are highly susceptible to warming ocean conditions (Aitken et al., 2014; Frederick et al., 2016; Noble et al., 2020). These basins are also subject to an active subglacial hydrology process (Wright et al., 245 2012), and evidence of ocean-driven dynamic ice loss has been documented along the ice-sheet margin (Li et al., 2016). The interplay between oceanic forcing, subglacial hydrology, and sedimentary geology significantly influences ice-sheet dynamics in these regions.

In the S1 experiment, enhanced oceanic forcing (Fig. 2), which is represented by higher basal melt rates, intensifies ice-shelf basal melting, leading to geometric thinning and reduced buttressing effect of upstream ice flow (Gudmundsson, 2013; Miles et al., 2022). This triggers grounding-line retreat, which accelerates ice flow, amplifies strain rates, and enhances dissipative heating (Cuffey &and Paterson, 2010; Dawson et al., 2022), thereby increasing temperatures at the basal ice layer (Fig. 6a7a-c). It then promotes basal melting (Fig. 6a7d-f) while reducing ice viscosity via thermal softening, collectively facilitating enhanced deformation and potentially increasing ice-sheet destabilization (Hindmarsh, 2006; Adams et al., 2021). Additionally, subglacial meltwater lubricates the ice-bed 250 interface, reducing basal friction through decreased effective pressure and accelerating ice flow (Fig. 6g7g-l). Enhanced sliding generates additional strain heating (Garbe et al., 2020), which promotes further basal melting and meltwater production. In this positive feedback process, termed the basal thermal-hydrological feedback, elevated basal water content persistently reduces resistance, thereby facilitating ice sliding and ultimately leading to ice thinning (Fowler et al., 2001; Clarke, 2005; van Pelt &and Oleremans, 2012; Zhao et al., 2025).



260

Figure 67: Spatial distribution of misfits (S1 relative to S2) in basal ice temperature field (Kelvin) (a-c), basal till water content (m) (d-f), basal friction (k Pa) (g-i), and basal velocity (m y^{-1}) (j-l) in the three basins. The blue and red lines indicate the grounding-line positions for S2 and S1, respectively, while the black line represents the observed grounding line from BedMachine v.3.

265 3.4 Grounding line location comparison

The impact of sub-ice shelf melt rates on ice-sheet initialization can also be seen at the locations of the grounding line. Particularly, in the WAIS, the retrograde bed topography amplifies the susceptibility to MISI (Pritchard et al., 2012;

Ritz et al., 2015), rendering it highly responsive to ocean forcing. During spin-up, these elevated basal melt rates (Fig. 2) trigger MISI more easily, causing the grounding line on retrograde bedrock to retreat continuously until reaching a new steady state (Rignot et al., 2019; Li et al., 2022). Cross-sectional analysis of Thwaites Glacier (Fig. 78) demonstrates this mechanism, with enhanced basal melting, causing an approximately 30 km grounding-line retreat from its stabilized position (S2, dashed grey line) to a new quasi-stable state (S1, solid grey line). The retreat in S1 increases ice discharge due to the reduced ice-shelf buttressing effect, resulting in roughly 40 m ice thinning proximal to the grounding line and an anomalous nearly twofold acceleration in ice surface velocity compared to S2 results. This feedback highlights how ocean-forced basal melting propagates through ice-sheet dynamics processes to alter initial ice geometry.





Figure 78: Comparison between S1 and S2 along the Thwaites Basin transect after spin-up. Ice elevation (blue line), ice surface velocity (brown line), and grounding-line positions for S1 and S2 are indicated by the grey solid and dashed lines, respectively. Sea level: black dashed line.

In fact, the grounding-line position varies across different ice streams depending on topography, neighbouring ice shelf basal melt rates, and ice velocities (Martin et al., 2011; Pittard et al., 2022). Consequently, the discrepancies between observed and simulated grounding-line positions differ across various regions of the AIS. For instance, the grounding line of the Siple Coast on the west side of the Ross Ice Shelf in S1 agrees with S2 but extends further to the nearshore compared to the observation (Fig. 546d-e). This discrepancy likely arises from a self-limiting, stabilizing mechanism inherent to prograde slopes. As the grounding-line retreats into shallower bedrock, the ice thins and ice flux decreases; this leads to ice re-accumulation that prompts grounding-line readvancement, creating reversible shifts around an equilibrium point (Huybers et al., 2017). Notably, under a consistent model parameter configuration, the inclusion of observed sub-ice shelf melt rates did not significantly alter the steady-state grounding-line migration position across the whole AIS, except within three marine-based regions.

4 Model Projection Results

4.1 Global mean sea-level contribution from AIS

Prognostic simulations (2015–2100) show that the divergent initial ice-sheet states of S1 and S2 lead to markedly different sea-level contributions across the AIS, even under identical climatic forcings and basal melting scheme (Figs. 8, 9, 10). Specifically, S1-based projections of a 0.20–0.52 m SLE total AIS contribution exceed the 0–0.32 m SLE range of the LOW21 ensemble projections (which includes predicted results from S2) by roughly 0.18 m SLE, representing a ~57% increase (Fig. 89). This discrepancy is driven primarily by enhanced ice loss from the West Antarctica (0.29–0.34 m SLE, Table 2), where high sub-ice shelf melt rates triggered MISI more readily, whereas the East Antarctica and the Antarctic Peninsula (AP) show minimal sea-level contributions by 2100, i.e., 0.01–0.02 m SLE and 0.0011–0.0045 m SLE, respectively (Table 2). However, the S1-based projections of AIS contributions to sea-level rise from 2015 to 2075 exhibit no significant dependence on emission scenarios, with substantial overlap in prediction ranges (Fig. 8e9c). This is consistent with the hysteretic response of ice-sheet dynamics, meaning that the ice sheet's state in the near-term (2015–2075) is largely determined by historical forcing, masking the influence of divergent future scenarios (Garbe et al., 2020). By 2100, the mean AIS contributions to sea-level rise based on S1 under SSP 5-8.5 reach 0.36 m SLE—12.5% higher than the RCP 8.5 equivalent (0.32 m SLE)—with RCP high-emission projections even matching SSP low-scenario results (Fig. 8e9d, Table 2). These differences between RCP 8.5 and SSP 5-8.5 projections are largely due to the SSP scenarios in CMIP6 climate models simulating higher warming magnitudes (averaging +0.14–0.25 °C) than RCP scenarios in CMIP5 at equivalent radiative forcing (Tokarska et al., 2020; Wyser et al., 2020; Rounce et al., 2023). Consequently, under anthropogenic warming, the sea-level commitment of AIS under SSP high-risk scenarios demands heightened scientific attention.







Figure 89: Ice sheet thickness misfits and projected contribution of the AIS to sea-level rise (m SLE). Spatial differences in the projected mean ice thickness between the multi-scenario (RCP 2.6 and RCP 8.5) ensemble means based on S1 and S2 in 2050 (a) and 2100 (b). (c) Predicted sea-level rise for “SGO scenario” to “SGF scenario” simulations under four scenarios (color shading) and mean values (color lines). The dashed lines represent projections from the S2 initial state—a set of results from the LOW21 ensemble projections: red for RCP 8.5, blue for RCP 2.6. (d) Projected contributions to sea-level rise (SLR) by 2100 based on S1, compared to other studies: CHU13 (Church et al., 2013), GOL15 (Golledge et al., 2015), DP16 (DeConto & Pollard, 2016), EDW19 (Edwards et al., 2019), SER20 (Seroussi et al., 2020), LOW21 (Lowry et al., 2021), VAN23 (van der Linden et al., 2023), KLO24 (Klose et al., 2024), and COU24 (Coulon et al., 2024).


4.2 Regional Contributions to Global Mean Sea-Level Rise

To explore the spatially variable response of the AIS, we partitioned the ice sheet into seven sectors based on their adjacency to surrounding oceans (Fig. 910). Following the IMBIE (Zwally et al., 2012), we further subdivided the East Antarctica into the East Indian Ocean (EIO) and West Indian Ocean (WIO) sectors. This regional breakdown reveals stark contrasts in the mechanisms and magnitudes of sea-level rise across the WAIS, EAIS, and AP (Fig. 910, Table 2), highlighting their distinct sensitivities to climate forcings. The WAIS emerges as the primary driver of AIS-related sea-level rise, contributing 0.29–0.34 m SLE by 2100. A mechanism similar to that observed in recent decades may be responsible for the projected mass loss. Specifically, anthropogenic warming could alter shelf-break wind patterns over the Amundsen and Bellingshausen Sea (AS/BS) embayments (Holland et al., 2019; Noble et al., 2020), potentially facilitating greater intrusion of warm water and intensifying ice melting beneath ice shelves (Dinniman et al., 2016; Noble et al., 2020; Li et al., 2023). The resulting reduction in ice-shelf buttressing accelerates ice discharge, which explains the dominance of the AS and BS sectors—accounting for ~55% of total WAIS mass loss in our projections (Table 2).

335 The EAIS presents a more complex picture, with a net contribution of 0.01–0.02 mSLE by 2100. While the integrated signal is small, it masks pronounced regional heterogeneity in mass changes. The net mass gain in the West Indian Ocean (WIO) sector shown in the S1-based projection may be linked to enhanced moisture transport from the Southern Ocean, a mechanism consistent with observational trends (Boening et al., 2012) that would promote increased ice surface accumulation. However, this marginal gain is counteracted by ice dynamical adjustments within the East
340 Indian Ocean (EIO) sector, specifically across the WL, where enhanced oceanic thermal forcing drives accelerated ice mass loss from the dynamically vulnerable Totten Glacier (Konrad et al., 2018). This contrast between surface mass gains and ice-dynamic losses underscores the spatially heterogeneous response of EAIS, modulated by regional bathymetry and ocean-driven melt.

The AP plays a comparatively minor role in sea-level rise, with contributions ranging from 0.0011–0.0043 m SLE by
345 2100. Peak mass loss occurs between 2075 and 2079, reaching 0.0061 m SLE under RCP 8.5 and 0.0045 m SLE under SSP 5-8.5, followed by a gradual decline (Fig. 910). The intensification of polar westerly winds could enhance snowfall in the northern AP, which may partially offset warming-induced ice discharge and thus generate a negative feedback that suppresses AIS mass loss (Goodwin et al., 2016). Given its limited ice volume, however, the AP's overall impact on sea-level rise remains marginal. The findings underscore the divergent climate responses of the
350 EAIS, AP, and the WAIS. While the EAIS and AP exhibit mass gain or loss depending on the balance between accumulation and ablation, the WAIS is primarily driven by dynamic mass loss resulting from changes in oceanographic conditions.





355

Figure 910: The mean contribution of the AIS seven sectors to sea-level rise from 2015 to 2100 (m SLE). The solid and dashed lines represent projections from the S1 and S2 initial states, respectively, under different climate scenarios, with the S2 predicted results being part of the LOW21 ensemble projections (red for RCP 8.5, blue for RCP 2.6).

360

Table 2: Sea-level contribution (m SLE) of Antarctic Ice Sheet Basins by 2100. The confidence intervals represent the range of sea-level contribution from the “SGO scenario” to the “SGF scenario” simulation across different RCP/SSP scenarios; the single value denotes the mean value of this range.

Region	RCP 2.6	RCP 8.5	SSP 1-2.6	SSP 5-8.5
Bellingshausen Sea	0.1138	0.1102	0.1256	0.1297
(BS)	(0.0857, 0.1419)	(0.0813, 0.1391)	(0.0982, 0.1530)	(0.1064, 0.1585)
Amundsen Sea (AS)	0.0499	0.0493	0.0701	0.0700
	(0.0405, 0.0593)	(0.0386, 0.0599)	(0.0553, 0.0850)	(0.0548, 0.0968)
Ross Sea (RS)	0.0034	0.0247	0.0163	0.0193
	(-0.0089, 0.0157)	(-0.0009, 0.0503)	(-0.0107, 0.0435)	(-0.0048, 0.0552)
Weddell Sea (WS)	0.1260	0.1249	0.1005	0.1025

	(0.0759, 0.1762)	(0.0734, 0.1764)	(0.0542, 0.1467)	(0.0582, 0.1638)
The West Antarctic ice sheet (WAIS)	0.2931 (0.1932, 0.3931)	0.3090 (0.1924, 0.4257)	0.3126 (0.197, 0.4282)	0.3444 (0.2146, 0.4743)
West Indian Ocean (WIO)	-0.0330 (-0.0448, -0.0211)	-0.0363 (-0.0560, -0.0166)	-0.0389 (-0.0531, -0.0247)	-0.0379 (-0.0471, -0.0227)
East Indian Ocean (EIO)	0.0423 (0.0269, 0.0578)	0.0473 (0.0277, 0.0669)	0.0467 (0.0284, 0.0650)	0.0559 (0.0467, 0.0867)
The East Antarctic ice sheet (EAIS)	0.0093 (0.0058, 0.0130)	0.0110 (0.0109, 0.0111)	0.0078 (0.0037, 0.0119)	0.0180 (0.0240, 0.0396)
Antarctica Peninsula (AP)	0.0043 (0.0024, 0.0062)	0.0028 (0.0007, 0.0049)	0.0022 (0.00004, 0.0044)	0.0011 (-0.0021, 0.0034)
The Antarctic ice sheet (AIS)	0.3067 (0.2014, 0.4123)	0.3228 (0.2042, 0.4415)	0.3226 (0.2007, 0.4445)	0.3635 (0.2365, 0.5173)

4.3 Comparison with previous projections

The S1-based simulation projects AIS contributions to sea-level rise by 2100 under the SSP 5-8.5 scenario diverge significantly from previous estimates, particularly for the WAIS. The Coupled Model Intercomparison Project Phase 6 models (CMIP6, Edwards et al., 2021) employed Gaussian process emulators—statistical approximations built upon ice-sheet simulations for ISMIP6 (Nowicki et al., 2016, 2020) and GlacierMIP Phase 2 (Hock et al., 2019)—to generate sea-level projections. While their ensemble projections suggest that the WAIS contributions range from -0.04 to 0.11 m SLE, S1-based projections show a significantly higher contribution of 0.20–0.47 m SLE. For the EAIS and AP, S1-derived projected sea-level contributions (0.03–0.04 m SLE and -0.0021–0.0034 m SLE, respectively; Table 2) align closely with the emulator results (-0.05–0.06 m SLE and -0.01–0.02 m SLE, respectively), with the WAIS emerging as the dominant divergence source.

Compared to the full ensemble results of ISMIP6 (Ice Sheet Model Intercomparison for CMIP6) Antarctic projections under RCP 8.5 (Seroussi et al., 2020), the S1-based projected sea-level contribution for the WAIS is approximately 0.15 m SLE higher, while the AP shows a slight increase (~0.002 m SLE), and the EAIS exhibits a minor reduction (~0.02 m SLE). The ISMIP6-Antarctica projections ~~improve~~ provide a more comprehensive representation of potential Antarctic sea-level contribution under climatic forcings, with the parameterizations of oceanic conditions into basal melt rates being the dominant source of uncertainty (Seroussi et al., 2020). However, these experiments cannot identify the specific physical mechanisms behind the inter-model differences. Although a key limitation of our single model experiment is its reliance on PISM-specific parameterizations, which restrict the range of projected sea-level contributions and provide limited statistical uncertainty. Nevertheless, by comparing observed and parameterized basal melt rates as represented in model simulations under a consistent single-model framework, our experiments identify the specific regions and dynamic mechanisms underlying the ISMIP6 projection uncertainty associated with the representation of oceanic conditions.

385 These results demonstrate significant deviations in the WAIS sea-level contributions compared to prior studies,
aligning with the disparity between projections based on the S1 and S2 initial states (Fig. 910). The pronounced
discrepancies in the WAIS primarily stem from its vulnerability to oceanic forcing and complex bed topography,
projecting an average ice thickness decrease of 50 m by 2100 in S1-based results relative to S2-based (Fig. 89). While
S2 experiment used parameterized melt rates in model spin-up, S1 incorporation of observational values reveals
390 enhanced basal melting in the critical WAIS regions like TB (Fig. 2), accelerating dynamic ice loss through processes
such as MISI. In contrast, the EAIS regions (WL and GVL) exhibit minor variations, contributing little to sea-level
rise during projection periods (2050 and 2100; Fig. 89). This regional contrast highlights the WAIS's dominant role
in creating divergence from earlier projections, demonstrating its heightened vulnerability to ocean-induced melt rates
at the initialization.

5 Model Uncertainties

395 The elevated sub-ice shelf melt rates in experiment S1 (Fig. 2), notably in the Thwaites and Pine Island shelves, modify
the initial AIS state. Through an iterative spin-up process, the model iteratively adjusts key variables such as basal ice
temperature field, basal friction coefficients, and grounding-line positions to minimize discrepancies between
simulation and observation. While the ice-sheet geometries for S2 and S1 are initialized with identical spin-up, distinct
sub-ice shelf melt rates produce divergent ice thermodynamic states, causing the ice sheet to follow unique
400 evolutionary trajectories in projection under identical external forcing, thus altering the potential contributions of sea
level rise.

The rationale for using the Rignot et al. (2013) basal melt rates in S1 was that the corresponding ocean thermal forcing
aligns with the 1975–2012 mean state of the Southern Ocean captured in Schmidtko et al. (2014) dataset (used in S2).
However, as these data reflect conditions from approximately a decade ago, they inherently represent a temporal
405 average and do not capture interannual variability in ocean forcing (Adusumilli et al., 2020). Furthermore, Paolo et al.
(2023) also observed a widespread slowdown in ice-shelf thinning across the Amundsen, Bellingshausen, and Wilkes
sectors, attributing it to changes in ocean forcing and internal ice-dynamic feedbacks. Therefore, S1 simulated results
should be interpreted as a response to a steady-state, general ice-shelf basal melting field. Future work should
incorporate time-evolving melt rates to better constrain the sensitivity of the AIS to oceanic variability on interannual
410 to decadal timescales.

Compared to other prior studies, our sea-level projections from the S1 differ due to variations in ice-sheet model
configurations, including model resolution, ice dynamics (particularly stress balance schemes), represented physical
processes (calving, hydrology, or bedrock uplift), and initialization methods (data assimilation or spin-up) (Seroussi
et al., 2019; Levermann et al., 2020; Klose et al., 2024). Of these factors, the parameterizations of ice melt dynamics
415 contribute most significantly to the uncertainty in sea-level estimates, surpassing those from climate forcing,
initialization methods, or the selected physical processes. Given this dominance, ice-model-related uncertainties
prevail throughout the entire simulation period (Seroussi et al., 2019, 2023). Therefore, continual model improvement
and further exploration of the broader parameter space covered by initial state ensembles are essential to reduce

420 uncertainties in future projections of dynamic mass loss from the AIS (Favier et al., 2019; Coulon et al., 2024; Klose et al., 2024).

Notably, the present-day AIS may not have been in a steady-state during the observational period (Martinet et al., 2011). While this inference is primarily based on model–observation discrepancies, it may also be influenced by uncertainties inherent in the validation datasets. For example, the BedMachine v3 dataset relies on approximate calculations in regions such as ice-free land, ocean bathymetry, and cavities under ice shelves, potentially introducing spatial biases in thickness estimates (Morlighem et al., 2019). Similarly, the MEaSURES velocity inevitably contains errors in flow direction derived from phase data and speckle tracking during SAR data processing (Mouginot et al., 2019). The apparent model–data mismatch reflects both a non-steady-state of AIS and the challenge of validating model simulations against uncertain modern records, which underscores the need for more accurate and extensive observations to better constrain ice-sheet models and improve the reliability of sea-level projections (Seroussi et al., 430 2020, 2023). Moreover, global climate models exhibit significant differences in projected global temperature increases, which in turn affect ice dynamics (Golledge et al., 2015; Schlegel et al., 2018; Klose et al., 2024). High-sensitivity climate models within the CMIP6 ensemble, such as IPSL-CM6A-LR (4.6°C), UKESM1-0-LL (5.3°C), and CESM2-WACCM (4.8°C), predict substantial warming over Antarctica, potentially driving extensive melting of the WAIS.

6 Conclusions

435 The ice-sheet model was initialized using two different basal melting schemes: experiment S1 used observed sub-ice shelf melt rates, while S2 employed the TF-linear parameterization to replicate the LOW21 study. Following spin-up, the modeled ice geometry is consistent with observations in two experiments. However, S1 simulated results reveal notable regional variations in ice-sheet dynamics across three marine ice-sheet regions compared to S2: the Thwaites Basin in the West Antarctica, Wilkes Land, and George V Land–Terre Adelie in the East Antarctica. In Thwaites 440 Basin, elevated sub-ice shelf melt rates progressively trigger MISI, driving grounding-line retreat that significantly weakens the ice-shelf buttressing effect for upstream glaciers. Experiment S1 demonstrates 3 m ice thickness discrepancies and 74 m y⁻¹ ice velocity deviations compared to S2 results under observational validation. Variations in ocean forcing conditions in Wilkes Land and George V Land–Terre Adelie may alter the thermomechanical features at the grounded ice sheet, which then induce dynamic adjustments, causing approximately 6 m and 44 m y⁻¹ differences 445 in ice thickness and surface velocity between S1 and S2, respectively.

Despite identical model configurations and future climate scenarios, projections from the S1 initial state estimate a 57% higher sea-level contribution (~0.18 m SLE) by 2100 compared to the LOW21 ensemble results, which includes the projection based on S2. This divergence stems from the different treatment in prescribing sub-ice shelf melt rates during the PISM spin-up. The majority contributor to this SLE discrepancy stems from the Amundsen Sea sector in 450 the West Antarctica, a region typical of MISI, which also aligns with comparisons to other previous model projection results. In future modeling efforts, we suggest further efforts in investigating the sensitivity of the Antarctic ice sheet model initializations to critical environmental factors before conducting fully prognostic AIS simulations, to better constrain the projected ranges of global sea-level rise.

Code and Data Availability

455 The Parallel Ice Sheet Model is freely available as open-source code from the PISM GitHub repository (<https://github.com/pism/pism>). Bedrock topography and ice thickness data are from the MEaSURES BedMachine Antarctica, Version 3 compilation, available at <https://nsidc.org/data/nsidc-0756/versions/3>. Air temperature, precipitation, and geothermal heat flux inputs were taken from the ALBMAP version 1 compilation and can be downloaded from <http://doi.pangaea.de/10.1594/PANGAEA.734145>. Ice surface velocity used in validation may be
460 obtained from MEaSURES Phase-Based Antarctica Ice Velocity Map, Version 1, available at <https://nsidc.org/data/nsidc-0754/versions/1>. The forcing data under RCP and SSP scenarios were sourced from the dataset published by Nowicki et al. (2021). The data preprocessing tool used is the publicly available scripts `pism-ais` (<https://github.com/pism/pism-ais>).

Author contribution

465 FG and TZ conceived and designed this experiment. FG performed data curation. QS, HW, and TZ acquired funding. HW provided resources. FG and TZ conducted the experiments. FG performed simulations. QS, LJ, and YL performed validation. CKS, YA, and XZ conducted the visualization. FG wrote the original manuscript draft, and all authors contributed to reviewing and editing the manuscript.

Competing interests

470 There are no real or perceived conflicts of interest for any author.

Acknowledgments

We express our heartfelt gratitude to Prof. Lowry for his invaluable mentorship and significant contributions to this study. His provision of critical simulation datasets and expert guidance throughout the research process has been indispensable. We also sincerely thank Prof. Rignot for generously sharing observational data on Antarctic sub-ice
475 shelf melt rates, which greatly enriched our work. Additionally, we extend our sincere appreciation to the Parallel Ice Sheet Model team for their continuous support and for developing and maintaining the PISM, which was essential to the success of this research. We ~~also thank~~ are grateful to the editor, Alexander Robinson, and to the three reviewers ~~co-reviewers Muruganandham Shivaprakash~~ Alexander Robel, Shivaprakash Muruganandham and an anonymous referee, for their help that significantly enhanced in significant improvements of this paper. this manuscript.

480 Financial support

This work is supported by the National Natural Science Foundation of China (42374042, 42271133, and 42374045) and the Natural Science Foundation of Wuhan (2024040701010065).

References

- 485 Adams, C. J. C., Iverson, N. R., Helanow, C., Zoet, L. K., & Bate, C. E. (2021). Softening of Temperate Ice by Interstitial Water. *Frontiers in Earth Science*, 9. <https://doi.org/10.3389/feart.2021.702761>, 1-11. <https://doi.org/10.3389/feart.2021.702761>, 2021.
- Adusumilli, S., Fricker, H. A., Medley, B., Padman, L., & Siegfried, M. R. (2020). Interannual variations in meltwater input to the Southern Ocean from Antarctic ice shelves. *Nature Geoscience*, 13(9), 616-620. <https://doi.org/10.1038/s41561-020-0616-z>, 2020.
- 490 Aitken, A. R. A., Young, D. A., Ferraccioli, F., Betts, P. G., Greenbaum, J. S., Richter, T. G., Roberts, J. L., Blankenship, D. D., & Siegert, M. J. (2014). The subglacial geology of Wilkes Land, East Antarctica. *Geophysical Research Letters*, 41(7), 2390-2400. <https://doi.org/10.1002/2014gl059405>, 2014.
- Albrecht, T., Martin, M., Haseloff, M., Winkelmann, R., & Levermann, A. (2011). Parameterization for subgrid-scale motion of ice shelf calving fronts. *The Cryosphere*, 5(1), 35-44. <https://doi.org/10.5194/tc-5-35-2011>
- 495 ~~Albrecht, T., Winkelmann, R., & Levermann, A. (2020). Glacial-cycle simulations of the Antarctic Ice Sheet with the Parallel Ice Sheet Model (PISM) – Part 2: Parameter ensemble analysis. *The Cryosphere*, 14(2), 633-656. <https://doi.org/10.5194/tc-14-633-2020>, 2020.~~
- ~~Albrecht, T., Martin, M., Haseloff, M., Winkelmann, R., and Levermann, A.: Parameterization for subgrid-scale motion of ice-shelf calving fronts, *The Cryosphere*, 5, 35-44, <https://doi.org/10.5194/tc-5-35-2011>, 2011.~~
- 500 Alevropoulos-Borrill, A. V., Nias, I. J., Payne, A. J., Golledge, N. R., & Bingham, R. J. (2020). Ocean-forced evolution of the Amundsen Sea catchment, West Antarctica, by 2100. *The Cryosphere*, 14(4), 1245-1258. <https://doi.org/10.5194/tc-14-1245-2020>, 2020.
- ~~Beckmann, A. and Goosse, H. (2003). A parameterization of ice shelf-ocean interaction for climate models, *Ocean Model.*, 5(2), 157-170. [https://doi.org/10.1016/S1463-5003\(02\)00019-7](https://doi.org/10.1016/S1463-5003(02)00019-7)~~
- 505 Barthel, A., Agosta, C., Little, C. M., Hattermann, T., Jourdain, N. C., Goelzer, H., Nowicki, S., Seroussi, H., Straneo, F., & Bracegirdle, T. J. (2020). CMIP5 model selection for ISMIP6 ice sheet model forcing: Greenland and Antarctica. *The Cryosphere*, 14(3), 855-879. <https://doi.org/10.5194/tc-14-855-2020>, 2020.
- ~~Beckmann, A. and Goosse, H.: A parameterization of ice shelf-ocean interaction for climate models, *Ocean Modeling*, 5, 157-170, [https://doi.org/10.1016/S1463-5003\(02\)00019-7](https://doi.org/10.1016/S1463-5003(02)00019-7), 2003.~~
- 510 Berdahl, M., Leguy, G., Lipscomb, W. H., Urban, N. M., & Hoffman, M. J. (2023). Exploring ice sheet model sensitivity to ocean thermal forcing and basal sliding using the Community Ice Sheet Model (CISM). *The Cryosphere*, 17(4), 1513-1543. <https://doi.org/10.5194/tc-17-1513-2023>, 2023.
- Bindschadler, R. A., Nowicki, S., Abe-Ouchi, A., Aschwanden, A., Choi, H., Fastook, J., Granzow, G., Greve, R., Gutowski, G., Herzfeld, U., Jackson, C., Johnson, J., Khroulev, C., Levermann, A., Lipscomb, W. H., Martin, M. A., Morlighem, M., Parizek, B. R., Pollard, D., Price, S. F., Ren, D., Saito, F., Sato, T., Seddik, H., Seroussi, H., Takahashi, K., Walker, R., & Wang, W. L. (2013). Ice-sheet model sensitivities to environmental forcing and their use in projecting future sea level (the SeaRISE project). *Journal of Glaciology*, 59(214), 195-224. <https://doi.org/10.3189/2013JoG12J125>, 2013.
- 515 Boening, C., Lebsack, M., Landerer, F., & Stephens, G. (2012). Snowfall - driven mass change on the East Antarctic ice sheet. *Geophysical Research Letters*, 39(24), 1-5. <https://doi.org/10.1029/2012gl053316>, 2012.
- 520 Bueler, E., & Brown, J. (2009). Shallow shelf approximation as a “sliding law” in a thermomechanically coupled ice sheet model. *Journal of Geophysical Research*, 114(F3), 1-21. <https://doi.org/10.1029/2008jf001179>, 2009.
- Bueler, E., Brown, J., & Lingle, C. (2007). Exact solutions to the thermomechanically coupled shallow-ice approximation: effective tools for verification. *Journal of Glaciology*, 53(182), 499-516. <https://doi.org/10.3189/002214307783258396>, 2007.
- 525 Calov, R., & Greve, R. (2017). A semi-analytical solution for the positive degree-day model with stochastic temperature variations. *Journal of Glaciology*, 51(172), 173-175. <https://doi.org/10.3189/172756505781829601>, 2005.
- 530 Chambers, C., Greve, R., Obase, T., Saito, F., & Abe-Ouchi, A. (2021). Mass loss of the Antarctic ice sheet until the year 3000 under a sustained late-21st-century climate. *Journal of Glaciology*, 1-13. <https://doi.org/10.1017/jog.2021.124>, 2021.
- Church, J., Clark, P., Cazenave, A., Gregory, J., Jevrejeva, S., Levermann, A., Merrifield, M., Milne, G., Nerem, R., & Nunn, P. (2013). “Chapter 13: Sea Level Change” in *Climate Change 2013: The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. 2013.
- 535 Clarke, G. K. C. (2005). Subglacial Processes. *Annual Review of Earth and Planetary Sciences*, 33(1), 247-276. <https://doi.org/10.1146/annurev.earth.33.092203.122621>, 2005.

540 Cornford, S. L., Martin, D. F., Payne, A. J., Ng, E. G., Le Brocq, A. M., Gladstone, R. M., Edwards, T. L., Shannon, S. R., Agosta, C., van den Broeke, M. R., Hellmer, H. H., Krinner, G., Ligtenberg, S. R. M., Timmermann, R., ~~&and~~ Vaughan, D. G. ~~(2015)~~.: Century-scale simulations of the response of the West Antarctic Ice Sheet to a warming climate. *The Cryosphere*, 9(4), 1579-1600. <https://doi.org/10.5194/tc-9-1579-2015>, 2015.

545 Cornford, S. L., Seroussi, H., Asay-Davis, X. S., Gudmundsson, G. H., Arthern, R., Borstad, C., Christmann, J., Dias dos Santos, T., Feldmann, J., Goldberg, D., Hoffman, M. J., Humbert, A., Kleiner, T., Leguy, G., Lipscomb, W. H., Merino, N., Durand, G., Morlighem, M., Pollard, D., Rückamp, M., Williams, C. R., ~~&and~~ Yu, H. ~~(2020)~~.: Results of the third Marine Ice Sheet Model Intercomparison Project (MISMIP+) ~~(+)~~. *The Cryosphere*, 14(7), 2283-2301. <https://doi.org/10.5194/tc-14-2283-2020>, 2020.

550 Coulon, V., Klose, A. K., Kittel, C., Edwards, T., Turner, F., Winkelmann, R., ~~&and~~ Pattyn, F. ~~(2024)~~.: Disentangling the drivers of future Antarctic ice loss with a historically calibrated ice-sheet model. *The Cryosphere*, 18(2), 653-681. <https://doi.org/10.5194/tc-18-653-2024>, 2024.

555 Cuffey, K. M., ~~&~~ ~~and~~ Paterson, W. S. B. ~~(2010)~~.: *The physics of glaciers*. Academic Press, Burlington, MA, USA, 704 pp., ISBN 9780123694614, 2010.

560 Dawson, E. J., Schroeder, D. M., Chu, W., Mantelli, E., ~~&and~~ Seroussi, H. ~~(2022)~~.: Ice mass loss sensitivity to the Antarctic ice sheet basal thermal state. *Nature Communications*, 13(4), <https://doi.org/10.1038/s41467-022-32632-2>, 2022.

565 DeConto, R. M., ~~&~~ ~~and~~ Pollard, D. ~~(2016)~~.: Contribution of Antarctica to past and future sea-level rise. *Nature*, 531(7596), 591-597. <https://doi.org/10.1038/nature17145>, 2016.

570 Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T. M., Ligtenberg, S. R. M., van den Broeke, M. R., ~~&and~~ Moholdt, G. ~~(2013)~~.: Calving fluxes and basal melt rates of Antarctic ice shelves. *Nature*, 502(7469), 89-92. <https://doi.org/10.1038/nature12567>, 2013.

575 Dinniman, M., Asay-Davis, X., Galton-Fenzi, B., Holland, P., Jenkins, A., ~~&and~~ Timmermann, R. ~~(2016)~~.: Modeling Ice Shelf/Ocean Interaction in Antarctica: A Review. *Oceanography*, 29(4), 144-153. <https://doi.org/10.5670/oceanog.2016.106>, 2016.

580 Edwards, T. L., Brandon, M. A., Durand, G., Edwards, N. R., Golledge, N. R., Holden, P. B., Nias, I. J., Payne, A. J., Ritz, C., ~~&and~~ Wernecke, A. ~~(2019)~~.: Revisiting Antarctic ice loss due to marine ice-cliff instability. *Nature*, 566(7742), 58-64. <https://doi.org/10.1038/s41586-019-0901-4>, 2019.

585 Edwards, T. L., Nowicki, S., Marzeion, B., Hock, R., Goelzer, H., Seroussi, H., Jourdain, N. C., Slater, D. A., Turner, F. E., Smith, C. J., McKenna, C. M., Simon, E., Abe-Ouchi, A., Gregory, J. M., Larour, E., Lipscomb, W. H., Payne, A. J., Shepherd, A., Agosta, C., Alexander, P., Albrecht, T., Anderson, B., Asay-Davis, X., Aschwanden, A., Barthel, A., Bliss, A., Calov, R., Chambers, C., Champollion, N., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Felikson, D., Fettweis, X., Fujita, K., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huss, M., Huybrechts, P., Immerzeel, W., Kleiner, T., Kraaijenbrink, P., Le Clec'h, S., Lee, V., Leguy, G. R., Little, C. M., Lowry, D. P., Malles, J. H., Martin, D. F., Maussion, F., Morlighem, M., O'Neill, J. F., Nias, I., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Radic, V., Reese, R., Rounce, D. R., Rückamp, M., Sakai, A., Shafer, C., Schlegel, N. J., Shannon, S., Smith, R. S., Straneo, F., Sun, S., Tarasov, L., Trusel, L. D., Van Breedam, J., van de Wal, R., van den Broeke, M., Winkelmann, R., Zekollari, H., Zhao, C., Zhang, T., ~~&and~~ Zwinger, T. ~~(2021)~~.: Projected land ice contributions to twenty-first-century sea level rise. *Nature*, 593(7857), 74-82. <https://doi.org/10.1038/s41586-021-03302-y>, 2021.

590 Favier, L., Jourdain, N. C., Jenkins, A., Merino, N., Durand, G., Gagliardini, O., Gillet-Chaulet, F., ~~&and~~ Mathiot, P. ~~(2019)~~.: Assessment of sub-shelf melting parameterisations using the ocean-ice-sheet coupled model NEMO(v3.6)-Elmer/Ice(v8.3). *Geoscientific Model Development*, 12(6), 2255-2283. <https://doi.org/10.5194/gmd-12-2255-2019>, 2019.

585 Feldmann, J. and Levermann, A.: Timescales of outlet-glacier flow with negligible basal friction: theory, observations and modeling, *The Cryosphere*, 17, 327-348, <https://doi.org/10.5194/tc-17-327-2023>, 2023.

585 Feldmann, J., Levermann, A., and Winkelmann, R.: Hysteresis of idealized, instability-prone outlet glaciers in response to pinning-point buttressing variation, *The Cryosphere*, 18, 4011-4028, <https://doi.org/10.5194/tc-18-4011-2024>, 2024.

590 Feldmann, J., Albrecht, T., Khroulev, C., Pattyn, F., ~~&and~~ Levermann, A. ~~(2017)~~.: Resolution-dependent performance of grounding line motion in a shallow model compared with a full-Stokes model according to the MISMIP3d intercomparison. *Journal of Glaciology*, 60(220), 353-360. <https://doi.org/10.3189/2014JG13J093>, 2014.

Feldmann, J., & Levermann, A. (2023). Timescales of outlet-glacier flow with negligible basal friction: theory, observations, and modeling. *The Cryosphere*, 17(1), 327-348. <https://doi.org/10.5194/tc-17-327-2023>

595 ~~Feldmann, J., Levermann, A., & Winkelmann, R. (2024). Hysteresis of idealized, instability-prone outlet glaciers in response to pinning-point buttressing variation. *The Cryosphere*, 18(9), 4011-4028. <https://doi.org/10.5194/te-18-4011-2024>~~

Fowler, A., Murray, T., & Ng, F. (2001). Thermally controlled glacier surging. *Journal of Glaciology*, 47(159), 527-538. <https://doi.org/10.3189/172756501781831792>, 2001.

600 Frederick, B. C., Young, D. A., Blankenship, D. D., Richter, T. G., Kempf, S. D., Ferraccioli, F., & Siegert, M. J. (2016). Distribution of subglacial sediments across the Wilkes Subglacial Basin, East Antarctica. *Journal of Geophysical Research: Earth Surface*, 121(4), 790-813. <https://doi.org/10.1002/2015jg003760>, 2016.

Gagliardini, O., Durand, G., Zwinger, T., Hindmarsh, R. C. A., & Le Meur, E. (2010). Coupling of ice - shelf melting and buttressing is a key process in ice - sheets dynamics. *Geophysical Research Letters*, 37(14). <https://doi.org/10.1029/2010gl043334>, 2010.

605 Garbe, J., Albrecht, T., Levermann, A., Donges, J. F., & Winkelmann, R. (2020). The hysteresis of the Antarctic Ice Sheet. *Nature*, 585(7826), 538-544. <https://doi.org/10.1038/s41586-020-2727-5>, 2020.

Goldberg, D., Holland, D. M., & Schoof, C. (2009). Grounding line movement and ice shelf buttressing in marine ice sheets. *Journal of Geophysical Research: Earth Surface*, 114(F4). <https://doi.org/10.1029/2008jg001227>, 2009.

610 Golledge, N. R., Kowalewski, D. E., Naish, T. R., Levy, R. H., Fogwill, C. J., & Gasson, E. G. (2015). The multi-millennial Antarctic commitment to future sea-level rise. *Nature*, 526(7573), 421-425. <https://doi.org/10.1038/nature15706>, 2015.

Goodwin, B. P., Mosley - Thompson, E., Wilson, A. B., Porter, S. E., & Sierra - Hernandez, M. R. (2016). Accumulation Variability in the Antarctic Peninsula: The Role of Large-Scale Atmospheric Oscillations and Their Interactions. *Journal of Climate*, 29(7), 2579-2596. <https://doi.org/10.1175/JCLI-D-15-0354.1>, 2016.

615 Greenbaum, J., Blankenship, D., Young, D., Richter, T., Roberts, J., Aitken, A., Legresy, B., Schroeder, D., Warner, R., & Van Ommen, T. (2015). Ocean access to a cavity beneath Totten Glacier in East Antarctica. *Nature Geoscience*, 8(4), 294-298. <https://doi.org/10.1038/ngeo2388>, 2015.

Gudmundsson, G. H. (2003). Transmission of basal variability to a glacier surface. *Journal of Geophysical Research: Solid Earth*, 108(B5). <https://doi.org/10.1029/2002jb002107>, 2003.

620 Gudmundsson, G. H. (2013). Ice-shelf buttressing and the stability of marine ice sheets. *The Cryosphere*, 7(2), 647-655. <https://doi.org/10.5194/te-7-647-2013>, 2013.

Hellmer, H. H. and Olbers, D. J. (1989). A Two-Dimensional Model for the Thermohaline Circulation Under an Ice Shelf. *Antarctic Science*, 1(04), 325-336. <https://doi.org/10.1017/S09541020890004901016/j.ocemod.2007.01.001>, 1989.

625 Hill, E. A., Gudmundsson, G. H., & Chandler, D. M. (2024). Ocean warming as a trigger for irreversible retreat of the Antarctic ice sheet. *Nature Climate Change*, 14(11), 1165-1171. <https://doi.org/10.1038/s41558-024-02134-8>, 2024.

Hindmarsh, R. C. (2006). The role of membrane-like stresses in determining the stability and sensitivity of the Antarctic ice sheets: back pressure and grounding line motion. *Philos. Trans. A Math. Phys. Eng. Sci.*, 364(1844), 1733-1767. <https://doi.org/10.1098/rsta.2006.1797>, 2006.

630 Hock, R., Bliss, A., Marzeion, B. E. N., Giesen, R. H., Hirabayashi, Y., Huss, M., Radić, V., & Slangen, A. B. A. (2019). GlacierMIP - A model intercomparison of global-scale glacier mass-balance models and projections. *Journal of Glaciology*, 65(251), 453-467. <https://doi.org/10.1017/jog.2019.22>, 2019.

Holland, D. M. and Jenkins, A. (1999). Modeling Thermodynamic Ice-Ocean Interactions at the Base of an Ice Shelf. *Journal of Physical Oceanography*, 29, 1787-1800. [https://doi.org/10.1175/1520-0485\(1999\)029<1787:MTIOIA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1999)029<1787:MTIOIA>2.0.CO;2), 1999.

635 Holland, P. R., Bracegirdle, T. J., Dutrieux, P., Jenkins, A., & Steig, E. J. (2019). West Antarctic ice loss influenced by internal climate variability and anthropogenic forcing. *Nature Geoscience*, 12(9), 718-724. <https://doi.org/10.1038/s41561-019-0420-9>, 2019.

640 Huybers, K., Roe, G., & Conway, H. (2017). Basal topographic controls on the stability of the West Antarctic ice sheet: lessons from Foundation Ice Stream. *Annals of Glaciology*, 58(75pt2), 193-198. <https://doi.org/10.1017/aog.2017.9>, 2017.

645 Jacobs, S. S., Jenkins, A., Giulivi, C. F., & Dutrieux, P. (2011). Stronger ocean circulation and increased melting under Pine Island Glacier ice shelf. *Nature Geoscience*, 4(8), 519-523. <https://doi.org/10.1038/ngeo1188>, 2011.

- Joughin, I., E. Smith, B., ~~& Brooke, M.~~ (2014). Marine Ice Sheet Collapse Potentially Under Way for the Thwaites Glacier Basin, West Antarctica. *Science*, 344(6185), 735-738. <https://doi.org/10.1126/science.124905>. 2014.
- 650 Jourdain, N. C., Asay-Davis, X., ~~& Hattermann, T.~~ (2020). A protocol for calculating basal melt rates in the ISMIP6 Antarctic ice sheet projections. *The Cryosphere*, 14, 3111-3134. <https://doi.org/10.5194/tc-14-3111-2020>. 2020.
- ~~Juarez-Martinez, A., Blasco, J., Robinson, A., Montoya, M., and Alvarez-Solas, J.: Antarctic sensitivity to oceanic melting parameterizations, *The Cryosphere*, 18, 4257-4283, <https://doi.org/10.5194/tc-18-4257-2024>, 2024.~~
- 655 Kamworapan, S., Bich Thao, P. T., Gheewala, S. H., Pimonsree, S., ~~& Prueksakorn, K.~~ (2021). Evaluation of CMIP6 GCMs for simulations of temperature over Thailand and nearby areas in the early 21st century. *Heliyon*, 7(11), e08263. <https://doi.org/10.1016/j.heliyon.2021.e08263>. 2021.
- Klose, A. K., Coulon, V., Pattyn, F., ~~& Winkelmann, R.~~ (2024). The long-term sea-level commitment from Antarctica. *The Cryosphere*, 18(9), 4463-4492. <https://doi.org/10.5194/tc-18-4463-2024>. 2024.
- 660 Konrad, H., Shepherd, A., Gilbert, L., Hogg, A. E., McMillan, M., Muir, A., ~~& Slater, T.~~ (2018). Net retreat of Antarctic glacier grounding lines. *Nature Geoscience*, 11(4), 258-262. <https://doi.org/10.1038/s41561-018-0082-z>. 2018.
- Lazeroms, W. M. J., Jenkins, A., Gudmundsson, G. H., ~~& van de Wal, R. S. W.~~ (2018). Modeling: Modelling present-day basal melt rates for Antarctic ice shelves using a parametrization of buoyant meltwater plumes. *The Cryosphere*, 12(1), 49-70. <https://doi.org/10.5194/tc-12-49-2018>. 2018.
- 665 Leguy, G. R., Asay-Davis, X. S., ~~& Lipscomb, W. H.~~ (2014). Parameterization of basal friction near grounding lines in a one-dimensional ice sheet model. *The Cryosphere*, 8(4), 1239-1259. <https://doi.org/10.5194/tc-8-1239-2014>. 2014.
- Levermann, A., Winkelmann, R., Albrecht, T., Goelzer, H., Golledge, N. R., Greve, R., Huybrechts, P., Jordan, J., Leguy, G., Martin, D., Morlighem, M., Pattyn, F., Pollard, D., Quiquet, A., Rodehacke, C., Seroussi, H., Sutter, J., Zhang, T., Van Breedam, J., Calov, R., DeConto, R., Dumas, C., Garbe, J., Gudmundsson, G. H., Hoffman, M. J., Humbert, A., Kleiner, T., Lipscomb, W. H., Meinshausen, M., Ng, E., Nowicki, S. M. J., Perego, M., Price, S. F., Saito, F., Schlegel, N.-J., Sun, S., ~~& van de Wal, R. S. W.~~ (2020). Projecting Antarctica's contribution to future sea level rise from basal ice shelf melt using linear response functions of 16 ice sheet models (LARMIP-2). *Earth System Dynamics*, 11(1), 35-76. <https://doi.org/10.5194/esd-11-35-2020>. 2020.
- 675 Li, L., Aitken, A. R., Lindsay, M. D., ~~& Kulesa, B.~~ (2022). Sedimentary basins reduce stability of Antarctic ice streams through groundwater feedbacks. *Nature Geoscience*, 15(8), 645-650. <https://doi.org/10.1038/s41561-022-00992-5>. 2022.
- Li, Q., England, M. H., Hogg, A. M., Rintoul, S. R., ~~& Morrison, A. K.~~ (2023). Abyssal ocean overturning slowdown and warming driven by Antarctic meltwater. *Nature*, 615(7954), 841-847. <https://doi.org/10.1038/s41586-023-05762-w>. 2023.
- ~~Li, X., Rignot, E., Mouginot, J., & Scheuchl, B.: Ice flow dynamics and mass loss of Totten Glacier, East Antarctica, from 1989 to 2015, *Geophysical Research Letters*, 43, 6366-6373, <https://doi.org/10.1002/2016gl069173>, 2016.~~
- 685 Lipscomb, W. H., Price, S. F., Hoffman, M. J., Leguy, G. R., Bennett, A. R., Bradley, S. L., Evans, K. J., Fyke, J. G., Kennedy, J. H., Perego, M., Ranken, D. M., Sacks, W. J., Salinger, A. G., Vargo, L. J., ~~& Worley, P. H.~~ (2019). Description and evaluation of the Community Ice Sheet Model (CISM) v2.1. *Geoscientific Model Development*, 12(1), 387-424. <https://doi.org/10.5194/gmd-12-387-2019>. 2019.
- ~~Li, X., Rignot, E., Mouginot, J., & Scheuchl, B. (2016). Ice flow dynamics and mass loss of Totten Glacier, East Antarctica, from 1989 to 2015. *Geophysical Research Letters*, 43(12), 6366-6373. <https://doi.org/10.1002/2016gl069173>~~
- 690 Lowry, D. P., Krapp, M., Golledge, N. R., ~~& Alevropoulos-Borrill, A.~~ (2021). The influence of emissions scenarios on future Antarctic ice loss is unlikely to emerge this century. *Communications Earth & Environment*, 2(1), <https://doi.org/10.1038/s43247-021-00289-2>. 2021.
- Martin, M. A., Winkelmann, R., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., ~~& Levermann, A.~~ (2011). The Potsdam Parallel Ice Sheet Model (PISM-PIK) – Part 2: Dynamic equilibrium simulation of the Antarctic ice sheet. *The Cryosphere*, 5(3), 727-740. <https://doi.org/10.5194/tc-5-727-2011>. 2011.
- 695 Mengel, M., ~~& Levermann, A.~~ (2014). Ice plug prevents irreversible discharge from East Antarctica. *Nature Climate Change*, 4(6), 451-455. <https://doi.org/10.1038/nclimate2226>. 2014.
- Miles, B. W. J., ~~& Bingham, R. G.~~ (2024). Progressive unanchoring of Antarctic ice shelves since 1973. *Nature*, 626(8000), 785-791. <https://doi.org/10.1038/s41586-024-07049-0>. 2024.
- 700

- Miles, B. W. J., Stokes, C. R., Jamieson, S. S. R., Jordan, J. R., Gudmundsson, G. H., ~~& Jenkins, A. (2022).~~ High spatial and temporal variability in Antarctic ice discharge linked to ice shelf buttressing and bed geometry. *Scientific Reports*, 12(1), 10968. <https://doi.org/10.1038/s41598-022-13517-2>, 2022.
- 705 Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., Goel, V., Greenbaum, J. S., Gudmundsson, H., Guo, J., Helm, V., Hofstede, C., Howat, I., Humbert, A., Jokat, W., Karlsson, N. B., Lee, W. S., Matsuoka, K., Millan, R., Mouginit, J., Paden, J., Pattyn, F., Roberts, J., Rosier, S., Ruppel, A., Seroussi, H., Smith, E. C., Steinhage, D., Sun, B., Broeke, M. R. v. d., Ommen, T. D. v., Wesse, M. v., ~~& Young, D. A. (2019).~~ Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet. *Nature Geoscience*, 13(2), 132-137. <https://doi.org/10.1038/s41561-019-0510-8>, 2019.
- 710 Mouginit, J., Rignot, E., ~~& Scheuchl, B. (2019).~~ Continent - Wide, Interferometric SAR Phase, Mapping of Antarctic Ice Velocity. *Geophysical Research Letters*, 46(16), 9710-9718. <https://doi.org/https://doi.org/10.1029/2019GL083826>, 2019.
- Noble, T. L., Rohling, E. J., Aitken, A. R. A., Bostock, H. C., Chase, Z., Gomez, N., Jong, L. M., King, M. A., Mackintosh, A. N., McCormack, F. S., McKay, R. M., Menviel, L., Phipps, S. J., Weber, M. E., Fogwill, C. J., Gayen, B., Golledge, N. R., Gwyther, D. E., Hogg, A. M., Martos, Y. M., Pena - Molino, B., Roberts, J., Flierdt, T., ~~& Williams, T. (2020).~~ The Sensitivity of the Antarctic Ice Sheet to a Changing Climate: Past, Present, and Future. *Reviews of Geophysics*, 58(4), <https://doi.org/10.1029/2019rg000663>, 2020.
- 715 ~~Nowicki, S. M. J., Payne, T., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., & Shepherd, A. (2016). Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6. *Geoscientific Model Development*, 9(12), 4521-4545. <https://doi.org/10.5194/gmd-9-4521-2016>~~
- 720 Nowicki, S., Goelzer, H., Seroussi, H., Payne, A. J., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Alexander, P., Asay-Davis, X. S., Barthel, A., Bracegirdle, T. J., Cullather, R., Felikson, D., Fettweis, X., Gregory, J. M., Hattermann, T., Jourdain, N. C., Kuipers Munneke, P., Larour, E., Little, C. M., Morlighem, M., Nias, I., Shepherd, A., Simon, E., Slater, D., Smith, R. S., Straneo, F., Trusel, L. D., van den Broeke, M. R., ~~& van de Wal, R. (2020).~~ Experimental protocol for sea level projections from ISMIP6 stand-alone ice sheet models. *The Cryosphere*, 14(7), 2331-2368. <https://doi.org/10.5194/tc-14-2331-2020>, 2020.
- 725 ~~Nowicki, S. M. J., Payne, T., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., and Shepherd, A.: Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6, *Geoscientific Model Development*, 9, 4521-4545, <https://doi.org/10.5194/gmd-9-4521-2016>, 2016.~~
- 730 Nowicki, S., Simon, E., ~~& the ISMIP6 Team. (2021).~~ ISMIP6 21st Century Forcing Datasets [Data set]. *The GitHub*. <https://doi.org/10.5281/zenodo.11176009>, 2021.
- Ohmura, A. (2001). Physical basis for the temperature-based melt-index method. *Journal of applied Meteorology*, 40(4), 753-761. [https://doi.org/10.1175/1520-0450\(2001\)040%3C0753:PBFTTB%3E2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040%3C0753:PBFTTB%3E2.0.CO;2), 2001.
- 735 Paolo, F. S., Fricker, H. A., and Padman, L.: Volume loss from Antarctic ice shelves is accelerating, *Science*, 348, 327-331, <https://doi.org/10.1126/science.aaa0940>, 2015.
- ~~Paolo, F. S., Gardner, A. S., Greene, C. A., Nilsson, J., Schodlok, M. P., Schlegel, N. J., and Fricker, H. A.: Widespread slowdown in thinning rates of West Antarctic ice shelves, *The Cryosphere*, 17, 3409-3433, <https://doi.org/10.5194/tc-17-3409-2023>, 2023.~~
- 740 Pattyn, F., Schoof, C., Perichon, L., Hindmarsh, R. C. A., Bueller, E., de Fleurian, B., Durand, G., Gagliardini, O., Gladstone, R., Goldberg, D., Gudmundsson, G. H., Huybrechts, P., Lee, V., Nick, F. M., Payne, A. J., Pollard, D., Rybak, O., Saito, F., ~~& Vieli, A. (2012).~~ Results of the Marine Ice Sheet Model Intercomparison Project, MISIP. *The Cryosphere*, 6(3), 573-588. <https://doi.org/10.5194/tc-6-573-2012>, 2012.
- ~~Paolo, F. S., Fricker, H. A., & Padman, L. (2015). Volume loss from Antarctic ice shelves is accelerating. *Science*, 348(6232), 327-331. <https://doi.org/10.1126/science.aaa0940>~~
- 745 ~~Paolo, F. S., Gardner, A. S., Greene, C. A., Nilsson, J., Schodlok, M. P., Schlegel, N. J., & Fricker, H. A. (2023). Widespread slowdown in thinning rates of West Antarctic ice shelves. *The Cryosphere*, 17(8), 3409-3433. <https://doi.org/10.5194/tc-17-3409-2023>~~
- Payne, A. J., Nowicki, S., Abe - Ouchi, A., Agosta, C., Alexander, P., Albrecht, T., Asay - Davis, X., Aschwanden, A., Barthel, A., Bracegirdle, T. J., Calov, R., Chambers, C., Choi, Y., Cullather, R., Cuzzone, J., Dumas, C., Edwards, T. L., Felikson, D., Fettweis, X., Galton - Fenzi, B. K., Goelzer, H., Gladstone, R., Golledge, N. R., Gregory, J. M., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huybrechts, P., Jourdain, N. C., Kleiner, T., Munneke, P. K., Larour, E., Le clec'h, S., Lee, V., Leguy, G., Lipscomb, W. H., Little, C. M., Lowry, D. P., Morlighem, M., Nias, I., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Reese, R., Rückamp, M., Schlegel, N. J., Seroussi, H., Shepherd, A., Simon, E., Slater, D., Smith, R. S., Straneo, F., Sun, S., Tarasov, L., Trusel, L. D., Van Breedam, J., Wal, R., Broeke, M., Winkelmann, R., Zhao, C., Zhang, T., ~~& Zwinger, T. (2021).~~ Future Sea Level Change Under Coupled Model

Intercomparison Project Phase 5 and Phase 6 Scenarios From the Greenland and Antarctic Ice Sheets. *Geophysical Research Letters*, 48(16). <https://doi.org/10.1029/2020gl091741>, 2021.

Pelle, T., Morlighem, M., & Bondzio, J. H. (2019). Brief communication: PICOP, a new ocean melt parameterization under ice shelves combining PICO and a plume model. *The Cryosphere*, 13(3), 1043-1049. <https://doi.org/10.5194/tc-13-1043-2019>, 2019.

Pittard, M. L., Whitehouse, P. L., Bentley, M. J., & Small, D. (2022). An ensemble of Antarctic deglacial simulations constrained by geological observations. *Quaternary Science Reviews*, 298. <https://doi.org/10.1016/j.quascirev.2022.107800>, 2022.

Pollard, D., & DeConto, R. M. (2012). Description of a hybrid ice sheet-shelf model, and application to Antarctica. *Geoscientific Model Development*, 5(5), 1273-1295. <https://doi.org/10.5194/gmd-5-1273-2012>, 2012.

Pritchard, H. D., Ligtenberg, S. R. M., Fricker, H. A., Vaughan, D. G., van den Broeke, M. R., & Padman, L. (2012). Antarctic ice-sheet loss driven by basal melting of ice shelves. *Nature*, 484(7395), 502-505. <https://doi.org/10.1038/nature10968>, 2012.

Reese, R., Albrecht, T., Mengel, M., Asay-Davis, X., & Winkelmann, R. (2018). Antarctic sub-shelf melt rates via PICO. *The Cryosphere*, 12(6), 1969-1985. <https://doi.org/10.5194/tc-12-1969-2018>, 2018.

Reese, R., Levermann, A., Albrecht, T., Seroussi, H., & Winkelmann, R. (2020). The role of history and strength of the oceanic forcing in sea level projections from Antarctica with the Parallel Ice Sheet Model. *The Cryosphere*, 14(9), 3097-3110. <https://doi.org/10.5194/tc-14-3097-2020>, 2020.

Rignot, E., Bamber, J. L., van den Broeke, M. R., Davis, C., Li, Y., van de Berg, W. J., & van Meijgaard, E. (2008). Recent Antarctic ice mass loss from radar interferometry and regional climate modeling. *Nature Geoscience*, 1(2), 106-110. <https://doi.org/10.1038/ngeo102>

Rignot, E., Jacobs, R., Mouginot, J., & Scheuchl, B. (2013). Ice-Shelf Melting Around Antarctica. *Science*, 341, 266. <https://doi.org/10.1126/science.1235798>, 2013.

Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., & Morlighem, M. (2019). Four decades of Antarctic Ice Sheet mass balance from 1979-2017. *Proc. Natl. Acad. Sci. USA, U.S.A.*, 116(4), 1095-1103. <https://doi.org/10.1073/pnas.1812883116>, 2019.

Rignot, E., Bamber, J. L., van den Broeke, M. R., Davis, C., Li, Y., van de Berg, W. J., and van Meijgaard, E.: Recent Antarctic ice mass loss from radar interferometry and regional climate modelling, *Nature Geoscience*, 1, 106-110, <https://doi.org/10.1038/ngeo102>, 2008.

Ritz, C., Edwards, T. L., Durand, G., Payne, A. J., Peyaud, V., & Hindmarsh, R. C. A. (2015). Potential sea-level rise from Antarctic ice-sheet instability constrained by observations. *Nature*, 528(7580), 115-118. <https://doi.org/10.1038/nature16147>, 2015.

Rounce, D. R., Regine Hock, Fabien Maussion, Romain Hugonnet, William Kochtitzky, Matthias Huss, Etienne Berthier, Douglas Brinkerhoff, Loris Compagno, Luke Copland, Daniel Farinotti, Brian Menounos, & McNabb, R. W. (2023). Global glacier change in the 21st century: Every increase in temperature matters. *Science*, 379(6627), 78-83. <https://doi.org/10.1126/science.abo1324>, 2023.

Schlegel, N.-J., Seroussi, H., Schodlok, M. P., Larour, E. Y., Boening, C., Limonadi, D., Watkins, M. M., Morlighem, M., & van den Broeke, M. R. (2018). Exploration of Antarctic Ice Sheet 100-year contribution to sea level rise and associated model uncertainties using the ISSM framework. *The Cryosphere*, 12(11), 3511-3534. <https://doi.org/10.5194/tc-12-3511-2018>, 2018.

Schlemm, T., Feldmann, J., Winkelmann, R., & Levermann, A. (2022). Stabilizing effect of mélange buttressing on the marine ice-cliff instability of the West Antarctic Ice Sheet. *The Cryosphere*, 16(5), 1979-1996. <https://doi.org/10.5194/tc-16-1979-2022>, 2022.

Schmidtko, S., Heywood, K. J., Thompson, A. F., & Aoki, S. (2014). Multidecadal warming of Antarctic waters. *Science*, 346(6214), 1227-1231. <https://doi.org/10.1126/science.1256117>, 2014.

Schoof, C. (2007). Ice sheet grounding line dynamics: Steady states, stability, and hysteresis. *Journal of Geophysical Research: Earth Surface*, 112(F3). <https://doi.org/10.1029/2006jf000664>, 2007.

Seroussi, H., Morlighem, M., Rignot, E., Mouginot, J., Larour, E., Schodlok, M., & Khazendar, A. (2014). Sensitivity of the dynamics of Pine Island Glacier, West Antarctica, to climate forcing for the next 50 years. *The Cryosphere*, 8(5), 1699-1710. <https://doi.org/10.5194/tc-8-1699-2014>, 2014.

Seroussi, H., Nowicki, S., Simon, E., Abe-Ouchi, A., Albrecht, T., Brondex, J., Cornford, S., Dumas, C., Gillet-Chaulet, E., Goelzer, H., Gollledge, N. R., Gregory, J. M., Greve, R., Hoffman, M. J., Humbert, A., Huybrechts, P., Kleiner, T., Larour, E., Leguy, G., Lipscomb, W. H., Lowry, D., Mengel, M., Morlighem, M., Pattyn, F., Payne, A. J., Pollard, D., Price, S. F., Quiquet, A., Reerink, T. J., Reese, R., Rodehacke, C. B., Schlegel, N.-J., Shepherd, A., Sun, S., Sutter, J., Van Breedam, J., van de Wal, R. S. W., Winkelmann, R., and Zhang, T.: initMIP-Antarctica: an ice

[sheet model initialization experiment of ISMIP6, The Cryosphere, 13, 1441-1471, https://doi.org/10.5194/tc-13-1441-2019, 2019.](https://doi.org/10.5194/tc-13-1441-2019)

815 Seroussi, H., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., Barthel, A., Calov, R., Cullather, R., Dumas, C., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Gregory, J. M., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huybrechts, P., Jourdain, N. C., Kleiner, T., Larour, E., Leguy, G. R., Lowry, D. P., Little, C. M., Morlighem, M., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Reese, R., Schlegel, N.-J., Shepherd, A., Simon, E., Smith, R. S., Straneo, F., Sun, S., Trusel, L. D., Van Breedam, J., van de Wal, R. S. W., Winkelmann, R., Zhao, C., Zhang, T., ~~& Zwinger, T. (2020).~~ ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st century. *The Cryosphere*, 14(9), 3033-3070. <https://doi.org/10.5194/tc-14-3033-2020>, 2020.

820 ~~Seroussi, H., Nowicki, S., Simon, E., Abe-Ouchi, A., Albrecht, T., Brondex, J., Cornford, S., Dumas, C., Gillet-Chaulet, F., Goelzer, H., Golledge, N. R., Gregory, J. M., Greve, R., Hoffman, M. J., Humbert, A., Huybrechts, P., Kleiner, T., Larour, E., Leguy, G., Lipscomb, W. H., Lowry, D., Mengel, M., Morlighem, M., Pattyn, F., Payne, A. J., Pollard, D., Price, S. F., Quiquet, A., Reerink, T. J., Reese, R., Rodehaeke, C. B., Schlegel, N. J., Shepherd, A., Sun, S., Sutter, J., Van Breedam, J., van de Wal, R. S. W., Winkelmann, R., & Zhang, T. (2019). initMIP Antarctica: an ice sheet model initialization experiment of ISMIP6. *The Cryosphere*, 13(5), 1441-1471. <https://doi.org/10.5194/tc-13-1441-2019>~~

830 Seroussi, H., Verjans, V., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., Barthel, A., Calov, R., Cullather, R., Dumas, C., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Gregory, J. M., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huybrechts, P., Jourdain, N. C., Kleiner, T., Larour, E., Leguy, G. R., Lowry, D. P., Little, C. M., Morlighem, M., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Reese, R., Schlegel, N.-J., Shepherd, A., Simon, E., Smith, R. S., Straneo, F., Sun, S., Trusel, L. D., Van Breedam, J., Van Katwyk, P., van de Wal, R. S. W., Winkelmann, R., Zhao, C., Zhang, T., ~~& Zwinger, T. (2023).~~ Insights into the vulnerability of Antarctic glaciers from the ISMIP6 ice sheet model ensemble and associated uncertainty. *The Cryosphere*, 17(12), 5197-5217. <https://doi.org/10.5194/tc-17-5197-2023>, 2023.

835 Tokarska, K. B., Stolpe, M. B., Sippel, S., Fischer, E. M., Smith, C. J., Lehner, F., ~~& Knutti, R. (2020).~~ Past warming trend constrains future warming in CMIP6 models. *Science Advances*, 6(12), eaaz9549. <https://doi.org/10.1126/sciadv.aaz9549>, 2020.

840 van der Linden, E. C., Le Bars, D., Lambert, E., ~~& Drijfhout, S. (2023).~~ Antarctic contribution to future sea level from ice shelf basal melt as constrained by ice discharge observations. *The Cryosphere*, 17(1), 79-103. <https://doi.org/10.5194/tc-17-79-2023>, 2023.

845 Van Der Veen, C. J., Stearns, L. A., Johnson, J., ~~& Csatho, B. (2014).~~ Flow dynamics of Byrd Glacier, East Antarctica. *Journal of Glaciology*, 60(224), 1053-1064. <https://doi.org/10.3189/2014JoG14J052>, 2014.

~~van Van Pelt, W. J. J., & Oerlemans, J. (2012).~~ Numerical simulations of cyclic behaviour in the Parallel Ice Sheet Model (PISM). *Journal of Glaciology*, 58(208), 347-360. <https://doi.org/10.3189/2012JoG11J217>, 2012.

850 van Wessem, J. M., van de Berg, W. J., Noël, B. P. Y., van Meijgaard, E., Amory, C., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts, J. T. M., Lhermitte, S., Ligtenberg, S. R. M., Medley, B., Reijmer, C. H., van Tricht, K., Trusel, L. D., van Ulf, L. H., Wouters, B., Wuite, J., ~~& van den Broeke, M. R. (2018).~~ Modeling: Modelling the climate and surface mass balance of polar ice sheets using RACMO2—Part 2: Antarctica (1979–2016). *The Cryosphere*, 12(4), 1479-1498. <https://doi.org/10.5194/tc-12-1479-2018>, 2018.

Winkelmann, R., Martin, M. A., Haseloff, M., Albrecht, T., Bueller, E., Khroulev, C., ~~& Levermann, A. (2011).~~ The Potsdam Parallel Ice Sheet Model (PISM-PIK)—Part 1: Model description. *The Cryosphere*, 5(3), 715-726. <https://doi.org/10.5194/tc-5-715-2011>, 2011.

855 Wright, A. P., Young, D. A., Roberts, J. L., Schroeder, D. M., Bamber, J. L., Dowdeswell, J. A., Young, N. W., Le Brocq, A. M., Warner, R. C., Payne, A. J., Blankenship, D. D., van Ommen, T. D., ~~& Siegert, M. J. (2012).~~ Evidence of a hydrological connection between the ice divide and ice sheet margin in the Aurora Subglacial Basin, East Antarctica. *Journal of Geophysical Research: Earth Surface*, 117(F1). <https://doi.org/10.1029/2011jg002066>, 2012.

860 Wyser, K., Kjellström, E., Koenig, T., Martins, H., ~~& Döscher, R. (2020).~~ Warmer climate projections in EC-Earth3-Veg: the role of changes in the greenhouse gas concentrations from CMIP5 to CMIP6. *Environmental Research Letters*, 15(5), 054020. <https://doi.org/10.1088/1748-9326/ab81c2>, 2020.

~~Zhang, T., Colgan, W., Wansing, A., Løkkegaard, A., Leguy, G., Lipscomb, W. H., and Xiao, C.: Evaluating different geothermal heat-flow maps as basal boundary conditions during spin-up of the Greenland ice sheet, *The Cryosphere*, 18, 387-402, <https://doi.org/10.5194/tc-18-387-2024>, 2024.~~

870

Zhao, C., Gladstone, R., Zwinger, T., Gillet-Chaulet, F., Wang, Y., Caillet, J., Mathiot, P., Saraste, L., Jager, E., Galton-Fenzi, B. K., Christoffersen, P., & King, M. A. (2025). Subglacial water amplifies Antarctic contributions to sea-level rise. *Nature Communications*, 16(1). <https://doi.org/10.1038/s41467-025-58375-4>, 2025.

~~Zhang, T., Colgan, W., Wansing, A., Løkkegaard, A., Leguy, G., Lipscomb, W. H., & Xiao, C. (2024). Evaluating different geothermal heat flow maps as basal boundary conditions during spin-up of the Greenland ice sheet. *The Cryosphere*, 18(1), 387–402. <https://doi.org/10.5194/te-18-387-2024>~~

Zwally, H., Jay, M. B. G., Matthew A. Beckley, & Jack L. Saba. (2012). Antarctic and Greenland Drainage Systems. GSFC Cryospheric Sciences Laboratory, 2012.