

1 Impact of lower atmospheric scattering on ground-based optical 2 thermospheric wind observations with spatially uneven airglow

3 Xiaolong Wei^{1,2,4}, Guoying Jiang^{1,3,4,*}, Yajun Zhu^{1,3,4,*}, Jiyao Xu^{1,4}, Weijun Liu^{1,4}, Tiancai Wang^{1,2,4},
4 Guangyi Zhu^{1,4}, Wei Yuan^{1,4}

5 ¹State Key Laboratory of Solar Activity and Space Weather, National Space Science Center, Chinese Academy of Sciences,
6 Beijing, 100190, China

7 ²College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China.

8 ³School of Astronomy and Space Science, University of Chinese Academy of Sciences, Beijing, 100049, China.

9 ⁴Hainan National Field Science Observation and Research Observatory for Space Weather, Hainan, 571734, China

10 *Correspondence to: Guoying Jiang (gyjiang@spaceweather.ac.cn), Yajun Zhu (y.zhu@swl.ac.cn)

11 **Abstract.** Scattered airglow emissions in the lower atmosphere can bias ground-based interferometer observations of
12 thermospheric winds, particularly when airglow brightness becomes spatially uneven due to auroras. During two geomagnetic
13 storms with visible auroras on May 10th and Oct. 10th, 2024, the Doppler Asymmetric Spatial Heterodyne (DASH) and Fabry-
14 Perot (FP) interferometers concurrently detected atypical winds at Siziwang (SIZW, 41.83° N, 111.93° E), suspected to be
15 caused by scattering. These atypical winds, characterized by horizontal differences exceeding 400 m·s⁻¹ between opposite
16 cardinal directions (N-S or E-W) and downwelling exceeding 100 m·s⁻¹, showed a strong temporal association with airglow
17 brightness. By modelling the transmission of scattered airglow emissions, we calculated post-scattering wind speeds as the
18 initial wind speeds weighted by both scattered and direct intensities. With fixed initial speeds, the simulation reproduced the
19 temporal characteristics of the atypical winds, demonstrating that scattering ~~may contribute to~~ ~~causes~~ these intense horizontal
20 differences and downwelling. The simulation also shows that the scattering-induced biases have directional inhomogeneity
21 with characteristics linked to the location and background line-of-sight speed of the brighter airglow region. ~~The commonly~~
22 ~~used horizontal average wind may experience numerical deviations due to directional inhomogeneity.~~ The accuracy of the
23 simulation is limited by the accuracy of airglow observations and atmospheric optical depth.

24 1 Introduction

25 Optical interferometers are widely utilized to observe thermospheric neutral wind (Burnside et al., 1981; Burnside and Tepley,
26 1989; Killeen et al., 1995; Emmert et al., 2001; Fejer et al., 2002; Emmert et al., 2006; Wu et al., 2014). Thermospheric wind
27 can be derived from measuring the Doppler shift of OI red-line airglow emission at 630.0 nm. This emission, primarily from
28 the collisional deactivation of O(¹D) generated by O₂⁺ dissociative recombination, peaks near 250 km altitude. The height-
29 integrated thermospheric wind around the peak altitude can be obtained (Biondi and Feibelman, 1968; Hernandez and Roble,
30 1976; Burnside et al., 1981; Biondi et al., 1995; Nakajima et al., 1995). For scanning interferometers, three-dimensional wind
31 vectors can be derived by observing the zenith and four cardinal directions at a specific elevation angle. The scanning range

covers a circular area about 500 km in diameter at airglow altitude. Given thermospheric wind uniformity at this scale, horizontal winds observed in two opposite cardinal directions (N-S or E-W) are typically similar. Averaging opposite cardinal directions improves accuracy, mitigates cloud effects, and is typically used to represent local meridional or zonal winds even during geomagnetic storms (Friedman and Herrero, 1982; Fejer et al., 2002; Sakanoi et al., 2002; Dhadly et al., 2017; Huang et al., 2018; Xu et al., 2019; Li et al., 2023; Wang et al., 2025).

However, horizontal winds in opposite cardinal directions occasionally show significant separation exceeding 100 m s^{-1} , with and strong vertical winds, ~~generally~~ deviating from typical thermospheric wind uniformity. These observations often occur near auroras, unaffected by clouds or moonlight, and have acceptable standard errors. They mainly occur in polar regions (Crickmore et al., 1991; Price et al., 1995; Smith and Hernandez, 1995; Innis et al., 1999; Ishii et al., 2001; Guo and McEwen, 2003; Anderson et al., 2012), but have also been seen at mid-latitudes during major geomagnetic storms (Hernandez and Roble, 1976; Makela et al., 2014).

Atmospheric scattering of airglow emissions is thought to be one of the factors that biases ground-based wind observations, potentially accounting for the atypical wind. ~~introduces errors to ground-based interferometers, potentially accounting for these atypical wind observations.~~ Initially, it was thought to impact airglow peak height measurements by photometers (Ashburn, 1954). Subsequent studies by Abreu et al. (1983) explored its impact on thermospheric wind speed measurements using a Fabry-Perot interferometer. Harding et al. (2017a; 2017b) later systematically modelled and estimated these effects, revealing that scattering was responsible for the anomalous vertical winds observed at mid-latitudes during geomagnetic storms by Makela et al. (2014). Light from brighter airglow regions scatters omnidirectionally in the lower atmosphere, primarily the troposphere and stratosphere, and is detectable outside its original direction. The additional Doppler shift of this scattered light can bias the retrieval of line-of-sight (LOS) speeds as well as the converted horizontal and vertical winds. Harding et al. (2017b) also investigated the impact of atmospheric scattering on interferometer wind and temperature measurements during quiet periods.

Scattering-induced biases are more pronounced during spatially uneven airglow brightness, such as during auroras (Harding et al., 2017a). Uneven airglow brightness refers specifically to inhomogeneous red-line emissions. At mid-latitudes, marked uneven red-line airglow usually comes from red aurora. Despite their distinct origins, the spectral and altitudinal overlap of airglow and aurora will let ground-based optical instruments conflate the two. ~~Airglow and auroral emissions have similar wavelengths and peak heights, though differing in mechanism, making them hard to distinguish in ground-based observations. Harding et al. (2017b) also investigated the impact of atmospheric scattering on interferometer wind and temperature measurements during quiet periods and attempted to correct for the associated errors.~~ For red-line observations, the aurora itself may also bias the derived winds. Aurora could elevate the red-line emission profile (Kataoka et al., 2024b), so the interferometer samples winds that are both higher and farther away. This makes the northward view sense winds deviate from the expected thermospheric wind at 250 km altitude when looking toward the aurora. Additionally, spectral contamination

64 from precipitating energetic ions ~~could~~ ~~can~~ also bias interferometers (Makela et al., 2014). They suggested that the enhanced
65 downwelling at mid-latitudes during storms might result from the contamination of the spectral profile by fast O atoms
66 associated with the influx of low-energy O⁺ ions.

67 From a dynamical perspective, wind differences in opposite cardinal directions are considered horizontal divergence, which
68 are often associated with changes in vertical winds. Near the aurora arc, these atypical winds are mainly caused by ion drag,
69 Joule heating, and energy particle precipitation (Hays et al., 1984; Rees et al., 1984; Conde and Smith, 1995; Conde et al.,
70 2001; Anderson et al., 2012). Generally, excessive horizontal divergence and vertical wind appear alongside rapidly changing
71 auroras and exhibit a matching spatial relationship that upward (downward) winds accompanied by divergences (convergences)
72 are often detected when aurora exists equatorward (poleward) of the observatory (Ishii et al., 2001; Guo and McEwen, 2003).
73 The combination of vertical wind and horizontal divergence is related to gravity waves excited by the above processes in polar
74 regions, presenting a wave-like structure and phase delay between vertical and horizontal wind components. (Price et al., 1995;
75 Smith and Hernandez, 1995; Ishii et al., 1999; Ishii et al., 2001; Shinagawa and Oyama, 2006). At mid-latitudes, which are
76 not primary regions for magnetospheric energy injection, atypical winds are instead related to the propagation of gravity waves
77 from polar regions. (Hernandez and Roble, 1976).

78 During two geomagnetic storms on May 10th and October 10th, 2024, with visible auroras, we observed similar atypical winds
79 in ground-based interferometers at Siziwang (SIZW, 41.83° N, 111.93° E), China. These winds showed intense differences
80 over 400 m·s⁻¹ in two opposite cardinal directions for both meridional and zonal components, along with downward wind
81 exceeding 100 m·s⁻¹. The observations were unaffected by moonlight or clouds, and the interferometer retrieval errors were
82 acceptable (see Section 3.1). These atypical winds at SIZW only occurred with auroras statistically and significantly deviated
83 from the regional climatological norms over the China region (Jiang et al., 2018; Yang et al., 2020). This raises the question
84 of whether the atypical winds arise from dynamical processes, are influenced by red aurora, or stem from scattering-induced
85 biases and other measurement-related factors. Unfortunately, most of these mechanisms could amplify the wind-speed contrast
86 between opposite cardinal directions, rendering them difficult to disentangle (Harding et al., 2017a). Given the scarcity of
87 additional thermospheric-wind or auroral instruments, we remain unable to quantify every potential mechanism. Motivated by
88 the observed phenomena, this study attempts to estimate how scattering modulates the atypical winds in these storms. While
89 prior studies focus on vertical wind biases of Fabry-Perot interferometers under auroral conditions (Harding et al., 2017a;
90 Harding et al., 2017b), we will analyze the formation and patterns of horizontal differences caused by scattering. We will also
91 incorporate Doppler Asymmetric Spatial Heterodyne (DASH) interferometer data to compare scattering impact across
92 different interferometer types. As red auroras now regularly appear at the low magnetic latitudes of Japan and China during
93 elevated solar activity (Kataoka et al., 2024a; Kataoka et al., 2024b; Ma et al., 2024), a deeper understanding of scattering-
94 induced biases is essential for the proper use of interferometer data collected in these regions. In the following text, a scattering
95 radiative transfer model is used to simulate interferometer observations in two cases with visible aurora. The presence and

96 ~~patterns of scattering-induced biases are analyzed by comparing simulations with observations. This raises the question of~~
97 ~~whether the atypical winds are driven by dynamical processes or by scattering-induced biases of interferometers, which causes~~
98 ~~the thermospheric wind to deviate from horizontal uniformity. Unfortunately, both factors can manifest as increased~~
99 ~~differences between opposite cardinal directions, complicating the distinction between them (Harding et al., 2017a). However,~~
100 ~~the simultaneous variations in vertical winds, horizontal differences, and red-line brightness show no phase lag, thus not~~
101 ~~providing evidence for the energy conversion process (Ishii et al., 1999). Instead, these variations resemble a systematic error,~~
102 ~~as they all involve negative LOS speeds. This suggests that scattering impact may be more significant than dynamic~~
103 ~~mechanisms in these cases.—~~

104 ~~Auroras are rare at mid-latitudes in East Asia due to low magnetic latitude. The magnetic latitude of SIZW is just 37.7° N. But~~
105 ~~increased solar activity has led to more frequent sightings of mid-latitude red auroras in Japan and China (Kataoka et al., 2024a;~~
106 ~~Kataoka et al., 2024b; Ma et al., 2024). Previously overlooked, the scattering-induced biases associated with auroras need~~
107 ~~further analysis to determine the accuracy and suitability of storm time observations for dynamical analysis. While prior~~
108 ~~studies have examined vertical wind biases of Fabry-Perot interferometers under auroral conditions (Abreu et al., 1983;~~
109 ~~Harding et al., 2017a; Harding et al., 2017b), we will incorporate Doppler Asymmetric Spatial Heterodyne (DASH)~~
110 ~~interferometer data to compare scattering impact across different interferometer types. Additionally, we will investigate the~~
111 ~~causes and patterns of horizontal differences and assess the reliability of individual directions and horizontal average winds~~
112 ~~under the influence of scattering. In the following text, a scattering radiative transfer model is used to simulate interferometer~~
113 ~~observations in two cases with visible aurora. The presence and patterns of scattering-induced biases are analyzed by~~
114 ~~comparing simulations with observations.~~

115 **2 Instruments and model**

116 This study was conducted at the Siziwang station (SIZW; 41.83° N, 111.93° E, and 37.7° N MLat) of the Chinese Meridian
117 Project Phase II (Wang et al., 2024), utilizing a Dual-Channel All-sky Airglow Imager (DCAI), a Dual-Channel Optical
118 Interferometer (DCOI), and a Fabry-Perot Interferometer (FPI). DCOI derives neutral winds by observing atomic oxygen
119 green-line (557.7 nm, around 96 km) and red-line (630.0 nm, around 250 km). DCAI observes hydroxyl (around 87 km) and
120 atomic oxygen red-line nightglow, respectively. FPI only works at the red-line. Our focus is on the red-line channel. Using
121 DCAI images as one of the inputs, wind biases from optical interferometers can be simulated by a scattering radiative transfer
122 model (scattering model for short). Instruments and the model are described in the following subsections.

123 **2.1 Dual-Channel All-Sky Airglow Imager**

124 Dual-Channel All-Sky Airglow Imager (DCAI) comprises a fisheye lens with an approximate 170 degree field of view, a 2
125 nm narrow-band filter, and a 1024×1024 pixel, 16 bit cooled CCD. DCAI exposure time of the red-line is 2 minutes. The

obtained airglow images are first calibrated to the local spherical coordinate system, then sequentially corrected for stray light, Van Rhijn effect, and atmospheric extinction, and finally projected onto the 250 km airglow plane. Due to DCAI not calibrating the Rayleigh unit (Shiokawa et al., 2000), observed brightness is only normalized to the full-well value. And because of fish-eye lens distortion and the lack of Rayleigh unit calibration, the edge brightness of the view is inaccurate. Thus, observations are restricted within a 70° zenith angle. For larger zenith angles, the brightness is obtained by radial zero-order extrapolation in airglow projection. Detailed image processing procedures are in Appendix B.

2.2 Dual-channel optical interferometer

Dual-channel optical interferometer (DCOI) is a scanning interferometer using Doppler Asymmetric Spatial Heterodyne (DASH) technology. DASH exhibits a wider field of view, better thermal stability, simplified mechanisms, and lower tolerance requirements than other interferometer structures (Englert et al., 2007; Englert et al., 2010; Harlander et al., 2017; Wei et al., 2020). DCOI consists of a 630 nm narrow-band filter (2 nm bandwidth), a 9 degree field-of-view lens (f/6), a DASH interferometer with a 25 mm aperture, a Neon lamp for calibration, and a 2048×2048 pixel CCD (13.5 μm per pixel) (Wei et al., 2020; Zhu et al., 2023; Liu et al., 2025). Its thermal stability is maintained within 0.1 K. DCOI measures three-dimensional wind speeds by scanning five directions (zenith and four cardinal directions at 45° zenith angle). Each direction is exposed for 5 minutes, completing a cycle roughly every 25 minutes. DCOI adopts an observation with the smallest error after evening as the reference zero wind speed. The slant LOS speeds are subtracted by the time-regressed projection of vertical speed and then converted to horizontal using the sine of zenith angles. It is worth noting that during auroral events, vertical winds with absolute values exceeding 50 m·s⁻¹ are excluded from the regression, as they contain scattering effects that could introduce additional biases to other directions. DCOI provides two series of meridional wind, two series of zonal wind, and one series of vertical wind.

2.3 Fabry-Perot Interferometer

Fabry-Perot Interferometer (FPI), as a mature solution, conducts comparative observations with DCOI. It features a 630 nm narrow-band filter (2 nm bandwidth), a 2.54 degree field-of-view lens (f/6), a 50 mm aperture etalon with a 7 mm gap, a frequency-stabilized laser for calibration, and a 1024×1024 pixel CCD (13 μm per pixel). FPI uses the same integration time and scanning method as DCOI to obtain horizontal and vertical winds for each cardinal direction and zenith. Details and historical results of FPI are in these references (Yuan et al., 2010; Wu et al., 2014; Yu et al., 2014; Huang et al., 2018; Jiang et al., 2018).

2.4 Scattering radiative transfer model

The model for estimating scattering impact is based on the scattering radiative transfer model and numerical solution by Harding (2017a). It assumes airglow emission undergoes elastic scattering, preserving its wavelength and initial Doppler shift. By specifying airglow brightness distribution, original Doppler shift, lower atmosphere scattering characteristics, and a

157 simplified geometric relationship, the radiation transfer equations (see Appendix A) can be solved to compute the distribution
158 of multiple scattered light and its associated Doppler shift. This enables the wind simulation with atmospheric scattering. A
159 schematic diagram (Fig. 1) illustrates the basic mechanism. To enhance applicability, we have refined several aspects: (1) The
160 upper boundary of the lower atmosphere is set at 40 km to improve the accuracy of the effective extinction path in the initial
161 source function. (2) ~~The LOS speed is used directly instead of the Doppler shift, assuming a constant background temperature.~~
162 The Doppler shift is replaced by LOS speed, with every incident ray from the airglow layer mapped directly to its
163 corresponding LOS speed. (3) ~~After binning different LOS speeds and computing the corresponding scattered light intensity,~~
164 ~~contaminated LOS speeds are calculated via weighted average, simplifying the wind simulation.~~ After slicing the airglow layer
165 into several bins by LOS-speed, the model illuminates one bin per run, records its scattered intensity, then merges all bins with
166 an intensity-weighted average to yield the post-scattered LOS speed. ~~We directly use LOS speed instead of Doppler shift,~~
167 ~~primarily neglecting the interference fringe recognition errors caused by spectral broadening due to temperature variations.~~
168 ~~During auroral events, FPI observations show similar neutral temperatures in all directions, with the northward direction~~
169 ~~occasionally being about 300 K higher (not shown here). Overall, the temperature at mid-latitudes is uniform at the 500 km~~
170 ~~spatial scale, and the variations caused by spectral line broadening can be neglected.~~ The detailed model description is provided
171 in Appendix A.

172 Additionally, the scattering characteristics of the lower atmosphere in our model, including the scattering phase function and
173 optical depth, were derived from Aerosol Robotic Network (AERONET) observations (Holben et al., 2001). We utilized data
174 from the Baotou site (40.9° N, 109.6° E), which is the nearest available site to SIZW, located approximately 180 km away.
175 The total optical depth, accounting for both aerosol and molecular scattering, was calculated using monthly averages and was
176 found to be 0.43 in May and 0.2 in October. The scattering phase function was determined based on AERONET data following
177 the previous method (Harding et al., 2017a). Further details regarding the scattering characteristics are described in Appendix
178 B.




Figure 1: The schematic diagram of the scattering radiative transfer model


The grey shading represents the lower atmospheric layer, with darker hues indicating greater optical depth. The yellow-green fillers represent the relative brightness from the red-line airglow layer. Yellow indicates higher light intensity. The blue-red fillers, which correspond to the relative brightness, represent the Doppler shift type (blue-shift or red-shift) of LOS wind speeds. (a) to (e) represent airglow emissions travelling along different paths, carrying Doppler shifts from outside the line of sight into the interferometer, thereby causing biases in the observations. The model estimates the biases by simulating the distribution of airglow emissions after scattering.

3 Results

Two storms with visible auroras on May 10th and Oct. 10th, 2024, respectively, are used to study the scattering impact. The storm from May 10th to 11th is characterized by its significant magnitude and prolonged duration. Multiple works report this event (Guo et al., 2024; Hajra et al., 2024; Themens et al., 2024), with particular focus on the variations of thermospheric winds (Wang et al., 2025; Zhang et al., 2025) and auroras (Gonzalez-Esparza et al., 2024; Kataoka et al., 2024b; Mikhalev, 2024; Nanjo and Shiokawa, 2024) at mid-latitudes. The storm commenced around 17:00 UT on May 10th and the main phase persisted until 02:00 UT on May 11th. After that, the local night of May 11th in the China region sank into a continuous recovery phase. Another storm from Oct. 10th to 11th is weaker than May's (Ranjan and Pallamraju, 2025; Singh et al., 2025), with the main phase from 18:00 UT on Oct. 10th to 02:00 UT on Oct. 11th. During the two geomagnetic storms with visible auroras, both the DCOI and FPI at SIZW observed atypical winds, characterized by intense horizontal wind differences and downward vertical winds.

197 **3.1 Storm-time wind speed statistics**

198 It is necessary to ascertain whether atypical winds originate from atmospheric scattering with spatially uneven airglow
199 brightness or dynamic processes during storms. To investigate the impact of visible auroras on atypical winds during storms,
200 we made the most of the available observations, tracking DCOI's storm-time observations for nearly a year and FPI's for almost
201 five months. We employed the planetary magnetic index Kp exceeding 3 to identify geomagnetic storms (Yang et al., 2020).
202 Besides, to rule out moonlight and cloud effects, we only used clear sky conditions, which means: (1) excluding cases where
203 the angle between the moon and the line of sight is less than 30 degrees, and (2) excluding cases where large-area thick cloud
204 coverage is visible in DCAI. Additionally, data with standard errors greater than 50 m·s⁻¹ were also excluded. A few aurora
205 events, including Nov. 5th, Dec. 1st, 2023, and Aug. 12th, 2024, that did not meet this criterion were excluded.



206
207 **Figure 2: Storm-time (Kp>3) thermospheric wind speed statistics at SIZW**

208 Figure 2a shows meridional winds observed along two opposite directions (N-S) by FPI, with north-looking in red and south-looking in
209 blue. Figure 2b shows zonal and vertical wind, with east-looking in yellow, west-looking in green, and zenith-ward in black. The
210 northward, eastward, and upward speeds are positive in coordinates. Observations without aurora are shown as points, while those with
211 visible auroras are shown as lines. Figures 2c and 2d are similar but show DCOI data.

212 Figure 2 shows thermospheric wind statistics during geomagnetic storms (Kp>3) at SIZW. The first two panels display FPI
213 data from May to Oct. 2024, while the rest display DCOI data from Nov. 2023 to Oct. 2024. FPI began operation on May 8th,
214 2024, with about half a year less data than DCOI. Observations with no aurora in the field of view are marked as points, while

the two cases with visible auroras are shown as lines. The five observation directions of the interferometer are marked by different colors. During typical storms, horizontal winds consistently increase to around $150 \text{ m}\cdot\text{s}^{-1}$ both equatorward and westward with no significant downward wind. However, under visible auroras, both DCOI and FPI have detected large wind speeds, such as a southward wind of about $600 \text{ m}\cdot\text{s}^{-1}$ and downwelling exceeding $100 \text{ m}\cdot\text{s}^{-1}$. The two series of winds observed along opposite cardinal directions (N-S or E-W) exhibit overt differences, with values exceeding $400 \text{ m}\cdot\text{s}^{-1}$ and contrary directions. This is markedly different from the wind patterns observed during non-aurora storms, where opposite-direction winds do not show significant divergence. Comparing the results of DCOI and FPI, the observations are largely consistent both with and without auroras. The atypical winds observed simultaneously by two interferometers with different principles suggest a systematic error from outside the instruments. Besides, these simultaneous changes appear in five observation directions, all characterized by enhanced negative LOS speeds, indicating likely LOS speed contamination. These factors point more towards scattering impact rather than dynamical processes as the cause. Next, the relationship between scattered light and atypical winds will be investigated through simulation.

3.2 Comparison of observations and simulations

Figure 3 shows the red-line airglow brightness from DCAI (Fig. 3a, 3e), the observed winds from DCOI and FPI (solid lines with different markers in Fig. 3b-3d, 3f-3h) and the simulated winds from the scattering model (dotted lines in Fig. 3b-3d, 3f-3h) during the two nights of May 10th and 11th, 2024, at SIZW, in which the different colors denote distinct directions. The grey lines in the horizontal wind plots represent the average values between opposite cardinal directions. The multi-directional brightness series from DCAI are extracted at 45° zenith angle, consistent with the scanning zenith angle of interferometers. Time intervals with visible auroras are highlighted in red, showing much higher brightness in northward directions than others. Figure 4 supplements the auroral distribution compared to Fig. 3a and 3e. Images from DCAI are projected onto the airglow layer at 250 km. The red circle encloses the actual observations with zenith angles less than 70° , while the values outside are extrapolated. The red dots represent the interferometer's pierce points on the airglow layer at 45° zenith angle.




Figure 3: Observations of aurora and wind speeds, and the scattering model simulation on the nights of May 10th and 11th, 2024, at SIZW

Figure 3a shows the brightness of 8 cardinal directions, all at 45° zenith angle, along with the zenith-ward, extracted from DCAI. The color coding is as follows: red for northern directions, green for east and west, blue for southern directions, black for the zenith, and yellow for the average brightness excluding the three northern directions. Figure 3b shows the meridional wind, with north-looking in red and south-looking in blue, and the average of the two directions in grey. DCOI observations are shown as solid lines with circular dots, FPI as solid lines with rhombus dots, and simulations as dotted lines. Figures 3c, 3d are similar to Fig. 3b, but for zonal and vertical wind, with east-looking in yellow, west-looking in green, and zenith-ward in black. For a more concise figure, if the standard error exceeds 100 m s⁻¹, the point will be filled with black instead of error bar. Figures 3a-3d show data from May 10th, and Fig. 3e-3h from May 11th. The time intervals with visible auroras are marked in red.




Figure 4: Auroral distribution observed by DCAI on the nights of May 10th and 11th, 2024, at SIZW

Images from DCAI (Fig. 4a-4d, 4f-4i) have been projected onto the airglow layer at 250 km. The red circle encloses actual observations with zenith angles $< 70^\circ$, while values outside are extrapolated using zero-order extrapolation. The red dots represent the interferometer's pierce points on the airglow layer at 45° zenith angle. Figures 4e, 4j are similar to Fig. 3a, 3e, with the corresponding images time labelled.

The coastlines in projected DCAI images are made with Natural Earth.


After 18:30 UT on May 10th, as the aurora intensified, both DCOI and FPI detected simultaneous changes in meridional, zonal, and vertical winds. The north-looking red-line brightness at 45° zenith angle exceeded three times that of other directions. The meridional and zonal wind differences between opposite cardinal directions (N-S or E-W) increased. And the winds detected in opposite directions reversed. The maximum meridional difference was close to $800 \text{ m}\cdot\text{s}^{-1}$, while that in zonal exceeded $500 \text{ m}\cdot\text{s}^{-1}$. The downward wind was enhanced by over $100 \text{ m}\cdot\text{s}^{-1}$. These four aforementioned variables, that red-line brightness, the meridional and zonal differences, and downwelling, increased almost simultaneously, peaked at 19:05 UT, and then decayed.

260 Moreover, the average meridional wind, derived from averaging opposite cardinal directions, continuously enhanced
261 equatorward to around 400 m/s, while the average zonal wind enhanced westward to around 100 m/s. Unlike the single-
262 direction results that peaked at 19:05 UT, the average wind varied steadily, consistent with storm-time circulation. Compared
263 to the average wind, the separated horizontal and enhanced vertical winds are atypical. Even with travelling atmospheric
264 disturbances (TADs) superimposed on storm-time circulation, phase lags between horizontal and vertical components would
265 be expected (Hernandez and Roble, 1976; Ishii et al., 1999), but none were observed. Thus, the atypical winds do not appear
266 to be the result of a dynamical process. During the recovery phase on May 11th, the aurora was present throughout the night
267 but much weaker than on May 10th, as seen in Fig. 4. Both DCOI and FPI showed westward and equatorward horizontal winds
268 with no significant downward wind. There was a meridional difference of about 100 to 300 m·s⁻¹ persisted throughout the
269 night, with no zonal difference.

270 Subsequently, we used the scattering model to explore the relationship between red-line brightness variations and atypical
271 winds through atmospheric scattering. On May 10th, a fixed wind vector of 100 m·s⁻¹ westward and 400 m·s⁻¹ southward was
272 set as the input. This assumed wind was kept constant over time and spatially uniform, with no vertical components. On May
273 11th, a fixed wind vector of 200 m·s⁻¹ westward and 100 m·s⁻¹ southward was used, again with no vertical component. These
274 values are chosen based on average observed wind speeds to approximate storm-time circulation. Although the specific values
275 may deviate, the main wind directions remain consistent. The storm-time enhancement of vertical winds may be caused by
276 scattering rather than representing real winds, so we set it to zero. It is worth noting that we neglect the variation of background
277 wind in the model inputs, due to uncertainty regarding whether the observed wind variations are biased. Moreover, using fixed
278 wind speeds allows us to highlight the impact of red-line brightness variations and determine the presence of scattering effects.

279 As shown by the dotted lines in Fig. 3b-3e, the simulations with scattering impact generally match the observations on May
280 10th. Simulated horizontal wind differences and downwelling increase initially with the aurora, peak around 19:15 UT, and
281 then decay. The simulated wind speed variations lag the observations by about 10 minutes. The lag may be due to the relatively
282 rough 25 minute scanning cycle or DCAI underestimating airglow brightness at the field of view's edge, leading to inaccurate
283 capture of scattering enhancement start time. Numerically, the simulated zonal and vertical winds match observations more
284 closely than the meridional wind. The simulated meridional difference is smaller than the observed difference, and the north-
285 looking simulation remains closer to the default value, unlike the equatorward-biased observation. The preset fixed wind may
286 influence the meridional simulation, as it does not follow the equatorward enhancement of the meridional wind. Other factors
287 beyond scattering impact might also have an impact, such as the lift of red-line emission profile and the spectral pollution
288 caused by auroras (Makela et al., 2014), to be discussed later. Additionally, the model simulates a similar intense downward
289 wind as observed under the preset zero vertical wind. This indicates that the vertical wind is significantly affected by scattering.
290 This is why the intense vertical wind is not subtracted when converting interferometer LOS speed to horizontal wind, to prevent
291 error propagation. For May 11th, due to the weak but continuous aurora, the simulation shows weak horizontal differences and

292 slight downward winds throughout the night. Compared with the observations, the simulation shows smaller meridional
 293 differences. It also indicates zonal differences and downward winds, which are not evident in the observations. The poorer
 294 simulation on May 11th may be due to misalignment between dominant horizontal winds and airglow brightness gradients,
 295 which will be discussed later. Additionally, there may be issues with the zero wind calibration. When auroras are present
 296 throughout the night, the vertical wind, which includes scattering biases, may have been used to calibrate zero wind speed. It
 297 likely masks the scattering impact in the observations and explains the discrepancies in the simulation.



298
 299 **Figure 5: Observations of aurora and wind speeds, and the scattering model simulation on the nights of Oct. 10th and 11th, 2024, at**
 300

SIZW

301 Figure 5 is analogous to Fig. 3, but for Oct. 10th and 11th.




Figure 6: Auroral distribution observed by DCAI on the nights of Oct. 10th and 11th, 2024, at SIZW

Figure 6 is analogous to Fig. 4, but for Oct. 10th and 11th.

302

303

304

305 Figures 5 and 6 show another case from Oct. 10th to 11th, analogous to Figures 3 and 4. On Oct. 10th, the aurora appeared at
 306 17:15 UT, expanded southward and increased in brightness, peaking first at 18:30 UT before decaying and then increasing
 307 again from 20:30 UT until sunrise. The second peak was brighter than the first (Fig. 6). Similar to the storm in May, once
 308 aurora appeared, both DCOI and FPI observed atypical winds, with synchronous meridional and zonal differences and
 309 downward enhancements in vertical wind. These atypical winds also exhibited two peaks, around 18:30 UT and 21:00 UT.
 310 The horizontal winds observed in opposite cardinal directions were basically in opposition. During visible aurora periods,
 311 DCOI and FPI showed a 50 to 100 m·s⁻¹ difference in vertical wind but consistent variation trends. Moreover, the average
 312 horizontal wind between opposite cardinal directions was dominantly equatorward and westward, which also had two peaks.
 313 On Oct. 11th, the storm had passed, and no visible aurora was present. The increase in brightness around 13:00 UT was due to
 314 moonset in the southwest. There were no significant horizontal differences or downward winds, consistent with geomagnetic
 315 quiet conditions.

316 We set a fixed $100 \text{ m}\cdot\text{s}^{-1}$ westward with $400 \text{ m}\cdot\text{s}^{-1}$ southward wind vector for Oct. 10th, and $100 \text{ m}\cdot\text{s}^{-1}$ westward with $200 \text{ m}\cdot\text{s}^{-1}$
317 southward wind vector for Oct. 11th in the model, respectively, with no vertical component. The simulation on Oct. 10th exhibits
318 two peaks. The second peak in the simulation is consistent with the observations better, while the first peak, although capturing
319 the trend, is significantly underestimated in magnitude. This discrepancy in the simulation may relate to optical depth, aurora
320 brightness, and background wind changes. The optical depth in Oct. is about half that of May, and simulations underestimate
321 observed values. As in previous studies (Harding et al., 2017b), optical depth can affect the scattering model response. On Oct.
322 10th, the first aurora brightening is weaker than the second. When red-line brightness differences are small, the model response
323 tends to be lower. The impact of optical depth and red-line brightness on the model will be discussed later. Additionally,
324 noticeable fluctuations in the average meridional wind on this day may also contribute to the deviation in the model with fixed
325 initial wind. The north-looking wind speed varied dramatically between 19:00 UT and 21:00 UT along with the aurora, which
326 may also be related to spectral contamination beyond scattering impact. This spectral contamination arises from fast O atoms
327 generated by low-energy O^+ ion precipitation in the auroral region, which occurs at higher altitudes. This issue introduces an
328 additional spectral shift that compromises wind retrieval (Makela et al., 2014). This exceeds the simulation range of the model,
329 thereby causing the discrepancy.


330 4 Discussions

331 In this study, we modelled the transmission of scattered airglow emission~~scattered airglow transmission~~ in the lower
332 atmosphere. Post-scattering wind speeds were calculated based on assumed initial wind speeds weighted by both scattered and
333 direct intensities. The model basically captured the temporal variations of horizontal wind differences and downward
334 enhancements associated with varying auroral brightness, suggesting the contribution of scattering mechanisms to atypical
335 winds. However, the simulation of scattering has certain limitations and characteristics: (1) The differences between simulation
336 and observation vary across different directions. (2) The simulated values sometimes exhibit significant numerical deviations
337 from observations. Could this be related to model errors? ~~(3) When using average wind speeds and individual direction wind~~
338 ~~speeds with scattering impact, what additional considerations are needed? Next, we will discuss these three issues in detail,~~
339 ~~focusing on the working principle of the model, the errors involved, and the insights that the simulation can provide regarding~~
340 ~~the impact of scattering. Next, we will discuss the working principle of the scattering model and the errors involved in the~~
341 simulation.

342 4.1 Core working principle of the scattering model

343 As shown in Fig. 3 and Fig. 5, observed winds respond differently to scattering across directions, especially on May 10th and
344 Oct. 10th with stronger auroras. Although the scattering model has numerical errors, the simulations also show directional
345 differences in scattering-induced biases. Both observations and simulations indicate that the meridional wind responds the
346 most, followed by the zonal wind, while the vertical wind responds the least. Since vertical and horizontal wind speeds are

347 derived from the projection of LOS wind speeds, this essentially reflects the non-uniform response of LOS speeds to scattering.
 348 This directional inhomogeneity of scattering impact aligns with previous studies. Harding et al. (2017a) simulated scattering
 349 effects under auroral conditions, using northward observations as the initial winds. They noted that this direction experiences
 350 minimal scattering contamination due to facing the brighter region. Abreu et al. (1983) used a meridional one-dimensional
 351 model, finding that LOS wind speeds near intense airglow brightness gradients and with weaker airglow intensity are more
 352 susceptible to contamination. They also showed that scattering-induced biases are minimal in the vertical direction, as the
 353 shorter atmospheric path length limits the opportunity for scattering. ~~because the light path through the atmosphere is shortest,~~
 354 ~~reducing scattering opportunities.~~ In this study, we further explore the scattering impact as a function of azimuth angle,
 355 revealing the formation of horizontal differences.



356
 357 **Figure 7: Post-scattering LOS speeds at 45° zenith angle**

358 The figure shows the post-scattering line-of-sight speeds at 45° zenith angle for various directions over time, with panels for May 10th,
 359 May 11th, and October 10th, respectively.

360 Figure 7 shows post-scattering LOS speeds at 45° zenith angle in the simulations on three aurora nights, with the vertical axis
 361 indicating cardinal directions derived from azimuth angles. Concerning the auroral variations in Fig. 4 and Fig. 8, LOS speeds
 362 show diffusion during auroral events. Negative LOS speeds initially concentrated northward, spread westward and eastward,

363 expanding horizontal coverage. Positive LOS speeds initially in the southward direction shrink. When converted to horizontal
 364 wind speeds, these changes lead to increased horizontal differences, or the false convergence caused by scattering, in other
 365 words. The northward LOS speed changes slightly, the southward speed changes the most and nearly reverses, while the
 366 eastward and westward speeds are intermediate.




Figure 8: Proportion of scattered light at 45° zenith angle

369 The proportion is the ratio of scattered light intensity to the total light intensity (both scattered and direct) in the simulation. The figure
 370 shows this proportion at 45° zenith angle for various directions over time, with panels for May 10th, May 11th, and October 10th,
 371 respectively.

372 Figure 8 shows the ratio of scattered light intensity to total light intensity at a fixed 45° zenith angle calculated by the scattering
 373 model. Consistent with the schematic diagram (Fig. 1), the scattered intensity is the sum of all injected directions, and the total
 374 light intensity includes the direct component on this basis. Without auroras, the scattering proportion is typically below 0.4
 375 and varies with atmospheric scattering capability. However, during auroral events, the scattering proportion in some directions
 376 can increase to 0.5 or higher. The northward scattering proportion increases the least and remains much lower than in other
 377 directions. In contrast, the scattering proportion is significantly enhanced in directions ranging from 135° to 180° away from

378 northward. The aurora appears in the north, resulting in much higher northern brightness. After atmospheric scattering, light
379 from the north diffuses into surrounding directions, increasing the scattering proportion. Because the north itself has strong
380 direct light, its scattering proportion remains small. In the model, we assume that ~~without considering temperature-related~~
381 ~~spectral broadening~~, stronger light rays dominate interference fringe identification (Wei et al., 2020), thereby determining the
382 Doppler shift or LOS speeds. The lower scattering proportion in the north allows northward observations to retain more LOS
383 speeds from themselves, while other directions experience greater LOS speed contamination from the north. Therefore, the
384 northward simulation is closest to the default inputs, the southward simulation deviates the most, and the eastward and
385 westward simulations lie in between. Additionally, the scattering impact should also consider the initial LOS speeds in the
386 brighter region. Despite the high scattering proportion on May 11th shown in Fig. 7, the simulated LOS speed changes in Fig.
387 8 are minimal. This is because of the smaller LOS speed in the auroral region on that day, resulting in less contamination
388 spread to other directions.

389 The core working principle of the scattering model relies on the relationship between airglow brightness and background LOS
390 wind speeds. As Harding (2017a) noted, scattering requires a bright sky region with large LOS wind speeds. Firstly, spatially
391 uneven airglow brightness is a prerequisite for significant scattering. The brightest airglow area contributes most to scattered
392 light intensity, and its Doppler shifts determine the LOS biases in other directions. This principle allows a rough assessment
393 of scattering impact without model computation when airglow is uneven. Locate the brightest region and its Doppler shift type,
394 as the same Doppler shift type will likely appear in other directions. Blue-shift dominance indicates increased downward wind
395 and horizontal deviations opposite to the line of sight, resembling convergence, while red-shift dominance resembles
396 divergence. Minimal scattering-induced biases occur if the scattering proportion is very low due to uniform airglow brightness,
397 or if the LOS velocity in the bright region is near zero (i.e., the wind speed is perpendicular to the line of sight). Previous
398 observations can be directly verified by this principle and are basically in line with it (Hernandez and Roble, 1976; Price et al.,
399 1995; Ishii et al., 1999; Ishii et al., 2001; Makela et al., 2014). Unfortunately, scattering impact can complicate dynamic
400 analysis. In polar regions, auroras are characterized by green-line emissions, and thermospheric winds are significantly
401 influenced by ion drag, where scattering effects may not be prominent. In contrast, mid-latitudes have mainly red-line auroras
402 with large-scale uniform circulation, making the scattering impact more pronounced and distinguishable.

403 4.2 Errors of the scattering model

404 The model also exhibits certain errors and limitations. Scattering-induced biases in observations have nearly similar
405 magnitudes on May 10th and Oct. 10th. However, with the same 100 m·s⁻¹ westward and 400 m·s⁻¹ southward wind input, the
406 scattering-induced biases in the May case are significantly larger in magnitude and closer to reality compared to October. In
407 Fig. 7, the simulated LOS speeds show a larger diffusion range for the May 10th case compared to October. In Fig. 8, the
408 scattering proportion for May 10th is consistently higher than that for October. We attribute this difference mainly to the distinct
409 optical depths in the two months, which are 0.43 and 0.2, respectively. Optical depth reflects atmospheric extinction capability

410 and is mainly related to aerosol content (see Appendix B). It primarily affects the extinction process and influences the
411 magnitude of scattering-induced biases by altering the proportion of scattered light. When the optical depth is artificially
412 increased to a higher value, such as around 0.6, the model more closely matches the Oct. observations. We find that the model
413 underestimates scattering effects when the optical depth is low.

414 In the October 10th event, the simulated scattering-induced biases are inconsistent between the two auroral brightness peaks.
415 In Fig. 7, the LOS speed variation is larger during the second peak, and in Fig. 8, the scattering proportion is greater. This is
416 because the scattering proportion is susceptible to errors in scattered and direct light intensities. DCAI does ~~not~~ correct for
417 Rayleigh units, leading to significant errors in regions with large zenith angles. The model ~~can't~~ cannot fully eliminate the stray
418 light caused by the glass dome when separating the initial direct and scattered light from DCAI images (Harding et al., 2017a),
419 resulting in errors. In our experiments, if the stray light effect is ~~not~~ subtracted as described in Appendix B, the model
420 becomes more inert, resulting in a smaller simulated scattering proportion.

421 We also considered the potential influence of thermospheric temperature. FPI data show uniformly elevated thermospheric
422 temperatures in these two storms, with the northward view occasionally about 300 K warmer than the others (not shown here).
423 Because our scattering model does not yet include temperature effects, we cannot quantify how much scattering biases the FPI
424 temperature measurements. In the study of Harding et al. (2017b), wind simulations are temperature-independent, while
425 temperature retrieval relies on the wind. Likewise, we substitute the LOS speed for the Doppler shift and ignore temperature-
426 induced spectral broadening. In principle, thermospheric temperature influences retrieval uncertainty, not the wind speed itself.
427 We remain cautious that ignoring this uncertainty could introduce extra bias if a horizontal temperature gradient is present, but
428 incorporating it would markedly raise the computational cost and remains a task for the future.

429 In our experiments, even after several refinements aimed at mitigating the underestimation, the scattering model still
430 underestimates the observed scattering impact: In our experiments, the scattering model consistently underestimates scattering
431 impact compared to observations, despite several applied enhancements: (1) A single-scattering albedo of 1 was used, ignoring
432 absorption. (2) Stray light effects were removed. (3) Attenuated airglow observations at the edge of DCAI images were
433 extrapolated, enhancing edge scattering. (4) Excessive edge extinction was reduced by correcting the extinction path geometry,
434 increasing the scattered light intensity integral. (5) Zero vertical wind was assumed when converting simulated LOS speeds to
435 geographical wind speeds. We suspect the underestimation arises from how the model integrates the optical depth. Since the
436 integral only includes 10 optical depth layers, with each light ray scattering once per layer and extinguishing once between
437 layers, it may be too crude compared to the real path, underestimating the scattered light. Simply increasing the number of
438 optical depth layers is not effective. We think this may be related to the non-linear variation of atmospheric density with
439 altitude, where optical depth may not vary linearly with height, and the scattering phase function may also change with altitude.
440 To address ~~this~~ these issues, future work should complete the DCAI correction. Additionally, introducing a model of optical

441 depth varying with altitude can increase the number of single-scattering nodes and ensure the geometric accuracy of the
442 extinction path, thereby improving the accuracy of scattered light intensity calculations.

443 Furthermore, these bright region observations do not necessarily reflect the usual 250 km thermospheric wind. ~~correspond to~~
444 ~~the wind speeds of the red-line airglow layer. They could also come from higher auroral layers and not even reflect the neutral~~
445 ~~thermospheric winds near 250 km (Makela et al., 2014).~~ In Fig. 3 and Fig. 5, the north-looking wind observations show
446 unusually high wind speeds, which are significantly different from the simulations. In particular, on October 10th, the north-
447 looking wind speed varied dramatically with the intensity of the northern aurora. During the two auroral peaks, the north-
448 looking wind direction also reversed. ~~The north-looking wind speed changed more than the south-looking wind speed,~~
449 ~~inconsistent with the simulation characteristics. When auroras occur, the north-looking wind speed in the bright region may~~
450 ~~be significantly higher than the average value over zenith and distinct from regions outside the aurora. This could enhance~~
451 ~~scattering effects, which the model fails to simulate.~~ This indicates that the interferometer receives an additional effect when
452 it looks toward the aurora. Kataoka et al. showed that red aurora lifted the red-line emission profile, raising its peak above 300
453 km and brightening the upper part on May 11th (Kataoka et al., 2024b). Consequently, the interferometer can sample winds
454 that are higher and more poleward. Because storm-time surges propagate from the polar region to the equator, these higher,
455 poleward regions are likely to carry stronger equatorward winds. The interferometer may record a larger wind speed toward
456 the aurora. Additionally, spectral contamination from precipitating energetic ions can also bias interferometers (Makela et al.,
457 2014). In other words, the interferometer is partly sensing the speed of non-neutral species, boosting the observed wind. These
458 issues lie beyond what scattering models can reproduce. From the observed pattern, we infer the presence of non-scattering
459 effects, especially in the poleward view. Due to the absence of nearby higher-latitude neutral-wind observations relative to
460 SIZW, quantifying their respective contributions remains challenging.

461 **4.3 Scattering effects on thermospheric wind observation**

462 ~~Given the directional inhomogeneity of scattering impact, the reliability of the commonly utilized average wind and individual~~
463 ~~direction wind observations needs to be considered. Wind speed detected from a less bright region is more doubtful, because~~
464 ~~these regions are more polluted by scattered light. The average speed of opposite cardinal directions will maintain the correct~~
465 ~~time-varying trend of the background thermospheric wind but may deviate in absolute value. Since the LOS speed deviations~~
466 ~~caused by scattering in each direction have the same type, averaging can cancel out the scattering deviations between opposite~~
467 ~~directions, making the average wind closer to the true background thermospheric wind. However, in directions with strong~~
468 ~~light intensity gradients, such as north-south, the deviations in the two directions are different, and averaging will cause the~~
469 ~~wind speed to deviate from the true value. Additionally, the wave-like structures appearing in each individual direction may~~
470 ~~originate from changes in scattering effects caused by variations in auroral intensity, and whether they come from TADs needs~~
471 ~~further verification. The wave-like structures superimposed on the average value may also have their amplitudes enhanced or~~
472 ~~cancelled due to the directionality of scattering deviations, which similarly requires additional validation.~~

473 5 Conclusions

474 This study has further proved that lower atmospheric scattering ~~can contribute to biases in~~ ~~can bias~~ thermospheric wind
475 observation on ground-based optical interferometers. The light scattered from the non-line-of-sight directions of the scope will
476 lead to additional LOS speeds and appear as atypical horizontal differences and vertical wind at geographic coordinates. With
477 a simplified scattering radiative transfer model, we simulate the distribution of airglow intensity after the multiple scattering
478 of the lower atmosphere and estimate the wind observation bias under scattering impact via a weighted average method.
479 Atypical winds under conditions of spatially uneven airglow have been generally captured.

480 We have refined the scattering model in previous research to enhance its computational efficiency. Specifically, we simplified
481 the LOS wind speed simulation and introduced an upper limit for the lower atmosphere to improve the accuracy of the
482 extinction length calculation. The scattering impact can be directly estimated through the relationship between the bright
483 airglow region and the LOS wind speed. The brightest airglow region contributes most to the scattering impact, of which the
484 Doppler shift type determines the LOS biases in other directions. ~~Due to the directional inhomogeneity of scattering effects,~~
485 ~~we propose that in directions with strong light intensity gradients, such as north-south, the average wind speed of opposite~~
486 ~~directions and its fluctuations may be biased.~~ Although the observed winds are affected by scattering when airglow is uneven,
487 they still retain dynamic information, such as the average wind being close to the storm-time circulation. Unfortunately, we
488 lack alternative observational methods to verify the accuracy of the interferometer results. It deserves further study to the
489 extent of scattering impact with more cases and additional instrumental observations.

490 Limited by the accuracy of the model inputs, the scattering model can only simulate the wind features associated with scattering
491 impact under clear sky conditions. It remains incapable of precisely picking out the speed alterations induced by scattering
492 impact. Given that scattering ~~significantly~~ impacts observations during geomagnetic storms accompanied by auroras, it is
493 necessary to quantify these biases to provide accurate data for dynamic studies. Therefore, future efforts can focus on refining
494 the model and its inputs or statistically analyzing scattering biases under different airglow brightness distributions from various
495 stations to gain a clearer understanding of the magnitude of scattering biases.

496 Appendix A

497 In the following appendices, we provide a concise description of the scattering model's operational principles, inputs, and
498 modifications employed in our works. For more detailed solutions, please refer to the article by Harding et al. (Harding et al.,
499 2017a).

500 Based on the radiation transfer theory, Hansen and Travis (1974) and Sobolev (1975) gave the multiple scattering solution.
501 Harding (2017a) extended the initial source function $J_0(\tau, u, \phi)$ to airglow surface source, and corrected the missing
502 normalization factor in the scattering phase function (Eq. (1) to Eq. (3)):

$$u \frac{dI(\tau, u, \phi)}{d\tau} = -I(\tau, u, \phi) + J(\tau, u, \phi) \quad (1)$$

$$J(\tau, u, \phi) = \frac{\omega}{4\pi} \int_0^{2\pi} \int_{-1}^1 P(u, u', \phi, \phi') I(\tau, u', \phi') du' d\phi' + J_0(\tau, u, \phi) \quad (2)$$

$$J_0(\tau, u, \phi) = \frac{\omega}{4\pi} \int_0^{2\pi} \int_0^1 P(u, u', \phi, \phi') \sec(\gamma') f(u', \phi') \exp[-\tau L(u')] du' d\phi' \quad (3)$$

$$L(u) = [(R_e + H_L) \cos(\gamma_\tau) - R_e u] H_L^{-1} \quad (4)$$

$$u = \cos(\zeta) \quad (5)$$

The equations are formulated within an improved local spherical coordinate system, including azimuth ϕ , zenith angle ζ which is represented in cosine form u , and vertical height which is expressed as optical depth τ .

In the case of scattering, as illustrated in Fig. 1, the light intensity along a line of sight, represented by $I(\tau, u, \phi)$, consists of two parts, the direct light (a in Fig. 1) from the same direction, and the aggregate of scattered light (b-e in Fig. 1) from other directions, which is represented by the source function $J(\tau, u, \phi)$. Based on the radiative transfer equation (Eq. (1)), at each optical depth layer, the scattered light intensity received from all directions will be integrated. Simultaneously, the original intensity in the line of sight will be added to the total scattered light. Besides, the extinction in the path will be calculated according to the optical depth.

There are two potential scattering paths in the lower atmosphere: single scattering (b, c in Fig. 1) and multiple scattering (d, e in Fig. 1). The model computes them sequentially via an iterative process. In the initial state, there is no light intensity in the lower atmosphere. Therefore, the single scattering will originate solely from the airglow layer, and the source function $J(\tau, u, \phi)$ will be equivalent to the initial source function $J_0(\tau, u, \phi)$ at this state. By solving Eq. (1), the model can obtain the single scattered intensity in each direction at every optical depth layer, which is the updated source function $J(\tau, u, \phi)$. Then, the multiple scattering can be calculated based on it. Typically, the total scattered intensity will remain relatively constant when accounting for the fourth scattering. By using this iteration, the scattered light and residual direct light in DCAI images can be effectively separated. The residual direct light will subsequently serve as the background intensity distribution for simulating speeds.

In the source function $J(\tau, u, \phi)$, the scattering phase function $P(u, u', \phi, \phi')$ quantifies the relative gain of an incident angle to an exit angle during the scattering process. The reference value is based on a unit-radius sphere, which necessitates the introduction of a factor $\frac{1}{4\pi}$. Furthermore, ω represents the single-scattering albedo, set as 1. The initial source function $J_0(\tau, u, \phi)$ is responsible for introducing the airglow distribution $f(u', \phi')$. Here, $\sec(\gamma')$ represents the secant of the zenith angle at the puncture point of the airglow layer, which helps eliminate the Van Rhijn effect. Additionally, the exponential term with base e is utilized to calculate the equivalent extinction length along an inclined path.

531 It is primarily the lower atmosphere that significantly scatters and absorbs light (He et al., 2021; Li et al., 2022). Therefore,
 532 when computing the extinction length, just employing the cosine of zenith angle u' will lead to an overestimation of the
 533 effective length, as illustrated on the right side of Fig. 1. To address it, we set an upper boundary H_L of the lower atmosphere
 534 at 40 km, assuming an optical depth of zero above this altitude. Using geometric relationships, an equivalent length factor
 535 $L(u)$ can be derived, where R_e means Earth radius, and γ_t represents the zenith angle at the penetration point of 40 km
 536 height. This value can be readily calculated by adjusting the target height of the formula for γ . Inside the lower atmosphere,
 537 we apply a thin-layer approximation, which also utilizes the geometric relationships at the upper boundary.


$$538 \quad V = \frac{V_{dr}I_{dr} + \sum_k V_{sc}(k)I_{sc}(k)}{I_{dr} + \sum_k I_{sc}(k)} \quad (6)$$

539 After working out the background intensity distribution, we partition the LOS speeds at 250 km into several bins. ~~Ignoring the~~
 540 ~~background temperature gradient, LOS speeds can directly correspond to Doppler shifts to simplify the simulation of the~~
 541 ~~Doppler distribution.~~ To simplify simulation, the model directly uses LOS speeds corresponding to Doppler shifts. As roughly
 542 illustrated in Fig. 1, all LOS speeds are categorized into $k=10$ bins valued from highest to lowest, assuming that the LOS
 543 speeds within each bin approximate their mean value V_{sc} . The scattered intensity distribution I_{sc} is computed by extracting
 544 the airglow brightness from the corresponding region of each bin. And, there will be no intensity from other areas during a
 545 single bin's computing. According to Eq. (6), the simulative LOS speed at a specific angle will be an averaged result, where
 546 the original speed V_{dr} is weighted by the direct light intensity I_{dr} , and the additional speed resulting from scattering V_{sc} is
 547 weighted by the scattered light intensity I_{sc} . The model directly uses the average within DCOI's 9 degree field of view as
 548 simulated post-scattering LOS speed of interferometers, since DCOI and FPI have ~~not~~ measured the reception gain of light
 549 outside their fields of view. We find that due to the coarse model grid, the 9 degree average is nearly the same as using the
 550 nearest single-sight observation. Finally, the LOS speed will be converted to horizontal wind, maintaining the assumption of
 551 zero vertical wind to prevent the propagation of scattering biases in the vertical direction.

552 Appendix B

553 This appendix details the scattering model's inputs from measurements, supplementing the second section. The background
 554 airglow brightness for the model comes from DCAI. Image processing includes: (1) dark field exclusion, (2) median filtering
 555 to remove starlight, (3) conversion to the local spherical coordinate, (4) stray light correction, and (5) radial zero-order
 556 extrapolation for regions beyond 70° zenith angle. Stray light results from the scattering of strong incident light by the glass
 557 dome. During quiet nights without auroras, it is weak and uniformly distributed across all LOS directions. We use the azimuthal
 558 average of the nearest quiet night at 45° zenith angle as a reference for weak stray light conditions. After the aurora onset,
 559 stray light brightens all directions. The difference between the darkest direction at 45° zenith angle and the reference value is
 560 considered the additional stray light caused by the aurora and is subtracted from the entire image. The shown airglow images

561 additionally mitigate the Van Rhijn effect through $\sec(\gamma')$ and the atmospheric extinction through $\exp[-\tau L(u')]$ (see
 562 Appendix A). Since the scattering model already includes these processes, no separate treatment is needed. The optical depth
 563 and scattering phase function inputs are shown in Fig. 9. Optical depth is calculated using AERONET's monthly averages,
 564 with interpolation at 630 nm. Since only daytime observations are available, local daytime values are used to represent
 565 nighttime values. Fig. 9a and 9b show monthly average optical depths at local daytime, with total averages of 0.43 and 0.2.
 566 The scattering phase function is a weighted average of molecular and aerosol scattering phase functions from AERONET at
 567 675 nm, which is weighted on the total optical depth of aerosols and molecules.



569 **Figure 9: Optical depth and scattering phase function in May and October**

570 The first two panels show daytime monthly average optical depths, with molecular optical depth amplified for clarity. The rest panels
 571 show the scattering phase functions.

572 **Code availability**

573 The code of DCAI images correction, the scattering model, and the visualization are not publicly available yet. If needed, they
 574 can be obtained by contacting the corresponding authors via email.

575 **Data availability**

576 The data of DCOI and DCAI from the Chinese Meridian Project can be obtained from <https://www.meridianproject.ac.cn/en/>.
577 The FPI data can be obtained by contacting the corresponding authors via email. The data of AERONET can be obtained from
578 <https://aeronet.gsfc.nasa.gov/>. The Kp index provided by GFZ, German Research Centre for Geosciences, can be obtained
579 from <https://kp.gfz-potsdam.de/en/>.

580 **Author contributions**

581 Conceptualization: XW; investigation: XW, GJ; instruments construction and maintenance: YZ, JX, WL, TW, GZ, WY; wind
582 data retrieval: YZ, WL, GZ, TW; airglow images correction: XW; model improvement and programming: XW; visualization:
583 XW; validation: GJ, YZ, JX, WL, TW; writing - original draft preparation: XW; writing-review and editing: GJ, YZ, JX, WL,
584 TW; supervision: GJ, YZ; project administration: GJ, YZ; funding acquisition: YZ. All authors contributed to the revision and
585 improvement of the paper.

586 **Competing interests**

587 The authors declare that they have no conflict of interest.

588 **Acknowledgements**

589 We appreciate all the funding from the National Key R&D program of China (2023YFB3905100), the Project of Stable Support
590 for Youth Team in Basic Research Field, CAS (YSBR-018), the National Natural Science Foundation of China (42174212),
591 the Chinese Meridian Project, and the Specialized Research Fund for State Key Laboratories. We acknowledge the use of data
592 from the Chinese Meridian Project. We thank all the builders and maintainers of DCOI, FPI, and DCAI of Siziwang station.
593 We thank Lingli Tang for their effort in establishing and maintaining AOE_Baotou site of AERONET.

594 **Financial support**

595 This work was supported by the National Key R&D program of China (2023YFB3905100), the Project of Stable Support for
596 Youth Team in Basic Research Field, CAS (YSBR-018), the National Natural Science Foundation of China (42174212), the
597 Chinese Meridian Project, and the Specialized Research Fund for State Key Laboratories.

- Abreu, V. J., Schmitt, G. A., Hays, P. B., Meriwether, J. W., Tepley, C. A., and Cogger, L. L.: Atmospheric Scattering Effects on Ground-Based Measurements of Thermospheric Winds, *Planet Space Sci*, 31, 303-310, doi:10.1016/0032-0633(83)90080-6, 1983.
- Anderson, C., Conde, M., and McHarg, M. G.: Neutral thermospheric dynamics observed with two scanning Doppler imagers: 2. Vertical winds, *Journal of Geophysical Research: Space Physics*, 117, doi:10.1029/2011ja017157, 2012.
- Ashburn, E. V.: The Effect of Rayleigh Scattering and Ground Reflection Upon the Determination of the Height of the Night Airglow, *J Atmos Terr Phys*, 5, 83-91, doi:10.1016/0021-9169(54)90012-4, 1954.
- Biondi, M. A. and Feibelman, W. A.: Twilight and nightglow spectral line shapes of oxygen λ 6300 and λ 5577 radiation, *Planet Space Sci*, 16, 431-443, doi:10.1016/0032-0633(68)90158-X, 1968.
- Biondi, M. A., Sipler, D. P., Zipf, M. E., and Baumgardner, J. L.: All-Sky Doppler Interferometer for Thermospheric Dynamics Studies, *Appl Optics*, 34, 1646-1654, doi:10.1364/Ao.34.001646, 1995.
- Burnside, R. G., Herrero, F. A., Meriwether, J. W., and Walker, J. C. G.: Optical observations of thermospheric dynamics at Arecibo, *Journal of Geophysical Research: Space Physics*, 86, 5532-5540, doi:10.1029/JA086iA07p05532, 1981.
- Burnside, R. G. and Tepley, C. A.: Optical Observations of Thermospheric Neutral Winds at Arecibo between 1980 and 1987, *J Geophys Res-Space*, 94, 2711-2716, doi:10.1029/JA094iA03p02711, 1989.
- Conde, M. and Smith, R. W.: Mapping Thermospheric Winds in the Auroral-Zone, *Geophysical Research Letters*, 22, 3019-3022, doi:10.1029/95gl02437, 1995.
- Conde, M., Craven, J. D., Immel, T., Hoch, E., Stenbaek-Nielsen, H., Hallinan, T., Smith, R. W., Olson, J., Sun, W., Frank, L. A., and Sigwarth, J.: Assimilated observations of thermospheric winds, the aurora, and ionospheric currents over Alaska, *Journal of Geophysical Research: Space Physics*, 106, 10493-10508, doi:10.1029/2000ja000135, 2001.
- Crickmore, R. I., Dudeney, J. R., and Rodger, A. S.: Vertical Thermospheric Winds at the Equatorward Edge of the Auroral Oval, *J Atmos Terr Phys*, 53, 485-492, doi:10.1016/0021-9169(91)90076-J, 1991.
- Dhadly, M., Emmert, J., Drob, D., Conde, M., Doornbos, E., Shepherd, G., Makela, J., Wu, Q., Niciejewski, R., and Ridley, A.: Seasonal dependence of northern high-latitude upper thermospheric winds: A quiet time climatological study based on ground-based and space-based measurements, *Journal of Geophysical Research: Space Physics*, 122, 2619-2644, doi:10.1002/2016JA023688, 2017.
- Emmert, J. T., Fejer, B. G., Fesen, C. G., Shepherd, G. G., and Solheim, B. H.: Climatology of middle- and low-latitude daytime region disturbance neutral winds measured by Wind Imaging Interferometer (WINDII), *J Geophys Res-Space*, 106, 24701-24712, doi:10.1029/2000ja000372, 2001.
- Emmert, J. T., Faivre, M. L., Hernandez, G., Jarvis, M. J., Meriwether, J. W., Niciejewski, R. J., Sipler, D. P., and Tepley, C. A.: Climatologies of nighttime upper thermospheric winds measured by ground-based Fabry-Perot interferometers during geomagnetically quiet conditions: 1. Local time, latitudinal, seasonal, and solar cycle dependence, *J Geophys Res-Space*, 111, Artn A12302, doi:10.1029/2006ja011948, 2006.
- Englert, C. R., Babcock, D. D., and Harlander, J. M.: Doppler asymmetric spatial heterodyne spectroscopy (DASH): concept and experimental demonstration, *Appl Optics*, 46, 7297-7307, doi:10.1364/Ao.46.007297, 2007.
- Englert, C. R., Harlander, J. M., Emmert, J. T., Babcock, D. D., and Roesler, F. L.: Initial ground-based thermospheric wind measurements using Doppler asymmetric spatial heterodyne spectroscopy (DASH), *Optics Express*, 18, 27416-27430, doi:10.1364/Oe.18.027416, 2010.
- Fejer, B. G., Emmert, J. T., and Sipler, D. P.: Climatology and storm time dependence of nighttime thermospheric neutral winds over Millstone Hill, *Journal of Geophysical Research: Space Physics*, 107, doi:10.1029/2001ja000300, 2002.
- Friedman, J. F. and Herrero, F. A.: Fabry-Perot interferometer measurements of thermospheric neutral wind gradients and reversals at Arecibo, *Geophysical Research Letters*, 9, 785-788, doi:10.1029/GL009i007p00785, 1982.
- Gonzalez-Esparza, J. A., Sanchez-Garcia, E., Sergeeva, M., Corona-Romero, P., Gonzalez-Mendez, L. X., Valdes-Galicia, J. F., Aguilar-Rodriguez, E., Rodriguez-Martinez, M., Ramirez-Pacheco, C., Castellanos, C. I., Pazos, M., Mendoza, B., Gatica-Acevedo, V. J., Melgarejo-Morales, A., Caraballo, R., Andrade-Mascote, E., Villanueva-Hernandez, P., Bonifaz-Alfonzo, R., Sierra, P., Romero-Hernandez, E., Peralta-Mendoza, I., Perez-Tijerina, E., Mejia-Ambriz, J. C., Guerrero-Peña, C., Caccavari, A., Cifuentes-Nava, G., and Hernandez-Quintero, E.: The Mother's Day Geomagnetic Storm on 10

646 May 2024: Aurora Observations and Low Latitude Space Weather Effects in Mexico, *Space Weather-the International*
 647 *Journal of Research and Applications*, 22, ARTN e2024SW0041111, doi:10.1029/2024SW004111, 2024.
 648 Guo, W. and McEwen, D. J.: Vertical winds in the central polar cap, *Geophysical Research Letters*, 30, Artn 1725,
 649 doi:10.1029/2003gl017124, 2003.
 650 Guo, X., Zhao, B. Q., Yu, T. T., Hao, H. L., Sun, W. J., Wang, G. J., He, M. S., Mao, T., Li, G. Z., and Ren, Z. P.: East-West
 651 Difference in the Ionospheric Response During the Recovery Phase of May 2024 Super Geomagnetic Storm Over the
 652 East Asian, *J Geophys Res-Space*, 129, ARTN e2024JA033170, doi:10.1029/2024JA033170, 2024.
 653 Hajra, R., Tsurutani, B. T., Lakhina, G. S., Lu, Q. M., and Du, A. M.: Interplanetary Causes and Impacts of the 2024 May
 654 Superstorm on the Geosphere: An Overview, *Astrophys J*, 974, ARTN 264, doi:10.3847/1538-4357/ad7462, 2024.
 655 Hansen, J. E. and Travis, L. D.: Light-Scattering in Planetary Atmospheres, *Space Science Reviews*, 16, 527-610, doi:
 656 10.1007/Bf00168069, 1974.
 657 Harding, B. J., Makela, J. J., Qin, J., Fisher, D. J., Martinis, C. R., Noto, J., and Wrasse, C. M.: Atmospheric scattering effects
 658 on ground-based measurements of thermospheric vertical wind, horizontal wind, and temperature, *Journal of Geophysical*
 659 *Research: Space Physics*, 122, 7654-7669, doi:10.1002/2017JA023942, 2017a.
 660 Harding, B. J., Qin, J., and Makela, J. J.: Ground-Based Optical Measurements of Quiet Time Thermospheric Wind and
 661 Temperature: Atmospheric Scattering Corrections, *Journal of Geophysical Research: Space Physics*, 122,
 662 doi:10.1002/2017JA024705, 2017b.
 663 Harlander, J. M., Englert, C. R., Brown, C. M., Marr, K. D., Miller, I. J., Zastera, V., Bach, B. W., and Mende, S. B.: Michelson
 664 Interferometer for Global High-Resolution Thermospheric Imaging (MIGHTI): Monolithic Interferometer Design and
 665 Test, *Space Science Reviews*, 212, 601-613, doi:10.1007/s11214-017-0374-4, 2017.
 666 Hays, P. B., Killeen, T. L., Spencer, N. W., Wharton, L. E., Roble, R. G., Emery, B. A., Fullerrowell, T. J., Rees, D., Frank,
 667 L. A., and Craven, J. D.: Observations of the Dynamics of the Polar Thermosphere, *J Geophys Res-Space*, 89, 5597-5612,
 668 doi:10.1029/JA089iA07p05597, 1984.
 669 He, Q. F., Fang, Z., Shoshanim, O., Brown, S. S., and Rudich, Y.: Scattering and absorption cross sections of atmospheric
 670 gases in the ultraviolet-visible wavelength range (307-725 nm), *Atmos Chem Phys*, 21, 14927-14940, doi:10.5194/acp-
 671 21-14927-2021, 2021.
 672 Hernandez, G. and Roble, R. G.: Direct Measurements of Nighttime Thermospheric Winds and Temperatures .2. Geomagnetic
 673 Storms, *J Geophys Res-Space*, 81, 5173-5181, doi:10.1029/JA081i028p05173, 1976.
 674 Holben, B. N., Tanré, D., Smirnov, A., Eck, T. F., Slutsker, I., Abuhassan, N., Newcomb, W. W., Schafer, J. S., Chatenet, B.,
 675 Lavenue, F., Kaufman, Y. J., Castle, J. V., Setzer, A., Markham, B., Clark, D., Frouin, R., Halthore, R., Karnieli, A.,
 676 O'Neill, N. T., Pietras, C., Pinker, R. T., Voss, K., and Zibordi, G.: An emerging ground-based aerosol climatology::
 677 Aerosol optical depth from AERONET, *J Geophys Res-Atmos*, 106, 12067-12097, doi:10.1029/2001jd900014, 2001.
 678 Huang, C., Xu, J. Y., Zhang, X. X., Liu, D. D., Yuan, W., and Jiang, G. Y.: Mid-latitude thermospheric wind changes during
 679 the St. Patrick's Day storm of 2015 observed by two Fabry-Perot interferometers in China, *Advances in Space Research*,
 680 61, 1873-1879, doi:10.1016/j.asr.2017.10.013, 2018.
 681 Innis, J. L., Greet, P. A., Murphy, D. J., Conde, M. G., and Dyson, P. L.: A large vertical wind in the thermosphere at the
 682 auroral oval/polar cap boundary seen simultaneously from Mawson and Davis, Antarctica, *J Atmos Sol-Terr Phy*, 61,
 683 1047-1058, doi:10.1016/S1364-6826(99)00060-7, 1999.
 684 Ishii, M., Oyama, S., Nozawa, S., Fujii, R., Sagawa, E., Watari, S., and Shinagawa, H.: Dynamics of Neutral Wind in the polar
 685 region observed with two Fabry-Perot interferometers, *Earth Planets Space*, 51, 833-844, doi:10.1186/Bf03353242, 1999.
 686 Ishii, M., Conde, M., Smith, R. W., Krynicki, M., Sagawa, E., and Watari, S.: Vertical wind observations with two Fabry-
 687 Perot interferometers at Poker Flat, Alaska, *Journal of Geophysical Research: Space Physics*, 106, 10537-10551,
 688 doi:10.1029/2000JA900148, 2001.
 689 Jiang, G., Xu, J., Wang, W., Yuan, W., Zhang, S., Yu, T., Zhang, X., Huang, C., Kerr, R. B., Noto, J., Li, J., Liu, W., and Li,
 690 Q.: A Comparison of Quiet Time Thermospheric Winds Between FPI Observations and Model Calculations, *Journal of*
 691 *Geophysical Research: Space Physics*, 123, 7789-7805, doi:10.1029/2018JA025424, 2018.
 692 Kataoka, R., Miyoshi, Y., Shiokawa, K., Nishitani, N., Keika, K., Amano, T., and Seki, K.: Magnetic Storm-Time Red Aurora
 693 as Seen From Hokkaido, Japan on 1 December 2023 Associated With High-Density Solar Wind, *Geophysical Research*
 694 *Letters*, 51, e2024GL108778, doi:10.1029/2024GL108778, 2024a.

Kataoka, R., Reddy, S. A., Nakano, S., Pettit, J., and Nakamura, Y.: Extended magenta aurora as revealed by citizen science, *Scientific Reports*, 14, 25849, doi:10.1038/s41598-024-75184-9, 2024b.

Killeen, T. L., Won, Y. I., Niciejewski, R. J., and Burns, A. G.: Upper Thermosphere Winds and Temperatures in the Geomagnetic Polar-Cap - Solar-Cycle, Geomagnetic-Activity, and Interplanetary Magnetic-Field Dependencies, *J Geophys Res-Space*, 100, 21327-21342, doi:10.1029/95ja01208, 1995.

Li, J., Carlson, B. E., Yung, Y. L., Lv, D. R., Hansen, J., Penner, J. E., Liao, H., Ramaswamy, Kahn, R. A., Zhang, P., Dubovik, O., Ding, A. J., Lacis, A. A., Zhang, L., and Dong, Y. M.: Scattering and absorbing aerosols in the climate system, *Nat Rev Earth Env*, 3, 363-379, doi:10.1038/s43017-022-00296-7, 2022.

Li, W. B., Liu, L. B., Chen, Y. D., Yang, Y. Y., Han, T. W., Ding, F., Le, H. J., and Zhang, R. L.: Multi-Instruments Observation of Ionospheric-Thermospheric Dynamic Coupling Over Mohe (53.5°N, 122.3°E) During the April 2023 Geomagnetic Storm, *J Geophys Res-Space*, 128, ARTN e2023JA032141, doi:10.1029/2023JA032141, 2023.

Liu, W. J., Zhu, Y. J., Xu, J. Y., Li, Q. Z., Yuan, W., Zhu, G. Y., Wang, T. C., Yang, G. T., Du, L. F., Liu, S. Y., and Li, F. Q.: Validation of Neutral Wind in the Mesopause Measured by a Dual-Channel Optical Interferometer (DCOI) Network of the Chinese Meridian Project, *Space Weather-the International Journal of Research and Applications*, 23, ARTN e2025SW004468, doi:10.1029/2025SW004468, 2025.

Ma, L. X., Yu, Y. Q., Ding, X. Q., Liu, X. Y., An, D. P., Zhou, C. L., Cao, J. B., and Shiokawa, K.: Mid-Latitude Auroras and Energetic Particle Precipitation Occurred Unusually in a Moderate Magnetic Storm on 1 December 2023, *Geophysical Research Letters*, 51, ARTN e2024GL110764, doi:10.1029/2024GL110764, 2024.

Makela, J. J., Harding, B. J., Meriwether, J. W., Mesquita, R., Sanders, S., Ridley, A. J., Castellez, M. W., Ciocca, M., Earle, G. D., Frisell, N. A., Hampton, D. L., Gerrard, A. J., Noto, J., and Martinis, C. R.: Storm time response of the midlatitude thermosphere: Observations from a network of Fabry-Perot interferometers, *Journal of Geophysical Research: Space Physics*, 119, 6758-6773, doi:10.1002/2014JA019832, 2014.

Mikhalev, A.: Auroras during extreme geomagnetic storms: Some features of mid-latitude aurora on February 11, 1958, *Solar-Terrestrial Physics*, 55-61, 10.12737/stp-102202406, 2024.

Nakajima, H., Okano, S., Fukunishi, H., and Ono, T.: Observations of thermospheric wind velocities and temperatures by the use of a Fabry-Perot Doppler imaging system at Syowa Station, Antarctica, *Appl Optics*, 34, 8382-8395, doi:10.1364/Ao.34.008382, 1995.

Nanjo, S. and Shiokawa, K.: Spatial structures of blue low-latitude aurora observed from Japan during the extreme geomagnetic storm of May 2024, *Earth Planets Space*, 76, ARTN 156, doi:10.1186/s40623-024-02090-9, 2024.

Price, G. D., Smith, R. W., and Hernandez, G.: Simultaneous Measurements of Large Vertical Winds in the Upper and Lower Thermosphere, *J Atmos Terr Phys*, 57, 631-643, doi:10.1016/0021-9169(94)00103-U, 1995.

Ranjan, A. K. and Pallamraju, D.: Latitudinal Distribution of Thermospheric Nitric Oxide (NO) Infrared Radiative Cooling During May and October 2024 Geomagnetic Storms, *J Geophys Res-Space*, 130, ARTN e2024JA033559, doi:10.1029/2024JA033559, 2025.

Rees, D., Smith, M. F., and Gordon, R.: The Generation of Vertical Thermospheric Winds and Gravity-Waves at Auroral Latitudes .2. Theory and Numerical Modeling of Vertical Winds, *Planet Space Sci*, 32, 685, doi:10.1016/0032-0633(84)90093-X, 1984.

Sakanoi, T., Fukunishi, H., Okano, S., Sato, N., Yamagishi, H., and Yukimatu, A. S.: Dynamical coupling of neutrals and ions in the high-latitude *F* region: Simultaneous FPI and HF radar observations at Syowa Station, Antarctica, *Journal of Geophysical Research: Space Physics*, 107, doi:10.1029/2001JA007530, 2002.

Shinagawa, H. and Oyama, S.: A two-dimensional simulation of thermospheric vertical winds in the vicinity of an auroral arc, *Earth Planets Space*, 58, doi:10.1186/BF03352007, 2006.

Shiokawa, K., Katoh, Y., Satoh, M., Ejiri, M. K., and Ogawa, T.: Integrating-sphere calibration of all-sky cameras for nightglow measurements, *Adv Space Res*, 26, 1025-1028, doi:10.1016/S0273-1177(00)00052-1, 2000.

Singh, R., Scipión, D. E., Kuyeng, K., Condor, P., Flores, R., Pacheco, E., de la Jara, C., and Manay, E.: Ionospheric Responses to an Extreme (G5-Level) Geomagnetic Storm Using Multi-Instrument Measurements at the Jicamarca Radio Observatory on 10-11 October 2024, *J Geophys Res-Space*, 130, ARTN e2024JA033642, doi:10.1029/2024JA033642, 2025.

Smith, R. W. and Hernandez, G.: Vertical Winds in the Thermosphere within the Polar-Cap, *J Atmos Terr Phys*, 57, 611-620, doi:10.1016/0021-9169(94)00101-S, 1995.

745 Sobolev, V. V., Irvine, W. M., Gendel, M., and Lane, A. P.: Light scattering in planetary atmospheres, 1st . International series
 746 of monographs in natural philosophy vol.76, Pergamon Press, Oxford, New York, doi:10.1016/C2013-0-05661-7, 1975.

747 Themens, D. R., Elvidge, S., McCaffrey, A., Jayachandran, P. T., Coster, A., Varney, R. H., Galkin, I., Goodwin, L. V., Watson,
 748 C., Maguire, S., Kavanagh, A. J., Zhang, S. R., Goncharenko, L., Bhatt, A., Dorrian, G., Groves, K., Wood, A. G., and
 749 Reid, B.: The High Latitude Ionospheric Response to the Major May 2024 Geomagnetic Storm: A Synoptic View,
 750 Geophysical Research Letters, 51, e2024GL111677, doi:10.1029/2024GL111677, 2024.

751 Wang, C., Xu, J., Chen, Z., Li, H., Feng, X., Huang, Z., and Wang, J.: China's Ground-Based Space Environment Monitoring
 752 Network—Chinese Meridian Project (CMP), Space Weather, 22, doi:10.1029/2024sw003972, 2024.

753 Wang, X., Aa, E., Chen, Y., Zhang, J., Zhu, Y., Cai, L., Lu, X., Luo, B., Liu, S., Li, M., Shen, H., and Yuan, T.: Midlatitude
 754 Neutral Wind Response During the Mother's Day Super-Intense Geomagnetic Storm in 2024 Using Observations From
 755 the Chinese Meridian Project, Journal of Geophysical Research: Space Physics, 130, e2024JA033574,
 756 doi:10.1029/2024JA033574, 2025.

757 Wei, D., Zhu, Y., Liu, J., Gong, Q., Kaufmann, M., Olschewski, F., Knieling, P., Xu, J., Koppmann, R., and Riese, M.:
 758 Thermally stable monolithic Doppler asymmetric spatial heterodyne interferometer: optical design and laboratory
 759 performance, Optics Express, 28, doi:10.1364/oe.394101, 2020.

760 Wu, Q., Yuan, W., Xu, J. Y., Huang, C., Zhang, X. X., Wang, J. S., and Li, T.: First U.S.-China joint ground-based Fabry-
 761 Perot interferometer observations of longitudinal variations in the thermospheric winds, J Geophys Res-Space, 119, 5755-
 762 5763, doi:10.1002/2014ja020089, 2014.

763 Xu, H., Shiokawa, K., Oyama, S., and Otsuka, Y.: Thermospheric wind variations observed by a Fabry-Perot interferometer
 764 at Tromsø, Norway, at substorm onsets, Earth Planets Space, 71, ARTN 93, doi:10.1186/s40623-019-1072-0, 2019.

765 Yang, C., Zhao, B., Jin, Y., Huang, C., Yao, X., and Wan, W.: Climatology of Nighttime Upper Thermospheric Winds From
 766 Fabry-Perot Interferometer 2011–2019 Measurements Over Kelan (38.7°N, 111.6°E), China: Local Time, Seasonal, Solar
 767 Cycle, and Geomagnetic Activity Dependence, Journal of Geophysical Research: Space Physics, 125, e2020JA027892,
 768 doi:10.1029/2020JA027892, 2020.

769 Yu, T., Huang, C., Zhao, G. X., Mao, T., Wang, Y. G., Zeng, Z. C., Wang, J. S., and Xia, C. L.: A preliminary study of
 770 thermosphere and mesosphere wind observed by Fabry-Perot over Kelan, China, J Geophys Res-Space, 119, 4981-4997,
 771 doi:10.1002/2013ja019492, 2014.

772 Yuan, W., Xu, J. Y., Ma, R. P., Wu, Q. A., Jiang, G. Y., Gao, H., Liu, X. A., and Chen, S. Z.: First observation of mesospheric
 773 and thermospheric winds by a Fabry-Perot interferometer in China, Chinese Sci Bull, 55, 4046-4051, doi:10.1007/s11434-
 774 010-4192-2, 2010.

775 Zhang, K. D., Wang, H., Liu, J., Song, H. M., and Xia, H.: The Quasi-Periodic Nighttime Traveling Ionospheric Disturbances
 776 on 13 May 2024 During the Recovery Phase of a SuperStorm, J Geophys Res-Space, 130, ARTN e2024JA033257,
 777 doi:10.1029/2024JA033257, 2025.

778 Zhu, G., Zhu, Y., Kaufmann, M., Wang, T., Liu, W., and Xu, J.: An Efficient Calibration System of Optical Interferometer for
 779 Measuring Middle and Upper Atmospheric Wind, Remote Sensing, 15, doi:10.3390/rs15071898, 2023.