

1 Impact of lower atmospheric scattering on ground-based optical 2 thermospheric wind observations with spatially uneven airglow

3 Xiaolong Wei^{1,2,4}, Guoying Jiang^{1,3,4,*}, Yajun Zhu^{1,3,4,*}, Jiyao Xu^{1,4}, Weijun Liu^{1,4}, Tiancai Wang^{1,2,4},
4 Guangyi Zhu^{1,4}, Wei Yuan^{1,4}

5 ¹State Key Laboratory of Solar Activity and Space Weather, National Space Science Center, Chinese Academy of Sciences,
6 Beijing, 100190, China

7 ²College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China.

8 ³School of Astronomy and Space Science, University of Chinese Academy of Sciences, Beijing, 100049, China.

9 ⁴Hainan National Field Science Observation and Research Observatory for Space Weather, Hainan, 571734, China

10 *Correspondence to: Guoying Jiang (gyjiang@spaceweather.ac.cn), Yajun Zhu (y.zhu@swl.ac.cn)

11 **Abstract.** Scattered airglow emissions in the lower atmosphere can bias ground-based interferometer observations of
12 thermospheric winds, particularly when airglow brightness becomes spatially uneven due to auroras. During two geomagnetic
13 storms with visible auroras on May 10th and Oct. 10th, 2024, the Doppler Asymmetric Spatial Heterodyne (DASH) and Fabry-
14 Perot (FP) interferometers concurrently detected atypical winds at Siziwang (SIZW, 41.83° N, 111.93° E), suspected to be
15 caused by scattering. These atypical winds, characterized by horizontal differences exceeding 400 m·s⁻¹ between opposite
16 cardinal directions (N-S or E-W) and downwelling exceeding 100 m·s⁻¹, showed a strong temporal association with airglow
17 brightness. By modelling the transmission of scattered airglow emissions, we calculate post-scattering wind speeds as the
18 initial wind speeds weighted by both scattered and direct intensities. With fixed initial winds (100 m·s⁻¹ westward, 400 m·s⁻¹
19 southward, zero vertical wind), the simulation reproduces horizontal differences of approximately 400 m·s⁻¹ on May 10th and
20 100 m·s⁻¹ on Oct. 10th, both capturing the temporal characteristics of the atypical winds. The simulation shows that scattering-
21 induced biases on line-of-sight speed take their sign from the brighter region, while their magnitude varies directionally with
22 the angle to that region: at 45° elevation, biases 135–180° azimuth away exceed those in the brighter region by more than 10
23 times. Limited by uncertainties in airglow images and optical depth of model inputs, the simulation incurs numerical errors of
24 roughly 75 % during some periods. Effective correction of the scattering impact will require improved accuracy of model
25 inputs in the future. ~~By modelling the transmission of scattered airglow emissions, we calculated post-scattering wind speeds~~
26 ~~as the initial wind speeds weighted by both scattered and direct intensities. With fixed initial speeds, the simulation reproduced~~
27 ~~the temporal characteristics of the atypical winds, demonstrating that scattering may contribute to these intense horizontal~~
28 ~~differences and downwelling. The simulation also shows that the scattering-induced biases have directional inhomogeneity~~

29 with characteristics linked to the location and background line-of-sight speed of the brighter airglow region. The accuracy of
30 the simulation is limited by the accuracy of airglow observations and atmospheric optical depth.

31 1 Introduction

32 Optical interferometers are widely utilized to observe thermospheric neutral wind (Burnside et al., 1981; Burnside and Tepley,
33 1989; Killeen et al., 1995; Emmert et al., 2001; Fejer et al., 2002; Emmert et al., 2006; Wu et al., 2014). Thermospheric wind
34 can be derived from measuring the Doppler shift of OI red-line airglow emission at 630.0 nm. This emission, primarily from
35 the collisional deactivation of $O(^1D)$ generated by O_2^+ dissociative recombination, peaks near 250 km altitude. The height-
36 integrated thermospheric wind around the peak altitude can be obtained (Biondi and Feibelman, 1968; Hernandez and Roble,
37 1976; Burnside et al., 1981; Biondi et al., 1995; Nakajima et al., 1995). For scanning interferometers, three-dimensional wind
38 vectors can be derived by observing the zenith and four cardinal directions at a specific elevation angle. The scanning range
39 covers a circular area about 500 km in diameter at airglow altitude. Given thermospheric wind uniformity at this scale,
40 horizontal winds observed in two opposite cardinal directions (N-S or E-W) are typically similar. Averaging opposite cardinal
41 directions improves accuracy, mitigates cloud effects, and is typically used to represent local meridional or zonal winds even
42 during geomagnetic storms (Friedman and Herrero, 1982; Fejer et al., 2002; Sakanoi et al., 2002; Dhady et al., 2017; Huang
43 et al., 2018; Xu et al., 2019; Li et al., 2023; Wang et al., 2025).

44 However, horizontal winds in opposite cardinal directions occasionally show significant separation exceeding $100\text{ m}\cdot\text{s}^{-1}$ and
45 strong vertical winds, deviating from typical thermospheric wind uniformity. These observations often occur near auroras,
46 unaffected by clouds or moonlight, and have acceptable standard errors. They mainly occur in polar regions (Crickmore et al.,
47 1991; Price et al., 1995; Smith and Hernandez, 1995; Innis et al., 1999; Ishii et al., 2001; Guo and McEwen, 2003; Anderson
48 et al., 2012), but have also been seen at mid-latitudes during major geomagnetic storms (Hernandez and Roble, 1976; Makela
49 et al., 2014).

50 Atmospheric scattering of airglow emissions is thought to be one of the factors that biases ground-based wind observations,
51 potentially accounting for the atypical wind. Initially, it was thought to impact airglow peak height measurements by
52 photometers (Ashburn, 1954). Subsequent studies by Abreu et al. (1983) explored its impact on thermospheric wind speed
53 measurements using a Fabry-Perot interferometer. Harding et al. (2017a; 2017b) later systematically modelled and estimated
54 these effects, revealing that scattering was responsible for the anomalous vertical winds observed at mid-latitudes during
55 geomagnetic storms by Makela et al. (2014). Light from brighter airglow regions scatters omnidirectionally in the lower
56 atmosphere, primarily the troposphere and stratosphere, and is detectable outside its original direction. The additional Doppler
57 shift of this scattered light can bias the retrieval of line-of-sight (LOS) speeds as well as the converted horizontal and vertical
58 winds. Harding et al. (2017b) also investigated the impact of atmospheric scattering on interferometer wind and temperature
59 measurements during quiet periods.

60 Scattering-induced biases are more pronounced during spatially uneven airglow brightness, such as during auroras (Harding
61 et al., 2017a). Uneven airglow brightness refers specifically to inhomogeneous red-line emissions. At mid-latitudes, marked
62 uneven red-line airglow usually comes from red aurora. Despite their distinct origins, the spectral and altitudinal overlap of
63 airglow and aurora will let ground-based optical instruments conflate the two. For red-line observations, the aurora itself may
64 also bias the derived winds. Aurora could elevate the red-line emission profile (Kataoka et al., 2024b), so the interferometer
65 samples winds that are both higher and farther away. This makes the northward view sense winds deviate from the expected
66 thermospheric wind at 250 km altitude when looking toward the aurora. Additionally, spectral contamination from
67 precipitating energetic ions could also bias interferometers (Makela et al., 2014). They suggested that the enhanced
68 downwelling at mid-latitudes during storms might result from the contamination of the spectral profile by fast O atoms
69 associated with the influx of low-energy O⁺ ions.

70 From a dynamical perspective, wind differences in opposite cardinal directions are considered horizontal divergence, which
71 are often associated with changes in vertical winds. Near the aurora arc, these atypical winds are mainly caused by ion drag,
72 Joule heating, and energy particle precipitation (Hays et al., 1984; Rees et al., 1984; Conde and Smith, 1995; Conde et al.,
73 2001; Anderson et al., 2012). Generally, excessive horizontal divergence and vertical wind appear alongside rapidly changing
74 auroras and exhibit a matching spatial relationship that upward (downward) winds accompanied by divergences (convergences)
75 are often detected when aurora exists equatorward (poleward) of the observatory (Ishii et al., 2001; Guo and McEwen, 2003).
76 The combination of vertical wind and horizontal divergence is related to gravity waves excited by the above processes in polar
77 regions, presenting a wave-like structure and phase delay between vertical and horizontal wind components. (Price et al., 1995;
78 Smith and Hernandez, 1995; Ishii et al., 1999; Ishii et al., 2001; Shinagawa and Oyama, 2006). At mid-latitudes, which are
79 not primary regions for magnetospheric energy injection, atypical winds are instead related to the propagation of gravity waves
80 from polar regions. (Hernandez and Roble, 1976).

81 During two geomagnetic storms on May 10th and October 10th, 2024, with visible auroras, we observed similar atypical winds
82 in ground-based interferometers at Siziwang (SIZW, 41.83° N, 111.93° E), China. These winds showed intense differences
83 over 400 m·s⁻¹ in two opposite cardinal directions for both meridional and zonal components, along with downward wind
84 exceeding 100 m·s⁻¹. The observations were unaffected by moonlight or clouds, and the interferometer retrieval errors were
85 acceptable (see Section 3.1). These atypical winds at SIZW only occurred with auroras statistically and significantly deviated
86 from the regional climatological norms over the China region (Jiang et al., 2018; Yang et al., 2020). This raises the question
87 of whether the atypical winds arise from dynamical processes, are influenced by red aurora, or stem from scattering-induced
88 biases and other measurement-related factors. Unfortunately, most of these mechanisms could amplify the wind-speed contrast
89 between opposite cardinal directions, rendering them difficult to disentangle (Harding et al., 2017a). Given the scarcity of
90 additional thermospheric-wind or auroral instruments, we remain unable to quantify every potential mechanism. Motivated by
91 the observed phenomena, this study attempts to estimate how scattering modulates the atypical winds in these storms. While

prior studies focus on vertical wind biases of Fabry-Perot interferometers under auroral conditions (Harding et al., 2017a; Harding et al., 2017b), we will analyze the formation and patterns of horizontal differences caused by scattering. We will also incorporate Doppler Asymmetric Spatial Heterodyne (DASH) interferometer data to compare scattering impact across different interferometer types. As red auroras now regularly appear at the low magnetic latitudes of Japan and China during elevated solar activity (Kataoka et al., 2024a; Kataoka et al., 2024b; Ma et al., 2024), a deeper understanding of scattering-induced biases is essential for the proper use of interferometer data collected in these regions. In the following text, a scattering radiative transfer model is used to simulate interferometer observations in two cases with visible aurora. The presence and patterns of scattering-induced biases are analyzed by comparing simulations with observations.

2 Instruments and model

This study was conducted at the Siziwang station (SIZW; 41.83° N, 111.93° E, and 37.7° N MLat) of the Chinese Meridian Project Phase II (Wang et al., 2024), utilizing a Dual-Channel All-sky Airglow Imager (DCAI), a Dual-Channel Optical Interferometer (DCOI), and a Fabry-Perot Interferometer (FPI). DCOI derives neutral winds by observing atomic oxygen green-line (557.7 nm, around 96 km) and red-line (630.0 nm, around 250 km). DCAI observes hydroxyl (around 87 km) and atomic oxygen red-line nightglow, respectively. FPI only works at the red-line. Our focus is on the red-line channel. Using DCAI images as one of the inputs, wind biases from optical interferometers can be simulated by a scattering radiative transfer model (scattering model for short). Instruments and the model are described in the following subsections.

2.1 Dual-Channel All-Sky Airglow Imager

Dual-Channel All-Sky Airglow Imager (DCAI) comprises a fisheye lens with an approximate 170 degree field of view, a 2 nm narrow-band filter, and a 1024×1024 pixel, 16 bit cooled CCD. DCAI exposure time of the red-line is 2 minutes. The obtained airglow images are first calibrated to the local spherical coordinate system, then sequentially corrected for stray light, Van Rhijn effect, and atmospheric extinction, and finally projected onto the 250 km airglow plane. Due to DCAI not calibrating the Rayleigh unit (Shiokawa et al., 2000), observed brightness is only normalized to the full-well value. And because of fish-eye lens distortion and the lack of Rayleigh unit calibration, the edge brightness of the view is inaccurate. Thus, observations are restricted within a 70° zenith angle. For larger zenith angles, the brightness is obtained by radial zero-order extrapolation in airglow projection. Detailed image processing procedures are in Appendix B.

2.2 Dual-channel optical interferometer

Dual-channel optical interferometer (DCOI) is a scanning interferometer using Doppler Asymmetric Spatial Heterodyne (DASH) technology. DASH exhibits a wider field of view, better thermal stability, simplified mechanisms, and lower tolerance requirements than other interferometer structures (Englert et al., 2007; Englert et al., 2010; Harlander et al., 2017; Wei et al., 2020). DCOI consists of a 630 nm narrow-band filter (2 nm bandwidth), a 9 degree field-of-view lens (f/6), a DASH

interferometer with a 25 mm aperture, a Neon lamp for calibration, and a 2048×2048 pixel CCD (13.5 μm per pixel) (Wei et al., 2020; Zhu et al., 2023; Liu et al., 2025). Its thermal stability is maintained within 0.1 K. DCOI measures three-dimensional wind speeds by scanning five directions (zenith and four cardinal directions at 45° zenith angle). Each direction is exposed for 5 minutes, completing a cycle roughly every 25 minutes. DCOI adopts an observation with the smallest error after evening as the reference zero wind speed. The slant LOS speeds are subtracted by the time-regressed projection of vertical speed and then converted to horizontal using the sine of zenith angles. It is worth noting that during auroral events, vertical winds with absolute values exceeding 50 m·s⁻¹ are excluded from the regression, as they contain scattering effects that could introduce additional biases to other directions. DCOI provides two series of meridional wind, two series of zonal wind, and one series of vertical wind.

2.3 Fabry-Perot Interferometer

Fabry-Perot Interferometer (FPI), as a mature solution, conducts comparative observations with DCOI. It features a 630 nm narrow-band filter (2 nm bandwidth), a 2.54 degree field-of-view lens (f/6), a 50 mm aperture etalon with a 7 mm gap, a frequency-stabilized laser for calibration, and a 1024×1024 pixel CCD (13 μm per pixel). FPI uses the same integration time and scanning method as DCOI to obtain horizontal and vertical winds for each cardinal direction and zenith. Details and historical results of FPI are in these references (Yuan et al., 2010; Wu et al., 2014; Yu et al., 2014; Huang et al., 2018; Jiang et al., 2018).

2.4 Scattering radiative transfer model

The model for estimating scattering impact is based on the scattering radiative transfer model and numerical solution by Harding et al. (2017a). It assumes airglow emission undergoes elastic scattering, preserving its wavelength and initial Doppler shift. By specifying airglow brightness distribution, original Doppler shift, lower atmosphere scattering characteristics, and a simplified geometric relationship, the radiation transfer equations (see Appendix A) can be solved to compute the distribution of multiple scattered light and its associated Doppler shift. This enables the wind simulation with atmospheric scattering. A schematic diagram (Fig. 1) illustrates the basic mechanism. To enhance applicability, we have refined several aspects: (1) The upper boundary of the lower atmosphere is set at 40 km to improve the accuracy of the effective extinction path in the initial source function. (2) The Doppler shift is replaced by LOS speed, with every incident ray from the airglow layer mapped directly to its corresponding LOS speed. (3) After slicing the airglow layer into several bins by LOS-speed, the model illuminates one bin per run, records its scattered intensity, then merges all bins with an intensity-weighted average to yield the post-scattered LOS speed. The detailed model description is provided in Appendix A.

Additionally, the scattering characteristics of the lower atmosphere in our model, including the scattering phase function and optical depth, were derived from Aerosol Robotic Network (AERONET) observations (Holben et al., 2001). We utilized data from the Baotou site (40.9° N, 109.6° E), which is the nearest available site to SIZW, located approximately 180 km away.

153 The total optical depth, accounting for both aerosol and molecular scattering, was calculated using monthly averages and was
 154 found to be 0.43 in May and 0.2 in October. The scattering phase function was determined based on AERONET data following
 155 the previous method (Harding et al., 2017a). Further details regarding the scattering characteristics are described in Appendix
 156 B.




Figure 1: The schematic diagram of the scattering radiative transfer model

159 The grey shading represents the lower atmospheric layer, with darker hues indicating greater optical depth. The yellow-green fillers
 160 represent the relative brightness from the red-line airglow layer. Yellow indicates higher light intensity. The blue-red fillers, which
 161 correspond to the relative brightness, represent the Doppler shift type (blue-shift or red-shift) of LOS wind speeds. (a) to (e) represent
 162 airglow emissions travelling along different paths, carrying Doppler shifts from outside the line of sight into the interferometer, thereby
 163 causing biases in the observations. The model estimates the biases by simulating the distribution of airglow emissions after scattering.

164 3 Results

165 Two storms with visible auroras on May 10th and Oct. 10th, 2024, respectively, are used to study the scattering impact. The
 166 storm from May 10th to 11th is characterized by its significant magnitude and prolonged duration. Multiple works report this
 167 event (Guo et al., 2024; Hajra et al., 2024; Themens et al., 2024), with particular focus on the variations of thermospheric
 168 winds (Wang et al., 2025; Zhang et al., 2025) and auroras (Gonzalez-Esparza et al., 2024; Kataoka et al., 2024b; Mikhalev,
 169 2024; Nanjo and Shiokawa, 2024) at mid-latitudes. The storm commenced around 17:00 UT on May 10th and the main phase
 170 persisted until 02:00 UT on May 11th. After that, the local night of May 11th in the China region sank into a continuous recovery
 171 phase. Another storm from Oct. 10th to 11th is weaker than May's (Ranjan and Pallamraju, 2025; Singh et al., 2025), with the

main phase from 18:00 UT on Oct. 10th to 02:00 UT on Oct. 11th. During the two geomagnetic storms with visible auroras, both the DCOI and FPI at SIZW observed atypical winds, characterized by intense horizontal wind differences and downward vertical winds.

3.1 Storm-time wind speed statistics

It is necessary to ascertain whether atypical winds originate from atmospheric scattering with spatially uneven airglow brightness or dynamic processes during storms. To investigate the impact of visible auroras on atypical winds during storms, we made the most of the available observations, tracking DCOI's storm-time observations for nearly a year and FPI's for almost five months. We employed the planetary magnetic index Kp exceeding 3 to identify geomagnetic storms (Yang et al., 2020). Besides, to rule out moonlight and cloud effects, we only used clear sky conditions, which means: (1) excluding cases where the angle between the moon and the line of sight is less than 30 degrees, and (2) excluding cases where large-area thick cloud coverage is visible in DCAI. Additionally, data with standard errors greater than 50 m·s⁻¹ were also excluded. A few aurora events, including Nov. 5th, Dec. 1st, 2023, and Aug. 12th, 2024, that did not meet this criterion were excluded.




Figure 2: Storm-time (Kp>3) thermospheric wind speed statistics at SIZW

Figure 2a shows meridional winds observed along two opposite directions (N-S) by FPI, with north-looking in red and south-looking in blue. Figure 2b shows zonal and vertical wind, with east-looking in yellow, west-looking in green, and zenith-ward in black. The northward, eastward, and upward speeds are positive in coordinates. Observations without aurora are shown as points, while those with visible auroras are shown as lines. Figures 2c and 2d are similar but show DCOI data.

Figure 2 shows thermospheric wind statistics during geomagnetic storms ($K_p > 3$) at SIZW. The first two panels display FPI data from May to Oct. 2024, while the rest display DCOI data from Nov. 2023 to Oct. 2024. FPI began operation on May 8th, 2024, with about half a year less data than DCOI. Observations with no aurora in the field of view are marked as points, while the two cases with visible auroras are shown as lines. The five observation directions of the interferometer are marked by different colors. During typical storms, horizontal winds consistently increase to around $150 \text{ m}\cdot\text{s}^{-1}$ both equatorward and westward with no significant downward wind. However, under visible auroras, both DCOI and FPI have detected large wind speeds, such as a southward wind of about $600 \text{ m}\cdot\text{s}^{-1}$ and downwelling exceeding $100 \text{ m}\cdot\text{s}^{-1}$. The two series of winds observed along opposite cardinal directions (N-S or E-W) exhibit overt differences, with values exceeding $400 \text{ m}\cdot\text{s}^{-1}$ and contrary directions. This is markedly different from the wind patterns observed during non-aurora storms, where opposite-direction winds do not show significant divergence. Comparing the results of DCOI and FPI, the observations are largely consistent both with and without auroras. The atypical winds observed simultaneously by two interferometers with different principles suggest a systematic error from outside the instruments. Besides, these simultaneous changes appear in five observation directions, all characterized by enhanced negative LOS speeds, a signature consistent with scattering-induced contamination, indicating likely LOS speed contamination. These factors point more towards scattering impact rather than dynamical processes as the cause. Next, the relationship between scattered light and atypical winds will be investigated through simulation.

3.2 Comparison of observations and simulations

Figure 3 shows the red-line airglow brightness from DCAI (Fig. 3a, 3e), the observed winds from DCOI and FPI (solid lines with different markers in Fig. 3b-3d, 3f-3h) and the simulated winds from the scattering model (dotted lines in Fig. 3b-3d, 3f-3h) during the two nights of May 10th and 11th, 2024, at SIZW, in which the different colors denote distinct directions. The grey lines in the horizontal wind plots represent the average values between opposite cardinal directions. The multi-directional brightness series from DCAI are extracted at 45° zenith angle, consistent with the scanning zenith angle of interferometers. Time intervals with visible auroras are highlighted in red, showing much higher brightness in northward directions than others. Figure 4 supplements the auroral distribution compared to Fig. 3a and 3e. Images from DCAI are projected onto the airglow layer at 250 km. The red circle encloses the actual observations with zenith angles less than 70° , while the values outside are extrapolated. The red dots represent the interferometer's pierce points on the airglow layer at 45° zenith angle.




Figure 3: Observations of aurora and wind speeds, and the scattering model simulation on the nights of May 10th and 11th, 2024, at SIZW

Figure 3a shows the brightness of 8 cardinal directions, all at 45° zenith angle, along with the zenith-ward, extracted from DCAI. The color coding is as follows: red for northern directions, green for east and west, blue for southern directions, black for the zenith, and yellow for the average brightness excluding the three northern directions. Figure 3b shows the meridional wind, with north-looking in red and south-looking in blue, and the average of the two directions in grey. DCOI observations are shown as solid lines with circular dots, FPI as solid lines with rhombus dots, and simulations as dotted lines. Figures 3c, 3d are similar to Fig. 3b, but for zonal and vertical wind, with east-looking in yellow, west-looking in green, and zenith-ward in black. For a more concise figure, if the standard error exceeds 100 m s⁻¹, the point will be filled with black instead of error bar. Figures 3a-3d show data from May 10th, and Fig. 3e-3h from May 11th. The time intervals with visible auroras are marked in red.




Figure 4: Auroral distribution observed by DCAI on the nights of May 10th and 11th, 2024, at SIZW

Images from DCAI (Fig. 4a-4d, 4f-4i) have been projected onto the airglow layer at 250 km. The red circle encloses actual observations with zenith angles $< 70^\circ$, while values outside are extrapolated using zero-order extrapolation. The red dots represent the interferometer's pierce points on the airglow layer at 45° zenith angle. Figures 4e, 4j are similar to Fig. 3a, 3e, with the corresponding images time labelled.

The coastlines in projected DCAI images are made with Natural Earth.


After 18:30 UT on May 10th, as the aurora intensified, both DCOI and FPI detected simultaneous changes in meridional, zonal, and vertical winds. The north-looking red-line brightness at 45° zenith angle exceeded three times that of other directions. The meridional and zonal wind differences between opposite cardinal directions (N-S or E-W) increased. And the winds detected in opposite directions reversed. The maximum meridional difference was close to $800 \text{ m}\cdot\text{s}^{-1}$, while that in zonal exceeded $500 \text{ m}\cdot\text{s}^{-1}$. The downward wind was enhanced by over $100 \text{ m}\cdot\text{s}^{-1}$. These four aforementioned variables that red-line brightness, the meridional and zonal differences, and downwelling, increased almost simultaneously, peaked at 19:05 UT, and then decayed.

238 Moreover, the average meridional wind, derived from averaging opposite cardinal directions, continuously enhanced
 239 equatorward to around 400 $\text{m}\cdot\text{s}^{-1}$, while the average zonal wind enhanced westward to around 100 $\text{m}\cdot\text{s}^{-1}$. Unlike the
 240 single-direction results that peaked at 19:05 UT, the average wind varied steadily, consistent with storm-time circulation.
 241 Compared to the average wind, the separated horizontal and enhanced vertical winds are atypical. Even with travelling
 242 atmospheric disturbances (TADs) superimposed on storm-time circulation, phase lags between horizontal and vertical
 243 components would be expected (Hernandez and Roble, 1976; Ishii et al., 1999), but none were observed. Thus, the atypical
 244 winds do not appear to be the result of a dynamical process. During the recovery phase on May 11th, the aurora was present
 245 throughout the night but much weaker than on May 10th, as seen in Fig. 4. Both DCOI and FPI showed westward and
 246 equatorward horizontal winds with no significant downward wind. There was a meridional difference of about 100 to 300
 247 $\text{m}\cdot\text{s}^{-1}$ persisted throughout the night, with no zonal difference.


248 Subsequently, we used the scattering model to quantitatively examine ~~explore~~ the relationship between red-line brightness
 249 variations and atypical winds through atmospheric scattering. On May 10th, a fixed wind vector of 100 $\text{m}\cdot\text{s}^{-1}$ westward and
 250 400 $\text{m}\cdot\text{s}^{-1}$ southward was set as the input. This assumed wind was kept constant over time and spatially uniform, with no
 251 vertical components. On May 11th, a fixed wind vector of 200 $\text{m}\cdot\text{s}^{-1}$ westward and 100 $\text{m}\cdot\text{s}^{-1}$ southward was used, again with
 252 no vertical component. These values are chosen based on average observed wind speeds to approximate storm-time circulation.
 253 Although the specific values may deviate, the main wind directions remain consistent. The storm-time enhancement of vertical
 254 winds may be caused by scattering rather than representing real winds, so we set it to zero. It is worth noting that we neglect
 255 the variation of background wind in the model inputs, due to uncertainty regarding whether the observed wind variations are
 256 biased. Moreover, using fixed wind speeds allows us to highlight the impact of red-line brightness variations and determine
 257 the presence of scattering effects.

258 As shown by the dotted lines in Fig. 3b-3e, the simulations with scattering impact generally match the observations on May
 259 10th. ~~Simulated horizontal wind differences and downwelling increase initially with the aurora, peak around 19:15 UT, and~~
 260 ~~then decay.~~ Assuming a uniform, constant wind field, the scattering model yields horizontal differences (about 400 $\text{m}\cdot\text{s}^{-1}$) and
 261 downwelling (about 100 $\text{m}\cdot\text{s}^{-1}$) that track the red-line auroral emission, peaking near 19:15 UT before fading. The simulated
 262 wind speed variations lag the observations by about 10 minutes. The lag may be due to the relatively rough 25 minute scanning
 263 cycle of interferometers or DCAI underestimating airglow brightness at the field of view's edge, leading to inaccurate capture
 264 of scattering enhancement start time. Numerically, the simulated zonal and vertical winds match observations more closely
 265 than the meridional wind. ~~The simulated meridional difference is smaller than the observed difference, and the north-looking~~
 266 ~~simulation remains closer to the default value, unlike the equatorward biased observation.~~ The simulated meridional difference
 267 falls short of the observed value, with the north-looking simulation remaining near the default 400 $\text{m}\cdot\text{s}^{-1}$ southward while the
 268 observation is equatorward-biased at approximately 600 $\text{m}\cdot\text{s}^{-1}$ southward. The preset fixed wind may influence the meridional
 269 simulation, as it does not follow the equatorward enhancement of the meridional wind. Other factors beyond scattering impact

270 might also have an impact, such as the lift of red-line emission profile and the spectral pollution caused by auroras (Makela et
 271 al., 2014), to be discussed later. Additionally, the model simulates a similar intense downward wind as observed under the
 272 preset zero vertical wind. This indicates that the vertical wind is significantly affected by scattering. This is why the intense
 273 vertical wind is not subtracted when converting interferometer LOS speed to horizontal wind, to prevent error propagation.
 274 For May 11th, due to the weak but continuous aurora, the simulation shows weak horizontal differences and slight downward
 275 winds throughout the night. Compared with the observations, the simulation underestimates the meridional differences by
 276 more than 50 %, and it produces zonal differences and downward winds that are not evident in the data. ~~the simulation shows~~
 277 ~~smaller meridional differences. It also indicates zonal differences and downward winds, which are not evident in the~~
 278 ~~observations.~~ The poorer simulation on May 11th may be due to misalignment between dominant horizontal winds and airglow
 279 brightness gradients, which will be discussed later. Additionally, there may be issues with the zero wind calibration. When
 280 auroras are present throughout the night, the vertical wind, which includes scattering biases, may have been used to calibrate
 281 zero wind speed. It likely masks the scattering impact in the observations and explains the discrepancies in the simulation.



283 **Figure 5: Observations of aurora and wind speeds, and the scattering model simulation on the nights of Oct. 10th and 11th, 2024, at**
 284 **SIZW**
 285 Figure 5 is analogous to Fig. 3, but for Oct. 10th and 11th.



286 **Figure 6: Auroral distribution observed by DCAI on the nights of Oct. 10th and 11th, 2024, at SIZW**
 287 **Figure 6 is analogous to Fig. 4, but for Oct. 10th and 11th.**
 288

289 Figures 5 and 6 show another case from Oct. 10th to 11th, analogous to Figures 3 and 4. On Oct. 10th, the aurora appeared at
 290 17:15 UT, expanded southward and increased in brightness, peaking first at 18:30 UT before decaying and then increasing
 291 again from 20:30 UT until sunrise. The second peak was brighter than the first (Fig. 6). Similar to the storm in May, once
 292 aurora appeared, both DCOI and FPI observed atypical winds, with synchronous meridional and zonal differences and
 293 downward enhancements in vertical wind. These atypical winds also exhibited two peaks, around 18:30 UT and 21:00 UT.
 294 The horizontal winds observed in opposite cardinal directions were basically in opposition. During visible aurora periods,
 295 DCOI and FPI showed a 50 to 100 m s⁻¹ difference in vertical wind but consistent variation trends. Moreover, the average

296 horizontal wind between opposite cardinal directions was dominantly equatorward and westward, which also had two peaks.
297 On Oct. 11th, the storm had passed, and no visible aurora was present. The increase in brightness around 13:00 UT was due to
298 moonset in the southwest. There were no significant horizontal differences or downward winds, consistent with geomagnetic
299 quiet conditions.


300 We set a fixed $100 \text{ m}\cdot\text{s}^{-1}$ westward with $400 \text{ m}\cdot\text{s}^{-1}$ southward wind vector for Oct. 10th, and $100 \text{ m}\cdot\text{s}^{-1}$ westward with $200 \text{ m}\cdot\text{s}^{-1}$
301 southward wind vector for Oct. 11th in the model, respectively, with no vertical component. The ~~simulation~~ simulated
302 horizontal wind differences on Oct. 10th ~~exhibits also exhibit~~ two peaks. The second simulated peak matches well with the
303 observations, whereas the first peak, though capturing the trend, is underestimated by approximately 75 % of its magnitude.
304 ~~The second peak in the simulation is consistent with the observations better, while the first peak, although capturing the trend,~~
305 ~~is significantly underestimated in magnitude.~~ This discrepancy in the simulation may relate to optical depth, aurora brightness,
306 and background wind changes. The optical depth in Oct. is about half that of May, and simulations underestimate observed
307 values. As in previous studies (Harding et al., 2017b), optical depth can affect the scattering model response. On Oct. 10th, the
308 first aurora brightening is weaker than the second. When red-line brightness spatial differences are small, the model response
309 tends to be lower. The impact of optical depth and red-line brightness on the model will be discussed later. Additionally,
310 noticeable fluctuations in the average meridional wind on this day may also contribute to the deviation in the model with fixed
311 initial wind. The north-looking wind speed varied dramatically between 19:00 UT and 21:00 UT along with the aurora, which
312 may also be related to spectral contamination beyond scattering impact. This spectral contamination arises from fast O atoms
313 generated by low-energy O^+ ion precipitation in the auroral region, which occurs at higher altitudes. This issue introduces an
314 additional spectral shift that compromises wind retrieval (Makela et al., 2014). This exceeds the simulation range of the model,
315 thereby causing the discrepancy.

316 4 Discussions

317 In this study, we modelled the transmission of scattered airglow emission in the lower atmosphere. Post-scattering wind speeds
318 were calculated based on assumed initial wind speeds weighted by both scattered and direct intensities. The model basically
319 captured the temporal variations of horizontal wind differences and downward enhancements associated with varying auroral
320 brightness, suggesting the contribution of scattering mechanisms to atypical winds. However, the simulation of scattering has
321 certain limitations and characteristics: (1) The differences between simulation and observation vary across different directions.
322 (2) The simulated values sometimes exhibit significant numerical deviations from observations. Could this be related to model
323 errors? Next, we will discuss the working principle of the scattering model and the errors involved in the simulation.

324 **4.1 Core working principle of the scattering model**

325 As shown in Fig. 3 and Fig. 5, observed winds respond differently to scattering across directions, especially on May 10th and
326 Oct. 10th with stronger auroras. Although the scattering model has numerical errors, the simulations also show directional
327 differences in scattering-induced biases. Both observations and simulations indicate that the meridional wind responds the
328 most, followed by the zonal wind, while the vertical wind responds the least. Since vertical and horizontal wind speeds are
329 derived from the projection of LOS wind speeds, this essentially reflects the non-uniform response of LOS speeds to scattering.
330 This directional inhomogeneity of scattering impact aligns with previous studies. Harding et al. (2017a) simulated scattering
331 effects under auroral conditions, using northward observations as the initial winds. They noted that this direction experiences
332 minimal scattering contamination due to facing the brighter region. Abreu et al. (1983) used a meridional one-dimensional
333 model, finding that LOS wind speeds near intense airglow brightness gradients and with weaker airglow intensity are more
334 susceptible to contamination. They also showed that scattering-induced biases are minimal in the vertical direction, as the
335 shorter atmospheric path length limits the opportunity for scattering. In this study, we further explore the scattering impact as
336 a function of azimuth angle, revealing the formation of horizontal differences.



337
338 **Figure 7: Post-scattering LOS speeds at 45° zenith angle**

339 The figure shows the post-scattering line-of-sight speeds at 45° zenith angle for various directions over time, with panels for May 10th,
340 May 11th, and October 10th, respectively.

341 Figure 7 shows post-scattering LOS speeds at 45° zenith angle in the simulations on three aurora nights, with the vertical axis
342 indicating cardinal directions derived from azimuth angles. Compared with the auroral variations in Fig. 4 and Fig. 6, LOS
343 speeds exhibit diffusion during auroras, most evident in Fig. 7a after 18:30 UT on May 10th. ~~Concerning the auroral variations~~
344 ~~in Fig. 4 and Fig. 8, LOS speeds show diffusion during auroral events.~~ Negative LOS speeds initially concentrated to the north,
345 ~~northward,~~ spread westward and eastward, expanding horizontal coverage. Positive LOS speeds initially in the southward
346 direction shrink. When converted to horizontal wind speeds, these changes lead to increased horizontal differences, or the false
347 convergence caused by scattering, in other words. Across all azimuths, the changes of LOS speed share the same sign, but
348 their amplitude scales with the angle to the northward direction. Relative to the roughly 300 m·s⁻¹ LOS speed change in the
349 southward direction, the eastward and westward directions each attain about 60 %, whereas the northward variation remains
350 below 10 %. The scattering model shows that LOS speed changes in dimmer airglow regions are more than 10 times those in
351 the brighter zone. ~~The northward LOS speed changes slightly, the southward speed changes the most and nearly reverses,~~
352 ~~while the eastward and westward speeds are intermediate.~~




Figure 8: Proportion of scattered light at 45° zenith angle

The proportion is the ratio of scattered light intensity to the total light intensity (both scattered and direct) in the simulation. The figure shows this proportion at 45° zenith angle for various directions over time, with panels for May 10th, May 11th, and October 10th, respectively.

Figure 8 shows the ratio of scattered light intensity to total light intensity at a fixed 45° zenith angle calculated by the scattering model. Consistent with the schematic diagram (Fig. 1), the scattered intensity is the sum of all injected directions, and the total light intensity includes the direct component on this basis. Without auroras, the scattering proportion is typically below 0.4 and increases with atmospheric scattering capability, rising as the optical depth increases. ~~varies with atmospheric scattering capability.~~ However, during auroral events, the scattering proportion in some directions can increase to 0.5 or higher. The northward scattering proportion increases the least and remains close to that under uniform airglow conditions. ~~remains much lower than in other directions.~~ In contrast, the scattering proportion is significantly enhanced in directions ranging from 135° to 180° away from northward. The aurora appears in the north, resulting in much higher northern brightness. After atmospheric scattering, light from the north diffuses into surrounding directions, increasing the scattering proportion. Because the north itself has strong direct light, its scattering proportion remains small. In the model, we assume that stronger light rays dominate interference fringe identification (Wei et al., 2020), thereby determining the Doppler shift or LOS speeds. The lower scattering

369 proportion in the north allows northward observations to retain more LOS speeds from themselves, while other directions
 370 experience greater LOS speed contamination from the north. Therefore, as shown in Fig. 7, the northward simulation is closest
 371 to the default initial speed~~inputs~~, the southward simulation deviates the most, and the eastward and westward simulations lie
 372 in between. Additionally, the scattering impact should also consider the initial LOS speeds in the brighter region. Despite the
 373 high scattering proportion on May 11th shown in Fig. 7b, the simulated LOS speed changes in Fig. 8b are minimal. This is
 374 because of the smaller LOS speed in the auroral region on that day, resulting in less contamination spread to other directions.

375 The core working principle of the scattering model relies on the relationship between airglow brightness and background LOS
 376 wind speeds. As Harding et al. (2017a) noted, scattering requires a bright sky region with large LOS wind speeds. Firstly,
 377 spatially uneven airglow brightness is a prerequisite for significant scattering. The brightest airglow area contributes most to
 378 scattered light intensity, and its Doppler shifts determine the LOS biases in other directions. This principle allows a rough
 379 assessment of scattering impact without model computation when airglow is uneven. Locate the brightest region and its
 380 Doppler shift type, as the same Doppler shift type will likely appear in other directions. Blue-shift dominance indicates
 381 increased downward wind and horizontal deviations opposite to the line of sight, resembling convergence, while red-shift
 382 dominance resembles divergence. Minimal scattering-induced biases occur if the scattering proportion is very low due to
 383 uniform airglow brightness, or if the LOS velocity in the bright region is near zero (i.e., the wind speed is perpendicular to the
 384 line of sight). Previous observations can be directly verified by this principle and are basically in line with it (Hernandez and
 385 Roble, 1976; Price et al., 1995; Ishii et al., 1999; Ishii et al., 2001; Makela et al., 2014). Unfortunately, scattering impact can
 386 complicate dynamic analysis. In polar regions, auroras are characterized by green-line emissions, and thermospheric winds are
 387 significantly influenced by ion drag, where scattering effects may not be prominent. In contrast, mid-latitudes have mainly
 388 red-line auroras with large-scale uniform circulation, making the scattering impact more pronounced and distinguishable.

389 4.2 Errors of the scattering model

390 The model also exhibits certain errors and limitations. Scattering-induced biases in observations have nearly similar
 391 magnitudes on May 10th and Oct. 10th. However, with the same 100 m·s⁻¹ westward and 400 m·s⁻¹ southward wind input, the
 392 scattering-induced biases in the May case are significantly larger in magnitude and closer to reality compared to October. In
 393 Fig. 7, the simulated LOS speeds show a larger diffusion range for the May 10th case compared to October. In Fig. 8, the
 394 scattering proportion for May 10th is consistently higher than that for October. We attribute this difference mainly to the distinct
 395 optical depths in the two months, which are 0.43 and 0.2, respectively. Optical depth reflects atmospheric extinction capability
 396 and is mainly related to aerosol content (see Appendix B). It primarily affects the extinction process and influences the
 397 magnitude of scattering-induced biases by altering the proportion of scattered light. When the optical depth is artificially raised
 398 to 0.5, the model produces a meridional wind difference exceeding 400 m·s⁻¹ at 18:30 UT, Oct. 10th, roughly triple the value
 399 obtained with an optical depth of 0.2 and in much closer agreement with the observations. ~~When the optical depth is artificially~~
 400 ~~increased to a higher value, such as around 0.6, the model more closely matches the Oct. observations.~~ We find that the model

underestimates scattering effects when the optical depth is low. Once the optical depth reaches 0.6 or higher, the simulated wind bias accelerates nonlinearly until the model diverges.

In the October 10th event, the simulated scattering-induced biases are inconsistent between the two auroral brightness peaks. In Fig. 7, the LOS speed variation is larger during the second peak, and in Fig. 8, the scattering proportion is greater. This is because the scattering proportion is susceptible to errors in scattered and direct light intensities. DCAI does not correct for Rayleigh units, leading to significant errors in regions with large zenith angles. In other words, when the aurora appears at the edge of the field of view, its intensity may be underestimated due to vignetting. It is an effect opposite to the Van Rhijn enhancement. Correcting for Van Rhijn would then further reduce the edge brightness. For now, we extrapolate the edge values of the DCAI imager to mitigate this issue (Appendix B), yet some uncertainty remains. Besides, the ~~The~~-model cannot fully eliminate the stray light caused by the glass dome when separating the initial direct and scattered light from DCAI images (Harding et al., 2017a), resulting in errors. In our experiments, if the stray light effect is not subtracted as described in Appendix B, the model becomes more inert, resulting in a smaller simulated scattering proportion. In our experiments, without the stray-light correction in Appendix B, the airglow brightness gradient flattens slightly, the model becomes more inert, and the simulated horizontal wind differences shrink by about 30 %.

We also considered the potential influence of thermospheric temperature. FPI data show uniformly elevated thermospheric temperatures in these two storms, with the northward view occasionally about 300 K warmer than the others (not shown here). Because our scattering model does not yet include temperature effects, we cannot quantify how much scattering biases the FPI temperature measurements. In the study of Harding et al. (2017b), wind simulations are temperature-independent, while temperature retrieval relies on the wind. Likewise, we substitute the LOS speed for the Doppler shift and ignore temperature-induced spectral broadening. In principle, thermospheric temperature influences retrieval uncertainty, not the wind speed itself. We remain cautious that ignoring this uncertainty could introduce extra bias if a horizontal temperature gradient is present, but incorporating it would markedly raise the computational cost and remains a task for the future.

On Oct. 10th, even after the refinements listed below to reduce the underestimation, the scattering model still accounts for only about 25 % of the observed horizontal wind difference: In our experiments, even after several refinements aimed at mitigating the underestimation, the scattering model still underestimates the observed scattering impact: (1) A single-scattering albedo of 1 was used, ignoring absorption. (2) Stray light effects were removed. (3) Attenuated airglow observations at the edge of DCAI images were extrapolated, enhancing edge scattering. (4) Excessive edge extinction was reduced by correcting the extinction path geometry, increasing the scattered light intensity integral. (5) Zero vertical wind was assumed when converting simulated LOS speeds to geographical wind speeds. Raising the optical depth from the observed 0.2 to 0.5 would reduce the underestimation, yet clear-sky DCAI images appear to rule out such high values. We suspect the underestimation arises from how the model integrates the optical depth. Since the integral only includes 10 optical depth layers, with each light ray scattering once per layer and extinguishing once between layers, it may be too crude compared to the real path, underestimating

433 the scattered light. Simply increasing the number of optical depth layers is not effective in our experiments. We think this may
434 be related to the non-linear variation of atmospheric density with altitude, where optical depth may not vary linearly with
435 height, and the scattering phase function may also change with altitude. To address these issues, future work should complete
436 the DCAI correction. Additionally, introducing a model of optical depth varying with altitude can increase the number of
437 single-scattering nodes and ensure the geometric accuracy of the extinction path, thereby improving the accuracy of scattered
438 light intensity calculations.

439 Furthermore, these bright region observations do not necessarily reflect the usual 250 km thermospheric wind. In Fig. 3 and
440 Fig. 5, the north-looking wind observations show unusually high wind speeds, which are significantly different from the
441 simulations. In particular, on October 10th, the north-looking wind speed varied dramatically with the intensity of the northern
442 aurora. During the two auroral peaks, the north-looking wind direction also reversed. This indicates that the interferometer
443 receives an additional effect when it looks toward the aurora. Kataoka et al. showed that red aurora lifted the red-line emission
444 profile, raising its peak above 300 km and brightening the upper part on May 11th (Kataoka et al., 2024b). Consequently, the
445 interferometer can sample winds that are higher and more poleward. Because storm-time surges propagate from the polar
446 region to the equator, these higher, poleward regions are likely to carry stronger equatorward winds. The interferometer may
447 record a larger wind speed toward the aurora. Additionally, spectral contamination from precipitating energetic ions can also
448 bias interferometers (Makela et al., 2014). In other words, the interferometer is partly sensing the speed of non-neutral species,
449 boosting the observed wind. These issues lie beyond what scattering models can reproduce. From the observed pattern, we
450 infer the presence of non-scattering effects, especially in the poleward view. Due to the absence of nearby higher-latitude
451 neutral-wind observations relative to SIZW, quantifying their respective contributions remains challenging.

452 5 Conclusions

453 This study has further proved that lower atmospheric scattering can contribute to biases in thermospheric wind observation on
454 ground-based optical interferometers. During two geomagnetic storms with spatially uneven airglow on May 10th and Oct. 10th,
455 2024, the the light scattered from the non-line-of-sight directions of the scope will lead to additional LOS speeds and appear
456 as atypical horizontal differences exceeding 400 m·s⁻¹ and downward vertical wind exceeding 100 m·s⁻¹ at geographic
457 coordinates. With a simplified scattering radiative transfer model, we simulate the distribution of airglow intensity after the
458 multiple scattering of the lower atmosphere and estimate the wind observation bias under scattering impact via a weighted
459 average method. ~~Atypical winds under conditions of spatially uneven airglow have been generally captured.~~ Starting from the
460 assumed zero vertical wind and spatially uniform, steady horizontal winds (100 m·s⁻¹ westward with 400 m·s⁻¹ southward),
461 the model produces approximately 400 m·s⁻¹ horizontal differences between opposite cardinal directions and 100 m·s⁻¹
462 downwelling on May 10th, in basic agreement with observations, whereas on Oct. 10th it yields only 100 m·s⁻¹ horizontal
463 differences, amounting to about 75 % underestimation in magnitude, yet it still captures the temporal trend.

464 We have refined the scattering model in previous research to enhance its computational efficiency. Specifically, we simplified
 465 the LOS wind speed simulation and capped the lower atmosphere at 40 km to refine the extinction-length calculation.
 466 ~~introduced an upper limit for the lower atmosphere to improve the accuracy of the extinction-length calculation.~~ The scattering
 467 impact can be directly estimated through the relationship between the bright airglow region and the LOS wind speed. The
 468 brightest airglow region contributes most to the scattering impact, of which the Doppler shift type determines the LOS biases
 469 in other directions. Although the observed winds are affected by scattering when airglow is uneven, they still retain dynamic
 470 information, such as the average wind being close to the storm-time circulation. Unfortunately, we lack alternative
 471 observational methods to verify the accuracy of the interferometer results. It deserves further study to the extent of scattering
 472 impact with more cases and additional instrumental observations. Limited by the accuracy of the model inputs, the scattering
 473 model can only simulate the wind features associated with scattering impact under clear sky conditions. It remains incapable
 474 of precisely picking out the speed biases induced by scattering impact. Either artificially doubling the optical depth or
 475 subtracting stray light can introduce uncertainties exceeding 30 % in simulated scattering-induced biases. Future model
 476 improvements could include in situ real-time optical depth measurements, airglow imager corrections, and incorporating
 477 vertical optical depth profiles into the model.

~~Limited by the accuracy of the model inputs, the scattering model can only simulate the wind features associated with scattering
 impact under clear sky conditions. It remains incapable of precisely picking out the speed alterations induced by scattering
 impact. Given that scattering impacts observations during geomagnetic storms accompanied by auroras, it is necessary to
 quantify these biases to provide accurate data for dynamic studies. Therefore, future efforts can focus on refining the model
 and its inputs or statistically analyzing scattering biases under different airglow brightness distributions from various stations
 to gain a clearer understanding of the magnitude of scattering biases.~~

484 Appendix A

485 In the following appendices, we provide a concise description of the scattering model's operational principles, inputs, and
 486 modifications employed in our works. For more detailed solutions, please refer to the article by Harding et al. (Harding et al.,
 487 2017a).

488 Based on the radiation transfer theory, Hansen and Travis (1974) and Sobolev (1975) gave the multiple scattering solution.
 489 Harding et al. (2017a) extended the initial source function $J_0(\tau, u, \phi)$ to airglow surface source, and corrected the missing
 490 normalization factor in the scattering phase function (Eq. (1) to Eq. (3)):

$$491 \quad u \frac{dI(\tau, u, \phi)}{d\tau} = -I(\tau, u, \phi) + J(\tau, u, \phi) \quad (1)$$

$$492 \quad J(\tau, u, \phi) = \frac{\omega}{4\pi} \int_0^{2\pi} \int_{-1}^1 P(u, u', \phi, \phi') I(\tau, u', \phi') du' d\phi' + J_0(\tau, u, \phi) \quad (2)$$

$$J_0(\tau, u, \phi) = \frac{\omega}{4\pi} \int_0^{2\pi} \int_0^1 P(u, u', \phi, \phi') \sec(\gamma') f(u', \phi') \exp[-\tau L(u')] du' d\phi' \quad (3)$$

$$L(u) = [(R_e + H_L) \cos(\gamma_\tau) - R_e u] H_L^{-1} \quad (4)$$

$$u = \cos(\zeta) \quad (5)$$

The equations are formulated within an improved local spherical coordinate system, including azimuth ϕ , zenith angle ζ which is represented in cosine form u , and vertical height which is expressed as optical depth τ .

In the case of scattering, as illustrated in Fig. 1, the light intensity along a line of sight, represented by $I(\tau, u, \phi)$, consists of two parts, the direct light (a in Fig. 1) from the same direction, and the aggregate of scattered light (b-e in Fig. 1) from other directions, which is represented by the source function $J(\tau, u, \phi)$. Based on the radiative transfer equation (Eq. (1)), at each optical depth layer, the scattered light intensity received from all directions will be integrated. Simultaneously, the original intensity in the line of sight will be added to the total scattered light. Besides, the extinction in the path will be calculated according to the optical depth.

There are two potential scattering paths in the lower atmosphere: single scattering (b, c in Fig. 1) and multiple scattering (d, e in Fig. 1). The model computes them sequentially via an iterative process. In the initial state, there is no light intensity in the lower atmosphere. Therefore, the single scattering will originate solely from the airglow layer, and the source function $J(\tau, u, \phi)$ will be equivalent to the initial source function $J_0(\tau, u, \phi)$ at this state. By solving Eq. (1), the model can obtain the single scattered intensity in each direction at every optical depth layer, which is the updated source function $J(\tau, u, \phi)$. Then, the multiple scattering can be calculated based on it. Typically, the total scattered intensity will remain relatively constant when accounting for the fourth scattering. By using this iteration, the scattered light and residual direct light in DCAI images can be effectively separated. The residual direct light will subsequently serve as the background intensity distribution for simulating speeds.

In the source function $J(\tau, u, \phi)$, the scattering phase function $P(u, u', \phi, \phi')$ quantifies the relative gain of an incident angle to an exit angle during the scattering process. The reference value is based on a unit-radius sphere, which necessitates the introduction of a factor $\frac{1}{4\pi}$. Furthermore, ω represents the single-scattering albedo, set as 1. The initial source function $J_0(\tau, u, \phi)$ is responsible for introducing the airglow distribution $f(u', \phi')$. Here, $\sec(\gamma')$ represents the secant of the zenith angle at the puncture point of the airglow layer, which helps eliminate the Van Rhijn effect. Additionally, the exponential term with base e is utilized to calculate the equivalent extinction length along an inclined path.

It is primarily the lower atmosphere that significantly scatters and absorbs light (He et al., 2021; Li et al., 2022). Therefore, when computing the extinction length, just employing the cosine of zenith angle u' will lead to an overestimation of the effective length, as illustrated on the right side of Fig. 1. To address it, we set an upper boundary H_L of the lower atmosphere

at 40 km, assuming an optical depth of zero above this altitude. Using geometric relationships, an equivalent length factor $L(u)$ can be derived, where R_e means Earth radius, and γ_t represents the zenith angle at the penetration point of 40 km height. This value can be readily calculated by adjusting the target height of the formula for γ . Inside the lower atmosphere, we apply a thin-layer approximation, which also utilizes the geometric relationships at the upper boundary.

$$V = \frac{V_{dr}I_{dr} + \sum_k V_{sc}(k)I_{sc}(k)}{I_{dr} + \sum_k I_{sc}(k)} \quad (6)$$

After working out the background intensity distribution, we partition the LOS speeds at 250 km into several bins. To simplify simulation, the model directly uses LOS speeds corresponding to Doppler shifts. As roughly illustrated in Fig. 1, all LOS speeds are categorized into $k=10$ bins valued from highest to lowest, assuming that the LOS speeds within each bin approximate their mean value V_{sc} . The scattered intensity distribution I_{sc} is computed by extracting the airglow brightness from the corresponding region of each bin. And, there will be no intensity from other areas during a single bin's computing. According to Eq. (6), the simulative LOS speed at a specific angle will be an averaged result, where the original speed V_{dr} is weighted by the direct light intensity I_{dr} , and the additional speed resulting from scattering V_{sc} is weighted by the scattered light intensity I_{sc} . The model directly uses the average within DCOI's 9 degree field of view as simulated post-scattering LOS speed of interferometers, since DCOI and FPI have not measured the reception gain of light outside their fields of view. We find that due to the coarse model grid, the 9 degree average is nearly the same as using the nearest single-sight observation. Finally, the LOS speed will be converted to horizontal wind, maintaining the assumption of zero vertical wind to prevent the propagation of scattering biases in the vertical direction.

Appendix B

This appendix details the scattering model's inputs from measurements, supplementing the second section. The background airglow brightness for the model comes from DCAI. Image processing includes: (1) dark field exclusion, (2) median filtering to remove starlight, (3) conversion to the local spherical coordinate, (4) stray light correction, and (5) radial zero-order extrapolation for regions beyond 70° zenith angle. Stray light results from the scattering of strong incident light by the glass dome. During quiet nights without auroras, it is weak and uniformly distributed across all LOS directions. We use the azimuthal average of the nearest quiet night at 45° zenith angle as a reference for weak stray light conditions. After the aurora onset, stray light brightens all directions. The difference between the darkest direction at 45° zenith angle and the reference value is considered the additional stray light caused by the aurora and is subtracted from the entire image. The shown airglow images additionally mitigate the Van Rhijn effect through $\sec(\gamma')$ and the atmospheric extinction through $\exp[-\tau L(u')]$ (see Appendix A). Since the scattering model already includes these processes, no separate treatment is needed. The optical depth and scattering phase function inputs are shown in Fig. 9. Optical depth is calculated using AERONET's monthly averages, with interpolation at 630 nm. Since only daytime observations are available, local daytime values are used to represent

nighttime values. Fig. 9a and 9b show monthly average optical depths at local daytime, with total averages of 0.43 and 0.2. The scattering phase function is a weighted average of molecular and aerosol scattering phase functions from AERONET at 675 nm, which is weighted on the total optical depth of aerosols and molecules.




Figure 9: Optical depth and scattering phase function in May and October

The first two panels show daytime monthly average optical depths, with molecular optical depth amplified for clarity. The rest panels show the scattering phase functions.

Code availability

The code of DCAI images correction, the scattering model, and the visualization are not publicly available yet. If needed, they can be obtained by contacting the corresponding authors via email.

Data availability

The data of DCOI and DCAI from the Chinese Meridian Project can be obtained from <https://www.meridianproject.ac.cn/en/>. The FPI data can be obtained by contacting the corresponding authors via email. The data of AERONET can be obtained from

565 <https://aeronet.gsfc.nasa.gov/>. The Kp index provided by GFZ, German Research Centre for Geosciences, can be obtained
566 from <https://kp.gfz-potsdam.de/en/>.

567 **Author contributions**

568 Conceptualization: XW; investigation: XW, GJ; instruments construction and maintenance: YZ, JX, WL, TW, GZ, WY; wind
569 data retrieval: YZ, WL, GZ, TW; airglow images correction: XW; model improvement and programming: XW; visualization:
570 XW; validation: GJ, YZ, JX, WL, TW; writing - original draft preparation: XW; writing-review and editing: GJ, YZ, JX, WL,
571 TW; supervision: GJ, YZ; project administration: GJ, YZ; funding acquisition: YZ. All authors contributed to the revision and
572 improvement of the paper.

573 **Competing interests**

574 The authors declare that they have no conflict of interest.

575 **Acknowledgements**

576 We appreciate all the funding from the National Key R&D program of China (2023YFB3905100), the Project of Stable Support
577 for Youth Team in Basic Research Field, CAS (YSBR-018), the National Natural Science Foundation of China (42174212),
578 the Chinese Meridian Project, and the Specialized Research Fund for State Key Laboratories. We acknowledge the use of data
579 from the Chinese Meridian Project. We thank all the builders and maintainers of DCOI, FPI, and DCAI of Siziwang station.
580 We thank Lingli Tang for their effort in establishing and maintaining AOE_Baotou site of AERONET.

581 **Financial support**

582 This work was supported by the National Key R&D program of China (2023YFB3905100), the Project of Stable Support for
583 Youth Team in Basic Research Field, CAS (YSBR-018), the National Natural Science Foundation of China (42174212), the
584 Chinese Meridian Project, and the Specialized Research Fund for State Key Laboratories.

585 **References**

586 Abreu, V. J., Schmitt, G. A., Hays, P. B., Meriwether, J. W., Tepley, C. A., and Cogger, L. L.: Atmospheric Scattering Effects
587 on Ground-Based Measurements of Thermospheric Winds, Planet Space Sci, 31, 303-310, doi:10.1016/0032-
588 0633(83)90080-6, 1983.
589 Anderson, C., Conde, M., and McHarg, M. G.: Neutral thermospheric dynamics observed with two scanning Doppler imagers:
590 2. Vertical winds, Journal of Geophysical Research: Space Physics, 117, doi:10.1029/2011ja017157, 2012.

591 Ashburn, E. V.: The Effect of Rayleigh Scattering and Ground Reflection Upon the Determination of the Height of the Night
592 Airglow, *J Atmos Terr Phys*, 5, 83-91, doi:10.1016/0021-9169(54)90012-4, 1954.

593 Biondi, M. A. and Feibelman, W. A.: Twilight and nightglow spectral line shapes of oxygen $\lambda 6300$ and $\lambda 5577$ radiation, *Planet*
594 *Space Sci*, 16, 431-443, doi:10.1016/0032-0633(68)90158-X, 1968.

595 Biondi, M. A., Sipler, D. P., Zipf, M. E., and Baumgardner, J. L.: All-Sky Doppler Interferometer for Thermospheric Dynamics
596 *Studies*, *Appl Optics*, 34, 1646-1654, doi:10.1364/Ao.34.001646, 1995.

597 Burnside, R. G., Herrero, F. A., Meriwether, J. W., and Walker, J. C. G.: Optical observations of thermospheric dynamics at
598 Arecibo, *Journal of Geophysical Research: Space Physics*, 86, 5532-5540, doi:10.1029/JA086iA07p05532, 1981.

599 Burnside, R. G. and Tepley, C. A.: Optical Observations of Thermospheric Neutral Winds at Arecibo between 1980 and 1987,
600 *J Geophys Res-Space*, 94, 2711-2716, doi:10.1029/JA094iA03p02711, 1989.

601 Conde, M. and Smith, R. W.: Mapping Thermospheric Winds in the Auroral-Zone, *Geophysical Research Letters*, 22, 3019-
602 3022, doi:10.1029/95gl02437, 1995.

603 Conde, M., Craven, J. D., Immel, T., Hoch, E., Stenbaek-Nielsen, H., Hallinan, T., Smith, R. W., Olson, J., Sun, W., Frank, L.
604 A., and Sigwarth, J.: Assimilated observations of thermospheric winds, the aurora, and ionospheric currents over Alaska,
605 *Journal of Geophysical Research: Space Physics*, 106, 10493-10508, doi:10.1029/2000ja000135, 2001.

606 Crickmore, R. I., Dudeney, J. R., and Rodger, A. S.: Vertical Thermospheric Winds at the Equatorward Edge of the Auroral
607 Oval, *J Atmos Terr Phys*, 53, 485-492, doi:10.1016/0021-9169(91)90076-J, 1991.

608 Dhadly, M., Emmert, J., Drob, D., Conde, M., Doornbos, E., Shepherd, G., Makela, J., Wu, Q., Niciejewski, R., and Ridley,
609 A.: Seasonal dependence of northern high-latitude upper thermospheric winds: A quiet time climatological study based
610 on ground-based and space-based measurements, *Journal of Geophysical Research: Space Physics*, 122, 2619-2644,
611 doi:10.1002/2016JA023688, 2017.

612 Emmert, J. T., Fejer, B. G., Fesen, C. G., Shepherd, G. G., and Solheim, B. H.: Climatology of middle- and low-latitude
613 daytime region disturbance neutral winds measured by Wind Imaging Interferometer (WINDII), *J Geophys Res-Space*,
614 106, 24701-24712, doi:10.1029/2000ja000372, 2001.

615 Emmert, J. T., Faivre, M. L., Hernandez, G., Jarvis, M. J., Meriwether, J. W., Niciejewski, R. J., Sipler, D. P., and Tepley, C.
616 A.: Climatologies of nighttime upper thermospheric winds measured by ground-based Fabry-Perot interferometers during
617 geomagnetically quiet conditions: 1. Local time, latitudinal, seasonal, and solar cycle dependence, *J Geophys Res-Space*,
618 111, ArtN A12302, doi:10.1029/2006ja011948, 2006.

619 Englert, C. R., Babcock, D. D., and Harlander, J. M.: Doppler asymmetric spatial heterodyne spectroscopy (DASH): concept
620 and experimental demonstration, *Appl Optics*, 46, 7297-7307, doi:10.1364/Ao.46.007297, 2007.

621 Englert, C. R., Harlander, J. M., Emmert, J. T., Babcock, D. D., and Roesler, F. L.: Initial ground-based thermospheric wind
622 measurements using Doppler asymmetric spatial heterodyne spectroscopy (DASH), *Optics Express*, 18, 27416-27430,
623 doi:10.1364/Oe.18.027416, 2010.

624 Fejer, B. G., Emmert, J. T., and Sipler, D. P.: Climatology and storm time dependence of nighttime thermospheric neutral
625 winds over Millstone Hill, *Journal of Geophysical Research: Space Physics*, 107, doi:10.1029/2001ja000300, 2002.

626 Friedman, J. F. and Herrero, F. A.: Fabry-Perot interferometer measurements of thermospheric neutral wind gradients and
627 reversals at Arecibo, *Geophysical Research Letters*, 9, 785-788, doi:10.1029/GL009i007p00785, 1982.

628 Gonzalez-Esparza, J. A., Sanchez-Garcia, E., Sergeeva, M., Corona-Romero, P., Gonzalez-Mendez, L. X., Valdes-Galicia, J.
629 F., Aguilar-Rodriguez, E., Rodriguez-Martinez, M., Ramirez-Pacheco, C., Castellanos, C. I., Pazos, M., Mendoza, B.,
630 Gatica-Acevedo, V. J., Melgarejo-Morales, A., Caraballo, R., Andrade-Mascote, E., Villanueva-Hernandez, P., Bonifaz-
631 Alfonso, R., Sierra, P., Romero-Hernandez, E., Peralta-Mendoza, I., Perez-Tijerina, E., Mejia-Ambroz, J. C., Guerrero-
632 Peña, C., Caccavari, A., Cifuentes-Nava, G., and Hernandez-Quintero, E.: The Mother's Day Geomagnetic Storm on 10
633 May 2024: Aurora Observations and Low Latitude Space Weather Effects in Mexico, *Space Weather-the International*
634 *Journal of Research and Applications*, 22, ARTN e2024SW004111, doi:10.1029/2024SW004111, 2024.

635 Guo, W. and McEwen, D. J.: Vertical winds in the central polar cap, *Geophysical Research Letters*, 30, ArtN 1725,
636 doi:10.1029/2003gl017124, 2003.

637 Guo, X., Zhao, B. Q., Yu, T. T., Hao, H. L., Sun, W. J., Wang, G. J., He, M. S., Mao, T., Li, G. Z., and Ren, Z. P.: East-West
638 Difference in the Ionospheric Response During the Recovery Phase of May 2024 Super Geomagnetic Storm Over the
639 East Asian, *J Geophys Res-Space*, 129, ARTN e2024JA033170, doi:10.1029/2024JA033170, 2024.

640 Hajra, R., Tsurutani, B. T., Lakhina, G. S., Lu, Q. M., and Du, A. M.: Interplanetary Causes and Impacts of the 2024 May
641 Superstorm on the Geosphere: An Overview, *Astrophys J*, 974, ARTN 264, doi:10.3847/1538-4357/ad7462, 2024.

642 Hansen, J. E. and Travis, L. D.: Light-Scattering in Planetary Atmospheres, *Space Science Reviews*, 16, 527-610, doi:
643 10.1007/Bf00168069, 1974.

644 Harding, B. J., Makela, J. J., Qin, J., Fisher, D. J., Martinis, C. R., Noto, J., and Wrasse, C. M.: Atmospheric scattering effects
645 on ground-based measurements of thermospheric vertical wind, horizontal wind, and temperature, *Journal of Geophysical*
646 *Research: Space Physics*, 122, 7654-7669, doi:10.1002/2017JA023942, 2017a.

647 Harding, B. J., Qin, J., and Makela, J. J.: Ground-Based Optical Measurements of Quiet Time Thermospheric Wind and
648 Temperature: Atmospheric Scattering Corrections, *Journal of Geophysical Research: Space Physics*, 122,
649 doi:10.1002/2017JA024705, 2017b.

650 Harlander, J. M., Englert, C. R., Brown, C. M., Marr, K. D., Miller, I. J., Zastera, V., Bach, B. W., and Mende, S. B.: Michelson
651 Interferometer for Global High-Resolution Thermospheric Imaging (MIGHTI): Monolithic Interferometer Design and
652 Test, *Space Science Reviews*, 212, 601-613, doi:10.1007/s11214-017-0374-4, 2017.

653 Hays, P. B., Killeen, T. L., Spencer, N. W., Wharton, L. E., Roble, R. G., Emery, B. A., Fullerrowell, T. J., Rees, D., Frank,
654 L. A., and Craven, J. D.: Observations of the Dynamics of the Polar Thermosphere, *J Geophys Res-Space*, 89, 5597-5612,
655 doi:10.1029/JA089iA07p05597, 1984.

656 He, Q. F., Fang, Z., Shoshanim, O., Brown, S. S., and Rudich, Y.: Scattering and absorption cross sections of atmospheric
657 gases in the ultraviolet-visible wavelength range (307-725 nm), *Atmos Chem Phys*, 21, 14927-14940, doi:10.5194/acp-
658 21-14927-2021, 2021.

659 Hernandez, G. and Roble, R. G.: Direct Measurements of Nighttime Thermospheric Winds and Temperatures .2. Geomagnetic
660 Storms, *J Geophys Res-Space*, 81, 5173-5181, doi:10.1029/JA081i028p05173, 1976.

661 Holben, B. N., Tanré, D., Smirnov, A., Eck, T. F., Slutsker, I., Abuhassan, N., Newcomb, W. W., Schafer, J. S., Chatenet, B.,
662 Lavenu, F., Kaufman, Y. J., Castle, J. V., Setzer, A., Markham, B., Clark, D., Frouin, R., Halthore, R., Karnieli, A.,
663 O'Neill, N. T., Pietras, C., Pinker, R. T., Voss, K., and Zibordi, G.: An emerging ground-based aerosol climatology::
664 Aerosol optical depth from AERONET, *J Geophys Res-Atmos*, 106, 12067-12097, doi:10.1029/2001jd900014, 2001.

665 Huang, C., Xu, J. Y., Zhang, X. X., Liu, D. D., Yuan, W., and Jiang, G. Y.: Mid-latitude thermospheric wind changes during
666 the St. Patrick's Day storm of 2015 observed by two Fabry-Perot interferometers in China, *Advances in Space Research*,
667 61, 1873-1879, doi:10.1016/j.asr.2017.10.013, 2018.

668 Innis, J. L., Greet, P. A., Murphy, D. J., Conde, M. G., and Dyson, P. L.: A large vertical wind in the thermosphere at the
669 auroral oval/polar cap boundary seen simultaneously from Mawson and Davis, Antarctica, *J Atmos Sol-Terr Phy*, 61,
670 1047-1058, doi:10.1016/S1364-6826(99)00060-7, 1999.

671 Ishii, M., Oyama, S., Nozawa, S., Fujii, R., Sagawa, E., Watari, S., and Shinagawa, H.: Dynamics of Neutral Wind in the polar
672 region observed with two Fabry-Perot interferometers, *Earth Planets Space*, 51, 833-844, doi:10.1186/Bf03353242, 1999.

673 Ishii, M., Conde, M., Smith, R. W., Krynicki, M., Sagawa, E., and Watari, S.: Vertical wind observations with two Fabry-
674 Perot interferometers at Poker Flat, Alaska, *Journal of Geophysical Research: Space Physics*, 106, 10537-10551,
675 doi:10.1029/2000JA900148, 2001.

676 Jiang, G., Xu, J., Wang, W., Yuan, W., Zhang, S., Yu, T., Zhang, X., Huang, C., Kerr, R. B., Noto, J., Li, J., Liu, W., and Li,
677 Q.: A Comparison of Quiet Time Thermospheric Winds Between FPI Observations and Model Calculations, *Journal of*
678 *Geophysical Research: Space Physics*, 123, 7789-7805, doi:10.1029/2018JA025424, 2018.

679 Kataoka, R., Miyoshi, Y., Shiokawa, K., Nishitani, N., Keika, K., Amano, T., and Seki, K.: Magnetic Storm-Time Red Aurora
680 as Seen From Hokkaido, Japan on 1 December 2023 Associated With High-Density Solar Wind, *Geophysical Research*
681 *Letters*, 51, e2024GL108778, doi:10.1029/2024GL108778, 2024a.

682 Kataoka, R., Reddy, S. A., Nakano, S., Pettit, J., and Nakamura, Y.: Extended magenta aurora as revealed by citizen science,
683 *Scientific Reports*, 14, 25849, doi:10.1038/s41598-024-75184-9, 2024b.

684 Killeen, T. L., Won, Y. I., Niciejewski, R. J., and Burns, A. G.: Upper Thermosphere Winds and Temperatures in the
685 Geomagnetic Polar-Cap - Solar-Cycle, Geomagnetic-Activity, and Interplanetary Magnetic-Field Dependencies, *J*
686 *Geophys Res-Space*, 100, 21327-21342, doi:10.1029/95ja01208, 1995.

687 Li, J., Carlson, B. E., Yung, Y. L., Lv, D. R., Hansen, J., Penner, J. E., Liao, H., Ramaswamy, Kahn, R. A., Zhang, P., Dubovik,
688 O., Ding, A. J., Lacis, A. A., Zhang, L., and Dong, Y. M.: Scattering and absorbing aerosols in the climate system, *Nat*
689 *Rev Earth Env*, 3, 363-379, doi:10.1038/s43017-022-00296-7, 2022.

Li, W. B., Liu, L. B., Chen, Y. D., Yang, Y. Y., Han, T. W., Ding, F., Le, H. J., and Zhang, R. L.: Multi-Instruments Observation of Ionospheric-Thermospheric Dynamic Coupling Over Mohe (53.5°N, 122.3°E) During the April 2023 Geomagnetic Storm, *J Geophys Res-Space*, 128, ARTN e2023JA032141, doi:10.1029/2023JA032141, 2023.

Liu, W. J., Zhu, Y. J., Xu, J. Y., Li, Q. Z., Yuan, W., Zhu, G. Y., Wang, T. C., Yang, G. T., Du, L. F., Liu, S. Y., and Li, F. Q.: Validation of Neutral Wind in the Mesopause Measured by a Dual-Channel Optical Interferometer (DCOI) Network of the Chinese Meridian Project, *Space Weather-the International Journal of Research and Applications*, 23, ARTN e2025SW004468, doi:10.1029/2025SW004468, 2025.

Ma, L. X., Yu, Y. Q., Ding, X. Q., Liu, X. Y., An, D. P., Zhou, C. L., Cao, J. B., and Shiokawa, K.: Mid-Latitude Auroras and Energetic Particle Precipitation Occurred Unusually in a Moderate Magnetic Storm on 1 December 2023, *Geophysical Research Letters*, 51, ARTN e2024GL110764, doi:10.1029/2024GL110764, 2024.

Makela, J. J., Harding, B. J., Meriwether, J. W., Mesquita, R., Sanders, S., Ridley, A. J., Castellez, M. W., Ciocca, M., Earle, G. D., Frissell, N. A., Hampton, D. L., Gerrard, A. J., Noto, J., and Martinis, C. R.: Storm time response of the midlatitude thermosphere: Observations from a network of Fabry-Perot interferometers, *Journal of Geophysical Research: Space Physics*, 119, 6758-6773, doi:10.1002/2014JA019832, 2014.

Mikhalev, A.: Auroras during extreme geomagnetic storms: Some features of mid-latitude aurora on February 11, 1958, *Solar-Terrestrial Physics*, 55-61, 10.12737/stp-102202406, 2024.

Nakajima, H., Okano, S., Fukunishi, H., and Ono, T.: Observations of thermospheric wind velocities and temperatures by the use of a Fabry-Perot Doppler imaging system at Syowa Station, Antarctica, *Appl Optics*, 34, 8382-8395, doi:10.1364/Ao.34.008382, 1995.

Nanjo, S. and Shiokawa, K.: Spatial structures of blue low-latitude aurora observed from Japan during the extreme geomagnetic storm of May 2024, *Earth Planets Space*, 76, ARTN 156, doi:10.1186/s40623-024-02090-9, 2024.

Price, G. D., Smith, R. W., and Hernandez, G.: Simultaneous Measurements of Large Vertical Winds in the Upper and Lower Thermosphere, *J Atmos Terr Phys*, 57, 631-643, doi:10.1016/0021-9169(94)00103-U, 1995.

Ranjan, A. K. and Pallamraju, D.: Latitudinal Distribution of Thermospheric Nitric Oxide (NO) Infrared Radiative Cooling During May and October 2024 Geomagnetic Storms, *J Geophys Res-Space*, 130, ARTN e2024JA033559, doi:10.1029/2024JA033559, 2025.

Rees, D., Smith, M. F., and Gordon, R.: The Generation of Vertical Thermospheric Winds and Gravity-Waves at Auroral Latitudes .2. Theory and Numerical Modeling of Vertical Winds, *Planet Space Sci*, 32, 685, doi:10.1016/0032-0633(84)90093-X, 1984.

Sakanai, T., Fukunishi, H., Okano, S., Sato, N., Yamagishi, H., and Yukimatu, A. S.: Dynamical coupling of neutrals and ions in the high-latitude *F* region: Simultaneous FPI and HF radar observations at Syowa Station, Antarctica, *Journal of Geophysical Research: Space Physics*, 107, doi:10.1029/2001JA007530, 2002.

Shinagawa, H. and Oyama, S.: A two-dimensional simulation of thermospheric vertical winds in the vicinity of an auroral arc, *Earth Planets Space*, 58, doi:10.1186/BF03352007, 2006.

Shiokawa, K., Katoh, Y., Satoh, M., Ejiri, M. K., and Ogawa, T.: Integrating-sphere calibration of all-sky cameras for nightglow measurements, *Adv Space Res*, 26, 1025-1028, doi:10.1016/S0273-1177(00)00052-1, 2000.

Singh, R., Scipión, D. E., Kuyeng, K., Condor, P., Flores, R., Pacheco, E., de la Jara, C., and Manay, E.: Ionospheric Responses to an Extreme (G5-Level) Geomagnetic Storm Using Multi-Instrument Measurements at the Jicamarca Radio Observatory on 10-11 October 2024, *J Geophys Res-Space*, 130, ARTN e2024JA033642, doi:10.1029/2024JA033642, 2025.

Smith, R. W. and Hernandez, G.: Vertical Winds in the Thermosphere within the Polar-Cap, *J Atmos Terr Phys*, 57, 611-620, doi:10.1016/0021-9169(94)00101-S, 1995.

Sobolev, V. V., Irvine, W. M., Gendel, M., and Lane, A. P.: Light scattering in planetary atmospheres, 1st . International series of monographs in natural philosophy vol.76, Pergamon Press, Oxford, New York, doi:10.1016/C2013-0-05661-7, 1975.

Themens, D. R., Elvidge, S., McCaffrey, A., Jayachandran, P. T., Coster, A., Varney, R. H., Galkin, I., Goodwin, L. V., Watson, C., Maguire, S., Kavanagh, A. J., Zhang, S. R., Goncharenko, L., Bhatt, A., Dorrian, G., Groves, K., Wood, A. G., and Reid, B.: The High Latitude Ionospheric Response to the Major May 2024 Geomagnetic Storm: A Synoptic View, *Geophysical Research Letters*, 51, e2024GL111677, doi:10.1029/2024GL111677, 2024.

Wang, C., Xu, J., Chen, Z., Li, H., Feng, X., Huang, Z., and Wang, J.: China's Ground-Based Space Environment Monitoring Network—Chinese Meridian Project (CMP), *Space Weather*, 22, doi:10.1029/2024sw003972, 2024.

740 Wang, X., Aa, E., Chen, Y., Zhang, J., Zhu, Y., Cai, L., Lu, X., Luo, B., Liu, S., Li, M., Shen, H., and Yuan, T.: Midlatitude
741 Neutral Wind Response During the Mother's Day Super-Intense Geomagnetic Storm in 2024 Using Observations From
742 the Chinese Meridian Project, *Journal of Geophysical Research: Space Physics*, 130, e2024JA033574,
743 doi:10.1029/2024JA033574, 2025.

744 Wei, D., Zhu, Y., Liu, J., Gong, Q., Kaufmann, M., Olschewski, F., Knieling, P., Xu, J., Koppmann, R., and Riese, M.:
745 Thermally stable monolithic Doppler asymmetric spatial heterodyne interferometer: optical design and laboratory
746 performance, *Optics Express*, 28, doi:10.1364/oe.394101, 2020.

747 Wu, Q., Yuan, W., Xu, J. Y., Huang, C., Zhang, X. X., Wang, J. S., and Li, T.: First U.S.-China joint ground-based Fabry-
748 Perot interferometer observations of longitudinal variations in the thermospheric winds, *J Geophys Res-Space*, 119, 5755-
749 5763, doi:10.1002/2014ja020089, 2014.

750 Xu, H., Shiokawa, K., Oyama, S., and Otsuka, Y.: Thermospheric wind variations observed by a Fabry-Perot interferometer
751 at Tromsø, Norway, at substorm onsets, *Earth Planets Space*, 71, ARTN 93, doi:10.1186/s40623-019-1072-0, 2019.

752 Yang, C., Zhao, B., Jin, Y., Huang, C., Yao, X., and Wan, W.: Climatology of Nighttime Upper Thermospheric Winds From
753 Fabry-Perot Interferometer 2011–2019 Measurements Over Kelan (38.7°N, 111.6°E), China: Local Time, Seasonal, Solar
754 Cycle, and Geomagnetic Activity Dependence, *Journal of Geophysical Research: Space Physics*, 125, e2020JA027892,
755 doi:10.1029/2020JA027892, 2020.

756 Yu, T., Huang, C., Zhao, G. X., Mao, T., Wang, Y. G., Zeng, Z. C., Wang, J. S., and Xia, C. L.: A preliminary study of
757 thermosphere and mesosphere wind observed by Fabry-Perot over Kelan, China, *J Geophys Res-Space*, 119, 4981-4997,
758 doi:10.1002/2013ja019492, 2014.

759 Yuan, W., Xu, J. Y., Ma, R. P., Wu, Q. A., Jiang, G. Y., Gao, H., Liu, X. A., and Chen, S. Z.: First observation of mesospheric
760 and thermospheric winds by a Fabry-Perot interferometer in China, *Chinese Sci Bull*, 55, 4046-4051, doi:10.1007/s11434-
761 010-4192-2, 2010.

762 Zhang, K. D., Wang, H., Liu, J., Song, H. M., and Xia, H.: The Quasi-Periodic Nighttime Traveling Ionospheric Disturbances
763 on 13 May 2024 During the Recovery Phase of a SuperStorm, *J Geophys Res-Space*, 130, ARTN e2024JA033257,
764 doi:10.1029/2024JA033257, 2025.

765 Zhu, G., Zhu, Y., Kaufmann, M., Wang, T., Liu, W., and Xu, J.: An Efficient Calibration System of Optical Interferometer for
766 Measuring Middle and Upper Atmospheric Wind, *Remote Sensing*, 15, doi:10.3390/rs15071898, 2023.