

Impact of lower atmospheric scattering on ground-based optical thermospheric wind observations with spatially uneven airglow

Xiaolong Wei^{1,2,4}, Guoying Jiang^{1,3,4,*}, Yajun Zhu^{1,3,4,*}, Jiyao Xu^{1,4}, Weijun Liu^{1,4}, Tiancai Wang^{1,2,4}, Guangyi Zhu^{1,4}, Wei Yuan^{1,4}

¹State Key Laboratory of Solar Activity and Space Weather, National Space Science Center, Chinese Academy of Sciences, Beijing, 100190, China

²College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China.

³School of Astronomy and Space Science, University of Chinese Academy of Sciences, Beijing, 100049, China.

⁴Hainan National Field Science Observation and Research Observatory for Space Weather, Hainan, 571734, China

*Correspondence to: Guoying Jiang (gyjiang@spaceweather.ac.cn), Yajun Zhu (y.zhu@swl.ac.cn)

Abstract. Scattered airglow emissions in the lower atmosphere can bias ground-based interferometer observations of thermospheric winds, particularly when airglow brightness becomes spatially uneven due to auroras. During two geomagnetic storms with visible auroras on May 10th and Oct. 10th, 2024, the Doppler Asymmetric Spatial Heterodyne (DASH) and Fabry-Perot (FP) interferometers concurrently detected atypical winds at Siziwang (SIZW, 41.83° N, 111.93° E), suspected to be caused by scattering. These atypical winds, characterized by horizontal differences exceeding 400 m·s⁻¹ between opposite cardinal directions (N-S or E-W) and downwelling exceeding 100 m·s⁻¹, showed a strong temporal association with airglow brightness. By modelling the transmission of scattered airglow emissions, we calculated post-scattering wind speeds as the initial wind speeds weighted by both scattered and direct intensities. With fixed initial speeds, the simulation reproduced the temporal characteristics of the atypical winds, demonstrating that scattering may contribute to these intense horizontal differences and downwelling. The simulation also shows that the scattering-induced biases have directional inhomogeneity with characteristics linked to the location and background line-of-sight speed of the brighter airglow region. The accuracy of the simulation is limited by the accuracy of airglow observations and atmospheric optical depth.

1 Introduction

Optical interferometers are widely utilized to observe thermospheric neutral wind (Burnside et al., 1981; Burnside and Tepley, 1989; Killeen et al., 1995; Emmert et al., 2001; Fejer et al., 2002; Emmert et al., 2006; Wu et al., 2014). Thermospheric wind can be derived from measuring the Doppler shift of OI red-line airglow emission at 630.0 nm. This emission, primarily from the collisional deactivation of O(¹D) generated by O₂⁺ dissociative recombination, peaks near 250 km altitude. The height-integrated thermospheric wind around the peak altitude can be obtained (Biondi and Feibelman, 1968; Hernandez and Roble, 1976; Burnside et al., 1981; Biondi et al., 1995; Nakajima et al., 1995). For scanning interferometers, three-dimensional wind vectors can be derived by observing the zenith and four cardinal directions at a specific elevation angle. The scanning range covers a circular area about 500 km in diameter at airglow altitude. Given thermospheric wind uniformity at this scale,

horizontal winds observed in two opposite cardinal directions (N-S or E-W) are typically similar. Averaging opposite cardinal directions improves accuracy, mitigates cloud effects, and is typically used to represent local meridional or zonal winds even during geomagnetic storms (Friedman and Herrero, 1982; Fejer et al., 2002; Sakanoi et al., 2002; Dhadly et al., 2017; Huang et al., 2018; Xu et al., 2019; Li et al., 2023; Wang et al., 2025).

However, horizontal winds in opposite cardinal directions occasionally show significant separation exceeding $100 \text{ m}\cdot\text{s}^{-1}$ and strong vertical winds, deviating from typical thermospheric wind uniformity. These observations often occur near auroras, unaffected by clouds or moonlight, and have acceptable standard errors. They mainly occur in polar regions (Crickmore et al., 1991; Price et al., 1995; Smith and Hernandez, 1995; Innis et al., 1999; Ishii et al., 2001; Guo and McEwen, 2003; Anderson et al., 2012), but have also been seen at mid-latitudes during major geomagnetic storms (Hernandez and Roble, 1976; Makela et al., 2014).

Atmospheric scattering of airglow emissions is thought to be one of the factors that biases ground-based wind observations, potentially accounting for the atypical wind. Initially, it was thought to impact airglow peak height measurements by photometers (Ashburn, 1954). Subsequent studies by Abreu et al. (1983) explored its impact on thermospheric wind speed measurements using a Fabry-Perot interferometer. Harding et al. (2017a; 2017b) later systematically modelled and estimated these effects, revealing that scattering was responsible for the anomalous vertical winds observed at mid-latitudes during geomagnetic storms by Makela et al. (2014). Light from brighter airglow regions scatters omnidirectionally in the lower atmosphere, primarily the troposphere and stratosphere, and is detectable outside its original direction. The additional Doppler shift of this scattered light can bias the retrieval of line-of-sight (LOS) speeds as well as the converted horizontal and vertical winds. Harding et al. (2017b) also investigated the impact of atmospheric scattering on interferometer wind and temperature measurements during quiet periods.

Scattering-induced biases are more pronounced during spatially uneven airglow brightness, such as during auroras (Harding et al., 2017a). Uneven airglow brightness refers specifically to inhomogeneous red-line emissions. At mid-latitudes, marked uneven red-line airglow usually comes from red aurora. Despite their distinct origins, the spectral and altitudinal overlap of airglow and aurora will let ground-based optical instruments conflate the two. For red-line observations, the aurora itself may also bias the derived winds. Aurora could elevate the red-line emission profile (Kataoka et al., 2024b), so the interferometer samples winds that are both higher and farther away. This makes the northward view sense winds deviate from the expected thermospheric wind at 250 km altitude when looking toward the aurora. Additionally, spectral contamination from precipitating energetic ions could also bias interferometers (Makela et al., 2014). They suggested that the enhanced downwelling at mid-latitudes during storms might result from the contamination of the spectral profile by fast O atoms associated with the influx of low-energy O^+ ions.

62 From a dynamical perspective, wind differences in opposite cardinal directions are considered horizontal divergence, which
63 are often associated with changes in vertical winds. Near the aurora arc, these atypical winds are mainly caused by ion drag,
64 Joule heating, and energy particle precipitation (Hays et al., 1984; Rees et al., 1984; Conde and Smith, 1995; Conde et al.,
65 2001; Anderson et al., 2012). Generally, excessive horizontal divergence and vertical wind appear alongside rapidly changing
66 auroras and exhibit a matching spatial relationship that upward (downward) winds accompanied by divergences (convergences)
67 are often detected when aurora exists equatorward (poleward) of the observatory (Ishii et al., 2001; Guo and McEwen, 2003).
68 The combination of vertical wind and horizontal divergence is related to gravity waves excited by the above processes in polar
69 regions, presenting a wave-like structure and phase delay between vertical and horizontal wind components. (Price et al., 1995;
70 Smith and Hernandez, 1995; Ishii et al., 1999; Ishii et al., 2001; Shinagawa and Oyama, 2006). At mid-latitudes, which are
71 not primary regions for magnetospheric energy injection, atypical winds are instead related to the propagation of gravity waves
72 from polar regions. (Hernandez and Roble, 1976).

73 During two geomagnetic storms on May 10th and October 10th, 2024, with visible auroras, we observed similar atypical winds
74 in ground-based interferometers at Siziwang (SIZW, 41.83° N, 111.93° E), China. These winds showed intense differences
75 over 400 m·s⁻¹ in two opposite cardinal directions for both meridional and zonal components, along with downward wind
76 exceeding 100 m·s⁻¹. The observations were unaffected by moonlight or clouds, and the interferometer retrieval errors were
77 acceptable (see Section 3.1). These atypical winds at SIZW only occurred with auroras statistically and significantly deviated
78 from the regional climatological norms over the China region (Jiang et al., 2018; Yang et al., 2020). This raises the question
79 of whether the atypical winds arise from dynamical processes, are influenced by red aurora, or stem from scattering-induced
80 biases and other measurement-related factors. Unfortunately, most of these mechanisms could amplify the wind-speed contrast
81 between opposite cardinal directions, rendering them difficult to disentangle (Harding et al., 2017a). Given the scarcity of
82 additional thermospheric-wind or auroral instruments, we remain unable to quantify every potential mechanism. Motivated by
83 the observed phenomena, this study attempts to estimate how scattering modulates the atypical winds in these storms. While
84 prior studies focus on vertical wind biases of Fabry-Perot interferometers under auroral conditions (Harding et al., 2017a;
85 Harding et al., 2017b), we will analyze the formation and patterns of horizontal differences caused by scattering. We will also
86 incorporate Doppler Asymmetric Spatial Heterodyne (DASH) interferometer data to compare scattering impact across
87 different interferometer types. As red auroras now regularly appear at the low magnetic latitudes of Japan and China during
88 elevated solar activity (Kataoka et al., 2024a; Kataoka et al., 2024b; Ma et al., 2024), a deeper understanding of scattering-
89 induced biases is essential for the proper use of interferometer data collected in these regions. In the following text, a scattering
90 radiative transfer model is used to simulate interferometer observations in two cases with visible aurora. The presence and
91 patterns of scattering-induced biases are analyzed by comparing simulations with observations.

92 2 Instruments and model

93 This study was conducted at the Siziwang station (SIZW; 41.83° N, 111.93° E, and 37.7° N MLat) of the Chinese Meridian
94 Project Phase II (Wang et al., 2024), utilizing a Dual-Channel All-sky Airglow Imager (DCAI), a Dual-Channel Optical
95 Interferometer (DCOI), and a Fabry-Perot Interferometer (FPI). DCOI derives neutral winds by observing atomic oxygen
96 green-line (557.7 nm, around 96 km) and red-line (630.0 nm, around 250 km). DCAI observes hydroxyl (around 87 km) and
97 atomic oxygen red-line nightglow, respectively. FPI only works at the red-line. Our focus is on the red-line channel. Using
98 DCAI images as one of the inputs, wind biases from optical interferometers can be simulated by a scattering radiative transfer
99 model (scattering model for short). Instruments and the model are described in the following subsections.

100 2.1 Dual-Channel All-Sky Airglow Imager

101 Dual-Channel All-Sky Airglow Imager (DCAI) comprises a fisheye lens with an approximate 170 degree field of view, a 2
102 nm narrow-band filter, and a 1024×1024 pixel, 16 bit cooled CCD. DCAI exposure time of the red-line is 2 minutes. The
103 obtained airglow images are first calibrated to the local spherical coordinate system, then sequentially corrected for stray light,
104 Van Rhijn effect, and atmospheric extinction, and finally projected onto the 250 km airglow plane. Due to DCAI not calibrating
105 the Rayleigh unit (Shiokawa et al., 2000), observed brightness is only normalized to the full-well value. And because of fish-
106 eye lens distortion and the lack of Rayleigh unit calibration, the edge brightness of the view is inaccurate. Thus, observations
107 are restricted within a 70° zenith angle. For larger zenith angles, the brightness is obtained by radial zero-order extrapolation
108 in airglow projection. Detailed image processing procedures are in Appendix B.

109 2.2 Dual-channel optical interferometer

110 Dual-channel optical interferometer (DCOI) is a scanning interferometer using Doppler Asymmetric Spatial Heterodyne
111 (DASH) technology. DASH exhibits a wider field of view, better thermal stability, simplified mechanisms, and lower tolerance
112 requirements than other interferometer structures (Englert et al., 2007; Englert et al., 2010; Harlander et al., 2017; Wei et al.,
113 2020). DCOI consists of a 630 nm narrow-band filter (2 nm bandwidth), a 9 degree field-of-view lens (f/6), a DASH
114 interferometer with a 25 mm aperture, a Neon lamp for calibration, and a 2048×2048 pixel CCD (13.5 μm per pixel) (Wei et
115 al., 2020; Zhu et al., 2023; Liu et al., 2025). Its thermal stability is maintained within 0.1 K. DCOI measures three-dimensional
116 wind speeds by scanning five directions (zenith and four cardinal directions at 45° zenith angle). Each direction is exposed for
117 5 minutes, completing a cycle roughly every 25 minutes. DCOI adopts an observation with the smallest error after evening as
118 the reference zero wind speed. The slant LOS speeds are subtracted by the time-regressed projection of vertical speed and then
119 converted to horizontal using the sine of zenith angles. It is worth noting that during auroral events, vertical winds with absolute
120 values exceeding 50 m·s⁻¹ are excluded from the regression, as they contain scattering effects that could introduce additional
121 biases to other directions. DCOI provides two series of meridional wind, two series of zonal wind, and one series of vertical
122 wind.

123 2.3 Fabry-Perot Interferometer

124 Fabry-Perot Interferometer (FPI), as a mature solution, conducts comparative observations with DCOI. It features a 630 nm
125 narrow-band filter (2 nm bandwidth), a 2.54 degree field-of-view lens (f/6), a 50 mm aperture etalon with a 7 mm gap, a
126 frequency-stabilized laser for calibration, and a 1024×1024 pixel CCD (13 μm per pixel). FPI uses the same integration time
127 and scanning method as DCOI to obtain horizontal and vertical winds for each cardinal direction and zenith. Details and
128 historical results of FPI are in these references (Yuan et al., 2010; Wu et al., 2014; Yu et al., 2014; Huang et al., 2018; Jiang
129 et al., 2018).

130 2.4 Scattering radiative transfer model

131 The model for estimating scattering impact is based on the scattering radiative transfer model and numerical solution by
132 Harding (2017a). It assumes airglow emission undergoes elastic scattering, preserving its wavelength and initial Doppler shift.
133 By specifying airglow brightness distribution, original Doppler shift, lower atmosphere scattering characteristics, and a
134 simplified geometric relationship, the radiation transfer equations (see Appendix A) can be solved to compute the distribution
135 of multiple scattered light and its associated Doppler shift. This enables the wind simulation with atmospheric scattering. A
136 schematic diagram (Fig. 1) illustrates the basic mechanism. To enhance applicability, we have refined several aspects: (1) The
137 upper boundary of the lower atmosphere is set at 40 km to improve the accuracy of the effective extinction path in the initial
138 source function. (2) The Doppler shift is replaced by LOS speed, with every incident ray from the airglow layer mapped
139 directly to its corresponding LOS speed. (3) After slicing the airglow layer into several bins by LOS-speed, the model
140 illuminates one bin per run, records its scattered intensity, then merges all bins with an intensity-weighted average to yield the
141 post-scattered LOS speed. The detailed model description is provided in Appendix A.

142 Additionally, the scattering characteristics of the lower atmosphere in our model, including the scattering phase function and
143 optical depth, were derived from Aerosol Robotic Network (AERONET) observations (Holben et al., 2001). We utilized data
144 from the Baotou site (40.9° N, 109.6° E), which is the nearest available site to SIZW, located approximately 180 km away.
145 The total optical depth, accounting for both aerosol and molecular scattering, was calculated using monthly averages and was
146 found to be 0.43 in May and 0.2 in October. The scattering phase function was determined based on AERONET data following
147 the previous method (Harding et al., 2017a). Further details regarding the scattering characteristics are described in Appendix
148 B.




Figure 1: The schematic diagram of the scattering radiative transfer model


The grey shading represents the lower atmospheric layer, with darker hues indicating greater optical depth. The yellow-green fillers represent the relative brightness from the red-line airglow layer. Yellow indicates higher light intensity. The blue-red fillers, which correspond to the relative brightness, represent the Doppler shift type (blue-shift or red-shift) of LOS wind speeds. (a) to (e) represent airglow emissions travelling along different paths, carrying Doppler shifts from outside the line of sight into the interferometer, thereby causing biases in the observations. The model estimates the biases by simulating the distribution of airglow emissions after scattering.

3 Results

Two storms with visible auroras on May 10th and Oct. 10th, 2024, respectively, are used to study the scattering impact. The storm from May 10th to 11th is characterized by its significant magnitude and prolonged duration. Multiple works report this event (Guo et al., 2024; Hajra et al., 2024; Themens et al., 2024), with particular focus on the variations of thermospheric winds (Wang et al., 2025; Zhang et al., 2025) and auroras (Gonzalez-Esparza et al., 2024; Kataoka et al., 2024b; Mikhalev, 2024; Nanjo and Shiokawa, 2024) at mid-latitudes. The storm commenced around 17:00 UT on May 10th and the main phase persisted until 02:00 UT on May 11th. After that, the local night of May 11th in the China region sank into a continuous recovery phase. Another storm from Oct. 10th to 11th is weaker than May's (Ranjan and Pallamraju, 2025; Singh et al., 2025), with the main phase from 18:00 UT on Oct. 10th to 02:00 UT on Oct. 11th. During the two geomagnetic storms with visible auroras, both the DCOI and FPI at SIZW observed atypical winds, characterized by intense horizontal wind differences and downward vertical winds.

167 **3.1 Storm-time wind speed statistics**

168 It is necessary to ascertain whether atypical winds originate from atmospheric scattering with spatially uneven airglow
169 brightness or dynamic processes during storms. To investigate the impact of visible auroras on atypical winds during storms,
170 we made the most of the available observations, tracking DCOI's storm-time observations for nearly a year and FPI's for almost
171 five months. We employed the planetary magnetic index Kp exceeding 3 to identify geomagnetic storms (Yang et al., 2020).
172 Besides, to rule out moonlight and cloud effects, we only used clear sky conditions, which means: (1) excluding cases where
173 the angle between the moon and the line of sight is less than 30 degrees, and (2) excluding cases where large-area thick cloud
174 coverage is visible in DCAI. Additionally, data with standard errors greater than 50 m·s⁻¹ were also excluded. A few aurora
175 events, including Nov. 5th, Dec. 1st, 2023, and Aug. 12th, 2024, that did not meet this criterion were excluded.



176
177 **Figure 2: Storm-time (Kp>3) thermospheric wind speed statistics at SIZW**

178 Figure 2a shows meridional winds observed along two opposite directions (N-S) by FPI, with north-looking in red and south-looking in
179 blue. Figure 2b shows zonal and vertical wind, with east-looking in yellow, west-looking in green, and zenith-ward in black. The
180 northward, eastward, and upward speeds are positive in coordinates. Observations without aurora are shown as points, while those with
181 visible auroras are shown as lines. Figures 2c and 2d are similar but show DCOI data.

182 Figure 2 shows thermospheric wind statistics during geomagnetic storms (Kp>3) at SIZW. The first two panels display FPI
183 data from May to Oct. 2024, while the rest display DCOI data from Nov. 2023 to Oct. 2024. FPI began operation on May 8th,
184 2024, with about half a year less data than DCOI. Observations with no aurora in the field of view are marked as points, while

the two cases with visible auroras are shown as lines. The five observation directions of the interferometer are marked by different colors. During typical storms, horizontal winds consistently increase to around $150 \text{ m}\cdot\text{s}^{-1}$ both equatorward and westward with no significant downward wind. However, under visible auroras, both DCOI and FPI have detected large wind speeds, such as a southward wind of about $600 \text{ m}\cdot\text{s}^{-1}$ and downwelling exceeding $100 \text{ m}\cdot\text{s}^{-1}$. The two series of winds observed along opposite cardinal directions (N-S or E-W) exhibit overt differences, with values exceeding $400 \text{ m}\cdot\text{s}^{-1}$ and contrary directions. This is markedly different from the wind patterns observed during non-aurora storms, where opposite-direction winds do not show significant divergence. Comparing the results of DCOI and FPI, the observations are largely consistent both with and without auroras. The atypical winds observed simultaneously by two interferometers with different principles suggest a systematic error from outside the instruments. Besides, these simultaneous changes appear in five observation directions, all characterized by enhanced negative LOS speeds, indicating likely LOS speed contamination. These factors point more towards scattering impact rather than dynamical processes as the cause. Next, the relationship between scattered light and atypical winds will be investigated through simulation.

3.2 Comparison of observations and simulations

Figure 3 shows the red-line airglow brightness from DCAI (Fig. 3a, 3e), the observed winds from DCOI and FPI (solid lines with different markers in Fig. 3b-3d, 3f-3h) and the simulated winds from the scattering model (dotted lines in Fig. 3b-3d, 3f-3h) during the two nights of May 10th and 11th, 2024, at SIZW, in which the different colors denote distinct directions. The grey lines in the horizontal wind plots represent the average values between opposite cardinal directions. The multi-directional brightness series from DCAI are extracted at 45° zenith angle, consistent with the scanning zenith angle of interferometers. Time intervals with visible auroras are highlighted in red, showing much higher brightness in northward directions than others. Figure 4 supplements the auroral distribution compared to Fig. 3a and 3e. Images from DCAI are projected onto the airglow layer at 250 km. The red circle encloses the actual observations with zenith angles less than 70° , while the values outside are extrapolated. The red dots represent the interferometer's pierce points on the airglow layer at 45° zenith angle.




Figure 3: Observations of aurora and wind speeds, and the scattering model simulation on the nights of May 10th and 11th, 2024, at SIZW

Figure 3a shows the brightness of 8 cardinal directions, all at 45° zenith angle, along with the zenith-ward, extracted from DCAI. The color coding is as follows: red for northern directions, green for east and west, blue for southern directions, black for the zenith, and yellow for the average brightness excluding the three northern directions. Figure 3b shows the meridional wind, with north-looking in red and south-looking in blue, and the average of the two directions in grey. DCOI observations are shown as solid lines with circular dots, FPI as solid lines with rhombus dots, and simulations as dotted lines. Figures 3c, 3d are similar to Fig. 3b, but for zonal and vertical wind, with east-looking in yellow, west-looking in green, and zenith-ward in black. For a more concise figure, if the standard error exceeds 100 m s⁻¹, the point will be filled with black instead of error bar. Figures 3a-3d show data from May 10th, and Fig. 3e-3h from May 11th. The time intervals with visible auroras are marked in red.




Figure 4: Auroral distribution observed by DCAI on the nights of May 10th and 11th, 2024, at SIZW

Images from DCAI (Fig. 4a-4d, 4f-4i) have been projected onto the airglow layer at 250 km. The red circle encloses actual observations with zenith angles $< 70^\circ$, while values outside are extrapolated using zero-order extrapolation. The red dots represent the interferometer's pierce points on the airglow layer at 45° zenith angle. Figures 4e, 4j are similar to Fig. 3a, 3e, with the corresponding images time labelled.

The coastlines in projected DCAI images are made with Natural Earth.


After 18:30 UT on May 10th, as the aurora intensified, both DCOI and FPI detected simultaneous changes in meridional, zonal, and vertical winds. The north-looking red-line brightness at 45° zenith angle exceeded three times that of other directions. The meridional and zonal wind differences between opposite cardinal directions (N-S or E-W) increased. And the winds detected in opposite directions reversed. The maximum meridional difference was close to $800 \text{ m}\cdot\text{s}^{-1}$, while that in zonal exceeded $500 \text{ m}\cdot\text{s}^{-1}$. The downward wind was enhanced by over $100 \text{ m}\cdot\text{s}^{-1}$. These four aforementioned variables that red-line brightness, the meridional and zonal differences, and downwelling, increased almost simultaneously, peaked at 19:05 UT, and then decayed.

Moreover, the average meridional wind, derived from averaging opposite cardinal directions, continuously enhanced equatorward to around 400 m/s, while the average zonal wind enhanced westward to around 100 m/s. Unlike the single-direction results that peaked at 19:05 UT, the average wind varied steadily, consistent with storm-time circulation. Compared to the average wind, the separated horizontal and enhanced vertical winds are atypical. Even with travelling atmospheric disturbances (TADs) superimposed on storm-time circulation, phase lags between horizontal and vertical components would be expected (Hernandez and Roble, 1976; Ishii et al., 1999), but none were observed. Thus, the atypical winds do not appear to be the result of a dynamical process. During the recovery phase on May 11th, the aurora was present throughout the night but much weaker than on May 10th, as seen in Fig. 4. Both DCOI and FPI showed westward and equatorward horizontal winds with no significant downward wind. There was a meridional difference of about 100 to 300 m·s⁻¹ persisted throughout the night, with no zonal difference.

Subsequently, we used the scattering model to explore the relationship between red-line brightness variations and atypical winds through atmospheric scattering. On May 10th, a fixed wind vector of 100 m·s⁻¹ westward and 400 m·s⁻¹ southward was set as the input. This assumed wind was kept constant over time and spatially uniform, with no vertical components. On May 11th, a fixed wind vector of 200 m·s⁻¹ westward and 100 m·s⁻¹ southward was used, again with no vertical component. These values are chosen based on average observed wind speeds to approximate storm-time circulation. Although the specific values may deviate, the main wind directions remain consistent. The storm-time enhancement of vertical winds may be caused by scattering rather than representing real winds, so we set it to zero. It is worth noting that we neglect the variation of background wind in the model inputs, due to uncertainty regarding whether the observed wind variations are biased. Moreover, using fixed wind speeds allows us to highlight the impact of red-line brightness variations and determine the presence of scattering effects.

As shown by the dotted lines in Fig. 3b-3e, the simulations with scattering impact generally match the observations on May 10th. Simulated horizontal wind differences and downwelling increase initially with the aurora, peak around 19:15 UT, and then decay. The simulated wind speed variations lag the observations by about 10 minutes. The lag may be due to the relatively rough 25 minute scanning cycle or DCAI underestimating airglow brightness at the field of view's edge, leading to inaccurate capture of scattering enhancement start time. Numerically, the simulated zonal and vertical winds match observations more closely than the meridional wind. The simulated meridional difference is smaller than the observed difference, and the north-looking simulation remains closer to the default value, unlike the equatorward-biased observation. The preset fixed wind may influence the meridional simulation, as it does not follow the equatorward enhancement of the meridional wind. Other factors beyond scattering impact might also have an impact, such as the lift of red-line emission profile and the spectral pollution caused by auroras (Makela et al., 2014), to be discussed later. Additionally, the model simulates a similar intense downward wind as observed under the preset zero vertical wind. This indicates that the vertical wind is significantly affected by scattering. This is why the intense vertical wind is not subtracted when converting interferometer LOS speed to horizontal wind, to prevent error propagation. For May 11th, due to the weak but continuous aurora, the simulation shows weak horizontal differences and

262 slight downward winds throughout the night. Compared with the observations, the simulation shows smaller meridional
 263 differences. It also indicates zonal differences and downward winds, which are not evident in the observations. The poorer
 264 simulation on May 11th may be due to misalignment between dominant horizontal winds and airglow brightness gradients,
 265 which will be discussed later. Additionally, there may be issues with the zero wind calibration. When auroras are present
 266 throughout the night, the vertical wind, which includes scattering biases, may have been used to calibrate zero wind speed. It
 267 likely masks the scattering impact in the observations and explains the discrepancies in the simulation.



268
 269 **Figure 5: Observations of aurora and wind speeds, and the scattering model simulation on the nights of Oct. 10th and 11th, 2024, at**
 270

271 **SIZW**

Figure 5 is analogous to Fig. 3, but for Oct. 10th and 11th.




Figure 6: Auroral distribution observed by DCAI on the nights of Oct. 10th and 11th, 2024, at SIZW

Figure 6 is analogous to Fig. 4, but for Oct. 10th and 11th.

272

273

274

275 Figures 5 and 6 show another case from Oct. 10th to 11th, analogous to Figures 3 and 4. On Oct. 10th, the aurora appeared at
 276 17:15 UT, expanded southward and increased in brightness, peaking first at 18:30 UT before decaying and then increasing
 277 again from 20:30 UT until sunrise. The second peak was brighter than the first (Fig. 6). Similar to the storm in May, once
 278 aurora appeared, both DCOI and FPI observed atypical winds, with synchronous meridional and zonal differences and
 279 downward enhancements in vertical wind. These atypical winds also exhibited two peaks, around 18:30 UT and 21:00 UT.
 280 The horizontal winds observed in opposite cardinal directions were basically in opposition. During visible aurora periods,
 281 DCOI and FPI showed a 50 to 100 m·s⁻¹ difference in vertical wind but consistent variation trends. Moreover, the average
 282 horizontal wind between opposite cardinal directions was dominantly equatorward and westward, which also had two peaks.
 283 On Oct. 11th, the storm had passed, and no visible aurora was present. The increase in brightness around 13:00 UT was due to
 284 moonset in the southwest. There were no significant horizontal differences or downward winds, consistent with geomagnetic
 285 quiet conditions.

286 We set a fixed $100 \text{ m}\cdot\text{s}^{-1}$ westward with $400 \text{ m}\cdot\text{s}^{-1}$ southward wind vector for Oct. 10th, and $100 \text{ m}\cdot\text{s}^{-1}$ westward with $200 \text{ m}\cdot\text{s}^{-1}$
287 southward wind vector for Oct. 11th in the model, respectively, with no vertical component. The simulation on Oct. 10th exhibits
288 two peaks. The second peak in the simulation is consistent with the observations better, while the first peak, although capturing
289 the trend, is significantly underestimated in magnitude. This discrepancy in the simulation may relate to optical depth, aurora
290 brightness, and background wind changes. The optical depth in Oct. is about half that of May, and simulations underestimate
291 observed values. As in previous studies (Harding et al., 2017b), optical depth can affect the scattering model response. On Oct.
292 10th, the first aurora brightening is weaker than the second. When red-line brightness differences are small, the model response
293 tends to be lower. The impact of optical depth and red-line brightness on the model will be discussed later. Additionally,
294 noticeable fluctuations in the average meridional wind on this day may also contribute to the deviation in the model with fixed
295 initial wind. The north-looking wind speed varied dramatically between 19:00 UT and 21:00 UT along with the aurora, which
296 may also be related to spectral contamination beyond scattering impact. This spectral contamination arises from fast O atoms
297 generated by low-energy O^+ ion precipitation in the auroral region, which occurs at higher altitudes. This issue introduces an
298 additional spectral shift that compromises wind retrieval (Makela et al., 2014). This exceeds the simulation range of the model,
299 thereby causing the discrepancy.


300 4 Discussions

301 In this study, we modelled the transmission of scattered airglow emission in the lower atmosphere. Post-scattering wind speeds
302 were calculated based on assumed initial wind speeds weighted by both scattered and direct intensities. The model basically
303 captured the temporal variations of horizontal wind differences and downward enhancements associated with varying auroral
304 brightness, suggesting the contribution of scattering mechanisms to atypical winds. However, the simulation of scattering has
305 certain limitations and characteristics: (1) The differences between simulation and observation vary across different directions.
306 (2) The simulated values sometimes exhibit significant numerical deviations from observations. Could this be related to model
307 errors? Next, we will discuss the working principle of the scattering model and the errors involved in the simulation.

308 4.1 Core working principle of the scattering model

309 As shown in Fig. 3 and Fig. 5, observed winds respond differently to scattering across directions, especially on May 10th and
310 Oct. 10th with stronger auroras. Although the scattering model has numerical errors, the simulations also show directional
311 differences in scattering-induced biases. Both observations and simulations indicate that the meridional wind responds the
312 most, followed by the zonal wind, while the vertical wind responds the least. Since vertical and horizontal wind speeds are
313 derived from the projection of LOS wind speeds, this essentially reflects the non-uniform response of LOS speeds to scattering.
314 This directional inhomogeneity of scattering impact aligns with previous studies. Harding et al. (2017a) simulated scattering
315 effects under auroral conditions, using northward observations as the initial winds. They noted that this direction experiences
316 minimal scattering contamination due to facing the brighter region. Abreu et al. (1983) used a meridional one-dimensional

317 model, finding that LOS wind speeds near intense airglow brightness gradients and with weaker airglow intensity are more
 318 susceptible to contamination. They also showed that scattering-induced biases are minimal in the vertical direction, as the
 319 shorter atmospheric path length limits the opportunity for scattering. In this study, we further explore the scattering impact as
 320 a function of azimuth angle, revealing the formation of horizontal differences.



321
 322 **Figure 7: Post-scattering LOS speeds at 45° zenith angle**

323 The figure shows the post-scattering line-of-sight speeds at 45° zenith angle for various directions over time, with panels for May 10th,
 324 May 11th, and October 10th, respectively.

325 Figure 7 shows post-scattering LOS speeds at 45° zenith angle in the simulations on three aurora nights, with the vertical axis
 326 indicating cardinal directions derived from azimuth angles. Concerning the auroral variations in Fig. 4 and Fig. 8, LOS speeds
 327 show diffusion during auroral events. Negative LOS speeds initially concentrated northward, spread westward and eastward,
 328 expanding horizontal coverage. Positive LOS speeds initially in the southward direction shrink. When converted to horizontal
 329 wind speeds, these changes lead to increased horizontal differences, or the false convergence caused by scattering, in other
 330 words. The northward LOS speed changes slightly, the southward speed changes the most and nearly reverses, while the
 331 eastward and westward speeds are intermediate.




Figure 8: Proportion of scattered light at 45° zenith angle

The proportion is the ratio of scattered light intensity to the total light intensity (both scattered and direct) in the simulation. The figure shows this proportion at 45° zenith angle for various directions over time, with panels for May 10th, May 11th, and October 10th, respectively.

Figure 8 shows the ratio of scattered light intensity to total light intensity at a fixed 45° zenith angle calculated by the scattering model. Consistent with the schematic diagram (Fig. 1), the scattered intensity is the sum of all injected directions, and the total light intensity includes the direct component on this basis. Without auroras, the scattering proportion is typically below 0.4 and varies with atmospheric scattering capability. However, during auroral events, the scattering proportion in some directions can increase to 0.5 or higher. The northward scattering proportion increases the least and remains much lower than in other directions. In contrast, the scattering proportion is significantly enhanced in directions ranging from 135° to 180° away from northward. The aurora appears in the north, resulting in much higher northern brightness. After atmospheric scattering, light from the north diffuses into surrounding directions, increasing the scattering proportion. Because the north itself has strong direct light, its scattering proportion remains small. In the model, we assume that stronger light rays dominate interference fringe identification (Wei et al., 2020), thereby determining the Doppler shift or LOS speeds. The lower scattering proportion in the north allows northward observations to retain more LOS speeds from themselves, while other directions experience

greater LOS speed contamination from the north. Therefore, the northward simulation is closest to the default inputs, the southward simulation deviates the most, and the eastward and westward simulations lie in between. Additionally, the scattering impact should also consider the initial LOS speeds in the brighter region. Despite the high scattering proportion on May 11th shown in Fig. 7, the simulated LOS speed changes in Fig. 8 are minimal. This is because of the smaller LOS speed in the auroral region on that day, resulting in less contamination spread to other directions.

The core working principle of the scattering model relies on the relationship between airglow brightness and background LOS wind speeds. As Harding (2017a) noted, scattering requires a bright sky region with large LOS wind speeds. Firstly, spatially uneven airglow brightness is a prerequisite for significant scattering. The brightest airglow area contributes most to scattered light intensity, and its Doppler shifts determine the LOS biases in other directions. This principle allows a rough assessment of scattering impact without model computation when airglow is uneven. Locate the brightest region and its Doppler shift type, as the same Doppler shift type will likely appear in other directions. Blue-shift dominance indicates increased downward wind and horizontal deviations opposite to the line of sight, resembling convergence, while red-shift dominance resembles divergence. Minimal scattering-induced biases occur if the scattering proportion is very low due to uniform airglow brightness, or if the LOS velocity in the bright region is near zero (i.e., the wind speed is perpendicular to the line of sight). Previous observations can be directly verified by this principle and are basically in line with it (Hernandez and Roble, 1976; Price et al., 1995; Ishii et al., 1999; Ishii et al., 2001; Makela et al., 2014). Unfortunately, scattering impact can complicate dynamic analysis. In polar regions, auroras are characterized by green-line emissions, and thermospheric winds are significantly influenced by ion drag, where scattering effects may not be prominent. In contrast, mid-latitudes have mainly red-line auroras with large-scale uniform circulation, making the scattering impact more pronounced and distinguishable.

4.2 Errors of the scattering model

The model also exhibits certain errors and limitations. Scattering-induced biases in observations have nearly similar magnitudes on May 10th and Oct. 10th. However, with the same 100 m·s⁻¹ westward and 400 m·s⁻¹ southward wind input, the scattering-induced biases in the May case are significantly larger in magnitude and closer to reality compared to October. In Fig. 7, the simulated LOS speeds show a larger diffusion range for the May 10th case compared to October. In Fig. 8, the scattering proportion for May 10th is consistently higher than that for October. We attribute this difference mainly to the distinct optical depths in the two months, which are 0.43 and 0.2, respectively. Optical depth reflects atmospheric extinction capability and is mainly related to aerosol content (see Appendix B). It primarily affects the extinction process and influences the magnitude of scattering-induced biases by altering the proportion of scattered light. When the optical depth is artificially increased to a higher value, such as around 0.6, the model more closely matches the Oct. observations. We find that the model underestimates scattering effects when the optical depth is low.

378 In the October 10th event, the simulated scattering-induced biases are inconsistent between the two auroral brightness peaks.
379 In Fig. 7, the LOS speed variation is larger during the second peak, and in Fig. 8, the scattering proportion is greater. This is
380 because the scattering proportion is susceptible to errors in scattered and direct light intensities. DCAI does not correct for
381 Rayleigh units, leading to significant errors in regions with large zenith angles. The model cannot fully eliminate the stray
382 light caused by the glass dome when separating the initial direct and scattered light from DCAI images (Harding et al., 2017a),
383 resulting in errors. In our experiments, if the stray light effect is not subtracted as described in Appendix B, the model becomes
384 more inert, resulting in a smaller simulated scattering proportion.

385 We also considered the potential influence of thermospheric temperature. FPI data show uniformly elevated thermospheric
386 temperatures in these two storms, with the northward view occasionally about 300 K warmer than the others (not shown here).
387 Because our scattering model does not yet include temperature effects, we cannot quantify how much scattering biases the FPI
388 temperature measurements. In the study of Harding et al. (2017b), wind simulations are temperature-independent, while
389 temperature retrieval relies on the wind. Likewise, we substitute the LOS speed for the Doppler shift and ignore temperature-
390 induced spectral broadening. In principle, thermospheric temperature influences retrieval uncertainty, not the wind speed itself.
391 We remain cautious that ignoring this uncertainty could introduce extra bias if a horizontal temperature gradient is present, but
392 incorporating it would markedly raise the computational cost and remains a task for the future.

393 In our experiments, even after several refinements aimed at mitigating the underestimation, the scattering model still
394 underestimates the observed scattering impact: (1) A single-scattering albedo of 1 was used, ignoring absorption. (2) Stray
395 light effects were removed. (3) Attenuated airglow observations at the edge of DCAI images were extrapolated, enhancing
396 edge scattering. (4) Excessive edge extinction was reduced by correcting the extinction path geometry, increasing the scattered
397 light intensity integral. (5) Zero vertical wind was assumed when converting simulated LOS speeds to geographical wind
398 speeds. We suspect the underestimation arises from how the model integrates the optical depth. Since the integral only includes
399 10 optical depth layers, with each light ray scattering once per layer and extinguishing once between layers, it may be too
400 crude compared to the real path, underestimating the scattered light. Simply increasing the number of optical depth layers is
401 not effective. We think this may be related to the non-linear variation of atmospheric density with altitude, where optical depth
402 may not vary linearly with height, and the scattering phase function may also change with altitude. To address these issues,
403 future work should complete the DCAI correction. Additionally, introducing a model of optical depth varying with altitude
404 can increase the number of single-scattering nodes and ensure the geometric accuracy of the extinction path, thereby improving
405 the accuracy of scattered light intensity calculations.

406 Furthermore, these bright region observations do not necessarily reflect the usual 250 km thermospheric wind. In Fig. 3 and
407 Fig. 5, the north-looking wind observations show unusually high wind speeds, which are significantly different from the
408 simulations. In particular, on October 10th, the north-looking wind speed varied dramatically with the intensity of the northern
409 aurora. During the two auroral peaks, the north-looking wind direction also reversed. This indicates that the interferometer

410 receives an additional effect when it looks toward the aurora. Kataoka et al. showed that red aurora lifted the red-line emission
411 profile, raising its peak above 300 km and brightening the upper part on May 11th (Kataoka et al., 2024b). Consequently, the
412 interferometer can sample winds that are higher and more poleward. Because storm-time surges propagate from the polar
413 region to the equator, these higher, poleward regions are likely to carry stronger equatorward winds. The interferometer may
414 record a larger wind speed toward the aurora. Additionally, spectral contamination from precipitating energetic ions can also
415 bias interferometers (Makela et al., 2014). In other words, the interferometer is partly sensing the speed of non-neutral species,
416 boosting the observed wind. These issues lie beyond what scattering models can reproduce. From the observed pattern, we
417 infer the presence of non-scattering effects, especially in the poleward view. Due to the absence of nearby higher-latitude
418 neutral-wind observations relative to SIZW, quantifying their respective contributions remains challenging.

419 **5 Conclusions**

420 This study has further proved that lower atmospheric scattering can contribute to biases in thermospheric wind observation on
421 ground-based optical interferometers. The light scattered from the non-line-of-sight directions of the scope will lead to
422 additional LOS speeds and appear as atypical horizontal differences and vertical wind at geographic coordinates. With a
423 simplified scattering radiative transfer model, we simulate the distribution of airglow intensity after the multiple scattering of
424 the lower atmosphere and estimate the wind observation bias under scattering impact via a weighted average method. Atypical
425 winds under conditions of spatially uneven airglow have been generally captured.

426 We have refined the scattering model in previous research to enhance its computational efficiency. Specifically, we simplified
427 the LOS wind speed simulation and introduced an upper limit for the lower atmosphere to improve the accuracy of the
428 extinction length calculation. The scattering impact can be directly estimated through the relationship between the bright
429 airglow region and the LOS wind speed. The brightest airglow region contributes most to the scattering impact, of which the
430 Doppler shift type determines the LOS biases in other directions. Although the observed winds are affected by scattering when
431 airglow is uneven, they still retain dynamic information, such as the average wind being close to the storm-time circulation.
432 Unfortunately, we lack alternative observational methods to verify the accuracy of the interferometer results. It deserves further
433 study to the extent of scattering impact with more cases and additional instrumental observations.

434 Limited by the accuracy of the model inputs, the scattering model can only simulate the wind features associated with scattering
435 impact under clear sky conditions. It remains incapable of precisely picking out the speed alterations induced by scattering
436 impact. Given that scattering impacts observations during geomagnetic storms accompanied by auroras, it is necessary to
437 quantify these biases to provide accurate data for dynamic studies. Therefore, future efforts can focus on refining the model
438 and its inputs or statistically analyzing scattering biases under different airglow brightness distributions from various stations
439 to gain a clearer understanding of the magnitude of scattering biases.

440 Appendix A

441 In the following appendices, we provide a concise description of the scattering model's operational principles, inputs, and
 442 modifications employed in our works. For more detailed solutions, please refer to the article by Harding et al. (Harding et al.,
 443 2017a).

444 Based on the radiation transfer theory, Hansen and Travis (1974) and Sobolev (1975) gave the multiple scattering solution.
 445 Harding (2017a) extended the initial source function $J_0(\tau, u, \phi)$ to airglow surface source, and corrected the missing
 446 normalization factor in the scattering phase function (Eq. (1) to Eq. (3)):

$$447 \quad u \frac{dI(\tau, u, \phi)}{d\tau} = -I(\tau, u, \phi) + J(\tau, u, \phi) \quad (1)$$

$$448 \quad J(\tau, u, \phi) = \frac{\omega}{4\pi} \int_0^{2\pi} \int_{-1}^1 P(u, u', \phi, \phi') I(\tau, u', \phi') du' d\phi' + J_0(\tau, u, \phi) \quad (2)$$

$$449 \quad J_0(\tau, u, \phi) = \frac{\omega}{4\pi} \int_0^{2\pi} \int_0^1 P(u, u', \phi, \phi') \sec(\gamma') f(u', \phi') \exp[-\tau L(u')] du' d\phi' \quad (3)$$

$$450 \quad L(u) = [(R_e + H_L) \cos(\gamma_\tau) - R_e u] H_L^{-1} \quad (4)$$

$$451 \quad u = \cos(\zeta) \quad (5)$$

452 The equations are formulated within an improved local spherical coordinate system, including azimuth ϕ , zenith angle ζ
 453 which is represented in cosine form u , and vertical height which is expressed as optical depth τ .

454 In the case of scattering, as illustrated in Fig. 1, the light intensity along a line of sight, represented by $I(\tau, u, \phi)$, consists of
 455 two parts, the direct light (a in Fig. 1) from the same direction, and the aggregate of scattered light (b-e in Fig. 1) from other
 456 directions, which is represented by the source function $J(\tau, u, \phi)$. Based on the radiative transfer equation (Eq. (1)), at each
 457 optical depth layer, the scattered light intensity received from all directions will be integrated. Simultaneously, the original
 458 intensity in the line of sight will be added to the total scattered light. Besides, the extinction in the path will be calculated
 459 according to the optical depth.

460 There are two potential scattering paths in the lower atmosphere: single scattering (b, c in Fig. 1) and multiple scattering (d, e
 461 in Fig. 1). The model computes them sequentially via an iterative process. In the initial state, there is no light intensity in the
 462 lower atmosphere. Therefore, the single scattering will originate solely from the airglow layer, and the source function
 463 $J(\tau, u, \phi)$ will be equivalent to the initial source function $J_0(\tau, u, \phi)$ at this state. By solving Eq. (1), the model can obtain
 464 the single scattered intensity in each direction at every optical depth layer, which is the updated source function $J(\tau, u, \phi)$.
 465 Then, the multiple scattering can be calculated based on it. Typically, the total scattered intensity will remain relatively constant
 466 when accounting for the fourth scattering. By using this iteration, the scattered light and residual direct light in DCAI images
 467 can be effectively separated. The residual direct light will subsequently serve as the background intensity distribution for
 468 simulating speeds.

469 In the source function $J(\tau, u, \phi)$, the scattering phase function $P(u, u', \phi, \phi')$ quantifies the relative gain of an incident angle
 470 to an exit angle during the scattering process. The reference value is based on a unit-radius sphere, which necessitates the
 471 introduction of a factor $\frac{1}{4\pi}$. Furthermore, ω represents the single-scattering albedo, set as 1. The initial source function
 472 $J_0(\tau, u, \phi)$ is responsible for introducing the airglow distribution $f(u', \phi')$. Here, $\sec(\gamma')$ represents the secant of the zenith
 473 angle at the puncture point of the airglow layer, which helps eliminate the Van Rhijn effect. Additionally, the exponential term
 474 with base e is utilized to calculate the equivalent extinction length along an inclined path.

475 It is primarily the lower atmosphere that significantly scatters and absorbs light (He et al., 2021; Li et al., 2022). Therefore,
 476 when computing the extinction length, just employing the cosine of zenith angle u' will lead to an overestimation of the
 477 effective length, as illustrated on the right side of Fig. 1. To address it, we set an upper boundary H_L of the lower atmosphere
 478 at 40 km, assuming an optical depth of zero above this altitude. Using geometric relationships, an equivalent length factor
 479 $L(u)$ can be derived, where R_e means Earth radius, and γ_t represents the zenith angle at the penetration point of 40 km
 480 height. This value can be readily calculated by adjusting the target height of the formula for γ . Inside the lower atmosphere,
 481 we apply a thin-layer approximation, which also utilizes the geometric relationships at the upper boundary.

$$482 \quad V = \frac{V_{dr}I_{dr} + \sum_k V_{sc}(k)I_{sc}(k)}{I_{dr} + \sum_k I_{sc}(k)} \quad (6)$$

483 After working out the background intensity distribution, we partition the LOS speeds at 250 km into several bins. To simplify
 484 simulation, the model directly uses LOS speeds corresponding to Doppler shifts. As roughly illustrated in Fig. 1, all LOS
 485 speeds are categorized into $k=10$ bins valued from highest to lowest, assuming that the LOS speeds within each bin
 486 approximate their mean value V_{sc} . The scattered intensity distribution I_{sc} is computed by extracting the airglow brightness
 487 from the corresponding region of each bin. And, there will be no intensity from other areas during a single bin's computing.
 488 According to Eq. (6), the simulative LOS speed at a specific angle will be an averaged result, where the original speed V_{dr} is
 489 weighted by the direct light intensity I_{dr} , and the additional speed resulting from scattering V_{sc} is weighted by the scattered
 490 light intensity I_{sc} . The model directly uses the average within DCOI's 9 degree field of view as simulated post-scattering LOS
 491 speed of interferometers, since DCOI and FPI have not measured the reception gain of light outside their fields of view. We
 492 find that due to the coarse model grid, the 9 degree average is nearly the same as using the nearest single-sight observation.
 493 Finally, the LOS speed will be converted to horizontal wind, maintaining the assumption of zero vertical wind to prevent the
 494 propagation of scattering biases in the vertical direction.

495 Appendix B

496 This appendix details the scattering model's inputs from measurements, supplementing the second section. The background
 497 airglow brightness for the model comes from DCAI. Image processing includes: (1) dark field exclusion, (2) median filtering

498 to remove starlight, (3) conversion to the local spherical coordinate, (4) stray light correction, and (5) radial zero-order
 499 extrapolation for regions beyond 70° zenith angle. Stray light results from the scattering of strong incident light by the glass
 500 dome. During quiet nights without auroras, it is weak and uniformly distributed across all LOS directions. We use the azimuthal
 501 average of the nearest quiet night at 45° zenith angle as a reference for weak stray light conditions. After the aurora onset,
 502 stray light brightens all directions. The difference between the darkest direction at 45° zenith angle and the reference value is
 503 considered the additional stray light caused by the aurora and is subtracted from the entire image. The shown airglow images
 504 additionally mitigate the Van Rhijn effect through $\sec(\gamma')$ and the atmospheric extinction through $\exp[-\tau L(u')]$ (see
 505 Appendix A). Since the scattering model already includes these processes, no separate treatment is needed. The optical depth
 506 and scattering phase function inputs are shown in Fig. 9. Optical depth is calculated using AERONET's monthly averages,
 507 with interpolation at 630 nm. Since only daytime observations are available, local daytime values are used to represent
 508 nighttime values. Fig. 9a and 9b show monthly average optical depths at local daytime, with total averages of 0.43 and 0.2.
 509 The scattering phase function is a weighted average of molecular and aerosol scattering phase functions from AERONET at
 510 675 nm, which is weighted on the total optical depth of aerosols and molecules.




Figure 9: Optical depth and scattering phase function in May and October

The first two panels show daytime monthly average optical depths, with molecular optical depth amplified for clarity. The rest panels
 show the scattering phase functions.

515 **Code availability**

516 The code of DCAI images correction, the scattering model, and the visualization are not publicly available yet. If needed, they
517 can be obtained by contacting the corresponding authors via email.

518 **Data availability**

519 The data of DCOI and DCAI from the Chinese Meridian Project can be obtained from <https://www.meridianproject.ac.cn/en/>.
520 The FPI data can be obtained by contacting the corresponding authors via email. The data of AERONET can be obtained from
521 <https://aeronet.gsfc.nasa.gov/>. The Kp index provided by GFZ, German Research Centre for Geosciences, can be obtained
522 from <https://kp.gfz-potsdam.de/en/>.

523 **Author contributions**

524 Conceptualization: XW; investigation: XW, GJ; instruments construction and maintenance: YZ, JX, WL, TW, GZ, WY; wind
525 data retrieval: YZ, WL, GZ, TW; airglow images correction: XW; model improvement and programming: XW; visualization:
526 XW; validation: GJ, YZ, JX, WL, TW; writing - original draft preparation: XW; writing-review and editing: GJ, YZ, JX, WL,
527 TW; supervision: GJ, YZ; project administration: GJ, YZ; funding acquisition: YZ. All authors contributed to the revision and
528 improvement of the paper.

529 **Competing interests**

530 The authors declare that they have no conflict of interest.

531 **Acknowledgements**

532 We appreciate all the funding from the National Key R&D program of China (2023YFB3905100), the Project of Stable Support
533 for Youth Team in Basic Research Field, CAS (YSBR-018), the National Natural Science Foundation of China (42174212),
534 the Chinese Meridian Project, and the Specialized Research Fund for State Key Laboratories. We acknowledge the use of data
535 from the Chinese Meridian Project. We thank all the builders and maintainers of DCOI, FPI, and DCAI of Siziwang station.
536 We thank Lingli Tang for their effort in establishing and maintaining AOE_Baotou site of AERONET.

537 **Financial support**

538 This work was supported by the National Key R&D program of China (2023YFB3905100), the Project of Stable Support for
539 Youth Team in Basic Research Field, CAS (YSBR-018), the National Natural Science Foundation of China (42174212), the
540 Chinese Meridian Project, and the Specialized Research Fund for State Key Laboratories.

541 **References**

542 Abreu, V. J., Schmitt, G. A., Hays, P. B., Meriwether, J. W., Tepley, C. A., and Cogger, L. L.: Atmospheric Scattering Effects
543 on Ground-Based Measurements of Thermospheric Winds, *Planet Space Sci*, 31, 303-310, doi:10.1016/0032-
544 0633(83)90080-6, 1983.

545 Anderson, C., Conde, M., and McHarg, M. G.: Neutral thermospheric dynamics observed with two scanning Doppler imagers:
546 2. Vertical winds, *Journal of Geophysical Research: Space Physics*, 117, doi:10.1029/2011ja017157, 2012.

547 Ashburn, E. V.: The Effect of Rayleigh Scattering and Ground Reflection Upon the Determination of the Height of the Night
548 Airglow, *J Atmos Terr Phys*, 5, 83-91, doi:10.1016/0021-9169(54)90012-4, 1954.

549 Biondi, M. A. and Feibelman, W. A.: Twilight and nightglow spectral line shapes of oxygen λ 6300 and λ 5577 radiation, *Planet*
550 *Space Sci*, 16, 431-443, doi:10.1016/0032-0633(68)90158-X, 1968.

551 Biondi, M. A., Sipler, D. P., Zipf, M. E., and Baumgardner, J. L.: All-Sky Doppler Interferometer for Thermospheric Dynamics
552 Studies, *Appl Optics*, 34, 1646-1654, doi:10.1364/Ao.34.001646, 1995.

553 Burnside, R. G., Herrero, F. A., Meriwether, J. W., and Walker, J. C. G.: Optical observations of thermospheric dynamics at
554 Arecibo, *Journal of Geophysical Research: Space Physics*, 86, 5532-5540, doi:10.1029/JA086iA07p05532, 1981.

555 Burnside, R. G. and Tepley, C. A.: Optical Observations of Thermospheric Neutral Winds at Arecibo between 1980 and 1987,
556 *J Geophys Res-Space*, 94, 2711-2716, doi:10.1029/JA094iA03p02711, 1989.

557 Conde, M. and Smith, R. W.: Mapping Thermospheric Winds in the Auroral-Zone, *Geophysical Research Letters*, 22, 3019-
558 3022, doi:10.1029/95gl02437, 1995.

559 Conde, M., Craven, J. D., Immel, T., Hoch, E., Stenbaek-Nielsen, H., Hallinan, T., Smith, R. W., Olson, J., Sun, W., Frank, L.
560 A., and Sigwarth, J.: Assimilated observations of thermospheric winds, the aurora, and ionospheric currents over Alaska,
561 *Journal of Geophysical Research: Space Physics*, 106, 10493-10508, doi:10.1029/2000ja000135, 2001.

562 Crickmore, R. I., Dudeney, J. R., and Rodger, A. S.: Vertical Thermospheric Winds at the Equatorward Edge of the Auroral
563 Oval, *J Atmos Terr Phys*, 53, 485-492, doi:10.1016/0021-9169(91)90076-J, 1991.

564 Dhadly, M., Emmert, J., Drob, D., Conde, M., Doornbos, E., Shepherd, G., Makela, J., Wu, Q., Niciejewski, R., and Ridley,
565 A.: Seasonal dependence of northern high-latitude upper thermospheric winds: A quiet time climatological study based
566 on ground-based and space-based measurements, *Journal of Geophysical Research: Space Physics*, 122, 2619-2644,
567 doi:10.1002/2016JA023688, 2017.

568 Emmert, J. T., Fejer, B. G., Fesen, C. G., Shepherd, G. G., and Solheim, B. H.: Climatology of middle- and low-latitude
569 daytime region disturbance neutral winds measured by Wind Imaging Interferometer (WINDII), *J Geophys Res-Space*,
570 106, 24701-24712, doi:10.1029/2000ja000372, 2001.

571 Emmert, J. T., Faivre, M. L., Hernandez, G., Jarvis, M. J., Meriwether, J. W., Niciejewski, R. J., Sipler, D. P., and Tepley, C.
572 A.: Climatologies of nighttime upper thermospheric winds measured by ground-based Fabry-Perot interferometers during
573 geomagnetically quiet conditions: 1. Local time, latitudinal, seasonal, and solar cycle dependence, *J Geophys Res-Space*,
574 111, Artn A12302, doi:10.1029/2006ja011948, 2006.

575 Englert, C. R., Babcock, D. D., and Harlander, J. M.: Doppler asymmetric spatial heterodyne spectroscopy (DASH): concept
576 and experimental demonstration, *Appl Optics*, 46, 7297-7307, doi:10.1364/Ao.46.007297, 2007.

577 Englert, C. R., Harlander, J. M., Emmert, J. T., Babcock, D. D., and Roesler, F. L.: Initial ground-based thermospheric wind
578 measurements using Doppler asymmetric spatial heterodyne spectroscopy (DASH), *Optics Express*, 18, 27416-27430,
579 doi:10.1364/Oe.18.027416, 2010.

580 Fejer, B. G., Emmert, J. T., and Sipler, D. P.: Climatology and storm time dependence of nighttime thermospheric neutral
581 winds over Millstone Hill, *Journal of Geophysical Research: Space Physics*, 107, doi:10.1029/2001ja000300, 2002.

582 Friedman, J. F. and Herrero, F. A.: Fabry-Perot interferometer measurements of thermospheric neutral wind gradients and
 583 reversals at Arecibo, *Geophysical Research Letters*, 9, 785-788, doi:10.1029/GL009i007p00785, 1982.

584 Gonzalez-Esparza, J. A., Sanchez-Garcia, E., Sergeeva, M., Corona-Romero, P., Gonzalez-Mendez, L. X., Valdes-Galicia, J.
 585 F., Aguilar-Rodriguez, E., Rodriguez-Martinez, M., Ramirez-Pacheco, C., Castellanos, C. I., Pazos, M., Mendoza, B.,
 586 Gatica-Acevedo, V. J., Melgarejo-Morales, A., Caraballo, R., Andrade-Mascote, E., Villanueva-Hernandez, P., Bonifaz-
 587 Alfonso, R., Sierra, P., Romero-Hernandez, E., Peralta-Mendoza, I., Perez-Tijerina, E., Mejia-Ambriz, J. C., Guerrero-
 588 Peña, C., Caccavari, A., Cifuentes-Nava, G., and Hernandez-Quintero, E.: The Mother's Day Geomagnetic Storm on 10
 589 May 2024: Aurora Observations and Low Latitude Space Weather Effects in Mexico, *Space Weather-the International*
 590 *Journal of Research and Applications*, 22, ARTN e2024SW004111, doi:10.1029/2024SW004111, 2024.

591 Guo, W. and McEwen, D. J.: Vertical winds in the central polar cap, *Geophysical Research Letters*, 30, ArtN 1725,
 592 doi:10.1029/2003gl017124, 2003.

593 Guo, X., Zhao, B. Q., Yu, T. T., Hao, H. L., Sun, W. J., Wang, G. J., He, M. S., Mao, T., Li, G. Z., and Ren, Z. P.: East-West
 594 Difference in the Ionospheric Response During the Recovery Phase of May 2024 Super Geomagnetic Storm Over the
 595 East Asian, *J Geophys Res-Space*, 129, ARTN e2024JA033170, doi:10.1029/2024JA033170, 2024.

596 Hajra, R., Tsurutani, B. T., Lakhina, G. S., Lu, Q. M., and Du, A. M.: Interplanetary Causes and Impacts of the 2024 May
 597 Superstorm on the Geosphere: An Overview, *Astrophys J*, 974, ARTN 264, doi:10.3847/1538-4357/ad7462, 2024.

598 Hansen, J. E. and Travis, L. D.: Light-Scattering in Planetary Atmospheres, *Space Science Reviews*, 16, 527-610, doi:
 599 10.1007/Bf00168069, 1974.

600 Harding, B. J., Makela, J. J., Qin, J., Fisher, D. J., Martinis, C. R., Noto, J., and Wrasse, C. M.: Atmospheric scattering effects
 601 on ground-based measurements of thermospheric vertical wind, horizontal wind, and temperature, *Journal of Geophysical*
 602 *Research: Space Physics*, 122, 7654-7669, doi:10.1002/2017JA023942, 2017a.

603 Harding, B. J., Qin, J., and Makela, J. J.: Ground-Based Optical Measurements of Quiet Time Thermospheric Wind and
 604 Temperature: Atmospheric Scattering Corrections, *Journal of Geophysical Research: Space Physics*, 122,
 605 doi:10.1002/2017JA024705, 2017b.

606 Harlander, J. M., Englert, C. R., Brown, C. M., Marr, K. D., Miller, I. J., Zastera, V., Bach, B. W., and Mende, S. B.: Michelson
 607 Interferometer for Global High-Resolution Thermospheric Imaging (MIGHTI): Monolithic Interferometer Design and
 608 Test, *Space Science Reviews*, 212, 601-613, doi:10.1007/s11214-017-0374-4, 2017.

609 Hays, P. B., Killeen, T. L., Spencer, N. W., Wharton, L. E., Roble, R. G., Emery, B. A., Fullerrowell, T. J., Rees, D., Frank,
 610 L. A., and Craven, J. D.: Observations of the Dynamics of the Polar Thermosphere, *J Geophys Res-Space*, 89, 5597-5612,
 611 doi:10.1029/JA089iA07p05597, 1984.

612 He, Q. F., Fang, Z., Shoshanim, O., Brown, S. S., and Rudich, Y.: Scattering and absorption cross sections of atmospheric
 613 gases in the ultraviolet-visible wavelength range (307-725 nm), *Atmos Chem Phys*, 21, 14927-14940, doi:10.5194/acp-
 614 21-14927-2021, 2021.

615 Hernandez, G. and Roble, R. G.: Direct Measurements of Nighttime Thermospheric Winds and Temperatures .2. Geomagnetic
 616 Storms, *J Geophys Res-Space*, 81, 5173-5181, doi:10.1029/JA081i028p05173, 1976.

617 Holben, B. N., Tanré, D., Smirnov, A., Eck, T. F., Slutsker, I., Abuhassan, N., Newcomb, W. W., Schafer, J. S., Chatenet, B.,
 618 Lavenu, F., Kaufman, Y. J., Castle, J. V., Setzer, A., Markham, B., Clark, D., Frouin, R., Halthore, R., Karnieli, A.,
 619 O'Neill, N. T., Pietras, C., Pinker, R. T., Voss, K., and Zibordi, G.: An emerging ground-based aerosol climatology::
 620 Aerosol optical depth from AERONET, *J Geophys Res-Atmos*, 106, 12067-12097, doi:10.1029/2001jd900014, 2001.

621 Huang, C., Xu, J. Y., Zhang, X. X., Liu, D. D., Yuan, W., and Jiang, G. Y.: Mid-latitude thermospheric wind changes during
 622 the St. Patrick's Day storm of 2015 observed by two Fabry-Perot interferometers in China, *Advances in Space Research*,
 623 61, 1873-1879, doi:10.1016/j.asr.2017.10.013, 2018.

624 Innis, J. L., Greet, P. A., Murphy, D. J., Conde, M. G., and Dyson, P. L.: A large vertical wind in the thermosphere at the
 625 auroral oval/polar cap boundary seen simultaneously from Mawson and Davis, Antarctica, *J Atmos Sol-Terr Phy*, 61,
 626 1047-1058, doi:10.1016/S1364-6826(99)00060-7, 1999.

627 Ishii, M., Oyama, S., Nozawa, S., Fujii, R., Sagawa, E., Watari, S., and Shinagawa, H.: Dynamics of Neutral Wind in the polar
 628 region observed with two Fabry-Perot interferometers, *Earth Planets Space*, 51, 833-844, doi:10.1186/Bf03353242, 1999.

629 Ishii, M., Conde, M., Smith, R. W., Krynicki, M., Sagawa, E., and Watari, S.: Vertical wind observations with two Fabry-
 630 Perot interferometers at Poker Flat, Alaska, *Journal of Geophysical Research: Space Physics*, 106, 10537-10551,
 631 doi:10.1029/2000JA900148, 2001.

632 Jiang, G., Xu, J., Wang, W., Yuan, W., Zhang, S., Yu, T., Zhang, X., Huang, C., Kerr, R. B., Noto, J., Li, J., Liu, W., and Li,
633 Q.: A Comparison of Quiet Time Thermospheric Winds Between FPI Observations and Model Calculations, *Journal of*
634 *Geophysical Research: Space Physics*, 123, 7789-7805, doi:10.1029/2018JA025424, 2018.

635 Kataoka, R., Miyoshi, Y., Shiokawa, K., Nishitani, N., Keika, K., Amano, T., and Seki, K.: Magnetic Storm-Time Red Aurora
636 as Seen From Hokkaido, Japan on 1 December 2023 Associated With High-Density Solar Wind, *Geophysical Research*
637 *Letters*, 51, e2024GL108778, doi:10.1029/2024GL108778, 2024a.

638 Kataoka, R., Reddy, S. A., Nakano, S., Pettit, J., and Nakamura, Y.: Extended magenta aurora as revealed by citizen science,
639 *Scientific Reports*, 14, 25849, doi:10.1038/s41598-024-75184-9, 2024b.

640 Killeen, T. L., Won, Y. I., Niciejewski, R. J., and Burns, A. G.: Upper Thermosphere Winds and Temperatures in the
641 Geomagnetic Polar-Cap - Solar-Cycle, Geomagnetic-Activity, and Interplanetary Magnetic-Field Dependencies, *J*
642 *Geophys Res-Space*, 100, 21327-21342, doi:10.1029/95ja01208, 1995.

643 Li, J., Carlson, B. E., Yung, Y. L., Lv, D. R., Hansen, J., Penner, J. E., Liao, H., Ramaswamy, Kahn, R. A., Zhang, P., Dubovik,
644 O., Ding, A. J., Lacis, A. A., Zhang, L., and Dong, Y. M.: Scattering and absorbing aerosols in the climate system, *Nat*
645 *Rev Earth Env*, 3, 363-379, doi:10.1038/s43017-022-00296-7, 2022.

646 Li, W. B., Liu, L. B., Chen, Y. D., Yang, Y. Y., Han, T. W., Ding, F., Le, H. J., and Zhang, R. L.: Multi-Instruments
647 Observation of Ionospheric-Thermospheric Dynamic Coupling Over Mohe (53.5°N, 122.3°E) During the April 2023
648 Geomagnetic Storm, *J Geophys Res-Space*, 128, ARTN e2023JA032141, doi:10.1029/2023JA032141, 2023.

649 Liu, W. J., Zhu, Y. J., Xu, J. Y., Li, Q. Z., Yuan, W., Zhu, G. Y., Wang, T. C., Yang, G. T., Du, L. F., Liu, S. Y., and Li, F.
650 Q.: Validation of Neutral Wind in the Mesopause Measured by a Dual-Channel Optical Interferometer (DCOI) Network
651 of the Chinese Meridian Project, *Space Weather-the International Journal of Research and Applications*, 23, ARTN
652 e2025SW004468, doi:10.1029/2025SW004468, 2025.

653 Ma, L. X., Yu, Y. Q., Ding, X. Q., Liu, X. Y., An, D. P., Zhou, C. L., Cao, J. B., and Shiokawa, K.: Mid-Latitude Auroras and
654 Energetic Particle Precipitation Occurred Unusually in a Moderate Magnetic Storm on 1 December 2023, *Geophysical*
655 *Research Letters*, 51, ARTN e2024GL110764, doi:10.1029/2024GL110764, 2024.

656 Makela, J. J., Harding, B. J., Meriwether, J. W., Mesquita, R., Sanders, S., Ridley, A. J., Castellez, M. W., Ciocca, M., Earle,
657 G. D., Frissell, N. A., Hampton, D. L., Gerrard, A. J., Noto, J., and Martinis, C. R.: Storm time response of the midlatitude
658 thermosphere: Observations from a network of Fabry-Perot interferometers, *Journal of Geophysical Research: Space*
659 *Physics*, 119, 6758-6773, doi:10.1002/2014JA019832, 2014.

660 Mikhalev, A.: Auroras during extreme geomagnetic storms: Some features of mid-latitude aurora on February 11, 1958, *Solar-*
661 *Terrestrial Physics*, 55-61, 10.12737/stp-102202406, 2024.

662 Nakajima, H., Okano, S., Fukunishi, H., and Ono, T.: Observations of thermospheric wind velocities and temperatures by the
663 use of a Fabry-Perot Doppler imaging system at Syowa Station, Antarctica, *Appl Optics*, 34, 8382-8395,
664 doi:10.1364/Ao.34.008382, 1995.

665 Nanjo, S. and Shiokawa, K.: Spatial structures of blue low-latitude aurora observed from Japan during the extreme geomagnetic
666 storm of May 2024, *Earth Planets Space*, 76, ARTN 156, doi:10.1186/s40623-024-02090-9, 2024.

667 Price, G. D., Smith, R. W., and Hernandez, G.: Simultaneous Measurements of Large Vertical Winds in the Upper and Lower
668 Thermosphere, *J Atmos Terr Phys*, 57, 631-643, doi:10.1016/0021-9169(94)00103-U, 1995.

669 Ranjan, A. K. and Pallamraju, D.: Latitudinal Distribution of Thermospheric Nitric Oxide (NO) Infrared Radiative Cooling
670 During May and October 2024 Geomagnetic Storms, *J Geophys Res-Space*, 130, ARTN e2024JA033559,
671 doi:10.1029/2024JA033559, 2025.

672 Rees, D., Smith, M. F., and Gordon, R.: The Generation of Vertical Thermospheric Winds and Gravity-Waves at Auroral
673 Latitudes .2. Theory and Numerical Modeling of Vertical Winds, *Planet Space Sci*, 32, 685, doi:10.1016/0032-
674 0633(84)90093-X, 1984.

675 Sakanoi, T., Fukunishi, H., Okano, S., Sato, N., Yamagishi, H., and Yukimatu, A. S.: Dynamical coupling of neutrals and ions
676 in the high-latitude *F* region: Simultaneous FPI and HF radar observations at Syowa Station, Antarctica, *Journal of*
677 *Geophysical Research: Space Physics*, 107, doi:10.1029/2001JA007530, 2002.

678 Shinagawa, H. and Oyama, S.: A two-dimensional simulation of thermospheric vertical winds in the vicinity of an auroral arc,
679 *Earth Planets Space*, 58, doi:10.1186/BF03352007, 2006.

680 Shiokawa, K., Katoh, Y., Satoh, M., Ejiri, M. K., and Ogawa, T.: Integrating-sphere calibration of all-sky cameras for
681 nightglow measurements, *Adv Space Res*, 26, 1025-1028, doi:10.1016/S0273-1177(00)00052-1, 2000.

682 Singh, R., Scipi3n, D. E., Kuyeng, K., Condor, P., Flores, R., Pacheco, E., de la Jara, C., and Manay, E.: Ionospheric Responses
683 to an Extreme (G5-Level) Geomagnetic Storm Using Multi-Instrument Measurements at the Jicamarca Radio
684 Observatory on 10-11 October 2024, *J Geophys Res-Space*, 130, ARTN e2024JA033642, doi:10.1029/2024JA033642,
685 2025.

686 Smith, R. W. and Hernandez, G.: Vertical Winds in the Thermosphere within the Polar-Cap, *J Atmos Terr Phys*, 57, 611-620,
687 doi:10.1016/0021-9169(94)00101-S, 1995.

688 Sobolev, V. V., Irvine, W. M., Gendel, M., and Lane, A. P.: Light scattering in planetary atmospheres, 1st . International series
689 of monographs in natural philosophy vol.76, Pergamon Press, Oxford, New York, doi:10.1016/C2013-0-05661-7, 1975.

690 Themens, D. R., Elvidge, S., McCaffrey, A., Jayachandran, P. T., Coster, A., Varney, R. H., Galkin, I., Goodwin, L. V., Watson,
691 C., Maguire, S., Kavanagh, A. J., Zhang, S. R., Goncharenko, L., Bhatt, A., Dorrian, G., Groves, K., Wood, A. G., and
692 Reid, B.: The High Latitude Ionospheric Response to the Major May 2024 Geomagnetic Storm: A Synoptic View,
693 *Geophysical Research Letters*, 51, e2024GL111677, doi:10.1029/2024GL111677, 2024.

694 Wang, C., Xu, J., Chen, Z., Li, H., Feng, X., Huang, Z., and Wang, J.: China's Ground-Based Space Environment Monitoring
695 Network—Chinese Meridian Project (CMP), *Space Weather*, 22, doi:10.1029/2024sw003972, 2024.

696 Wang, X., Aa, E., Chen, Y., Zhang, J., Zhu, Y., Cai, L., Lu, X., Luo, B., Liu, S., Li, M., Shen, H., and Yuan, T.: Midlatitude
697 Neutral Wind Response During the Mother's Day Super-Intense Geomagnetic Storm in 2024 Using Observations From
698 the Chinese Meridian Project, *Journal of Geophysical Research: Space Physics*, 130, e2024JA033574,
699 doi:10.1029/2024JA033574, 2025.

700 Wei, D., Zhu, Y., Liu, J., Gong, Q., Kaufmann, M., Olschewski, F., Knieling, P., Xu, J., Koppmann, R., and Riese, M.:
701 Thermally stable monolithic Doppler asymmetric spatial heterodyne interferometer: optical design and laboratory
702 performance, *Optics Express*, 28, doi:10.1364/oe.394101, 2020.

703 Wu, Q., Yuan, W., Xu, J. Y., Huang, C., Zhang, X. X., Wang, J. S., and Li, T.: First U.S.-China joint ground-based Fabry-
704 Perot interferometer observations of longitudinal variations in the thermospheric winds, *J Geophys Res-Space*, 119, 5755-
705 5763, doi:10.1002/2014ja020089, 2014.

706 Xu, H., Shiokawa, K., Oyama, S., and Otsuka, Y.: Thermospheric wind variations observed by a Fabry-Perot interferometer
707 at Troms3, Norway, at substorm onsets, *Earth Planets Space*, 71, ARTN 93, doi:10.1186/s40623-019-1072-0, 2019.

708 Yang, C., Zhao, B., Jin, Y., Huang, C., Yao, X., and Wan, W.: Climatology of Nighttime Upper Thermospheric Winds From
709 Fabry-Perot Interferometer 2011–2019 Measurements Over Kelan (38.7°N, 111.6°E), China: Local Time, Seasonal, Solar
710 Cycle, and Geomagnetic Activity Dependence, *Journal of Geophysical Research: Space Physics*, 125, e2020JA027892,
711 doi:10.1029/2020JA027892, 2020.

712 Yu, T., Huang, C., Zhao, G. X., Mao, T., Wang, Y. G., Zeng, Z. C., Wang, J. S., and Xia, C. L.: A preliminary study of
713 thermosphere and mesosphere wind observed by Fabry-Perot over Kelan, China, *J Geophys Res-Space*, 119, 4981-4997,
714 doi:10.1002/2013ja019492, 2014.

715 Yuan, W., Xu, J. Y., Ma, R. P., Wu, Q. A., Jiang, G. Y., Gao, H., Liu, X. A., and Chen, S. Z.: First observation of mesospheric
716 and thermospheric winds by a Fabry-Perot interferometer in China, *Chinese Sci Bull*, 55, 4046-4051, doi:10.1007/s11434-
717 010-4192-2, 2010.

718 Zhang, K. D., Wang, H., Liu, J., Song, H. M., and Xia, H.: The Quasi-Periodic Nighttime Traveling Ionospheric Disturbances
719 on 13 May 2024 During the Recovery Phase of a SuperStorm, *J Geophys Res-Space*, 130, ARTN e2024JA033257,
720 doi:10.1029/2024JA033257, 2025.

721 Zhu, G., Zhu, Y., Kaufmann, M., Wang, T., Liu, W., and Xu, J.: An Efficient Calibration System of Optical Interferometer for
722 Measuring Middle and Upper Atmospheric Wind, *Remote Sensing*, 15, doi:10.3390/rs15071898, 2023.