

1 Attribution of changes in winds over the
2 Southern Ocean from 1950 to 2100
3

4 Tereza Jarníková¹, Colin Jones², Steven Rumbold³, Corinne Le Quéré¹

5
6 1. School of Environmental Sciences, University of East Anglia, UK

7 2. National Centre for Atmospheric Science, University of Leeds, UK.

8 3 National Centre for Atmospheric Science, University of Reading, UK

9 Correspondence to: Tereza Jarníková (T.Jarnikova@uea.ac.uk)

Deleted: Jarníková

Deleted: Jones

Deleted: Rumbold

Deleted: Quéré

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font: Arial, Font color: Auto

Formatted: Heading 1, Space After: 0 pt, Line spacing: single

Formatted: Font: Italic

1 Abstract

2
3 Strong near-surface westerly winds drive the Southern Ocean circulation and play a key role in
4 setting regional and global climate. In the latter half of the 20th century, depletion of stratospheric ozone
5 over Antarctica has caused these winds to accelerate and move polewards, particularly in austral summer.
6 However, the future evolution of these winds remains uncertain. We use reanalysis data and the UK
7 Earth System Model (UKESM1), with full atmospheric chemistry, to assess the drivers of winds over the
8 recent past and coming century. We first characterize the wind mean state, distribution, and trends over
9 1980-2019 in the most commonly used atmospheric reanalyses (ERA5, JRA3Q, and MERRA2) to gain
10 insight into observed wind behaviour in the past. We show that while the representation of the mean wind
11 is similar among reanalyses, MERRA2 shows stronger wind acceleration trends that persist year-round,
12 while JRA3Q and ERA5 show weaker acceleration, primarily in austral summer. Using an observational
13 Southern Annular Mode (SAM) index, we show that the weaker, summer-focused trends of JRA3Q and
14 ERA5 are likely more realistic. UKESM1 represents historical trends in winds accurately compared to
15 ERA5 and is within the range of other CMIP6 models for wind and SAM trends over the historical
16 period. Targeted simulations with UKESM1 show ozone depletion is overwhelmingly responsible for the
17 wind acceleration observed in 1980-2020, primarily in austral summer. The effect of ozone depletion on
18 wind speeds peaks in 1980-2000, when it is roughly double that for the entire 40-year period. Ozone
19 recovery is then associated with a slowdown of winds from 2000 to 2050. Beyond 2050, the ozone effect
20 becomes minimal and winds accelerate primarily due to greenhouse gas induced warming, with this trend
21 more evenly distributed across seasons.
22

Deleted: , and NCEP-NCAR

Deleted: and NCEP-NCAR show

23 1. Introduction

24
25 Strong near-surface westerly winds are a dominant feature of the midlatitude atmospheric
26 circulation over the Southern Ocean, driving the Antarctic Circumpolar Current (ACC), the largest ocean
27 current on Earth. The ACC is an important control on the Atlantic Meridional Overturning Circulation
28 (AMOC; J. Marshall & Speer, 2012) and plays a fundamental role in controlling Southern Ocean heat
29 (Huguenin et al., 2022) and carbon uptake (Le Quéré et al., 2007).

30 Over the last four decades, substantial changes in the Southern Hemisphere wind regime have
31 been seen in both reanalyses and observations. Chiefly, the wind jet (i.e., the location of the strongest
32 westerly winds) has intensified and moved poleward (Swart & Fyfe, 2012), with substantial longitudinal
33 variation (Goyal et al., 2021; Waugh et al., 2020). These changes are manifested in significant trends in
34 the Southern Annular Mode (SAM; Fogt & Marshall, 2020).
35

36 The observed changes in southern hemisphere winds over the latter half of the 20th century have
37 been linked to the depletion of stratospheric ozone, particularly the Antarctic ozone hole (Thompson &
38 Solomon, 2002). The absorption of incoming solar UV radiation by ozone warms the stratosphere, so
39 ozone depletion leads to a relative stratospheric cooling. This cooling strengthens the north-south
40 temperature gradient, which in turn enhances the vertical wind shear and deepens the polar vortex,

Formatted: Outline numbered + Level: 1 + Numbering
Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at:
0.63 cm + Indent at: 1.27 cm

1 especially in austral spring and summer (for an overview, see, e.g. Previdi & Polvani, 2014; Thompson et
2 al., 2011). This acceleration results in an overall strengthening and poleward shift of Southern Ocean
3 winds throughout the troposphere, particularly in austral summer (DJF), with an accompanying shift
4 towards a more positive index of the SAM, the principal mode of atmospheric variability in the region (G.
5 J. Marshall, 2003; Swart & Fyfe, 2012; Thompson et al., 2011).

6
7 Substantial modeling efforts have sought to understand the relative contributions of ozone
8 depletion versus greenhouse gas forcing to the observed Southern Hemisphere wind changes. Past work,
9 using several generations of CMIP (Coupled Model Intercomparison Project) and CCMI (Chemistry-
10 Climate Model Initiative) models, has shown the recent Southern Hemisphere atmospheric circulation
11 changes are primarily attributable to stratospheric ozone depletion (e.g. S. Son et al., 2009; S.-W. Son et
12 al., 2018). However, inter-model spread is often large, and the quantitative responses of the winds and
13 other atmospheric responses differ substantially between models (e.g. Gerber & Son, 2014). For example,
14 the poleward migration of the westerly jet tends to be stronger in models whose mean jet position is
15 biased toward lower latitudes (S. -W. Son et al., 2010). The representation of ozone also plays a role in
16 the character of the atmospheric response. For example, CMIP5 models with interactive ozone have a
17 larger spread in the historical and future changes in jet position than those with prescribed ozone (Eyring
18 et al., 2013). Simultaneously, CMIP6 models with interactive ozone tend to have a stronger role for ozone
19 in the SAM strengthening than models with prescribed ozone (Morgenstern, 2021). Together, these
20 results highlight the importance of correctly representing ozone loss and recovery for capturing
21 circulation changes.

22
23 Ozone-driven shifts in the jet stream and wind intensification are expected to reverse as Southern
24 Hemisphere ozone recovers (Polvani et al., 2011; Solomon et al., 2017). However, this recovery will be
25 simultaneously opposed by greenhouse gas-driven warming, which acts in the opposite direction
26 (Arblaster et al., 2011; McLandress et al., 2011; Zambri et al., 2021). The future evolution of the overall
27 wind patterns thus depends on which forcing dominates. Barnes et al. (2014) showed that ozone recovery
28 delays the effect of greenhouse gas driven climate change on multiple Southern Hemisphere climate
29 indicators, including the position of the jet stream, and that the historical ozone-driven circulation changes
30 are larger than those projected to the end of the twenty-first century. Other studies observe that under
31 ozone recovery, the westerly jet that had previously moved poleward in response to ozone depletion
32 remains largely stationary, as the effects of ozone recovery are opposed by greenhouse gas forcing
33 (Gerber & Son, 2014; S. -W. Son et al., 2010).

34
35 Considering trends in the SAM index, which can be taken as a proxy for wind speed, over the
36 period ozone recovery is expected (~2000–2050) Simpkins and Karpechko (2012) show the effects of
37 greenhouse gas forcing, pushing the SAM index to be more positive, opposes ozone recovery acting to
38 make the SAM more negative. In the latter half of the century, they show that the evolution of the SAM is
39 sensitive to the magnitude of greenhouse gas emissions. This sensitivity of atmospheric circulation
40 features to the strength of greenhouse gas emissions is seen in multiple studies (e.g. Eyring et al., 2013,
41 Barnes et al., 2014), highlighting that uncertainties in both future emissions and model response play a
42 role in the uncertainty in future atmospheric circulation changes.

1 In this paper, we examine both recent changes in the Southern Ocean winds in reanalysis products
2 and quantify the relative contribution of stratospheric ozone depletion and rising greenhouse gas (GHG)
3 concentrations in driving these changes. While the general trends in the Southern Ocean winds are well
4 understood, differences in the representation of wind trends exist between reanalysis products. These may
5 translate into differences in ocean responses when global ocean models are forced by different reanalyses
6 (Friedlingstein et al., 2023; Tsujino et al., 2020) or have implications for energy production, as wind
7 energy infrastructure depends on accurate estimates of regional wind state (Gualtieri, 2022). Furthermore,
8 most studies to date have focused on the mean trend in the wind jet and omitted variability and extreme
9 winds, which are disproportionately important for a number of ocean and climate processes, for example
10 in rapid sea-ice loss (Jena et al., 2022), ice shelf calving events (Francis et al., 2020), and air-sea CO₂
11 exchange (Gu et al., 2021).

12
13 We first intercompare a suite of the most commonly used reanalysis products over the period
14 1980-2020, assessing the wind climatology and wind speed frequency distribution, as well as time trends
15 in the mean wind speed, in extreme wind speeds, the position and variability of the wind jet, and the SAM
16 index. Our general aim is to determine which reanalyses are most suitable for wind applications in the
17 Southern Ocean, for example for forcing offline ocean models.

18
19 In the second part of the paper, we assess the representation of the winds from the UK Earth
20 System Model (UKESM1) and then use this model to attribute relative drivers of historical and future
21 Southern Ocean winds. We first evaluate the UKESM1 wind and SAM representation against the
22 reanalyses and then compare wind speeds, SAM, and total column ozone (TCO) in UKESM1 to other
23 CMIP6 models that include interactive atmospheric chemistry and have relevant output available. We
24 have two main applications in mind. First, we want to understand how suitable UKESM1 is for studying
25 the wider climate effects of changes in the wind distribution. For example, we recently used UKESM1 to
26 study the role of wind changes in modifying the uptake of carbon dioxide by the Southern Ocean
27 (Jarníková et al., 2025); here we aim to provide an evaluation of the robustness of the wind changes
28 reported there. Second, we want to understand to what extent this model can be used to attribute the
29 relative contribution of ozone and greenhouse gases to past and future changes in the wind distribution.

30
31 Finally, following this evaluation, we attribute the relative drivers of historical changes in
32 Southern Ocean winds and project their future evolution through to the end of the century. To do this, we
33 perform a set of experiments with UKESM1 covering the period 1950-2100, modifying the surface
34 mixing ratios of Ozone Depleting Substances (ODS) to generate two stratospheric ozone scenarios (one
35 with essentially no ozone loss and a second with historically accurate ozone evolution, i.e. Antarctic
36 ozone depletion, followed by a subsequent recovery). These two ozone scenarios are each combined with
37 two CMIP6 SSP (Shared Socioeconomic Pathway) scenarios that represent a high and low GHG emission
38 scenario. The resulting four simulations, covering 1950 to 2100, allow us to assess both past and future
39 wind speed changes over the Southern Ocean and attribute these changes to one or both of ozone and
40 greenhouse gas forcing.

41 2. Methods

Formatted: Outline numbered + Level: 1 + Numbering
Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at:
0.63 cm + Indent at: 1.27 cm

2.1 Selection of reanalyses and previous evaluation

Meteorological reanalyses provide a gridded estimate of the complete atmospheric state by assimilating satellite and in-situ observations within a numerical weather prediction model. A large number of global atmospheric reanalysis products are available (see <https://reanalyses.org/> for an overview). Reanalyses are commonly used in climate monitoring and policy applications, as well as in fundamental research more broadly. For example, global and regional carbon cycle models are often forced by reanalysis products (Friedlingstein et al., 2023), so any differences between reanalyses may influence such studies.

We use a subset of the latest generation of products that are commonly used in earth system research: ERA5, JRA3Q and MERRA2. The main characteristics of these reanalyses are given in Table 1. We excluded the NCEP-NCAR reanalysis as it is planned to be discontinued in 2026. We initially also included NCEP-DOE2, an update of NCEP-NCAR, but ultimately excluded it because of anomalously large wind speed biases relative to the other products. Spurious high winds in NCEP-DOE2 have been noted previously. For example, Lucio-Eceiza et al. (2019) found that NCEP-DOE2 has worse performance than its predecessor NCEP-NCAR in estimating surface wind speeds in the North Atlantic, while Dong et al. (2020) found it had markedly worst performance in estimating Antarctic Ice Sheet winds compared to five other contemporary products.

Here we briefly survey some existing assessments of our chosen reanalyses and their predecessors, focusing on the Antarctic and Southern Ocean. Though a number of past studies have evaluated reanalysis winds in the Southern Ocean, the extent of past evaluation in the near-Antarctic region remains relatively limited because of the sparsity of available in-situ measurements (Caton Harrison et al., 2022; Jones et al., 2016).

Broadly, most analysis found ERA5 performed better than, or as well as, other products when evaluated against in-situ observations. Li et al. (2013) found that ERA-Interim (a predecessor to ERA5) generally had a low bias in 10m wind speed compared to in-situ shipboard observations across the entire Southern Ocean (0.06 m s^{-1} , as compared to a high bias of 1.37 m s^{-1} for NCEP-DOE (the predecessor to NCEP-DOE2)). When Jones et al. (2016) evaluated the ERA-Interim, JRA-55 (predecessor to JRA3Q), and MERRA1 (predecessor to MERRA2) reanalyses against research vessel observations, radiosonde automatic weather station (AWS) measurements, and radiosondes in the Amundsen sea, the group found JRA-55 had the smallest wind speed biases compared to AWS and research vessel observations, while ERA-Interim showed lowest biases compared to radiosonde profiles. Overall, Jones et al. showed that all three products represented open-ocean wind speeds reasonably against observations. Caton Harrison et al. (2022) found ERA5 slightly outperformed JRA-55 and MERRA2 winds in an analysis against coastal station and Advanced Scatterometer (ASCAT) measurements, though performance was similar between reanalyses. Similarly, Dong et al. (2020) found ERA5 outperformed MERRA2, ERA-Interim, and JRA-55 against 56 meteorological stations over the Antarctic Ice Sheet. Li, Jones, and Caton Harrison all note that all reanalysis products overestimate in-situ winds at low-wind speed conditions ($< \sim 4 \text{ m s}^{-1}$) and underestimate them at high-wind conditions ($> \sim 25 \text{ m s}^{-1}$).

Deleted: We also include the older, but still commonly used, NCEP-NCAR reanalysis (R1), as a comparison.

Deleted: analyzed

Deleted: R1

Deleted: four

Deleted: R1

Deleted: , R1

1 A global intercomparison of near-surface winds that included ERA5, MERRA2, and JRA-55 (a
2 predecessor of JRA3Q) found ERA5 substantially outperformed the other reanalyses in reproducing in-
3 situ observations from (land-based) tall tower platforms (Ramon et al., 2019). Based on these findings
4 and the studies outlined above, we treat ERA5 as our benchmark reanalysis, following other Southern
5 Ocean wind studies (e.g. Goyal et al., 2021).

6 2.2 Spatiotemporal standardization

7
8 We account for differing spatial and temporal resolution when comparing reanalysis products. 10-
9 m wind speed calculated from u and v components at hourly resolution and then averaged to daily
10 resolution is typically higher than wind speed calculated from the same u and v components that have
11 been first averaged to daily resolution. We therefore average all u and v components to daily resolution
12 before deriving wind speed. For similar reasons, we interpolate all three fields (u-component, v-
13 component, and wind speed) to a standard $1^\circ \times 1^\circ$ grid using the cdo package (Schulzweida, 2023). We are
14 interested primarily in the mean state, trends, and extremes of the open-ocean circumpolar winds. As the
15 wind jet is typically found between 48°S and 54°S (Swart & Fyfe, 2012), we focus our analysis on the
16 open water winds in the region 40°S to 60°S .

17
18 Annual and seasonal means are reported by first calculating area-weighted daily mean wind
19 speeds from the $1^\circ \times 1^\circ$ gridded product for the region 40°S to 60°S , then calculating the seasonal mean
20 from these daily means. To calculate extreme high (low) wind speeds, we first calculate the daily
21 weighted 95th (5th) percentile of winds from the $1^\circ \times 1^\circ$ product over 40°S to 60°S , then take the weighted
22 average of all cells above (below) this percentile. The seasonal extreme winds are then the average of
23 these daily extreme winds for each season in each year. We calculate linear decadal trends in mean and
24 extreme winds from annual and seasonal values for the time periods 1980-2019 and 1980-1999, testing
25 for significance at the 5% level using the Wald test (Wald, 1943). We calculate the interannual variability
26 (IAV) of each reanalysis from the annual (seasonal) means as the unbiased standard deviation of the time
27 series, expressed as a percentage of the mean wind speed. When calculating the frequency distribution of
28 wind speeds, we consider the full time series of area-weighted, daily mean open water wind speeds from
29 the $1^\circ \times 1^\circ$ gridded product for the region 40°S to 60°S , using 100 evenly spaced bins between 0 and 20 m
30 s^{-1} .

31
32 We use the $1^\circ \times 1^\circ$ gridded products to calculate the wind jet position. At each longitude, for each
33 day, we record the jet position as the location of the maximum of the u-component of the 10-meter wind
34 speed between 30°S and 70°S , following Bracegirdle et al. (2013). We then use this daily wind jet
35 position at each longitude to calculate the zonal average and the seasonal average.

36
37 We consider a forty-year time period (Jan 1, 1980 – Dec 31, 2019), available for all considered
38 reanalyses; prior to 1979, satellite measurements are limited. We further separately consider the first half
39 of the time period (1980-1999), which corresponds to the period of maximum Antarctic ozone loss
40 (Solomon et al., 2017), and which is expected to drive wind speed increases and a poleward shift of the
41 westerly jet.

1 2.3 SAM index

2 In each reanalysis, we calculate and then evaluate the SAM index at monthly resolution against the
3 observational SAM index (G. J. Marshall, 2003). Following Velasquez-Jimenez and Abram (2024), the
4 “natural”, or non-normalized, SAM index is calculated as:

$$5 \text{ SAM} = P^*_{40^\circ \text{ S}} - P^*_{65^\circ \text{ S}},$$

7
8 where $P^*_{40^\circ \text{ S}}$ and $P^*_{65^\circ \text{ S}}$ are the zonal MSLP anomalies at 40°S and 65°S, respectively, relative to the time
9 period 1980-2019.

10 In the observational SAM index, these MSLP anomalies are calculated from the mean of six
11 station records near each of the two latitudes for which good long-term records exist. In the reanalysis
12 SAM index we simply use the zonal mean at both latitudes as in Gong and Wang (1999). We use the
13 zonal mean rather than subsampling the reanalysis to the six station locations to maintain consistency
14 with standard reanalysis-based SAM calculations (e.g. Morgenstern, 2021) and to avoid sampling biases
15 introduced by point-location extractions from gridded fields. The zonal mean better represents the large-
16 scale circulation patterns that define the SAM, whereas point samples can be influenced by local
17 topographic effects and sub-grid-scale processes that are not fully resolved in reanalysis products.

18 Unlike the Marshall and Gong and Wang indices, the natural SAM index presented here is not
19 normalized by dividing by the reference interval standard deviation. It is therefore not dimensionless and
20 given in units of hPa. This approach has the advantage of making the trends and magnitude of the index
21 less sensitive to sampling frequency.
22

23 2.4 Modelled Ozone Depletion

24
25 We perform simulations with UKESM1 to quantify the relative contribution of ozone depletion
26 and greenhouse gas induced warming on the wind field trends. UKESM is a well-established earth system
27 model (ESM, Sellar et al., 2020; Yool et al., 2021) based on the HadGEM3-GC3.1 coupled physical
28 atmosphere-ocean model (Kuhlbrodt et al., 2018). We refer readers to these articles for a general
29 evaluation of UKESM1, and only highlight here components of the model that are central to our work.
30 The atmospheric component of UKESM1 is the Global Atmosphere 7.1 (GA7.1) science configuration of
31 the Unified Model (Walters et al., 2019), with horizontal resolution of approximately 135 km ($1.25^\circ \times$
32 1.875°), 85 vertical levels and a model top located at 85km altitude. Unusually for a CMIP6 model
33 (Coupled Model Intercomparison Project Phase 6; Eyring et al., 2016), UKESM1 simulates full
34 atmosphere ozone chemistry through the U.K. Chemistry and Aerosols (UKCA) model (Archibald et al.,
35 2020; Mulcahy et al., 2018), interactively coupled to the model’s physics and dynamics. Ozone is
36 therefore prognostic, with its evolution dependent on, and influencing, the simulated atmospheric
37 thermal, dynamical and chemical states (Keeble et al., 2021).
38
39
40

1 The ability of UKESM1 to simulate the historical and future evolution of ozone has been
2 discussed in Keeble et al. (2021), Morgenstern et al. (2022) and Zeng et al., (2022), while representation
3 of the SAM in CMIP6 models (including UKESM1) has been analysed by Morgenstern (2021) .
4 UKESM1 overestimates the observed global mean total column ozone (TCO) over the period 1980 to
5 2015, with this overestimation reduced over the Antarctic region (60°S to 90°S). In addition, UKESM1
6 has a stronger negative TCO trend globally than observed, with this overestimate also reduced for the
7 60°S to 90°S region . The bias in UKESM1 trends in TCO (discussed further in section 3.3) should be
8 kept in mind when the impact of ozone depletion and recovery on surface winds is discussed.
9

10 2.5 CMIP6 model intercomparison

11 To contextualize UKESM1 results, we report trends in the winds and the SAM index, as well as
12 in total column ozone (TCO), for other CMIP6 models with interactive chemistry for which SAM, TCO,
13 and (at least) daily-resolution wind fields are available (Table 2). Spatiotemporal standardization and
14 SAM index calculation is performed as described in sections 2.2 and 2.3. For the intercomparison of the
15 subset of CMIP6 models, we use the historical run for the period 1980-2014 and the SSP 3-7.0 run for the
16 period 2015-2019. We calculate trends in SAM, 10-m wind speed, and total column ozone (TCO) for the
17 band 70°S- 90°S, for 1980-1999 and 1980-2019. We compare CMIP6 TCO trends to an observational
18 dataset (Bodeker et al., 2021). For this intercomparison, to maintain consistency in method, only one
19 ensemble member of each model, including UKESM1, is used.

20 2.6 Experimental Design

21 Morgenstern (2020), Revell et al. (2022) and Zeng et al., (2022) emphasize that to simulate
22 forced trends in Southern Ocean winds models need to interactively simulate ozone chemistry, GHGs and
23 ODS. This is to capture multiple interactions between GHGs (e.g. CO₂, CH₄, N₂O), ozone, and their
24 combined impact on model dynamics. Prescribing an externally generated ozone field risks it being
25 chemically inconsistent with other time-evolving GHGs (Zeng et al., 2022), and potentially be offset with
26 respect to the model thermodynamical fields (Morgenstern, 2020). Morgenstern (2020) shows that the
27 CMIP6 DAMIP (Detection and Attribution Model Intercomparison Project) experiments (Gillett et al.,
28 2016); *hist-GHG* (historical GHG forcings with all other forcings pre-industrial) and *hist-stratO3*
29 (prescribed historical ozone and all other forcings pre-industrial) do not combine to give the (forced) time
30 evolution of the SAM index seen in full historical simulations. This is due to feedbacks between the time-
31 varying GHGs and ozone leading to a different impact on model dynamics compared to the combination
32 of the two individual drivers. This feedback is captured in models that run with interactive chemistry.
33

34 Motivated by these studies, we use UKESM1 with interactive chemistry to study the role of
35 stratospheric ozone loss and recovery on Southern Ocean winds. We control the evolution of stratospheric
36 ozone in UKESM1 by modifying surface mixing ratios of ozone depleting substances (ODS, e.g.
37 chlorofluorocarbons and hydrochlorofluorocarbons), which play a central role in driving stratospheric ozone
38 loss (Farman et al., 1985; Solomon et al., 1986). We perform simulations for 1950 to 2100 using two
39 ODS surface mixing ratio scenarios: (i) ODS use the standard CMIP6 surface mixing ratios (historical
40 followed by a projection), and (ii) ODS are fixed at 1950 values. We refer to these two experiments as

1 OZONE-HIST and OZONE-1950. OZONE-HIST results in ozone loss from approximately 1970 to 2000,
2 followed by a slow recovery through to 2100 (Keeble et al. 2021, Fig. 7). OZONE-1950 minimizes
3 stratospheric ozone loss throughout the simulation as only trace amounts of ODS, emitted or produced in
4 the atmosphere before 1950, are available for ozone destruction in the stratosphere. The two ODS
5 scenarios are combined with two CMIP6 SSP scenarios (SSP 3-7.0 and SSP 1-2.6; Gidden et al., 2019)
6 that represent a high and low GHG emission scenario.

7
8 This configuration results in four experiments that allow us to isolate the effects of simulated
9 stratospheric ozone and GHG on wind trends; see Table 3 for a summary. Following McLandress et al.
10 (2011), we assume that ozone-driven trends (resulting from the different ODS surface mixing ratios) and
11 GHG-driven trends are additive; that is, the OZONE-HIST run demonstrates a linear addition of ODS-
12 driven trends and GHG-driven trends, so [OZONE-HIST – OZONE-1950] will isolate the ODS-driven
13 trend, while any trends in the OZONE-1950 runs are due to GHG emissions alone. We acknowledge the
14 potential for some non-linear interactions that may weaken this assumption but suggest to first order it is a
15 reasonable approximation for attributing the primary differences identified either to ODS or GHG
16 differences in the respective experiments.

17
18 For each UKESM1 experiment, we run three ensemble members, each branched in 1850 from the
19 CMIP6 UKESM1 piControl following the procedure for generating initial conditions outlined in section 4
20 of Sellar et al., (2020). With respect to means and trends, we report the ensemble mean value, calculated
21 at daily $1^\circ \times 1^\circ$ resolution. When reporting extreme values, interannual variability, jet position, and
22 standard deviation in the jet position, we calculate them for each ensemble member separately and give
23 the mean of these calculations.

24 3. Results

25 3.1 Reanalysis Intercomparison

26 3.1.1 Wind climatology and distribution evaluation

27
28 We intercompare the reanalyses both to identify their utility for applications such as forcing
29 ocean-only models, and to provide a baseline against which to evaluate the UKESM1 historical
30 simulation. All the reanalysis products feature a prominent band of high winds between 40°S and 60°S ,
31 with intensification over the Indian Sector (Fig. 1). MERRA2 agrees with ERA5 at high latitudes, but is
32 evenly low compared to ERA5 over the open ocean. JRA3Q is generally higher than ERA5 (Fig. 1).

33
34 The reanalysis products generally agree when considering only open-ocean winds between 40°S
35 and 60°S . Most reanalyses are relatively similar to ERA5, ranging from a difference of -0.20 m s^{-1}
36 (MERRA2) to $+0.28 \text{ m s}^{-1}$ (JRA3Q) for annual-average winds. In comparison, NCEP-DOE2, excluded
37 from this analysis for high bias, has a bias of $+1.31 \text{ m s}^{-1}$ for yearly-averaged winds in the same region.
38 Interannual variability (IAV) is relatively consistent but varies somewhat between reanalyses, ranging

Formatted: Outline numbered + Level: 1 + Numbering
Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at:
0.63 cm + Indent at: 1.27 cm

Deleted: R1 strongly underestimates wind speeds at high
latitudes near the continent relative to the 1980-2019 ERA5
climatology but shows a less clear difference over the mid-
latitude (40°S and 60°S) open ocean that varies zonally and
averages to near-zero (Table 4).

1 from 1.2% in ERA5 to 1.80% in MERRA2 (Supplementary Table ST1). No strong seasonal differences in
2 interannual variability are seen in any reanalysis.

Deleted: 2.4%, or roughly double,

Deleted: R1

3
4 The frequency distribution of the daily winds is remarkably similar between the reanalyses (Fig.
5 2, Supplementary Table ST2), with differences mostly visible at the tails of the distribution. ERA5 has the
6 lowest 'low' (5th percentile) winds, but the other reanalyses have 'low' winds only slightly ($\sim 0.1 \text{ m s}^{-1}$)
7 higher than ERA5. There is more spread at the high end of the distribution, where ERA5 sits in the
8 middle of the reanalysis set and year-round differences from ERA5 range from -0.35 m s^{-1} (MERRA2) to
9 0.59 m s^{-1} (JRA3Q).

Deleted: and R1 has anomalously high "low" winds (0.44 m s^{-1} higher than ERA5). The

10 3.1.2 Wind speed trends

11 Trends in open-ocean wind speed differ between the reanalyses much more than the
12 climatological mean fields (Fig. 2a-b, Fig. 3, Table 5). JRA3Q and ERA5 behave similarly over the whole
13 time period (1980-2019), having a statistically significant year-round mean trend of $\sim 0.04 \text{ m s}^{-1} \text{ dec}^{-1}$ that
14 is slightly stronger in austral summer and autumn and not statistically significant in winter (JJA) and
15 spring (SON). In contrast, MERRA2 has much stronger trends $\sim 2.5\text{x}$ the magnitude of the ERA5 annual
16 mean tren. The seasonality in this reanalysis is also very different from ERA5, with relatively strong,
17 statistically significant trends persisting year round. Trends in MERRA2 are also statistically significant
18 over a much larger area of the Southern Ocean, as well as in the JJA season (Fig. 3).

Deleted: R1 and

Deleted: have

Deleted: 3.5x and ~

Deleted: trend, respectively.

Deleted: these two reanalyses

Deleted: R1 and

Deleted: R1 and

19 Summer (DJF) mean wind speed trends in the period of maximum ozone depletion (1980-1999)
20 are approximately double those for the full time period in all reanalyses (Fig. 2, Table 5). Trends in the
21 extreme winds follow a similar pattern to the mean winds, with MERRA2 trends much stronger than
22 those of JRA3Q and ERA5 (Fig. 2, e-h)). Trends in extreme winds are not systematically stronger than
23 trends in mean winds and are less likely to be statistically significant.
24
25
26

27 3.1.3 Jet position and poleward trend

28
29 The annual average zonal mean wind jet sits at 52.1°S in ERA5 and is similar between products,
30 (Fig. 4, Sup. Table ST3). The jet position exhibits a substantial seasonal cycle, being at its most southerly
31 location in austral autumn and most northerly location in austral spring. All reanalyses represent this
32 seasonal cycle in the jet position, with an approximate amplitude (of the seasonal averages) of $\sim 1.6^\circ$.
33 Interannual variability of the mean jet position over the time period 1980-2019 is comparable between
34 reanalyses \sim with a standard deviation of the zonal mean of approximately 0.8° (MERRA2). The zonally-
35 varying standard deviation of the jet position (shown in Fig. 4 only for ERA5, for clarity) is also
36 comparable. The strongest poleward trend in the wind jet occurs in austral summer and autumn, and is
37 much stronger ($1.8\text{x} - 3.8\text{x}$) during the period of maximum ozone depletion (1980-1999) than over the
38 whole time period (Sup. Table ST4). Trends in the position of the jet are not always statistically
39 significant in the reanalyses because of the large interannual variability.
40

Deleted: , ranging from 51.6°S (R1) to 52.3°S (JRA3Q) (Fig.

Deleted: which ranges in

Deleted: from

Deleted: $^\circ$ (MERRA2) to $\sim 2.4^\circ$ (R1).

Deleted: the

Deleted: ranges from

Deleted:) to 0.9° (R1

1 3.1.4 SAM index

2
3 The timeseries of the station-based SAM index (see Methods) provides a possible observational
4 constraint on the accuracy of the reanalysis products. The SAM indices derived from reanalyses exhibit a
5 high temporal coherence with the observational SAM index, with some deviations – for example in
6 JRA3Q during 2005-2010 (Fig. 5). The trend in the SAM index has a seasonal character both in the
7 observations and in the reanalyses. It is strongest in austral summer (DJF) and autumn (MAM), and
8 typically much weaker and not statistically significant in the other seasons (Fig. 5, Sup. Table ST5). The
9 observational trend is much stronger in the period of maximum ozone loss than in the whole time period
10 (factor of ~3x), which is reflected in the reanalyses (factor of ~2.3-3.6x).

11
12 In all seasons, the trend in the SAM index in the reanalyses correlates tightly with the trend in the
13 estimated wind speed ($r=0.72-0.93$, $p<0.05$). Because the wind acceleration is mechanically linked to
14 the trend in the SAM, evaluating the accuracy of the reanalysis-derived SAM indices against the
15 observational index may provide a constraint on the accuracy of reanalyzed wind speed trends; i.e. a trend
16 in the SAM should result in a trend in the wind speed. Notably, this observational SAM constraint
17 suggests that the strong trends in the MERRA2 reanalysis in austral winter and spring over the period of
18 maximum ozone depletion may be spurious, as they are accompanied by strong SAM index trends that
19 are not present in the observations (Fig. 5, panels d) and e)). Additionally, in autumn, the ERA5 and
20 JRA3Q time series appear to underestimate the SAM trend during 1980-1999, while MERRA2
21 overestimates it over the course of the whole time series, suggesting possible corresponding
22 overestimations/underestimations of the respective wind trends. In summer, the observational constraint
23 appears to have less power, as the reanalyses show a variety of wind speed trends while all representing
24 the SAM relatively accurately (Fig. 5b).

Deleted: R1 and

Deleted: reanalyses

Deleted: R1 and

Deleted: overestimate

26 3.1.5 Reanalysis intercomparison summary

27
28 All four reanalyses show relative coherence in mean wind speed climatology, distribution, jet
29 position and jet seasonal cycle. Moderate differences in extreme winds, interannual variability and jet
30 latitudinal trend are observed. Reanalyses tend to exhibit systematic biases in capturing the tails of the
31 observational wind distribution (i.e., they overestimate extremely low winds and overestimate extremely
32 high winds relative to observations; see summary of previous in-situ evaluation in Methods). Thus, it is
33 plausible that ERA5, which has the lowest extreme low winds, is the best representation of the low tail of
34 the distribution, while JRA3Q may be the best at representing high winds.

35
36 The most important differences between the reanalysis products are in the representation of the
37 wind speed trends over the time period 1980-2019. MERRA2 has considerably stronger trends than
38 ERA5 and JRA3Q, and moreover these trends persist year round. To evaluate which trends seem more
39 plausible, we consider an observational constraint in the form of the observational SAM index. MERRA2
40 appears to overestimate SAM index trends, especially in austral winter and spring, suggesting that their
41 strong wind trends in these time periods may be spurious. This provides additional rationale for using
42 ERA5 as the baseline reanalysis against which to evaluate the UKESM1 model at southern high latitudes.

Deleted: R1 and

Deleted: have

Deleted: R1 and

Deleted: appear

1 3.2 UKESM1 evaluation against reanalyses

2

3 We next evaluate the main Southern Ocean wind features in UKESM1 against our default
4 reanalysis, ERA5, to assess the model's suitability for answering scientific questions related to changing
5 wind patterns in this region. The UKESM1 model has an overall year-average positive bias of $\sim 0.3 \text{ m s}^{-1}$
6 relative to the ERA5 climatology in the band between 40°S and 60°S (Fig. 1), which does not vary
7 substantially by season (Table 4). UKESM1 underestimates winds at high latitudes compared to ERA5
8 while showing the strongest positive bias over the mid-latitude open ocean. However, on average, the
9 UKESM1 distribution is similar to that of ERA5, though slightly shifted towards higher winds than the
10 reanalyses in the middle of the distribution (Fig. 2). Extreme low (5th percentile) winds are nearly
11 identical to ERA5 values, while the highest winds are somewhat higher ($\sim 0.3 \text{ m s}^{-1}$, ST2).
12 Interannual variability in UKESM1 is comparable to that of ERA5, with an average ratio of 0.8
13 (IAV_{UKESM1}/IAV_{ERA5}) for annual values (ST1).

14

15 UKESM1 has similar decadal trends in wind speed to ERA5, with strongest acceleration in
16 austral summer (0.04 and $0.06 \text{ s}^{-1} \text{ dec}^{-1}$ for UKESM1 and ERA5, respectively, over 1980-2019, as
17 compared to 0.11 and $0.14 \text{ s}^{-1} \text{ dec}^{-1}$ over 1980-1999; Table 5). The general pattern of significant wind
18 trend strength peaking in austral summer is consistent between the model and ERA5. Spatial patterns
19 (significant acceleration occurring mostly in the 40 - 60°S band, and more visible in austral summer, Fig.
20 3), are also reproduced by the model.

21

22 The UKESM1 jet position is similar to that of ERA5 (Fig. 4, ST3) and accurately captures the
23 observed seasonal cycle in position (most southerly in MAM, most northerly in SON). The majority of
24 the temporal variability in jet location is a result of natural variability (as opposed to from a forced signal,
25 such as ozone depletion). We would therefore not expect a free running model like UKESM1 to capture
26 the temporal evolution of this variability. (We would, however, expect the model to capture any
27 externally forced, longer-term trends.) The magnitude of variability in the jet location is comparable
28 between UKESM1 and ERA5, as seen in the zonally-varying standard deviation for 1980-2019 (Fig. 4).
29 The UKESM1 trends in the poleward position of the jet are much stronger than in the reanalyses over the
30 whole time period ($\sim 1.8x$ ERA5 in the annual mean, $\sim 1.7x$ ERA5 in the DJF mean, Sup. Table ST4), a
31 feature that is amplified in the period of maximum ozone loss (1980-1999, ST4). This feature is likely due
32 to Antarctic ozone loss being higher in UKESM1 than observed (Keeble et al., 2021, and section 3.3 of
33 this paper).

34

35 The variability and trend in the SAM index is comparable to that of ERA5 and observations (Fig.
36 5, Sup. Table ST5). As in the case with the jet position, UKESM1 does not reproduce the time variation
37 in the SAM index seen in the reanalyses because of the significant role played by natural variability.
38 Trends in the SAM index during 1980-1999 are much (up to $\sim 3x$) stronger than those during 1980-2019,
39 both in the ERA5 reanalysis and in UKESM1. Similar to ERA5 and JRA3Q, in both time periods,
40 UKESM1 has strongest, most significant SAM trends during DJF and MAM, accompanied by stronger
41 wind acceleration in those seasons (Fig. 5). In particular, UKESM1 accurately captures the SAM trend in
42 DJF, including the differential trends between the full period and the shorter 1980-1999 period. This

1 differential response is less accurately captured in MAM, although the SAM trend for the full period is
2 well simulated.

3
4 Overall, the UKESM1 Southern Ocean wind climatology and distribution, including extremes, as
5 well as decadal trends wind speeds and the SAM index, are remarkably consistent with that of ERA5;
6 however, mean wind speeds are somewhat higher in the model. The mean jet position is also reproduced,
7 though UKESM1 demonstrates stronger poleward movement than all reanalyses.

8 3.3 UKESM1 in the context of CMIP6 models with interactive 9 chemistry

10 The CMIP6 models with interactive ozone show substantial spread (~1 m/s) in their
11 representation of the mean wind speed over the Southern Ocean (Fig. 6 a)-c)). All models, except GISS-
12 E2-1-G and EC-Earth3-AerChem, overestimate the climatological mean wind speed with respect to
13 ERA5, with UKESM1 within the spread of other models. Agreement across the models in wind speed
14 trends is also generally weak. Most models (except MRI-ESM2-0) show larger DJF trends for 1980–
15 1999 than 1980–2019, with GFDL_ESM4 and UKESM1 being notably accurate for both time periods.
16 Furthermore, all models except UKESM1 underestimate the JJA wind speed trend during 1980-1999,
17 though trends in this season are not expected to be strongly forced by ozone trends. UKESM1 performs
18 well in capturing wind trends compared to other CMIP6 models, reproducing relatively well the wind
19 speed trends seen in ERA5, for DJF and JJA, for both time periods considered, including differences
20 between the two periods.

21
22 On average, the models overestimate TCO depletion (for the area 70°S- 90°S), though with
23 considerable inter-model variation: MRI-ESM2-0 underestimates depletion, GFDL-ESM4, EC-Earth3-
24 AerChem, and CNRM-ESM2-1 are quite accurate, while UKESM1 and GISS-E2-1-G substantially
25 overestimate ozone loss across all seasons and time periods (Fig. 6 lower panels, Fig. S1). Though
26 stronger ozone depletion is expected to correlate with larger wind speed trends, this relationship holds
27 only partially in austral summer. GISS-E2-1-G substantially overestimates ozone depletion in 1980–1999,
28 but fails to produce a correspondingly strong wind speed trend, even in summer. Note that GISS-E2-1-G
29 is known to have biases in simulating the recovery of volcanic eruptions, which amplifies the influence of
30 ODS on TCO, so while we include it in the comparison of models, we keep this bias in mind when
31 interpreting results. Similarly, CNRM-ESM2-1 reasonably captures the observed ozone loss but
32 underestimates wind speed trends. GFDL-ESM4 best matches both metrics in austral summer but not
33 winter. While the strongest ozone loss typically occurs in austral spring (SON), inter-model TCO trends
34 and their relationship with wind speed trends are consistent whether wind speed trends in DJF are plotted
35 relative to TCO trends in SON or DJF (Fig. S2). To maintain consistency with the rest of this study, we
36 therefore present only DJF results in Figure 6.

37
38 UKESM1 overestimates ozone depletion, primarily in the 1980-1999 period, but accurately
39 reproduces the ERA5 wind speed trends for both periods, suggesting the dynamical link between
40 Antarctic ozone loss and near-surface westerlies is weaker in UKESM1 than in reality. The weak TCO–
41 wind relationship across models in austral winter (JJA) for 1980–1999 indicates observed wind increases
42 during this period are likely not ozone-driven. As in the reanalyses (Fig. 5), SAM trends correlate

1 strongly with wind speed trends across CMIP6 models (Fig. S3), due to the mechanistic link between
2 these metrics. While acknowledging recent trends in Antarctic TCO are larger than observed in
3 UKESM1, our analysis of wind speed and SAM trends, and the implied response to ozone forcing,
4 suggests UKESM1 is suitable for studying (and attributing) past and future changes in Southern Ocean
5 winds, and by extension the drivers of changes in Southern ocean carbon uptake, as carried out in our
6 earlier study (Jarníková et al., 2025).
7

8 3.4 Attribution of wind speed trends to GHG and ozone forcing with 9 UKESM1

10 Finally, we use UKESM1 to attribute past, and potential future, changes in the Southern Ocean
11 wind regimes to either stratospheric ozone changes or greenhouse gas forcing. We do this by combining
12 two ODS scenarios (ODS-HIST and ODS-1950) with two CMIP6 emission scenarios: SSP 1-2.6 and SSP
13 3-7.0 for the period 1950 to 2100 (see Methods). For each experiment, we show all three model ensemble
14 members, as well as the ensemble mean.
15

16 Over the full time period, winds increase in all seasons and scenarios (Fig. 7). The most
17 prominent feature in the time series is a sharp increase in wind speeds in the scenario with historical
18 ozone depletion (OZONE-HIST), approximately during 1980-2000, occurring predominantly in austral
19 summer (DJF) and to a lesser extent in austral spring (SON; Fig. 7). During the same time period, we
20 observe no wind increase in the scenario without ozone depletion – in fact, a minor decrease is seen (Fig.
21 7, Sup. Table ST6). We first partition the attribution of the observed historical wind changes to
22 greenhouse gas emissions and ozone forcing, comparing them with those seen in ERA5, and then
23 consider potential changes over the 21st century, again attributing changes to ozone forcing and/or
24 greenhouse gas emissions.
25

26 UKESM1 exhibits the same positive relationship between the trend in the SAM index and wind
27 speed that was seen in the reanalyses. Wind acceleration (deceleration) corresponds to a positive
28 (negative trend) in the SAM index in all ensemble members of the two experiments over the historical
29 period (Fig. 8). Though absolute wind speeds are higher in UKESM1 than in ERA5, the magnitude of the
30 trends in the SAM index and wind speeds, for both periods, are very similar in both datasets for all
31 seasons (Fig. 8, Table 5), suggesting the forced signal of ozone depletion in UKESM1 captures real-world
32 dynamics.
33

34 The key role of ozone in driving the historical SAM and wind speed changes becomes further
35 apparent when comparing the OZONE-HIST and OZONE-1950 runs: all three ensemble members of the
36 OZONE-1950 experiment show a much weaker wind speed trend over the whole time period (Fig. 8,
37 filled green squares) and a *negative* trend over the period of maximum ozone depletion (empty green
38 squares). Meanwhile, the OZONE-HIST runs (black squares) agree with both ERA5 and the
39 observational SAM index over both the whole time period (filled black squares) and the period of
40 maximum ozone depletion (empty black squares), with the differential response in DJF between these two
41 periods, in both the SAM index and wind speeds, particularly well captured.
42

1 The general congruence of the historical UKESM1 runs with ERA5 suggests that UKESM1
2 captures both the sensitivity of the wind and SAM response to the time-varying trend in the ozone
3 forcing, as well as the seasonality of this response. Both wind and SAM trends are stronger over the
4 period of maximum ozone depletion than over the full time period, and changes in the austral summer
5 mean are larger than in the annual mean, with minimal changes in austral winter.
6

7 We next attribute the wind changes throughout the whole simulation period 1950-2100 to either
8 ozone or GHG forcing, subdividing the time series into three 50-year periods. In the latter half of the 20th
9 century (1950-1999; Table 6), winds accelerated in all seasons but most strongly in DJF, followed by
10 SON. This acceleration was driven entirely by ozone loss; winds actually slowed down slightly over
11 1950-2000 in the OZONE-1950 run, in all seasons except SON (Table 6).
12

13 In the first half of the 21st century (2000-2049), in the realistic (ODS-HIST) runs, winds stay
14 approximately constant in most seasons in both SSP scenarios, though they increase substantially in
15 austral winter (Table 6). This lack of trend is due to the (accelerating) GHG effect countering the
16 (decelerating) ozone recovery effect, which can be seen when decomposing the overall signal into the two
17 forcing factors (Table 6). Though the acceleration due to GHG is stronger under the higher-emissions
18 (SSP 3-7.0) scenario, the effect of ozone recovery is also stronger under this scenario and the two effects
19 cancel out (Table 6). The sensitivity of ozone recovery to SSP pathway reflects the interaction of other
20 GHGs (e.g. CH₄ and N₂O) with ozone chemistry. SSP 3-7.0 has higher CH₄ concentrations in the future
21 than SSP 1-2.6, and UKESM1 has a significant ozone response to increasing CH₄ (see figure 14 in Zeng
22 et al., 2022). Ozone recovery will therefore be accelerated in SSP 3-7.0 relative to SSP 1-2.6, with a
23 concomitant stronger forcing of surface winds. The overall acceleration seen in austral winter (JJA) is a
24 result of the different seasonalities of the two forcing factors: while ozone depletion and recovery is
25 predominantly a summer phenomenon, these simulations show that GHG-driven wind increase is
26 strongest in winter, resulting in an unbalanced GHG forcing of wind trends during austral winter. Using a
27 regression analysis of CMIP6 simulations, Morgenstern (2021) finds a similar seasonality in CO₂ forcing
28 of the SAM. Simpkins and Karpechko (2012) also see this strong winter increase in the SAM, which is
29 related to the winds, also concluding that the difference is due to the lack of ozone recovery in winter.
30

31 Finally, in the latter half of the 21st century (2050-2099), the simulations show a difference in
32 wind trends based on GHG emission level. Under SSP 1-2.6 with historical ozone recovery, wind speed
33 stagnates in all seasons, while under SSP 3-7.0, winds accelerate in austral winter and spring, both with
34 and without ozone depletion (Fig. 7). However, these trends are typically not statistically significant, due
35 to large interannual variability. Analogous results were found by Simpkins and Karpechko (2012), who
36 noted the dependence of SAM trends on emissions scenario and also noted the lack of statistical
37 significance.
38

39 4. Discussion and Conclusions

40 The main aims of this work were to evaluate the representation of recent-past Southern Ocean
41 winds in the most commonly used contemporary reanalysis products, in a free-running Earth system
42

Formatted: Outline numbered + Level: 1 + Numbering
Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at:
0.63 cm + Indent at: 1.27 cm

1 model (UKESM1) and other CMIP6 models with interactive ozone, and finally to use UKESM1 to
2 attribute the drivers of changes in winds over the recent past and through to 2100. The initial
3 intercomparison of the reanalyses shows that, while the general features of the Southern Ocean winds
4 (mean, distribution, jet location, and extremes) are relatively congruent, the reanalyses vary considerably
5 in wind speed trends. In all reanalyses, trends are strongest in the period of maximum ozone depletion
6 (1980-1999), and comparatively strongest in austral summer, consistent with the existing understanding
7 of surface wind forcing by ozone loss, which is primarily a summer phenomenon. However, trends in
8 MERRA2 are comparatively much stronger and tend to persist year-round, while trends in ERA5 and
9 JRA3Q are weaker and concentrated in austral summer.

Deleted: and R1

10
11 We assess which trend representation is more realistic using a novel observational constraint: we
12 compare trends in the observational SAM index to their representation in the reanalyses, and then relate
13 them to wind speed trends. Because a positive (negative) SAM index trend is mechanistically linked to a
14 positive (negative) wind trend, performance in one metric can provide information about performance in
15 the other. Applying this constraint suggests the large winter trends seen in winter and spring in the
16 MERRA2 reanalysis may be spurious, as they are accompanied by strong SAM trends not seen in the
17 observational record. Spuriously large trends in the previously commonly used NCEP-NCAR reanalysis
18 were also observed by Morgenstern (2021); here we suggest that this problem persists in MERRA2. Thus,
19 we conclude that ERA5 and JRA3Q are best suited for studies of wind and SAM changes in this region,
20 as well as for forcing ocean models.

Deleted: and R1 reanalyses

Deleted: R1

21
22 UKESM1 winds and SAM index evaluate well against our baseline reanalysis (ERA5). Notably,
23 UKESM1 performs well in reproducing the intensity distribution of daily wind speeds seen in ERA5,
24 including the high wind speed tail of the distribution, though the overall mean of the distribution is
25 somewhat higher in UKESM1 ($\sim 0.3 \text{ m s}^{-1}$). UKESM1 also reproduces the ERA5 historical decadal trends
26 in both wind speed and the SAM index, including the differential response between the 1980-2000 period
27 and the full 40 year period, indicating a realistic sensitivity of winds to ozone changes. The seasonality of
28 the SAM and wind speed trends is also well reproduced, with strongest trends in austral summer.
29 However, the poleward progression of the jet latitude is stronger in UKESM1 than in any of the
30 reanalyses, somewhat in contrast to the correctly reproduced behaviour of the other metrics.

31
32 We find that UKESM1 represents wind trends well relative to similar CMIP6 models with
33 interactive chemistry (section 3.3), but highlight again that the accurate wind speed trends in UKESM1
34 are realized in combination with too strong ozone decline, particularly in the period 1980-1990. We note
35 that the relationship between TCO depletion and wind trends in the CMIP6 models holds only partially in
36 austral summer. No relationship is observable in austral winter, consistent with the understood low role of
37 ozone in forcing wind changes in this season.

38
39 Using future projections made with UKESM1, with and without ozone depletion, we attribute the
40 observed changes to the relative influence of ozone depletion and greenhouse gas emissions, both
41 historically over 1980-2020 and as projected by the model to the end of the 21st century. The UKESM1
42 simulations show the observed acceleration of winds in the latter half of the 20th century is entirely
43 attributable to ozone depletion. In the first half of the 21st century, summer winds stagnate, as the
44 decrease of wind speeds due to ozone recovery competes with wind speed increase due to GHG

1 emissions. In contrast, wind speeds increase in austral winter and spring, as the GHG effect outcompetes
2 the ozone recovery effect, which is predominantly a summer phenomenon. In the latter half of the 21st
3 century, a more clear bifurcation between wind speed trends under the low and high SSP scenario is
4 visible, especially in austral winter and spring. Winds accelerate more under the high SSP scenario, while
5 stagnating under the low SSP scenario. In this time period, the role of ozone recovery is minor.
6

7 We show that ozone depletion overwhelmingly drove the observed summer wind acceleration in
8 the latter half of the 20th century, and we expect this trend to reverse in the first half of the 21st century.
9 By the second half of the 21st century, greenhouse gas loading is the main factor driving wind changes,
10 with more emissions leading to more acceleration, but it is strongly active only in austral winter and
11 spring. Thus, we demonstrate a shift in the dynamical controls on the wind behaviour over the Southern
12 Ocean from ozone in the latter half of the 20th century to greenhouse gas emissions in the latter half of
13 the 21st century.
14

15 This control shift is consistent with the general understanding. As in Barnes et al. (2014), we find
16 historical wind speed trends due to ozone depletion are stronger than greenhouse gas driven trends seen in
17 the twenty first century. Similar to Barnes et al (2014), Gerber and Son (2014), McLandress et al.
18 (2011), and Simpkins and Karpechko (2012), we observe a cancellation of effects in the first half of the
19 twenty first century, as ozone recovery competes with greenhouse gas emissions, leading to only weak
20 wind trends in this period. We also reproduce the dependence of future trends on greenhouse gas emission
21 strength seen in Simpkins and Karpechko (2012) and other studies. Our work, focusing on near-surface
22 winds, extends this understanding by providing an estimate of the changing wind trends in response to the
23 shift in dominant forcing. We do this using an Earth system model (UKESM1), with full-atmosphere
24 interactive chemistry, allowing a large set of interactions and feedbacks to be simulated; chemically
25 between a range of GHGs and stratospheric ozone, between both sets of gases and atmospheric dynamics,
26 and between the atmosphere and the underlying ocean and sea ice. The ability of UKESM1 to represent
27 these interactions, and how they shape surface winds over the Southern Ocean, was validated against both
28 reanalyses and an observational SAM index and further contextualized against other CMIP6 models with
29 interactive ozone chemistry. Our analysis suggests UKESM1 is suitable for investigating links between
30 ozone loss and recovery, GHG scenarios, and Southern Ocean uptake of heat and carbon.
31

1 Code and data availability

2 ~~Analysis code is provided~~ at <https://github.com/tjarnikova/windEval>.

3

Deleted: Code and data are provided on DRYAD: xx. For convenience, analysis code is also

Formatted: Not Highlight

4 Author contribution

5 Conceptualization: T.J., C.J., and C.L.Q. Methodology: C.J., T.J., S.R., and C.L.Q. Investigation: T.J.
6 Visualization: T.J. Funding acquisition: C.L.Q. and C.J. Project administration: C.L.Q. Supervision: C.J.
7 and C.L.Q. Writing—original draft: T.J. Formal analysis: T.J. Software: T.J. and S.R. Data curation: T.J.
8 Validation: T.J., S.R., and C.L.Q. Writing—review and editing: C.J., T.J., and C.L.Q.

9 Competing interests

10 The authors declare that they have no conflict of interest.

11 Acknowledgements


12 We thank all people who contributed to the development of the UKESM1 model and the atmospheric
13 reanalyses used in this analysis, and the Research and Specialist Computing Support service of the High
14 Performance Computing Cluster at the University of East Anglia. We would like to thank Bodeker
15 Scientific, funded by the New Zealand Deep South National Science Challenge, for providing the
16 combined NIWA-BS total column ozone database and James Keeble (University of Lancaster) for advice
17 on using this data set.
18

19 Funding

20 U. K. Natural Environment Research Council CELOS project (grant NE/T01086X/1) (CJ, CLQ)
21 U. K. Natural Environment Research Council TerraFIRMA: Future Impacts, Risks and Mitigation
22 Actions in a changing Earth System project, (grant NE/W004895/1) (CJ, SR)
23 U.K. Royal Society (grant RSRP\R\241002) (CLQ)
24
25

1 Figures

2 Fig. 1




3
4

5 **Fig. 1: Climatological wind speed for 1980-2019.** For ERA5, the climatological wind speed for the full
6 year and austral summer (DJF) and winter (JJA) is shown. For the other two reanalyses and UKESM1,
7 differences from ERA5 are shown as [product x - ERA5]; i.e. positive (red) values indicate higher winds
8 than ERA5. See also Table 4.

Moved (insertion) [1]

Moved up [1]: Figures

Deleted: 1




Deleted:


Deleted: 3

Deleted: 2

1 **Fig. 2**



Deleted: []




Deleted:


2
3
4
5
6
7
8
9
10
11
12

Fig. 2: Summary statistics for wind distributions in four reanalysis products and UKESM1. a-b): mean 10-m wind speed trends relative to climatology for a) full year and b) DJF only. Thick lines have been smoothed by a three-point running filter. c-d): windspeed frequency distribution (100 bins, 0-20 $m s^{-1}$) for c) full year and d) DJF, with high tails shown in inset. e-h) Trends in mean and extreme winds for full year and DJF vs. climatological means, with colours the same as in panel a). Filled symbols represent 1980-2019, while open symbols represent 1980-1999; circles represent reanalyses while squares represent UKESM1. Colours from panel a) are repeated throughout the figure. Summary statistics shown in panels e)-h) are also given in Tables 4 and 5. All figure statistics are calculated for daily winds at $1^\circ \times 1^\circ$ resolution, $40^\circ S$ - $60^\circ S$.


1 Fig. 3



2
3 **Fig. 3: Decadal trends in 10-m wind speed (1980-2019).** Hatching shows trends significant at the 95%
4 confidence level. See also Table 5.
5
6




1 Fig. 4





2

3 **Fig. 4: Summary statistics for the wind jet.** The wind jet is calculated as the location of the maximum of
 4 the *u*-component of the 10-m wind speed between 30°S and 70°S (see Methods). a): the climatological
 5 mean annual jet position, with one standard deviation of the annual mean shown for ERA5 and UKESM1.
 6 b): The zonal mean of the austral summer jet position. Thick lines indicate smoothing by a 3-point
 7 running mean filter. c-d): Trends in the annual (c) and summer (d) jet position mean vs. climatological
 8 means. Colours from panel a) are repeated throughout the figure. See also Sup. Tables ST3 and ST4.



Deleted:

1 Fig. 5




Deleted:

Deleted: S5

Formatted: Normal

2
3 **Fig. 5: Summary statistics for the SAM index.** a) Timeseries of the austral summer (DJF) SAM index. To
4 highlight coherence of interannual variability, no running-mean filter has been applied. b-e) Trends in
5 the wind speed vs trends in the SAM index, subdivided by season. Filled symbols represent 1980-2019,
6 while open symbols represent 1980-1999. Vertical lines show the observational SAM index (full line
7 represents 1980-2019, dotted line represents 1980-1999). Colours from panel a) are repeated throughout
8 the figure. See also Sup. Table [S75](#).


1 Fig. 6.



2
3
4
5
6
7
8

3 **Fig. 6: 10-m wind speed, decadal wind speed trend, and decadal TCO trend (70°S- 90°S) for CMIP**
4 **models with interactive chemistry.** For each model, one ensemble member is used (see Table 2). *Wind*
5 *speeds and trends are calculated from daily winds at 1°x1° resolution, 40°S- 60°S. Note the caveats*
6 *regarding TCO trends in GISS-E2-I-G discussed in section 3.4.*

Deleted:




Deleted:

Formatted: Font: Italic

Formatted: Font: Not Bold

1

2 Fig. 7



3


4

5

6

7

Fig. 7: 10-m wind speed evolution from 1950 to 2100 in the OZONE-1950 and OZONE-HIST experiments. a) Full year, b-e): seasonally subdivided means. All means are calculated for daily winds at $1^\circ \times 1^\circ$ resolution, 40°S - 60°S . For each experiment, three ensemble members are shown, with the ensemble mean shown in bold. All lines are smoothed with a 10-year running filter. See also Table 6.




Deleted:

Formatted: Normal

Deleted: 6Table 4. 1

1 Fig. 8



2
3
4
5
6
7
8
9

Fig. 8: Trends in mean wind speed vs. trends in the SAM index for the OZONE-1950 and OZONE-HIST experiments. ERA5 is shown for comparison (blue circles), and the observational SAM index is shown as vertical lines (full for 1980-2019, dotted for 1980-1999). For the OZONE-1950 and OZONE-HIST experiments, individual ensemble members are shown as smaller squares, while the ensemble mean is shown as a larger square. As in Figs. 2,4, and 5, filled symbols represent 1980-2019, while open symbols represent 1980-1999.

1

2 Tables

3 Table 1

4

Name	ERA5	JRA3Q	MERRA2	
Institution	ECMWF	JMA	NASA GMAO	
Period coverage	1940 to present	1947 to present	1980 to present	
Time resolution	1-hr	6-hr	1-hr	
Horizontal grid (lat. × lon.)	0.25° × 0.25°	0.37° × 0.37°	0.5° × 0.625°	
Reference	Hersbach et al., 2020	Kosaka et al., 2024	Molod et al., 2015	

Formatted: Heading 2

Deleted: R1 (NCEP-NCAR)

Deleted Cells

Formatted Table

Deleted: NOAA/NCEP and NCAR

Deleted: 1948 to present

Deleted: 6-hr

Deleted: 1.875° × 2°

Deleted: Kalnay et al., 1996

5

6 Table 1: Basic characteristics of the reanalysis products used.

Formatted: Font: Not Bold, Not Italic

7

Formatted: Font: Times New Roman

Formatted: Font: 16 pt, Not Bold, Not Italic

1 Table 2

2

Model	Reference
UKESM1-0-LL	Sellar et al., 2019
CNRM-ESM2-1	Séférian et al., 2019
GISS-E2-1-G	Kelley et al., 2020
MRI-ESM2-0	Yukimoto, Kawai, et al., 2019
GFDL-ESM4	Dunne et al., 2020
EC-Earth3-AerChem	(Van Noije et al., 2021)

Formatted Table

3

4 **Table 2: CMIP6 models with interactive chemistry used in the intercomparison.** Only models with both
5 interactive chemistry and 10-m wind fields available at (at least) daily resolution are used. One ensemble
6 member is used per model.

1 Table 3

2
3

Experiment	Ozone forcing	GHG forcing
OZONE-HIST SSP 1-2.6	Standard CMIP-6 surface mixing ratios for ODS (historical to 2015, followed by specific SSP pathway projection) (Meinshausen et al., 2017)	SSP 1-2.6 (Gidden et al., 2019)
OZONE-HIST SSP 3-7.0	as above	SSP 3-7.0 (Gidden et al., 2019)
OZONE-1950 SSP 1-2.6	ODS surface mixing ratios fixed at 1950 values from 1950 onwards	SSP 1-2.6 (Gidden et al., 2019)
OZONE-1950 SSP 3-7.0	as above	SSP 3-7.0 (Gidden et al., 2019)

Formatted Table

4
5 *Table 3: Summary of experiments with description of ozone and GHG forcing.*

1 Table 4

climatological 10-m wind speed, m s ⁻¹ , 1980-2019					
	ERA5		MERRA 2	JRA3Q	UKESM 1
full year	9.05	▼	-0.20	0.28	0.30
DJF	8.41	▼	-0.23	0.27	0.33
MAM	9.13	▼	-0.28	0.24	0.36
JJA	9.5	▼	-0.13	0.31	0.19
SON	9.16	▼	-0.18	0.31	0.32

Formatted Table

Deleted: R1

Deleted Cells

Deleted: -0.05

Deleted: -0.04

Deleted: -0.02

Deleted: -0.08

Deleted: -0.07

2
3
4
5

Table 4: Climatological 10-m wind speed (1980-2019). Seasonally subdivided open-ocean wind speed between 40°S and 60° S for ERA5 , and differences from ERA5 for the other reanalyses and UKESM1.

1 Table 5

trend in 10-m wind speed, m s ⁻¹ decade ⁻¹ , 1980-2019					
	ERA5	▼	MERRA2	JRA3Q	UKESM1
full year	0.04	▼	0.10	0.04	0.03
DJF	0.06	▼	0.12	0.05	0.04
MAM	0.05	▼	0.13	0.06	0.02
JJA	0.02	▼	0.08	0.03	0.02
SON	0.01	▼	0.06	0.02	0.02
trend in 10-m wind speed, m s ⁻¹ decade ⁻¹ , 1980-1999					
full year	0.09	▲	0.26	0.12	0.06
DJF	0.14	▲	0.22	0.14	0.11
MAM	0.09	▼	0.26	0.14	0.02
JJA	0.08	▼	0.28	0.12	0.08
SON	0.07	▼	0.25	0.09	0.04

2

3 **Table 5: Decadal trends in 10-m wind speed.** Seasonally subdivided trends in open-ocean wind speed
 4 between 40°S and 60° S are shown for 1980-2019 and 1980-1999. Trends significant at the 5% level are
 5 given in bold. ▼ ▲

Formatted Table

Deleted: R1

Deleted Cells

Deleted: 0.14

Deleted: 0.17

Deleted: 0.17

Deleted: 0.13

Deleted: 0.11

Deleted: 0.32

Deleted Cells

Deleted: 0.35

Deleted: 0.31

Deleted: 0.30

Deleted: 0.31

Formatted: Normal

Deleted: ¶

Formatted: Font: 10 pt, Italic

1 Table 6 - attribution 1950-2099

2
3

1950-1999	GHG + ozone	GHG only	ozone only			
FY	0.03	-0.01	0.04			
DJF	0.06	-0.02	0.07			
MAM	0.01	0.00	0.01			
JJA	0.02	-0.01	0.03			
SON	0.04	0.01	0.03			
	SSP 1-2.6			SSP 3-7.0		
2000-2049	GHG + ozone	GHG only	ozone only	GHG + ozone	GHG only	ozone only
FY	0.01	0.02	-0.01	0.01	0.03	-0.02
DJF	0.00	0.01	-0.01	0.00	0.02	-0.03
MAM	0.00	0.01	-0.01	0.00	0.02	-0.02
JJA	0.03	0.03	0.00	0.03	0.04	-0.01
SON	0.02	0.03	-0.01	0.02	0.04	-0.02
	SSP 1-2.6			SSP 3-7.0		
2050-2099	GHG + ozone	GHG only	ozone only	GHG + ozone	GHG only	ozone only
FY	-0.01	0.01	-0.01	0.00	0.01	0.00

Formatted Table

DJF	-0.02	0.01	-0.03	-0.02	-0.01	-0.01
MAM	0.00	-0.01	0.00	-0.02	-0.01	-0.01
JJA	-0.01	0.01	-0.02	0.02	0.02	0.00
SON	0.00	0.01	-0.01	0.03	0.02	0.01

1
2
3
4
5
6
7
8

Table 6: Attribution of 10-m wind trends ($m s^{-1} dec^{-1}$) to GHG and ozone forcing in three 50-year time-periods in the UKESM1 model runs. Values for GHG + ozone refer to the ensemble mean trend of the OZONE-HIST run, while the GHG only column refers to the OZONE-1950 run. The effect of ozone only is then obtained by subtracting OZONE-1950 from OZONE-HIST. Statistically significant trends are given in bold.

1 Bibliography

- 2
3 Arblaster, J. M., Meehl, G. A., & Karoly, D. J. (2011). Future climate change in the Southern
4 Hemisphere: Competing effects of ozone and greenhouse gases: SH CLIMATE CHANGE-
5 OZONE VERSUS GHGS. *Geophysical Research Letters*, 38(2), n/a-n/a.
6 <https://doi.org/10.1029/2010GL045384>
- 7 Archibald, A. T., O'Connor, F. M., Abraham, N. L., Archer-Nicholls, S., Chipperfield, M. P., Dalvi, M.,
8 Folberth, G. A., Dennison, F., Dhomse, S. S., Griffiths, P. T., Hardacre, C., Hewitt, A. J., Hill, R.
9 S., Johnson, C. E., Keeble, J., Köhler, M. O., Morgenstern, O., Mulcahy, J. P., Ordóñez, C., ...
10 Zeng, G. (2020). Description and evaluation of the UKCA stratosphere–troposphere chemistry
11 scheme (StratTrop v1.0) implemented in UKESM1. *Geoscientific Model Development*, 13(3),
12 1223–1266. <https://doi.org/10.5194/gmd-13-1223-2020>
- 13 Bishop, S. P., Gent, P. R., Bryan, F. O., Thompson, A. F., Long, M. C., & Abernathy, R. (2016).
14 Southern Ocean Overturning Compensation in an Eddy-Resolving Climate Simulation. *Journal of*
15 *Physical Oceanography*, 46(5), 1575–1592. <https://doi.org/10.1175/JPO-D-15-0177.1>
- 16 Bodeker, G. E., Nitzbon, J., Tradowsky, J. S., Kremser, S., Schwertheim, A., & Lewis, J. (2021). A global
17 total column ozone climate data record. *Earth System Science Data*, 13(8), 3885–3906.
18 <https://doi.org/10.5194/essd-13-3885-2021>
- 19 Bracegirdle, T. J., Shuckburgh, E., Sallee, J.-B., Wang, Z., Meijers, A. J. S., Bruneau, N., Phillips, T., &
20 Wilcox, L. J. (2013). Assessment of surface winds over the Atlantic, Indian, and Pacific Ocean
21 sectors of the Southern Ocean in CMIP5 models: Historical bias, forcing response, and state
22 dependence: CMIP5 SOUTHERN OCEAN SURFACE WINDS. *Journal of Geophysical*
23 *Research: Atmospheres*, 118(2), 547–562. <https://doi.org/10.1002/jgrd.50153>
- 24 Caton Harrison, T., Biri, S., Bracegirdle, T. J., King, J. C., Kent, E. C., Vignon, É., & Turner, J. (2022).
25 *Reanalysis representation of low-level winds in the Antarctic near-coastal region* [Preprint].
26 Dynamical processes in polar regions, incl. polar–midlatitude interactions.
27 <https://doi.org/10.5194/egusphere-2022-693>
- 28 Dong, X., Wang, Y., Hou, S., Ding, M., Yin, B., & Zhang, Y. (2020). Robustness of the Recent Global
29 Atmospheric Reanalyses for Antarctic Near-Surface Wind Speed Climatology. *Journal of*
30 *Climate*, 33(10), 4027–4043. <https://doi.org/10.1175/JCLI-D-19-0648.1>
- 31 Dunne, J. P., Horowitz, L. W., Adcroft, A. J., Ginoux, P., Held, I. M., John, J. G., Krasting, J. P.,
32 Malyshev, S., Naik, V., Paulot, F., Shevliakova, E., Stock, C. A., Zadeh, N., Balaji, V., Blanton,
33 C., Dunne, K. A., Dupuis, C., Durachta, J., Dussin, R., ... Zhao, M. (2020). The GFDL Earth
34 System Model Version 4.1 (GFDL-ESM 4.1): Overall Coupled Model Description and
35 Simulation Characteristics. *Journal of Advances in Modeling Earth Systems*, 12(11),
36 e2019MS002015. <https://doi.org/10.1029/2019MS002015>
- 37 Eyring, V., Arblaster, J. M., Cionni, I., Sedláček, J., Perlwitz, J., Young, P. J., Bekki, S., Bergmann, D.,
38 Cameron-Smith, P., Collins, W. J., Faluvegi, G., Gottschaldt, K. -D., Horowitz, L. W., Kinnison,
39 D. E., Lamarque, J. -F., Marsh, D. R., Saint-Martin, D., Shindell, D. T., Sudo, K., ... Watanabe,
40 S. (2013). Long-term ozone changes and associated climate impacts in CMIP5 simulations.
41 *Journal of Geophysical Research: Atmospheres*, 118(10), 5029–5060.
42 <https://doi.org/10.1002/jgrd.50316>
- 43 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2016).
44 Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design
45 and organization. *Geoscientific Model Development*, 9(5), 1937–1958.
46 <https://doi.org/10.5194/gmd-9-1937-2016>
- 47 Farman, J. C., Gardiner, B. G., & Shanklin, J. D. (1985). *Large losses of total ozone in Antarctica reveal*
48 *seasonal ClONO₂ interaction*.

Deleted: ¶

Formatted: Indent: Left: -1.27 cm

1 Fogt, R. L., & Marshall, G. J. (2020). The Southern Annular Mode: Variability, trends, and climate
2 impacts across the Southern Hemisphere. *WIREs Climate Change*, 11(4), e652.
3 <https://doi.org/10.1002/wcc.652>

4 Francis, D., Mattingly, K. S., Lhermitte, S., Temimi, M., & Heil, P. (2020). *Atmospheric extremes*
5 *triggered the biggest calving event in more than 50 years at the Amery Ice shelf in September*
6 *2019*. <https://doi.org/10.5194/tc-2020-219>

7 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J.,
8 Landschützer, P., Le Quéré, C., Lujikx, I. T., Peters, G. P., Peters, W., Pongratz, J.,
9 Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., ...
10 Zheng, B. (2023). Global Carbon Budget 2023. *Earth System Science Data*, 15(12), 5301–5369.
11 <https://doi.org/10.5194/essd-15-5301-2023>

12 Gerber, E. P., & Son, S.-W. (2014). Quantifying the Summertime Response of the Austral Jet Stream and
13 Hadley Cell to Stratospheric Ozone and Greenhouse Gases. *Journal of Climate*, 27(14), 5538–
14 5559. <https://doi.org/10.1175/JCLI-D-13-00539.1>

15 Gidden, M. J., Riahi, K., Smith, S. J., Fujimori, S., Luderer, G., Kriegler, E., Van Vuuren, D. P., Van Den
16 Berg, M., Feng, L., Klein, D., Calvin, K., Doelman, J. C., Frank, S., Fricko, O., Harmsen, M.,
17 Hasegawa, T., Havlik, P., Hilaire, J., Hoesly, R., ... Takahashi, K. (2019). Global emissions
18 pathways under different socioeconomic scenarios for use in CMIP6: A dataset of harmonized
19 emissions trajectories through the end of the century. *Geoscientific Model Development*, 12(4),
20 1443–1475. <https://doi.org/10.5194/gmd-12-1443-2019>

21 Gillett, N. P., Shiogama, H., Funke, B., Hegerl, G., Knutti, R., Matthes, K., Santer, B. D., Stone, D., &
22 Tebaldi, C. (2016). The Detection and Attribution Model Intercomparison Project (DAMIP
23 v1.0) contribution to CMIP6. *Geoscientific Model Development*, 9(10), 3685–3697.
24 <https://doi.org/10.5194/gmd-9-3685-2016>

25 Goyal, R., Sen Gupta, A., Jucker, M., & England, M. H. (2021). Historical and Projected Changes in the
26 Southern Hemisphere Surface Westerlies. *Geophysical Research Letters*, 48(4), e2020GL090849.
27 <https://doi.org/10.1029/2020GL090849>

28 Gu, Y., Katul, G. G., & Cassar, N. (2021). The Intensifying Role of High Wind Speeds on Air-Sea
29 Carbon Dioxide Exchange. *Geophysical Research Letters*, 48(5), e2020GL090713.
30 <https://doi.org/10.1029/2020GL090713>

31 Gualtieri, G. (2022). Analysing the uncertainties of reanalysis data used for wind resource assessment: A
32 critical review. *Renewable and Sustainable Energy Reviews*, 167, 112741.
33 <https://doi.org/10.1016/j.rser.2022.112741>

34 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey,
35 C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G.,
36 Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., ... Thépaut, J. (2020). The ERA5 global
37 reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999–2049.
38 <https://doi.org/10.1002/qj.3803>

39 Huguenin, M. F., Holmes, R. M., & England, M. H. (2022). Drivers and distribution of global ocean heat
40 uptake over the last half century. *Nature Communications*, 13(1), 4921.
41 <https://doi.org/10.1038/s41467-022-32540-5>

42 Jarníková, T., Quéré, C. L., Rumbold, S., & Jones, C. (2025). Decreasing importance of carbon-climate
43 feedbacks in the Southern Ocean in a warming climate. *Science Advances*, 11(20).
44 <https://doi.org/10.1126/sciadv.adr3589>

45 Jena, B., Bajish, C. C., Turner, J., Ravichandran, M., Anilkumar, N., & Kshitija, S. (2022). Record low
46 sea ice extent in the Weddell Sea, Antarctica in April/May 2019 driven by intense and explosive
47 polar cyclones. *Npj Climate and Atmospheric Science*, 5(1), 19. <https://doi.org/10.1038/s41612-022-00243-9>

48
49 Jones, R. W., Renfrew, I. A., Orr, A., Webber, B. G. M., Holland, D. M., & Lazzara, M. A. (2016).
50 Evaluation of four global reanalysis products using in situ observations in the Amundsen Sea

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

1 Embayment, Antarctica. *Journal of Geophysical Research: Atmospheres*, 121(11), 6240–6257.
2 <https://doi.org/10.1002/2015JD024680>

3 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White,
4 G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W.,
5 Janowiak, J., Mo, K. C., Ropelewski, C., ... Joseph, D. (1996). The NCEP/NCAR 40-Year
6 Reanalysis Project. *Bulletin of the American Meteorological Society*, 77(3), 437–471.
7 [https://doi.org/10.1175/1520-0477\(1996\)077%253C0437:TNYRP%253E2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077%253C0437:TNYRP%253E2.0.CO;2)

8 Keeble, J., Hassler, B., Banerjee, A., Checa-Garcia, R., Chiodo, G., Davis, S., Eyring, V., Griffiths, P. T.,
9 Morgenstern, O., Nowack, P., Zeng, G., Zhang, J., Bodeker, G., Burrows, S., Cameron-Smith, P.,
10 Cugnet, D., Danek, C., Deushi, M., Horowitz, L. W., ... Wu, T. (2021). Evaluating stratospheric
11 ozone and water vapour changes in CMIP6 models from 1850 to 2100. *Atmospheric Chemistry
12 and Physics*, 21(6), 5015–5061. <https://doi.org/10.5194/acp-21-5015-2021>

13 Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S.,
14 Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I.,
15 Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., ... Yao, M. (2020). GISS-E2.1:
16 Configurations and Climatology. *Journal of Advances in Modeling Earth Systems*, 12(8),
17 e2019MS002025. <https://doi.org/10.1029/2019MS002025>

18 Kosaka, Y., Kobayashi, S., Harada, Y., Kobayashi, C., Naoe, H., Yoshimoto, K., Harada, M., Goto, N.,
19 Chiba, J., Miyaoka, K., Sekiguchi, R., Deushi, M., Kamahori, H., Nakaegawa, T., Tanaka, T. Y.,
20 Tokuyoshi, T., Sato, Y., Matsushita, Y., & Onogi, K. (2024). The JRA-3Q Reanalysis. *Journal of
21 the Meteorological Society of Japan. Ser. II*, 102(1), 49–109. <https://doi.org/10.2151/jmsj.2024-004>

22

23 Kuhlbrodt, T., Jones, C. G., Sellar, A., Storkey, D., Blockley, E., Stringer, M., Hill, R., Graham, T.,
24 Ridley, J., Blaker, A., Calvert, D., Copsey, D., Ellis, R., Hewitt, H., Hyder, P., Ineson, S.,
25 Mulcahy, J., Siahann, A., & Walton, J. (2018). The Low-Resolution Version of HadGEM3
26 GC3.1: Development and Evaluation for Global Climate. *Journal of Advances in Modeling Earth
27 Systems*, 10(11), 2865–2888. <https://doi.org/10.1029/2018MS001370>

28 Le Quéré, C., Rödenbeck, C., Buitenhuis, E. T., Conway, T. J., Langenfelds, R., Gomez, A.,
29 Labuschagne, C., Ramonet, M., Nakazawa, T., Metzl, N., Gillett, N., & Heimann, M. (2007).
30 Saturation of the Southern Ocean CO₂ Sink Due to Recent Climate Change. *Science*, 316(5832),
31 1735–1738. <https://doi.org/10.1126/science.1136188>

32 Li, M., Liu, J., Wang, Z., Wang, H., Zhang, Z., Zhang, L., & Yang, Q. (2013). Assessment of Sea Surface
33 Wind from NWP Reanalyses and Satellites in the Southern Ocean. *Journal of Atmospheric and
34 Oceanic Technology*, 30(8), 1842–1853. <https://doi.org/10.1175/JTECH-D-12-00240.1>

35 Lucio-Eceiza, E. E., González-Rouco, J. F., García-Bustamante, E., Navarro, J., & Beltrami, H. (2019).
36 Multidecadal to centennial surface wintertime wind variability over Northeastern North America
37 via statistical downscaling. *Climate Dynamics*, 53(1–2), 41–66. <https://doi.org/10.1007/s00382-018-4569-5>

38

39 Marshall, G. J. (2003). Trends in the Southern Annular Mode from Observations and Reanalyses. *Journal
40 of Climate*, 16(24), 4134–4143. [https://doi.org/10.1175/1520-0442\(2003\)016%253C4134:TITSAM%253E2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016%253C4134:TITSAM%253E2.0.CO;2)

41

42 Marshall, J., & Speer, K. (2012). Closure of the meridional overturning circulation through Southern
43 Ocean upwelling. *Nature Geoscience*, 5(3), 171–180. <https://doi.org/10.1038/ngeo1391>

44 McLandress, C., Shepherd, T. G., Scinocca, J. F., Plummer, D. A., Sigmond, M., Jonsson, A. I., &
45 Reader, M. C. (2011). Separating the Dynamical Effects of Climate Change and Ozone
46 Depletion. Part II: Southern Hemisphere Troposphere. *Journal of Climate*, 24(6), 1850–1868.
47 <https://doi.org/10.1175/2010JCLI3958.1>

48 Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P. J.,
49 Montzka, S. A., Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G.,
50 Daniel, J. S., Enting, I. G., Law, R. M., Lunder, C. R., O'Doherty, S., Prinn, R. G., ... Weiss, R.

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

1 (2017). Historical greenhouse gas concentrations for climate modelling (CMIP6). *Geoscientific*
2 *Model Development*, 10(5), 2057–2116. <https://doi.org/10.5194/gmd-10-2057-2017>

3 Molod, A., Takacs, L., Suarez, M., & Bacmeister, J. (2015). Development of the GEOS-5 atmospheric
4 general circulation model: Evolution from MERRA to MERRA2. *Geoscientific Model*
5 *Development*, 8(5), 1339–1356. <https://doi.org/10.5194/gmd-8-1339-2015>

6 Morgenstern, O. (2021). The Southern Annular Mode in 6th Coupled Model Intercomparison Project
7 Models. *Journal of Geophysical Research: Atmospheres*, 126(5), e2020JD034161.
8 <https://doi.org/10.1029/2020JD034161>

9 Morgenstern, O., Kinnison, D. E., Mills, M., Michou, M., Horowitz, L. W., Lin, P., Deushi, M., Yoshida,
10 K., O'Connor, F. M., Tang, Y., Abraham, N. L., Keeble, J., Dennison, F., Rozanov, E., Egorova,
11 T., Sukhodolov, T., & Zeng, G. (2022). Comparison of Arctic and Antarctic Stratospheric
12 Climates in Chemistry Versus No-Chemistry Climate Models. *Journal of Geophysical Research:*
13 *Atmospheres*, 127(20), e2022JD037123. <https://doi.org/10.1029/2022JD037123>

14 Morrison, A. K., Hogg, A. M., & Ward, M. L. (2011). Sensitivity of the Southern Ocean overturning
15 circulation to surface buoyancy forcing: SOUTHERN OCEAN OVERTURNING. *Geophysical*
16 *Research Letters*, 38(14), n/a-n/a. <https://doi.org/10.1029/2011GL048031>

17 Mulcahy, J. P., Jones, C., Sellar, A., Johnson, B., Boutle, I. A., Jones, A., Andrews, T., Rumbold, S. T.,
18 Mollard, J., Bellouin, N., Johnson, C. E., Williams, K. D., Grosvenor, D. P., & McCoy, D. T.
19 (2018). Improved Aerosol Processes and Effective Radiative Forcing in HadGEM3 and
20 UKESM1. *Journal of Advances in Modeling Earth Systems*, 10(11), 2786–2805.
21 <https://doi.org/10.1029/2018MS001464>

22 Polvani, L. M., Previdi, M., & Deser, C. (2011). Large cancellation, due to ozone recovery, of future
23 Southern Hemisphere atmospheric circulation trends: OZONE RECOVERY AND SH
24 CIRCULATION TRENDS. *Geophysical Research Letters*, 38(4), n/a-n/a.
25 <https://doi.org/10.1029/2011GL046712>

26 Previdi, M., & Polvani, L. M. (2014). Climate system response to stratospheric ozone depletion and
27 recovery. *Quarterly Journal of the Royal Meteorological Society*, 140(685), 2401–2419.
28 <https://doi.org/10.1002/qj.2330>

29 Ramon, J., Lledó, L., Torralba, V., Soret, A., & Doblas-Reyes, F. J. (2019). What global reanalysis best
30 represents near-surface winds? *Quarterly Journal of the Royal Meteorological Society*, 145(724),
31 3236–3251. <https://doi.org/10.1002/qj.3616>

32 Revell, L. E., Robertson, F., Douglas, H., Morgenstern, O., & Frame, D. (2022). Influence of Ozone
33 Forcing on 21st Century Southern Hemisphere Surface Westerlies in CMIP6 Models.
34 *Geophysical Research Letters*, 49(6), e2022GL098252. <https://doi.org/10.1029/2022GL098252>

35 Schulzweida, U. (2023, October). *CDO User Guide* (Version 2.3.0). Zenodo.
36 <https://doi.org/10.5281/zenodo.10020800>

37 S  f  rian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C.,
38 Berthet, S., Chevallier, M., S  n  si, S., Franchisteguy, L., Vial, J., Mallet, M., Joetjzer, E.,
39 Geoffroy, O., Gu  r  my, J., Moine, M., Msadek, R., ... Madec, G. (2019). Evaluation of CNRM
40 Earth System Model, CNRM-ESM2-1: Role of Earth System Processes in Present-Day and
41 Future Climate. *Journal of Advances in Modeling Earth Systems*, 11(12), 4182–4227.
42 <https://doi.org/10.1029/2019MS001791>

43 Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer,
44 M., Hill, R., Palmieri, J., Woodward, S., Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I.,
45 Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., ... Zerroukat, M. (2019). UKESM1:
46 Description and Evaluation of the U.K. Earth System Model. *Journal of Advances in Modeling*
47 *Earth Systems*, 11(12), 4513–4558. <https://doi.org/10.1029/2019MS001739>

48 Sellar, A. A., Walton, J., Jones, C. G., Wood, R., Abraham, N. L., Andrejczuk, M., Andrews, M. B.,
49 Andrews, T., Archibald, A. T., De Mora, L., Dyson, H., Elkington, M., Ellis, R., Florek, P.,
50 Good, P., Gohar, L., Haddad, S., Hardiman, S. C., Hogan, E., ... Griffiths, P. T. (2020).

Field Code Changed

Field Code Changed

1 Implementation of U.K. Earth System Models for CMIP6. *Journal of Advances in Modeling*
2 *Earth Systems*, 12(4), e2019MS001946. <https://doi.org/10.1029/2019MS001946>

3 Shi, J.-R., Talley, L. D., Xie, S.-P., Liu, W., & Gille, S. T. (2020). Effects of Buoyancy and Wind Forcing
4 on Southern Ocean Climate Change. *Journal of Climate*, 33(23), 10003–10020.
5 <https://doi.org/10.1175/JCLI-D-19-0877.1>

6 Simpkins, G. R., & Karpechko, A. Yu. (2012). Sensitivity of the southern annular mode to greenhouse
7 gas emission scenarios. *Climate Dynamics*, 38(3–4), 563–572. [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-011-1121-2)
8 011-1121-2

9 Solomon, S., Garcia, R. R., Rowland, F. S., & Wuebbles, D. J. (1986). *On the depletion of Antarctic*
10 *ozone*.

11 Solomon, S., Ivy, D., Gupta, M., Bandoro, J., Santer, B., Fu, Q., Lin, P., Garcia, R. R., Kinnison, D., &
12 Mills, M. (2017). Mirrored changes in Antarctic ozone and stratospheric temperature in the late
13 20th versus early 21st centuries. *Journal of Geophysical Research: Atmospheres*, 122(16), 8940–
14 8950. <https://doi.org/10.1002/2017JD026719>

15 Son, S., Tandon, N. F., Polvani, L. M., & Waugh, D. W. (2009). Ozone hole and Southern Hemisphere
16 climate change. *Geophysical Research Letters*, 36(15), 2009GL038671.
17 <https://doi.org/10.1029/2009GL038671>

18 Son, S.-W., Gerber, E. P., Perlwitz, J., Polvani, L. M., Gillett, N. P., Seo, K.-H., Eyring, V., Shepherd,
19 T. G., Waugh, D., Akiyoshi, H., Austin, J., Baumgaertner, A., Bekki, S., Braesicke, P., Brühl, C.,
20 Butchart, N., Chipperfield, M. P., Cugnet, D., Dameris, M., ... Yamashita, Y. (2010). Impact of
21 stratospheric ozone on Southern Hemisphere circulation change: A multimodel assessment.
22 *Journal of Geophysical Research: Atmospheres*, 115(D3), 2010JD014271.
23 <https://doi.org/10.1029/2010JD014271>

24 Son, S.-W., Han, B.-R., Garfinkel, C. I., Kim, S.-Y., Park, R., Abraham, N. L., Akiyoshi, H., Archibald,
25 A. T., Butchart, N., Chipperfield, M. P., Dameris, M., Deushi, M., Dhomse, S. S., Hardiman, S.
26 C., Jöckel, P., Kinnison, D., Michou, M., Morgenstern, O., O'Connor, F. M., ... Zeng, G. (2018).
27 Tropospheric jet response to Antarctic ozone depletion: An update with Chemistry-Climate
28 Model Initiative (CCMI) models. *Environmental Research Letters*, 13(5), 054024.
29 <https://doi.org/10.1088/1748-9326/aabf21>

30 Swart, N. C., & Fyfe, J. C. (2012). Observed and simulated changes in the Southern Hemisphere surface
31 westerly wind-stress: CHANGES IN THE S.H. WESTERLIES. *Geophysical Research Letters*,
32 39(16), n/a-n/a. <https://doi.org/10.1029/2012GL052810>

33 Thompson, D. W. J., & Solomon, S. (2002). Interpretation of Recent Southern Hemisphere Climate
34 Change. *Science*, 296(5569), 895–899. <https://doi.org/10.1126/science.1069270>

35 Thompson, D. W. J., Solomon, S., Kushner, P. J., England, M. H., Grise, K. M., & Karoly, D. J. (2011).
36 Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change. *Nature*
37 *Geoscience*, 4(11), 741–749. <https://doi.org/10.1038/ngeo1296>

38 Tsujino, H., Urakawa, L. S., Griffies, S. M., Danabasoglu, G., Adcroft, A. J., Amaral, A. E., Arsouze, T.,
39 Bentsen, M., Bernardello, R., Böning, C. W., Bozec, A., Chassignet, E. P., Danilov, S., Dussin,
40 R., Exarchou, E., Fogli, P. G., Fox-Kemper, B., Guo, C., Ilicak, M., ... Yu, Z. (2020). Evaluation
41 of global ocean–sea-ice model simulations based on the experimental protocols of the Ocean
42 Model Intercomparison Project phase 2 (OMIP-2). *Geoscientific Model Development*, 13(8),
43 3643–3708. <https://doi.org/10.5194/gmd-13-3643-2020>

44 [Van Noije, T., Bergman, T., Le Sager, P., O'Donnell, D., Makkonen, R., Gonçalves-Ageitos, M.,](https://doi.org/10.5194/gmd-14-5637-2021)
45 [Döscher, R., Fladrich, U., Von Hardenberg, J., Keskinen, J.-P., Korhonen, H., Laakso, A.,](https://doi.org/10.5194/gmd-14-5637-2021)
46 [Myriokefalitakis, S., Ollinaho, P., Pérez García-Pando, C., Reerink, T., Schrödner, R., Wyser, K.,](https://doi.org/10.5194/gmd-14-5637-2021)
47 [& Yang, S. \(2021\). EC-Earth3-AerChem: A global climate model with interactive aerosols and](https://doi.org/10.5194/gmd-14-5637-2021)
48 [atmospheric chemistry participating in CMIP6. *Geoscientific Model Development*, 14\(9\), 5637–](https://doi.org/10.5194/gmd-14-5637-2021)
49 [5668. <https://doi.org/10.5194/gmd-14-5637-2021>](https://doi.org/10.5194/gmd-14-5637-2021)

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

1 Velasquez-Jimenez, L., & Abram, N. J. (2024). Technical note: An improved methodology for calculating
 2 the Southern Annular Mode index to aid consistency between climate studies. *Climate of the*
 3 *Past*, 20(5), 1125–1139. <https://doi.org/10.5194/cp-20-1125-2024>
 4 Wald, A. (1943). Tests of statistical hypotheses concerning several parameters when the number of
 5 observations is large. *Transactions of the American Mathematical Society*, 54(3), 426–482.
 6 Walters, D., Baran, A. J., Boutle, I., Brooks, M., Earnshaw, P., Edwards, J., Furtado, K., Hill, P., Lock,
 7 A., Manners, J., Morcrette, C., Mulcahy, J., Sanchez, C., Smith, C., Stratton, R., Tennant, W.,
 8 Tomassini, L., Van Weverberg, K., Vosper, S., ... Zerroukat, M. (2019). The Met Office Unified
 9 Model Global Atmosphere 7.0/7.1 and JULES Global Land 7.0 configurations. *Geoscientific*
 10 *Model Development*, 12(5), 1909–1963. <https://doi.org/10.5194/gmd-12-1909-2019>
 11 Waugh, D. W., Banerjee, A., Fyfe, J. C., & Polvani, L. M. (2020). Contrasting Recent Trends in Southern
 12 Hemisphere Westerlies Across Different Ocean Basins. *Geophysical Research Letters*, 47(18),
 13 e2020GL088890. <https://doi.org/10.1029/2020GL088890>
 14 Yool, A., Palmiéri, J., Jones, C. G., de Mora, L., Kuhlbrodt, T., Popova, E. E., Nurser, A. J. G., Hirschi,
 15 J., Blaker, A. T., Coward, A. C., Blockley, E. W., & Sellar, A. A. (2021). Evaluating the physical
 16 and biogeochemical state of the global ocean component of UKESM1 in CMIP6 historical
 17 simulations. *Geoscientific Model Development*, 14(6), 3437–3472. <https://doi.org/10.5194/gmd-14-3437-2021>
 18 Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M.,
 19 Tanaka, T., Hosaka, M., Yabu, S., Yoshimura, H., Shindo, E., Mizuta, R., Obata, A., Adachi, Y.,
 20 & Ishii, M. (2019). The Meteorological Research Institute Earth System Model Version 2.0,
 21 MRI-ESM2.0: Description and Basic Evaluation of the Physical Component. *Journal of the*
 22 *Meteorological Society of Japan. Ser. II*, 97(5), 931–965. <https://doi.org/10.2151/jmsj.2019-051>
 23 Zambri, B., Solomon, S., Thompson, D. W. J., & Fu, Q. (2021). Emergence of Southern Hemisphere
 24 stratospheric circulation changes in response to ozone recovery. *Nature Geoscience*, 14(9), 638–
 25 644. <https://doi.org/10.1038/s41561-021-00803-3>
 26 Zeng, G., Morgenstern, O., Williams, J. H. T., O'Connor, F. M., Griffiths, P. T., Keeble, J., Deushi, M.,
 27 Horowitz, L. W., Naik, V., Emmons, L. K., Abraham, N. L., Archibald, A. T., Bauer, S. E.,
 28 Hassler, B., Michou, M., Mills, M. J., Murray, L. T., Oshima, N., Sentman, L. T., ... Young, P. J.
 29 (2022). Attribution of Stratospheric and Tropospheric Ozone Changes Between 1850 and 2014 in
 30 CMIP6 Models. *Journal of Geophysical Research: Atmospheres*, 127(16), e2022JD036452.
 31 <https://doi.org/10.1029/2022JD036452>
 32
 33
 34




Fig. S1: Timeseries of Total Column Ozone (TCO) for 70-90°S, SAM index, and 10-m wind speed (40-60°S) for CMIP6 models with interactive chemistry. Relevant observations and reanalyses are shown as follows: the Bodecker observational TCO is shown in dashed blue in the top row. The Marshall station-based SAM index is shown in dashed blue in the middle row, and the ERA5 wind speed is shown in light blue in the bottom row. ¶
 Fig. S2 ¶

