



The role of atmospheric circulation changes in Western European warm season heat extremes

Douwe Sierk Noest^{1,2}, Izidine Pinto¹, Vikki Thompson^{1,2}, Dim Coumou²

¹Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands

²Institute for Environmental Studies, Vrije Universiteit Amsterdam, Amsterdam, Netherlands

Correspondence to: Izidine Pinto (izidine.pinto@knmi.nl)

Abstract. Climate change has led to an intensification of summer heat extremes, with especially pronounced warming over Western Europe. Here, the maximum and mean of the daily maximum summer temperatures have warmed 3.3 and 2.4 times faster than global mean temperatures. A large part of this enhanced warming can be attributed to dynamical changes. The effects of climate change on springtime heat extremes and circulation changes are less well understood, though changes in spring can influence summer via soil moisture memory. Here we show that between 1950 and 2023, the maximum and mean of the daily maximum spring temperatures in Western Europe have intensified 2.2 and 2.0 times faster than global warming respectively. We show that most of this enhanced warming can be attributed to thermodynamical effects. However, using circulation analogues, we show that locally more than a third of the total temperature trends can be attributed to changes in atmospheric circulation. Specifically, southerly flow patterns, characterized by high pressure over Western Europe and low pressure over the Eastern Atlantic, are found to become more frequent and intense in spring, contributing to the warming trend. Finally, individual ensemble members from large ensemble historical climate model simulations show that those models are capable of simulating temperature trends nearly as extreme as observed, but the model mean underestimates the Western European trends.

1. Introduction

Human-induced climate change has led to the intensification of heat extremes, and their impacts are expected to increase even more with global warming in the future (IPCC, 2021). Extreme heat can deteriorate health (Kjellstrom et al., 2010), cause excess mortality (Yang et al., 2021), affect ecosystems, and threaten food security (Horton et al., 2016). Moreover, extreme heat can cause additional economic damages through reduced productivity, with some heatwave events in Europe causing losses of up to 0.5% of the European gross domestic product (García-León et al., 2021).

Although most land areas have experienced an increase in heat extremes, some regions are warming faster than others. Globally, the strongest warming of the hottest days is expected to be around 1.5 to 2 times the rate of global-mean, annual-mean warming (IPCC, 2021). However, Vautard et al. (2023) show that summer heat extremes for Western Europe are warming much faster, up to 5 times faster than the global mean temperature trend.

Such trends in extremes can partly be attributed to thermodynamical effects, e.g. warming caused by increased atmospheric greenhouse gas concentrations, enhanced warming over land as compared to oceans, and increased evaporation contributing to soil moisture depletion. However, although these processes are typically less well understood, trends in extremes can also partly be explained by atmospheric dynamical changes (IPCC, 2021; Shaw et al., 2024). Vautard et al. (2023) focus on “Southerly Flow” (SF) patterns, whose increase in both frequency and



persistence is the main driver of the dynamical component of the summer heat extreme trends. These patterns contain an anticyclonic component over Central Europe and are characterized by the transport of warm air from more southern regions towards Western Europe. This can lead to extreme temperatures, as seen during the
40 heatwaves in June and July 2019, when advection of air from North Africa, caused by subtropical ridges, resulted in temperatures of 46 °C in France and other record-breaking temperatures in the Netherlands, Belgium, and Germany (Sousa et al., 2020; Vautard et al., 2020).

Climate models underestimate trends in heat extremes for Western Europe, especially the trend in the maximum of the daily maximum summer temperatures (Vautard et al., 2023). Models also underestimate the increase in
45 frequency of SF days and the dynamical contribution to the temperature trends, which explains a large part of the mismatch (Vautard et al., 2023). Similar underestimations of Western European heat extremes by climate models are found in other studies (e.g., Kornhuber et al., 2024; Lorenz et al., 2019; Patterson, 2023). Climate models play an essential role in understanding future changes, regional climate impact assessments, and weighing possible adaptation and mitigation strategies (e.g., IPCC, 2023; Lopez et al., 2009). Therefore, it is important to analyse the
50 performance and limitations of these models and understand what causes the differences between simulations and observations.

One factor that can influence summer heat extremes is the springtime soil moisture (e.g., Whan et al., 2015). Soil moisture deficits in spring can persist into the summer, where they can influence heat extremes by limiting the latent heat flux, thereby increasing the amount of energy that is available for the sensible heat flux (Seneviratne et al., 2010; Wu & Zhang, 2015). Spring temperatures, in turn, can influence spring soil moisture in different ways. Extremely warm springs can lead to an earlier start of the growing season, resulting in prolonged evapotranspiration and reduced soil moisture (Fischer et al., 2007; Liu & Zhang, 2020). Warmer springs can also lead to a higher evaporative demand, potentially leading to more evapotranspiration and decreased soil moisture (Seneviratne et al., 2010). However, spring heat extremes are less studied compared to summer heat extremes
60 (Sulikowska & Wypych, 2021).

To better understand how these processes, and their influence on summer heat extremes, might have changed due to global warming, this study investigates spring heat extremes in Western Europe. We analyse how fast Western European spring heat extremes are intensifying and what part of this intensification can be attributed to atmospheric dynamical changes. Since the analysis of spring heat extremes in this study applies a similar approach
65 as used by Vautard et al. (2023), their findings on summer heat extremes will be reproduced as well to validate the results. Moreover, it is investigated how Southerly Flow patterns over Western Europe have changed in spring and how they are distributed over the longer warm season. Finally, we analyse how climate models perform in reproducing this seasonal behaviour and long-term trends.

2. Data and methods

70 2.1 Total temperature trends and dynamical contributions

Trends in spring, March–April–May (MAM), and summer, June–July–August (JJA), heat extremes have been calculated for the period 1950–2023, using the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 reanalysis dataset (Hersbach et al., 2020). The intensification of the maximum (TXx) and the mean (TXm) of the daily maximum temperatures in a season are compared to the rate of global warming, expressed in degrees
75 Celsius of warming per global warming degree (°C / GWD), using the method applied in Vautard et al. (2023).



Part of these trends can be explained by changes in atmospheric circulation patterns. For example, more occurrences of circulation patterns associated with warm weather can increase temperatures, even without the thermodynamical effects of global warming. To isolate this dynamical contribution and remove the thermodynamical component, trends have also been calculated from a temperature time series in which all
80 temperatures have been scaled to an arbitrary reference warming level, using 2023 as a reference year, and in which temperature fields have been replaced by the fields of days with a similar circulation pattern. Details on the calculation of the trends, the thermodynamical correction, and the ‘shuffling’ of the temperature time series are described in Vautard et al. (2023). For both the total and dynamical temperature trends, the trend at a specific grid point is considered to be significant when the absolute slope of the regression is larger than 2 times the standard
85 error of the slope. The average trends for Western Europe are calculated as the area-weighted average over the land areas within 5° W to 15° E and 45° N to 55° N (Fig. 1, box B). A land mask was derived from the E-OBS dataset (Cornes et al., 2018).




Figure 1: The 500 hPa streamfunction for the Southerly Flow event on 29 June 2019. Box (A) shows the domain used to select analogues. Box (B) shows the area over which trends for Western Europe are averaged.

2.2 Circulation analogues

To analyse the dynamical component of the temperature trends and the changes in SF patterns, circulation analogues have been used. An analogue is a day that has a similar atmospheric circulation as a given event, which for example allows for the analyses of temperatures conditioned to different circulation patterns (Jézéquel et al.,
95 2018). In this study, analogue days were selected based on their 500 hPa streamfunction within 30° W to

20° E and 30° N to 60° N (Fig. 1, box A), as in Vautard et al. (2023). To select the best analogues, the Euclidian distance (ED) between the streamfunction fields of a selected event and each day in the season of interest is calculated as described in Thompson et al. (2024a). A lower ED indicates a better analogue, but the event itself (ED = 0) cannot be its own analogue. Moreover, analogues have to be separated by at least 6 days to prevent
100 selecting more analogues from events that have already been selected. To ensure the analogue selection is focused on the actual circulation pattern, rather than absolute streamfunction values, all daily streamfunction fields were prepared by subtracting the mean of all streamfunction values within the domain from each individual value within the domain. This preserves the gradients, and thus the wind field pattern as described by the contour lines, whilst removing any differences in absolute values.



105 2.3 Southerly Flow patterns

To analyse whether and how SF patterns occurring in spring have changed over time, different characteristics of the analogues for a selected event have been investigated using the ERA5 reanalysis dataset. Figure 1 shows the 500 hPa streamfunction field for the selected SF event on 29 June 2019. This event was part of the hottest June on record for Europe and its circulation pattern is the most representative of days on which the maximum summer
110 temperature is recorded in central France (Vautard et al., 2020, 2023).

The first methods to assess changes in SF patterns analyse differences between analogue sets from two 30-year periods. From both the past period (1950–1979) and the present period (1994–2023), the 30 spring days with circulation patterns closest to the selected SF event are selected as analogues. Between these two periods, changes in typicality and persistence of both the event (t_{event} & p_{event}) and analogues ($t_{analogue}$ & $p_{analogue}$), as well as changes
115 in the intensity of the analogues have been analysed using the methods described in Thompson et al. (2024a). The typicality can also be used as a quality check for the analogues. If the t_{event} value falls below the distribution of the $t_{analogue}$ values, this suggests that the analogues are more like the other analogues than like the event itself. This would mean that the event is too unique and that therefore, the dynamical changes associated with the event cannot be analysed using the analogues (Thompson et al., 2024a). To test whether the means of the distributions of the
120 $t_{analogue}$ and $p_{analogue}$ values differ significantly between the two time periods, a t-test is performed. The significance of changes in intensity, defined as the difference between the means of the streamfunction fields from the 30 selected analogues for both periods, is also assessed for each location using a t-test.

To test whether the SF analogues are also changing when using a different method, the similarity approach as used by Thompson et al. (2024b) has been adapted. Rather than using only the 30 best analogues over an entire period,
125 the best analogue day within each year is selected. This method has been applied to the same 30-year periods, and a t-test was performed to analyse the statistical difference between the mean of the values from both periods.

As a final method to assess changes in SF patterns, the frequency has been analysed using a Euclidian distance threshold. This method has also been used to analyse how SF patterns are distributed over different months of a longer warm season (MAMJJAS). This time, the entire dataset is divided into a past (1950–1986) and a present
130 (1987–2023) period. For each month, the number of analogues with a Euclidian distance below a certain threshold in that period is counted. The threshold is taken as the fifth percentile of all Euclidian distances in the analysed months of both periods combined. The streamfunction field corresponding to the threshold is shown in the Supplementary Material (Fig. S1).

2.4 Climate models

135 To analyse the ability of climate models to simulate the observed temperature trends and changes in SF days, part of the analysis has been repeated with climate model output data from the Large Ensemble Single Forcing Model Intercomparison Project (LESFMIP) (Smith et al., 2022). This was done using two global climate models that are also part of the Coupled Model Intercomparison Project Phase 6 (CMIP6), the HadGEM3 and MIROC6 models. These models were chosen due to the high number of ensembles available at the time of analysis. For the
140 HadGEM3 model, more details on the exact simulation that was used (HadGEM3-GC31-LL) and specifics of the model are described in Andrews et al. (2020) and Ridley et al. (2019). For the MIROC6 model, more information on the simulation, model components, and their resolution are described by Shiogama et al. (2023) and Tatebe et al. (2019). From both models, data covering the 1950–2014 period from the historical-forcing runs were used.



To compare the ERA5 and model results, the data have been regridded to the HadGEM3 model resolution ($1.875^\circ \times 1.25^\circ$). Moreover, since the model data only covers up until 2014, the ERA5 analyses have also been repeated with the 1950–2014 period to make comparisons possible. For the total temperature trends, the analysis has been repeated using 55 ensemble members from the HadGEM3 model and 50 ensemble members from the MIROC6 model. Due to missing data in the daily wind fields from the models, the analogue analyses that use streamfunction have only been repeated with five ensemble members from the HadGEM3 model and three ensemble members from the MIROC6 model. These eight ensemble members have been used to repeat the analysis of the dynamical component of the temperature trend and the typicality, persistence, and intensity of SF days. For the analyses investigating SF days using the model data, the past period remains 1950–1979 and the present period becomes 1985–2014.

3. Results

3.1 Total temperature trends and the dynamical contributions

Figure 2 shows the total trends in spring and summer temperature extremes, as well as the dynamical contributions to these trends. For spring, the largest TXx trends are found in Norway and Spain, with trends in Norway reaching more than 7°C per GWD (Fig. 2A). For Western Europe, the warming is somewhat slower, but still significantly faster than global warming with an average spring TXx trend for land areas equalling 2.2°C per GWD and maximum trends of more than 3°C per GWD around the north of Italy. The total spring TXm trend is found to have smaller regional differences with generally lower trends as well (Fig. 2C). The maximum trends reach up to 2.5°C per GWD and are found in for example Spain and Sweden. For Western Europe, the average spring TXm trend is 2.0°C per GWD. Both the total TXx and TXm trends are found to be significant on a 95% confidence level for most of Europe.

For the dynamical spring TXx trend, values of around -1°C to 1°C per GWD are found (Fig. 2B). For Western Europe, this results in an average dynamical contribution to the TXx trend of 0.2°C per GWD. However, note that most of these trends are insignificant on a 95% confidence level, apart from the warming in some small regions like the south of the United Kingdom and Italy, the west of France, and the east of Spain. A small area in the east of Romania is the only region with a significant cooling trend. The dynamical contribution to the spring TXm trend is a warming over all of Western Europe, with an average trend of 0.4°C per GWD and significant trends between 0.5°C and 1°C per GWD over parts of France and Spain (Fig. 2D). Note that, since European temperatures typically increase throughout spring, the spring TXx trend could be biased towards May rather than equally representing the entire season. However, by repeating the analyses with anomalies from the daily climatological mean temperature instead of absolute temperatures, it has been shown that this has little influence on the results, as shown in the Supplementary Material (Fig. S2).

For summer, the largest TXx trends are found in Western Europe, with a maximum of more than 5°C per GWD and an average Western European trend of 3.3°C per GWD (Fig. 2E). The dynamical contributions to the TXx trends are largest in Western Europe as well, with an average of 1.0°C per GWD (Fig. 2F). The summer TXm trends result in a lower Western European average of 2.4°C per GWD with an average dynamical contribution of 0.7°C per GWD (Fig. 2G and 2H). In general, both the spatial pattern and the magnitude of the summer trends are similar to the trends up to 2022 as found by Vautard et al. (2023).




Figure 2: The total trends in the maximum (TXx) and the mean (TXm) of the daily maximum temperatures in a season, for spring and summer, and the dynamical contributions to these trends, expressed in the amount of warming, in degrees Celsius, per global warming degree (GWD). Dotted areas represent regions where the trend is not significant on a 95% confidence level.


3.2 Changes in Southerly Flow days

To assess the changes in frequency and persistence of SF events, we analyse the difference in typicality and persistence between the past and present time periods for the selected SF event (Fig. 3). For both periods, the t_{event} falls within the distribution of the $t_{analogue}$ values, showing the 30 analogues are of sufficient quality to investigate dynamical changes associated with the event (Fig. 3A). The typicality of the event in the present period is higher compared to the past period, showing that analogues in the present period are more similar to the event, suggesting an increase in typicality – and thus frequency – of SF days like the event. The $t_{analogue}$ distributions support this finding, the higher mean value in the present period indicates that the analogues, that became more similar to the



195 event, also become more typical to each other. However, note that the difference between the means of the $t_{analogue}$ distributions is not significant on a 95% confidence level.

Whereas the event itself only persisted for 1 day, Fig. 3B shows that most of the analogues persist longer, with one analogue from the past period persisting for up to 8 days. Note that, since a persistence of 8 days exceeds the analogue separation range of 6 days, it has been checked that not two analogues were selected from this single
200 event. However, apart from this single analogue, the distributions of the $p_{analogue}$ values did not change a lot over time. There seems to be a slight shift from analogues persisting for 2 days towards analogues persisting for 3 days, but the means of both distributions are almost identical.



205 **Figure 3: The changes in typicality (A) and persistence (B) for spring between the past (1950–1979) and present (1994–2023) periods. Red dots indicate the typicality and persistence of the event. The violins show the distribution of the $t_{analogue}$ and $p_{analogue}$ values, whose means are represented by horizontal lines. The p-value indicates the statistical significance of the difference between these means.**

Figure 4 shows the similarity value for each year in the two time periods. The highest similarity values are found in the present period. Moreover, the mean similarity of the present period is higher compared to the past period,
210 again suggesting that analogues are becoming more similar to the event. However, note that with a p-value of 0.095, the difference between the means is not significant on a 95% confidence level.




Figure 4: The similarity value of the best analogue in a year, for each year within the past and present time periods. Dashed grey lines represent the mean of all similarity values within a time period.

215 Figure 5 shows the SF event, the SF analogue composites for both the past and present periods, and the changes in intensity. Over most of Western Europe, there is an increase in streamfunction gradient, indicating a



strengthening of the high pressures and a deepening of low pressures. For the higher pressure over Europe, this would result in stronger subsidence of air, and therefore an increase in the intensity of potential heat events. This increase is found to be even stronger, and statistically significant, for parts of France and Spain. Moreover, the increase in streamfunction in the centre of the domain, combined with the decrease in both lower corners of the domain, increases the gradient of the streamfunction. This reflects an increased advection of air from southern regions towards the north, further intensifying the SF days. When comparing the streamfunction pattern from the event with the past composite, the event seems to have higher values over Southern and Western European land, but lower values southeast of Italy and in the southwest corner of the domain. This matches the pattern found in the difference map between the present and past composites, once again confirming that the analogues are becoming more similar to the event over time, as also seen in Fig. 3A, whilst also showing how this happens spatially.




Figure 5: The difference in 500 hPa streamfunction between analogue composites from the present (1994–2023) and past (1950–1979) periods. Hashed areas show regions where the difference is significant on a 95% confidence level.

Finally, Fig. 6 shows the distribution of SF days over the longer warm season and how their frequency changes over time. It becomes clear that the frequency of SF days in spring has slightly increased over time, but that the increase is much bigger for the summer season. Interestingly, the frequencies in July and September have decreased over time. When looking at the distribution over the months, by far the largest amount of SF days occur in summer, especially for the present period. For both time periods, there seems to be a rapid increase in frequency after April.




Figure 6: The distribution of analogues within the fifth percentile of Euclidian distances over different months, for two time periods.

3.3 The performance of climate models

240 The total temperature trends as found when using ERA5 data until 2014 are shown in Fig. 7A and B. Although the spatial patterns are comparable to the trends up to 2023, the trends until 2014 actually tend to be larger, with a new maximum of 9 °C per GWD in Norway. Figure 7 also visualizes the performance of the HadGEM3 and MIROC6 models to reproduce these total temperature trends. When looking at the average trends found by the 105 model ensemble members, the models seem to correctly capture the general spatial patterns, with the largest TXx trends

245 found in Norway and Spain and relatively high TXm trends simulated in Eastern Europe (Fig. 7C and D). However, the average modelled trends, especially the TXx trends, are much lower compared to the ERA5 trends with a maximum average TXx trend of 3.4 °C per GWD. The standard deviations of the modelled trends are largest in areas with the highest temperature trends and are especially large for the TXx trends (Fig. 7E and F), indicating that individual ensemble members could reach higher trends. Indeed, the largest TXx trend found in an individual

250 ensemble member equals 8.8 °C per GWD, showing that, although the models on average underestimate the trends, extreme cases as found for ERA5 can be captured by the models. To investigate how many of the individual ensemble members find a large enough trend, Fig. 7G and H show the percentage of ensemble members that find a trend as high as the trend found in ERA5 or higher, for each location. For some areas, like Greece and the south of Italy, most of the ensemble members seem to simulate a large enough trend. However, the trends in most parts

255 of Western Europe are underestimated by more than 70% of the ensemble members. This is also the case for Spain and parts of Eastern Europe. In general, the TXx trend seems to have the most extreme underestimated trends, like in Norway and the north of Italy.





Figure 7: The ERA5 total temperature trends in the maximum (TXx) and the mean (TXm) of the daily maximum spring temperatures up to 2014, the mean trends found by the 105 model ensemble members from the HadGEM3 and MIROC6 models, the standard deviation of the modelled trends and the percentage of the ensemble members that find a trend as high as the ERA5 trend or higher.

To further analyse the models' performances for Western Europe, Fig. 8 shows the distribution of the Western European average trends as found by all individual ensemble members. Again, most of the ensemble members underestimate the trend as found in ERA5. For the Western European TXx trend, the mean of the ensemble members equals 1.7 °C per GWD and only 10 out of 105 ensemble members simulate a trend as high as the ERA5 trend. For the TXm trend, the mean is 1.9 °C per GWD and 26 ensemble members find a large enough trend. However, with standard deviations of 0.8 and 0.7 °C per GWD for the TXx and TXm trends respectively, the ERA5 trends of 2.9 and 2.4 °C per GWD do fall within the 95% confidence interval of the ensemble means. Interestingly, there is a large difference between the two different models. When only taking into account the HadGEM3 model, only one ensemble member finds a TXm trend as high as ERA5, and no ensemble member finds a large enough TXx trend. With respective TXx and TXm means equalling 1.6 and 1.4 °C per GWD and



standard deviations of 0.6 and 0.4 °C per GWD, both the ERA5 trends fall outside of the 95% confidence interval of the HadGEM3 mean trends.



275

Figure 8: The distribution of the for Western Europe averaged trends in the maximum (TXx) and the mean (TXm) of the daily maximum spring temperatures, as found by 105 ensemble members from the HadGEM3 and MIROC6 models. The dashed grey lines represent the trends found in ERA5.

To investigate whether the underestimation of the total temperature trends is caused by an underestimation of the dynamical changes, the analysis of the dynamical trends has been repeated with ERA5 and model data until 2014. The results for the eight ensemble members are shown in the Supplementary Material (Fig. S3 & S4). The result for the TXx trend shows large differences between ensemble members, with half warming and half cooling over Western Europe. All four ensemble members that show a warming find trends larger than observed in ERA5, with maximum trends of 0.73 and 0.86 °C per GWD for the HadGEM3 and MIROC6 models respectively. Therefore, although the average model trend (0.11 °C per GWD) is lower than the observed ERA5 trend (0.26 °C per GWD), the models do not seem to systematically underestimate the dynamical component of the TXx trend. The results for the TXm trend are a little less clear, with an observed ERA5 trend showing a warming over almost all of Europe, and a larger Western European trend compared to the 2023 trend of 0.64 °C per GWD. From the models, only three ensemble members show a warming trend over Western Europe, with a maximum trend of 0.52 °C per GWD, resulting in an average Western European trend of 0.00 °C per GWD. Only one member shows a warming over most of Europe, and three members show a cooling trend over most of Europe.

The typicality and persistence analysis have been repeated for the model ensembles, and the results are shown in the Supplementary Material (Fig. S5). Only two out of the eight ensemble members find an increase in both the t_{event} and the mean of the $t_{analogue}$ distribution, and for one of them, the difference between the means is even significant on a 95% confidence level. However, their increase in the typicality of the event and the absolute typicality values are still lower compared to the typicality found in ERA5. One HadGEM3 ensemble even finds a statistically significant decrease in the mean $t_{analogue}$ value. There also seems to be a difference between the performance of the models in simulating similar SF days, with each mean of the $t_{analogue}$ distributions found by the MIROC6 model being similar or higher than the ERA5 means and higher than all means found by the HadGEM3 model. Apart from an analogue persisting up to 10 days, the persistence does not seem to change over time for all ensemble members, who find similar means as found in ERA5.

Finally, the analysis of the intensity of SF days has been repeated, and results are shown in Fig. 9. When looking at the composites of the analogues, all ensemble members still show clear SF patterns. However, the difference



between time periods differs a lot between the members. Only one ensemble member shows a pattern similar to
305 ERA5, with an increase in streamfunction magnitude over the north of Africa, Spain, and France and decreases in
both lower corners of the domain. For the remaining ensembles, changes are either small and insignificant or show
an opposite change with large decreases over European land.




Figure 9: The difference in 500 hPa streamfunction between analogue composites from the past (1950–1979) and present (1985–2014) periods, for ERA5 and the different model ensembles. Hashed areas show regions where the difference is significant on a 95% confidence level.



4. Discussion

The increase in the typicality of the selected SF event can partly be attributed to a small increase in the frequency of SF days, although the increase in spring is much smaller compared to the increase for the summer season.

315 Vautard et al. (2023) also found a large increase in the frequency of summer SF days. However, they also found an increase in the persistence of these days (24% between 1950 and 2022), whereas this study shows no clear increase in the persistence of SF days can be detected for springtime. The remaining increase in typicality can be attributed to the fact that SF days are becoming more similar to the selected SF event. With higher pressure over Western European land and a stronger advection of warm air from the south, caused by an increase in the gradient
320 of the streamfunction field, the SF days are becoming more intense. Both changes are part of the dynamical contributions to the total temperature trend and could be important for spring heat extremes.

The changes in SF days may partly be caused by changes in land-sea temperature contrast, which is found to be increasing with global warming (e.g., Shaw & Voigt, 2015). A larger temperature contrast also results in a geopotential height contrast. The anticyclonic anomalies over land and cyclonic anomalies over the ocean that can
325 result from this (Kamae et al., 2014), could contribute to the intensification of the Southerly flow patterns. SF patterns related to heat events over Europe can also be part of a larger Rossby wave (White et al., 2022). For example, several heat extremes over Western Europe were related to a wave 7 pattern, a wave pattern that might also be favoured by a large land-sea temperature contrast in the mid-latitudes (Kornhuber et al., 2019). With an increasing temperature contrast caused by global warming, as well as from spring to summer, this could influence
330 the increases in the frequency of SF days during the warm season and between time periods. Furthermore, it has been hypothesized that a larger land-sea temperature contrast, that can favour a double-jet structure, can amplify such planetary waves as well as lead to more frequent blocking events that influence heat extremes (He et al., 2018; Rousi et al., 2022).

The results show that the Western European spring TXx has warmed with 2.2 °C per GWD. Although not as fast
335 as the summer trend of 3.3 °C per GWD, it is more than twice the rate of global warming. For the TXm trends, there is a smaller difference between seasons, with 2.4 °C per GWD in summer and 2.0 °C per GWD in spring. This rapid spring warming could directly contribute to spring soil moisture deficits and therefore influence the summer extremes in Western Europe.

Other studies have shown that there is a connection between Southern European spring soil moisture deficits and
340 Western European summer heat extremes (e.g., Vautard et al., 2007; Zampieri et al., 2009). Dry soils in Southern Europe result in local warming due to the increase in the sensible heat flux. Moreover, a soil moisture deficit results in drier air as well, which in turn leads to a reduction in cloudiness and therefore amplifies the dry and hot conditions through an increase in radiation. These dry conditions can be propagated towards more northern parts of Europe by southerly winds, where they increase the temperature and evaporative demand, again resulting in
345 drier conditions. Here, the dry soils can influence extreme heat through the previously described feedbacks, as well as by favouring anticyclonic conditions (Vautard et al., 2007; Zampieri et al., 2009). Whereas for summer, the highest TXx trends are found in Western Europe, the highest trends in spring are found in Norway and Spain, the latter of which also has relatively high TXm trends in both spring and summer. Rapid Southern European warming trends in spring, like those found for Spain, could therefore potentially influence Western European



350 summer heat extremes by contributing to the initial soil moisture deficits in Southern Europe needed to start the propagation of drought and heat towards Western Europe.

The most extreme spring trends are found in Norway, with TXx trends of more than 7 °C per GWD. Similar results are found when using different methods to calculate trends, as shown by Sulikowska & Wypych (2021). They find that spring is the season with the largest TXx trend for Western Scandinavia, with an average warming of 0.5 °C
355 per decade between 1950 and 2019. Moreover, they show that when dividing most of Europe into five study domains, and considering all seasons, the Western Scandinavian spring trend is tied for the largest TXx trend with the summer trend in the British Isles domain, covering the Netherlands, Belgium, Ireland, the United Kingdom, and part of France. However, note that the highest trends in Scandinavia found in their study are located in southwestern Finland, an area that shows average trends in this study, and in the north of Finland, an area that is
360 not investigated in this study. Finally, Sulikowska & Wypych (2021) show that the trends in average maximum spring temperatures are higher for Central Europe and Iberia than for the British Isles domain, which seems to match the pattern in the TXm trends found in this study.

The dynamical components of the summer TXx trends are especially high and significant for Western Europe. For spring on the other hand, there are lower and mostly insignificant dynamical components over Western Europe,
365 with even a cooling trend over parts of Germany. However, there are still some small regions in Spain, France, and the United Kingdom that show a significant warming caused by dynamical changes, with in the United Kingdom dynamical components reaching more than 1 °C per GWD. The spring TXm trends show a more homogenous warming over most of Europe, which corresponds better to the summer TXm trends. Although spring trends again tend to be slightly lower and more insignificant, parts of Spain and France still show a significant warming caused
370 by dynamical changes of 0.5 to 1 °C per GWD, which can locally explain more than a third of the total temperature trend.

Although some of the individual model ensemble members reproduce extreme total temperature trends close to those observed in ERA5, most of the ensemble members underestimate the trends over large parts of Europe, including Western Europe. There are also large differences between the performance of the different models, with
375 the HadGEM3 model significantly underestimating the average temperature trends for Western Europe. Vautard et al. (2023) attribute a large part of the underestimation of summer trends to the underestimation of the dynamical component, with 0 out of 170 simulations finding a dynamical TXx component as large as in ERA5. D'Andrea et al. (2024) also show that CMIP6 models are unable to reproduce the large observed increase in occurrences of summertime mid-tropospheric deep depressions over the Eastern North Atlantic, a pattern that could also be linked
380 to Western European summer heat extremes. For spring, the underestimation of dynamical changes seems to be less of an issue, with lower observed dynamical trends to start with and four out of eight ensemble members finding a dynamical component larger than observed in ERA5. However, on average the dynamical components of both the TXx and TXm trends are still underestimated, and zero out of eight ensemble members simulate a dynamical TXm trend as high as observed in ERA5. A similar result is found for the models their ability to simulate the
385 changes in SF days, with some ensemble members finding similar changes in intensity and typicality, although smaller, but most ensemble members finding little change or opposite changes.



5. Conclusion

This study shows that Southerly Flow days are becoming more frequent and more intense in spring, contributing to the observed pronounced warming trends. The maximum and the mean of the daily maximum spring temperatures in Western Europe are found to intensify 2.2 and 2.0 times faster than global warming respectively. By influencing soil moisture deficits, these warming trends can drive changes in summer heat extremes in Western Europe. Although the dynamical contributions to the TXx trends are less significant than in summer, locally still more than a third of the TXm trend can be attributed to changes in atmospheric circulation patterns only. Future research could focus on quantifying the effect of changes in different circulation patterns on the spring temperature trends, and how they in turn influence summer heat extremes. Finally, this study shows that, although climate models are capable of simulating temperature trends as extreme as observed, on average they underestimate the trends. Similar results are found for the models their ability to reproduce the changes in Southerly Flow days. Although the underestimation of the total temperature trends does not seem to be caused by a systematic underestimation of the dynamical component, as is the case for summer trends, future research should further expand on this analysis by analysing more model data.

Code and data availability

ERA5 and E-OBS datasets can be previewed or downloaded through the KNMI Climate Explorer (KNMI, 2022). The HadGEM3 model data that was used in this study is available through Ridley et al. (2019) and the MIROC6 data is available through Tatebe & Watanabe (2018). All code used for this study can be accessed at https://github.com/douwe3/Western_EU_warm_season_heat_extremes (Noest, 2025).

Author contribution

DN: conceptualization, formal analysis, writing (original draft preparation, review and editing). IP: conceptualization, supervision, writing (review and editing). DC: conceptualization, supervision, writing (review and editing). VT: conceptualization, supervision, writing (review and editing).

410 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We would like to thank our colleagues from the KNMI and Institute for Environmental Studies (IVM) for useful discussions during this project. Moreover, the authors acknowledge the KNMI for the use of their computational resources.

References

- Andrews, M. B., Ridley, J. K., Wood, R. A., Andrews, T., Blockley, E. W., Booth, B., Burke, E., Dittus, A. J., Florek, P., Gray, L. J., Haddad, S., Hardiman, S. C., Hermanson, L., Hodson, D., Hogan, E., Jones, G. S., Knight, J. R., Kuhlbrodt, T., Misios, S., Mizielinski, M. S., Ringer, M. A., Robson, J., and Sutton, R. T.: Historical Simulations With HadGEM3-GC3.1 for CMIP6, *J. Adv. Model. Earth Syst.*, 12, e2019MS001995, <https://doi.org/10.1029/2019MS001995>, 2020.
- Cornes, R. C., Van Der Schrier, G., Van Den Besselaar, E. J. M., and Jones, P. D.: An Ensemble Version of the E-OBS Temperature and Precipitation Data Sets, *J. Geophys. Res. Atmospheres*, 123, 9391–9409, <https://doi.org/10.1029/2017JD028200>, 2018.



- 425 D'Andrea, F., Duvel, J., Rivière, G., Vautard, R., Cassou, C., Cattiaux, J., Coumou, D., Faranda, D., Happé, T.,
Jézéquel, A., Ribes, A., and Yiou, P.: Summer Deep Depressions Increase Over the Eastern North Atlantic,
Geophys. Res. Lett., 51, e2023GL104435, <https://doi.org/10.1029/2023GL104435>, 2024.
- Fischer, E. M., Seneviratne, S. I., Vidale, P. L., Lüthi, D., and Schär, C.: Soil Moisture–Atmosphere Interactions
during the 2003 European Summer Heat Wave, J. Clim., 20, 5081–5099,
430 <https://doi.org/10.1175/JCLI4288.1>, 2007.
- García-León, D., Casanueva, A., Standardi, G., Burgstall, A., Flouris, A. D., and Nybo, L.: Current and projected
regional economic impacts of heatwaves in Europe, Nat. Commun., 12, 5807,
<https://doi.org/10.1038/s41467-021-26050-z>, 2021.
- He, Y., Huang, J., Li, D., Xie, Y., Zhang, G., Qi, Y., Wang, S., and Totz, S.: Comparison of the effect of land-sea
435 thermal contrast on interdecadal variations in winter and summer blockings, Clim. Dyn., 51, 1275–1294,
<https://doi.org/10.1007/s00382-017-3954-9>, 2018.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu,
R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G.,
Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming,
440 J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M.,
Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F.,
Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, Q. J. R. Meteorol. Soc., 146, 1999–2049,
<https://doi.org/10.1002/qj.3803>, 2020.
- Horton, R. M., Mankin, J. S., Lesk, C., Coffel, E., and Raymond, C.: A Review of Recent Advances in Research
445 on Extreme Heat Events, Curr. Clim. Change Rep., 2, 242–259, <https://doi.org/10.1007/s40641-016-0042-x>, 2016.
- IPCC: Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment
Report of the Intergovernmental Panel on Climate Change, 1st ed., Cambridge University Press,
<https://doi.org/10.1017/9781009157896>, 2021.
- 450 IPCC: IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the
Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee
and J. Romero (eds.)]. IPCC, Geneva, Switzerland., Intergovernmental Panel on Climate Change (IPCC),
<https://doi.org/10.59327/IPCC/AR6-9789291691647>, 2023.
- Jézéquel, A., Yiou, P., and Radanovics, S.: Role of circulation in European heatwaves using flow analogues, Clim.
455 Dyn., 50, 1145–1159, <https://doi.org/10.1007/s00382-017-3667-0>, 2018.
- Kamae, Y., Watanabe, M., Kimoto, M., and Shiogama, H.: Summertime land–sea thermal contrast and atmospheric
circulation over East Asia in a warming climate—Part I: Past changes and future projections, Clim. Dyn.,
43, 2553–2568, <https://doi.org/10.1007/s00382-014-2073-0>, 2014.



- 460 Kjellstrom, T., Butler, A. J., Lucas, R. M., and Bonita, R.: Public health impact of global heating due to climate change: potential effects on chronic non-communicable diseases, *Int. J. Public Health*, 55, 97–103, <https://doi.org/10.1007/s00038-009-0090-2>, 2010.
- KNMI, Climate Explorer: Select a daily field: <https://climexp.knmi.nl/selectdailyfield2.cgi?id=someone@somewhere>, last access: 20 May 2024, 2022.
- 465 Kornhuber, K., Osprey, S., Coumou, D., Petri, S., Petoukhov, V., Rahmstorf, S., and Gray, L.: Extreme weather events in early summer 2018 connected by a recurrent hemispheric wave-7 pattern, *Environ. Res. Lett.*, 14, 054002, <https://doi.org/10.1088/1748-9326/ab13bf>, 2019.
- Kornhuber, K., Bartusek, S., Seager, R., Schellnhuber, H. J., and Ting, M.: Global emergence of regional heatwave hotspots outpaces climate model simulations, *Proc. Natl. Acad. Sci.*, 121, e2411258121, <https://doi.org/10.1073/pnas.2411258121>, 2024.
- 470 Liu, L. and Zhang, X.: Effects of temperature variability and extremes on spring phenology across the contiguous United States from 1982 to 2016, *Sci. Rep.*, 10, 17952, <https://doi.org/10.1038/s41598-020-74804-4>, 2020.
- Lopez, A., Fung, F., New, M., Watts, G., Weston, A., and Wilby, R. L.: From climate model ensembles to climate change impacts and adaptation: A case study of water resource management in the southwest of England, *Water Resour. Res.*, 45, 2008WR007499, <https://doi.org/10.1029/2008WR007499>, 2009.
- 475 Lorenz, R., Stalhandske, Z., and Fischer, E. M.: Detection of a Climate Change Signal in Extreme Heat, Heat Stress, and Cold in Europe From Observations, *Geophys. Res. Lett.*, 46, 8363–8374, <https://doi.org/10.1029/2019GL082062>, 2019.
- Noest, D.S.: Western_EU_warm_season_heat_extremes, GitHub [code], https://github.com/douwe3/Western_EU_warm_season_heat_extremes, 2025.
- 480 Patterson, M.: North-West Europe Hottest Days Are Warming Twice as Fast as Mean Summer Days, *Geophys. Res. Lett.*, 50, e2023GL102757, <https://doi.org/10.1029/2023GL102757>, 2023.
- Ridley, J., Menary, M., Kuhlbrodt, T., Andrews, M., and Andrews, T.: MOHC HadGEM3-GC31-LL model output prepared for CMIP6 CMIP historical (20230220), <https://doi.org/10.22033/ESGF/CMIP6.6109>, 2019.
- 485 Rousi, E., Kornhuber, K., Beobide-Arsuaga, G., Luo, F., and Coumou, D.: Accelerated western European heatwave trends linked to more-persistent double jets over Eurasia, *Nat. Commun.*, 13, 3851, <https://doi.org/10.1038/s41467-022-31432-y>, 2022.
- Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B., and Teuling, A. J.: Investigating soil moisture–climate interactions in a changing climate: A review, *Earth-Sci. Rev.*, 99, 125–161, <https://doi.org/10.1016/j.earscirev.2010.02.004>, 2010.
- 490 Shaw, T. A. and Voigt, A.: Tug of war on summertime circulation between radiative forcing and sea surface warming, *Nat. Geosci.*, 8, 560–566, <https://doi.org/10.1038/ngeo2449>, 2015.



- Shaw, T. A., Arias, P. A., Collins, M., Coumou, D., Diedhiou, A., Garfinkel, C. I., Jain, S., Roxy, M. K., Kretschmer, M., Leung, L. R., Narsey, S., Martius, O., Seager, R., Shepherd, T. G., Sörensson, A. A., Stephenson, T.,
495 Taylor, M., and Wang, L.: Regional climate change: consensus, discrepancies, and ways forward, *Front. Clim.*, 6, 1391634, <https://doi.org/10.3389/fclim.2024.1391634>, 2024.
- Shiogama, H., Tatebe, H., Hayashi, M., Abe, M., Arai, M., Koyama, H., Imada, Y., Kosaka, Y., Ogura, T., and Watanabe, M.: MIROC6 Large Ensemble (MIROC6-LE): experimental design and initial analyses, *Earth Syst. Dynam.*, 14, 1107–1124, <https://doi.org/10.5194/esd-2023-12>, 2023.
- 500 Smith, D. M., Gillett, N. P., Simpson, I. R., Athanasiadis, P. J., Baehr, J., Bethke, I., Bilge, T. A., Bonnet, R., Boucher, O., Findell, K. L., Gastineau, G., Gualdi, S., Hermanson, L., Leung, L. R., Mignot, J., Müller, W. A., Osprey, S., Otterå, O. H., Persad, G. G., Scaife, A. A., Schmidt, G. A., Shiogama, H., Sutton, R. T., Swingedouw, D., Yang, S., Zhou, T., and Ziehn, T.: Attribution of multi-annual to decadal changes in the climate system: The Large Ensemble Single Forcing Model Intercomparison Project (LESFMIP),
505 *Front. Clim.*, 4, 955414, <https://doi.org/10.3389/fclim.2022.955414>, 2022.
- Sousa, P. M., Barriopedro, D., García-Herrera, R., Ordóñez, C., Soares, P. M. M., and Trigo, R. M.: Distinct influences of large-scale circulation and regional feedbacks in two exceptional 2019 European heatwaves, *Commun. Earth Environ.*, 1, 48, <https://doi.org/10.1038/s43247-020-00048-9>, 2020.
- Sulikowska, A. and Wypych, A.: Seasonal Variability of Trends in Regional Hot and Warm Temperature Extremes
510 in Europe, *Atmosphere*, 12, 612, <https://doi.org/10.3390/atmos12050612>, 2021.
- Tatebe, H. and Watanabe, M.: MIROC MIROC6 model output prepared for CMIP6 CMIP historical (20230220), <https://doi.org/10.22033/ESGF/CMIP6.5603>, 2018.
- Tatebe, H., Ogura, T., Nitta, T., Komuro, Y., Ogochi, K., Takemura, T., Sudo, K., Sekiguchi, M., Abe, M., Saito, F., Chikira, M., Watanabe, S., Mori, M., Hirota, N., Kawatani, Y., Mochizuki, T., Yoshimura, K., Takata, K., O’ishi, R., Yamazaki, D., Suzuki, T., Kurogi, M., Kataoka, T., Watanabe, M., and Kimoto, M.:
515 Description and basic evaluation of simulated mean state, internal variability, and climate sensitivity in MIROC6, *Geosci. Model Dev.*, 12, 2727–2765, <https://doi.org/10.5194/gmd-12-2727-2019>, 2019.
- Thompson, V., Coumou, D., Galfi, V. M., Happé, T., Kew, S., Pinto, I., Philip, S., De Vries, H., and Van Der Wiel, K.: Changing dynamics of Western European summertime cut-off lows: A case study of the July 2021
520 flood event, *Atmospheric Sci. Lett.*, 25, e1260, <https://doi.org/10.1002/asl.1260>, 2024a.
- Thompson, V., Philip, S. Y., Pinto, I., and Kew, S. F.: The influence of the Atlantic Multidecadal Variability on Storm Babet-like events, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-1136>, 2024b.
- Vautard, R., Yiou, P., D’Andrea, F., De Noblet, N., Viovy, N., Cassou, C., Polcher, J., Ciais, P., Kageyama, M., and Fan, Y.: Summertime European heat and drought waves induced by wintertime Mediterranean rainfall
525 deficit, *Geophys. Res. Lett.*, 34, 2006GL028001, <https://doi.org/10.1029/2006GL028001>, 2007.
- Vautard, R., Van Aalst, M., Boucher, O., Drouin, A., Hausteijn, K., Kreienkamp, F., Van Oldenborgh, G. J., Otto, F. E. L., Ribes, A., Robin, Y., Schneider, M., Soubeyroux, J.-M., Stott, P., Seneviratne, S. I., Vogel, M. M.,



- and Wehner, M.: Human contribution to the record-breaking June and July 2019 heatwaves in Western Europe, *Environ. Res. Lett.*, 15, 094077, <https://doi.org/10.1088/1748-9326/aba3d4>, 2020.
- 530 Vautard, R., Cattiaux, J., Hap  , T., Singh, J., Bonnet, R., Cassou, C., Coumou, D., D'Andrea, F., Faranda, D., Fischer, E., Ribes, A., Sippel, S., and Yiou, P.: Heat extremes in Western Europe increasing faster than simulated due to atmospheric circulation trends, *Nat. Commun.*, 14, 6803, <https://doi.org/10.1038/s41467-023-42143-3>, 2023.
- 535 Whan, K., Zscheischler, J., Orth, R., Shongwe, M., Rahimi, M., Asare, E. O., and Seneviratne, S. I.: Impact of soil moisture on extreme maximum temperatures in Europe, *Weather Clim. Extrem.*, 9, 57–67, <https://doi.org/10.1016/j.wace.2015.05.001>, 2015.
- White, R. H., Kornhuber, K., Martius, O., and Wirth, V.: From Atmospheric Waves to Heatwaves: A Waveguide Perspective for Understanding and Predicting Concurrent, Persistent, and Extreme Extratropical Weather, *Bull. Am. Meteorol. Soc.*, 103, E923–E935, <https://doi.org/10.1175/BAMS-D-21-0170.1>, 2022.
- 540 Wu, L. and Zhang, J.: The relationship between spring soil moisture and summer hot extremes over North China, *Adv. Atmos. Sci.*, 32, 1660–1668, <https://doi.org/10.1007/s00376-015-5003-0>, 2015.
- Yang, J., Zhou, M., Ren, Z., Li, M., Wang, B., Liu, D. L., Ou, C.-Q., Yin, P., Sun, J., Tong, S., Wang, H., Zhang, C., Wang, J., Guo, Y., and Liu, Q.: Projecting heat-related excess mortality under climate change scenarios in China, *Nat. Commun.*, 12, 1039, <https://doi.org/10.1038/s41467-021-21305-1>, 2021.
- 545 Zampieri, M., D'Andrea, F., Vautard, R., Ciais, P., De Noblet-Ducoudr  , N., and Yiou, P.: Hot European Summers and the Role of Soil Moisture in the Propagation of Mediterranean Drought, *J. Clim.*, 22, 4747–4758, <https://doi.org/10.1175/2009JCLI2568.1>, 2009.